Preliminary Supporting Reference List for
2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD): Mode/Mechanism Of Action, Toxicokinetics, *In Vitro* Study Data, Modeling-Methodology

Abbott, B.D., A.R. Buckalew, M.J. DeVito, D. Ross, P.L. Bryant and J.E. Schmid. 2003. EGF and TGF-alpha expression influence the developmental toxicity of TCDD: dose response and AhR phenotype in EGF, TGF-alpha, and EGF + TGF-alpha knockout mice. Toxicol. Sci. 71(1):84-95.

Abbott, B.D., A.R. Buckalew and K.E. Leffler. 2005. Effects of epidermal growth factor (EGF), transforming growth factor-alpha (TGFalpha), and 2,3,7,8-tetrachlorodibenzo-p-dioxin on fusion of embryonic palates in serum-free organ culture using wild-type, EGF knockout, and TGFalpha knockout mouse strains. Birth Defects Res. A Clin. Mol. Teratol. 73(6):447-454.

Abdelrahim, M., R. Smith, III and S. Safe. 2003. Aryl hydrocarbon receptor gene silencing with small inhibitory RNA differentially modulates Ah-responsiveness in MCF-7 and HepG2 cancer cells. Mol. Pharmacol. 63(6):1373-1381.

Abe, Y., H. Sinozaki, T. Takagi, T. Minegishi, K. Kokame, K. Kangawa, M. Uesaka and K. Miyamoto. 2006. Identification of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-inducible genes in human amniotic epithelial cells. Reprod. Biol. Endocrinol. 4:27.

Aboutabl, M.E. and A.O. El-Kadi. 2007. Constitutive expression and inducibility of CYP1A1 in the H9c2 rat cardiomyoblast cells. Toxicol. In Vitro. 21(8):1686-1691.

Abrahams, V.M., J.E. Collins, C.R. Wira, M.W. Fanger and G.R. Yeaman. 2003. Inhibition of human polymorphonuclear cell oxidative burst by 17-beta-estradiol and 2,3,7,8-tetrachlorodibenzo-p-dioxin. Am. J. Reprod. Immunol. 50(6):463-472.

Adachi, J., Y. Mori, S. Matsui and T. Matsuda. 2004. Comparison of gene expression patterns between 2,3,7,8-tetrachlorodibenzo-p-dioxin and a natural arylhydrocarbon receptor ligand, indirubin. Toxicol. Sci. 80(1):161-169.

Ahmed, S., M. Shibazaki, T. Takeuchi and H. Kikuchi. 2005. Protein kinase Ctheta activity is involved in the 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced signal transduction pathway leading to apoptosis in L-MAT, a human lymphoblastic T-cell line. FEBS J. 272(4):903-915.

Ahn, N.S., H. Hu, J.S. Park, J.S. Park, J.S. Kim, S. An, G. Kong, O.I. Aruoma, Y.S. Lee and K.S. Kang. 2005. Molecular mechanisms of the 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced inverted U-shaped dose responsiveness in anchorage independent growth and cell proliferation of human breast epithelial cells with stem cell characteristics. Mutat. Res. 579(1-2):189-199.

Akahoshi, E., S. Yoshimura and M. Ishihara-Sugano. 2006. Over-expression of AhR (aryl hydrocarbon receptor) induces neural differentiation of Neuro2a cells: neurotoxicology study. Environ. Health. 5:24.

Akintobi, A.M., C.M. Villano and L.A. White. 2007. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) exposure of normal human dermal fibroblasts results in AhR-dependent and -independent changes in gene expression. Toxicol. Appl. Pharmacol. 220(1):9-17.

Al-Anizy, M., N.J. Horley, C.W. Kuo, L.C. Gillett, C.A. Laughton, D. Kendall, D.A. Barrett, T. Parker and D.R. Bell. 2006. Cytochrome P450 Cyp4x1 is a major P450 protein in mouse brain. FEBS J. 273(5):936-947.

Allan, L.L. and D.H. Sherr. 2005. Constitutive activation and environmental chemical induction of the aryl hydrocarbon receptor/transcription factor in activated human B lymphocytes. Mol. Pharmacol. 67(5):1740-1750.

Allgeier, S.H., T.M. Lin, C.M. Vezina, R.W. Moore, W.A. Fritz, S.Y. Chiu, C. Zhang and R.E. Peterson. 2008. WNT5A selectively inhibits mouse ventral prostate development. Dev. Biol. 324(1):10-17.

Allis, J.W., B.P. Anderson, G. Zhao, T.M. Ross and R.A. Pegram. 2002. Evidence for the involvement of CYP1A2 in the metabolism of bromodichloromethane in rat liver. Toxicology. 176(1-2):25-37.

Amakura, Y., T. Tsutsumi, K. Sasaki, T. Yoshida and T. Maitani. 2003. Screening of the inhibitory effect of vegetable constituents on the aryl hydrocarbon receptor-mediated activity induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin. Biol. Pharm. Bull. 26(12):1754-1760.

An, J., D.Y. Yang, Q.Z. Xu, S.M. Zhang, Y.Y. Huo, Z.F. Shang, Y. Wang, D.C. Wu and P.K. Zhou. 2008. DNA-dependent protein kinase catalytic subunit modulates the stability of c-Myc oncoprotein. Mol. Cancer. 7:32.

Andersen, M.E. 2002. The use of quantitative histological and molecular data for risk assessment and biologically based model development. Toxicol. Pathol. 30(1):106-111.

Andersen, M.E. and J.E. Dennison. 2001. Mode of action and tissue dosimetry in current and future risk assessments. Sci. Total Environ. 274(1-3):3-14.

Andersson, P., J. McGuire, C. Rubio, K. Gradin, M.L. Whitelaw, S. Pettersson, A. Hanberg and L. Poellinger. 2002. A constitutively active dioxin/aryl hydrocarbon receptor induces stomach tumors. Proc. Natl. Acad. Sci. U. S. A. 99(15):9990-9995.

Andersson, P., A. Ridderstad, J. McGuire, S. Pettersson, L. Poellinger and A. Hanberg. 2003. A constitutively active aryl hydrocarbon receptor causes loss of peritoneal B1 cells. Biochem. Biophys. Res. Commun. 302(2):336-341.

Andersson, P., C. Rubio, L. Poellinger and A. Hanberg. 2005. Gastric hamartomatous tumours in a transgenic mouse model expressing an activated dioxin/Ah receptor. Anticancer Res. 25(2A):903-911.

Anderson, G.R., A. Hasan, H. Yin, I. Qadri and L.C. Quattrochi. 2006. Regulation of the CYP1A1 gene by 2,3,7,8-tetrachlorodibenzo-p-dioxin but not by beta-naphthoflavone or 3-methylcholanthrene is altered in hepatitis C virus replicon-expressing cells. Mol. Pharmacol. 70(3):1062-1070.

Angus, W.G., M. Campaigne Larsen and C.R. Jefcoate. 2000. TCDD elevates erbB2 expression and signaling in T47D cells by reversing serum potentiation of estrogen receptor activity, independent of estrogen levels and enhanced ER down-regulation. Mol. Cell Endocrinol. 170(1-2):1-13.

Anwar-Mohamed, A. and A.O. El-Kadi. 2008. Down-regulation of the carcinogen-metabolizing enzyme cytochrome P450 1a1 by vanadium. Drug Metab Dispos. 36(9):1819-1827.

Aoki, Y. 2000. [Effects of dioxins on protein kinases and disruption of signal transduction pathways]. Nippon Rinsho. 58(12):2458-2463.

Aozasa, O., T. Tetsumi, S. Ohta, T. Nakao, H. Miyata and T. Nomura. 2003. Fecal excretion of dioxin in mice enhanced by intake of dietary fiber bearing chlorophyllin. Bull. Environ. Contam Toxicol. 70(2):359-366.

Arehart, E., G. Giasson, M.T. Walsh and H. Patterson. 2004. Dioxin alters the human low-density and very low-density lipoprotein structure with evidence for specific quenching of Trp-48 in apolipoprotein C-II. Biochemistry. 43(26):8503-8509.

Arpiainen, S., F. Raffalli-Mathieu, M.A. Lang, O. Pelkonen and J. Hakkola. 2005. Regulation of the Cyp2a5 gene involves an aryl hydrocarbon receptor-dependent pathway. Mol. Pharmacol. 67(4):1325-1333.

Arsenescu, V., R.I. Arsenescu, V. King, H. Swanson and L.A. Cassis. 2008. Polychlorinated biphenyl-77 induces adipocyte differentiation and proinflammatory adipokines and promotes obesity and atherosclerosis. Environ. Health Perspect. 116(6):761-768.

Ashida, H. 2000. Suppressive effects of flavonoids on dioxin toxicity. Biofactors. 12(1-4):201-206.

Augustowska, K., E.L. Gregoraszczuk, T. Milewicz, J. Krzysiek, A. Grochowalski and R. Chrzaszcz. 2003a. Effects of dioxin (2,3,7,8-TCDD) and PCDDs/PCDFs congeners mixture on steroidogenesis in human placenta tissue culture. Endocr. Regul. 37(1):11-19.

Augustowska, K., E.L. Gregoraszczuk, A. Grochowalski, T. Milewicz, M. Mika, J. Krzysiek and R. Chrzaszcz. 2003b. Comparison of accumulation and altered steroid secretion by placental tissue treated with TCDD and natural mixture of PCDDs-PCDFs. Reproduction. 126(5):681-687.

Augustowska, K., Z. Magnowska, M. Kapiszewska and E.L. Gregoraszczuk. 2007. Is the natural PCDD/PCDF mixture toxic for human placental JEG-3 cell line? The action of the toxicants on hormonal profile, CYP1A1 activity, DNA damage and cell apoptosis. Hum. Exp. Toxicol. 26(5):407-417.

Augereau, P., E. Badia, M. Fuentes, F. Rabenoelina, M. Corniou, D. Derocq, P. Balaguer and V. Cavailles. 2006. Transcriptional regulation of the human NRIP1/RIP140 gene by estrogen is modulated by dioxin signalling. Mol. Pharmacol. 69(4):1338-1346.

Aylward, L.L. and S.M. Hays. 2002. Temporal trends in human TCDD body burden: decreases over three decades and implications for exposure levels. J. Expo. Anal. Environ. Epidemiol. 12(5):319-328.

Aylward, L., C. Kirman, D. Cher and S. Hays. 2003a. Re: analysis of dioxin cancer threshold. Environ. Health Perspect. 111(10):A510.

Aylward, L.L., S.M. Hays, J.S. LaKind and J.J. Ryan. 2003b. Rapid communication: partitioning of persistent lipophilic compounds, including dioxins, between human milk lipid and blood lipid: an initial assessment. J. Toxicol. Environ. Health A. 66(1):1-5.

Aylward, L.L., J.C. Lamb and S.C. Lewis. 2005a. Issues in risk assessment for developmental effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin and related compounds. Toxicol. Sci. 87(1):3-10.

Aylward, L.L., R.C. Brunet, T.B. Starr, G. Carrier, E. Delzell, H. Cheng and C. Beall. 2005b. Exposure reconstruction for the TCDD-exposed NIOSH cohort using a concentration- and age-dependent model of elimination. Risk Anal. 25(4):945-956.

Aylward, L.L., J.S. LaKind and S.M. Hays. 2008a. Derivation of biomonitoring equivalent (BE) values for 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and related compounds: a screening tool for interpretation of biomonitoring data in a risk assessment context. J. Toxicol. Environ. Health A. 71(22):1499-1508.

Aylward, L.L., G. Charnley, J. Goodman and L. Rhomberg. 2008b. Comment on "Chronic disease and early exposure to air-borne mixtures. 2. Exposure assessment". Environ. Sci. Technol. 42(6):2201-2203.

Aylward, L.L., J.E. Goodman, G. Charnley and L.R. Rhomberg. 2008c. A margin-of-exposure approach to assessment of noncancer risks of dioxins based on human exposure and response data. Environ. Health Perspect. 116(10):1344-1351.

Baba, T., Y. Shima, A. Owaki, J. Mimura, M. Oshima, Y. Fujii-Kuriyama and K.I. Morohashi. 2008. Disruption of aryl hydrocarbon receptor (AhR) induces regression of the seminal vesicle in aged male mice. Sex Dev. 2(1):1-11.

Baccarelli, A., R. Pfeiffer, D. Consonni, A.C. Pesatori, M. Bonzini, D.G. Patterson, Jr., P.A. Bertazzi and M.T. Landi. 2005. Handling of dioxin measurement data in the presence of non-detectable values: overview of available methods and their application in the Seveso chloracne study. Chemosphere. 60(7):898-906.

Backlund, M. and M. Ingelman-Sundberg. 2004. Different structural requirements of the ligand binding domain of the aryl hydrocarbon receptor for high- and low-affinity ligand binding and receptor activation. Mol. Pharmacol. 65(2):416-425.

Backlund, M. and M. Ingelman-Sundberg. 2005. Regulation of aryl hydrocarbon receptor signal transduction by protein tyrosine kinases. Cell Signal. 17(1):39-48.

Bagchi, D., J. Balmoori, M. Bagchi, X. Ye, C.B. Williams and S.J. Stohs. 2000. Role of p53 tumor suppressor gene in the toxicity of TCDD, endrin, naphthalene, and chromium (VI) in liver and brain tissues of mice. Free Radic. Biol. Med. 28(6):895-903.

Bajanowski, T., P. Furst, K. Wilmers, J. Beike, H. Kohler and B. Brinkmann. 2002. Dioxin in infants--an environmental hazard? Int. J. Legal Med. 116(1):27-32.

Baldridge, M.G. and R.J. Hutz. 2007. Autoradiographic localization of aromatic hydrocarbon receptor (AHR) in rhesus monkey ovary. Am. J. Primatol. 69(6):681-691.

Bapiro, T.E., T.B. Andersson, C. Otter, J.A. Hasler and C.M. Masimirembwa. 2002. Cytochrome P450 1A1/2 induction by antiparasitic drugs: dose-dependent increase in ethoxyresorufin O-deethylase activity and mRNA caused by quinine, primaquine and albendazole in HepG2 cells. Eur. J. Clin. Pharmacol. 58(8):537-542.

Barnes-Ellerbe, S., K.E. Knudsen and A. Puga. 2004. 2,3,7,8-Tetrachlorodibenzo-p-dioxin blocks androgen-dependent cell proliferation of LNCaP cells through modulation of pRB phosphorylation. Mol. Pharmacol. 66(3):502-511.

Barouki, R. 2003. [A fatal kiss between estrogen receptor and dioxin]. Med. Sci. (Paris). 19(11):1068-1069.

Barouki, R. and Y. Morel. 2001. Repression of cytochrome P450 1A1 gene expression by oxidative stress: mechanisms and biological implications. Biochem. Pharmacol. 61(5):511-516.

Barrow, L.L., M.E. Wines, P.A. Romitti, B.C. Holdener and J.C. Murray. 2002. Aryl hydrocarbon receptor nuclear translocator 2 (ARNT2): structure, gene mapping, polymorphisms, and candidate evaluation for human orofacial clefts. Teratology. 66(2):85-90.

Baumgart, A., M. Schmidt, H.J. Schmitz and D. Schrenk. 2005. Natural furocoumarins as inducers and inhibitors of cytochrome P450 1A1 in rat hepatocytes. Biochem. Pharmacol. 69(4):657-667.

Beischlag, T.V. and G.H. Perdew. 2005. ER alpha-AHR-ARNT protein-protein interactions mediate estradiol-dependent transrepression of dioxin-inducible gene transcription. J. Biol. Chem. 280(22):21607-21611.

Beischlag, T.V., S. Wang, D.W. Rose, J. Torchia, S. Reisz-Porszasz, K. Muhammad, W.E. Nelson, M.R. Probst, M.G. Rosenfeld and O. Hankinson. 2002. Recruitment of the NCoA/SRC-1/p160 family of transcriptional coactivators by the aryl hydrocarbon receptor/aryl hydrocarbon receptor nuclear translocator complex. Mol. Cell Biol. 22(12):4319-4333.

Beischlag, T.V., R.T. Taylor, D.W. Rose, D. Yoon, Y. Chen, W.H. Lee, M.G. Rosenfeld and O. Hankinson. 2004. Recruitment of thyroid hormone receptor/retinoblastoma-interacting protein 230 by the aryl hydrocarbon receptor nuclear translocator is required for the transcriptional response to both dioxin and hypoxia. J. Biol. Chem. 279(52):54620-54628.

Beischlag, T.V., M.J. Luis, B.D. Hollingshead and G.H. Perdew. 2008. The aryl hydrocarbon receptor complex and the control of gene expression. Crit Rev. Eukaryot. Gene Expr. 18(3):207-250.

Belpomme, D., P. Irigaray, L. Hardell, R. Clapp, L. Montagnier, S. Epstein and A.J. Sasco. 2007. The multitude and diversity of environmental carcinogens. Environ. Res. 105(3):414-429.

Bemis, J.C., D.A. Nazarenko and T.A. Gasiewicz. 2005. Coplanar polychlorinated biphenyls activate the aryl hydrocarbon receptor in developing tissues of two TCDD-responsive lacZ mouse lines. Toxicol. Sci. 87(2):529-536.

Berg, P. and I. Pongratz. 2002. Two parallel pathways mediate cytoplasmic localization of the dioxin (aryl hydrocarbon) receptor. J. Biol. Chem. 277(35):32310-32319.

Berge, G., S. Mollerup, S. OVrebo, A. Hewer, D.H. Phillips, E. Eilertsen and A. Haugen. 2004. Role of estrogen receptor in regulation of polycyclic aromatic hydrocarbon metabolic activation in lung. Lung Cancer. 45(3):289-297.

Bermudez de Leon, M., P. Gomez, G. Elizondo, R. Zatarain-Palacios, F. Garcia-Sierra and B. Cisneros. 2006. Beta-naphthoflavone represses dystrophin Dp71 expression in hepatic cells. Biochim. Biophys. Acta. 1759(3-4):152-158.

Bhathena, A., C. Lee and D.S. Riddick. 2002. Suppression of cytochrome P450 2C11 by aromatic hydrocarbons: mechanistic insights from studies of the 5'-flanking region of the CYP2C11 gene. Drug Metab Dispos. 30(12):1385-1392.

Bielefeld, K.A., C. Lee and D.S. Riddick. 2008. Regulation of aryl hydrocarbon receptor expression and function by glucocorticoids in mouse hepatoma cells. Drug Metab Dispos. 36(3):543-551.

Biswas, G., S. Srinivasan, H.K. Anandatheerthavarada and N.G. Avadhani. 2008. Dioxin-mediated tumor progression through activation of mitochondria-to-nucleus stress signaling. Proc. Natl. Acad. Sci. U. S. A. 105(1):186-191.

Black, V.H. and L.C. Quattrochi. 2004. Molecular cloning of the guinea pig CYP1A2 gene 5'-flanking region: identification of functional aromatic hydrocarbon response element and characterization of CYP1A2 expression in GPC16 cells. Drug Metab Dispos. 32(6):595-602.

Bock, K.W. and C. Kohle. 2005. Ah receptor- and TCDD-mediated liver tumor promotion: clonal selection and expansion of cells evading growth arrest and apoptosis. Biochem. Pharmacol. 69(10):1403-1408.

Bock, K.W. and C. Kohle. 2006. Ah receptor: dioxin-mediated toxic responses as hints to deregulated physiologic functions. Biochem. Pharmacol. 72(4):393-404.

Bock, K.W., T. Eckle, M. Ouzzine and S. Fournel-Gigleux. 2000a. Coordinate induction by antioxidants of UDP-glucuronosyltransferase UGT1A6 and the apical conjugate export pump MRP2 (multidrug resistance protein 2) in Caco-2 cells. Biochem. Pharmacol. 59(5):467-470.

Bock, K.W., H. Gschaidmeier, B.S. Bock-Hennig and L.C. Eriksson. 2000b. Density-dependent growth of normal and nodular hepatocytes. Toxicology. 144(1-3):51-56.

Boffetta, P., J.K. McLaughlin, C. La Vecchia, R.E. Tarone, L. Lipworth and W.J. Blot. 2008. False-positive results in cancer epidemiology: a plea for epistemological modesty. J. Natl. Cancer Inst. 100(14):988-995.

Bofinger, D.P., L. Feng, L.H. Chi, J. Love, F.D. Stephen, T.R. Sutter, K.G. Osteen, T.G. Costich, R.E. Batt, S.T. Koury and J.R. Olson. 2001. Effect of TCDD exposure on CYP1A1 and CYP1B1 expression in explant cultures of human endometrium. Toxicol. Sci. 62(2):299-314.

Bohonowych, J.E. and M.S. Denison. 2007. Persistent binding of ligands to the aryl hydrocarbon receptor. Toxicol. Sci. 98(1):99-109.

Bonzo, J.A., S. Chen, A. Galijatovic and R.H. Tukey. 2005. Arsenite inhibition of CYP1A1 induction by 2,3,7,8-tetrachlorodibenzo-p-dioxin is independent of cell cycle arrest. Mol. Pharmacol. 67(4):1247-1256.

Borak, J. and G. Sirianni. 2005. Hormesis: implications for cancer risk assessment. Dose. Response. 3(3):443-451.

Borlak, J. and H.S. Jenke. 2008. Cross-talk between aryl hydrocarbon receptor and mitogen-activated protein kinase signaling pathway in liver cancer through c-raf transcriptional regulation. Mol. Cancer Res. 6(8):1326-1336.

Bortot, P., K. Thomaseth and A. Salvan. 2002. Population toxicokinetic analysis of 2,3,7,8-tetrachlorodibenzo-p-dioxin using Bayesian techniques. Stat. Med. 21(4):533-547.

Boutros, P.C., I.D. Moffat, M.A. Franc, N. Tijet, J. Tuomisto, R. Pohjanvirta and A.B. Okey. 2004. Dioxin-responsive AHRE-II gene battery: identification by phylogenetic footprinting. Biochem. Biophys. Res. Commun. 321(3):707-715.

Boutros, P.C., R. Yan, I.D. Moffat, R. Pohjanvirta and A.B. Okey. 2008. Transcriptomic responses to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in liver: comparison of rat and mouse. BMC. Genomics. 9:419.

Boverhof, D.R., E. Tam, A.S. Harney, R.B. Crawford, N.E. Kaminski and T.R. Zacharewski. 2004. 2,3,7,8-Tetrachlorodibenzo-p-dioxin induces suppressor of cytokine signaling 2 in murine B cells. Mol. Pharmacol. 66(6):1662-1670.

Boverhof, D.R., L.D. Burgoon, C. Tashiro, B. Sharratt, B. Chittim, J.R. Harkema, D.L. Mendrick and T.R. Zacharewski. 2006a. Comparative toxicogenomic analysis of the hepatotoxic effects of TCDD in Sprague Dawley rats and C57BL/6 mice. Toxicol. Sci. 94(2):398-416.

Boverhof, D.R., J.C. Kwekel, D.G. Humes, L.D. Burgoon and T.R. Zacharewski. 2006b. Dioxin induces an estrogen-like, estrogen receptor-dependent gene expression response in the murine uterus. Mol. Pharmacol. 69(5):1599-1606.

Bradley, C., R. van der Meer, N. Roodi, H. Yan, M.B. Chandrasekharan, Z.W. Sun, R.L. Mernaugh and F.F. Parl. 2007. Carcinogen-induced histone alteration in normal human mammary epithelial cells. Carcinogenesis. 28(10):2184-2192.

Brauze, D. and D. Malejka-Giganti. 2000. A novel 4 S [3H]beta-naphthoflavone-binding protein in liver cytosol of female Sprague-Dawley rats treated with aryl hydrocarbon receptor agonists. Biochem. J. 347 Pt 3:787-795.

Brauze, D., M. Widerak, J. Cwykiel, K. Szyfter and W. Baer-Dubowska. 2006. The effect of aryl hydrocarbon receptor ligands on the expression of AhR, AhRR, ARNT, Hif1alpha, CYP1A1 and NQO1 genes in rat liver. Toxicol. Lett. 167(3):212-220.

Bredhult, C., B.M. Backlin and M. Olovsson. 2007. Effects of some endocrine disruptors on the proliferation and viability of human endometrial endothelial cells in vitro. Reprod. Toxicol. 23(4):550-559.

Brosens, J.J. and M.G. Parker. 2003. Gene expression: Oestrogen receptor hijacked. Nature. 423(6939):487-488.

Bruno, M.E., C.H. Borchers, J.M. Dial, N.J. Walker, J.E. Hartis, B.A. Wetmore, J.C. Barrett, K.B. Tomer and B.A. Merrick. 2002. Effects of TCDD upon IkappaB and IKK subunits localized in microsomes by proteomics. Arch. Biochem. Biophys. 406(2):153-164.

Bryant, P.L., L.M. Reid, J.E. Schmid, A.R. Buckalew and B.D. Abbott. 2001. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on fetal mouse urinary tract epithelium in vitro. Toxicology. 162(1):23-34.

Bukowska, B. 2004. Damage to erythrocytes caused by 2,3,7,8-tetrachloro-dibenzo-p-dioxin (in vitro). Cell Mol. Biol. Lett. 9(2):261-270.

Bulun, S.E., K.M. Zeitoun and G. Kilic. 2000. Expression of dioxin-related transactivating factors and target genes in human eutopic endometrial and endometriotic tissues. Am. J. Obstet. Gynecol. 182(4):767-775.

Bunger, M.K., S.M. Moran, E. Glover, T.L. Thomae, G.P. Lahvis, B.C. Lin and C.A. Bradfield. 2003. Resistance to 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity and abnormal liver development in mice carrying a mutation in the nuclear localization sequence of the aryl hydrocarbon receptor. J. Biol. Chem. 278(20):17767-17774.

Bunger, M.K., E. Glover, S.M. Moran, J.A. Walisser, G.P. Lahvis, E.L. Hsu and C.A. Bradfield. 2008. Abnormal liver development and resistance to 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity in mice carrying a mutation in the DNA-binding domain of the aryl hydrocarbon receptor. Toxicol. Sci. 106(1):83-92.

Burgoon, L.D. and T.R. Zacharewski. 2008. Automated quantitative dose-response modeling and point of departure determination for large toxicogenomic and high-throughput screening data sets. Toxicol. Sci. 104(2):412-418.

Butler, D. 2003. Agent Orange health investigation stuck at square one. Nature. 422(6934):793.

Butler, D. 2008. Further delays to full Agent Orange study. Nature. 452(7189):786-787.

Cacciari, P. 2007. [Dioxins and sarcomas in Veneto: the risk of confusing the citizens]. Epidemiol. Prev. 31(2-3):74-75.

Camacho, I.A., M. Nagarkatti and P.S. Nagarkatti. 2002. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) induces Fas-dependent activation-induced cell death in superantigen-primed T cells. Arch. Toxicol. 76(10):570-580.

Camacho, I.A., N. Singh, V.L. Hegde, M. Nagarkatti and P.S. Nagarkatti. 2005. Treatment of mice with 2,3,7,8-tetrachlorodibenzo-p-dioxin leads to aryl hydrocarbon receptor-dependent nuclear translocation of NF-kappaB and expression of Fas ligand in thymic stromal cells and consequent apoptosis in T cells. J. Immunol. 175(1):90-103.

Canistro, D., G. Cantelli-Forti, G.L. Biagi and M. Paolini. 2002. Re: dioxin increases reactive oxygen production in mouse liver mitochondria (Toxicol. Appl. Pharmacol.178, 15-21, 2002). To the editor. Toxicol. Appl. Pharmacol. 185(1):74-75.

Carpenter, D.O. 2008. Environmental contaminants as risk factors for developing diabetes. Rev. Environ. Health. 23(1):59-74.

Caudill, S.P., L.Y. Wong, W.E. Turner, R. Lee, A. Henderson and D.G. Patterson, Jr. 2007. Percentile estimation using variable censored data. Chemosphere. 68(1):169-180.

Celius, T., S. Roblin, P.A. Harper, J. Matthews, P.C. Boutros, R. Pohjanvirta and A.B. Okey. 2008. ARYL HYDROCARBON RECEPTOR (AHR)-DEPENDENT INDUCTION OF FLAVIN-CONTAINING MONOOXYGENASE mRNAs IN MOUSE LIVER. Drug Metab Dispos.

Chan, C.Y., P.M. Kim and L.M. Winn. 2004a. TCDD-induced homologous recombination: the role of the Ah receptor versus oxidative DNA damage. Mutat. Res. 563(1):71-79.

Chan, C.Y., P.M. Kim and L.M. Winn. 2004b. TCDD affects DNA double strand-break repair. Toxicol. Sci. 81(1):133-138.

Chang, S.F., Y.Y. Sun, L.Y. Yang, S.Y. Hu, S.Y. Tsai, W.S. Lee and Y.H. Lee. 2005. Bcl-2 gene family expression in the brain of rat offspring after gestational and lactational dioxin exposure. Ann. N. Y. Acad. Sci. 1042:471-480.

Chang, H., L.W. Chang, Y.H. Cheng, W.T. Tsai, M.X. Tsai and P. Lin. 2006. Preferential induction of CYP1A1 and CYP1B1 in CCSP-positive cells. Toxicol. Sci. 89(1):205-213.

Chang, J.T., H. Chang, P.H. Chen, S.L. Lin and P. Lin. 2007. Requirement of aryl hydrocarbon receptor overexpression for CYP1B1 up-regulation and cell growth in human lung adenocarcinomas. Clin. Cancer Res. 13(1):38-45.

Chao, H.R., T.C. Tsou, L.A. Li, F.Y. Tsai, Y.F. Wang, C.H. Tsai, E.E. Chang, Z.F. Miao, C.H. Wu and W.J. Lee. 2006. Arsenic inhibits induction of cytochrome P450 1A1 by 2,3,7,8-tetrachlorodibenzo-p-dioxin in human hepatoma cells. J. Hazard. Mater. 137(2):716-722.

Chaudhary, A. and K.L. Willett. 2006. Inhibition of human cytochrome CYP 1 enzymes by flavonoids of St. John's wort. Toxicology. 217(2-3):194-205.

Chen, G. and N.J. Bunce. 2003. Polybrominated diphenyl ethers as Ah receptor agonists and antagonists. Toxicol. Sci. 76(2):310-320.

Chen, G. and N.J. Bunce. 2004. Interaction between halogenated aromatic compounds in the Ah receptor signal transduction pathway. Environ. Toxicol. 19(5):480-489.

Chen, I., T. Hsieh, T. Thomas and S. Safe. 2001. Identification of estrogen-induced genes downregulated by AhR agonists in MCF-7 breast cancer cells using suppression subtractive hybridization. Gene. 262(1-2):207-214.

Chen, J., T.L. Thirkill, J.W. Overstreet, B.L. Lasley and G.C. Douglas. 2003a. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on chorionic gonadotropin secretion by human trophoblasts. Reprod. Toxicol. 17(1):87-93.

Chen, J.J., G.S. Chen and N.J. Bunce. 2003b. Inhibition of CYP 1A2-dependent MROD activity in rat liver microsomes: an explanation of the hepatic sequestration of a limited subset of halogenated aromatic hydrocarbons. Environ. Toxicol. 18(2):115-119.

Chen, Z.H., Y.J. Hurh, H.K. Na, J.H. Kim, Y.J. Chun, D.H. Kim, K.S. Kang, M.H. Cho and Y.J. Surh. 2004. Resveratrol inhibits TCDD-induced expression of CYP1A1 and CYP1B1 and catechol estrogen-mediated oxidative DNA damage in cultured human mammary epithelial cells. Carcinogenesis. 25(10):2005-2013.

Chen, S., T. Operana, J. Bonzo, N. Nguyen and R.H. Tukey. 2005a. ERK kinase inhibition stabilizes the aryl hydrocarbon receptor: implications for transcriptional activation and protein degradation. J. Biol. Chem. 280(6):4350-4359.

Chen, S., D. Beaton, N. Nguyen, K. Senekeo-Effenberger, E. Brace-Sinnokrak, U. Argikar, R.P. Remmel, J. Trottier, O. Barbier, J.K. Ritter and R.H. Tukey. 2005b. Tissue-specific, inducible, and hormonal control of the human UDP-glucuronosyltransferase-1 (UGT1) locus. J. Biol. Chem. 280(45):37547-37557.

Chen, J.W., S.L. Wang, P.C. Liao, H.Y. Chen, Y.C. Ko and C.C. Lee. 2008. Relationship between insulin sensitivity and exposure to dioxins and polychlorinated biphenyls in pregnant women. Environ. Res. 107(2):245-253.

Cheng, S.B., S. Kuchiiwa, X.Q. Ren, H.Z. Gao, T. Kuchiiwa and S. Nakagawa. 2003a. Dioxin exposure down-regulates nitric oxide synthase and NADPH-diaphorase activities in the hypothalamus of Long-Evans rat. Neurosci. Lett. 345(1):5-8.

Cheng, S.B., S. Kuchiiwa, A. Kawachi, H.Z. Gao, A. Gohshi, T. Kozako, T. Kuchiiwa and S. Nakagawa. 2003b. Up-regulation of methionine-enkephalin-like immunoreactivity by 2,3,7,8-tetrachlorodibenzo-p-dioxin treatment in the forebrain of the Long-Evans rat. J. Chem. Neuroanat. 25(2):73-82.

Cheng, X., J. Maher, M.Z. Dieter and C.D. Klaassen. 2005. Regulation of mouse organic anion-transporting polypeptides (Oatps) in liver by prototypical microsomal enzyme inducers that activate distinct transcription factor pathways. Drug Metab Dispos. 33(9):1276-1282.

Cheng, H., L. Aylward, C. Beall, T.B. Starr, R.C. Brunet, G. Carrier and E. Delzell. 2006. TCDD exposure-response analysis and risk assessment. Risk Anal. 26(4):1059-1071.

Cheng, Y., L.W. Chang, L.C. Cheng, M.H. Tsai and P. Lin. 2007. 4-Methoxyestradiol-induced oxidative injuries in human lung epithelial cells. Toxicol. Appl. Pharmacol. 220(3):271-277.

Cheon, H., Y.S. Woo, J.Y. Lee, H.S. Kim, H.J. Kim, S. Cho, N.H. Won and J. Sohn. 2007. Signaling pathway for 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced TNF-alpha production in differentiated THP-1 human macrophages. Exp. Mol. Med. 39(4):524-534.

Chiaro, C.R., R.D. Patel, C.B. Marcus and G.H. Perdew. 2007. Evidence for an aryl hydrocarbon receptor-mediated cytochrome p450 autoregulatory pathway. Mol. Pharmacol. 72(5):1369-1379.

Cho, J.S., M.S. Chang and H.M. Rho. 2001. Transcriptional activation of the human Cu/Zn superoxide dismutase gene by 2,3,7,8-tetrachlorodibenzo-p-dioxin through the xenobiotic-responsive element. Mol. Genet. Genomics. 266(1):133-141.

Cho, S.J., J.S. Jung, I. Jin, Y.W. Jung, B.H. Ko, K.S. Nam, I.K. Park and I.S. Moon. 2002. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the expression of synaptic proteins in dissociated rat cortical cells. Mol. Cells. 14(2):238-244.

Cho, Y.C., W. Zheng and C.R. Jefcoate. 2004. Disruption of cell-cell contact maximally but transiently activates AhR-mediated transcription in 10T1/2 fibroblasts. Toxicol. Appl. Pharmacol. 199(3):220-238.

Cho, Y.C., W. Zheng, M. Yamamoto, X. Liu, P.R. Hanlon and C.R. Jefcoate. 2005. Differentiation of pluripotent C3H10T1/2 cells rapidly elevates CYP1B1 through a novel process that overcomes a loss of Ah Receptor. Arch. Biochem. Biophys. 439(2):139-153.

Choi, J.Y., J.A. Oughton and N.I. Kerkvliet. 2003. Functional alterations in CD11b(+)Gr-1(+) cells in mice injected with allogeneic tumor cells and treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin. Int. Immunopharmacol. 3(4):553-570.

Chramostova, K., J. Vondracek, L. Sindlerova, B. Vojtesek, A. Kozubik and M. Machala. 2004. Polycyclic aromatic hydrocarbons modulate cell proliferation in rat hepatic epithelial stem-like WB-F344 cells. Toxicol. Appl. Pharmacol. 196(1):136-148.

Chun, H.S., P.A. Kuzmicky, L. Rucoba, N.Y. Kado and R.H. Rice. 2000. Cytotoxicity and keratinocyte microsome-mediated mutagenic activation of carcinogens in cultured epidermal cells. Toxicol. Lett. 115(2):165-172.

Chun, Y.J., S.K. Lee and M.Y. Kim. 2005. Modulation of human cytochrome P450 1B1 expression by 2,4,3',5'-tetramethoxystilbene. Drug Metab Dispos. 33(12):1771-1776.

Cimafranca, M.A., P.R. Hanlon and C.R. Jefcoate. 2004. TCDD administration after the pro-adipogenic differentiation stimulus inhibits PPARgamma through a MEK-dependent process but less effectively suppresses adipogenesis. Toxicol. Appl. Pharmacol. 196(1):156-168.

Ciolino, H.P., M. Dankwah and G.C. Yeh. 2002. Resistance of MCF-7 cells to dimethylbenz(a)anthracene-induced apoptosis is due to reduced CYP1A1 expression. Int. J. Oncol. 21(2):385-391.

Collier, A.C., K.L. Pritsos and C.A. Pritsos. 2006. TCDD as a biological response modifier for Mitomycin C: oxygen tension affects enzyme activation, reactive oxygen species and cell death. Life Sci. 78(13):1499-1507.

Collins, J.J., K. Bodner, C.J. Burns, R.A. Budinsky, L.L. Lamparski, M. Wilken, G.D. Martin and M.L. Carson. 2007. Body mass index and serum chlorinated dibenzo-p-dioxin and dibenzofuran levels. Chemosphere. 66(6):1079-1085.

Cooney, C.M. 2001. How serious is dioxin's cancer risk? Environ. Sci. Technol. 35(9):180A-181A.

Coumoul, X., M. Diry, C. Robillot and R. Barouki. 2001. Differential regulation of cytochrome P450 1A1 and 1B1 by a combination of dioxin and pesticides in the breast tumor cell line MCF-7. Cancer Res. 61(10):3942-3948.

Crawford, R.B., C.E. Sulentic, B.S. Yoo and N.E. Kaminski. 2003. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) alters the regulation and posttranslational modification of p27kip1 in lipopolysaccharide-activated B cells. Toxicol. Sci. 75(2):333-342.

Cunha, G.R., P.S. Cooke and T. Kurita. 2004. Role of stromal-epithelial interactions in hormonal responses. Arch. Histol. Cytol. 67(5):417-434.

Dale, Y.R. and S.E. Eltom. 2006. Calpain mediates the dioxin-induced activation and down-regulation of the aryl hydrocarbon receptor. Mol. Pharmacol. 70(5):1481-1487.

Dasmahapatra, A.K., B.A. Wimpee, A.L. Trewin, C.F. Wimpee, J.K. Ghorai and R.J. Hutz. 2000. Demonstration of 2,3,7,8-tetrachlorodibenzo-p-dioxin attenuation of P450 steroidogenic enzyme mRNAs in rat granulosa cell in vitro by competitive reverse transcriptase-polymerase chain reaction assay. Mol. Cell Endocrinol. 164(1-2):5-18.

Dasmahapatra, A.K., B.A. Wimpee, A.L. Trewin and R.J. Hutz. 2001. 2,3,7,8-Tetrachlorodibenzo-p-dioxin increases steady-state estrogen receptor-beta mRNA levels after CYP1A1 and CYP1B1 induction in rat granulosa cells in vitro. Mol. Cell Endocrinol. 182(1):39-48.

Dasmahapatra, A.K., A.L. Trewin and R.J. Hutz. 2002. Estrous cycle-regulated expression of CYP1B1 mRNA in the rat ovary. Comp Biochem. Physiol B Biochem. Mol. Biol. 133(1):127-134.

Davis, J.W., K. Melendez, V.M. Salas, F.T. Lauer and S.W. Burchiel. 2000. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) inhibits growth factor withdrawal-induced apoptosis in the human mammary epithelial cell line, MCF-10A. Carcinogenesis. 21(5):881-886.

Davis, J.W., F.T. Lauer, A.D. Burdick, L.G. Hudson and S.W. Burchiel. 2001. Prevention of apoptosis by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the MCF-10A cell line: correlation with increased transforming growth factor alpha production. Cancer Res. 61(8):3314-3320.

Davis, J.W., Jr., A.D. Burdick, F.T. Lauer and S.W. Burchiel. 2003. The aryl hydrocarbon receptor antagonist, 3'methoxy-4'nitroflavone, attenuates 2,3,7,8-tetrachlorodibenzo-p-dioxin-dependent regulation of growth factor signaling and apoptosis in the MCF-10A cell line. Toxicol. Appl. Pharmacol. 188(1):42-49.

de Oliveira, S.K., M. Hoffmeister, S. Gambaryan, W. Muller-Esterl, J.A. Guimaraes and A.P. Smolenski. 2007. Phosphodiesterase 2A forms a complex with the co-chaperone XAP2 and regulates nuclear translocation of the aryl hydrocarbon receptor. J. Biol. Chem. 282(18):13656-13663.

de Waard, P.W., A.A. Peijnenburg, H. Baykus, J.M. Aarts, R.L. Hoogenboom, F.J. van Schooten and T.M. de Kok. 2008a. A human intervention study with foods containing natural Ah-receptor agonists does not significantly show AhR-mediated effects as measured in blood cells and urine. Chem. Biol. Interact. 176(1):19-29.

de Waard, P.W., T.M. de Kok, L.M. Maas, A.A. Peijnenburg, R.L. Hoogenboom, J.M. Aarts and F.J. van Schooten. 2008b. Influence of TCDD and natural Ah receptor agonists on benzo[a]pyrene-DNA adduct formation in the Caco-2 human colon cell line. Mutagenesis. 23(1):67-73.

de Waard, W.J., J.M. Aarts, A.A. Peijnenburg, H. Baykus, E. Talsma, A. Punt, T.M. de Kok, F.J. van Schooten and L.A. Hoogenboom. 2008c. Gene expression profiling in Caco-2 human colon cells exposed to TCDD, benzo[a]pyrene, and natural Ah receptor agonists from cruciferous vegetables and citrus fruits. Toxicol. In Vitro. 22(2):396-410.

Degner, S.C., M.Q. Kemp, J.K. Hockings and D.F. Romagnolo. 2007. Cyclooxygenase-2 promoter activation by the aromatic hydrocarbon receptor in breast cancer mcf-7 cells: repressive effects of conjugated linoleic acid. Nutr. Cancer. 59(2):248-257.

Dere, E., D.R. Boverhof, L.D. Burgoon and T.R. Zacharewski. 2006. In vivo-in vitro toxicogenomic comparison of TCDD-elicited gene expression in Hepa1c1c7 mouse hepatoma cells and C57BL/6 hepatic tissue. BMC. Genomics. 7:80.

Dietrich, C., D. Faust, S. Budt, M. Moskwa, A. Kunz, K.W. Bock and F. Oesch. 2002. 2,3,7,8-tetrachlorodibenzo-p-dioxin-dependent release from contact inhibition in WB-F344 cells: involvement of cyclin A. Toxicol. Appl. Pharmacol. 183(2):117-126.

Dietrich, C., D. Faust, M. Moskwa, A. Kunz, K.W. Bock and F. Oesch. 2003. TCDD-dependent downregulation of gamma-catenin in rat liver epithelial cells (WB-F344). Int. J. Cancer. 103(4):435-439.

Diry, M., C. Tomkiewicz, C. Koehle, X. Coumoul, K.W. Bock, R. Barouki and C. Transy. 2006. Activation of the dioxin/aryl hydrocarbon receptor (AhR) modulates cell plasticity through a JNK-dependent mechanism. Oncogene. 25(40):5570-5574.

Dixit, R., J. Riviere, K. Krishnan and M.E. Andersen. 2003. Toxicokinetics and physiologically based toxicokinetics in toxicology and risk assessment. J. Toxicol. Environ. Health B Crit Rev. 6(1):1-40.

Doi, H., T. Baba, C. Tohyama and K. Nohara. 2003. Functional activation of arylhydrocarbon receptor (AhR) in primary T cells by 2,3,7,8-tetrachlorodibenzo-p-dioxin. Chemosphere. 52(4):655-662.

Dong, B. and F. Matsumura. 2008. Roles of cytosolic phospholipase A2 and Src kinase in the early action of 2,3,7,8-tetrachlorodibenzo-p-dioxin through a nongenomic pathway in MCF10A cells. Mol. Pharmacol. 74(1):255-263.

Dong, L., N.J. Tang and L.J. Zhao. 2008. [Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on epidermal cells proliferation in vitro]. Zhonghua Lao. Dong. Wei Sheng Zhi. Ye. Bing. Za Zhi. 26(3):178-180.

Drukteinis, J.S., T. Medrano, E.A. Ablordeppey, J.M. Kitzman and K.T. Shiverick. 2005. Benzo[a]pyrene, but not 2,3,7,8-TCDD, induces G2/M cell cycle arrest, p21CIP1 and p53 phosphorylation in human choriocarcinoma JEG-3 cells: a distinct signaling pathway. Placenta. 26 Suppl A:S87-S95.

Du, L., M.M. Neis, P.A. Ladd and D.S. Keeney. 2006. Differentiation-specific factors modulate epidermal CYP1-4 gene expression in human skin in response to retinoic acid and classic aryl hydrocarbon receptor ligands. J. Pharmacol. Exp. Ther. 319(3):1162-1171.

Dvorak, Z., I. Sovadinova, L. Blaha, J.P. Giesy and J. Ulrichova. 2006a. Quaternary benzo[c]phenathridine alkaloids sanguinarine and chelerythrine do not affect transcriptional activity of aryl hydrocarbon receptor: analyses in rat hepatoma cell line H4IIE.luc. Food Chem. Toxicol. 44(9):1466-1473.

Dvorak, Z., R. Vrzal, J. Ulrichova, J.M. Pascussi, P. Maurel and M. Modriansky. 2006b. Involvement of cytoskeleton in AhR-dependent CYP1A1 expression. Curr. Drug Metab. 7(3):301-313.

Dvorak, Z., A. Zdarilova, L. Sperlikova, E. Anzenbacherova, V. Simanek and J. Ulrichova. 2006c. Cytotoxicity of sanguinarine in primary rat hepatocytes is attenuated by dioxin and phenobarbital. Toxicol. Lett. 165(3):282-288.

Dvorak, Z., R. Vrzal, P. Pavek and J. Ulrichova. 2008a. An evidence for regulatory cross-talk between aryl hydrocarbon receptor and glucocorticoid receptor in HepG2 cells. Physiol Res. 57(3):427-435.

Dvorak, Z., R. Vrzal, P. Henklova, P. Jancova, E. Anzenbacherova, P. Maurel, L. Svecova, P. Pavek, J. Ehrmann, R. Havlik, P. Bednar, K. Lemr and J. Ulrichova. 2008b. JNK inhibitor SP600125 is a partial agonist of human aryl hydrocarbon receptor and induces CYP1A1 and CYP1A2 genes in primary human hepatocytes. Biochem. Pharmacol. 75(2):580-588.

Ebert, B., A. Seidel and A. Lampen. 2005. Identification of BCRP as transporter of benzo[a]pyrene conjugates metabolically formed in Caco-2 cells and its induction by Ah-receptor agonists. Carcinogenesis. 26(10):1754-1763.

Elbekai, R.H. and A.O. El-Kadi. 2004. Modulation of aryl hydrocarbon receptor-regulated gene expression by arsenite, cadmium, and chromium. Toxicology. 202(3):249-269.

Elbekai, R.H. and A.O. El-Kadi. 2005. The role of oxidative stress in the modulation of aryl hydrocarbon receptor-regulated genes by As3+, Cd2+, and Cr6+. Free Radic. Biol. Med. 39(11):1499-1511.

Elferink, C.J., N.L. Ge and A. Levine. 2001. Maximal aryl hydrocarbon receptor activity depends on an interaction with the retinoblastoma protein. Mol. Pharmacol. 59(4):664-673.

Elizondo, G., P. Fernandez-Salguero, M.S. Sheikh, G.Y. Kim, A.J. Fornace, K.S. Lee and F.J. Gonzalez. 2000. Altered cell cycle control at the G(2)/M phases in aryl hydrocarbon receptor-null embryo fibroblast. Mol. Pharmacol. 57(5):1056-1063.

Emond, C., J.E. Michalek, L.S. Birnbaum and M.J. DeVito. 2005. Comparison of the use of a physiologically based pharmacokinetic model and a classical pharmacokinetic model for dioxin exposure assessments. Environ. Health Perspect. 113(12):1666-1668.

Emond, C., L.S. Birnbaum and M.J. DeVito. 2006. Use of a physiologically based pharmacokinetic model for rats to study the influence of body fat mass and induction of CYP1A2 on the pharmacokinetics of TCDD. Environ. Health Perspect. 114(9):1394-1400.

Erichsen, T.J., U. Ehmer, S. Kalthoff, T.O. Lankisch, T.M. Muller, P.A. Munzel, M.P. Manns and C.P. Strassburg. 2008. Genetic variability of aryl hydrocarbon receptor (AhR)-mediated regulation of the human UDP glucuronosyltransferase (UGT) 1A4 gene. Toxicol. Appl. Pharmacol. 230(2):252-260.

Eskenazi, B., P. Mocarelli, M. Warner, S. Samuels, L. Needham, D. Patterson, P. Brambilla, P.M. Gerthoux, W. Turner, S. Casalini, M. Cazzaniga and W.Y. Chee. 2001. Seveso Women's Health Study: does zone of residence predict individual TCDD exposure? Chemosphere. 43(4-7):937-942.

Esser, C., V. Temchura, M. Majora, C. Hundeiker, C. Schwarzler and U. Gunthert. 2004. Signaling via the AHR leads to enhanced usage of CD44v10 by murine fetal thymic emigrants: possible role for CD44 in emigration. Int. Immunopharmacol. 4(6):805-818.

Fan, M.Q., A.R. Bell, D.R. Bell, S. Clode, A. Fernandes, P.M. Foster, J.R. Fry, T. Jiang, G. Loizou, A. MacNicoll, B.G. Miller, M. Rose, O. Shaikh-Omar, L. Tran and S. White. 2008. Recombinant expression of aryl hydrocarbon receptor for quantitative ligand-binding analysis. Anal. Biochem.

Faulconer, L., I. Camacho, M. Nagarkatti and P.S. Nagarkatti. 2006. Superantigen-primed T cells exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) replicate poorly following recall encounter. Arch. Toxicol. 80(3):134-145.

Filbrandt, C.R., Z. Wu, B. Zlokovic, L. Opanashuk and T.A. Gasiewicz. 2004. Presence and functional activity of the aryl hydrocarbon receptor in isolated murine cerebral vascular endothelial cells and astrocytes. Neurotoxicology. 25(4):605-616.

Finn, R.D., A.W. McLaren, D. Carrie, C.J. Henderson and C.R. Wolf. 2007. Conditional deletion of cytochrome P450 oxidoreductase in the liver and gastrointestinal tract: a new model for studying the functions of the P450 system. J. Pharmacol. Exp. Ther. 322(1):40-47.

Fiorito, F., U. Pagnini, L. De Martino, S. Montagnaro, R. Ciarcia, S. Florio, M. Pacilio, A. Fucito, A. Rossi, G. Iovane and A. Giordano. 2008a. 2,3,7,8-Tetrachlorodibenzo-p-dioxin increases Bovine Herpesvirus type-1 (BHV-1) replication in Madin-Darby Bovine Kidney (MDBK) cells in vitro. J. Cell Biochem. 103(1):221-233.

Fiorito, F., G. Marfe, E. De Blasio, G.E. Granato, M. Tafani, L. De Martino, S. Montagnaro, S. Florio and U. Pagnini. 2008b. 2,3,7,8-tetrachlorodibenzo-p-dioxin regulates bovine herpesvirus type 1 induced apoptosis by modulating Bcl-2 family members. Apoptosis. 13(10):1243-1252.

Fisher, M.T., M. Nagarkatti and P.S. Nagarkatti. 2004. Combined screening of thymocytes using apoptosis-specific cDNA array and promoter analysis yields novel gene targets mediating TCDD-induced toxicity. Toxicol. Sci. 78(1):116-124.

Fisher, M.T., M. Nagarkatti and P.S. Nagarkatti. 2005. Aryl hydrocarbon receptor-dependent induction of loss of mitochondrial membrane potential in epididydimal spermatozoa by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Toxicol. Lett. 157(2):99-107.

Fletcher, N., D. Wahlstrom, R. Lundberg, C.B. Nilsson, K.C. Nilsson, K. Stockling, H. Hellmold and H. Hakansson. 2005. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) alters the mRNA expression of critical genes associated with cholesterol metabolism, bile acid biosynthesis, and bile transport in rat liver: a microarray study. Toxicol. Appl. Pharmacol. 207(1):1-24.

Franc, M.A., R. Pohjanvirta, J. Tuomisto and A.B. Okey. 2001a. Persistent, low-dose 2,3,7,8-tetrachlorodibenzo-p-dioxin exposure: effect on aryl hydrocarbon receptor expression in a dioxin-resistance model. Toxicol. Appl. Pharmacol. 175(1):43-53.

Franc, M.A., R. Pohjanvirta, J. Tuomisto and A.B. Okey. 2001b. In vivo up-regulation of aryl hydrocarbon receptor expression by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in a dioxin-resistant rat model. Biochem. Pharmacol. 62(12):1565-1578.

Franc, M.A., I.D. Moffat, P.C. Boutros, J.T. Tuomisto, J. Tuomisto, R. Pohjanvirta and A.B. Okey. 2008. Patterns of dioxin-altered mRNA expression in livers of dioxin-sensitive versus dioxin-resistant rats. Arch. Toxicol.

Frericks, M., L.D. Burgoon, T.R. Zacharewski and C. Esser. 2008. Promoter analysis of TCDD-inducible genes in a thymic epithelial cell line indicates the potential for cell-specific transcription factor crosstalk in the AhR response. Toxicol. Appl. Pharmacol. 232(2):268-279.

Fried, K.W., R. Bazzi, L.W. Levy, C. Corsten, K.W. Schramm, D.R. Bell and K.K. Rozman. 2007a. Relationship between aryl hydrocarbon receptor-affinity and the induction of EROD activity by 2,3,7,8-tetrachlorinated phenothiazine and derivatives. Toxicol. Appl. Pharmacol. 224(2):147-155.

Fried, K.W., C.M. Schneider, K.W. Schramm, A. Datta, N. Chahbane, C. Corsten, D.R. Powell, D. Lenoir, A. Kettrup, P. Terranova, G.I. Georg and K.K. Rozman. 2007b. From dioxin to drug lead--the development of 2,3,7,8-tetrachlorophenothiazine. ChemMedChem. 2(6):890-897.

Fritz, W.A., T.M. Lin, R.D. Cardiff and R.E. Peterson. 2007. The aryl hydrocarbon receptor inhibits prostate carcinogenesis in TRAMP mice. Carcinogenesis. 28(2):497-505.

Frowein, J. 2000. Hypothesis: chemical carcinogenesis mediated by a transiently active carcinogen receptor. Cytogenet. Cell Genet. 91(1-4):102-104.

Frueh, F.W., K.C. Hayashibara, P.O. Brown and J.P. Whitlock, Jr. 2001. Use of cDNA microarrays to analyze dioxin-induced changes in human liver gene expression. Toxicol. Lett. 122(3):189-203.

Fujita, H., R. Kosaki, H. Yoshihashi, T. Ogata, M. Tomita, T. Hasegawa, T. Takahashi, N. Matsuo and K. Kosaki. 2002. Characterization of the aryl hydrocarbon receptor repressor gene and association of its Pro185Ala polymorphism with micropenis. Teratology. 65(1):10-18.

Fujita, H., H. Samejima, N. Kitagawa, T. Mitsuhashi, T. Washio, J. Yonemoto, M. Tomita, T. Takahashi and K. Kosaki. 2006. Genome-wide screening of dioxin-responsive genes in fetal brain: bioinformatic and experimental approaches. Congenit. Anom. (Kyoto). 46(3):135-143.

Fukuda, I., I. Sakane, Y. Yabushita, R. Kodoi, S. Nishiumi, T. Kakuda, S. Sawamura, K. Kanazawa and H. Ashida. 2004a. Pigments in green tea leaves (Camellia sinensis) suppress transformation of the aryl hydrocarbon receptor induced by dioxin. J. Agric. Food Chem. 52(9):2499-2506.

Fukuda, I., I. Sakane, Y. Yabushita, S. Sawamura, K. Kanazawa and H. Ashida. 2004b. Black tea extract suppresses transformation of aryl hydrocarbon receptor induced by dioxin. Biofactors. 21(1-4):367-369.

Fukuda, I., I. Sakane, Y. Yabushita, S. Sawamura, K. Kanazawa and H. Ashida. 2005. Black tea theaflavins suppress dioxin-induced transformation of the aryl hydrocarbon receptor. Biosci. Biotechnol. Biochem. 69(5):883-890.

Fukuzawa, N.H., S. Ohsako, Q. Wu, M. Sakaue, Y. Fujii-Kuriyama, T. Baba and C. Tohyama. 2004. Testicular cytochrome P450scc and LHR as possible targets of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the mouse. Mol. Cell Endocrinol. 221(1-2):87-96.

Funatake, C.J., N.B. Marshall and N.I. Kerkvliet. 2008. 2,3,7,8-Tetrachlorodibenzo-p-dioxin alters the differentiation of alloreactive CD8+ T cells toward a regulatory T cell phenotype by a mechanism that is dependent on aryl hydrocarbon receptor in CD4+ T cells. J. Immunotoxicol. 5(1):81-91.

Funseth, E., M. Pahlman, M.L. Eloranta, G. Friman and N.G. Ilback. 2002. Effects of coxsackievirus B3 infection on the acute-phase protein metallothionein and on cytochrome P-4501A1 involved in the detoxification processes of TCDD in the mouse. Sci. Total Environ. 284(1-3):37-47.

Furness, S.G., M.J. Lees and M.L. Whitelaw. 2007. The dioxin (aryl hydrocarbon) receptor as a model for adaptive responses of bHLH/PAS transcription factors. FEBS Lett. 581(19):3616-3625.

Furst, P. 2006. Dioxins, polychlorinated biphenyls and other organohalogen compounds in human milk. Levels, correlations, trends and exposure through breastfeeding. Mol. Nutr. Food Res. 50(10):922-933.

Furue, M. and T. Uenotsuchi. 2005. [Steps for establishment of the diagnostic standard in Yusho patients]. Fukuoka Igaku Zasshi. 96(5):124-134.

Gasiewicz, T.A., T.S. Thurmond, J.E. Staples, F.G. Murante and A.E. Silverstone. 2000. Use of bone marrow chimeras to identify cell targets in the immune system for the actions of chemicals. Ann. N. Y. Acad. Sci. 919:300-303.

Gasiewicz, T.A., E.C. Henry and L.L. Collins. 2008. Expression and activity of aryl hydrocarbon receptors in development and cancer. Crit Rev. Eukaryot. Gene Expr. 18(4):279-321.

Gaylor, D.W. and L.L. Aylward. 2004. An evaluation of benchmark dose methodology for non-cancer continuous-data health effects in animals due to exposures to dioxin (TCDD). Regul. Toxicol. Pharmacol. 40(1):9-17.

Genter, M.B., S. Apparaju and P.B. Desai. 2002. Induction of olfactory mucosal and liver metabolism of lidocaine by 2,3,7,8-tetrachlorodibenzo-p-dioxin. J. Biochem. Mol. Toxicol. 16(3):128-134.

Gerbal-Chaloin, S., L. Pichard-Garcia, J.M. Fabre, A. Sa-Cunha, L. Poellinger, P. Maurel and M. Daujat-Chavanieu. 2006. Role of CYP3A4 in the regulation of the aryl hydrocarbon receptor by omeprazole sulphide. Cell Signal. 18(5):740-750.

Geusau, A., E. Tschachler, M. Meixner, O. Papke, G. Stingl and M. McLachlan. 2001. Cutaneous elimination of 2,3,7,8-tetrachlorodibenzo-p-dioxin. Br. J. Dermatol. 145(6):938-943.

Gill, B.C., C.H. Jeon, H.N. Sung, H.L. Kim, D.W. Jin and J.H. Park. 2008. 2,3,7,8-Tetrachlorodibenzo-p-dioxin modulates the expression of cKrox and Runx3, transcription regulatory factors controlling the lineage commitment of CD4+CD8+ into CD4 and CD8 thymocytes, respectively. Toxicol. Lett. 180(3):189-195.

Gomez-Duran, A., J.M. Carvajal-Gonzalez, S. Mulero-Navarro, B. Santiago-Josefat, A. Puga and P.M. Fernandez-Salguero. 2008. Fitting a xenobiotic receptor into cell homeostasis: How the dioxin receptor interacts with TGFbeta signaling. Biochem. Pharmacol.

Gonzalez, C.A., M. Kogevinas, E. Gadea, G. Pera and O. Papke. 2001. Increase of dioxin blood levels over the last 4 years in the general population in Spain. Epidemiology. 12(3):365.

Gouedard, C., R. Barouki and Y. Morel. 2004. Dietary polyphenols increase paraoxonase 1 gene expression by an aryl hydrocarbon receptor-dependent mechanism. Mol. Cell Biol. 24(12):5209-5222.

Gray, L.E., J. Ostby, J. Furr, C.J. Wolf, C. Lambright, L. Parks, D.N. Veeramachaneni, V. Wilson, M. Price, A. Hotchkiss, E. Orlando and L. Guillette. 2001. Effects of environmental antiandrogens on reproductive development in experimental animals. Hum. Reprod. Update. 7(3):248-264.

Greene, J.F., S. Hays and D. Paustenbach. 2003. Basis for a proposed reference dose (RfD) for dioxin of 1-10 pg/kg-day: a weight of evidence evaluation of the human and animal studies. J. Toxicol. Environ. Health B Crit Rev. 6(2):115-159.

Greenlee, W.E., L.J. Hushka and D.R. Hushka. 2001. Molecular basis of dioxin actions: evidence supporting chemoprotection. Toxicol. Pathol. 29(1):6-7.

Gregoraszczuk, E.L. 2002. Dioxin exposure and porcine reproductive hormonal activity. Cad. Saude Publica. 18(2):453-462.

Gregoraszczuk, E.L., A.K. Wojtowicz, E. Zabielny and A. Grochowalski. 2000. Dose-and-time dependent effect of 2,3,7,8-tetrachlorodibenzo-P-dioxin (TCDD) on progesterone secretion by porcine luteal cells cultured in vitro. J. Physiol Pharmacol. 51(1):127-135.

Gregoraszczuk, E.L., E. Zabielny and D. Ochwat. 2001. Aryl hydrocarbon receptor (AhR)-linked inhibition of luteal cell progesterone secretion in 2,3,7,8-tetrachlorodibenzo-p-dioxin treated cells. J. Physiol Pharmacol. 52(2):303-311.

Grochowalski, A., R. Pieklo, A. Gasinska, R. Chrzaszcz and E.L. Gregoraszczuk. 2000. Accumulation of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in porcine preovulatory follicles after in vitro exposure to TCDD: effects on steroid secretion and cell proliferation. Cytobios. 102(399):21-31.

Grochowalski, A., R. Chrzaszcz, R. Pieklo and E.L. Gregoraszczuk. 2001. Estrogenic and antiestrogenic effect of in vitro treatment of follicular cells with 2,3,7,8-tetrachlorodibenzo-p-dioxin. Chemosphere. 43(4-7):823-827.

Guo, M., A. Joiakim and J.J. Reiners, Jr. 2000. Suppression of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-mediated aryl hydrocarbon receptor transformation and CYP1A1 induction by the phosphatidylinositol 3-kinase inhibitor 2-(4-morpholinyl)-8-phenyl-4H-1- benzopyran-4-one (LY294002). Biochem. Pharmacol. 60(5):635-642.

Guo, L., Y.Y. Zhao, L.Y. Dong, Q. Ding, K.L. Sun, S.J. Ji and L.Z. Wang. 2003. [Antiestrogenic effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the gene expression of insulin-like growth factors family in osteoblast cells]. Zhonghua Yi. Xue. Yi. Chuan Xue. Za Zhi. 20(6):490-494.

Guo, J., M. Sartor, S. Karyala, M. Medvedovic, S. Kann, A. Puga, P. Ryan and C.R. Tomlinson. 2004. Expression of genes in the TGF-beta signaling pathway is significantly deregulated in smooth muscle cells from aorta of aryl hydrocarbon receptor knockout mice. Toxicol. Appl. Pharmacol. 194(1):79-89.

Guo, L., Y.Y. Zhao, S.L. Zhang, K. Liu and X.Y. Gao. 2008. [Effect of dioxin on apoptosis of osteogenic sarcoma cells and regulation on gene expression of insulin-like growth factor binding protein 6]. Zhonghua Lao. Dong. Wei Sheng Zhi. Ye. Bing. Za Zhi. 26(4):223-226.

Haarmann-Stemmann, T. and J. Abel. 2006. The arylhydrocarbon receptor repressor (AhRR): structure, expression, and function. Biol. Chem. 387(9):1195-1199.

Haarmann-Stemmann, T., H. Bothe, A. Kohli, U. Sydlik, J. Abel and E. Fritsche. 2007. Analysis of the transcriptional regulation and molecular function of the aryl hydrocarbon receptor repressor in human cell lines. Drug Metab Dispos. 35(12):2262-2269.

Haarmann-Stemmann, T., H. Bothe and J. Abel. 2008. Growth factors, cytokines and their receptors as downstream targets of arylhydrocarbon receptor (AhR) signaling pathways. Biochem. Pharmacol.

Haddad, S., P. Poulin and K. Krishnan. 2000. Relative lipid content as the sole mechanistic determinant of the adipose tissue:blood partition coefficients of highly lipophilic organic chemicals. Chemosphere. 40(8):839-843.

Hamada, M., H. Satsu, Y. Natsume, S. Nishiumi, I. Fukuda, H. Ashida and M. Shimizu. 2006. TCDD-induced CYP1A1 expression, an index of dioxin toxicity, is suppressed by flavonoids permeating the human intestinal Caco-2 cell monolayers. J. Agric. Food Chem. 54(23):8891-8898.

Han, H.J., M.J. Lim, Y.J. Lee, E.J. Kim, Y.J. Jeon and J.H. Lee. 2005. Effects of TCDD and estradiol-17beta on the proliferation and Na+/glucose cotransporter in renal proximal tubule cells. Toxicol. In Vitro. 19(1):21-30.

Han, E.H., J.Y. Kim and H.G. Jeong. 2006. Effect of biochanin A on the aryl hydrocarbon receptor and cytochrome P450 1A1 in MCF-7 human breast carcinoma cells. Arch. Pharm. Res. 29(7):570-576.

Hankinson, O. 2005. Role of coactivators in transcriptional activation by the aryl hydrocarbon receptor. Arch. Biochem. Biophys. 433(2):379-386.

Hanlon, P.R., L.G. Ganem, Y.C. Cho, M. Yamamoto and C.R. Jefcoate. 2003. AhR- and ERK-dependent pathways function synergistically to mediate 2,3,7,8-tetrachlorodibenzo-p-dioxin suppression of peroxisome proliferator-activated receptor-gamma1 expression and subsequent adipocyte differentiation. Toxicol. Appl. Pharmacol. 189(1):11-27.

Hanlon, P.R., W. Zheng, A.Y. Ko and C.R. Jefcoate. 2005a. Identification of novel TCDD-regulated genes by microarray analysis. Toxicol. Appl. Pharmacol. 202(3):215-228.

Hanlon, P.R., M.A. Cimafranca, X. Liu, Y.C. Cho and C.R. Jefcoate. 2005b. Microarray analysis of early adipogenesis in C3H10T1/2 cells: cooperative inhibitory effects of growth factors and 2,3,7,8-tetrachlorodibenzo-p-dioxin. Toxicol. Appl. Pharmacol. 207(1):39-58.

Hansen, D.A. 2008. Paternal environmental exposures and gene expression during spermatogenesis: research review to research framework. Birth Defects Res. C Embryo. Today. 84(2):155-163.

Haque, M., J. Francis and I. Sehgal. 2005. Aryl hydrocarbon exposure induces expression of MMP-9 in human prostate cancer cell lines. Cancer Lett. 225(1):159-166.

Hardell, L. 2008. Pesticides, soft-tissue sarcoma and non-Hodgkin lymphoma - historical aspects on the precautionary principle in cancer prevention. Acta Oncol. 47(3):347-354.

Harper, P.A., D.S. Riddick and A.B. Okey. 2006. Regulating the regulator: factors that control levels and activity of the aryl hydrocarbon receptor. Biochem. Pharmacol. 72(3):267-279.

Harrad, S., Y. Wang, S. Sandaradura and A. Leeds. 2003. Human dietary intake and excretion of dioxin-like compounds. J. Environ. Monit. 5(2):224-228.

Harrigan, J.A., C.M. Vezina, B.P. McGarrigle, N. Ersing, H.C. Box, A.E. Maccubbin and J.R. Olson. 2004. DNA adduct formation in precision-cut rat liver and lung slices exposed to benzo[a]pyrene. Toxicol. Sci. 77(2):307-314.

Harrigan, J.A., B.P. McGarrigle, T.R. Sutter and J.R. Olson. 2006. Tissue specific induction of cytochrome P450 (CYP) 1A1 and 1B1 in rat liver and lung following in vitro (tissue slice) and in vivo exposure to benzo(a)pyrene. Toxicol. In Vitro. 20(4):426-438.

Harvey, J.L., A.J. Paine, P. Maurel and M.C. Wright. 2000. Effect of the adrenal 11-beta-hydroxylase inhibitor metyrapone on human hepatic cytochrome P-450 expression: induction of cytochrome P-450 3A4. Drug Metab Dispos. 28(1):96-101.

Hassoun, E.A., J. Vodhanel, B. Holden and A. Abushaban. 2006. The effects of ellagic acid and vitamin E succinate on antioxidant enzymes activities and glutathione levels in different brain regions of rats after subchronic exposure to TCDD. J. Toxicol. Environ. Health A. 69(5):381-393.

Hayes, K.R., G.M. Zastrow, M. Nukaya, K. Pande, E. Glover, J.P. Maufort, A.L. Liss, Y. Liu, S.M. Moran, A.L. Vollrath and C.A. Bradfield. 2007. Hepatic transcriptional networks induced by exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. Chem. Res. Toxicol. 20(11):1573-1581.

Hays, S.M. and L.L. Aylward. 2003. Dioxin risks in perspective: past, present, and future. Regul. Toxicol. Pharmacol. 37(2):202-217.

Hays, L.E., C.D. Carpenter and S.L. Petersen. 2002. Evidence that GABAergic neurons in the preoptic area of the rat brain are targets of 2,3,7,8-tetrachlorodibenzo-p-dioxin during development. Environ. Health Perspect. 110 Suppl 3:369-376.

Heilier, J.F., J. Donnez and D. Lison. 2008. Organochlorines and endometriosis: a mini-review. Chemosphere. 71(2):203-210.

Heinzl, H., M. Mittlbock and L. Edler. 2007. On the translation of uncertainty from toxicokinetic to toxicodynamic models--the TCDD example. Chemosphere. 67(9):S365-S374.

Heinzl, H., M. Mittlbock and L. Edler. 2008. Technical uncertainty in the back-calculation of occupational exposure to dioxins. Stat. Med. 27(12):2214-2233.

Henklova, P., R. Vrzal, J. Ulrichova and Z. Dvorak. 2008. Role of mitogen-activated protein kinases in aryl hydrocarbon receptor signaling. Chem. Biol. Interact. 172(2):93-104.

Henley, D.V., C.J. Bellone, D.A. Williams, T.S. Ruh and M.F. Ruh. 2004a. Aryl hydrocarbon receptor-mediated posttranscriptional regulation of IL-1beta. Arch. Biochem. Biophys. 422(1):42-51.

Henley, D.V., C.J. Bellone, D.A. Williams and M.F. Ruh. 2004b. MAPK signaling pathways modulate IL-1beta expression in human keratinocytes. Arch. Biochem. Biophys. 424(1):112-118.

Henry, E.C. and T.A. Gasiewicz. 2008. Molecular determinants of species-specific agonist and antagonist activity of a substituted flavone towards the aryl hydrocarbon receptor. Arch. Biochem. Biophys. 472(2):77-88.

Hillegass, J.M., K.A. Murphy, C.M. Villano and L.A. White. 2006. The impact of aryl hydrocarbon receptor signaling on matrix metabolism: implications for development and disease. Biol. Chem. 387(9):1159-1173.

Hirai, K., J.H. Pan, Y.B. Shui, E. Simamura, H. Shimada, T. Kanamaru and J. Koyama. 2002. Alpha-tocopherol protects cultured human cells from the acute lethal cytotoxicity of dioxin. Int. J. Vitam. Nutr. Res. 72(3):147-153.

Hirakawa, T., T. Minegishi, K. Abe, H. Kishi, Y. Ibuki and K. Miyamoto. 2000a. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the expression of luteinizing hormone receptors during cell differentiation in cultured granulosa cells. Arch. Biochem. Biophys. 375(2):371-376.

Hirakawa, T., T. Minegishi, K. Abe, H. Kishi, K. Inoue, Y. Ibuki and K. Miyamoto. 2000b. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the expression of follicle-stimulating hormone receptors during cell differentiation in cultured granulosa cells. Endocrinology. 141(4):1470-1476.

Hiramatsu, N., A. Kasai, K. Hayakawa, K. Nagai, T. Kubota, J. Yao and M. Kitamura. 2006. Secreted protein-based reporter systems for monitoring inflammatory events: critical interference by endoplasmic reticulum stress. J. Immunol. Methods. 315(1-2):202-207.

Ho, H.M., K. Ohshima, G. Watanabe, K. Taya, E.Y. Strawn and R.J. Hutz. 2006. TCDD increases inhibin A production by human luteinized granulosa cells in vitro. J. Reprod. Dev. 52(4):523-528.

Ho-Jun, K., B.N. Kang, S.W. Cho, H.Y. Son, K.S. Jeong, S.J. Park, S.H. Kim, S.R. Kim, T.H. Kim, M.Y. An and S.Y. Ryu. 2002. Effects of benzo[a]pyrene, 2-bromopropane, phenol and 2,3,7,8-tetrachlorodibenzo-p-dioxin on proinflammatory cytokines gene expression by mice spleen cells. J. Vet. Sci. 3(4):247-254.

Hockings, J.K., P.A. Thorne, M.Q. Kemp, S.S. Morgan, O. Selmin and D.F. Romagnolo. 2006. The ligand status of the aromatic hydrocarbon receptor modulates transcriptional activation of BRCA-1 promoter by estrogen. Cancer Res. 66(4):2224-2232.

Hockley, S.L., V.M. Arlt, D. Brewer, P.R. Te, P. Workman, I. Giddings and D.H. Phillips. 2007. AHR- and DNA-damage-mediated gene expression responses induced by benzo(a)pyrene in human cell lines. Chem. Res. Toxicol. 20(12):1797-1810.

Hoegberg, P., C.K. Schmidt, H. Nau, A.C. Ross, R. Zolfaghari, N. Fletcher, C. Trossvik, C.B. Nilsson and H. Hakansson. 2003. 2,3,7,8-tetrachlorodibenzo-p-dioxin induces lecithin: retinol acyltransferase transcription in the rat kidney. Chem. Biol. Interact. 145(1):1-16.

Hoelper, P., D. Faust, F. Oesch and C. Dietrich. 2004. Transforming growth factor beta1 is not involved in 2,3,7,8-tetrachlorodibenzo- p-dioxin-dependent release from contact-inhibition in WB-F344 cells. Arch. Toxicol. 78(11):643-648.

Hoelper, P., D. Faust, F. Oesch and C. Dietrich. 2005a. Evaluation of the role of c-Src and ERK in TCDD-dependent release from contact-inhibition in WB-F344 cells. Arch. Toxicol. 79(4):201-207.

Hoelper, P., D. Faust, F. Oesch and C. Dietrich. 2005b. Transforming growth factor-beta1 is not involved in TCDD-dependent release from contact inhibition in WB-F344 cells. Arch. Toxicol. 79(1):31-36.

Hokanson, R., S. Miller, M. Hennessey, M. Flesher, W. Hanneman and D. Busbee. 2004. Disruption of estrogen-regulated gene expression by dioxin: downregulation of a gene associated with the onset of non-insulin-dependent diabetes mellitus (type 2 diabetes). Hum. Exp. Toxicol. 23(12):555-564.

Hombach-Klonisch, S., P. Pocar, J. Kauffold and T. Klonisch. 2006. Dioxin exerts anti-estrogenic actions in a novel dioxin-responsive telomerase-immortalized epithelial cell line of the porcine oviduct (TERT-OPEC). Toxicol. Sci. 90(2):519-528.

Hough, R.B. and J. Piatigorsky. 2004. Preferential transcription of rabbit Aldh1a1 in the cornea: implication of hypoxia-related pathways. Mol. Cell Biol. 24(3):1324-1340.

Hsu, E.L., D. Yoon, H.H. Choi, F. Wang, R.T. Taylor, N. Chen, R. Zhang and O. Hankinson. 2007. A proposed mechanism for the protective effect of dioxin against breast cancer. Toxicol. Sci. 98(2):436-444.

Hu, S.W., C.C. Chen, C.Y. Kuo, W.H. Lin and P. Lin. 2006. Increased cytochrome P4501B1 gene expression in peripheral leukocytes of municipal waste incinerator workers. Toxicol. Lett. 160(2):112-120.

Huang, G. and C.J. Elferink. 2005. Multiple mechanisms are involved in Ah receptor-mediated cell cycle arrest. Mol. Pharmacol. 67(1):88-96.

Huang, P., A. Rannug, E. Ahlbom, H. Hakansson and S. Ceccatelli. 2000. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the expression of cytochrome P450 1A1, the aryl hydrocarbon receptor, and the aryl hydrocarbon receptor nuclear translocator in rat brain and pituitary. Toxicol. Appl. Pharmacol. 169(2):159-167.

Huang, P., S. Ceccatelli, H. Hakansson, L. Grandison and A. Rannug. 2002. Constitutive and TCDD-induced expression of Ah receptor-responsive genes in the pituitary. Neurotoxicology. 23(6):783-793.

Huang, P., S. Ceccatelli, P. Hoegberg, T.J. Sten Shi, H. Hakansson and A. Rannug. 2003. TCDD-induced expression of Ah receptor responsive genes in the pituitary and brain of cellular retinol-binding protein (CRBP-I) knockout mice. Toxicol. Appl. Pharmacol. 192(3):262-274.

Huang, P., R. Tofighi, M. Emgard and S. Ceccatelli. 2005. Cell death induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin (2,3,7,8-TCDD) in AtT-20 pituitary cells. Toxicology. 207(3):391-399.

Hukkanen, J., A. Lassila, K. Paivarinta, S. Valanne, S. Sarpo, J. Hakkola, O. Pelkonen and H. Raunio. 2000. Induction and regulation of xenobiotic-metabolizing cytochrome P450s in the human A549 lung adenocarcinoma cell line. Am. J. Respir. Cell Mol. Biol. 22(3):360-366.

Hurst, C.H., M.J. DeVito and L.S. Birnbaum. 2000. Tissue disposition of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in maternal and developing long-evans rats following subchronic exposure. Toxicol. Sci. 57(2):275-283.

Hwang, S.G., H. Sasagawa and F. Matsumura. 2007. Effect of in vitro administered 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on DNA-binding activities of nuclear transcription factors in NIH-3T3 mouse fibroblasts. J. Environ. Sci. Health B. 42(1):115-123.

Iba, M.M. and J. Fung. 2001. Pulmonary Cyp1A1 and CYP1A2 levels and activities in adult male and female offspring of rats exposed during gestation and lactation to 2,3,7, 8-tetrachlorodibenzo-p-dioxin. Biochem. Pharmacol. 62(5):617-626.

Igarashi, T.M., K.L. Bruner-Tran, G.R. Yeaman, B.A. Lessey, D.P. Edwards, E. Eisenberg and K.G. Osteen. 2005. Reduced expression of progesterone receptor-B in the endometrium of women with endometriosis and in cocultures of endometrial cells exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. Fertil. Steril. 84(1):67-74.

Ilvesaro, J., R. Pohjanvirta, J. Tuomisto, M. Viluksela and J. Tuukkanen. 2005. Bone resorption by aryl hydrocarbon receptor-expressing osteoclasts is not disturbed by TCDD in short-term cultures. Life Sci. 77(12):1351-1366.

Imaida, K. and T. Shirai. 2000. [Endocrine disrupting chemicals and carcinogenesis--breast, testis and prostate cancer]. Nippon Rinsho. 58(12):2527-2532.

Inadera, H., H.Y. Dong and K. Matsushima. 2002. WISP-2 is a secreted protein and can be a marker of estrogen exposure in MCF-7 cells. Biochem. Biophys. Res. Commun. 294(3):602-608.

Inouye, K., R. Shinkyo, T. Takita, M. Ohta and T. Sakaki. 2002. Metabolism of polychlorinated dibenzo-p-dioxins (PCDDs) by human cytochrome P450-dependent monooxygenase systems. J. Agric. Food Chem. 50(19):5496-5502.

Irigaray, P., L. Mejean and F. Laurent. 2005. Behaviour of dioxin in pig adipocytes. Food Chem. Toxicol. 43(3):457-460.

Iseki, M., T. Ikuta, T. Kobayashi and K. Kawajiri. 2005. Growth suppression of Leydig TM3 cells mediated by aryl hydrocarbon receptor. Biochem. Biophys. Res. Commun. 331(4):902-908.

Ishii, Y., T. Ishida, J. Mutoh, H. Yamada and K. Oguri. 2005. [Possible candidates for the compound which is expected to attenuate dioxin toxicity]. Fukuoka Igaku Zasshi. 96(5):204-213.

Ishimura, R., S. Ohsako, T. Kawakami, M. Sakaue, Y. Aoki and C. Tohyama. 2002. Altered protein profile and possible hypoxia in the placenta of 2,3,7,8-tetrachlorodibenzo-p-dioxin-exposed rats. Toxicol. Appl. Pharmacol. 185(3):197-206.

Ito, T., S. Tsukumo, N. Suzuki, H. Motohashi, M. Yamamoto, Y. Fujii-Kuriyama, J. Mimura, T.M. Lin, R.E. Peterson, C. Tohyama and K. Nohara. 2004. A constitutively active arylhydrocarbon receptor induces growth inhibition of jurkat T cells through changes in the expression of genes related to apoptosis and cell cycle arrest. J. Biol. Chem. 279(24):25204-25210.

Ito, T., H. Nagai, T.M. Lin, R.E. Peterson, C. Tohyama, T. Kobayashi and K. Nohara. 2006. Organic Chemicals Adsorbed onto Diesel Exhaust Particles Directly Alter the Differentiation of Fetal Thymocytes Through Arylhydrocarbon Receptor but Not Oxidative Stress Responses. J. Immunotoxicol. 3(1):21-30.

Ivnitski-Steele, I. and M.K. Walker. 2005. Inhibition of neovascularization by environmental agents. Cardiovasc. Toxicol. 5(2):215-226.

Jackson, W.G., Jr. and J.E. Michalek. 2001. Temporal changes in TCDD levels in 1419 Air Force Vietnam-era veterans not occupationally exposed to herbicides. J. Expo. Anal. Environ. Epidemiol. 11(1):50-55.

Jana, N.R., S. Sarkar, M. Ishizuka, J. Yonemoto, C. Tohyama and H. Sone. 2000. Comparative effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on MCF-7, RL95-2, and LNCaP cells: role of target steroid hormones in cellular responsiveness to CYP1A1 induction. Mol. Cell Biol. Res. Commun. 4(3):174-180.

Jaruchotikamol, A., K. Jarukamjorn, W. Sirisangtrakul, T. Sakuma, Y. Kawasaki and N. Nemoto. 2007. Strong synergistic induction of CYP1A1 expression by andrographolide plus typical CYP1A inducers in mouse hepatocytes. Toxicol. Appl. Pharmacol. 224(2):156-162.

Jeffy, B.D., R.B. Chirnomas, E.J. Chen, J.M. Gudas and D.F. Romagnolo. 2002. Activation of the aromatic hydrocarbon receptor pathway is not sufficient for transcriptional repression of BRCA-1: requirements for metabolism of benzo[a]pyrene to 7r,8t-dihydroxy-9t,10-epoxy-7,8,9,10-tetrahydrobenzo[a]pyrene. Cancer Res. 62(1):113-121.

Jensen, B.A., R.J. Leeman, J.J. Schlezinger and D.H. Sherr. 2003. Aryl hydrocarbon receptor (AhR) agonists suppress interleukin-6 expression by bone marrow stromal cells: an immunotoxicology study. Environ. Health. 2(1):16.

Jeon, M.S. and C. Esser. 2000. The murine IL-2 promoter contains distal regulatory elements responsive to the Ah receptor, a member of the evolutionarily conserved bHLH-PAS transcription factor family. J. Immunol. 165(12):6975-6983.

Jeong, H.G. and J.Y. Kim. 2002. Effects of o,p'-DDT on the 2,3,7,8-tetrachlorodibenzo-p-dioxin-inducible CYP1A1 expression in murine Hepa-1c1c7 cells. Food Chem. Toxicol. 40(11):1685-1692.

Jeong, H.G., J.Y. Kimand and C.Y. Choi. 2000. Down-regulation of murine Cyp1a-1 in mouse hepatoma Hepa-1c1c7 cells by bisphenol A. Biochem. Biophys. Res. Commun. 277(3):594-598.

Jeong, H.G., J.Y. Kim, C.Y. Choi, H.J. You and K. Hahm. 2001. Suppression of CYP1A1 expression by 4-nonylphenol in murine Hepa-1c1c7 cells. Cancer Lett. 165(1):95-101.

Ji, H.J., H.K. Yeo, N.H. Lee, J.S. Hwang, K.A. Koo, S.W. Cheong, J.H. Park, G.S. Oh, C.S. Yoon and H.J. Youn. 2005. A carbohydrate fraction, AIP1, from Artemisia iwayomogi down-regulates Fas gene expression and suppresses apoptotic death of the thymocytes induced by 2,3,7,8-tectrachlorodibenzo-p-dioxin. Biotechnol. Lett. 27(4):253-257.

Jiang, H., Y.M. Shen, A.M. Quinn and T.M. Penning. 2005. Competing roles of cytochrome P450 1A1/1B1 and aldo-keto reductase 1A1 in the metabolic activation of (+/-)-7,8-dihydroxy-7,8-dihydro-benzo[a]pyrene in human bronchoalveolar cell extracts. Chem. Res. Toxicol. 18(2):365-374.

Jiang, H., D.K. Vudathala, I.A. Blair and T.M. Penning. 2006. Competing roles of aldo-keto reductase 1A1 and cytochrome P4501B1 in benzo[a]pyrene-7,8-diol activation in human bronchoalveolar H358 cells: role of AKRs in P4501B1 induction. Chem. Res. Toxicol. 19(1):68-78.

Jigorel, E., M. Le Vee, C. Boursier-Neyret, Y. Parmentier and O. Fardel. 2006. Differential regulation of sinusoidal and canalicular hepatic drug transporter expression by xenobiotics activating drug-sensing receptors in primary human hepatocytes. Drug Metab Dispos. 34(10):1756-1763.

Jin, B. and D.Y. Ryu. 2004. Regulation of CYP1A2 by histone deacetylase inhibitors in mouse hepatocytes. J. Biochem. Mol. Toxicol. 18(3):131-132.

Jin, B., G. Kim, D.W. Park and D.Y. Ryu. 2004a. Microarray analysis of gene regulation in the Hepa1c1c7 cell line following exposure to the DNA methylation inhibitor 5-aza-2'-deoxycytidine and 2,3,7,8-tetrachlorodibenzo-p-dioxin. Toxicol. In Vitro. 18(5):659-664.

Jin, D.Q., J.W. Jung, Y.S. Lee and J.A. Kim. 2004b. 2,3,7,8-Tetrachlorodibenzo-p-dioxin inhibits cell proliferation through arylhydrocarbon receptor-mediated G1 arrest in SK-N-SH human neuronal cells. Neurosci. Lett. 363(1):69-72.

Joiakim, A., P.A. Mathieu, A.A. Elliott and J.J. Reiners, Jr. 2004. Superinduction of CYP1A1 in MCF10A cultures by cycloheximide, anisomycin, and puromycin: a process independent of effects on protein translation and unrelated to suppression of aryl hydrocarbon receptor proteolysis by the proteasome. Mol. Pharmacol. 66(4):936-947.

Jones, L.C. and J.P. Whitlock, Jr. 2001. Dioxin-inducible transactivation in a chromosomal setting. Analysis of the acidic domain of the Ah receptor. J. Biol. Chem. 276(27):25037-25042.

Ju, Q., S. Fimmel, R. Stahlmann and C.C. Zouboulis. 2008. 2,3,7,8-tetrachlorodibenzo-p-dioxin alters the differentiation of SZ95 sebocytes in vitro. Exp. Dermatol. 17(7):632.

Jung, K.K., S.Y. Kim, T.G. Kim, J.H. Kang, S.Y. Kang, J.Y. Cho and S.H. Kim. 2007. Differential regulation of thyroid hormone receptor-mediated function by endocrine disruptors. Arch. Pharm. Res. 30(5):616-623.

Kamiguchi, Y. and H. Tateno. 2002. Radiation- and chemical-induced structural chromosome aberrations in human spermatozoa. Mutat. Res. 504(1-2):183-191.

Kang, H.K., N.A. Dalager, L.L. Needham, D.G. Patterson, Jr., G.M. Matanoski, S. Kanchanaraksa and P.S. Lees. 2001. US Army Chemical Corps Vietnam veterans health study: preliminary results. Chemosphere. 43(4-7):943-949.

Kang, M.J., D.Y. Lee, W.A. Joo and C.W. Kim. 2005. Plasma protein level changes in waste incineration workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. J. Proteome. Res. 4(4):1248-1255.

Kang, H.J., H.J. Kim, S.K. Kim, R. Barouki, C.H. Cho, K.K. Khanna, E.M. Rosen and I. Bae. 2006. BRCA1 modulates xenobiotic stress-inducible gene expression by interacting with ARNT in human breast cancer cells. J. Biol. Chem. 281(21):14654-14662.

Kang, H.J., H.J. Kim, C.H. Cho, Y. Hu, R. Li and I. Bae. 2008. BRCA1 transcriptional activity is enhanced by interactions between its AD1 domain and AhR. Cancer Chemother. Pharmacol. 62(6):965-975.

Kanno, J. 2000. [In vivo test for endocrine disruptors]. Nippon Rinsho. 58(12):2495-2501.

Karyala, S., J. Guo, M. Sartor, M. Medvedovic, S. Kann, A. Puga, P. Ryan and C.R. Tomlinson. 2004. Different global gene expression profiles in benzo[a]pyrene- and dioxin-treated vascular smooth muscle cells of AHR-knockout and wild-type mice. Cardiovasc. Toxicol. 4(1):47-73.

Kasai, A., J. Yao, K. Yamauchi, N. Hiramatsu, K. Hayakawa, Y. Meng, S. Maeda and M. Kitamura. 2006. Influence of cAMP on reporter bioassays for dioxin and dioxin-like compounds. Toxicol. Appl. Pharmacol. 211(1):11-19.

Kasai, A., N. Hiramatsu, K. Hayakawa, J. Yao and M. Kitamura. 2008. Direct, continuous monitoring of air pollution by transgenic sensor mice responsive to halogenated and polycyclic aromatic hydrocarbons. Environ. Health Perspect. 116(3):349-354.

Kashuba, E.V., K. Gradin, M. Isaguliants, L. Szekely, L. Poellinger, G. Klein and A. Kazlauskas. 2006. Regulation of transactivation function of the aryl hydrocarbon receptor by the Epstein-Barr virus-encoded EBNA-3 protein. J. Biol. Chem. 281(2):1215-1223.

Kato, S. 2000. [Nuclear steroid hormone receptors as possible target molecule for endocrine disruptant]. Nippon Rinsho. 58(7):1535-1543.

Kato, S. and F. Otake. 2004. [A molecular structure involved in the estrogenic disrupting action caused by dioxin receptors]. Nippon Naika Gakkai Zasshi. 93(4):808-815.

Kayano, S., Y. Suzuki, K. Kanno, Y. Aoki, S. Kure, A. Yamada and Y. Matsubara. 2004. Significant association between nonsyndromic oral clefts and arylhydrocarbon receptor nuclear translocator (ARNT). Am. J. Med. Genet. A. 130A(1):40-44.

Kerger, B.D., H.W. Leung, P. Scott, D.J. Paustenbach, L.L. Needham, D.G. Patterson, Jr., P.M. Gerthoux and P. Mocarelli. 2006. Age- and concentration-dependent elimination half-life of 2,3,7,8-tetrachlorodibenzo-p-dioxin in Seveso children. Environ. Health Perspect. 114(10):1596-1602.

Kerger, B.D., H.W. Leung, P.K. Scott and D.J. Paustenbach. 2007a. Refinements on the age-dependent half-life model for estimating child body burdens of polychlorodibenzodioxins and dibenzofurans. Chemosphere. 67(9):S272-S278.

Kerger, B.D., H.W. Leung, P.K. Scott and D.J. Paustenbach. 2007b. An adaptable internal dose model for risk assessment of dietary and soil dioxin exposures in young children. Toxicol. Sci. 100(1):224-237.

Kerkvliet, N.I. 2002. Recent advances in understanding the mechanisms of TCDD immunotoxicity. Int. Immunopharmacol. 2(2-3):277-291.

Kern, P.A., R.B. Fishman, W. Song, A.D. Brown and V. Fonseca. 2002a. The effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on oxidative enzymes in adipocytes and liver. Toxicology. 171(2-3):117-125.

Kern, P.A., A. Dicker-Brown, S.T. Said, R. Kennedy and V.A. Fonseca. 2002b. The stimulation of tumor necrosis factor and inhibition of glucose transport and lipoprotein lipase in adipose cells by 2,3,7,8-tetrachlorodibenzo-p-dioxin. Metabolism. 51(1):65-68.

Khan, S., R. Barhoumi, R. Burghardt, S. Liu, K. Kim and S. Safe. 2006a. Molecular mechanism of inhibitory aryl hydrocarbon receptor-estrogen receptor/Sp1 cross talk in breast cancer cells. Mol. Endocrinol. 20(9):2199-2214.

Khan, S., A. Konstantinov, B. Chittim, A. McAlees, B. Yeo and S. Safe. 2006b. Structure-dependent Ah receptor agonist activities of chlorinated biphenylenes. Toxicol. In Vitro. 20(7):1234-1237.

Khan, S., S. Liu, M. Stoner and S. Safe. 2007. Cobaltous chloride and hypoxia inhibit aryl hydrocarbon receptor-mediated responses in breast cancer cells. Toxicol. Appl. Pharmacol. 223(1):28-38.

Kietz, S., J.S. Thomsen, J. Matthews, K. Pettersson, A. Strom and J.A. Gustafsson. 2004. The Ah receptor inhibits estrogen-induced estrogen receptor beta in breast cancer cells. Biochem. Biophys. Res. Commun. 320(1):76-82.

Kikuchi, H., M. Shibazaki, S. Ahmed and T. Baba. 2001. Method for evaluation of immunotoxicity of dioxin compounds using human T-lymphoblastic cell line, L-MAT. Chemosphere. 43(4-7):815-818.

Kim, J.E. and Y.Y. Sheen. 2000a. Inhibition of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-stimulated Cyp1a1 promoter activity by hypoxic agents. Biochem. Pharmacol. 59(12):1549-1556.

Kim, J.E. and Y.Y. Sheen. 2000b. Nitric oxide inhibits dioxin action for the stimulation of Cyp1a1 promoter activity. Biol. Pharm. Bull. 23(5):575-580.

Kim, S.Y. and J.H. Yang. 2005. Neurotoxic effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin in cerebellar granule cells. Exp. Mol. Med. 37(1):58-64.

Kim, J.Y., E.H. Han, D.W. Shin, T.C. Jeong, E.S. Lee, E.R. Woo and H.G. Jeong. 2004. Suppression of CYP1A1 expression by naringenin in murine Hepa-1c1c7 cells. Arch. Pharm. Res. 27(8):857-862.

Kim, W.K., Y.J. In, J.H. Kim, H.J. Cho, J.H. Kim, S. Kang, C.Y. Lee and S.C. Lee. 2006. Quantitative relationship of dioxin-responsive gene expression to dioxin response element in Hep3B and HepG2 human hepatocarcinoma cell lines. Toxicol. Lett. 165(2):174-181.

Kim, S.Y., H.G. Lee, E.J. Choi, K.Y. Park and J.H. Yang. 2007. TCDD alters PKC signaling pathways in developing neuronal cells in culture. Chemosphere. 67(9):S421-S427.

Kim, J.H., Y.J. In, W.K. Kim, K.H. Bae, S. Kang and S.C. Lee. 2008. Differential signatures of protein glycosylation and phosphorylation in human Chang liver cells induced by TCDD treatment. Toxicol. Lett. 178(1):20-28.

Kimata, H. 2003. 2,3,7,8-tetrachlorodibenzo-p-dioxin selectively enhances spontaneous IgE production in B cells from atopic patients. Int. J. Hyg. Environ. Health. 206(6):601-604.

Kishi, R., F. Sata, E. Yoshioka, S. Ban, S. Sasaki, K. Konishi and N. Washino. 2008. Exploiting gene-environment interaction to detect adverse health effects of environmental chemicals on the next generation. Basic Clin. Pharmacol. Toxicol. 102(2):191-203.

Kitamura, K., M. Nagao, T. Yamada, M. Sunaga, J. Hata and S. Watanabe. 2001. Dioxins in bile in relation to those in the human liver and blood. J. Toxicol. Sci. 26(5):327-336.

Kitamura, K., M. Nagao, H. Hayatsu and M. Morita. 2005. Effect of chlorophyllin-chitosan on excretion of dioxins in a healthy man. Environ. Sci. Technol. 39(4):1084-1091.

Kiukkonen, A., C. Sahlberg, P.L. Lukinmaa, S. Alaluusua, E. Peltonen and A.M. Partanen. 2006a. 2,3,7,8-tetrachlorodibenzo-p-dioxin specifically reduces mRNA for the mineralization-related dentin sialophosphoprotein in cultured mouse embryonic molar teeth. Toxicol. Appl. Pharmacol. 216(3):399-406.

Kiukkonen, A., C. Sahlberg, A.M. Partanen, S. Alaluusua, R. Pohjanvirta, J. Tuomisto and P.L. Lukinmaa. 2006b. Interference by 2,3,7,8-tetrachlorodibenzo-p-dioxin with cultured mouse submandibular gland branching morphogenesis involves reduced epidermal growth factor receptor signaling. Toxicol. Appl. Pharmacol. 212(3):200-211.

Klinge, C.M., K. Kaur and H.I. Swanson. 2000. The aryl hydrocarbon receptor interacts with estrogen receptor alpha and orphan receptors COUP-TFI and ERRalpha1. Arch. Biochem. Biophys. 373(1):163-174.

Klinge, C.M., S.C. Jernigan, K.E. Risinger, J.E. Lee, V.V. Tyulmenkov, K.C. Falkner and R.A. Prough. 2001. Short heterodimer partner (SHP) orphan nuclear receptor inhibits the transcriptional activity of aryl hydrocarbon receptor (AHR)/AHR nuclear translocator (ARNT). Arch. Biochem. Biophys. 390(1):64-70.

Knerr, S. and D. Schrenk. 2006. Carcinogenicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in experimental models. Mol. Nutr. Food Res. 50(10):897-907.

Knerr, S., J. Schaefer, S. Both, A. Mally, W. Dekant and D. Schrenk. 2006. 2,3,7,8-Tetrachlorodibenzo-p-dioxin induced cytochrome P450s alter the formation of reactive oxygen species in liver cells. Mol. Nutr. Food Res. 50(4-5):378-384.

Kobayashi, S., H. Okamoto, T. Iwamoto, Y. Toyama, T. Tomatsu, H. Yamanaka and S. Momohara. 2008. A role for the aryl hydrocarbon receptor and the dioxin TCDD in rheumatoid arthritis. Rheumatology. (Oxford). 47(9):1317-1322.

Kohle, C., I. Hassepass, B.S. Bock-Hennig, K. Walter Bock, L. Poellinger and J. McGuire. 2002. Conditional expression of a constitutively active aryl hydrocarbon receptor in MCF-7 human breast cancer cells. Arch. Biochem. Biophys. 402(2):172-179.

Kohn, M.C. 2000. Effects of TCDD on thyroid hormone homeostasis in the rat. Drug Chem. Toxicol. 23(1):259-277.

Kohn, M.C., N.J. Walker, A.H. Kim and C.J. Portier. 2001. Physiological modeling of a proposed mechanism of enzyme induction by TCDD. Toxicology. 162(3):193-208.

Koliopanos, A., J. Kleeff, Y. Xiao, S. Safe, A. Zimmermann, M.W. Buchler and H. Friess. 2002. Increased arylhydrocarbon receptor expression offers a potential therapeutic target for pancreatic cancer. Oncogene. 21(39):6059-6070.

Kollara, A. and T.J. Brown. 2006. Functional interaction of nuclear receptor coactivator 4 with aryl hydrocarbon receptor. Biochem. Biophys. Res. Commun. 346(2):526-534.

Kolluri, S.K., C. Balduf, M. Hofmann and M. Gottlicher. 2001. Novel target genes of the Ah (dioxin) receptor: transcriptional induction of N-myristoyltransferase 2. Cancer Res. 61(23):8534-8539.

Korashy, H.M. and A.O. El-Kadi. 2004. Differential effects of mercury, lead and copper on the constitutive and inducible expression of aryl hydrocarbon receptor (AHR)-regulated genes in cultured hepatoma Hepa 1c1c7 cells. Toxicology. 201(1-3):153-172.

Korashy, H.M. and A.O. El-Kadi. 2005. Regulatory mechanisms modulating the expression of cytochrome P450 1A1 gene by heavy metals. Toxicol. Sci. 88(1):39-51.

Korashy, H.M. and A.O. El-Kadi. 2008. Modulation of TCDD-mediated induction of cytochrome P450 1A1 by mercury, lead, and copper in human HepG2 cell line. Toxicol. In Vitro. 22(1):154-158.

Korkalainen, M., J. Tuomisto and R. Pohjanvirta. 2000. Restructured transactivation domain in hamster AH receptor. Biochem. Biophys. Res. Commun. 273(1):272-281.

Korkalainen, M., J. Tuomisto and R. Pohjanvirta. 2003. Identification of novel splice variants of ARNT and ARNT2 in the rat. Biochem. Biophys. Res. Commun. 303(4):1095-1100.

Korkalainen, M., J. Tuomisto and R. Pohjanvirta. 2004. Primary structure and inducibility by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) of aryl hydrocarbon receptor repressor in a TCDD-sensitive and a TCDD-resistant rat strain. Biochem. Biophys. Res. Commun. 315(1):123-131.

Korkalainen, M., J. Linden, J. Tuomisto and R. Pohjanvirta. 2005. Effect of TCDD on mRNA expression of genes encoding bHLH/PAS proteins in rat hypothalamus. Toxicology. 208(1):1-11.

Korner, W., G. Golor, T. Schulz, T. Wiesmuller, H. Hagenmaier and D. Neubert. 2002. Tissue concentrations and induction of a hepatic monooxygenase in male Wistar rats after repeated doses of defined polychlorinated dibenzo-p-dioxin and dibenzofuran (PCDDs and PCDFs) mixtures. Arch. Toxicol. 75(11-12):653-664.

Kosina, P., P. Maurel, J. Ulrichova and Z. Dvorak. 2005. Effect of silybin and its glycosides on the expression of cytochromes P450 1A2 and 3A4 in primary cultures of human hepatocytes. J. Biochem. Mol. Toxicol. 19(3):149-153.

Krig, S.R. and R.H. Rice. 2000. TCDD suppression of tissue transglutaminase stimulation by retinoids in malignant human keratinocytes. Toxicol. Sci. 56(2):357-364.

Krig, S.R., R.A. Chandraratna, M.M. Chang, R. Wu and R.H. Rice. 2002. Gene-specific TCDD suppression of RARalpha- and RXR-mediated induction of tissue transglutaminase. Toxicol. Sci. 68(1):102-108.

Kronenberg, S., C. Esser and C. Carlberg. 2000. An aryl hydrocarbon receptor conformation acts as the functional core of nuclear dioxin signaling. Nucleic Acids Res. 28(12):2286-2291.

Kumagai, S. and S. Koda. 2005. Polychlorinated dibenzo-p-dioxin and dibenzofuran concentrations in serum samples of workers at an infectious waste incineration plant in Japan. J. Occup. Environ. Hyg. 2(2):120-125.

Kumar, M.B., P. Ramadoss, R.K. Reen, J.P. Vanden Heuvel and G.H. Perdew. 2001. The Q-rich subdomain of the human Ah receptor transactivation domain is required for dioxin-mediated transcriptional activity. J. Biol. Chem. 276(45):42302-42310.

Kurachi, M., S. Hashimoto, A. Obata, S. Nagai, T. Nagahata, H. Inadera, H. Sone, C. Tohyama, S. Kaneko, K. Kobayashi and K. Matsushima. 2002. Identification of 2,3,7,8-tetrachlorodibenzo-p-dioxin-responsive genes in mouse liver by serial analysis of gene expression. Biochem. Biophys. Res. Commun. 292(2):368-377.

Kuroda, M., K. Oikawa, T. Ohbayashi, K. Yoshida, K. Yamada, J. Mimura, Y. Matsuda, Y. Fujii-Kuriyama and K. Mukai. 2005. A dioxin sensitive gene, mammalian WAPL, is implicated in spermatogenesis. FEBS Lett. 579(1):167-172.

Kuznetsov, N.V., P. Andersson, K. Gradin, P. Stein, A. Dieckmann, S. Pettersson, A. Hanberg and L. Poellinger. 2005. The dioxin/aryl hydrocarbon receptor mediates downregulation of osteopontin gene expression in a mouse model of gastric tumourigenesis. Oncogene. 24(19):3216-3222.

Kwon, M.J., K.S. Jeong, E.J. Choi and B.H. Lee. 2003. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)-induced activation of mitogen-activated protein kinase signaling pathway in Jurkat T cells. Pharmacol. Toxicol. 93(4):186-190.

Lahvis, G.P., S.L. Lindell, R.S. Thomas, R.S. McCuskey, C. Murphy, E. Glover, M. Bentz, J. Southard and C.A. Bradfield. 2000. Portosystemic shunting and persistent fetal vascular structures in aryl hydrocarbon receptor-deficient mice. Proc. Natl. Acad. Sci. U. S. A. 97(19):10442-10447.

Lai, Z.W., N.C. Fiore, P.J. Hahn, T.A. Gasiewicz and A.E. Silverstone. 2000. Differential effects of diethylstilbestrol and 2,3,7,8-tetrachlorodibenzo-p-dioxin on thymocyte differentiation, proliferation, and apoptosis in bcl-2 transgenic mouse fetal thymus organ culture. Toxicol. Appl. Pharmacol. 168(1):15-24.

Lai, K.P., M.H. Wong and C.K. Wong. 2004. Modulation of AhR-mediated CYP1A1 mRNA and EROD activities by 17beta-estradiol and dexamethasone in TCDD-induced H411E cells. Toxicol. Sci. 78(1):41-49.

Lai, K.P., M.H. Wong and C.K. Wong. 2005a. Inhibition of CYP450scc expression in dioxin-exposed rat Leydig cells. J. Endocrinol. 185(3):519-527.

Lai, K.P., M.H. Wong and C.K. Wong. 2005b. Effects of TCDD in modulating the expression of Sertoli cell secretory products and markers for cell-cell interaction. Toxicology. 206(1):111-123.

Lai, K.P., N.K. Mak, X. Wei, R.N. Wong, M.H. Wong and C.K. Wong. 2006. Bifunctional modulating effects of an indigo dimer (bisindigotin) to CYP1A1 induction in H4IIE cells. Toxicology. 226(2-3):188-196.

Laiosa, M.D., A. Wyman, F.G. Murante, N.C. Fiore, J.E. Staples, T.A. Gasiewicz and A.E. Silverstone. 2003. Cell proliferation arrest within intrathymic lymphocyte progenitor cells causes thymic atrophy mediated by the aryl hydrocarbon receptor. J. Immunol. 171(9):4582-4591.

Lakhman, S.S., X. Chen, V. Gonzalez-Covarrubias, E.G. Schuetz and J.G. Blanco. 2007. Functional characterization of the promoter of human carbonyl reductase 1 (CBR1). Role of XRE elements in mediating the induction of CBR1 by ligands of the aryl hydrocarbon receptor. Mol. Pharmacol. 72(3):734-743.

LaKind, J.S., S.M. Hays, L.L. Aylward and D.Q. Naiman. 2008. Perspective on serum dioxin levels in the United States: an evaluation of the NHANES data. J. Expo. Sci. Environ. Epidemiol.

Lamartiniere, C.A. 2002. Timing of exposure and mammary cancer risk. J. Mammary. Gland. Biol. Neoplasia. 7(1):67-76.

Landi, M.T. and A. Baccarelli. 2003. Correspondence re: K. Toide et al., Aryl hydrocarbon hydroxylase represents CYP1B1, and not CYP1A1, in human freshly isolated white cells: trimodal distribution of Japanese population according to induction of CYP1B1 mRNA by environmental dioxins. 12: 219-222, 2003. Cancer Epidemiol. Biomarkers Prev. 12(10):1116-1117.

Landi, M.T., P.A. Bertazzi, A. Baccarelli, D. Consonni, S. Masten, G. Lucier, P. Mocarelli, L. Needham, N. Caporaso and J. Grassman. 2003. TCDD-mediated alterations in the AhR-dependent pathway in Seveso, Italy, 20 years after the accident. Carcinogenesis. 24(4):673-680.

Landi, M.T., A.W. Bergen, A. Baccarelli, D.G. Patterson, Jr., J. Grassman, M. Ter-Minassian, P. Mocarelli, N. Caporaso, S.A. Masten, A.C. Pesatori, G.S. Pittman and D.A. Bell. 2005. CYP1A1 and CYP1B1 genotypes, haplotypes, and TCDD-induced gene expression in subjects from Seveso, Italy. Toxicology. 207(2):191-202.

Langer, P., A. Kocan, M. Tajtakova, Z. Radikova, J. Petrik, J. Koska, L. Ksinantova, R. Imrich, M. Huckova, J. Chovancova, B. Drobna, S. Jursa, A. Bergman, M. Athanasiadou, L. Hovander, D. Gasperikova, T. Trnovec, E. Sebokova and I. Klimes. 2007a. Possible effects of persistent organochlorinated pollutants cocktail on thyroid hormone levels and pituitary-thyroid interrelations. Chemosphere. 70(1):110-118.

Langer, P., M. Tajtakova, A. Kocan, J. Petrik, J. Koska, L. Ksinantova, Z. Radikova, J. Ukropec, R. Imrich, M. Huckova, J. Chovancova, B. Drobna, S. Jursa, M. Vlcek, A. Bergman, M. Athanasiadou, L. Hovander, Y. Shishiba, T. Trnovec, E. Sebokova and I. Klimes. 2007b. Thyroid ultrasound volume, structure and function after long-term high exposure of large population to polychlorinated biphenyls, pesticides and dioxin. Chemosphere. 69(1):118-127.

Larsen, J.C. 2006. Risk assessments of polychlorinated dibenzo- p-dioxins, polychlorinated dibenzofurans, and dioxin-like polychlorinated biphenyls in food. Mol. Nutr. Food Res. 50(10):885-896.

Larsen, M.C., P.B. Brake, R.S. Pollenz and C.R. Jefcoate. 2004. Linked expression of Ah receptor, ARNT, CYP1A1, and CYP1B1 in rat mammary epithelia, in vitro, is each substantially elevated by specific extracellular matrix interactions that precede branching morphogenesis. Toxicol. Sci. 82(1):46-61.

Laupeze, B., L. Amiot, L. Sparfel, E. Le Ferrec, R. Fauchet and O. Fardel. 2002. Polycyclic aromatic hydrocarbons affect functional differentiation and maturation of human monocyte-derived dendritic cells. J. Immunol. 168(6):2652-2658.

Laurent, C., C. Feidt, N. Grova, D. Mpassi, E. Lichtfouse, F. Laurent and G. Rychen. 2002. Portal absorption of 14C after ingestion of spiked milk with 14C-phenanthrene, 14C-benzo[a]pyrene or 14C-TCDD in growing pigs. Chemosphere. 48(8):843-848.

Laurent, C., P. Marchand, C. Feidt, B. Le Bizec and G. Rychen. 2005. Tissue distribution and bioconcentration factors of PCDD/Fs in the liver and adipose tissue following chronic ingestion of contaminated milk in rats. Chemosphere. 60(7):929-938.

Lawrence, B.P. 2007. Environmental toxins as modulators of antiviral immune responses. Viral Immunol. 20(2):231-242.

Lecureur, V., E.L. Ferrec, M. N'Diaye, M.L. Vee, C. Gardyn, D. Gilot and O. Fardel. 2005. ERK-dependent induction of TNFalpha expression by the environmental contaminant benzo(a)pyrene in primary human macrophages. FEBS Lett. 579(9):1904-1910.

Ledirac, N., C. Delescluse, P. Lesca, M.P. Piechocki, R.N. Hines, G. de Sousa, M. Pralavorio and R. Rahmani. 2000. Diflubenzuron, a benzoyl-urea insecticide, is a potent inhibitor of TCDD-induced CYP1A1 expression in HepG2 cells. Toxicol. Appl. Pharmacol. 164(3):273-279.

Lee, J.E. and S. Safe. 2000. 3',4'-dimethoxyflavone as an aryl hydrocarbon receptor antagonist in human breast cancer cells. Toxicol. Sci. 58(2):235-242.

Lee, H.M., Q. He, E.W. Englander and G.H. Greeley, Jr. 2000. Endocrine disruptive effects of polychlorinated aromatic hydrocarbons on intestinal cholecystokinin in rats. Endocrinology. 141(8):2938-2944.

Lee, H.C., S.G. Hwang, Y.G. Lee, H.O. Sohn, D.W. Lee, S.Y. Hwang, Y.S. Kwak, J.J. Wee, W.H. Joo, Y.K. Cho and J.Y. Moon. 2002. In vivo effects of Panax ginseng extracts on the cytochrome P450-dependent monooxygenase system in the liver of 2,3,7,8-tetrachlorodibenzo-p-dioxin-exposed guinea pig. Life Sci. 71(7):759-769.

Lee, Y.S., D.Q. Jin, S.H. Park, S.Y. Han, H.S. Kim, T.C. Jeong, K. Huh and J.A. Kim. 2002. 2,3,7,8-tetrachlorobenzo-p-dioxin inhibits proliferation of SK-N-SH human neuronal cells through decreased production of reactive oxygen species. Free Radic. Res. 36(12):1283-1289.

Lee, S.H., D.Y. Lee, W.K. Son, W.A. Joo and C.W. Kim. 2005. Proteomic characterization of rat liver exposed to 2,3,7,8-tetrachlorobenzo-p-dioxin. J. Proteome. Res. 4(2):335-343.

Lee, K.A., L.D. Burgoon, L. Lamb, E. Dere, T.R. Zacharewski, J.B. Hogenesch and J.J. LaPres. 2006. Identification and characterization of genes susceptible to transcriptional cross-talk between the hypoxia and dioxin signaling cascades. Chem. Res. Toxicol. 19(10):1284-1293.

Lee, H.G., S.Y. Kim, E.J. Choi, K.Y. Park and J.H. Yang. 2007a. Translocation of PKC-betaII is mediated via RACK-1 in the neuronal cells following dioxin exposure. Neurotoxicology. 28(2):408-414.

Lee, J.A., J.A. Hwang, H.N. Sung, C.H. Jeon, B.C. Gill, H.J. Youn and J.H. Park. 2007b. 2,3,7,8-Tetrachlorodibenzo-p-dioxin modulates functional differentiation of mouse bone marrow-derived dendritic cells Downregulation of RelB by 2,3,7,8-tetrachlorodibenzo-p-dioxin. Toxicol. Lett. 173(1):31-40.

Lees, M.J., D.J. Peet and M.L. Whitelaw. 2003. Defining the role for XAP2 in stabilization of the dioxin receptor. J. Biol. Chem. 278(38):35878-35888.

Legare, M.E., W.H. Hanneman, R. Barhoumi, R.C. Burghardt and E. Tiffany-Castiglioni. 2000. 2,3,7,8-Tetrachlorodibenzo-p-dioxin alters hippocampal astroglia-neuronal gap junctional communication. Neurotoxicology. 21(6):1109-1116.

Leijs, M.M., J.G. Koppe, K. Olie, W.M. van Aalderen, P. Voogt, T. Vulsma, M. Westra and G.W. ten Tusscher. 2008. Delayed initiation of breast development in girls with higher prenatal dioxin exposure; a longitudinal cohort study. Chemosphere. 73(6):999-1004.

Lekas, P., K.L. Tin, C. Lee and R.D. Prokipcak. 2000. The human cytochrome P450 1A1 mRNA is rapidly degraded in HepG2 cells. Arch. Biochem. Biophys. 384(2):311-318.

Leung, H.W., B.D. Kerger and D.J. Paustenbach. 2006. Elimination half-lives of selected polychlorinated dibenzodioxins and dibenzofurans in breast-fed human infants. J. Toxicol. Environ. Health A. 69(6):437-443.

Levin, M., B. Morsey, C. Mori, P.R. Nambiar and S. De Guise. 2005. PCBs and TCDD, alone and in mixtures, modulate marine mammal but not B6C3F1 mouse leukocyte phagocytosis. J. Toxicol. Environ. Health A. 68(8):635-656.

Levin, M., B. Morsey and S. De Guise. 2007. Modulation of the respiratory burst by organochlorine mixtures in marine mammals, humans, and mice. J. Toxicol. Environ. Health A. 70(1):73-83.

Levine-Fridman, A., L. Chen and C.J. Elferink. 2004. Cytochrome P4501A1 promotes G1 phase cell cycle progression by controlling aryl hydrocarbon receptor activity. Mol. Pharmacol. 65(2):461-469.

Levitt, D.G. 2007. Heterogeneity of human adipose blood flow. BMC. Clin. Pharmacol. 7:1.

Li, W., C.F. Vogel and F. Matsumura. 2007. Studies on the cell treatment conditions to elicit lipolytic responses from 3T3-L1 adipocytes to TCDD, 2,3,7,8-tetrachlorodibenzo-p-dioxin. J. Cell Biochem. 102(2):389-402.

Li, W., C.F. Vogel, P. Fujiyoshi and F. Matsumura. 2008. Development of a human adipocyte model derived from human mesenchymal stem cells (hMSC) as a tool for toxicological studies on the action of TCDD. Biol. Chem. 389(2):169-177.

Lin, P., Y.C. Chang, C.H. Chen, W.J. Yang, Y.H. Cheng and L.W. Chang. 2004a. A comparative study on the effects of 2,3,7,8,-tetrachlorodibenzo-p-dioxin polychlorinated biphenyl126 and estrogen in human bronchial epithelial cells. Toxicol. Appl. Pharmacol. 195(1):83-91.

Lin, P., J.T. Chang, J.L. Ko, S.H. Liao and W.S. Lo. 2004b. Reduction of androgen receptor expression by benzo[alpha]pyrene and 7,8-dihydro-9,10-epoxy-7,8,9,10-tetrahydrobenzo[alpha]pyrene in human lung cells. Biochem. Pharmacol. 67(8):1523-1530.

Lin, T.M., N.T. Rasmussen, R.W. Moore, R.M. Albrecht and R.E. Peterson. 2004c. 2,3,7,8-tetrachlorodibenzo-p-dioxin inhibits prostatic epithelial bud formation by acting directly on the urogenital sinus. J. Urol. 172(1):365-368.

Lin, P.H., C.H. Lin, C.C. Huang, M.C. Chuang and P. Lin. 2007. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) induces oxidative stress, DNA strand breaks, and poly(ADP-ribose) polymerase-1 activation in human breast carcinoma cell lines. Toxicol. Lett. 172(3):146-158.

Lin, C.H., S.H. Juan, C.Y. Wang, Y.Y. Sun, C.M. Chou, S.F. Chang, S.Y. Hu, W.S. Lee and Y.H. Lee. 2008a. Neuronal activity enhances aryl hydrocarbon receptor-mediated gene expression and dioxin neurotoxicity in cortical neurons. J. Neurochem. 104(5):1415-1429.

Lin, P.H., C.H. Lin, C.C. Huang, J.P. Fang and M.C. Chuang. 2008b. 2,3,7,8-Tetrachlorodibenzo-p-Dioxin Modulates the Induction of DNA Strand Breaks and Poly(ADP-ribose) Polymerase-1 Activation by 17beta-Estradiol in Human Breast Carcinoma Cells Through Alteration of CYP1A1 and CYP1B1 Expression. Chem. Res. Toxicol. 21(7):1337-1347.

Link, B., T. Gabrio, I. Zoellner, I. Piechotowski, O. Paepke, T. Herrmann, A. Felder-Kennel, V. Maisner, K.H. Schick, M. Schrimpf, M. Schwenk and J. Wuthe. 2005. Biomonitoring of persistent organochlorine pesticides, PCDD/PCDFs and dioxin-like PCBs in blood of children from South West Germany (Baden-Wuerttemberg) from 1993 to 2003. Chemosphere. 58(9):1185-1201.

Lipponen, A., S. Lensu, J. Tuomisto, A. Yamatodani, J. Tarhanen and L. Tuomisto. 2004. The role of the chorda tympani nerve and histaminergic system in TCDD toxicity. Inflamm. Res. 53 Suppl 1:S59-S60.

Liu, P.C. and F. Matsumura. 2006. TCDD suppresses insulin-responsive glucose transporter (GLUT-4) gene expression through C/EBP nuclear transcription factors in 3T3-L1 adipocytes. J. Biochem. Mol. Toxicol. 20(2):79-87.

Liu, X. and C. Jefcoate. 2006. 2,3,7,8-tetrachlorodibenzo-p-dioxin and epidermal growth factor cooperatively suppress peroxisome proliferator-activated receptor-gamma1 stimulation and restore focal adhesion complexes during adipogenesis: selective contributions of Src, Rho, and Erk distinguish these overlapping processes in C3H10T1/2 cells. Mol. Pharmacol. 70(6):1902-1915.

Liu, Y.Q. and Y.K. Zhou. 2006. In vitro cytotoxicity of TCDD on SPC-A1 cells. Biomed. Environ. Sci. 19(1):21-26.

Liu, N., Q.Y. Zhang, D. Vakharia, D. Dunbar and L.S. Kaminsky. 2001. Induction of CYP1A by benzo[k]fluoranthene in human hepatocytes: CYP1A1 or CYP1A2? Arch. Biochem. Biophys. 389(1):130-134.

Liu, P.C., M.J. Moreno-Aliaga, D.Y. Dunlap, X.M. Hu, M.S. Denison and F. Matsumura. 2002. Correlation between the high expression of C/EBPbeta protein in F442A cells and their relative resistance to antiadipogenic action of TCDD in comparison to 3T3-L1 cells. J. Biochem. Mol. Toxicol. 16(2):70-83.

Liu, Y.R., N.J. Tang and D.L. Ren. 2003. [Induction of aryl hydrocarbon receptor and CYP1A1 mRNA by 2,3,7,8-tetrachlorodibenzo-p-dioxin in rat liver]. Zhonghua Lao. Dong. Wei Sheng Zhi. Ye. Bing. Za Zhi. 21(6):417-419.

Liu, S., M. Abdelrahim, S. Khan, E. Ariazi, V.C. Jordan and S. Safe. 2006. Aryl hydrocarbon receptor agonists directly activate estrogen receptor alpha in MCF-7 breast cancer cells. Biol. Chem. 387(9):1209-1213.

Loertscher, J.A., C.A. Sattler and B.L. Allen-Hoffmann. 2001a. 2,3,7,8-Tetrachlorodibenzo-p-dioxin alters the differentiation pattern of human keratinocytes in organotypic culture. Toxicol. Appl. Pharmacol. 175(2):121-129.

Loertscher, J.A., C.S. Sadek and B.L. Allen-Hoffmann. 2001b. Treatment of normal human keratinocytes with 2,3,7,8-tetrachlorodibenzo-p-dioxin causes a reduction in cell number, but no increase in apoptosis. Toxicol. Appl. Pharmacol. 175(2):114-120.

Lu, F., M. Zahid, M. Saeed, E.L. Cavalieri and E.G. Rogan. 2007. Estrogen metabolism and formation of estrogen-DNA adducts in estradiol-treated MCF-10F cells. The effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin induction and catechol-O-methyltransferase inhibition. J. Steroid Biochem. Mol. Biol. 105(1-5):150-158.

Luco, J.M., J. Galvez, R. Garcia-Domenech and J.V. de Julian-Ortiz. 2004. Structural invariants for the prediction of relative toxicities of polychloro dibenzo-p-dioxins and dibenzofurans. Mol. Divers. 8(4):331-342.

Luebke, R.W., C.B. Copeland and D.L. Andrews. 2000. Aging and resistance to Trichinella spiralis infection following xenobiotic exposure. Ann. N. Y. Acad. Sci. 919:221-229.

Ma, Q. 2002. Induction and superinduction of 2,3,7,8-tetrachlorodibenzo-rho-dioxin-inducible poly(ADP-ribose) polymerase: role of the aryl hydrocarbon receptor/aryl hydrocarbon receptor nuclear translocator transcription activation domains and a labile transcription repressor. Arch. Biochem. Biophys. 404(2):309-316.

Ma, Q. 2007. Aryl hydrocarbon receptor degradation-promoting factor (ADPF) and the control of the xenobiotic response. Mol. Interv. 7(3):133-137.

Ma, Q. and K.T. Baldwin. 2000. 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced degradation of aryl hydrocarbon receptor (AhR) by the ubiquitin-proteasome pathway. Role of the transcription activaton and DNA binding of AhR. J. Biol. Chem. 275(12):8432-8438.

Ma, Q. and K.T. Baldwin. 2002. A cycloheximide-sensitive factor regulates TCDD-induced degradation of the aryl hydrocarbon receptor. Chemosphere. 46(9-10):1491-1500.

Ma, Q., A.J. Renzelli, K.T. Baldwin and J.M. Antonini. 2000. Superinduction of CYP1A1 gene expression. Regulation of 2,3,7, 8-tetrachlorodibenzo-p-dioxin-induced degradation of Ah receptor by cycloheximide. J. Biol. Chem. 275(17):12676-12683.

Ma, Q., K. Kinneer, Y. Bi, J.Y. Chan and Y.W. Kan. 2004. Induction of murine NAD(P)H:quinone oxidoreductase by 2,3,7,8-tetrachlorodibenzo-p-dioxin requires the CNC (cap 'n' collar) basic leucine zipper transcription factor Nrf2 (nuclear factor erythroid 2-related factor 2): cross-interaction between AhR (aryl hydrocarbon receptor) and Nrf2 signal transduction. Biochem. J. 377(Pt 1):205-213.

MacDonald, C.J., H.P. Ciolino and G.C. Yeh. 2004. The drug salicylamide is an antagonist of the aryl hydrocarbon receptor that inhibits signal transduction induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin. Cancer Res. 64(1):429-434.

Machemer, D.E. and R.H. Tukey. 2005. The role of protein kinase C in regulation of TCDD-mediated CYP1A1 gene expression. Toxicol. Sci. 87(1):27-37.

Maher, J.M., X. Cheng, A.L. Slitt, M.Z. Dieter and C.D. Klaassen. 2005. Induction of the multidrug resistance-associated protein family of transporters by chemical activators of receptor-mediated pathways in mouse liver. Drug Metab Dispos. 33(7):956-962.

Maier, A., T.P. Dalton and A. Puga. 2000. Disruption of dioxin-inducible phase I and phase II gene expression patterns by cadmium, chromium, and arsenic. Mol. Carcinog. 28(4):225-235.

Maines, M.D. 2003. Bile pigments: newcomers to the cell signaling arena. Toxicol. Sci. 71(1):9-10.

Majora, M., M. Frericks, V. Temchura, G. Reichmann and C. Esser. 2005. Detection of a novel population of fetal thymocytes characterized by preferential emigration and a TCRgammadelta+ T cell fate after dioxin exposure. Int. Immunopharmacol. 5(12):1659-1674.

Mandal, P.K. 2005. Dioxin: a review of its environmental effects and its aryl hydrocarbon receptor biology. J. Comp Physiol [B]. 175(4):221-230.

Mandal, P.K., L.R. McDaniel, R.A. Prough and B.J. Clark. 2001. 7,12-Dimethylbenz[a]anthracene inhibition of steroid production in MA-10 mouse Leydig tumor cells is not directly linked to induction of CYP1B1. Toxicol. Appl. Pharmacol. 175(3):200-208.

Marchand, A., R. Barouki and M. Garlatti. 2004. Regulation of NAD(P)H:quinone oxidoreductase 1 gene expression by CYP1A1 activity. Mol. Pharmacol. 65(4):1029-1037.

Marchand, A., C. Tomkiewicz, J.P. Marchandeau, E. Boitier, R. Barouki and M. Garlatti. 2005. 2,3,7,8-Tetrachlorodibenzo-p-dioxin induces insulin-like growth factor binding protein-1 gene expression and counteracts the negative effect of insulin. Mol. Pharmacol. 67(2):444-452.

Marlowe, J.L., E.S. Knudsen, S. Schwemberger and A. Puga. 2004. The aryl hydrocarbon receptor displaces p300 from E2F-dependent promoters and represses S phase-specific gene expression. J. Biol. Chem. 279(28):29013-29022.

Marquez-Bravo, L.G. and J.F. Gierthy. 2008. Differential expression of estrogen receptor alpha (ERalpha) protein in MCF-7 breast cancer cells chronically exposed to TCDD. J. Cell Biochem. 103(2):636-647.

Marshall, N.B., W.R. Vorachek, L.B. Steppan, D.V. Mourich and N.I. Kerkvliet. 2008. Functional characterization and gene expression analysis of CD4+ CD25+ regulatory T cells generated in mice treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin. J. Immunol. 181(4):2382-2391.

Martin, R., J. O'Shea, L.S. Birnbaum and R. Luebke. 2008. Community corner. Striking the balance in multiple sclerosis. Nat. Med. 14(5):491.

Martinez, J.M., C.A. Afshari, P.R. Bushel, A. Masuda, T. Takahashi and N.J. Walker. 2002. Differential toxicogenomic responses to 2,3,7,8-tetrachlorodibenzo-p-dioxin in malignant and nonmalignant human airway epithelial cells. Toxicol. Sci. 69(2):409-423.

Martinez, J.M., S.J. Baek, D.M. Mays, P.K. Tithof, T.E. Eling and N.J. Walker. 2004. EGR1 is a novel target for AhR agonists in human lung epithelial cells. Toxicol. Sci. 82(2):429-435.

Maruyama, W. and Y. Aoki. 2006. Estimated cancer risk of dioxins to humans using a bioassay and physiologically based pharmacokinetic model. Toxicol. Appl. Pharmacol. 214(2):188-198.

Maruyama, W., K. Yoshida, T. Tanaka and J. Nakanishi. 2002. Determination of tissue-blood partition coefficients for a physiological model for humans, and estimation of dioxin concentration in tissues. Chemosphere. 46(7):975-985.

Massaad, C., F. Entezami, L. Massade, M. Benahmed, F. Olivennes, R. Barouki and S. Hamamah. 2002. How can chemical compounds alter human fertility? Eur. J. Obstet. Gynecol. Reprod. Biol. 100(2):127-137.

Massart, F., R. Parrino, P. Seppia, G. Federico and G. Saggese. 2006. How do environmental estrogen disruptors induce precocious puberty? Minerva Pediatr. 58(3):247-254.

Mastin, J.P. 2005. Environmental cardiovascular disease. Cardiovasc. Toxicol. 5(2):91-94.

Matikainen, T., G.I. Perez, A. Jurisicova, J.K. Pru, J.J. Schlezinger, H.Y. Ryu, J. Laine, T. Sakai, S.J. Korsmeyer, R.F. Casper, D.H. Sherr and J.L. Tilly. 2001. Aromatic hydrocarbon receptor-driven Bax gene expression is required for premature ovarian failure caused by biohazardous environmental chemicals. Nat. Genet. 28(4):355-360.

Matsumura, F. and C.F. Vogel. 2006. Evidence supporting the hypothesis that one of the main functions of the aryl hydrocarbon receptor is mediation of cell stress responses. Biol. Chem. 387(9):1189-1194.

Matthews, J. and J.A. Gustafsson. 2006. Estrogen receptor and aryl hydrocarbon receptor signaling pathways. Nucl. Recept. Signal. 4:e016.

Matthews, M., I. Heimler, M. Fahy, E. Radwanska, R. Hutz, A. Trewin and R. Rawlins. 2001. Effects of dioxin, an environmental pollutant, on mouse blastocyst development and apoptosis. Fertil. Steril. 75(6):1159-1162.

Matthews, J., B. Wihlen, J. Thomsen and J.A. Gustafsson. 2005. Aryl hydrocarbon receptor-mediated transcription: ligand-dependent recruitment of estrogen receptor alpha to 2,3,7,8-tetrachlorodibenzo-p-dioxin-responsive promoters. Mol. Cell Biol. 25(13):5317-5328.

Mazina, O., S. Park, H. Sano, P. Wong and F. Matsumura. 2004. Studies on the mechanism of rapid activation of protein tyrosine phosphorylation activities, particularly c-Src kinase, by TCDD in MCF10A. J. Biochem. Mol. Toxicol. 18(6):313-321.

McGarry, M.A., G.D. Charles, T. Medrano, M.R. Bubb, M.B. Grant, M. Campbell-Thompson and K.T. Shiverick. 2002. Benzo(a)pyrene, but not 2,3,7,8-tetrachlorodibenzo-p-dioxin, alters cell adhesion proteins in human uterine RL95-2 cells. Biochem. Biophys. Res. Commun. 294(1):101-107.

McHale, C.M., L. Zhang, A.E. Hubbard, X. Zhao, A. Baccarelli, A.C. Pesatori, M.T. Smith and M.T. Landi. 2007. Microarray analysis of gene expression in peripheral blood mononuclear cells from dioxin-exposed human subjects. Toxicology. 229(1-2):101-113.

McMillan, B.J. and C.A. Bradfield. 2007. The aryl hydrocarbon receptor sans xenobiotics: endogenous function in genetic model systems. Mol. Pharmacol. 72(3):487-498.

McMillan, B.J., S.N. McMillan, E. Glover and C.A. Bradfield. 2007. 2,3,7,8-Tetrachlorodibenzo-p-dioxin induces premature activation of the KLF2 regulon during thymocyte development. J. Biol. Chem. 282(17):12590-12597.

Mead, M.N. 2008. Cancer and TCDD: the mitochondrial connection. Environ. Health Perspect. 116(3):A112.

Miao, W., L. Hu, P.J. Scrivens and G. Batist. 2005. Transcriptional regulation of NF-E2 p45-related factor (NRF2) expression by the aryl hydrocarbon receptor-xenobiotic response element signaling pathway: direct cross-talk between phase I and II drug-metabolizing enzymes. J. Biol. Chem. 280(21):20340-20348.

Mimura, J. and Y. Fujii-Kuriyama. 2000. [Transcriptional regulation by Ah (dioxin) receptor]. Tanpakushitsu Kakusan Koso. 45(9 Suppl):1526-1533.

Mimura, J. and Y. Fujii-Kuriyama. 2003. Functional role of AhR in the expression of toxic effects by TCDD. Biochim. Biophys. Acta. 1619(3):263-268.

Minami, K., M. Nakajima, Y. Fujiki, M. Katoh, F.J. Gonzalez and T. Yokoi. 2008. Regulation of insulin-like growth factor binding protein-1 and lipoprotein lipase by the aryl hydrocarbon receptor. J. Toxicol. Sci. 33(4):405-413.

Minegishi, T., T. Hirakawa, K. Abe, H. Kishi and K. Miyamoto. 2003. Effect of IGF-1 and 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on the expression of LH receptors during cell differentiation in cultured granulosa cells. Mol. Cell Endocrinol. 202(1-2):123-131.

Minegishi, T., T. Hirakawa, K. Abe, H. Kishi and K. Miyamoto. 2004. Effect of insulin-like growth factor-1 and 2,3,7,8-tetrachlorodibenzo-p-dioxin on the expression of luteinizing hormone receptors in cultured granulosa cells. Environ. Sci. 11(1):57-71.

Miniero, R., E. De Felip, F. Ferri and A. di Domenico. 2001. An overview of TCDD half-life in mammals and its correlation to body weight. Chemosphere. 43(4-7):839-844.

Minsavage, G.D., D.P. Vorojeikina and T.A. Gasiewicz. 2003. Mutational analysis of the mouse aryl hydrocarbon receptor tyrosine residues necessary for recognition of dioxin response elements. Arch. Biochem. Biophys. 412(1):95-105.

Minsavage, G.D., S.K. Park and T.A. Gasiewicz. 2004. The aryl hydrocarbon receptor (AhR) tyrosine 9, a residue that is essential for AhR DNA binding activity, is not a phosphoresidue but augments AhR phosphorylation. J. Biol. Chem. 279(20):20582-20593.

Mitsushima, D., T. Funabashi and F. Kimura. 2003. Estrogen increases messenger RNA and immunoreactivity of aryl-hydrocarbon receptor nuclear translocator 2 in the rat mediobasal hypothalamus. Biochem. Biophys. Res. Commun. 307(2):248-253.

Miyabara, Y. and J. Yonemoto. 2000. [Polychlorinated biphenyl(PCBs, PCDDs, PCDFs)]. Nippon Rinsho. 58(12):2422-2427.

Miyamoto, K. 2004. Effects of dioxin on gene expression in female reproductive system in the rat. Environ. Sci. 11(1):47-55.

Miyazaki, W., T. Iwasaki, A. Takeshita, C. Tohyama and N. Koibuchi. 2008. Identification of the functional domain of thyroid hormone receptor responsible for polychlorinated biphenyl-mediated suppression of its action in vitro. Environ. Health Perspect. 116(9):1231-1236.

Mizutani, T., M. Yoshino, T. Satake, M. Nakagawa, R. Ishimura, C. Tohyama, K. Kokame, K. Kangawa and K. Miyamoto. 2004. Identification of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-inducible and -suppressive genes in the rat placenta: induction of interferon-regulated genes with possible inhibitory roles for angiogenesis in the placenta. Endocr. J. 51(6):569-577.

Moffat, I.D., P.C. Boutros, T. Celius, J. Linden, R. Pohjanvirta and A.B. Okey. 2007a. microRNAs in adult rodent liver are refractory to dioxin treatment. Toxicol. Sci. 99(2):470-487.

Moffat, I.D., S. Roblin, P.A. Harper, A.B. Okey and R. Pohjanvirta. 2007b. Aryl hydrocarbon receptor splice variants in the dioxin-resistant rat: tissue expression and transactivational activity. Mol. Pharmacol. 72(4):956-966.

Mollerup, S., S. OVrebo and A. Haugen. 2001. Lung carcinogenesis: resveratrol modulates the expression of genes involved in the metabolism of PAH in human bronchial epithelial cells. Int. J. Cancer. 92(1):18-25.

Monk, S.A., M.S. Denison and R.H. Rice. 2001. Transient expression of CYP1A1 in rat epithelial cells cultured in suspension. Arch. Biochem. Biophys. 393(1):154-162.

Monk, S.A., M.S. Denison and R.H. Rice. 2003. Reversible stepwise negative regulation of CYP1A1 in cultured rat epidermal cells. Arch. Biochem. Biophys. 419(2):158-169.

Monteiro, P., D. Gilot, E. LeFerrec, C. Rauch, D. Lagadic-Gossmann and O. Fardel. 2008a. Dioxin-mediated up-regulation of aryl hydrocarbon receptor target genes is dependent on the calcium/calmodulin/CaMKIalpha pathway. Mol. Pharmacol. 73(3):769-777.

Monteiro, P., D. Gilot, S. Langouet and O. Fardel. 2008b. Activation of the aryl hydrocarbon receptor by the calcium/calmodulin-dependent protein kinase kinase inhibitor STO-609. Drug Metab Dispos.

Moran, F.M., A.J. Conley, C.J. Corbin, E. Enan, C. VandeVoort, J.W. Overstreet and B.L. Lasley. 2000. 2,3,7,8-tetrachlorodibenzo-p-dioxin decreases estradiol production without altering the enzyme activity of cytochrome P450 aromatase of human luteinized granulosa cells in vitro. Biol. Reprod. 62(4):1102-1108.

Moran, F.M., C.A. VandeVoort, J.W. Overstreet, B.L. Lasley and A.J. Conley. 2003a. Molecular target of endocrine disruption in human luteinizing granulosa cells by 2,3,7,8-tetrachlorodibenzo-p-dioxin: inhibition of estradiol secretion due to decreased 17alpha-hydroxylase/17,20-lyase cytochrome P450 expression. Endocrinology. 144(2):467-473.

Moran, F.M., P. Lohstroh, C.A. VandeVoort, J. Chen, J.W. Overstreet, A.J. Conley and B.L. Lasley. 2003b. Exogenous steroid substrate modifies the effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on estradiol production of human luteinized granulosa cells in vitro. Biol. Reprod. 68(1):244-251.

Mori, C., B. Morsey, M. Levin, P.R. Nambiar and S. De Guise. 2006. Immunomodulatory effects of in vitro exposure to organochlorines on T-cell proliferation in marine mammals and mice. J. Toxicol. Environ. Health A. 69(3-4):283-302.

Mori, C., B. Morsey, M. Levin, T.S. Gorton and S. De Guise. 2008. Effects of organochlorines, individually and in mixtures, on B-cell proliferation in marine mammals and mice. J. Toxicol. Environ. Health A. 71(4):266-275.

Morita, K. and T. Nakano. 2002. Seaweed accelerates the excretion of dioxin stored in rats. J. Agric. Food Chem. 50(4):910-917.

Morita, K. and K. Tobiishi. 2002. Increasing effect of nori on the fecal excretion of dioxin by rats. Biosci. Biotechnol. Biochem. 66(11):2306-2313.

Morita, K., M. Ogata and T. Hasegawa. 2001. Chlorophyll derived from Chlorella inhibits dioxin absorption from the gastrointestinal tract and accelerates dioxin excretion in rats. Environ. Health Perspect. 109(3):289-294.

Morrow, D., C. Qin, R. Smith, III and S. Safe. 2004. Aryl hydrocarbon receptor-mediated inhibition of LNCaP prostate cancer cell growth and hormone-induced transactivation. J. Steroid Biochem. Mol. Biol. 88(1):27-36.

Moser, G.A. and M.S. McLachlan. 2001. The influence of dietary concentration on the absorption and excretion of persistent lipophilic organic pollutants in the human intestinal tract. Chemosphere. 45(2):201-211.

Moser, G.A. and M.S. McLachlan. 2002. Modeling digestive tract absorption and desorption of lipophilic organic contaminants in humans. Environ. Sci. Technol. 36(15):3318-3325.

Mueller, M.D., J.L. Vigne, M. Streich, M.K. Tee, L. Raio, E. Dreher, N.A. Bersinger and R.N. Taylor. 2005. 2,3,7,8-Tetrachlorodibenzo-p-dioxin increases glycodelin gene and protein expression in human endometrium. J. Clin. Endocrinol. Metab. 90(8):4809-4815.

Mukai, R., I. Fukuda, S. Nishiumi, K. Hosokawa, K. Kanazawa and H. Ashida. 2004. Anthocyan does not suppress transformation of aryl hydrocarbon receptor induced by dioxin. Biofactors. 21(1-4):371-373.

Mukherjee, S., B.C. Koner, S. Ray and A. Ray. 2006. Environmental contaminants in pathogenesis of breast cancer. Indian J. Exp. Biol. 44(8):597-617.

Muller, G.F., O. Dohr, C. El-Bahay, R. Kahl and J. Abel. 2000. Effect of transforming growth factor-beta1 on cytochrome P450 expression: inhibition of CYP1 mRNA and protein expression in primary rat hepatocytes. Arch. Toxicol. 74(3):145-152.

Mullerova, D. and J. Kopecky. 2007. White adipose tissue: storage and effector site for environmental pollutants. Physiol Res. 56(4):375-381.

Munzel, P.A., S. Schmohl, F. Buckler, J. Jaehrling, F.T. Raschko, C. Kohle and K.W. Bock. 2003. Contribution of the Ah receptor to the phenolic antioxidant-mediated expression of human and rat UDP-glucuronosyltransferase UGT1A6 in Caco-2 and rat hepatoma 5L cells. Biochem. Pharmacol. 66(5):841-847.

Murayama, T., M. Inoue, T. Nomura, S. Mori and Y. Eizuru. 2002. 2,3,7,8-Tetrachlorodibenzo-p-dioxin is a possible activator of human cytomegalovirus replication in a human fibroblast cell line. Biochem. Biophys. Res. Commun. 296(3):651-656.

Murphy, K.A., C.M. Villano, R. Dorn and L.A. White. 2004. Interaction between the aryl hydrocarbon receptor and retinoic acid pathways increases matrix metalloproteinase-1 expression in keratinocytes. J. Biol. Chem. 279(24):25284-25293.

Murray, I.A., R.K. Reen, N. Leathery, P. Ramadoss, L. Bonati, F.J. Gonzalez, J.M. Peters and G.H. Perdew. 2005. Evidence that ligand binding is a key determinant of Ah receptor-mediated transcriptional activity. Arch. Biochem. Biophys. 442(1):59-71.

Mutoh, J., J. Taketoh, K. Okamura, T. Kagawa, T. Ishida, Y. Ishii and H. Yamada. 2006. Fetal pituitary gonadotropin as an initial target of dioxin in its impairment of cholesterol transportation and steroidogenesis in rats. Endocrinology. 147(2):927-936.

N'Diaye, M., E. LeFerrec, D. Lagadic-Gossmann, S. Corre, D. Gilot, V. Lecureur, P. Monteiro, C. Rauch, M.D. Galibert and O. Fardel. 2006. Aryl hydrocarbon receptor- and calcium-dependent induction of the chemokine CCL1 by the environmental contaminant benzo[a]pyrene. J. Biol. Chem. 281(29):19906-19915.

N'Jai, A., D.R. Boverhof, E. Dere, L.D. Burgoon, Y.S. Tan, J.C. Rowlands, R.A. Budinsky, K.E. Stebbins and T.R. Zacharewski. 2008. Comparative temporal toxicogenomic analysis of TCDD- and TCDF-mediated hepatic effects in immature female C57BL/6 mice. Toxicol. Sci. 103(2):285-297.

Nagai, H., T. Takei, C. Tohyama, M. Kubo, R. Abe and K. Nohara. 2005. Search for the target genes involved in the suppression of antibody production by TCDD in C57BL/6 mice. Int. Immunopharmacol. 5(2):331-343.

Nagashima, H. and F. Matsumura. 2002. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)-induced down-regulation of glucose transporting activities in mouse 3T3-L1 preadipocyte. J. Environ. Sci. Health B. 37(1):1-14.

Nakajima, M., M. Iwanari and T. Yokoi. 2003. Effects of histone deacetylation and DNA methylation on the constitutive and TCDD-inducible expressions of the human CYP1 family in MCF-7 and HeLa cells. Toxicol. Lett. 144(2):247-256.

Nakano, S., T. Noguchi, H. Takekoshi, G. Suzuki and M. Nakano. 2005. Maternal-fetal distribution and transfer of dioxins in pregnant women in Japan, and attempts to reduce maternal transfer with Chlorella (Chlorella pyrenoidosa) supplements. Chemosphere. 61(9):1244-1255.

Natsume, Y., H. Satsu, K. Kitamura, N. Okamoto and M. Shimizu. 2004. Assessment system for dioxin absorption in the small intestine and prevention of its absorption by food factors. Biofactors. 21(1-4):375-377.

Nayyar, T., N.H. Zawia and D.B. Hood. 2002. Transplacental effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the temporal modulation of Sp1 DNA binding in the developing cerebral cortex and cerebellum. Exp. Toxicol. Pathol. 53(6):461-468.

Nayyar, T., J. Wu and D.B. Hood. 2003. Downregulation of hippocampal NMDA receptor expression by prenatal exposure to dioxin. Cell Mol. Biol. (Noisy. -le-grand). 49(8):1357-1362.

Nazarenko, D.A., S.D. Dertinger and T.A. Gasiewicz. 2001. In vivo antagonism of AhR-mediated gene induction by 3'-methoxy-4'-nitroflavone in TCDD-responsive lacZ mice. Toxicol. Sci. 61(2):256-264.

Ndebele, K., P.B. Tchounwou and R.W. McMurray. 2004. Coumestrol, bisphenol-A, DDT, and TCDD modulation of interleukin-2 expression in activated CD+4 Jurkat T cells. Int. J. Environ. Res. Public Health. 1(1):3-11.

Nebert, D.W. and C.L. Karp. 2008. Endogenous functions of the Aryl hydrocarbon receptor: Intersection of cytochromeP450 (CYP1)-metabolized eicosanoids and AHR biology. J. Biol. Chem.

Neri, T., V. Merico, S. Garagna, C.A. Redi and M. Zuccotti. 2008. Expression of phase I and phase II genes in mouse embryonic stem cells cultured in the presence of 2,3,7,8-tetrachlorodibenzo-para-dioxin. Biochim. Biophys. Acta. 1780(5):826-836.

Niermann, T., S. Schmutz, P. Erne and T. Resink. 2003. Aryl hydrocarbon receptor ligands repress T-cadherin expression in vascular smooth muscle cells. Biochem. Biophys. Res. Commun. 300(4):943-949.

Nishiumi, S., Y. Yabushita, I. Fukuda, R. Mukai, K. Yoshida and H. Ashida. 2006. Molokhia (Corchorus olitorius L.) extract suppresses transformation of the aryl hydrocarbon receptor induced by dioxins. Food Chem. Toxicol. 44(2):250-260.

Nohara, K., X. Pan, S. Tsukumo, A. Hida, T. Ito, H. Nagai, K. Inouye, H. Motohashi, M. Yamamoto, Y. Fujii-Kuriyama and C. Tohyama. 2005. Constitutively active aryl hydrocarbon receptor expressed specifically in T-lineage cells causes thymus involution and suppresses the immunization-induced increase in splenocytes. J. Immunol. 174(5):2770-2777.

Nohara, K., K. Ao, Y. Miyamoto, T. Ito, T. Suzuki, H. Toyoshiba and C. Tohyama. 2006. Comparison of the 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-induced CYP1A1 gene expression profile in lymphocytes from mice, rats, and humans: most potent induction in humans. Toxicology. 225(2-3):204-213.

North, C.M., B.S. Kim, N. Snyder, R.B. Crawford, M.P. Holsapple and N.E. Kaminski. 2008. TCDD-Mediated Suppression of the in vitro Anti-Sheep Erythrocyte IgM Antibody Forming Cell Response is Reversed by Interferon Gamma. Toxicol. Sci.

Novak, J., M. Benisek and K. Hilscherova. 2008. Disruption of retinoid transport, metabolism and signaling by environmental pollutants. Environ. Int. 34(6):898-913.

Nowak, E., G. Wyrwicz, Z. Dabrowski, O. Smolenski and K. Spodaryk. 2002. Rheological and biochemical examination of red blood cells after treatment with different pharmacological and toxic substances. Clin. Hemorheol. Microcirc. 26(2):91-97.

Oberemm, A., C. Meckert, L. Brandenburger, A. Herzig, Y. Lindner, K. Kalenberg, E. Krause, C. Ittrich, A. Kopp-Schneider, R. Stahlmann, H.B. Richter-Reichhelm and U. Gundert-Remy. 2005. Differential signatures of protein expression in marmoset liver and thymus induced by single-dose TCDD treatment. Toxicology. 206(1):33-48.

Oenga, G.N., D.C. Spink and D.O. Carpenter. 2004. TCDD and PCBs inhibit breast cancer cell proliferation in vitro. Toxicol. In Vitro. 18(6):811-819.

Oesch-Bartlomowicz, B., K. Janssen, O. Wiss and F. Oesch. 2004. Guanine 6-O-methylation pattern within the dioxin responsive element of the CYP1A1 enhancer shows two critical guanines for AhR/ARNT binding. Chem. Biodivers. 1(3):473-480.

Oesch-Bartlomowicz, B., A. Huelster, O. Wiss, P. Antoniou-Lipfert, C. Dietrich, M. Arand, C. Weiss, E. Bockamp and F. Oesch. 2005. Aryl hydrocarbon receptor activation by cAMP vs. dioxin: divergent signaling pathways. Proc. Natl. Acad. Sci. U. S. A. 102(26):9218-9223.

Ogawa, T., Y. Asai, M. Yamashita and T. Takasuga. 2003. Detectable dioxins in human saliva and their effects on gingival epithelial cells. J. Dent. Res. 82(10):849-853.

Ogiso, H., N. Kagi, E. Matsumoto, M. Nishimoto, R. Arai, M. Shirouzu, J. Mimura, Y. Fujii-Kuriyama and S. Yokoyama. 2004. Phosphorylation analysis of 90 kDa heat shock protein within the cytosolic arylhydrocarbon receptor complex. Biochemistry. 43(49):15510-15519.

Ohbayashi, T., K. Oikawa, R. Iwata, A. Kameta, K. Evine, T. Isobe, Y. Matsuda, J. Mimura, Y. Fujii-Kuriyama, M. Kuroda and K. Mukai. 2001. Dioxin induces a novel nuclear factor, DIF-3, that is implicated in spermatogenesis. FEBS Lett. 508(3):341-344.

Ohtake, F., K. Takeyama, T. Matsumoto, H. Kitagawa, Y. Yamamoto, K. Nohara, C. Tohyama, A. Krust, J. Mimura, P. Chambon, J. Yanagisawa, Y. Fujii-Kuriyama and S. Kato. 2003. Modulation of oestrogen receptor signalling by association with the activated dioxin receptor. Nature. 423(6939):545-550.

Ohtake, F., Y. Fujii-Kuriyama and S. Kato. 2008a. AhR acts as an E3 ubiquitin ligase to modulate steroid receptor functions. Biochem. Pharmacol.

Ohtake, F., A. Baba, Y. Fujii-Kuriyama and S. Kato. 2008b. Intrinsic AhR function underlies cross-talk of dioxins with sex hormone signalings. Biochem. Biophys. Res. Commun. 370(4):541-546.

Ohyama, K. 2004. [Disorders of sex differentiation caused by exogenous hormones]. Nippon Rinsho. 62(2):379-384.

Oikawa, K., T. Ohbayashi, J. Mimura, R. Iwata, A. Kameta, K. Evine, K. Iwaya, Y. Fujii-Kuriyama, M. Kuroda and K. Mukai. 2001. Dioxin suppresses the checkpoint protein, MAD2, by an aryl hydrocarbon receptor-independent pathway. Cancer Res. 61(15):5707-5709.

Oikawa, K., T. Ohbayashi, J. Mimura, Y. Fujii-Kuriyama, S. Teshima, K. Rokutan, K. Mukai and M. Kuroda. 2002. Dioxin stimulates synthesis and secretion of IgE-dependent histamine-releasing factor. Biochem. Biophys. Res. Commun. 290(3):984-987.

Oikawa, K., Y. Kosugi, T. Ohbayashi, A. Kameta, K. Isaka, M. Takayama, M. Kuroda and K. Mukai. 2003. Increased expression of IgE-dependent histamine-releasing factor in endometriotic implants. J. Pathol. 199(3):318-323.

Oikawa, K., K. Yoshida, M. Takanashi, H. Tanabe, T. Kiyuna, M. Ogura, A. Saito, A. Umezawa and M. Kuroda. 2008. Dioxin interferes in chromosomal positioning through the aryl hydrocarbon receptor. Biochem. Biophys. Res. Commun. 374(2):361-364.

Okey, A.B. 2007. An aryl hydrocarbon receptor odyssey to the shores of toxicology: the Deichmann Lecture, International Congress of Toxicology-XI. Toxicol. Sci. 98(1):5-38.

Okey, A.B., M.A. Franc, I.D. Moffat, N. Tijet, P.C. Boutros, M. Korkalainen, J. Tuomisto and R. Pohjanvirta. 2005. Toxicological implications of polymorphisms in receptors for xenobiotic chemicals: The case of the aryl hydrocarbon receptor. Toxicol. Appl. Pharmacol. 207(2 Suppl):43-51.

Okino, S.T., D. Pookot, L.C. Li, H. Zhao, S. Urakami, H. Shiina, M. Igawa and R. Dahiya. 2006. Epigenetic inactivation of the dioxin-responsive cytochrome P4501A1 gene in human prostate cancer. Cancer Res. 66(15):7420-7428.

Okino, S.T., L.C. Quattrochi, D. Pookot, M. Iwahashi and R. Dahiya. 2007. A dioxin-responsive enhancer 3' of the human CYP1A2 gene. Mol. Pharmacol. 72(6):1457-1465.

Oliver, J.D. and R.A. Roberts. 2002. Receptor-mediated hepatocarcinogenesis: role of hepatocyte proliferation and apoptosis. Pharmacol. Toxicol. 91(1):1-7.

Ovando, B.J., C.M. Vezina, B.P. McGarrigle and J.R. Olson. 2006. Hepatic gene downregulation following acute and subchronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. Toxicol. Sci. 94(2):428-438.

Palermo, C.M., J.I. Hernando, S.D. Dertinger, A.S. Kende and T.A. Gasiewicz. 2003. Identification of potential aryl hydrocarbon receptor antagonists in green tea. Chem. Res. Toxicol. 16(7):865-872.

Parfett, C.L. 2003. Combined effects of tumor promoters and serum on proliferin mRNA induction: a biomarker sensitive to saccharin, 2,3,7,8-TCDD, and other compounds at minimal concentrations promoting C3H/10T1/2 cell transformation. J. Toxicol. Environ. Health A. 66(20):1943-1966.

Park, E.Y. and H.M. Rho. 2002. The transcriptional activation of the human copper/zinc superoxide dismutase gene by 2,3,7,8-tetrachlorodibenzo-p-dioxin through two different regulator sites, the antioxidant responsive element and xenobiotic responsive element. Mol. Cell Biochem. 240(1-2):47-55.

Park, J.H. and S.W. Lee. 2002. Up-regulated expression of genes encoding Hrk and IL-3R beta subunit by TCDD in vivo and in vitro. Toxicol. Lett. 129(1-2):1-11.

Park, S. and F. Matsumura. 2006. Characterization of anti-apoptotic action of TCDD as a defensive cellular stress response reaction against the cell damaging action of ultra-violet irradiation in an immortalized normal human mammary epithelial cell line, MCF10A. Toxicology. 217(2-3):139-146.

Park, S., E.C. Henry and T.A. Gasiewicz. 2000. Regulation of DNA binding activity of the ligand-activated aryl hydrocarbon receptor by tyrosine phosphorylation. Arch. Biochem. Biophys. 381(2):302-312.

Park, J.H., S.W. Lee, I.T. Kim, B.S. Shin, S.W. Cheong, U.H. Cho, M.J. Huh and G.S. Oh. 2001. TCDD-up-regulation of IGFBP-6 and IL-5R alpha subunit genes in vivo and in vitro. Mol. Cells. 12(3):372-379.

Park, J.H., E.J. Hahn, J.H. Kong, H.J. Cho, C.S. Yoon, S.W. Cheong, G.S. Oh and H.J. Youn. 2003. TCDD-induced apoptosis in EL-4 cells deficient of the aryl hydrocarbon receptor and down-regulation of IGFBP-6 prevented the apoptotic cell death. Toxicol. Lett. 145(1):55-68.

Park, S., O. Mazina, A. Kitagawa, P. Wong and F. Matsumura. 2004a. TCDD causes suppression of growth and differentiation of MCF10A, human mammary epithelial cells by interfering with their insulin receptor signaling through c-Src kinase and ERK activation. J. Biochem. Mol. Toxicol. 18(6):322-331.

Park, Y.K., I. Fukuda, H. Ashida, S. Nishiumi, J.P. Guzman, H.H. Sato and G.M. Pastore. 2004b. Suppression of dioxin mediated aryl hydrocarbon receptor transformation by ethanolic extracts of propolis. Biosci. Biotechnol. Biochem. 68(4):935-938.

Park, S.J., W.K. Yoon, H.J. Kim, H.Y. Son, S.W. Cho, K.S. Jeong, T.H. Kim, S.H. Kim, S.R. Kim and S.Y. Ryu. 2005. 2,3,7,8-Tetrachlorodibenzo-p-dioxin activates ERK and p38 mitogen-activated protein kinases in RAW 264.7 cells. Anticancer Res. 25(4):2831-2836.

Park, S., B. Dong and F. Matsumura. 2007. Rapid activation of c-Src kinase by dioxin is mediated by the Cdc37-HSP90 complex as part of Ah receptor signaling in MCF10A cells. Biochemistry. 46(3):899-908.

Partanen, A.M., A. Kiukkonen, C. Sahlberg, S. Alaluusua, I. Thesleff, R. Pohjanvirta and P.L. Lukinmaa. 2004. Developmental toxicity of dioxin to mouse embryonic teeth in vitro: arrest of tooth morphogenesis involves stimulation of apoptotic program in the dental epithelium. Toxicol. Appl. Pharmacol. 194(1):24-33.

Partridge, N.C., G.J. Fiacco, H.W. Walling, O.Y. Barmina, J.J. Jeffrey and M.F. Ruh. 2000. Effects of dioxin and estrogen on collagenase-3 in UMR 106-01 osteosarcoma cells. Arch. Biochem. Biophys. 382(2):182-188.

Pastorelli, R., D. Carpi, R. Campagna, L. Airoldi, R. Pohjanvirta, M. Viluksela, H. Hakansson, P.C. Boutros, I.D. Moffat, A.B. Okey and R. Fanelli. 2006. Differential expression profiling of the hepatic proteome in a rat model of dioxin resistance: correlation with genomic and transcriptomic analyses. Mol. Cell Proteomics. 5(5):882-894.

Patel, R.D., D.J. Kim, J.M. Peters and G.H. Perdew. 2006. The aryl hydrocarbon receptor directly regulates expression of the potent mitogen epiregulin. Toxicol. Sci. 89(1):75-82.

Patel, R.D., B.D. Hollingshead, C.J. Omiecinski and G.H. Perdew. 2007. Aryl-hydrocarbon receptor activation regulates constitutive androstane receptor levels in murine and human liver. Hepatology. 46(1):209-218.

Patterson, R.M. and D.R. Germolec. 2006. Gene expression alterations in immune system pathways following exposure to immunosuppressive chemicals. Ann. N. Y. Acad. Sci. 1076:718-727.

Paustenbach, D. and D. Galbraith. 2006. Biomonitoring: is body burden relevant to public health? Regul. Toxicol. Pharmacol. 44(3):249-261.

Pesatori, A.C. and F. Forastiere. 2003. [Methodology in occupational and environmental epidemiology]. G. Ital. Med. Lav. Ergon. 25(3):290-291.

Pestka, J. and H.R. Zhou. 2006. Toll-like receptor priming sensitizes macrophages to proinflammatory cytokine gene induction by deoxynivalenol and other toxicants. Toxicol. Sci. 92(2):445-455.

Peter Guengerich, F., M.V. Martin, W.A. McCormick, L.P. Nguyen, E. Glover and C.A. Bradfield. 2004. Aryl hydrocarbon receptor response to indigoids in vitro and in vivo. Arch. Biochem. Biophys. 423(2):309-316.

Peters, A.K., J.T. Sanderson, A. Bergman and M. Van den Berg. 2006a. Antagonism of TCDD-induced ethoxyresorufin-O-deethylation activity by polybrominated diphenyl ethers (PBDEs) in primary cynomolgus monkey (Macaca fascicularis) hepatocytes. Toxicol. Lett. 164(2):123-132.

Peters, A.K., P.E. Leonards, B. Zhao, A. Bergman, M.S. Denison and M. Van den Berg. 2006b. Determination of in vitro relative potency (REP) values for mono-ortho polychlorinated biphenyls after purification with active charcoal. Toxicol. Lett. 165(3):230-241.

Petersen, S.L., M.A. Curran, S.A. Marconi, C.D. Carpenter, L.S. Lubbers and M.D. McAbee. 2000. Distribution of mRNAs encoding the arylhydrocarbon receptor, arylhydrocarbon receptor nuclear translocator, and arylhydrocarbon receptor nuclear translocator-2 in the rat brain and brainstem. J. Comp Neurol. 427(3):428-439.

Petersen, S.L., S. Krishnan and E.D. Hudgens. 2006. The aryl hydrocarbon receptor pathway and sexual differentiation of neuroendocrine functions. Endocrinology. 147(6 Suppl):S33-S42.

Petreas, M., D. Smith, S. Hurley, S.S. Jeffrey, D. Gilliss and P. Reynolds. 2004. Distribution of persistent, lipid-soluble chemicals in breast and abdominal adipose tissues: lessons learned from a breast cancer study. Cancer Epidemiol. Biomarkers Prev. 13(3):416-424.

Petrick, J.S. and C.D. Klaassen. 2007. Importance of hepatic induction of constitutive androstane receptor and other transcription factors that regulate xenobiotic metabolism and transport. Drug Metab Dispos. 35(10):1806-1815.

Petroff, B.K., K.F. Roby, X. Gao, D. Son, S. Williams, D. Johnson, K.K. Rozman and P.F. Terranova. 2001. A review of mechanisms controlling ovulation with implications for the anovulatory effects of polychlorinated dibenzo-p-dioxins in rodents. Toxicology. 158(3):91-107.

Petroff, B.K. and K.M. Mizinga. 2003. Pharmacokinetics of ovarian steroids in Sprague-Dawley rats after acute exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Reprod. Biol. 3(2):131-141.

Petrulis, J.R. and N.J. Bunce. 2000. Competitive behavior in the interactive toxicology of halogenated aromatic compounds. J. Biochem. Mol. Toxicol. 14(2):73-81.

Petrulis, J.R., G. Chen, S. Benn, J. LaMarre and N.J. Bunce. 2001. Application of the ethoxyresorufin-O-deethylase (EROD) assay to mixtures of halogenated aromatic compounds. Environ. Toxicol. 16(2):177-184.

Pflieger-Bruss, S. and W.B. Schill. 2000. Effects of chlorinated hydrocarbons on sperm function in vitro. Andrologia. 32(4-5):311-315.

Piaggi, S., M. Novelli, L. Martino, M. Masini, C. Raggi, E. Orciuolo, P. Masiello, A. Casini and V. De Tata. 2007. Cell death and impairment of glucose-stimulated insulin secretion induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the beta-cell line INS-1E. Toxicol. Appl. Pharmacol. 220(3):333-340.

Pieklo, R., A. Grochowalski and E.L. Gregoraszczuk. 2000. 2,3,7,8-Tetrachlorodibenzo-p-dioxin alters follicular steroidogenesis in time- and cell-specific manner. Exp. Clin. Endocrinol. Diabetes. 108(4):299-304.

Pitt, J.A., L. Feng, B.D. Abbott, J. Schmid, R.E. Batt, T.G. Costich, S.T. Koury and D.P. Bofinger. 2001. Expression of AhR and ARNT mRNA in cultured human endometrial explants exposed to TCDD. Toxicol. Sci. 62(2):289-298.

Pliskova, M., J. Vondracek, B. Vojtesek, A. Kozubik and M. Machala. 2005. Deregulation of cell proliferation by polycyclic aromatic hydrocarbons in human breast carcinoma MCF-7 cells reflects both genotoxic and nongenotoxic events. Toxicol. Sci. 83(2):246-256.

Pocar, P., T.A. Brevini, B. Fischer and F. Gandolfi. 2003. The impact of endocrine disruptors on oocyte competence. Reproduction. 125(3):313-325.

Pocar, P., B. Fischer, T. Klonisch and S. Hombach-Klonisch. 2005. Molecular interactions of the aryl hydrocarbon receptor and its biological and toxicological relevance for reproduction. Reproduction. 129(4):379-389.

Pocar, P., T. Klonisch, C. Brandsch, K. Eder, C. Frohlich, C. Hoang-Vu and S. Hombach-Klonisch. 2006. AhR-agonist-induced transcriptional changes of genes involved in thyroid function in primary porcine thyrocytes. Toxicol. Sci. 89(2):408-414.

Pohl, C., F. Will, H. Dietrich and D. Schrenk. 2006. Cytochrome P450 1A1 expression and activity in Caco-2 cells: modulation by apple juice extract and certain apple polyphenols. J. Agric. Food Chem. 54(26):10262-10268.

Pohl, H.R., H.G. Abadin, D.E. Jones and C.T. De Rosa. 2007. The role of exposure versus body burden data in deriving health guidance values. J. Toxicol. Environ. Health B Crit Rev. 10(6):401-415.

Pollenz, R.S. 2007. Specific blockage of ligand-induced degradation of the Ah receptor by proteasome but not calpain inhibitors in cell culture lines from different species. Biochem. Pharmacol. 74(1):131-143.

Pollenz, R.S. and E.R. Barbour. 2000. Analysis of the complex relationship between nuclear export and aryl hydrocarbon receptor-mediated gene regulation. Mol. Cell Biol. 20(16):6095-6104.

Pollenz, R.S. and E.J. Dougherty. 2005. Redefining the role of the endogenous XAP2 and C-terminal hsp70-interacting protein on the endogenous Ah receptors expressed in mouse and rat cell lines. J. Biol. Chem. 280(39):33346-33356.

Pollenz, R.S., J. Popat and E.J. Dougherty. 2005. Role of the carboxy-terminal transactivation domain and active transcription in the ligand-induced and ligand-independent degradation of the mouse Ahb-1 receptor. Biochem. Pharmacol. 70(11):1623-1633.

Porter, W., F. Wang, R. Duan, C. Qin, E. Castro-Rivera, K. Kim and S. Safe. 2001. Transcriptional activation of heat shock protein 27 gene expression by 17beta-estradiol and modulation by antiestrogens and aryl hydrocarbon receptor agonists. J. Mol. Endocrinol. 26(1):31-42.

Portier, C. 2000. Risk ranges for various endpoints following exposure to 2,3,7,8-TCDD. Food Addit. Contam. 17(4):335-346.

Portier, C.J. 2001. Linking toxicology and epidemiology: the role of mechanistic modelling. Stat. Med. 20(9-10):1387-1393.

Portier, C.J. 2002. Endocrine dismodulation and cancer. Neuro. Endocrinol. Lett. 23 Suppl 2:43-47.

Procopio, M., A. Lahm, A. Tramontano, L. Bonati and D. Pitea. 2002. A model for recognition of polychlorinated dibenzo-p-dioxins by the aryl hydrocarbon receptor. Eur. J. Biochem. 269(1):13-18.

Przyrembel, H., B. Heinrich-Hirsch and B. Vieth. 2000. Exposition to and health effects of residues in human milk. Adv. Exp. Med. Biol. 478:307-325.

Puga, A., M.A. Sartor, M.Y. Huang, J.K. Kerzee, Y.D. Wei, C.R. Tomlinson, C.S. Baxter and M. Medvedovic. 2004. Gene expression profiles of mouse aorta and cultured vascular smooth muscle cells differ widely, yet show common responses to dioxin exposure. Cardiovasc. Toxicol. 4(4):385-404.

Puga, A., C.R. Tomlinson and Y. Xia. 2005. Ah receptor signals cross-talk with multiple developmental pathways. Biochem. Pharmacol. 69(2):199-207.

Puga, A., C. Ma and J.L. Marlowe. 2008. The aryl hydrocarbon receptor cross-talks with multiple signal transduction pathways. Biochem. Pharmacol.

Qu, X., R.P. Metz, W.W. Porter, V.M. Cassone and D.J. Earnest. 2007. Disruption of clock gene expression alters responses of the aryl hydrocarbon receptor signaling pathway in the mouse mammary gland. Mol. Pharmacol. 72(5):1349-1358.

Quinn, A.M. and T.M. Penning. 2008. Comparisons of (+/-)-benzo[a]pyrene-trans-7,8-dihydrodiol activation by human cytochrome P450 and aldo-keto reductase enzymes: effect of redox state and expression levels. Chem. Res. Toxicol. 21(5):1086-1094.

Quintana, F.J., A.S. Basso, A.H. Iglesias, T. Korn, M.F. Farez, E. Bettelli, M. Caccamo, M. Oukka and H.L. Weiner. 2008. Control of T(reg) and T(H)17 cell differentiation by the aryl hydrocarbon receptor. Nature. 453(7191):65-71.

Racky, J., H.J. Schmitz, H.M. Kauffmann and D. Schrenk. 2004. Single nucleotide polymorphism analysis and functional characterization of the human Ah receptor (AhR) gene promoter. Arch. Biochem. Biophys. 421(1):91-98.

Rajkovic, A., M. Uyttendaele and J. Debevere. 2007. Computer aided boar semen motility analysis for cereulide detection in different food matrices. Int. J. Food Microbiol. 114(1):92-99.

Ramadoss, P. and G.H. Perdew. 2005. The transactivation domain of the Ah receptor is a key determinant of cellular localization and ligand-independent nucleocytoplasmic shuttling properties. Biochemistry. 44(33):11148-11159.

Ramadoss, P., J.R. Petrulis, B.D. Hollingshead, A. Kusnadi and G.H. Perdew. 2004. Divergent roles of hepatitis B virus X-associated protein 2 (XAP2) in human versus mouse Ah receptor complexes. Biochemistry. 43(3):700-709.

Rattenborg, T., I. Gjermandsen and E.C. Bonefeld-Jorgensen. 2002. Inhibition of E2-induced expression of BRCA1 by persistent organochlorines. Breast Cancer Res. 4(6):R12.

Ray, S.S. and H.I. Swanson. 2004. Dioxin-induced immortalization of normal human keratinocytes and silencing of p53 and p16INK4a. J. Biol. Chem. 279(26):27187-27193.

Reen, R.K., A. Cadwallader and G.H. Perdew. 2002. The subdomains of the transactivation domain of the aryl hydrocarbon receptor (AhR) inhibit AhR and estrogen receptor transcriptional activity. Arch. Biochem. Biophys. 408(1):93-102.

Reichard, J.F., T.P. Dalton, H.G. Shertzer and A. Puga. 2005. Induction of Oxidative Stress Responses by Dioxin and other Ligands of the Aryl Hydrocarbon Receptor. Dose. Response. 3(3):306-331.

Remillard, R.B. and N.J. Bunce. 2002. Use of Haber's Rule to estimate the risk of diabetes from background exposures to dioxin-like compounds. Toxicol. Lett. 131(3):161-166.

Renner, R. 2004. Redrawing the dose-response curve. Environ. Sci. Technol. 38(5):90A-95A.

Reynolds, P., S.E. Hurley, M. Petreas, D.E. Goldberg, D. Smith, D. Gilliss, M.E. Mahoney and S.S. Jeffrey. 2005. Adipose levels of dioxins and risk of breast cancer. Cancer Causes Control. 16(5):525-535.

Ricci, P. 2007. [Venetian sarcomas and exposure to dioxins]. Epidemiol. Prev. 31(1):2.

Richter, E. 2003. The article of Shertzer et al. (2002) has several short comings. Toxicol. Appl. Pharmacol. 188(3):199-200.

Richter, C.A., D.E. Tillitt and M. Hannink. 2001. Regulation of subcellular localization of the aryl hydrocarbon receptor (AhR). Arch. Biochem. Biophys. 389(2):207-217.

Riddick, D.S., C. Lee, A. Bhathena and Y.E. Timsit. 2003. The 2001 Veylien Henderson Award of the Society of Toxicology of Canada. Positive and negative transcriptional regulation of cytochromes P450 by polycyclic aromatic hydrocarbons. Can. J. Physiol Pharmacol. 81(1):59-77.

Riecke, K., A. Schmidt and R. Stahlmann. 2003. Effects of 2,3,7,8-TCDD and PCB 126 on human thymic epithelial cells in vitro. Arch. Toxicol. 77(6):358-364.

Rier, S.E. 2002. The potential role of exposure to environmental toxicants in the pathophysiology of endometriosis. Ann. N. Y. Acad. Sci. 955:201-212.

Rier, S.E. 2008. Environmental immune disruption: a comorbidity factor for reproduction? Fertil. Steril. 89(2 Suppl):e103-e108.

Rifkind, A.B. 2006. CYP1A in TCDD toxicity and in physiology-with particular reference to CYP dependent arachidonic acid metabolism and other endogenous substrates. Drug Metab Rev. 38(1-2):291-335.

Rivera, S.P., S.T. Saarikoski and O. Hankinson. 2002. Identification of a novel dioxin-inducible cytochrome P450. Mol. Pharmacol. 61(2):255-259.

Rivera, S.P., H.H. Choi, B. Chapman, M.J. Whitekus, M. Terao, E. Garattini and O. Hankinson. 2005. Identification of aldehyde oxidase 1 and aldehyde oxidase homologue 1 as dioxin-inducible genes. Toxicology. 207(3):401-409.

Rivera, S.P., F. Wang, S.T. Saarikoski, R.T. Taylor, B. Chapman, R. Zhang and O. Hankinson. 2007a. A novel promoter element containing multiple overlapping xenobiotic and hypoxia response elements mediates induction of cytochrome P4502S1 by both dioxin and hypoxia. J. Biol. Chem. 282(15):10881-10893.

Rivera, S.P., S.T. Saarikoski, W. Sun and O. Hankinson. 2007b. Identification of novel dioxin-responsive genes by representational difference analysis. Xenobiotica. 37(3):271-279.

Roberts, E.A., P.A. Harper, J.M. Wong, Y. Wang and S. Yang. 2000. Failure of Ah receptor to mediate induction of cytochromes P450 in the CYP1 family in the human hepatoma line SK-Hep-1. Arch. Biochem. Biophys. 384(1):190-198.

Robinson, S.W., B. Clothier, R.A. Akhtar, A.L. Yang, I. Latour, C. Van Ijperen, M.F. Festing and A.G. Smith. 2002. Non-ahr gene susceptibility Loci for porphyria and liver injury induced by the interaction of 'dioxin' with iron overload in mice. Mol. Pharmacol. 61(3):674-681.

Rogers, J.M. and M.S. Denison. 2002. Analysis of the antiestrogenic activity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in human ovarian carcinoma BG-1 cells. Mol. Pharmacol. 61(6):1393-1403.

Roy, T.A., K. Hammerstrom and J. Schaum. 2008. Percutaneous absorption of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) from soil. J. Toxicol. Environ. Health A. 71(23):1509-1515.

Rozman, K.K. 2004. Area under the curve as a dose metric for promotional responses following 2,3,7,8-tetrachlorodibenzo-p-dioxin exposure. Toxicol. Appl. Pharmacol. 195(1):126-128.

Ruby, C.E., M. Leid and N.I. Kerkvliet. 2002a. 2,3,7,8-Tetrachlorodibenzo-p-dioxin suppresses tumor necrosis factor-alpha and anti-CD40-induced activation of NF-kappaB/Rel in dendritic cells: p50 homodimer activation is not affected. Mol. Pharmacol. 62(3):722-728.

Ruby, M.V., K.A. Fehling, D.J. Paustenbach, B.D. Landenberger and M.P. Holsapple. 2002b. Oral bioaccessibility of dioxins/furans at low concentrations (50-350 ppt toxicity equivalent) in soil. Environ. Sci. Technol. 36(22):4905-4911.

Ruby, M., S. Casteel, T. Evans, K. Fehling, D. Paustenbach, R. Budinsky, J. Giesy, L. Aylward and B. Landenberger. 2004. Background concentrations of dioxins, furans, and PCBs in Sprague-Dawley rats and juvenile swine. J. Toxicol. Environ. Health A. 67(11):845-850.

Ruby, C.E., C.J. Funatake and N.I. Kerkvliet. 2005. 2,3,7,8 Tetrachlorodibenzo-p-Dioxin (TCDD) Directly Enhances the Maturation and Apoptosis of Dendritic Cells In Vitro. J. Immunotoxicol. 1(3):159-166.

Ruegg, J., E. Swedenborg, D. Wahlstrom, A. Escande, P. Balaguer, K. Pettersson and I. Pongratz. 2008. The transcription factor aryl hydrocarbon receptor nuclear translocator functions as an estrogen receptor beta-selective coactivator, and its recruitment to alternative pathways mediates antiestrogenic effects of dioxin. Mol. Endocrinol. 22(2):304-316.

Runge-Morris, M. and T.A. Kocarek. 2005. Regulation of sulfotransferases by xenobiotic receptors. Curr. Drug Metab. 6(4):299-307.

Rushing, S.R. and M.S. Denison. 2002. The silencing mediator of retinoic acid and thyroid hormone receptors can interact with the aryl hydrocarbon (Ah) receptor but fails to repress Ah receptor-dependent gene expression. Arch. Biochem. Biophys. 403(2):189-201.

Rushton, L., S. Hutchings and T.P. Brown. 2008. The Burden of Cancer at Work: Estimation as the first step to prevention. Occup. Environ. Med.

Ryan, E.P., J.D. Holz, M. Mulcahey, T.J. Sheu, T.A. Gasiewicz and J.E. Puzas. 2007. Environmental toxicants may modulate osteoblast differentiation by a mechanism involving the aryl hydrocarbon receptor. J. Bone Miner. Res. 22(10):1571-1580.

Rychen, G., C. Laurent, C. Feidt, N. Grova, P.E. Lafargue, A. Hachimi and F. Laurent. 2002. Milk-arterial plasma transfer of PCDDs and PCDFs in pigs. J. Agric. Food Chem. 50DV(6):1695-1699.

Saarikoski, S.T., S.P. Rivera, O. Hankinson and K. Husgafvel-Pursiainen. 2005a. CYP2S1: a short review. Toxicol. Appl. Pharmacol. 207(2 Suppl):62-69.

Saarikoski, S.T., H.A. Wikman, G. Smith, C.H. Wolff and K. Husgafvel-Pursiainen. 2005b. Localization of cytochrome P450 CYP2S1 expression in human tissues by in situ hybridization and immunohistochemistry. J. Histochem. Cytochem. 53(5):549-556.

Safe, S. 2001. Molecular biology of the Ah receptor and its role in carcinogenesis. Toxicol. Lett. 120(1-3):1-7.

Safe, S. and A. McDougal. 2002. Mechanism of action and development of selective aryl hydrocarbon receptor modulators for treatment of hormone-dependent cancers (Review). Int. J. Oncol. 20(6):1123-1128.

Safe, S. and M. Wormke. 2003. Inhibitory aryl hydrocarbon receptor-estrogen receptor alpha cross-talk and mechanisms of action. Chem. Res. Toxicol. 16(7):807-816.

Safe, S., M. Wormke and I. Samudio. 2000. Mechanisms of inhibitory aryl hydrocarbon receptor-estrogen receptor crosstalk in human breast cancer cells. J. Mammary. Gland. Biol. Neoplasia. 5(3):295-306.

Saghir, S.A., M. Lebofsky, D.M. Pinson and K.K. Rozman. 2005. Validation of Haber's Rule (dose x time = constant) in rats and mice for monochloroacetic acid and 2,3,7,8-tetrachlorodibenzo-p-dioxin under conditions of kinetic steady state. Toxicology. 215(1-2):48-56.

Sahlberg, C., E. Peltonen, P.L. Lukinmaa and S. Alaluusua. 2007. Dioxin alters gene expression in mouse embryonic tooth explants. J. Dent. Res. 86(7):600-605.

Sakai, R., T. Kajiume, H. Inoue, R. Kanno, M. Miyazaki, Y. Ninomiya and M. Kanno. 2003. TCDD treatment eliminates the long-term reconstitution activity of hematopoietic stem cells. Toxicol. Sci. 72(1):84-91.

Sakata, Y., W. Yoshioka, C. Tohyama and S. Ohsako. 2007. Internal genomic sequence of human CYP1A1 gene is involved in superinduction of dioxin-induced CYP1A1 transcription by cycloheximide. Biochem. Biophys. Res. Commun. 355(3):687-692.

Sakurai, K., E. Todaka, Y. Suzuki and C. Mori. 2002. An experimental model using guinea pigs to reduce accumulated dioxins in the body. Congenit. Anom. (Kyoto). 42(4):323-326.

Sakurai, K., E. Todaka, Y. Saito and C. Mori. 2004. Pilot study to reduce dioxins in the human body. Intern. Med. 43(9):792-795.

Sand, S., A.F. Filipsson and K. Victorin. 2002. Evaluation of the benchmark dose method for dichotomous data: model dependence and model selection. Regul. Toxicol. Pharmacol. 36(2):184-197.

Sanderson, J.T., L. Slobbe, G.W. Lansbergen, S. Safe and M. Van den Berg. 2001. 2,3,7,8-Tetrachlorodibenzo-p-dioxin and diindolylmethanes differentially induce cytochrome P450 1A1, 1B1, and 19 in H295R human adrenocortical carcinoma cells. Toxicol. Sci. 61(1):40-48.

Santiago-Josefat, B., S. Mulero-Navarro, S.L. Dallas and P.M. Fernandez-Salguero. 2004. Overexpression of latent transforming growth factor-beta binding protein 1 (LTBP-1) in dioxin receptor-null mouse embryo fibroblasts. J. Cell Sci. 117(Pt 6):849-859.

Santini, R.P., S. Myrand, C. Elferink and J.J. Reiners, Jr. 2001. Regulation of Cyp1a1 induction by dioxin as a function of cell cycle phase. J. Pharmacol. Exp. Ther. 299(2):718-728.

Sarafian, T., N. Habib, J.T. Mao, I.H. Tsu, M.L. Yamamoto, E. Hsu, D.P. Tashkin and M.D. Roth. 2005. Gene expression changes in human small airway epithelial cells exposed to Delta9-tetrahydrocannabinol. Toxicol. Lett. 158(2):95-107.

Sarioglu, H., S. Brandner, C. Jacobsen, T. Meindl, A. Schmidt, J. Kellermann, F. Lottspeich and U. Andrae. 2006. Quantitative analysis of 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced proteome alterations in 5L rat hepatoma cells using isotope-coded protein labels. Proteomics. 6(8):2407-2421.

Sarioglu, H., S. Brandner, M. Haberger, C. Jacobsen, J. Lichtmannegger, M. Wormke and U. Andrae. 2008. Analysis of 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced proteome changes in 5L rat hepatoma cells reveals novel targets of dioxin action including the mitochondrial apoptosis regulator VDAC2. Mol. Cell Proteomics. 7(2):394-410.

Sarkar, S., N.R. Jana, J. Yonemoto, C. Tohyama and H. Sone. 2000. Estrogen enhances induction of cytochrome P-4501A1 by 2,3,7, 8-tetrachlorodibenzo-p-dioxin in liver of female Long-Evans rats. Int. J. Oncol. 16(1):141-147.

Sarkar, P., K. Shiizaki, J. Yonemoto and H. Sone. 2006. Activation of telomerase in BeWo cells by estrogen and 2,3,7,8-tetrachlorodibenzo-p-dioxin in co-operation with c-Myc. Int. J. Oncol. 28(1):43-51.

Sartori, N., K. Thomaseth and A. Salvan. 2004. Local influence analysis when interfacing toxicokinetic and proportional hazard models. Stat. Med. 23(15):2399-2412.

Sasamoto, T., S. Horii, A. Ibe, N. Takada and K. Shirota. 2006. Concentration changes of PCDDs, PCDFs, and dioxin-like PCBs in human breast milk samples as shown by a follow-up survey. Chemosphere. 64(4):642-649.

Sassi, G., A.M. Vernai and B. Ruggeri. 2007. Quantitative estimation of uncertainty in human risk analysis. J. Hazard. Mater. 145(1-2):296-304.

Sato, S., H. Shirakawa, S. Tomita, Y. Ohsaki, K. Haketa, O. Tooi, N. Santo, M. Tohkin, Y. Furukawa, F.J. Gonzalez and M. Komai. 2008. Low-dose dioxins alter gene expression related to cholesterol biosynthesis, lipogenesis, and glucose metabolism through the aryl hydrocarbon receptor-mediated pathway in mouse liver. Toxicol. Appl. Pharmacol. 229(1):10-19.

Savouret, J.F., M. Antenos, M. Quesne, J. Xu, E. Milgrom and R.F. Casper. 2001. 7-ketocholesterol is an endogenous modulator for the arylhydrocarbon receptor. J. Biol. Chem. 276(5):3054-3059.

Sawaya, R.M. and D.S. Riddick. 2008. Cytochrome P450 2C11 5'-flanking region and promoter: regulation by aromatic hydrocarbons in vitro. Toxicology. 248(2-3):104-112.

Schaufler, K., P. Haslmayer, W. Jager, M. Pec and T. Thalhammer. 2002. The environmental toxin 2,3,7,8-tetrachlorodibenzo-p-dioxin induces cytochrome P450 activity in high passage PC 3 and DU 145 human prostate cancer cell lines. Int. J. Mol. Med. 9(4):411-416.

Schecter, A., M. Pavuk, O. Papke and J.J. Ryan. 2003. Dioxin, dibenzofuran, and coplanar PCB levels in Laotian blood and milk from agent orange-sprayed and nonsprayed areas, 2001. J. Toxicol. Environ. Health A. 66(21):2067-2075.

Schmidt, C.K., P. Hoegberg, N. Fletcher, C.B. Nilsson, C. Trossvik, H. Hakansson and H. Nau. 2003. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) alters the endogenous metabolism of all-trans-retinoic acid in the rat. Arch. Toxicol. 77(7):371-383.

Schmidt, I., M. Fahling, B. Nafz, A. Skalweit and B.J. Thiele. 2007. Induction of translationally controlled tumor protein (TCTP) by transcriptional and post-transcriptional mechanisms. FEBS J. 274(20):5416-5424.

Schneider, D., M.A. Manzan, R.B. Crawford, W. Chen and N.E. Kaminski. 2008. 2,3,7,8-Tetrachlorodibenzo-p-dioxin-mediated impairment of B cell differentiation involves dysregulation of paired box 5 (Pax5) isoform, Pax5a. J. Pharmacol. Exp. Ther. 326(2):463-474.

Schrader, T.J. and G.M. Cooke. 2000. Examination of selected food additives and organochlorine food contaminants for androgenic activity in vitro. Toxicol. Sci. 53(2):278-288.

Schreiber, T.D., C. Kohle, F. Buckler, S. Schmohl, A. Braeuning, A. Schmiechen, M. Schwarz and P.A. Munzel. 2006. Regulation of CYP1A1 gene expression by the antioxidant tert-butylhydroquinone. Drug Metab Dispos. 34(7):1096-1101.

Schrenk, D., H.J. Schmitz, S. Bohnenberger, B. Wagner and W. Worner. 2004. Tumor promoters as inhibitors of apoptosis in rat hepatocytes. Toxicol. Lett. 149(1-3):43-50.

Schultz, R., J. Suominen, T. Varre, H. Hakovirta, M. Parvinen, J. Toppari and M. Pelto-Huikko. 2003. Expression of aryl hydrocarbon receptor and aryl hydrocarbon receptor nuclear translocator messenger ribonucleic acids and proteins in rat and human testis. Endocrinology. 144(3):767-776.

Schulz, T.G., R. Thiel, D. Neubert, P.J. Brassil, T. Schulz-Utermoehl, A.R. Boobis and R.J. Edwards. 2001. Assessment of P450 induction in the marmoset monkey using targeted anti-peptide antibodies. Biochim. Biophys. Acta. 1546(1):143-155.

Schwanekamp, J.A., M.A. Sartor, S. Karyala, D. Halbleib, M. Medvedovic and C.R. Tomlinson. 2006. Genome-wide analyses show that nuclear and cytoplasmic RNA levels are differentially affected by dioxin. Biochim. Biophys. Acta. 1759(8-9):388-402.

Schwarz, M. and K.E. Appel. 2005. Carcinogenic risks of dioxin: mechanistic considerations. Regul. Toxicol. Pharmacol. 43(1):19-34.

Schwarz, M., A. Buchmann, S. Stinchcombe, A. Kalkuhl and K. Bock. 2000. Ah receptor ligands and tumor promotion: survival of neoplastic cells. Toxicol. Lett. 112-113:69-77.

Sciullo, E.M., C.F. Vogel, W. Li and F. Matsumura. 2008. Initial and extended inflammatory messages of the nongenomic signaling pathway of the TCDD-activated Ah receptor in U937 macrophages. Arch. Biochem. Biophys. 480(2):143-155.

Seidel, S.D., G.M. Winters, W.J. Rogers, M.H. Ziccardi, V. Li, B. Keser and M.S. Denison. 2001. Activation of the Ah receptor signaling pathway by prostaglandins. J. Biochem. Mol. Toxicol. 15(4):187-196.

Seifert, A., D.M. Katschinski, S. Tonack, B. Fischer and A. Navarrete Santos. 2008. Significance of prolyl hydroxylase 2 in the interference of aryl hydrocarbon receptor and hypoxia-inducible factor-1 alpha signaling. Chem. Res. Toxicol. 21(2):341-348.

Selmin, O., P.A. Thorne, F.M. Blachere, P.D. Johnson and D.F. Romagnolo. 2005. Transcriptional activation of the membrane-bound progesterone receptor (mPR) by dioxin, in endocrine-responsive tissues. Mol. Reprod. Dev. 70(2):166-174.

Senft, A.P., T.P. Dalton, D.W. Nebert, M.B. Genter, R.J. Hutchinson and H.G. Shertzer. 2002. Dioxin increases reactive oxygen production in mouse liver mitochondria. Toxicol. Appl. Pharmacol. 178(1):15-21.

Sergent, T., L. Ribonnet, A. Kolosova, S. Garsou, A. Schaut, S. De Saeger, C. Van Peteghem, Y. Larondelle, L. Pussemier and Y.J. Schneider. 2008. Molecular and cellular effects of food contaminants and secondary plant components and their plausible interactions at the intestinal level. Food Chem. Toxicol. 46(3):813-841.

Shaban, Z., S. El-Shazly, S. Abdelhady, I. Fattouh, K. Muzandu, M. Ishizuka, K. Kimura, A. Kazusaka and S. Fujita. 2004. Down regulation of hepatic PPARalpha function by AhR ligand. J. Vet. Med. Sci. 66(11):1377-1386.

Shen, D., T.P. Dalton, D.W. Nebert and H.G. Shertzer. 2005. Glutathione redox state regulates mitochondrial reactive oxygen production. J. Biol. Chem. 280(27):25305-25312.

Shertzer, H.G., C.D. Clay, M.B. Genter, M.C. Chames, S.N. Schneider, G.G. Oakley, D.W. Nebert and T.P. Dalton. 2004a. Uncoupling-mediated generation of reactive oxygen by halogenated aromatic hydrocarbons in mouse liver microsomes. Free Radic. Biol. Med. 36(5):618-631.

Shertzer, H.G., C.D. Clay, M.B. Genter, S.N. Schneider, D.W. Nebert and T.P. Dalton. 2004b. Cyp1a2 protects against reactive oxygen production in mouse liver microsomes. Free Radic. Biol. Med. 36(5):605-617.

Shertzer, H.G., M.B. Genter, D. Shen, D.W. Nebert, Y. Chen and T.P. Dalton. 2006. TCDD decreases ATP levels and increases reactive oxygen production through changes in mitochondrial F(0)F(1)-ATP synthase and ubiquinone. Toxicol. Appl. Pharmacol. 217(3):363-374.

Shi, Y.L., X.Z. Luo, X.Y. Zhu, K.Q. Hua, Y. Zhu and D.J. Li. 2006. Effects of combined 17beta-estradiol with TCDD on secretion of chemokine IL-8 and expression of its receptor CXCR1 in endometriotic focus-associated cells in co-culture. Hum. Reprod. 21(4):870-879.

Shi, Y.L., X.Z. Luo, X.Y. Zhu and D.J. Li. 2007. Combination of 17beta-estradiol with the environmental pollutant TCDD is involved in pathogenesis of endometriosis via up-regulating the chemokine I-309-CCR8. Fertil. Steril. 88(2):317-325.

Shi, Z., Y. Chen, H. Dong, R.M. Amos-Kroohs and D.W. Nebert. 2008. Generation of a 'humanized' hCYP1A1_1A2_Cyp1a1/1a2(-/-)_Ahrd mouse line harboring the poor-affinity aryl hydrocarbon receptor. Biochem. Biophys. Res. Commun. 376(4):775-780.

Shibazaki, M., T. Takeuchi, S. Ahmed and H. Kikuchia. 2004a. Blockade by SB203580 of Cyp1a1 induction by 2,3,7,8-tetrachlorodibenzo-p-dioxin, and the possible mechanism: possible involvement of the p38 mitogen-activated protein kinase pathway in shuttling of Ah receptor overexpressed in COS-7 cells. Ann. N. Y. Acad. Sci. 1030:275-281.

Shibazaki, M., T. Takeuchi, S. Ahmed and H. Kikuchi. 2004b. Suppression by p38 MAP kinase inhibitors (pyridinyl imidazole compounds) of Ah receptor target gene activation by 2,3,7,8-tetrachlorodibenzo-p-dioxin and the possible mechanism. J. Biol. Chem. 279(5):3869-3876.

Shiizaki, K., S. Ohsako, T. Koyama, R. Nagata, J. Yonemoto and C. Tohyama. 2005. Lack of CYP1A1 expression is involved in unresponsiveness of the human hepatoma cell line SK-HEP-1 to dioxin. Toxicol. Lett. 160(1):22-33.

Shimba, S., M. Hayashi, H. Sone, J. Yonemoto and M. Tezuka. 2000. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) induces binding of a 50 kDa protein on the 3' untranslated region of urokinase-type plasminogen activator mRNA. Biochem. Biophys. Res. Commun. 272(2):441-448.

Shinkyo, R., T. Sakaki, M. Ohta and K. Inouye. 2003. Metabolic pathways of dioxin by CYP1A1: species difference between rat and human CYP1A subfamily in the metabolism of dioxins. Arch. Biochem. Biophys. 409(1):180-187.

Shipley, J.M. and D.J. Waxman. 2006. Aryl hydrocarbon receptor-independent activation of estrogen receptor-dependent transcription by 3-methylcholanthrene. Toxicol. Appl. Pharmacol. 213(2):87-97.

Sienkiewicz, P., H.P. Ciolino, B.J. Leslie, P.J. Hergenrother, K. Singletary and G.C. Yeh. 2007. A novel synthetic analogue of a constituent of Isodon excisus inhibits transcription of CYP1A1, -1A2 and -1B1 by preventing activation of the aryl hydrocarbon receptor. Carcinogenesis. 28(5):1052-1057.

Silkworth, J.B., A. Koganti, K. Illouz, A. Possolo, M. Zhao and S.B. Hamilton. 2005. Comparison of TCDD and PCB CYP1A induction sensitivities in fresh hepatocytes from human donors, sprague-dawley rats, and rhesus monkeys and HepG2 cells. Toxicol. Sci. 87(2):508-519.

Silkworth, J.B., E.A. Carlson, C. McCulloch, K. Illouz, S. Goodwin and T.R. Sutter. 2008. Toxicogenomic analysis of gender, chemical, and dose effects in livers of TCDD- or aroclor 1254-exposed rats using a multifactor linear model. Toxicol. Sci. 102(2):291-309.

Singh, N.P., M. Nagarkatti and P.S. Nagarkatti. 2007. Role of dioxin response element and nuclear factor-kappaB motifs in 2,3,7,8-tetrachlorodibenzo-p-dioxin-mediated regulation of Fas and Fas ligand expression. Mol. Pharmacol. 71(1):145-157.

Singh, M.N., H.F. Stringfellow, S.E. Taylor, K.M. Ashton, M. Ahmad, K.R. Abdo, O.M. El-Agnaf, P.L. Martin-Hirsch and F.L. Martin. 2008a. Elevated expression of CYP1A1 and {gamma}-SYNUCLEIN in human ectopic (ovarian) endometriosis compared to eutopic endometrium. Mol. Hum. Reprod.

Singh, N.P., M. Nagarkatti and P. Nagarkatti. 2008b. Primary peripheral T cells become susceptible to 2,3,7,8-tetrachlorodibenzo-p-dioxin-mediated apoptosis in vitro upon activation and in the presence of dendritic cells. Mol. Pharmacol. 73(6):1722-1735.

Sogawa, K. 2000. [Regulatory mechanism of genes by the Ah receptor which mediates toxic effects of dioxins]. Nippon Rinsho. 58(12):2446-2451.

Son, D.S. and K.K. Rozman. 2002. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) induces plasminogen activator inhibitor-1 through an aryl hydrocarbon receptor-mediated pathway in mouse hepatoma cell lines. Arch. Toxicol. 76(7):404-413.

Son, D.S., K.F. Roby, K.K. Rozman and P.F. Terranova. 2002. Estradiol enhances and estriol inhibits the expression of CYP1A1 induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin in a mouse ovarian cancer cell line. Toxicology. 176(3):229-243.

Son, W.K., D.Y. Lee, S.H. Lee, W.A. Joo and C.W. Kim. 2003. Analysis of proteins expressed in rat plasma exposed to dioxin using 2-dimensional gel electrophoresis. Proteomics. 3(12):2393-2401.

Soneda, S., M. Fukami, M. Fujimoto, T. Hasegawa, Y. Koitabashi and T. Ogata. 2005. Association of micropenis with Pro185Ala polymorphism of the gene for aryl hydrocarbon receptor repressor involved in dioxin signaling. Endocr. J. 52(1):83-88.

Song, Z. and R.S. Pollenz. 2002. Ligand-dependent and independent modulation of aryl hydrocarbon receptor localization, degradation, and gene regulation. Mol. Pharmacol. 62(4):806-816.

Song, Z. and R.S. Pollenz. 2003. Functional analysis of murine aryl hydrocarbon (AH) receptors defective in nuclear import: impact on AH receptor degradation and gene regulation. Mol. Pharmacol. 63(3):597-606.

Sonneveld, E., A. Jonas, O.C. Meijer, A. Brouwer and B. van der Burg. 2007. Glucocorticoid-enhanced expression of dioxin target genes through regulation of the rat aryl hydrocarbon receptor. Toxicol. Sci. 99(2):455-469.

Spinelli, J. 2008. Pesticides, soft tissue sarcoma and non-Hodgkin lymphoma. Acta Oncol. 47(3):335-336.

Spink, B.C., B.H. Katz, M.M. Hussain, S. Pang, S.P. Connor, K.M. Aldous, J.F. Gierthy and D.C. Spink. 2000. SULT1A1 catalyzes 2-methoxyestradiol sulfonation in MCF-7 breast cancer cells. Carcinogenesis. 21(11):1947-1957.

Spink, B.C., M.M. Hussain, B.H. Katz, L. Eisele and D.C. Spink. 2003a. Transient induction of cytochromes P450 1A1 and 1B1 in MCF-7 human breast cancer cells by indirubin. Biochem. Pharmacol. 66(12):2313-2321.

Spink, D.C., B.H. Katz, M.M. Hussain, B.T. Pentecost, Z. Cao and B.C. Spink. 2003b. Estrogen regulates Ah responsiveness in MCF-7 breast cancer cells. Carcinogenesis. 24(12):1941-1950.

Spitaler, M., C. Iben and H. Tausch. 2005. Dioxin residues in the edible tissue of finishing pigs after dioxin feeding. J. Anim Physiol Anim Nutr. (Berl). 89(3-6):65-71.

Staskal, D.F., J.J. Diliberto, M.J. DeVito and L.S. Birnbaum. 2005. Inhibition of human and rat CYP1A2 by TCDD and dioxin-like chemicals. Toxicol. Sci. 84(2):225-231.

Steenland, K. and J.A. Deddens. 2004. A practical guide to dose-response analyses and risk assessment in occupational epidemiology. Epidemiology. 15(1):63-70.

Steenland, K. and S. Bartell. 2007. Caution in drawing conclusions from PBPK models based on sparse human data. Risk Anal. 27(1):7-8.

Sterling, K.M., Jr. and K.R. Cutroneo. 2002. Differentiation-dependent induction of CYP1A1 in cultured rat small intestinal epithelial cells, colonocytes, and human colon carcinoma cells: basement membrane-mediated apoptosis. J. Cell Biochem. 86(3):440-450.

Stevens, E.A. and C.A. Bradfield. 2008. Immunology: T cells hang in the balance. Nature. 453(7191):46-47.

Sugihara, K., S. Kitamura, T. Yamada, T. Okayama, S. Ohta, K. Yamashita, M. Yasuda, Y. Fujii-Kuriyama, K. Saeki, S. Matsui and T. Matsuda. 2004. Aryl hydrocarbon receptor-mediated induction of microsomal drug-metabolizing enzyme activity by indirubin and indigo. Biochem. Biophys. Res. Commun. 318(2):571-578.

Suh, J., Y.J. Jeon, H.M. Kim, J.S. Kang, N.E. Kaminski and K.H. Yang. 2002. Aryl hydrocarbon receptor-dependent inhibition of AP-1 activity by 2,3,7,8-tetrachlorodibenzo-p-dioxin in activated B cells. Toxicol. Appl. Pharmacol. 181(2):116-123.

Sulentic, C.E., M.P. Holsapple and N.E. Kaminski. 2000. Putative link between transcriptional regulation of IgM expression by 2,3,7,8-tetrachlorodibenzo-p-dioxin and the aryl hydrocarbon receptor/dioxin-responsive enhancer signaling pathway. J. Pharmacol. Exp. Ther. 295(2):705-716.

Sulentic, C.E., J.S. Kang, Y.J. Na and N.E. Kaminski. 2004a. Interactions at a dioxin responsive element (DRE) and an overlapping kappaB site within the hs4 domain of the 3'alpha immunoglobulin heavy chain enhancer. Toxicology. 200(2-3):235-246.

Sulentic, C.E., W. Zhang, Y.J. Na and N.E. Kaminski. 2004b. 2,3,7,8-tetrachlorodibenzo-p-dioxin, an exogenous modulator of the 3'alpha immunoglobulin heavy chain enhancer in the CH12.LX mouse cell line. J. Pharmacol. Exp. Ther. 309(1):71-78.

Sun, Y.V., D.R. Boverhof, L.D. Burgoon, M.R. Fielden and T.R. Zacharewski. 2004. Comparative analysis of dioxin response elements in human, mouse and rat genomic sequences. Nucleic Acids Res. 32(15):4512-4523.

Suzuki, T. and K. Nohara. 2007. Regulatory factors involved in species-specific modulation of arylhydrocarbon receptor (AhR)-dependent gene expression in humans and mice. J. Biochem. 142(4):443-452.

Suzuki, G., M. Nakano and S. Nakano. 2005. Distribution of PCDDs/PCDFs and Co-PCBs in human maternal blood, cord blood, placenta, milk, and adipose tissue: dioxins showing high toxic equivalency factor accumulate in the placenta. Biosci. Biotechnol. Biochem. 69(10):1836-1847.

Svensson, C. and K. Lundberg. 2001. Immune-specific up-regulation of adseverin gene expression by 2,3,7,8-tetrachlorodibenzo-p-dioxin. Mol. Pharmacol. 60(1):135-142.

Sycheva, L.P., T.E. Mozhaeva, N.V. Umnova, N.A. Zhuchenko, V.H. Diep and H.A. Tuyet. 2008. [Cytogenetic and other cariological parameters of exfoliative buccal cells in Vietnamese children from areas where dioxin-containing herbicides were applied]. Vestn. Ross. Akad. Med. Nauk(1):19-23.

Takahashi, S., S. Hirano, N. Ito and T. Tamaya. 2005. Effects of dioxin and nutrition on cellular proliferation and dioxin- and estrogen-linked genes in ovarian cancer cell lines. Eur. J. Gynaecol. Oncol. 26(2):175-180.

Takanaga, H., M. Kunimoto, T. Adachi, C. Tohyama and Y. Aoki. 2001. Inhibitory effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on cAMP-induced differentiation of rat C6 glial cell line. J. Neurosci. Res. 64(4):402-409.

Takasugi, N. 2000. [Research trace of endocrine-disrupting chemicals]. Nippon Rinsho. 58(12):2387-2392.

Takenaka, S., T. Todaka, M. Nakamura, T. Hori, T. Iida, T. Yamada and J. Hata. 2002. Polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans and non-ortho, mono-ortho chlorine substituted biphenyls in Japanese human liver and adipose tissue. Chemosphere. 49(2):161-172.

Taketoh, J., J. Mutoh, T. Takeda, T. Ogishima, S. Takeda, Y. Ishii, T. Ishida and H. Yamada. 2007. Suppression of fetal testicular cytochrome P450 17 by maternal exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin: a mechanism involving an initial effect on gonadotropin synthesis in the pituitary. Life Sci. 80(14):1259-1267.

Tamaki, A., H. Hayashi, H. Nakajima, T. Takii, D. Katagiri, K. Miyazawa, K. Hirose and K. Onozaki. 2004. Polycyclic aromatic hydrocarbon increases mRNA level for interleukin 1 beta in human fibroblast-like synoviocyte line via aryl hydrocarbon receptor. Biol. Pharm. Bull. 27(3):407-410.

Tan, Z., X. Chang, A. Puga and Y. Xia. 2002. Activation of mitogen-activated protein kinases (MAPKs) by aromatic hydrocarbons: role in the regulation of aryl hydrocarbon receptor (AHR) function. Biochem. Pharmacol. 64(5-6):771-780.

Tan, Z., M. Huang, A. Puga and Y. Xia. 2004. A critical role for MAP kinases in the control of Ah receptor complex activity. Toxicol. Sci. 82(1):80-87.

Tanabe, S. 2005. [Contamination of dioxins and related compounds in human and wildlife: evidences with focus on our recent studies]. Chudoku. Kenkyu. 18(4):319-331.

Tanaka, G., S. Kanaji, A. Hirano, K. Arima, A. Shinagawa, C. Goda, S. Yasunaga, K. Ikizawa, Y. Yanagihara, M. Kubo, Y. Kuriyama-Fujii, Y. Sugita, A. Inokuchi and K. Izuhara. 2005. Induction and activation of the aryl hydrocarbon receptor by IL-4 in B cells. Int. Immunol. 17(6):797-805.

Tanaka, J., J. Yonemoto, H. Zaha, R. Kiyama and H. Sone. 2007. Estrogen-responsive genes newly found to be modified by TCDD exposure in human cell lines and mouse systems. Mol. Cell Endocrinol. 272(1-2):38-49.

Tang, N.J., L. Dong, J. Liu, S.W. Ma, L.J. Zhao, Z.P. Bai, W.Q. Zhang, P.J. Coenraads and C.M. Zhang. 2008. [Effect of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin on the expression of AhR and TGF-alpha]. Zhonghua Yu Fang Yi. Xue. Za Zhi. 42(1):21-24.

Taylor, R.T., F. Wang, E.L. Hsu and O. Hankinson. 2008. Roles of Coactivator Proteins in Dioxin Induction of CYP1A1 and CYP1B1 in Human Breast Cancer Cells. Toxicol. Sci.

Temchura, V.V., M. Frericks, W. Nacken and C. Esser. 2005. Role of the aryl hydrocarbon receptor in thymocyte emigration in vivo. Eur. J. Immunol. 35(9):2738-2747.

Teneng, I., V. Stribinskis and K.S. Ramos. 2007. Context-specific regulation of LINE-1. Genes Cells. 12(10):1101-1110.

Thackaberry, E.A., E.J. Bedrick, M.B. Goens, L. Danielson, A.K. Lund, D. Gabaldon, S.M. Smith and M.K. Walker. 2003. Insulin regulation in AhR-null mice: embryonic cardiac enlargement, neonatal macrosomia, and altered insulin regulation and response in pregnant and aging AhR-null females. Toxicol. Sci. 76(2):407-417.

Thackaberry, E.A., Z. Jiang, C.D. Johnson, K.S. Ramos and M.K. Walker. 2005. Toxicogenomic profile of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the murine fetal heart: modulation of cell cycle and extracellular matrix genes. Toxicol. Sci. 88(1):231-241.

Thomae, T.L., E.A. Stevens and C.A. Bradfield. 2005. Transforming growth factor-beta3 restores fusion in palatal shelves exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. J. Biol. Chem. 280(13):12742-12746.

Thomas, R.D., M.R. Green, C. Wilson, A.L. Weckle, Z. Duanmu, T.A. Kocarek and M. Runge-Morris. 2006. Cytochrome P450 expression and metabolic activation of cooked food mutagen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in MCF10A breast epithelial cells. Chem. Biol. Interact. 160(3):204-216.

Thomsen, J.S., S. Kietz, A. Strom and J.A. Gustafsson. 2004. HES-1, a novel target gene for the aryl hydrocarbon receptor. Mol. Pharmacol. 65(1):165-171.

Tian, Y., A.B. Rabson and M.A. Gallo. 2002. Ah receptor and NF-kappaB interactions: mechanisms and physiological implications. Chem. Biol. Interact. 141(1-2):97-115.

Tian, Y., S. Ke, M. Chen and T. Sheng. 2003. Interactions between the aryl hydrocarbon receptor and P-TEFb. Sequential recruitment of transcription factors and differential phosphorylation of C-terminal domain of RNA polymerase II at cyp1a1 promoter. J. Biol. Chem. 278(45):44041-44048.

Tijet, N., P.C. Boutros, I.D. Moffat, A.B. Okey, J. Tuomisto and R. Pohjanvirta. 2006. Aryl hydrocarbon receptor regulates distinct dioxin-dependent and dioxin-independent gene batteries. Mol. Pharmacol. 69(1):140-153.

Timsit, Y.E., F.S. Chia, A. Bhathena and D.S. Riddick. 2002. Aromatic hydrocarbon receptor expression and function in liver of hypophysectomized male rats. Toxicol. Appl. Pharmacol. 185(2):136-145.

Toide, K., H. Yamazaki, R. Nagashima, K. Itoh, S. Iwano, Y. Takahashi, S. Watanabe and T. Kamataki. 2003. Aryl hydrocarbon hydroxylase represents CYP1B1, and not CYP1A1, in human freshly isolated white cells: trimodal distribution of Japanese population according to induction of CYP1B1 mRNA by environmental dioxins. Cancer Epidemiol. Biomarkers Prev. 12(3):219-222.

Tokizane, T., H. Shiina, M. Igawa, H. Enokida, S. Urakami, T. Kawakami, T. Ogishima, S.T. Okino, L.C. Li, Y. Tanaka, N. Nonomura, A. Okuyama and R. Dahiya. 2005. Cytochrome P450 1B1 is overexpressed and regulated by hypomethylation in prostate cancer. Clin. Cancer Res. 11(16):5793-5801.

Tokunaga, S., T. Iida and M. Furue. 2005. [Concept on the statistically approached new diagnostic standard for Yusho patients]. Fukuoka Igaku Zasshi. 96(5):135-145.

Tonack, S., K. Kind, J.G. Thompson, A.M. Wobus, B. Fischer and A.N. Santos. 2007. Dioxin affects glucose transport via the arylhydrocarbon receptor signal cascade in pluripotent embryonic carcinoma cells. Endocrinology. 148(12):5902-5912.

Toyoshiba, H., T. Yamanaka, H. Sone, F.M. Parham, N.J. Walker, J. Martinez and C.J. Portier. 2004. Gene interaction network suggests dioxin induces a significant linkage between aryl hydrocarbon receptor and retinoic acid receptor beta. Environ. Health Perspect. 112(12):1217-1224.

Trewin, A.L., M.J. Woller, B.A. Wimpee, L.K. Conley, M.G. Baldridge and R.J. Hutz. 2007. Short-term hormone release from adult female rat hypothalamic and pituitary explants is not altered by 2,3,7,8-tetrachlorodibenzo-p-dioxin. J. Reprod. Dev. 53(4):765-775.

Tsuchiya, Y., M. Nakajima and T. Yokoi. 2003a. Critical enhancer region to which AhR/ARNT and Sp1 bind in the human CYP1B1 gene. J. Biochem. 133(5):583-592.

Tsuchiya, Y., M. Nakajima, S. Itoh, M. Iwanari and T. Yokoi. 2003b. Expression of aryl hydrocarbon receptor repressor in normal human tissues and inducibility by polycyclic aromatic hydrocarbons in human tumor-derived cell lines. Toxicol. Sci. 72(2):253-259.

Tsuchiya, Y., M. Nakajima and T. Yokoi. 2004. [Regulation of human CYP1B1 and its role in carcinogenesis and estrogen homeostasis]. Seikagaku. 76(12):1565-1569.

Tsukumo, S., M. Iwata, C. Tohyama and K. Nohara. 2002. Skewed differentiation of thymocytes toward CD8 T cells by 2,3,7,8-tetrachlorodibenzo- p-dioxin requires activation of the extracellular signal-related kinase pathway. Arch. Toxicol. 76(5-6):335-343.

Tsutsumi, O. 2005. Assessment of human contamination of estrogenic endocrine-disrupting chemicals and their risk for human reproduction. J. Steroid Biochem. Mol. Biol. 93(2-5):325-330.

Tuomisto, J. and T. Vartiainen. 2004. [Dioxin and health--from molecular biology to preventive care]. Duodecim. 120(13):1664-1672.

Tuomisto, J., J. Pekkanen, H. Kiviranta, E. Tukiainen, T. Vartiainen, M. Viluksela and J.T. Tuomisto. 2006. Dioxin Cancer Risk - Example of Hormesis? Dose. Response. 3(3):332-341.

Uemura, H., K. Arisawa, M. Hiyoshi, H. Satoh, Y. Sumiyoshi, K. Morinaga, K. Kodama, T. Suzuki, M. Nagai and T. Suzuki. 2008. Associations of environmental exposure to dioxins with prevalent diabetes among general inhabitants in Japan. Environ. Res. 108(1):63-68.

Umannova, L., J. Zatloukalova, M. Machala, P. Krcmar, Z. Majkova, B. Hennig, A. Kozubik and J. Vondracek. 2007. Tumor necrosis factor-alpha modulates effects of aryl hydrocarbon receptor ligands on cell proliferation and expression of cytochrome P450 enzymes in rat liver "stem-like" cells. Toxicol. Sci. 99(1):79-89.

Uno, S., T.P. Dalton, H.G. Shertzer, M.B. Genter, D. Warshawsky, G. Talaska and D.W. Nebert. 2001. Benzo[a]pyrene-induced toxicity: paradoxical protection in Cyp1a1(-/-) knockout mice having increased hepatic BaP-DNA adduct levels. Biochem. Biophys. Res. Commun. 289(5):1049-1056.

Uno, S., N. Dragin, M.L. Miller, T.P. Dalton, F.J. Gonzalez and D.W. Nebert. 2008. Basal and inducible CYP1 mRNA quantitation and protein localization throughout the mouse gastrointestinal tract. Free Radic. Biol. Med. 44(4):570-583.

Vaiman, A.V., N.P. Shcherbak and V.A. Kobliakov. 2004. Cytochrome P-450 family 1 in rat embryo cell culture immortalized by Rausher leukemia virus. Biochemistry (Mosc.). 69(3):306-310.

Valdez, K.E. and B.K. Petroff. 2004. Potential roles of the aryl hydrocarbon receptor in female reproductive senescence. Reprod. Biol. 4(3):243-258.

Valet, G. and C. Esser. 2006. Data sieving analysis as a novel method to asses immunotoxic exposure to dioxins retrospectively. Int. Immunopharmacol. 6IM(8):1374-1375.

van Birgelen, A.P. and M. Van den Berg. 2000. Toxicokinetics. Food Addit. Contam. 17(4):267-273.

van den Hazel, P., M. Zuurbier, W. Babisch, A. Bartonova, M.L. Bistrup, G. Bolte, C. Busby, M. Butter, S. Ceccatelli, A. Fucic, W. Hanke, C. Johansson, M. Kohlhuber, M. Leijs, C. Lundqvist, H. Moshammer, R. Naginiene, A. Preece, R. Ronchetti, G. Salines, M. Saunders, G. Schoeters, N. Stilianakis, G. ten Tusscher and J.G. Koppe. 2006. Today's epidemics in children: possible relations to environmental pollution and suggested preventive measures. Acta Paediatr. Suppl. 95(453):18-25.

van Duursen, M.B., J.T. Sanderson, M. van der Bruggen, J. van der Linden and M. Van den Berg. 2003. Effects of several dioxin-like compounds on estrogen metabolism in the malignant MCF-7 and nontumorigenic MCF-10A human mammary epithelial cell lines. Toxicol. Appl. Pharmacol. 190(3):241-250.

van Duursen, M.B., J.T. Sanderson and M. Van den Berg. 2005. Cytochrome P450 1A1 and 1B1 in human blood lymphocytes are not suitable as biomarkers of exposure to dioxin-like compounds: polymorphisms and interindividual variation in expression and inducibility. Toxicol. Sci. 85(1):703-712.

van Grevenynghe, J., L. Sparfel, M. Le Vee, D. Gilot, B. Drenou, R. Fauchet and O. Fardel. 2004. Cytochrome P450-dependent toxicity of environmental polycyclic aromatic hydrocarbons towards human macrophages. Biochem. Biophys. Res. Commun. 317(3):708-716.

van Grevenynghe, J., M. Bernard, S. Langouet, C. Le Berre, T. Fest and O. Fardel. 2005. Human CD34-positive hematopoietic stem cells constitute targets for carcinogenic polycyclic aromatic hydrocarbons. J. Pharmacol. Exp. Ther. 314(2):693-702.

van Loveren, H., J. Vos, E. Putman and A. Piersma. 2003. Immunotoxicological consequences of perinatal chemical exposures: a plea for inclusion of immune parameters in reproduction studies. Toxicology. 185(3):185-191.

Van Maele-Fabry, G., S. Duhayon, C. Mertens and D. Lison. 2008. Risk of leukaemia among pesticide manufacturing workers: a review and meta-analysis of cohort studies. Environ. Res. 106(1):121-137.

Vasiliou, V., T. Buetler, D.L. Eaton and D.W. Nebert. 2000. Comparison of oxidative stress response parameters in newborn mouse liver versus simian virus 40 (SV40)-transformed hepatocyte cell lines. Biochem. Pharmacol. 59(6):703-712.

Vasquez, A., N. Atallah-Yunes, F.C. Smith, X. You, S.E. Chase, A.E. Silverstone and K.L. Vikstrom. 2003. A role for the aryl hydrocarbon receptor in cardiac physiology and function as demonstrated by AhR knockout mice. Cardiovasc. Toxicol. 3(2):153-163.

Veldhoen, M., K. Hirota, A.M. Westendorf, J. Buer, L. Dumoutier, J.C. Renauld and B. Stockinger. 2008. The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to environmental toxins. Nature. 453(7191):106-109.

Vezina, C.M., N.J. Walker and J.R. Olson. 2004. Subchronic exposure to TCDD, PeCDF, PCB126, and PCB153: effect on hepatic gene expression. Environ. Health Perspect. 112(16):1636-1644.

Vezina, C.M., T.M. Lin and R.E. Peterson. 2008. AHR signaling in prostate growth, morphogenesis, and disease. Biochem. Pharmacol.

Vidal, J.D., C.A. VandeVoort, C.B. Marcus, N.R. Lazarewicz and A.J. Conley. 2005. 2,3,7,8-tetrachlorodibenzo-p-dioxin induces CYP1B1 expression in human luteinized granulosa cells. Arch. Biochem. Biophys. 439(1):53-60.

Villano, C.M., K.A. Murphy, A. Akintobi and L.A. White. 2006. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) induces matrix metalloproteinase (MMP) expression and invasion in A2058 melanoma cells. Toxicol. Appl. Pharmacol. 210(3):212-224.

Vinggaard, A.M., C. Hnida and J.C. Larsen. 2000. Environmental polycyclic aromatic hydrocarbons affect androgen receptor activation in vitro. Toxicology. 145(2-3):173-183.

Vogel, C.F. and F. Matsumura. 2003. Interaction of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) with induced adipocyte differentiation in mouse embryonic fibroblasts (MEFs) involves tyrosine kinase c-Src. Biochem. Pharmacol. 66(7):1231-1244.

Vogel, C.F. and F. Matsumura. 2008. A new cross-talk between the aryl hydrocarbon receptor and RelB, a member of the NF-kappaB family. Biochem. Pharmacol.

Vogel, C., A.M. Boerboom, C. Baechle, C. El-Bahay, R. Kahl, G.H. Degen and J. Abel. 2000. Regulation of prostaglandin endoperoxide H synthase-2 induction by dioxin in rat hepatocytes: possible c-Src-mediated pathway. Carcinogenesis. 21(12):2267-2274.

Vogel, C.F., E. Sciullo, S. Park, C. Liedtke, C. Trautwein and F. Matsumura. 2004. Dioxin increases C/EBPbeta transcription by activating cAMP/protein kinase A. J. Biol. Chem. 279(10):8886-8894.

Vogel, C.F., W. Li, E. Sciullo, J. Newman, B. Hammock, J.R. Reader, J. Tuscano and F. Matsumura. 2007a. Pathogenesis of aryl hydrocarbon receptor-mediated development of lymphoma is associated with increased cyclooxygenase-2 expression. Am. J. Pathol. 171(5):1538-1548.

Vogel, C.F., E. Sciullo and F. Matsumura. 2007b. Involvement of RelB in aryl hydrocarbon receptor-mediated induction of chemokines. Biochem. Biophys. Res. Commun. 363(3):722-726.

Vogel, C.F., E. Sciullo, W. Li, P. Wong, G. Lazennec and F. Matsumura. 2007c. RelB, a new partner of aryl hydrocarbon receptor-mediated transcription. Mol. Endocrinol. 21(12):2941-2955.

Vogel, C.F., S.R. Goth, B. Dong, I.N. Pessah and F. Matsumura. 2008. Aryl hydrocarbon receptor signaling mediates expression of indoleamine 2,3-dioxygenase. Biochem. Biophys. Res. Commun. 375(3):331-335.

Vrzal, R., J. Ulrichova and Z. Dvorak. 2004. Aromatic hydrocarbon receptor status in the metabolism of xenobiotics under normal and pathophysiological conditions. Biomed. Pap. Med. Fac. Univ Palacky. Olomouc. Czech. Repub. 148(1):3-10.

Vrzal, R., M. Daujat-Chavanieu, J.M. Pascussi, J. Ulrichova, P. Maurel and Z. Dvorak. 2008. Microtubules-interfering agents restrict aryl hydrocarbon receptor-mediated CYP1A2 induction in primary cultures of human hepatocytes via c-jun-N-terminal kinase and glucocorticoid receptor. Eur. J. Pharmacol. 581(3):244-254.

Walisser, J.A., E. Glover, K. Pande, A.L. Liss and C.A. Bradfield. 2005. Aryl hydrocarbon receptor-dependent liver development and hepatotoxicity are mediated by different cell types. Proc. Natl. Acad. Sci. U. S. A. 102(49):17858-17863.

Walker, N.J. 2007. Unraveling the complexities of the mechanism of action of dioxins. Toxicol. Sci. 95(2):297-299.

Wang, S. and O. Hankinson. 2002. Functional involvement of the Brahma/SWI2-related gene 1 protein in cytochrome P4501A1 transcription mediated by the aryl hydrocarbon receptor complex. J. Biol. Chem. 277(14):11821-11827.

Wang, S., K. Ge, R.G. Roeder and O. Hankinson. 2004a. Role of mediator in transcriptional activation by the aryl hydrocarbon receptor. J. Biol. Chem. 279(14):13593-13600.

Wang, S.L., C.Y. Lin, Y.L. Guo, L.Y. Lin, W.L. Chou and L.W. Chang. 2004b. Infant exposure to polychlorinated dibenzo-p-dioxins, dibenzofurans and biphenyls (PCDD/Fs, PCBs)--correlation between prenatal and postnatal exposure. Chemosphere. 54(10):1459-1473.

Warner, M., B. Eskenazi, D.G. Patterson, G. Clark, W.E. Turner, L. Bonsignore, P. Mocarelli and P.M. Gerthoux. 2005. Dioxin-Like TEQ of women from the Seveso, Italy area by ID-HRGC/HRMS and CALUX. J. Expo. Anal. Environ. Epidemiol. 15(4):310-318.

Watanabe, T., I. Imoto, Y. Kosugi, Y. Fukuda, J. Mimura, Y. Fujii, K. Isaka, M. Takayama, A. Sato and J. Inazawa. 2001. Human arylhydrocarbon receptor repressor (AHRR) gene: genomic structure and analysis of polymorphism in endometriosis. J. Hum. Genet. 46(6):342-346.

Watanabe, H., A. Suzuki, M. Goto, S. Ohsako, C. Tohyama, H. Handa and T. Iguchi. 2004a. Comparative uterine gene expression analysis after dioxin and estradiol administration. J. Mol. Endocrinol. 33(3):763-771.

Watanabe, M., K. Sueoka, I. Sasagawa, A. Nakabayashi, Y. Yoshimura and T. Ogata. 2004b. Association of male infertility with Pro185Ala polymorphism in the aryl hydrocarbon receptor repressor gene: implication for the susceptibility to dioxins. Fertil. Steril. 82 Suppl 3:1067-1071.

Wei, C., R.J. Caccavale, E.H. Weyand, S. Chen and M.M. Iba. 2002. Induction of CYP1A1 and CYP1A2 expressions by prototypic and atypical inducers in the human lung. Cancer Lett. 178(1):25-36.

Wei, X.Y., C.Y. Leung, C.K. Wong, X.L. Shen, R.N. Wong, Z.W. Cai and N.K. Mak. 2005. Bisindigotin, a TCDD antagonist from the Chinese medicinal herb Isatis indigotica. J. Nat. Prod. 68(3):427-429.

Weiss, J., O. Papke, A. Bignert, S. Jensen, E. Greyerz, C. Agostoni, R. Besana, E. Riva, M. Giovannini and R. Zetterstrom. 2003. Concentrations of dioxins and other organochlorines (PCBs, DDTs, HCHs) in human milk from Seveso, Milan and a Lombardian rural area in Italy: a study performed 25 years after the heavy dioxin exposure in Seveso. Acta Paediatr. 92(4):467-472.

Weiss, C., D. Faust, H. Durk, S.K. Kolluri, A. Pelzer, S. Schneider, C. Dietrich, F. Oesch and M. Gottlicher. 2005. TCDD induces c-jun expression via a novel Ah (dioxin) receptor-mediated p38-MAPK-dependent pathway. Oncogene. 24(31):4975-4983.

Weiss, C., D. Faust, I. Schreck, A. Ruff, T. Farwerck, A. Melenberg, S. Schneider, B. Oesch-Bartlomowicz, J. Zatloukalova, J. Vondracek, F. Oesch and C. Dietrich. 2008. TCDD deregulates contact inhibition in rat liver oval cells via Ah receptor, JunD and cyclin A. Oncogene. 27(15):2198-2207.

Wejheden, C., S. Brunnberg, A. Hanberg and P.M. Lind. 2006. Osteopontin: a rapid and sensitive response to dioxin exposure in the osteoblastic cell line UMR-106. Biochem. Biophys. Res. Commun. 341(1):116-120.

Werner, K., W.R. Schaefer, H. Schweer, W.R. Deppert, U. Karck and H.P. Zahradnik. 2002. Characterization and identification of cytochrome P450 metabolites of arachidonic acid released by human peritoneal macrophages obtained from the pouch of Douglas. Prostaglandins Leukot. Essent. Fatty Acids. 67(6):397-404.

Wiberg, K., A. Aberg, T.E. McKone, M. Tysklind, A. Hanberg and M. MacLeod. 2007. Model selection and evaluation for risk assessment of dioxin-contaminated sites. Ambio. 36(6):458-466.

Widerak, M., C. Ghoneim, M.F. Dumontier, M. Quesne, M.T. Corvol and J.F. Savouret. 2006. The aryl hydrocarbon receptor activates the retinoic acid receptoralpha through SMRT antagonism. Biochimie. 88(3-4):387-397.

Wiesmuller, G.A., R. Eckard, L. Dobler, A. Gunsel, M. Oganowski, C. Schroter-Kermani, C. Schluter, A. Gies and F.H. Kemper. 2007. The Environmental Specimen Bank for Human Tissues as part of the German Environmental Specimen Bank. Int. J. Hyg. Environ. Health. 210(3-4):299-305.

Wigle, D.T., T.E. Arbuckle, M. Walker, M.G. Wade, S. Liu and D. Krewski. 2007. Environmental hazards: evidence for effects on child health. J. Toxicol. Environ. Health B Crit Rev. 10(1-2):3-39.

Williams, S.R., D.S. Son and P.F. Terranova. 2004. Protein kinase C delta is activated in mouse ovarian surface epithelial cancer cells by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Toxicology. 195(1):1-17.

Williamson, M.A., T.A. Gasiewicz and L.A. Opanashuk. 2005. Aryl hydrocarbon receptor expression and activity in cerebellar granule neuroblasts: implications for development and dioxin neurotoxicity. Toxicol. Sci. 83(2):340-348.

Wilson, C.L. 2004. Molecular switch circuits in toxicology: a dimmer switch for dioxin. Toxicol. Sci. 78(2):178-180.

Wittsiepe, J., P. Schrey, U. Ewers, F. Selenka and M. Wilhelm. 2000. Decrease of PCDD/F levels in human blood from Germany over the past ten years (1989-1998). Chemosphere. 40(9-11):1103-1109.

Wittsiepe, J., P. Furst, P. Schrey, F. Lemm, M. Kraft, G. Eberwein, G. Winneke and M. Wilhelm. 2007. PCDD/F and dioxin-like PCB in human blood and milk from German mothers. Chemosphere. 67(9):S286-S294.

Wojtowicz, A., M. Tomanek, K. Augustowska and E.L. Gregoraszczuk. 2005. Aromatic hydrocarbon receptor (AhR) in the porcine theca and granulosa cells: effect of TCDD, PCB 126 and PCB 153 on the expression of AhR. Endocr. Regul. 39(4):109-118.

Wolfle, D. 2003. Enhancement of carcinogen-induced malignant cell transformation by prostaglandin F(2 alpha). Toxicology. 188(2-3):139-147.

Wolfle, D., S. Marotzki, D. Dartsch, W. Schafer and H. Marquardt. 2000. Induction of cyclooxygenase expression and enhancement of malignant cell transformation by 2,3,7,8-tetrachlorodibenzo-p-dioxin. Carcinogenesis. 21(1):15-21.

Wong, J.M., P.A. Harper, U.A. Meyer, K.W. Bock, K. Morike, J. Lagueux, P. Ayotte, R.F. Tyndale, E.M. Sellers, D.K. Manchester and A.B. Okey. 2001a. Ethnic variability in the allelic distribution of human aryl hydrocarbon receptor codon 554 and assessment of variant receptor function in vitro. Pharmacogenetics. 11(1):85-94.

Wong, J.M., A.B. Okey and P.A. Harper. 2001b. Human aryl hydrocarbon receptor polymorphisms that result in loss of CYP1A1 induction. Biochem. Biophys. Res. Commun. 288(4):990-996.

Wormke, M., E. Castro-Rivera, I. Chen and S. Safe. 2000a. Estrogen and aryl hydrocarbon receptor expression and crosstalk in human Ishikawa endometrial cancer cells. J. Steroid Biochem. Mol. Biol. 72(5):197-207.

Wormke, M., M. Stoner, B. Saville and S. Safe. 2000b. Crosstalk between estrogen receptor alpha and the aryl hydrocarbon receptor in breast cancer cells involves unidirectional activation of proteasomes. FEBS Lett. 478(1-2):109-112.

Wormke, M., M. Stoner, B. Saville, K. Walker, M. Abdelrahim, R. Burghardt and S. Safe. 2003. The aryl hydrocarbon receptor mediates degradation of estrogen receptor alpha through activation of proteasomes. Mol. Cell Biol. 23(6):1843-1855.

Wormley, D.D., A. Ramesh and D.B. Hood. 2004. Environmental contaminant-mixture effects on CNS development, plasticity, and behavior. Toxicol. Appl. Pharmacol. 197(1):49-65.

Wu, Q., S. Ohsako, T. Baba, K. Miyamoto and C. Tohyama. 2002. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on preimplantation mouse embryos. Toxicology. 174(2):119-129.

Wu, R., L. Zhang, M.S. Hoagland and H.I. Swanson. 2007. Lack of the aryl hydrocarbon receptor leads to impaired activation of AKT/protein kinase B and enhanced sensitivity to apoptosis induced via the intrinsic pathway. J. Pharmacol. Exp. Ther. 320(1):448-457.

Wu, Q., J.S. Suzuki, H. Zaha, T.M. Lin, R.E. Peterson, C. Tohyama and S. Ohsako. 2008. Differences in gene expression and benzo[a]pyrene-induced DNA adduct formation in the liver of three strains of female mice with identical AhR(b2) genotype treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin and/or benzo[a]pyrene. J. Appl. Toxicol. 28(6):724-733.

Wyman, A., A.L. Lavin, G.E. Wilding and T.A. Gasiewicz. 2002. 2,3,7,8-tetrachlorodibenzo-p-dioxin does not directly alter the phenotype of maturing B cells in a murine coculture system. Toxicol. Appl. Pharmacol. 180(3):164-177.

Xie, A., N.J. Walker and D. Wang. 2006. Dioxin (2,3,7,8-tetrachlorodibenzo-p-dioxin) enhances triggered afterdepolarizations in rat ventricular myocytes. Cardiovasc. Toxicol. 6(2):99-110.

Xu, L., A.P. Li, D.L. Kaminski and M.F. Ruh. 2000. 2,3,7,8 Tetrachlorodibenzo-p-dioxin induction of cytochrome P4501A in cultured rat and human hepatocytes. Chem. Biol. Interact. 124(3):173-189.

Yamada-Okabe, T., T. Aono, H. Sakai, Y. Kashima and H. Yamada-Okabe. 2004. 2,3,7,8-tetrachlorodibenzo-p-dioxin augments the modulation of gene expression mediated by the thyroid hormone receptor. Toxicol. Appl. Pharmacol. 194(3):201-210.

Yamano, Y., K. Ohyama, M. Ohta, J. Nakamura and I. Morishima. 2005a. Expression of small stress protein hsp20 gene in the maturing rat testis. J. Vet. Med. Sci. 67(11):1181-1184.

Yamano, Y., K. Ohyama, M. Ohta, T. Sano, A. Ritani, J. Shimada, N. Ashida, E. Yoshida, K. Ikehara and I. Morishima. 2005b. A novel spermatogenesis related factor-2 (SRF-2) gene expression affected by TCDD treatment. Endocr. J. 52(1):75-81.

Yang, F. and D. Bleich. 2004. Transcriptional regulation of cyclooxygenase-2 gene in pancreatic beta-cells. J. Biol. Chem. 279(34):35403-35411.

Yang, D., Y. Li, X. Yuan, L. Matoney and B. Yan. 2001. Regulation of rat carboxylesterase expression by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD): a dose-dependent decrease in mRNA levels but a biphasic change in protein levels and activity. Toxicol. Sci. 64(1):20-27.

Yang, C., F. Boucher, A. Tremblay and J.L. Michaud. 2004. Regulatory interaction between arylhydrocarbon receptor and SIM1, two basic helix-loop-helix PAS proteins involved in the control of food intake. J. Biol. Chem. 279(10):9306-9312.

Yang, Y.M., D.Y. Huang, G.F. Liu, J.C. Zhong, K. Du, Y.F. Li and X.H. Song. 2005. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on vitamin A metabolism in mice. J. Biochem. Mol. Toxicol. 19(5):327-335.

Yang, X., S. Solomon, L.R. Fraser, A.F. Trombino, D. Liu, G.E. Sonenshein, E.V. Hestermann and D.H. Sherr. 2008. Constitutive regulation of CYP1B1 by the aryl hydrocarbon receptor (AhR) in pre-malignant and malignant mammary tissue. J. Cell Biochem. 104(2):402-417.

Yerokun, T. and J. Stewart. 2006. Novel functional association of serine palmitoyltransferase subunit 1-A peptide in sphingolipid metabolism with cytochrome P4501A1 transactivation and proliferative capacity of the human Glioma LN18 brain tumor cell line. Int. J. Environ. Res. Public Health. 3(3):252-261.

Yim, S., M. Oh, S.M. Choi and H. Park. 2004. Inhibition of the MEK-1/p42 MAP kinase reduces aryl hydrocarbon receptor-DNA interactions. Biochem. Biophys. Res. Commun. 322(1):9-16.

Yoo, B.S., D.R. Boverhof, D. Shnaider, R.B. Crawford, T.R. Zacharewski and N.E. Kaminski. 2004. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) alters the regulation of Pax5 in lipopolysaccharide-activated B cells. Toxicol. Sci. 77(2):272-279.

Yoshida, K. and J. Nakanishi. 2003. Estimation of dioxin risk to Japanese from the past to the future. Chemosphere. 53(4):427-436.

Young, A.L. 2002. The volunteers: the first human biopsy studies of TCDD from Agent Orange exposure. Environ. Sci. Pollut. Res. Int. 9(3):157.

Young, J.G., N.P. Jewell and S.J. Samuels. 2008. Regression analysis of a disease onset distribution using diagnosis data. Biometrics. 64(1):20-28.

Yu, J., Y. Wang, W.H. Zhou, L. Wang, Y.Y. He and D.J. Li. 2008. Combination of estrogen and dioxin is involved in the pathogenesis of endometriosis by promoting chemokine secretion and invasion of endometrial stromal cells. Hum. Reprod. 23(7):1614-1626.

Yueh, M.F., Y.H. Huang, A. Hiller, S. Chen, N. Nguyen and R.H. Tukey. 2003. Involvement of the xenobiotic response element (XRE) in Ah receptor-mediated induction of human UDP-glucuronosyltransferase 1A1. J. Biol. Chem. 278(17):15001-15006.

Yueh, M.F., J.A. Bonzo and R.H. Tukey. 2005. The role of Ah receptor in induction of human UDP-glucuronosyltransferase 1A1. Methods Enzymol. 400:75-91.

Zambon, P. 2007. [Dioxins and sarcomas in Veneto: the context, the method and the peer review]. Epidemiol. Prev. 31(2-3):76-77.

Zatloukalova, J., L. Svihalkova-Sindlerova, A. Kozubik, P. Krcmar, M. Machala and J. Vondracek. 2007. beta-Naphthoflavone and 3'-methoxy-4'-nitroflavone exert ambiguous effects on Ah receptor-dependent cell proliferation and gene expression in rat liver 'stem-like' cells. Biochem. Pharmacol. 73(10):1622-1634.

Zeytun, A., R.J. McKallip, M. Fisher, I. Camacho, M. Nagarkatti and P.S. Nagarkatti. 2002. Analysis of 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced gene expression profile in vivo using pathway-specific cDNA arrays. Toxicology. 178(3):241-260.

Zhang, N. and M.K. Walker. 2007. Crosstalk between the aryl hydrocarbon receptor and hypoxia on the constitutive expression of cytochrome P4501A1 mRNA. Cardiovasc. Toxicol. 7(4):282-290.

Zhang, Z.R., S.Q. Xu, Y.K. Zhou, S. Ren and B. Lu. 2001. [Construction of luciferase reporter plasmid which is under the control of dioxin-responsive enhancers]. Sheng Wu Gong. Cheng Xue. Bao. 17(2):170-174.

Zhang, L., W. Zheng and C.R. Jefcoate. 2003a. Ah receptor regulation of mouse Cyp1B1 is additionally modulated by a second novel complex that forms at two AhR response elements. Toxicol. Appl. Pharmacol. 192(2):174-190.

Zhang, S., C. Qin and S.H. Safe. 2003b. Flavonoids as aryl hydrocarbon receptor agonists/antagonists: effects of structure and cell context. Environ. Health Perspect. 111(16):1877-1882.

Zhang, Z.R., S.Q. Xu, X. Sun, Y.J. Xu, X.K. Cai, Z.W. Liu, X.L. Tan, Y.K. Zhou, J.Y. Zhang and H. Yan. 2003c. Characteristics and application of established luciferase hepatoma cell line that responds to dioxin-like chemicals. World J. Gastroenterol. 9(7):1460-1464.

Zhang, W., B. Moorthy, M. Chen, K. Muthiah, R. Coffee, A.F. Purchio and D.B. West. 2004. A Cyp1a2-luciferase transgenic CD-1 mouse model: responses to aryl hydrocarbons similar to the humanized AhR mice. Toxicol. Sci. 82(1):297-307.

Zhang, Z.Y., R.D. Pelletier, Y.N. Wong, M. Sugawara, N. Zhao and B.A. Littlefield. 2006. Preferential inducibility of CYP1A1 and CYP1A2 by TCDD: differential regulation in primary human hepatocytes versus transformed human cells. Biochem. Biophys. Res. Commun. 341(2):399-407.

Zhang, L., R. Wu, R.W. Dingle, C.G. Gairola, J. Valentino and H.I. Swanson. 2007. Cigarette smoke condensate and dioxin suppress culture shock induced senescence in normal human oral keratinocytes. Oral Oncol. 43(7):693-700.

Zhang, S., C. Rowlands and S. Safe. 2008. Ligand-dependent interactions of the Ah receptor with coactivators in a mammalian two-hybrid assay. Toxicol. Appl. Pharmacol. 227(2):196-206.

Zhao, D., E.A. Pritts, V.A. Chao, J.F. Savouret and R.N. Taylor. 2002. Dioxin stimulates RANTES expression in an in-vitro model of endometriosis. Mol. Hum. Reprod. 8(9):849-854.

Zheng, W., P.B. Brake, K.K. Bhattacharyya, L. Zhang, D. Zhao and C.R. Jefcoate. 2003. Cell selective cAMP induction of rat CYP1B1 in adrenal and testis cells. Identification of a novel cAMP-responsive far upstream enhancer and a second Ah receptor-dependent mechanism. Arch. Biochem. Biophys. 416(1):53-67.

Zhou, J. and T.A. Gasiewicz. 2003. 3'-methoxy-4'-nitroflavone, a reported aryl hydrocarbon receptor antagonist, enhances Cyp1a1 transcription by a dioxin responsive element-dependent mechanism. Arch. Biochem. Biophys. 416(1):68-80.

Zhou, J.G., E.C. Henry, C.M. Palermo, S.D. Dertinger and T.A. Gasiewicz. 2003. Species-specific transcriptional activity of synthetic flavonoids in guinea pig and mouse cells as a result of differential activation of the aryl hydrocarbon receptor to interact with dioxin-responsive elements. Mol. Pharmacol. 63(4):915-924.