

Draft Selection Criteria to identify Key *In Vivo* Mammalian Studies that Inform Dose-Response Modeling for 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)^a

Study Feature	Selection Rationale		
	<i>Primary^b</i>	<i>Secondary^c</i>	<i>Currently Excluded</i>
Chemical, purity, matrix/medium	TCDD-only doses included, purity specified, matrix in which TCDD is administered is identified	TCDD purity or matrix not clearly identified	Studies of dioxin-like compounds (DLCs) or mixtures
Peer review	Independently peer-reviewed, publicly available	Supplementary materials accompanying peer-reviewed publication	Not formally peer-reviewed; literature not publicly available
Study design, execution, and reporting	Clearly documented and consistent with standard toxicological principles, testing protocols, and practice (i.e., endpoint-appropriate, particularly for negative findings)	Testing protocol provides incomplete coverage of relevant endpoint-specific measures, particularly for negative findings	Studies not meeting standard principles and practices
Study subject: species, strain, and sensitivity for given endpoint; litter; life stage; gender	Mammalian species Strain and gender identified Animal age at beginning of treatment identified Litter confounders (within/between) accounted for	Mammalian species, <i>in vivo</i> , but only studying an artificially sensitive subject (e.g., knockout mouse)	Non-mammalian or not <i>in vivo</i>
Exposure route	Oral	Parenteral (e.g., intravenous, intramuscular, intraperitoneal, subcutaneous)	Inhalation, dermal, ocular
Dose level	Lowest dose ≤ 200 ng/kg-d for noncancer endpoints and $\leq 1\mu\text{g}$ /kg-d for cancer	Lowest dose > 200 ng/kg-d for noncancer endpoints, or > 1.0 μg /kg-d for cancer	
Exposure frequency, duration, and timing	Dosing regimen characterized and explained		Characterization/explanation missing or cannot be determined
Controls	Appropriate and well characterized	Effect reported, but with no negative control	
Response	Effect relevant to human health Magnitude outside range of normal variability	Precursor effects, or adaptive responses potentially relevant to human health	Lethality
Statistical evaluation	Clearly described and appropriate to the endpoint and study design (e.g., per error variance, magnitude of effect)	Limited statistical context	

^a NAS (2006) commented that the selection of data sets for quantitative dose-response modeling needed to be more transparent. These draft criteria are offered for consideration at the kickoff workshop. These criteria would be used to identify candidate studies of non-human mammals that would be used to define the point-of-departure (POD). These criteria are not designed for hazard identification or weight-of-evidence determinations. Studies addressing data other than direct TCDD dose-response in mammals (including toxicokinetic data on absorption, distribution, metabolism, or elimination; information on physiologically-based pharmacokinetic [PBPK] modeling, and mode of action data) will be evaluated separately.

^b Presents preliminary draft criteria for evaluating a study being considered for estimating a POD in a TCDD dose-response model.

^c Presents preliminary draft criteria that could qualify a study as primary with support from other lines of evidence (e.g., PBPK modeling), when no study for an endpoint meets the "primary" criteria.