



## **APPENDIX F**

### **Comparison of Arthropod and Non-Arthropod Invertebrate SSDs**


## Comparing arthropod and non-arthropod invertebrate cadmium LC50 SSDs in moderately hard water over moderate (1-3 days) exposure at T>15C




### Comparing arthropod and non-arthropod invertebrate cadmium LC50 SSDs in soft water over long (3-30 days) exposure at T>15C




**Comparing arthropod and non-arthropod invertebrate cadmium LC50 SSDs  
in soft water over moderate (1-3 days) exposure at T>15C**




## Comparing arthropod and non-arthropod invertebrate cadmium LC50 SSDs in soft water over short ( $\leq 1$ day) exposure at $T > 15^{\circ}\text{C}$




### Comparing arthropod and non-arthropod invertebrate chromium LC50 SSDs in soft water over long (3-30 days) exposure at T>15C



## Comparing arthropod and non-arthropod invertebrate chromium LC50 SSDs in soft water over moderate (1-3 days) exposure at T>15C




### Comparing arthropod and non-arthropod invertebrate chromium LC50 SSDs in soft water over short ( $\leq 1$ day) exposure at $T > 15C$







# Comparing arthropod and non-arthropod invertebrate copper LC50 SSDs in moderately hard water over moderate (1-3 days) exposure at T>15C




## Comparing arthropod and non-arthropod invertebrate copper LC50 SSDs in soft water over long (3-30 days) exposure at T>15C




# Comparing arthropod and non-arthropod invertebrate copper LC50 SSDs in soft water over moderate (1-3 days) exposure at T>15C




# Comparing arthropod and non-arthropod invertebrate copper LC50 SSDs in soft water over short ( $\leq 1$ day) exposure at $T > 15^{\circ}\text{C}$




# Comparing arthropod and non-arthropod invertebrate zinc LC50 SSDs in moderately hard water over moderate (1-3 days) exposure at T>15C




# Comparing arthropod and non-arthropod invertebrate zinc LC50 SSDs in soft water over long (3-30 days) exposure at T>15C



# Comparing arthropod and non-arthropod invertebrate zinc LC50 SSDs in soft water over moderate (1-3 days) exposure at T<=15C



# Comparing arthropod and non-arthropod invertebrate zinc LC50 SSDs in soft water over moderate (1-3 days) exposure at T>15C





# Comparing arthropod and non-arthropod invertebrate zinc LC50 SSDs in soft water over short ( $\leq 1$ day) exposure at $T > 15C$

