



APPENDIX G
Comparison of Hardness Effect on SSDs


Cadmium LC50 SSDs - comparing hardness
for Arthropod species at T<=15C over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing hardness for Arthropod species at T>15C over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing hardness for Arthropod species at T>15C over moderate (1-3 days) exposure




Cadmium LC50 SSDs - comparing hardness
for Invertebrate species at $T \leq 15^\circ\text{C}$ over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing hardness for Invertebrate species at T>15C over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over moderate (1-3 days) exposure




Cadmium LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over short (<=1 day) exposure




Cadmium LC50 SSDs - comparing hardness for Non-Arthropod Invertebrate species at T>15C over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing hardness for Non-Arthropod Invertebrate species at T>15C over moderate (1-3 days) exposure




Cadmium LC50 SSDs - comparing hardness
for Non-Arthropod Invertebrate species at T>15C over short (<=1 day) exposure




Cadmium LC50 SSDs - comparing hardness for Vertebrate species at T<=15C over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing hardness for Vertebrate species at T>15C over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing hardness for Vertebrate species at T>15C over moderate (1-3 days) exposure




Cadmium LC50 SSDs - comparing hardness
for Vertebrate species at T>15C over short (<=1 day) exposure




Chromium LC50 SSDs - comparing hardness
for Arthropod species at T>15C over short (<=1 day) exposure




Chromium LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over moderate (1-3 days) exposure




Chromium LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over short (<=1 day) exposure




Chromium LC50 SSDs - comparing hardness for Vertebrate species at T>15C over long (3-30 days) exposure




Copper LC50 SSDs - comparing hardness for Arthropod species at T>15C over moderate (1-3 days) exposure




Copper LC50 SSDs - comparing hardness for Invertebrate species at T>15C over long (3-30 days) exposure




Copper LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over moderate (1-3 days) exposure




Copper LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over short (<=1 day) exposure




Copper LC50 SSDs - comparing hardness for Non-Arthropod Invertebrate species at T>15C over long (3-30 days) exposure




Copper LC50 SSDs - comparing hardness for Non-Arthropod Invertebrate species at T>15C over moderate (1-3 days) exposure




Copper LC50 SSDs - comparing hardness
for Non-Arthropod Invertebrate species at T>15C over short (<=1 day) exposure




Copper LC50 SSDs - comparing hardness for Vertebrate species at T<=15C over long (3-30 days) exposure




Copper LC50 SSDs - comparing hardness for Vertebrate species at T>15C over long (3-30 days) exposure




Copper LC50 SSDs - comparing hardness for Vertebrate species at T>15C over moderate (1-3 days) exposure




Copper LC50 SSDs - comparing hardness
for Vertebrate species at T>15C over short (<=1 day) exposure




Lead LC50 SSDs - comparing hardness for Vertebrate species at T>15C over long (3-30 days) exposure




Mercury LC50 SSDs - comparing hardness for Invertebrate species at T>15C over long (3-30 days) exposure




Mercury LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over moderate (1-3 days) exposure




Mercury LC50 SSDs - comparing hardness
for Invertebrate species at T>15C over short (<=1 day) exposure




Mercury LC50 SSDs - comparing hardness for Vertebrate species at T>15C over long (3-30 days) exposure




Mercury LC50 SSDs - comparing hardness
for Vertebrate species at T>15C over moderate (1-3 days) exposure




Mercury LC50 SSDs - comparing hardness
for Vertebrate species at T>15C over short (<=1 day) exposure




Nickel LC50 SSDs - comparing hardness for Invertebrate species at T>15C over long (3-30 days) exposure




Nickel LC50 SSDs - comparing hardness for Invertebrate species at T>15C over short (<=1 day) exposure




Nickel LC50 SSDs - comparing hardness
for Non-Arthropod Invertebrate species at T>15C over short (<=1 day) exposure




Nickel LC50 SSDs - comparing hardness for Vertebrate species at T>15C over long (3-30 days) exposure




Nickel LC50 SSDs - comparing hardness for Vertebrate species at T>15C over moderate (1-3 days) exposure




Nickel LC50 SSDs - comparing hardness for Vertebrate species at T>15C over short (<=1 day) exposure




Zinc LC50 SSDs - comparing hardness for Arthropod species at T>15C over moderate (1-3 days) exposure




Zinc LC50 SSDs - comparing hardness for Invertebrate species at T>15C over long (3-30 days) exposure




Zinc LC50 SSDs - comparing hardness for Invertebrate species at T>15C over moderate (1-3 days) exposure




Zinc LC50 SSDs - comparing hardness for Invertebrate species at T>15C over short (<=1 day) exposure




Zinc LC50 SSDs - comparing hardness for Non-Arthropod Invertebrate species at T>15C over long (3-30 days) exposure




Zinc LC50 SSDs - comparing hardness for Non-Arthropod Invertebrate species at T>15C over moderate (1-3 days) exposure




Zinc LC50 SSDs - comparing hardness
for Non-Arthropod Invertebrate species at T>15C over short (<=1 day) exposure



Zinc LC50 SSDs - comparing hardness for Vertebrate species at T>15C over long (3-30 days) exposure



Zinc LC50 SSDs - comparing hardness for Vertebrate species at T>15C over moderate (1-3 days) exposure



Zinc LC50 SSDs - comparing hardness
for Vertebrate species at T>15C over short (<=1 day) exposure

