



APPENDIX H
Comparison of Temperature Effect on SSDs


Arsenic LC50 SSDs - comparing temperature
for Arthropod species in water over moderate (1-3 days) exposure




Arsenic LC50 SSDs - comparing temperature for Invertebrate species in water over moderate (1-3 days) exposure




Arsenic LC50 SSDs - comparing temperature for Vertebrate species in water over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing temperature for Arthropod species in moderately hard water over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing temperature for Arthropod species in soft water over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing temperature
for Invertebrate species in moderately hard water over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing temperature
for Invertebrate species in moderately hard water over moderate (1-3 days) exposure




Cadmium LC50 SSDs - comparing temperature
for Invertebrate species in soft water over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing temperature for Vertebrate species in moderately hard water over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing temperature for Vertebrate species in soft water over long (3-30 days) exposure




Cadmium LC50 SSDs - comparing temperature
for Vertebrate species in very hard water over long (3-30 days) exposure




Chromium LC50 SSDs - comparing temperature
for Invertebrate species in soft water over moderate (1-3 days) exposure




Chromium LC50 SSDs - comparing temperature
for Invertebrate species in soft water over short (≤ 1 day) exposure




Copper LC50 SSDs - comparing temperature for Arthropod species in soft water over long (3-30 days) exposure




Copper LC50 SSDs - comparing temperature for Arthropod species in soft water over moderate (1-3 days) exposure




Copper LC50 SSDs - comparing temperature
for Invertebrate species in soft water over long (3-30 days) exposure




Copper LC50 SSDs - comparing temperature for Invertebrate species in soft water over moderate (1-3 days) exposure




Copper LC50 SSDs - comparing temperature
for Invertebrate species in soft water over short (≤ 1 day) exposure




Copper LC50 SSDs - comparing temperature for Vertebrate species in hard water over long (3-30 days) exposure




Copper LC50 SSDs - comparing temperature
for Vertebrate species in moderately hard water over long (3-30 days) exposure




Copper LC50 SSDs - comparing temperature for Vertebrate species in soft water over long (3-30 days) exposure




Copper LC50 SSDs - comparing temperature for Vertebrate species in soft water over moderate (1-3 days) exposure




Copper LC50 SSDs - comparing temperature
for Vertebrate species in soft water over short (≤ 1 day) exposure




Copper LC50 SSDs - comparing temperature for Vertebrate species in very hard water over long (3-30 days) exposure




Lead LC50 SSDs - comparing temperature
for Arthropod species in soft water over long (3-30 days) exposure




Lead LC50 SSDs - comparing temperature
for Invertebrate species in soft water over long (3-30 days) exposure




Mercury LC50 SSDs - comparing temperature
for Vertebrate species in moderately hard water over long (3-30 days) exposure




Selenium LC50 SSDs - comparing temperature for Vertebrate species in water over long (3-30 days) exposure




Zinc LC50 SSDs - comparing temperature for Arthropod species in soft water over moderate (1-3 days) exposure




Zinc LC50 SSDs - comparing temperature for Invertebrate species in soft water over long (3-30 days) exposure




Zinc LC50 SSDs - comparing temperature for Invertebrate species in soft water over moderate (1-3 days) exposure




Zinc LC50 SSDs - comparing temperature
for Invertebrate species in soft water over short (≤ 1 day) exposure




Zinc LC50 SSDs - comparing temperature
for Non-Arthropod Invertebrate species in soft water over moderate (1-3 days) exposure




Zinc LC50 SSDs - comparing temperature for Non-Arthropod Invertebrate species in soft water over short (≤ 1 day) exposure



Zinc LC50 SSDs - comparing temperature for Vertebrate species in soft water over long (3-30 days) exposure



Zinc LC50 SSDs - comparing temperature for Vertebrate species in soft water over moderate (1-3 days) exposure



Zinc LC50 SSDs - comparing temperature for Vertebrate species in soft water over short (≤ 1 day) exposure

