



TOXICOLOGICAL REVIEW

OF

DICHLOROMETHANE
(METHYLENE CHLORIDE)

(CAS No. 75-09-2)

**In Support of Summary Information on the
Integrated Risk Information System (IRIS)**

March 2010

NOTICE

This document is an **External Review draft**. This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by EPA. It does not represent and should not be construed to represent any Agency determination or policy. It is being circulated for review of its technical accuracy and science policy implications.

U.S. Environmental Protection Agency
Washington, DC

DISCLAIMER

This document is a preliminary draft for review purposes only. This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by EPA. It does not represent and should not be construed to represent any Agency determination or policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

CONTENTS—TOXICOLOGICAL REVIEW OF DICHLOROMETHANE
(CAS No. 75-09-2)

LIST OF TABLES	viii
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS AND ACRONYMS	xxi
FOREWORD	xxiii
AUTHORS, CONTRIBUTORS, AND REVIEWERS	xxiv
1. INTRODUCTION	1
2. CHEMICAL AND PHYSICAL INFORMATION	3
3. TOXICOKINETICS	5
3.1. ABSORPTION.....	5
3.1.1. Oral—Gastrointestinal Tract Absorption	5
3.1.2. Inhalation—Respiratory Tract Absorption.....	5
3.2. DISTRIBUTION.....	7
3.3. METABOLISM	9
3.3.1. The CYP2E1 Pathway.....	11
3.3.2. The GST Pathway.....	14
3.4. ELIMINATION	20
3.5. PHYSIOLOGICALLY BASED TOXICOKINETIC MODELS.....	21
3.5.1. Probabilistic Mouse PBPK Dichloromethane Model (Marino et al., 2006).....	27
3.5.2. Probabilistic Human PBPK Dichloromethane Model (David et al., 2006).....	31
3.5.3. Evaluation of Rat PBPK Dichloromethane Models	40
3.5.4. Comparison of Mouse, Rat, and Human PBPK Models	41
3.5.5. Uncertainties in PBPK Model Structure for the Mouse, Rat and Human	45
4. HAZARD IDENTIFICATION.....	50
4.1. STUDIES IN HUMANS.....	50
4.1.1. Introduction—Case Reports, Epidemiologic, and Clinical Studies	50
4.1.2. Noncancer Studies	50
4.1.2.1. Case Reports of Acute, High-dose Exposures	50
4.1.2.2. Controlled Experiments Examining Acute Effects.....	51
4.1.2.3. Observational Studies Focusing on Clinical Chemistries, Clinical Examinations, and Symptoms	52
4.1.2.4. Observational Studies Using Workplace Medical Program Data.....	59
4.1.2.5. Studies of Ischemic Heart Disease Mortality Risk	62
4.1.2.6. Studies of Suicide Risk.....	63
4.1.2.7. Studies of Infectious Disease Risk.....	64
4.1.2.8. Studies of Reproductive Outcomes.....	64
4.1.2.9. Summary of Noncancer Studies	67
4.1.3. Cancer Studies	69
4.1.3.1. Identification and Selection of Studies for Evaluation of Cancer Risk	69
4.1.3.2. Description of the Selected Studies	70
4.1.3.3. Cellulose Triacetate Film Base Production Cohorts.....	70
4.1.3.4. Cellulose Triacetate Fiber Production Cohorts.....	78
4.1.3.5. Solvent-Exposed Workers—Hill Air Force Base, Utah	83
4.1.3.6. Case-Control Studies of Specific Cancers and Dichloromethane	84
4.1.3.7. Summary of Cancer Studies by Type of Cancer.....	94

4.2. SUBCHRONIC AND CHRONIC STUDIES AND CANCER BIOASSAYS IN ANIMALS—ORAL AND INHALATION	100
4.2.1. Oral Exposure: Overview of Noncancer and Cancer Effects	100
4.2.1.1. Toxicity Studies of Subchronic Oral Exposures: Hepatic Effects	101
4.2.1.2. Toxicity Studies of Chronic Oral Exposures: Hepatic Effects and Carcinogenicity	105
4.2.2. Inhalation Exposure: Overview of Noncancer and Cancer Effects	112
4.2.2.1. Toxicity Studies of Subchronic Inhalation Exposures: General, Renal, and Hepatic Effects	113
4.2.2.2. Toxicity Studies from Chronic Inhalation Exposures	117
4.3. REPRODUCTIVE/DEVELOPMENTAL STUDIES—ORAL AND INHALATION	137
4.3.1. Reproductive Toxicity Studies	139
4.3.1.1. Gavage and Subcutaneous Injection Studies	139
4.3.1.2. Inhalation Studies	140
4.3.2. Developmental Toxicity Studies	141
4.3.2.1. Gavage Studies and Culture Studies	141
4.3.2.2. Inhalation Studies	142
4.4. OTHER DURATION- OR ENDPOINT-SPECIFIC STUDIES	144
4.4.1. Short-term (2-Week) Studies of General and Hepatic Effects in Animals	144
4.4.2. Immunotoxicity Studies in Animals	145
4.4.3. Neurotoxicology Studies in Animals	147
4.4.3.1. Neurotoxicology Studies—Oral Exposures	155
4.4.3.2. Neurotoxicology Studies—Inhalational Exposure	156
4.5. MECHANISTIC DATA AND OTHER STUDIES IN SUPPORT OF THE MODE OF ACTION	163
4.5.1. Genotoxicity Studies	163
4.5.1.1. In Vitro Genotoxicity Assays	163
4.5.1.2. In Vivo Genotoxicity Assays	174
4.5.2. Mechanistic Studies of Liver Effects	183
4.5.3. Mechanistic Studies of Lung Effects	187
4.5.4. Mechanistic Studies of Neurological Effects	192
4.6. SYNTHESIS OF MAJOR NONCANCER EFFECTS	194
4.6.1. Oral	194
4.6.1.1. Summary of Human Data	194
4.6.1.2. Summary of Animal Data	194
4.6.2. Inhalation	198
4.6.2.1. Summary of Human Data	198
4.6.2.2. Summary of Animal Studies	199
4.6.3. Mode-of-Action Information	207
4.6.3.1. Mode of Action for Nonneoplastic Liver Effects	207
4.6.3.2. Mode of Action for Nonneoplastic Lung Effects	208
4.6.3.3. Mode of Action for Neurological Effects	208
4.6.3.4. Mode of Action for Reproductive and Developmental Effects	209
4.6.3.5. Mode of Action for Immunotoxicity	210
4.7. EVALUATION OF CARCINOGENICITY	211
4.7.1. Summary of Overall Weight of Evidence	211
4.7.2. Synthesis of Human, Animal, and Other Supporting Evidence	212
4.7.3. Mode-of-Action Information	223

4.7.3.1. Hypothesized Mode of Action	223
4.7.3.2. General Conclusions About the Mode of Action for Tumors in Rodents and Relevance to Humans	230
4.8. SUSCEPTIBLE POPULATIONS AND LIFE STAGES	232
4.8.1. Possible Childhood Susceptibility	232
4.8.2. Possible Gender Differences	234
4.8.3. Other	234
5. DOSE-RESPONSE ASSESSMENTS	236
5.1. ORAL REFERENCE DOSE (RfD)	236
5.1.1. Choice of Principal Study and Critical Effect—with Rationale and Justification	236
5.1.2. Derivation Process for Noncancer Reference Values.....	239
5.1.3. Evaluation of Dose Metrics for Use in Noncancer Reference Value Derivations	243
5.1.4. Methods of Analysis—including Models (PBPK, BMD, etc.)	244
5.1.5. RfD Derivation—including Application of Uncertainty Factors (UFs).....	249
5.1.6. Previous RfD Assessment	250
5.1.7. RfD Comparison Information.....	250
5.2. INHALATION REFERENCE CONCENTRATION (RfC).....	253
5.2.1. Choice of Principal Study and Critical Effect—with Rationale and Justification	253
5.2.2. Derivation Process for RfC Values	258
5.2.3. Methods of Analysis—including Models (PBPK, BMD, etc.)	258
5.2.4. RfC Derivation—including Application of Uncertainty Factors (UFs).....	263
5.2.5. Previous RfC Assessment.....	265
5.2.6. RfC Comparison Information.....	265
5.3. UNCERTAINTIES IN THE ORAL REFERENCE DOSE AND INHALATION REFERENCE CONCENTRATION.....	269
5.4. CANCER ASSESSMENT	279
5.4.1. Cancer OSF	279
5.4.1.1. Choice of Study/Data—with Rationale and Justification	279
5.4.1.2. Derivation of OSF.....	280
5.4.1.3. Dose-Response Data	282
5.4.1.4. Dose Conversion and Extrapolation Methods: Cancer OSF	283
5.4.1.5. Oral Cancer Slope Factor.....	289
5.4.1.6. Alternative Derivation Based on Route-to-Route Extrapolation	289
5.4.1.7. Alternative Based On Administered Dose.....	291
5.4.1.8. Previous IRIS Assessment: Cancer OSF	292
5.4.1.9. Comparison of Cancer OSFs Using Different Methodologies	292
5.4.2. Cancer IUR.....	294
5.4.2.1. Choice of Study/Data—with Rationale and Justification	294
5.4.2.2. Derivation of the Cancer IUR.....	295
5.4.2.3. Dose-Response Data	295
5.4.2.4. Dose Conversion and Extrapolation Methods: Cancer IUR	296
5.4.2.5. Cancer IUR	304
5.4.2.6. Comparative Derivation Based on Rat Mammary Tumor Data	307
5.4.2.7. Alternative Based on Administered Concentration	307
5.4.2.8. Previous IRIS Assessment: Cancer IUR.....	309
5.4.2.9. Comparison of Cancer IUR Using Different Methodologies	309

5.4.3. Differences Between Current Assessment and Previous IRIS PBPK-based Assessment.....	311
5.4.4. Application of Age-Dependent Adjustment Factors (ADAFs)	313
5.4.4.1. Application of ADAFs in Oral Exposure Scenarios	313
5.4.4.2. Application of ADAFs in Inhalation Exposure Scenarios	314
5.4.5. Uncertainties in Cancer Risk Values	315
6. MAJOR CONCLUSIONS IN THE CHARACTERIZATION OF HAZARD AND DOSE RESPONSE	331
6.1. HUMAN HAZARD POTENTIAL	331
6.2. DOSE RESPONSE	334
6.2.1. Oral RfD	334
6.2.2. Inhalation RfC	335
6.2.3. Uncertainties in RfD and RfC Values	337
6.2.4. Oral Cancer Slope Factor	339
6.2.5. Cancer IUR.....	343
6.2.6. Uncertainties in Cancer Risk Values	346
7. REFERENCES	349
APPENDIX A: SUMMARY OF EXTERNAL PEER REVIEW AND PUBLIC COMMENTS AND DISPOSITION	A-1
APPENDIX B: HUMAN PBPK DICHLOROMETHANE MODEL	B-1
B.1. HUMAN MODEL DESCRIPTION.....	B-1
B.2. REVISIONS TO PARAMETER DISTRIBUTIONS OF DAVID ET AL. (2006).....	B-3
B.3. CY2E1 AND GST-T1	B-5
B.4. ANALYSIS OF HUMAN PHYSIOLOGICAL DISTRIBUTIONS FOR PBPK MODELING	B-10
B.4.1. Age.....	B-10
B.4.2. Gender.....	B-11
B.4.3. BW	B-12
B.4.4. Alveolar Ventilation	B-14
B.4.5. QCC	B-15
B.4.6. Fat Fraction	B-16
B.4.7. Liver Fraction	B-17
B.4.8. Tissue Volume Normalization.....	B-18
B.5. SUMMARY OF REVISED HUMAN PBPK MODEL	B-18
APPENDIX C. RAT DICHLOROMETHANE PBPK MODELS	C-1
C.1. METHODS OF ANALYSIS	C-1
C.1.1. Selection of Evaluation Data Sets and PBPK Models.....	C-1
C.1.2. Analysis	C-3
C.2. RESULTS.....	C-4
C.2.1. Evaluation of Model Structure for Description of Carboxyhemoglobin Levels.....	C-4
C.2.2. Evaluation of Prediction of Uptake, Blood and Liver Concentrations, and Expiration of Dichloromethane.....	C-7
C.2.3. Evaluation of Relative Flux of CYP and GST Metabolism of Dichloromethane.....	C-13
C.2.4. Evaluation of Model Predictions of Oral Absorption of Dichloromethane.....	C-17

C.3. MODEL OPTION SUMMARY.....	C-19
APPENDIX D. SUMMARY OF BENCHMARK DOSE (BMD) MODELING OF NONCANCER ENDPOINTS	D-1
D.1. ORAL RfD: BMD MODELING OF LIVER LESION INCIDENCE DATA FOR RATS EXPOSED TO DICHLOROMETHANE IN DRINKING WATER FOR 2 YEARS (SEROTA ET AL., 1986A)	D-1
D.2. INHALATION RfC: BMD MODELING OF LIVER LESION INCIDENCE DATA FOR RATS EXPOSED TO DICHLOROMETHANE VIA INHALATION FOR 2 YEARS (NITSCHKE ET AL., 1988A)	D-5
APPENDIX E: SUMMARY OF BENCHMARK DOSE (BMD) MODELING OF CANCER ENDPOINTS	E-1
E.1. ORAL CANCER SLOPE FACTORS: BMD MODELING OF LIVER TUMOR INCIDENCE DATA FOR MICE EXPOSED TO DICHLOROMETHANE IN DRINKING WATER FOR 2 YEARS (SEROTA ET AL., 1986B; HAZLETON LABORATORIES, 1983).....	E-1
E.1.1. Modeling Results for the Internal Liver Metabolism Metric	E-3
E.1.2. Modeling Results for the Whole Body Metabolism Metric.....	E-6
E.2. CANCER IUR: BMD MODELING OF LIVER AND LUNG TUMOR INCIDENCE DATA FOR MALE MICE EXPOSED TO DICHLOROMETHANE VIA INHALATION FOR 2 YEARS (MENNEAR ET AL., 1988; NTP, 1986)	E-9
E.2.1. Modeling Results for the Internal Liver Metabolism Metric, Liver Tumors. Mennear et al. (1988); NTP (1986): Internal Liver Dose-Response for Liver Tumors in Male Mice.....	E-11
E.2.2. Modeling Results for the Internal Lung Metabolism Metric, Lung Tumors. Mennear et al. (1988); NTP (1986): Internal Lung Dose-Response for Lung Tumors in Male Mice.....	E-14
E.2.3. Modeling Results for the Whole Body Metabolism Metric, Liver Tumors. Mennear et al. (1988); NTP (1986): Internal Whole-Body Metabolism Dose-Response for Liver Tumors in Male Mice	E-16
E.2.4. Modeling Results for the Whole Body Metabolism Metric, Lung Tumors. Mennear et al. (1988); NTP (1986): Internal Whole-Body Metabolism Dose-Response for Lung Tumors in Male Mice.....	E-19
APPENDIX F. COMPARATIVE CANCER IUR BASED ON FEMALE MICE DATA	F-1
APPENDIX G. COMPARATIVE CANCER IUR BASED ON BENIGN MAMMARY GLAND TUMORS IN RATS	G-1
APPENDIX H: SOURCE CODE AND COMMAND FILES FOR DICHLOROMETHANE PBPK MODELS	H-1

LIST OF TABLES

Table 2-1. Physical properties and chemical identity of dichloromethane	3
Table 3-1. Distribution of radioactivity in tissues 48 hours after inhalation exposure of mature male Sprague-Dawley rats (n = 3) for 6 hours	7
Table 3-2. Brain and perirenal fat dichloromethane and blood CO concentrations in male Wistar rats exposed by inhalation to dichloromethane at constant exposure concentrations compared with intermittently high exposure concentrations.....	9
Table 3-3. Mean prevalences of the GST-T1 null (-/-) genotype in human ethnic groups.....	16
Table 3-4. GST-T1 enzyme activities toward dichloromethane in human, rat, mouse, and hamster tissues (liver, kidney, and erythrocytes).....	18
Table 3-5. Values for parameter distributions in a B6C3F ₁ mouse probabilistic PBPK model for dichloromethane compared with associated values for point parameters in earlier deterministic B6C3F ₁ mouse PBPK models for dichloromethane	29
Table 3-6. Internal daily doses for B6C3F ₁ mice exposed to dichloromethane for 2 years (6 hours/day, 5 days/week) calculated with different PBPK models	31
Table 3-7. Results of calibrating metabolic parameters in a human probabilistic PBPK model for dichloromethane with individual kinetic data for 42 exposed volunteers and MCMC analysis	33
Table 3-8. Parameter distributions used in human Monte Carlo analysis for dichloromethane by David et al. (2006).....	35
Table 3-9. Parameter distributions for the human PBPK model for dichloromethane used by EPA.....	38
Table 3-10. Parameter values for the rat PBPK model for dichloromethane used by EPA	41
Table 3-11. Parameters in the mouse, rat, and human PBPK model for dichloromethane used by EPA.....	43
Table 4-1. Percentage of male General Electric plastic polymer workers reporting neurologic symptoms or displaying abnormal values in measures of neurological function, hepatic function, and cardiac function.....	60
Table 4-2. Ischemic heart disease mortality risk in four cohorts of dichloromethane-exposed workers	63
Table 4-3. Suicide risk in two cohorts of dichloromethane-exposed workers.....	64

Table 4-4. Mortality risk in Eastman Kodak cellulose triacetate film base production workers, Rochester, New York.....	73
Table 4-5. Mortality risk by cumulative exposure in Eastman Kodak cellulose triacetate film base production workers, Rochester, New York	75
Table 4-6. Mortality risk in Imperial Chemical Industries cellulose triacetate film base production workers, Brantham, United Kingdom: 1,473 men employed 1946–1988, followed through 1994.....	78
Table 4-7. Mortality risk in Hoechst Celanese Corporation cellulose triacetate fiber production workers, Rock Hill, South Carolina: 1,271 men and women employed 1954–1977, followed through 1990.....	80
Table 4-8. Cancer mortality risk in Hoechst Celanese Corporation cellulose triacetate fiber production workers, Cumberland, Maryland: 2,909 men and women employed 1970–1981, followed through 1989.....	82
Table 4-9. Summary of cohort studies of cancer risk and dichloromethane exposure	85
Table 4-10. Summary of case-control studies of cancer risk and dichloromethane exposure.....	95
Table 4-11. Incidences of histopathologic changes in livers of male and female F344 rats exposed to dichloromethane in drinking water for 90 days.....	102
Table 4-12. Incidences of histopathologic changes in livers of male and female B6C3F ₁ mice exposed to dichloromethane in drinking water for 90 days.....	104
Table 4-13. Studies of chronic oral dichloromethane exposures (up to 2 years).....	105
Table 4-14. Incidences of nonneoplastic liver changes and liver tumors in male and female F344 rats exposed to dichloromethane in drinking water for 2 years.....	107
Table 4-15. Incidences for focal hyperplasia and tumors in the liver of male B6C3F ₁ mice exposed to dichloromethane in drinking water for 2 years.....	110
Table 4-16. Studies of chronic inhalation dichloromethane exposures	118
Table 4-17. Incidences of nonneoplastic histologic changes in male and female F344/N rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	120
Table 4-18. Incidences of selected neoplastic lesions in male and female F344/N rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	122
Table 4-19. Incidences of nonneoplastic histologic changes in B6C3F ₁ mice exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	125
Table 4-20. Incidences of neoplastic lesions in male and female B6C3F ₁ mice exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	127

Table 4-21. Incidences of selected nonneoplastic and neoplastic histologic changes in male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	131
Table 4-22. Incidences of selected nonneoplastic histologic changes in male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	134
Table 4-23. Incidences of selected neoplastic histologic changes in male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	135
Table 4-24. Summary of studies of reproductive and developmental effects of dichloromethane exposure in animals	138
Table 4-25. Reproductive outcomes in F344 rats exposed to dichloromethane by inhalation for 14 weeks prior to mating and from GDs 0–21	140
Table 4-26. Studies of neurobehavioral changes from dichloromethane, by route of exposure and type of effect	149
Table 4-27. Studies of neurophysiological changes as measured by evoked potentials resulting from dichloromethane, by route of exposure.....	151
Table 4-28. Studies of neurochemical changes from dichloromethane, by route of exposure..	153
Table 4-29. Results from in vitro genotoxicity assays of dichloromethane with bacteria, yeast, or fungi.....	164
Table 4-30. Results from in vitro genotoxicity assays of dichloromethane with mammalian systems, by type of test.....	168
Table 4-31. Results from in vivo genotoxicity assays of dichloromethane in insects.....	174
Table 4-32. Results from in vivo genotoxicity assays of dichloromethane in mice	175
Table 4-33. Results from in vivo genotoxicity assays of dichloromethane in rats and hamsters	179
Table 4-34. Comparison of in vivo dichloromethane genotoxicity assays targeted to lung or liver cells, by species	181
Table 4-35. NOAELs and LOAELs in selected animal studies involving oral exposure to dichloromethane for short-term, subchronic, or chronic durations	196
Table 4-36. NOAELs and LOAELs in animal studies involving inhalation exposure to dichloromethane for subchronic or chronic durations, hepatic, pulmonary, and neurologic effects.....	201

Table 4-37. NOAELs and LOAELs in selected animal studies involving inhalation exposure to dichloromethane, reproductive and developmental effects	205
Table 4-38. Incidence of liver tumors in male B6C3F ₁ mice exposed to dichloromethane in a 2-year oral exposure (drinking water) study ^a	215
Table 4-39. Incidences of liver tumors in male and female F344 rats exposed to dichloromethane in drinking water for 2 years.....	216
Table 4-40. Incidences of selected neoplastic lesions in B6C3F ₁ mice exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	218
Table 4-41. Incidences of selected neoplastic lesions in F344/N rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	219
Table 4-42. Incidences of mammary gland tumors in two studies of male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years	221
Table 4-43. Comparison of internal dose metrics in inhalation and oral exposure scenarios in male mice and rats	223
Table 5-1. Incidence data for liver lesions and internal liver doses based on various metrics in male and female F344 rats exposed to dichloromethane in drinking water for 2 years	245
Table 5-2. BMD modeling results for incidence of liver lesions in male and female F344 rats exposed to dichloromethane in drinking water for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolism via CYP pathway per liter liver tissue per day)	247
Table 5-3. RfD for dichloromethane based on PBPK model-derived probability distributions of human drinking water exposures extrapolated from liver lesion incidence data for male rats exposed via drinking water for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day)	248
Table 5-4. Potential points of departure with applied UFs and resulting candidate RfDs.....	251
Table 5-5. Incidence data for liver lesions (hepatic vacuolation) and internal liver doses based on various metrics in female Sprague-Dawley rats exposed to dichloromethane via inhalation for 2 years	259
Table 5-6. BMD modeling results for incidence of noncancer liver lesions in female Sprague-Dawley rats exposed to dichloromethane by inhalation for 2 years, based on liver specific CYP metabolism metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day).....	261

Table 5-7. Inhalation RfC for dichloromethane based on PBPK model-derived probability distributions of human inhalation exposure extrapolated from liver lesion data for female rats exposed via inhalation for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day)	262
Table 5-8. Potential points of departure with applied UFs and resulting candidate RfCs.....	267
Table 5-9. Statistical characteristics of human equivalent applied doses in specific populations of the GST-T1 ^{-/-} group.....	276
Table 5-10. Statistical characteristics of HECs in specific populations of the GST-T1 ^{-/-} group.....	278
Table 5-11. Incidence data for liver tumors and internal liver doses, based on GST metabolism dose metrics in male B6C3F ₁ mice exposed to dichloromethane in drinking water for 2 years.....	283
Table 5-12. BMD modeling results and tumor risk factors for internal dose metric associated with 10% extra risk for liver tumors in male B6C3F ₁ mice exposed to dichloromethane in drinking water for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics.....	285
Table 5-13. Cancer OSFs for dichloromethane based on PBPK model-derived internal liver doses in B6C3F ₁ mice exposed via drinking water for 2 years, based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype	288
Table 5-14. Alternative route-to-route cancer OSFs for dichloromethane extrapolated from male B6C3F ₁ mouse inhalation liver tumor incidence data using a tissue-specific GST metabolism dose metric, by population genotype	290
Table 5-15. Cancer OSF based on a human BMDL ₁₀ using administered dose for liver tumors in male B6C3F ₁ mice exposed to dichloromethane in drinking water for 2 years	291
Table 5-16. Comparison of OSFs derived using various assumptions and metrics, based on tumors in male mice.....	293
Table 5-17. Incidence data for liver and lung tumors and internal doses based on GST metabolism dose metrics in male and female B6C3F ₁ mice exposed to dichloromethane via inhalation for 2 years	296
Table 5-18. BMD modeling results and tumor risk factors associated with 10% extra risk for liver and lung tumors in male and female B6C3F ₁ mice exposed by inhalation to dichloromethane for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics	300
Table 5-19. IURs for dichloromethane based on PBPK model-derived internal liver and lung doses in B6C3F ₁ male mice exposed via inhalation for 2 years, based on liver-	

specific GST metabolism and whole body metabolism dose metrics, by population genotype.....	303
Table 5-20. Upper bound estimates of combined human IURs for liver and lung tumors resulting from lifetime exposure to 1 µg/m ³ dichloromethane based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype....	306
Table 5-21. Inhalation units risks based on human BMDL ₁₀ values using administered concentration for liver and lung tumors in B6C3F ₁ mice exposed by inhalation to dichloromethane for 2 years	308
Table 5-22. Comparison of IURs derived by using various assumptions and metrics	310
Table 5-23. Comparison of key B6C3F ₁ mouse parameters differing between prior and current PBPK model application.....	311
Table 5-24. Application of ADAFs to dichloromethane cancer risk following a lifetime (70-year) oral exposure	314
Table 5-25. Application of ADAFs to dichloromethane cancer risk following a lifetime (70-year) inhalation exposure.....	315
Table 5-26. Summary of uncertainty in the derivation of cancer risk values for dichloromethane	316
Table 5-27. Statistical characteristics of human internal doses for 1 mg/kg-day oral exposures in specific populations	328
Table 5-28. Statistical characteristics of human internal doses for 1 mg/m ³ inhalation exposures in specific subpopulations.....	329
Table 6-1. Comparison of OSFs derived by using various assumptions and metrics, based on liver tumors in male mice	342
Table 6-2. Comparison of IURs derived by using various assumptions and metrics	345
Table B-1. Parameter distributions used in human Monte Carlo analysis for dichloromethane by David et al. (2006)	B-2
Table B-2. Parameters for BW distributions as functions of age and gender.....	B-13
Table B-3. Parameter distributions for the human PBPK model for dichloromethane used by EPA.....	B-20
Table C-1. Parameter values used in rat PBPK models.....	C-4
Table C-2. Effect of PBPK model configuration on predicted dichloromethane metabolite production in the liver of (male) rats from inhalation exposures ^a	C-14

Table C-3. Observations and predictions of total expired dichloromethane resulting from gavage doses in rats ^a	C-19
Table D-1. Incidence data for liver lesions and internal liver doses based on various metrics in male and female F344 rats exposed to dichloromethane in drinking water for 2 years (Serota et al., 1986a).....	D-1
Table D-2. BMD modeling results for incidence of liver lesions in male and female F344 rats exposed to dichloromethane in drinking water for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolism via CYP pathway per liter liver tissue per day)	D-2
Table D-3. Incidence data for liver lesions (hepatic vacuolation) and internal liver doses based on various metrics in female Sprague-Dawley rats exposed to dichloromethane via inhalation for 2 years (Nitschke et al., 1988a).....	D-5
Table D-4. BMD modeling results for incidence of liver lesions in female Sprague-Dawley rats exposed to dichloromethane by inhalation for 2 years, based on liver specific CYP metabolism metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day).....	D-6
Table E-1. Incidence data for liver tumors and internal liver doses, based on GST metabolism dose metrics, in male B6C3F ₁ mice exposed to dichloromethane in drinking water for 2 years	E-2
Table E-2. BMD modeling results and tumor risk factors for internal dose metric associated with 10% extra risk for liver tumors in male B6C3F ₁ mice exposed to dichloromethane in drinking water for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics	E-2
Table E-3. Incidence data for liver and lung tumors and internal doses based on GST metabolism dose metrics in male B6C3F ₁ mice exposed to dichloromethane via inhalation for 2 years	E-9
Table E-4. BMD modeling results and tumor risk factors associated with 10% extra risk for liver and lung tumors in male B6C3F ₁ mice exposed by inhalation to dichloromethane for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics	E-10
Table F-1. Incidence data for liver and lung tumors and internal doses based on GST metabolism dose metrics in female B6C3F ₁ mice exposed to dichloromethane via inhalation for 2 years	F-1
Table F-2. BMD modeling results and tumor risk factors associated with 10% extra risk for liver and lung tumors in female B6C3F ₁ mice exposed by inhalation to dichloromethane for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics.....	F-3

Table F-3. IURs for dichloromethane based on PBPK model-derived internal liver and lung doses in B6C3F₁ female mice exposed via inhalation for 2 years, based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype.....F-5

Table F-4. Upper bound estimates of combined human IURs for liver and lung tumors resulting from lifetime exposure to 1 µg/m³ dichloromethane based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype, using female mouse data for derivation of risk factorsF-7

Table G-1. Incidence data for mammary gland tumors and internal doses based on different dose metrics in male and female F344 rats exposed to dichloromethane via inhalation for 2 years G-1

Table G-2. BMD modeling results associated with 10% extra risk for mammary gland tumors in F344 rats exposed by inhalation to dichloromethane for 2 years based on AUC for dichloromethane in slowly perfused tissue..... G-3

Table G-3. IURs for dichloromethane based on benign mammary tumors and PBPK model-derived internal doses in F344N rats exposed via inhalation for 2 years based on AUC for dichloromethane in slowly perfused tissue dose metric G-5

LIST OF FIGURES

Figure 3-1. Proposed pathways for dichloromethane metabolism.....	10
Figure 3-2. Schematics of PBPK models (1986–2006) used in the development of estimates for dichloromethane internal dosimetry.	23
Figure 3-3. Schematic of mouse PBPK model used by Marino et al. (2006).....	28
Figure 3-4. Schematic of human PBPK used by David et al. (2006).	32
Figure 3-5. Schematic of rat PBPK model used in current assessment.	40
Figure 3-6. Comparison of dichloromethane oxidation rate data with alternate kinetic models.	48
Figure 5-1. Exposure response array for oral exposure to dichloromethane.	238
Figure 5-2. Process for deriving noncancer oral RfDs and inhalation RfCs using rodent and human PBPK models.	240
Figure 5-3. PBPK model-derived internal doses (mg dichloromethane metabolized via the CYP pathway per liter liver per day) in rats and humans and their associated external exposures (mg/kg-day), used for the derivation of RfDs.	246
Figure 5-4. Comparison of candidate RfDs derived from selected points of departure for endpoints presented in Table 5-4.	252
Figure 5-5. Exposure response array for chronic (animal) or occupational (human) inhalation exposure to dichloromethane (log Y axis).....	254
Figure 5-6. Exposure response array for subacute to subchronic inhalation exposure to dichloromethane (log Y axis).	256
Figure 5-7. PBPK model-derived internal doses (mg dichloromethane metabolized via the CYP pathway per liter liver per day) in rats and humans versus external exposures (ppm).....	260
Figure 5-8. Comparison of candidate RfCs derived from selected points of departure for endpoints presented in Table 5-8.	268
Figure 5-9. Sensitivity coefficients for long-term mass CYP- and GST-mediated metabolites per liver volume from a daily drinking water concentration of 10 mg/L in rats.	273
Figure 5-10. Sensitivity coefficients for long-term mass CYP- and GST-mediated metabolites per liver volume from a long-term average daily inhalation concentration of 500 ppm in rats.	273

Figure 5-11. Frequency density of human equivalent doses in specific populations in comparison to a general population (0.5- to 80-year-old males and females) estimate for an internal dose of 15.1 mg dichloromethane metabolized by CYP per liter liver per day.....	275
Figure 5-12. Frequency density of HECs in specific populations in comparison to a general population (0.5- to 80-year-old males and females) estimate for an internal dose of 128.1 mg dichloromethane metabolized by CYP per liter liver per day.	277
Figure 5-13. Process for deriving cancer OSFs and IURs by using rodent and human PBPK models.....	281
Figure 5-14. PBPK model-derived internal doses (mg dichloromethane metabolized via the GST pathway per liter liver per day) in mice and humans and their associated external exposures (mg/kg-day) used for the derivation of cancer OSFs based on liver tumors in mice.	284
Figure 5-15. PBPK model-derived internal doses (mg dichloromethane metabolized via the GST pathways per liter tissue per day) for liver (A) and lung (B) in mice and humans and their associated external exposures (ppm) used for the derivation of cancer IURs.	298
Figure 5-16. PBPK-model-predicted exposure-response relationships for hepatic CYP and GST metabolism for continuous inhalation exposure to dichloromethane in 30-year-old GST +/+ women.	321
Figure 5-17. Sensitivity coefficients for long-term mass GST-mediated metabolites per liver volume from a long-term average daily inhalation concentration of 2,000 ppm in mice.....	324
Figure 5-18. Sensitivity coefficients for long-term mass GST-mediated metabolites per liver volume from a long-term average daily drinking water concentration of 500 mg/L in mice.....	325
Figure 5-19. Sensitivity coefficients for long-term mass GST-mediated metabolites per lung volume from a long-term average daily inhalation concentration of 500 ppm in mice.	326
Figure 5-20. Histograms for a liver-specific dose of GST metabolism (mg GST metabolites per liter liver per day) for the general population (0.5- to 80-year-old males and females), and specific age/gender groups within the population of GST-T1 ^{+/+} genotypes, given a daily oral dose-rate of 1 mg/kg-day dichloromethane.	328
Figure 5-21. Histograms for liver-specific dose of GST metabolism (mg GST metabolites per liter liver per day) for the general population (0.5- to 80-year-old males and females), and specific age/gender groups within the population of GST-T1 ^{+/+} genotypes, given a continuous inhalation exposure to 1 mg/m ³ dichloromethane.....	329
Figure B-1. Schematic of the David et al. (2006) PBPK model for dichloromethane in the human.	B-1

Figure B-2. Total CYP2E1 activity (V_{\max}) normalized to the average total activity in 14–18 year-old individuals ($V_{\max}[14–18]$) plotted against normalized BW for individuals ranging from 6 months to 18 years of age.	B-7
Figure B-3. Body-weight scaled CYP2E1 activity ($V_{\max C}$) normalized to the average scaled activity in 14–18 year-old individuals ($V_{\max C}[14–18]$) plotted against age individuals ranging from 6 months to 18 years of age.	B-9
Figure B-4. U.S. age distribution, 6 months to 80 years (values from U.S. Census Bureau)..	B-11
Figure B-5. U.S. age-specific gender distribution (values from U.S. Census Bureau).	B-11
Figure B-6. Function fits to age-dependent data for BW mean and SDs for males and females in the United States (values from Portier et al. [2007]).	B-13
Figure B-7. Example BW histogram from Monte Carlo simulation for 0.5- to 80-year-old males and females in the United States (simulated n = 10,000).	B-14
Figure B-8. Mean value respiration rates for males and females as a function of age (values from Clewell et al. [2004]).	B-15
Figure B-9. GSDs for respiration rates for males and females as a function of age (values from Arcus-Arth and Blaisdell [2007]).	B-15
Figure B-10. Fraction body fat (VFC) over various age ranges in males and females (data from Clewell et al. [2004]).	B-17
Figure C-1. Schematic of the Andersen et al. (1991) PBPK model (model B) for dichloromethane in the rat.	C-2
Figure C-2. A: Observations of exhaled [^{14}C]-labelled dichloromethane (left y-axis) and CO (right y-axis) after a bolus oral dose of 200 mg/kg [^{14}C]-dichloromethane in rats (data of Angelo et al., 1986b). B: Blood COHb (percent of total hemoglobin) from a single gavage dose of 526 mg/kg dichloromethane in rats.	C-6
Figure C-3. Observations of Gargas et al. (1986; data) and predictions (models A–D) for respiratory uptake by three rats of 100–3,000 ppm dichloromethane in a 9-L closed chamber.	C-8
Figure C-4. Observations (points) of Angelo et al. (1986b) and predictions (curves, denoted “sim” in legend) for models A–D.	C-9
Figure C-5. Observations of Angelo et al. (1986b) and predictions (curves, denoted “sim” in legend) for models A–D following 10 and 50 mg/kg intravenous injection in rats.	C-11

Figure C-6. Observations of Andersen et al. (1987; data points) and simulations for models A–D (curves, denoted “sim” in legend) for dichloromethane in rat blood from inhalation of 200 and 1,000 ppm dichloromethane for 4 hours..... C-12

Figure C-7. Simulation results using models B and D for weekly average metabolic rates by the GST and CYP pathways for 6 hours/day, 5 days/week inhalation exposures..... C-15

Figure C-8. Observations of Angelo et al. (1986b; data) and model D predictions for: (A) percent dose expired as dichloromethane; (B) blood dichloromethane; (C) percent expired as CO; and (D) liver DCM in rats given a single dichloromethane gavage dose of 50 and 200 mg/kg, using a numerically fitted k_a of 1.8 hours⁻¹ (see Section C.2.4). C-16

Figure C-9. Model predictions of blood COHb (percent of total hemoglobin) from a single gavage dose of 526 mg/kg dichloromethane in rats, compared to the data of Pankow et al. (1991a). C-18

Figure D-1. Predicted (logistic model) and observed incidence of noncancer liver lesions in male F344 rats exposed to dichloromethane in drinking water for 2 years (Serota et al., 1986a). D-3

Figure D-2. Predicted (log-probit model) and observed incidence of noncancer liver lesions in female Sprague-Dawley rats inhaling dichloromethane for 2 years (Nitschke 1988a). D-7

Figure E-1. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years, using liver-specific metabolism dose metric (Serota et al., 1986b; Hazleton Laboratories, 1983). E-3

Figure E-2. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years, using whole-body metabolism dose metric (Serota et al., 1986b; Hazleton Laboratories, 1983). E-6

Figure E-3. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F₁ mice exposed by inhalation to dichloromethane for 2 years, using liver-specific metabolism dose metric (Mennear et al., 1988; NTP, 1986). E-11

Figure E-4. Predicted and observed incidence of animals with carcinoma or adenoma in the lung of male B6C3F₁ mice exposed by inhalation to dichloromethane for 2 years, using liver-specific metabolism dose metric (Mennear et al., 1988; NTP, 1986). E-14

Figure E-5. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F₁ mice exposed by inhalation to dichloromethane for 2 years, using whole-body metabolism dose metric (Mennear et al., 1988; NTP, 1986). E-16

Figure E-6. Predicted and observed incidence of animals with carcinoma or adenoma in the lung of male B6C3F₁ mice exposed by inhalation to dichloromethane for 2 years, using whole-body metabolism dose metric (Mennear et al., 1988; NTP, 1986).....E-19

Figure G-1. PBPK model-derived internal doses (daily average AUC for dichloromethane in blood) in rats and humans and their associated external exposures (ppm) used for the derivation of cancer IURs based on mammary tumors in rats..... G-2

LIST OF ABBREVIATIONS AND ACRONYMS

A1	ratio of lung $V_{\max C}$ to liver $V_{\max C}$
A2	ratio of lung k_{fC} to liver k_{fC}
ABCOC	background amount of CO
ADAF	age-dependent adjustment factor
AIC	Akaike's Information Criterion
ALT	alanine aminotransferase
AP	alkaline phosphatase
AST	aspartate aminotransferase
AUC	area under the curve
BAER	brainstem-auditory evoked response
BMD	benchmark dose
BMDL₁₀	95% lower bound on the BMD
BMDS	benchmark dose software
BMR	benchmark response
BW	body weight
CAEP	cortical-auditory-evoked potential
CASRN	Chemical Abstracts Service Registry Number
CHO	Chinese hamster ovary
CI	confidence interval
CNS	central nervous system
CO	carbon monoxide
COHb	carboxyhemoglobin
CV	coefficient of variation
CYP	cytochrome P450
DNA	deoxyribonucleic acid
FEP	flash-evoked potential
FOB	functional observational battery
FracR	fraction of $V_{\max C}$ in rapidly perfused tissues
GD	gestational day
GM	geometric mean
GSD	geometric standard deviation
GSH	reduced glutathione
GST	glutathione S-transferase
GST-T1	GST-theta1-1
HEC	human equivalent concentration
HPRT	hypoxanthine-guanine phosphoribosyl transferase
ICD-9	International Classification of Diseases 9 th ed.
IgM	immunoglobulin M
IRIS	Integrated Risk Information System
IUR	inhalation unit risk
k_a	first-order oral absorption rate constant
K_m	Michaelis-Menten kinetic constant
k_{fC}	first-order GST metabolic rate constant
LOAEL	lowest-observed-adverse-effect level
LOH	loss of heterozygosity
MCHC	mean corpuscular hemoglobin concentration

MCMC	Markov Chain Monte Carlo
mRNA	messenger ribonucleic acid
NADPH	nicotinamide adenine dinucleotide phosphate
NIOSH	National Institute of Occupational Safety and Health
NOAEL	no-observed-adverse-effect level
NRC	National Research Council
NTP	National Toxicology Program
OR	odds ratio
OSF	oral slope factor
OSHA	Occupational Safety and Health Administration
P₅₀	partial oxygen pressure
PB	blood:air partition coefficient
PBPK	physiologically based pharmacokinetic
POD	point of departure
PND	postnatal day
QA_{lvC}	allometric alveolar ventilation constant
QCC	cardiac output
QSC	fractional flow rate of slowly perfused tissues (fraction of QCC)
REnCO_C	endogenous rate of CO production
RfC	reference concentration
RfD	reference dose
SD	standard deviation
SEM	standard error of the mean
SEP	somatosensory-evoked potential
SMR	standardized mortality ratio
SSB	single strand break
TWA	time-weighted average
UF	uncertainty factor
U.S. EPA	U.S. Environmental Protection Agency
VFC	fractional tissue volume of fat (fraction of BW)
VLC	fractional tissue volume of liver (fraction of BW)
V_{maxC}	CYP maximum velocity
VPR	ventilation:perfusion ratio
VSC	fractional tissue volume of slowly perfused tissues (fraction of BW)

FOREWORD

The purpose of this Toxicological Review is to provide scientific support and rationale for the hazard and dose-response assessment in IRIS pertaining to exposure to dichloromethane. It is not intended to be a comprehensive treatise on the chemical or toxicological nature of dichloromethane.

The intent of Section 6, *Major Conclusions in the Characterization of Hazard and Dose Response*, is to present the major conclusions reached in the derivation of the reference dose, reference concentration and cancer assessment, where applicable, and to characterize the overall confidence in the quantitative and qualitative aspects of hazard and dose response by addressing the quality of data and related uncertainties. The discussion is intended to convey the limitations of the assessment and to aid and guide the risk assessor in the ensuing steps of the risk assessment process.

For other general information about this assessment or other questions relating to IRIS, the reader is referred to EPA's IRIS Hotline at (202) 566-1676 (phone), (202) 566-1749 (fax), or hotline.iris@epa.gov (email address).

AUTHORS, CONTRIBUTORS, AND REVIEWERS

CHEMICAL MANAGERS

Glinda S. Cooper, Ph.D.
Ambuja S. Bale, Ph.D., DABT
Office of Research and Development, IRIS Program
U.S. Environmental Protection Agency
Washington, DC

AUTHORS

Glinda S. Cooper, Ph.D.
Ambuja S. Bale, Ph.D., DABT
Andrew Rooney, Ph.D.
Paul Schlosser, Ph.D.
Allan Marcus, Ph.D.
Gene (Ching-Hung) Hsu, Ph.D., DABT
National Center for Environmental Assessment
Office of Research and Development
U.S. Environmental Protection Agency
Washington, DC

John C. Lipscomb, Ph.D., DABT
National Center for Environmental Assessment
U.S. Environmental Protection Agency
Cincinnati, OH

Peter McClure, Ph.D., DABT
Michael Lumpkin, Ph.D.
Fernando Lladós, Ph.D.
Mark Osier, Ph.D., DABT
Daniel Plewak, B.S.
Syracuse Research Corporation
Syracuse, NY

Elizabeth Dupree Ellis, Ph.D.
Oak Ridge Institute for Science and Education
Center for Epidemiologic Research
Oak Ridge, TN

REVIEWERS

This document has been provided for review to EPA scientists, interagency reviewers from other federal agencies and White House offices.

INTERNAL EPA REVIEWERS

Ghazi Dannan, Ph.D.

Karen Hogan, M.S.

Jennifer Jinot, Ph.D.

Paul White, Ph.D.

Samantha Jones, Ph.D.

Jamie Strong, Ph.D.

National Center for Environmental Assessment

Office of Research and Development

U.S. Environmental Protection Agency

David Herr, Ph.D.

National Health and Environmental Effect Research Laboratory

Office of Research and Development

U.S. Environmental Protection Agency

1. INTRODUCTION

This document presents background information and justification for the Integrated Risk Information System (IRIS) Summary of the hazard and dose-response assessment of dichloromethane. IRIS Summaries may include oral reference dose (RfD) and inhalation reference concentration (RfC) values for chronic and other exposure durations, and a carcinogenicity assessment.

The RfD and RfC, if derived, provide quantitative information for use in risk assessments for health effects known or assumed to be produced through a nonlinear (presumed threshold) mode of action. The RfD (expressed in units of mg/kg-day) is defined as an estimate (with uncertainty spanning perhaps an order of magnitude) of a daily exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious effects during a lifetime. The inhalation RfC (expressed in units of mg/m³) is analogous to the oral RfD, but provides a continuous inhalation exposure estimate. The inhalation RfC considers toxic effects for both the respiratory system (portal-of-entry) and for effects peripheral to the respiratory system (extrapulmonary or systemic effects). Reference values are generally derived for chronic exposures (up to a lifetime), but may also be derived for acute (≤ 24 hours), short-term (>24 hours up to 30 days), and subchronic (>30 days up to 10% of lifetime) exposure durations, all of which are derived based on an assumption of continuous exposure throughout the duration specified. Unless specified otherwise, the RfD and RfC are derived for chronic exposure duration.

The carcinogenicity assessment provides information on the carcinogenic hazard potential of the substance in question and quantitative estimates of risk from oral and inhalation exposure may be derived. The information includes a weight-of-evidence judgment of the likelihood that the agent is a human carcinogen and the conditions under which the carcinogenic effects may be expressed. Quantitative risk estimates may be derived from the application of a low-dose extrapolation procedure. If derived, the oral slope factor is a plausible upper bound on the estimate of risk per mg/kg-day of oral exposure. Similarly, an inhalation unit risk is a plausible upper bound on the estimate of risk per $\mu\text{g}/\text{m}^3$ air breathed.

Development of these hazard identification and dose-response assessments for dichloromethane has followed the general guidelines for risk assessment as set forth by the National Research Council (1983). EPA Guidelines and Risk Assessment Forum Technical Panel Reports that may have been used in the development of this assessment include the following: *Guidelines for the Health Risk Assessment of Chemical Mixtures* (U.S. EPA, 1986a), *Guidelines for Mutagenicity Risk Assessment* (U.S. EPA, 1986b), *Recommendations for and Documentation of Biological Values for Use in Risk Assessment* (U.S. EPA, 1988a), *Guidelines for Developmental Toxicity Risk Assessment* (U.S. EPA, 1991), *Interim Policy for Particle Size*

and Limit Concentration Issues in Inhalation Toxicity Studies (U.S. EPA, 1994a), *Methods for Derivation of Inhalation Reference Concentrations and Application of Inhalation Dosimetry* (U.S. EPA, 1994b), *Use of the Benchmark Dose Approach in Health Risk Assessment* (U.S. EPA, 1995), *Guidelines for Reproductive Toxicity Risk Assessment* (U.S. EPA, 1996), *Guidelines for Neurotoxicity Risk Assessment* (U.S. EPA, 1998), *Science Policy Council Handbook: Risk Characterization* (U.S. EPA, 2000a), *Benchmark Dose Technical Guidance Document* (U.S. EPA, 2000b), *Supplementary Guidance for Conducting Health Risk Assessment of Chemical Mixtures* (U.S. EPA, 2000c), *A Review of the Reference Dose and Reference Concentration Processes* (U.S. EPA, 2002), *Guidelines for Carcinogen Risk Assessment* (U.S. EPA, 2005a), *Supplemental Guidance for Assessing Susceptibility from Early-Life Exposure to Carcinogens* (U.S. EPA, 2005b), *Science Policy Council Handbook: Peer Review* (U.S. EPA, 2006a), and *A Framework for Assessing Health Risk of Environmental Exposures to Children* (U.S. EPA, 2006b).

The literature search strategy employed for this compound was based on the Chemical Abstracts Service Registry Number (CASRN) and at least one common name. Any pertinent scientific information submitted by the public to the IRIS Submission Desk was also considered in the development of this document. The relevant literature was reviewed through April 2009.

2. CHEMICAL AND PHYSICAL INFORMATION

Dichloromethane is a colorless liquid with a penetrating, ether-like odor (Lewis, 1997).¹ Selected chemical and physical properties of dichloromethane are listed in Table 2-1.

Table 2-1. Physical properties and chemical identity of dichloromethane

	Physical property/chemical identity	Reference
CAS number	75-09-2	Lide (2000)
Synonyms	Methylene chloride, methylene dichloride, methyl bichloride	O'Neil et al. (2001)
Molecular weight	84.93	O'Neil et al. (2001)
Chemical formula	CH ₂ Cl ₂	O'Neil et al. (2001)
Boiling point	40°C	Lide (2000)
Melting point	-95.1°C	Lide (2000)
Vapor pressure	1.15 × 10 ² mm Hg at 25°C	Boublik et al. (1984)
Density	1.3266 g/mL at 20°C	Lide (2000)
Vapor density	2.93 (air = 1.02)	Holbrook (2003)
Water solubility	1.30 × 10 ⁴ mg/L at 25°C	Horvath (1982)
Other solubility	Miscible in ethanol, ether, and dimethylformamide; soluble in carbon tetrachloride	IARC (1999)
Partition coefficient	log K _{ow} = 1.25	Hansch et al. (1995)
Flash point	Not flammable	U.S. Coast Guard (1999)
Auto ignition temperature	640°C	Holbrook (2003)
Latent heat of vaporization	3.30 × 10 ⁵ J/kg	U.S. Coast Guard (1999)
Heat of fusion	16.89 cal/g	U.S. Coast Guard (1999)
Critical temperature	245.0°C	Holbrook (2003)
Critical pressure	6.171 × 10 ⁶ Pa	Holbrook (2003)
Viscosity	0.430 cP at 20°C	Lewis (1997)
Henry's law constant	3.25 × 10 ⁻³ atm m ³ /mol at 25°C	Leighton and Calo (1981)
OH reaction rate constant	1.42 × 10 ⁻¹³ cm ³ /molecule sec at 25°C	Atkinson (1989)
Chemical structure	$\begin{array}{c} \text{H} \\ \\ \text{Cl}-\text{C}-\text{Cl} \\ \\ \text{H} \end{array}$	

Dichloromethane is produced by two methods of manufacturing (IARC, 1999). The older method involves the direct reaction of methane with chlorine either at high temperatures or

¹To avoid confusion, "dichloromethane" is used throughout this summary even if a specific paper used the term "methylene chloride."

at lower temperatures under catalytic or photolytic conditions (Holbrook, 2003). The more common method used today involves an initial reaction of hydrochloric acid with methanol to yield methyl chloride. Excess methyl chloride is then reacted in the gas phase thermally with chlorine to produce dichloromethane (Holbrook, 2003). This process can also be carried out catalytically or photolytically.

Dichloromethane became an important industrial chemical in the United States during World War II (Hardie, 1964). Dichloromethane has been used in paint strippers and removers, as a propellant in aerosols, in the manufacture of drugs, pharmaceuticals, film coatings, electronics, and polyurethane foam, and as a metal-cleaning solvent. Dichloromethane can also be used in the decaffeination process of coffee and tea (ATSDR, 2000). The U.S. production was 3.8 million pounds in 1941 and 8.3 million pounds in 1944 (Searles and McPhail, 1949). Dichloromethane production rose sharply in the decades following the war due to the increased demand for this substance for use mainly in paint strippers (Hardie, 1964; Searles and McPhail, 1949). U.S. production in 1947, 1955, 1960, and 1962 was approximately 19, 74, 113, and 144 million pounds, respectively (Hardie, 1964; Searles and McPhail, 1949). As other solvent uses and its use in aerosol propellants became important, demand for this substance increased further (Anthony, 1979). Dichloromethane production continued to rise dramatically through the 1970s; production capacities were 520 million pounds in 1973 and 830 million pounds in 1979 (CMR, 1979, 1973).

After 1980, production of dichloromethane began to decline. Production capacities fell from 722 million pounds in 1982 to 465 million pounds in 1997 (CMR, 1997, 1982). The total U.S. production capacity for dichloromethane in 2000 was 535 million pounds (CMR, 2000). The demand for dichloromethane decreased from 600 million pounds in 1979 to 200 million pounds in 1999 (CMR, 2000, 1979). The decline in production of and demand for dichloromethane over the past 2 decades has been attributed to increased regulation, the use of alternative chemicals in aerosol spray cans, and concern over dichloromethane carcinogenicity (Holbrook, 2003; ATSDR, 2000).

Dichloromethane in the environment will partition mainly to air (NLM, 2003). In air, dichloromethane exists as a vapor. Some of the dichloromethane released to soil or water is expected to volatilize to air. In soil, dichloromethane is expected to be highly mobile and may migrate to groundwater. The potential for dichloromethane to bioconcentrate in aquatic or marine organisms is low. Dichloromethane may biodegrade in soil or water under both aerobic and anaerobic conditions.

3. TOXICOKINETICS

3.1. ABSORPTION

3.1.1. Oral—Gastrointestinal Tract Absorption

There are currently no data available on absorption of dichloromethane following oral intake in humans. However, after oral administration in animals, dichloromethane is rapidly and nearly completely absorbed in the gastrointestinal tract (Angelo et al., 1986a, b; McKenna and Zempel, 1981). Angelo et al. (1986b) reported that, following administration of single radiolabeled oral doses (10, 50, or 200 mg/kg) to mature male F344 rats, 97% of the label was detected in the exhaled air within 24 hours, indicating nearly complete absorption. At several time points within 40 minutes of dose administration, <2% of the dose was found in the lower part of the gastrointestinal tract, indicating that the majority of dichloromethane absorption occurs in the upper gastrointestinal tract (Angelo et al., 1986b). Similar results were reported in mature male B6C3F₁ mice exposed to up to 50 mg/kg (Angelo et al., 1986a). In mature male Sprague-Dawley rats administered a single dose (1 or 50 mg/kg) of radiolabeled dichloromethane, <1% of the label was found in feces collected for 48 hours after dose administration (McKenna and Zempel, 1981). Absorption of dichloromethane generally follows first-order kinetics (Angelo et al., 1986a), and no evidence for a dichloromethane-specific carrier has been presented. The vehicle appears to affect the rate but not the extent of gastrointestinal absorption, with an aqueous vehicle resulting in a more rapid absorption of dichloromethane than an oil-based vehicle (Angelo et al., 1986a).

3.1.2. Inhalation—Respiratory Tract Absorption

Several studies in humans have demonstrated the absorption of dichloromethane following inhalation exposure. In a study by Astrand et al. (1975), 14 male volunteers (ages 19–29) were exposed to about 870 mg/m³ (250 ppm) or 1,740 mg/m³ (500 ppm) for 30 minutes while resting or exercising on a bicycle ergometer. There was a pause of about 20 minutes without exposure between rest and exercise periods. Uptake of dichloromethane was estimated at about 55% while resting and about 40, 30, and 35% at respective workloads of 50, 100, and 150 watts. Blood levels of dichloromethane correlated directly with exposure concentrations, and did not appear to increase when a workload was applied (Astrand et al., 1975). Similar reports of rapid uptake and a direct correlation between dichloromethane exposure level and blood levels in humans have been presented by other groups (DiVincenzo and Kaplan, 1981; DiVincenzo et al., 1971).

With extended (≥1–2 hours) exposure, uptake tends to reach a steady-state level, at which point blood dichloromethane levels remain more or less constant (DiVincenzo and Kaplan, 1981; DiVincenzo et al., 1972; Riley et al., 1966). DiVincenzo et al. (1972) reported that in humans

exposed to 100 or 200 ppm of dichloromethane for 2 hours (without physical exercise), dichloromethane was rapidly absorbed, reaching an approximate steady state, as assessed by levels of unchanged dichloromethane in the expired air, within the first 15–30 minutes of exposure. A later study by the same group (DiVincenzo and Kaplan, 1981) similarly reported a rapid absorption of dichloromethane in volunteers exposed to 50–200 ppm for 7.5 hours on each of 5 consecutive days. A steady-state level, as assessed by levels of unchanged dichloromethane in the expired air, was reached quickly (1–2 hours), with exhaled dichloromethane levels increasing with increasing exposure level. A similar pattern was seen with blood dichloromethane levels. Estimated pulmonary uptake was 69–75% and did not vary appreciably with exposure concentration. In another experiment in which one of the investigators was seated during exposure to 100 ppm dichloromethane for 2 hours, concentrations of dichloromethane in expired air reached an apparent plateau of about 70 ppm within the first hour of exposure (Riley et al., 1966).

Body fat may influence absorption of dichloromethane, as evidenced by data from an experiment involving 12 men ages 21–35, divided into two groups (n = 6 per group) based on percent body fat (Engström and Bjurström, 1977). The mean percent body fat in the leaner group was 7.8% (standard error of the mean [SEM] 1.9), range 2.3–13.6%, compared with 25.1% (SEM 2.8), range 18.3–36.2%, in the more overweight group. Total uptake of dichloromethane during a light exercise period (50 watts²) for 1 hour with an exposure level of 750 ppm was positively correlated with percent body fat ($r = 0.81$), and the estimated amount of dichloromethane in fat storage was also correlated with percent body fat ($r = 0.84$).

A pattern of absorption similar to that seen in humans has been seen in animals. Initially, dichloromethane is readily absorbed following inhalation exposure, as evidenced by rapid appearance of dichloromethane in blood, tissues, and expired air (Withey and Karpinski, 1985; Stott and McKenna, 1984; Anders and Sunram, 1982; Carlsson and Hultengren, 1975; Roth et al., 1975). For example, absorption of inhaled 500 ppm dichloromethane in anesthetized, mature male F344 rats reached an apparent plateau within 10–20 minutes and was relatively constant for up to 2 hours (Stott and McKenna, 1984). In these experiments, absorption was calculated from measurements of exposure (nose only) and effluent concentrations and ventilation flow rate in intact animals; double tracheostomized rats were used to measure absorption in the isolated upper respiratory tract and the lower respiratory tract. At a ventilation rate of 53 mL/minute, absorption expressed as mean percentage of dichloromethane available for absorption was 44% (standard deviation [SD] 10) in intact rats, 13.2% (SD 3.6) in the upper respiratory tract, and 37% (SD 4.1) in the lower respiratory tract.

²A watt is the International System Unit of power and is equal to 1 joule of energy per second. It is a measure of the rate of energy use or production (i.e., the exercise effort that was exerted by the individuals in the study).

3.2. DISTRIBUTION

Results from studies of animals show that, following absorption, dichloromethane is rapidly distributed throughout the body and has been detected in all tissues that have been evaluated. Twenty minutes after a single intravenous dose of 10 mg [¹⁴C]-dichloromethane/kg to mature male B6C3F₁ mice (Angelo et al., 1986a), total label was greatest in the liver (6.72 µg-equivalents/g tissue), with lower levels reported in the lung (1.82 µg-equivalents/g tissue), kidney (1.84 µg-equivalents/g tissue), and the remainder of the carcass (1.90 µg-equivalents/g tissue). By 4 hours post administration, levels in the liver had fallen to 3.08 µg-equivalents/g tissue, lung levels were 0.64 µg-equivalents/g tissue, and carcass levels were 0.23 µg-equivalents/g tissue. The levels in the kidney rose sharply in the first hour postexposure but then fell and remained steady at ~1.60 µg-equivalents/g tissue for the remaining 3 hours of the study (Angelo et al., 1986a). McKenna et al. (1982) exposed groups of mature male Sprague-Dawley rats to 50, 500, or 1,500 ppm [¹⁴C]-labeled dichloromethane for 6 hours and examined tissues at 48 hours for presence of radiolabel; results are shown in Table 3-1. The greatest concentration of label was found in the liver, followed by the kidney and lung.

Table 3-1. Distribution of radioactivity in tissues 48 hours after inhalation exposure of mature male Sprague-Dawley rats (n = 3) for 6 hours

Tissue	Mean ± SD, µg-equivalent dichloromethane/g tissue, by exposure level		
	50 ppm	500 ppm	1,500 ppm
Liver	8.4 ± 1.5	35.6 ± 7.5	44.2 ± 3.5
Kidney	3.3 ± 0.1	16.2 ± 2.4	30.5 ± 0.2
Lung	1.9 ± 0.2	11.0 ± 1.3	16.5 ± 1.6
Brain	0.8 ± 0.3	4.2 ± 1.3	6.7 ± 0.2
Epididymal fat	0.5 ± 0.2	6.5 ± 0.5	4.1 ± 0.9
Skeletal muscle	1.1 ± 0.1	4.4 ± 1.9	7.7 ± 0.7
Testes	1.1 ± 0.2	5.5 ± 1.3	8.1 ± 0.5
Whole blood	1.1 ± 0.2	8.1 ± 1.9	8.9 ± 1.7
Remaining carcass	1.3 ± 0.2	5.9 ± 0.9	8.6 ± 1.4

Source: McKenna et al. (1982).

As noted in the preceding section, body fat may affect the uptake of dichloromethane, and there is also evidence of a relationship between adiposity and dichloromethane storage. In the study by Engström and Bjurström (1977) involving 12 men ages 21–35 exposed to 750 ppm dichloromethane during a 1-hour light exercise (50 watts) period, dichloromethane was measured in body fat biopsy specimens at 1, 2, 3, and 4 hours postexposure. All specimens were taken from the buttocks. The concentration of dichloromethane (per gram tissue) was negatively

correlated with percent body fat, but the total estimated amount of dichloromethane in fat tissue 4 hours postexposure was higher in subjects with a higher amount of fat ($r = 0.84$).

Carlsson and Hultengren (1975) exposed groups of 10 mature male Sprague-Dawley rats to [^{14}C]-dichloromethane for 1 hour at a mean concentration of $1,935 \text{ mg/m}^3$ (557 ppm) and SD of 90 mg/m^3 (26 ppm). The initial levels were highest in the white adipose tissue (approximately $80 \text{ }\mu\text{g}$ dichloromethane per gram tissue) compared with approximately 35, 20, and $5 \text{ }\mu\text{g}$ -equivalent dichloromethane/g tissue in the liver, kidney and adrenal glands, and brain, respectively. These initial levels in the adipose quickly fell to $<10 \text{ }\mu\text{g}$ -equivalent dichloromethane/g tissue; more moderate declines were seen in the other tissues.

With acute 6-hour exposure scenarios, peak exposure concentrations may have a greater influence on dichloromethane levels in the brain and perirenal fat than time-weighted average (TWA) concentrations during the exposure period (Savolainen et al., 1981). In rats exposed over a 6-hour period for 5 days/week to a TWA of 1,000 ppm dichloromethane consisting of two 1-hour peak concentrations (2,800 ppm) interspersed with exposure to 100 ppm, levels of dichloromethane in the brain was 3-fold higher ($p < 0.001$) than corresponding levels in rats exposed to constant levels of 1,000 ppm; a two-fold increased risk was seen in the dichloromethane levels in perirenal fat after one week of exposure ($p < 0.001$), but this difference was much smaller after two weeks of exposure. This difference was not seen with blood carbon monoxide (CO) levels (Table 3-2). With constant exposure concentrations of 500 or 1,000 ppm, perirenal fat levels of dichloromethane approximately doubled following 2 weeks of exposure compared with 1 week of exposure, indicating that some storage of dichloromethane in fat tissue can occur with repeated exposure scenarios (Table 3-2). In contrast, brain levels of dichloromethane in rats exposed for 1 week were higher than brain levels in rats exposed for 2 weeks. One possible explanation of these observations is that there is an induction of enzymes involved in dichloromethane metabolism in liver and other tissues with repeated exposure, and dichloromethane in fat is poorly metabolized.

Table 3-2. Brain and perirenal fat dichloromethane and blood CO concentrations in male Wistar rats exposed by inhalation to dichloromethane at constant exposure concentrations compared with intermittently high exposure concentrations

Exposure level ^a (TWA, ppm)	Exposure wks					
	1	2	1	2	1	2
	Brain (nmol/g)		Perirenal fat (nmol/g)		Blood CO (nmol/g)	
Control	0	0	0	0	40 ± 15	30 ± 10
500, constant	30 ± 7	9 ± 3	436 ± 47	918 ± 215	675 ± 195	781 ± 62
1,000, constant	33 ± 2	14 ± 3	1,316 ± 209	2,171 ± 219	876 ± 80	825 ± 56
1,000, with two 1-hr peaks of 2,800 ppm	111 ± 18 ^b	50 ± 15 ^b	2,295 ± 147 ^b	2,431 ± 146	728 ± 84	873 ± 90

^aGroups of five rats were exposed to 0, 50, or 1,000 ppm 6 hrs/d or 100 ppm interspersed with two 1-hr peaks of 2,800 ppm for 5 d/wk for 1 or 2 wks. Tissue concentration values are mean ± SD.

^bDifference between 1,000 ppm TWA constant exposure, $p < 0.001$; t-test calculated by EPA using sample size, mean and standard deviation as provided by Savolainen et al. (1981).

Source: Savolainen et al. (1981).


Placental transfer. Dichloromethane is capable of crossing the placental barrier and entering the fetal circulation. Anders and Sunram (1982) reported that when pregnant Sprague-Dawley rats ($n = 3$) were exposed to 500 ppm dichloromethane for 1 hour on gestational day (GD) 21, mean maternal blood levels were 176 nmol/mL (SEM 50), while fetal levels were 115 nmol/mL (SEM 40); interestingly, the levels of CO, a metabolite of dichloromethane, were similar in both the maternal blood (167 nmol/mL, SEM 12) and fetal blood (160 nmol/mL, SEM 31). Withey and Karpinski (1985) also reported higher maternal compared with fetal dichloromethane levels based on a study of five pregnant Sprague-Dawley rats exposed to 107–2,961 ppm of dichloromethane. Maternal blood levels of dichloromethane were 2–2.5-fold higher than those found in the fetal circulation.

Blood-brain barrier transfer. Dichloromethane is thought to readily transfer across the blood-brain barrier, as evidenced by the detection of radioactivity in brain tissue 48 hours after exposures of rats to radiolabeled dichloromethane at concentrations of 50, 500, or 1,500 ppm for 6 hours (McKenna et al., 1982) (see Table 3-1), and the historical demonstrations that dichloromethane has transient sedative and anesthetic properties in humans (for review of these reports, see Mattsson et al. [1990] and Winneke [1974]). Dichloromethane is no longer used as an anesthetic gas because the margin between anesthetic and lethal doses is narrow (Winneke, 1974).

3.3. METABOLISM

Metabolism of dichloromethane involves two primary pathways, outlined in Figure 3-1 (ATSDR, 2000; Guengerich, 1997; Hashmi et al., 1994; Gargas et al., 1986). Dichloromethane

is metabolized to CO in a cytochrome P450 (CYP)-dependent oxidative pathway that is predominant at low exposure levels. The CYP-related pathway results in the addition of oxygen, followed by spontaneous rearrangement to formyl chloride, and then to CO; each spontaneous rearrangement releases H⁺ and Cl⁻ ions. At higher exposure levels, the CYP pathway becomes saturated and a second pathway begins to predominate. Glutathione S-transferase (GST)-catalyzed addition of glutathione (GSH) is the initial step in this pathway. The replacement of one of the chlorine atoms with the S-glutathione group results in formation of S-(chloromethyl)glutathione and the release of H⁺ and Cl⁻ ions. Hydration of S-(chloromethyl)glutathione results in an S-glutathionyl methanol molecule, which can spontaneously form formaldehyde or rearrange to form an S-glutathione formaldehyde molecule, and then further rearrange to formate. Both formaldehyde and formate can then be further metabolized to CO₂.



Adapted from: ATSDR (2000); Guengerich (1997); Hashmi et al. (1994); Gargas et al. (1986).

Figure 3-1. Proposed pathways for dichloromethane metabolism.

As described in the following discussion of the two pathways, a metabolic balance appears to exist between them, with the CYP pathway tending to be relatively more active at lower doses and the fraction of dichloromethane metabolized by the GST pathway increasing at higher exposure levels, as the CYP pathway becomes saturated. Both pathways are assumed to be operating, however, even at low exposures. Exposure to other agents may shift the balance

between the pathways. For example, pretreatment with compounds that deplete GSH (e.g., buthionine sulfoximine, diethylmaleate, phorone) resulted in an increase in blood carboxyhemoglobin (COHb) levels following a single injection of dichloromethane relative to animals that did not receive GSH depletion, indicating a shift to the CYP pathway (Oh et al., 2002). Similarly, co-exposure to agents that compete for CYP2E1 results in a shift toward the GST pathway and away from CO production (Lehnebach et al., 1995; Pankow and Jagielki, 1993; Pankow et al., 1991a, b; Glatzel et al., 1987; Roth et al., 1975).

3.3.1. The CYP2E1 Pathway

There is considerable evidence of the importance of the CYP2E1 metabolic pathway in studies in animals (Oh et al., 2002; Wirkner et al., 1997; Kim and Kim, 1996; Lehnebach et al., 1995; Pankow et al., 1991a, b; Pankow and Hoffmann, 1989; Pankow, 1988; Glatzel et al., 1987; Angelo et al., 1986a, b; Landry et al., 1983; Anders and Sunram, 1982; McKenna et al., 1982; McKenna and Zempel, 1981; Rodkey and Collison, 1977; Carlsson and Hultengren, 1975; Roth et al., 1975; Fodor et al., 1973) and humans (Takeshita et al., 2000; DiVincenzo and Kaplan, 1981; Astrand et al., 1975). These studies demonstrate that exposure to dichloromethane, regardless of exposure route, results in the formation of CO, as assessed by direct measurements of elevated levels of CO in expired air and increased levels of COHb in the blood.

The first step in the CYP2E1 pathway is the formation of formyl chloride (Figure 3-1). Watanabe and Guengerich (2006) conducted a series of studies to investigate the downstream metabolites of formyl chloride and reported only marginal (3% maximum at pH 9) formation of *S*-formyl GSH from formyl chloride in the presence of GSH. Therefore, most (>97%) of the formyl chloride is metabolized further to CO. Furthermore, CO formation from formyl chloride was independent of GSH presence in the assay.

Results from numerous studies in rats in which CYP2E1 metabolism was blocked or induced indicate that the generation of CO occurs as a result of metabolism of dichloromethane by the CYP2E1 pathway (Figure 3-1). Co-exposure of rats to a high dose of ethanol (174 mmol/kg), which is metabolized by CYP2E1, and dichloromethane (1.6, 6.2, 15.6 mmol/kg) resulted in no increase in blood COHb, indicating that the metabolic pathway for CO formation had been either blocked or saturated (Glatzel et al., 1987). Similar results have been seen with coadministration of other known CYP substrates, including diethyldithiocarbamate (Lehnebach et al., 1995), methanol (Pankow and Jagielki, 1993), benzene, toluene, and three xylene isomers (Pankow et al., 1991b). Pretreatment of animals with CYP inducers (e.g., benzene, toluene, xylenes, methanol, isoniazid), particularly those that induce CYP2E1, resulted in an increased level of CO formation, as assessed by COHb formation or measurement in expired air, following single exposures to dichloromethane (Kim and Kim, 1996; Pankow and Jagielki, 1993; Pankow et al., 1991b; Pankow and Hoffmann, 1989; Pankow, 1988). Pretreatment with disulfuram, a CYP2E1 blocker, resulted in a complete lack of formation of

COHb following dichloromethane exposure, indicating that CYP2E1 is the isozyme responsible for metabolism of dichloromethane (Kim and Kim, 1996).

Evidence in hamster and rat studies suggests that the CYP2E1 pathway becomes saturated at high dichloromethane exposure levels; comparable data from studies in mice were not found. In hamsters, mean COHb percentages were elevated to a similar degree (about 28–30%, compared with <1% in controls) in three groups exposed by inhalation to 500, 1,500, or 3,500 ppm dichloromethane for 6 hours (Burek et al., 1984). After 21 months of exposure by this protocol, mean COHb percentages in the three exposure groups remained similarly elevated, indicative of saturation of the CYP2E1 pathway in hamsters at exposure levels ≥ 500 ppm and a lack of accumulation of dichloromethane and CYP2E1 metabolites with chronic exposure. McKenna et al. (1982) found that blood COHb levels in rats increased when inhalation exposure concentration was increased from 50 to 500 ppm but that similar levels of COHb were reported following exposure to 1,500 ppm as following exposure to 500 ppm; the peak blood COHb percentages were approximately 10%. In rats exposed to 0, 50, 200, or 500 ppm for 6 hours/day, 5 days/week for 2 years, mean COHb percentages were 2.2, 6.5, 12.5, and 13.7%, respectively, suggesting that saturation of the CYP2E1 pathway is approached at 200 ppm (Nitschke et al., 1988a). In male F344 rats exposed for 4 hours to dichloromethane concentrations of about 150, 300, 600, 1,000, and 2,000 ppm, mean COHb percentages (estimated from a figure) were about 4% at 150 ppm and about 8% at each of the four higher exposure concentrations (Gargas et al., 1986). McKenna and Zempel (1981) reported that increasing the oral dose of labeled dichloromethane from 1 to 50 mg/kg in rats resulted in a lower fraction of the total dose being metabolized to CO. Single injections of 3 and 6 mmol/kg of dichloromethane in rats resulted in nearly identical levels of blood COHb (Oh et al., 2002).

In human subjects exposed to dichloromethane in the workplace, saturation of CYP metabolism appears to be approached by the 400–500 ppm range (Ott et al., 1983e). Blood samples were drawn during working hours from 136 fiber production workers who were exposed to dichloromethane, acetone, and methanol. TWA exposure concentrations for the workers were determined by personal monitoring techniques, and percent COHb levels in the blood samples were determined. Estimated TWA concentrations in the exposed workers followed a bimodal distribution, with a lower mode of exposure concentrations in the 150–200 ppm range and the higher mode in the range of 450–500 ppm; only 21% (29 out of 136 workers) were in the 200–400 ppm range. Plots of percent COHb against TWA exposure concentrations showed the appearance of saturation at around 400 ppm.

The liver is the tissue most enriched in CYP2E1 catalytic activity, but CYP2E1 protein and messenger ribonucleic acid (mRNA) have been detected in other human tissues, including the lung, brain, kidney, pancreas, bladder, small intestine, and blood lymphocytes (Nishimura et al., 2003). As such, the liver is expected to be the main site of CYP metabolism of dichloromethane, but other tissues are also expected to metabolize dichloromethane via this

pathway. Of particular relevance given the neurologic effects seen with dichloromethane are the distribution and inducibility of CYP2E1 in different areas of the brain (Miksys and Tyndale, 2004). Individuals with decreased CYP2E1 activity may experience decreased generation of CO and an increased level of GST-related metabolites following exposure to dichloromethane. As a result, these individuals may be more susceptible to the chronic effects of dichloromethane from GST-related metabolites than individuals with higher levels of CYP2E1 activity. Conversely, individuals with higher CYP2E1 activity may experience relatively increased generation of CO at a given dichloromethane exposure level and, therefore, may be more susceptible to the acute toxicity of dichloromethane (from CO).

Results from studies examining human interindividual variation in CYP2E1 activities (e.g., catalytic activities, protein levels, or mRNA levels) indicate that individuals may vary in their ability to metabolize dichloromethane through the CYP2E1 pathway. In a study of liver samples from 30 Japanese and 30 Caucasian individuals, two- to threefold variation was found in the levels of CYP2E1 protein, whereas catalytic activity toward substrates associated with CYP2E1 (e.g., 7-ethoxycoumarin) displayed a wider range of values, approximately 25-fold; no clear gender-specific or ethnic differences were found in hepatic levels of CYP2E1 protein or enzymatic activities associated with CYP2E1 (Shimada et al., 1994). In a study of interindividual variation in 70 healthy human subjects (40 men and 30 women) given an oral dose of chlorzoxazone, a therapeutic agent whose metabolism and blood clearance has been related to CYP2E1 levels, a three- to fourfold range in plasma half-life and clearance values was observed, with no clear or dramatic age- or gender-specific differences (Kim et al., 1995). A six- to sevenfold range in chlorzoxazone hydroxylation activity was reported for a group of 69 healthy, smoking and nonsmoking male and female volunteers with mixed ethnic backgrounds; the range was markedly increased when a group of 72 alcoholic inpatients was included (Lucas et al., 1999). In studies of human liver microsomes, four- to sixfold ranges in CYP2E1-dependent oxidation of trichloroethylene have been reported (Lipscomb et al., 2003, 1997). CYP2E1 protein levels in 50 specimens of human lymphocytes from healthy individuals showed an approximate fivefold range (Bernauer et al., 2000), and a 3.7-fold range in liver CYP2E1 mRNA levels was reported for a group of 24 patients with chronic hepatitis (Haufrond et al., 2003). More recently, a threefold range was reported for maximal rates of hepatic CYP2E1-catalyzed metabolism of dichloromethane, which were estimated with a modified physiologically based pharmacokinetic (PBPK) model originally developed by Andersen et al. (1987) and kinetic data (e.g., dichloromethane breath and blood concentrations) for 13 volunteers (10 males and 3 females) exposed to one or more concentrations of dichloromethane by inhalation for 7.5 hours (Sweeney et al., 2004). In summary, most studies indicate a three- to sevenfold variability in CYP2E1 activity, as assessed by various types of measurements, among “healthy” volunteers. However, various clinical factors (i.e., obesity, alcoholism, use of specific

medications) or co-exposures (i.e., to various solvents) (Lucas et al., 1999) may result in greater variation and thus the potential for saturation at lower exposures within the general population.

Several genetic polymorphisms for the human CYP2E1 gene have been described, but clear and consistent correlations with interindividual variation in CYP2E1 protein levels or associated enzyme activities have not been identified (Ingelman-Sundberg, 2004; Lucas et al., 2001; Kim et al., 1995; Shimada et al., 1994). The most frequently studied CYP2E1 polymorphisms, *RsaI/PstI*, are located in the 5'-flanking region of the gene, and mutations are thought to lead to increased CYP2E1 protein expression via transcription (Lucas et al., 2001). Available data indicate that the frequency of this polymorphism, as well as other CYP2E1 polymorphisms, varies among ethnic groups. For example, Stephens et al. (1994) examined blood samples from 126 African-Americans, 449 European Americans, and 120 Taiwanese subjects and found frequencies for a rare *RsaI* allele (C2) of 0.01 in African-Americans, 0.04 in European Americans, and 0.28 in Taiwanese subjects. In a study of 102 Mexicans, the reported mutation frequency at the *RsaI* C2 allele was 0.30 (Mendoza-Cantú et al., 2004).

3.3.2. The GST Pathway

The other major pathway for dichloromethane metabolism involves the conjugation of dichloromethane to GSH, catalyzed by GST. This results in the formation of a GSH conjugate that is eventually metabolized to CO₂ (Figure 3-1). The conjugation of dichloromethane to GSH results in formation of two reactive intermediates that have been proposed to be involved in dichloromethane toxicity, S-(chloromethyl)glutathione and formaldehyde. In studies with rat, mouse, and human liver cytosol preparations in the presence of GSH, examination of metabolites with [¹³C]-NMR indicated that S-(chloromethyl)glutathione was an intermediate in the pathway to formaldehyde (Hashmi et al., 1994). Formaldehyde formation from dichloromethane has been noted in human (Bruhn et al., 1998; Hallier et al., 1994; Hashmi et al., 1994), rat, and mouse (Casanova et al., 1997; Hashmi et al., 1994) cells in vitro. Formation of a free hydrogen ion is also hypothesized, although no direct evidence supporting this has been presented.

The GST pathway has approximately a 10-fold lower affinity for dichloromethane than the CYP pathway (Reitz et al., 1989; Andersen et al., 1987). Although both pathways are assumed to be operating at all exposures, at lower exposure concentrations the CYP pathway is expected to predominate, and as exposure concentrations increase, the GST pathway is expected to gain in relative importance as a dispositional pathway for absorbed dichloromethane. Based on in vitro studies with liver preparations, the estimated Michaelis-Menten kinetic constant (K_m) values in GST assays with dichloromethane were about 137 mM in a B6C3F₁ mouse preparation and about 44 mM in two human preparations (Reitz et al., 1989). In contrast, estimated K_m values in CYP assays were about 1.8, 1.4, and 2.0 mM in B6C3F₁ mouse, F344 rat, and Syrian golden hamster preparations, respectively. In four human liver preparations, estimated CYP K_m values were about 2.6, 2.0, 0.9, and 2.8 mM (Reitz et al., 1989). A possible resolution of these

apparent in vitro vs. in vivo discrepancies is discussed in Section 3.5.5 (in particular, see Figure 3-6).

Early investigations indicated that in humans, GSTs of the α -, μ -, and π -classes were not responsible for the metabolism of dichloromethane (Bogaards et al., 1993). Tissue samples that metabolized substrates specific to those GST classes did not conjugate dichloromethane to GSH. Later investigations identified the recently-characterized GST theta class (Meyer et al., 1991), specifically GST-theta1-1 (GST-T1), as the GST isoenzyme responsible for the metabolism of dichloromethane (Mainwaring et al., 1996; Blocki et al., 1994). In the absence of the GST-T1 gene, no deoxyribonucleic acid (DNA)-protein cross-links were formed by human liver cells exposed to dichloromethane (Casanova et al., 1997), and formaldehyde production was not detected in human erythrocytes (Hallier et al., 1994). In a mouse model with a disrupted GST-T1 gene, GST activity with dichloromethane in liver and kidney cytosol samples was substantially lower compared with wild-type GST mice (Fujimoto et al., 2007).

A polymorphism of the GST-T1 gene has been demonstrated in humans. People with two functional copies of the gene (+/+) readily conjugate GSH to dichloromethane. Individuals having only one working copy of the gene (+/-) display relatively decreased conjugation ability. Individuals with no functional copy of the gene (-/-) do not express active GST-T1 protein and do not metabolize dichloromethane via a GST-related pathway (Thier et al., 1998). Results from studies of GST-T1 genotypes in human blood samples indicate that average prevalences of the GST-T1 null (-/-) genotype are higher in Asian ethnic groups (47–64%) than in other groups, including Caucasians (19–20%), African-Americans (22%), and mixed groups (19%) (Raimondi et al., 2006; Garte et al., 2001; Nelson et al., 1995) (see Table 3-3). Although information on the age distribution of study subjects was not generally reported in these analyses, there is little reason to expect effect modification by age since this is not a gene linked to early mortality. Based on data collected by Nelson et al. (1995) and U.S. 2000 census data (and assuming Hardy-Weinberg equilibrium), Haber et al. (2002) calculated U.S. average distributions of GST-T1 genotypes as follows: 32% +/+; 48% +/-; and 20% -/-.

Table 3-3. Mean prevalences of the GST-T1 null (-/-) genotype in human ethnic groups

Ethnic group	Reference		
	Nelson et al. (1995) ^a	Garte et al. (2001) ^b	Raimondi et al. (2006) ^c
Chinese	64.4% (n = 45)	Not reported	Not reported
Korean	60.2% (n = 103)	Not reported	Not reported
Caucasian	20.4% (n = 442)	19.7% (n = 5,577)	19.0% (n = 6,875)
Asian	Not reported	47.0% (n = 575)	53.6% (n = 1,727)
African-American	21.8% (n = 119)	Not reported	Not reported
Mexican American	9.7% (n = 73)	Not reported	Not reported
Other	Not reported	Not reported	19.4% (n = 1,485)

^aNelson et al. (1995) examined prevalence of the null GST-T1 genotype from analysis of blood samples from subjects of various ethnicities as noted above.

^bGarte et al. (2001) collected GST-T1 genotype data in Caucasian (29 studies; 5,577 subjects) and Asian (3 studies, 575 subjects) ethnic groups; subjects were controls in case-control studies of cancer and various polymorphisms in genes for bioactivating enzymes.

^cRaimondi et al. (2006) collected GST-T1 genotype data from 35 case-control studies of cancer and GST-T1 genotype; data in this table are for control subjects. The “other” group in this study is defined as Latino, African-American, and mixed ethnicities.

Results from a study of the distribution of activity levels for in vitro conjugation of dichloromethane with GSH in 22 human liver samples are roughly reflective of these estimates of the distribution of this polymorphism (Bogaards et al., 1993). No activity was found in 3/22 of the liver samples. Eleven of the samples showed low activity levels (0.21–0.41 nmol product/minute/mg protein), and eight samples showed high activity levels ranging from 0.82 to 1.23 nmol/minute/mg protein. In another study of seven human subjects, lysates of erythrocytes showed high activities for producing formaldehyde from dichloromethane (presumably via GST-T1) in three subjects (15.4, 17.7, and 17.8 nmol product/minute/mg hemoglobin) and lower activity in the other four subjects (4.3, 6.0, 7.2, and 7.6 nmol product/minute/mg hemoglobin) (Hallier et al., 1994).

Comparisons of mice, rats, humans, and hamsters for the ability to metabolize dichloromethane via the GST pathway in liver and lung tissues indicate that mice appear to be the most active at metabolizing dichloromethane (Sherratt et al., 2002; Thier et al., 1998; Casanova et al., 1997, 1996; Hashmi et al., 1994; Reitz et al., 1989). Reitz et al. (1989) reported mean (\pm SD) GST enzymatic activity levels with dichloromethane as substrate (in units of nmol product formed/minute/mg protein) in liver cytosol preparations to be: 25.9 ± 4.2 units in B6C3F₁ mice (n = 15 determinations per preparation); 7.05 ± 1.7 nmol/minute/mg in F344 rats (n = 6); and 1.27 ± 0.21 nmol/minute/mg in Syrian golden hamsters (n = 6). Mean GST activity levels in liver preparations from four human subjects (accident victims screened for human immunodeficiency virus and hepatitis B and C and obtained through a transplant center) were 2.62 ± 0.44 (n = 10), -0.01 ± 0.04 (n = 6), 2.71 ± 0.45 (n = 6), and 3.03 ± 0.44 nmol/minute/mg

(n = 6) (Reitz et al., 1989). The finding that one of the four individuals was unable to conjugate dichloromethane with GST was reflective of the estimated frequency of the GST-T1 null genotype in the U.S. population (approximately 20% in Caucasians and African-Americans see Table 3-3 and Haber et al., 2002). Mean GST activity levels in lung cytosol preparations showed a similar rank order among species: 7.3 ± 1.4 nmol/minute/mg in mice (n = 4), 1.0 ± 0.1 nmol/minute/mg in rats (n = 4), 0.0 ± 0.2 nmol/minute/mg in hamsters (n = 4), and 0.37 ± 0.25 nmol/minute/mg in a pooled lung preparation from the same four human subjects (n = 2).

Thier et al. (1998) conducted a study evaluating the activity of GST-T1 after treatment of dichloromethane in the cytosol of liver and kidney homogenates from hamsters (pooled male and females), rats (pooled male and female), male mice, and female mice and for humans classified as nonconjugators, low conjugators, or high conjugators of GST to dichloromethane. Little information is provided about the human samples other than that 13 kidney cancer patients were the source of the kidney samples; normal tissue identified by pathological exam was used. Blood samples from 10 of these patients were collected, and enzyme activities measured in erythrocytes from 9 of these samples were reported. Results of conjugation of dichloromethane to GSH from these studies are presented in Table 3-4. As can be seen from the table, activity levels (expressed as nmol/minute per mg of cytosolic protein) of humans varied considerably, with nonconjugators (presumed to be GST-T1^{-/-}) having no detectable activity, low conjugators (presumed to be GST-T1^{+/-}) having moderate activity, and high conjugators (presumed to be GST-T1^{+/+}) having approximately twice the activity seen in low conjugators. In the liver, the activity of rat GST conjugation was over twofold that seen in human high conjugators, while levels in mice were >11-fold (males) or 18-fold (females) greater than those of human high conjugators. In the kidney, the activity of high-conjugator humans was approximately 1.8-fold that of rats and was comparable to the activity of both male and female mice. The data in Table 3-4 show the following order for GST-T1 activities with dichloromethane as substrate: in liver preparations, mouse >> rat > human high conjugators > human low conjugators > hamster > human nonconjugators and, in kidney preparations, female mouse \approx male mouse \approx human high conjugators > rat \approx human low conjugators > hamster > human nonconjugators. In addition, the data indicate that activity levels in liver, kidney, and erythrocytes of human subjects are in correspondence with the nonconjugator, low conjugator, and high conjugator designations.

Table 3-4. GST-T1 enzyme activities toward dichloromethane in human, rat, mouse, and hamster tissues (liver, kidney, and erythrocytes)

	Activity (nmol/min per mg protein) ^a		Activity (nmol/min per mL) ^a
	Liver	Kidney	Erythrocytes
Human, nonconjugators	Not detectable (2)	Not detectable (1)	Not detectable (1)
Human, low conjugators	0.62 ± 0.30 (11)	1.38 ± 0.52 (8)	9.67 ± 2.49 (5)
Human, high conjugators	1.60 ± 0.48 (12)	3.05 ± 0.72 (4)	18.28 ± 0.46 (3)
Rat	3.71 ± 0.28 (8)	1.71 ± 0.28 (8)	Not measured
Mouse, male	18.2 ± 2.22 (5)	3.19 ± 0.46 (5)	Not measured
Mouse, female	29.7 ± 6.31 (5)	3.88 ± 0.90 (5)	Not measured
Hamster	0.27 ± 0.20 (6)	0.25 ± 0.21 (6)	Not measured

^aMean ± SD with number of samples noted in parentheses.

Source: Adapted from Thier et al. (1998).

Sherratt et al. (2002) reported that, on a per mg basis, native recombinant mouse GST-T1 (purified after expression in *Escherichia coli*) was approximately twofold more active toward dichloromethane than native recombinant human enzyme, as well as being approximately fivefold more efficient (as assessed by the ratio of k_{cat}/K_m), where k_{cat} is the maximum rate of the reaction catalyzed by the enzyme *per enzyme molecule*; i.e., V_{max}/E_t where E_t is the total enzyme concentration).

The distribution of GST-T1 in human tissues has been examined with antibodies raised against recombinant human GST-T1 (Sherratt et al., 2002, 1997). Immunoblotting of sodium dodecyl sulfate polyacrylamide gel electrophoresis gels loaded with tissue extracts from a 73-year-old man who had died with bronchopneumonia and atherosclerosis indicated the following order of expression of GST-T1: liver ≈ kidney > prostate ≈ small intestine > cerebrum ≈ pancreas ≈ skeletal muscle > lung ≈ spleen ≈ heart ≈ testis (Sherratt et al., 1997). It was estimated that the levels of cross-reacting materials in the cerebrum, pancreas, or skeletal muscle extracts were about 10% of those in the liver, whereas levels in the lung, spleen, heart, and testis were <5% of the levels in the liver. Comparison of the amounts of cross-reacting material in soluble liver extracts from a B6C3F₁ mouse and five human subjects (i.e., normal liver tissue samples from biopsies of secondary liver tumors) found that levels of GST-T1 protein were higher in the mouse extracts than in any of the human liver extracts (Sherratt et al., 2002). Densitometer analysis indicated that the GST-T1 level in the mouse liver extract was about fivefold higher than those in human liver extracts displaying the highest level. Cross-reacting material was not detectable in liver extracts from one of the five human subjects, indicating that this individual may have been GST-T1 null (Sherratt et al., 2002).

Results from *in situ* hybridization with oligonucleotide antisense probes for GST-T1 mRNA levels and immunohistochemical studies with antibodies to GST-T1 have indicated that there may be subtle differences between mice and humans in the intracellular localization of GST-T1 in the liver. Mainwaring et al. (1996) reported that staining for GST-T1 mRNA was higher in liver slices from B6C3F₁ mice than in liver slices from F344 rats and that staining in human liver samples was very low. Although the number of mouse and rat liver samples examined in this study was not indicated in the available report, it was reported that slices from five human liver samples were examined. No information was provided regarding the clinical history of the sources of the human samples. In mouse liver, staining for GST-T1 mRNA was enhanced in the limiting plate hepatocytes, in nuclei, in bile-duct epithelial cells, and in lesser amounts in the centrilobular cells in general. In rat liver, a similar pattern was observed, except no enhanced staining was observed in the limiting plate hepatocytes or in nuclei. Staining for GST-T1 mRNA in the human liver samples showed an even distribution throughout the liver lobule, and no mention of a specific nuclear localization was made (Mainwaring et al., 1996). Quondamatteo et al. (1998), using antibodies to GST-T1, subsequently reported a similar localization of GST-T1 protein in nuclei of cells in mouse liver slices. In another study using antibodies raised against recombinant human GST-T1 or a peptide derived from the deduced mouse GST-T1 primary sequence, Sherratt et al. (2002) reported that nuclear staining was observed in all cells in mouse liver slices (from five individual B6C3F₁ mice) showing the presence of mouse GST-T1; staining in the cytoplasm was only detected in cells with very high levels of GST-T1. In liver slices obtained from two human subjects (males, ages 60 and 61 years, with a secondary liver tumor and what was described as a “cavernous hemangioma” without malignancy, respectively), the most intense nuclear staining was associated with bile duct epithelial cells, but there was heterogeneity of staining within hepatocytes; some cells showed nuclear staining, but others only exhibited cytoplasmic staining (Sherratt et al., 2002).

In summary, the relative amount of dichloromethane metabolized via the GST pathway increases with increasing exposure concentrations. As the high affinity CYP pathway becomes saturated (either from high exposure levels of genetic or other factors that decrease CYP2E1 activity), the GST pathway increases in relative importance as a dispositional pathway for dichloromethane. Two reactive metabolites (*S*-(chloromethyl)glutathione and formaldehyde) resulting from this pathway have been identified. GST-T1 is the GST isozyme that catalyzes conjugation of dichloromethane with GST. Interindividual variation in the ability to metabolize dichloromethane via GST-T1 is associated with genetic polymorphisms in humans. Estimated U.S. population prevalence of nonconjugators (-/- at the GST-T1 locus) is about 20%, but higher prevalences (47–64%) have been reported for Asians (Raimondi et al., 2006; Haber et al., 2002; Garte et al., 2001; Nelson et al., 1995). The prevalences for low (+/- at the GST-T1 locus) and high (+/+) conjugators have been estimated at 48 and 32%, respectively (Haber et al., 2002). The liver and kidney are the most enriched tissues in GST-T1, but evidence is available for the

presence of GST-T1 in other tissues at lower levels, including the brain and lung. In humans, GST-T1 expression in the brain is lower than that seen in the liver or kidney but higher than in the lung. Comparisons of mice, rats, humans, and hamsters for the ability to metabolize dichloromethane via the GST pathway in liver (based on measurement of tissue-specific enzyme activity) indicate the following rank order: mice > rats > or \approx humans > hamsters. In mouse liver tissue, GST-T1 appears to be localized in the nuclei of hepatocytes and bile-duct epithelium, but rat liver does not show preferential nuclear localization of GST-T1. In human liver tissue, some hepatocytes show nuclear localization of GST-T1 and others show localization in cytoplasm, as well as in bile duct epithelial cells. The apparent species differences in intracellular localization of GST-T1 may play a role in species differences in susceptibility to dichloromethane carcinogenicity if nuclear production of S-(chloromethyl)glutathione is more likely to lead to DNA alkylation than cytoplasmic production.

3.4. ELIMINATION

Dichloromethane is eliminated mainly through exhalation either of the parent compound or as the two primary metabolites CO₂ and CO (Angelo et al., 1986a, b; McKenna et al., 1982; DiVincenzo and Kaplan, 1981; DiVincenzo et al., 1972, 1971). In human studies, dichloromethane is rapidly eliminated from the body following the cessation of exposure, with much of the parent compound completely removed from the bloodstream and expired air by 5 hours postexposure in experiments using exposure levels of 90, 100, or 210 ppm (DiVincenzo et al., 1972, 1971; Riley et al., 1966). Studies in rats have similarly demonstrated that elimination from the blood is rapid, with elimination half-times in F344 rats on the order of 4–6 minutes following intravenous doses in the range of 10–50 mg/kg (Angelo et al., 1986a). In a study using Sprague-Dawley rats, Carlsson and Hultengren (1975) demonstrated variability in elimination rates between different types of tissues, with the most rapid elimination seen in the adipose and brain tissue, while elimination from liver, kidneys, and adrenals proceeded more slowly.

In a study using human volunteers, DiVincenzo and Kaplan (1981) reported a dose-related increase in CO in the expired breath after inhalation exposure to 50–200 ppm of dichloromethane, with a net elimination as CO on the order of 25–35% of the absorbed dose. Similar results have been reported in animal studies. Following gavage administration of 50 or 200 mg/kg-day doses of [¹⁴C]-labeled dichloromethane in water to groups of six mature male F344 rats for up to 14 days, >90% of the label was recovered in the expired air within 24 hours of dose administration (Angelo et al., 1986b). Following administration of the first of 14 daily 50 mg/kg-day doses, radioactivity in parent compound, CO₂, and CO in the 24-hour expired breath accounted for 66, 17, and 16% of the administered radioactivity, respectively; similar patterns were reported for 24-hour periods following administration of the seventh and fourteenth 50 mg/kg-day dose. Following administration of the first 200 mg/kg-day dose,


radioactivity in parent compound, CO₂, and CO in the 24-hour expired breath accounted for 77, 9, and 6%, respectively, of the administered radioactivity (Angelo et al., 1986b). In mature, male Sprague-Dawley rats given a smaller dose (1 mg/kg) of [¹⁴C]-labeled dichloromethane, radioactivity in parent compound, CO₂, and CO in 48-hour expired breath accounted for 12, 35, and 31%, respectively; these data indicate that, at lower dose levels, a greater percentage of the administered dose was metabolized by the CYP pathway and eliminated in the expired breath, compared with higher dose levels (McKenna and Zempel, 1981). Similar patterns of radioactivity distribution in parent compound, CO₂, and CO in expired breath were found in mature, male B6C3F₁ mice following gavage administration of 50 mg/kg-day (in water) or 500 or 1,000 mg/kg-day (in corn oil) [¹⁴C]-labeled dichloromethane (Angelo et al., 1986a). For example, radioactivity in parent compound, CO₂, and CO in 24-hour expired breath accounted for 61, 18, and 11% of the administered radioactivity, following administration of a single 50 mg/kg dose to a group of six mice (Angelo et al., 1986a). Exhalation rates were similarly high following inhalation exposure of mature, male Sprague-Dawley rats (>90%) (McKenna et al., 1982) or following intravenous administration of dichloromethane to mature, male F344 rats (Angelo et al., 1986b).

Elimination of dichloromethane in the urine of exposed humans is generally small, with total urinary dichloromethane levels on the order of 20–25 or 65–100 µg in 24 hours following a 2-hour inhalation exposure to 100 or 200 ppm, respectively (DiVincenzo et al., 1972). However, a direct correlation between urinary dichloromethane and dichloromethane exposure levels was found in volunteers, despite the comparatively small urinary elimination (Sakai et al., 2002). Following administration of a labeled dose in animals, regardless of exposure route, generally <5–8% of the label is found in the urine and <2% in the feces (McKenna et al., 1982; McKenna and Zempel, 1981; DiVincenzo et al., 1972, 1971).

3.5. PHYSIOLOGICALLY BASED TOXICOKINETIC MODELS

Several PBPK models for dichloromethane in animals and humans have been developed from 1986 to 2006. These models are mathematical representations of the body and its absorption, distribution, metabolism, and elimination of dichloromethane and select metabolites, based on the structure of the Ramsey and Andersen (1984) model for styrene. The models' equations are designed to mimic actual biological behavior of dichloromethane, incorporating *in vitro* and *in vivo* data to define physiological and metabolic equation parameters. As such, the models can simulate animal or human dichloromethane exposures and predict a variety of dichloromethane and metabolite internal dosimeters (i.e., instantaneous blood and tissue concentration, area under the curve [AUC] of concentration versus time plots, rate of metabolite formation), allowing for the extrapolation of toxicity data across species, route of exposure, and high to low exposure levels. The development of dichloromethane PBPK models has resulted in either increased biological detail and functionality or refinement of model parameters with newly

available data. The former type of development provides more options for toxicity data extrapolation, while the latter serves to increase confidence in model predictions and decrease uncertainty in risk assessments for which the models were, or will be, applied. This section of the document describes each of the models reported in the scientific literature and/or used by the regulatory community (i.e., Occupational Safety and Health Administration [OSHA], EPA) and their contribution to the advancement of predictive dosimetry and data extrapolation for dichloromethane. In some instances, model development was accomplished by the addition of new biological compartments (e.g., tissue systems). Diagrams of the compartmental structure of the models are shown in Figure 3-2. Significant statistical advances in parameter estimation also have been incorporated in model development. For this reason, some animal and human PBPK models may be described as deterministic (Sweeney et al., 2004; Casanova et al., 1996; Reitz et al., 1988a, b; U.S. EPA, 1988b, 1987a, b; Andersen et al., 1987; Gargas et al., 1986) in which point estimates for each model parameter are used, resulting in point estimates for dosimetry. Others may be described as probabilistic (Jonsson and Johanson, 2001; El-Masri et al., 1999; OSHA, 1997), in which probability distributions for each parameter were defined, resulting in probability distributions for dosimetry. The latter approach, particularly utilizing a Bayesian hierarchical statistical model structure (described below) (David et al., 2006; Marino et al., 2006) to estimate parameter values, allows for the introduction of intra- and interspecies variability into model predictions and quantitative assessment of model uncertainty. Both deterministic (U.S. EPA, 1988b, 1987a, b) and probabilistic (OSHA, 1997) applications have been used to develop regulatory values. As discussed below, subsequent applications of the developed models for cancer risk assessment have resulted in significantly different estimates of human cancer risk.



Models C–G all build on the structure in model B. Models E and G have been applied in humans; all others have been applied in humans and rodents (mice and/or rats).

CYP = CYP pathway metabolites; GST = GST pathway metabolites

Adapted from: Model A—Gargas et al. (1986); B—Andersen et al. (1987); C—Andersen et al. (1991); D—Casanova et al. (1996); E—Sweeney et al. (2004); F—OSHA (1997); G—Jonsson and Johanson (2001).

Figure 3-2. Schematics of PBPK models (1986–2006) used in the development of estimates for dichloromethane internal dosimetry.

The deterministic rat model of Gargas et al. (1986), based on previous work by Ramsey and Andersen (1984) examining inhalation pharmacokinetics of styrene in rats, was the first PBPK model for dichloromethane. It comprised four compartments (fat, liver, richly perfused tissues, and slowly perfused tissues [Figure 3-2A]) and described flows and partitioning of parent material and metabolites through the compartments with differential equations. Metabolism, which was restricted to the liver compartment, was described as two competing pathways: the GST pathway, described with a linear first-order kinetic model, and the CYP pathway, described with a saturable Michaelis-Menten kinetic model. Rate constants for the CYP and GST pathways in rats were determined by optimization of the model with in vivo gas uptake data. COHb production was modeled both endogenously and from CYP-mediated metabolism of dichloromethane. This model demonstrated the dose-dependent flux through the competing CYP and GST metabolic pathways and the effect of CYP inhibition on COHb generation.

Andersen et al. (1987) extended the rat model of Gargas et al. (1986) to include a lung compartment, including CYP and GST metabolism pathways within the lung, in rats, mice, hamsters, and humans (Figure 3-2B). Physiological flow rates were allometrically scaled among species by $\frac{3}{4}$ power of body weight (BW). Rate constants for the CYP and GST pathways in rodents were determined by optimization of the model with in vivo gas uptake data. CYP rate constants for humans were derived from data on dichloromethane uptake in human subjects (number of subjects not reported). Human GST rate constants were derived by allometric scaling of the animal GST rate constants. Model predictions compared favorably with kinetic data for human subjects exposed by inhalation to dichloromethane (Andersen et al., 1987). Using the mouse cancer bioassay data from NTP (1986), Andersen et al. (1987) compared the linear body surface area-derived or the PBPK model-derived human liver and lung dose surrogates associated with tumor development (mg dichloromethane metabolized via GST pathway/volume tissue/day). They reported that PBPK model-extrapolated human liver and lung internal doses were 167- and 144-fold lower for inhalation exposure and 45- and 213-fold lower for drinking water exposure, respectively, than body surface area scaled internal doses. The study authors suggested that the lower model-predicted human internal dose surrogates were due to the need to saturate the CYP pathway before appreciable tumorigenic metabolite levels could be attained, which is not captured by extrapolation based on body surface area.

U.S. EPA (1988b, 1987a, b) slightly modified the Andersen et al. (1987) model for mice by using different alveolar ventilation and cardiac flow rates and used the mouse and human models to derive human cancer risks from animal tumor incidence data. The flow rate parameters in the Andersen et al. (1987) model were based on a human breathing rate of 12.5 m³/day (reflecting a resting rate), compared with the EPA value of 20 m³/day (reflecting average daily activity level), and a mouse breathing rate of 0.084 m³/day (based on allometric scaling of bioassay-specific BWs), compared with the rate commonly used by EPA, 0.043 m³/day (U.S. EPA, 1987a). The internal dose metric used in the applications of the model

to cancer risk assessment was reflective of the amount of dichloromethane metabolized by the GST pathway. In addition to using the mouse and human PBPK models to account for species differences in dosimetry, a body surface area correction factor of 12.7 was applied to low-dose slopes of estimated dose-response relationships for liver and lung tumors in mice to account for presumed higher human responsiveness, relative to mice, to dichloromethane-induced cancer (U.S. EPA, 1987a). The factor of 12.7 is the cube root of the ratio of human to mouse reference BWs; this BW scaling factor was applied to adjust for interspecies toxicodynamic variability (i.e., presumed differences in the lifetime impact in mice and humans of a given daily amount of dichloromethane metabolically activated per liter of tissue) (Rhomberg, 1995). A human cancer IUR of 4.7×10^{-7} per ($\mu\text{g}/\text{m}^3$), based on this analysis, was placed on IRIS in September 1990.

The Andersen et al. (1987) models were also modified by addition of submodel structures for estimation of new dosimeters of interest. Andersen et al. (1991) added the capability to specifically describe the kinetics of dichloromethane, CO, and COHb in rats and humans with the addition of the Coburn-Forster-Kane equation to describe CO and COHb kinetics (Figure 3-2C). However, equations were not added for metabolism of dichloromethane to CO in the lung. Casanova et al. (1996) extended the Andersen et al. (1987) mouse model to include a submodel that predicted the formation of formaldehyde and DNA-protein cross-links in the liver (Figure 3-2D).

Further refinements of the Andersen et al. (1987) models allowed for incorporation of new data. New in vitro measurements of metabolic rate constants in human and animal tissues were incorporated into the Andersen et al. (1987) models by Reitz and coworkers (Reitz, 1991; Reitz et al., 1988a, b). Sweeney et al. (2004) modified the Andersen et al. (1987) human PBPK model, adding extrahepatic CYP metabolism in richly perfused tissues (Figure 3-2E) to obtain a better fit of the model to kinetics data for humans. Data for 13 volunteers (10 men and 3 women) who were exposed to one or more concentrations of dichloromethane for 7.5 hours included dichloromethane concentrations in breath and blood, COHb concentrations in blood, and CO concentrations in exhaled breath. Individual CYP maximal velocity (V_{maxC}) values were obtained by optimizing model predictions to match time-course data simultaneously for dichloromethane concentrations in blood and exhaled breath for each individual. Resultant individual values of CYP V_{maxC} ranged from 7.4 to 23.6 mg/hour/kg^{0.7}, indicating an approximate threefold range in maximal CYP metabolic activity.

The significance of metabolic variability for the kinetics of dichloromethane in animals and humans was explored by several investigators using PBPK models. Dankovic and Bailer (1994) used the updated human model presented by Reitz et al. (1988a, b) to explore the consequences of interindividual variability in in vitro kinetic constants for the CYP and GST pathways (based on data for four human subjects) and reported that predicted GST-metabolized doses to the lung and liver could range from about zero to up to fivefold greater than those predicted with the values of these rate constants used in the Reitz et al. (1988a, b) model.

El-Masri et al. (1999) replaced parameter estimates in the mouse and human PBPK models presented by Casanova et al. (1996) with probability distributions, including published information on the distribution of GST-T1 polymorphism in human populations, and used Monte Carlo simulations to estimate distributions of cancer potency of dichloromethane in mice, distributions of the amount of DNA-protein cross-links formed in the liver of humans, and distributions of human cancer risks at given exposure levels of dichloromethane. The analysis showed that, at exposure levels of 1, 10, 100, and 1,000 ppm dichloromethane, average and median cancer risk estimates were 23–30% higher when GST-T1 polymorphism was not included in the model.

Given the demonstrated influence of population variability in dichloromethane metabolism on PBPK model-derived cancer risk estimates (El-Masri et al., 1999; Dankovic and Bailer, 1994), PBPK model development has included a more formal statistical treatment of data for physiological and metabolic variability. Bayesian statistical approaches have been applied to develop probabilistic PBPK models for dichloromethane. Probabilistic models account for variability between individuals in model parameters by replacing point estimates for the model parameters with probability distributions. Calibration or fitting of probabilistic PBPK models to experimental toxicokinetic data is facilitated by a Bayesian technique called Markov Chain Monte Carlo (MCMC) simulation, which quantitatively addresses both variability and uncertainty in PBPK modeling (Jonsson and Johanson, 2003).

OSHA (1997) used MCMC simulation to fit probabilistic versions of the Reitz et al. (1988a, b) and Andersen et al. (1991, 1987) mouse and human models, which included probability distributions for all model parameters. GST- and CYP-mediated metabolism occurred in the liver and lung compartments (see Figure 3-2F). The model parameters were modified to focus on occupational exposure scenarios; that is, a parameter distribution for work intensity (using data from Astrand et al. [1975]) was added, which adjusted physiological flow rates as a function of work intensity as measured in watts. In addition, updated measurements of blood:air and tissue:air partition coefficients (Clewell et al., 1993) were used to describe distributions for these parameters. The Clewell et al. (1993) blood:air partition coefficient (PB) of 23 is higher than the value of 8.29 reported by Andersen et al. (1987) and used by EPA (1988b, 1987a, b). The newer Clewell et al. (1993) value for mice is the preferred value, since it is much closer to the values for rats (19.4) and hamsters (22.5) rather than humans (9.7), as reported by Andersen et al. (1987). Distributions of metabolic, physiological, and partitioning parameters in the mouse and human models were updated by using Bayesian methods with data for mice and humans in published studies of mouse and human physiology and dichloromethane kinetic behavior.

Jonsson et al. (2001) used additional human kinetics data to expand the PBPK model of Reitz et al. (1988a, b) and added new model compartments (Figure 3-2G). These investigators used MCMC simulation to develop a probabilistic model from the Reitz et al. (1988a, b) human

model by using published in vitro measurements of liver $V_{\max C}$ for the CYP pathway (Reitz et al., 1989) and kinetic data for five human subjects exposed by inhalation to dichloromethane (Astrand et al., 1975). A working muscle compartment was added to the basic Andersen et al. (1987) and Reitz et al. (1988a, b) structure (see Figure 3-2G). Jonsson and Johanson (2001) refined and extended this probabilistic model by including an additional fat compartment (to provide a better description of the experimental data for the time course of dichloromethane in subcutaneous fat), incorporating (with MCMC simulation) kinetic data for dichloromethane in an additional 21 human subjects and including three GST-T1 genotypes/phenotypes (nonconjugators -/-, low conjugators +/-, high conjugators +/+). Monte Carlo simulations were then used with the refined probabilistic model to predict human liver cancer risk estimates at several dichloromethane exposure levels using an algorithm similar to the one used by El-Masri et al. (1999), using DNA-protein cross-links as the internal dose metric. The mean, 50th, 90th, and 95th percentile human cancer risk values from Jonsson et al. (2001) and El-Masri et al. (1999) were very similar, within onefold of one another for simulated exposure levels up to 100 ppm.

The most statistically rigorous and data-intensive PBPK model development was performed by Marino et al. (2006) for mice and David et al. (2006) for humans. Development of these models used multiple mouse and human data sets in a Bayesian hierarchical statistical structure to quantitatively capture population variability and reduce uncertainty in model dosimetry and the resulting risk values. EPA used these models in the derivation of reference values and cancer risk estimates in the current assessment, and these models are described in more detail below.

3.5.1. Probabilistic Mouse PBPK Dichloromethane Model (Marino et al., 2006)

Marino et al. (2006) used MCMC analysis to develop a probabilistic PBPK model for dichloromethane in mice, using the Andersen et al. (1987) model structure as a starting point (Figure 3-3). Metabolic kinetic parameters ($V_{\max C}$, K_m , k_{fC} , ratio of lung V_{\max} to liver V_{\max} [A1], and ratio of lung k_{fC} to liver k_{fC} [A2]) (Table 3-5) were calibrated with this Bayesian methodology by using several experimental data sets. Distribution parameters (i.e., means and coefficients of variation [CVs]) for other physiological parameters (i.e., BW, fractional flow rates, and fractional tissue volumes) and partition coefficients were taken from the general literature as noted by Clewell et al. (1993). Marino et al. (2006) noted that using distributions for these latter parameters from the general literature (based on a large number of animals) was better than updating them based on the relatively smaller number of animals in the available dichloromethane kinetic studies. Clewell et al. (1993) determined blood:air and tissue:air partition coefficients (means and CVs) with tissues from groups of male and female B6C3F₁ mice. These partition coefficients were derived by using a vial equilibration method similar to that used by prior investigators (Andersen et al., 1987; Gargas et al., 1986). Tissue:air partition

coefficients were approximately 2–3 times lower than previously utilized values with the exception of the liver coefficient, which was similar to previous values (Table 3-5). The PB (23) from Clewell et al. (1993) is higher than the previously reported value of 8.3 (Gargas et al., 1986). The higher value is more in line with values measured in rats (19.4) and hamsters (22.5) and, thus, is more reasonable than the older value of 8.3. Table 3-5 shows mean and CVs for physiological parameters and partition coefficients in the Marino et al. (2006) mouse model as well as values used in earlier deterministic PBPK mouse models for dichloromethane.




Figure 3-3. Schematic of mouse PBPK model used by Marino et al. (2006).

Table 3-5. Values for parameter distributions in a B6C3F₁ mouse probabilistic PBPK model for dichloromethane compared with associated values for point parameters in earlier deterministic B6C3F₁ mouse PBPK models for dichloromethane

Parameter	Marino et al. (2006) ^a				EPA (1988b, 1987a, b)	Andersen et al. (1987)	
	Prior mean	Prior CV	Final posterior mean	Final posterior CV			
<i>Fractional flow rates (fraction of QCC)^b</i>				These parameters were taken from an extensive literature database derived from a large number of animals; therefore, further Bayesian updating does not inform on the true mean and variance for these values.			
QFC	Fat	0.05	0.60			0.05	0.05
QLC	Liver	0.24	0.96			0.24	0.24
QRC	Rapidly perfused tissues	0.52	0.50			0.52	0.52
QSC	Slowly perfused tissues	0.19	0.40			0.19	0.19
<i>Fractional tissue volumes (fraction of BW)^b</i>							
VFC	Fat	0.04	0.30	0.04	0.04		
VLC	Liver	0.04	0.06	0.04	0.04		
VLuC	Lung	0.0115	0.27	0.0119	0.0119		
VRC	Rapidly perfused tissues	0.05	0.30	0.05	0.05		
VSC	Slowly perfused tissues	0.78	0.30	0.78	0.78		
<i>Partition coefficients^c</i>							
PB	Blood:air	23	0.15	8.29	8.29		
PF	Fat:blood	5.1	0.30	14.5	14.5		
PL	Liver:blood	1.6	0.20	1.71	1.71		
PLu	Lung:blood	0.46	0.27	1.71	1.71		
PR	Rapidly perfused:blood	0.52	0.20	1.71	1.71		
PS	Slowly perfused:blood	0.44	0.20	0.96	0.96		
<i>Flow rates</i>							
QCC	Cardiac output (L/hr/kg ^{0.74})	28.0	0.58	24.2	0.19	14.3 ^d	28.0 ^e
VPR	ventilation:perfusion ratio	1.52	0.75	1.45	0.20	1.0	1.0
<i>Metabolism parameters</i>							
V _{maxC}	Maximum CYP metabolic rate (mg/hr/kg ^{0.7})	11.1	2	9.27	0.21	11.1	11.1
K _m	CYP affinity (mg/L)	0.396	2	0.574	0.42	0.396	0.396
k _{fC}	First-order GST metabolic rate constant (kg ^{0.3} /hr)	1.46	2	1.41	0.28	1.46	1.46
A1	Ratio of lung V _{maxC} to liver V _{maxC}	0.462	0.55	0.207	0.36	0.416	0.416
A2	Ratio of lung k _{fC} to liver k _{fC}	0.322	0.55	0.196	0.37	0.137	0.137

^aMCMC analysis was used to update prior distributions (means and CVs) for flow rate and metabolic parameters in a sequential process with three sets of kinetic data from mouse studies, as explained further in the text. Final values for posterior distributions are given in this table.

^bSource: Andersen et al. (1991, 1987).

^cSource: Clewell et al. (1993).

^dBased on a mouse breathing rate of 0.043 m³/d.

^eBased on a mouse breathing rate of 0.084 m³/d.

The Bayesian calibration of the cardiac output (QCC) constant, ventilation:perfusion ratio (VPR), and metabolic parameters was divided into three sequential steps: using kinetic data from closed chamber studies with mice treated with an inhibitor of CYP2E1 (trans-1,2-dichloroethylene) in order to minimize the oxidative pathway and enable a more precise estimate of parameters for the GST pathway, followed by kinetic data for mice given intravenous injections of dichloromethane to estimate metabolism parameters in the absence of pulmonary absorption processes and, finally, kinetic data for naïve mice exposed to dichloromethane in closed chambers (Marino et al., 2006). The initial prior distributions were based on mean values used by Andersen et al. (1987) for the metabolic parameters and by OSHA (1997) for the parameters for VPR, A1, and A2. Posterior distributions from the first Bayesian analysis were used as prior distributions for the second step, and posterior distributions from the second step were used as prior distributions for the final updating. Final results from the Bayesian calibration of the mouse probabilistic model are shown in Table 3-5.

Marino et al. (2006) used the Bayesian-calibrated mouse model to calculate internal dose metrics associated with exposure conditions in the NTP (1986) B6C3F₁ mouse cancer inhalation bioassay. The internal dose metric selected was mg dichloromethane metabolized by the GST pathway per liter tissue per day. This is the same dose metric used in earlier applications of PBPK models to derive human cancer IUR estimates based on cancer responses in mice (OSHA, 1997; Andersen et al., 1987; U.S. EPA, 1987a, b). Its use is consistent with evidence that dichloromethane metabolism via GST-T1 results in the formation of a reactive metabolite that damages DNA and results in the formation of tumors (see Section 4.7). The model was used to calculate values for this internal dose metric in the lung and liver of mice in the NTP (1986) study, using the mean values of the final distributions for the parameters in the model. Resultant values were three- to fourfold higher than values calculated with the Andersen et al. (1987) and U.S. EPA (1987a, b) versions of the model (Table 3-6). Marino et al. (2006) noted that the difference could be primarily attributed to the changes in the partition coefficients based on Clewell et al. (1993) as well as to the Bayesian updating of the metabolic parameters (see Table 3-5).

Table 3-6. Internal daily doses for B6C3F₁ mice exposed to dichloromethane for 2 years (6 hours/day, 5 days/week) calculated with different PBPK models

Target organ	NTP (1986) exposure level ^a	PBPK model		
		Marino et al. (2006)	EPA (1987a, b)	Andersen et al. (1987)
Liver ^b	Control	0	0	0
	2,000 ppm	2,359.99	727.8	851
	4,000 ppm	4,869.85	1,670	1,811
Lung ^b	Control	0	0	0
	2,000 ppm	474.991	111.4	123
	4,000 ppm	973.343	243.7	256

^a2,000 ppm = 6,947 mg/m³; 4,000 ppm = 13,894 mg/m³.

^bInternal dose expressed as mg dichloromethane metabolized by the GST pathway per liter tissue per d.

Marino et al. (2006) noted that inclusion of extrahepatic CYP metabolism in the slowly perfused tissue compartment in the mouse model had little impact on the formation of GST metabolites in the liver and lung, especially at exposure levels used in the mouse NTP (1986) bioassay. To support this contention, the Andersen et al. (1987) model was modified to include 10% of the liver rate of oxidative metabolism in the slowly perfused tissue compartment (as suggested by Sweeney et al. [2004]), and the modified model was used to calculate the formation of GST metabolites. If extrahepatic metabolism was included in the slowly perfused tissue compartment, there was a 5–6% reduction in the formation of GST metabolites in the lung and liver at an exposure level of 50 ppm. At 2,000 or 4,000 ppm, however, there was only a 0.77 or 0.37% reduction, respectively. Marino et al. (2006) did not discuss the impact of including extrahepatic metabolism in the rapidly perfused tissue compartment; the same group of investigators developed a human PBPK model that included CYP metabolism in the richly perfused compartment (David et al., 2006).

3.5.2. Probabilistic Human PBPK Dichloromethane Model (David et al., 2006)

The basic model structure used by David et al. (2006) was that of Andersen et al. (1987) with the addition of the CO submodel of Andersen et al. (1991), refinements from the Marino et al. (2006) mouse model, and an inclusion of CYP metabolism in richly perfused tissue (Figure 3-4). David et al. (2006) used Bayesian analysis to develop and calibrate metabolic parameters in a human probabilistic PBPK model for dichloromethane, using kinetic data from several studies of volunteers exposed to dichloromethane (n = 13 from DiVincenzo and Kaplan [1981]; n = 12 from Engström and Bjurström [1977]; n = 14 from Astrand et al. [1975]; n = 3 from Stewart et al. [1972a], and group means for metabolism parameters from Andersen et al. [1991]). Exhaled dichloromethane and CO and blood levels of dichloromethane and COHb were available in the studies by Andersen et al. (1991) and DiVincenzo and Kaplan (1981). The other three studies included two or three of these measures. The only available data for levels of

dichloromethane in fat came from the study of Engström and Bjurström (1977) (described in Section 3.2 within adipose tissue).




Figure 3-4. Schematic of human PBPK used by David et al. (2006).

Values (means and SDs or CVs) for the model parameter distributions were selected from multiple sources considered to provide the most current scientific evidence for each parameter (David et al., 2006). Mean values for QCC, VPR, and all fractional tissue volumes and blood flow rates were based on mean values used by U.S. EPA (2000d) in a PBPK model for vinyl chloride, as were values for CVs for all physiological parameters, except CVs for VPR and fractional lung volume, which were set to those used by OSHA (1997). Means for the CO submodel parameters were set equal to those in Andersen et al. (1991), except for those for the endogenous rate of CO production (REnCO) and the background amount of CO (ABCOC), which were based on data collected by DiVincenzo and Kaplan (1981). Means for partition coefficients, the A1 ratio and the A2 ratio were those used by Andersen et al. (1987), whereas prior means for $V_{\max C}$ and K_m were those used by Andersen et al. (1991). The prior mean for the metabolic parameter for CYP metabolism in the rapidly perfused tissue was set at 0.03, slightly lower than the value suggested by Sweeney et al. (2004). Prior CVs for the metabolic parameters were set at 200%.

MCMC analysis was used to calibrate metabolic parameters in the human model in a two-step approach: (1) posterior distributions were estimated separately by using data from each of the five studies with kinetic data for humans exposed to dichloromethane (with durations ranging from 1 to 8 hours and concentrations ranging from 50 to 1,000 ppm), and (2) posterior distributions were estimated with combined data from the 42 individual subjects from the four studies with individual subject data (DiVincenzo and Kaplan, 1981; Engström and Bjurström, 1977; Astrand et al., 1975; Stewart et al., 1972a). Estimates of the population mean values for the fitted parameters from the Bayesian calibration with the combined kinetic data for individual subjects are shown in Table 3-7. This analysis resulted in a narrowing of the distribution for the

CYP2E1 metabolism parameters $V_{\max C}$ and K_m from a fairly broad prior distribution with a CV of 200% for both parameters to 13.1 and 33.6%, respectively, for $V_{\max C}$ and K_m . It should be noted that the CV values only represent the uncertainties in the corresponding population mean values and do not include the estimated interindividual variability (Harvey Clewell, to Paul Schlosser, U.S. EPA, email dated October 14, 2009). Thus, that narrowing should only be interpreted as indicating a high degree of confidence in the population mean. As will be discussed in detail later, other data which better characterize the variability in CYP2E1 activity among the human population should then be used in conjunction with these uncertainties to characterize the full range of uncertainty and variability.

Table 3-7. Results of calibrating metabolic parameters in a human probabilistic PBPK model for dichloromethane with individual kinetic data for 42 exposed volunteers and MCMC analysis

Parameter	Prior distributions		Posterior distributions	
	Mean (arithmetic)	CV	Mean (arithmetic)	CV
$V_{\max C}$ —maximal CYP metabolic rate (mg/hr/kg ^{0.7})	6.25	2	9.42	0.131
K_m —CYP affinity (mg/L)	0.75	2	0.433	0.336
k_{fC} —first-order GST metabolic rate (kg ^{0.3} /hr)	2	2	0.852	0.711
A1—ratio of lung $V_{\max C}$ to liver $V_{\max C}$	0.00143	2	0.000993	0.399
A2—ratio of lung k_{fC} to liver k_{fC}	0.0473	2	0.0102	0.728
FracR—fraction of $V_{\max C}$ in rapidly perfused tissues	0.03	2	0.0193	0.786

Source: David et al. (2006).

A component of quantitative uncertainty arises in examining the results of David et al. (2006), specifically for the GST metabolic parameter, k_{fC} . The authors reported Bayesian posterior statistics for the population mean parameters when calibration was performed either with specific published data sets or the entire combined data set. While one would generally expect that the values obtained from the combined data set should be a weighted average of the values from individual data sets, the population mean for the liver GST activity (coefficient), k_{fC} , was 0.852 from the combined data set while the values from the individual data sets ranged from 1.92 to 34.0 kg^{0.3}/hour.

A clarification provided by Marino (Dale Marino, to Glinda Cooper, U.S. EPA, email dated April 25, 2007) is that the parameter bounds stated in the text of David et al. (2006) were only applied for the analysis of the DiVincenzo and Kaplan (1981) and the combined data set. But according to the text and distribution prior statistics specified, the upper bound for k_{fC} would have been 12 kg^{0.3}/hour (mean + 2.5 SDs, with mean = 2 and SD = mean × CV = 2 × 2 = 4). The data of Andersen et al. (1991) were not used in the combined analysis because only group average values were available from that source, rather than individual data. Since the remaining

study-specific mean k_{fC} values were 7.95, 5.87, 34.0, and 1.92, with CVs of <2 , it seems unlikely that application of this upper bound would result in a value of $k_{fC} = 0.852 \text{ kg}^{0.3}/\text{hour}$. Given that there had been convergence problems with the combined data set when parameter values were unbounded, it is possible that convergence had not actually been reached after parameter bounds were introduced, and a higher value for k_{fC} would have been obtained had the chain been continued longer. The implications of this parameterization uncertainty are discussed further in Section 5.3 for noncancer toxicity modeling and Section 5.4.5 for cancer dose-response modeling.

Setting this uncertainty aside, since the parameter statistics shown in Table 3-7 (values reported by David et al., 2006) represent population means and the level of uncertainty in those means, their correct interpretation requires further consideration. As noted above, there is a known range of variability in CYP2E1 expression among the human population which should be incorporated when estimating overall population variability. For the remaining parameters except k_{fC} (i.e., for K_m , A1, A2, and FracR), there is not a known equivalent level of variability and it will be assumed that there is, in fact, a single true value for the population which is estimated as the population mean by David et al. (2006). In that case one needs only to include the uncertainty in the mean represented by the CV values in Table 3-7 in a statistical (Monte Carlo) sampling in order to fully characterize model uncertainty and variability. While the analysis of David et al. (2006) may have included variability among individual-specific estimates for those parameters, this treatment effectively assumes that this variability was only an apparent artifact of the limited data, measurement noise, etc.

David et al. (2006) further refined the human probabilistic model to reflect polymorphisms in the GST pathway: homozygous positive (+/+) GST-T1, heterozygous (+/-) GST-T1, and homozygous negative (-/-) GST-T1 individuals with no GST activity. Distributions of GST activities for these genotypes in a group of 208 healthy male and female subjects from Sweden were scaled to obtain distributions of k_{fC} for each genotype (Warholm et al., 1994). When weighted by estimated frequencies of the genotypes in the U.S. population and appropriately scaled, these genotype-specific activity distributions would result in an overall population mean equal to the k_{fC} mean for the posterior distribution shown in Table 3-7 ($0.852 \text{ kg}^{0.3}/\text{hour}$). The resultant mean k_{fC} values were $0.676 \text{ kg}^{0.3}/\text{hour}$ (SD 0.123) for heterozygous individuals and $1.31 \text{ kg}^{0.3}/\text{hour}$ (SD 0.167) for homozygous positive individuals, as indicated in Table 2 of David et al. (2006). The final parameter distributions used by David et al. (2006) are summarized in Table 3-8.

Table 3-8. Parameter distributions used in human Monte Carlo analysis for dichloromethane by David et al. (2006)

Parameter		Distribution		Source
		Mean (arithmetic)	SD	
BW	Body weight (kg)	70.0	21.0	Humans ^a
QCC	Cardiac output (L/hr/kg ^{0.74})	16.5	1.49	Humans ^a
VPR	Ventilation:perfusion ratio	1.45	0.203	Humans ^a
QFC	Fat	0.05	0.0150	Humans ^a
QLC	Liver	0.26	0.0910	Humans ^a
QRC	Rapidly perfused tissues	0.50	0.10	Humans ^a
QSC	Slow perfused tissues	0.19	0.0285	Humans ^a
Tissue volumes (fraction BW)				
VFC	Fat	0.19	0.0570	Humans ^a
VLC	Liver	0.026	0.00130	Humans ^a
VLuC	Lung	0.0115	0.00161	Humans ^a
VRC	Rapidly perfused tissues	0.064	0.00640	Humans ^a
VSC	Slowly perfused tissues (muscle)	0.63	0.189	Humans ^a
Partition coefficients				
PB	Blood:air	9.7	0.970	Humans ^b
PF	Fat:blood	12.4	3.72	Rats ^b
PL	Liver:blood	1.46	0.292	Rats ^b
PLu	Lung:arterial blood	1.46	0.292	Rats ^b
PR	Rapidly perfused tissue:blood	1.46	0.292	Rats ^b
PS	Slowly perfused tissue (muscle:blood)	0.82	0.164	Rats ^b
Metabolism parameters				
V _{maxC}	Maximum metabolism rate (mg/hr/kg ^{0.7})	9.42	1.23	Calibration ^c
K _m	Affinity (mg/L)	0.433	0.146	Calibration ^c
A1	Ratio of lung V _{max} to liver V _{max}	0.000993	0.000396	Calibration ^c
A2	Ratio of lung KF to liver KF	0.0102	0.00739	Calibration ^c
FracR	Fractional CYP2E1 capacity in rapidly perfused tissue	0.0193	0.0152	Calibration ^c
First order metabolism rate (/hr/kg ^{0.3})				
k _{fC}	Homozygous (-/-)	0	0	Hybrid ^d
	Heterozygous (+/-)	0.676	0.123	Hybrid ^d
	Homozygous (+/+)	1.31	0.167	Hybrid ^d

SD = standard deviation.

^aU.S. EPA, 2000d. Human PBPK model used for vinyl chloride.

^bAndersen et al. (1987). Blood:air partition measured using human samples; other partition coefficients based on estimates from tissue measures in rats.

^cBayesian calibration based on five data sets (see text for description); posterior distributions presented in this table.

^dThe overall population mean for k_{fC} as determined by Bayesian calibration; the distribution of activity among the three genotypes and variability in activity for each genotype (SD values) were then scaled from the ex vivo data of Warholm et al. (1994).

Source: David et al. (2006).

Since the measurements of GST-T1 activity by Warholm et al. (1994) were performed *ex vivo* using blood samples, it is reasonable to assume that those measurements are accurate and, hence, that the characterization of the distributions for each genotype as being normal with the reported level of variance represents the true level of human variability for each genotype. However that characterization does not include the uncertainty in the overall population mean for the rate of hepatic GST activity towards dichloromethane, indicated by the CV for k_{fC} in Table 3-7 (from Table 4 of David et al. [2006]). Thus, to fully account for both the population variability and parameter uncertainty, a Monte Carlo statistical sampling should first sample the population mean from a distribution with mean = 0.852 kg^{0.3}/hour and CV = 0.711 (thus accounting for uncertainty) and then reweight the population-specific distributions listed in Table 3-8 to have that sampled population mean, before selecting (sampling) an individual value of k_{fC} from those weighted distributions (thus accounting for variability). EPA incorporated this change in the PBPK modeling used in this assessment.

As described in detail in Appendix B, EPA evaluated the adequacy of all the parameter distributions used by David et al. (2006) to characterize variability among the full human (U.S.) population. EPA's conclusion is that the reported distributions for many of the physiological parameters in particular, as well as V_{maxC} (CYP2E1) as described above, only represented a narrow set of adults and did not represent the full range of variability. EPA therefore chose to use supplemental data sources to define these distributions in a way that should fully characterize the variability in the human population for individuals between 6 months and 80 years of age. Specifically, while the BW distribution used in the David et al. (2006) PBPK model used ranges from 7 to 130 kg, thus covering 6-month-old children to obese adults, there are age-dependent changes and gender-dependent differences in ventilation rates and body fat that are not explicitly included. To more accurately reflect the distribution of physiological parameters in the entire population, EPA replaced the unstructured distributions of David et al. (2006) with distributions based on available information that specifically account for population variability in age, gender, and age- and gender-specific distributions or functions for BW, QCC, alveolar ventilation, body fat (fraction), and liver fraction (see Appendix B for more details of the evaluation of each of these parameters).

For V_{maxC} (CYP2E1), EPA also incorporated additional data for the variability in CYP2E1 activity among humans based on Lipscomb et al. (2003). The Lipscomb et al. (2003) study used *in vitro* analysis of liver samples from 75 human tissue donors (activity towards trichloroethylene and measurements of protein content) to estimate a distribution of activity in the population. These data support a wider distribution in CYP2E1 activity than had been used in the David et al. (2006) dose-metric and unit-risk calculations, with approximately a sixfold range observed for CYP2E1 in Lipscomb et al. (2003) and a twofold range used by David et al. (2006). After sampling the population mean for CYP2E1 from the distribution indicated by the parameters in Table 3-8 to capture the uncertainty in the population mean, EPA assumed a log-

normal distribution for human variability around that mean and sampled the individual value using that population mean with geometric standard deviation (GSD) = 1.73. Further, since even the data available to Lipscomb et al. (2003) were limited, and the log-normal distribution is naturally bounded to be greater than zero, EPA chose to use a nontruncated log-normal distribution in the second (variability) sampling step for this parameter. (In sampling the population-mean value for $V_{\max C}$ from its range of uncertainty, EPA did truncate the distribution as indicated by David et al. [2006], since that mean is expected to be bounded away from zero.) Finally, the scaling of CYP2E1 for individuals under the age of 18 was adjusted based on the data of Johnsrud et al. (2003); EPA's analysis of these data indicate CYP2E1 activity in children is better predicted when assumed to scale with BW raised to the 0.88 power, as compared to the more general power of 0.74, used by David et al. (2006). CYP2E1 activity for individuals over the age of 18 is still assumed to scale as $BW^{0.74}$.

The resulting set of parameter distribution characteristics, including those used as defined by David et al. (2006) are described in Table 3-9. Using this revised set of distributions, including the CYP and GST activity distributions, and other distributions used as defined by David et al. (2006) or revised by EPA, the model as applied should reflect the full variability in the (U.S.) human population.

Table 3-9. Parameter distributions for the human PBPK model for dichloromethane used by EPA

Parameter		Distribution					Section or source
		Shape	(Geometric) mean ^a	SD/GSD ^a	Lower bound	Upper bound	
BW	Body weight (kg)	Normal	$f(\text{age}, \text{gender})$		1 st %tile	99 th %tile	B.4.3; NHANES IV
Flow rates							
QAlvC	Alveolar ventilation (L/hr/kg ^{0.75})	Normal	$f(\text{age}, \text{gender})$	$f(\text{age})$	5 th %tile	95 th %tile	B.4.4; mean: Clewell et al. (2004); SD: Arcus-Arth and Blaisdell (2007)
vprv	Variability in ventilation:perfusion ratio	Log-normal	1.00	0.203	0.69	1.42	VPR/VPR _{mean} of David et al. (2006)
QCC	Cardiac output (L/hr/kg ^{0.75})	QCC _{mean} = $f(\text{QAlvC})$		QCC = QCC _{mean} /vprv			B.4.5; Clewell et al. (2004) (mean)
Fractional flow rates (fraction of QCC)							
QFC	Fat	Normal	0.05	0.0150	0.0050	0.0950	David et al. (2006); <i>after sampling from these distributions, normalize:</i> $Q_i = \frac{QC \cdot Q_iC}{\sum Q_jC}$
QLC	Liver	Normal	0.26	0.0910	0.010	0.533	
QRC	Rapidly perfused tissues	Normal	0.50	0.10	0.20	0.80	
QSC	Slow perfused tissues	Normal	0.19	0.0285	0.105	0.276	
Tissue volumes (fraction BW)							
VFC	Fat	Normal	$f(\text{age}, \text{gender})$	0.3·mean	0.1·mean	1.9·mean	Fat mean: B.4.6 (Clewell et al., 2004); liver mean: B.4.7 (Clewell et al., 2004); otherwise, David et al. (2006); <i>after sampling from these distributions, normalize:</i> $V_i = \frac{0.9215 \cdot BW \cdot V_iC}{\sum V_jC}$
VLC	Liver	Normal	$f(\text{age})$	0.05·mean	0.85·mean	1.15·mean	
VLuC	Lung	Normal	0.0115	0.00161	0.00667	0.0163	
VRC	Rapidly perfused tissues	Normal	0.064	0.00640	0.0448	0.0832	
VSC	Slowly perfused tissues	Normal	0.63	0.189	0.431	0.829	
Partition coefficients							
PB	Blood:air	Log-normal	9.7	1.1	7.16	13.0	Geometric mean (GM) & GSD values listed here, converted from arithmetic mean and SD values of David et al. (2006)
PF	Fat:blood	Log-normal	11.9	1.34	4.92	28.7	
PL, PLu, & PR	Liver:blood, lung:arterial blood, and rapidly perfused tissue:blood	Log-normal	1.43	1.22	0.790	2.59	
PS	Slowly perfused tissue (muscle):blood	Log-normal	0.80	1.22	0.444	1.46	

(Table 3-9 continues on next page)

Table 3-9. Parameter distributions for the human PBPK model for dichloromethane used by EPA

Parameter		Distribution					Section or source
		Shape	(Geometric) mean ^a	SD/GSD ^a	Lower bound	Upper bound	
Metabolism parameters (based on Monte Carte calibration from five human data sets)							
$V_{\max C_{\text{mean}}}$ / $V_{\max C}$	Population mean / individual maximum metabolism rate (mg/hr/kg ^{X_{vmax}})	Log-normal	9.34	1.14	6.96	11.88	B.3 mean: David et al. (2006); Individual GSD: Lipscomb et al. (2003); X _{vmax} = 0.88 for age <18; X _{vmax} = 0.70 for age ≥18
		Log-normal	$V_{\max C_{\text{mean}}}$	1.73	(none)	(none)	
K_m	Affinity (mg/L)	Log-normal	0.41	1.39	0.154	1.10	Geometric mean (GM) & GSD values listed here, converted from arithmetic mean and SD values of David et al. (2006)
A1	Ratio of lung V_{\max} to liver V_{\max}	Log-normal	0.00092	1.47	0.000291	0.00292	
A2	Ratio of lung KF to liver KF	Log-normal	0.0083	1.92	0.00116	0.0580	
FracR	Fractional MFO capacity in rapidly perfused tissue	Log-normal	0.0152	2.0	0.00190	0.122	
First order metabolism rate ($[\text{hr}/\text{kg}^{0.3}]^{-1}$)							
$k_{fC_{\text{mean}}}$	Population average	Log-normal	0.6944	1.896	0.1932	2.496	Adapted from David et al. (2006); $k_{fC_{\text{mean}}}$ is first sampled, then the relative individual value, $k_{fC}/k_{fC_{\text{mean}}}$, given the genotype; k_{fC} is then the product
$k_f C / k_f C_{\text{mean}}$	Homozygous (-/-)	Normal	0	0	0	0	
	Heterozygous (+/-)	Normal	0.8929	0.1622	0	1.704	
	Homozygous (+/+)	Normal	1.786	0.2276	0	2.924	

^aArithmetic mean and SD listed for normal distributions; GM and GSD listed for log-normal distributions.

3.5.3. Evaluation of Rat PBPK Dichloromethane Models

Several deterministic PBPK rat models have been reported in the scientific literature (Sweeney et al., 2004; Andersen et al., 1991, 1987; Reitz, 1991; Reitz et al., 1988a, b; U.S. EPA 1988b, 1987a, b; Gargas et al., 1986). Unlike the mouse (Marino et al., 2006) and human (David et al., 2006), no hierarchical population model for dichloromethane in the rat exists in which parameter uncertainty is quantitatively integrated into model calibration. Rat data are not available that would allow for Bayesian calibration of individual metabolic parameters for the CYP or GST pathways. Thus, EPA assessed modified versions of deterministic rat PBPK models to select the most appropriate model for use in extrapolating internal dosimetry from rats to humans, for example in the determination of RfDs and RfCs based on effects seen in the rat. This work is described in detail in Appendix C and is based on evaluation of blood levels of dichloromethane, the percent saturation of hemoglobin as COHb (%COHb), and expired dichloromethane following intravenous injection (Angelo et al., 1986b) and closed chamber gas uptake (Gargas et al., 1986), as well as evaluation of dichloromethane and %COHb blood levels from a 4-hour inhalation exposure (Andersen et al., 1991, 1987). Based on this work, the basic model structure of Andersen et al. (1991) was chosen, with the inclusion of lung dichloromethane metabolism via CYP (4% of liver metabolite production) and GST (14% of liver metabolite production) pathways (estimated from Reitz et al., 1989) (Figure 3-5) with metabolic parameters recalibrated against data of Andersen et al. (1991), based on prediction agreement of the various parameters with the available rat data sets. Table 3-10 presents the parameter distribution data for this model.




Figure 3-5. Schematic of rat PBPK model used in current assessment.

Table 3-10. Parameter values for the rat PBPK model for dichloromethane used by EPA

Parameter	Mean
Flow rates	
QCC (L/hr/kg ^{0.74})	15.9
VPR	0.94
Fractional flow rates (percent of QCC)	
Fat	9
Liver	20
Rapidly perfused tissues	56
Slowly perfused tissues	15
Tissue volumes (percent BW)	
Fat	7
Liver	4
Lung (scaled as BW ^{0.99})	1.15
Rapidly perfused tissues	5
Slowly perfused tissues	75
Partition coefficients	
Blood:air	19.4
Fat:blood	6.19
Liver:blood	0.732
Lung:arterial blood	0.46
Rapidly perfused tissue:blood	0.732
Slowly perfused tissue (muscle):blood	0.408
Metabolism parameters	
Maximum metabolism rate (mg/hr/kg ^{0.7})	3.93
Affinity (mg/L)	0.524
Ratio of lung V _{max} to liver V _{max}	0.04
Ratio of lung KF to liver KF	0.14
1st order metabolism rate (liver KF) ([hr/kg ^{0.3}] ⁻¹)	2.46
First-order oral absorption rate constant, k _a (1/hr)	1.80

3.5.4. Comparison of Mouse, Rat, and Human PBPK Models

The comparison of various parameters across species (Table 3-11) primarily shows the modest interspecies differences that are known to occur in physiological parameters, also including the approximately twofold differences in partition coefficients which occur because of differences in rodent versus human blood lipid content. The 2.5-fold lower V_{maxC} (CYP activity) in rats versus mice is also typical. The most striking difference is the variation in A1 and A2. Those values, however, reflect the in vitro differences originally quantified by Lorenz et al. (1984) and used in the dichloromethane PBPK modeling of Andersen et al. (1987). Thus, these differences are based on independent measurements of tissue-specific metabolic capacity, and

while the specific values for mouse and human were refined through Bayesian analysis, the ultimate (posterior) values used are within a reasonable range of the in vitro measurements and so do not appear to be artifactual. (Since in vivo kinetics often indicate some differences from what would be predicted without adjustment from in vitro, it is not surprising that such differences occur here.) These differences do explain why lung-specific metrics in particular lead to lower internal dose and hence risk predictions in humans compared to whole-body metrics.

Table 3-11. Parameters in the mouse, rat, and human PBPK model for dichloromethane used by EPA

Parameter	Mouse ^a mean	Rat ^b value	Human ^c		
			Mean	CV/GSD (shape, bounds)	Sources
<i>Fractional flow rates (fraction of cardiac output)^b</i>					
QFC Fat	0.05	0.09	0.05	0.3 (N, 0.1–1.9)	David et al. (2006); then normalized: $Q_i = \frac{QC \cdot QiC}{\sum QjC}$
QLC Liver	0.24	0.20	0.26	0.35 (N, 0.0385–2.05)	
QRC Rapidly perfused tissues	0.52	0.56	0.50	0.2 (N, 0.4–1.6)	
QSC Slowly perfused tissues	0.19	0.15	0.19	0.15 (N, 0.553–1.453)	
<i>Fractional tissue volumes (fraction of BW)^b</i>					
VFC Fat	0.04	0.07	<i>f(age, gender)</i>	0.3 (N, 0.1–1.9)	Fat mean: §2.2.3.6; Liver mean: §2.2.3.7; otherwise David et al. (2006); then normalized: $V_i = \frac{0.9215 \cdot BW \cdot ViC}{\sum VjC}$
VLC Liver	0.04	0.04		<i>f(age)</i>	
VluC Lung	0.0115	0.0115	0.0115	0.14 (N, 0.58–1.42)	
VRC Rapidly perfused tissues	0.05	0.05	0.064	0.1 (N, 0.7–1.3)	
VSC Slowly perfused tissues	0.78	0.75	0.63	0.3 (N, 0.684–1.32)	
<i>Partition coefficients^c</i>					
PB Blood/air	23.0	19.4	9.7	1.1 (LN, 0.738–1.34)	Geometric mean (GM) & GSD/GM values converted from arithmetic mean & SDs of David et al. (2006)
PF Fat/blood	5.1	6.19	11.9	1.34 (LN, 0.413–2.41)	
PL Liver/blood	1.6	0.73	1.43	1.22 (LN, 0.552–1.81)	
PLu Lung/blood	0.46	0.46	1.43	"	
PR Rapidly perfused/blood	0.52	0.73	1.43	"	
PS Slowly perfused/blood	0.44	0.41	0.80	1.22 (LN, 0.555–1.83)	
<i>Flow rates</i>					
QCC Cardiac output (L/hr/kg ^{0.74})	24.2	14.99	QCC _{mean} = <i>f</i> (QAlvC) (variable)	QCC = QCC _{mean} /vprv (varies) (LN, 0.69–1.42)	QCC: §2.2.3.5; vprv = VPR/VPR _{mean} ; David et al. (2006); QAlvC: §2.2.3.4;
VPR Ventilation/perfusion ratio	1.45	0.94			
QAlvC	QCC/VPR	QCC/VPR			

(Table 3-11 continues on next page)

Table 3-11. Parameters in the mouse, rat, and human PBPK model for dichloromethane used by EPA

Parameter	Mouse ^a mean	Rat ^b value	Human ^c		
			Mean	CV/GSD (shape, bounds)	Sources
<i>Metabolism parameters</i>					
V _{maxC} Maximum CYP metabolic rate (mg/hr/kg ^{Xvmax})	9.27	3.93	<i>LN(m = 9.34, s = 1.14 / lb = 6.96, ub = 11.88)</i>	1.73 (LN, [unbounded])	V _{maxC} : §2.2.2; others: David et al. (2006) (GM & GSD/GM values converted from arithmetic mean & SDs) k _{fC(mean)} : Combined data set posterior, Table 4 of David et al. (2006) k _{fC} / k _{fC(mean)} : rescaled from Table 2 of David et al. (2006)
Xvmax CYP allometric scaling power	0.7	0.7	0.88 for age <18; 0.7 for age ≥18)	1.39 (LN, 0.376–2.68)	
K _m CYP affinity (mg/L)	0.574	0.524	0.41		
k _{fC(mean)} First-order GST metabolic rate constant (kg ^{0.3} /hr)	1.41	2.46	<i>LN(m = 0.6944, s = 1.896 / lb = 0.1932, ub = 2.496)</i>		
k _{fC} /k _{fC(mean)}	1	1	0 (-/-) ^e	-/-: NA	
			0.8929 (+/-) ^e 1.7896 (+/+) ^e	+/-: 0.182 (N, 0–1.91) +/: 0.127 (N, 0–1.64)	
A1 Ratio of lung V _{maxC} to liver V _{maxC}	0.207	0.04	0.00092	1.47 (LN, 0.316–3.17)	
A2 Ratio of lung k _{fC} to liver k _{fC}	0.196	0.14	0.0083	1.92 (LN, 0.140–6.99)	

^aBased on Marino et al. (2006) (source for all mouse parameters).

^bBased on Andersen et al. (1991), with the addition of lung metabolism of dichloromethane via the CYP (4% of liver metabolite production) and GST (14% of liver metabolite production) pathways. Physiological parameters and partition coefficients are from Andersen et al. (1991). The values for dichloromethane metabolism in the lung (as a fractional yield of liver metabolism for each pathway) were estimated from the in vitro ratios of enzyme activity (nmol/min/mg protein) in lung and liver cytosolic (GST) and microsomal (CYP) tissue fractions (Reitz et al., 1989). Metabolic parameters were re-optimized against the inhalation data of Andersen et al. (1991) using a heteroscedasticity parameter value of 2, which uses relative error for the model fitting algorithm. See Appendix C for further details.

^cBased on David et al. (2006), with changes as noted. Additional sources include Clewell et al. (2004), Arcus-Arth and Blaisdell (2007), and Lipscomb et al. (2003). See identified sections for details. Distribution values (mean and a measure of dispersion) are provided with the CV (mean/SD) presented for normal (N) distributions and the GSD (italicized) presented for log-normal (LN) distributions. Distributions were truncated, bounds are (upper-lower bound)/mean.

^eValues for the homozygous (-/-), heterozygous (+/-), and homozygous (+/+) GST-T1 genotypes, respectively.

3.5.5. Uncertainties in PBPK Model Structure for the Mouse, Rat and Human

There is uncertainty in the dichloromethane PBPK modeling because there are parts of the entire data set for which the existing model and parameters fit very poorly and where the discrepancies appear to be structural (i.e., cannot be resolved simply by re-fitting model parameters unless fits to other data are degraded). The data used for model parameter estimation are primarily measurements of parent dichloromethane kinetics (e.g., blood or closed-chamber air concentrations over time), rather than measurements of metabolite levels which can be unambiguously attributed to one of the two principal metabolic pathways (GST and CYP). For the mouse model in particular, only parent dichloromethane data were used, though exhaled amounts of CO₂ and CO are available. Because only dichloromethane measurements are used, estimation of the fraction of dichloromethane metabolized by the GST vs. CYP pathway depends strongly on the assumed equations describing those rates and how the pharmacokinetic data are interpreted. Specifically, if there is some degree of saturation in the GST kinetics or the CYP kinetics are not accurately described by Michaelis-Menten kinetics (see next paragraph), then the estimation of the fraction of metabolism going down each pathway would shift. Further, while Marino et al. (2006) used data from mice pretreated with *trans*-1,2-dichloroethylene (tDCE), a specific CYP2E1 inhibitor (mice were exposed to 100 ppm tDCE for 1.5 h prior to dichloromethane exposure), the authors assumed without verification that 100% of the CYP2E1 activity was eliminated by the inhibitor when using those data. In contrast, Mathews et al. (1997) found that pretreatment of F344 rats by tDCE (100 mg/kg intraperitoneally) only yielded 65% inhibition of CYP2E1. If a significant fraction of the CYP2E1 activity was not eliminated in the dichloromethane experiments, then that activity is erroneously assigned to the GST pathway in the parameter estimation of Marino et al. (2006).

In addition to the possibility of incomplete inhibition of CYP2E1 affecting the data interpretation, the Michaelis-Menten rate equation used in all of the published PBPK models for dichloromethane, including that of Marino et al. (2006), has in fact not been shown to accurately describe the CYP2E1-mediated metabolism of dichloromethane in the relevant concentration range. While the Michaelis-Menten equation usually describes CYP-mediated oxidation data quite well, if there is some departure of the actual kinetics from this equation and incomplete inhibition from the tDCE treatment, the model parameters obtained under those assumptions would be compromised. If pathway-specific metabolite data were used to define or bound the ratio of GST to CYP metabolism, the resulting estimates would be less sensitive to errors in the CYP rate equation.

EPA compared model predictions of total CYP metabolism in mice, which should match with CO elimination at 24 hours after bolus exposures to dichloromethane as measured by Angelo et al. (1986a), since only the CYP pathway produces CO. At 50 mg/kg (in water), the model predicts that 10% of the dichloromethane is metabolized by the CYP pathway, which agrees with the observed values (11–12%; Angelo et al., 1986a). However, at 500 and 1,000


mg/kg, the model predicts that only 1.8 and 1.0% will be metabolized by the CYP pathway, while Angelo et al. (1986a) observed 9–10 and 3.6–7%, respectively. Thus, the extent of CYP saturation predicted by the model does not match with these pathway-specific metabolite data. However, as shown in Appendix C, the rat model is able to predict total exhaled CO quite well, indicating an error in the fraction of metabolism via the GST pathway of less than 13% in that species. That the *mouse* model does not describe well the dose-dependent shift in metabolism shown by those CO data suggests that the dose-dependence of the CYP Michaelis-Menten rate-equation may not be adequate. As will be shown, an alternative equation for CYP kinetics may fit the existing dichloromethane data better than Michaelis-Menten kinetics, with the result that a higher portion of total dichloromethane metabolism would be interpreted as being CYP-mediated. Thus, there is some uncertainty in the choice of equation for the CYP pathway, which leads to some uncertainty in the estimated GST:CYP metabolic ratio, upon which current risk predictions are based. However the extent of the error appears quite limited in the rat and more predominant at high exposures vs. low exposures in the mouse.

The potential error in assuming Michaelis-Menten kinetics for CYP-mediated oxidation of dichloromethane is reinforced by examining the *in vitro* oxidative (i.e., CYP-specific) kinetics of dichloromethane reported by Reitz et al. (1989). When extrapolated from *in vitro* to *in vivo*, the apparent values of the oxidative saturation constant, K_m , identified by Reitz et al. (1989) for mice, rats, and humans are over 2 orders of magnitude greater than those obtained *in vivo* with the PBPK model. This apparent discrepancy is partly explained by the disparate concentration ranges investigated: Reitz et al. (1989) used much higher dichloromethane concentrations *in vitro* than those observed in or predicted for the various *in vivo* pharmacokinetic studies. In particular, the oxidation of dichloromethane could involve two oxidative processes, one with a high affinity (low K_m) corresponding to the nonlinearity observed *in vivo* and one with a low affinity (high K_m) corresponding to the nonlinearity observed *in vitro*. Such a low-affinity process might account for the higher CO production observed *in vivo* (see above) than predicted by the current model. Further, a low-affinity process would have nearly linear kinetics in the exposure range used for the *in vivo* dosimetry studies and hence would be difficult to distinguish from GST-mediated metabolism unless pathway-specific metabolite data are used. If this second oxidative process is not inhibited by tDCE, then it may correspond to the 35% of oxidative metabolism which was observed to remain in rats after tDCE treatment by Mathews et al. (1997).

The data of Reitz et al. (1989) could simply indicate a second CYP with low-affinity dichloromethane activity. However that possibility is contradicted by the results of Kim and Kim (1996) who observed that another CYP2E1-specific inhibitor, disulfiram, completely abolished dichloromethane-induced increases on COHb in rats. Another possible explanation which would support the findings observed in Kim and Kim (1996) as well as Reitz et al. (1989) and the various *in vivo* data is that a number of CYPs exhibit “atypical” kinetics, not described by the classic Michaelis-Menten equation, consistent with the enzymes having dual binding sites

as proposed by Korzekwa et al. (1998). (Korzekwa et al. [1998] demonstrated atypical kinetics for several CYP-isozyme/substrate pairs, but not specifically for CYP2E1.) The application of this alternate kinetic model to dichloromethane dosimetry in mice has been explored by Evans and Caldwell (2010), who demonstrate that all the dichloromethane gas-uptake data in mice can be explained with this model in the hypothetical case where the GST pathway is not included. The alternate PBPK model of Evans and Caldwell (2010) is not considered further here because GST-mediated metabolism of dichloromethane clearly occurs in mice, rats, and humans based on the in-vitro observations of Reitz et al. (1989) and is mechanistically linked to dichloromethane-induced cancer as discussed in Section 4.5. Thus a model which excludes GST-mediated metabolism is not consistent with the overall database concerning dichloromethane metabolism and carcinogenesis research.

Figure 3-6 shows kinetic model fits to the in vitro mouse dichloromethane oxidation kinetic data of Reitz et al. (1989), after expressing those data on a per gram of liver basis. Both the standard Michaelis-Menten kinetic equation (solid line) and the dual-binding equation (dashed line) given by Korzekwa et al. (1998) are shown. In particular, the high-affinity (low) K_m for the dual-binding equation was set equal to that obtained by Marino et al. (2006) from their PBPK modeling. This figure shows that the dual binding model is not only consistent with the apparent high-affinity saturation obtained from in vivo PBPK modeling (K_m of Marino et al. [2006]), but also with the apparent low-affinity (high K_m) data of Reitz et al. (1989), and describes those in vitro data better than the standard Michaelis-Menten equation. Reitz et al. (1989) used classic Lineweaver-Burk plots to display their kinetic data; i.e., $1/\text{reaction rate}$ vs. $1/\text{concentration}$. The systematic discrepancy between their data and Michaelis-Menten kinetics evident in Figure 3-6 is much less obvious with that scaling, which likely explains why they made no note of it.



Dichloromethane oxidation data obtained with mouse liver microsomes by Reitz et al. (1989) (points), expressed on a per gram of liver basis, are shown with a fitted Michaelis-Menten equation (solid line) or a fitted dual-binding-site equation as described by Korzekwa et al. (1998) (dashed line), where the high affinity saturation constant of the dual-binding-site equation set equal to the mean K_m determined for mice via PBPK modeling by Marino et al. (2006). The K_m for the Michaelis-Menten equation (108 mg/L) is inconsistent with the in vivo dichloromethane dosimetry data, while the in vitro data shown here are inconsistent with the K_m estimated in vivo (0.42 mg/L) if that equation is used.

Figure 3-6. Comparison of dichloromethane oxidation rate data with alternate kinetic models.

In summary regarding model equations, the current PBPK model used the standard Michaelis-Menten equation to describe CYP2E1-catalyzed oxidation of small volatile organic compounds. Analysis of the dichloromethane (pharmaco)kinetic data and evaluation of the inconsistencies described above suggest that an alternate equation, which would impact risk predictions, may better represent CYP2E1-induced oxidation of dichloromethane. The analysis provided here demonstrates shortcomings in the existing model which the alternate model may address, indicating that this is a substantial model uncertainty. However, the hypothesis that CYP2E1 kinetics for dichloromethane should be described by this alternate rate equation requires further laboratory testing. For example, dichloromethane oxidation in a bacterial expression system where only CYP2E1 is expressed could be measured over a concentration range sufficient to firmly distinguish between the two kinetic forms indicated in the figure above. Such experiments would clearly show that the metabolic kinetics are due to atypical kinetics occurring with a single enzyme (CYP2E1), vs. involvement of a second, low-affinity enzyme. Also, the alternate equation would need to be incorporated into a PBPK model which also

included the GST pathway, and the resulting model calibrated not only for the mouse, but also the rat and the human. Until such additional experiments and modeling are available, the existing PBPK model remains the best available science for dose- and risk-extrapolation from rodents to humans despite this uncertainty. Analysis of the GST-mediated metabolism of dichloromethane measured by Reitz et al. (1989) shows that those results are within a factor of three of the GST kinetic parameters used in the current PBPK model, indicating that any error in the GST:CYP balance is in that range.

4. HAZARD IDENTIFICATION

4.1. STUDIES IN HUMANS

4.1.1. Introduction—Case Reports, Epidemiologic, and Clinical Studies

There has been considerable interest in the influence of occupational exposure to dichloromethane in relation to a variety of conditions. The recognition that dichloromethane can be metabolized and bound to hemoglobin to form COHb, resulting in a reduction in the oxygen carrying capacity of the blood (Stewart et al., 1972b), prompted investigations into risk of ischemic heart disease and other cardiovascular effects. Reports of neurological effects from acute, high-exposure situations contributed to concern about neurological effects of chronic exposure to lower levels of dichloromethane. A general interest in potential cancer risk became more focused on lung and liver cancer because of the observation of these specific tumors in the NTP (1986) experiments in mice. Details of the studies pertaining to the experimental and epidemiologic studies of noncancer outcomes (e.g., cardiac, neurologic, hepatic, reproductive) are presented in Section 4.1.2, and studies of cancer risk are presented in Section 4.1.3.

4.1.2. Noncancer Studies

4.1.2.1. *Case Reports of Acute, High-dose Exposures*

Numerous case reports have been published that describe health effects resulting from acute exposure to dichloromethane. Most of the reports describe health effects resulting from inhalation of dichloromethane or dermal contact, but a few involve ingestion. The COHb levels in some of these cases were relatively low (7.5–13%), so the initial toxic effects of acute dichloromethane exposure appear to be due to its anesthetic properties as opposed to metabolic conversion of dichloromethane to CO.

Bakinson and Jones (1985) reported on a series of 33 cases of acute inhalation exposures to dichloromethane that occurred in the workplace over the period 1961–1980. Thirteen had lost consciousness, and one of the workers died. Nineteen cases reported general neurological effects, 13 reported gastrointestinal symptoms, 4 reported respiratory symptoms, and 1 reported hepatic symptoms. Of the 19 with general neurological symptoms, all reported headache, and dizziness was reported by 11 workers. Five workers reported one of the following symptoms: drunkenness, confusion, lack of coordination, or paresthesia.

Rioux and Myers (1988) summarized the health effects reported for 26 cases of dichloromethane poisoning published in the literature between 1936 and 1986. Three cases resulted from abuse-related exposures, 2 from chronic exposures, and 21 from acute exposures. The most common effects involved the central nervous system (CNS) (unconsciousness, drowsiness, headache, and behavioral symptoms), pulmonary edema and dyspnea, and dermatologic symptoms. Even severe symptoms could be reversed, but four deaths occurred.

More than 10 other case reports of fatalities or poisonings have been published since the summaries by Rioux and Myers (1988) and Bakinson and Jones (1985), and many of these incidents involve inadequately ventilated occupational settings (Jacobovich et al., 2005; Raphael et al., 2002; Fechner et al., 2001; Zarrabeitia et al., 2001; Gouille et al., 1999; Mahmud and Kales, 1999; Kim et al., 1996; Tay et al., 1995; Manno et al., 1992; Leikin et al., 1990; Shusterman et al., 1990). CNS depression and resulting narcosis, respiratory failure, and heart failure are common features of these reports. In a survey of workers in furniture stripping shops, 10 of the 21 workers stated that they sometimes experienced dizziness, nausea, or headache during furniture stripping operations (Hall and Rumack, 1990).

Chang et al. (1999) reported details of six patients who had ingested dichloromethane (four in a suicide attempt and two from accidental ingestion during a state of intoxication). The estimated amounts ingested were ≤ 350 mL. COHb levels, which were measured in only two of the cases, were 8.4 and 35% (with the latter being seen in a fatal case). As in exposures resulting from inhalation, the most common symptoms involved CNS depression, ranging from somnolence and weakness to deep coma. Tachypnea ($n = 6$) and corrosive gastrointestinal tract injury ($n = 3$) were also reported. Hepatic and renal failure and pancreatitis were found in the two most severe cases.

4.1.2.2. *Controlled Experiments Examining Acute Effects*

Several controlled experiments were conducted in the 1970s examining neurophysiological effects and levels of COHb resulting from short-term (1–4 hours) exposures to dichloromethane at levels up to 1,000 ppm, or longer-term exposures at levels up to 500 ppm. The 8-hour threshold limit value before 1975 was 500 ppm (NIOSH, 1986). These studies are described below. With the exception of Putz et al. (1979), there is no description in the published reports of the informed consent and other human subjects research ethics procedures undertaken in these studies, but there is no evidence that the conduct of the research was fundamentally unethical or significantly deficient relative to the ethical standards prevailing at the time the research was conducted.

In 1972, Stewart et al. (1972a, b) reported results from four experiments that were initiated because of the chance observation of an elevation in COHb saturation levels in an individual (one of the investigators) the morning after he had spent 2 hours working with varnish remover. Participants were medical students and faculty (including at least one of the coauthors). A total of 11 healthy nonsmoking volunteers were placed in an exposure chamber with mean concentrations of dichloromethane ranging from 213 to 986 ppm for 1 or 2 hours. These experiments indicated that dichloromethane exposure at these levels resulted in COHb saturation levels that exceeded and were more prolonged than those seen with threshold limit value exposures to CO. The exposures also resulted in symptoms of CNS depression indicated by visual evoked response changes and reports of light-headedness. Although return of COHb

levels to background levels could take >24 hours, all of the other symptoms were reversible within a few hours after exposure ceased.

Winneke (1974) measured auditory vigilance, visual flicker fusion frequency, and 14 psychomotor tasks in a total of 38 women exposed to dichloromethane levels of 300–800 ppm for 4 hours in an exposure chamber. A comparison group (nine females, nine males) exposed to 100 ppm CO for 5 hours was also included. Exposure to 800 ppm dichloromethane resulted in a statistically significant decrease in the performance of 10 of the 14 psychomotor tasks. In tests of auditory vigilance and visual flicker fusion, depressed response was seen at 300 ppm and was further depressed at 800 ppm. These effects were not seen with CO exposure.

Forster et al. (1974) exposed four healthy young men to dichloromethane levels ranging from 0 to 500 ppm for 7.5 hours/day for a total of 26 days over a 6-week period to investigate alterations in hemoglobin affinity for oxygen and altered pulmonary function. While no changes were observed in pulmonary function, hemoglobin affinity for oxygen was increased with no indication of adaptation to restore this affinity for oxygen to normal.

Putz et al. (1979) examined the behavioral effects seen after exposure to dichloromethane and to CO. Twelve healthy volunteers (six men and six women) each acted as his/her own control in separate 4-hour exposures to 70 ppm CO and 200 ppm dichloromethane. These levels were chosen so that the COHb level would reach 5% from both the CO and dichloromethane exposures. The experiments were conducted in a double-blind manner so that neither the investigators nor the participant knew the exposure condition under study at any particular time. Informed consent was obtained, and the study was reviewed by the National Institute of Occupational Safety and Health (NIOSH) Human Subject Review Board. The performance tests were dual tasks (an eye-hand coordination task in conjunction with a tracking task), with five measures of performance assessed at six time points over the 4-hour test period and an auditory vigilance task. Two levels of difficulty were assessed for each task to allow assessment of whether the exposure effect was similar in low and high difficulty tasks. The tests of eye-hand coordination, tracking tasks, and auditory vigilance revealed significant impairment with both exposures under the more difficult task conditions. Effects were similar or stronger in magnitude for dichloromethane compared with CO.

4.1.2.3. *Observational Studies Focusing on Clinical Chemistries, Clinical Examinations, and Symptoms*

Studies in currently exposed workers. Ott et al. (1983a, c, d) evaluated several parameters of hepatic, hematopoietic, and cardiac function in workers exposure to dichloromethane in a triacetate fiber production plant in Rock Hill, South Carolina. Two hundred sixty-six Rock Hill workers and a comparison group of 251 workers in an acetate fiber production plant in Narrows, Virginia, were included in the examination of urinary and blood measures. These groups included men and women, blacks and whites, and smokers and

nonsmokers. The median 8-hour TWA exposure for dichloromethane ranged from 60 to 475 ppm in Rock Hill. Acetone at levels up to >1,000 ppm was present in both plants, but dichloromethane and acetone exposures were inversely related.

There were differences in blood collection procedures between the two plants and in the age, sex, race, and smoking history distribution of the study groups. The demographic and smoking differences were accounted for in the analysis by stratification. Statistically significant differences were seen between the workers in the two plants for COHb, serum alanine aminotransferase (ALT), total bilirubin, and mean corpuscular hemoglobin concentration (MCHC) (although the direction and magnitude of these differences were not reported, and the authors stated that the difference in serum ALT could be due to the differences in blood collection procedures, which involved a sitting versus recumbent position of the subjects at the exposed and nonexposed plants, respectively) (Ott et al., 1983c). Within the Rock Hill plant, analyses were also conducted to examine associations between dichloromethane exposure and the clinical parameters within specific race-sex groups by using multiple regression to control for smoking status, age, and time of venipuncture. Positive associations were seen with COHb in all race-sex groups (increases of 0.7–2.1% per 100 ppm increase in dichloromethane) and with total bilirubin (increases of 0.05–0.08 mg/dL per 100 ppm increase in dichloromethane) in all groups except nonwhite men (which was a much smaller group, $n = 20$, than the other groups). Red cell count, hematocrit, hemoglobin, and aspartate aminotransferase (AST) were also positively associated with dichloromethane exposure in white females. The increase in total bilirubin level was not supported by parallel changes in other measures of liver function or red blood cell turnover, suggesting that this measure was not reflecting liver damage or hemolysis.

The increased red cell count, hemoglobin, and hematocrit in women exposed to high levels of dichloromethane (up to 475 ppm, 8-hour TWA) may indicate a compensatory hematopoietic effect. The fact that these changes were not significant among men may be due to higher baseline hemoglobin, which was observed when comparisons were made between nonsmoking men and women. No such difference in the baseline values was observed among the smoking men and women, suggesting that the compensatory advantage may be lost among smokers.

Ott et al. (1983e) presented results from a further investigation of changes in COHb, alveolar CO, and oxygen half-saturation pressure in relation to dichloromethane exposure. Blood samples were collected before and after shifts from 136 Rock Hill and 132 Narrows workers. For the Rock Hill workers, personal monitoring for dichloromethane exposure was done during the shift. The TWA for dichloromethane ranged from 0 to 900 ppm, with a bimodal distribution (peaks around 150 and 500 ppm) resulting from the layout of the plant. The blood samples were used to determine blood COHb, alveolar CO levels, and the partial oxygen pressure (P_{50} ; that is, the pressure required to keep 50% of the blood oxygen-carrying capacity saturated with oxygen at pH 7.4 and 37°C). Separate analyses were conducted for smokers and

nonsmokers to account for the smoking-related effects on COHb. Linear relationships were seen between dichloromethane exposure and the before-shift COHb and alveolar CO levels, reflecting residual CO metabolism from the previous day's exposure. There were significant quadratic relationships between dichloromethane exposure and the postshift COHb and alveolar CO levels, indicating a partial saturation of the enzyme system metabolizing dichloromethane. The P₅₀ group means were lower among the exposed compared with the referents, among smokers compared with nonsmokers, and among men compared with women. Given the relationship between COHb and P₅₀, an expected decrease in P₅₀ during the shift was observed among the exposed.

Continuous 24-hour cardiac monitoring was also evaluated in a smaller sample of 24 dichloromethane-exposed workers from the triacetate fiber production plant in Rock Hill, South Carolina, and 26 workers from the comparison plant in Narrows, Virginia. This study (Ott et al., 1983d) was limited to white men ages ≥ 35 years. Special efforts were made to recruit men with a history of heart disease, because this group was postulated to be most likely to demonstrate positive findings. The estimated TWA dichloromethane exposure ranged from 60 to 475 ppm in the exposed group. The evaluation examined ventricular and supraventricular ectopic activity and S-T segment depression in the exposed and nonexposed groups. Comparisons were also made between cardiac performance during work hours and nonwork hours to discern possible short-term effects of recent exposure. Comparing the findings for the 24 exposed and 26 referent volunteers indicated no difference in ventricular or supraventricular ectopic activity or S-T-segment depression. There was no difference comparing work and nonwork hours among exposed volunteers.

Soden et al. (1996) studied all active male workers exposed to dichloromethane at a Hoechst Celanese triacetate film production plant in Belgium. The production process was the same as the process at the Hoechst Celanese Rock Hill plant, except the Belgium plant was newer with better engineering controls to significantly reduce overall levels of the dichloromethane, acetone, and methanol used in the process. The objectives of the study were to determine the impact of varying levels of dichloromethane exposure on COHb levels, whether successive days of dichloromethane exposure affected the COHb levels, and what impact smoking had on COHb levels in conjunction with dichloromethane exposure. Workers were monitored semiannually for COHb at the end of the work shift and were personally monitored for exposure to the three solvents. Smoking status was defined based on a health assessment questionnaire, with smokers smoking at least one cigarette per day. Among nonsmokers, a dose response was found among COHb levels and average dichloromethane exposure levels in the range of 7–90 ppm. The maximum COHb was 4.00% at an average exposure of 90 ppm (correlation coefficient = 0.58, $p < 0.05$). Smokers' COHb levels were elevated when compared with those of nonsmokers with similar dichloromethane air levels, but the dose-response correlation between dichloromethane air levels and COHb levels was weaker and not statistically

significant (correlation coefficient 0.20). The maximum COHb level for smokers was 6.35% at an average dichloromethane air level of 99 ppm. The authors concluded that dichloromethane exposures up to the levels observed did not produce COHb levels that are likely to cause cardiac symptoms.

Cherry et al. (1983, 1981) reported the results of health evaluations of two studies of triacetate film production workers. Cherry et al. (1981) recruited 46 of the 76 male workers at a triacetate film factory, where workers were exposed to dichloromethane and methanol in a ratio of 9:1 at air levels of dichloromethane ranging from 75 to 100 ppm. A small comparison group ($n = 12$) of workers at this factory who worked a similar shift pattern (rapidly rotating shifts) but who were not exposed to dichloromethane was also included. The men were asked whether they had ever experienced cardiac symptoms (pain in the arms, chest pain sitting or lying, or chest pain when walking or hurrying) and were asked about the presence in the past 12 months of neurological disorders (frequent headaches, dizziness, loss of balance, difficulty remembering things, numbness and tingling in the hands or feet), affective symptoms (irritability, depression, tiredness), and stomachache (as an indicator of symptom overreporting). No difference in response was found in history of stomachache (reported by 15% of exposed workers compared with 17% nonexposed workers). Six of the exposed and none of the unexposed men responded positively to the cardiac symptoms. The exposed group reported an excess of neurological symptoms; the number (and proportion) reporting zero, one, two, and three or more symptoms were 26 (0.56), 8 (0.17), 9 (0.20), and 3 (0.07), respectively, in exposed workers compared with 11 (0.92), 1 (0.12), 0 (0.00), and 0 (0.00), respectively, in controls ($p < 0.02$ for χ^2 test of linear trend). With respect to affective symptoms, the number (and proportion) reporting zero, one, two, and three symptoms were 28 (0.61), 6 (0.13), 7 (0.15), and 5 (0.11), respectively, among the exposed workers, and 9 (0.75), 2 (0.17), 1 (0.08), and 0 (0.0), respectively, among the unexposed workers. The authors concluded that there was no difference between exposed and nonexposed in reporting of affective symptoms based on a χ^2 test of linear trend. There was no discussion of the statistical power of this test or of tests of the proportion reporting a specified number of symptoms (which may be a more appropriate test given the sample size), but it is clear that the statistical power of this test was very low. For example, taking the simple case of the comparison of the proportion reporting two or more symptoms and using the approximate estimates from this study (25 and 10% in the exposed and unexposed, respectively), approximately 75 exposed and 300 unexposed workers would be needed for a power of 0.80 (i.e., an 80% chance of rejecting the null hypothesis when the null hypothesis was false); the actual power with the sample size of 46 and 12 is <0.10 .

Based on these results, a follow-up study was conducted which included a larger referent group. This study included the symptom list described in the previous paragraph, a standardized clinical exam (including an electrocardiograph), and neurological and psychological tests of nerve conduction, motor speed and accuracy, intelligence, reading, and memory (Cherry et al.,

1981). Twenty-nine of the original 46 exposed workers participated in the follow-up. The men who did not participate in the follow-up were similar in age and symptoms to the men who did. The new referent group was recruited from another plant with the same owner and a very similar process but without dichloromethane exposure. One control, age-matched within 3 years, was selected for each exposed worker. No differences between the groups were found in the clinical exam, electrocardiogram, or nerve conduction tests. A statistically significant ($p < 0.05$) deficit among the exposed workers was found for coarse motor speed. On two tests of overall intelligence, the exposed group did significantly better than the referent, but on a reading ability test designed to assess premorbid educational level, scores for the exposed group were slightly lower than for the referent group. (Only one of these three differences, the trail making intelligence test, was statistically significant.) With respect to the report of neurological symptoms in the past year, the number (and proportion) reporting zero, one, two, and three symptoms were 17 (0.59), 4 (0.14), 6 (0.21), and 2 (0.07), respectively, among the exposed workers, and 21 (0.72), 6 (0.21), 0 (0.0), and 2 (0.07), respectively, among the unexposed workers, with a test of linear trend that was not statistically significant. The authors interpret the results as indicating that the differences in neurological symptoms seen in the initial study were due to chance and that, taken as a whole, the exposed workers had no detrimental effect attributable to dichloromethane exposure. Again, the limitations of the statistical power of the analysis and alternative interpretations that might have resulted from approaches taken to improve the power were not discussed. These approaches include combining the unexposed groups from the two analyses, using the full sample of the exposed group instead of the subset of 29 who completed the clinical exam, or using a different test (i.e., of a proportion rather than a linear trend).

Cherry et al. (1983) compared dichloromethane-exposed workers at an acetate film factory to nonexposed workers (from the same plant but from areas without solvent contact or from another film production factory in which solvents were not used). The 56 exposed and 36 unexposed workers were matched to within 3 years of age. Both factories were on rapid rotating shifts. Exposure to dichloromethane ranged from 28 to 173 ppm, using individual air sampling pumps. Blood samples were taken to monitor dichloromethane levels at the beginning and end of the shift. Study participants were asked to rate sleepiness, physical and mental tiredness, and general health on visual analog scales with the extreme responses at either end. Participants were also given a digit symbol substitution test and a test of simple reaction time. No differences were seen between exposed and unexposed groups at the beginning of the shift on the four visual analog scales, but the exposed deteriorated more on each of the scales than did the controls. This difference in deterioration was statistically significant ($p \leq 0.05$) during the morning shift but was not statistically significant during the afternoon or night shifts. A significant correlation was shown between change in mood over the course of the shift and level of dichloromethane in the blood. No difference was seen between the exposed and referents on

the tests of reaction time or digit substitution. However, among the exposed, deterioration in the digit substitution tests at the end of the shift was significantly related to blood dichloromethane levels.

Anundi et al. (1993) studied 12 men who worked in a graffiti-removing company. Each worker filled out a questionnaire about previous occupational and nonoccupational exposure to solvents and use of protective equipment. Half-day breathing zone samples were taken for each of the 12 workers, and 15-minute samples were also taken for 10 workers. On the day the air sampling was done, a structured interview pertaining to recent diseases or symptoms related to allergies, asthma, diseases of the skin, respiratory organs, gastrointestinal tract, urinary organs, neurological trauma and disease, and neuropsychiatric symptoms was conducted by a physician, and blood and urine samples were collected. The results were compared with those of 233 men from the area population. The 12 men (mean age 23 years) had worked between 3 months and 4.5 years cleaning graffiti from underground stations. No respiratory protection was used, and the leather gloves were frequently soaked with solvent. While mixed solvent was used to do the cleaning, dichloromethane was the predominant component, as confirmed by the air samples. The geometric mean (GM) of the TWA calculated from the half-day samples was 127 mg/m^3 (range 18–1,188 mg/m^3), with half of the samples exceeding the Swedish permissible exposure limit of 120 mg/m^3 . The GM of the 15-minute samples was 400 mg/m^3 (range 6–5,315 mg/m^3), with most samples exceeding the Swedish short-time exposure limit of 300 mg/m^3 . Two workers had clinical laboratory data outside the normal range (urinary α_1 - or β_2 -microglobulin, serum ALT, γ -glutamyl transpeptidase), which could indicate possible kidney and liver damage. The authors stated that in both cases, factors other than the solvent exposure (i.e., urinary tract medical condition preceding employment, history of renal stones) could have influenced these laboratory results. The prevalence of irritation of the eyes and upper respiratory tract (blocked nose and nasal catarrh) was much higher in the graffiti-cleaning workers compared with the referent group (e.g., >70% of the workers compared with 18% of the comparison group reported a blocked nose; ~50% of workers and 15% of the comparison group reported eye irritation), but there were no or much smaller differences in abnormal tiredness, headache, nausea, or irritative cough. No acute effects on the CNS were noted.

Studies in retired workers. Lash et al. (1991) examined the hypothesis that long-term exposure to dichloromethane produces lasting CNS effects as measured by long-term impairment on memory and attention centers. Retired aircraft maintenance workers employed in at least 1 of 14 targeted jobs with dichloromethane exposure for ≥ 6 years between 1970 and 1984 were compared to a like group of workers without dichloromethane exposure. The unexposed workers were also retired aircraft mechanics at the same base and held 1 of 10 jobs in the jet shop where little solvent was used. The exposed group made up of painters and mechanics in the overhaul department was chosen to maximize the exposure contrast yet minimize differences in potential

confounders between exposed and nonexposed groups. Exposures were typically within state and federal guidelines for dichloromethane exposure. From 1974 to 1986, when 155 measurements for dichloromethane exposure were made, mean breathing zone TWAs ranged from 82 to 236 ppm and averaged 225 ppm for painters and 100 ppm for mechanics.

Data collection occurred in three phases: (1) an initial questionnaire was given to all retired members of the airline mechanics union to identify eligible workers, (2) a telephone survey was conducted to collect medical, demographic, and general employment criteria, and (3) subjects who qualified were then recruited to participate in the medical evaluation. Sixty percent of the 1,758 retirees responded to the questionnaire, and 259 of these retirees met the eligibility criteria. Ninety-one men qualified for the medical evaluation based on the telephone survey; 25 retirees exposed to solvents and 21 unexposed retirees participated in the evaluation. All were men between the ages of 55 and 75 without a history of alcoholism or any neurological disorder. The 25 exposed participants worked an average of 11.6 years in dichloromethane-exposed jobs during the target period and 23.8 years in the industry.

The medical evaluation included a questionnaire about the occurrence of 33 different symptoms in the past year, physiological measurement of odor and color vision senses, auditory response potential, hand grip strength, and measures of reaction time (simple, choice, and complex), short-term visual memory and visual retention, attention, and spatial ability. The only large differences (i.e., effect size, or mean difference between groups divided by the SD of the outcome measure, of ≥ 0.4) between the two groups were a higher score on verbal memory tasks (effect size approximately 0.45, $p = 0.11$) and lower score on attention tasks (effect size approximately -0.55, $p = 0.08$) and complex reaction time (effect size approximately -0.40, $p = 0.18$) in the exposed compared with the control group. (Although not noted by the authors, the power to detect a statistically significant difference between the groups given this sample size was low [i.e., approximately 0.30 for an effect size of 0.40, using a two-tailed alpha of 0.05]) (Cohen, 1987). The authors investigated the possibility of response bias, given the low initial response to the mailed questionnaire recruiting retirees and the small number of workers from the entire pool of eligible participants who actually participated in the medical evaluation. Attempts were made to contact 30% of the questionnaire nonrespondents, with 46% contacted and 31% completing the telephone interview. The only difference found between those who responded to the mailed questionnaire and those who did not was a higher percentage of diagnosed heart disease among the nonrespondents who were 2.5 years older and had been retired 1.7 more years than the respondents. Those who were eligible but did not participate in the medical evaluation were similar to the exam participants on all characteristics included in the interview. The only difference was a higher prevalence of gout among the unexposed who did not participate compared to the unexposed who did participate.

4.1.2.4. *Observational Studies Using Workplace Medical Program Data*

Kolodner et al. (1990) investigated the effect of occupational exposure to dichloromethane on six health outcomes identified in the literature or based on biological plausibility. Participants in the study were male workers at least 19 years old at two General Electric plastic polymer plants where dichloromethane was one of the chemicals used. Four dichloromethane exposure categories were established based on full-shift personal air monitoring data (8-hour TWA) collected in 1979–1985, job titles, and industrial hygienists' knowledge of plant operations. The mean 8-hour TWA and number of workers in each of the four exposure groups were 49.0 ppm for the 19 workers in the highest, 10.9 ppm for the 49 workers in the intermediate, 3.3 ppm for the 56 workers in the low, and <1.0 ppm for the 772 workers in the minimal/no exposure group.

Data from 1984 annual medical exams and 1985 absence data from payroll records were evaluated for possible health effects resulting from occupational exposure to dichloromethane. A high percentage of workers participated in the annual medical exams, with only 5 of the 896 eligible for inclusion in the study refusing the exam completely in 1984. Six hypotheses were specifically tested regarding dichloromethane exposure in relation to different health outcomes: absence due to illness, hepatotoxicity (manifested by nausea, weakness and fatigue, palpable liver, abdominal tenderness, jaundice, hepatomegaly, abnormal serum γ -glutamyl transferase, ALT, AST, or bilirubin), diabetes mellitus (manifested by weight loss, weakness and fatigue, polydipsia, polyuria, impaired vision, excessive weight loss, elevated fasting blood sugar, and abnormal urinary glucose or urinary acetone), CNS toxicity (manifested by headache, lightheadedness, dizziness and vertigo, ataxia, weakness and fatigue, and abnormalities detected in the central motor, central sensory, cranial nerve, gait, neurocoordination, or Bibinski reflex examinations), cardiovascular abnormalities (manifested by fatigue, dyspnea, chest pain with exertion, palpitations, or abnormalities detected in the point maximum impulse exam, blood pressure measurements, or electrocardiogram), and neoplastic breast changes (154 women were included in this portion of the study—manifested by painful breast, breast swelling, lump, nipple discharge, or abnormalities detected in the breast examination).

Workers were placed in exposure categories based on their current jobs. In addition, exposure to high noise levels occurred in both plants, and workers in each plant had exposure to another chemical, either phenol or phosgene. The authors noted that workers tended to move from entry-level jobs with high dichloromethane exposure to supervisory jobs with lower dichloromethane exposure, based on the seniority system in place at both plants. Thus, current exposure levels reported did not necessarily reflect cumulative exposure. Because of the way the seniority system moved workers through jobs and the fact that workers were assigned to dichloromethane exposure categories based on their current job, age was inversely related to exposure and was controlled in the analysis of some of the continuous variables using analysis of covariance. Age adjustment was not employed in the analysis of dichotomous variables. The

mean age was 35.3, 39.7, 37.1, and 29.5 years in the minimal/no, low, medium³, and high exposure groups, respectively. The small number of workers in the exposed groups limited the ability to evaluate the effects of dichloromethane exposure on health outcomes related to age, since age had to be adjusted in these analyses. The racial distribution did not differ among the exposure groups.

The authors indicated that all the hypotheses were accepted with the exception of CNS symptoms. However, it should be noted that the small size and younger age distribution in the high exposure group and the lack of adjustment for age in most of the analyses make it difficult to interpret the statistical testing that was performed. Data pertaining to neurological, hepatic, and cardiac function are shown in Table 4-1. Among the six neurological symptoms evaluated, a statistically significant positive exposure-effect relationship between dizziness/vertigo and dichloromethane exposure was identified. This trend was driven most strongly by the low frequency of this reported symptom in the minimal/no exposure group (1.2%), but there was no linear trend across the higher levels of exposure (7.5, 2.1, and 5.3% in the low, medium, and high exposure groups, respectively).

Table 4-1. Percentage of male General Electric plastic polymer workers reporting neurologic symptoms or displaying abnormal values in measures of neurological function, hepatic function, and cardiac function

	Exposure group ^a			
	Minimal/no (n = 772)	Low (n = 56)	Medium (n = 49)	High (n = 19)
Neurological				
Headache	8.7	7.5	10.4	5.3
Lightheadedness	2.9	3.8	4.2	5.3
Dizziness/vertigo	1.2	7.5	2.1	5.3
Ataxia	0.0	1.9	0.0	0.0
Babinsky	0.0	0.0	0.0	0.0
Gait	0.0	0.0	0.0	0.0
Faintness/syncope ^b	0.1	0.0	2.1	0.0
Seizures ^b	0.4	0.0	2.1	0.0
Paresis/paralysis ^b	0.7	0.0	0.0	0.0
Parasthesis ^b	4.0	7.5	14.6	0.0
Head trauma/concussion ^b	0.8	1.9	0.0	0.0
Peripheral motor exam ^{b,c}	0.5	0.0	0.0	0.0
Peripheral sensory exam ^{b,c}	1.1	2.4	5.1	0.0
Rhomberg exam ^{b,c}	0.0	0.0	2.6	0.0

³The “medium” exposure group is also referred to as the “intermediate” exposure group in Kolodner et al. (1990).

Table 4-1. Percentage of male General Electric plastic polymer workers reporting neurologic symptoms or displaying abnormal values in measures of neurological function, hepatic function, and cardiac function

	Exposure group ^a			
	Minimal/no (n = 772)	Low (n = 56)	Medium (n = 49)	High (n = 19)
Hepatic				
Serum gamma glutamyl transferase	8.0	16.1	12.2	5.3
Serum total bilirubin	3.0	1.8	2.0	10.0
Serum AST	1.8	3.6	4.1	0.0
Serum ALT	9.1	10.7	8.2	5.3
Cardiac^d				
Palpitations: percent abnormal	1.2	9.1	2.1	0.0
Electrocardiogram				
Borderline/abnormal	18.5	16.7	19.1	8.3
Bradycardia/tachycardia abnormalities ^b	20.2	16.7	25.5	0.0
General rhythm abnormalities	12.0	11.1	17.0	8.3
Atrial, atrioventricular, or sinus abnormalities	0.8	0.0	0.0	0.0
Bundle blocks or ventricular abnormalities	3.9	5.6	10.6	8.3
Axis deviations	2.6	1.9	2.1	8.3
Wave abnormalities	4.0	3.7	10.6	0.0
Hypertrophy	3.8	3.7	6.4	0.0
Evidence of infarction	2.3	5.6	2.1	0.0

^aMean 8-hr TWA exposure was <1.0, 3.3, 0.9, and 49.0 ppm in the minimal/no, low, medium, and high groups, respectively; mean age 35.3, 39.7, 37.1, and 29.5 yrs in the minimal/no, low, medium, and high groups, respectively.

^bThe authors considered these to be screening variables rather than hypothesis-testing variables.

^cn = 629, 42, 39, and 14 in the minimal/no, low, medium, and high groups, respectively.

^dFor all cardiac outcomes except bradycardia/tachycardia, n = 728, 54, 47, and 12 in the minimal/no, low, medium, and high groups, respectively. For bradycardia/tachycardia, n = 727 in the minimal/no group.

Source: Kolodner et al. (1990).

Soden (1993) compared health-monitoring data from dichloromethane-exposed workers in the Rock Hill triacetate fiber production plant to workers from another plant making polyester fibers owned by the same company in the same geographic area. Exposed and control workers were chosen from among workers who had worked at least 10 years in their respective areas and who participated in the company's health-monitoring program between 1984 and 1986 and were still employed on December 31, 1986. Controls were matched by race, age, and gender to each Rock Hill worker for a sample size of 150 and 260 in the exposed and control groups, respectively. (The aim of the study had been 1:2 matching.) The 8-hour TWAs among the Rock Hill workers were those reported by Lanes et al. (1990), namely 475 ppm for dichloromethane, 900 ppm for acetone, and 100 ppm for methanol. None of these exposures occurred at the polyester plant. There was a 90% participation rate in the health-monitoring program. Six

questions in the health history portion of the health-monitoring program concerned cardiac and neurological symptoms (chest discomfort with exercise; racing, skipping, or irregular heartbeat; recurring severe headaches; numbness/tingling in hands or feet; loss of memory; dizziness). Part of this program included blood samples used for standard clinical hepatic and hematologic parameters: serum ALT, AST, total bilirubin, and hematocrit. The clinical measures were available for 90 (60%) of the exposed and 120 (46%) of the control group; some participants declined this part of the health-monitoring program because similar tests had been part of recent personal medical care.

There was little difference in the frequency of reported symptoms between exposed workers and controls: chest discomfort reported by 2.0% of exposed and 4.0% of the controls, irregular heartbeat reported by 5.5% of exposed and 6.0% of the controls, recurring severe headaches reported by 3.5% of exposed and 5.5% of the controls, numbness/tingling in hands and feet reported by 6.4% of exposed and 8.1% of the controls, loss of memory reported by 1.3% of exposed and 0.4% of the controls, and dizziness reported by 2.7% of exposed and 4.8% of the controls (Soden, 1993). The levels of the blood values were similar in the exposed and control groups, except for a 3.1 IU/L decrease in serum AST activity ($p = 0.06$). The authors concluded that this difference was not clinically significant, but they did not discuss the potential bias introduced by the selective participation in this part of the study.

4.1.2.5. *Studies of Ischemic Heart Disease Mortality Risk*

Several studies have examined the relation between dichloromethane exposure and risk of cardiovascular-related mortality. The methodological details of these studies are described in Section 4.1.3.2). No evidence of increased risk of ischemic heart disease mortality was seen in two triacetate film production cohort studies (Hearne and Pifer, 1999; Tomenson et al., 1997) or in two triacetate fiber production cohort studies (Gibbs et al., 1996; Lanes et al., 1993). Information on this outcome was not included in the dichloromethane analysis of civilian Air Force base workers (Blair et al., 1998). The standardized mortality ratios (SMRs) for ischemic heart disease mortality were <1.0 in all of the cohorts and dose groups examined (Table 4-2). The “healthy worker effect” may have contributed to these observations. There are no case-control studies of ischemic heart disease and dichloromethane exposure.

Table 4-2. Ischemic heart disease mortality risk in four cohorts of dichloromethane-exposed workers

		Obs	Exp	SMR	95% CI
Triacetate film production					
Hearne and Pifer (1999)	Cohort 1 (men)	117	136.7	0.86	0.71–1.03
	Cohort 2 (men)	122	143.3	0.85	0.71–1.02
Tomenson et al. (1997)	Men	114	123.9	0.92	0.76–1.10
Triacetate fiber production					
Lanes et al. (1993)	Men and women	43	47.8	0.90	65–121
Gibbs et al. (1996)	Men				
	50–100 ppm	96	100.1	0.96	0.78–1.2
	350–700 ppm	98	106.8	0.92	0.75–1.1
	Women				
	50–100 ppm	32	45.8	0.70	0.48–0.99
	350–700 ppm	0	3.4	–	0.0–1.1

CI = confidence interval; Exp = number of expected deaths; Obs = number of observed deaths

4.1.2.6. Studies of Suicide Risk

Suicide risk is not an outcome that was a primary hypothesis or motivation of the cohort studies, but it may be relevant given the potential neuropsychological effects of dichloromethane, as evidenced from studies of acute and chronic exposure scenarios described previously. In a triacetate film production cohort in Rochester, New York, Hearne and Pifer (1999) reported 14 observed deaths from suicide compared with 7.8 expected, for an SMR of 1.8 (95% confidence interval [CI] 0.98–3.0) (Table 4-3). This cohort (“Cohort 1”) consisted of 1,311 men who were first employed between 1946 and 1970 and were followed through 1994. Similar results were seen in a different, but somewhat overlapping, cohort in this study (“Cohort 2”) of 1,013 men employed between 1964 and 1970 and followed through 1994 (see Section 4.1.3.3.1). There was also evidence of increasing suicide risk with dichloromethane exposure, particularly in the highest exposure group, in the study of triacetate fiber production workers in Maryland (Gibbs, 1992). The triacetate fiber production cohort study in Rock Hill, South Carolina, has published what appears to be erroneous information about suicide risk. In the 1993 paper (Lanes et al., 1993), 4 observed and 5.21 expected cases were reported (SMR 0.77), but the SMR that was reported with these data was 1.19 (95% CI 0.39–2.8). This ratio would correspond to 6 observed and around 5.2 expected cases. Information on suicide was not included in the other film and fiber cohort studies (Tomenson et al., 1997) or in the analysis of civilian Air Force base workers (Blair et al., 1998). There are no case-control studies of suicide risk and dichloromethane exposure.

Table 4-3. Suicide risk in two cohorts of dichloromethane-exposed workers

		Obs ^a	Exp ^b	SMR	95% CI
Triacetate film production					
Hearne and Pifer (1999)	Cohort 1	14	7.8	1.8	0.98–3.0
	Cohort 2	9	5.1	1.8	0.81–3.4
Triacetate fiber production^c					
Gibbs (1992)	50–100 ppm	8	6.4	1.3	0.54–2.5
	350–700 ppm	8	4.4	1.8	0.78–3.6

^aObs = number of observed deaths.

^bExp = number of expected deaths.

^cOne additional study provided data on suicide risk, but some kind of error seems to be present: 4 observed and 5.21 expected cases were reported in Lanes et al. (1993), which would be an SMR of 0.77, but the SMR reported with these data was 1.19 (95% CI 0.39–2.8). This ratio would correspond to 6 observed and around 5.2 expected cases.

4.1.2.7. Studies of Infectious Disease Risk

There is limited information pertaining to infectious disease risk in relation to dichloromethane exposure. Only one of the cohort studies (Hearne and Pifer, 1999) reported data for the broad category of infectious and parasitic disease mortality. In Cohort 1 of this analysis, there were no observed deaths in this category (5.6 expected), and in Cohort 2 there were 3 observed and 4.7 expected deaths, for an SMR of 0.64. The detailed report by Gibbs (1992) of the cellulose triacetate fiber production cohorts in Maryland (Gibbs et al., 1996) also contained information on the facility in South Carolina that was the site of the report by Lanes et al. (1993, 1990). Slightly elevated risks of mortality due to influenza and pneumonia were seen among the male workers in the high exposure group in Maryland (7 observed, 5.62 expected, SMR 1.25) and in South Carolina (3 observed, 1.33 expected, SMR 2.26). Among females, there were few observed or expected cases (in Maryland, 1 observed, 0.23 expected, SMR 4.36; in South Carolina, 0 observed, 0.74 expected).

4.1.2.8. Studies of Reproductive Outcomes

Pregnancy outcomes in women exposed to dichloromethane have been investigated in two studies. Taskinen et al. (1986) studied spontaneous abortions among women employed in eight pharmaceutical factories between 1973 and 1980. Data on pregnancy outcomes were collected from a national hospital and clinic discharge registry in Finland from 1973 to 1981 by matching the worker rosters to the registry. Exposure to dichloromethane was one of eight solvents or classes of solvents included in the study. The study consisted of two parts. The first investigated the rate of spontaneous abortions (number of spontaneous abortions divided by the sum of spontaneous abortions and births) during, before, or after employment in the pharmaceutical industry. One hundred and forty-two spontaneous abortions and 1,179 births were identified among the female workers at the eight plants. Employment hire and termination

dates were obtained from plant records. The spontaneous abortion rate was 10.9% during employment compared with 10.6% before and after employment. These results compared to a rate of 8.5% in the general population in the geographic area where the factories were located. The rate of spontaneous abortions among workers declined over the period of the study, with a 3-year moving average of 15% at the beginning declining to 9.5% at the end of the study. Over the same period, the industrial hygiene allegedly improved in the plants. Ten congenital malformations of different types were identified among the women (five among those who were employed in the pharmaceutical industry during the pregnancy and five among those whose pregnancies occurred before or after this employment).

The second part of the study by Taskinen et al. (1986) was a case-control study of the risk of spontaneous abortions in relation to workplace exposures during pregnancy. The source population consisted of women who were employed in one of the eight Finnish pharmaceutical factories during at least 1 week of the first trimester of pregnancy during the study period. Cases ($n = 44$) were selected from this population based on hospital or clinic records indicating a spontaneous abortion, and 130 controls (women who had given birth) were age-matched (3:1 matching; age within 2.5 years) to each case. Occupational exposure data were obtained by questionnaires completed by the plant physician or the nursing staff, blinded to the case status of the study member, in consultation with labor protection chiefs and department foremen. The questionnaire requested information about job history and job tasks, exposure to eight specific solvents or classes of solvents (aliphatic solvents, alicyclic solvents, toluene, xylene, benzene, chloroform, dichloromethane, and other solvents), antineoplastic agents, carcinogens, hormones, antibiotics, heavy lifting, known chronic diseases, acute diseases during pregnancy, smoking status, and previous pregnancies. Exposure frequency to each solvent was based on the cumulative weighted sum of the number of days/week the woman was exposed to the solvent. While overall response to the questionnaire was 93%, less than half the questionnaires contained information about smoking or previous pregnancies, precluding inclusion of these variables in the analysis. The distribution of broad categories of occupations (i.e., pharmaceutical workers and packers, laboratory assistants) was similar in both groups. However, exposure to each of the solvents was higher in the cases compared with controls, and the results for dichloromethane were relatively strong. For dichloromethane, the prevalence of exposure was 28.9 and 14.3% in cases and controls, respectively, resulting in an odds ratio (OR) of 2.3 (95% CI 1.0–5.7). There was also evidence of an increasing risk with higher exposure frequency, with an OR of 2.0 (95% CI 0.6–6.6) with exposures of less than once a week and 2.8 (95% CI 0.8–9.5) with exposures of once a week or more. An association was also seen with exposure to four or more solvents (OR 3.4, [95% CI 1.0–12.5]), and weaker associations were seen with other specific solvents (e.g., chloroform, toluene).

Bell et al. (1991) investigated the relation between birth weight and maternal exposure to airborne dichloromethane as a result of living around the triacetate film facility in Rochester,

New York. For this population-based cross-sectional study, birth certificates were obtained for all births in 1976–1987 in Monroe County, where the triacetate film facility is located. Multiple births and births of infants weighing <750 grams were excluded. Data abstracted from the certificate included date of birth, census tract of residence, age, race, educational level of the mother and father, sex, gestational age, multiple births, month of the pregnancy that prenatal care began, total previous births, total previous live births, and conditions present during the pregnancy. An air dispersion modeling system for 250 air emissions, including dichloromethane and predicting average annual ground level concentrations in the surrounding community, was used to assign dichloromethane exposure levels to each birth mother. One of four levels of exposure was assigned to each census tract based on the isopleth of exposure in which more than half of the census tract population resided. Because of the few births among nonwhites that occurred in areas of higher exposure, the study was restricted to whites (n = 91,302). The number of births that occurred in each of the four exposure levels was n = 1,085 in the high-exposure group (50 µg/m³ [0.014 ppm]), n = 1,795 in the moderate-exposure group (25 µg/m³ [0.007 ppm]), n = 6,044 in the low-exposure group (10 µg/m³ [0.003 ppm]), and n = 82,076 in the no-exposure group. At the levels of dichloromethane exposure in this population, no significant adverse effect on birth weight was found. There was an 18.7 g decrease in birthweight (95% CI 51.6–14.2) in the high- compared with the no-exposure group, adjusting for maternal age, maternal education, parity, previous pregnancy loss, late start of prenatal care, sex of the child, and pregnancy complications. No significant association was found between any combination of exposure levels and birth weight. There was no association between exposure group and risk of a low birthweight infant (i.e., <2,500 g, OR 1.0 [95% CI 0.81–1.2] in the high-compared with the no-exposure group). The authors point out a number of problems with assignment of dichloromethane exposure. It is possible that the dichloromethane exposure was overestimated using the model. Comparisons to ambient air sampling levels collected 6 times/year resulted in the dichloromethane exposure derived from the model being twice as high as the ambient air samples. There was also inaccuracy in the assignment of dichloromethane exposure level to each birth because the exposure assignment was made using the predominant value of the isopleth for a census tract.

Two studies have investigated the occurrence of oligospermia among men occupationally exposed to dichloromethane exposure. Kelly (1988) studied 34 men employed in an automotive plant as bonders, finishers, and press operators. These men were self-referred to a health center for a variety of complaints, including neurological symptoms, musculoskeletal symptoms, and shortness of breath. Twenty-six of the men were bonders and eight were finishers or press operators. The job as bonder consisted of dipping hands into an open bucket of dichloromethane and splashing it onto plastic automobile parts. The dichloromethane exposure for bonders averaged 68 ppm with a range of 3.3–154.4 ppm. Eight men, all of whom were bonders, reported symptoms of testicular and epididymal tenderness, with confirmation on medical exam.

They ranged in age from 20 to 47 years old and had been bonders for up to 2.9 years. The COHb levels for the eight workers with genital symptoms ranged from 1.2 to 17.3%, with an average of 6.9% anywhere from 4 to 90 hours postexposure. The COHb levels for the two men who smoked were among the highest, namely 7.3 and 17.3%. Four of the eight workers agreed to provide semen samples; their sperm counts were 2–26 × 10⁶/cm³. The authors stated that men with sperm counts as low as 25 × 10⁶/cm³ may still be fertile, but none of these men had had any children since working with dichloromethane despite not using contraceptives. There was one miscarriage. All four men reported dipping their hands into open buckets of dichloromethane without any protective equipment, and two men reported feeling dizzy, giddy, and high at work.

Based on the results of the Kelly (1988) case report, Wells et al. (1989) planned to do a study of oligospermia among 20 exposed workers and 20 unexposed workers to dichloromethane. The exposed workers were unvasectomized men who had worked for the 3 months prior to recruitment in furniture stripping shops. Eleven men were recruited from among 14 eligible workers at six different shops where dichloromethane was utilized. Names of acquaintances of the exposed were solicited as potential referents. Only one exposed man provided any names. Therefore, the study was redirected as a case report on the 11 exposed men. The mean TWA dichloromethane exposure was 122 ppm (range 15–366 ppm) with a mean COHb of 5.8% (range 2.2–13.5%). The mean COHb for smokers, 10.2% (range 8.1–13.5), was higher than for nonsmokers, 3.9% (range 2.2–5.9), and the nonsmoker levels were higher than the 2% level considered to be the upper limit of normal in nonsmoking populations. The mean sperm count was 54 × 10⁶/cm³ (range 23–128 × 10⁶/cm³) compared to a population value of 47 × 10⁶/cm³ for the same geographic area based on samples analyzed at the same laboratory. Using the standard definition for oligospermia of 20 × 10⁶/cm³, none of the 11 workers had oligospermia.

4.1.2.9. Summary of Noncancer Studies

The clinical and workplace studies of noncancer health effects of dichloromethane exposure have examined markers of disease and specific clinical endpoints relating to cardiac, neurological disease, hepatic function, and reproductive health.

Cardiac effects. The effect of dichloromethane on the formation of COHb (Stewart et al., 1972b) raised concerns about potential risk of cardiovascular damage. To date, there is little evidence of cardiac damage related to dichloromethane exposure in the cohort studies of dichloromethane-exposed workers that examined ischemic heart disease mortality risk (Hearne and Pifer, 1999; Tomenson et al., 1997; Gibbs et al., 1996; Lanes et al., 1993) or in two small cardiac monitoring studies (Ott et al., 1983d; Cherry et al., 1981). However, limitations in these studies should be noted, including the healthy worker effect and the absence of data pertaining to

workers who died before the establishment of the analytic cohort (Gibbs et al., 1996; Gibbs, 1992).

Neurological effects. The acute effects of dichloromethane exposure on neurological function seen in numerous case reports have also been established in experimental studies in humans (Putz et al., 1979; Winneke, 1974; Stewart et al., 1972a, b). Relatively less is known about the long-term effects of chronic exposures in humans. Some data from studies of workers suggest that the effects of dichloromethane are relatively short-lived. For example, in the study by Cherry et al. (1983) of 56 exposed and 36 unexposed workers, alterations in mood or in digit substitution test results were seen during the course of a work shift but were not seen at the beginning of a shift. No difference in four neurological symptoms was seen in an analysis of exposed workers (average exposure 475 ppm, ≥ 10 -year duration) and an unexposed comparison group by Soden (1993). Other data suggest an increase in prevalence of neurological symptoms among workers (Cherry et al., 1981) and possible detriments in attention and reaction time in complex tasks among retired workers (Lash et al., 1991). These latter two studies are limited by the small sample size. Thus, Cherry et al. (1981) and Lash et al. (1991) have low power for detecting statistically significant results and consequently should not be interpreted as definitive analyses showing no effects. Rather, these analyses provide some evidence of an increased prevalence of neurological symptoms among workers with average exposures of 75–100 ppm (Cherry et al., 1981) and long-term effects on specific neurological measures (i.e., attention and reaction time) in workers whose past exposures, at least for part of their work history, were in the 100–200 ppm range (Lash et al., 1991). The increased risk of suicide (approximately a twofold increased risk) seen in two of the worker cohort studies (Hearne and Pifer, 1999; Gibbs, 1992) is an additional indication of potential neurological consequences of dichloromethane exposure. Adequate studies addressing these specific issues are not available. Thus, given the suggestions from the currently available studies, the statement that there are no long-term neurological effects of chronic exposures to dichloromethane cannot be made with confidence.

Hepatic effects. Three studies provide data pertaining to markers of hepatic damage (i.e., serum enzymes and bilirubin levels) (Soden, 1993; Kolodner et al., 1990; Ott et al., 1983c). Two of these studies were based in the Rock Hill, South Carolina, cellulose triacetate fiber plant (Soden, 1993; Ott et al., 1983c), with the most recent of the studies focusing on workers with >10 years duration in a high exposure area (average exposure estimated as 475 ppm). There is some evidence of increasing levels of serum bilirubin with increasing dichloromethane exposure in Ott et al. (1983c) and Kolodner et al. (1990), but there are no consistent patterns with respect to the other hepatic enzymes examined (serum γ -glutamyl transferase, serum AST, serum ALT). These studies do not provide clear evidence of hepatic damage in dichloromethane-exposed workers, to the extent that this damage could be detected by these serologic measures; however,

these data are limited and thus the absence, presence, or extent of hepatic damage is not known with certainty.

Immune effects. Only limited and somewhat indirect evidence pertaining to immune-related effects of dichloromethane in humans is available. No risk was seen in the broad category of infectious and parasite-related mortality reported by Hearne and Pifer (1999), but there was some evidence of an increased risk for influenza and pneumonia-related mortality at two cellulose triacetate fiber production work sites in Maryland and South Carolina (Gibbs, 1992). In the Maryland facility, an increased risk of cervical cancer was seen among the 938 female workers, with an SMR of 3.0 (95% CI 0.96–6.9) in the 50–100 ppm group and 5.4 (95% CI 0.13–30.1) in the 350–700 ppm group (Gibbs et al., 1996). Cervical cancer is viral mediated (human papilloma virus), and immunosuppression is a risk factor for development of this disease, as seen by the increased risk in immunocompromised patients and people taking immunosuppressant medications (Leitao et al., 2008; Ognenovski et al., 2004).

Reproductive effects. Studies pertaining to various reproductive effects and dichloromethane exposure from workplace settings (Wells et al., 1989; Kelly, 1988; Taskinen et al., 1986) or environmental settings (Bell et al., 1991) have examined possible associations with spontaneous abortion (Taskinen et al., 1986), low birth weight (Bell et al., 1991), and oligospermia (Wells et al., 1989; Kelly, 1988). Of these, the data pertaining to spontaneous abortion provide the strongest evidence of an adverse effect of dichloromethane exposure, particularly with respect to the case-control study in which the strongest association was seen specifically with the higher frequency category of dichloromethane exposure. However, it is a small study (44 cases, 130 controls) with limited quantitative exposure assessment and multiple exposures (although the association seen with dichloromethane was among the highest seen among the solvents) and so cannot be considered to firmly establish the role of dichloromethane in induction of miscarriage. Nevertheless, the high exposure scenario, including the potential for substantial dermal exposure in the study of Kelly (1988), also suggests the potential for adverse male reproductive effects.

4.1.3. Cancer Studies

4.1.3.1. Identification and Selection of Studies for Evaluation of Cancer Risk

Twelve epidemiologic studies of cancer risk were identified and included in this evaluation: four cohorts for which the primary solvent exposure was to dichloromethane (two in film production settings and two in cellulose triacetate fiber production), one large cohort of civilian employees at a military base with exposures to a variety of solvents but that included an assessment specifically of dichloromethane exposure, and seven case-control studies of specific cancers with data on dichloromethane exposure. One additional study (Ott et al., 1985), a cohort

of 1,919 men employed at Dow Chemical facilities, was identified but was not included in the summary. The analysis was based on exposure to a combined group of chlorinated methanes (e.g., carbon tetrachloride, chloroform, methyl chloride, and dichloromethane), and it was not possible from the data presented to assess the individual effects of dichloromethane.

4.1.3.2. *Description of the Selected Studies*

In this section, the study setting, methods (including exposure assessment techniques), results pertaining to incidence of mortality from specific cancers, and a brief summary of primary strengths and limitations are summarized for each of the 12 selected studies. When two papers of the same cohort were available, the results from the longer period of follow-up are emphasized in the summary. Information from earlier reports is used when these reports contain more details regarding working conditions, study design, and exposure assessment. The description of individual studies is followed by a summary of the evidence available from these studies relating to specific types of cancer.

4.1.3.3. *Cellulose Triacetate Film Base Production Cohorts*

4.1.3.3.1. *Cellulose triacetate film base production—Rochester, New York (Eastman Kodak).*

Friedlander et al. (1978) reported a cohort mortality study of workers in an Eastman Kodak facility in Rochester, New York. This study was expanded and extended several times during the next 20 years (Hearne and Pifer, 1999; Hearne et al., 1990, 1987). The latest analysis provided data on two overlapping cohorts. The first cohort (Cohort 1) consisted of 1,311 male workers employed in the roll coating division ($n = 1,070$) or the dope and distilling departments ($n = 241$) of the Eastman Kodak facility in Rochester, New York. Men who began working in these areas after 1945 and were employed in these areas for at least 1 year (including seasonal or part-time work that equaled 1 full-time year equivalent) from 1946 to 1970 were included. Follow-up time was calculated from the end of the first year of employment in the study area through December 31, 1994. The mean duration of work in Cohort 1 was 17 years. The total number of person-years of follow-up was 46,112, and the mean duration of follow-up was 35.2 years (range 25–49 years). The second cohort (Cohort 2) included 1,013 male workers in the roll coating division who were employed for at least 1 year in this division between 1964 and 1970. Follow-up time was calculated from January 1, 1964, for those who were employed there before 1964, or the date of first employment in the roll coating division for those who began in 1964 or later. Follow-up continued through December 31, 1994. The mean duration of work in Cohort 2 was 24 years. Total follow-up time was 26,251 person-years, and the mean duration of follow-up was 25.9 years (range 25–31 years). Cohort 2 was the focus of previous analyses by Friedlander et al. (1978) and Hearne et al. (1990, 1987).

For both cohorts, causes of death were based on the underlying causes of death recorded on the death certificates, which were routinely obtained by the company for the processing of life

insurance claims. The expected number of deaths was calculated using appropriate age-, sex-, calendar time-, and cause-specific death rates for men in New York State (excluding New York City). In addition, another referent group was also used in the analysis of the second cohort. This other referent was based on the age-, sex-, calendar time-, and cause-specific death rates of other hourly male workers employed at the Eastman Kodak plant in Rochester, New York. (An internal referent group was also described for Cohort 1, but data for that analysis were not presented.)

Dichloromethane was first used in the film production process at the Eastman Kodak facility around 1944 (Hearne et al., 1987). Cellulose triacetate was dissolved in dichloromethane and then cast into a thin film onto revolving wheels. The film was then cured by circulating hot air in the coating machines, and the solvent was recovered and redistilled. 1,2-Dichloropropane and 1,2-dichloroethane were also used as solvents from the 1930s to the 1960s, but dichloromethane was predominant (ratio 17:2:1 for dichloromethane:1,2-dichloropropane:1,2-dichloroethane in general workplace air measurements) (Hearne et al., 1987).

The exposure assessment in the Rochester, New York, Eastman Kodak cohort studies was based on employment records (start and stop dates for specific jobs in the relevant areas of the company) in combination with air monitoring data used to estimate the exposure level for a given job, location, and time period (Hearne et al., 1987). Air monitoring began in the 1940s, but few data are available before 1959. In the most recent update (Hearne and Pifer, 1999), more than 1,500 area and 2,500 breathing zone air samples were used in the exposure assessment process. Reductions in exposures in the dope department and the distilling department began after 1965. The highest exposure jobs were operator and maintenance workers (dope department) and filter washing and waste operator (distilling department), with estimated 8-hour TWA exposures of 100–520 ppm between 1946 and 1985. There was little change in estimated exposures for jobs in the roll coating division from the 1940s through 1985, but some reduction was seen from 1986 to 1994. The mean 8-hour TWA exposures were 39 ppm for Cohort 1 and 26 ppm for Cohort 2.

These data were used to estimate a cumulative exposure index (i.e., the summation across all jobs held by an individual of the product of the average dichloromethane concentration as ppm and the duration of employment in that job). The authors refer to this as a “career exposure index.” Additional adjustment in these estimates was made for respiratory protection, but the details of this adjustment were not described. For Cohort 1, the cumulative exposure categories used in exposure-effect analyses were <150, 150–349, 350–799, and \geq 800 ppm. For Cohort 2, the cumulative exposure categories were <400, 401–799, 800–1,199, and \geq 1,200 ppm. The cut points were chosen to produce an approximately equal number of expected total deaths in these categories. Two different methods to calculate expected number of deaths within each exposure category were used for each cohort analysis. For Cohort 1, an internal comparison was made based on the distribution of person-years within each exposure category, and an external

comparison was made applying New York State mortality rates. For Cohort 2, the internal comparison using the distribution of person-years within each exposure category was also used, but the external comparison was based on mortality rates in other hourly workers at the Rochester, New York, Eastman Kodak work site.

There was no increased risk of mortality for all sites of cancer or for lung cancer in either cohort analysis (Table 4-4). Data pertaining to smoking history, obtained from a survey of workers in the New York film production facility, indicate that smoking rates were similar in the exposed group, the internal comparison group, and the general population; therefore, it is unlikely that differences in smoking could be masking an effect of dichloromethane.

Table 4-4. Mortality risk in Eastman Kodak cellulose triacetate film base production workers, Rochester, New York

Cancer type	Cohort 1: 1,311 men employed 1946–1970, followed through 1994				Cohort 2: 1,013 men employed 1964–1970, followed through 1994						
	New York referent group				New York referent group				Kodak referent group		
	Obs ^b	Exp ^b	SMR	95% CI	Obs	Exp	SMR	95% CI	Exp	SMR	95% CI
Cancer, all sites	93	105.8	0.88	0.71–1.08	91	102.0	0.89	0.72–1.10	94.7	0.96	0.77–1.18
Liver ^a	1	2.4	0.42	0.01–2.36	1	2.4	0.42	0.01–2.33	1.8	0.55	0.01–3.07
Pancreas	5	5.5	0.92	0.30–2.14	8	5.3	1.51	0.65–2.98	5.1	1.55	0.67–3.06
Lung ^a	27	36.0	0.75	0.49–1.09	28	34.2	0.82	0.55–1.19	31.3	0.89	0.59–1.29
Brain ^a	6	2.8	2.16	0.79–4.69	4	2.1	1.88	0.51–4.81	2.7	1.46	0.39–3.75
Lymphatic system	5	6.6	0.75	0.24–1.78	6	5.7	1.06	0.39–2.30	5.7	1.05	0.38–2.28
Non-Hodgkin's	2	4.1	0.49	0.06–1.76	3	3.5	0.85	0.17–2.50	3.6	0.84	0.17–2.46
Hodgkin's	2	1.1	1.82	0.20–6.57	2	0.6	3.13	0.35–11.30	0.9	2.23	0.25–8.05
Multiple myeloma	1	1.5	0.68	0.01–3.79	1	1.5	0.65	0.01–3.62	1.3	0.79	0.01–4.39
Leukemia	8	3.9	2.04	0.88–4.03	6	3.5	1.73	0.63–3.76	4.4	1.37	0.50–2.98

^aLiver includes liver and biliary duct; lung includes lung, trachea, and bronchus; brain includes brain and CNS.

^bObs = number observed deaths, Exp = number of expected deaths.

Source: Hearne and Pifer (1999).

The only specific sites for which there were increased SMRs in both cohorts were brain and CNS cancer, Hodgkin's lymphoma, and leukemia. Pancreatic cancer mortality risk was increased in Cohort 2 but not in Cohort 1. None of these associations were statistically significant, and the Hodgkin's lymphoma observations were based on a total of only four cases in both cohorts combined, and so were imprecise. Within Cohort 2, there was little difference in results for most sites using the different referent groups, but the point estimates for the SMRs for brain and CNS cancer, Hodgkin's lymphoma, and leukemia were somewhat higher using the New York State referent group compared with the internal Eastman Kodak referent group. An attenuation of the dichloromethane association seen in the analyses using the internal Kodak referent group would be expected if the risk of specific cancers was increased in this comparison group, possibly because of other workplace exposures.

The authors presented the exposure-effect analysis based on the estimated cumulative dichloromethane exposure groups for all sites of cancer, pancreatic cancer, lung cancer, brain cancer, and leukemia (Table 4-5).

Table 4-5. Mortality risk by cumulative exposure in Eastman Kodak cellulose triacetate film base production workers, Rochester, New York

Cohort, cancer, referent group	SMRs (number of observed deaths)					
	Cohort 1 ^a					
Cumulative exposure (ppm yrs)	<150		150–349	350–799		≥800
Cancer, all sites						
Internal New York	0.81 (20) 0.67		1.02 (19) 0.93	1.10 (28) 0.95		1.07 (26) 1.00
Pancreas						
Internal New York	0.74 (1) 0.68		0.00 (0) 0.00	0.77 (1) 0.65		2.34 (3) 2.18
Lung ^b						
Internal New York	0.78 (5) 0.52		1.07 (6) 0.90	1.25 (9) 0.86		0.90 (7) 0.77
Brain ^b						
Internal New York	0.58 (1) 1.10		0.78 (1) 1.77	1.65 (3) 3.99		0.85 (1) 1.78
Leukemia						
Internal New York	0.83 (2) 1.61		0.00 (0) 0.00	0.48 (1) 0.98		2.73 (5) 5.79
Cohort 2 ^c						
Cumulative exposure (ppm yrs)		<400		400–799	800–1,199	≥1,200
Cancer, all sites						
Internal New York		0.89 (18) 0.76		0.96 (33) 0.93	1.11 (23) 1.13	1.08 (17) 1.12
Pancreas						
Internal New York		2.58 (4) 2.86		0.00 (0) 0.00	0.95 (2) 1.83	1.43 (2) 2.67
Lung ^b						
Internal New York		0.95 (6) 0.80		1.15 (12) 1.00	0.94 (6) 0.89	0.82 (4) 0.79
Brain ^b						
Internal New York		0.00 (0) 0.00		1.13 (2) 2.02	1.37 (1) 1.75	1.49 (1) 2.50
Leukemia						
Internal New York		0.00 (0) 0.00		0.84 (2) 1.26	0.75 (1) 1.10	2.70 (3) 4.84

^aCohort 1: 1,311 men employed 1946–1970 in the roll coating division, dope department, or distilling department, followed through 1994; mean exposure (cumulative exposure yrs) 66, 244, 543, and 1,782 ppm-yrs in the four dose groups, respectively.

^bLung includes lung, trachea, and bronchus; brain includes brain and CNS.

^cCohort 2: 1,013 men employed 1964–1970 in the roll coating division, followed through 1994; mean exposure (cumulative exposure yrs) 168, 581, 981, and 1,670 ppm-yrs in the four dose groups, respectively.

Source: Hearne and Pifer (1999).

There is no evidence of an exposure-effect for all site cancer mortality or lung cancer mortality risk. The relatively sparse number of deaths for the other specific cancer types makes it difficult to interpret the data. The patterns for pancreatic cancer differ between the two

cohorts, with increased risk at the higher dose in Cohort 1 and a U-shaped curve seen in Cohort 2. For brain cancer mortality, a higher SMR was seen in the groups with cumulative exposure levels of ≥ 800 ppm-years compared with lower exposure groups. For leukemia in both cohorts, an increased mortality risk is seen in the highest exposure group (mean approximately 1,700 ppm-years).

A strength of the Eastman Kodak cohort studies was the sampling data for dichloromethane that allowed an assessment of each worker's exposure using the monitoring data and the worker's job history, making exposure-effect analyses possible. Follow-up of the vital status of the cohort was $>99\%$ (Hearne and Pifer, 1999). There was also some information on smoking history for workers in the plant, based on a survey conducted in 1986 (Hearne et al., 1987). A difficulty in interpreting the data, however, is that there was some overlap between the cohorts: 707 of the men were included in both Cohort 1 and Cohort 2. Data are not presented in a way that would allow the reader to eliminate duplicate cases and person-years so that cases are only counted once when examining both cohorts. A strength of the Cohort 1 sampling strategy, compared with that of Cohort 2, is that Cohort 1 is limited to workers who began work at the plant after 1945. These workers would not have had workplace exposure to methanol and acetone, which were used at the plant in the film production process prior to that time. Also, follow-up began with the beginning of employment in the relevant area. In contrast, Cohort 2 was limited to workers who were employed from 1964 to 1970, so exposed workers who left or died before 1964 were not included. The relatively small number of cases with specific low incidence cancers (e.g., brain, leukemia) is also a limitation of the analyses of both of the cohorts in this study. In addition, the exposure levels in both cohorts (mean 8-hour TWA 39 and 26 ppm in Cohorts 1 and 2, respectively) is relatively low compared with values seen in other workplaces, including the cellulose triacetate fiber production cohorts described in Ott et al. (1983a) and Gibbs et al. (1996). Also, the outcome assessment is based on mortality (underlying cause from death certificates) rather than incidence data, and, because the Kodak studies were limited to men, there is no information on risk of breast cancer or other female reproductive cancers.

4.1.3.3.2. Cellulose triacetate film base production—Brantham, United Kingdom (Imperial Chemical Industries). Tomenson et al. (1997) reported the results of a retrospective cohort mortality study of 1,473 men who worked at a film-base production facility in Brantham, England, anytime between 1946 and 1988 in jobs that were considered to have dichloromethane exposure. The start of the follow-up period was not specified by the authors but is likely to have been 1946 or the date of first employment at the plant. Follow-up of the cohort continued through December 31, 1994, and vital status was based on national records (United Kingdom National Health Service Central Register and the Department of Social Security). Causes of death were based on the underlying causes of death recorded on the death certificates. The

expected number of deaths was calculated using age-, sex-, calendar time-, and cause-specific death rates for England and Wales. In addition, a comparison using mortality rates for the local areas (Tendring and Samford) for 1968–1978 and analyses limited to workers who had been employed for at least 3 months were also made, but the results of these analyses were not presented. The mean duration of work in the cohort was 9 years, the total number of person-years was 39,759, and the mean duration of follow-up was 27.0 years (7–49 years).

This facility produced cellulose diacetate film from 1950 to 1988, with other types of films also manufactured beginning in the 1960s. Dichloromethane was the solvent used in this process, and exposure occurred in the production of the triacetate film base and the casting of the film into rolls. The exposure assessment was based on >2,700 personal or air monitoring samples collected since 1975. An exposure matrix was constructed, assigning jobs to 1 of 20 work groups with similar exposure potential for each of four different time periods (before 1960, 1960–1969, 1970–1979, and 1980–1988). For the 1980–1988 period, exposure estimates for specific jobs were based on about 330 personal monitoring samples. For the earlier time periods, information about work tasks and location was used in combination with the information about the number of, use of, speed of, and problems with casting machines at different times from their initial introduction in 1950. The highest exposures were estimated to be in the casting machine operators and cleaners. Lifetime cumulative exposure to dichloromethane was calculated as the product of the mean level of exposure for the assigned work group and the duration of employment in that job summed across all jobs. Three categories of cumulative exposure were used for the analysis of ever-exposed workers: <400, 400–700, and \geq 800 ppm-years. Approximately 30% of the workers in the cohort were classified as “unassigned” for the calculation of exposure group because sufficient information needed to determine exposures (i.e., the location and tasks assigned to laborers and maintenance workers) was not available. The mean 8-hour TWA exposure was estimated at 19 ppm for the cohort.

There was no increased risk of mortality for all sites of cancer (Table 4-6), and the SMRs for most of the specific cancer sites examined (stomach, colon, rectum, liver, pancreas, lung, and prostate) were <1.0. The only specific sites for which there was an increased SMR (i.e., 1.1 or higher) were brain and CNS cancer and leukemia, and these estimates were based on few (less than five) observed cases (Table 4-6). Tomenson et al. (1997) present the exposure-effect analysis based on the estimated cumulative dichloromethane exposure groups for all sites of cancer, pancreatic cancer, and lung cancer, and there is no evidence of an increasing effect with increasing exposure level in these analyses. A formal exposure-effect analysis for brain cancer or leukemia was not presented. However, the authors described two of the brain cancer cases as having “minimal” exposure to dichloromethane (and thus presumably would have been in the <400 ppm-year cumulative exposure group). One case was estimated as having 572 ppm-years cumulative exposure, and the other case was an electrician classified in the unassigned exposure

group. He had worked for 21 years at an exposure level “that was unlikely to have exceeded 15 ppm 8-hour TWA.”

Table 4-6. Mortality risk in Imperial Chemical Industries cellulose triacetate film base production workers, Brantham, United Kingdom: 1,473 men employed 1946–1988, followed through 1994

Cancer type	Observed	Expected ^a	SMR	95% CI
Cancer, all sites	68	104.6	0.65	0.51–0.82
Liver and biliary duct	0	1.5	–	–
Pancreas	3	4.4	0.68	0.14–1.99
Lung, trachea, bronchus	19	41.3	0.46	0.29–0.75
Brain and CNS system	4	2.8	1.45	0.40–3.72
Lymphatic and hematopoietic	6	7.1	0.85	0.31–1.84
Leukemia	3	2.7	1.11	0.23–1.84

^aExpected, calculated from observed and SMR data reported by the authors by using the following formula: $\text{expected} = 100 \times \text{observed} \div \text{SMR}$; SMRs and CIs were not calculated for categories with zero observed cases.

Source: Tomenson et al. (1997).

A strength of this study was the monitoring data available that allowed assignment of cumulative exposure categories for use in exposure-effect analyses. However, 30% (439) of exposed workers had insufficient work histories to determine lifetime cumulative exposure. Air measurements were not available until 1975, and personal measures were not available until 1980. In addition, the duration of exposure was relatively low (mean, 9 years), the mean exposure level was relatively low (mean 8-hour TWA, 19 ppm), and there were very few deaths from specific types of cancer, which limit the statistical power of the study to examine associations among dichloromethane and specific cancers. Other limitations, as were also noted in the Kodak cohort studies, include the use of mortality rather than incidence to define risk, the reliance solely on underlying causes of death from death certificates to classify specific cancer types, and the lack of information on breast cancer risk.

4.1.3.4. Cellulose Triacetate Fiber Production Cohorts

4.1.3.4.1. Cellulose triacetate fiber production—Rock Hill, South Carolina (Hoechst Celanese Corporation). Two cohorts of cellulose triacetate fiber workers have been studied in Rock Hill, South Carolina (Lanes et al., 1993, 1990; Ott et al., 1983a, b), and Cumberland, Maryland (Gibbs et al., 1996; Gibbs, 1992). Workers were exposed to dichloromethane, methanol, and acetone in both facilities.

Ott et al. (1983a, b) conducted a retrospective cohort mortality study of 1,271 acetate fiber production workers (551 men and 720 women) employed at least 3 months from 1954 to 1977 at Dow Chemical Company, Rock Hill, South Carolina. This analysis focused on ischemic

heart disease mortality risk, and there was no presentation of cancer risk. The Rock Hill cohort study was updated through September 30, 1986 (Lanes et al., 1990), and December 31, 1990 (Lanes et al., 1993), and analyses of cancer mortality risks were included in these later reports. Causes of death information was obtained from death certificates with coding based on the underlying and contributing causes (Ott et al., 1983a). The referent used in the updates was the general population of York County, South Carolina, and analyses were adjusted for age, race, gender, and calendar period. Because the results of the mortality risk analyses were similar for both updates, those from the 1993 paper are presented here. The mean duration of work in the cohort was not reported, but 56% worked for <5 years (calculated from Tables 3 and 4 of Ott et al., 1983b). The mean duration of follow-up was 23.6 years in the analysis through 1986 (Lanes et al., 1990) but was not reported in the later paper (Lanes et al., 1993). The 1993 report added approximately 4.25 years of follow-up, which would result in an estimate of approximately 28 years of follow-up for this report.

The Rock Hill, South Carolina, plant began producing cellulose triacetate fiber in 1954. Dichloromethane was used as the solvent for the initial mixing with cellulose triacetate flakes. This mixture was then filtered and transferred to the extrusion area for drying and winding. Air measurements taken in 1977–1978 were assumed to be representative of operations since dichloromethane use began in 1954, based on review of processing operations. The median 8-hour TWA exposures were estimated at 140, 280, and 475 ppm in the low, moderate, and high categories of exposure (Ott et al., 1983a). Employment records provided information on jobs held and dates employed, and this was used in conjunction with the exposure estimates for specific jobs and work areas to classify individual exposures. However, detailed work history information was only available for 475 (37%) of the workers (Lanes et al., 1990), and it is not clear how the exposure assessment was applied to workers with missing job history data.

Methanol was also used in the cellulose triacetate fiber production process, and methanol exposure was estimated as 1/10 that of dichloromethane. Acetone exposure was used in the production of acetate (cellulose diacetate) fiber at an adjacent part of the plant. The exposure to acetone was inversely related to that of dichloromethane, with estimated median 8-hour TWAs of 1,080 ppm acetone in the low dichloromethane exposure group and 110 ppm acetone in the moderate and high dichloromethane groups in the Rock Hill plant (Ott et al., 1983a).

In the latest follow-up (Lanes et al., 1993), there was no increase in mortality risk from cancer (all sites) or from cancer of the lung or pancreas (Table 4-7). The SMR for liver and bile duct cancer, based on four observed cases, was 2.98 (95% CI 0.81–7.63). This was lower than the SMR of 5.75 (95% CI 1.82–13.8) that was reported in the 1990 analysis based on these same four cases but on a shorter follow-up period (and thus lower number of expected cases). Three of these cases were bile duct cancers. This was the first cohort study that included women, and this study provided data on breast cancer risk. There were 3 observed breast cancer deaths compared with 5.59 expected, yielding an SMR of 0.54 (95% CI 0.11–1.57). No data were

provided pertaining to reproductive risk factors (e.g., pregnancy history) for breast cancer among the women in this cohort, so it is difficult to assess whether these potential confounders are likely to have been distributed differently in the cohort compared with the referent group. Information about brain cancer, Hodgkin’s lymphoma, and leukemia (Table 4-7) was not included in this report but was included in the report by Gibbs (1992) (see Table 11 of that report).

Table 4-7. Mortality risk in Hoechst Celanese Corporation cellulose triacetate fiber production workers, Rock Hill, South Carolina: 1,271 men and women employed 1954–1977, followed through 1990

Cancer type	Observed	Expected	SMR ^a	95% CI ^{a,b}
Cancer, all sites	39	47.7	0.82	0.58–1.52
Liver and biliary duct	4	1.34	2.98	0.81–7.63
Pancreas	2	2.42	0.83	0.10–2.99
Lung, trachea, bronchus	13	16.21	0.80	0.43–1.37
Brain and CNS ^c	1	1.5	0.67	0.2–3.71
Hodgkin’s lymphoma ^c	0	0.24	–	–
Leukemia ^c	1	1.11	0.90	0.02–5.0
Breast cancer (women)	3	5.59	0.54	0.11–1.57

^aSMRs and CIs were not calculated for categories with zero observed cases.

^bCIs were calculated from Breslow and Day (1987, Table 2.10).

^cData for brain and CNS cancer, Hodgkin’s lymphoma, and leukemia were reported in Gibbs (1992).

Source: Lanes et al. (1993).

There are a number of limitations in this study including the small size of the cohort, small number of observed cancer deaths, availability of detailed work history information for only 37% of the workers, and use of mortality rather than incidence data. The exposure levels at this plant were high, but the duration of exposure for most of the cohort was relatively short (<5 years). It is the first cohort study, however, that included women and provided information on breast cancer risk.

4.1.3.4.2. Cellulose triacetate fiber production—Cumberland, Maryland (Hoechst Celanese Corporation). Gibbs et al. (1996) studied a cohort of 2,909 cellulose triacetate fiber production workers (1,931 men and 978 women) at a Hoechst Celanese plant in Cumberland, Maryland. This retrospective cohort mortality study included all workers who were employed on or after January 1, 1970, and who worked at least 3 months. This study also included a very small comparison group (256 men, 46 women) that was described as a “0” or “no” exposure group of workers at the plant who worked in jobs that were not considered to have had dichloromethane exposure; totals for this study were 2,187 men and 1,024 women in the exposed and nonexposed groups combined.

The plant closed in 1981, and mortality was followed through 1989. Since 1955, employees of this plant were exposed to dichloromethane, methanol, acetone, and finishing oils used as lubricants. Before 1955, acetone was the only exposure. Industrial hygiene monitoring focusing on dichloromethane, acetone, and methanol began in the late 1960s. Exposure groupings (low = 50–100 ppm and high = 350–700 ppm) were assigned by area in which employees worked. The extrusion and spinning workers and jet wipers were among the high exposure group (300–1,250 ppm 8-hour TWA). The SMR analysis that was reported used Allegany County, Maryland, as the comparison group. Cause of death information was obtained from death certificates, but the authors did not state whether they used underlying or underlying and contributing cause of death information. The mean duration of work in the cohort was not reported. The total follow-up period included 49,828 person-years (16,292 years in the high exposure group and 33,536 years in the low exposure group), and the mean duration of follow-up was 17.2 years (range 8–20 years). These data were found in Hearne and Pifer (1999, Table 7).

There was little evidence of an increase in mortality risk from cancer (all sites) or from cancer of the liver and bile duct, pancreas, or brain in men or women (Table 4-8). An increasing risk with increasing exposure level was seen for prostate cancer mortality in men. The *p*-value for the trend was not given, but the authors describe it as a “nonstatistically significant dose-response relationship.” A statistically significant SMR for prostate cancer death was seen in the 350–700 ppm group when latency (at least 20 years since first exposure) was included in the analysis (SMR = 2.08, *p* < 0.05). Cervical cancer mortality risk was increased, but the small number of cases in the high exposure group did not allow a precise assessment of the pattern with respect to exposure level. There was no increased risk of breast cancer.

Table 4-8. Cancer mortality risk in Hoechst Celanese Corporation cellulose triacetate fiber production workers, Cumberland, Maryland: 2,909 men and women employed 1970–1981, followed through 1989

Cancer type, exposure level ^a	Men (n = 1,931)				Women (n = 978)			
	Obs ^b	Exp ^b	SMR	95% CI ^c	Obs ^b	Exp ^b	SMR	95% CI ^c
Cancer, all sites	121				42			
50–100 ppm	64	70.0	0.91	0.70–1.2	37	44.79	0.83	0.58–1.1
350–700 ppm	57	75.6	0.75	0.57–0.98	5	4.61	1.1	0.35–2.5
Liver	2				0			
50–100 ppm	1	1.33	0.75	0.02–4.2	0	1.04		–
350–700 ppm	1	1.24	0.81	0.02–4.5	0	0.10		–
Pancreas	3				1			
50–100 ppm	2	2.24	0.89	0.1–3.2	1	1.73	0.58	0.01–3.2
350–700 ppm	1	2.90	0.35	0.01–1.9	0	0.18		–
Lung	35				11			
50–100 ppm	20	25.7	0.78	0.48–1.2	9	8.24	1.1	0.50–2.1
350–700 ppm	15	27.3	0.55	0.31–0.91	2	0.87	2.3	0.28–8.3
Brain ^a	2				2			
50–100 ppm	1	1.88	0.53	0.01–2.96	2	0.66	3.1	0.37–10.9
350–700 ppm	1	1.94	0.52	0.01–2.87	0	0.07		
Hodgkin's ^a								
50–100 ppm	1	0.4	2.5	0.06–13.9	0	0.23		
350–700 ppm	0	0.41			0	0.02		
Leukemia ^a								
50–100 ppm	4	2.14	1.9	0.51–4.8	0	1.25		
350–700 ppm	1	2.28	0.44	0.01–2.4	0	0.13		
Prostate	22				Not applicable			
50–100 ppm	9	6.41	1.4	0.64–2.7				
350–700 ppm	13	7.26	1.8	0.95–3.1				
Cervical	Not applicable				6			
50–100 ppm					5	1.69	3.0	0.96–6.9
350–700 ppm					1	0.19	5.4	0.13–30.1
Breast ^a	0				10			
50–100 ppm	0	0.03			9	9.8	0.92	0.42–1.7
350–700 ppm	0	0.02			1	1.07	0.93	0.02–5.2

^aData for brain and CNS cancer, Hodgkin's lymphoma, leukemia, and breast cancer reported in Gibbs (1992).

^bObs = number of observed deaths, Exp = number of expected deaths. Referent group = Allegany County, Maryland. SMRs and CIs were not calculated for categories with zero observed cases.

^cCIs were calculated from Breslow and Day (1987, Table 2.10).

Sources: Gibbs et al. (1996); Gibbs (1992).

A primary limitation of this study is that workers who were exposed before 1970 but were not working at the plant in 1970 were not included in the cohort. The authors had

attempted to create a cohort of all workers who were employed on or after January 1, 1954, but problems with the completeness of the personnel file made it impossible to use this study design. From what the author (Gibbs, 1992) was able to determine, the records of workers who had died, left the company, or retired before the mid to late 1960s (when a new personnel system was developed) were not available. Additional limitations include the small size of the cohort, small number of observed cancer deaths, and use of mortality (death certificate) data. This is particularly problematic for cancers with relatively high survival rates (such as prostate cancer and cervical cancer), since incidence rates are not estimated well by mortality rates in this situation.

4.1.3.5. Solvent-Exposed Workers—Hill Air Force Base, Utah

Spirtas et al. (1991) and Blair et al. (1998) evaluated exposure to dichloromethane in relation to mortality risk in successive retrospective cohort studies of 14,457 civilian workers employed at Hill Air Force Base in Utah for at least 1 year from 1952 to 1956. The analysis was limited to the workers that were white or who had missing data on race, resulting in a sample size of 14,066 (10,461 men and 3,605 women). Spirtas et al. (1991) examined mortality through 1982 (3,832 deaths), and Blair et al. (1998) updated mortality through 1990 (4,195 deaths). The underlying and contributing causes of death information from death certificates was used to classify cause-specific mortality. SMRs were calculated by using mortality rates from the Utah population, and an internally standardized life table method was used to adjust for age at entry into the cohort and competing causes of death. In the Blair et al. (1998) analysis, adjusted relative risks (rate ratios) were estimated from a Poisson regression analysis with unexposed workers as the referent. The mean duration of work was not reported. In the analysis through 1982 (Spirtas et al., 1991), there were 22,770 person-years of follow-up in men and 3,091 person-years of follow-up in women who were classified as exposed to dichloromethane. The total number of workers classified as exposed to dichloromethane was 1,222 (Stewart et al., 1991), which would yield an estimated mean of approximately 21 years of follow-up through 1982. The total number of person-years included in the later report (Blair et al., 1998), with the addition of 8 more years of follow-up, was not reported but would be expected to increase the mean follow-up time to approximately 29 years.

Two industrial hygienists developed the exposure assessment based on walkthrough surveys, interviews with management and labor representatives, review of historical records, job descriptions, monitoring data and other information pertaining to chemicals used, and organization of the work site (Blair et al., 1998; Spirtas et al., 1991). Each worker was assigned exposure by using information on the worker's job history, which included job titles, department codes, and dates of employment. The most detailed exposure assessment was done for trichloroethylene, the primary focus of the study. Dichloromethane, 1 of 25 other exposures analyzed, was classified as a dichotomous exposure (ever exposed, never exposed).

Blair et al. (1998) presented the mortality risk for three specific cancers in relation to 15 of the 25 chemicals classified as dichotomized exposures. The rate ratios for non-Hodgkin's lymphoma and multiple myeloma in relation to dichloromethane in men were 3.0 (95% CI 0.9–10.0, based on six observed cases) and 3.4 (95% CI 0.9–13.2, based on five observed cases), respectively. These rate ratios (particularly those for multiple myeloma) were considerably higher than the rate ratios for any of the other chemicals examined in which the next highest observed rate ratio was 1.8 for Freon. No cases of either of these cancers were observed in women with dichloromethane exposure, but the rate ratio for breast cancer in these women was 3.0 (95% CI 1.0–8.8, based on four observed cases). Associations of similar magnitude (rate ratios of 3.0–4.0) were also seen among breast cancer and some other exposures (Freon, solder flux, isopropyl alcohol, and trichloroethane).

This is the largest of the cohort studies that were identified that included women and specifically reported data pertaining to breast cancer risk. The major limitation of this study is that the exposure assessment for dichloromethane was based on a dichotomized classification. In addition, exposure to many different types of solvents was common; thus, it is difficult to completely separate the effects of individual exposures. Some aspects of reproductive history, such as age at first pregnancy, are known risk factors for breast cancer. Reproductive history was not included in this analysis, but Blair et al. (1998) noted that it is unlikely that these factors would confound the results of a few specific chemicals, since the referent group was an internal group within the cohort (and thus would be expected to be similar in terms of socioeconomic status) and there was no association overall between solvent exposures and breast cancer mortality.

4.1.3.6. Case-Control Studies of Specific Cancers and Dichloromethane

Seven site-specific cancer case-control studies included dichloromethane as an exposure of interest. These studies involve six cancer sites: brain and CNS (Cocco et al., 1999; Heineman et al., 1994), breast (Cantor et al., 1995), kidney (Dosemeci et al., 1999), pancreas (Kernan et al., 1999), rectum (Dumas et al., 2000), and childhood leukemia (Infante-Rivard et al., 2005). A synopsis of cohort studies in humans is provided in Table 4-9.

Table 4-9. Summary of cohort studies of cancer risk and dichloromethane exposure

Reference and Cohort	Total n, exposure level^a and duration, length of follow-up	Inclusion criteria^b	Exposure assessment; outcome assessment	Results^c
Hearne and Pifer (1999) Cellulose triacetate film base production; New York Cohort 1	n = 1,311 men Mean, 39 ppm mean duration, 17 yr mean follow-up, 35 yr	Began working after 1945; worked at least 1 yr	Work history (job records) and personal/air monitoring; death certificate (underlying causes)	See Table 4-4. Brain cancer SMR 2.16 (95% CI 0.79–4.69); leukemia SMR 2.04 (95% CI 0.88–4.03). SMRs <1.0 for lung cancer, liver cancer, and pancreatic cancers
Cohort 2	n = 1,013 men mean, 26 ppm mean duration, 24 yr mean follow-up, 26 yr	Employed at least 1 yr between 1964 and 1970 (potential exposure began 1946)	Work history (job records) and personal/air monitoring; death certificate (underlying causes)	See Table 4-4. Results similar to Cohort 1 except for pancreatic cancer, SMR 1.55 (95% CI 0.67–3.06)
Tomenson et al. (1997) Cellulose triacetate film base production; United Kingdom	n = 1,473 men mean, 19 ppm mean duration, 9 yr mean follow-up, 27 yr	Employed anytime between 1946 and 1988	Work history (job records) and personal/air monitoring; death certificate (underlying causes)	See Table 4-6. Brain cancer SMR 1.45 (95% CI 0.40–3.72). Lung cancer SMR 0.46 (95% CI 0.29–0.75)
Lanes et al. (1993) Cellulose triacetate fiber production; South Carolina	n = 551 men and 720 women (total n = 1,271); median 140, 280, and 475 ppm in low, moderate, and high, respectively; 56% <5 yr work duration; mean follow-up, ~28 yr	Worked at least 3 mo in the preparation or extrusion areas from 1954 to 1977	Job history data and personal/air monitoring of specific areas (but job history data available for 37%); death certificate (underlying and contributing causes)	See Table 4-7. Liver cancer SMR 2.98 (95% CI 0.81–7.63), estimate from earlier follow-up SMR 5.75 (95% CI 1.82–13.8); lung cancer SMR 0.80 (95% CI 0.43–1.37)
Gibbs et al. (1996) Cellulose triacetate fiber production; Maryland	n = 1,931 men and 978 women (total n = 2,909); 50–100 ppm in low and 350–700 ppm in high exposure; duration not reported; mean follow-up 17 yr	Employed on or after January 1, 1970, for at least 3 mo (potential exposure began 1955)	Work history (job records) and personal/air monitoring; death certificate (fields used not stated)	See Table 4-8. Increasing risk across exposure groups seen for prostate cancer and cervical cancer. In men, SMRs ~1.0 or <1.0 for lung cancer and, combining exposure groups, leukemia. In women, SMRs ~1.0 or <1.0 for breast cancer and, combining exposure groups, lung cancer

(Table 4-9 continues on next page)

Table 4-9. Summary of cohort studies of cancer risk and dichloromethane exposure

Reference and Cohort	Total n, exposure level^a and duration, length of follow-up	Inclusion criteria^b	Exposure assessment; outcome assessment	Results^c
Blair et al. (1998) Air Force Base, Utah	n = 10,461 men and 3,605 women (total n = 14,066) ^d dichotomized (yes, no) exposure duration not reported mean follow-up ~29 yr	Employed at least 1 yr from 1952 to 1956 (potential exposure began 1939)	Work history (job records) and industrial hygiene assessment based on work site review (dichotomized exposure); (underlying and contributing causes)	See Section 4.1.3.5. In men, non-Hodgkin's lymphoma RR 3.0 (95% CI 0.9–10.0) and multiple myeloma RR 3.4 (95% CI 0.9–13.2). In women, breast cancer RR 3.0 (95% CI 1.0–8.8)

^a8-hr TWA.

^bIf dichloromethane was used at the plant before the first date of entrance into the cohort, the yr that potential exposure began is noted.

^cResults for all studies are described as SMR and 95% CI except for Blair et al. (1998), in which results are presented as RR (relative risks) (95% CI). Results are presented cancers based on a minimum of four observed cases. More comprehensive information, when available for other cancers, is shown in the summary tables for each study.

^dIncludes whites and unknown race.

4.1.3.6.1. Case-control studies of brain cancer. Heineman et al. (1994) studied the association between astrocytic brain cancer (International Classification of Diseases 9th ed. [ICD-9] codes 191, 192, 225, and 239.7) and occupational exposure to chlorinated aliphatic hydrocarbons. Cases were identified by using death certificates from southern Louisiana, northern New Jersey, and the Philadelphia area. This analysis was limited to white males who died between 1978 and 1981. Controls were randomly selected from the death certificates of white males who died of causes other than brain tumors, cerebrovascular disease, epilepsy, suicide, and homicide. The controls were frequency matched to cases by age, year of death, and study area.

Next of kin were successfully located for interview for 654 cases and 612 controls, which represents 88 and 83% of the identified cases and controls, respectively. Interviews were completed for 483 cases (74%) and 386 controls (63%). There were 300 cases of astrocytic brain cancer (including astrocytoma, glioblastoma, mixed glioma with astrocytic cells). The ascertainment of type of cancer was based on review of hospital records, which included pathology reports for 229 cases and computerized tomography reports for 71 cases. After the exclusion of 66 controls with a possible association between cause of death and occupational exposure to chlorinated aliphatic hydrocarbons (some types of cancer, cirrhosis of the liver), the final analytic sample consisted of 300 cases and 320 controls.

In the next-of-kin interviews, the work history included information about each job held since the case (or control) was 15 years old (job title, description of tasks, name and location of company, kinds of products, employment dates, and hours worked per week). Occupation and industry were coded based on four-digit Standard Industrial Classification and Standard Occupational Classification (Department of Commerce) codes. The investigators developed matrices linked to jobs with likely exposure to dichloromethane, five other chlorinated aliphatic hydrocarbons (carbon tetrachloride, chloroform, methyl chloroform, tetrachloroethylene, and trichloroethylene), and organic solvents (Gomez et al., 1994). This assessment was done blinded to case-control status. Exposure was defined as the probability of exposure to a substance (the highest probability score for that substance among all jobs), duration of employment in the exposed occupation and industry, specific exposure intensity categories, average intensity score (the three-level semiquantitative exposure concentration assigned to each job multiplied by duration of employment in the job, summed across all jobs), and cumulative exposure score (weighted sum of years in all exposed jobs with weights based on the square of exposure intensity [1, 2, 3] assigned to each job). Secular trends in the use of specific chemicals were considered in the assignment of exposure potential. Exposures were lagged 10 or 20 years to account for latency. Thus, this exposure assessment procedure was quite detailed.

Adjusting for age and study area, the OR for the association between any exposure to dichloromethane and risk of astrocytic brain cancer was 1.3 (95% CI 0.9–1.8). There was a statistically significant trend ($p < 0.05$) with increasing probability of exposure to dichloromethane with an OR = 1.0 (95% CI 0.7–1.6) for low probability, OR = 1.6 (95% CI 0.8–

3.0) for medium probability, and OR = 2.4 (95% CI 1.0–5.9) for high probability compared with the referent group of unexposed men. An increased risk with higher duration of exposure was also observed with OR = 1.7 (95% CI 0.9–3.6) for ≥ 21 years of work in exposed jobs for all exposed workers and OR = 6.1 (95% CI 1.1–43.8) for the combination of ≥ 21 years of work in a high probability of exposure job. Similar results were seen in additional analyses controlling for age, study area, employment in electronics occupations and industries, and exposure to carbon tetrachloride, tetrachloroethylene, and trichloroethylene. There was also evidence of an association between astrocytic brain cancer risk and dichloromethane exposure, based on the average intensity score, with an OR = 1.1 (95% CI 0.7–1.7) for the low-medium intensity group and an OR = 2.2 (95% CI 1.1–4.1) for the high intensity group, and trend p -value < 0.05. The combination of high intensity and high duration (≥ 21 years) was strongly associated with risk (OR = 6.1 [95% CI 1.5–28.3]), and a weaker association (OR = 1.4 [95% CI 0.6–3.2]) was seen for high intensity and shorter duration (2–20 years). The association between cumulative exposure score (low, medium, and high) and astrocytic brain cancer risk was nonlinear (ORs of 0.9, 1.9, and 1.2 in the low, medium, and high exposure categories, respectively).

The strengths of this case-control study include a large sample size, detailed work histories (including information not just about usual or most recent industry and occupation but also about tasks and products for all jobs held since age 15), and comprehensive exposure assessment and analysis along several different dimensions of exposure. The major limitations were the lack of direct exposure information and potential inaccuracy of the descriptions of work histories that were obtained from next-of-kin interviews. Heineman et al. (1994) acknowledge these limitations in the report, and in response to a letter by Norman and Boggs (1996) criticizing the methodology and interpretation of the study, Heineman et al. (1996) noted that while the lack of direct exposure information must be interpreted cautiously, it does not invalidate the results. Differential recall bias between cases and controls was unlikely because work histories came from next-of-kin for both groups, the industrial hygienists made their judgments blinded to disease status, and the strong associations that were seen with the exposure measures for dichloromethane were not seen with the other solvents included in the analysis. The relatively strong and statistically significant associations between dichloromethane and astrocytic brain tumors were seen along multiple measures of exposure, suggesting that the results were unlikely to be spurious. Nondifferential misclassification would, on average, attenuate true associations and would be unlikely to result in the types of exposure-response relationships that were observed in this study.

Norman and Boggs (1996) described an apparent inconsistency in the estimated trends in dichloromethane and carbon tetrachloride exposure based on the methodology used in this case-control study (described in more detail in Gomez et al. [1994]). In response, Gomez (1996) noted that the apparent inconsistency was actually due to an error in the labeling of the lines on one of the figures in the report rather than an inconsistency with the estimated trends. Another

point raised by Norman and Boggs (1996) was that the Heineman et al. (1994) findings were surprising in light of the lack of brain carcinogenesis in animals. In response, Heineman et al. (1996) pointed out that carcinogens commonly cause different cancers in animals and humans. It can also be noted that brain tumors are exceedingly rare in animal bioassays (Sills et al., 1999). Norman and Boggs (1996) also suggested that the results of the Heineman et al. (1994) study be given no weight when compared with the results of the cohort studies. The authors responded by pointing out that the cohort studies had low statistical power and large CIs around their point estimates but were not inconsistent with an association between dichloromethane and brain cancer (Heineman et al., 1996). This point is strengthened further by the more recent results from the Rochester, New York, Eastman Kodak cohort (Hearne and Pifer, 1999), described previously, since an increased SMR for brain and CNS cancers was seen in the longer follow-up period of this cohort.

In another case-control study of brain cancer and dichloromethane exposure, Cocco et al. (1999) identified 12,980 female cases of cancer of the brain and CNS through the underlying cause of death listing (ICD-9 codes 191 and 192) on death certificates from 24 states from 1984 to 1992. (This collection of death certificates is a data set created by the National Center for Health Statistics, NIOSH, and the National Cancer Institute to facilitate research on occupational exposures and mortality risk.) The cases included 161 women with meningioma (ICD-9 codes 192.1, 192.3). Four women who died of nonmalignant diseases, excluding neurological disorders, were chosen as controls for each case. The controls were frequency matched to the cases by state, race, and 5-year age group. Occupation data were based on the occupation fields in the death certificates. This job was coded based on the three-digit industry and three-digit occupation (Department of Census) codes. The investigators developed job exposure matrices that were applied to these industry/occupation codes. The job exposure matrices included probability and intensity scores for 11 occupational hazards, one of which was dichloromethane, but also included other solvents, electromagnetic fields, chlorinated aliphatic hydrocarbons, benzene, lead, nitrosamines, insecticides, herbicides, and public contact. The investigators used logistic regression models to estimate ORs, adjusting for each workplace exposure, marital status, three levels of socioeconomic status (based on occupation), and age at death. For each chemical, four levels of intensity and probability were defined (unexposed, low, medium, and high).

A weak association between dichloromethane exposure and brain/CNS cancer was seen (OR 1.2 [95% CI 1.1–1.3]) (Cocco et al., 1999). There was no exposure-related trend in the association between probability or intensity of exposure and brain cancer. A similar but more imprecise association was seen with meningioma cancer (OR 1.2 [95% CI 0.7–2.2]). There were too few cases of meningioma to stratify by exposure probability and intensity.

The major limitations of this study are the use of mortality rather than incidence data and the reliance on occupation data from death certificates. The death certificate occupation data are

based on “usual” occupation, which may be more prone to misclassification in studies of women because of gender-related differences in work patterns (i.e., shorter duration jobs for women compared with men). A relatively broad job exposure matrix was applied to the job information, and typically more generic job exposure matrices result in less sensitive assessment with limited ability to detect exposure-response trends (Teschke et al., 2002). Nondifferential misclassification of outcome and exposure would generally result in attenuated effect estimates.

4.1.3.6.2. Case-control studies of breast cancer. Cantor et al. (1995) conducted a case-control study of occupational exposures and breast cancer using the 24-state (1984–1989) death certificate data described in the previous section. Cases were women with breast cancer coded as the underlying cause of death (ICD-9 code 174). Four female controls per case were selected from all noncancer deaths, frequency matched by age (5-year age groups) and ethnicity (black, white). The occupation listed on the death certificate was coded based on the three-digit industry and three-digit occupation (Department of Census) codes, and this was used with a job exposure matrix developed by the investigators to assess 31 workplace exposures, one of which was dichloromethane. Four exposure probability and three exposure level scores were assigned. ORs for probability and level were calculated for each ethnic group, adjusting for age at death and a measure of socioeconomic status (based on occupation). After excluding subjects whose death certificate occupations were listed as homemaker, there were 29,397 white cases and 4,112 black cases (total 33,509) and 102,955 white controls and 14,839 black controls (total 117,794).

There was little evidence of an association between exposure probability and breast cancer mortality using the probability exposure metric. The ORs were 1.05 (95% CI 0.97–1.1) and 0.76 (95% CI 0.3–2.0) in probability level 3 and level 4, respectively, for white women and 1.13 (95% CI 0.9–1.4) in probability level 3 for black women. (There were too few black women in exposure probability level 4 for analysis.) Weak associations were seen with exposure level. In white women, an OR of 1.17 (95% CI 1.1–1.3) was seen with the highest exposure level, and in black women the OR in this exposure group was 1.46 (95% CI 1.2–1.7). In the analysis that jointly considered exposure level and probability ratings but excluded the lowest probability of exposure, the OR for the highest category of exposure level was 1.28 in whites ($p < 0.05$) and 1.21 in blacks.

As with the Cocco et al. (1999) case-control study that used a similar methodology, the limitations of this study include the use of an outcome defined by mortality rather than incidence, use of usual occupation information as recorded in death certificates, and use of a very broad job exposure matrix to classify 31 different exposures. Although information on pregnancy and lactation history (known risk factors for breast cancer) was not available, the authors did adjust for socioeconomic status by using the occupation data, which may have corrected for some of the potential confounding due to reproductive history.

4.1.3.6.3. Case-control studies of pancreatic cancer. Kernan et al. (1999) conducted a case-control study of 63,097 pancreatic cancer cases using the 24-state (1984–1993) death certificate data. The diagnosis of pancreatic cancer was based on underlying cause of death (ICD-9 code 157). Four controls who had died during the same time period of causes other than cancer were selected for each case, frequency-matched by state, race, gender, and 5-year age group (n = 252,386). Usual occupation and industry, based on the occupation data in the death certificate, were coded by using the three-digit (Department of Census) codes. A job-exposure matrix was used with the industry and occupation codes to evaluate exposure intensity and probability (each categorized as high, medium, or low) for formaldehyde, dichloromethane, 10 other solvents, and a combined “organic solvents” measure. Race- and gender-specific analyses were conducted by using logistic regression to estimate ORs and 95% CIs, adjusting for age, marital status (ever, never married), residential area (metropolitan, nonmetropolitan), and region (east, south central, south, and west).

The point estimates for the ORs in the low, medium, and high intensity categories in the four race-gender groups ranged from 0.8 to 1.3, with no exposure-effect trend seen in any group. The only statistically significant OR was for high exposure intensity in white females (OR 1.3 [95% CI 1.1–1.6]), with ORs of 1.0 (95% CI 0.9–1.1) for medium intensity and 1.1 (95% CI 1.0–1.2) for low intensity in this group. An elevated OR was seen with high exposure probability in black males (OR 2.2 [95% CI 1.0–4.8]) but not in white females (OR 1.0 [95% CI 0.8–1.4]) or white males (OR 1.0 [95% CI 0.8–1.3]), and the ORs were 0.9 for medium exposure probability in these three groups. There were relatively few black females in this study, resulting in imprecise estimates (OR 2.0 [95% CI 0.8–5.4] for medium exposure and OR 1.5 [95% CI 0.6–3.6] for high exposure).

The limitations of this study, as with the other case-control studies that used the 24-state death certificate data set, include the reliance on cause of death data from death certificates rather than medical-record validated incidence data and the use of death certificate occupation data. The job exposure matrix used with the occupation data was more focused than those used in Cocco et al. (1999) and Cantor et al. (1995). Although the analysis adjusted for some sociodemographic characteristics, it did not include measures of smoking history or diabetes, which are known risk factors for pancreatic cancer (Lowenfels and Maisonneuve, 2005).

4.1.3.6.4. Case-control studies of renal cancer. Dosemeci et al. (1999) reported data from a population-based case-control study of the association between occupational exposures and renal cancer risk. The investigators identified newly diagnosed patients with histologically confirmed renal cell carcinoma from the Minnesota Cancer Surveillance System from July 1, 1988, to December 31, 1990. The study was limited to white cases, and age and gender-stratified controls were ascertained by using random digit dialing (for subjects ages 20–64) and from Medicare

records (for subjects 65–85 years). Of the 796 cases and 796 controls initially identified, 438 cases (273 men, 165 women) and 687 controls (462 men, 225 women) with complete personal interviews were included in the occupational analysis.

Data were obtained through in-person interviews that included demographic variables, residential history, diet, smoking habits, medical history, and drug use. The occupational history included information about the most recent and usual industry and occupation (coded using the standard industrial and occupation codes, Department of Commerce), job activities, hire and termination dates, and full- and part-time status. A job exposure matrix developed by the National Cancer Institute was used with the coded job data to estimate exposure status to dichloromethane and eight other chlorinated aliphatic hydrocarbons.

ORs were adjusted for age, smoking, hypertension and use of drugs for hypertension, and body mass index. No association between renal cell carcinoma and exposure to dichloromethane was observed in men (OR 0.85 [95% CI 0.6–1.2]), women (OR 0.95 [95% CI 0.4–2.2]), or both sexes combined (OR 0.87 [95% CI 0.6–1.2]).

Strengths of this study include the use of incident cases of renal cancer from a defined population area and confirmation of the diagnosis using histology reports. The occupation history was based on usual and most recent job in combination with a relatively focused job exposure matrix. In contrast to the type of exposure assessment that can be conducted in cohort studies within a specific workplace, however, exposure measurements based on personal or workplace measurements were not used, and a full lifetime job history was not obtained.

4.1.3.6.5. Case-control studies of rectal cancer. Dumas et al. (2000) reported data from a case-control study of occupational exposures and rectal cancer conducted in Montreal, Quebec, Canada. The investigators identified 304 newly diagnosed cases of primary rectal cancer, confirmed on the basis of histology reports, between 1979 and 1985; 257 of these participated in the study interview. One control group (n = 1,295) consisted of patients with other forms of cancer (excluding lung cancer and other intestinal cancers), recruited through the same study procedures and time period as the rectal cancer cases. A population-based control group (n = 533), frequency matched by age strata, was drawn by using electoral lists and random digit dialing. The occupational assessment consisted of a detailed description of each job held during the working lifetime, including the company, products, nature of work at site, job activities, and any additional information from the interviews that could furnish clues about exposure. The percentage of proxy respondents was 15.2% for cases, 19.7% for other cancer controls, and 12.6% for the population controls.

A team of industrial hygienists and chemists blinded to subjects' disease status translated jobs into potential exposure to 294 substances with three dimensions (degree of confidence that exposure occurred, frequency of exposure, and concentration of exposure). Each of these exposure dimensions was categorized into none, any, or substantial exposure. Logistic

regression models adjusted for age, education, proxy versus subject responder status, cigarette smoking, beer drinking, and body mass index. Using the cancer control group, the OR for any exposure to dichloromethane was 1.2 (95% CI 0.5–2.8) and the OR for substantial exposure (confident that exposure occurred with ≥ 5 years of exposure at medium or high frequency and concentration) was 3.8 (95% CI 1.1–12.2). The results using the population-based control group for this exposure were not presented.

The strengths of this study were the large number of incident cases, specific information about job duties for all jobs held, and a definitive diagnosis of rectal cancer. However, the use of the general population (rather than a known cohort of exposed workers) reduced the likelihood that subjects were exposed to dichloromethane, resulting in relatively low statistical power for the analysis. The job exposure matrix applied to the job information was very broad since it was used to evaluate 294 chemicals.

4.1.3.6.6. Case-control studies of childhood leukemia. Infante-Rivard et al. (2005) examined the association between maternal occupational exposures, before and during pregnancy, and risk of childhood acute lymphoblastic leukemia (ICD-9 code 204.0) by using data from a population-based case-control study in Quebec, Canada. Incident cases diagnosed from 1980 to 2000 were identified from the cancer hospitals in the province, and diagnosis was confirmed based on clinical records from an oncologist or hematologist. Between 1980 and 1993, cases ages 0–9 years at diagnosis were included, and from 1994 to 2000 the age range was expanded to 14 years. The number of eligible cases identified was 848 and of these, 790 parents (93%) participated in the study. Population-based controls, individually matched to the sex and age at diagnosis of the cases, were identified from government registries of all children in the province (1980–1993) and the universal health insurance files (1994–2000). The parents of 790 (86%) of the 916 eligible controls who were identified participated in the study.

Data were collected by using a structured telephone interview. Some information (i.e., job title, dates, type of industry, industry name and address) was obtained for all jobs held since age 18, and additional information (e.g., materials and machines used, typical activities) was obtained for jobs held by the mother from 2 years before the pregnancy through the birth of the child. Specialized exposure modules were also used to collect information about specific jobs (e.g., nurse, waitress, hair dresser, textile dry cleaner). All of this information was reviewed by chemists and industrial hygienists, blinded to case-control status, to classify exposure to over 300 chemicals, although the primary focus of the study was on solvents (21 individual substances, including dichloromethane, and six mixtures). The exposure assessment included ratings of confidence (possible, probable, and definite), frequency of exposure during a normal workweek (<5, 5–30, or >30% of the time), and level of concentration (low = slightly above background, high = highest possible exposure in the study population, and medium for in-between levels).

A weak association was seen between any dichloromethane exposure during the 2 years before pregnancy up to the birth and risk of leukemia in the child (OR 1.34 [95% CI 0.54–3.34]), and results were similar when limited to exposures during pregnancy. Stronger associations were seen with probable or definite exposure (OR 3.22 [95% CI 0.88–11.7]) compared with possible or no exposure. The estimates for categories based on concentration and frequency were similar but there was no evidence for an increasing risk with increasing exposure level.

4.1.3.7. Summary of Cancer Studies by Type of Cancer

The cohort and case-control studies with data relevant to the issue of dichloromethane exposure and cancer risk are summarized in Tables 4-9 and 4-10, respectively. The strongest of the cohort studies in terms of design are two of the triacetate film base production cohorts (Cohort 1 in New York and the United Kingdom cohort, reported in Hearne and Pifer [1999] and Tomenson et al. [1997], respectively). These are the cohorts with the most extensive exposure assessment information. The start of eligibility for cohort entrance corresponds with the beginning of the time when the exposure potential at the work site began, and the follow-up period is relatively long (mean >25 years). Although Cohort 2 of the New York film base production study has similar exposure data and follow-up, this cohort was limited to workers employed between 1964 and 1970 and therefore would have missed anyone leaving (possibly because of illness or death) before this time. In addition, because of the overlap between Cohort 1 and Cohort 2, including both cohorts in an evaluation would be double-counting experiences of some individuals. Several limitations of the triacetate film base production cohorts should be noted, however. One of these limitations concerns the generalizability of the results given the relatively low exposure level (mean 8-hour TWA <40 ppm). Exposures in small, poorly ventilated work areas are also often much higher than those seen in these film base production cohorts (Estill and Spencer, 1996; Anundi et al., 1993). Other limitations include the limited power to detect a risk of low-incidence cancers (including brain and leukemia), the lack of women and thus lack of data pertaining to breast cancer, and the use of mortality rather than incidence data. Although the exposure levels in the cohorts involved in cellulose triacetate fiber production were much higher than those of the film production cohorts, the duration of exposure was relatively short in the South Carolina cohort (Lanes et al., 1993), and the majority of workers were missing job history data. In the Maryland triacetate fiber production plant, duration of exposure was not reported and the length of follow-up was relatively short (mean 17 years) (Gibbs et al., 1996). Also, the cohort began in 1970, even though production began in 1955, and the missing personnel records made it impossible to recreate an inception cohort. The exposure assessment in the study of civilian Air Force base workers (Blair et al., 1998) allowed for only a dichotomized classification of exposure, and there was considerable exposure to other solvents among these workers. This Air Force base study was the largest of the cohort studies that included women and presented data pertaining to breast cancer.

Table 4-10. Summary of case-control studies of cancer risk and dichloromethane exposure

Cancer type, reference	Location n cases, n controls (source), demographic group	Exposure assessment	Results^a
Brain Heineman et al. (1994)	Louisiana, New Jersey, Philadelphia 300 cases, 320 controls (death certificates); cancer confirmed by hospital records; white men	Job exposure matrix applied to detailed information on all jobs held (at least 1 yr) since age 15, as obtained from next-of-kin interviews; probability, duration, intensity, and cumulative exposure scores; six solvents evaluated	See Section 4.1.3.6.1. OR 1.3 (0.9, 1.8) for any exposure; increased risk with increased probability (trend <i>p</i> -value < 0.05, OR 2.4 [1.0, 5.9] for high probability), increased duration, increased intensity; strongest effects seen in high probability plus high duration, OR 6.1 (1.1, 43.8) or high intensity and high duration, OR 6.1 (1.5, 28.3) combinations; no association with cumulative exposure score
Brain Cocco et al. (1999)	24 states, U.S. 12,980 cases, 51,920 controls (death certificates); women	Job exposure matrix applied to death certificate occupation; probability, and intensity scores; 11 exposures evaluated	See Section 4.1.3.6.1. Weak association overall, OR 1.2 (1.1, 1.3), no trend with probability or intensity scores
Breast Cantor et al. (1995)	24 states, U.S. 33,509 cases, 117,794 controls (death certificates); black and white women	Job exposure matrix applied to death certificate job data, probability, and exposure level; 31 substances evaluated	See Section 4.1.3.6.2. Little evidence of association with exposure probability; weak association with highest exposure level in whites, OR 1.17 (1.1, 1.3) and in blacks, OR 1.46 (1.2, 1.7)
Pancreas Kernan et al. (1999)	24 states, U.S. 63,037 cases, 252,386 controls (death certificates); black and white men and women	Job exposure matrix applied to death certificate occupation, probability, and intensity scores; 11 chlorinated solvents and formaldehyde evaluated	See Section 4.1.3.6.3. Little evidence of associations with intensity or probability
Kidney Dosemeci et al. (1999)	Minnesota 438 incident cases (Minnesota cancer registry), 687 controls (random digit dialing and Medicare records); cancer confirmed by histology; men and women	Job exposure matrices applied to most recent and usual job, as ascertained from interviews; nine solvents evaluated	See Section 4.1.3.6.4. No evidence of increased risk associated with dichloromethane in men, OR 0.85 (0.6, 1.2) or women, OR 0.95 (0.4, 2.2)

(Table 4-10 continues on next page)

Table 4-10. Summary of case-control studies of cancer risk and dichloromethane exposure

Cancer type, reference	Location n cases, n controls (source), demographic group	Exposure assessment	Results^a
Rectum Dumas et al. (2000)	Montreal, Canada 257 incident cases, 1,295 other cancer controls from 19 hospitals; 533 population-based controls (electoral rolls and random digit dialing), cancer confirmed by histology; men	Job exposure matrix applied to detailed information on all jobs held, as ascertained from interviews; 294 substances evaluated	See Section 4.1.3.6.5. Little evidence of an association with any exposure, OR 1.2 (0.5, 2.8), but increased risk in a small, “substantial exposure” group, OR 3.8 (1.1, 12.2) (using cancer controls; analysis of population controls not given for this exposure)
Childhood leukemia (acute lymphoblastic leukemia) Infante-Rivard et al. (2005)	Quebec, Canada 790 incident cases (hospitals—all provinces), 790 population-based controls (government population registries); cancer based on oncologist or hematologist diagnosis ages 0–14, ^b both sexes	Systematic review of detailed information on all jobs held by the mother from 2 yrs before pregnancy through birth of the child; 21 individual substances and six mixtures evaluated (mostly solvents); confidence, frequency, and concentration of exposure rated	See Section 4.1.3.6.6. Little evidence of association with any exposure, OR 1.34 (0.54, 3.34), but stronger associations with probable or definite, OR 3.22 (0.88, 11.7) (referent group = possible/no exposure) and with combinations of frequency and concentration

^aResults given as OR and (95% CI).

^bFrom 1980 to 1993, study was limited to diagnoses of ages 0–9, but this was expanded between 1994 and 2000 to ages 0–14.

Case-control studies offer the potential for increased statistical power for assessing associations with relatively rare cancers such as brain cancer and leukemia. Case-control studies are often designed to examine incidence rather than mortality, which is of particular importance in etiologic research for diseases with relatively high survival rates and diseases in which survival may be strongly related to factors that are difficult to adjust for without detailed data collection (e.g., access to health care). There is a considerable range in the detail and quality of the exposure assessment used in case-control studies, however. Case-control studies rarely include specific measurements taken at specific work sites of individual study participants. Although it is more difficult to determine absolute exposure levels without these individual measurements, the exposure assessment methodology used in case-control studies can result in useful between-group comparisons of risk if the intra-group variability is less than the intergroup variability in potential exposure levels. Among the case-control studies with data pertaining to cancer risk and dichloromethane exposure, the two studies with the strongest designs are the study of brain cancer by Heineman et al. (1994) and the study of childhood leukemia by Infante-Rivard et al. (2005). These are the studies that obtained detailed information about all jobs held (rather than just the usual or most recent job), focused on a relatively small number of exposures, and used medical record data to confirm the diagnosis. Heineman et al. (1994) obtained the work history from interviews with next-of-kin, however, which is most likely to have resulted in nondifferential misclassification of exposure, and thus attenuation in the observed associations. The use of death certificate data to classify disease and occupational exposures in the three studies using the large 24-state death certificate database (brain cancer, Cocco et al. [1999]; breast cancer, Cantor et al. [1995]; pancreatic cancer, Kernan et al. [1999]) is also likely to have resulted in nondifferential misclassification of both outcome and exposure (and thus attenuated associations).

Considering the issues described above with respect to the strengths and limitations of the available epidemiologic studies, a summary of the epidemiologic evidence relating to dichloromethane exposure and specific types of cancer can be made, as described below. The available epidemiologic data suggest an association between dichloromethane and brain cancer and liver cancer, but not lung cancer.

4.1.3.7.1. Brain and CNS cancer. An increased risk of brain and CNS cancers was seen in the strongest cohort studies; SMRs were 2.16 in Cohort 1 in New York (Hearne and Pifer, 1999) and 1.45 in the United Kingdom cohort (Tomenson et al., 1997). These estimates are based on a small number of observations (six cases in New York and four in the United Kingdom) and so are relatively imprecise. It is only in the latest follow-up of the New York film base production cohort that an elevated SMR was observed, further suggesting that the statistical power of the other cohort studies for examining risk of this disease may be quite low. Two case-control studies of dichloromethane exposure and brain cancer have been conducted (Cocco et al., 1999;

Heineman et al., 1994). The Heineman et al. (1994) study, which is the stronger study in terms of exposure assessment strategy and confirmation of diagnosis, reported relatively strong trends ($p < 0.05$) with increasing probability, duration, and intensity measures of exposure, but a nonlinear trend was seen with the cumulative exposure metric. This difference could reflect a more valid measure of relevant exposures in the brain from the intensity measure, as suggested by the study in rats reported by Savolainen et al. (1981) in which dichloromethane levels in the brain were much higher with a higher intensity exposure scenario compared with a constant exposure period with an equivalent TWA (see Section 3.2). The combination of high probability of exposure and long (> 20 years) duration of employment in exposed jobs was strongly associated with brain cancer risk (OR 6.1, 95% CI 1.1-43.8) in the Heineman et al. (1994) study; similar associations were seen with the high intensity in combination with long duration measures. The available epidemiologic studies provide some evidence of an association between dichloromethane and brain cancer, and this area of research represents a data gap in the understanding of the carcinogenic potential of dichloromethane.

4.1.3.7.2. *Liver and biliary duct cancer.* Liver and biliary duct cancer are relatively uncommon (age-adjusted incidence 6.2 per 100,000 person-years) (SEER website, seer.cancer.gov, accessed April 2006), so it is difficult to study in most occupational cohorts of limited size. The cohort study with the higher exposures, the Rock Hill triacetate fiber production plant, suggested an increased risk of liver cancer (Lanes et al., 1993, 1990). The SMR for liver and bile duct cancer was 2.98 (95% CI 0.81–7.63) in the latest update of this cohort. This observation was based on four cases; two of these cases were biliary duct cancers. As the follow-up period has increased, the strength of this association has decreased, although it is relatively strong (albeit with wide CIs). The decrease in the SMR with increasing follow-up reflects the increase in number of expected cases because the four observed cases were seen earlier in the follow-up period. No other cohort study has reported an increased risk of liver cancer mortality, although it should be noted that there is no other inception cohort study of a population with exposure levels similar to those of the Rock Hill plant, and no data from a case-control study of liver cancer are available pertaining to dichloromethane exposure. The available epidemiologic studies, with biological plausibility inferred from the results from studies in mice and female rats (see Section 4.2) (NTP, 1986; Serota et al., 1986a, b; Nitschke, 1988a), provide evidence of an association between dichloromethane and liver and biliary duct cancer, although it should be noted that this evidence is based on very limited epidemiologic data.

4.1.3.7.3. *Lung cancer.* In the stronger cohort studies (Cohort 1 in the New York Eastman Kodak Company triacetate film production study reported by Hearne and Pifer [1999] and the United Kingdom triacetate film production study reported by Tomenson et al. [1997]), the SMRs for lung cancer were well below 1.0. The New York study had also obtained data on smoking

history that indicated it was unlikely that differences in smoking could be masking an effect of dichloromethane (Hearne et al., 1987). Lung cancer is a common cancer (age-adjusted incidence 61 per 100,000 person-years) (SEER website, seer.cancer.gov, accessed April 2006) so the expected rates, even in small cohorts, are based on relatively robust estimates. The only group in any study that had an increased risk for lung cancer was the high-exposure women in the triacetate fiber production cohort in Maryland (Gibbs et al., 1996). However, this was based on only two cases and was a highly imprecise estimate (SMR 2.3 [95% CI 0.28–8.3]). No case-control study of dichloromethane exposure and lung cancer risk is available. The available epidemiologic studies do not provide evidence for an association between dichloromethane and lung cancer, although it should be noted that the studies with the best designs are limited to relatively low exposure levels.

4.1.3.7.4. Pancreatic cancer. An early study (Hearne et al., 1990) of Cohort 2 of the New York triacetate film production cohort had reported 8 observed and 4.2 expected pancreatic cancer deaths, for a twofold increased SMR ($p = 0.13$). This association was reduced in the subsequent follow-up (SMR 1.5 [95% CI 0.7–3.0]) (Hearne and Pifer, 1999) but was not seen in the more methodologically sound Cohort 1 (SMR 0.92) or in any of the other cohorts. A meta-analysis of the cohort studies (using the data of Hearne et al. [1990]) reported a summary association of 1.42 (95% CI 0.80–2.53) (Ojajärvi et al., 2001). This summary measure would be further reduced with the updated data for Cohort 2 and the addition of Cohort 1 from Hearne and Pifer (1999). The only case-control study of pancreatic cancer mortality risk and dichloromethane exposure (based on death certificate data) did not report consistent patterns with respect to intensity or exposure among the race-sex groups studied. The available epidemiologic studies do not provide evidence for an association between dichloromethane and pancreatic cancer.

4.1.3.7.5. Leukemia and lymphoma. Each of the individual hematopoietic cancers is relatively uncommon, with age-adjusted incidence rates of 5 per 100,000 person-years or less (SEER website, seer.cancer.gov, accessed April 2006). The relatively inconsistent (point estimates ranging from ≤ 0.50 to ≥ 2.0) and imprecise measures of association between dichloromethane exposure and non-Hodgkin's lymphoma, Hodgkin's lymphoma, myeloma, and leukemia are thus expected, given the relatively small size of the available cohort studies. Only one case-control study of any of these diseases and dichloromethane is available, and this is a study of childhood leukemia (acute lymphoblastic leukemia) in relation to maternal occupational history (Infante-Rivard et al., 2005). This is a large, population-based study of confirmed incident cases of leukemia, with a detailed exposure assessment pertaining to the period before and during pregnancy. A threefold increased risk was seen with probable or definite exposure (OR 3.22 [95% CI 0.88–11.7]) compared with possible or no exposure. The available epidemiologic studies do not provide an adequate basis for the evaluation of the role of dichloromethane in any

of the specific hematopoietic cancers because of the small size of the cohort studies and the relative lack of case-control studies pertaining to these outcomes.

4.1.3.7.6. Breast cancer. Only one large cohort study included women and reported data pertaining to breast cancer risk (Blair et al., 1998), and this is a cohort with a limited exposure assessment (dichotomized) and multiple exposures. A relatively strong association was seen between dichloromethane exposure and breast cancer mortality in this study (rate ratio 3.0 [95% CI 1.0–8.8]). Similar associations were seen with several other chemicals, and the potential effect of confounding and misclassification of these exposures may have biased the estimate in either direction. The only case-control study of breast cancer risk and dichloromethane exposure used the 24-state death certificate data to classify exposure and disease. The available epidemiologic studies do not provide an adequate basis for the evaluation of the role of dichloromethane in breast cancer because there are currently no cohort studies with adequate statistical power and no case-control studies with adequate exposure methodology to examine this relationship.

4.2. SUBCHRONIC AND CHRONIC STUDIES AND CANCER BIOASSAYS IN ANIMALS—ORAL AND INHALATION

4.2.1. Oral Exposure: Overview of Noncancer and Cancer Effects

Results from studies of animals exposed by the oral route for short-term, subchronic, and chronic durations identify the liver and the nervous system as the most sensitive targets for noncancer toxicity from repeated oral exposure to dichloromethane. In a 90-day exposure study, nonneoplastic histopathologic changes in the liver were observed in F344 rats exposed to drinking water doses of ≥ 166 mg/kg-day (males) or ≥ 209 mg/kg-day (females) (Kirschman et al., 1986). Similar changes were seen in F344 rats in a 2-year exposure of ≥ 50 mg/kg-day (Serota et al., 1986a).

The 2-year oral exposure study in F344 rats did not produce evidence of increasing incidence of liver tumors across all of the dose groups in males or females (Serota et al., 1986a). In females, however, a jagged stepped pattern of increasing incidence was observed. In a parallel study in B6C3F₁ mice (Serota et al., 1986b; Hazleton Laboratories, 1983), a clearer trend with respect to hepatic cancer was seen in males but not females.

None of the chronic oral exposure studies included a systematic measurement of potential neurological effects. One 14-day study focusing on neurobehavioral changes is available, however. Changes in autonomic, neuromuscular, and sensorimotor functions were observed in F344 rats exposed for 14 days to gavage doses ≥ 337 mg/kg-day (Moser et al., 1995) (see Section 4.4.3 for more details).

No effects on reproductive parameters were observed in Charles River CD rats exposed for 90 days to gavage doses as high as 225 mg/kg-day (General Electric Company, 1976) or in

pregnant F344 rats exposed to gavage doses of up to 450 mg/kg-day on GDs 6–19 (Narotsky and Kavlock, 1995). However, no oral exposure studies examining developmental neurobehavioral effects have been conducted (see Section 4.3 for more details).

4.2.1.1. Toxicity Studies of Subchronic Oral Exposures: Hepatic Effects

Kirschman et al. (1986) examined the toxicity of dichloromethane in a 90-day drinking water study in F344 rats (20/sex/dose level). The nominal concentration of dichloromethane in the water was 0.15, 0.45, or 1.5%. Based on BW and water consumption data, average intakes were reported to be 0, 166, 420, or 1,200 mg/kg-day for males and 0, 209, 607, or 1,469 mg/kg-day for females. Clinical chemistry tests (hematological and chemical variables in samples of blood and urine) and tissue histopathology were evaluated in groups of five rats/sex/dose level after 1 month of treatment. These endpoints were also evaluated in the remaining rats sacrificed after 90 days of exposure.

Exposure to dichloromethane did not affect mortality or cause adverse clinical signs of toxicity. Gross necropsy was also unremarkable. Reported changes in mean values for clinical chemistry variables compared with controls included elevated serum ALT activities for all treated males at 1 month and for the high-dose females at 3 months, elevated serum AST activity in high-dose females at 3 months, elevated serum lactate dehydrogenase activities in mid- and high-dose females at 3 months, and decreases in serum concentrations of fasting glucose, cholesterol, and triglycerides in all exposed groups of both sexes at 1 and 3 months. Actual values for clinical chemistry variables, however, were not presented in the report.

No histopathologic alterations were seen in tissues after 1 month of treatment (a detailed description of tissues examined was not presented). In rats exposed for 3 months, exposure-related histopathologic changes were restricted to the liver. Elevated, statistically significant incidences of hepatocytic vacuolation were observed in all exposed male and female groups (see Table 4-11). The most frequently observed vacuolation was described as generalized and occurring throughout the lobule, and Oil Red-O-staining indicated that most were lipid-containing vacuoles. The incidences of generalized vacuolation scored as mild or moderate were higher in all of the female dose groups compared with the controls. The authors stated that the no-observed-adverse-effect level (NOAEL) based on this study is <200 mg/kg-day and the lowest-observed-adverse-effect level (LOAEL) for males was 166 mg/kg-day. The authors did not explicitly provide a LOAEL for females. The results indicate that 166 mg/kg-day and 209 mg/kg-day were the LOAELs for liver effects in male and female rats, respectively.

Table 4-11. Incidences of histopathologic changes in livers of male and female F344 rats exposed to dichloromethane in drinking water for 90 days

Lesion, by sex	Controls	Low dose	Mid dose	High dose
Males—n per group ^a	15	15	15	15
Estimated mean intake (mg/kg-d)	0	166	420	1,200
Number (%) with: Hepatocyte vacuolation (generalized, centrilobular, or periportal)	1 (7)	10 ^b (67)	9 ^b (60)	7 ^b (47)
Generalized vacuolation severity:				
minimal	0 (0)	5 ^b (33)	8 ^b (53)	6 ^b (40)
mild	0	4	7	6
moderate	0	0	1	0
moderate	0	1	0	0
Centrilobular severity:				
minimal	0 (0)	1 (7)	0 (0)	2 (13)
mild	0	1	0	0
moderate	0	0	0	2
moderate	0	0	0	0
Hepatocyte degeneration	0 (0)	0 (0)	0 (0)	2 (13)
Focal granuloma	1 (7)	0 (0)	0 (0)	1 (7)
Females—n per group ^a	15	15	15	15
Estimated mean intake (mg/kg-d)	0	209	607	1,469
Number (%) with: Hepatocyte vacuolation (generalized, centrilobular, or periportal)	6 (40)	13 ^b (87)	15 ^b (100)	15 ^b (100)
Generalized vacuolation severity:				
minimal	5 (33)	13 ^b (87)	15 ^b (100)	15 ^b (100)
mild	5	8	6	8
moderate	0	4	5	6
moderate	0	1	4	1
Centrilobular severity:				
minimal	0 (0)	0 (0)	1 (7)	11 ^b (28)
mild	0	0	0	2
moderate	0	0	1	4
moderate	0	0	0	3
marked	0	0	0	2
Hepatocyte degeneration	0 (0)	0 (0)	0 (0)	12 ^b (80)
Focal granuloma	0 (0)	0 (0)	4 (27) ^c	6 ^b (40)

^a20 per group; 5 sacrificed at 1 mo; these endpoints for the remaining 15 per group.

^bStatistical significance testing not reported by authors; Fisher's exact test for comparison with control *p*-value < 0.05 (two-sided).

^cStatistical significance testing not reported by authors; Fisher's exact test for comparison with control *p*-value < 0.10 (two-sided). Authors stated LOAEL = 166 mg/kg-d in males but did not explicitly provide LOAEL for females; NOAEL is <200 mg/kg-d.

Source: Kirschman et al. (1986).

Kirschman et al. (1986) conducted a similar 90-day study in B6C3F₁ mice (20/sex/dose level). The estimated average intakes were 0, 226, 587, or 1,911 mg/kg-day for males and 231, 586, or 2,030 mg/kg-day for females. Six mice (two controls, two low dose, and two mid dose) died during the study from unknown causes. Administration of dichloromethane did not cause

adverse clinical signs of toxicity or affect food consumption, ophthalmology, or serum ALT activity. Gross necropsy examinations were also unremarkable.

Histopathologic evaluation of tissues from mice killed after 1 month of treatment did not reveal any compound-related effects. Evaluation at 3 months showed subtle generalized or centrilobular changes in the liver (characterized as increased vacuolation with fat deposition), which was evident in all exposed groups and most prominent in mid- and high-dose female groups (Table 4-12). The most frequently detected change was characterized as a generalized vacuolation. Some evidence was found for an increase in severity of the generalized vacuolation with increasing exposure level, but the incidence of this lesion in the control mice was substantial, especially in females (Table 4-12). Incidences for centrilobular vacuolation were significantly increased only for the mid-dose female group. No other changes were found.

Table 4-12. Incidences of histopathologic changes in livers of male and female B6C3F₁ mice exposed to dichloromethane in drinking water for 90 days

Lesion, by sex	Controls	Low dose	Mid dose	High dose
Males—n per group ^a	14	14	14	15
Estimated mean intake (mg/kg-d)	0	226	587	1,911
Number (%) with:				
Hepatocyte vacuolation (generalized, centrilobular, or periportal)	9 (64)	12 (86)	13 (93)	12 (80)
Generalized vacuolation, severity:	7 (50)	12 ^b (86)	13 ^b (93)	10 (67)
minimal	4	3	9	7
mild	2	7	5	3
moderate	1	2	0	0
marked	0	0	0	0
Centrilobular severity:	2 (14)	0 (0)	1 (7)	5 (33)
minimal	2	0	0	1
mild	0	0	0	3
moderate	0	0	1	1
Females—n per group ^a	14	11	13	15
Estimated mean intake (mg/kg-d)	0	231	586	2,030
Number (%) with:				
Hepatocyte vacuolation (generalized, centrilobular, or periportal)	13 (93)	11 (100)	13 (100)	13 (87)
Generalized vacuolation severity:	13 (93)	11 (100)	13 (100)	13 (87)
minimal	1	3	5	3
mild	8	7	6	6
moderate	4	1	2	1
marked	0	0	0	3
Centrilobular severity:	0 (0)	0 (0)	5 ^c (39)	1 (7)
minimal	0	0	0	0
mild	0	0	2	1
moderate	0	0	3	0
marked	0	0	0	0

^a20 per group; 5 sacrificed at 1 mo.

^bStatistical significance testing not reported by authors; Fisher's exact test for comparison with control p -value = 0.10 for low dose group and p = 0.032 for mid-dose group (two-sided).

^cStatistical significance testing not reported by authors; Fisher's exact test for comparison with control p -value = 0.016 (two-sided). Authors say LOAEL = 587 mg/kg-d; NOAEL between 226 and 587 mg/kg-d for males; not explicitly stated for females.

Source: Kirschman et al. (1986).

Using the results from this study to select doses for a chronic study, Kirschman et al. (1986) expressed the opinion that the mid-dose level (587 mg/kg-day) was the LOAEL in this study. Although incidences for generalized vacuolation were increased in the low- and mid-dose male groups, the incidences in the high-dose groups were not significantly increased compared with controls (Table 4-12). The study authors identified a LOAEL of 587 mg/kg-day for centrilobular vacuolation in male B6C3F₁ mice. The NOAEL for males was considered by the investigators to be between 226 and 587 mg/kg-day.

4.2.1.2. Toxicity Studies of Chronic Oral Exposures: Hepatic Effects and Carcinogenicity

Longer-term (up to 2-year) oral exposure studies in mice and rats are summarized in Table 4-13 and described in more detail below. These studies provide additional information pertaining to hepatotoxicity and carcinogenicity.

Table 4-13. Studies of chronic oral dichloromethane exposures (up to 2 years)

Reference, strain/species	Number per group	Exposure information	Comments
Serota et al. (1986a) F344 rats	85/sex/dose + 135 controls	Drinking water, 2 yrs, target dose 0, 5, 50, 125, 250 mg/kg-d Mean intake: males: 0, 6, 52, 125, 235 mg/kg-d females: 0, 6, 58, 136, 263 mg/kg-d	Nonneoplastic liver effects (foci/areas of alteration) in males and females (see Table 4-14); jagged stepped pattern of increasing incidence of neoplastic nodules or hepatocellular carcinoma in females (i.e., increased in the 50 and 250 mg/kg-d but not 125 mg/kg-d groups) (see Table 4-14)
Serota et al. (1986b); Hazleton Laboratories (1983) B6C3F ₁ mice	Males: 125, 200, 100, 100, 125 Females: 100, 100, 50, 50, 50	Drinking water, 2 yrs, target dose 0, 60, 125, 185, 250 mg/kg-d Mean intake: males: 0, 61, 124, 177, 234 mg/kg-d females: 0, 59, 118, 172, 238 mg/kg-d	Increasing trend of liver cancer (hepatocellular adenoma or carcinoma) in males (see Table 4-15)
Maltoni et al. (1988) Sprague-Dawley rats	50/sex/dose	Gavage, up to 64 wks 0, 100, 500 mg/kg-d, 4–5 d/wk	High mortality in high dose group led to termination of study at 64 wks; statistically nonsignificant increase in malignant mammary tumors in female rats
Maltoni et al. (1988) Swiss mice	50/sex/dose + 60 controls	Gavage, up to 64 wks 0, 100, 500 mg/kg-d, 4–5 d/wk	High mortality in high dose group led to termination of study at 64 wks

4.2.1.2.1. Chronic oral exposure in F344 rats (Serota et al., 1986a). Treatment with dichloromethane did not induce adverse clinical signs or affect survival in the F344 rats (Serota et al., 1986a). BWs of rats in the 125 and 250 mg/kg-day groups were generally lower than in controls throughout the study. The authors stated that the differences, although small, were statistically significant, but the data were not shown in the published report. Water consumption was lower throughout the study in both sexes of rats from the 125 and 250 mg/kg-day groups relative to controls; food consumption was also lower in these groups during the first 13 weeks of treatment. Mean hematocrit, hemoglobin, and red blood cell count were increased in both sexes at dichloromethane levels of 50, 125, and 250 mg/kg-day for 52 and 78 weeks. Half of these increases were reported to be statistically significant, but the report did not provide the numerical values or specify which parameters were significant. Clinical chemistry results

showed decreases in alkaline phosphatase (AP), creatinine, blood urea nitrogen, total protein, and cholesterol in both sexes at 250 mg/kg-day, and most of these changes were statistically significant at one or both of the intervals evaluated. (Significant parameters not specified and the mean group values were not presented in the published report.) No significant deviations in urinary parameters were observed. Organ weights were not significantly affected by treatment with dichloromethane.

No treatment-related histopathological effects were noted in the tissues examined except for the liver (Serota et al., 1986a). Examination of liver sections showed a dose-related positive trend (positive Cochran-Armitage trend test) in the incidences of foci/areas of cellular alteration in treated F344 rats (Table 4-14). Comparisons of incidences with control incidences indicated statistically significant elevations at all dose levels except 5 mg/kg-day. These liver changes were first noted after treatment for 78 weeks and progressed until week 104. Livers of animals treated with dichloromethane also showed an increased incidence of fatty change, but incidence data for this lesion were not presented in the published report. The recovery group also showed an increased incidence of areas of cellular alterations, but the fatty changes were less pronounced than in the 250 mg/kg-day group dosed for 104 weeks. The authors indicate that 5 mg/kg-day was a NOAEL and 50 mg/kg-day was a LOAEL for liver changes (foci/areas of cellular alteration) in male and female F344 rats exposed to dichloromethane in drinking water for 2 years.

Table 4-14. Incidences of nonneoplastic liver changes and liver tumors in male and female F344 rats exposed to dichloromethane in drinking water for 2 years

	Target dose (mg/kg-d)						
	0 ^a (Controls)	5	50	125	250	Trend <i>p</i> -value ^b	250 with recovery ^c
Males							
Estimated mean intake (mg/kg-d)	0	6	52	125	235		232
total n	135	85	85	85	85		25
n at terminal kill ^d	76	34	38	35	41		15
Number (%) with: Liver foci/areas of alteration	52 (70)	22 (65)	35 (92) ^e	34 (97) ^e	40 (98) ^e	<0.0001	15 (100) ^e
Neoplastic nodules	9 (12)	1 (3)	0 (0)	2 (6)	1 (2)	Not reported	2 (13)
Hepatocellular carcinoma	3 (4)	0 (0)	0 (0)	0 (0)	1 (2)	Not reported	0 (0)
Neoplastic nodules and hepatocellular carcinoma	12 (16)	1 (3)	0 (0)	2 (6)	2 (5)	Not reported	2 (13)
Females							
Estimated mean intake (mg/kg-d)	0	6	58	136	263		239
total n	135	85	85	85	85		25
n at terminal kill ^d	67	29	41	38	34		20
Number (%) with: Liver foci/areas of alteration	34 (51)	12 (41)	30 (73) ^e	34 (89) ^e	31 (91) ^e	<0.0001	17 (85) ^e
Neoplastic nodules	0(0)	1 (3)	2 (5)	1 (3)	3 (9)	Not reported	2 (10)
Hepatocellular carcinoma	0 (0)	0 (0)	2 (5)	0 (0)	2 (6)	Not reported	0 (0)
Neoplastic nodules and hepatocellular carcinoma	0 (0)	1 (3)	4 (10) ^f	1 (3)	5 (14) ^f	<i>p</i> < 0.01	2 (10) ^f

^aTwo control groups combined. Sample size (incidence of liver foci) in group 1 and 2, respectively, was 36 (75%) and 40 (63%) in males and 31 (55%) and 36 (47%) in females.

^bCochran-Armitage trend test was used for trend test of liver foci/areas of alteration. For tumor mortality-unadjusted analysis, a Cochran-Armitage trend test was used, and for tumor mortality-adjusted analyses, tumor prevalence analytic method by Dinse and Lagakos (1982) was used. Similar results were seen in these two analyses.

^cRecovery group was exposed for 78 wks and then had a 26-wk period without dichloromethane exposure; n = 15 for nonneoplastic lesions and n = 17 for neoplastic lesions.

^dExcludes 5, 10, and 20 per group sacrificed at 25, 52, and 78 wks, respectively, and unscheduled deaths, which ranged from 5 to 19 per group.

^eSignificantly (*p* < 0.05) different from control with Fisher's exact test.

^fSignificantly (*p* < 0.05) different from controls with Fisher's exact test, mortality-unadjusted and mortality-adjusted analyses.

Source: Serota et al. (1986a).

Dichloromethane-exposed male rats showed no statistically significant increased incidence of liver tumors. In females, there was a positive trend for increasing incidence of hepatocellular carcinoma or neoplastic nodules with increasing dose (Table 4-14) (Serota et al., 1986a). Statistically significant increases in tumor incidences were observed in the 50 and 250 mg/kg-day groups (incidence rates of 10 and 14%, respectively) but not in the 125 mg/kg-day group (incidence rate of 3%). Incidence was also increased (10%) in a group exposed for 78 weeks followed by a 26-week period of no exposure. The characterization of malignant potential of the nodules was not described however, and no trend was seen in the data limited to hepatocellular carcinomas. The incidence of hepatocellular carcinoma or neoplastic nodules in this control group (0%) was lower than that seen in historical controls from the same laboratory (324 female F344 rats; 4 with carcinoma, 21 with neoplastic nodules; $25/324 = 7.7\%$).

4.2.1.2.2. Chronic oral exposure in B6C3F₁ mice (Serota et al., 1986b; Hazleton Laboratories, 1983). A 2-year drinking water study similar to the previously described study in F344 rats was also conducted in B6C3F₁ mice (Serota et al., 1986b; Hazleton Laboratories, 1983). The mice received target doses of 0, 60, 125, 185, or 250 mg/kg-day of dichloromethane in deionized drinking water for 24 months. Treatment groups consisted of 100 female mice in the low-dose (60 mg/kg-day) group and 50 in the remaining treatment groups; larger sample sizes were used in the male bioassay, with 200, 100, 100, and 125 male mice in the 60, 125, 185 and 250 mg/kg-day groups, respectively. One hundred females (in two groups of 50) and 125 males (in two groups of 60 and 65 mice) served as controls. The authors indicate that this study design involving two groups of control mice was used because of the high and erratic incidence of liver tumors in historical control B6C3F₁ mice; when the results were similar in the two control groups, the groups could be combined to provide a more statistically precise estimate for comparisons with the exposed groups. Based on water consumption and BW measurements, mean intakes were reported to be 61, 124, 177, and 234 mg/kg-day for males and 59, 118, 172, and 238 mg/kg-day for females. Endpoints examined included clinical signs, BW and water consumption, hematology at weeks 52 and 104, and gross and microscopic examinations of tissues and organs at termination. All tissues from the control and 250 mg/kg-day groups were examined microscopically, as well as the livers and neoplasms from all groups and the eyes of all males from all groups.

Throughout the 2-year study, mice from both control and treatment groups exhibited a high incidence of convulsions (Serota et al., 1986b; Hazleton Laboratories, 1983). The convulsions were noted only during handling for BW determinations, and efforts to establish a basis for this response were unsuccessful. The incidence of convulsions did not correlate with an increased mortality rate. Survival to 104 weeks was high (82% in males and 78% in females), and no evidence for exposure-related negative effects on survival were found. Exposure had no significant effect on BW or water consumption. Mean leukocyte count was significantly

elevated in males and females dosed with 250 mg/kg-day dichloromethane for 52 weeks, but the authors indicated that the mean values were within the normal historical range for the laboratory. Treatment-related nonproliferative histopathologic effects were restricted to the liver and consisted of a marginal increase in the amount of Oil Red O-positive material in the liver of males and females dosed with 250 mg/kg-day (group incidences for this lesion, however, were not presented in the published report). The results indicate that 185 mg/kg-day was a NOAEL and 250 mg/kg-day was a LOAEL for marginally increased amounts of fat in livers of male and female B6C3F₁ mice.

Incidences of liver tumors in female mice were not presented in the published reports (Serota et al., 1986b; Hazleton Laboratories, 1983), but it was reported that exposed female mice did not show increased incidences of proliferative hepatocellular lesions. In the male B6C3F₁ mice, incidences for hepatic focal hyperplasia showed no evidence of an exposure-related effect (Table 4-15). The incidence of hepatocellular adenomas or carcinomas was 18 and 20% in each of the two control groups, and the combined group is presented in this table and used as the comparison group for the analysis. The incidence of hepatocellular adenomas or carcinomas across exposure groups was 26, 30, 31, and 28% in the 60, 125, 185 and 250 mg/kg-day groups, respectively. Similar patterns are seen with the mortality-adjusted incidences (Table 4-15). The trend tests and the tests of the comparisons between individual exposure groups and the controls were not reported by Serota et al. (1986b) but were reported in Hazleton Laboratories (1983). Exposed male mice showed a marginally increased combined incidence of hepatocellular adenomas and carcinomas, with a linear trend p -value = 0.058; the individual p -values for the 60, 125, 185, and 250 mg/kg-day dose groups were 0.071, 0.023, 0.019, and 0.036, respectively.

Table 4-15. Incidences for focal hyperplasia and tumors in the liver of male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years

	Target dose (mg/kg-d)					Trend <i>p</i> -value ^b
	0 ^a (Controls)	60	125	185	250	
n per group ^c	125	200	100	99	125	
Estimated mean intake (mg/kg-d)	0	61	124	177	234	
Number (%) with: Focal hyperplasia ^d	10 (8)	14 (7)	4(4)	10 (10)	13 (10)	Not reported
Hepatocellular adenoma mortality-adjusted percent and <i>p</i> -value ^e	10 (8) (9)	20 (10) (12) <i>p</i> = 0.24	14 (14) (17) <i>p</i> = 0.064	14 (14) (16) <i>p</i> = 0.076	15 (12) (12) <i>p</i> = 0.13	0.172
Hepatocellular carcinoma mortality-adjusted percent and <i>p</i> -value ^e	14 (11) (13)	33 (17) (19) <i>p</i> = 0.082	18 (18) (21) <i>p</i> = 0.073	17 (17) (19) <i>p</i> = 0.11	23 (18) (21) <i>p</i> = 0.044	0.147
Hepatocellular adenoma or carcinoma mortality-adjusted percent and <i>p</i> -value ^e	24 (19) (21)	51 (26) (29) <i>p</i> = 0.071	30 (30) (34) <i>p</i> = 0.023	31 (31) (34) <i>p</i> = 0.019	35 (28) (32) 0.036	0.058

^aTwo control groups combined. Sample size (incidence of hepatocellular adenoma or carcinoma) in group 1 and 2, respectively, was 60 (18%) and 65 (20%). Two additional sets of analyses using the individual control groups were also presented in Hazleton Laboratories (1983).

^bCochran-Armitage trend test (source: Hazleton Laboratories [1983]).

^cNumber at start of treatment.

^dSome mice with hyperplasia also had hepatocellular neoplasms, but the exact number was unspecified by Serota et al. (1986b).

^ePercent calculated based on number at risk, using Kaplan-Meier estimation, taking into account mortality losses; *p*-value for comparison with control group, using asymptotic normal test (source: Hazleton Laboratories [1983]).

Sources: Serota et al. (1986b); Hazleton Laboratories (1983).

Serota et al. (1986b) summarized these results as showing slight increases in proliferative hepatocellular lesions in exposed male B6C3F₁ mice that were not dose-related and were within the range of historical controls, with no effect seen in female B6C3F₁ mice. Serota et al. (1986b) concluded that dichloromethane “did not induce a treatment-related carcinogenic response in B6C3F₁ mice” under the conditions of this study. An alternative conclusion, as determined by EPA based on the results of the analysis shown in Hazleton Laboratories (1983), is that dichloromethane induced a carcinogenic response in male B6C3F₁ mice as evidenced by small but statistically significant (*p* < 0.05) increases in hepatocellular adenomas and carcinomas at dose levels of 125, 185, and 250 mg/kg-day, and by a marginally increased trend test (*p* = 0.058) for combined hepatocellular adenomas and carcinomas. The incidence in the control groups was almost identical to the mean seen in the historical controls (17.8%, based on 354 male B6C3F₁ mice), so there is no indication that the observed trend is being driven by an artificially low incidence in controls. There is also no indication that the experimental conditions resulted in a systematic increase in the incidence of hepatocellular adenomas and carcinomas. Given the information provided regarding the incidence in historical controls (mean 17.8%, range 5 to

40%), the pattern of results (increased incidence in all four dose groups, with three of these increases significant at a p -value < 0.05) indicates a treatment-related increase.

One reason for the difference between Serota et al. (1986b) and EPA in the interpretation of these data is the difference in the significance level used to evaluate the between-group comparisons. EPA used the standard two-tailed significance level of $p = 0.05$. Serota et al. (1986b) indicate that a two-tailed significance level of $p = 0.05$ was used for all tests. However, Hazleton Laboratories (1983) indicated that a correction factor for multiple comparisons was used specifically for the liver cancer data, reducing the nominal p -value from 0.05 to 0.0125. None of the group comparisons shown in Table 4-15 are statistically significant when a p -value of 0.0125 is used. A multiple comparisons correction is sometimes advocated in situations examining many different types of effects (e.g., ≥ 20 individual causes of death) or many different types of exposures (≥ 20 different chemicals) to protect against inappropriately focusing on spurious findings. Thus, EPA concluded that the use of this multiple comparisons correction factor in the 2-year mouse oral carcinogenicity study is not warranted.

4.2.1.2.3. Chronic oral exposure in Sprague-Dawley rats and Swiss mice (Maltoni et al., 1988). Maltoni et al. (1988) conducted gavage carcinogenicity studies in Sprague-Dawley rats and in Swiss mice. Groups of rats (50/sex/dose level) were gavaged with dichloromethane (99.9% pure) in olive oil at dose levels of 0 (olive oil), 100, or 500 mg/kg-day 4–5 days/week for 64 weeks. This dosing regime was also used for groups of Swiss mice (50/sex/dose level plus 60/sex as controls). Endpoints monitored included clinical signs, BW, and full necropsy at sacrifice (when spontaneous death occurred). For each animal sacrificed, histopathologic examinations were performed on the following organs: brain and cerebellum, zymbal glands, interscapular brown fat, salivary glands, tongue, thymus and mediastinal lymph nodes, lungs, liver, kidneys, adrenals, spleen, pancreas, esophagus, stomach, intestine, bladder, uterus, gonads, and any other organs with gross lesions. High mortality was observed in male and female high-dose rats (data not shown) and achieved significance ($p < 0.01$) in males. The increased mortality became evident after 36 weeks of treatment and led to the termination of treatment at week 64. Explanation of the mortality was not provided by the study authors. As with the rats, high mortality occurred in male and female mice from the high-dose group ($p < 0.01$), and the exposure was terminated after 64 weeks.

Little information is provided regarding nonneoplastic effects (Maltoni et al., 1988). Treatment with dichloromethane did not affect BW in the Sprague-Dawley rats. A reduction in BW was apparent in treated mice after 36–40 weeks of treatment, but no data were shown to determine the magnitude of the effect. The lack of reporting of nonneoplastic findings from the histopathologic examinations precludes assigning NOAELs and LOAELs for possible nonneoplastic effects in these studies.

The Maltoni et al. (1988) studies of Sprague-Dawley rats and Swiss mice did not find distinct exposure-related carcinogenic responses following gavage exposure to dichloromethane at dose levels up to 500 mg/kg-day, although the early termination of the study (at 64 weeks) limits the interpretation of this finding. Dichloromethane exposure was not related to the percentage of either study animal bearing benign and/or malignant tumors or to the number of total malignant tumors per 100 animals. High-dose female rats showed an increased incidence in malignant mammary tumors, mainly due to adenocarcinomas (8, 6, and 18% in the control, 100, and 500 mg/kg dose groups, respectively; the number of animals examined was not provided), but the increase was not statistically significant. A dose-related increase, although not statistically significant, in pulmonary adenomas was observed in male mice (5, 12, and 18% in control, 100, and 500 mg/kg-day groups, respectively). When mortality was taken into account, high-dose male mice that died in the period ranging from 52 to 78 weeks were reported to show a statistically significantly ($p < 0.05$) elevated incidence for pulmonary tumors (1/14, 4/21, and 7/24 in control, 100, and 500 mg/kg-day groups, respectively). Details of this analysis were not provided. EPA applied a Fisher's exact test to these incidences and determined a p -value of 0.11 for the comparison of the 500 mg/kg-day group (7/24) to the controls (1/14).

4.2.2. Inhalation Exposure: Overview of Noncancer and Cancer Effects

Inhalation dichloromethane exposure studies in rats and mice using subchronic and chronic durations identify the CNS, liver, and lungs as potential toxicity targets. Data from other studies indicate that hamsters are less susceptible to the nonneoplastic and neoplastic effects of dichloromethane than are rats and mice.

Increased incidences of nonneoplastic liver lesions were observed in Sprague-Dawley rats exposed to ≥ 500 ppm for 2 years (Nitschke et al., 1988a; Burek et al., 1984), F344 rats exposed to $\geq 1,000$ ppm for 2 years (Mennear et al., 1988; NTP, 1986), and B6C3F₁ mice exposed to $\geq 2,000$ ppm for 2 years (Mennear et al., 1988; NTP, 1986).

Two-year inhalation exposure studies at concentrations of 2,000 or 4,000 ppm dichloromethane produced increased incidences of lung and liver tumors in B6C3F₁ mice (Mennear et al., 1988; NTP, 1986). Additional studies examining mechanistic issues regarding this effect are described in Sections 4.5.2 and 4.5.3 (Maronpot et al., 1995; Foley et al., 1993; Kari et al., 1993). Significantly increased incidences of benign mammary tumors (primarily fibroadenomas) were also observed in male and female F344/N rats exposed by inhalation to 2,000 or 4,000 ppm for 2 years (Mennear et al., 1988; NTP, 1986). In the male rats, the incidence of fibromas or sarcomas originating from the subcutaneous tissue around the mammary gland was also increased, but the difference was not statistically significant. In other studies in Sprague-Dawley rats with exposures of 50–500 ppm (Nitschke et al., 1988a) and 500–3,500 ppm (Burek et al., 1984), the incidence of benign mammary tumors was not increased, but in females, the number of tumors per tumor-bearing rat increased at the higher dose levels.

No obvious clinical signs of neurological impairment were observed in the 2-year bioassays involving exposure concentrations up to 2,000 ppm in F344 rats (Mennear et al., 1988; NTP, 1986) or 3,500 ppm in Sprague-Dawley rats (Nitschke et al., 1988a; Burek et al., 1984). In B6C3F₁ mice exposed to 4,000 ppm there was some evidence of hyperactivity during the first year of the study and lethargy during the second year, with female mice appearing to be more sensitive (Mennear et al., 1988; NTP, 1986). Studies that evaluated batteries of neurobehavioral endpoints following subchronic or chronic inhalation exposure are restricted to one in F344 rats exposed to concentrations up to 2,000 ppm for 13 weeks (Mattsson et al., 1990). No effects were observed >64 hours postexposure in an observational battery, a test of hind-limb grip strength, a battery of evoked potentials, or histologic examinations of brain, spinal cord, or peripheral nerves (see Table 4-26 and Section 4.4.3).

No effects on reproductive performance were found in a two-generation reproductive toxicity study with F344 rats exposed to concentrations up to 1,500 ppm for 14 and 17 weeks before mating of the F0 and F1 generations, respectively (Nitschke et al., 1988b) (described more completely in Section 4.3). Developmental effects following exposure of Long-Evans rats to 4,500 ppm for 14 days prior to mating and during gestation (or during gestation alone) included decreased offspring weight at birth and changed behavioral habituation of the offspring to novel environments (Bornschein et al., 1980; Hardin and Manson, 1980) (see Section 4.3 for more details). In standard developmental toxicity studies involving exposure to 1,250 ppm on GDs 6–15, no adverse effects on fetal development were found in Swiss-Webster mice or Sprague-Dawley rats (Schwetz et al., 1975) (see Section 4.3).

4.2.2.1. Toxicity Studies of Subchronic Inhalation Exposures: General, Renal, and Hepatic Effects

Data pertaining to general (e.g., BW, mortality), hepatic, and renal effects from several inhalation exposure studies in various species with exposure periods of 3–6 months are described below. (Studies providing detailed neurological data are described separately in Section 4.4.3) The earliest study involved several different species with exposures of 5,000 ppm for up to 6 months (Heppel et al., 1944). Two 14-week studies in dogs, monkeys, rats, and mice were conducted with exposures at 0, 1,000, and 5,000 ppm (Haun et al., 1972, 1971; Weinstein et al., 1972) and at 0, 25, and 100 ppm (Haun et al., 1972). Neurological effects and hepatic degeneration were seen at the 1,000 ppm dose. In the lower-dose portion of the Haun et al. (1972) study in mice, decreased CYP levels in liver microsomes and some histopathologic liver changes (fat stains and cytoplasmic vacuolation) were seen at 100 ppm, but more obvious adverse effects were not observed. Leuschner et al. (1984) reported data from a high exposure (10,000 ppm) 90-day study of rats; beagle dogs were also included in this study at an exposure level of 5,000 ppm. No evidence of toxicity was reported by the authors of this study. In a 13-week exposure study conducted by NTP (1986), decreased BWs and increased incidence of

foreign body pneumonia were seen at 8,400 ppm in F344 rats, and histologic changes in the liver in B6C3F₁ mice were seen at 4,200 ppm.

The first experimental study of dichloromethane exposure included dogs, rabbits, guinea pigs, and rats with an exposure of approximately 5,000 ppm for 7 hours/day, 5 days/week for up to 6 months (Heppel et al., 1944). The strains of the animals, the comparability between exposed and unexposed groups (in terms of sex distribution and other attributes), and the process by which animals were chosen for histologic examination are not clearly described in the report. Exposed animals included adult dogs (1 male and 5 females), juvenile dogs (1 male and 1 female born in the exposure chamber and exposed daily from birth), adult rabbits (2 males and 2 females), guinea pigs (14 males), and rats (15 males and 6 females). The nonexposed control group included 14 guinea pigs, 28 rats, 4 rabbits, and an unspecified number of dogs. Exposure produced no significant effects on BWs except in the guinea pigs; after 131 exposures, average BWs were 0.820 and 1.025 kg for exposed and control guinea pigs, respectively. Three exposed guinea pigs died after 35, 90, and 96 exposures. No other deaths occurred except for one exposed female rat that died after 22 exposures and giving birth to a litter. Autopsy showed thrombi in the renal vessels associated with marked cortical infarction. No adverse clinical signs of toxicity (such as decreased activity or incoordination) were observed in exposed animals during the study. Urinalysis, hematology tests, and tests of liver function performed on dogs during the study showed no treatment-related effects. At termination, gross and microscopic examination of the major organs showed no pathological changes after exposure to 5,000 ppm dichloromethane, with the exception that two of the exposed guinea pigs that died showed extensive pneumonia associated with moderate centrilobular fatty degeneration of the liver. The results indicate that 5,000 ppm was a NOAEL for systemic effects in dogs, rabbits, and rats exposed 7 hours/day, 5 days/week for up to 6 months. The findings of three deaths (two with pulmonary congestion and centrilobular fatty degeneration) and 20% decreased average BW among the 14 exposed guinea pigs indicates that 5,000 ppm was a LOAEL in this species.

Haun et al. (1972, 1971) and Weinstein et al. (1972) reported results from studies in which groups of 8 female beagle dogs, 4 female rhesus monkeys, 20 male Sprague-Dawley rats, and 380 female ICR mice were continuously exposed to 0, 1,000, or 5,000 ppm dichloromethane for up to 14 weeks in whole-body exposure chambers. Gross and histopathologic examinations were scheduled to be made on animals that died or were sacrificed during or at termination of the study. At 5,000 ppm, obvious nervous system effects (e.g., incoordination, lethargy) were observed in dogs, monkeys, and mice. At 1,000 ppm, these effects were most apparent in dogs and monkeys (Haun et al., 1971). Food consumption was reduced in all species at 5,000 ppm and in dogs and monkeys at 1,000 ppm. All exposed animals either lost weight or showed markedly decreased BW gains compared with controls. For example, rats exposed to 1,000 or 5,000 ppm for 14 weeks showed average BWs that were roughly 10 and 20% lower than control values. Significant numbers of dogs (4) and mice (123), as well as 1 monkey, died within the

first 3 weeks of exposure to 5,000 ppm. Because of this high mortality, all surviving 5,000 ppm animals were sacrificed at 4 weeks of exposure, except for one half (10) of the rats that went on to survive the 14-week exposure period. At 1,000 ppm, 6/8 dogs died by 7 weeks, at which time the remaining two were sacrificed. Monkeys, rats, and all but a few mice survived exposure to 1,000 ppm for 14 weeks.

Gross examination of tissues showed yellow, fatty livers in dogs that died during exposure to 1,000 or 5,000 ppm, “borderline” liver changes in 3 monkeys exposed to 5,000 ppm, and mottled liver changes in 4/10 rats exposed to 5,000 ppm for 14 weeks (Haun et al., 1971). Comprehensive reporting of the histologic findings from this study were not available, but Haun et al. (1972) reported that the primary target organ was the liver and that in some exposed animals, the kidney was also affected. Light and electron microscopy of liver sections from groups of 4–10 mice sacrificed after 1, 4, 8, and 12 hours and 1, 2, 3, 4, 6, and 7 days of exposure to 5,000 ppm showed hepatocytes with balloon degeneration (dissociation of polyribosomes and swelling of rough endoplasmic reticulum) as early as 12 hours of exposure (Weinstein et al., 1972). The degeneration peaked in severity after 2 days of exposure and, subsequently, partially reversed in severity. Information on possible histopathologic changes in mice exposed to 1,000 ppm was not provided.

The results from this study demonstrate that dogs and mice were more sensitive than were rats and monkeys to lethal effects, nervous system depression, and possibly liver effects from continuous exposure to 1,000 or 5,000 ppm. The results indicate that continuous exposure to 1,000 ppm was an adverse effect level for mortality and effects on the nervous system and liver in dogs (exposed for up to 4 weeks) and for BW changes in rats (exposed for 14 weeks). The 5,000 ppm level induced mortality in beagle dogs, ICR mice, and rhesus monkeys (but not in Sprague-Dawley rats); obvious nervous system effects in dogs, mice, monkeys, and rats; and gross liver changes in dogs, mice, monkeys, and rats.

Haun et al. (1972) also conducted studies with groups of 20 mice, 20 rats, 16 dogs, and 4 monkeys exposed continuously to 0, 25, or 100 ppm dichloromethane for 100 days (14 weeks). The animals presumably were of the same strains and sexes as those used in the studies involving exposure to 1,000 or 5,000 ppm dichloromethane (Haun et al., 1972, 1971; Weinstein et al., 1972). All animals underwent necropsy and histopathologic evaluation at termination of the exposure, but a list of the tissues examined and incidence or severity data were not presented in the report. Hematology and clinical chemistry variables (including COHb levels) were measured in blood samples collected from dogs and monkeys at biweekly or monthly intervals during exposure. COHb levels were elevated in a dose-related manner in monkeys and peaked at about 5% (approximately 0.8% preexposure) after 6 weeks of exposure. COHb levels in dogs were unaffected by the 25 ppm exposure level and rose to about 2% (from about 0.6%) from week 4 in high-dose dogs. Additional groups of mice were included for assessment of hexobarbital sleep

times at monthly intervals; levels of cytochromes P-450, P-420, and b₅ in liver microsomes at monthly intervals; and spontaneous physical activity at several intervals during the study.

No clinical signs of toxicity or alterations in weight gain were seen in any of the species examined. In dogs and monkeys, hematology and clinical chemistry results throughout the study and at termination were unremarkable, as were the results of the gross and histopathologic examinations. In mice exposed to 100 ppm, CYP levels in liver microsomes were significantly decreased (compared with control values) after 30, 60, and 90 days of exposure to 100 ppm, whereas levels of cytochrome b₅ and P-420 decreased after 30 days and increased after 90 days of exposure. At 25 ppm, no significant differences from controls were seen in mouse liver levels of cytochromes. Mice exposed to 25 ppm showed no histopathologic changes, while histologic changes in mice at 100 ppm were restricted to positive fat stains and some cytoplasmic vacuolation in the liver. In rats at both exposure levels, the livers showed positive staining for increased fat, and the kidneys showed evidence of nonspecific tubular degenerative and regenerative changes. Haun et al. (1972) indicate that no distinctively adverse effects were found in monkeys, dogs, rats, or mice continuously exposed to 25 or 100 ppm for up to 14 weeks. Decreased CYP levels in liver microsomes and some histopathologic liver changes (fat stains and cytoplasmic vacuolation) were seen at the 100 ppm dose.

Leuschner et al. (1984) exposed Sprague-Dawley rats (20/sex/dose level) to 0 or 10,000 ppm and beagle dogs (3/sex/dose level) to 0 or 5,000 ppm dichloromethane in whole-body exposure chambers. Exposure periods were 6 hours/day for 90 consecutive days. Endpoints evaluated in both species included clinical signs, food and water consumption, BW, hematology, clinical chemistry, urinalysis, and gross and microscopic evaluation of 27 organs at termination. Electrocardiography and blood pressure measurements were also done in dogs. The only significant effect observed in rats was a slight redness of the conjunctiva 1–10 hours after each exposure. In dogs, compound-related effects were restricted to slight sedation throughout the exposure period and slight erythema lasting up to 10 hours after exposure. In this 90-day study involving daily 6-hour exposures, 10,000 and 5,000 ppm were NOAELs for behavioral, clinical chemistry, hematologic, and histologic signs of toxicity in Sprague-Dawley rats and beagle dogs, respectively.

NTP (1986) exposed groups of F344 rats and B6C3F₁ mice (10/sex/dose level) to target concentrations of 0, 525, 1,050, 2,100, 4,200, or 8,400 ppm dichloromethane 6 hours/day, 5 days/week for 13 weeks in whole-body exposure chambers. Endpoints monitored included clinical signs, BW, and necropsy at termination. Comprehensive sets of tissues and organs in control and high-dose animals were histologically examined; tissues from the lower dose groups were examined to determine the no-observed-effect level. One male and one female rat from the 8,400 ppm exposure group died before the end of the study, but the cause of death was not discussed. The final mean BWs of 8,400 ppm male and female rats were reduced by 23 and 11%, respectively, relative to controls. Foreign-body pneumonia was present in 4/10 male and

6/10 female rats exposed to 8,400 ppm and in 1/10 female rats from the 4,200 ppm exposure group. The liver lipid/liver weight ratios for 8,400 ppm rats of both sexes and 4,200 ppm female rats were significantly lower than in controls. In mice, 4/10 males and 2/10 females exposed to 8,400 ppm died before the end of the study, and these deaths were considered treatment-related. Histologic changes in exposed mice consisted of hepatic centrilobular hydropic degeneration (of minimal to mild severity) in 3/10 males and 8/10 females at 8,400 ppm and in 9/10 females from the 4,200 ppm exposure group. Histologic changes in the 2,100 ppm mouse group were not mentioned. The liver lipid/liver weight ratio for the high-dose female mice was significantly lower than in controls. In this 13-week study involving 6-hour exposure periods for 5 days/week, 4,200 ppm was a NOAEL and 8,400 ppm was a LOAEL for decreased BWs and increased incidence of foreign-body pneumonia in F344 rats. In B6C3F₁ mice, 2,100 ppm was a NOAEL and 4,200 ppm was a LOAEL for histologic changes in the liver.

4.2.2.2. Toxicity Studies from Chronic Inhalation Exposures

Chronic inhalation exposure studies are summarized in Table 4-16. Details of each study are described below, with the results pertaining to nonneoplastic and neoplastic effects summarized in the following sections.

Table 4-16. Studies of chronic inhalation dichloromethane exposures

Reference, strain/species	Number per group	Exposure information	Comments
Mennear et al. (1988); NTP (1986) F344 rats	50/sex/dose	2 yrs, 6 hrs/d, 5 d/wk 0, 1,000, 2,000, 4,000 ppm	Nonneoplastic liver effects and hemosiderosis in males and females (see Table 4-17) Weak trend for neoplastic nodule or hepatocellular carcinoma in females, benign mammary tumors in males and females (see Table 4-18)
Mennear et al. (1988); NTP (1986) B6C3F ₁ mice	50/sex/dose	2 yrs, 6 hrs/d, 5 d/wk 0, 2,000, 4,000 ppm	Varied nonneoplastic effects (see Table 4-19) Liver and lung tumors (adenomas or carcinomas) in males and females (see Table 4-20)
Burek et al. (1984) Syrian hamsters	95/sex/dose	2 yrs, 6 hrs/d, 5 d/wk 0, 500, 1,500, 3,500 ppm	Decreased mortality Increased CoHb at 500 ppm (see Section 4.2.2.2.3)
Burek et al. (1984) Sprague-Dawley rats	92–97/sex/dose	2 yrs, 6 hrs/d, 5 d/wk 0, 500, 1,500, 3,500 ppm	Nonneoplastic liver effects in males and females (see Table 4-21) Increased CoHb at 500 ppm Increased number of benign mammary tumors per tumor bearing rat (females) (see Table 4-21)
Nitschke et al. (1988a) Sprague-Dawley rats	90/dose/sex	2 yrs, 6 hrs/d, 5 d/wk 0, 50, 200, 500 ppm	Nonneoplastic liver effects in males and females (statistically significant in females) (see Table 4-22) Increased CoHb at 50 ppm Increased number of benign mammary tumors per animal in females (see Table 4-23)
Maltoni et al. (1988) Sprague-Dawley rats, female	54–60/dose	2 yrs, 4 hrs/d, 5 d/wk for 7 wks; 7 hrs/d, 5 d/wk for 97 wks 0, 100 ppm	No effects seen on total number of benign or malignant cancers

4.2.2.2.1. Chronic inhalation exposure in F344/N rats (Mennear et al., 1988; NTP, 1986).

NTP conducted a 2-year inhalation exposure study in F344/N rats (Mennear et al., 1988; NTP, 1986). The rats (50/sex/exposure level) were exposed to dichloromethane (>99% pure) by inhalation in exposure chambers 6 hours/day, 5 days/week for 2 years. Exposure concentrations were 0, 1,000, 2,000, or 4,000 ppm. Mean daily concentrations never exceeded 110% of target and were <90% of target in only 23 of 1,476 analyses. Endpoints monitored included clinical signs, mortality, and gross and microscopic examinations of 32 tissues at study termination. Clinical examinations were conducted weekly for 3.5 months and biweekly until month 8. After 8 months, the animals were clinically examined and palpated for tumors and masses monthly until the end of the study.

Dichloromethane exposure did not significantly alter BW gain or terminal BWs (Mennear et al., 1988; NTP, 1986). Survival of male rats was low in all exposed groups and the

control group, and no significant exposure-related differences were apparent. Most deaths occurred during the last 16 weeks of the study. Survival at week 86 was 36/50, 39/50, 37/50, and 33/50 for the control, 1,000, 2,000, and 4,000 ppm groups, respectively. In female rats, there was a trend towards decreased survival, and the survival of high-dose female rats was significantly reduced, possibly due to leukemia. Survival in the females at 86 weeks was 30/50, 22/50, 22/50, and 15/50 for the control, 1,000, 2,000, and 4,000 ppm groups, respectively. Nonneoplastic lesions with statistically significantly elevated incidences compared with controls included hepatocyte cytoplasmic vacuolation and necrosis and liver hemosiderosis in males and females, renal tubular cell degeneration in males and females, splenic fibrosis in males, and nasal cavity squamous metaplasia in females (Table 4-17). The results indicate that 1,000 ppm (6 hours/day, 5 days/week) was a LOAEL for liver changes (hepatocyte cytoplasmic vacuolation and necrosis, hepatic hemosiderosis) in male and female F344/N rats. A NOAEL was not established because effects were observed at the lowest dose.

Table 4-17. Incidences of nonneoplastic histologic changes in male and female F344/N rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Lesion, by sex	Exposure (ppm) ^a			
	Controls			
	0	1,000	2,000	4,000
Males				
n per group ^b	50	50	50	50
Number (%) ^c with: Liver changes				
Hepatocyte cytoplasmic vacuolation	8 (16)	26 (53) ^d	22 (44) ^d	25 (50) ^d
Hepatocyte focal necrosis	7 (14)	23 (47) ^d	6 (12)	16 (32) ^d
Hepatocytomegaly	2 (4)	10 (20)	6 (12)	5 (10)
Hemosiderosis	8 (16)	29 (59) ^d	37 (74) ^d	42 (84) ^d
Bile duct fibrosis	8 (16)	10 (20)	17 (34)	23 (46) ^d
Renal tubular cell degeneration	11 (22)	13 (26)	23 (46) ^d	10 (20) ^d
Splenic fibrosis	2 (4)	6 (12)	11 (22) ^d	8 (16) ^d
Females				
n per group ^c	50	50	50	50
Number (%) ^c with: Liver changes				
Hepatocyte cytoplasmic vacuolation	10 (20)	43 (86) ^d	44 (88) ^d	43 (86) ^d
Hepatocyte focal necrosis	2 (4)	32 (64) ^d	19 (38) ^d	9 (18) ^d
Hepatocytomegaly	3 (6)	10 (20) ^d	18 (36) ^d	5 (10)
Hemosiderosis	19 (38)	29 (58) ^d	38 (76) ^d	45 (90) ^d
Bile duct fibrosis	4 (8)	3 (6)	10 (20) ^d	3 (6)
Renal tubular cell degeneration	14 (28)	20 (40)	22 (44)	25 (51) ^d
Splenic fibrosis	0 (0)	2 (4)	4 (8)	4 (8)
Nasal cavity squamous metaplasia	1 (2)	2 (4)	3 (6)	9 (18) ^d

^a1,000 ppm = 3,474 mg/m³, 2,000 ppm = 6,947 mg/m³, 4,000 ppm = 13,894 mg/m³.

^bNumber of male rats necropsied per group; only 49 1,000 ppm livers were examined microscopically.

^cPercentages were based on the number of tissues examined microscopically per group.

^dStatistical significance not reported in publications but significantly ($p \leq 0.05$) different from controls as calculated by Fisher's exact test.

^eNumber of females necropsied per group; only 49 4,000 ppm kidneys and spleens were examined microscopically.

Sources: Mennear et al. (1988); NTP (1986); Appendix B, Tables C1 and C2 of the NTP (1986) report.

Incidences of mammary fibroadenomas were significantly increased in 4,000 ppm males and 2,000 and 4,000 ppm females compared with controls (Table 4-18). Similar patterns were seen with the combination of fibroadenomas and adenomas (not shown in Table 4-18). In males, subcutaneous tissue fibroma or sarcoma was seen in 1/50, 1/50, 2/50, and 5/50 rats in the 0, 1,000, 2,000, and 4,000 ppm groups, respectively, but these lesions were not seen in females. Incidences of female rats with liver neoplastic nodules or carcinomas (combined) showed a

significant trend test after survival adjustment only, but the incidences at the two highest dose levels were not significantly increased relative to the control (Table 4-18).

Table 4-18. Incidences of selected neoplastic lesions in male and female F344/N rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Neoplastic lesion, by sex	Exposure (ppm) ^a												Trend <i>p</i> -value ^d
	0 (Controls)			1,000			2,000			4,000			
	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	
Males													
n per group	50			50			50			50			
Liver—neoplastic nodule or hepatocellular carcinoma	2	(4)	(10)	3	(6)	(13)	4	(8)	(19)	1	(2)	(6)	0.55
Liver—hepatocellular carcinoma	2	(4)	(10)	1	(2)	(4)	2	(4)	(10)	1	(2)	(6)	Not reported
Lung—bronchoalveolar adenoma or carcinoma	1			1	(2)		2	(4)		1	(2)		
Mammary gland													
Adenoma, adenocarcinoma, or carcinoma	0	(0)		0	(0)		0	(0)		1	(2)		
Subcutaneous tissue fibroma or sarcoma	1	(2)	(6)	1	(2)	(6)	2	(4)	(9)	5	(10)	(23)	0.008
Fibroadenoma	0	(0)	(0)	0	(0)	(0)	2	(4)	(12)	1	(2)	(8)	<0.001
Mammary gland or subcutaneous tissue adenoma, fibroadenoma, fibroma, or sarcoma	1	(2)	(6)	1	(2)	(6)	4	(8)	(21)	9 ^e	(18)	(49)	<0.001
Brain (carcinoma, not otherwise specified, invasive)	0	(0)		1	(2)		0	(0)		0	(0)		
Females													
n per group	50			50			50			50			
Liver—neoplastic nodule or hepatocellular carcinoma	2	(4)	(7)	1	(2)	(2)	4	(8)	(14)	5	(10)	(20)	0.08
Liver—hepatocellular carcinoma	0	(0)	(0)	0	(0)	(0)	1	(2)	(4)	0	(0)	(0)	Not reported
Lung—bronchoalveolar adenoma or carcinoma	1	(2)		1	(2)		0	(0)		0	(0)		
Mammary gland													
Adenocarcinoma or carcinoma	1	(2)		2	(4)		2	(4)		0	(0)		
Adenoma, adenocarcinoma, or carcinoma	1	(2)		2	(4)		2	(4)		1	(2)		
Fibroadenoma	5	(10)	(16)	11 ^e	(22)	(41)	13 ^e	(26)	(44)	22 ^e	(44)	(79)	<0.001
Mammary gland adenoma, fibroadenoma, or adenocarcinoma	6	(12)	(18)	13	(26)	(44)	14 ^e	(28)	(45)	23 ^e	(46)	(86)	<0.001
Brain (carcinoma, not otherwise specified, invasive, and oligodendroglioma) ^f	1	(2)		0	(0)		2	(4)		0	(0)		

^a1,000 ppm = 3,474 mg/m³, 2,000 ppm = 6,947 mg/m³, 4,000 ppm = 13,894 mg/m³.

^bPercentages based on the number of tissues examined microscopically per group; for males, 49 livers and lungs were examined microscopically in the 1,000 ppm groups and only 49 brains were examined microscopically in the 4,000 ppm group. For comparison, incidence in historical controls reported in NTP (1986) were 1% for female liver tumors and 16% for female mammary fibroadenomas.

^cMortality-adjusted percentage.

^dLife-table trend test, as reported by NTP (1986).

^eLife-table test comparison dose group with control <0.05, as reported by NTP (1986).

^fThe oligodendroglioma occurred in the 2,000 ppm group.

Sources: Mennear et al. (1988); NTP (1986); Appendix A and Appendix E, Tables E1 and E2 of the NTP (1986) report.

Incidences for mononuclear cell leukemias in mid- and high-dose female rats were statistically significant after a survival-adjustment analysis. However, Mennear et al. (1988) considered the relationship between exposure to dichloromethane and mononuclear cell leukemia to be equivocal, based on the fact that most male rats had leukemia (34/50, 26/50, 32/50, and 35/50 in controls, 1,000, 2,000, and 4,000 ppm rats, respectively). Other neoplasms that had increased incidences included mesotheliomas (predominantly in the tunica vaginalis) in males (0/50, 2/50, 5/50, and 4/50 in controls, 1,000, 2,000, and 4,000 ppm rats, respectively). This lesion was not considered to be related to dichloromethane exposure because the concurrent control incidence (0/50) for this neoplasm was low relative to earlier inhalation studies conducted at this laboratory (4/100, 4%) and in other NTP studies with male F344/N rats (44/1,727) (mean historical percentage across NTP studies = $3 \pm 2\%$).

NTP (1986) concluded that there was “some evidence of carcinogenicity of dichloromethane” in male F344/N rats as shown by increased incidence of benign mammary gland tumors and “clear evidence of carcinogenicity” of dichloromethane in female F344/N rats as shown by increased incidence of benign mammary gland tumors. The summary of the hepatic effects in rats in the NTP (1986) report also notes the positive trend in the incidence of hepatocellular neoplastic nodules or carcinomas in females which “may have been due to dichloromethane exposure.”

4.2.2.2.2. Chronic inhalation exposure in B6C3F₁ mice (Mennear et al., 1988; NTP, 1986). A 2-year inhalation exposure study in B6C3F₁ mice, similar to that in F344/N rats, was also conducted by NTP (Mennear et al., 1988; NTP, 1986). The mice (50/sex/exposure level) were exposed to dichloromethane (>99% pure) by inhalation at concentrations of 0, 2,000, or 4,000 ppm in exposure chambers 6 hours/day, 5 days/week for 2 years. As with the study in rats, mean daily concentrations in the mice never exceeded 110% of target and were <90% of target in only 23 of 1,476 analyses. Endpoints monitored included clinical signs, mortality, and gross and microscopic examinations of 32 tissues at study termination. Clinical examinations were conducted weekly for 3.5 months and biweekly until month 8. After 8 months, the animals were clinically examined and palpated monthly for tumors and masses until the end of the study.

The BW of 4,000 ppm males was comparable to controls until week 90 and 8–11% below controls thereafter. The BW of 4,000 ppm females was 0–8% lower than that of controls from week 51 to 95 and 17% lower at study termination. No information was provided regarding food consumption during the study. Male and female mice from the high-dose groups (4,000 ppm) were hyperactive during the first year of the study; during the second year, high-dose females appeared lethargic. Exposure was associated with decreased survivability of both male and female mice (males: 39/50, 24/50, and 11/50 and females: 25/50, 25/50, and 8/50 in controls, 2,000, and 4,000 ppm at 104 weeks, respectively). In 4,000 ppm mice, statistically significant incidences of nonneoplastic lesions were found in the liver (cytologic degeneration), testes

(atrophy), ovary and uterus (atrophy), kidneys (tubule casts in males only), stomach (dilatation), and spleen (splenic follicles in males only) (Table 4-19). In 2,000 ppm mice, the only nonneoplastic lesions showing statistically significantly elevated incidences were ovarian atrophy, renal tubule casts, and hepatocyte degeneration in female mice (Table 4-19). The results indicate that 2,000 ppm, the lowest exposure level, was a LOAEL for nonneoplastic changes in the ovaries, kidneys, and livers of female B6C3F₁ mice. A NOAEL was not established because effects occurred at the lowest exposure level.

Table 4-19. Incidences of nonneoplastic histologic changes in B6C3F₁ mice exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Lesion, by sex	Exposure (ppm) ^a		
	Controls		
	0	2,000	4,000
Males: n per group ^b	50	50	50
Number (%) ^c with:			
Liver changes			
Hepatocyte cytoplasmic vacuolation	Not reported	Not reported	Not reported
Hepatocyte focal necrosis	0 (0)	0 (0)	2 (4)
Cytologic degeneration	0 (0)	0 (0)	22 (45) ^d
Testicular atrophy	0 (0)	4 (8)	31 (62) ^d
Renal tubule casts	6 (12)	11 (22)	20 (40) ^d
Stomach dilatation	3 (6)	7 (15)	9 (18) ^d
Splenic follicular atrophy	0 (0)	3 (6)	7 (15) ^d
Females: n per group ^e	50	50	50
Number (%) ^c with:			
Liver changes			
Hepatocyte cytoplasmic vacuolation	Not reported	Not reported	Not reported
Hepatocyte focal necrosis	Not reported	Not reported	Not reported
Cytologic degeneration	0 (0)	23 (48) ^d	21 (44) ^d
Ovarian atrophy	6 (12)	28 (60) ^d	32 (74) ^d
Uterus atrophy	0 (0)	1 (2)	8 (17) ^d
Renal tubule casts	8 (16)	23 (48) ^d	23 (49) ^d
Glandular stomach dilatation	1 (2)	2 (4)	10 (20) ^d
Splenic follicular atrophy	0 (0)	0 (0)	1 (2)

^a2,000 ppm = 6,947 mg/m³, 4,000 ppm = 13,894 mg/m³.

^bNumber of male mice necropsied per group. The number biopsied in the 0, 2,000, and 4,000 ppm dose groups was 50, 49, and 49 for liver; 50, 49, and 50 for renal tubules; 49, 47, and 49 for stomach; and 49, 49, and 48 for spleen.

^cPercentages were based on the number of tissues examined microscopically per group.

^dStatistical significance not reported in publications but significantly different ($p \leq 0.05$) from control as calculated by EPA using Fisher's exact test.

^eNumber of females necropsied per group. The number biopsied in the 0, 2,000, and 4,000 ppm dose groups was 50, 48, and 48 for liver; 50, 47, and 43 for ovaries; 50, 48, and 47 for uterus; 49, 48, and 47 for renal tubule; 49, 47, and 48 for stomach; and 49, 48, and 47 for spleen.

Sources: Mennear et al. (1988); NTP (1986); Appendix C, Tables D1 and D2 of the NTP (1986) report.

At both exposure levels, statistically significantly elevated incidences were found for hepatocellular adenomas (males only), hepatocellular carcinomas, hepatocellular adenomas and carcinomas combined, bronchoalveolar adenomas, bronchoalveolar carcinomas, and bronchoalveolar adenomas and carcinomas combined (Table 4-20). Statistically significant positive trend tests were found for each of these tumor types in female mice. The trend tests were significant for the liver tumors in male mice after life-table adjustment for reduced survival. The only other statistically significant carcinogenic response was for increased incidence of

hemangiosarcomas or combined hemangiomas and hemangiosarcomas in male mice exposed to 4,000 ppm. NTP (1986) concluded that the elevated incidences of liver and lung tumors provided clear evidence of carcinogenicity in male and female B6C3F₁ mice.

Table 4-20. Incidences of neoplastic lesions in male and female B6C3F₁ mice exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Neoplastic lesion, by sex	Exposure (ppm) ^a									Trend p-value ^d
	0 (Controls)			2,000			4,000			
	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	
Males										
Liver										
Hepatocellular adenoma	10	(20)	(23)	14	(29)	(47)	14	(29)	(68)	0.19
Hepatocellular	13	(26)	(30)	15	(30)	(44)	26 ^e	(53)	(76)	0.004
Hepatocellular adenoma or carcinoma	22	(44)	(48)	24	(49)	(67)	33 ^e	(67)	(93)	0.013
Lung										
Bronchoalveolar adenoma	3	(6)	(8)	19 ^e	(38)	(56)	24 ^e	(48)	(79)	<0.001
Bronchoalveolar carcinoma	2	(4)	(5)	10 ^e	(20)	(34)	28 ^e	(56)	(93)	<0.001
Bronchoalveolar adenoma or carcinoma	5	(10)	(12)	27 ^e	(54)	(74)	40 ^e	(80)	(100)	<0.001
Mammary adenocarcinoma ^f	–			–			–			
Hemangioma or hemangiosarcoma, combined	2	(4)	(5)	2	(4)	(8)	6	(12)	(26)	0.08
Females										
Liver										
Hepatocellular adenoma	2	(4)	(7)	6	(13)	(21)	22 ^e	(46)	(83)	<0.001
Hepatocellular carcinoma	1	(1)	(4)	11	(23)	(34)	32 ^e	(67)	(97)	<0.001
Hepatocellular adenoma or carcinoma	3	(6)	(10)	16 ^e	(33)	(48)	40 ^e	(83)	(100)	<0.001
Lung										
Bronchoalveolar adenoma	2	(4)	(7)	23 ^e	(48)	(58)	28 ^e	(58)	(91)	<0.001
Bronchoalveolar carcinoma	1	(1)	(4)	13 ^e	(27)	(46)	29 ^e	(60)	(92)	<0.001
Bronchoalveolar adenoma or carcinoma	3	(6)	(11)	30 ^e	(63)	(83)	41 ^e	(85)	(100)	<0.001
Mammary adenocarcinoma	2	(4)	(8)	3	(6)	(10)	0	(0)	(0)	0.21
Hemangioma or hemangiosarcoma, combined ^f	–			–			–			

^a2,000 ppm = 6,947 mg/m³, 4,000 ppm = 13,894 mg/m³.

^bPercentages based on the number of tissues examined microscopically per group; for males, 49 livers were examined in the 2,000 and 4,000 ppm groups; for females, only 48 livers and lungs and 49 mammary glands were microscopically examined in the 2,000 and 4,000 ppm groups. For comparison, incidence in historical controls reported in NTP (1986) were 28% for male liver tumors, 31% for male lung tumors, 5% for female liver tumors, and 10% for female lung tumors.

^cMortality-adjusted percentage.

^dLife-table trend test, as reported by NTP (1986).

^eLife-table test comparison dose group with control <0.05, as reported by NTP (1986).

^fData not reported.

Sources: Mennear et al. (1988); NTP (1986); Appendix E, Tables E3 and E4)of the NTP (1986) report.

4.2.2.2.3. Chronic inhalation exposure in Syrian hamsters (Burek et al., 1984). Burek et al. (1984) conducted a chronic toxicity and carcinogenicity study in rats and hamsters. In the hamster study, groups of 95 Syrian golden hamsters of each sex were exposed to 0 (filtered air), 500, 1,500, or 3,500 ppm dichloromethane (>99% pure) under dynamic airflow conditions in whole-body exposure chambers 6 hours/day, 5 days/week for 2 years. Exposure started when the animals were approximately 8 weeks of age. Interim sacrifices were conducted at 6, 12, and 18 months. The hamsters were observed daily during exposure days and were palpated monthly for palpable masses starting the third month of the study. BWs were monitored weekly for the first 8 weeks of the study and monthly thereafter. Hematologic determinations included packed cell volume, total erythrocyte counts, total red blood cells, differential leukocyte counts, and hemoglobin concentration. The mean corpuscular volume, mean corpuscular hemoglobin, and MCHC were also determined. A reticulocyte count was also performed on all animals at the 18-month kill and on 10 animals/sex/dose at 24 months. Clinical chemistry determinations included serum AP and ALT activities, blood urea nitrogen levels, and total protein and albumin. Urinary parameters measured were specific gravity, pH, glucose, ketones, bilirubin, occult blood, protein, and urobilinogen. Hematology, clinical chemistries, and urinalysis were performed at interim sacrifices and at termination. COHb was measured after a single 6-hour exposure and following 22 months of exposure. Gross and microscopic examinations were conducted on all tissues. In addition, the weights of the brain, heart, liver, kidneys, and testes were recorded.

In the study using Syrian hamsters (Burek et al., 1984), hamsters were exposed to analytical concentrations of dichloromethane of 510 ± 27 , $1,510 \pm 62$, and $3,472 \pm 144$ ppm for the target concentrations of 500, 1,500, and 3,500 ppm, respectively. No exposure-related clinical signs were observed in the hamsters throughout the study. Significantly decreased mortality was observed in females exposed to 3,500 ppm from the 13th through the 24th month and from the 20th to the 24th month in females exposed to 1,500 ppm. Exposure to dichloromethane had no significant effect on BW or on mean organ weights. Regarding hematology parameters (actual data were not shown), Burek et al. (1984) stated that a few statistically significant changes occurred, but no obvious pattern could be discerned and most values were within the expected range for the animals. There were no exposure-related alterations in clinical chemistry or urinalysis values. Male and female hamsters in all dose groups had significantly elevated COHb values after a single 6-hour exposure and after 22 months of exposure, but at both time points there was no dose-response relationship above the first dose level and no apparent significant differences in the magnitude of the changes between the two time points. For example, mean values (\pm SD) for percentage COHb in male hamsters after 22 months of exposure were $3.3 (\pm 3.5)$, $28.4 (\pm 5.9)$, $27.8 (\pm 2.9)$, and $30.2 (\pm 4.9)$, for the control through 3,500 ppm groups, respectively. Similar values were obtained for females at 22 months and for males and females after the first day of exposure. Pathological evaluation of hamsters showed a lack of evidence of definite target organ toxicity. Specific observations

mentioned by the authors included a trend of increasing hemosiderin in the liver of male hamsters at 6 and 12 months; decreased amyloid deposit in organs, such as the liver, kidneys, adrenal, and thyroid glands in exposed animals; and fewer biliary cysts in the liver. Increased hepatic hemosiderin at the 12-month sacrifice was observed in 1/5, 1/5, 3/5, and 5/5 male hamsters in the control through 3,500 ppm groups, respectively. No exposure-related increased incidences of hepatic hemosiderin or other liver effects were reported for the terminal sacrifice. The exposure-related decreases in geriatric changes (i.e., amyloid deposits and biliary cysts) were more prominent in females and were associated with the increased survivability in the exposed female hamsters compared with controls. The results indicate that 3,500 ppm was a NOAEL for adverse changes in clinical chemistry and hematological variables, as well as for histologic changes in tissues in male and female Syrian golden hamsters. A LOAEL was not established based on the lack of adverse changes in clinical chemistry and hematological variables as well as the absence of histologic changes in tissues in male and female Syrian golden hamsters.

Evaluation of the total number of hamsters with a tumor, the number with a benign tumor, or the number with a malignant tumor revealed no exposure-related differences in male hamsters. In the high-dose female group, there was a statistically significant increase in the total number of benign tumors at any tissue site (the report did not specify which sites), but this was considered to be secondary to the increased survival of this group. Incidences of male or female hamsters with tumors in specific tissues were not statistically significantly elevated in exposed groups compared with control incidences. The results indicate that no statistically significant, exposure-related carcinogenic responses occurred in male or female Syrian golden hamsters exposed (6 hours/day, 5 days/week) to up to 3,500 ppm dichloromethane for 2 years.

4.2.2.2.4. Chronic inhalation exposure in Sprague-Dawley rats (Burek et al., 1984). In the rat study, groups of 92–97 Sprague-Dawley rats of each sex were exposed (similar to the hamster study described in the previous section) to 0, 500, 1,500, or 3,500 ppm dichloromethane 6 hours/day, 5 days/week for 2 years (Burek et al., 1984). Rats were approximately 8 weeks old when exposure started. Interim sacrifices were conducted at 6, 12, 15, and 18 months. Endpoints monitored in rats were the same as in hamsters except that total protein and albumin in blood were not determined in rats. In addition to measurement at scheduled sacrifices, serum ALT activity was also measured after 30 days of exposure. COHb was measured after 6, 11, 18, and 21 months of exposure. Bone marrow cells were collected for cytogenetic studies from 5 rats/sex/dose after 6 months of exposure. The scope of the pathological examinations of the rats was the same as in the hamster study.

No significant exposure-related signs of toxicity were observed in the rats during the study. A significant increase in mortality was seen in high-dose female rats from the 18th to the 24th month of exposure, and this appeared to be exposure-related. Exposure to dichloromethane

had no significant effect on BW gain in either males or females. The only exposure-related alterations in organ weights was a significant increase in both absolute and relative liver weight in high-dose males at the 18-month interim kill and a significant increase in relative liver weight in high-dose females also at 18 months. Statistically significant changes in hematologic parameters were restricted to increased mean corpuscular volume and mean corpuscular hemoglobin values at 15 months in males. The clinical chemistry tests revealed no significant exposure-related effects. Male and female rats in all exposed groups had significantly elevated COHb values at all time points, but no dose-response relationship was apparent. For example, mean (\pm SD) values for percentage COHb after 21 months of exposure were 0.4 (\pm 0.7), 12.8 (\pm 2.6), 14.8 (\pm 4.4), and 12.2 (\pm 5.7) for the control through 3,500 ppm female rat groups, respectively. Exposure-related statistically significant increases in incidences of nonneoplastic lesions were restricted to the liver (Table 4-21). The incidences of males or females with hepatocellular vacuolation consistent with fatty change increased as the exposure concentration increased. Hepatocellular necrosis occurred at elevated incidences in male rats exposed to 1,500 or 3,500 ppm compared with controls, but this endpoint was not reported in the female data. Liver lesions were initially observed after 12 months of treatment. There was some evidence that exposure at the two highest levels provided some inhibition of the age-related glomerulonephropathy observed in the control rats at termination. The results indicate that the lowest exposure level, 500 ppm, was a LOAEL for fatty changes in the liver of male and female Sprague-Dawley rats and that exposure to \geq 1,500 ppm induced hepatocellular necrosis in males.

Table 4-21. Incidences of selected nonneoplastic and neoplastic histologic changes in male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Lesion, by sex	Exposure (ppm) ^a			
	0 (Controls)	500	1,500	3,500
Males—n per group	92	95	95	97
Number (%) with:				
Liver changes				
Hepatocellular necrosis	2 ^b (2)	8 (8)	10 (10) ^c	11 (11) ^c
Coagulation necrosis	— ^d	—	—	—
Hepatic vacuolation (fatty change)	16 ^b (17)	36 (38) ^c	43 (45) ^c	52 (54) ^c
Foci of altered hepatocytes	—	—	—	—
Foci of altered hepatocytes, basophilic	—	—	—	—
Area of altered hepatocytes	—	—	—	—
Multinucleated hepatocytes	—	—	—	—
Glomerulonephropathy				
Severe	70 ^b (76)	62 (65)	53 (56) ^c	39 (40) ^c
Any degree	92 ^{b,e} (100)	91 (96)	93 (98)	90 (93)
Mammary changes				
Rats with benign mammary tumors	7 ^b (8)	3 (3)	7 (7)	14 (14)
Total number of benign mammary tumors	8	6	11	17
Number of tumors per tumor-bearing rat ^f	1.1	2.0	1.6	1.2
Females—n per group	96	95	96	97
Number (%) with:				
Liver changes				
Hepatocellular necrosis	—	—	—	—
Coagulation necrosis	1 ^b (1)	0 (0)	2 (2)	7 (7)
Hepatic vacuolation (fatty change)	33 ^b (34)	49 (52) ^c	56 (58) ^c	63 (65) ^c
Foci of altered hepatocytes	35 ^b (37)	36 (38)	27 (28)	50 (52) ^c
Foci of altered hepatocytes, basophilic	3 ^b (3)	0 (0)	4 (4)	10 (10)
Area of altered hepatocytes	19 ^b (20)	24 (25)	28 (29)	35 (36) ^c
Multinucleated hepatocytes	7 ^b (7)	36 (38) ^c	34 (35) ^c	29 (30) ^c
Glomerulonephropathy				
Severe	5 (5)	3 (3)	4 (4)	5 (5)
Any degree	62 ^b (65)	64 (67)	59 (62)	48 (50) ^c
Mammary changes				
Rats with benign mammary tumors	79 (82)	81 (85)	80 (83)	83 (86)
Total number of benign mammary tumors	165	218	245	287
Number of tumors per tumor-bearing rat ^f	2.1	2.7	3.1	3.5

^a500 ppm = 1,737 mg/m³, 1,500 ppm = 5,210 mg/m³, 3,500 ppm = 12,158 mg/m³.

^bSignificant dose-related trend—Cochran-Armitage trend test $p < 0.05$.

^cSignificantly higher than control incidence by Fisher's exact test.

^d— Reported as “no exposure effect” by Burek et al. (1984); data not given.

^eBurek et al. (1984) reported that 93/92 male mice had glomerulonephropathy in the kidney in the control group; the incidence was corrected to 92/92.

^fCalculated by EPA.

Source: Burek et al. (1984).

In females, an increasing trend was seen in the incidence of foci or areas of altered hepatocytes. Female rats in all exposed groups showed increased incidence of multinucleated hepatocytes in the centrilobular region compared with controls, but there was no evidence of increasing incidence or severity with increasing exposure level (Table 4-21). The foci and areas were apparent after 12 months, and their number and size increased thereafter, but incidences for neoplastic nodules in the liver or hepatocellular carcinomas were not increased in any exposure group. A statistically significant increased incidence of salivary gland sarcomas was reported for male rats exposed to 3,500 ppm. Burek et al. (1984) considered this finding unusual and inconsistent with other existing data because the primary target organ for dichloromethane seems to be the liver. Incidences of rats with benign mammary gland tumors were not statistically significantly higher in exposed male or female groups compared with controls, and exposed male and female groups showed no significantly increased incidences for malignant mammary gland tumors. The average number of benign mammary tumors per tumor-bearing rat increased with increasing exposure level. In females, the values were 2.1, 2.7, 3.1, and 3.5 in the control through 3,500 ppm groups, respectively; males showed a similar response with increasing exposure level, albeit to a lesser extent (Table 4-21). Burek et al. (1984) concluded that the significance of this benign mammary tumor response (i.e., increase in number of tumors per tumor-bearing rat) was unknown but speculated that the predisposition of this strain of rats (historical control incidences of females with benign mammary tumors normally exceeded 80%) plus the high exposure to dichloromethane may have resulted in the response.

4.2.2.2.5. *Chronic inhalation exposure in Sprague-Dawley rats (Nitschke et al., 1988a).*

Nitschke et al. (1988a) examined the toxicity and carcinogenicity of lower concentrations of dichloromethane in Sprague-Dawley rats. Groups of 90 male and 90 female rats were exposed to 0, 50, 200, or 500 ppm dichloromethane (>99.5% pure) 6 hours/day, 5 days/week for 2 years. Interim sacrifices were conducted at 6, 12, 15, and 18 months (five rats/sex/interval). An additional group of 30 female rats was exposed to 500 ppm for 12 months and then exposed to room air for up to an additional 12 months, and another group of 30 female rats was exposed to room air for the first 12 months, followed by exposure to 500 ppm for the last 12 months of the study. These latter groups were included to examine temporal relationships between exposure and potential carcinogenic response. All groups of rats were examined daily for signs of toxicity and all rats were examined for palpable masses prior to the initial exposure and at monthly intervals after the first 12 months. BW was checked twice a month for the first 3 months and monthly thereafter. Blood samples were collected at interim sacrifices and analyzed for total bilirubin, cholesterol, triglycerides, potassium, estradiol, follicle-stimulating hormone, and luteinizing hormone levels. In addition, COHb was determined at multiple times in blood collected from the tail vein. DNA synthesis (incorporation of ³H-thymidine as a measure of cellular proliferation) was measured in the liver of separate groups of female rats after exposure

to the various concentrations for 6 and 12 months (four females/exposure group/interval). All rats were subjected to a complete necropsy, and sections from most tissues were processed for microscopic examination.

Exposure to dichloromethane at any of the exposure levels did not significantly alter mortality rates, BWs, organ weights, clinical chemistry values, or plasma hormone levels (Nitschke et al., 1988a). Blood COHb was elevated in a dose-related manner but not in an exposure duration-related fashion, suggesting lack of accumulation with repeated exposures. For example, mean (\pm SD) values for percentage COHb were 2.2 (\pm 1.3), 6.5 (\pm 1.1), 12.5 (\pm 0.8), and 13.7 (\pm 0.6) for male rats in the control through 500 ppm groups, respectively, at the terminal sacrifice. These values were similarly affected at the 6-month and 12-month intervals (e.g., respective values for males were 0.3 [\pm 0.7], 2.8 [\pm 0.3], 9.6 [\pm 1.2], and 12.7 [\pm 1.6] at the 12-month sacrifice).

The results of the thymidine incorporation experiment revealed no detectable alteration in the rate of liver DNA synthesis in the exposed groups compared with controls. Statistically significantly increased incidences of nonneoplastic liver lesions (hepatic vacuolation and multinucleated hepatocytes) occurred only in females in the 500 ppm group (Table 4-22). Male rat incidence for hepatocyte vacuolation was elevated at 500 ppm but not to a statistically significant degree. In the group of female rats exposed for only 12 months to 500 ppm, significantly increased incidences of nonneoplastic lesions compared with controls were restricted to liver cytoplasmic vacuolization (16/25 = 64%) and multinucleated hepatocytes (9/25 = 36%) in rats exposed during the first 12 months of the study; rats exposed only during the last 12 months of the study showed no elevated incidences of the liver lesions.

Table 4-22. Incidences of selected nonneoplastic histologic changes in male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Lesion, by sex	Exposure (ppm) ^a						
	0 (Controls)	50	200	500	Trend <i>p</i> -value ^b	Late 500 ^c	Early 500 ^c
Males—n per group	70	70	70	70		NA ^d	NA
Number (%) with:							
Hepatic vacuolation (fatty change)	22 (31)	— ^e	—	28 (40)			
Multinucleated hepatocytes	—	—	—	—			
Females—n per group	70	70	70	70		25	25
Number (%) with:							
Hepatic vacuolation (fatty change)	41 (59)	42 (60)	41 (59)	53 (76) ^f	0.01	15 (60)	16 (64) ^f
Multinucleated hepatocytes	8 (11)	6 (9)	12 (17)	27 (39) ^f	<0.0001	3 (12)	9 (36) ^f

^a50 ppm = 174 mg/m³, 200 ppm = 695 mg/m³, 500 ppm = 1,737 mg/m³.

^bCochran-Armitage trend test.

^cLate 500 = no exposure for first 12 mo followed by 500 ppm for last 12 mo; early 500 = 500 ppm for first 12 mo followed by no exposure for last 12 mo.

^dNA = there were no male rats in these exposure groups.

^e— = Incidences not reported.

^fSignificantly ($p \leq 0.05$) higher than control incidence by Fisher's exact test (Nitschke et al., 1988a).

Source: Nitschke et al. (1988a).

A few fibrosarcomas or undifferentiated sarcomas in the mammary gland were seen in the exposed rats, but these incidences were not statistically significant (Table 4-23). Significantly increased incidences of rats with neoplastic lesions were restricted to benign mammary tumors in female rats exposed for 2 years to 200 ppm compared with controls (61/69 = 88%) (Table 4-23). However, significantly elevated incidences of this tumor type were not observed in 500 ppm females, and the 200 ppm incidence was within the range of historical control values for benign mammary tumors in female Sprague-Dawley rats (79–82%) from two other chronic toxicity/carcinogenicity studies from the same laboratory. A slight but statistically significant increase in the number of palpable masses in subcutaneous or mammary regions (at 23 months) per tumor-bearing rat was observed only in the 500 ppm female group. The numbers of benign mammary tumors per tumor-bearing rat were slightly elevated in the exposed groups compared with control groups, but no statistical analysis of this variable was performed. In female rats exposed to 500 ppm (during the first or second 12 months of the study), slight but statistically significant elevations were found in the number of palpable masses in subcutaneous or mammary regions per tumor-bearing rat; the numbers of benign mammary tumors per tumor-bearing rat were slightly elevated compared with those of controls, but statistical analysis of this variable was not performed.

Table 4-23. Incidences of selected neoplastic histologic changes in male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Lesion, by sex	Exposure (ppm) ^a					
	0 (Controls)	50	200	500	Late 500 ^b	Early 500 ^b
Males—n per group	70	70	70	70	0	0
Number (%) ^c with:						
Liver tumors	0 (0)	0 (0)	0 (0)	0 (0)		
Lung tumors	0 (0)	0 (0)	0 (0)	0 (0)		
Mammary gland tumors						
Adenocarcinoma or carcinoma	0 (0)	0 (0)	0 (0)	0 (0)		
Fibroadenoma	2 (4)	0 (0)	2 (3)	2 (3)		
Fibroma	6 (11)	1 (6)	6 (11)	10 (16)		
Fibrosarcoma	0 (0)	1 (6)	1 (6)	0 (0)		
Undifferentiated sarcoma	0 (0)	2 (4)	0 (0)	0 (0)		
Fibroma, fibrosarcoma, or undifferentiated sarcoma ^d	6 (11)	4 (6)	7 (12)	10 (16)		
Brain tumors						
Astrocytoma or glial cell	0 (0)	1 (1)	2 (3)	1 (1)		
Granular cell	0 (0)	0 (0)	0 (0)	1 (1)		
Females—n per group	70	70	70	70	25	25
Number (%) ^c with:						
Liver tumors						
Neoplastic nodule(s)	4 (6)	4 (6)	3 (4)	4 (6)	0 (0)	1 (4)
Hepatocellular carcinoma	1 (1)	0 (0)	2 (3)	1 (1)	0 (0)	0 (0)
Lung tumors	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Mammary gland tumors						
Adenocarcinoma or carcinoma	6 (9)	5 (7)	4 (6)	4 (6)	3 (12)	2 (8)
Adenoma	1 (1)	1 (1)	2 (3)	1 (1)	2 (8)	0 (0)
Fibroadenoma	51 (74)	57 (83)	60 (87)	55 (80)	22 (88)	23 (92)
Fibroma	0 (0)	1 (1)	0 (0)	1 (1)	1 (4)	1 (1)
Fibrosarcoma	1 (1)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Number with palpable masses in subcutaneous or mammary region	55 (78)	56 (81)	60 (87)	59 (86)	22 (88)	23 (92)
Number of palpable masses in subcutaneous or mammary region per tumor-bearing rat	1.8	2.1	2.0	2.2 ^e	2.3 ^e	2.7 ^e
Number with benign tumors	52 (75)	58 (84)	61 ^f (88)	55 (80)	23 (92)	23 (92)
Number of benign tumors per tumor-bearing rat	2.0	2.3	2.2	2.7	2.2	2.6

(Table 4-23 continues on next page)

Table 4-23. Incidences of selected neoplastic histologic changes in male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Lesion, by sex	Exposure (ppm) ^a					
	0 (Controls)	50	200	500	Late 500 ^b	Early 500 ^b
Brain tumors						
Astrocytoma or glial cell	0 (0)	0 (0)	0 (0)	2 (3)	0 (0)	0 (0)
Granular cell	1 (1)	0 (0)	0 (0)	1 (1)	0 (0)	0 (0)

^a50 ppm = 174 mg/m³, 200 ppm = 695 mg/m³, 500 ppm = 1,737 mg/m³.

^bLate 500 = no exposure for first 12 mo followed by 500 ppm for last 12 mo; early 500 = 500 ppm for first 12 mo followed by no exposure for last 12 mo. No males were included in these exposure groups.

^cPercentages were based on the number of tissues examined microscopically per group. In males, 69 lungs were examined microscopically in the 50 ppm groups, and only 57, 65, 59, and 64 mammary glands were examined in the control, 50, 200, and 500 ppm groups, respectively. In females, 69 mammary glands were examined microscopically in the control, 50, 200, and 500 ppm groups.

^dEPA summed across these three tumors, assuming no overlap.

^eSignificantly ($p \leq 0.05$) higher than control by Haseman's test (Nitschke et al., 1988a).

^fSignificantly ($p \leq 0.05$) higher than control incidence by Fisher's exact test (Nitschke et al., 1988a).

Source: Nitschke et al. (1988a).

A statistically significant increased incidence of brain or CNS tumors was not observed, but six astrocytoma or glioma (mixed glial cell) tumors were seen in the exposed groups (four in males, two in females). The authors concluded that there was no distinct exposure-related malignant carcinogenic response in male or female Sprague-Dawley rats exposed (6 hours/day, 5 days/week) to up to 500 ppm dichloromethane for 2 years (Nitschke et al., 1988a).

4.2.2.2.6. Chronic inhalation exposure in Sprague-Dawley rats (Maltoni et al., 1988). Maltoni et al. (1988) conducted an inhalation exposure study in Sprague-Dawley rats. Two groups of female rats (54–60/dose) were exposed to 0 or 100 ppm dichloromethane for 104 weeks. The exposure period was 4 hours/day, 4 days/week for 7 weeks and then 7 hours/day, 5 days/week for 97 weeks. Endpoints monitored included clinical signs, BW, and full necropsy at sacrifice (when spontaneous death occurred). For each animal sacrificed, histopathologic examinations were performed on the following organs: brain and cerebellum, zymbal glands, interscapular brown fat, salivary glands, tongue, thymus and mediastinal lymph nodes, lungs, liver, kidneys, adrenals, spleen, pancreas, esophagus, stomach, intestine, bladder, uterus, gonads, and any other organs with gross lesions.

There was no evidence of increased mortality in the exposed group, and there was no effect on BW (Maltoni et al., 1988). Little information was provided regarding nonneoplastic effects, precluding identification of NOAELs and LOAELs for nonneoplastic effects in this study. Dichloromethane exposure was not related to the percentage of rats with benign tumors

and malignant tumors, malignant tumors, or the number of total malignant tumors per 100 animals. The percentage of rats with benign mammary tumors was 40.0% in controls and 64.8% in the exposed group, and the percentage of malignant mammary tumors was 3.3 and 5.5% in controls and exposed rats, respectively. Neither of these differences was statistically significant.

4.3. REPRODUCTIVE/DEVELOPMENTAL STUDIES—ORAL AND INHALATION

Reproductive and development studies of dichloromethane exposure are summarized in Table 4-24 and described in detail below. No effects on reproductive performance were observed in a 90-day gavage study in Charles River CD rats with doses up to 225 mg/kg-day (General Electric Company, 1976) or in a two-generation reproductive toxicity study with F344 rats exposed to concentrations up to 1,500 ppm for 14 or 17 weeks before mating of the F0 and F1 generations, respectively, as well as during the F1 gestational period (GDs 0–21) (Nitschke et al., 1988b). Reproductive parameters (e.g., number of litters, implants/litter, live fetuses/litter, percent dead/litter, percent resorbed/litter, or fertility index⁴) were also examined in a study of male Swiss-Webster mice administered dichloromethane (250 or 500 mg/kg) by subcutaneous injection 3 times/week for 4 weeks, and in a similar study involving inhalation exposure to 0, 100, 150, or 200 ppm dichloromethane; no statistically significant effects were seen in either protocol, although some evidence of a decrease in fertility index was seen in the 150 and 200 ppm groups (Raje et al., 1988).

⁴Fertility index defined as number of females impregnated divided by total number of females mated times 100.

Table 4-24. Summary of studies of reproductive and developmental effects of dichloromethane exposure in animals

Species and n	Exposure dose	Exposure period	Results	Reference
Gavage or subcutaneous				
Charles River rats (males and females), 10 per sex per dose group	0, 25, 75, 225 mg/kg (gavage)	90 d before mating (10 d between last exposure and mating period)	No effects on fertility index, number of pups per litter, pup survival, or F1 BW, hematology, and clinical chemistry tests (up to 90 d of age)	General Electric Company (1976)
Swiss-Webster mice (males), 20 per group	0, 250, 500 mg/kg (subcutaneous injection), 3 × per wk	4 wks prior to mating (1 wk between last exposure and mating period)	No effects on fertility index, number of litters, implants per litter, live fetuses per litter, resorption rate; no testicular effects	Raje et al. (1988)
F344 rats (females), 17–21 per dose group	0, 337.5, 450 mg/kg-d (gavage)	GDs 6–19	Decreased maternal weight gain; no effect on resorption rate, number of live litters, implants, live pups, or pup weight	Narotsky and Kavlock (1995)
Inhalation				
F344 rats (males and females, two generation), 30 per sex per dose group (F0 and F1)	0, 100, 500, 1,500 ppm, 6 hrs/d	14 wks prior to mating (F0), GDs 0–21, and 17 wks prior to mating, beginning PND 4, (F1)	No effect on fertility index, litter size, neonatal survival, growth rates, or histopathologic lesions	Nitschke et al. (1988b)
Swiss-Webster mice (males), 20 per group	0, 100, 150, 200 ppm, 2 hrs/d	6 wks, prior to mating (2 d between last exposure and mating period)	Fertility index decreased in 150 and 200 ppm group (statistical significance depends on test used); no effects on number of litters, implants per litter, live fetuses per litter, resorption rate; no testicular effects	Raje et al. (1988)
Long-Evans rats (female), 16–21 per dose group	0, 4,500 ppm	12–14 d before mating and/or GDs 1–17	Gestational exposure resulted in increased absolute and relative maternal liver weight, decreased fetal BW	Hardin and Manson (1980)
Long-Evans rats (female), 16–21 per dose group	0, 4,500 ppm	12–14 d before mating and/or GDs 1–17	Altered rate of behavioral habituation to novel environment (at 4 d of age). No effect on crawling (at 10 d), movement in photocell cage (15 d), use of running wheel (45–108 d), and shock avoidance (4 mo)	Bornschein et al. (1980)
Swiss-Webster mice (females), 30–40 per group	0, 1,250 ppm, 7 hrs/d	GDs 6–15	Increased incidence of extra center of ossification in sternum, increased (~10%) maternal blood COHb, increased maternal weight, increased maternal absolute liver weight	Schwetz et al. (1975)
Sprague-Dawley rats (females), 20–35 per group	0, 1,250 ppm, 7 hrs/d	GDs 6–15	Decreased incidence of lumbar ribs or spurs, increased incidence of delayed ossification of sternbrae, increased (~10%) maternal blood COHb, increased maternal absolute liver weight	Schwetz et al. (1975)

PND = postnatal day

Following exposure of pregnant F344 rats to gavage doses of up to 450 mg/kg-day on GDs 6–19, maternal weight gain was decreased, but no effects were found on the number of resorption sites, pup survivability, or pup weights at postnatal days (PNDs) 1 or 6 (Narotsky and Kavlock, 1995). The developmental effects following exposure of Long-Evans rats to 4,500 ppm for 14 days prior to mating and during gestation (or during gestation alone) were decreased offspring weight at birth and changed behavioral habituation of the offspring to novel environments (Bornschein et al., 1980; Hardin and Manson, 1980) (see Section 4.3.2 for more details). In standard developmental toxicity studies involving exposure to 1,250 ppm on GDs 6–15, no adverse effects on fetal development were found in Swiss-Webster mice or Sprague-Dawley rats, but the incidence of minor skeletal variants (e.g., delayed ossification of sternbrae) was increased. (Schwetz et al., 1975) (see Section 4.3.2).

4.3.1. Reproductive Toxicity Studies

4.3.1.1. *Gavage and Subcutaneous Injection Studies*

In a study sponsored by the General Electric Company (1976), Charles River CD rats (10/sex/dose level) were administered 0, 25, 75, or 225 mg/kg-day dichloromethane by gavage in water for 90 days. The test material was dichloromethane (of unspecified purity) purchased from Dow Chemical Company. At approximately 100 days of age, the rats were mated 1 to 1 to produce the F1 generation. F1 rats (15/sex/dose level) received the same treatment as F0 for 90 days, at which time they were sacrificed and necropsied. Comprehensive sets of 24 tissues from 10 male and 10 female F1 rats from the control and 225 mg/kg-day groups were examined microscopically after embedding, sectioning, and staining. F1 rats were monitored for clinical signs, BW effects, and food consumption. Reproductive parameters examined were fertility index, number of pups per litter, and pup survival. F1 rats also underwent hematology and clinical chemistry tests and urinalysis at 1, 2, and 3 months of the study and ophthalmoscopic examination at 3 months. There were no significant compound-related alterations in any of the endpoints monitored.

Raje et al. (1988) administered dichloromethane (250 or 500 mg/kg) by subcutaneous injection 3 times/week for 4 weeks to male Swiss-Webster mice (20/group). Mating with unexposed females started 1 week after the last exposure and continued for 2 weeks. After the mating period, the males were sacrificed and the testes were examined microscopically. On GD 17, the females were sacrificed and the uterine horns examined for live, dead, or resorbed fetuses. The authors reported that exposure to dichloromethane had no statistically significant effects on number of litters, implants/litter, live fetuses/litter, percent dead/litter, percent resorbed/litter, or fertility index. Examination of the testes showed no significant alterations compared with controls.

4.3.1.2. Inhalation Studies

Nitschke et al. (1988b) conducted a two-generation reproductive toxicity study in rats. Groups of F344 rats (30/sex/dose level) were exposed by inhalation in whole-body chambers to 0, 100, 500, or 1,500 ppm dichloromethane (99.86% pure) 6 hours/day, 5 days/week for 14 weeks and then mated to produce the F1 generation. Exposure of dams continued after mating on GDs 0–21 but was interrupted until PND 4. After weaning, 30 randomly selected F1 pups/sex/dose level were exposed as the parental generation for 17 weeks and subsequently mated to produce the F2 generation. The results showed no statistically significant exposure-related changes in reproductive performance indices (fertility, litter size), neonatal survival, growth rates, or histopathologic lesions in F1 (Table 4-25) or F2 weanlings sacrificed at time of weaning. According to the authors, none of the values in Table 4-25 were significantly different from control values ($\alpha = 0.05$).

Table 4-25. Reproductive outcomes in F344 rats exposed to dichloromethane by inhalation for 14 weeks prior to mating and from GDs 0–21

	Exposure (ppm) ^a			
	0	100	500	1,500
Fertility index ^b	77%	77%	63%	87%
Gestation index ^c	100%	100%	100%	100%
Gestation survival index ^d	99.6%	100%	100%	96.6%
4-d survival index ^e	91.0%	95.2%	98.5%	98.6%
28-d survival index ^f	99.4%	99.4%	100%	99.5%
Sex ratio on d 1 (M:F)	48:52	50:50	50:50	52:48
Litter size				
D 0	11 ± 2	10 ± 2	10 ± 3	11 ± 2
D 28	7 ± 2	7 ± 2	7 ± 2	8 ± 2
Pup BWs, g				
D 1	5.2 ± 0.4	5.3 ± 0.5	5.3 ± 0.4	5.2 ± 0.4
D 4	7.4 ± 0.7	7.5 ± 1.1	7.7 ± 0.7	7.3 ± 0.7
D 28, male	44.6 ± 5.8	45.9 ± 5.0	47.0 ± 5.4	45.0 ± 5.9
D 28, female	43.2 ± 4.3	43.8 ± 4.5	44.4 ± 5.7	43.0 ± 4.8

^a100 ppm = 347 mg/m³, 500 ppm = 1,737 mg/m³, 1,500 ppm = 5,210 mg/m³.

^bNumber of females delivering a litter expressed as a percentage of females placed with a male.

^cNumber of females delivering a live litter expressed as a percentage of the number of females delivering a litter.

^dPercentage of newborn pups that were alive at birth.

^ePercentage of pups surviving to d 4.

^fPercentage of pups alive on d 4 and surviving to d 28.

Source: Nitschke et al. (1988b).

Raje et al. (1988) exposed groups of male Swiss-Webster mice (20/group) to 0, 100, 150, or 200 ppm dichloromethane (HPLC grade, JT Baker Chemical Co.) in inhalation chambers

2 hours/day, 5 days/week for 6 weeks. Mating with unexposed females started 2 days after the last exposure. As in the subcutaneous injection protocol described in the previous section, after the 2-week mating period, the males were sacrificed and the females were sacrificed on GD 17. Exposure of the male mice to dichloromethane had no statistically significant effects on number of litters, implants/litter, live fetuses/litter, percent dead/litter, or percent resorbed/litter, and no significant alterations in the testes were noted. The fertility index was 95, 95, 80, and 80% in the control, 100, 150 and 200 ppm groups, respectively. This decrease was not statistically significant as reported by the authors. Details of the statistical analyses were not provided. The overall χ^2 *p*-value was 0.27. Using a Cochran-Armitage exact trend test on these data, EPA calculated a one-sided *p*-value of 0.059. Individual *p*-values for the comparison of each group with the control group were 0.97, 0.17, and 0.17 for the 100, 150, and 200 ppm groups, respectively. The results for the combined 150 and 200 ppm groups were statistically different from the combined controls and 100 ppm group (Fisher's exact test, one-sided *p*-value = 0.048), suggesting a NOAEL of 100 ppm and LOAEL of 150 ppm.

4.3.2. Developmental Toxicity Studies

The metabolism of dichloromethane into CO by CYP2E1 raises concerns pertaining to developmental neurotoxicity. Gestational exposure to CO results in developmental toxicity and there are reports indicating that exposures as low as 75 ppm CO can result in significant neurological effects in offspring (Giustino et al., 1999). Neurobehavioral deficits in offspring include impaired avoidance behavior (De Salvia et al., 1995) and memory (Giustino et al., 1999). Neurochemical changes, such as abnormal dopaminergic function (Cagiano et al., 1998) and disruption of neuronal proliferation (Fechter, 1987), have also been observed. Oral and inhalation dichloromethane exposure studies have demonstrated increased blood CO levels (see Section 3.3). In addition, increased blood CO levels were seen in rat fetuses exposed through maternal inhalation to 500 ppm dichloromethane on GD 21 (Anders and Sunram, 1982), and placental transfer of dichloromethane also occurs (Withey and Karpinski, 1985; Anders and Sunram, 1982).

4.3.2.1. Gavage Studies and Culture Studies

Narotsky and Kavlock (1995) evaluated developmental effects of dichloromethane (99.9% pure) in F344 rats (17–21/dose group) treated with 0, 337.5, or 450 mg/kg-day dichloromethane by gavage in corn oil on GDs 6–19. Dams were weighed on GDs 6, 8, 10, 13, 16, and 20 and allowed to deliver naturally. They were sacrificed on PND 6 to count uterine implantation sites. Pups were grossly examined for developmental abnormalities and weighed on PNDs 1, 3, and 6. Dead pups or pups with no gross abnormalities were sacrificed and examined for soft tissue abnormalities. Maternal weight gain during pregnancy was significantly reduced in high-dose dams (by 33%, as estimated from Figure 5 of the paper); this group also

exhibited rales and nasal congestion. Treatment with dichloromethane did not induce resorptions or alter the number of live litters on PND 1 or 6, the number of implants, the number of live pups on PND 1 or 6, or pup weight per litter. No gross or soft tissue abnormalities were observed.

Rat embryos in culture medium were exposed to 0, 3.46, 6.54, 9.79, or 11.88 $\mu\text{mol/mL}$ dichloromethane for 40 hours. At the end of the exposure, embryos were observed for development of yolk sac vasculature, crown-rump length, total embryonic protein content, and number of somite pairs. A concentration of dichloromethane of 6.54 $\mu\text{mol/mL}$ of culture medium resulted in decreased crown-rump length, decreased somite number, and decreased amount of protein per embryo, whereas no effects were seen at 3.46 $\mu\text{mol/mL}$ (Brown-Woodman et al., 1998). A time-course experiment conducted with a concentration of dichloromethane of 9.22 $\mu\text{mol/mL}$ showed that marked differences in growth and development from controls were not significant until about 8 hours of culture. Brown-Woodman et al. (1998) noted that the concentrations that caused embryotoxicity in this study were much higher than those found in individuals studied under controlled exposure conditions and comparable to those found in postmortem blood after fatal inhalation.

4.3.2.2. Inhalation Studies

Schwetz et al. (1975) exposed pregnant Swiss-Webster mice (30–40/group) and Sprague-Dawley rats (20–35/group) by inhalation in whole-body chambers to 0 or 1,250 ppm dichloromethane (97.86% pure) 7 hours/day on GDs 6–15. Maternal BWs were recorded on GDs 6, 10, and 16 and on the day of sacrifice (GD 18 for mice, GD 21 for rats). At sacrifice, uterine horns were excised and examined for fetal position and number of live, dead, or absorbed fetuses. Fetuses were observed for gross, soft tissue, and skeletal abnormalities. The only effects seen on developing fetuses were changes in the incidence of minor skeletal variants. In rats, the incidence of lumbar ribs or spurs was significantly decreased compared with controls, whereas the incidence of delayed ossification of sternebrae was significantly greater than in controls. In mice, a significant number of litters contained pups with a single extra center of ossification in the sternum. Exposure to dichloromethane produced significantly elevated blood COHb content in dams of both species (approximately 9–10% after 10 exposures versus 1–2% in controls). BWs in exposed mouse dams were significantly increased (11–15%) compared with those in controls but were not affected in exposed rat dams. Mean absolute liver weights of exposed dams of both species were significantly elevated compared with controls, but mean relative liver weights were not affected. The results indicate that 1,250 ppm was a LOAEL for minimal maternal effects (increased COHb and increased absolute liver weight) and adverse effects on the fetuses.

Hardin and Manson (1980) conducted a study in female Long-Evans rats to determine whether exposure before and during gestation is more detrimental to reproductive outcome than exposure either before or during gestation alone. Four groups of 16–21 rats were formed in

which the rats were exposed by inhalation in whole-body chambers to 4,500 ppm dichloromethane (technical grade, >97% pure) 6 hours/day for 12–14 days before breeding and/or on GDs 1–17 or were exposed to filtered air. Maternal BWs were measured every 4 days. Dams were euthanized on GD 21, and livers and uteri were removed. Livers were weighed, and uterine horns were examined for fetal position and number of live, dead, or absorbed fetuses. Fetuses were observed for gross, soft-tissue, and skeletal abnormalities. Exposure during gestation (with or without pregestational exposure) significantly increased maternal absolute and relative liver weights by about 10–12 and 9–12%, respectively, and decreased fetal BW by about 9–10% relative to those exposed to filtered air during gestation. None of the groups showed significant alterations in the incidence of gross, external, skeletal, or soft-tissue anomalies. Using the same study design and exposure level, Bornschein et al. (1980) observed behavioral activities at various ages. Assessed activities included head movement/pivoting when placed in a novel environment (4 days of age), limited crawling (10 days), movement in a photocell cage (15 days), use of running wheel (45–108 days), and shock avoidance (4 months). Exposure during gestation (with or without pregestational exposure) caused altered rates of behavioral habituation to novel environments in the pups tested as early as 10 days of age; these altered rates were still present at 150 days of age. Growth, food and water consumption, wheel running activity, and avoidance learning were not significantly affected by exposure to dichloromethane. The results indicate that 4,500 ppm was a LOAEL for maternal effects (10% increased absolute and relative liver weight) and for effects on the fetuses (10% decreased fetal BW and altered behavioral habituation to novel environments).

In a study of early-life (including gestational) exposures, Maltoni et al. (1988) exposed 54 pregnant Sprague-Dawley rats to 100 ppm dichloromethane via inhalation 4 hours/day, 5 days/week for 7 weeks, followed by 7 hours/day, 5 days/week for 97 weeks. Exposure apparently started on GD 12. Groups of 60 male and 69 female newborns continued to be exposed after birth to 60 ppm dichloromethane 4 hours/day, 5 days/week for 7 weeks, followed by exposure 7 hours/day, 5 days/week for 97 weeks. Additional groups of 60 male and 70 female newborn were exposed after birth to 60 ppm dichloromethane 4 hours/day, 5 days/week for 7 weeks and then for 7 hours/day, 5 days/week for 8 weeks. BWs were measured every 2 weeks during exposure and every 8 weeks thereafter. At the end of exposure, animals were sacrificed and histologic examinations were performed on 20 tissue types.

Early life exposures of Sprague-Dawley rats to dichloromethane (Maltoni et al., 1988) did not affect mortality or BW in any group. Also, there was no significant effect of exposure to dichloromethane on the percentage of animals with benign and malignant tumors and malignant tumors, the number of malignant tumors per 100 animals, or the percentage of animals with benign mammary tumors, malignant mammary tumors, leukemias, pheochromocytomas, and pheochromoblastomas. The results provide no evidence that gestational exposure to 100 ppm dichloromethane during early life stages of development increases the susceptibility of Sprague-

Dawley rats to the potential carcinogenicity of dichloromethane, but further conclusions from these results are precluded because the study included only one exposure level that was below the maximum tolerated dose for adult Sprague-Dawley rats. Experiments comparing cancer responses from early-life exposures with adult exposures are not available for F344 rats or B6C3F₁ mice, the strains of animals in which carcinogenic responses to dichloromethane have been observed.

In summary, the potential for gestational exposure to CO, resulting from maternal dichloromethane exposure via oral and inhalation routes, raises concerns regarding neurodevelopmental effects. In addition, dichloromethane transfer across the placenta has also been seen in inhalation exposure studies in rats (Withey and Karpinski, 1985; Anders and Sunram, 1982). Although few developmental effects were observed at high exposures of dichloromethane (Bornschein et al., 1980; Schwetz et al., 1975), there are no studies that have thoroughly evaluated neurobehavioral and neurochemical changes resulting from gestational dichloromethane exposure. The available data identify changes of behavior habituation at 4,500 ppm (Bornschein et al., 1980) and increases in COHb at 1,250 ppm (Schwetz et al., 1975). The behavioral changes observed at 4,500 ppm indicate developmental neurotoxic effects. No other neurological endpoints have been evaluated in the available developmental studies of dichloromethane, but increases in blood COHb strongly suggest that dichloromethane is being metabolized to CO. Gestational exposure to CO can result in significant neurological effects in offspring, including neurobehavioral deficits (De Salvia et al., 1995), memory effects (Giustino et al., 1999), and neurochemical changes (Cagiano et al., 1998; Fechter, 1987). As a result, it is unknown if developmental neurotoxicity could occur at lower exposures to dichloromethane.

4.4. OTHER DURATION- OR ENDPOINT-SPECIFIC STUDIES

4.4.1. Short-term (2-Week) Studies of General and Hepatic Effects in Animals

Two short-term (2-week) studies examined hepatic and renal effects of dichloromethane exposure in F344 rats (Berman et al., 1995) and CD-1 mice (Condie et al., 1983). Berman et al. (1995) administered dichloromethane by gavage in corn oil for up to 14 days to groups of eight female F344 rats at dose levels of 0, 34, 101, 337, or 1,012 mg/kg-day. Starting at day 4, deaths occurred in the 1,012 mg/kg-day exposure group, with seven of eight rats dying before the end of the 14-day exposure period. In the dose groups that did not experience this high mortality, incidences of increased necrotic hepatocytes were 0/8, 0/8, 0/8, and 3/8 for the 0, 34, 101, and 337 mg/kg-day groups, respectively. The increase in liver lesions was not accompanied by increases in serum activities of ALT or AST. Kidneys, spleen, and thymus were also histopathologically examined in this study, but none showed exposure-related lesions. The results indicate that 101 mg/kg-day was a NOAEL and 337 mg/kg-day was a LOAEL for increased incidence of degenerative lesions in female rats exposed for 14 days. In a companion study with groups of eight female F344 rats that were given single doses of 0, 101, 337, 1,012, or

1,889 mg/kg-day, incidences of rats with increased necrotic hepatocytes were 1/8, 0/8, 8/8, 7/8, and 8/8, respectively (Berman et al., 1995).

Condie et al. (1983) detected exposure-related liver lesions in a 14-day gavage study in which dichloromethane in corn oil was administered to male CD-1 mice at dose levels of 0, 133, 333, or 665 mg/kg-day. Incidences of mice with minimal or slight cytoplasmic vacuolation were 1/16, 0/5, 3/5, and 4/5 for the control through high-dose groups, respectively. The kidneys were also examined histopathologically in this study but showed no exposure-related lesions. No other tissues were prepared for histologic examination. Blood urea nitrogen, serum creatinine, and serum ALT activities were not significantly altered by exposure. All dose levels significantly reduced to the same extent the active transport of *p*-aminohippurate into renal cortical slices in vitro, a measure of proximal tubule function. The results most clearly identify 133 mg/kg-day as a NOAEL and 333 mg/kg-day as a LOAEL for increased incidence of hepatocyte vacuolation in male mice.

4.4.2. Immunotoxicity Studies in Animals

Aranyi et al. (1986) studied the effects of acute inhalation exposures to 50 or 100 ppm dichloromethane on two measures of immune response (susceptibility to respiratory infection and mortality due to *Streptococcus zooepidemicus* exposure and ability of pulmonary macrophages to clear infection with *Klebsiella pneumoniae*). Female CD1 mice that were 5–7 weeks of age at the start of the exposure portion of the experiment were used for both assays. Up to five replicate groups of about 30 mice were challenged with viable *S. zooepidemicus* during simultaneous exposure to dichloromethane or to filtered air. Deaths were recorded over a 14-day observation period. Clearance of ³⁵S-labeled *K. pneumoniae* by pulmonary macrophages was determined by measuring the ratio of the viable bacterial counts to the radioactive counts in each animal's lungs 3 hours after infection; 18 animals were used per dose group. A single 3-hour exposure to 100 ppm dichloromethane significantly increased the susceptibility to respiratory infection and greater mortality following exposure to *S. zooepidemicus* ($p \leq 0.01$). Twenty-six deaths occurred in 140 (18.6%) mice challenged during a 3-hour exposure to 100 ppm dichloromethane; in contrast, nine deaths occurred in 140 mice (6.4%) exposed to filtered air. The 3-hour exposure to 100 ppm dichloromethane was associated with a statistically significant ($p \leq 0.001$) 12% decrease in pulmonary bactericidal activity (91.6 and 79.6% of bacteria killed in controls and 100 ppm group, respectively). No difference was seen in either mortality rate or bactericidal activity in experiments using a single 3-hour exposure to 50 ppm or 3-hour exposures to 40 ppm dichloromethane repeated daily for 5 days compared with control animals exposed to filtered air. These results suggest that 3-hour exposure to 50 ppm dichloromethane was a NOAEL and 100 ppm was a LOAEL for decreased immunological competence (immunosuppression) in CD-1 mice.

Aranyi et al. (1986) also conducted a similar set of experiments with 13 other chemicals (acetaldehyde, acrolein, propylene oxide, chloroform, methyl chloroform, carbon tetrachloride, allyl chloride, benzene, phenol, monochlorobenzene, benzyl chloride, perchloroethylene, and ethylene trichloride). Perchloroethylene and ethylene trichloride were the only chemicals in this group for which an increased mortality risk from streptococcal pneumonia was seen (mortality risk 15.0 and 31.4% in controls and 50 ppm exposure groups, respectively, for perchloroethylene and 13.4 and 58.1% in controls and 50 ppm exposure groups, respectively, for ethylene trichloride). Decreased bactericidal activity was also seen with acetaldehyde, acrolein, methyl chloroform, allyl chloride, benzene, benzyl chloride, perchloroethylene, and ethylene trichloride at one or more exposures. Results from several chemicals suggest that 5 days of exposure results in greater decrease in bactericidal activity (i.e., acetaldehyde, acrolein, and benzene), and others (e.g., perchloroethylene) suggest that 5 days of exposure does not result in greater suppression than a single exposure period.

There was considerable variation in both measures of immune response among the controls in the experiments (Aranyi et al., 1986). Among the controls in the experiments with the 13 chemicals other than dichloromethane, mortality in the streptococcal infectivity model ranged from 5.7 to 22.1%, with a mean of 12.7%.⁵ Bactericidal activity in the klebsiella model among controls ranged from 67.9 to 94.7%, with a mean of 81.8%. The number of bacteria deposited in the lung in an inhalation bacterial infectivity model can show considerable variation, (i.e., between 750 to 1,500 viable streptococcus or klebsiella organisms, [Ehrlich, 1980]). Therefore, concurrent controls are particularly important due to the variation in preparation and aerosol administration of the bacteria in these assays.

Warbrick et al. (2003) evaluated immunocompetence in male and female Sprague-Dawley rats by measuring the immunoglobulin M (IgM) antibody responses following immunization with sheep red blood cells in addition to hematological parameters and histopathology of the spleen, thymus, lungs, and liver. Groups of rats (8/sex/dose level) were exposed to 0 or 5,000 ppm dichloromethane 6 hours/day, 5 days/week for 28 days. Rats injected with cyclophosphamide served as positive controls. Five days before sacrifice (day 23 of exposure) all rats were injected with sheep red blood cells. IgM levels in response to the sheep red blood cells were comparable between dichloromethane-exposed and air-exposed rats, indicating that dichloromethane did not produce immunosuppression in the animals under these exposure conditions. Cyclophosphamide-treated animals had significantly lower levels of IgM in the blood serum, indicating immunosuppression. Rats exposed to dichloromethane showed reduced response to sound, piloerection, and hunched posture during exposures. Neither BW gain nor the hematological parameters monitored were significantly affected by exposure to dichloromethane. Relative and absolute liver weights were significantly increased in females but

⁵EPA did not include the duplicate assay of perchloroethylene in calculating this summary statistic. If this additional assay is included, the mortality risk ranges from 5.7 to 45.7%, with a mean of 15.0%.

not in males. Relative spleen weight was reduced in females, and no significant changes were seen in the weight of the thymus and lungs. Histopathology of the tissues examined was unremarkable. Exposure to 5,000 ppm dichloromethane did not affect antibody production to the challenge with sheep red blood cells.

In the 2-year drinking water study (Serota et al., 1986a, b) and 2-year inhalation study (Nitschke et al., 1988a), histopathologic analyses were conducted on the lymph nodes, thymus, and spleen among several other organs, and no significant changes were noted.

In summary, one study (Aranyi et al., 1986) demonstrated evidence of immunosuppression, including increased risk of streptococcal-pneumonia-related mortality and decreased clearance of klebsiella bacteria following a single dichloromethane exposure at 100 ppm for 3 hours in CD-1 mice. The streptococcal and klebsiella bacterial inhalation assays are models of respiratory infection that test for local immune effects associated with inhalation exposure rather than systemic immunosuppression. The NOAEL identified in this study was 50 ppm. In contrast, in a functional immune assay of systemic immunosuppression conducted in rats, Warbrick et al. (2003) did not observe changes in the antibody response to sheep red blood cells in a 28-day inhalation exposure to 5,000 ppm dichloromethane. Histopathologic analyses of immune system organs in chronic exposure studies for B6C3F₁ mice and F344 rats (Nitschke et al., 1988a; Serota et al., 1986a, b) revealed no changes from controls. However, no assays of functional immunity were included in these chronic studies. These two studies for dichloromethane do not suggest systemic immunosuppression, but the Aranyi et al. (1986) study provides evidence of route-specific local immunosuppression from acute inhalation exposure in CD1 mice. Due to the acute exposure duration used in Aranyi et al. (1986), the immune effects of short-term or chronic exposure to dichloromethane are unclear.

4.4.3. Neurotoxicology Studies in Animals

Neurological evaluations in animals during and after exposure to dichloromethane have resulted in CNS depressant effects similar to other chlorinated solvents (e.g., trichloroethylene, perchloroethylene) and ethanol. Overall, there are decreased motor activity, impaired memory, and changes in responses to sensory stimuli. Neurobehavioral, neurophysiological, and neurochemical/neuropathological studies have been used to characterize the effects of dichloromethane on the CNS. A brief overview of these types of studies is provided below, followed by a detailed description of individual studies.

Neurobehavioral studies with dichloromethane used protocols to measure changes in spontaneous motor activity, a functional observational battery (FOB) test (to evaluate gross neurobehavioral deficits), and a task developed to assess learning and memory. The FOB protocol includes various autonomic parameters, neuromuscular parameters, sensorimotor parameters, excitability measures, and activity. Learning and memory changes with

dichloromethane were studied by using a passive avoidance task. The oral and inhalation studies that examined neurobehavioral endpoints are summarized in Table 4-26.

Table 4-26. Studies of neurobehavioral changes from dichloromethane, by route of exposure and type of effect

Species	Exposure(s)	Duration	Neurobehavioral effect	Reference
Gavage exposure				
<i>FOB</i>				
F344 rat, female	101, 337, 1,012, 1,889 mg/kg, gavage	Acute—evaluated 4 and 24 hrs after dosing	FOB neuromuscular and sensorimotor parameters significantly different from controls at 1,012 and 1,889 mg/kg (337 mg/kg = NOAEL)	Moser et al. (1995)
F344 rat, female	34, 101, 337, 1,012 mg/kg-d, gavage	14 d—evaluated on d 4, 9, and 15	All FOB parameters (except activity) significantly affected from d 4 at doses of 337 and 1,012 mg/kg-d	Moser et al. (1995)
Inhalation exposure				
<i>Spontaneous activity</i>				
NMRI mouse, male	400–2,500 ppm	1 hr	Initial increase in activity followed by a pronounced decrease at exposures ≥ 600 ppm	Kjellstrand et al. (1985)
Rat, male	5,000 ppm	1 hr, every other d for 10 d	Decreased spontaneous locomotor activity	Heppel and Neal (1944)
Wistar rat, male	500 ppm	6 hrs/d, 6 d	Increased preening frequency	Savolainen et al. (1977)
ICR mouse, female	5,000 ppm	Continuous, 7 d	Increased spontaneous activity in first few hrs and then decreased activity	Weinstein et al. (1972)
Sprague-Dawley rat, male	1,000, 5,000 ppm	Continuous, 14 wks	No neurobehavioral changes	Haun et al. (1971)
ICR mouse, female	1,000, 5,000 ppm	Continuous, 14 wks	Incoordination, lethargy	Haun et al. (1971)
Beagle dog, female	1,000, 5,000 ppm	Continuous, 14 wks	Incoordination, lethargy	Haun et al. (1971)
Rhesus monkey, female	1,000, 5,000 ppm	Continuous, 14 wks	Incoordination, lethargy	Haun et al. (1971)
ICR mouse, female	25, 100 ppm	Continuous, 14 wks	Increased spontaneous activity at 25 ppm	Thomas et al. (1972)
<i>FOB</i>				
F344 rat, male and female	50, 200, 2,000 ppm	6 hrs/d, 5 d/wk, 13 wks + 65 hrs exposure free	No effects observed on FOB, grip strength	Mattsson et al. (1990)
<i>Learning and memory</i>				
Swiss-Webster mouse, male	47,000 ppm	Approximately 20 sec + 1 hr exposure free before training; retested at d 1, 2, and 4	Significant decrease in learning and recall ability	Alexeef and Kilgore (1983)

Neurophysiological studies with dichloromethane exposure consisted of measuring evoked responses in response to sensory stimuli. In these studies, animals were implanted with electrodes over the brain region that responds to the particular stimuli. For example, an electrode would be implanted over the visual cortex in an animal presented with a visual stimulus. Once the stimulus is presented to the animal, an evoked response is elicited from the brain region and transmitted to the implanted electrode. During administration of a chemical, if there is a significant change in the magnitude, shape, and latency (among other measures) in the evoked response, then the chemical is considered to produce neurological effects. A summary of studies examining dichloromethane exposure and neurophysiological changes is shown in Table 4-27.

Table 4-27. Studies of neurophysiological changes as measured by evoked potentials resulting from dichloromethane, by route of exposure

Species	Exposure(s)	Duration	SEPs measured	Effect	Reference
Intraperitoneal					
Long-Evans rat, male	57.5, 115, 230, 460 mg/kg	Acute; tested at 15 min, 1 hr, and 5 hrs after dosing	FEP	Significant changes in FEPs were noted in animals dosed \geq 115 mg/kg; FEP changes time and dose dependent	Herr and Boyes (1997)
Inhalation exposure					
F344 rat, male	5,000, 10,000, 15,000 ppm	Acute, 1 hr; tested during exposure	Electroencephalogram, BAER, CAEP, FEP, SEP	Significant changes in SEP, FEP, BAER, and CAEP responses at all exposures; slight recovery noted at 1 hr after exposure	Rebert et al. (1989)
F344 rat, male and female	50, 200, 2,000 ppm	Subchronic, 6 hr/d, 5 d/wk, 13 wks; tested 65 hrs after last exposure	FEP, CAEP, BAER, SEP	No significant changes noted in any evoked potential measurements	Mattsson et al. (1990)

BAER = brainstem-auditory-evoked response; CAEP = cortical-auditory-evoked potential; FEP = flash-evoked potential; SEP = somatosensory-evoked potential

In neurochemical/neuropathological studies with dichloromethane, animals were first exposed to dichloromethane (orally or via inhalation or injection), and then the brains were removed. Changes in excitatory neurotransmitters, such as glutamate and acetylcholine and the inhibitory neurotransmitter, GABA, were measured. Additionally, dopamine and serotonin levels, which are associated with addiction and mood, were also measured. Other parameters that were measured included DNA/protein content and regional brain changes in the cerebellum and hippocampus. Measurement of neurochemical changes provides mechanistic information, and neurobehavioral and neurophysiological effects can be correlated to these results. Table 4-28 summarizes studies of neurochemical changes and dichloromethane.

Table 4-28. Studies of neurochemical changes from dichloromethane, by route of exposure

Species and sex	Exposure	Duration	Regions	Effect ^a	Reference
Oral exposure					
Sprague-Dawley rat, male	534 mg/kg	Acute, single dose; evaluated 2 hrs after dosed	Hippocampus, medulla, midbrain, hypothalamus	↑ acetylcholine in hippocampus ↑ dopamine and serotonin in medulla ↓ norepinephrine in midbrain ↓ norepinephrine and serotonin in hypothalamus	Kanada et al. (1994)
Inhalation exposure					
Wistar rat, male	1,000 ppm TWA (basal exposure of 100 ppm + 2,800 ppm, 1 hr peak exposures at hrs 1 and 4)	6 hrs/d, 5 d/wk, 2 wks	Cerebrum, cerebellum	↑ NADPH diaphorase, succinate dehydrogenase in cerebrum ↑ cerebral RNA ↓ succinate dehydrogenase in cerebellum	Savolainen et al. (1981)
Wistar rat, male	1,000 ppm TWA	6 hrs/d, 5 d/wk, 2 wks + 7 d exposure free	Cerebrum, cerebellum	↓ succinate dehydrogenase in both regions	Savolainen et al. (1981)
Wistar rat, male	1,000 ppm	6 hrs/d, 5 d/wk, 2 wks	Cerebrum, cerebellum	↑ acid proteinase ↓ succinate dehydrogenase in cerebellum	Savolainen et al. (1981)
Wistar rat, male	1,000 ppm	6 hrs/d, 5 d/wk, 2 wks + 7 d exposure free	Cerebrum	↓ cerebral RNA	Savolainen et al. (1981)
Sprague-Dawley rat, male	70, 300, 1,000 ppm	6 hrs/d, 3 d	Caudate nucleus—medial	↑ catecholamine levels (70 ppm) ↓ catecholamine levels (300 and 1,000 ppm) No effect on luteinizing hormone release	Fuxe et al. (1984)
Mongolian gerbil, male and female	210, 350 ppm	Continuous (24 hrs/d), 3 mo + 4 mo exposure free	Hippocampus, cerebellum, cerebral cortex	↓ DNA concentration per wet weight in hippocampus (210, 350 ppm) and cerebellar hemispheres (350 ppm) ↑ astroglial proteins in frontal and sensory motor cerebral cortex	Rosengren et al. (1986)
Mongolian gerbil, male and female	210 ppm	Continuous (24 hrs/d), 3 mo	Frontal cortex, cerebellum	↓ glutamate, GABA, phosphoethanolamine in frontal cortex ↑ glutamate, GABA in posterior cerebellar vermis	Briving et al. (1986)

Table 4-28. Studies of neurochemical changes from dichloromethane, by route of exposure

Species and sex	Exposure	Duration	Regions	Effect^a	Reference
Mongolian gerbil, male and female	210 ppm	Continuous (24 hrs/d), 3 mo + 4 mo exposure free	Hippocampus, olfactory bulbs, cerebral cortex	↓ DNA concentration per wet weight in hippocampus only	Karlsson et al. (1987)

^aAll effects shown in this table were statistically significant.

↑ = increase; ↓ = decrease; NADPH = nicotinamide adenine dinucleotide phosphate

4.4.3.1. Neurotoxicology Studies—Oral Exposures

Three studies evaluated the neurotoxic potential of dichloromethane by either administering the solvent orally or by injection; two of these studies (Herr and Boyes, 1997; Kanada et al., 1994) only evaluated acute effects (2–5 hours) from single-dose exposures. Observed neurological effects included decreased spontaneous activity (Moser et al., 1995), changes in flash-evoked potential (FEP) measurements (Herr and Boyes, 1997), and changes in catecholamine levels in the brain (Kanada et al., 1994).

Moser et al. (1995) conducted neurobehavioral evaluations in female F344 rats following an acute or 14-day oral administration of dichloromethane. A FOB protocol was utilized to determine changes in autonomic parameters (lacrimation, salivation, pupil response, urination, defecation), neuromuscular parameters (gait, righting reflex, forelimb and hind-limb grip strength, landing foot splay), sensorimotor parameters (tail pinch, click response, touch response), excitability measures (handling reactivity, arousal, clonic, and/or tonic movements), and activity (rearing, motor activity). A baseline FOB was performed on all rats prior to initial dichloromethane administration. After dichloromethane administration, a FOB was conducted at selected time points followed by a motor activity test in a maze. In the acute study, rats were dosed with 0, 101, 337, 1,012, or 1,889 mg/kg dichloromethane. At 4 and 24 hours after the administered dose, rats were tested for the neurological parameters. Significant changes in the neuromuscular and sensorimotor parameters were observed and occurred mostly in rats administered with the highest dose. These significant changes were only observed at the 4-hour time point and not when measured at 24 hours. The NOAEL identified by the authors for this study was 337 mg/kg based on no observable changes in the FOB. In the 14-day study, rats were administered 0, 34, 101, 337, or 1,012 mg/kg-day. FOB testing was conducted on days 4 and 9 (before the daily dose) and approximately 24 hours after the last (14th) dose. With the exception of the activity measurements, all other neurobehavioral parameters (neuromuscular, sensorimotor, autonomic, excitability) were significantly affected from the 4th day through the entire 14-day exposure cycle. The NOAEL identified for the 14-day study was 101 mg/kg-day based on FOB changes associated with the dichloromethane exposure.

A single dose acute neurophysiology study by Herr and Boyes (1997) evaluated the effect of dichloromethane on FEPs in adult male Long-Evans rats. Rats were implanted with epidural electrodes over the visual cortex area. After placement in an enclosed rectangular mirror chamber, FEPs were stimulated with a 10 µsec flash. Baseline FEPs were collected and rats were injected intraperitoneally with 0 (corn oil, n = 16), 57.5 (n = 15), 115 (n = 15), 230 (n = 14), or 460 (n = 15) mg/kg dichloromethane. Animals were retested at 15 minutes, 1 hour, and 5 hours after injection. Amplitude decreases in the early FEP components were observed. The FEP amplitude changes were time- and dose-dependent with maximal effects at 15 minutes after dichloromethane dosage. All of the waveform amplitudes returned to control levels when

measured at the 1-hour time point for all doses tested. Response latencies were still different from controls when measured 5 hours after dosing, but the effect was less pronounced than at the 15-minute and 1-hour time points. In this study, 57.5 mg/kg did not produce any significant changes in the FEP measures as compared to controls and was considered this study's NOAEL. The LOAEL was 115 mg/kg based on changes in the FEP amplitudes.

Kanada et al. (1994) examined the effect of dichloromethane on acetylcholine and catecholamines (dopamine, norepinephrine, serotonin) and their metabolites in the midbrain, hypothalamus, hippocampus, and medulla from male Sprague-Dawley rats (4–5/group) in a neurochemical/neuropathology study. The rats were sacrificed 2 hours after a single gavage dose of 0 or 534 mg/kg of undiluted dichloromethane. Administration of dichloromethane significantly increased the concentration of acetylcholine in the hippocampus by approximately 10% and increased dopamine and serotonin levels in the medulla by approximately 75%. Dichloromethane decreased norepinephrine levels in the midbrain and hypothalamus by 12–15%, and serotonin levels were decreased in the hypothalamus by approximately 30%. There was a trend toward decreased dopamine in the hypothalamus, but the variability between the animals was so high that the effect was not significant. (These values for the percent changes were estimated by EPA from the figures presented in the paper.) The authors speculated that increased acetylcholine release associated with exposure to dichloromethane and other solvents may originate from the nerve terminals.

4.4.3.2. Neurotoxicology Studies—Inhalational Exposure

The database pertaining to neurotoxic effects from inhalation exposure to dichloromethane is considerably larger than the oral exposure database. Acute (<1 day) and short-term (1–14 days) exposures resulted in an initial increase in spontaneous activity followed by a decrease for exposures between 500 and 2,500 ppm (Kjellstrand et al., 1985; Savolainen et al., 1977). Higher (5,000 ppm) acute and short-term exposures resulted in decreased spontaneous activity and lethargy (Weinstein et al., 1972; Heppel and Neal, 1944). Longer-term exposures (up to 14 weeks) produced decreased motor activity and lethargy in several animals at 1,000 and 5,000 ppm (Haun et al., 1971), and exposures at 25 ppm for 14 weeks produced significant increases in activity in mice, starting at week 9. CNS depression was evidenced by decreased responses in the auditory, visual, and somatosensory regions of the brain in a study of sensory-evoked potential effects in 12 adult male F344 rats exposed to 0, 5,000, 10,000, and 15,000 ppm for 1 hour periods (Rebert et al., 1989). Altered learning and memory abilities were demonstrated in young (3-, 5-, and 8-week-old) male Swiss-Webster mice exposed to 168 mg/L (~47,000 ppm) dichloromethane for approximately 20 seconds (until there was a loss of the righting reflex) (Alexeef and Kilgore, 1983).

4.4.3.2.1. *Inhalational exposure—neurobehavioral studies.*

Spontaneous motor activity—acute and short-term studies. Heppel and Neal (1944) evaluated the neurological effects of 5,000 ppm dichloromethane in five male rats by measuring changes in spontaneous activity during and after exposure. The five rats were not randomly selected, since the investigators chose to pick out the most active animals in the litter. During the 1-hour testing runs, rats were placed in a rotating drum. Spontaneous activity was reported as the number of drum revolutions/hour. Twenty control test runs (1 run/day) were conducted prior to dichloromethane exposure runs. After the preexposure period, rats were exposed to 5,000 ppm dichloromethane every other day for 1 hour, and activity was measured in the same manner as in the control runs. Once dichloromethane exposure was stopped, the animals were allowed to recover for 30 minutes and a second 1-hour test run was performed to evaluate spontaneous activity during recovery. On nonexposure days, spontaneous activity was also measured in 1-hour intervals to compare to the preexposure period. A total of five dichloromethane exposures, five postexposure, and five nonexposure trials were conducted over 10 days. Spontaneous activity significantly declined ($p < 0.01$, Fisher's t-test) during exposure to 5,000 ppm dichloromethane in comparison to nonexposure days. The average number of revolutions for all five rats over the test runs was 576 on nonexposure days and 59 revolutions during dichloromethane exposure.

Weinstein et al. (1972) continuously exposed female ICR mice to 5,000 ppm dichloromethane for up to 7 days. Clinical behavioral observations of the mice were made during dichloromethane exposure. Within the first few hours of exposure, spontaneous activity increased in comparison to control animals. After 24 hours of continuous exposure, there was a considerable decrease in spontaneous activity as noted by observation only. The mice also appeared to be very lethargic and had a hunched posture and a rough hair coat, which are all signs of CNS depressive effects in rodents. These effects became progressively worse until after 96 hours of exposure, where many mice resumed normal activity. After the 7-day exposure, mice were nearly as active as the control animals but had a rougher coat and were judged to be emaciated and dehydrated.

Male Wistar rats exposed to 500 ppm dichloromethane 6 hours/day for 6 days exhibited an increase in preening frequency and time 1 hour after the last exposure relative to controls (Savolainen et al., 1977). However, there were no significant changes in other types of spontaneous activity.

In the study by Kjellstrand et al. (1985), male NMRI mice were exposed to dichloromethane concentrations ranging from 400 to 2,500 ppm. At concentrations of ≥ 600 ppm, exposures for 1 hour produced a biphasic pattern of activity characterized by an initial increase in activity (as high as 200% of preexposure motor activity at 2,200 ppm, as estimated from Figure 6 in Kjellstrand et al. [1985]) during exposure followed by a decrease that reached the lowest point 1–2 hours after the end of exposure (as low as 40% motor activity at

2,200 ppm in comparison to preexposure, as estimated from Figure 6 in Kjellstrand et al. [1985]). Motor activity returned to normal levels after the decreased activity observed 1–2 hours after exposure was stopped, indicating that the effect was reversible in this study design.

Spontaneous motor activity—subchronic (14 week) studies. Haun et al. (1971) reported results from studies in which female beagle dogs, female rhesus monkeys, male Sprague-Dawley rats, and female ICR mice were continuously exposed to 0, 1,000, or 5,000 ppm dichloromethane for up to 14 weeks in whole-body exposure chambers. Gross and histopathologic examinations were made on animals that died or were sacrificed during or at termination of the study. At 5,000 ppm, obvious nervous system effects (e.g., incoordination, lethargy) were most apparent in dogs and also observed in monkeys and mice. Rats did not demonstrate any of these sedative effects. At 1,000 ppm, these effects were observed to a lesser extent in monkeys and mice, but dogs still displayed prominent CNS depressive behavior. Histopathologic analysis revealed edema of the brain in three dogs that died during exposure to 5,000 ppm dichloromethane. No other gross brain-related changes were reported. The results indicate that continuous exposure to 1,000 ppm was an adverse effect level for mortality and effects on the nervous system and liver in dogs (exposed for up to 4 weeks) and for BW changes in rats (exposed for 14 weeks). The 5,000 ppm level induced mortality in beagle dogs, ICR mice, and rhesus monkeys (but not Sprague-Dawley rats); obvious nervous system effects in dogs, mice, monkeys, and rats; and gross liver changes in dogs, mice, monkeys, and rats.

In the study by Thomas et al. (1972), female ICR mice were exposed continuously to 0, 25, or 100 ppm dichloromethane for 14 weeks. Spontaneous activity of mice was evaluated by using closed circuit television for monitoring. Mice were evaluated in daily 2-hour testing sessions. The 25 and 100 ppm exposure groups were tested for 2 weeks prior to the onset of dichloromethane exposure. Starting at week 9, mice exposed to 25 ppm dichloromethane exhibited increases in spontaneous activity, but no quantitative measurements or statistical analysis were reported. The authors stated that no significant effect was observed in the group exposed to 100 ppm.

FOB—subchronic (13 week) study. Only one study, a 13-week inhalation study in F344 rats (Mattsson et al., 1990), has conducted an FOB testing paradigm following a subchronic exposure to dichloromethane. Groups of rats (12/sex/exposure level) were exposed to 0, 50, 200, or 2,000 ppm dichloromethane 6 hours/day, 5 days/week for 13 weeks. An additional group of rats was exposed to 135 ppm CO to induce approximately 10% COHb, approximately the level produced by saturation of oxidative metabolism of dichloromethane. After the 13 weeks of exposure (beginning 65 hours after the last exposure), rats were subject to an FOB to evaluate any neurobehavioral changes from the dichloromethane exposure. Autonomic parameters were first characterized. Then the rat was placed in a clear plastic box to evaluate locomotor activity

and then responsiveness to touch, sharp noise, and tail pinch. Hind-limb grip strength was also measured by using a strain gauge. All animals were examined clinically at weekly intervals and were tested at the end of the exposure period by FOB, grip strength, BW, temperature, and sensory-evoked potentials. No exposure-related effects were observed on the FOB, grip strength, or sensory-evoked potentials. No histopathologic changes were noted in brains, spinal cords, or peripheral nerves from the high-dose dichloromethane group compared with control animals. In the absence of changes, lower concentrations were not examined.

Learning and memory—acute study. In a study by Alexeef and Kilgore (1983), a learning and memory evaluation was conducted following acute exposure to dichloromethane. Mice were exposed to 168 mg/L (~47,000 ppm) dichloromethane and were tested for learning ability by using a passive-avoidance conditioning task. Male Swiss-Webster mice (3, 5, and 8 weeks old) were used in this study. In the passive avoidance task, mice were placed on a metal platform that extended into a hole. If the mouse went into the hole (a darkened area, which would be the preferred area for the mouse), it received a foot shock. Prior to the training session, mice were exposed to either air or ~47,000 ppm dichloromethane. Animals were exposed to dichloromethane until there was a loss of the righting reflex, which would take about 20 seconds on average, and then placed back in their home cage. One hour after exposure, animals were trained to learn the passive avoidance task. A mouse was considered to have learned the task once it remained on the platform for at least 30 seconds without entering the hole. Mice were then tested for recollection of the task at either 1, 2, or 4 days after the initial training session. In the learning phase of the task, 74% of the control mice retained the task in comparison to 59% of the dichloromethane-exposed group, indicating the significant effect of dichloromethane on learning. There was also an age-related effect since exposed 3-week-old mice were less likely to recall the task than 5- or 8-week-old mice. There was no difference in task recall between the 5- and 8-week-old mice. Dichloromethane at the exposure used in the study was demonstrated to be nonanalgesic, since pain-response times were comparable to those in air-exposed animals in the hot-plate pain test, and therefore, the results of the passive avoidance test were not confounded by potential analgesic effects. As a result, it is demonstrated that exposure to an acute and high concentration of dichloromethane alters learning ability in mice.

4.4.3.2.2. Inhalational exposure—neurophysiological studies. The effect of dichloromethane on sensory stimuli was evaluated by measuring sensory-evoked responses during an acute exposure (Rebert et al., 1989) and following a subchronic (13-week) exposure (Mattsson et al., 1990). Rebert et al. (1989) evaluated the effects of dichloromethane on sensory-evoked potentials (auditory, visual, and somatosensory) in F344 rats exposed to 0, 5,000, 10,000, and 15,000 ppm dichloromethane for 1 hour in a head-only exposure chamber. Twelve adult male rats were implanted with chronic epidural electrodes placed over the visual and somatosensory

cortices. Each rat served as its own control, with a 1-week recovery period between testing sessions. During each testing session, spontaneous electroencephalograms were recorded. Additionally, brainstem-auditory-evoked responses (BAERs) (tone stimulus), cortical-auditory-evoked potentials (CAEPs) (click stimulus), FEPs (flash stimulus), and somatosensory-evoked potentials (SEPs) (tail current stimulus) were measured in response to the stimuli. Dichloromethane decreased the SEP response to the tail current stimulus, and earlier components of the FEP response were attenuated and eventually eliminated with increasing exposures. The BAER response profile was also significantly altered. Dichloromethane completely abolished the CAEP at all concentrations tested. Slight recovery of this response was noted approximately 1 hour after exposure. The collective results strongly suggest a CNS depressive profile for dichloromethane and indicate that this chemical affects the auditory, visual, and somatosensory regions of the brain.

In a subchronic exposure study, male and female F344 rats were exposed to dichloromethane 6 hours/day, 5 days/week for 13 weeks (Mattsson et al., 1990). Twelve animals of each sex were selected for exposure to 0, 50, 200, or 2,000 ppm dichloromethane or 135 ppm CO. For electrophysiological measures, rats were surgically implanted with epidural electrodes 10 weeks after the onset of exposure. Electrodes were placed over the somatosensory, visual, and cerebellar region. Electrophysiological measures that were recorded included FEP measurements, cortical flick fusion responses, CAEPs, BAERs, and SEPS recorded from the sensory (SEP-S) and cerebellar (SEP-C) regions. None of these measures were significantly altered by any dichloromethane or CO treatment in this study. However, it should be noted that all of the electrophysiological measures were conducted at least 65 hours after the last dichloromethane exposure. As a result, it can be concluded that a subchronic exposure to dichloromethane did not result in persistent changes in any of the neurophysiological measures that were evaluated in this study. It is not known if any neurological compensation occurred, since SEP measurements were not taken during actual dichloromethane exposure in this subchronic study.

Based on these two studies, the significant changes noted in several SEP measures during dichloromethane exposure were not observed after a subchronic exposure where animals were tested at least 65 hours after the last exposure. As a result, it is difficult to ascertain if tolerance is developed to the dichloromethane-mediated changes in sensory potentials during an acute exposure or if these effects are still maintained during repeated exposure, since measurements were not taken during the subchronic exposure.

4.4.3.2.3. Inhalational exposure—neurochemistry and neuropathology studies. The studies evaluating specific neurochemical changes in relation to dichloromethane exposure include studies of effects of short-term (3-day to 2-week) exposures (Fuxe et al., 1984; Savolainen et al.,

1981) and subchronic (3-month) exposures (Karlsson et al., 1987; Briving et al., 1986; Rosengren et al., 1986).

Savolainen et al. (1981) examined three different exposure schemes in male Wistar rats. The rats were exposed to 500, 1,000, or 1,000 ppm TWA dichloromethane 6 hours/day, 5 days/week for 2 weeks. (Note: The abstract of this paper describes the exposures as 500, 1,000, and 100 ppm TWA, but, based on information in the body of the paper, the abstract appears to be incorrect.) The 1,000 ppm TWA exposure consisted of a basal 100 ppm exposure with two 2,800 ppm 1-hour peak concentrations (at 1 and 4 hours) resulting in a time-weighted exposure of 1,000 ppm. Brains were removed from rats at the end of study and analyzed. The 1,000 ppm TWA group displayed increases in cerebral RNA. Other changes noted for this group in the cerebrum included significant increases in nicotinamide adenine dinucleotide phosphate (NADPH) diaphorase and succinate dehydrogenase activity. In the 1,000 ppm constant exposure group, acid proteinase activity was below the levels observed in control animals in the first week but increased to levels above control animals in the second week. In the cerebellum, there were no changes in RNA concentration, and there was a decrease in succinate dehydrogenase activity in both the 1,000 and 1,000 ppm TWA groups. After a 7-day withdrawal, RNA levels in the cerebrum were significantly lower in the 1,000 ppm group. Succinate dehydrogenase levels remained lowered in the 1,000 ppm TWA group after the 7-day exposure-free period. No significant effects were seen at 500 ppm.

Fuxe et al. (1984) evaluated changes in brain catecholamine levels after a 3-day exposure to dichloromethane using male Sprague-Dawley rats. Rats were exposed to 70, 300, and 1,000 ppm dichloromethane 6 hours/day for 3 consecutive days. Additional groups of rats were exposed to the same levels of dichloromethane and given intraperitoneal injections of the tyrosine hydroxylase inhibitor, α -methyl-dl-*p*-tyrosine methyl ester (H44/68), 2 hours prior to sacrifice. Brains were removed, stained, and evaluated for catecholamine changes 16–18 hours after the last exposure. Catecholamine levels were measured in the hypothalamus, frontal cortex, and caudate nucleus among other brain regions. At all exposures, there was a significant decrease by approximately 10–15% of catecholamine concentrations in the posterior periventricular region of the hypothalamus. In the medial part of the caudate nucleus, which is involved in memory processes, catecholamine levels were significantly higher (12%) in the 70 ppm group but significantly lower in the 300 ppm (1%) and 1,000 ppm (8%) groups compared with controls. The impact of dichloromethane was also evaluated on the hypothalamic-pituitary gonadal axis. The hypothalamus regulates secretion of reproductive hormones, such as follicle-stimulating hormone and luteinizing hormone. The levels of the hormone release were not significantly changed with dichloromethane exposure. However, when rats were dosed concurrently with H44/68 and dichloromethane, statistically significant, inversely dose-related increases in luteinizing hormone levels were observed (330, 233, and 172% higher than controls in the 70, 300, and 1,000 ppm groups, respectively). The study

overall demonstrates significant changes in catecholamine levels in the hypothalamus and caudate nucleus. No significant changes in catecholamine levels in the frontal cortex were reported. Catecholamine level changes in the hypothalamus did not appear to significantly affect hormone release; however, decreased catecholamine levels in the caudate nucleus at higher exposures may lead to memory and learning impairment.

A series of studies were conducted in male and female Mongolian gerbils exposed continuously to 210 (Karlsson et al., 1987; Briving et al., 1986), 350, or 700 ppm (Rosengren et al., 1986) dichloromethane for 3 months, followed by a 4-month exposure-free period. High mortality rates occurred at 350 ppm (6/10 males and 3/10 females by 71 days) and 700 ppm (10/10 males and 9/10 females by 52 days). Rosengren et al. (1986) monitored two astroglial proteins, S-100 and GFA, as well as DNA concentrations in the brain. Decreased DNA concentrations were noted in the hippocampus at both the 210 and 350 ppm exposures. At 350 ppm, there was also decreased DNA concentration in the cerebellar hemispheres, indicating a decreased cell density in these regions, probably due to cell loss. Increased astroglial proteins were found in the frontal and sensory motor cerebral cortex, which directly correlated to the astrogliosis that was observed in those areas. Up-regulation of these astroglial proteins is a good indicator of neuronal injury (Rosengren et al., 1986).

Karlsson et al. (1987) measured DNA concentrations in different regions of the gerbil brain. After the solvent-free exposure period, brains were removed and the olfactory bulbs and cerebral cortices were dissected. Brain weights and weights of the dissected brain regions were the same between control and dichloromethane-exposed animals. The total protein concentration per wet weight was not significantly different between dichloromethane-exposed and control animals. However, DNA concentrations per wet weight were significantly decreased in the hippocampus after dichloromethane exposure. No other examined regions demonstrated significant changes in DNA concentrations after dichloromethane exposure. This selective DNA concentration decrease observed in the hippocampus is a sign of neurotoxicity and may possibly explain why some studies have noted memory and learning deficits with dichloromethane exposure. In a companion paper, in which only the 210 ppm level was tested, it was found that exposure to dichloromethane decreased the levels of glutamate, γ -aminobutyric acid, and phosphoethanolamine in the frontal cortex, while glutamine and γ -aminobutyric acid were increased in the posterior cerebellar vermis (Briving et al., 1986). Increased levels of glutamate in the posterior cerebellar vermis could reflect an activation of astrocytic glia, since glutamine synthetase is localized exclusively in astrocytes. The gerbils did not have a solvent-free exposure period as in the other two studies (Karlsson et al., 1987; Rosengren et al., 1986). The exposure regime in these studies did not affect BW or brain weight. Furthermore, the neurochemical changes observed in these studies were not attributed to formation of CO.

Neurological changes have been investigated by measuring changes in neurotransmitter levels and changes in neurotransmitter localization. Changes in catecholamine levels in the

caudate nucleus after an acute exposure (Fuxe et al., 1984) as well as decreased DNA content in the hippocampus after a subchronic dichloromethane exposure (Rosengren et al., 1986) suggest that memory functions are altered since both brain regions are associated with learning and memory. The results from Fuxe et al. (1984) directly correlated with the finding that learning and memory were impaired in mice after an acute (single) and very high exposure (47,000 ppm) to dichloromethane (Alexeef and Kilgore, 1983). Additionally, changes in the hippocampus also suggest memory effects after a long-term, continual exposure to dichloromethane, although no conclusive evidence has been presented to date. In another subchronic, continuous exposure to 350 ppm dichloromethane for 3 months, decreased DNA concentration was observed in the cerebellar hemispheres of Mongolian gerbils and is suggestive of cell loss (Rosengren et al., 1986). However, in a 2-week exposure study in male Wistar rats, RNA changes were not noted in the cerebellum, although enzyme activity was significantly decreased in this region (but was increased in the cerebrum) (Savolainen et al., 1981). These results suggest that the cerebellum is a target for dichloromethane. Noted neurobehavioral effects that may be linked to impaired cerebellar function include changes in motor activity and impaired neuromuscular function (Moser et al., 1995).

4.5. MECHANISTIC DATA AND OTHER STUDIES IN SUPPORT OF THE MODE OF ACTION

4.5.1. Genotoxicity Studies

4.5.1.1. *In Vitro* Genotoxicity Assays

Bacterial, yeast, and fungi mutagenicity assays. Numerous in vitro studies have demonstrated dichloromethane as being mutagenic in bacterial assays, yeast, and fungi, and several studies provide evidence that the genotoxic action of dichloromethane in bacterial systems is enhanced in the presence of GSH (e.g., DeMarini et al., 1997; Pegram et al., 1997; Oda et al., 1996; Thier et al., 1993; Dillon et al., 1992) (Table 4-29). Considering the results are primarily dependent on the presence of GSH, activation likely involves the GST-T1 metabolic pathway, which produces two proposed DNA-reactive metabolites, S-(chloromethyl)glutathione and formaldehyde.

Table 4-29. Results from in vitro genotoxicity assays of dichloromethane with bacteria, yeast, or fungi

Assay	Test system	Concentration(s)	Results		Reference
			Without metabolic activation -S9	With metabolic activation +S9	
Bacteria					
Reverse mutation	<i>Salmonella typhimurium</i> TA98 ^a , TA100 ^a	6-hr exposure to 0, 7,000, and 14,000 ppm	+	+	Jongen et al. (1978)
Reverse mutation	<i>S. typhimurium</i> TA98, TA100	Up to 3,600 µg/plate	+	++	Gocke et al. (1981)
Reverse mutation	<i>S. typhimurium</i> TA1535 ^b , TA1537 ^b , TA1538 ^b	Up to 3,600 µg/plate	-	-	Gocke et al. (1981)
Reverse mutation	<i>S. typhimurium</i> TA100	6-hr exposure to 0, 7,000, and 14,000 ppm	+	++	Jongen et al. (1982)
Reverse mutation	<i>S. typhimurium</i> TA100	Up to 84,000 ppm, 3-d exposure	+	+	Green (1983)
Reverse mutation	<i>S. typhimurium</i> TA100, TA1535, TA1950 ^a , <i>E. coli</i> WU361089 ^a	10 µL/plate	+ for TA100, TA1950, WU361089 - for TA1535	Not determined	Osterman-Golkar et al. (1983)
Reverse mutation	<i>S. typhimurium</i> TA100		+	Not determined	Zeiger (1990)
Reverse mutation	<i>S. typhimurium</i> TA100, NG54 ^c	2- and 6-hr exposures to 0, 2,500, 5,000, 7,500, 10,000 ppm	+	+	Dillon et al. (1992)
Reverse mutation	<i>S. typhimurium</i> TA100, TA1535 and TA1538 (+GSTA1-1 and GSTP1-1)	0, 50, 100, and 200 µL/plate	+ for TA100 - for TA1535, TA1538	Not determined	Simula et al. (1993)
Reverse mutation	<i>S. typhimurium</i> TA1535 (+GST5-5), TA1535 (wild type)	0-2.0 mM/plate	+ for TA1535 (+GST 5-5) - for TA1535 (wild type)	Not determined	Pegram et al. (1997); Thier et al. (1993)

(Table 4-29 continues on next page)

Table 4-29. Results from in vitro genotoxicity assays of dichloromethane with bacteria, yeast, or fungi

Assay	Test system	Concentration(s)	Results		Reference
			Without metabolic activation -S9	With metabolic activation +S9	
Reverse mutation	<i>S. typhimurium</i> TA100, TA100/NG-11 ^d	0, 30, 60, 130 mM/plate	++ for TA100 + for TA100/NG-11	Not determined	Graves et al. (1994a)
Reverse mutation	<i>S. typhimurium</i> TA100, RSJ100 ^c	Up to 24,000 ppm	+ for TA100 + for RSJ100	+ for TA100 + for RSJ100	DeMarini et al. (1997)
Forward mutation	<i>S. typhimurium</i> BA13	0–130 µmol/plate	+++	+	Roldán-Arjona and Pueyo (1993)
Gene mutation	<i>S. typhimurium</i> TA1535/pSK1002 ^c , NM5004 ^c	0, 2.5, 5.0, 10, 20 mM	+ NM5004 – TA1535/ pSK100 2	Not determined	Oda et al. (1996)
Prophage induction	<i>E. coli</i> K-39 (λ)	10 µL/plate	+	Not determined	Osterman-Golkar et al. (1983)
Reverse mutation	<i>E. coli</i> WP2 uvra pKM101	2- and 6-hr exposures to 6,300, 12,500, 25,000, and 50,000 ppm	+	+	Dillon et al. (1992)
Forward mutation	<i>E. coli</i> K12	0, 30, 60, 130 mM/plate	–	+	Graves et al. (1994a)
Forward mutation	<i>E. coli</i> Uvr ⁺ , UvrB ⁻	20,000 ppm	+	Not determined	Zielenska et al. (1993)
Fungi and yeasts					
Mitotic segregation	<i>Aspergillus nidulans</i>	Up to 8,000 ppm	+ only at 4,000 ppm; no dose-response relationship established	Not determined	Crebelli et al. (1988)
Gene conversion and recombination	<i>Saccharomyces cerevisiae</i>	Up to 209 mM	+	Not determined	Callen et al. (1980)

^aBacterial strains that have GSH (e.g., TA100, TA 98).

^bBacterial strains that do not have GSH (e.g., TA1535).

^cBacterial strains engineered to have more GSH activity than wild type.

^dBacterial strains engineered to have less GSH activity than wild type.

Dichloromethane induced mutations in *Salmonella typhimurium* strains containing GSH (e.g., TA100, TA98). These effects were not markedly influenced by the addition of exogenous mammalian liver fractions, suggesting that endogenous metabolism in these strains was sufficient to activate dichloromethane (Green, 1983; Jongen et al., 1982; 1978; Gocke et al., 1981). In support of this hypothesis, dichloromethane exposure of NG-11, a glutathione-deficient variant of *S. typhimurium* strain TA100, produced twofold fewer base-pair mutations compared with exposure of strain TA100, which produces normal levels of GSH. Furthermore, this difference was not apparent when the culture medium contained 1 mM GSH (Graves et al., 1994a).

In contrast to strain TA100, *S. typhimurium* strains TA1535, TA1537, and TA1538 (strains deficient in GSH) did not develop base-pair mutations in response to dichloromethane exposure (Pegram et al., 1997; Simula et al., 1993; Thier et al., 1993; Osterman-Golkar et al., 1983; Gocke et al., 1981). However, when strain TA1535 was transfected with rat GST-T1, dichloromethane induced base-pair reverse mutations (DeMarini et al., 1997; Pegram et al., 1997; Thier et al., 1993). A 60-fold higher concentration of dichloromethane was needed to induce a response (i.e., a sixfold increase over background levels in reverse mutations) in *S. typhimurium* strain TA100 than in TA1535 transfected with rat GST-T1 (DeMarini et al., 1997). This study also included several trihalomethanes; dichloromethane was several-fold less genotoxic than dibromochloromethane or bromoform, but was similar in potency to bromodichloromethane (DeMarini et al., 1997; Pegram et al., 1997). The authors suggest that these results support a role of GST-T1 in the mutagenicity of the trihalomethanes.

The mutagenic effects of dichloromethane have also been examined in fungi and yeast assays with both systems reporting positive results. Fungi assays were positive for mitotic segregation in *Asperigillus nidulans* (Crebelli et al., 1988), but there was not a dose response relationship as only the 4,000 ppm dichloromethane exposure was positive (exposure up to 8,000 ppm). A yeast assay was positive for gene conversion and recombination in *Saccharomyces cerevisiae* for concentrations up to 209 mM (Callen et al., 1980).

Mammalian assays. In the in vitro mammalian system studies conducted with murine cell lines (Table 4-30), dichloromethane was negative for producing point mutations in the mouse lymphoma L5178Y cell line (Thilagar et al., 1984) but was positive in producing DNA single stranded breaks (SSBs) in mouse Clara cells (Graves et al., 1995) and mouse hepatocytes (Graves et al., 1994b). Given that exposure to dichloromethane results specifically in lung and liver tumors, this pattern is not surprising. Additionally, GST is localized in the nucleus of hepatocytes and lung cells in the mouse (Mainwaring et al., 1996), which would also increase sensitivity of these particular cell fractions to genotoxic effects of dichloromethane. DNA SSBs were induced at lower concentrations in mouse hepatocytes (0.5 mM) than in rat hepatocytes (30 mM). The extent of DNA damage was shown to be reduced to the background level seen in control (no exposure) conditions by pretreating the cells with buthionine sulfoxime to deplete

cellular levels of GSH and thus inhibit dichloromethane metabolism via the GST pathway (Graves et al., 1995, 1994b). Similar results were seen in mouse lung Clara cells. Freshly isolated Clara cells from the lungs of B6C3F₁ mice also showed significantly increased, concentration-dependent amounts of DNA SSBs when incubated in vitro for 2 hours in the presence of 5–60 mM dichloromethane. Pretreatment with buthionine sulphoximine before Clara-cell isolation or the presence of buthionine sulphoximine in the culture medium decreased the amount of in vitro DNA damage induced.

Table 4-30. Results from in vitro genotoxicity assays of dichloromethane with mammalian systems, by type of test

Assay	Test system	Concentrations	Results	Reference
Mouse				
Point mutation	Mouse lymphoma L5178Y cells	Not provided	Negative	Thilagar et al. (1984)
DNA SSBs by alkaline elution	Mouse hepatocytes (B6C3F ₁)	0, 0.4, 3.0, 5.5 mM	Positive at 0.4 mM	Graves et al. (1994b)
DNA SSBs by alkaline elution	Mouse Clara cells (B6C3F ₁)	0, 5, 10, 30, 60 mM	Positive, but DNA damage was reduced by incubating in the presence of GSH depletory	Graves et al. (1995)
DNA-protein cross-links	Mouse hepatocytes (B6C3F ₁)	0.5–5 mM	Positive	Casanova et al. (1997)
Rat				
Unscheduled DNA synthesis	Rat hepatocytes	Up to 16 mM (measured); 30 mM (nominal)	Negative	Andrae and Wolff (1983)
Unscheduled DNA synthesis	Rat hepatocytes	Not provided	Marginally positive	Thilagar et al. (1984)
DNA SSBs by alkaline elution	Rat hepatocytes	0, 30, 90, 90 mM	Positive at 30 mM	Graves et al. (1994b)
DNA-protein cross-links	Rat hepatocytes	0.5–5 mM	Negative	Casanova et al. (1997)
Hamster with GST activity from mouse				
DNA-protein cross-links	Chinese hamster ovary cells	60 mM	Positive with mouse liver cytosol (negative without) at much higher concentrations of dichloromethane (60 mM) than formaldehyde (0.5–4 mM)	Graves et al. (1994b)
HPRT mutation analysis	Chinese hamster ovary cells	2,500 ppm	Positive with mouse liver cytosol	Graves et al. (1996)
DNA SSBs and DNA-protein cross-links	Chinese hamster ovary cells	3,000 ppm (0.3%, volume per volume [v/v]) and 5,000 ppm (0.5%, v/v)	Positive at concentration of 0.5% (v/v) for SSBs in presence of mouse liver cytosol, but increase in DNA-protein cross-links marginal; formaldehyde (in absence of mouse liver cytosol) was positive at 0.5 mM for both DNA SSBs and DNA-protein cross-links; Chinese hamster ovary cell cultures were suspended	Graves and Green (1996)
DNA-protein cross-links	Syrian golden hamster hepatocytes	0.5–5 mM	Negative	Casanova et al. (1997)
Comet assay	V79 hamster cells transfected with mouse GST-T1	2.5, 5, 10 mM	A significant, dose-dependent increase in DNA damage resulting from DNA-protein cross-links in V79 cells transfected with mouse GST-T1 compared to parental cells	Hu et al. (2006)

Table 4-30. Results from in vitro genotoxicity assays of dichloromethane with mammalian systems, by type of test

Assay	Test system	Concentrations	Results	Reference
Hamster without GST activity from mouse				
Forward mutation	Chinese hamster epithelial cells	5,000, 10,000, 30,000, 50,000 ppm	Negative	Jongen et al. (1981)
Unscheduled DNA synthesis	Chinese hamster epithelial cells	5,000, 10,000, 30,000, 50,000 ppm	Negative	Jongen et al. (1981)
Sister chromatid exchange	Chinese hamster epithelial cells	5,000, 10,000, 20,000, 30,000, and 40,000 ppm	Weak positive with or without rat-liver microsomal system	Jongen et al. (1981)
Chromosomal aberrations	Chinese hamster ovary cells	Not provided	Positive, independent of rat liver S9	Thilagar and Kumaroo (1983)
Sister chromatid exchange	Chinese hamster ovary cells	Not provided	Negative with or without rat liver S9	Thilagar and Kumaroo (1983)
DNA and protein synthesis	Chinese hamster ovary cells	Up to 1,000 µg/mL	Negative	Garrett and Lewtas (1983)
DNA SSBs by alkaline elution	Hamster hepatocytes	0.4–90 mM	Negative	Graves et al. (1995)
Calf				
DNA Adducts	Calf thymus DNA	50 mM	Positive in the presence of bacterial GST DM11 and dichloromethane dehalogenase; adducts primarily formed with the guanine residues	Kayser and Vuilleumier (2001)
DNA Adducts	Calf thymus DNA	0–8.0 µmol (0–60 mM)	Positive in the presence of bacterial GST DM11, rat GST5-5, and human GSTT11; adducts primarily formed with the guanine residues	Marsch et al. (2004)
Human				
Unscheduled DNA synthesis	Human peripheral lymphocytes	250, 500, 1,000 ppm	Negative with or without rat liver S9	Perocco and Prodi (1981)
Unscheduled DNA synthesis	Primary human fibroblast	5,000, 10,000, 30,000, 50,000 ppm	Negative	Jongen et al. (1981)
Sister chromatid exchange	Human peripheral lymphocytes	Not provided	Weak positive	Thilagar et al. (1984)
Chromosomal aberrations	Human peripheral lymphocytes	Not provided	Positive	Thilagar et al. (1984)
DNA SSBs by alkaline elution	Human hepatocytes	Up to 120 mM	Negative at concentrations between 5 and 120 mM	Graves et al. (1995)

Table 4-30. Results from in vitro genotoxicity assays of dichloromethane with mammalian systems, by type of test

Assay	Test system	Concentrations	Results	Reference
Micronucleus test	Human AHH-1, MCL-5, h2E1 cell lines	Up to 10 mM	Positive in all three cell lines	Doherty et al. (1996)
DNA-protein cross-links	Mouse, rat, hamster, human hepatocytes	0.5–5 mM	Negative	Casanova et al. (1997)
DNA damage by comet assay	Primary human lung epithelial cells	10, 100, 1,000 μ M	Weak trend, independent of GST activity (GST enzymatic activity not present in the cultured cells)	Landi et al. (2003)

HPRT = hypoxanthine-guanine phosphoribosyl transferase

In a series of experiments with freshly isolated hepatocytes from multiple species (Table 4-30), DNA-protein cross-links were detected in hepatocytes of B6C3F₁ mice but not in hepatocytes of F344 rats, Syrian golden hamsters, or three human subjects following 2-hour in vitro exposures to concentrations ranging from 0.5 to 5 mM dichloromethane (Casanova et al., 1997). Within the range of concentrations tested, DNA-protein cross-links in mouse hepatocytes appeared to increase with increasing concentration of dichloromethane.

Negative results for dichloromethane were predominantly seen in in vitro test systems that used rat or hamster cell lines with low or no GST activity (Table 4-30). Several genotoxic endpoints including DNA and protein synthesis (Garrett and Lewtas, 1983), chromosomal aberrations or sister chromatid exchanges (Thilagar et al., 1984; Thilagar and Kumaroo, 1983; Jongen et al., 1981), unscheduled DNA synthesis (Thilagar et al., 1984; Andrae and Wolff, 1983; Jongen et al., 1981), and mutations (Thilagar et al., 1984; Jongen et al., 1981) were evaluated in these cell lines. In contrast, positive results (DNA-protein cross-links and DNA SSBs) were observed when mouse liver cytosol was included in Chinese hamster ovary (CHO) cells (Graves et al., 1995, 1994b). Dichloromethane also induced hypoxanthine-guanine phosphoribosyl transferase (HPRT) gene mutations in CHO cells when they were incubated with GST-competent mouse liver cytosol preparations (Graves et al., 1996).

The instability of the S-(chloromethyl)glutathione-adducts presents considerable challenges to studies of these products (Hashmi et al., 1994). Kayser and Vuilleumier (2001), however, demonstrated the formation of DNA adducts with radiolabeled dichloromethane in calf thymus DNA in the presence of dichloromethane dehalogenase/GST purified from a bacterial source (*Methylophilus* sp. strain DM11) and GSH (Table 4-30). The type of adduct could not be identified because of low yield, but it was determined that guanine was more actively incorporated than cytosine, adenine, or thymine by at least twofold in the presence of GST-activated dichloromethane, indicating a base specificity for these adducts. Incubation of calf thymus DNA with formaldehyde and GSH, however, did not result in detectable DNA adduct formation. In another study, Marsch et al. (2004) further evaluated the presence of adducts in calf thymus DNA in the presence of dichloromethane and human (GST-T1), rat (GST5-5), or bacterial (DM11) GST (Marsch et al., 2004). This study found that all three enzymes yielded a similar pattern of adduct formation, forming primarily with guanine and to a lesser extent with cytosine, adenine, and thymine (two- to threefold less than guanine), consistent with the results reported by Kayser and Vuilleumier (2001). High levels of guanosine-specific adducts were also seen with S-(1-acetoxymethyl)glutathione, a compound that is structurally similar but more stable than S-(chloromethyl)glutathione (Marsch et al., 2001). These findings indicate that the S-(chloromethyl)glutathione intermediate formed by GSH conjugation has mutagenic potential and is likely responsible, at least in part, for the mutagenic response observed following dichloromethane exposure.

In studies with human cell lines or isolated cells, positive results were reported for sister chromatid exchanges and chromosomal aberrations (Thilagar et al., 1984) and in the micronucleus test (Doherty et al., 1996). Negative results with human cells were seen in the unscheduled DNA synthesis assays (Jongen et al., 1981; Perocco and Prodi, 1981), DNA SSBs, and DNA-protein cross-links (Casanova et al., 1997; Graves et al., 1995).

Dichloromethane-induced DNA damage (comet assay) was examined in primary cultures of human lung epithelial cells collected by brush biopsy from four healthy volunteers (Landi et al., 2003). This study was designed to assess the genotoxicity of four trihalomethanes (chloroform, bromodichloromethane, dibromochloromethane, and bromoform), with dichloromethane included because of its known activation by GST-T1. Two of the subjects were of the GST-T1⁺ genotype, and two were of the GST-T1⁻ genotype.⁶ The cells had been frozen, and GST activity was not detected in the cultured cells. DNA damage was reported to occur in the combined GST-T1⁻ samples (tail extent moment 7.1, 13.7, and 15.3 in the 10, 100, and 1,000 μ M dichloromethane groups, respectively) but not in the combined GST-T1⁺ samples (tail extent moment 8.1, 11.5, and 10.4 in the 10, 100, and 1,000 μ M dichloromethane groups, respectively). This pattern was not seen across the individual samples, however, as only one sample exhibited a clear dose-response gradient. Given the absence of GST activity, an analysis combining the four samples could provide a more informative picture of the dose-response relationship between dichloromethane (and the other compounds studied) and DNA damage. For dichloromethane, values of 9.4, 7.6, 12.6, and 12.9 were seen in the 0, 10, 100, and 1,000 μ M groups, respectively. This pattern was similar to that seen with chloroform (9.4, 6.9, 11.4, and 12.7 in the 0, the 10, 100, and 1,000 μ M groups, respectively) but weaker than the pattern for bromoform (9.4, 12.5, 15.8, and 18.2 in the 0, the 10, 100, and 1,000 μ M groups, respectively), and much weaker than the pattern for bromodichloromethane (9.4, 25.2, 28.5, and 39.1 in the 0, the 10, 100, and 1,000 μ M groups, respectively).⁷ No dose-response gradient was seen with dibromochloromethane (9.4, 6.5, 8.1, and 8.0 in the 0, 10, 100, and 1,000 μ M groups, respectively). This relative pattern is also seen in the estimated slopes (beta coefficient for the change in tail extent moment per unit increase in μ M concentration): 0.0, 0.003, 0.004, 0.006, and 0.02 for dibromochloromethane, dichloromethane, chloroform, bromoform, and bromodichloromethane, respectively (statistical significance not reported).

A stronger and more consistent response was seen under the same experimental conditions with bromodichloromethane, but dibromochloromethane resulted in no increase in DNA damage in any of the donor cells at any concentration tested.

⁶Landi et al. (2003) did not clearly describe their treatment of GST-T1^{+/-} heterozygote genotypes; EPA considers it likely that they were included in the pool from which the GST-T1⁺ samples were drawn. In addition, there is a discrepancy in the paper regarding the coding of the GST-T1 genotypes. Samples A and C are noted to be the GST-T1⁻ samples in one part of the paper, and C and D are described as the GST-T1⁻ samples in another part of the paper.

⁷These values are based on the mean of the GST-T1⁺ and the GST-T1⁻ samples from Table 1 of Landi et al. (2003).

Several studies have examined patterns of mutations or DNA damage with dichloromethane and formaldehyde to assess the relative role of S-(chloromethyl)glutathione and formaldehyde in the observed genotoxicity. In a study in CHO cells incubated with dichloromethane (0.3% plus mouse liver cytosol), 2.5-fold increases in DNA-protein cross-links that are indicative of formaldehyde exposure were observed, compared with a 25-fold increase when 1 mM formaldehyde was added directly to cultures. Both treatments induced a comparable degree of DNA SSBs (Graves and Green, 1996). In a subsequent study, Graves et al. (1996) compared the mutational spectra induced by dichloromethane to that induced by direct addition of formaldehyde or 1,2-dibromoethane (a chemical known to act through a glutathionyl conjugate metabolite) at the HPRT locus in CHO cells. The mutations induced by dichloromethane and 1,2-dibromoethane were predominantly GC to AT transitions, while all six formaldehyde-induced mutants sequenced were single base transversions. This provided further evidence that the S-(chloromethyl)glutathione intermediate may be primarily responsible for dichloromethane genotoxicity. In contrast, Hu et al. (2006) found evidence of significant amounts of formaldehyde formation following dichloromethane exposure in the cytosol of V79 (hamster) cells transfected with the murine GST-T1 gene compared to the parent cell line. In accordance with this, they observed concentration-dependent increases in DNA-protein cross-links in the GST-T1 transfected cells using the comet assay with and without proteinase K treatment that frees DNA from cross-links and allows DNA migration. These findings are consistent with those by Casanova et al. (1997), who performed a comparison of the amounts of DNA-protein and RNA-formaldehyde cross-links formed following dichloromethane exposure in hepatocytes isolated from mice, rats, hamsters, and human GST-T1 genetic variants. Only DNA-protein cross-links were observed in mouse hepatocytes, but RNA-formaldehyde cross-links were found in all species and were highest in the mouse hepatocytes (4-, 7-, and 14-fold higher than rats, humans, and hamsters, respectively). These results showed that human hepatocytes can metabolize dichloromethane to formaldehyde, resulting in RNA-formaldehyde cross-links. In addition, the results indicate that there is considerable variation among species and that the human variation in the GST-T1 gene can also affect the amount of formaldehyde produced. The authors also noted that comparing results following ectopic addition of formaldehyde directly to cells with results following dichloromethane metabolism in situ can be misleading, as the formaldehyde produced internally may reside in different locations intracellularly, potentially affecting the capability of interacting with DNA. These results show that, while most studies indicate the importance of the S-(chloromethyl)glutathione intermediate in mediating genotoxic damage following dichloromethane exposure, DNA damage resulting from formaldehyde formation should also be considered.

4.5.1.2. *In Vivo* Genotoxicity Assays

Genotoxicity findings in *Drosophila melanogaster* assays are mixed (Table 4-31). A study of gene mutation in *D. melanogaster* showed a marginal increase in sex-linked recessive deaths following oral exposure (Gocke et al., 1981). An additional feeding study (Rodriguez-Arnaiz, 1998) reported a positive response in the somatic w/w⁺ assay. A third study of *D. melanogaster* (Kramers et al., 1991) found no evidence of increased sex-linked recessive deaths, somatic mutations, or recombinations following exposure to airborne dichloromethane.

Table 4-31. Results from in vivo genotoxicity assays of dichloromethane in insects

Assay	Test system	Doses	Result	Reference
Gene mutation (sex-linked recessive lethal)	<i>Drosophila</i>	125, 620 mM	Positive (feeding exposure)	Gocke et al. (1981)
Gene mutation (sex-linked recessive lethal, somatic mutation and recombination)	<i>Drosophila</i>	6 hrs—1,850, 5,500 ppm 1 wk—2,360, 4,660 ppm 2 wks—1,370, 2,360 ppm (all approximate)	Negative (inhalation exposure)	Kramers et al. (1991)
Somatic w/w ⁺ assay	<i>Drosophila</i>	50, 100, 250, 500 mM	Positive (feeding exposure)	Rodriguez-Arnaiz (1998)

Some in vivo studies investigating certain genotoxic endpoints in mice exposed to dichloromethane produced negative results (Table 4-32). Unscheduled DNA synthesis was not induced in hepatocytes from mice (and rats) after 2- or 6-hour inhalation exposures to concentrations that were carcinogenic in the NTP (1986) mouse bioassay (Trueman and Ashby, 1987) or after other exposure routes (Lefevre and Ashby, 1989). Although positive results were not observed in the unscheduled DNA synthesis studies, it is generally recognized that this assay is not sensitive for detecting genotoxic chemicals (Eastmond et al., 2009; Madle et al., 1994). Distinct, unequivocal cytogenetic effects (e.g., induction of micronuclei, sister chromatid exchanges, or chromosome aberrations) were not consistently found in bone marrow or erythrocytes in several studies of mice after acute oral exposures (Sheldon et al., 1987) or parenteral exposures (Westbrook-Collins et al., 1990; Gocke et al., 1981). However, tumorigenic effects in mice are generally localized to the liver and lung (due to high GST activity) and therefore, it is not surprising that genotoxic effects were for the most part not observed in the bone marrow or erythrocytes (cell types with minimal GST activity). Crebelli et al. (1999) stated that genotoxic effects induced by halogenated hydrocarbons (such as dichloromethane) are not very effective in inducing micronucleus formation in mouse bone marrow, and a negative bone marrow micronucleus assay should not offset the consistently positive in vitro results (Dearfield and Moore, 2005).

Table 4-32. Results from in vivo genotoxicity assays of dichloromethane in mice

Assay	Test system	Route and dose	Duration	Results	Reference
Micronucleus test	Mouse bone marrow (NMRI)	425, 850, or 1,700 mg/kg	Two doses	Negative at all doses	Gocke et al. (1981)
Micronucleus test	Mouse bone marrow (C57BL/6J/A1pk)	Gavage, 1,250, 2,500, and 4,000 mg/kg	Single dose	Negative at all doses	Sheldon et al. (1987)
Micronucleus test	Mouse peripheral red blood cells (B6C3F ₁)	Inhalation 6 hr/d, 5 d/wk, 0, 4,000, 8,000 ppm	2 wk	Positive at 4,000 and 8,000 ppm	Allen et al. (1990)
Micronucleus test	Mouse peripheral red blood cells (B6C3F ₁)	Inhalation, 6 hr/d, 5 d/wk, 0, 2,000 ppm	12 wks	Positive at 2,000 ppm	Allen et al. (1990)
DNA synthesis	Mouse liver (B6C3F ₁)	Gavage, 1,000 mg/kg; inhalation, 4,000 ppm	Single dose; 2 hrs	Negative in both oral and inhalation studies	Lefevre and Ashby (1989)
Unscheduled DNA synthesis	Mouse hepatocytes (B6C3F ₁)	Inhalation, 2,000 and 4,000 ppm.	2 or 6 hrs	Negative	Trueman and Ashby (1987)
Sister chromatid exchange	Mouse bone marrow (C57BL/6J)	Intraperitoneal, 100, 1,000, 1,500, 2,000 mg/kg	Single dose	Negative	Westbrook-Collins et al. (1990)
Sister chromatid exchange	Mouse bone marrow (B6C3F ₁)	Subcutaneous, 0, 2,500, 5,000 mg/kg	Single dose	Negative at all doses	Allen et al. (1990)
Sister chromatid exchange	Mouse lung cells and peripheral lymphocytes (B6C3F ₁)	Inhalation 6 hr/d, 5 d/wk, 0, 4,000, 8,000 ppm	2 wks	Positive at 8,000 ppm	Allen et al. (1990)
Sister chromatid exchange	Mouse lung cells (B6C3F ₁)	Inhalation 6 hr/d, 5 d/wk, 0, 2,000 ppm	12 wks	Positive at 2,000 ppm	Allen et al. (1990)
Chromosome aberrations	Mouse bone marrow (C57BL/6J)	Intraperitoneal, 100, 1,000, 1,500, 2,000 mg/kg	Single dose	Negative	Westbrook-Collins et al. (1990)
Chromosome aberrations	Mouse bone marrow (B6C3F ₁)	Subcutaneous, 0, 2,500, 5,000 mg/kg	Single dose	Negative	Allen et al. (1990)
Chromosome aberrations	Mouse lung and bone marrow cells (B6C3F ₁)	Inhalation, 6 hr/d, 5 d/wk, 0, 4,000, 8,000 ppm	2 wks	Positive at 8,000 ppm	Allen et al. (1990)
DNA-protein cross-links	Mouse liver and lung cells (B6C3F ₁)	Inhalation, 6 hr/d, 3 d, 4,000 ppm	3 d	Positive in mouse liver cells at 4,000 ppm; negative in mouse lung cells	Casanova et al. (1992)

(Table 4-32 continues on next page)

Table 4-32. Results from in vivo genotoxicity assays of dichloromethane in mice

Assay	Test system	Route and dose	Duration	Results	Reference
DNA-protein cross-links	Mouse liver and lung cells (B6C3F ₁)	Inhalation, 6 hr/d, 150, 500, 1,500, 3,000, 4,000 ppm	3 d	Positive in mouse liver cells at 500–4,000 ppm; negative in mouse lung cells	Casanova et al. (1996)
DNA SSBs by alkaline elution	Mouse hepatocytes (B6C3F ₁)	Inhalation, 2,000 and 4,000 ppm	3 or 6 hrs	Positive at 4,000 ppm at 3 and 6 hrs	Graves et al. (1994b)
DNA SSBs by alkaline elution	Mouse liver and lung homogenate (B6C3F ₁)	Liver: inhalation, 2,000, 4,000, 6,000, 8,000 ppm Lung: inhalation, 1,000, 2,000, 4,000, 6,000 ppm	3 hrs 3 hrs	Liver: positive at 4,000–8,000 ppm Lung: positive at 2,000–4,000 ppm	Graves et al. (1995)
DNA damage by comet assay	Mouse liver and lung cells (CD-1)	Gavage, 1,720 mg/kg; organs harvested at 0 (control), 3, and 24 hrs	Single dose	Positive only at 24 hrs after dosing	Sasaki et al. (1998)
DNA damage by comet assay	Mouse stomach, urinary bladder, kidney, brain, bone marrow (CD-1)	Gavage, 1,720 mg/kg; organs harvested at 0 (control), 3, and 24 hrs	Single dose	Negative 3 or 24 hr after dosing	Sasaki et al. (1998)
DNA adducts	Mouse liver and kidney cells (B6C3F ₁)	Intraperitoneal, 5 mg/kg	Single dose	Negative	Watanabe et al. (2007)
Kras and Hras oncogenes	Mouse liver and lung tumors (B6C3F ₁)	0, 2,000 ppm	Up to 104 wks	No difference in mutation profile between control and dichloromethane-induced liver tumors; number of spontaneous lung tumors (n = 4) limits comparison at this site	Devereux et al. (1993)
p53 tumor suppressor gene	Mouse liver and lung tumors (B6C3F ₁)	0, 2,000 ppm	Up to 104 wks	Loss of heterozygosity infrequently seen	Hegi et al. (1993)

When genotoxic endpoints were examined in the cancer target tissues (liver and lung) in mice exposed to dichloromethane, positive results were consistently reported (Table 4-32). These findings provide supporting evidence that GST-pathway metabolites may be key actors in the genotoxic effects and carcinogenic mode of action for dichloromethane. Increased sister chromatid exchanges were found in lung cells and peripheral lymphocytes from mice exposed by inhalation for 2 weeks to 8,000 ppm or for 12 weeks to 2,000 ppm (Allen et al., 1990). Under the same exposure conditions, increased chromosomal aberrations in lung and bone cells and micronuclei in peripheral red blood cells also were found (Allen et al., 1990). DNA-protein cross-links were detected in mouse hepatocytes but not in lung cells after a 3-day inhalation exposure to 4,000 ppm (Casanova et al., 1992) and between 500 and 4,000 ppm (Casanova et al., 1996). DNA damage, detected as increased DNA SSBs, was observed in liver and lung tissue of B6C3F₁ mice immediately following 3-hour exposures (Graves et al., 1995). The DNA damage was not detectable 2 hours after in vivo exposure, indicating that DNA repair occurs rapidly. Pretreatment of mice with buthionine sulphoximine, a GSH depletor, caused a decrease to levels seen in controls in the amount of DNA damage detected immediately after in vivo exposure in liver and lung tissue, indicating GSH involvement in the genotoxic process. DNA damage (detected by the comet assay) was also reported in liver and lung tissues from male CD-1 mice sacrificed 24 hours after administration of a single oral dose of 1,720 mg/kg of dichloromethane (Sasaki et al., 1998). In this study, DNA damage in lung and liver was not detected 3 hours after dose administration, and no DNA damage occurred at either time point in several other tissues in which a carcinogenic response was not seen in chronic animal cancer bioassays (e.g., stomach, kidney, bone marrow).

Formation of DNA adducts was evaluated in male and female B6C3F₁ mice as well as in male F344 rats (Watanabe et al., 2007). Animals were administered 5 mg/kg intraperitoneally of radiolabeled dichloromethane and sacrificed at 1 or 8 hours after administration. The kidneys and livers were removed and the DNA was isolated from these tissues to evaluate formation of DNA adducts. At the administered dose, DNA adducts were not detected.

Other studies in mice have looked for mutations in specific oncogenes (*K-ras* or *H-ras*) (Devereux et al., 1993) or in a tumor suppressor gene (*p53*) (Hegi et al., 1993) in liver or lung tumors from dichloromethane-exposed mice. These studies have not demonstrated exposure-related patterns of mutations in these genes, although it should be noted that the statistical power of this analysis for the lung tumors is limited (discussed further in Sections 4.5.2 and 4.5.3).

Results from in vivo studies in other mammals (i.e., rats and hamsters) of hepatocyte sensitivity to dichloromethane induction of DNA SSBs (Table 4-33) are consistent with interspecies differences in the induction of liver tumors in the inhalation cancer bioassays. A gavage study in rats reported the presence of DNA SSBs with a dose of 1,275 mg/kg (Kitchin and Brown, 1989). The other available studies, however, did not find any genotoxicity following dichloromethane exposure. No increase in unscheduled DNA synthesis in rat hepatocytes was

seen following inhalation of dichloromethane for 2–6 hours at 2,000 or 4,000 ppm (Trueman and Ashby, 1987), exposure by gavage up to 1,000 mg/kg (Trueman and Ashby, 1987), or intraperitoneal exposure of 400 mg/kg (Mirsalis et al., 1989). DNA adducts were not detected in the livers and kidneys of male F344 rats dosed with 5 mg/kg dichloromethane intraperitoneally (Watanabe et al., 2007). DNA SSBs were significantly increased in hepatocytes isolated from B6C3F₁ mice exposed to 4,831 ppm (4,000 ppm nominal) for 6 hours but were not increased in hepatocytes from Sprague-Dawley rats exposed to 4,527 ppm (4,000 ppm nominal) for 6 hours (Graves et al., 1994b). Results from *in vivo* interspecies comparisons of dichloromethane induction of DNA-protein cross-links in hepatocytes (expected products of the GSH pathway) are also consistent with the hypothesis that the mouse is more sensitive than other mammalian species due to greater activity of the GST pathway. DNA-protein cross-links were formed in the liver of mice but not hamsters following *in vivo* exposure to air concentrations ranging from 500 to 4,000 ppm, 6 hours/day for 3 days (Casanova et al., 1996). The absence of a genotoxic response in the rat and hamster is consistent with considerably lower GST activity and therefore, these mammalian systems would be expected to be less sensitive at detecting genotoxic effects than the studies conducted in mice.

Table 4-33. Results from in vivo genotoxicity assays of dichloromethane in rats and hamsters

Assay	Test system	Route and dose	Duration	Results	Reference
Unscheduled DNA synthesis	Rat hepatocytes	Gavage, 100, 500, 1,000 mg/kg	Liver harvested 4 and 12 hrs after dosing	Negative 4 or 12 hrs after dosing	Trueman and Ashby (1987)
Unscheduled DNA synthesis	Rat hepatocytes	Inhalation, 2 or 6 hrs, 2,000 and 4,000 ppm	2 or 6 hrs	Negative at both concentrations and exposure durations	Trueman and Ashby (1987)
Unscheduled DNA synthesis	Rat hepatocytes	Intraperitoneal, single dose, 400 mg/kg	Single dose	Negative 48 hrs after dosing	Mirsalis et al. (1989)
DNA SSBs by alkaline elution	Rat hepatocytes	Inhalation, 3 or 6 hrs, 2,000 and 4,000 ppm	3 or 6 hrs	Negative at all concentrations and time points	Graves et al. (1994b)
DNA SSBs by alkaline elution	Rat liver homogenate	Gavage, 2 doses, 425 mg/kg and 1,275 mg/kg, administered 4 and 21 hrs before liver harvesting	4 or 21 hrs (time between dosing and liver harvesting)	Positive at 1,275 mg/kg	Kitchin and Brown (1989)
DNA SSBs by alkaline elution	Rat liver and lung homogenate	Liver: inhalation, 4,000, 5,000 ppm Lung: inhalation, 4,000 ppm	3 hrs 3 hrs	Negative for both liver and lung at all concentrations	Graves et al. (1995)
DNA-protein cross-links	Hamster liver and lung cells	Inhalation, 6 hr/d, 500, 1,500, 4,000 ppm	3 d	Negative at all concentrations	Casanova et al. (1996)
DNA adducts	Rat liver and kidney cells	Intraperitoneal, 5 mg/kg	Single dose	Negative	Watanabe et al. (2007)

Table 4-34 compares results from studies of mice and rats in which comparable tissue-specific endpoints were examined in in vivo genotoxicity assays. Several of the endpoints that were positive in mice (e.g., sister chromatid exchange, DNA-protein cross-links, comet assay) have not been examined in the rat. Unscheduled DNA synthesis has been demonstrated in mouse but not in rat hepatocytes. In contrast to the positive results seen in mouse inhalation exposure studies, DNA SSB induction was not seen in rat inhalation studies but was seen in an gavage study.

Table 4-34. Comparison of in vivo dichloromethane genotoxicity assays targeted to lung or liver cells, by species

Studies in B6C3F ₁ mice					Studies in rats			
Assay	Test system	Route, dose (duration)	Results	Reference	Test system	Route, dose (duration)	Results	Reference
DNA synthesis	Liver	Gavage, 1,000 mg/kg; inhalation, 4,000 ppm (2 hrs)	Negative in oral and inhalation studies	Lefevre and Ashby (1989)				No studies
Unscheduled DNA synthesis	Hepatocytes	Inhalation, 2,000 and 4,000 ppm (2 or 6 hrs)	Negative	Trueman and Ashby (1987)	Hepatocytes	Inhalation, 2,000 and 4,000 ppm (2 or 6 hrs)	Negative	Trueman and Ashby (1987)
Unscheduled DNA synthesis					Hepatocytes	Intraperitoneal, 400 mg/kg	Negative	Mirsalis et al. (1989)
Sister chromatid exchange	Lung cells	Inhalation 6 hr/d, 5 d/wk, 0, 4,000, 8,000 ppm (2 wks) Inhalation 6 hr/d, 5 d/wk, 0, 2,000 ppm (12 wks)	Positive at 8,000 ppm Positive at 2,000 ppm	Allen et al. (1990)				No studies
Chromosome aberrations	Lung cells	Inhalation, 6 hr/d, 5 d/wk, 0, 4,000, 8,000 ppm (2 wks)	Positive at 8,000 ppm	Allen et al. (1990)				No studies
DNA-protein cross-links	Liver and lung cells	Inhalation, 6 hr/d, 3 d, 4,000 ppm (3 d) Inhalation, 6 hr/d, 150, 500, 1,500, 3,000, 4,000 ppm (3 d)	Positive in liver 4,000 ppm Positive in liver at 500–4,000 ppm; both studies negative in lung	Casanova et al. (1992)				No studies
DNA SSBs by alkaline elution	Hepatocytes	Inhalation, 2,000 and 4,000 ppm (3 or 6 hrs)	Positive at 4,000 ppm	Graves et al. (1994b)	Hepatocytes	Inhalation, 3 or 6 hrs, 2,000 and 4,000 ppm	Negative at all concentrations and time points	Graves et al. (1994b)
DNA SSBs by alkaline elution	Liver and lung homogenate	Liver: inhalation, 2,000, 4,000, 6,000, 8,000 ppm (3 hrs) Lung: inhalation, 1,000, 2,000, 4,000, 6,000 ppm (3 hrs)	Liver: Positive at 4,000–8,000 ppm Lung: Positive at 2,000–4,000 ppm	Graves et al. (1995)	Liver and lung homogenate	Liver: inhalation, 4,000, 5,000 ppm Lung: inhalation, 4,000 ppm	Negative in liver and lung at all concentrations and time points	Graves et al. (1995)

(Table 4-34 continues on next page)

Table 4-34. Comparison of in vivo dichloromethane genotoxicity assays targeted to lung or liver cells, by species

Studies in B6C3F ₁ mice					Studies in rats			
Assay	Test system	Route, dose (duration)	Results	Reference	Test system	Route, dose (duration)	Results	Reference
DNA SSBs by alkaline elution					Liver homogenate	Gavage, 425 mg/kg and 1,275 mg/kg	Positive at 1,275 mg/kg	Kitchin and Brown (1989)
DNA damage by comet assay	Liver and lung cells	Gavage, 1,720 mg/kg; organs harvested at 0 (control), 3, and 24 hrs	Positive only at 24 hrs after dosing	Sasaki et al. (1998)				No studies
DNA adducts	Liver and kidney cells	Intraperitoneal, 5 mg/kg	Negative	Watanabe et al. (2007)	Liver and kidney cells	Intraperitoneal, 5 mg/kg	Negative	Watanabe et al. (2007)

In summary, the available data provide evidence for mutagenicity of dichloromethane. Most of the *in vitro* bacterial assays showed positive results when there was GST activity. Nonpositive results were reported only in bacterial assays with low GST activity; in experiments where GST was added, positive results were then observed. Evaluation of the *in vitro* mammalian studies also demonstrates consistency of the requirement for GST for observation of genotoxic effects. In rat and hamster cell lines where GST activity is significantly less than mouse, primarily negative results were reported following dichloromethane exposure. However, when mouse liver cytosol or transfected mouse GST were included in these same cell lines, mutagenic effects were reported after dichloromethane exposure. In mouse cell lines, positive results were obtained in Clara cells, but no effects were observed in a mouse lymphoma cell line, which is consistent with the absence of tumors in this site for mice. The results of *in vivo* mutagenicity in mice also provide support for the site-specificity of the observed tumors. Assays using mouse bone marrow were all negative. However, micronuclei and sister chromatid exchange tests in peripheral blood produced a positive response at high doses. With the exception of one study of unscheduled DNA synthesis in hepatocytes, numerous site-specific studies in either the liver or lung were also positive at various doses. These liver and lung studies included chromosomal aberrations, SSBs, sister chromatid exchanges, and DNA-protein cross-links and correspond to genotoxic and mutagenic effects associated with metabolites from the GST pathway.

4.5.2. Mechanistic Studies of Liver Effects

One of the major target organs from dichloromethane exposure is the liver, and several studies have focused on examining the potential mechanisms producing liver tumors. This section summarizes the primary mechanistic studies that were conducted in order to examine the hepatic tumors produced by dichloromethane in mice. A parallel set of studies, discussed in the next section, focus on potential mechanisms that produce lung tumors. Briefly, dichloromethane-induced liver tumors first appeared in mice after 52 weeks of exposure (Maronpot et al., 1995; Kari et al., 1993), which was when tumors began to appear in control mice, indicating a similar time course in tumor formation between treated and untreated groups. Onset of liver tumor formation is not preceded by liver cell proliferation (Casanova et al., 1996; Foley et al., 1993). Further mechanistic studies were conducted to assay the tumor for significant changes in proto-oncogene activation and tumor suppressor gene inactivation (Maronpot et al., 1995; Devereux et al., 1993; Hegi et al., 1993). A second subset of mechanistic studies was conducted to elucidate the reason that mice are the most sensitive species to liver tumors and if other species exhibited changes in liver function (Thier et al., 1998; Reitz et al., 1989). It was found that mice have the highest level of GST-T1 catalytic activity but that humans, rats, and hamsters, among other species, also metabolize dichloromethane in the liver to a GST conjugate. In contrast, there has been little research focusing on the mechanisms through

which noncancer hepatic effects (seen most strongly in the rat) are produced, and the role of the parent material, metabolites of the CYP2E1 pathway, metabolites of the GST pathway, or some combination of parent material and metabolites is not known.

Liver tumor characterization studies. Several studies have examined the time course of appearance of liver tumors in B6C3F₁ mice exposed to 2,000 or 4,000 ppm and possible links between hepatic nonneoplastic cytotoxicity, enhanced hepatic cell proliferation, and the development of liver tumors (Casanova et al., 1996; Maronpot et al., 1995; Foley et al., 1993; Kari et al., 1993). The studies provide no clear evidence for a sustained liver cell proliferation response to dichloromethane that can be linked to the development of dichloromethane-induced liver tumors. Additionally, a few studies have examined if dichloromethane-induced liver tumors are the result of proto-oncogene activation and tumor suppressor gene inactivation (Maronpot et al., 1995; Devereux et al., 1993; Hegi et al., 1993).

Kari et al. (1993) (also summarized by Maronpot et al. [1995]) reported data from six groups of 68 female B6C3F₁ mice exposed to six “stop-exposure” protocols of differing durations and sequences, with each exposure concentration standardized at 2,000 ppm for 6 hours/day, 5 days/week. The six stop-exposure protocols were 26 weeks of exposure followed by 78 weeks without exposure, 78 weeks without exposure followed by 26 weeks of exposure, 52 weeks without exposure followed by 52 weeks with exposure, 52 weeks of exposure followed by 52 weeks without exposure, 78 weeks of exposure followed by 26 weeks without exposure, and 26 weeks without exposure followed by 78 weeks of exposure. A control group (no exposure, 104 weeks duration) and a maximum exposure (104 weeks duration) group were also included. Exposure for 26 weeks did not result in an increased incidence of liver tumors (adenomas or carcinomas). Respective percentages of animals with liver tumors were 27 (18/67), 40 (27/67), and 34% (23/67) for the controls, early 26-week exposure, and late 26-week exposure groups, respectively. Exposure to 2,000 ppm for 52 weeks (followed by no exposure until 104 weeks), 78 weeks (either early or late exposure periods), or 104 weeks produced increased incidences of mice with liver tumors ($p < 0.05$), but this increase was not seen in the 52-week late exposure group. Respective percentages of animals with liver tumors (adenomas and carcinomas combined) were 44 (28/64), 31 (21/67), 62 (42/68), 48 (32/67), and 69% (47/68) for the 52 (early exposure), 52 (late exposure), 78 (early exposure), 78 (late exposure), and 104 week exposure periods, respectively. With the 78 week exposures, the difference in the liver tumor incidence between the early and late exposure periods was statistically significant ($p < 0.01$). A greater increase in multiplicity of liver tumors was also seen with the early 78-week exposure period. These data suggest that 52 weeks of exposure was required to increase the incidence of liver tumors in mice, that early exposure was more effective than late exposure, and that the increased risk continued after cessation of exposure.

Histopathologic examination of liver tissue at interim killings at eight time periods (13, 26, 52, 68, 75, 78, 83, or 91 weeks) of exposure to 2,000 ppm (n = 20 mice per killing) found no evidence of nonneoplastic cytotoxicity that preceded the appearance of proliferative neoplastic liver lesions. Incidences of mice with liver adenomas or carcinomas were elevated (between 40 and 60%) at five of the six interim killings after 52 weeks. The incidence rates at each time period were 0/20 (0%) at 13 weeks, 1/20 (5%) at 26 weeks, 8/20 (40%) at 52 weeks, 4/26 (15%) at 68 weeks, 13/20 (65%) at 75 weeks, 12/19 (63%) at 78 weeks, 8/20 (40%) at 83 weeks, and 20/30 (66%) at 91 weeks. The collected liver lesion data identify no exposure-related increased incidence of nonneoplastic liver lesions that could be temporally linked to liver tumor development. Liver tumors first appeared at about the same time in control and exposed animals (52 weeks).

Foley et al. (1993) examined indices of cell proliferation in livers of female B6C3F₁ mice exposed to 1,000, 2,000, 4,000, or 8,000 ppm dichloromethane (6 hours/day, 5 days/week) for 1, 2, 3, or 4 weeks or to 2,000 ppm for 13, 26, 52, or 78 weeks but found no evidence for sustained cell proliferation with prolonged exposure to dichloromethane. To label liver cells in S phase, tritiated thymidine (1- to 4-week exposure protocols) or bromodeoxyuridine (13- to 78-week protocols) was administered subcutaneously via an osmotic mini-pump for 6 days prior to killing. Labeled hepatocytes in liver sections (from 10 mice in each exposure/duration group) were counted to assess the number of cells in S-phase per 1,000 cells. S-phase labeling indices in livers of exposed mice at most killings were equivalent to or less than those in control mice. A transient increase in S-phase labeling index of about two- to fivefold over controls was observed at the 2-week killing of mice exposed to 1,000, 4,000, or 8,000 ppm. Because of the transient nature and small magnitude of the response, it is not expected to be of significance to the promotion of liver tumors in chronically exposed mice. Foley et al. (1993) also compared cell proliferation labeling indices in foci of cellular alteration and nonaffected liver regions in control and exposed mice but found no significant difference between control and exposed mice. S-phase labeling was accomplished by immunohistologic staining for proliferating cell nuclear antigen in liver sections from 24 control mice and 15 exposed mice, with livers showing foci of cellular alteration. In both control and exposed livers, the labeling index was about four- to fivefold higher in foci of cellular alteration than in surrounding unaffected liver tissue.

In mice exposed to 2,000 ppm for 13–78 weeks, relative liver weights were statistically significantly elevated compared with controls; about 10% increased at 13 and 26 weeks and about 30–40% increased at 52 and 78 weeks. Histologic changes in liver sections of 2,000 ppm mice exposed for 13–78 weeks were restricted to hepatocellular hypertrophy (observed at all killing intervals) and preneoplastic (foci of cellular alteration) and neoplastic (adenoma and carcinoma) lesions. No signs of liver tissue degeneration were found. Adenoma and focus of alteration were first detected at 26 weeks (2/10 versus 0/10 in controls). At 52 weeks, 4/10 exposed mice had proliferative lesions (one focus, one adenoma, and two carcinomas),

compared with 1/10 in controls (one adenoma). At 78 weeks, 7/10 exposed mice had proliferative lesions (two foci, three adenomas, six carcinomas) compared with 1/10 in controls (one adenoma). In summary, the results indicate that inhalation exposure to 2,000 ppm dichloromethane produced an increase incidence of liver tumors in female B6C3F₁ mice. No evidence was found for sustained cell proliferation or liver tissue degeneration in response to dichloromethane exposure, but exposure was associated with relative liver weight increases and hepatocellular hypertrophy.

Casanova et al. (1996) found no clear evidence of increased cell proliferation in the livers of male B6C3F₁ mice exposed to dichloromethane concentrations >1,500 ppm 6 hours/day for 3 days. Three or four groups of three mice were exposed to 146, 498, 1,553, or 3,923 ppm unlabeled dichloromethane for 2 days and then exposed to [¹⁴C]-labeled dichloromethane for 6 hours on the third day. Radiolabel incorporated into liver DNA deoxyribonucleosides was measured as an index of cell proliferation. Radiolabel incorporated into liver DNA deoxyribonucleosides increased approximately fivefold from 146 to 1,553 ppm, but further increases were not apparent at 3,923 ppm. (In contrast, as described in Section 4.5.3, radiolabel incorporation into lung DNA deoxyribonucleosides displayed a 27-fold increase over this concentration range.) The small magnitude of the increase in radiolabel incorporation into liver DNA deoxyribonucleosides with increasing exposure concentration suggests that little if any enhanced cell proliferation occurred in the liver in response to dichloromethane exposure.

Devereux et al. (1993) (also reported in Maronpot et al. [1995]) analyzed liver tumors in female B6C3F₁ mice for the presence of activated *H-ras* oncogenes. Fifty dichloromethane-induced and 49 spontaneous liver tumors were screened for *H-ras* mutations. There was a relatively high frequency of activated *H-ras* among the nonexposed B6C3F₁ mice: 67% of the spontaneous tumors and 76% of the dichloromethane-induced tumors contained mutations in the *H-ras* gene. Overall, the mutation profile of the dichloromethane-induced tumors did not significantly differ from the spontaneous tumors.

Similarly, Hegi et al. (1993) analyzed the liver tumors from female B6C3F₁ mice for inactivation of the tumor suppressor genes, *p53* and *Rb-1*. Half of the liver tumors used for this study had an activated *H-ras* oncogene. Twenty liver tumors (15 carcinomas and 5 adenomas) were screened for loss of heterozygosity (LOH) on chromosome 11 and 14, which is associated with malignant conversion of the *p53* gene (chromosome 11) and the *Rb-1* gene (chromosome 14). Only one tumor out of 20 contained a LOH at chromosome 14, and no dichloromethane-induced liver tumors contained a LOH at chromosome 11.

Liver metabolic studies. As described in detail in Section 3.3, GST-T1 enzymatic activity and distribution is variable among species, and there is also considerable intraspecies variability among humans. In summary, Reitz et al. (1989) demonstrated a greater metabolic activity with respect to dichloromethane in livers of B6C3F₁ mice compared with F344 rats, Syrian golden

hamsters, and humans. The rates of in vitro metabolism by the GST pathway were about 4-, 12-, and 20-fold greater in B6C3F₁ mouse liver samples than in F344 rat, human, and Syrian golden hamster samples, respectively (Reitz et al., 1989). A more recent study characterized the dichloromethane metabolic capacity specifically of hepatic GST-T1 (Thier et al., 1998). Enzymatic activities of GST-T1 in liver from F344 rats, B6C3F₁ mice, Syrian golden hamsters, and humans with three different GST-T1 phenotypes (nonconjugators, low conjugators, high conjugators) showed the following order with dichloromethane as a substrate: mouse >> rat > human high conjugators > human low conjugators > hamster > human nonconjugators.

4.5.3. Mechanistic Studies of Lung Effects

The finding of increased lung tumors in B6C3F₁ mice exposed to dichloromethane (Menear et al., 1988; NTP, 1986) has stimulated a number of studies designed to examine the mechanism for dichloromethane-induced lung tumors in this animal. The lung tumor mechanism studies were conducted with B6C3F₁ mice, and the frequency of lung tumors in control animals was very low. Time-course studies for lung tumor development were conducted, and it was found that the onset of lung tumor development was much shorter than liver tumors (Kari et al., 1993) (reported in Maronpot et al., 1995). As a result, it is hypothesized that a potential common mechanism independent of liver metabolism is producing tumors in the lung. As with the liver tumors, there were no significant increases in mutations for the *K-ras* oncogene (Devereux et al., 1993) or the *p53* and *Rb-1* tumor suppressor genes (Hegi et al., 1993). Additionally, the Clara cells, which are nonciliary secretory cells found in the primary bronchioles of the lung, are selectively targeted after dichloromethane exposure. Acute dichloromethane exposure produces Clara cell vacuolization, which is not sustained with long-term exposure (Foster et al., 1992). There is a correlation between the acute effects on the Clara cell and the lung tumors from chronic exposure to dichloromethane (Kari et al., 1993). However, the exact mechanism for producing these lung effects is not completely understood at this point. Provided below is a summary of the studies examining the potential mechanisms for producing lung tumors resulting from dichloromethane exposure.

Lung tumor characterization studies. Kari et al. (1993) (also summarized in Maronpot et al. [1995]) demonstrated that only 26 weeks of exposure to 2,000 ppm was necessary to produce significantly increased incidences of female B6C3F₁ mice with lung tumors. In the six “stop-exposure” protocol experiments (26 weeks exposure followed by 78 weeks without exposure, 78 weeks without exposure followed by 26 weeks exposure, 52 weeks without exposure followed by 52 weeks with exposure, 52 weeks exposure followed by 52 weeks without exposure, 78 weeks exposure followed by 26 weeks without exposure, and 26 weeks without exposure followed by 78 weeks exposure), early but not late exposure for 26 or 52 weeks resulted in an increased incidence of animals with lung tumors (adenoma or carcinomas).

Respective percentages of animals with lung tumors were 7.5 (5/67), 31 (21/68), 4 (3/67), 63 (40/63), and 15% (10/67) for the controls, early 26-, late 26-, early 52-, and late 52-week exposure groups, respectively. With the 78-week exposures, both the early and late exposure regimens produced an increased incidence of lung tumors compared with controls (56 [38/68] and 19% [13/68], respectively), compared with the incidence of 63% (42/67) seen in the group exposed for the full 104 weeks. Thus a plateauing of risk was seen, with similar incidence rates seen with the early 52-week, early 78-week, and 104-week exposure periods. The difference in the lung tumor incidence between the early and late exposure periods of similar duration was statistically significant ($p < 0.01$) for the 26-, 52-, and 78-week duration protocols. A greater increase in multiplicity of lung tumors was also seen with the early 78-week exposure period. As with the liver tumor data from the same series of experiments, these data suggest that early exposure was more effective than late exposure and that the increased risk continued after cessation of exposure.

Histopathologic examination of lung tissue from mice killed at 13, 26, 52, 68, 75, 78, 83, or 91 weeks of exposure to 2,000 ppm ($n = 20$ mice per killing) found no evidence of nonneoplastic cytotoxicity that preceded the appearance of proliferative neoplastic lung lesions. In contrast, incidences of mice with lung adenomas or carcinomas (combined) were elevated at interim killings ≥ 52 weeks; incidences for the interim killings of mice exposed to 2,000 ppm (6 hours/day, 5 days/week) between 13 and 91 weeks were 0/20 (0%) at 13 weeks, 0/20 (0%) at 26 weeks, 6/20 (30%) at 52 weeks, 6/26 (23%) at 68 weeks, 8/20 (40%) at 75 weeks, 9/19 (47%) at 78 weeks, 10/20 (50%) at 83 weeks, and 14/30 (47%) at 91 weeks. Lung hyperplasia was found at an increased incidence only at 91 weeks, well after the 26- and 52-week periods that induced increased incidences of mice with lung tumors.

Kanno et al. (1993) found no evidence for histologic changes or increased cell proliferation in lung tissue of female B6C3F₁ mice exposed to 2,000 or 8,000 ppm dichloromethane for 1, 2, 3, or 4 weeks compared with control mice, or in mice exposed to 2,000 ppm for 13 or 26 weeks. Osmotic mini-pumps were used to deliver tritiated thymidine and label cells undergoing replicative DNA synthesis over 6-day periods before killing. Labeled cells undergoing rapid DNA synthesis and cell proliferation were assessed in sections of proximal and terminal bronchioles and alveoli of lungs from groups of 5 mice exposed for 1–4 weeks or 10 mice exposed for 13 or 26 weeks. There were no exposure-related histopathologic or labeling index changes in the alveoli, but lower labeling indices were found in the bronchiolar epithelium of exposed mice compared with controls.

The combined results from the Kari et al. (1993) and Kanno et al. (1993) studies are consistent with the hypothesis that dichloromethane-induced lung tumors in B6C3F₁ mice are not preceded by overt cytotoxicity, enhanced and sustained cell proliferation, or hyperplasia in the lung. Two other studies (Casanova et al., 1996; Foster et al., 1992), however, have reported evidence for enhanced cell proliferation in lungs of B6C3F₁ mice exposed for acute durations to

airborne dichloromethane. Only one of these studies (Foster et al., 1992), however, looked for sustained cell proliferation in the lung with prolonged exposure. In agreement with the results from Kanno et al. (1993), no evidence was found for sustained cell proliferation in lungs with prolonged exposure to dichloromethane at concentrations demonstrated to induce lung tumors in mice.

Casanova et al. (1996) detected evidence of increased cell proliferation in the lungs of male B6C3F₁ mice exposed to dichloromethane concentrations >1,500 ppm 6 hours/day for 3 days. Three or four groups of three mice were exposed to 146, 498, 1,553, or 3,923 ppm unlabeled dichloromethane for 2 days and then exposed to [¹⁴C]-labeled dichloromethane for 6 hours on the third day. Radiolabel incorporated into lung DNA deoxyribonucleosides (after removal of DNA-protein cross-links containing radiolabeled formaldehyde) was measured as an index of cell proliferation. Radiolabel incorporation into lung DNA deoxyribonucleosides increased with increasing exposure concentration, with the amount increasing by about 27-fold between 146 and 3,923 ppm. In hamsters that did not develop tumors in response to chronic inhalation exposure to 3,500 ppm dichloromethane (Burek et al., 1984), no evidence for enhanced radiolabel incorporation into lung DNA deoxyribonucleosides was found following acute exposure (Casanova et al., 1996).

Devereux et al. (1993) (also summarized in Maronpot et al. [1995]) analyzed lung tumors in female B6C3F₁ mice for the presence of activated *K-ras* oncogenes. Fifty-four dichloromethane-induced and 17 spontaneous lung tumors (7 from the NTP [1986] study and 10 from a study in C57BL/6 × C34F1 mice reported by Candrian et al. [1991]) were screened for *K-ras* mutations. Twenty percent of the dichloromethane-induced tumors and 24% of the spontaneous tumors contained mutations in the *K-ras* gene. Devereux et al. (1993) stated that there may be a significant difference in the incidence of *K-ras* activation between spontaneous and dichloromethane-induced tumors. However, the small number of the spontaneous tumors that were available for the study limits the conclusions that can be made from the results.

Hegi et al. (1993) analyzed the lung tumors from female B6C3F₁ mice for inactivation of the tumor suppressor genes, *p53* and *Rb-1*. Forty-nine dichloromethane-induced lung carcinomas, five lung adenomas, and seven spontaneous lung carcinomas were screened for LOH on mouse chromosome 11 and 14, which is associated with malignant conversion of the *p53* gene (chromosome 11) and the *Rb-1* gene (chromosome 14). Fourteen percent (n = 7) of the dichloromethane-induced lung carcinomas exhibited LOH at chromosome 11. No *p53* mutations were detected in the seven spontaneous lung tumors or the five dichloromethane-induced lung adenomas. Only three dichloromethane-induced tumors exhibited LOH at chromosome 14. The authors noted that inactivation of the *p53* and *Rb-1* tumor suppressor genes infrequently occur in lung and liver tumors.

Clara cell studies. Foster et al. (1992) found enhanced cell proliferation in bronchiolar cells and, to a lesser degree, alveolar cells in the lungs of male B6C3F₁ mice exposed for acute durations (2, 5, 8, or 9 days) to 4,000 ppm dichloromethane (6 hours/day, 5 days/week), but the response was less distinct after subchronic durations of exposure (89, 92, or 93 days). To measure cell proliferation, mice (n = 5 per exposure-duration group) were given subcutaneous doses of tritiated thymidine for 5 consecutive days prior to killing. Labeled cells in bronchiolar or alveolar epithelium in lung sections were counted to assess the number of cells in S phase per 1,000 cells. Counts of bronchiolar epithelium cells in S phase in exposed mice sacrificed on days 2, 5, 8, and 9 were approximately 2-, 15-, 3-, and 5-fold higher, respectively, than those of unexposed mice at day 0 of the experiment. In exposed mice sacrificed on days 89, 92, and 93, less than twofold increases in bronchiolar epithelium cell labeling were observed. Increased cell labeling was found in alveolar epithelium only on day 8 (about seven- to eightfold increase) and day 9 (about fourfold increase). Vacuolation of the Clara cells of the bronchiolar epithelium was observed on day 2 (scored as ++, majority of cells affected), day 9 (+++, virtually all the cells affected), and day 44 (+, moderate effect to cells) but was not apparent on days 5, 8, 40, 43, 89, 92, or 93. No hyperplasia of the bronchiolar epithelium or changes to Type II alveolar epithelial cells were observed in the lungs of any of the exposed mice at any time point. The appearance and disappearance of the Clara cell vacuolation was generally correlated with the appearance and disappearance of enhanced cell proliferation in the bronchiolar epithelium; enhanced cell proliferation was observed on days 2, 5, 8, and 9 (along with appearance of Clara cell vacuolation on days 2 and 9) but was not observed on days 89, 92, and 93 when Clara cell lesions also were not observed. This suggests that cell proliferation was enhanced in response to Clara cell damage but was not sustained with repeated exposure to dichloromethane.

Currently, a mechanistic connection has not been established between the acute effects of dichloromethane on Clara cells in the bronchiolar epithelium and the development of lung tumors in B6C3F₁ mice exposed by inhalation to concentrations $\geq 2,000$ ppm dichloromethane for 2 years (NTP, 1986) or for 26 weeks followed by no exposure through 2 years (Maronpot et al., 1995; Kari et al., 1993). There appears to be a concordance between exposure concentrations inducing acute Clara cell vacuolation ($\geq 2,000$ ppm) and those inducing lung tumors ($\geq 2,000$ ppm). However, transient acute Clara cell vacuolation does not appear to progress to necrosis or lead to sustained cell proliferation (which could promote the growth of tumor-initiated cells) and appears to be dependent on CYP metabolism of dichloromethane (see the following paragraphs discussing pertinent findings reported by Foster et al. [1994, 1992]). In contrast, there is consistent and specific evidence for an association between the formation of DNA-reactive GST-pathway metabolites and the formation of lung and liver tumors from inhalation exposure (see Sections 4.5.2 and 4.7.3).

Foster et al. (1992) noted that the appearance and disappearance of Clara cell vacuolation in mouse lungs showed concordance with temporal patterns for immunologic staining for

CYP2B1 and 2B2 levels in lung sections. A similar temporal pattern was reported for CYP2B1 and 2B2 monooxygenase activities (ethoxycoumarin O-dealkylation or aldrin epoxidation) assayed in lung microsomes. When there was a marked decrease in CYP2B1 and 2B2 staining (e.g., on day 5) or monooxygenase activities, the lesion was not present. Similarly, the appearance of the lesion was preceded (the day before) by the recovery of monooxygenase activities or immunologic staining close to control levels. These patterns suggested to Foster et al. (1992) that Clara cells may have developed tolerance to dichloromethane due to inactivation of a CYP isozyme.

In subsequent studies, increased percentages of vacuolated bronchiolar epithelium cells were noted in mice exposed to 2,000 ppm ($26.3 \pm 6.7\%$) or 4,000 ppm ($64.8 \pm 12.8\%$), but vacuolated cells were not observed in bronchiolar epithelium of lung sections from control mice or mice exposed to 125, 250, 500, or 1,000 ppm (Foster et al., 1994). Pretreatment with the CYP inhibitor, piperonyl butoxide, counteracted the 2,000 ppm effect ($2.4 \pm 3.6\%$ vacuolated cells), whereas GSH-depleted mice showed no statistically significant change in percentage of vacuolated cells ($32.7 \pm 16.9\%$) compared with the mean percentage in mice exposed to 2,000 ppm without pretreatment. No consistent, statistically significant, exposure-related changes were found in cytosolic GST metabolic activities (with dichloromethane as substrate) or microsomal CYP monooxygenase activities (ethoxycoumarin O-dealkylation), but mean cytosolic levels of nonprotein sulfhydryl compounds were elevated in lungs of mice exposed to 1,000 and 2,000 ppm (134.6 ± 17.1 and 146.4 ± 6.7 nmol/mg protein, respectively) compared with control levels (109.5 ± 7.6 nmol/mg protein). Increased cell proliferation was found in cultured Clara cells isolated from 4,000 ppm mice compared with nonexposed mice; respective values for percentage of cells in S phase were 18.97 ± 1.18 and $2.02 \pm 0.86\%$ (Foster et al., 1994).

Results from the studies by Foster et al. (1994, 1992) indicate that 6-hour exposures of B6C3F₁ mice to dichloromethane concentrations $\geq 2,000$ ppm caused transient Clara cell vacuolation in the bronchiolar epithelium, which was not consistently observed following repeated exposures. With repeated exposure to 4,000 ppm, the Clara cell vacuolation did not progress to necrosis, and no hyperplasia of the bronchiolar epithelium was found after up to 13 weeks of exposure. The transient Clara cell vacuolation was decreased by CYP inhibition with piperonyl butoxide and was unaffected by GSH depletion, indicating that a CYP metabolite was involved. Clara cell vacuolation was not found after five consecutive, daily 6-hour exposures to 4,000 ppm but reappeared after 2 days without exposure followed by two additional consecutive, daily exposures (day 9). With repeated exposure, the lesion was detected at a diminished severity on day 44 (but was not found on day 40 or 43) and on day 93 (but was not found on day 89 or 92). The temporal pattern of Clara cell vacuolation with repeated exposure was reflected in the occurrence of transiently decreased CYP metabolic activity after the appearance of vacuolation. Foster et al. (1994, 1992) proposed that the diminishment of severity

or the disappearance of the Clara cell vacuolation with repeated exposure was due to the development of a tolerance to dichloromethane, linked with a decrease of CYP metabolism of dichloromethane.

4.5.4. Mechanistic Studies of Neurological Effects

Several neurobehavioral studies (see Section 4.4.3 for a complete summary) have demonstrated that dichloromethane exposure results in decreased spontaneous motor activity with pronounced lethargy at high concentrations ($\geq 1,000$ ppm). These effects, combined with the observation that dichloromethane impairs learning and memory (Alexeef and Kilgore, 1983) and affects production of evoked responses to sensory stimuli (Herr and Boyes, 1997; Rebert et al., 1989), indicate that dichloromethane produces significant neurological effects. The mechanisms behind producing these changes have been examined by measuring changes in neurotransmitter levels and changes in neurotransmitter localization. Specific brain regions (e.g., hippocampus, caudate nucleus, cerebellum) were analyzed to determine if dichloromethane-induced behavioral effects, such as learning and memory (hippocampus, caudate nucleus) and movement (cerebellum), are resulting from pathological changes in these regions. Changes in neurotransmitter levels were also monitored to see if there was any correlation in behavior and neurochemical changes. Summaries of these studies are provided below. It is not yet known if dichloromethane directly interacts with neuronal receptors, as has been demonstrated for toluene and ethanol, two other solvents with neurobehavioral and neurophysiological profiles that are similar to those of dichloromethane (for a review see Bowen et al. [2006]).

Kanada et al. (1994) examined the effect of dichloromethane on acetylcholine and catecholamines (dopamine, norepinephrine, serotonin) and their metabolites in the midbrain, hypothalamus, hippocampus, and medulla from male Sprague-Dawley rats (four to five per group). The rats were sacrificed 2 hours after a single gavage dose of 0 or 534 mg/kg of undiluted dichloromethane. Administration of dichloromethane significantly increased the concentration of acetylcholine in the hippocampus and increased dopamine and serotonin levels in the medulla. Dichloromethane decreased norepinephrine levels in the midbrain, and hypothalamus and serotonin levels were decreased in the hypothalamus. There was a trend toward decreased dopamine in the hypothalamus, but the variability between the animals was so high that the effect was not significant. The authors speculated that increased acetylcholine release from dichloromethane administration may be due to decreased acetylcholine release from the nerve terminals. It is unclear as to how these neurochemical changes could be correlated to the neurobehavioral changes observed after dichloromethane exposure.

In a 2-week exposure study, male Wistar rats were exposed to dichloromethane at 500 or 1,000 ppm (6 hours/day, 5 days/week for 1 or 2 weeks) or 1,000 ppm TWA (1 hour at 100 ppm, 1 hour peak at 2,800 ppm, 1 hour at 100 ppm, repeated immediately, 5 days/week for 1 or 2 weeks) (Savolainen et al., 1981). Brains were removed from rats at the end of the study and

analyzed. The 1,000 ppm TWA group displayed increases in cerebral RNA. Other changes noted for this group in the cerebrum included significant increases in NADPH-diaphorase and succinate dehydrogenase activity. These changes suggest increased neural activity to possibly offset the overall inhibitory effect of dichloromethane in the CNS. It could also possibly explain why lethargic effects decrease with continued dichloromethane exposure, and this result demonstrates a neuroprotective mechanism resulting from dichloromethane exposure. After a 7-day withdrawal, RNA levels in the cerebrum were significantly lower in the 1,000 ppm group. Succinate dehydrogenase levels remained lowered in the 1,000 ppm TWA group after the 7-day exposure-free period.

Changes in brain catecholamine levels after a subacute exposure to dichloromethane were evaluated using male Sprague-Dawley rats (Fuxe et al., 1984). Rats were exposed to 70, 300, and 1,000 ppm dichloromethane 6 hours/day for 3 consecutive days. At all exposures, there was a significant decrease of catecholamine concentrations in the posterior periventricular region of the hypothalamus. The impact of dichloromethane was also evaluated on the hypothalamic-pituitary gonadal axis. The hypothalamus regulates secretion of reproductive hormones such as follicle-stimulating hormone and luteinizing hormone. The levels of the hormone release were not significantly changed with dichloromethane exposure. In the caudate nucleus, which is involved in memory processes, the catecholamine level initially increased (at 70 ppm) and then was lower (1,000 ppm) in comparison to the control. The study demonstrates significant changes in catecholamine levels in the hypothalamus and caudate nucleus. Catecholamine level changes in the hypothalamus did not have any significant effect on hormonal release, and decreased catecholamine levels in the caudate nucleus at higher exposures may lead to memory and learning impairment.

A series of studies were conducted in male and female Mongolian gerbils exposed continuously to ≥ 210 ppm dichloromethane for 3 months, followed by a 4-month exposure-free period (Karlsson et al., 1987; Briving et al., 1986; Rosengren et al., 1986). Decreased DNA concentrations were noted in the hippocampus at both the 210 and 350 ppm exposures. At 350 ppm, there was also decreased DNA concentration in the cerebellar hemispheres, indicating a decreased cell density in these regions probably due to cell loss (Rosengren et al., 1986). These findings indicate that the cerebellum, which is the section of the brain that regulates motor control, is a target for dichloromethane. In the same study, increased astroglial proteins were found in the frontal and sensory motor cerebral cortex, which directly correlated to the astrogliosis that was observed in those areas. Up-regulation of these astroglial proteins is a good indicator of neuronal injury (Rosengren et al., 1986).

Karlsson et al. (1987) measured DNA concentrations in different regions of the gerbil brain. The total brain protein concentration per wet weight was not significantly different between dichloromethane-exposed and control animals. However, DNA concentrations per wet weight were significantly decreased in the hippocampus after dichloromethane exposure. No

other examined regions demonstrated significant changes in DNA concentrations after dichloromethane exposure. Therefore, this result indicates that the hippocampus, which plays a role in the formation of new memories, is another target for dichloromethane in the CNS. This selective DNA concentration decrease observed in the hippocampus is a sign of neurotoxicity as noted by the authors and may possibly explain why some studies have noted memory and learning deficits with dichloromethane exposure.

At a 210 ppm exposure, Briving et al. (1986) observed that dichloromethane decreased glutamate, γ -aminobutyric acid, and phosphoethanolamine levels in the frontal cortex, while glutamate and γ -aminobutyric acid were increased in the posterior cerebellar vermis. Increased levels of glutamate in the posterior cerebellar vermis could reflect an activation of astrocytic glia, since glutamine synthetase is localized exclusively in astrocytes.

4.6. SYNTHESIS OF MAJOR NONCANCER EFFECTS

4.6.1. Oral

4.6.1.1. *Summary of Human Data*

Information on noncancer effects in humans exposed orally to dichloromethane are restricted to case reports of neurological impairment, liver and kidney effects (as severe as organ failure), and gastrointestinal irritation in individuals who ingested amounts ranging from about 25 to 300 mL (Chang et al., 1999; Hughes and Tracey, 1993). Neurological effects with these individuals consisted of general CNS depressive symptoms, such as drowsiness, confusion, headache, and dizziness. Hemoglobinuria has been noted as a kidney effect associated with ingestions.

4.6.1.2. *Summary of Animal Data*

Acute oral or intraperitoneal administration of dichloromethane in animals has resulted in several significant effects. General activity and function were affected as evidenced by decreased neuromuscular activity (Moser et al., 1995). Additionally, decreased sensorimotor function was detected through measurement of evoked potentials (Herr and Boyes, 1997) and by using the FOB (Moser et al., 1995). Neurochemical changes (e.g., acetylcholine, dopamine, norepinephrine, serotonin) were detected 2 hours after oral dosage of dichloromethane within specific parts of the brain. It should be noted that all the acute effects that were observed after oral or intraperitoneal administration occurred within 5 hours after dosage. No other significant organ effects were noted after a single acute oral exposure, but in oral pharmacokinetic studies, it is known that dichloromethane is primarily distributed to the liver, lungs, and kidneys (Angelo et al., 1986a).

Results from short-term, subchronic, and chronic oral toxicity studies in laboratory animals are summarized in Table 4-35. The data indicate that rats may be more sensitive than mice to nonneoplastic liver effects from orally administered dichloromethane, as evidenced by

lower NOAELs and LOAELs with more severe liver effects in rats. The most frequently observed liver effect was hepatocyte vacuolation, seen with drinking water exposure (90 days) in F344 rats at ≥ 166 mg/kg-day and B6C3F₁ mice at 586 mg/kg-day (Kirschman et al., 1986) and with gavage exposure (14 days) in CD-1 mice at 333 mg/kg-day (Condie et al., 1983). Hepatocyte degeneration or necrosis was observed in female F344 rats exposed by drinking water for 90 days to 1,469 mg/kg-day (Kirschman et al., 1986) and in female F344 rats exposed by gavage for 14 days to 337 mg/kg-day (Berman et al., 1995) but was not seen in a 90-day drinking water study in B6C3F₁ mice exposed to doses as high as 2,030 mg/kg-day (Kirschman et al., 1986). In the chronic-duration (2-year) study, liver effects were described as foci and areas of alteration in F344 rats exposed to drinking water doses between 50 and 250 mg/kg-day; an increased incidence of fatty changes in the liver was also noted but the incidence was not provided (Serota et al., 1986a). These effects were considered to be nonneoplastic for several reasons. Serota et al. (1986b) observed a dose-related increased incidence of 0, 65, 92, 97, 98, and 100% in male rats and 51, 41, 73, 89, 91, and 85% in female rats for the 0, 5, 50, 125, 250, and 250 mg/kg-day with recovery groups, respectively. Evidence for liver tumors has been reported in female rats only. Specifically, evidence for liver tumors in rats includes a small number of hepatocellular carcinomas observed in female rats at 50 and 250 mg/kg-day, which reached statistical significance (for trend and for individual pairwise comparisons) only with the combined grouping of neoplastic nodules and hepatocellular carcinomas. In male rats, only one hepatocellular carcinoma was observed in all of the exposure groups (compared to 4 in the controls), and the incidence of neoplastic nodules and hepatocellular carcinomas was higher in controls (16%) than in any exposure group (16, 3, 0, 6, 5, and 13% for the 0, 5, 50, 125, 250 mg/kg-day, and 250 mg/kg-day with recovery groups, respectively). The authors (Serota et al., 1986a) did not elaborate on the characterization of the altered foci. However, the characterization of altered foci could range from a focal change in fat distribution (nonneoplastic effect) to enzyme altered foci which are generally considered a precursor to tumor formation (Goodman et al., 1994). Serota et al. (1986a) reported an increased incidence of fatty change in the liver at doses of ≥ 50 mg/kg-day, but the incidence was not reported. In addition, a 90-day study (Kirschman et al., 1986) demonstrated that increased fatty deposits were present in the hepatocyte vacuoles. Therefore, the altered foci (i.e., change in fat distribution) observed by Serota et al. (1986b) may represent a precursor to fatty liver changes which is considered a nonneoplastic effect. Taken together, the data support the conclusion that the altered foci were nonneoplastic.

Table 4-35. NOAELs and LOAELs in selected animal studies involving oral exposure to dichloromethane for short-term, subchronic, or chronic durations

Type of effect and exposure, reference	Species and exposure details	Results	NOAEL	LOAEL
			(mg/kg-d)	
Hepatic, 14-d gavage				
Berman et al. (1995)	F344 rat, female, 8/dose group 0, 34, 101, 337, 1,012 mg/kg-d	Hepatocyte necrosis	101	337
Condie et al. (1983)	CD-1 mouse, male, 5/group for histological examinations; 8/group for blood urea nitrogen, serum creatinine, and serum glutamate-pyruvate transaminase; 0, 133, 333, 665 mg/kg-d	Hepatocyte vacuolation (minimal to mild in 3/5)	133	333
Hepatic, 90-d drinking water				
Kirschman et al. (1986)	F344 rat, male and female; 15/sex/group; males 0, 166, 420, 1,200 mg/kg-d females 0, 209, 607, 1,469 mg/kg-d	Hepatic vacuolation (generalized, centrilobular, or periportal, at lowest dose, in 10/15 males and 13/15 females compared with 1/15 males and 6/15 females in controls)	Not identified	166
Kirschman et al. (1986)	B6C3F ₁ mouse, male and female, males 0, 226, 587, 1,911 mg/kg-d females 0, 231, 586, 2,030 mg/kg-d	Hepatic vacuolation (increased severity of centrilobular fatty change in mid- and high-dose groups compared with controls)	231	586
Hepatic, 104-wk drinking water				
Serota et al. (1986a)	F344 rat, male and female, 0, 5, 50, 125, 250 mg/kg-d	Liver foci/areas of alteration (considered nonneoplastic histologic changes); fatty liver changes also seen at same doses but incidence data not reported; no evidence that increased altered foci progresses to liver tumor formation	5	50
Serota et al. (1986b); Hazleton Laboratories (1983)	B6C3F ₁ mouse, male and female, 0, 60, 125, 185, 250 mg/kg-d	Some evidence of fatty liver; marginal increase in the Oil Red-O-positive material in the liver	185	250
Neurologic, 14 d				
Moser et al. (1995)	F344 rat, female, 0, 34, 101, 337, 1,012 mg/kg-d	FOB 24 hrs postexposure: altered autonomic, neuromuscular, and sensorimotor and excitability measures	101	337

(Table 4-35 continues on next page)

Table 4-35. NOAELs and LOAELs in selected animal studies involving oral exposure to dichloromethane for short-term, subchronic, or chronic durations

Type of effect and exposure, reference	Species and exposure details	Results	NOAEL	LOAEL
			(mg/kg-d)	
Reproductive				
General Electric Company (1976)	Charles River CD rat, male and female, gavage for 90 d before mating (10 d between last exposure and mating period); 0, 25, 75, 225 mg/kg-d; F1 offspring received same treatment as parents for 90 d	Reproductive performance of F0 and histologic examination of tissues from F1 offspring	225	Not identified
Raje et al. (1988)	Swiss-Webster mouse, male, 0, 250, 500 mg/kg (subcutaneous injection), 3 × per wk, 4 wks prior to mating with nonexposed females (1 wk between last exposure and mating period)	No statistically significant effects on testes, number of litters, live fetuses/litter, percent dead fetuses/litter, percent resorbed/litter, or fertility index	500	Not identified
Developmental				
Narotsky and Kavlock (1995)	F344 rat, pregnant female, gavage on GDs 6–19; 0, 338, 450 mg/kg-d	Maternal: weight gain depression	338	450
		Fetal: no effects on pup survival, resorptions, pup weight	450	Not identified

The NOAEL and LOAEL, 101 and 337 mg/kg-day, respectively, for altered neurological functions in female F344 rats (as reported by Moser et al. [1995]) were identical to those reported by Berman et al. (1995) for hepatocyte necrosis in female F344 rats. In the 90-day (Kirschman et al., 1986) and 104-week (Serota et al., 1986a, b) drinking water studies, no obvious clinical signs of neurological impairment were observed in rats or mice at exposure levels that induced liver effects (see Table 4-35), but these studies did not include standardized neurological testing batteries.

Results from a limited number of studies do not provide evidence for effects on reproductive or developmental endpoints (Table 4-35). No effects on pup survival, resorptions, or pup weight were found following exposure of pregnant F344 rats to doses as high as 450 mg/kg-day on GDs 6–19, a dose that depressed maternal weight gain (Narotsky and Kavlock, 1995), and no effects on reproductive performance endpoints (fertility index, number of pups per litter, pup survival) were found in Charles River CD rats exposed for 90 days before mating to doses as high as 225 mg/kg-day. There are no oral exposure studies focusing on neurobehavioral effects or other developmental outcomes.

4.6.2. Inhalation

4.6.2.1. Summary of Human Data

As discussed in Section 4.1.2, acute inhalation exposure of humans to dichloromethane has been associated with cardiovascular impairments due to decreased oxygen availability from COHb formation and neurological impairment from interaction of dichloromethane with nervous system membranes. Results from studies of acutely exposed human subjects indicate that acute neurobehavioral deficits measured, for example, by psychomotor tasks, tests of hand-eye coordination, visual evoked response changes, and auditory vigilance, may occur at concentrations >200 ppm with 4–8 hours of exposure (Bos et al., 2006; ACGIH, 2001; ATSDR, 2000; Cherry et al., 1983; Putz et al., 1979; Gamberale et al., 1975; Winneke, 1974).

The clinical and workplace studies of noncancer health effects of chronic dichloromethane exposure have examined markers of disease and specific clinical endpoints relating to cardiac, neurological disease, hepatic function, and reproductive health. As summarized in Section 4.1.2.9, the limited available data do not provide evidence of cardiac damage related to dichloromethane exposure in occupationally exposed workers (Hearne and Pifer, 1999; Tomenson et al., 1997; Gibbs et al., 1996; Lanes et al., 1993; Ott et al., 1983d; Cherry et al., 1981). Relatively little is known about the long-term neurological effects of chronic exposures, although there are studies that provide some evidence of an increased prevalence of neurological symptoms among workers with average exposures of 75–100 ppm (Cherry et al., 1981), long-term effects on some neurological measures (i.e., possible detriments in attention and reaction time in complex tasks) in workers whose past exposures were in the 100–200 ppm range (Lash et al., 1991), and an increased risk of suicide in worker cohort studies

(Hearne and Pifer, 1999; Gibbs, 1992). Given the suggestions from these studies and their limitations (particularly with respect to sample size and power considerations), the statement that there are no long-term neurological effects of chronic exposures to dichloromethane cannot be made with confidence. With respect to markers of hepatic damage, three studies measured serum enzyme and bilirubin levels in workers exposed to dichloromethane (Soden, 1993; Kolodner et al., 1990; Ott et al., 1983c). There is some evidence of increasing levels of serum bilirubin with increasing dichloromethane exposure (Kolodner et al., 1990; Ott et al., 1983c), but there are no consistent patterns with respect to the other hepatic enzymes examined (serum γ -glutamyl transferase, serum AST, serum ALT). Thus, these studies do not provide clear evidence of hepatic damage in dichloromethane exposed workers to the extent that this damage could be detected by these serologic measures.

Only limited and somewhat indirect evidence pertaining to immune-related effects of dichloromethane in humans is available. No risk of the broad category of infection- and parasite-related mortality was reported by Hearne and Pifer (1999), but there was some evidence of an increased risk of influenza and pneumonia-related mortality at two cellulose triacetate fiber production work sites in Maryland and South Carolina (Gibbs, 1992).

Few studies have been conducted pertaining to reproductive effects (i.e., spontaneous abortion, low birth weight, or oligospermia) of dichloromethane exposure from workplace settings (Wells et al., 1989; Kelly, 1988; Taskinen et al., 1986) or environmental settings (Bell et al., 1991). Of these, the data pertaining to spontaneous abortion provide the strongest evidence of an adverse effect of dichloromethane exposure. The limitations of the only study pertaining to this outcome (Taskinen et al., 1986), however, do not allow firm conclusions to be made regarding dichloromethane and risk of spontaneous abortion in humans.

4.6.2.2. *Summary of Animal Studies*

Acute and short-term (up to 7 days) inhalational exposure to dichloromethane has resulted in neurological and hepatocellular changes. Several neurological-mediated parameters were reported, including decreased activity (Kjellstrand et al., 1985; Weinstein et al., 1972; Heppel and Neal, 1944), impairment of learning and memory (Alexeef and Kilgore, 1983), and changes in responses to sensory stimuli (Rebert et al., 1989). Although learning and memory properties were impaired in one acute exposure (47,000 ppm until loss of righting reflex), it should be noted that this effect has not been characterized by using other learning and memory tasks nor any other exposure paradigms. In a 3-day exposure to dichloromethane (70, 300, or 1,000 ppm 6 hours/day), it was found that in the rat brain, there were changes in catecholamine (dopamine, serotonin, norepinephrine) in the hypothalamus and caudate nucleus (Fuxe et al., 1984). The catecholamine level changes did not affect hormonal release which is a primary function of the hypothalamus.

Another acute exposure study examined immunological response as measured by increased streptococcal pneumonia-related mortality and decreased bactericidal activity of pulmonary macrophages in CD-1 mice following a single 3-hour exposure to dichloromethane at 100 ppm (Aranyi et al., 1986). No effects were seen at 50 ppm. A 4-week inhalation exposure to 5,000 ppm dichloromethane in rats did not result in changes in immune response as measured by the sheep red blood cell assay (Warbrick et al., 2003). These studies suggest a localized, portal-of-entry effect within the lung without evidence of systemic immunosuppression.

Mouse hepatocytes showed balloon degeneration (dissociation of polyribosomes and swelling of rough endoplasmic reticulum) within 12 hours of exposure to 5,000 ppm (Weinstein et al., 1972). A subacute exposure in Wistar rats to 500 ppm dichloromethane 6 hours/day for 6 days resulted in increased hemochrome content in liver microsomal CYP (Savolainen et al., 1977).

Results pertaining to liver, lung, and neurological effects from longer (>7 days) subchronic and chronic inhalation toxicity studies in laboratory animals are summarized in Table 4-36; reproductive and developmental studies are summarized in Table 4-37.

Table 4-36. NOAELs and LOAELs in animal studies involving inhalation exposure to dichloromethane for subchronic or chronic durations, hepatic, pulmonary, and neurologic effects

Type of effect and exposure period, reference	Species and exposure details	Results	NOAEL	LOAEL
			ppm	
Hepatic, subchronic (13–14 wks)				
Haun et al. (1971)	Beagle, female (n = 8); Rhesus monkeys, female (n = 4); Sprague-Dawley rat, male (n = 20); ICR mouse, female (n = 380) 0, 1,000, 5,000 ppm (continuous exposure; 14 wks)	Fatty liver at 1,000 ppm in dogs: “borderline” liver changes in monkey at 5,000 ppm; mottled liver changes in rats at 5,000 ppm; hepatocytes degeneration at 5,000 ppm in mice, no information about liver effects in mice at 1,000 ppm; decreased movement and lethargy at 1,000 ppm in dogs, mice, and monkey	Not identified (dog) Not identified (monkey) 1,000 (rat) Not identified (Mouse)	1,000 (dog) 5,000 (monkey) 5,000 (rat) 5,000 (mouse)
Haun et al. (1972)	Beagle (n = 16); Rhesus monkey (n = 4); Sprague-Dawley rat (n = 20), ICR mouse (n = 20) 0, 25, 100 ppm (continuous exposure; 14 wks)	Increased hepatic cytoplasmic vacuolation and decreased CYP levels in liver microsomes in mice at 100 ppm; increased fatty liver content at 25, 100 ppm in rats	100 (dog) 100 (monkey) Not identified (rat) 25 (mouse)	Not identified (dog) Not identified (monkey) 25 (rat) 100 (mouse)
Leuscher et al. (1984)	Sprague-Dawley rat, male and female, (20/sex/group) - 0, 1,000 ppm (6 hrs/d; 90 d); Beagle, male and female (3/sex/group) - 0, 5,000 ppm	No liver effects noted	1,000 (rat) 5,000 (dog)	Not identified (rat) Not identified (dog)
NTP (1986)	F344/N rat, male and female (10/sex/group) 0, 525, 1,050, 2,100, 4,200, 8,400 ppm (6 hrs/d, 5 d/wk, 13 wks)	Decreased lipid:liver weight ratios at 4,200 (females); 8,400 (males); decreased BW by 23 and 11% in males and females at 8,400 ppm compared with controls; one male and one female died at 8,400 ppm before the end of the study	4,200	8,400
NTP (1986)	B6C3F ₁ mouse, male and female (10/sex/group) 0, 525, 1,050, 2,100, 4,200, 8,400 ppm (6 hrs/d, 5 d/wk, 13 wks)	Hepatocyte centrilobular degeneration at 4,200 females) and 8,400 (males); decreased lipid:liver weight ratios at 8,400 (females); at 8,400 ppm, 4/10 males and 2/10 females died before end of study	2,100	4,200

(Table 4-36 continues on next page)

Table 4-36. NOAELs and LOAELs in animal studies involving inhalation exposure to dichloromethane for subchronic or chronic durations, hepatic, pulmonary, and neurologic effects

Type of effect and exposure period, reference	Species and exposure details	Results	NOAEL	LOAEL
			ppm	
Hepatic, 2 yrs (6 hrs/d, 5 d/wk)				
Mennear et al. (1988); NTP (1986)	F344/N rat, male and female 0, 1,000, 2,000, 4,000 ppm	Hepatocyte vacuolation and necrosis Hemosiderosis in liver Renal tubular degeneration	Not identified Not identified 1,000	1,000 1,000 2,000
Mennear et al. (1988); NTP (1986)	B6C3F ₁ mouse, male and female 0, 2,000, 4,000 ppm	Hepatocyte degeneration Renal tubule casts	Not identified Not identified	2,000 2,000
Burek et al. (1984)	Syrian golden hamster, male and female 0, 500, 1,500, 3,500 ppm	No effects on histologic, clinical chemistry, urinalytic, and hematologic variables no obvious clinical signs of toxicity	3,500	Not identified
Burek et al. (1984)	Sprague-Dawley rat, male and female 0, 500, 1,500, 3,500 ppm	Hepatocyte vacuolation (males and females) Hepatocyte necrosis (males only), no obvious clinical signs of toxicity)	Not identified 500	500 1,500
Nitschke et al. (1988a)	Sprague-Dawley rat, male and female 0, 50, 200, 500 ppm	Hepatocyte vacuolation significantly increased in females; nonsignificant increase in males at 500 ppm (31% in controls and 40% in 500 ppm group)	200	500
Pulmonary, 13 wks (6 hrs/d, 5 d/wk)				
NTP (1986)	F344 rat, male and female 0, 525, 1,050, 2,100, 4,200, 8,400 ppm	Foreign body pneumonia	4,200	8,400
NTP (1986)	B6C3F ₁ mouse, male and female (10/sex/group) 0, 525, 1,050, 2,100, 4,200, 8,400 ppm	No nonneoplastic pulmonary lesions	8,400	Not identified
Foster et al. (1992)	B6C3F ₁ mouse, male and female 0, 4,000 ppm	Clara cell vacuolation	Not identified	4,000
Neurological, 14 d				
Savolainen et al. (1981)	Wistar rat, male 500, 1,000, 1,000 TWA (100 + 2,800 1-hr peaks ^a) ppm (6 hrs/d, 5 d/wk, 2 wks)	Increased RNA in cerebrum at 1,000 ppm; increased enzymatic activities ^b in cerebrum and cerebellum at 1,000 ppm TWA	500	1,000 for brain RNA concentration; 1,000 TWA for brain enzymatic activity

(Table 4-36 continues on next page)

Table 4-36. NOAELs and LOAELs in animal studies involving inhalation exposure to dichloromethane for subchronic or chronic durations, hepatic, pulmonary, and neurologic effects

Type of effect and exposure period, reference	Species and exposure details	Results	NOAEL	LOAEL
			ppm	
Neurological, 13–14 wks				
Mattsson et al. (1990)	F344 rat, male and female 0, 50, 200, 2,000 ppm (6 hrs/d, 5 d/wk)	No exposure-related effects on an observational battery, hind-limb grip strength, a battery of evoked potentials, or histology of brain, spinal cord, peripheral nerves; measured 64 hrs postexposure	2,000	Not identified
Haun et al. (1971)	Beagle dog (female); Rhesus monkey (female); Sprague-Dawley rat (male); ICR mouse (female) 0, 1,000, 5,000 ppm (continuous exposure)	Clinical signs (incoordination, lethargy) of CNS depression most evident in dogs	Not identified (dog) Not identified (monkey) 1,000 (rat) Not identified (mouse)	1,000 (dog) 1,000 (monkey) 5,000 (rat) 1,000 (mouse)
Karlsson et al. (1987) Briving et al. (1986) Rosengren et al. (1986)	Mongolian gerbil, male and female 210, 350, 700 ppm (continuous exposure, followed by 4 mo exposure-free period)	Astrogliosis in frontal and sensory motor cerebral cortex suggested by increases in astroglial proteins; cell loss in cerebellar regions; decreased DNA in hippocampus; neurochemical changes observed at all exposures	Not identified	210
Thomas et al. (1972)	ICR mouse, female 0, 25, 100 ppm, continuous	Increased spontaneous activity observed at 25 ppm but not 100 ppm	Not identified	25
CoHb, 13–14 wks				
Haun et al. (1972)	Beagle (n = 16); Rhesus monkey (n = 4); Sprague-Dawley rat (n = 20), ICR mouse (n = 20) 0, 25, 100 ppm (continuous exposure; 14 wks)	CoHb levels significantly higher at 25, 100 ppm for monkeys and 100 ppm for beagles	Not identified	25

(Table 4-36 continues on next page)

Table 4-36. NOAELs and LOAELs in animal studies involving inhalation exposure to dichloromethane for subchronic or chronic durations, hepatic, pulmonary, and neurologic effects

Type of effect and exposure period, reference	Species and exposure details	Results	NOAEL	LOAEL
			ppm	
COHb, 2 yrs (6 hrs/d, 5 d/wk)				
Burek et al. (1984)	Syrian golden hamster, male and female 0, 500, 1,500, 3,500 ppm	About 30% COHb in each exposed group	Not identified	500
Burek et al. (1984)	Sprague-Dawley rat, male and female 0, 500, 1,500, 3,500 ppm	About 12–14% COHb in each exposed group	Not identified	500
Nitschke et al. (1988a)	Sprague-Dawley rat, male and female 0, 50, 200, 500 ppm	COHb values at 2 yrs: about 2, 7, 13, 14%	Not identified	500

^aEquivalent to 1,000 ppm TWA.

^bDecreased GSH, γ -aminobutyric acid, and phosphoethanolamine in frontal cortex; GSH and γ -aminobutyric acid increased in posterior cerebellar vermis.

Table 4-37. NOAELs and LOAELs in selected animal studies involving inhalation exposure to dichloromethane, reproductive and developmental effects

Type of effect and exposure period, reference	Species and exposure details	Results	NOAEL	LOAEL
			ppm	
Reproductive				
Nitschke et al. (1988b)	F344 rat, male and female, F0: 6 hr/d, 5 d/wk for 14 wk before mating and GDs 0 to 21; F1: 6 hr/d, 5 d/wk, beginning PND 4 for 17 wk before mating; 0, 100, 500, 1,500 ppm	No statistically significant effects on fertility or litter size, neonatal survival, growth rates, or histopathologic lesions in F1 or F2 weanlings	1,500	Not identified
Menear et al. (1988); NTP (1986)	B6C3F ₁ mouse; 0, 2,000 or 4,000 ppm, 6 hrs/d, 5 d/wk for 2 yrs	Testicular atrophy Ovarian atrophy (considered secondary to hepatic effects)	2,000 Not identified	4,000 2,000
Raje et al. (1988)	Swiss-Webster mouse, male, 2 hr/d, 5 d/wk for 6 wk before mating with nonexposed females; 0, 100, 150, 200 ppm	No statistically significant effects on testes, number of litters, live fetuses/litter, percent dead fetuses/litter, percent resorbed/litter Fertility index was lower in 150 and 200 ppm groups (80%) compared with controls and 100 ppm groups (95%) (statistical significance depends on test used)	200 100	Not identified 150
Developmental				
Schwetz et al. (1975)	Swiss-Webster mouse, pregnant female, 7 hr/d, GDs 6–15; 0, 1,250 ppm	Maternal effects: 9–10% COHb; increased absolute, not relative, liver weight, increased maternal weight (11–15%).	Not identified	1,250
		Fetal effects: increased litters with extra center of ossification in sternum	1,250	Not identified
Schwetz et al. (1975)	Sprague-Dawley rat, pregnant female, 7 hr/d, GDs 6–15; 0, 1,250 ppm	Maternal: 9–10% COHb; increased absolute, not relative, liver weight	Not identified	1,250
		Fetal: increased incidence of delayed ossification of sternebrae	1,250	Not identified
Other developmental				
Bornschein et al. (1980); Hardin and Manson (1980)	Long-Evans rat, female, 6 hr/d for 12–14 d before breeding and GDs 1–17; 6 hr/d on GDs 1–15; 0, 4,500 ppm	Maternal (both protocols): increased absolute and relative liver weight (~10%)	Not identified	4,500
		Fetal/offspring: decreased fetal BW (~10%); changed behavioral habituation to novel environments; no changes in gross, skeletal, or soft-tissue anomalies	Not identified	4,500

Hepatic centrilobular degeneration was observed in several studies containing different species and inhalational exposures. This effect was observed in guinea pigs exposed to 5,000 ppm (7 hours/day) for 6 months (Heppel et al., 1944). Monkeys, rats, and mice continuously exposed (24 hours/day) to 5,000 ppm dichloromethane for 14 weeks also had increased centrilobular degeneration (Haun et al., 1972, 1971). This effect was also observed at lower exposures when mice were exposed to 4,200 ppm 6 hours/day for 13 weeks (NTP, 1986) and in dogs exposed to 1,000 ppm 24 hours/day for up to 14 weeks (Haun et al., 1972, 1971).

Increased incidences of histologic hepatic lesions were not found in F344 rats exposed to 4,200 or 8,400 ppm 6 hours/day for 13 weeks (NTP, 1986) or in Sprague-Dawley rats exposed to 10,000 ppm 6 hours/day for 90 days (Leuschner et al., 1984). Hepatic lesions were also not observed in beagle dogs exposed to 5,000 ppm 6 hours/day for 90 days (Leuschner et al., 1984) or in dogs, monkeys, rats, and mice exposed to 25 or 100 ppm 24 hours/day for up to 14 weeks (Haun et al., 1972). Heppel et al. (1944) also demonstrated absence of hepatic lesions in unspecified strains of monkeys, rabbits, and rats exposed to 10,000 ppm 4 hours/day for up to 8 weeks and in unspecified strains of dogs, rabbits, and rats exposed to 5,000 ppm 7 hours/day for up to 6 months.

Gross neurological impairments were observed in several laboratory species with repeated exposure to 10,000 ppm 4 hours/day for 8 weeks (Heppel et al., 1944) or to 1,000 or 5,000 ppm 24 hours/day for 14 weeks (Haun et al., 1972, 1971). Dogs exposed to 5,000 ppm 6 hours/day for 90 days showed slight sedation during exposures, but Sprague-Dawley rats exposed to 10,000 ppm for 90 days did not (Leuschner et al., 1984). In F344 rats exposed to concentrations up to 2,000 ppm 6 hours/day for 13 weeks, no effects were observed on an observational battery, hind-limb grip strength, a battery of evoked potentials, or histology of the brain, spinal cord, or peripheral nerves; these tests were conducted beginning ≥ 65 hours after the last exposure (Mattsson et al., 1990).

Exposure-related nonneoplastic effects on the lungs reported in the subchronic studies were restricted to foreign body pneumonia in rats exposed to 8,400 ppm 6 hours/day for 13 weeks (NTP, 1986), Clara cell vacuolation in mice exposed to 4,000 ppm 6 hours/day for 13 weeks (Foster et al., 1992), and pulmonary congestion in guinea pigs exposed to 5,000 ppm 7 hours/day for 6 months (Heppel et al., 1944).

The chronic duration inhalation studies were conducted at lower exposure levels than the short-term and subchronic studies and provide results indicating that the liver is the most sensitive target for noncancer toxicity in rats and mice (Table 4-36). Life-time exposure was associated with hepatocyte vacuolation and necrosis in F344 rats exposed to 1,000 ppm 6 hours/day (Mennear et al., 1988; NTP, 1986), hepatocyte vacuolation in Sprague-Dawley rats exposed to 500 ppm 6 hours/day (Nitschke et al., 1988a; Burek et al., 1984), and hepatocyte degeneration in B6C3F₁ mice exposed to 2,000 ppm 6 hours/day (lower concentrations were not tested in mice) (Mennear et al., 1988; NTP, 1986). As shown in Tables 4-36 and 4-37, other

effects observed include renal tubular degenerations in F344 rats and B6C3F₁ mice at 2,000 ppm, testicular atrophy in B6C3F₁ mice at 4,000 ppm, and ovarian atrophy in B6C3F₁ mice at 2,000 ppm (considered secondary to hepatic effects). No exposure-related increased incidences of nonneoplastic lung lesions were found in any of the chronic studies (Table 4-36).

In comparison to rats and mice, Syrian golden hamsters are less sensitive to the chronic inhalation toxicity of dichloromethane. No exposure-related changes were found in comprehensive sets of histologic, clinical chemistry, urinalytic, and hematologic variables measured in hamsters exposed for 2 years to 500, 1,500, or 3,500 ppm for 6 hours/day, with the exception that mean COHb percentages were about 30% in each of these groups compared with a mean value of about 3% for the controls (Burek et al., 1984).

The reproductive and developmental studies are limiting in terms of the exposure regimen used, with two of the developmental studies using only a single, relatively high daily exposure over the gestational period (1,250 ppm, GDs 6–15 in Schwetz et al. [1975] and 4,500 ppm, GDs 1–17 in Bornschein [1980] and Hardin and Manson [1980]). No significant effects on reproductive performance variables were found in a two-generation reproduction assay with F344 rats exposed to concentrations as high as 1,500 ppm (Nitschke et al., 1988b). No effects on most of the measures of reproductive performance were observed in male mice exposed to 200 ppm 2 hours/day for 6 weeks before mating to nonexposed females. Fertility index was reduced in the 150 and 200 ppm groups, but the statistical significance of this effect varied considerably depending on the statistical test used in this analysis (Raje et al., 1988). No adverse effects on fetal development were found following exposure of pregnant Swiss-Webster mice or Sprague-Dawley rats to 1,250 ppm 6 hours/day on GDs 6–15 (Schwetz et al., 1975). Following exposure of female Long-Evans rats to 4,500 ppm (6 hours/day) for 14 days before breeding plus during gestation or during gestation alone, a 10% decrease in fetal BW and changed behavioral habituation of the offspring to novel environments were seen (Bornschein et al., 1980; Hardin and Manson, 1980). No exposure-related changes in gross, skeletal, or soft-tissue anomalies were found.

4.6.3. Mode-of-Action Information

4.6.3.1. *Mode of Action for Nonneoplastic Liver Effects*

Studies of chronically exposed rats, both by the oral route and the inhalation route, identified liver changes as the most sensitive exposure-related noncancer effect associated with exposure to dichloromethane (Tables 4-35 to 4-37). The liver changes included increased incidence of liver foci/areas of alteration and hepatocyte vacuolation in rats and degenerative liver effects in rats, guinea pigs, monkeys, and mice.

The mode of action by which dichloromethane induces these nonneoplastic hepatic effects is unknown. The determination of whether or not these effects are due to the parent material, metabolites of the CYP2E1 pathway, metabolites of the GST pathway, or some

combination of parent material and metabolites has not been elucidated. The available data indicate that rats may be more sensitive than mice to the noncancer hepatotoxicity, but a mechanistic explanation of this possible interspecies difference is not currently available.

4.6.3.2. Mode of Action for Nonneoplastic Lung Effects

Single 6-hour inhalation exposures to concentrations $\geq 2,000$ ppm dichloromethane produced a transient vacuolation of Clara cells in the bronchiolar epithelium of B6C3F₁ mice. Vacuolization of the Clara cells disappeared or was diminished with repeated exposure and was correlated with subsequent transient diminishment of CYP metabolic activity. CYP inhibition with piperonyl butoxide counteracted the vacuolation observed in the Clara cells (Foster et al., 1994, 1992). With repeated exposure to 4,000 ppm (up to 13 weeks), the Clara cell vacuolation did not appear to progress to necrosis, and no hyperplasia of the bronchiolar epithelium was found. Foster et al. (1994, 1992) proposed that the diminished severity or disappearance of Clara cell vacuolation with repeated exposure was due to the development of tolerance to dichloromethane, linked with a transient decrease of CYP metabolism of dichloromethane. The available data suggests that CYP metabolism of dichloromethane may be involved in the mode of action for the acute effects of dichloromethane on the bronchiolar epithelium of mice.

Mode of action research attention on lung effects from chronic exposure to dichloromethane has focused on neoplastic effect; nonneoplastic lung effects have received relatively little attention. No exposure-related increased incidences of nonneoplastic lung lesions (including epithelial hyperplasia) were found in any of the chronic studies listed in Table 4-36, but chronic inhalation exposure of B6C3F₁ mice to concentrations $\geq 2,000$ ppm has consistently been shown to induce lung tumors in several studies (Kari et al., 1993; NTP, 1986). In a study that included interim sacrifices at 13, 26, 52, 68, 75, 78, 83, and 91 weeks of B6C3F₁ mice exposed to 2,000 ppm, hyperplasia of lung epithelium (the only nonneoplastic lung lesion found) was found in only three of the eight interim sacrifices (68, 78, and 91 weeks) and was only statistically significantly elevated at 91 weeks (5/30 versus 0/15 in controls) (Kari et al., 1993).

4.6.3.3. Mode of Action for Neurological Effects

Results from studies of acutely exposed human subjects indicate that mild neurobehavioral deficits may occur at air concentrations >200 ppm with 4–8 hours of exposure (Bos et al., 2006; ACGIH, 2001; ATSDR, 2000; Cherry et al., 1983; Putz et al., 1979; Gamberale et al., 1975; Winneke, 1974). Acute high-dose exposures also resulted in gross neurological impairments in several laboratory species (Haun et al., 1972, 1971; Heppel et al., 1944). Exposure of F344 rats to concentrations up to 2,000 ppm 6 hours/day for 13 weeks produced no effects on an observational battery, hind-limb grip strength, a battery of evoked potentials, or histology of the brain, spinal cord, or peripheral nerves (Mattsson et al., 1990). However, oral

exposures have been shown to alter autonomic, neuromuscular, and sensorimotor functions in F344 rats exposed to gavage doses ≥ 337 mg/kg-day for 14 days (Moser et al., 1995).

Dichloromethane may be metabolized by the CYP2E1 enzyme to CO (Guengerich, 1997; Hashmi et al., 1994; Gargas et al., 1986). Many of the acute human exposure studies evaluated if CO was the primary metabolite responsible for producing the CNS depressant effects observed during dichloromethane exposure. Overall, at lower exposures and acute durations, it appears that CO is the primary mediator of the neurobehavioral effects. Putz et al. (1979) demonstrated that similar neurobehavioral deficits were present when an equivalent COHb blood level (and CO exposure) was achieved between CO and dichloromethane exposures. Incidentally, after a longer duration, neurobehavioral deficits are more pronounced with dichloromethane exposure in comparison to CO exposure alone. This additional increase in the CNS depressive effects is most likely due to the saturation of the CYP2E1 metabolic pathway. In humans, saturation of the CYP2E1 metabolic pathway was seen at approximately 400–500 ppm after a 1-hour exposure (Ott et al., 1983e). CYP2E1 pathway saturation with dichloromethane has also been noted in hamsters (Burek et al., 1984) and in rats (Nitschke et al., 1988a; McKenna et al., 1982). It is highly probable that initially, CYP2E1 is metabolizing dichloromethane to CO, which results in the neurological effects. However, at higher concentrations (>500 ppm) and for longer durations (>3 hours), the CYP2E1 pathway is most likely saturated. As a result, either the remaining dichloromethane could be metabolized by the GST pathway or the parent compound is producing the effects itself.

Once the CYP2E1 enzyme is saturated, it is unknown whether dichloromethane or a GST-T1 pathway metabolite (e.g., formaldehyde) mediates the resulting neurological effects. Based on the available literature on other solvents, such as toluene and perchloroethylene (for a review see Bowen et al. [2006]), it can be hypothesized that once the CYP2E1 enzyme is saturated, dichloromethane or a GST metabolite may interact directly with excitatory and inhibitory receptors such as the NMDA, GABA, dopamine, and serotonin receptors, among other targets, to produce the resulting neurobehavioral effects. This hypothesis is supported by the evidence that changes in relation to dichloromethane exposures in glutamate, GABA, dopamine, serotonin, acetylcholine, and other neurotransmitters are found in the brain (Kanada et al., 1994; Briving et al., 1986; Fuxe et al., 1984). Additionally, several neurobehavioral effects such as decreased spontaneous motor activity, deficits in learning and memory, and deficits in FOB parameters are similar to other more characterized solvents such as toluene. However, more comprehensive studies specifically designed to determine the mode of action for dichloromethane-induced impairment of neurological functions have not been conducted.

4.6.3.4. Mode of Action for Reproductive and Developmental Effects

No significant effects on reproductive performance variables were found in a two-generation reproduction assay with F344 rats exposed to concentrations as high as 1,500 ppm

(Nitschke et al., 1988b), and no effects were seen on most of the measures of reproductive performance examined in a study of male mice exposed to 200 ppm 2 hours/day for 6 weeks before mating to nonexposed females (Raje et al., 1988). In the mouse study, fertility index (number of females impregnated/total number of females mated \times 100) was reduced in the 150 and 200 ppm groups (Raje et al., 1988), but the statistical significance of this effect varied considerably depending on the statistical test used in the analysis. Mechanistic studies of dichloromethane or its metabolites that would provide mode of action information on reproductive effects in the male are not available.

The mode of action for developmental effects can be hypothesized to involve the CYP2E1 pathway and, specifically, the production of CO. CO is a known developmental neurotoxicant. Demonstrated effects include neurobehavioral deficits and neurochemical changes (Giustino et al., 1999; Cagiano et al., 1998; De Salvia et al., 1995; Fechter, 1987). In addition, placental transfer of dichloromethane has been demonstrated with inhalation exposure (Withey and Karpinski, 1985; Anders and Sunram, 1982). Pups exposed in utero to high concentrations of dichloromethane (4,500 ppm) demonstrated neurobehavioral-related changes in comparison to air-exposed animals (Bornschein et al., 1980). This observed effect coupled with the known developmental neurotoxicological effects produced by CO suggests that the CYP2E1 metabolic pathway is involved in producing observed and suspected neurodevelopmental effects. In humans, CYP2E1 activity in the brain occurs earlier in gestation than it does in the liver, with activity in the brain seen in the first trimester (Johnsrud et al., 2003; Brzezinski et al., 1999). Thus, the direct effects of dichloromethane in fetal circulation, as well as the effects of CO and the effects of the CYP2E1-related metabolism in the fetal liver and the fetal brain, may be relevant to the risk of developmental effects in humans. Mechanistic studies of dichloromethane or its metabolites that would provide mode of action information on other noted developmental effects such as delayed ossification (Schwetz et al., 1975) are not available.

4.6.3.5. *Mode of Action for Immunotoxicity*

Evidence of a localized immunosuppressive effect in the lung resulting from inhalation dichloromethane exposure was seen in an acute exposure (3 hours, 100 ppm) study in CD-1 mice (Aranyi et al., 1986). The lung infectivity assay used in this study examined response to bacterial challenges (i.e., risk of streptococcal-pneumonia-related mortality and clearance of *Klebsiella* bacteria). The innate immune response plays an important role in limiting the initial lung burden of bacteria through the activity of macrophages, neutrophils, and dendritic cells, and alveolar macrophages are particularly important in the response to respiratory infections (Marriott and Dockrell, 2007). The adaptive response develops from several days up to several weeks following infection so that an effective immune response in a lung infectivity assay requires multiple immune mechanisms and, in particular, cooperation of macrophages, neutrophils, and T cells along with the appropriate cytokines (Selgrade and Gilmour, 2006).

Although immunosuppression in the Streptococcal and Klebsiella infectivity models has been reported in the acute exposure scenarios tested in Aranyi et al. (1986), mechanistic studies of dichloromethane or its metabolites that would provide mode of action information on the immune system cells or function have not been performed.

4.7. EVALUATION OF CARCINOGENICITY

4.7.1. Summary of Overall Weight of Evidence

Following U.S. EPA (2005a) *Guidelines for Carcinogen Risk Assessment*, dichloromethane is “likely to be carcinogenic in humans”, based predominantly on evidence of carcinogenicity at two sites in 2-year bioassays in male and female B6C3F₁ mice (liver and lung tumors) with inhalation exposure (NTP, 1986) and at one site in male B6C3F₁ mice (liver tumors) with drinking water exposure (Serota et al., 1986b; Hazleton Laboratories, 1983). The incidence rates for liver tumors in female mice were not presented (Serota et al., 1986b; Hazleton Laboratories, 1983), but it was reported that exposed female mice did not show increased incidences of proliferative hepatocellular lesions. Evidence of a trend for increased risk of liver tumors (described as neoplastic nodule or hepatocellular carcinoma) was seen in female F344 rats exposed via drinking water ($p < 0.01$) (Serota et al., 1986a) or inhalation ($p = 0.08$) (NTP, 1986). However, the potential malignant characterization of the nodules was not described, and the data for hepatocellular carcinomas are much more limited. Additional evidence of the tumorigenic potential of dichloromethane in rats comes from the observation of an increase in benign mammary tumors following inhalation exposure (Nitschke et al., 1988a; Burek et al., 1986b; NTP, 1986). A gavage study in female Sprague-Dawley rats reported an increased incidence of malignant mammary tumors, mainly adenocarcinomas (8, 6, and 18% in the control, 100, and 500 mg/kg dose groups, respectively), but the increase was not statistically significant; data were not provided to allow an analysis that accounts for differing mortality rates (Maltoni et al., 1988). An inhalation study (exposures of 0, 50, 200, and 500 ppm) also reported the presence of another relatively rare tumor in rats, astrocytoma or glioma (mixed glial cell) tumors (Nitschke et al., 1988a). Taken together, the rat data provide supporting evidence of carcinogenicity. Studies in humans found some evidence linking occupational exposure to dichloromethane and increased risk for some specific cancers, including brain cancer (Hearne and Pifer, 1999; Tomenson et al., 1997; Heineman et al., 1994) and liver cancer (Lanes et al., 1993, 1990).

The proposed mode of action for dichloromethane-induced liver tumors is through a mutagenic mode of carcinogenic action. Mode of action data indicate that dichloromethane-induced DNA damage in cancer target tissues of mice involves DNA-reactive metabolites produced via a metabolic pathway initially catalyzed by GST-T1. Evidence of mutagenicity includes in vitro bacterial and mammalian assays as well as in vivo mammalian system assays, although mutational events in critical genes (tumor suppressor genes, oncogenes) leading to

tumor initiation and tumor promotion have not been established. This metabolic pathway has been found in human tissues, albeit at lower activities than in mouse tissues; therefore, the cancer results in animals are considered relevant to humans.

U.S. EPA's *Guidelines for Carcinogen Risk Assessment* (U.S. EPA, 2005a) indicate that for tumors occurring at a site other than the initial point of contact the weight of evidence for carcinogenic potential may apply to all routes of exposure that have not been adequately tested at sufficient doses. An exception occurs when there is convincing toxicokinetic data that absorption does not occur by other routes. For dichloromethane, systemic tumors were observed in mice following inhalation and oral exposure. No animal cancer bioassay data following dermal exposure to dichloroemthane are available. Based on the observance of systemic tumors following oral exposure and inhalation exposure, and in the absence of information to indicate otherwise, it is assumed that an internal dose will be achieved regardless of the route of exposure. Therefore, dichloromethane is "likely to be carcinogenic to humans" by all routes of exposure.

4.7.2. Synthesis of Human, Animal, and Other Supporting Evidence

Section 4.1.3 reviewed the results, strengths, and limitations of epidemiological research of dichloromethane and cancer, including cohort and case-control studies. The available epidemiologic studies provide some evidence of an association between dichloromethane and brain cancer and liver cancer, but the available data are limited.

Two small cohort studies with relatively good exposure metrics and relatively long follow-up periods (mean over 25 years) reported an increased risk of brain cancer, with SMRs of 1.45 (95% CI 0.40–3.72) in Tomenson et al. (1997) and 2.2 (95% CI 0.79–4.69) in Cohort 1 of Hearne and Pifer (1999). Cohort 1 is an inception cohort, following workers from the beginning of employment, which is methodologically more robust than Cohort 2, which only included workers who were working between 1964 and 1970. These observations are supported by the data from a case-control study of brain cancer that reported relatively strong trends ($p < 0.05$) with increasing probability, duration, and intensity measures of exposure but not with a cumulative exposure measure (Heineman et al., 1994). This difference could reflect a relatively more valid measure of relevant exposures in the brain from the intensity measure, as suggested by the study in rats reported by Savolainen et al. (1981) in which dichloromethane levels in the brain were much higher with a higher intensity exposure scenario compared with a constant exposure period with an equivalent TWA (see Section 3.2). The combination of high probability of exposure and long (> 20 years) duration of employment in exposed jobs was strongly associated with brain cancer risk (OR 6.1, 95% CI 1.1-43.8) in the Heineman et al. (1994) study; similar associations were seen with the high intensity in combination with long duration measures. A statistically significant increased incidence of brain or CNS tumors has not been observed in any of the animal cancer bioassays, but a 2-year study using relatively low exposure

levels (0, 50, 200, and 500 ppm) in Sprague-Dawley rats observed a total of six astrocytoma or glioma (mixed glial cell) tumors in the exposed groups (in females, the incidence was 0, 0, 0, and 2 in the 0, 50, 200, and 500 ppm exposure groups, respectively; in males, the incidence was 0, 1, 2, and 1 in the 0, 50, 200, and 500 ppm exposure groups, respectively; sample size of each group was 70 rats). These tumors are exceedingly rare in rats, and there are few examples of statistically significant trends in animal bioassays (Sills et al., 1999). These cancers were not seen in two other studies in rats, both involving higher doses (1,000–4,000 ppm) (NTP, 1986; Burek et al., 1984), or in a high dose (2,000–4,000 ppm) study in mice (NTP, 1986).

With respect to epidemiologic studies of liver and biliary duct cancer, the highest exposure cohort, based in the Rock Hill, South Carolina, triacetate fiber production plant, suggested an increased risk of liver cancer with an SMR of 2.98 (95% CI 0.81–7.63) in the latest study update (Lanes et al., 1993). This observation was based on four cases; an earlier analysis in this cohort reported an SMR of 5.75 (95% CI 1.82–13.8), based on these same four cases but with a shorter follow-up period (and thus a lower number of expected cases) (Lanes et al., 1990). No other cohort study has reported an increased risk of liver cancer mortality, although it should be noted that there is no other inception cohort study of a population with exposure levels similar to those of the Rock Hill plant, and no data from a case-control study of liver cancer are available pertaining to dichloromethane exposure.

The primary limitation of all of the available dichloromethane cohort studies is the limited statistical power for the estimation of effects relating to relatively rare cancers (such as brain cancer, liver cancer, and leukemia). Limitations with respect to studies of other cancers can also be noted. With respect to breast cancer, the only cohort that included a significant percentage of women had limited exposure information (analysis was based on a dichotomous exposure variable) and co-exposure to other solvents (Blair et al., 1998). The only breast cancer case-control study available used death certificate data to classify disease and occupational exposure (Cantor et al., 1995), which is likely to result in significant misclassification; exposure misclassification in particular would be expected to result in an attenuated measure of association (Rothman and Greenland, 1998). No studies of adult leukemia and dichloromethane exposure and only one study of childhood leukemia (acute lymphoblastic leukemia) in relation to maternal occupational dichloromethane exposure were found.

In addition to the epidemiologic studies, several dichloromethane cancer bioassays in animals are available. In the only oral exposure cancer bioassay involving lifetime exposure, increases in incidence of liver adenomas and carcinomas were observed in male but not female B6C3F₁ mice exposed for 2 years (Table 4-38 for males; female data not presented in the summary reports) (Serota et al., 1986b; Hazleton Laboratories, 1983). (The trend *p*-value and pairwise test *p*-values were not given in the Serota et al. [1986b] paper but can be found in the full report [Hazleton Laboratories, 1983]). The authors concluded that these increases were “within the normal fluctuation of this type of tumor incidence,” noting that there was no dose-

related trend and that there were no significant differences comparing the individual dose groups with the combined control group.⁸ Although Serota et al. (1986b) state that a two-tailed significance level of $p = 0.05$ was used for all tests, Hazleton Laboratories (1983) indicated that a correction factor for multiple comparisons was used specifically for the liver cancer data, reducing the nominal p -value from 0.05 to 0.0125; none of these individual group comparisons are statistically significant when a p -value of 0.0125 is used.

Based on the Hazleton Laboratories (1983) statistical analysis, EPA concluded that dichloromethane induced a carcinogenic response in male B6C3F₁ mice as evidenced by a marginally increased trend test ($p = 0.058$) for combined hepatocellular adenomas and carcinomas, and by small but statistically significant ($p < 0.05$) increases in hepatocellular adenomas and carcinomas at dose levels of 125 ($p=0.021$), 185 ($p=0.019$), and 250 mg/kg-day ($p=0.036$). EPA did not consider the use of a multiple comparisons correction factor for the evaluation of the liver tumor data (a primary a priori hypothesis) to be warranted.

With respect to comparisons with historical controls, the incidence in the control groups was almost identical to the mean seen in the historical controls from this laboratory (17.8% based on 354 male B6C3F₁ mice), so there is no indication that the observed trend is being driven by an artificially low rate in controls and no indication that the experimental conditions resulted in a systematic increase in the incidence of hepatocellular adenomas and carcinomas. Although the occurrence of one elevated rate in an exposed group may be within the normal fluctuations of this type of tumor incidence (described for this laboratory as 5–40%, with a mean of 17.8%, based on 354 male controls), the pattern of incidence rates (increased incidence in all four dose groups, with three of these increases significant at a p -value < 0.05) suggest a treatment-related increase.

⁸ Two control groups were used because of the potential for high and erratic liver tumor incidence in B6C3F₁ mice. The incidence of hepatocellular adenomas or carcinomas was 18 and 20% in the two control groups, and the combined group is used for the subsequent analysis because of the improved statistical precision of estimates based on the larger sample size ($n=125$ compared with $n=60$ and 65 for the individual control groups).

Table 4-38. Incidence of liver tumors in male B6C3F₁ mice exposed to dichloromethane in a 2-year oral exposure (drinking water) study^a

Estimated mean intake (mg/kg-d) ^a	0 (Controls)	61	124	177	234	Trend <i>p</i> -value ^c
Number of male mice	125 ^b	200	100	99	125	
	Number of cancers (%)					
Hepatocellular adenoma or carcinoma						
Mortality-adjusted percent ^d	24 (19)	51 (26)	30 (30)	31 (31)	35 (28)	0.058
Mortality-adjusted <i>p</i> -value ^d	(22)	(29) <i>p</i> = 0.071	(34) <i>p</i> = 0.023	(35) <i>p</i> = 0.019	(32) <i>p</i> = 0.036	

^aTarget doses were 60, 125, 185, and 250 mg/kg-d from the lowest dose group (excluding controls) to the highest dose group, respectively.

^bTwo control groups combined (n=60 and 65 in the individual groups). The mortality-adjusted incidence in control groups 1 and 2 were 20 and 23%, respectively. Two additional sets of analyses using the individual control groups were also presented in Hazleton Laboratories (1983).

^cCochran-Armitage trend test (Hazleton Laboratories [1983]).

^dMortality-adjusted percent calculated based on number at risk, using Kaplan-Meier estimation, taking into account mortality losses; *p*-value for comparison with control group using asymptotic normal test (source: Hazleton Laboratories [1983]).

Sources: Serota et al. (1986b); Hazleton Laboratories (1983).

In a similar study in F344 rats (Serota et al., 1986a), no increased incidence of liver tumors was seen in male rats, and the pattern in female rats was characterized by a jagged stepped pattern of increasing incidence of hepatocellular carcinoma or neoplastic nodule (Table 4-39). Information was not provided which would allow characterization of the nodules as benign or malignant. Statistically significant increases in incidences were observed in the 50 and 250 mg/kg-day groups (incidence rates of 0, 3, 10, 3, and 14%, respectively, for the 0, 5, 50, 125, and 250 mg/kg-day groups) and in the group exposed to 250 mg/kg-day for 78 weeks followed by a 26-week period of no exposure (incidence rate 10%). A similar pattern, but with more sparse data, was seen for hepatocellular carcinomas, with two incidences in the 50 mg/kg-day and two in the 250 mg/kg-day groups. The authors concluded that dichloromethane exposure did not result in an increased incidence of liver tumors because the increase was based on a low rate (0%) in the controls and because of a lack of monotonicity.

Table 4-39. Incidences of liver tumors in male and female F344 rats exposed to dichloromethane in drinking water for 2 years

	Target dose (mg/kg-d)						Trend <i>p</i> -value ^b	250 with recovery ^c
	0 ^a (Controls)	5	50	125	250			
Males								
Estimated mean intake (mg/kg-d)	0	6	52	125	235			232
total n	135	85	85	85	85			25
n at terminal kill ^d	76	34	38	35	41			15
Number (%) with neoplastic nodules	9 (12)	1 (3)	0 (0)	2 (6)	1 (2)	Not reported		2 (13)
Number (%) with hepatocellular carcinoma	3 (4)	0 (0)	0 (0)	0 (0)	1 (2)	Not reported		0 (0)
Number (%) with neoplastic nodules and hepatocellular carcinoma	12 (16)	1 (3)	0 (0)	2 (6)	2 (5)	Not reported		2 (13)
Females								
Estimated mean intake (mg/kg-d)	0	6	58	136	263			239
total n	135	85	85	85	85			25
n at terminal kill ^d	67	29	41	38	34			20
Number (%) with neoplastic nodules	0 (0)	1 (3)	2 (5)	1 (3)	3 (9)			2 (10) ^e
Number (%) with hepatocellular carcinoma	0 (0)	0 (0)	2 (5)	0 (0)	2 (6)	Not reported		0 (0)
Number (%) with neoplastic nodules and hepatocellular carcinoma	0 (0)	1 (3)	4 (10) ^e	1 (3)	5 (14) ^e	<i>p</i> < 0.01		2 (10) ^e

^aTwo control groups combined.

^bCochran-Armitage trend test was used for trend test of liver foci/areas of alteration. For tumor mortality-unadjusted analyses, a Cochran-Armitage trend test was used, and for tumor mortality-adjusted analyses, a tumor prevalence analytic method by Dinse and Lagakos (1982) was used. Similar results were seen in these two analyses.

^cRecovery group was exposed for 78 wks and then had a 26-wk period without dichloromethane exposure; n = 17 for neoplastic lesions.

^dExcludes 5, 10, and 20 per group sacrificed at 25, 52, and 78 wks, respectively, and unscheduled deaths, which ranged from 5 to 19 per group.

^eSignificantly (*p* < 0.05) different from controls with Fisher's exact test, mortality-unadjusted and mortality-adjusted analyses.

Source: Serota et al. (1986a).

Another gavage exposure study in Sprague-Dawley rats and in Swiss mice provides limited data concerning cancer incidence because the study was terminated early (at 64 weeks) due to high treatment-related mortality (Maltoni et al., 1988). Exposure groups included controls (olive oil), 100, or 500 mg/kg-day 4–5 days/week. High-dose female rats showed an increased incidence of malignant mammary tumors, mainly adenocarcinomas (8, 6, and 18% in the control, 100, and 500 mg/kg dose groups, respectively), but the increase was not statistically significant. Data were not provided to allow an analysis accounting for differing mortality rates. A dose-

related increase, although not statistically significant, in pulmonary adenomas was observed in male mice (5, 12, and 18% in control, 100, and 500 mg/kg-day groups, respectively). When mortality was taken into account, high-dose male mice that died in the period ranging from 52 to 78 weeks were reported to show a statistically significantly ($p < 0.05$) elevated incidence for pulmonary tumors (1/14, 4/21, and 7/24 in control, 100, and 500 mg/kg-day groups, respectively). Details of this analysis were not provided. EPA applied a Fisher's exact test to these incidences and determined a p -value of 0.11 for the comparison of the 500 mg/kg-day group (7/24) to the controls (1/14).

As discussed in Section 4.2, repeated inhalation exposure to concentrations of 2,000 or 4,000 ppm dichloromethane produced increased incidences of lung and liver tumors in B6C3F₁ mice (Mennear et al., 1988; NTP, 1986). The incidence of mortality-adjusted liver tumors across dose groups of 0, 2,000, and 4,000 ppm increased from 48 to 67 and 93%, respectively, in male mice (trend p -value = 0.013) and from 10 to 48 and 100% in female mice (trend p -values < 0.001) (Table 4-40). For lung tumors, the mortality-adjusted incidence was 12, 74, and 100% in males and 11, 83, and 100% in females in the 0, 2,000, and 4,000 ppm groups, respectively (trend p -values < 0.001). Elevated incidences of lung and liver tumors in B6C3F₁ mice were observed with 52 weeks of exposure to 2,000 ppm, and lung tumors were also elevated by week 104 in mice exposed for only 26 weeks to 2,000 ppm, followed by 78 weeks without exposure (Maronpot et al., 1995; Kari et al., 1993).

Table 4-40. Incidences of selected neoplastic lesions in B6C3F₁ mice exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Sex and neoplastic lesion	Exposure (ppm) ^a									Trend <i>p</i> -value ^d
	0 (Controls)			2,000			4,000			
	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	
Males										
Liver—hepatocellular adenoma or carcinoma	22	(44)	(48)	24	(49)	(67)	33 ^e	(67)	(93)	0.013
Lung—bronchoalveolar adenoma or carcinoma	5	(10)	(12)	27 ^e	(54)	(74)	40 ^e	(80)	(100)	<0.001
Females										
Liver— hepatocellular adenoma or carcinoma	3	(6)	(10)	16 ^e	(33)	(48)	40 ^e	(83)	(100)	<0.001
Lung—bronchoalveolar adenoma or carcinoma	3	(6)	(11)	30 ^e	(63)	(83)	41 ^e	(85)	(100)	<0.001

^a2,000 ppm = 6,947 mg/m³, 4,000 ppm = 13,894 mg/m³.

^bTotal sample size was 50 per sex and dose group. Percentages based on the number of tissues examined microscopically per group; for male mice, 49 livers were examined in the 2,000 and 4,000 ppm groups; for female mice, 48 liver and lungs were examined. For comparison, incidences in historical controls reported in NTP (1986) were 28% for male liver tumors, 31% for male lung tumors, 5% for female liver tumors, and 10% for female lung tumors.

^cMortality-adjusted percentage.

^dLife-table trend test, as reported by NTP (1986).

^eLife-table test comparison dose group with control <0.05, as reported by NTP (1986).

Sources: Mennear et al. (1988); NTP (1986).

Liver tumors are relatively rare in F344 rats, and a moderate trend of increasing incidence of what was described as neoplastic nodules or hepatocellular carcinoma was seen in the females (trend *p*-value = 0.08) but not the males in the NTP (1986) study (Table 4-41). As with the rat oral exposure study by Serota et al. (1986a), these nodules were not characterized as benign or malignant. There was no evidence of an increasing trend in incidence when hepatocellular carcinomas only were considered.

Table 4-41. Incidences of selected neoplastic lesions in F344/N rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Sex and neoplastic lesion	Exposure (ppm) ^a												Trend <i>p</i> -value ^d
	0 (Controls)			1,000			2,000			4,000			
	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	n	(%) ^b	(%) ^c	
Males													
Liver—Neoplastic nodule or hepatocellular carcinoma	2	(4)	(10)	3	(6)	(13)	4	(8)	(19)	1	(2)	(6)	0.55
Liver—hepatocellular carcinoma	2	(4)	(10)	1	(2)	(4)	2	(4)	(10)	1	(2)	(6)	nr
Lung—bronchoalveolar adenoma or carcinoma	1			1	(2)		2	(4)		1	(2)		
Mammary gland													
Adenoma, adenocarcinoma, or carcinoma	0	(0)		0	(0)		0	(0)		1	(2)		
Subcutaneous tissue fibroma or sarcoma	1	(2)	(6)	1	(2)	(6)	2	(4)	(9)	5	(10)	(23)	0.008
Fibroadenoma	0	(0)	(0)	0	(0)	(0)	2	(4)	(12)	1	(2)	(8)	<0.001
Mammary gland or subcutaneous tissue adenoma, fibroadenoma, fibroma, or sarcoma	1	(2)	(6)	1	(2)	(6)	4	(8)	(21)	9 ^d	(18)	(49)	<0.001
Females													
Liver—neoplastic nodule or hepatocellular carcinoma	2	(4)	(7)	1	(2)	(2)	4	(8)	(14)	5	(10)	(20)	0.08
Liver—hepatocellular carcinoma	0	(0)	(0)	0	(0)	(0)	1	(2)	(4)	0	(0)	(0)	nr
Lung—bronchoalveolar adenoma or carcinoma	1	(2)		1	(2)		0	(0)		0	(0)		
Mammary gland													
Adenocarcinoma or carcinoma	1	(2)		2	(4)		2	(4)		0	(0)		
Adenoma, adenocarcinoma, or carcinoma	1	(2)		2	(4)		2	(4)		1	(2)		
Fibroadenoma	5	(10)	(16)	11 ^d	(22)	(41)	13 ^d	(26)	(44)	22 ^d	(44)	(79)	<0.001
Mammary gland adenoma, fibroadenoma, or adenocarcinoma	6	(12)	(18)	13	(26)	(44)	14 ^d	(28)	(45)	23 ^e	(46)	(86)	<0.001

^a1,000 ppm = 3,474 mg/m³, 2,000 ppm = 6,947 mg/m³, 4,000 ppm = 13,894 mg/m³.

^bTotal sample size was 50 per sex and dose group. Percentages based on the number of tissues examined microscopically per group; for male rats, 49 livers were examined in the 2,000 and 4,000 ppm groups; for females, only 48 liver and lungs and 49 mammary glands were microscopically examined in the 2,000 and 4,000 ppm groups. For comparison, incidence in historical controls reported in NTP (1986) were 1% for female liver tumors and 16% for female mammary fibroadenomas.

^cMortality-adjusted percentage.

^dLife-table trend test, as reported by NTP (1986); nr = not reported.

^eLife-table test comparison dose group with control < 0.05, as reported by NTP (1986).

Sources: Mennear et al. (1988); NTP (1986).

Female F344 rats exposed by inhalation to 2,000 or 4,000 ppm showed significantly increased incidences of benign mammary tumors (adenomas or fibroadenomas) (Table 4-41); the number of benign mammary tumors per animal also increased with dichloromethane exposure in studies in Sprague-Dawley rats at levels of 50–500 ppm (Nitschke et al., 1988a) and 500–3,500 ppm (Burek et al., 1984) (Table 4-42). Male rats in two of these studies (Nitschke et al., 1988a; NTP, 1986) also exhibited a low rate of sarcoma or fibrosarcoma in mammary gland or subcutaneous tissue around the mammary gland.

Table 4-42. Incidences of mammary gland tumors in two studies of male and female Sprague-Dawley rats exposed to dichloromethane by inhalation (6 hours/day, 5 days/week) for 2 years

Study, lesion	0 (Controls)	Exposure (ppm) ^a						
		50	200	500	Late 500 ^b	Early 500 ^b	1,500	3,500
Nitschke et al. (1988a)								
Males—n per group	57	65	59	64	c	c	c	c
Number (%) with:								
Mammary gland tumors								
Adenocarcinoma or carcinoma	0 (0)	0 (0)	0 (0)	0 (0)				
Fibroadenoma	2 (4)	0 (0)	2 (3)	2 (3)				
Fibroma	6 (11)	1 (6)	6 (11)	10 (16)				
Fibrosarcoma	0 (0)	1 (6)	1 (6)	0 (0)				
Undifferentiated sarcoma	0 (0)	2 (4)	0 (0)	0 (0)				
Fibroma, fibrosarcoma, or undifferentiated sarcoma ^d	6 (11)	4 (6)	7 (12)	10 (16)				
Females—n per group	69	69	69	69	25	25	c	c
Number (%) with:								
Mammary gland tumors								
Adenocarcinoma or carcinoma	6 (9)	5 (7)	4 (6)	4 (6)	3 (12)	2 (8)		
Adenoma	1 (1)	1 (1)	2 (3)	1 (1)	2 (8)	0 (0)		
Fibroadenoma	51 (74)	57 (83)	60 (87)	55 (80)	22 (88)	23 (92)		
Fibroma	0 (0)	1 (1)	0 (0)	1 (1)	1 (4)	1 (1)		
Fibrosarcoma	1 (1)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)		
Number with benign tumors ^e	52 (74)	58 (83)	61(87) ^f	55 (79)	23 (92)	23 (92)		
Number of benign tumors per tumor-bearing rat ^e	2.0	2.3	2.2	2.7	2.2	2.6		
Burek et al. (1984)								
Males—n per group	92	c	c	95	c	c	96	97
Number (%) with benign tumors	7 (8)			3 (3)			7 (7)	14 (14)
Total number of benign tumors	8			6			11	17
Number of tumors per tumor-bearing rat ^g	1.1			2.0			1.6	1.2
Females—n per group	96	c	c	95	c	c	96	97
Number (%) with benign tumors	79 (82)			81 (85)			80 (83)	83 (86)
Total number of benign tumors	165			218			245	287
Number of tumors per tumor-bearing rat ^f	2.1			2.7			3.1	3.5

^a50 ppm = 174 mg/m³, 200 ppm = 695 mg/m³, 500 ppm = 1,737 mg/m³, 1,500 ppm = 5,210 mg/m³, 3,500 ppm = 12,158 mg/m³.

^bLate 500 = no exposure for first 12 mo followed by 500 ppm for last 12 mo; early 500 = 500 ppm for first 12 mo followed by no exposure for last 12 mo.

^cNo data for this exposure level in this study.

^dEPA summed across these tumor types, assuming no overlap.

^eIn historical controls, percent with benign tumors reported was 79–82% and number per tumor-bearing rat was 2.1.

^fSignificantly ($p \leq 0.05$) higher than control incidence by Fisher's exact test (Nitschke et al., 1988a).

^gCalculated by EPA.

Sources: Nitschke et al. (1988a); Burek et al. (1984).

In Syrian golden hamsters exposed to 500, 1,500, or 3,500 ppm for 2 years, no statistically significantly increased incidences of tumors were found in any tissues (Burek et al., 1984).

Supporting evidence for the carcinogenicity of dichloromethane comes from the results of genotoxicity and mode of action studies discussed in Section 4.5. A mutagenic mode of carcinogenic action for dichloromethane involves metabolic activation by GST, as evidenced by several observations, including the enhancement of dichloromethane mutagenic activity in normally unresponsive *S. typhimurium* strain TA1535 after it is transfected with the gene for rat GST-T1 (DeMarini et al., 1997; Thier et al., 1993); increased HPRT gene mutations and DNA damage (DNA SSBs) in CHO cells when they are incubated with dichloromethane in the presence of mouse liver cytosol preparations rich in GST enzymatic activities (Graves and Green, 1996; Graves et al., 1996, 1994b); the detection of DNA damage (DNA SSBs) in liver and lung tissue of B6C3F₁ mice immediately following 6-hour inhalation exposure to dichloromethane (2,000–8,000 ppm); and a suppression of the DNA damage when mice were pretreated with buthionine sulphoximine, a GSH depletor (Graves et al., 1995).

Additional data from several studies indicate that dichloromethane genotoxicity is expressed in cancer target tissues in mice following in vivo exposure. Increased sister chromatid exchanges were observed in lung cells of B6C3F₁ mice after 90 days of inhalation exposure to 2,000 ppm or 10 days of exposure to 4,000 or 8,000 ppm (Allen et al., 1990). DNA damage (comet assay) was detected in liver and lung tissue (but not stomach, kidney, brain, or bone marrow) 24 hours after oral administration of 1,720 mg/kg dichloromethane to CD-1 mice (Sasaki et al., 1998). DNA-protein cross-links were observed in the liver of B6C3F₁ mice but not hamsters following inhalation exposure to concentrations ranging from 500 to 4,000 ppm 6 hours/day for 3 days (Casanova et al., 1996, 1992). Much less is known about genotoxicity in the liver in rats. Studies of DNA SSBs in rat hepatocytes or liver homogenate were negative with inhalation exposures up to 5,000 ppm for 3 hours (Graves et al., 1995, 1994b), but positive results were seen in a high-dose gavage study (1,275 mg/kg) (Kitchin and Brown, 1989). Few other specific types of genotoxicity endpoints (e.g., sister chromatid exchange, DNA-protein cross-links) have been studied in the rat liver.

Since there are limited data on mutagenic events following oral exposure, EPA conducted a pharmacokinetic analysis to evaluate how comparable the internal doses to the liver in the oral bioassay (Serota et al., 1986b; Hazleton Laboratories, 1983) were to the internal doses to the liver in the inhalation bioassay (Mennear et al., 1988; NTP, 1986). The PBPK model of Marino et al. (2006) predicted that the average daily amount of dichloromethane metabolized via GST per liter of liver was about 14-fold lower in mice exposed to the highest dose of 234 mg/kg-day in the drinking water bioassay than in mice exposed to the lowest inhalation exposure of 2,000 ppm inducing liver tumors (Table 4-43). Thus, the lower incidence of liver tumors induced by oral doses of 234 mg/kg-day compared with the higher incidence induced by inhalation exposure

to 2,000 ppm is consistent with the predicted lower liver dose of GST metabolites (and hence lower probability of DNA modification) with oral exposure. 317

While the amount metabolized by the GST pathway for inhalation exposure shown in Table 4-43 is lower in the rat vs. the mouse, as one would expect based on the enzyme expression level, for oral exposure a higher amount of GST metabolism is predicted for the rat than the mouse. This difference occurs because for oral exposure 100% of the dose is absorbed, rather than absorption being limited by metabolism as it is for inhalation, and because the ratio of GST to CYP activity is higher in the rat than in the mouse. Specifically k_{TC}/V_{maxc} is 0.626 for the rat and is 0.152 for the mouse, so for the rat the fraction of the absorbed dose going to GST is roughly four times that in the mouse, hence for the same oral dose per kg body weight per day (with 100% absorbed), approximately four times more is metabolized by GST in the rat than in the mouse.

Table 4-43. Comparison of internal dose metrics in inhalation and oral exposure scenarios in male mice and rats

External dose	Internal exposure in liver (mg metabolized through GST pathway/L liver tissue/d) ^a	
	Male	
	Mouse	Rat
Inhalation (ppm)		
2,000	2,364	1,509
4,000	4,972	3,124
Oral (mg/kg-d) ^b		
61	17.5	77.1
124	63.3	233.6
177	112.0	385.7
234	169.5	559.0

^aMouse values derived by EPA from the PBPK model of Marino et al. (2006); rat values derived from EPA based on the modified PBPK model of Andersen et al. (1991) (see Appendix C for model details).

^bActual doses administered to mice (Serota et al., 1986a); BWs not given for males and females, so simulation results only provided for one gender.

4.7.3. Mode-of-Action Information

4.7.3.1. Hypothesized Mode of Action

The hypothesized mode of action for dichloromethane-induced tumors is through a mutagenic mode of carcinogenic action. Specifically, the data indicate that dichloromethane is metabolized by GST to reactive metabolites that induce mutations in DNA leading to carcinogenicity. Much of the experimental mode of action research has focused on the liver and lung, the sites of tumor formation in chronic bioassays (Mennear et al., 1988; NTP, 1986; Serota

et al., 1986b, Hazleton Laboratories, 1983). The mode of action is potentially relevant to other sites, particularly those in which GST-T1 is expressed, such as mammary tissue (Lehmann and Wagner, 2008) and the brain (Juronen et al., 1996).

Support for the importance of GST in the hypothesized mutagenic mode of action has been demonstrated in in vitro bacterial and mammalian assays as well as in vivo mammalian system assays. Dichloromethane is consistently mutagenic in *S. typhimurium* strains with GST capability but did not produce mutagenic effects in non-GST *S. typhimurium* strains (summarized in Section 4.5.1.1 and Table 4-29). In vitro mammalian cell studies (see Table 4-30) have consistently demonstrated genotoxic effects in mouse Clara cells and in CHO cell lines when a mouse liver cytosol fraction was exogenously added; positive responses were seen in studies measuring DNA-protein cross-links, HPRT mutation analysis, and DNA SSBs. Other studies have demonstrated DNA adducts with dichloromethane exposure in calf thymus DNA in the presence of bacterial GST DM11. Negative results were seen in most of the other in vitro cell studies using rat hepatocytes or CHO cells without mouse liver cytosol incubation. These studies were conducted in cell lines where GST activity is considerably lower than in mouse cell lines and therefore, these results are not unexpected.

In studies with human cell lines or isolated cells, positive results were reported for sister chromatid exchanges, chromosomal aberrations, and in the micronucleus test. In vivo studies in mice (Section 4.5.1.2 and Table 4-32) consistently showed genotoxic effects following dichloromethane exposure in the liver and lung where tumors are observed. Other organs in the mouse were evaluated and mutagenic changes were not consistently observed. The specificity of the observed effects support the hypothesized mode of action since these positive mutagenic responses are seen in organs where tumor formation occurs (i.e., liver and lung) rather than in areas that were not the site of tumors in the mouse bioassays (e.g., stomach, bladder, kidney). In vivo genotoxicity studies in rats and hamsters (the other test systems used, see Table 4-33) were predominantly nonpositive. However, rats and hamsters have considerably lower GST activity than the mouse and may be less sensitive to dichloromethane-induced genotoxic effects.

In vivo binding of S-(chloromethyl)glutathione, dichloromethane's reactive GST metabolite, to DNA was not demonstrated in one study in rats and mice using a relatively low dose (5 mg/kg). The reactivity of the postulated DNA-reactive species and the instability of the derived adducts presents considerable challenges to the ability to provide direct evidence of adduct formation. Thus, this lack of in vivo evidence of S-(chloromethyl)glutathione binding to DNA does not in itself represent a basis for invalidating the proposed mode of action.

4.7.3.1.1. *Experimental support for the hypothesized mode of action*

Strength, consistency, and specificity of association. It is hypothesized that mutagenic events lead to the development of liver and lung tumors following dichloromethane exposure. Several observations from experimental studies support the mutagenicity of dichloromethane and

the key role of GST metabolism and the formation of DNA-reactive GST-pathway metabolites. The GST pathway produces two metabolites of dichloromethane, S-(chloromethyl)glutathione and formaldehyde, which are potentially reactive with DNA and other cell macromolecules. Enhanced dichloromethane genotoxicity in bacterial and mammalian *in vitro* assays with the introduction of GST metabolic capacity provides support that GST metabolism and metabolites are involved (DeMarini et al., 1997; Graves and Green, 1996; Graves et al., 1996, 1995, 1994b; Thier et al., 1993).

In bacterial strains where GST activity was not present (e.g., TA1535, TA1538), mutagenic effects were not reported following dichloromethane exposure (Oda et al., 1996; Simula et al., 1993; Osterman-Golkar et al., 1983; Gocke et al., 1981). Further tests of GST-dependent mutagenicity were evaluated by transfecting GST into non-GST bacterial strains or decreasing GST activity in GST bacterial strains (e.g., TA100). When GST-T1 was cloned into bacterial strain TA1535, dichloromethane treatment resulted in reverse mutations in this new GST⁺ TA1535 strain, and these mutations were independent of rat S9 metabolic activation (DeMarini et al., 1997; Pegram et al., 1997; Thier et al., 1993). Similarly, TA100/NG-11, a bacterial strain with decreased GST activity in comparison to the wild-type TA100 strain, showed significantly decreased mutagenicity (reverse mutations) following dichloromethane treatment (Graves et al., 1994a).

In vitro mammalian genotoxicity studies also support the importance of the GST pathway in relation to the positive effects observed following dichloromethane exposure. Positive results in the *in vitro* assays were limited to experiments with the presence of GST in the cell system. When mouse liver cytosol was added to hamster cell lines, dichloromethane induced DNA-protein cross-links, DNA SSBs, and HPRT gene mutations (Graves and Green, 1996; Graves et al., 1996, 1994b). Additionally, in mouse Clara cells (GST is localized in the lung cells of mice), DNA SSBs were reported following dichloromethane treatment, and the extent of DNA damage was significantly decreased when the cells were pretreated with a glutathione depletor (Graves et al., 1995). Other studies evaluating similar genotoxic endpoints in rat or CHO cells without modification of the low GST activity in the test system generally reported no evidence of genotoxic events (Graves et al., 1995; Andrae and Wolff, 1983; Garrett and Lewtas, 1983; Thilagar and Kumaroo, 1983; Jongen et al., 1981). A study evaluating the genotoxic effects of dichloromethane (up to 6 mM) in freshly isolated mouse, rat, hamster, and human hepatocytes provides additional supporting evidence of the influence of GST activity on mutagenicity (Casanova et al., 1997). Positive results were only observed in hepatocytes from B6C3F₁ mice; the interspecies variability in effects correlated proportionally with the enhanced GST metabolic capacity in mice (Reitz et al., 1989). In studies with human cell lines or isolated cells, positive results were reported for sister chromatid exchanges, chromosomal aberrations, DNA damage, and in the micronucleus test. Negative results were obtained with human cells in unscheduled DNA synthesis assays (Jongen et al., 1981; Perocco and Prodi, 1981) and

dichloromethane was not demonstrated to be genotoxic in studies of human hepatocytes (Casanova et al., 1997; Graves et al., 1995).

Two of three *in vivo* genotoxicity studies in insects reported positive results. Genotoxicity was observed in *Drosophila* for the gene mutation assay (Gocke et al., 1981) and the somatic assay (Rodriguez-Arnaiz, 1998) when dichloromethane was administered through the food. When *Drosophila* were exposed to dichloromethane via inhalation, genotoxic effects were negative as measured through gene mutation assays (sex-linked recessive lethal, somatic mutation and recombination) (Kramers et al., 1991).

In vivo genotoxicity studies reported DNA-protein cross-links, DNA SSBs, chromosomal aberrations, and sister chromatid exchanges in liver cells of B6C3F₁ mice following acute inhalation exposure to concentrations producing liver tumors with chronic exposure (Casanova et al., 1996, 1992; Graves et al., 1995, 1994b). The formation of DNA SSBs was suppressed when the mice were pretreated with a GSH depletor (Graves et al., 1995), providing additional support for the involvement of GST metabolism. Increased sister chromatid exchanges and chromosomal aberrations were found in the lungs of mice exposed to dichloromethane for 2 weeks to 8,000 ppm or for 12 weeks to 2,000 ppm. In this study, however, there was evidence of damage at other sites, too: sister chromatid exchanges were also seen in peripheral lymphocytes, chromosomal aberrations were seen in bone marrow, and micronuclei were seen in peripheral red blood cells under the same exposure protocol (Allen et al., 1990). As was seen in the liver, DNA SSBs were seen in lungs of B6C3F₁ mice following acute inhalation exposure to concentrations producing lung tumors with chronic exposure, and this effect was suppressed with pretreatment with a GSH depletor, buthionine sulfoximine (Graves et al., 1995). Other studies of sister chromatid exchange (Allen et al., 1990) or DNA damage detected by the comet assay (Sasaki et al., 1998) also provide evidence of genotoxic effects specifically in lung cells of mice. These *in vivo* mammalian genotoxicity studies demonstrate site-specific effects correlating to the dichloromethane-induced tumors in animals. Additional evidence for site specificity comes from a study in which DNA damage (detected by the comet assay) was enhanced in liver tissue but not stomach, kidney, brain, or bone marrow 24 hours after oral administration of 1,720 mg/kg dichloromethane to CD-1 mice (Sasaki et al., 1998).

DNA reaction products (e.g., DNA adducts) produced by GST metabolites, such as S-(chloromethyl)glutathione, have not been identified in *in vivo* studies (Watanabe et al., 2007). The authors speculated that these results are due to the instability of the reaction products (Hashmi et al., 1994). DNA adducts, however, have been observed in *in vitro* studies in which calf thymus DNA was incubated with dichloromethane and GST or was incubated with S-(1-acetoxymethyl)glutathione, a compound structurally similar to S-(chloromethyl)glutathione (Marsch et al., 2004; Kayser and Vuilleumier, 2001). These findings indicate that the S-(chloromethyl)glutathione intermediate formed by GSH conjugation has mutagenic potential and is likely responsible, at least in part, for the mutagenic response observed following

dichloromethane exposure. However, other studies (Hu et al., 2006; Casanova et al., 1996) provide evidence of formaldehyde-related DNA-protein cross-links in relation to dichloromethane exposure. These results show that, while most studies indicate the importance of the S-(chloromethyl)glutathione intermediate in mediating genotoxic damage following dichloromethane exposure, DNA damage resulting from formaldehyde formation should also be considered.

Mutagenic data in critical genes leading to the initiation of dichloromethane-induced liver or lung tumors are not available. In vivo assays evaluating mutations in tumor suppressor genes and oncogenes reported similar frequencies of activated *H-ras* genes and inactivation of the tumor suppressor genes, *p53* and *Rb-1* in the liver tumors seen in the nonexposed and dichloromethane-exposed B6C3F₁ mice (Devereaux et al., 1993; Hegi et al., 1993). There were too few lung tumors (n = 4) in controls to provide a conclusive comparison of mutation patterns between exposed and nonexposed tumors.

Dose-response concordance. Statistically significant increases in liver tumor incidences in male and female (2,000 and 4,000 ppm) mice were observed in the inhalation bioassay in B6C3F₁ mice (NTP, 1986). Several studies provide evidence of an association between mutagenic events mediated by GST-pathway metabolites and the exposure levels inducing liver tumors in B6C3F₁ in this study, and concentration-dependent increases in genotoxicity have been observed in in vitro and in vivo assays.

In vitro mammalian genotoxicity studies were positive and demonstrated a dose-response relationship for DNA-protein cross-links, DNA SSBs, and DNA damage as measured by the comet assay at concentrations ranging from 2.5 to 60 mM when mouse liver cytosol was added or if mouse GST-T1 was transfected into hamster cell lines (Hu et al., 2006; Graves et al., 1996, 1994b). In mouse hepatocytes, DNA-protein cross-links were observed following dichloromethane exposures ranging between 0.5 and 6.0 mM (Casanova et al., 1997). DNA-protein cross-links were detected in mouse hepatocytes incubated with 1.9 mM dichloromethane (Casanova et al., 1997), a concentration chosen based on its correspondence to the TWA liver concentration of dichloromethane that was predicted by the Andersen et al. (1987) PBPK model for mice exposed by inhalation to 4,000 ppm for 6 hours (a dose that resulted in increased liver tumor incidence in the 2-year bioassay reported by NTP, 1986). Consistent with the relative lack of liver tumor responses in Syrian golden hamsters (Burek et al., 1984) and F344 rats (NTP, 1986) with chronic exposure to 3,500 or 4,000 ppm, hepatocytes from these strains of animals did not form detectable DNA-protein cross-links when incubated with 1.9 mM dichloromethane (Casanova et al., 1997).

DNA-protein cross-links were not detected in livers of mice exposed to 146 ppm 6 hours/day for 3 days, but a concentration-dependent increase in DNA-protein cross-links was observed in DNA from livers of mice exposed to several concentrations between 500 and

4,000 ppm (Casanova et al., 1996). Following exposure under similar conditions (concentrations of 498, 1,553, or 3,923 ppm), DNA-protein cross-links were not detected in the livers of Syrian golden hamsters, a species that did not develop tumors after chronic inhalation exposure to dichloromethane (Casanova et al., 1996, 1992). Increased DNA SSBs were detected in liver tissue of B6C3F₁ mice immediately following a 6-hour inhalation exposure to dichloromethane at concentrations ranging from 2,000 to 8,000 ppm (Graves et al., 1995), and in mouse hepatocytes after a 3-hour exposure to 4,000 (but not 2,000) ppm (Graves et al., 1994b).

Statistically significant increases in the incidence of lung tumors were observed in the inhalation chronic bioassay in male and female B6C3F₁ mice exposed to 2,000 or 4,000 ppm dichloromethane (Mennear et al., 1988; NTP, 1986). Evidence of mutagenicity at these exposure levels comes from two inhalation studies (Graves et al., 1995; Allen et al., 1990). Increased DNA SSBs were detected in lung tissue of B6C3F₁ mice immediately following a 6-hour inhalation exposure to dichloromethane at concentrations ranging from 2,000 to 8,000 ppm (Graves et al., 1995). In the study by Allen et al. (1990), increased presence of sister chromatid exchanges was observed in mouse lung cells following a 12-week exposure at 2,000 ppm; shorter durations of exposure (2 weeks) were positive for measures of sister chromatid exchange and chromosome aberrations at 8,000 ppm, but not at 2,000 or 4,000 ppm.

DNA adducts were observed and increased with dose in an *in vitro* preparation of calf thymus DNA when treated with dichloromethane (5–60 mM) and bacterial, rat, or human GST (Marsch et al., 2004).

Temporal relationship. Dichloromethane-induced liver and lung tumors first appeared in mice after 52 weeks of exposure (Maronpot et al., 1995; Kari et al., 1993). The detection of DNA-protein cross-links in the livers of B6C3F₁ mice following short-term inhalation exposures to dichloromethane concentrations that induced tumors with chronic exposure (Casanova et al., 1996, 1992) provides temporal support for the proposed mutagenic mode of action. Additional supporting evidence comes from observations that increased levels of DNA SSBs were detected in the liver and lungs of B6C3F₁ mice immediately following 3-hour inhalation exposure to 2,000–8,000 ppm dichloromethane (Graves et al., 1995; 1994b). Single dose and inhalation exposure studies of ≤6 hours did not detect an effect on DNA synthesis (Lefevre and Ashby, 1989) or unscheduled DNA synthesis (Trueman and Ashby, 1987) in mouse liver cells.

Biological plausibility and coherence. Bioactivation of a parent compound into a mutagenic metabolite resulting in cancer is a plausible mode of action of carcinogenicity in humans and is a generally accepted mode of action. Dichloromethane-induced carcinogenicity is hypothesized to be due to metabolism of the parent compound by the GST pathway (GST-T1) to a metabolite that is tumorigenic. The GST metabolite, S-(chloromethyl)glutathione, formed from dichloromethane, has been characterized as labile and highly reactive through *in vitro* evaluation of dichloromethane metabolism in hepatocytes using [¹³C]-NMR techniques (Hashmi

et al., 1994) and through an enzyme digestion assay using calf thymus DNA and GST-T1 enzyme (Marsch et al., 2004). The hypothesis that the formation of a mutagenic metabolite is a preliminary step resulting in carcinogenicity is based on evidence that malignant tumors are primarily located in areas where dichloromethane is highly metabolized by GST-T1, such as the liver and the lung, and on mutagenicity studies indicating the importance of the GST pathway and that the lung and liver are more prone to mutagenic effects of dichloromethane (Sasaki et al., 1998; Casanova et al., 1996, 1992; Graves et al., 1995, 1994b). The site selectivity of the mutagenicity in liver and lung tissue as evidenced by several studies suggests that the GST reactive metabolite remains in the tissue where it is formed. Collectively, the studies support the hypothesis that dichloromethane-mediated carcinogenicity results from a GST metabolite that produces selective DNA damage in the tissues where the metabolite is formed, but this hypothesis is based in part on assumptions regarding metabolite clearance and reactivity. DNA damage in the liver and lung, as well as the increased incidence of tumor formation resulting from dichloromethane exposure, indicates coherence of the mutagenic and carcinogenic effects and is evidence supporting a mutagenic mode of action.

Differences in GST activity in mice compared with other species, and the interspecies variability in genotoxic effects corresponding to interspecies variability in tumor response, support the mode of action hypothesis. DNA SSBs were not detected in liver or lung cells in rats exposed to similar inhalation exposures that induce strand breaks in mice (Graves et al., 1995; Graves et al., 1994b) and were detected at much lower in vitro concentrations in isolated hepatocytes from B6C3F₁ mice (0.4 mM) than in hepatocytes from Alpk:APfSD rats (30 mM) (Graves et al., 1995, Figure 3). The difference in susceptibility to carcinogenic response between mice and rats likely reflects differences in GST metabolism. Toxicokinetic studies indicate that with increasing exposure levels, increasing amounts of dichloromethane are metabolized via GST metabolism.

4.7.3.1.2. *Other possible modes of action for liver or lung tumors in rodents.* Data are not available to support other possible modes of action for the liver and lung tumors in rodents. Efforts to observe sustained cell proliferation in liver following dichloromethane exposure of B6C3F₁ mice have been unsuccessful. Groups of female B6C3F₁ mice that were exposed to 0 or 2,000 ppm dichloromethane 6 hours/day, 5 days/week for up to 78 weeks did not exhibit enhanced cell proliferation in the liver when assessed at various intervals during exposure (Foley et al., 1993).

Indices of enhanced cell proliferation have been measured in the lungs of male B6C3F₁ mice following acute duration exposure at concentrations of about 1,500, 2,500, or 4,000 ppm dichloromethane (6 hours/day for 2 days) but not at exposure concentrations of 150 or 500 ppm and not in lungs of Syrian golden hamsters exposed to concentrations up to 4,000 ppm (Casanova et al., 1996). Earlier studies showed somewhat consistent findings in that the

numbers of bronchiolar cells undergoing DNA synthesis (thymidine incorporation labeling) were markedly increased (about 6- to 15-fold) in bronchiolar cells of B6C3F₁ mice exposed to 4,000 ppm dichloromethane 6 hours/day on days 5, 8, and 9 of exposure, but no evidence for increased cell proliferation was found after 89, 92, or 93 days of exposure (Foster et al., 1992). The results suggest that enhanced cell proliferation is not sustained in the lung with longer-term exposure to dichloromethane concentrations associated with lung tumor development in mice, and that this mode of tumor promotion is not important in the development of dichloromethane-induced lung tumors.

4.7.3.2. General Conclusions About the Mode of Action for Tumors in Rodents and Relevance to Humans

The mode of action for dichloromethane is hypothesized to involve mutagenicity via reactive metabolites. Mechanistic evidence indicates that dichloromethane-induced DNA damage in cancer target tissues of mice involves DNA-reactive metabolites produced via a metabolic pathway initially catalyzed by GST. Although mutational events in critical genes leading to tumor initiation have not been established, evidence supporting a mutagenic mode of action includes the identification of mutagenic response (reverse mutations) in short-term bacterial assays (with microsomal activation) and induced DNA-protein cross-links and DNA SSBs in mammalian cell assays. There are numerous positive *in vivo* genotoxicity studies specifically examining responses in the liver and/or lung; these studies included evidence of chromosomal aberrations, SSBs, sister chromatid exchanges, and DNA-protein cross-links. The negative *in vivo* genotoxicity assays are generally those that were based on a micronucleus test using mouse bone marrow, which is expected, as halogenated hydrocarbons (such as dichloromethane) are not very effective in this type of assay (Dearfield and Moore, 2005; Crebelli et al., 1999).

Is the hypothesized mode of action sufficiently supported in test animals? Consistent and specific evidence for the association between the formation of DNA-reactive GST-pathway metabolites and the formation of liver and lung tumors from inhalation includes: (1) enhanced GST metabolic capacity in the liver and lung and enhanced localization of GST-T1 in hepatic cell nuclei in B6C3F₁ mice compared with rats and hamsters, which do not show strong tumor responses to chronic inhalation exposure, (2) the detection of DNA-protein cross-links, or DNA SSBs in livers and lungs of B6C3F₁ mice following acute inhalation exposure to concentrations that produce tumors with chronic exposure, (3) suppression of the formation of DNA SSBs in livers and lungs of B6C3F₁ mice pretreated with a GSH depletory, (4) the inability to detect DNA-protein cross-links or DNA SSBs in livers or lungs of similarly exposed rats or hamsters, (5) detection of DNA SSBs at much lower *in vitro* concentrations in isolated hepatocytes from B6C3F₁ mice than in hepatocytes from Alpk:APfSD rats, (6) dose-response concordance and a

temporal relationship for the formation of DNA-protein cross-links and DNA SSBs with the formation of liver and lung tumors in B6C3F₁ mice exposed to dichloromethane, (7) the detection of increased sister chromatid exchanges in lung cells from CD-1 mice exposed by inhalation to dichloromethane, and (8) enhancement of dichloromethane genotoxicity in bacterial and mammalian in vitro assays with the introduction of GST metabolic capacity. However, mutations in critical genes linked to initiation of tumor cells have not been identified.

The much weaker carcinogenic response in the liver of rats and mice to chronic drinking water exposure (Serota et al., 1986a, b) than that noted in mice exposed by inhalation (Kari et al., 1993; NTP, 1986) is correlated with much smaller amounts of GST metabolites produced in the liver under the exposure conditions of the oral bioassay than in the inhalation bioassay (Andersen et al., 1987).

In conclusion, there is sufficient evidence supporting a mutagenic mode of action and indicating the involvement of GST metabolism in the lung and liver carcinogenicity of dichloromethane in mice.

Is the hypothesized mode of action relevant to humans? The postulated mode of action that dichloromethane is metabolized by GST to reactive metabolites that induce mutations in DNA leading to carcinogenicity is possible in humans. Mutagenicity as a mode of action for carcinogenicity in humans is generally accepted and is a biologically plausible mechanism for tumor induction. The toxicokinetic and toxicodynamic processes that would enable reactive metabolites to produce mutations in animal models are biologically plausible in humans. Furthermore, the detection of the GST pathway in human tissues indicates that the hypothesized mode of action involving reactive metabolites from this pathway, S-(chloromethyl)glutathione and formaldehyde, is relevant to humans.

Another factor that may play a role in the apparent species differences in carcinogenicity resulting from dichloromethane exposure is species differences in intracellular localization of GST-T1 (Sherratt et al., 2002; Mainwaring et al., 1996). In mouse liver tissue, GST-T1 appears to be localized in the nuclei of hepatocytes and bile-duct epithelium, but rat liver does not show preferential nuclear localization of GST-T1. In human liver tissue, some hepatocytes show nuclear localization of GST-T1 and others show localization in cytoplasm. Nuclear production of S-(chloromethyl)glutathione catalyzed by GST-T1 in the nucleus is more likely than cytoplasmic production to lead to DNA alkylation. The finding of some nuclear localization of GST-T1 in human liver tissue supports the relevance of the hypothesized mode of action to humans.

Comparisons in mice, rats, humans, and hamsters of GST enzyme activity in liver and lung tissues have indicated the following rank order: mice > rats > or ≈ humans > hamsters (Thier et al., 1998; Reitz et al., 1989). This relative ranking does not preclude the relevance of the hypothesized mode of action to humans, however.

Which populations or lifestyles can be particularly susceptible to the hypothesized mode of action? As discussed in Section 3.3, a polymorphism of the GST-T1 gene is present in humans. People with two functional copies of the gene (+/+) readily conjugate GSH to dichloromethane. Individuals having only one working copy of the gene (+/-) display relatively decreased conjugation ability. Individuals with no functional copy of the gene (-/-) do not express active GST-T1 protein and do not metabolize dichloromethane via a GST-related pathway (Thier et al., 1998). Thus, the GST-T1^{+/+} (wild-type) genotype would be considered to be the more “at risk” population; this subgroup represents approximately 30% of the U.S. population (Haber et al., 2002) but would be expected to be more common among Caucasians and African-Americans than among Asians (Raimondi et al., 2006; Garte et al., 2001; Nelson et al., 1995) (see Table 3-3).

According to the *Supplemental Guidance for Assessing Susceptibility from Early-Life Exposure to Carcinogens* (U.S. EPA, 2005b), children exposed to carcinogens with a mutagenic mode of action are assumed to have increased early-life susceptibility. The *Supplemental Guidance* (U.S. EPA, 2005b) recommends the application of age-dependent adjustment factors (ADAFs) for carcinogens that act through a mutagenic mode of action. Although the database is lacking in vivo evidence of specific mutagenic events following chronic exposure to dichloromethane, the weight of the available evidence indicates that dichloromethane is acting through a mutagenic mode of carcinogenic action. Application of ADAFs is recommended for both the oral and inhalation routes of exposure when risks are assessed that are associated with early-life exposure.

4.8. SUSCEPTIBLE POPULATIONS AND LIFE STAGES

4.8.1. Possible Childhood Susceptibility

In humans, hepatic CYP2E1 begins to be expressed in the second trimester (Johnsrud et al., 2003), increases significantly in the third trimester, and continues to increase during the first year of life (Hines, 2007; Johnsrud et al., 2003; Treluyer et al., 1996; Vieira et al., 1996). In the fetal brain, however, CYP2E1 activity is seen as early as GD 50, with increasing levels seen until at least the end of the first trimester (Brzezinski et al., 1999). Neurobehavioral effects of dichloromethane are seen with acute exposures in adults, and the available data regarding neurological symptom history and standardized testing suggest the possibility of longer-term effects. The relatively high activity of CYP2E1 in the brain compared to the liver of the developing human fetus raises the potential for neurodevelopmental effects from dichloromethane exposure. Results from a developmental toxicity study in rats also raise concern for possible neurodevelopmental effects. Decreased offspring weight at birth and changed behavioral habituation of the offspring to novel environments were seen following exposure of adult Long-Evans rats to 4,500 ppm for 14 days prior to mating and during gestation (or during gestation alone) (Bornschein et al., 1980; Hardin and Manson, 1980). In the only

other animal study examining possible early-life susceptibility to dichloromethane toxicity, Alexeef and Kilgore (1983) found that exposure of young male mice to approximately 47,000 ppm for about 20 seconds significantly impaired the ability to learn using a passive-avoidance conditioning task. Three-week-old mice were more affected than 5- or 8-week-old mice. The broad issue of childhood susceptibility to chronic neurobehavioral effects of early life exposure represents a data gap in the understanding of the health effects of dichloromethane.

The relatively low CYP2E1 activity in the liver of infants would tend to shift metabolism of dichloromethane to the GST pathway. This shift could affect cancer risk, given the evidence of genotoxicity through this metabolic pathway. However, the available data in humans are not sufficient to address the question of whether in utero or early life exposures represent a period of increased susceptibility to potential carcinogenic effects of dichloromethane. A threefold increased risk of childhood leukemia (acute lymphoblastic leukemia) was seen in relation to maternal occupational exposure in the year before and during pregnancy in one population-based case-control study (OR 3.22 [95% CI 0.88–11.7]) for ratings of “probable or definite” exposure compared with possible or no exposure (Infante-Rivard et al., 2005). The estimates for categories based on concentration and frequency were similar, but there was no evidence for an increasing risk with increasing exposure level.

Experiments comparing cancer responses from early-life exposures with those from adult exposures are not available for F344 rats or B6C3F₁ mice, the strains of animals in which carcinogenic responses to dichloromethane have been observed (mammary gland tumors in F344 rats and liver and lung tumors in B6C3F₁ mice exposed by inhalation; liver tumors in female F344 rats and male B6C3F₁ mice exposed via drinking water). Animal data evaluating the effect of age on the susceptibility to dichloromethane carcinogenicity are restricted to a bioassay in which 54 pregnant Sprague-Dawley rats were exposed starting on GD 12 to 100 ppm dichloromethane 4 hours/day, 5 days/week for 7 weeks, followed by 7 hours/day, 5 days/week for 97 weeks (Maltoni et al., 1988). Groups of 60 male and 69 female newborns continued to be exposed after birth to 60 ppm dichloromethane 4 hours/day, 5 days/week for 7 weeks, followed by exposure 7 hours/day, 5 days/week for 97 weeks. Additional groups of 60 male and 70 female newborns were exposed after birth to 60 ppm dichloromethane 4 hours/day, 5 days/week for 7 weeks and then for 7 hours/day, 5 days/week for 8 weeks. Endpoints monitored included clinical signs, BW, and full necropsy at sacrifice (when spontaneous death occurred). For each animal sacrificed, histopathologic examinations were performed on the following organs: brain and cerebellum, zymbal glands, interscapular brown fat, salivary glands, tongue, thymus and mediastinal lymph nodes, lungs, liver, kidneys, adrenals, spleen, pancreas, esophagus, stomach, intestine, bladder, uterus, gonads, and any other organs with gross lesions. There was no significant effect of exposure to dichloromethane on the incidence of benign or malignant tumors among adults or the progeny. The results provide no evidence that Sprague-Dawley rats would be more sensitive to potential carcinogenic activity of dichloromethane

during early life stages. Further conclusions from these results are precluded because the study included only one exposure level, which was below the maximum tolerated dose for adult Sprague-Dawley rats.

4.8.2. Possible Gender Differences

The limited data available from studies in humans do not indicate that there are large differences by gender in sensitivity to cardiovascular, neurologic, cancer, or other effects; studies have not been conducted specifically to examine this question and so do not provide information pertaining to smaller or more subtle differences. The available animal studies similarly do not establish whether either gender may be more susceptible to the toxic effects of dichloromethane. Studies of the carcinogenic effects of dichloromethane, either by inhalation or by the oral route, have not suggested an increased susceptibility of either male or female animals.

4.8.3. Other

As discussed in Section 3.3, a polymorphism exists within the GST-T1 gene in humans, resulting in individuals with diminished or a lack of ability to conjugate GSH to dichloromethane. While the possible effects of this polymorphism on the toxicity of dichloromethane have not been directly demonstrated, it can be inferred from the proposed mode of action that a decrease in the GST-T1 metabolic pathway would result in a decreased generation of reactive metabolites and a decrease in any chronic effects mediated through those metabolites (Jonsson and Johanson, 2001; El-Masri et al., 1999).

Interindividual variation in the ability to metabolize dichloromethane via GST-T1 is associated with genetic polymorphisms in humans. Estimated U.S. population prevalence of nonconjugators (-/- at the GST-T1 locus) is about 20%, but higher prevalences (47–64%) have been reported for Asians (Raimondi et al., 2006; Haber et al., 2002; Garte et al., 2001; Nelson et al., 1995). Although nonconjugators are expected to have negligible extra risk for dichloromethane-induced cancer, the U.S. prevalences for low (+/- at the GST-T1 locus) and high (+/+) conjugators have been estimated at 48 and 32%, respectively (Haber et al., 2002). The liver and kidney are the most enriched tissues in GST-T1, but evidence is available for the presence of GST-T1 in other tissues including the brain and lung at lower levels (Sherratt et al., 2002, 1997).

Individuals may vary in their ability to metabolize dichloromethane through the CYP2E1 pathway. Individuals with decreased CYP2E1 activity may experience decreased generation of CO and an increased level of GST-related metabolites following exposure to dichloromethane, which may result in increased susceptibility to the chronic effects of dichloromethane from GST-related metabolites. Conversely, individuals with higher CYP2E1 activity may experience relatively increased generation of CO at a given dichloromethane exposure level and therefore, may be more susceptible to the acute CO-related toxicity or other chronic effects of

dichloromethane. Several studies indicate a three- to sevenfold variability in CYP2E1 activity among humans, as assessed by various types of measurements among “healthy” volunteers (Sweeney et al., 2004; Haufroid et al., 2003; Lipscomb et al., 2003; Lucas et al., 2001, 1999; Bernauer et al., 2000; Kim et al., 1995; Shimada et al., 1994). This variability is incorporated into the PBPK models for dichloromethane. Factors that may induce or inhibit CYP2E1 activity (e.g., obesity, alcohol use, diabetes) or co-exposures (i.e., to various solvents or medications) (Lucas et al., 1999) may result in greater variation within segments of the population. This variation in CYP2E1 activity may result in earlier saturation of this pathway and greater exposure to the parent compound, which would be of particular relevance to neurological effects.

5. DOSE-RESPONSE ASSESSMENTS

5.1. ORAL REFERENCE DOSE (RfD)

5.1.1. Choice of Principal Study and Critical Effect—with Rationale and Justification

As discussed in Section 4.6.1, human data for oral exposures to dichloromethane are limited to case reports involving intentional (i.e., suicidal) or accidental, acute ingestion exposures (Chang et al., 1999; Hughes and Tracey, 1993). Reported effects reflect frank toxicity from very high doses such as marked CNS depression, injury to the gastrointestinal tract, liver and kidney failure, coma, and death. No studies of human chronic oral exposures are available. In the absence of adequate studies evaluating possible health effects in humans repeatedly exposed to dichloromethane via the oral route, the results from the chronic laboratory animal studies are assumed to be relevant to humans.

The database of laboratory animal oral exposure studies includes 90-day (Kirschman et al., 1986) and 2-year drinking water toxicity studies in F344 rats (Serota et al., 1986a) and B6C3F₁ mice (Serota et al., 1986b; Hazleton Laboratories, 1983). A reproductive study exposed Charles River CD rats via gavage before mating (General Electric Company, 1976), and a developmental study exposed F344 rats via gavage during GDs 6–19 (Narotsky and Kavlock, 1995). A 14-day gavage study examined neurotoxicity in F344 rats (Moser et al., 1995).

Hepatic effects (hepatic vacuolation, liver foci) are the primary dose-dependent noncancer effects associated with oral exposure to dichloromethane (see Table 4-35). The 90-day drinking water toxicity study in F344 rats (Kirschman et al., 1986) reported significant increases in hepatocyte vacuolation and necrosis in animals dosed between 166 and 1,200 mg/kg-day (males) or 200 and 1,469 mg/kg-day (females). These doses were used to develop dosing levels for the 104-week drinking water study (Serota et al., 1986a). The 104-week drinking water study of F344 rats (Serota et al., 1986a) provides adequate data to describe dose-response relationships for liver lesions from chronic oral exposure to dichloromethane (e.g., includes four exposure levels and a control group). In this study, rats dosed at ≥ 50 mg/kg-day in both sexes had increased fatty livers, but quantitative data were not provided by the authors. Liver lesions, described as foci or areas of cellular alteration, were also seen in this study in the same dose groups in which the fatty changes had occurred. A limitation of this study is that Serota et al. (1986a) did not describe the evaluation of the altered foci in detail. However, increases in altered foci did not correspond to tumor rate incidences in either male or female rats. Instead, the altered foci correlated more closely to fatty liver incidence changes for both sexes in the rats. Altered foci could range from a focal fatty change (nonneoplastic) to an enzymatic altered foci change (neoplastic) (Goodman et al., 1994). Several lines of evidence were considered in determining whether the lesions should be characterized as nonneoplastic or neoplastic: (1) there is a congruence between the incidence of this lesion and the incidence of

the fatty liver in the study by Serota et al. (1986a), (2) at higher doses, hepatocyte vacuolation and hepatocyte necrosis were seen (Berman et al., 1995; Kirschman et al., 1986), and (3) there is no clear indication that these altered foci progress to liver tumors since the rate of increased foci did not correlate with liver tumor increases in either male or female rats. Based on these observations, EPA concluded that the altered foci were more likely to be representative of a focal fatty change (nonneoplastic) than a neoplastic event.

The LOAELs for the liver lesions in rodents following repeated oral exposure (50–586 mg/kg-day) (Table 4-35) are in the same range or below the NOAELs of 225 mg/kg-day for reproductive performance in Charles River CD rats exposed for 90 days before mating (General Electric Company, 1976) and 450 mg/kg-day for developmental toxicity in pregnant F344 rats exposed during gestation (Narotsky and Kavlock, 1995). The LOAEL (337 mg/kg-day) and NOAEL (101 mg/kg-day) for mild neurological impairment in a 14-day gavage exposure study of F344 rats (Moser et al., 1995) indicates that the threshold for neurological effects may be similar to the threshold for liver effects. A limitation of the Moser et al. (1995) study, however, is that the observed effects were limited to measures taken within 4 hours of exposure.

The subchronic (i.e., ≤ 90 -day study) data were not considered in the selection of a principal study for deriving the chronic RfD because the database contains reliable dose-response data from a chronic study at lower doses than the 90-day study (Kirschman et al., 1986) (conducted to provide data pertaining to relevant doses to use in the chronic study). The data from the subchronic studies are, however, used to corroborate the findings in the chronic studies with respect to relevant endpoints (i.e., hepatic and neurological effects). The neurotoxicity study was not selected as the principal study due to the limited measurements to inform the chronic exposure to dichloromethane. The rat rather than the mouse chronic bioassay (Serota et al., 1986a) was selected as the principal study for the RfD because of the consistent evidence that rats may be more sensitive than mice to noncancer liver effects from orally administered dichloromethane; available rat LOAELs for liver lesions are lower than mouse LOAELs (see Table 4-35). Figure 5-1 is an exposure-response array that presents NOAELs, LOAELs, and the dose range tested, corresponding to selected health effects from the short-term (neurotoxicological) and subchronic studies, and from the chronic, reproductive, and developmental toxicity studies that were evaluated for use in the derivation of the RfD.




Figure 5-1. Exposure response array for oral exposure to dichloromethane.

5.1.2. Derivation Process for Noncancer Reference Values

The toxicity values (oral RfD and inhalation RfC) for noncancer endpoints were derived by using rat and human PBPK models to calculate internal doses in rats from experimental exposures and extrapolate points of departure to human equivalent exposures. Figure 5-2 illustrates the process of using the PBPK models for toxicity value derivation. The process for the RfD and RfC is summarized below, using the example of a noncancer liver effect.




Figure 5-2. Process for deriving noncancer oral RfDs and inhalation RfCs using rodent and human PBPK models.

A deterministic PBPK model for dichloromethane in rats was first used to convert rat drinking water or inhalation exposures to values of an internal liver dose metric (see Appendix C for details of the rat PBPK model). Available models in EPA benchmark dose (BMD) software (BMDS) version 2.0 were then fit to the liver lesion incidence data, and internal liver dose data for rats and BMD₁₀ values and their lower 95% confidence limits associated with a 10% extra risk (BMDL₁₀) were calculated from each of the models. Adequacy of model fit was assessed by overall χ^2 goodness of fit ($p > 0.10$) and examination of residuals, particularly in the region of the benchmark response (BMR). The choice of best-fitting model was based on the lowest Akaike's Information Criterion (AIC) among models with adequate fits (U.S. EPA, 2000b).⁹

The use of a PBPK model can replace the use of the BW^{0.75} scaling factor to account for interspecies differences in toxicokinetics. The decision with respect to use of a scaling factor depends on the dose metric that is used. Where PBPK models predict the concentration (in particular, the AUC) of the proximate causative agent, a scaling factor to account for interspecies differences is not typically used. That is, it is assumed that if the time-averaged (or steady-state) concentration of the proximate causative agent predicted by the PBPK model in the target tissue is the same in the test species as in humans, and the test species was exposed for an equivalent portion of its lifetime (2 years in rats and mice being equivalent to a 70-year lifetime in humans), then the resulting risks in the two species are the same. However, when the PBPK model predicts the rate of production of the agent rather than its concentration, then a BW^{0.75} scaling factor may be appropriate, depending on what is known or expected regarding the rate of clearance of the agent or metabolite of interest. Two different scenarios can be considered. If the metabolite formed is considered to be highly reactive and is unlikely to involve processes or cofactors for which the rate or availability can be expected to scale allometrically, then it can be assumed that the rate of clearance (i.e., disappearance due to local reactivity) for this metabolite per volume tissue is equal in rodents and humans. Thus, in that situation as with the AUC dose metric, no BW^{0.75} scaling factor is necessary, although differences in tissue volume fraction in humans versus rats (as occurs for liver) should be and are accounted for by the PBPK model. However, if the metabolite is removed by processes that scale allometrically (including enzymatic reactions or reactions with cofactors whose supply is limited by overall metabolism) then it is expected that interspecies differences in clearance or removal of the toxic metabolite follow the generally assumed BW^{0.75} scaling for rates of metabolism and blood circulation. In this case, or in situations in which the reactivity or rate of removal of the metabolite has not been established, it is appropriate to use a scaling factor based on BW ratios to account for this difference. In the case of the noncancer liver effects of dichloromethane, very limited

⁹ If more than one model shares the lowest AIC value, BMDL₁₀ values from these models may be averaged to obtain a POD. However, this average is not a well-defined lower bound, and should be referred to only as averages of BMDL₁₀s. U.S. EPA does not support averaging BMDLs in situations in which AIC values are similar, but not identical, because the level of statistical confidence is lost and because there is no consensus regarding a cut-off between similar and dissimilar AIC values.

information is available on the mechanism(s) involved in creating the type of hepatic damage seen. The dose metric used in the PBPK modeling is a rate of metabolism rather than the concentration of putative toxic metabolites, and the clearance of these metabolites may be slower per volume tissue in the human compared with the rat. Thus, the rat internal dose metric for noncancer effects was adjusted by dividing by a pharmacokinetic scaling factor to obtain a human-equivalent internal BMDL₁₀.

A probabilistic PBPK model for dichloromethane in humans, adapted from the model of David et al. (2006) as described in Appendix B, was then used to calculate distributions of chronic exposures associated with the human equivalent internal BMDL₁₀, based on the responses in rats. Parameters in the human PBPK model are described by distributions that incorporate information about dichloromethane toxicokinetic and physiological variability and uncertainty among humans, with additional information on human variability for both the CYP2E1 and GST-T1 metabolic pathways (see Table 3-9 and Appendix B). Monte Carlo sampling was performed in which each human model parameter was defined by a value randomly drawn from its respective parameter distribution. The model was then executed by using the human internal BMDL₁₀ as input, and the resulting human equivalent dose or human equivalent concentration (HEC) was recorded. This process was repeated for 10- to 20,000 iterations to generate a distribution of human equivalent doses or concentrations.

As discussed in Section 3.5.2, the statistics reported for the fitted metabolic parameters by David et al. (2006; Table 4 in that publication) only represent the population mean and uncertainty in that mean for each parameter. For the parameters other than $V_{\max C}$ and k_{fC} , EPA considers it reasonable to assume that there is little true interindividual variability in the values, so the distributions were used as published in David et al. (2006). For the physiological parameters, the distributions presented by David et al. (2006) were supposed to represent a known range of interindividual variability, but EPA found that these did not adequately describe the full population, so many of the distributions were changed as discussed in Appendix B. For $V_{\max C}$, an independent data set (Lipscomb et al., 2003) where CYP2E1 levels were measured in vitro using liver samples from 75 human donors was used to estimate the degree of interindividual variability, and a “two-dimensional” sampling routine was used to incorporate the uncertainty as estimated by David et al. (2006) from those in vitro data. Finally, EPA concluded that the trivariate distribution (based on GST-T1 genotype), which David et al. (2006) used in place of the observed parameter uncertainty (based on ex vivo data from Warholm et al., 1994), adequately represented interindividual variability but neglected the uncertainty in the population mean. Therefore, a two-dimensional sampling routine was used: first a specific value for the population mean was sampled from the mean and variance (uncertainty) indicated for k_{fC} in Table 4 of David et al. (2006); second, given that value for the population mean, an individual value was sampled using the trivariate distribution as indicated in Table 2 of David et al. (2006), but re-scaled to the (sampled) population mean.

The sampling routine used by EPA effectively assumes that the parameters are distributed independently, ignoring the covariance that was likely represented in the actual posterior chains, hence will tend to over-estimate the overall range of parameters and distribution of dose metrics in the population compared to what one would obtain if the covariance were explicitly included. (This is offset to some extent by the assumption that there is no interindividual variability in the metabolic parameters other than $V_{\max C}$ and k_{fC} .) Thus if the covariance (i.e., the variance-covariance matrix) for the set of parameters had been reported by David et al. (2006) it could have been used to narrow the predicted distribution of internal doses or equivalent applied doses. Lacking such information, the approach used will not underestimate risk or overestimate lower bounds on human equivalent exposure levels.

From these distributions of human equivalent doses (or concentrations), candidate RfDs or RfCs were derived by dividing the first percentile value (point of departure) by uncertainty factors (UFs) to account for uncertainty about potential interspecies toxicodynamic variability, human toxicodynamic variability, and database deficiencies. The first percentile was chosen because it allowed generation of a stable estimate for the lower end of the distribution while being protective of the overall human population, including sensitive individuals. Choosing this lower point replaces the use of an additional UF to account for human toxicokinetic variability.

5.1.3. Evaluation of Dose Metrics for Use in Noncancer Reference Value Derivations

There are no data to support the role of a specific metabolite in the development of the noncancer liver lesions seen in oral and inhalation exposure studies. Four dose metrics were examined as potential metrics for the internal dose of interest: rate of hepatic metabolism through the CYP pathway, rate of hepatic metabolism through the GST pathway, the combined rate of hepatic metabolism through the CYP and GST pathways, and the concentration (AUC) of dichloromethane (the parent compound) in the liver. The dose-response patterns for each of these metrics in the oral study in rats (Serota et al., 1986a) and in two inhalation studies in rats (Nitscke et al., 1988a; Burek et al., 1984) were examined for fit and congruence.

Using the oral exposure data, only one of the seven models, the log-logistic model, produced an adequate fit ($p > 0.10$) for the GST metabolism metric and the dichloromethane AUC metrics. Adequate model fit was seen in all of the models using the CYP dose metric with the oral data and using the GST, CYP, and AUC dose metrics for the inhalation data.

A limitation in using the GST metric can be observed when comparing the oral and inhalation responses at various exposure levels. At 200 ppm, where the GST metric is predicted by the PBPK model to be 93 mg metabolism per liter liver per day, no liver effects were seen. In contrast, liver responses were elevated at an oral dose of 50 mg/kg-day, where the GST metric is predicted to be 60 mg metabolism per liter liver per day (see Tables 5-1 and 5-5, respectively, for the oral and inhalation internal metrics). Thus the liver GST metric produces an inconsistency in the dose-response relationship with very different responses observed depending on the route of

exposure. A similar inconsistency occurs with the AUC metric. These differences are not observed, however, when using the CYP metric. At the 200 ppm inhalation exposure, where no hepatotoxicity was observed, the CYP metric is predicted to be 660 mg per liter liver per day. This internal CYP metabolism metric is less than that predicted for the oral dose for the 50 mg/kg-day group (i.e., 872 mg metabolism per liter liver per day) in which liver effects were observed. Thus, the CYP internal metric is consistent with the observed responses seen in the oral and inhalation exposure studies.

The GST metabolism and the AUC dose metrics did not present reasonable choices based on model fit and consistency of response across studies at comparable dose levels. Given these results, the combination of hepatic metabolism through the GST and the CYP pathways would not be expected to result in an improvement to a metric based only on CYP metabolism. Thus the CYP-metabolism dose metric is the most consistent with the data and this metric was selected for the subsequent RfD and RfC derivations. The lack of information on mechanisms with respect to noncancer health effects represents data gaps in the understanding of the health effects of dichloromethane.

5.1.4. Methods of Analysis—Including Models (PBPK, BMD, etc.)

PBPK models for dichloromethane in rats were described previously in Section 3.5. From the evaluation described in Appendix C, a modified model of Andersen et al. (1991) was selected for the calculation of internal dosimetry of ingested dichloromethane in the rats in the principal study (Serota et al., 1986a).

PBPK model simulations of the drinking water study of Serota et al. (1986a) (Table 5-1) were performed to calculate average lifetime daily internal liver doses in male and female F344 rats. In the absence of data for group- and sex-specific BWs, reference values were used for male and female F344 rats in chronic studies (U.S. EPA, 1988a). The mode of action by which dichloromethane induces noncancer liver effects in rodents has not received research attention to determine the role of the parent material, metabolites of the CYP2E1 pathway, metabolites of the GST pathway, or some combination of parent material and metabolites. In the absence of this kind of knowledge, and considering the pattern of response seen in the oral and inhalation studies (as described in Section 5.1.3), an internal dose metric based on the amount of dichloromethane metabolized via the CYP pathway in the liver (mg dichloromethane metabolized via CYP pathway per liter liver per day) was used. Figure 5-3 shows the comparison between oral external and internal doses using this dose metric for the rat and for the human.

Table 5-1. Incidence data for liver lesions and internal liver doses based on various metrics in male and female F344 rats exposed to dichloromethane in drinking water for 2 years


Sex	Nominal (actual) daily intake (mg/kg-d)	Rat liver lesion incidence ^a	Rat internal liver dose ^b			
			CYP	GST	GST and CYP	Parent AUC
Male (BW = 380 g)	0 (0)	52/76 (68%)	0	0	0	0
	5 (6)	22/34 (65%)	133.9	2.1	136.1	0.5
	50 (52)	35/38 (92%) ^c	872.7	58.8	931.4	13.1
	125 (125)	34/35 (97%) ^c	1,433.1	236.0	1,669.1	52.6
	250 (235)	40/41 (98%) ^c	1,868.6	561.5	2,430.0	125.0
Female (BW = 229 g)	0 (0)	34/67 (51%)	0	0	0	0
	5 (6)	12/29 (41%)	134.5	2.1	136.6	0.4
	50 (58)	30/41 (73%) ^c	977.8	66.0	1,043.8	12.6
	125 (136)	34/38 (89%) ^c	1,577.0	258.7	1,835.7	49.5
	250 (263)	31/34 (91%) ^c	2,070.0	642.4	2,712.3	122.9

^aLiver foci/areas of cellular alteration; number affected divided by total sample size.

^bInternal doses were estimated using a rat PBPK model from simulations of actual daily doses reported by the study authors. CYP dose is in units of mg dichloromethane metabolized via CYP pathway/L tissue/d; GST dose is in units of mg dichloromethane metabolized via GST pathway/L tissue/d; GST and CYP dose is in units of mg dichloromethane metabolized via CYP and GST pathways/L tissue/d; and parent AUC dose is in units of mg dichloromethane × hrs/L tissue.

^cSignificantly ($p < 0.05$) different from control with Fisher's exact test.

Source: Serota et al. (1986a).



Six simulated daily drinking water episodes are described by Reitz et al. (1997). The human metabolism rates were estimated using a computational sample of 1,000 individuals per dose, including random samples of the three GST-T1 polymorphisms (+/+, +/-, -/-) in the current U.S. population based on data from Haber et al. (2002). Since a different set of samples was used for each dose, some stochasticity is evident as the human points (values) do not fall on smooth curves.

Figure 5-3. PBPK model-derived internal doses (mg dichloromethane metabolized via the CYP pathway per liter liver per day) in rats and humans and their associated external exposures (mg/kg-day), used for the derivation of RfDs.

The seven dichotomous dose-response models in BMDS version 2.0 were fit to the rat liver lesion incidence data and PBPK model-derived internal dose data to derive a rat internal BMD₁₀ and corresponding BMDL₁₀ associated with 10% extra risk (Table 5-2). The quantal model is identical to the one-stage multistage model and so is not included in this set of models. A BMR of 10% was selected because, in the absence of information regarding the magnitude of change in a response that is thought to be minimally biologically significant, a BMR of 10% is generally recommended since it provides a consistent basis of comparison across assessments. There are no additional data to suggest that the critical response has a greater sensitivity that would warrant a lower BMR. The male rats exhibited a greater sensitivity compared to the female rats (based on lower BMDL₁₀ values for all of the models examined), and thus the male data are used as the basis for the RfD derivation. The logistic model was the best fitting model for the male incidence data based on AIC value among models with adequate fit (U.S. EPA, 2000b). Modeling results are shown in detail in Appendix D-1.

Table 5-2. BMD modeling results for incidence of liver lesions in male and female F344 rats exposed to dichloromethane in drinking water for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolism via CYP pathway per liter liver tissue per day)

Sex and model ^a	BMD ₁₀	BMDL ₁₀	χ^2 goodness of fit p-value	AIC
Males				
Gamma ^a	151.73	48.93	0.62	185.33
Logistic^b	85.17	61.78	0.75	183.61
Log-logistic ^a	213.73	37.06	0.83	184.79
Multistage (1) ^a	68.62	47.58	0.71	183.74
Probit	98.87	75.49	0.69	183.81
Log-probit ^a	197.65	77.56	0.81	184.84
Weibull ^a	117.29	48.39	0.57	185.49
Females				
Gamma ^a	336.38	98.70	0.52	233.07
Logistic	169.77	134.87	0.59	231.70
Log-logistic ^a	404.87	101.15	0.60	232.80
Multistage (1) ^a	123.59	91.46	0.47	232.32
Probit	179.59	146.27	0.59	231.70
Log-probit ^a	400.95	173.57	0.60	232.80
Weibull ^a	283.24	97.31	0.47	233.27

^aThese models in EPA BMDS version 2.0 were fit to the rat dose-response data shown in Table 5-1 by using internal dose metrics calculated with the rat PBPK model. Details of the models are as follows: Gamma and Weibull models restrict power ≥ 1 ; Log-logistic and Log-probit models restrict to slope > 1 , multistage model restrict betas ≥ 0 ; lowest degree polynomial with an adequate fit is reported (degree of polynomial noted in parentheses).

^bBolded model is the best-fitting model in the most sensitive sex (males), which is used in the RfD derivation.

Source: Serota et al. (1986a).

The BMDL₁₀ from the logistic model was used as the point of departure (POD) for the RfD calculations (Table 5-3). This rat internal dose metric for noncancer effects was adjusted to obtain a human-equivalent internal BMDL₁₀ by dividing by a pharmacokinetic scaling factor based on a ratio of BWs ($BW_{\text{human}}/BW_{\text{rat}})^{0.25} = 4.09$). This scaling factor was used because the metric is a rate of metabolism rather than the concentration of putative toxic metabolites, and the clearance of these metabolites may be slower per volume tissue in the human compared with the rat (that is, total rate of removal may scale as $BW^{0.75}$, while tissue volume scales as BW^1).

Table 5-3. RfD for dichloromethane based on PBPK model-derived probability distributions of human drinking water exposures extrapolated from liver lesion incidence data for male rats exposed via drinking water for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day)

Model ^a	Rat internal BMDL ₁₀ ^b	Human internal BMDL ₁₀ ^c	Human equivalent dose (mg/kg-d) ^d			Human RfD (mg/kg-d) ^e
			1 st percentile	5 th percentile	Mean	
Logistic	61.78	15.11	0.216	0.253	0.399	7×10^{-3}

^aBased on the best-fitting model from Table 5-2.

^bRat dichloromethane PBPK model-derived internal liver dose associated with the lower bound on 10% extra risk for developing liver foci/areas of cellular alteration.

^cHuman dichloromethane internal liver dose, derived by dividing the rat internal BMDL₁₀ by a scaling factor of 4.09 [(BW_{human}/BW_{rat})^{0.25}] to account for potential interspecies pharmacokinetic differences in the clearance of metabolites.

^dPBPK model-derived distributions of daily average dichloromethane drinking water doses predicted by the PBPK model to yield an internal dose in humans equal to the dichloromethane internal BMDL₁₀.

^eHuman RfD, based on male rat data, derived by dividing the 1st percentile of human equivalent dose value by a total UF of 30: 3 (10^{0.5}) for possible toxicodynamic differences between species, 3 (10^{0.5}) for variability in human toxicodynamic response, and 3 (10^{0.5}) for database deficiencies. The 1st percentile POD is a stable estimate of the lower end of the distribution. Use of this value in the lower tail replaces use of a UF for human toxicokinetic variability. See Section 5.1.5 for discussion of UFs.

Source: Serota et al. (1986a).

The human PBPK model (adapted from David et al. [2006], as described in Appendix B), using Monte Carlo sampling techniques, was used to calculate quantiles of human equivalent administered oral daily doses (in mg/kg-day) associated with the internal BMDL₁₀ values (Table 5-3), as described above in Section 5.1.2. The human model used parameter values derived from Monte Carlo sampling of probability distributions for each parameter, including MCMC-derived distributions for the metabolic parameters for the metabolism through the CYP2E1 pathway (V_{max} and K_m), and a distribution of GST metabolic rate constants that is weighted to reflect the estimated frequency of GST-T1 genotypes (20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+}) in the current U.S. population based on data from Haber et al. (2002). All simulations also included a distribution of CYP activity based on data from Lipscomb et al. (2003). The drinking water exposures comprised six discrete drinking water episodes for specified times and percentages of total daily intake (Reitz et al., 1997). The mean and two lower points on the distributions of human equivalent administered daily doses derived from the Serota et al. (1986a) data for male rats, using the BMDL₁₀ from the logistic model, are shown in Table 5-3. Although a lower value in this distribution could be calculated, this would require proportionately greater iterations (i.e., up to 10,000) to achieve numerical stability.

5.1.5. RfD Derivation—Including Application of Uncertainty Factors (UFs)

The 1st percentile POD is a numerically stable estimate of the lower end of the distribution. Use of this value associated with a sensitive human population addresses the uncertainty associated with human toxicokinetic variability. To derive the candidate RfD based on data from male rats, the first percentile value of the distribution of human equivalent dose associated with the male rat-derived BMDL₁₀ was divided by a composite UF of 30 (3 [10^{0.5}] to account for uncertainty about interspecies toxicodynamic equivalence, 3 [10^{0.5}] to account for uncertainty about toxicodynamic variability in humans, and 3 [10^{0.5}] for database deficiencies) (Table 5-3). The resulting RfD recommended for dichloromethane is 7×10^{-3} mg/kg-day.

In deriving this RfD, factors for the following areas of uncertainty were considered:

- Uncertainty in extrapolating from laboratory animals to humans (UF_A). The use of PBPK models to extrapolate internal doses from rats to humans reduces toxicokinetic uncertainty in extrapolating from the rat liver lesion data but does not account for the possibility that humans may be more sensitive than rats to dichloromethane due to toxicodynamic differences. A default UF of 3 (10^{0.5}) to account for this toxicodynamic uncertainty was used, as shown in Table 5-3.
- Uncertainty about variation from average humans to sensitive humans (UF_H). The probabilistic human PBPK model used in this assessment incorporates the best available information about variability in toxicokinetic disposition of dichloromethane in humans but does not account for humans who may be sensitive due to toxicodynamic factors. Thus, a default UF of 3 (10^{0.5}) was applied to account for possible toxicodynamic differences in sensitive humans.
- Uncertainty in extrapolating from LOAELs to NOAELs (UF_L). A UF for extrapolation from a LOAEL to a NOAEL was not applied because BMD modeling was used to determine the POD, and this factor was addressed as one of the considerations in selecting the BMR. The BMR was selected based on the assumption that it represents a minimum biologically significant change.
- Uncertainty in extrapolating from subchronic to chronic durations (UF_S). The derived RfD is based on results from a chronic-duration drinking water toxicity study. No cross-duration UF is necessary.
- Uncertainty reflecting database deficiencies (UF_D). The oral database for dichloromethane includes well-conducted lifetime drinking water studies in rats (Serota et al., 1986a) and mice (Serota et al., 1986b) and a supporting subchronic study in rats and mice (Kirschman et al., 1986). These studies provided dose-response data for the hepatic effects of dichloromethane. The database also includes one-generation oral reproductive toxicity (General Electric Company, 1976) and developmental toxicity (Narotsky and Kavlock, 1995) studies that found no reproductive or developmental effects at dose levels in the range of doses associated with liver lesions. A two-generation oral exposure study is not available; however, a two-generation inhalation exposure study by Nitschke et al. (1988a) reported no

effect on fertility index, litter size, neonatal survival, growth rates, or histopathologic lesions at exposures of ≥ 100 ppm. Neurodevelopmental outcomes were not evaluated in this study, and there have been no oral exposure studies that evaluated neurobehavioral effects in offspring. This is a relevant endpoint given the increase in blood CO (a known developmental neurotoxicant) that occurs through the CYP2E1 metabolic pathway for dichloromethane after oral and inhalation exposures. There are no oral exposure studies that include functional immune assays; however, there is a 4-week inhalation study of potential systemic immunotoxicity that found no effect of dichloromethane exposure at concentrations up to 5,000 ppm on the antibody response to sheep red blood cells (Warbrick et al., 2003). The Warbrick et al. (2003) data suggest that systemic immunosuppression is not a concern for dichloromethane exposure. Because of concern regarding the adequacy of available data pertaining to possible neurodevelopmental toxicity and the lack of a two-generation reproductive study, a UF_D of 3 was applied.

5.1.6. Previous RfD Assessment

The previous IRIS assessment derived an RfD of 0.06 mg/kg-day based on the NOAELs of 5.85 and 6.47 mg/kg-day for liver toxicity (foci/areas of cellular alteration) in male and female rats, respectively, in a 2-year drinking water study (Serota et al., 1986a). The LOAELs associated with these NOAELs were 52.58 and 58.32 mg/kg-day for males and females, respectively. The RfD of 0.06 mg/kg-day was derived by dividing the average NOAEL of 6 mg/kg-day (for male and female rats) by a UF of 100 (10 for intraspecies variability and 10 for interspecies variability).

5.1.7. RfD Comparison Information

Use of the mean value (3.99×10^{-1} mg/kg-day) of the human equivalent dose distribution instead of the 1st percentile, with an additional UF of 3 ($10^{0.5}$) to account for human toxicokinetic variability, would yield an RfD of 4×10^{-3} mg/kg-day.

Additional comparisons between the derived RfD and values developed from other endpoints or data sets using NOAEL/LOAEL methods are shown in Table 5-4 and Figure 5-4. NOAELs were used as comparison points of departure and were not scaled allometrically.

Table 5-4. Potential points of departure with applied UFs and resulting candidate RfDs

Endpoint	POD (mg/kg-d)	POD type and description	UFs applied ^a					Candidate RfD (mg/kg-d)	Reference	
			Total UF	UF _A	UF _H	UF _L	UF _S			UF _D
Alterations in liver foci, male rats^b	61.78	BMDL; 10% increase in incidence of liver lesion	30	3	3	1	1	3	7×10^{-3}	Serota et al. (1986a)
Neurological changes (FOB), female rats	101	NOAEL; No effect at POD, approximate doubling of severity score of neuromuscular and sensorimotor domains	3,000	10	10	1	10	3	3.4×10^{-2}	Moser et al. (1995)
Maternal weight gain, female rats	338	NOAEL; No effect at POD, approximate 33% decrease in weight gain seen at next dose	300	10	10	1	1	3	1.1	Narotsky and Kavlock (1995)

^aA UF for extrapolation from a LOAEL to NOAEL (UF_L) was not used for any of these studies. For the Serota et al. (1986a) study, the use of the first percentile of the human equivalent dose distribution as the POD replaces the use of a UF_H for human toxicokinetic variability.

^bBolded value is the basis for the RfD of 7×10^{-3} mg/kg-d.

POD = point of departure; UF_A = uncertainty in extrapolating from laboratory animals to humans; UF_H = uncertainty about variation from average humans to sensitive humans; UF_L = uncertainty about extrapolating from LOAEL to NOAEL; UF_S = uncertainty in extrapolating from subchronic to chronic durations; UF_D = uncertainty reflecting database deficiencies




Figure 5-4. Comparison of candidate RfDs derived from selected points of departure for endpoints presented in Table 5-4.

5.2. INHALATION REFERENCE CONCENTRATION (RfC)

5.2.1. Choice of Principal Study and Critical Effect—with Rationale and Justification

Figure 5-5 includes exposure-response arrays from some of the human studies that were evaluated for use in the derivation of the RfC. Several acute-duration controlled exposure studies (Section 4.1.2.2) and cross-sectional occupational studies (Sections 4.1.2.3 and 4.1.2.4) in humans are available that show neurological effects from dichloromethane exposure. These effects include an increase in prevalence of neurological symptoms among workers (Cherry et al., 1981) and possible detriments in attention and reaction time in complex tasks among retired workers (Lash et al., 1991). However, these studies have inadequate power for the detection of effects with an acceptable level of precision. In addition, the Cherry et al. (1981) study is limited by the definition and documentation of neurological symptom history, and the Lash et al. (1991) study has exposure measurements from 1974 to 1986, but the work histories of exposed workers go back to the 1940s. Ott et al. (1983c) reported an increase in serum bilirubin among exposed workers, but there was no association seen with respect to the other hepatic enzymes examined (serum γ -glutamyl transferase, serum AST, serum ALT), and no evidence of hepatic effects was seen in a later study of the same cohort (Soden, 1993). Because of these limitations, these human studies of chronic exposures do not serve as an adequate basis for RfC derivation. As discussed in Section 5.2.6, however, the quantitative measures of neurological function from Cherry et al. (1983) and Lash et al. (1991) were used to derive a comparative RfC.




Figure 5-5. Exposure response array for chronic (animal) or occupational (human) inhalation exposure to dichloromethane (log Y axis).

The database of experimental animal dichloromethane inhalation studies includes numerous 90-day and 2-year studies, with data on hepatic, pulmonary, and neurological effects, (see Table 4-36) and reproductive and developmental studies (Table 4-37) (see summary in Section 4.6.2). NOAELs, LOAELs, and the dose range tested corresponding to selected health effects from the chronic studies are shown in Figure 5-5, and effects seen in subchronic, reproductive, and developmental studies are shown in Figure 5-6. The subchronic (i.e., ≤ 90 -day study) data were not considered in the selection of a principal study for deriving the RfC because the database contains reliable dose-response data from the chronic study at lower doses than the 90-day study. The data from the subchronic studies are, however, used to corroborate the findings with respect to relevant endpoints (i.e., hepatic and neurological effects).




Figure 5-6. Exposure response array for subacute to subchronic inhalation exposure to dichloromethane (log Y axis).

Hepatic effects (hepatic vacuolation and necrosis, hemosiderosis, hepatocyte degeneration) are the primary dose-dependent noncancer effects associated with inhalation exposure to dichloromethane. These effects were seen in mice (Mennear et al., 1988; NTP, 1986) and rats (Mennear et al., 1988; Nitschke et al., 1988a; NTP, 1986; Burek et al., 1984) but not in Syrian golden hamsters (Burek et al., 1984). Inhalation bioassays with Sprague-Dawley rats identified the lowest inhalation LOAEL for liver lesions in the database: 500 ppm (6 hours/day, 5 days/week for 2 years) (Nitschke et al., 1988a; Burek et al., 1984); Nitschke et al. (1988a) identified a NOAEL of 200 ppm in female rats. Based on the results reviewed above, liver lesions (specifically, hepatic vacuolation) in rats are identified as the critical noncancer effect from chronic dichloromethane inhalation in animals. Because Nitschke et al. (1988a) examined a range of exposures that included doses at the low end of the range compared with the range examined in Burek et al. (1984), the former study was selected as the principal study for derivation of a chronic inhalation RfC.

Reproductive performance (e.g., as assessed by number of litters, resorption rate, fetal survival, and growth) was not affected in two generations of F344 rats exposed to up to 1,500 ppm for 14 or 17 weeks before mating of the F0 and F1 generations, respectively (Nitschke et al., 1988b), or in a study of Swiss-Webster mice or Sprague-Dawley rats exposed to 1,250 ppm on GDs 6–15 (Schwetz et al., 1975). A decrease in fertility index was seen in the 150 and 200 ppm groups in a study of male Swiss-Webster mice exposed via inhalation for 6 weeks prior to mating (Raje et al., 1988), but the statistical significance of this effect varied considerably depending on the statistical test used in this analysis. Two types of developmental effects (decreased offspring weight at birth and changed behavioral habituation of the offspring to novel environments) were seen in Long-Evans rats following exposure to 4,500 ppm for 14 days prior to mating and during gestation (or during gestation alone) (Bornschein et al., 1980; Hardin and Manson, 1980). This dose was the only exposure dose used in this study. Schwetz et al. (1975) did not observe an adverse effect on gross development or soft tissue abnormalities in a study involving exposure to 1,250 ppm on GD 6 in Swiss-Webster mice or Sprague-Dawley rats, but an increase in delayed ossification of the sternebrae was seen.

Neurological impairment was not seen in lifetime rodent bioassays involving exposure to airborne dichloromethane concentrations of $\leq 2,000$ ppm in F344 rats (Mennear et al., 1988; NTP, 1986), $\leq 3,500$ ppm in Sprague-Dawley rats (Nitschke et al., 1988a; Burek et al., 1984), or $\leq 4,000$ ppm in B6C3F₁ mice (Mennear et al., 1988; NTP, 1986). It should be noted, however, that these studies did not include standardized neurological or neurobehavioral testing. The sole subchronic or chronic study in which neurobehavioral batteries were utilized found no effects in an observational battery, a test of hind-limb grip strength, a battery of evoked potentials, or brain, spinal cord, or peripheral nerve histology in F344 rats exposed to concentrations up to 2,000 ppm for 13 weeks, with the tests performed beginning 65 hours after the last exposure (Mattsson et al., 1990).

Other effects associated with lifetime inhalation exposure to dichloromethane include renal tubular degeneration in F344 rats exposed to $\geq 2,000$ ppm, testicular atrophy in male B6C3F₁ mice exposed to 4,000 ppm, and ovarian atrophy in female B6C3F₁ mice exposed to $\geq 2,000$ ppm (Mennear et al., 1988; NTP, 1986). No effects on histologic, clinical chemistry, urinalysis, or hematologic variables were found in Syrian golden hamsters exposed to concentrations up to 3,500 ppm for 2 years, with the exception that the mean COHb percentage of exposed hamsters was about 30% compared with values of about 3% in controls (Burek et al., 1984).

5.2.2. Derivation Process for RfC Values

The derivation process used for the RfC parallels the process described in Section 5.1.2 for the RfD derivation; consideration of dose metrics was described in Section 5.1.3. As was noted in the RfD discussion, the mechanistic issues with respect to noncancer health effects represent data gaps in the understanding of the health effects of dichloromethane.

5.2.3. Methods of Analysis—Including Models (PBPK, BMD, etc.)

The modified rat PBPK model of Andersen et al. (1991), described in Appendix C and also used in the derivation of the RfD (Figure 5-2), was used for calculating internal dosimetry of inhaled dichloromethane in Sprague-Dawley rats. Simulations of 6 hours/day, 5 days/week inhalation exposures used in the Nitschke et al. (1988a) study were performed to calculate average daily internal liver doses (Table 5-5). In the absence of data for group- and sex-specific BWs, reference values for male and female Sprague-Dawley rats in chronic studies were used (U.S. EPA, 1988a).

Table 5-5. Incidence data for liver lesions (hepatic vacuolation) and internal liver doses based on various metrics in female Sprague-Dawley rats exposed to dichloromethane via inhalation for 2 years

Sex	Exposure (ppm)	Liver lesion incidence ^a	Rat internal liver dose ^b			
			CYP	GST	GST and CYP	Parent AUC
Male	0	22/70 (31)	Not modeled because results from male rats were not provided for the 50 and 200 ppm groups			
	50	Not reported				
	200	Not reported				
	500	28/70 (40)				
Female (BW = 229 g)	0	41/70 (59%)	0	0	0	0
	50	42/70 (60%)	280.3	6.3	286.6	1.2
	200	41/70 (58%)	656.5	93.2	749.7	17.8
	500	53/70 (76%) ^c	772.6	359.0	1,131.6	68.7


^aNumber affected divided by total sample size.

^bInternal doses were estimated using a rat PBPK model using exposures reported by study authors (50 ppm = 174 mg/m³, 200 ppm = 695 mg/m³, and 500 ppm = 1,737 mg/m³) and are weighted-average daily values for 1 wk of exposure at 6 hr/d, 5 d/wk. CYP dose is in units of mg dichloromethane metabolized via CYP pathway/L tissue/d; GST dose is in units of mg dichloromethane metabolized via GST pathway/L tissue/d.; GST and CYP dose is in units of mg dichloromethane metabolized via CYP and GST pathways/L tissue/d; and Parent AUC dose is in units of mg dichloromethane × hrs/L tissue.

^cSignificantly ($p < 0.05$) different from control with Fisher's exact test.

Source: Nitschke et al. (1988a).

As described in Section 5.1.2, the internal dose metric used was based on total hepatic metabolism through the CYP2E1 pathway (mg dichloromethane metabolized via CYP pathway per liter liver per day). Figure 5-7 shows the comparison between inhalation external and internal doses, using this dose metric for the rat and the human.



Average daily doses were calculated from simulated rat exposures of 6 hours/day, 5 days/week, while simulated human exposures were continuous. The human metabolism rates were estimated using a computational sample of 1,000 individuals per dose, including random samples of the three GST-T1 polymorphisms (+/+, +/-, -/-) in the current U.S. population based on data from Haber et al. (2002). Since a different set of samples was used for each dose, some stochasticity is evident as the human points (values) do not fall on smooth curves.

Figure 5-7. PBPK model-derived internal doses (mg dichloromethane metabolized via the CYP pathway per liter liver per day) in rats and humans versus external exposures (ppm).

The seven dichotomous dose-response models available in EPA BMDS version 2.0 were fit to the female rat liver lesion incidence of Nitschke et al. (1988a) and PBPK model-derived internal dose data to derive rat internal BMD_{10} and the associated $BMDL_{10}$ values (Table 5-6). The quantal model is identical to the one-stage multistage model; therefore, it is not included in this set of models. A BMR of 10% was selected because, in the absence of information regarding the magnitude of change in a response that is thought to be minimally biologically significant, a BMR of 10% is generally recommended, as it provides a consistent basis of comparison across assessments. There are no additional data to suggest that the critical response has a greater sensitivity that would warrant a lower BMR. The log-probit model was the best fitting model for the female incidence data based on AIC value among models with adequate fit. Modeling results are shown in detail in Appendix D-2.

Table 5-6. BMD modeling results for incidence of noncancer liver lesions in female Sprague-Dawley rats exposed to dichloromethane by inhalation for 2 years, based on liver specific CYP metabolism metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day)

Model ^a	BMD ₁₀	BMDL ₁₀	χ^2 goodness of fit <i>p</i> -value	AIC
Gamma ^a	614.27	225.96	0.48	367.22
Logistic	274.58	150.43	0.14	369.77
Log-logistic ^a	697.90	499.42	0.94	365.90
Multistage (3) ^a	506.94	153.13	0.25	368.53
Probit	275.49	152.52	0.14	369.75
Log-probit^{a,b}	728.96	523.94	0.98	365.82
Weibull ^a	706.45	487.45	0.95	365.87

^aThese models in EPA BMDS version 2.0 were fit to the rat dose-response data shown in Table 5-5 by using internal dose metrics calculated with the rat PBPK model. Gamma and Weibull models restrict power ≥ 1 ; Log-logistic and Log-probit models restrict to slope > 1 , multistage model restrict betas ≥ 0 ; lowest degree polynomial with an adequate fit reported (degree of polynomial in parentheses).

^bBolded model is the best-fitting model in the most sensitive sex (females), which is used in the RfC derivation.

Source: Nitschke et al. (1988a).

As with the RfD derivation, the human-equivalent internal BMDL₁₀ was obtained by dividing this rat internal dose metric by a pharmacokinetic scaling factor based on a ratio of BWs (scaling factor = 4.09) (Table 5-7). This scaling factor was used because the metric is a rate of metabolism rather than the concentration of putative toxic metabolites, and the clearance of these metabolites may be slower per volume tissue in the human compared with the rat. A probabilistic PBPK model for dichloromethane in humans, adapted from the model of David et al. (2006) as described in Appendix B, was then used with Monte Carlo sampling to calculate distributions of chronic HECs (in units of mg/m³) associated with the internal BMDL₁₀ based on the responses in female Sprague-Dawley rats. Estimated mean, first, and fifth percentiles of this distribution are shown in Table 5-7.

Table 5-7. Inhalation RfC for dichloromethane based on PBPK model-derived probability distributions of human inhalation exposure extrapolated from liver lesion data for female rats exposed via inhalation for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day)

Model ^a	Rat internal BMDL ₁₀ ^b	Human internal BMDL ₁₀ ^c	HEC (mg/m ³) ^d			Human RfC (mg/m ³) ^e
			1 st percentile	5 th percentile	Mean	
Log-probit	523.94	128.10	16.85	20.97	47.24	0.2

^aBased on the best-fitting model from Table 5-6.

^bRat dichloromethane PBPK model-derived internal liver dose associated with lower bound on 10% extra risk for developing hepatocyte vacuolation.

^cHuman dichloromethane internal liver dose, derived by dividing the rat internal BMDL₁₀ by a scaling factor of 4.09 [(BW_{human}/BW_{rat})^{0.25}] to account for potential interspecies pharmacokinetic differences in the clearance of metabolites.

^dPBPK model-derived distributions of long-term, daily average airborne dichloromethane concentrations predicted by the PBPK model to yield an internal dose in humans equal to the dichloromethane internal BMDL₁₀.

^eHuman candidate RfC, based on female rat data, derived by dividing the 1st percentile of HEC values by a total UF of 100: 3 (10^{0.5}) for possible toxicodynamic differences between species, 3 (10^{0.5}) for variability in human toxicodynamic response, and 10 for database deficiencies. The 1st percentile POD is a stable estimate of the lower end of the distribution. Use of this value in the lower tail replaces use of a UF for human toxicokinetic variability. See Section 5.2.4 for discussion of UFs.

5.2.4. RfC Derivation—Including Application of Uncertainty Factors (UFs)

The 1st percentile POD is a numerically stable estimate of the lower end of the distribution. Use of this value associated with a sensitive human population addresses the uncertainty associated with human toxicokinetic variability. The RfC was calculated by dividing the first percentile of the HEC distribution in Table 5-7 by a composite UF of 100 (3 [$10^{0.5}$] to account for uncertainty about interspecies toxicodynamic equivalence, 3 [$10^{0.5}$] to account for uncertainty about toxicodynamic variability in humans, and 10 for database deficiencies). The resulting RfC was 0.2 mg/m³ based on liver lesions in female Sprague-Dawley rats in Nitschke et al. (1988a). In deriving this RfC, factors for the following areas of uncertainty were considered:

- Uncertainty in extrapolating from laboratory animals to humans (UF_A). The use of PBPK models to extrapolate internal doses from rats to humans reduces toxicokinetic uncertainty in extrapolating from the rat liver lesion data but does not account for the possibility that humans may be more sensitive than rats to dichloromethane due to toxicodynamic differences. A default UF of 3 ($10^{0.5}$) to account for this toxicodynamic uncertainty was applied, as shown previously in Table 5-7.
- Uncertainty about variation in human toxicokinetics (UF_H). The probabilistic human PBPK model used in this assessment incorporates the best available information about variability in toxicokinetic disposition of dichloromethane in humans but does not account for humans who may be sensitive due to toxicodynamic factors. Thus, a UF of 3 ($10^{0.5}$) was applied to account for possible toxicodynamic differences in sensitive humans.
- Uncertainty in extrapolating from LOAELs to NOAELs (UF_L). A UF for extrapolation from a LOAEL to a NOAEL was not applied because BMD modeling was used to determine the POD, and this factor was addressed as one of the considerations in selecting the BMR. The BMR was selected based on the assumption that it represents a minimum biologically significant change.
- Uncertainty in extrapolating from subchronic to chronic durations (UF_S). The derived RfD is based on results from a chronic-duration drinking water toxicity study. No cross-duration UF is necessary.
- Uncertainty reflecting database deficiencies (UF_D). A UF of 10 was selected to address the deficiencies in the dichloromethane toxicity database. The inhalation database for dichloromethane includes several well-conducted chronic inhalation studies. In these chronic exposure studies, the liver was identified as the most sensitive noncancer target organ in rats (Nitschke et al., 1988a; NTP, 1986; Burek et al., 1984). The critical effect of hepatocyte vacuolation was corroborated in the two principal studies (Nitschke et al., 1988a; Burek et al., 1984), which identified 500 ppm as the lowest inhalation LOAEL for noncancer liver lesions. Gross signs of neurologic impairment were not seen in lifetime rodent inhalation bioassays for dichloromethane at exposure levels up to 4,000 ppm (see Section 4.2.2.2 for

references), and no exposure-related effects were observed in an observational battery, a test of hind-limb grip strength, a battery of evoked potentials, or histologic examinations of nervous tissues in F344 rats exposed to dichloromethane concentrations as high as 2,000 ppm (Mattson et al., 1990). A two-generation reproductive study in F344 rats reported no effect on fertility index, litter size, neonatal survival, growth rates, or histopathologic lesions at exposures ≥ 100 ppm dichloromethane (Nitschke et al., 1988b). Fertility index (measured by number of unexposed females impregnated by exposed males per total number of unexposed females mated) was reduced following inhalation exposure of male mice to 150 and 200 ppm dichloromethane 2 hours/day for 6 weeks, but the statistical significance of this effect varied considerably depending on the statistical test used in this analysis (Raje et al., 1988). The available developmental studies include single-dose studies that use relatively high exposure concentrations (1,250 ppm in Schwetz et al. [1975]; 4,500 ppm in Hardin and Manson [1980]; and 4,500 ppm in Bornschein et al. [1980]). In one of the single-dose studies, decreased offspring weight at birth and changed behavioral habituation of the offspring to novel environments were seen following exposure of adult Long-Evans rats to 4,500 ppm for 14 days prior to mating and during gestation (or during gestation alone) (Bornschein et al., 1980; Hardin and Manson, 1980). CO, a known developmental neurotoxicant, is produced through the CYP2E1 metabolic pathway for dichloromethane. Schwetz et al. (1975) reported increased concentrations (~10% higher compared with controls) in maternal blood COHb levels in mice and rats exposed during GDs 6–15. A chronic exposure study in F344 rats reported a dose-related increase in blood COHb in females exposed to 50, 200, and 500 ppm, beginning with the first measure taken after 6 months of exposure (Nitschke et al., 1988a). The increase was seen at the lowest exposure group (50 ppm). Anders and Sunram (1982) reported elevated CO levels in maternal and fetal blood in rats following exposure to 500 ppm for 1 hour on GD 21; levels were similar in the maternal and fetal samples. Placental transfer of dichloromethane was also seen, although levels were lower in the fetus. The results from the single dose developmental toxicity study in rats (Bornschein et al., 1980; Hardin and Manson, 1980), in addition to the known increase in CO, the placental transfer of dichloromethane, and the relatively high activity of CYP2E1 in the brain compared to the liver of the developing human fetus (Hines, 2007; Johnsrud et al., 2003; Brzezinski et al., 1999), raise uncertainty regarding possible neurodevelopmental toxicity from gestational exposure to inhaled dichloromethane. In addition, Aranyi et al. (1986) demonstrated evidence of immunosuppression following a single 100 ppm dichloromethane exposure for 3 hours in CD-1 mice. This study used a functional immune assay that is directly relevant to humans (i.e., increased risk of Streptococcal pneumonia-related mortality and decreased clearance of Klebsiella bacteria). No effects were seen with 50 ppm exposure for either 1 or 5 days. Systemic immunosuppression was not seen in a 4-week, 5,000 ppm inhalation exposure study measuring the antibody response to sheep red blood cells in Sprague-Dawley rats (Warbrick et al., 2003). These studies suggest a localized, portal-of-entry effect within the lung rather than a systemic immunosuppression. Therefore, in consideration of the entire database for dichloromethane, a database UF of 10 was selected. This UF accounts for the lack of neurodevelopmental toxicity studies and lack of adequate immunotoxicity and developmental studies following inhalation exposure at low concentrations.

5.2.5. Previous RfC Assessment

No RfC was derived in the previous IRIS assessment.

5.2.6. RfC Comparison Information

A candidate RfC, based on a different approach to accounting for human toxicokinetic variability is similar to the derived RfC of 0.2 mg/m³. Use of the mean value on the HEC distribution (47.24) with an additional UF of 3 (10^{0.5}) to account for human toxicokinetic variability would yield an RfC of 0.2 mg/m³.

For an additional comparison, an RfC was derived based on neurological endpoints from human occupational exposures. Cherry et al. (1983) compared 56 exposed and 36 unexposed workers at an acetate film manufacturing plant for dichloromethane inhalation exposure, blood levels of dichloromethane, subjective self-reporting of general health, and two objective, quantitative measurements of neurological function (digit symbol substitution and simple reaction time). The exposed and unexposed individuals were matched to within 3 years of age. The measured dichloromethane concentrations from personal breathing zone sampling of the exposed workers ranged from 28 to 173 ppm. No information on exposure duration was given, and Cherry et al. (1983) did not indicate if the exposure measurements were indicative of historical exposure levels. There were no significant differences between exposed and unexposed workers in subjective or objective measurements collected at the beginning of the work shift on a Monday (after 2 nonworking days). Exposed workers showed a slightly slower (but not statistically significant) score than the control workers on a reaction time test, but the scores did not deteriorate during the shift. These findings suggest that repeated inhalation exposures in the range of 28–173 ppm do not result in significant effects, but the actual duration of exposure of the workers is uncertain. In the absence of data for the mean exposure levels, the exposure range midpoint of 101 ppm serves as a NOAEL for chronic neurological effects from dichloromethane exposure. Thus, a candidate RfC of 3.5 mg/m³ was derived by dividing the NOAEL of 351 mg/m³ (101 ppm) by a composite UF of 100. A UF of 10 was applied to account for potentially susceptible individuals in the absence of quantitative information on the variability of neurological response to dichloromethane in the human population. A UF of 10 was applied for database deficiencies. The duration of exposures of acetate film workers (Cherry et al., 1983) was not reported, and a limited number of endpoints was evaluated. Further, definitive neurological batteries were not administered in chronic-duration animal bioassays.

Another candidate RfC was developed by using the neurological data from the study by potential long-term CNS effects in a study of retired aircraft maintenance workers (Lash et al., 1991). Retired aircraft maintenance workers, ages 55–75 years, employed in at least 1 of 14 targeted jobs (e.g., paint strippers) with dichloromethane exposure for ≥6 years between 1970 and 1984 (n = 25) were compared to a like group of workers without dichloromethane exposure (n = 21). From 1974 to 1986, when 155 measurements for dichloromethane exposure

were made, mean breathing zone TWAs ranged from 82 to 236 ppm and averaged 225 ppm for painters and 100 ppm for mechanics; information on exposure levels prior to this time was not provided although the work histories of exposed workers goes back to the 1940s. The evaluation included several standard neurological tests, including physiological measurement of odor and color vision senses, auditory response potential, hand grip strength, measures of reaction time (simple, choice, and complex), short-term visual memory and visual retention, attention, and spatial ability. The exposed group had a higher score on verbal memory tasks (effect size approximately 0.45, $p = 0.11$) and lower score on attention tasks (effect size approximately -0.55, $p = 0.08$) and complex reaction time (effect size approximately -0.40, $p = 0.18$) compared with the control group. None of these differences were statistically significant. Given the sample size, however, the power to detect a statistically significant difference between the groups was very low (i.e., approximately 0.30 for an effect size of 0.40 using a two-tailed alpha of 0.05) (Cohen, 1987), and these results cannot be taken as evidence of no effect. An estimated exposure level from the study can be generated from the midpoint value from the exposure range (82–236 ppm; mean = 159 ppm), converted to 552 mg/m^3 . If these results are viewed as a LOAEL and this estimated mean exposure level of 552 mg/m^3 was used, a composite UF of 1,000 would be applied for interspecies toxicodynamics (10), extrapolation from a LOAEL to a NOAEL (10), and database uncertainties (10), resulting in an RfC of 0.55 mg/m^3 .

The value of the candidate RfC based on the data from Cherry et al. (1983), 3.5 mg/m^3 , is approximately 15-fold higher, and the value of the candidate RfC based on the data from Lash et al. (1991), 0.55 mg/m^3 , is approximately 3 times higher than the derived RfC of 0.2 mg/m^3 based on liver lesions in rats. The animal-derived RfC is preferable to the human-derived RfC because of the uncertainties about the exposure durations, statistical analysis, and statistical power in Cherry et al. (1983) and the uncertainties regarding the exposure levels, effect sizes, and statistical power in Lash et al. (1991), and because the RfC based on the rat data is more health protective.

Additional comparisons among the RfC and candidate values developed from other endpoints or data sets using NOAEL/LOAEL methods are shown in Table 5-8 and Figure 5-8.

Table 5-8. Potential points of departure with applied UFs and resulting candidate RfCs

Endpoint	POD (mg/m ³) ^a	POD Type and Description ^b	UFs ^c						RfC (mg/m ³)	Reference
			Total UF	UF _A	UF _H	UF _L	UF _S	UF _D		
Hepatocyte vacuolation, female rat^d	523	BMDL, 10% increase in incidence of liver lesion	100	3	3	1	1	10	0.2	Nitschke et al. (1988a)
Renal tubular degeneration; NOAEL, male rat	620	NOAEL	1,000	3	10	1	1	10	2.07	Mennear et al. (1988); NTP (1986)
Reproductive - fertility index; NOAEL, male mouse	20.7	No effect at POD, 16% decrease in fertility index seen at LOAEL dose	300	3	10	1	1	10	0.071	Raje et al. (1988)
Increased infection susceptibility (mortality risk), female mouse	15.5	NOAEL	1,000	3	10	1	10	3	0.015	Aranyi et al. (1986)
Increased IgM production, male and female rat	17,366	NOAEL	3,000	3	10	1	10	10	1.03	Warbrick et al. (2003)
Chronic CNS effects, human male	351	NOAEL	100	1	10	1	1	10	3.51	Cherry et al. (1983)
CNS changes, human male	552	LOAEL	1,000	1	10	10	1	10	0.55	Lash et al. (1991)

^aPOD = point of departure. For Nitschke et al. (1988a), this is based on BMD modeling of a 10% increase in liver lesions using internal liver dose metric (mg dichloromethane metabolism via CYP pathway per liter liver tissue per d) derived from a rat PBPK model. After an allometric scaling factor of 4.09 was applied, the human internal BMDL₁₀ was 128 mg/m³. A probabilistic human PBPK model adapted from David et al. (2006) was used to generate a distribution of HECs from the human internal BMDL₁₀ and the first percentile of this distribution was used as the POD. For other rodent studies, the NOAEL or LOAEL concentration, in mg/m³, was adjusted to a continuous exposure taking into account hrs/d and d/wk of exposure. This adjusted exposure was then converted to an HEC by multiplying the value by a dosimetric adjustment factor (DAF). PBs were 8.24 for humans, 19.8 for rats, and 23 for mice. Since the PBs for both the mice and rats were greater than for humans, a DAF of 1 is recommended and was used. NOAELs or LOAELs were used as points of departure in human studies since the concentrations were already human exposures.

^bExtra risk defined for incidence data as $(\text{Incidence}_1 - \text{Incidence}_0) / (1 - \text{Incidence}_0)$, where 1 = dose at observed increased and 0 = background incidence.

^cUF_A = uncertainty in extrapolating from laboratory animals to humans, UF_H = uncertainty about variation from average humans to sensitive humans, UF_L = uncertainty about extrapolating from LOAEL to NOAEL, and UF_D = uncertainty reflecting incompleteness of the overall database. A UF extrapolating from subchronic to chronic durations (UF_S) was not used for any of these studies.

^dBolded value is the basis of the RfC of 0.2 mg/m³.




Figure 5-8. Comparison of candidate RfCs derived from selected points of departure for endpoints presented in Table 5-8.

5.3. UNCERTAINTIES IN THE ORAL REFERENCE DOSE AND INHALATION REFERENCE CONCENTRATION

Risk assessments need to include a discussion of uncertainties associated with the derived toxicity values. For dichloromethane, uncertainties related to inter- and intraspecies differences in toxicodynamics and database deficiencies are treated quantitatively via the UF approach (U.S. EPA, 1994b). Uncertainties in the toxicokinetic differences of dichloromethane between species and within humans are reduced by application of the PBPK models for rats and humans. These and other areas of uncertainty of the derived RfD and RfC are discussed below.

Adequacy of database for derivation of RfD and RfC. As summarized in Sections 4.6.1.1 and 4.6.2.1, data from the available human studies on the health effects from occupational inhalation exposures provide some but not conclusive evidence of long-term health consequences of chronic dichloromethane exposure, specifically with respect to neurologic and hepatic damage. These data are not adequate for derivation of an RfD or RfC. However, a broad range of animal toxicology data is available for the hazard assessment of dichloromethane, as described in Section 4. The database of oral (Table 4-35) and inhalation (Tables 4-36 and 4-37) toxicity studies includes numerous chronic, subchronic, acute, reproductive, and developmental studies. Liver toxicity in multiple rodent species is consistently identified as the most sensitive noncancer effect from oral and inhalation exposure to dichloromethane. In addition to the oral and inhalation toxicity data, there are numerous studies describing the toxicokinetics of dichloromethane. Consideration of the available dose-response data to determine an estimate of oral exposure that is likely to be without an appreciable risk of adverse noncancer health effects over a lifetime has led to the selection of noncancer liver lesions in the 2-year drinking water study in F344 rats (Serota et al., 1986a) as the critical effect and principal study for deriving the RfD for dichloromethane. The critical effect selected for the derivation of the chronic RfC is also hepatic lesions; two different studies in Sprague-Dawley rats (Nitschke et al., 1988a; Burek et al., 1984) spanning overlapping exposures reported data on hepatic vacuolation, and the lower exposure study was chosen as the principal study (Nitschke et al., 1988a).

A critical data uncertainty was identified for neurodevelopmental effects. Animal bioassays have not identified gross or microscopic effects on neural tissues from long-term exposures or single (Schwetz et al., 1975) or multigenerational (Nitschke et al., 1988b) developmental toxicity studies. However, behavioral changes were observed in pups born to rats exposed to high levels (4,500 ppm) of dichloromethane (Bornschein et al., 1980; Hardin and Manson, 1980); lower exposures were not examined in this study. Uncertainty exists as to the development of neurological effects from lower gestational exposures in animals or humans. In addition, a critical data uncertainty has been identified that relates to potential immunotoxicity, specifically immunosuppression seen as a localized portal-of-entry effect within the lung with an acute inhalation exposure (Aranyi et al., 1986). The lack of data on immune effects from longer-

term exposure represents a significant data gap and is of particular importance because of the potential importance of immunosuppression with respect to response to infections and tumor surveillance. The weight of evidence for noncancer effects in humans and animals suggests that the development of liver lesions is the most sensitive effect, with a UF applied because of the lack of reproductive and neurodevelopmental studies for the RfD and, for the RfC, the uncertainty regarding developmental, neurodevelopmental, and immune system toxicity.

Dose-response modeling. The selection of the BMD model(s) for the quantitation of the RfD and RfC does not lead to significant uncertainty in estimating the POD. It should be noted, however, that a level of uncertainty is inherent given the lack of data in the region of the BMR.

Interspecies extrapolation of dosimetry and risk. The extrapolation of internal dichloromethane dosimetry from liver lesions in rats to human risk was accomplished using PBPK models for dichloromethane in rats and humans. Uncertainties in rat and human dosimetry used for RfD and RfC derivation can arise from uncertainties in the PBPK models with regard to accurately simulating the toxicokinetics of dichloromethane for animals under bioassay conditions and humans experiencing relatively low, chronic environmental exposures. Specific uncertainties regarding the model structure are described in detail in Section 3.5.5. A structural uncertainty previously discussed arises from the indication by various data that the standard Michaelis-Menten equation used in the existing model may not accurately describe the CYP2E1-catalyzed oxidation of dichloromethane. An alternate equation described by Korzekwa et al. (1998) may better represent CYP2E1-induced oxidation of dichloromethane, which would lead to a higher fraction of total dichloromethane predicted to be metabolized by CYP2E1 at higher dichloromethane doses (or exposures). Since this shift in predicted metabolism would occur for both the human and rodent PBPK models, if the alternate equation was applied, it is difficult to estimate the net impact of using this equation on risk predictions. As described in Section 3.5.5 and Appendix C, the error in the ratio of GST:CYP metabolism at low concentrations appears to be less than 13% based on comparison of model predictions to CO metabolism data. Further, analysis of the GST-mediated metabolism of dichloromethane measured by Reitz et al. (1989) shows that those results are within a factor of three of the GST kinetic parameters used in the current PBPK model, indicating that any error in the GST:CYP balance is no greater than that.

Also as discussed in detail in Section 3.5.2, there appears to be inconsistency in the numerical results of David et al. (2006) for the liver GST activity (coefficient), k_{fC} , between that obtained for each published data set when analyzed separately and that obtained for the combined data set. Since the numerical average of the mean k_{fC} values for the four data sets included in the combined data set was 12.4 and the upper bound was 12, the impact of using an intermediate value of k_{fC} , specifically the DiVincenzo and Kaplan value of $5.87 \text{ kg}^{0.3}/\text{hour}$ was

explored. Changing only the k_{fC} is not realistic since the dichloromethane data effectively define total metabolism (sum of CYP and GST pathways), and there is naturally a negative correlation between the predicted CYP metabolic rate and the GST metabolic rate required to describe this total. Therefore, it would be inconsistent with the dichloromethane data to increase k_{fC} without adjusting the CYP metabolic rate downward and likewise all other parameters. Therefore, for consistency, the distributions for all of the fitted parameters were rescaled by the ratio of the mean for DiVincenzo and Kaplan (1981) to the mean for the combined data set (e.g., the distribution for k_{fC} was multiplied by 5.87/0.852, the ratio of the two posterior means). The resulting HEC and HED calculations increased by 10–30% for the mixed GST-T1 population, depending on the route of exposure and distribution statistic compared. Thus the impact of this model uncertainty appears to be modest for the noncancer assessment.

The dose metric used in the models is the rate of metabolism to a putative toxic metabolite rather than an average or AUC of the metabolite concentration, so the model specifically fails to account for rodent-human differences in clearance or removal of the toxic metabolite. Therefore, a scaling factor based on BW ratios was used to account for this difference.

Sensitivity analysis of rat model parameters. The rat model was modified and utilized in a deterministic manner. Data were not available to perform a hierarchical Bayesian calibration in the rat. Thus, uncertainties in the rat model predictions had to be assessed qualitatively. To address these uncertainties, a sensitivity analysis was conducted to determine which model parameters most influence the predictions for a given dose metric and exposure scenario.

Sensitivity is a measure of the degree to which a given model output variable (i.e., dose metric) is influenced by perturbation in the value of model parameters. The approach implemented was a univariate analysis in which the value of an individual model parameter was perturbed by an amount (Δ) in the forward and reverse direction (i.e., an increase and decrease from the nominal value), and the change in the output variable was determined. Sensitivity coefficients were calculated as follows:

$$f'(x) \approx \frac{f(x + \Delta x) - f(x)}{\Delta x} \cdot \frac{x}{f(x)} \quad (\text{Eq. 5-1})$$

where x is the model parameter, $f(x)$ is the output variable, Δx is the perturbation of the parameter from the nominal value, and $f'(x)$ is the sensitivity coefficient. In equation 5-1, the sensitivity coefficients are scaled to the nominal value of x and $f(x)$ to eliminate the potential effect of units of expression. Therefore, the sensitivity coefficient is a measure of the proportional (unitless) change in the output variable produced by proportional change in the

parameter value. Parameters that have higher sensitivity coefficients have greater influence on the output variable. They are considered more sensitive than parameters with lower values. The results of the sensitivity analysis are useful for assessing uncertainty in model predictions, based on the level of confidence or uncertainty in the model parameter(s) to which the dose metric is most sensitive.

Sensitivity coefficients for the noncancer dose metric (mg dichloromethane metabolized via CYP-mediated pathway per liter liver per day) were determined for each of the model parameters. Sensitivity analyses for both oral and inhalation exposures were performed. The exposure conditions were set to be near or just below the lowest bioassay exposure resulting in significant increases in the critical effect.

For the CYP-mediated metabolism from oral exposure, the liver volume (VLC) and slowly perfused tissue volume (VSC) parameters exert the largest influence (Figure 5-9). The high influence of these two parameters was due to the fact that the dose metric is a tissue-specific rate of metabolism, the majority of CYP metabolism is attributed to the liver, and changes in liver volume have a greater impact on the total CYP metabolism than the individual $V_{\max C}$ value. For inhalation exposures $V_{\max C}$, VLC, and VSC have the highest sensitivity coefficients (Figure 5-10). The physiological parameters (VLC and VSC) are known with a high degree of confidence (Brown et al., 1997). $V_{\max C}$ for the rat was estimated by fitting to the pharmacokinetic data as described in Chapter 3 and Appendix C, subject to model structure/equation uncertainties as detailed above, and hence is known with less certainty than the physiological parameters. That total exhaled CO, which is specific to the CYP pathway, is within 50% of measured levels (Figure C-8, panel C), however, provides a similar level of confidence in the balance between CYP and GST pathways predicted by the rat PBPK model.





Figure 5-9. Sensitivity coefficients for long-term mass CYP- and GST-mediated metabolites per liver volume from a daily drinking water concentration of 10 mg/L in rats.



(KA is not included since it has no impact on inhalation dosimetry.)

Figure 5-10. Sensitivity coefficients for long-term mass CYP- and GST-mediated metabolites per liver volume from a long-term average daily inhalation concentration of 500 ppm in rats.

In summary, the uncertainties associated with use of the rat PBPK model should not markedly affect the values (i.e., an effect of no more than 30%) of the RfD and RfC based on the metrics considered. An additional uncertainty results from the lack of knowledge concerning the most relevant dose metric (e.g., a specific metabolite) for the noncancer endpoints considered. This basic research question represents a data gap. This uncertainty was addressed by


considering different dose metrics (CYP metabolism alone, GST metabolism alone, sum of GST and CYP, and the AUC of the parent compound). The GST metabolism and the AUC dose metrics did not present reasonable choices based on model fit and consistency of response across studies at comparable dose levels. Given these results, the combination of hepatic metabolism through the GST and the CYP pathways would not be expected to result in an improvement to a metric based only on CYP metabolism. The CYP-metabolism dose metric seems to be most consistent with the data, and so is the metric chosen for the RfD and RfC derivations.

Sensitive human populations. The potential for sensitivity to dichloromethane in a portion of the human population due to pharmacokinetic differences was addressed quantitatively by using a human probabilistic PBPK model, as modified by EPA, to generate distributions of human exposures likely to result in a specified internal BMDL₁₀. The model and resulting distributions take into account the known nonchemical-specific variability in human physiology as well as total variability and uncertainty in dichloromethane-specific metabolic capability. The first percentile values of the distributions of human equivalent doses (Table 5-3) and HECs (Tables 5-7) served as points of departure for candidate RfDs and RfCs, respectively, to protect toxicokinetically sensitive individuals. Selection of the first percentile allows generation of a numerically stable estimate for the lower end of the distribution. The mean value of the human equivalent oral dose in Table 5-3 was about twofold higher than the corresponding first percentile values, and the mean value of human equivalent inhalation concentration in Table 5-7 was approximately threefold higher than the first percentile value. The internal dose metric in the analyses described in these tables was the mg dichloromethane metabolized via the CYP pathway per liter liver per day, and thus the comparisons of the first percentile and mean values give estimates of the amount of variability in the population to metabolize dichloromethane by the CYP metabolic pathways on a liver-specific basis. The mean:1st percentile ratios for these distributions is attributed to the dependence of the dose metric on hepatic blood flow rate (metabolism being flow-limited). This blood flow is expected to be highly and tightly correlated with liver volume, resulting in very similar delivery of dichloromethane per volume liver across the population. The mean:1st percentile ratio for the oral distribution is 1.85, which is less than the default intra-human toxicokinetic UF of 3. The population-structured distributions for physiological parameters and broadened distributions for metabolic parameters used here provide a good degree of confidence that the population variability has not been underestimated.

The internal dose metric used in the RfD and RfC derivations was based on the rate of CYP metabolism. GST-T1 polymorphisms could affect this rate, as the GST-T1 null genotype would be expected to result in an increase in the metabolism through the CYP pathway, resulting in a greater sensitivity to a CYP-related effect. The effect of GST variability on the RfD and RfC values was examined by comparing results obtained specifically for the GST-T1 null

genotype to those obtained for the population of mixed genotypes. The values for human equivalent doses and HECs were very similar for these two groups (e.g., mean HEC 47.36 and 47.49 for the mixed and the GST-T1^{-/-} null genotypes, respectively; 1st percentile HEC 16.63 and 16.69 for the mixed and the GST-T1^{-/-} null genotypes, respectively), and use of this population would not result in a change in the recommended RfD or RfC.

As a further level of sensitivity analysis, model predictions of the human equivalent dose for the general population, as listed in Table 5-3 (estimates covered 0.5- to 80-year-old male and female individuals), were compared to three subpopulations: 1-year-old children (males and females), 70-year-old men, and 70-year-old women. For the general population and each subpopulation, a Monte Carlo simulation representing 10,000 individuals was conducted, and histograms of the resulting distribution of human equivalent doses are shown in Figure 5-11, with corresponding statistics in Table 5-9. All groups used in these comparisons were limited to the GST-T1^{-/-}.



All groups were restricted to the GST-T1^{-/-} population.

Figure 5-11. Frequency density of human equivalent doses in specific populations in comparison to a general population (0.5- to 80-year-old males and females) estimate for an internal dose of 15.1 mg dichloromethane metabolized by CYP per liter liver per day.

Table 5-9. Statistical characteristics of human equivalent doses in specific populations of the GST-T1^{-/-} group


Population	Human equivalent dose (mg/kg-d) ^a		
	Mean	5 th percentile	1 st percentile
All ages ^b	3.98×10^{-1}	2.53×10^{-1}	2.14×10^{-1}
1-yr-old children	6.35×10^{-1}	4.88×10^{-1}	4.52×10^{-1}
70-yr-old men	3.20×10^{-1}	2.48×10^{-1}	2.30×10^{-1}
70-yr-old women	2.64×10^{-1}	2.02×10^{-1}	1.84×10^{-1}

^aExposure levels predicted to result in 15.1 mg dichloromethane metabolized via CYP pathway per liter liver per d (based on BMDL₁₀ from the best-fitting model from Table 5-2; human dichloromethane internal liver dose, derived by dividing the rat internal BMDL₁₀ by a scaling factor of 4.09 [(BW_{human}/BW_{rat})^{0.25}] to account for potential interspecies pharmacokinetic differences in the clearance of metabolites).

^b0.5- to 80-yr-old males and females.

The results shown above for differences in human equivalent dose values in different populations are qualitatively what would be expected: a relatively broad distribution for the general population with specific populations representing narrower components of that distribution. There are some differences between men and women at 70 years of age, but neither of these would be greatly misrepresented by the general population estimate. While 1-year-old children represent more of a distinct tail in the general population, in this case, the distribution of HECs in the general estimate is lower than that seen in what would otherwise be considered a more sensitive population. This difference most likely results from the higher specific respiration rate in children versus adults, which allows them to eliminate more of orally ingested dichloromethane by exhalation, leading to lower internal metabolized doses.

A similar comparison was made for inhalation HEC values, as shown in Figure 5-12 and Table 5-10. For HEC values, the distributions for 70-year-old men and women are both virtually indistinguishable from the general population, and while 1-year-old children are clearly distinct, they are less different than in the human equivalent dose comparison and, in this case, are more sensitive than the population in general. As described in detail in Appendix B, the allometric alveolar ventilation constant (QAlvC) is about 28 L/hour·kg^{0.75} in a 1-year-old child but averages around 14 L/hour·kg^{0.75} in an adult. Combining this with the difference between a BW of 10 kg in that child and 70 kg in an “average” adult, the respiration rate per kg BW is about threefold higher in the child versus adult. As noted above, for oral exposures, this leads to faster elimination by respiration in children, while for inhalation exposures it leads to higher uptake for a given air concentration.



All groups restricted to the GST-T1^{-/-} population.

Figure 5-12. Frequency density of HECs in specific populations in comparison to a general population (0.5- to 80-year-old males and females) estimate for an internal dose of 128.1 mg dichloromethane metabolized by CYP per liter liver per day.

Table 5-10. Statistical characteristics of HECs in specific populations of the GST-T1^{-/-} group

Population	HEC (mg/m ³) ^a		
	Mean	5 th percentile	1 st percentile
All ages ^b	47.7	21.0	16.8
1-yr-old children	24.8	14.3	12.2
70-yr-old men	46.6	21.6	17.7
70-yr-old women	50.1	22.1	18.1

^aExposure levels predicted to result in 128.1 mg dichloromethane metabolized via CYP pathway per liter liver per d (based on BMDL₁₀ from the best-fitting model from Table 5-6; Human dichloromethane internal liver dose, derived by dividing the rat internal BMDL₁₀ by a scaling factor of 4.09 [(BW_{human}/BW_{rat})^{0.25}] to account for potential interspecies pharmacokinetic differences in the clearance of metabolites).

^b0.5- to 80-yr-old males and females.

The lack of difference in elderly adults versus the general population in HEC values is likely due to the fact that the rate of exposure and rates of metabolism (the latter being the key dose metric) both scale as BW^{0.75}, with the scaling coefficients being either similar (respiration) or identical (metabolism) among adults who comprise the majority of the population. For oral exposures, the exposure rate is normalized to total BW and scales as BW¹, while elimination routes increase as BW^{0.75}. Moreover, oral exposures are simulated as occurring in a series of bolus exposures (drinking episodes) during the day, and the higher body-fat content occurring in the elderly (see Appendix B) means that such a dose that might saturate metabolism and therefore have a higher fraction exhaled in a leaner individual will tend to be more sequestered in fat and slowly released, resulting in a higher fraction metabolized (less saturation of metabolism) in a more obese individual. The difference among adults of different ages for dosimetry from oral ingestion (bolus exposure) will be greater than the difference for inhalation exposures. More careful examination of Figure 5-12 shows that the distribution for 70-year-old women, for whom the fat fraction is estimated to be greatest, has a lower peak and higher upper tail than for the general population. Thus, the physiological differences have some impact that is qualitatively consistent with what is seen from oral exposure, given the mechanistic considerations described here. But the impact of those differences is far less for inhalation exposure.

No data are available regarding toxicodynamic differences within a human population. Therefore, a UF of 3 for possible differences in human toxicodynamic responses is intended to be protective for sensitive individuals.

5.4. CANCER ASSESSMENT

5.4.1. Cancer OSF

5.4.1.1. Choice of Study/Data—with Rationale and Justification

No human data are available for the quantification of potential neoplastic effects from oral exposures to dichloromethane. In the only chronic (2-year) oral exposure cancer bioassay, significant increases in the incidence of liver adenomas and carcinomas were observed in male B6C3F₁ mice exposed by drinking water, with incidence rates of 19, 26, 30, 31, and 28% in groups with estimated mean intakes of 0, 61, 124, 177, and 234 mg/kg-day, respectively (trend p -value = 0.058) (Table 4-38) (Serota et al., 1986b; Hazleton Laboratories, 1983). Incidences of liver tumors in female mice were not presented in the summary reports, but it was reported that exposed female mice did not show increased incidences of proliferative hepatocellular lesions (Serota et al., 1986b; Hazleton Laboratories, 1983). Evidence of a trend for increased risk of liver tumors (described as neoplastic nodule or hepatocellular carcinoma) was seen in female F344 rats but not males exposed via drinking water ($p < 0.01$) (Serota et al., 1986a). However, the potential malignant characterization of the nodules was not described, and no trend was seen in the data limited to hepatocellular carcinomas.

The derivation of the cancer OSF is based on the male mouse data (Serota et al., 1986b; Hazleton Laboratories, 1983) because of their greater sensitivity compared to female mice and to male and female rats. (The trend p -value and pairwise test p -values were not given in the Serota et al. [1986b] paper but can be found in the full report [Hazleton Laboratories, 1983]). The study authors concluded that there was no dose-related trend and that there were no significant differences comparing the individual dose groups with the combined control group, and that the observed incidences were “within the normal fluctuation of this type of tumor incidence.” Although Serota et al. (1986b) state that a two-tailed significance level of $p = 0.05$ was used for all tests, Hazleton Laboratories (1983) indicated that a correction factor for multiple comparisons was used specifically for the liver cancer data, reducing the nominal p -value from 0.05 to 0.0125; none of these individual group comparisons are statistically significant when a p -value of 0.0125 is used.

Based on the Hazleton Laboratories (1983) statistical analysis, EPA concluded that dichloromethane induced a carcinogenic response in male B6C3F₁ mice as evidenced by a marginally increased trend test ($p = 0.058$) for combined hepatocellular adenomas and carcinomas, and by small but statistically significant ($p < 0.05$) increases in hepatocellular adenomas and carcinomas at dose levels of 125 ($p=0.023$), 185 ($p=0.019$), and 250 mg/kg-day ($p = 0.036$). EPA did not consider the use of a multiple comparisons correction factor for the evaluation of the liver tumor data (a primary a priori hypothesis) to be warranted.

With respect to comparisons with historical controls, the incidence in the control groups (19%) was almost identical to the mean seen in the historical controls from this laboratory (17.8% based on 354 male B6C3F₁ mice), so there is no indication that the observed trend is

being driven by an artificially low rate in controls and no indication that the experimental conditions resulted in a systematic increase in the incidence of hepatocellular adenomas and carcinomas. Although the occurrence of one elevated rate in an exposed group may reflect normal fluctuations in the incidence of these tumors (described for this laboratory as 5–40%, with a mean of 17.8%, based on 354 male controls), the pattern of incidence rates (increased incidence in all four dose groups, with three of these increases significant at a p -value of < 0.05) suggest a treatment-related increase.

The development of liver tumors in B6C3F₁ mice is associated with metabolite production in this tissue via the GST metabolic pathway (Section 4.7.3), a pathway that also exists in humans. Modeling intake, metabolism, and elimination of dichloromethane in mice and humans is feasible. Thus, it is reasonable to apply the best available PBPK models to estimate equivalent internal doses in mice and humans.

5.4.1.2. Derivation of OSF

In a manner similar to the derivation of the noncancer toxicity values, PBPK models for dichloromethane in mice and humans were used in the derivation of toxicity values (cancer OSF and IUR) for cancer endpoints based on lung (for inhalation) and liver (for oral and inhalation) tumor data in the mouse (Figure 5-13). A deterministic PBPK model for dichloromethane in mice was first used to convert mouse drinking water or inhalation exposures to long-term daily average values of internal lung-specific GST metabolism (GST metabolism in lung/lung volume) or liver-specific GST metabolism (GST metabolism in liver/liver volume). The choice of these dose metrics was made based on data pertaining to the mechanism(s) involved in the carcinogenic response, specifically data supporting the involvement of a GST metabolite(s). The evidence pertaining to the GST pathway is discussed in Section 4.7 and includes the enhanced genotoxicity seen in bacterial and mammalian in vitro assays with the introduction of GST metabolic capacity (Graves et al., 1994a) and the suppression of the production of DNA SSBs by pretreatment with a GSH depletory seen in acute inhalation exposure to dichloromethane in mice (Graves et al., 1995). Although the GST metabolic pathway takes on a greater role as the CYP pathway is saturated, both the GST and CYP pathways are operating even at low exposures. The PBPK model incorporates the metabolic shift and expected nonlinearity (GST dose attenuation with low exposures) in the exposure-dose relationship across exposure levels.




Figure 5-13. Process for deriving cancer OSFs and IURs by using rodent and human PBPK models.

The multistage cancer model (using BMDS version 2.0) was fit to the tumor incidence data and internal dose data for rodents, and BMD₁₀ and associated BMDL₁₀ values (for a BMR of 10% extra risk) were estimated. A probabilistic PBPK model for dichloromethane in humans, adapted from David et al. (2006) (see Appendix B), was used with Monte Carlo sampling to calculate distributions of internal lung or liver doses associated with chronic unit oral (1 mg/kg-day) or inhalation (1 µg/m³) exposures. The resulting distribution of human internal doses was multiplied by a human internal dose tumor risk factor (in units of reciprocal internal dose) to generate a distribution of OSFs or IURs associated with a chronic unit oral or inhalation exposure, respectively.

As discussed in Section 3.5.2, the statistics reported for the fitted metabolic parameters by David et al. (2006; Table 4 in that publication) only represent the population mean and uncertainty in that mean for each parameter. EPA's revision of the model parameter distributions are generally described in Section 5.1.2, with details provided in Appendix B. Subsequent to this revision, the human PBPK parameter distributions are expected to appropriately account for both parametric uncertainty and interindividual variability, with sampling weighted to represent the full population from 6 months to 80 years of age. The model code also allows estimation of risk for subpopulations defined by a specific age in that range, gender, and/or GST-T1 genotype (e.g., the GST-T1 +/+ subpopulation).

5.4.1.3. Dose-Response Data

Data for liver tumors in male B6C3F₁ mice following exposure to dichloromethane in drinking water were used to develop oral cancer slope factors (Serota et al., 1986b; Hazleton Laboratories, 1983). Significant increases in incidence of liver adenomas and carcinomas were observed in male but not female B6C3F₁ mice exposed for 2 years (Table 5-11). No significant decreases in survival were observed in the treated groups of either sex compared with controls. The at-risk study populations (represented by the denominators in the incidence data) were determined by excluding all animals dying prior to 52 weeks.

Table 5-11. Incidence data for liver tumors and internal liver doses, based on GST metabolism dose metrics in male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years

Sex	Nominal (actual) daily intake (mg/kg-d)	Mouse liver tumor incidence ^a	Mouse internal liver metabolism dose ^b	Mouse whole body metabolism dose ^c
Male (BW = 37.3 g)	0 (0)	24/125 (19%)	0	0
	60 (61)	51/199 (26%)	17.5	0.73
	125 (124)	30/99 (30%)	63.3	2.65
	185 (177)	31/98 (32%)	112.0	4.68
	250 (234)	35/123 (28%)	169.5	7.1

^aHepatocellular carcinoma or adenoma, combined. Mice dying prior to 52 wks, as estimated from the survival data shown in Figure 1 of Hazleton Laboratories (1983), were excluded from the denominators. Cochran-Armitage trend *p*-value = 0.058. *P*-values for comparisons with the control group were 0.071, 0.023, 0.019, and 0.036 in the 60, 125, 185, and 250 mg/kg-d groups, respectively, based on statistical analyses reported by Hazleton Laboratories (1983).

^bmg dichloromethane metabolized via GST pathway/L liver/d. Internal doses were estimated from simulations of actual daily doses reported by the study authors.


^cBased on the sum of dichloromethane metabolized via the GST pathway in the lung plus the liver, normalized to total BW (i.e., [lung GST metabolism (mg/d) + liver GST metabolism (mg/d)]/kg BW). Units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

Sources: Serota et al. (1986b); Hazleton Laboratories (1983).

5.4.1.4. Dose Conversion and Extrapolation Methods: Cancer OSF

Dose conversion. The mouse PBPK model of Marino et al. (2006) was based on the PBPK model for dichloromethane by Andersen et al. (1987), which was modified to include dichloromethane metabolism in the lung compartment and kinetics of CO and COHb (Andersen et al., 1991). For the mouse, physiological parameters and partition coefficients were adjusted to match those reported in Andersen et al. (1991, 1987) and Clewell et al. (1993), respectively, while QCC, VPR, and metabolic parameter distribution mean values were derived via MCMC model calibration reported by Marino et al. (2006) (Appendix B). The model of Marino et al. (2006) was used to simulate daily drinking water exposures comprising six discrete drinking water episodes for specified times and percentage of total daily intake (Reitz et al., 1997), and to calculate average lifetime daily internal doses for the male mouse data shown in Table 5-11. A first-order oral absorption rate constant (k_a) of 5 hours⁻¹ was taken from Reitz et al. (1997) to describe the uptake of dichloromethane from the gastrointestinal tract to the liver. Study-specific BWs were not available, so reference BWs for male B6C3F₁ mice in chronic studies (U.S. EPA, 1988a) were used. Based on evidence that metabolites of dichloromethane produced via the GST pathway are primarily responsible for dichloromethane carcinogenicity in mouse liver (summarized in Section 4.7.3) and the assumption that these metabolites are sufficiently reactive that they do not have substantial distribution outside the liver, the recommended selected internal dose metric for liver tumors was daily mass of dichloromethane metabolized via the GST pathway per unit volume of liver (Table 5-11). Figure 5-14 shows the comparison between

internal and external doses in the liver in mice and humans. The whole-body metabolism metric was also examined; however, this metric would be more relevant under a scenario of slowly cleared metabolites that undergo general circulation.



Six simulated drinking water episodes are described by Reitz et al. (1997). The human metabolism rates were estimated using a computational sample of 1,000 individuals per dose, including random samples of the three GST-T1 polymorphisms (+/+, +/-, -/-; “Human mixed” curve) or samples restricted to the GST +/+ or +/- populations in the current U.S. population based on data from Haber et al. (2002). Since a different set of samples was used for each dose, some stochasticity is evident as the human points (values) do not fall on smooth curves. Error bars indicate the range of 5th–95th percentile for the subpopulations sampled at select concentrations.

Figure 5-14. PBPK model-derived internal doses (mg dichloromethane metabolized via the GST pathway per liter liver per day) in mice and humans and their associated external exposures (mg/kg-day) used for the derivation of cancer OSFs based on liver tumors in mice.

Dose-response modeling and extrapolation. The multistage dose-response model was fit to the mouse liver tumor incidence and PBPK model-derived internal dose data to derive a mouse internal BMD₁₀ and BMDL₁₀ associated with 10% extra risk (Table 5-12). Different polynomial models and models dropping dose groups starting with the highest dose group were compared based on adequacy of model fit as assessed by overall χ^2 goodness of fit (p -value >

0.10) and examination of residuals at the 0 dose exposure (controls) and in the region of the BMR. Appendix E-1 provides details of the BMD modeling results. The mouse liver tumor risk factor (extra risk per unit internal dose) was calculated by dividing 0.1 by the mouse BMDL₁₀ for liver tumors.

Table 5-12. BMD modeling results and tumor risk factors for internal dose metric associated with 10% extra risk for liver tumors in male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics

Internal dose metric	BMDS model ^b	χ^2 goodness of fit <i>p</i> -value	Mouse BMD ₁₀ ^c	Mouse BMDL ₁₀ ^c	Allometric-scaled human BMDL ₁₀ ^d	Tumor risk factor ^e	
						Scaling = 1.0	Allometric-scaled
Liver-specific	MS (1,1)	0.56	73.0	39.6	5.66	2.53×10^{-3}	1.77×10^{-2}
Whole-body	MS (1,1)	0.56	3.05	1.65	0.24	–	4.24×10^{-1}

^aLiver specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue per d; whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

^bThe multistage (MS) model in EPA BMDS version 2.0 was fit to the mouse dose-response data shown in Table 5-11 using internal dose metrics calculated with the mouse PBPK model. Numbers in parentheses indicate (1) the number of dose groups dropped in order to obtain an adequate fit; and (2) the degree polynomial of the model.

^cBMD₁₀ and BMDL₁₀ refer to the BMD-model-predicted mouse internal and its 95% lower confidence limit, associated with a 10% extra risk for the incidence of tumors.

^dMouse BMDL₁₀ divided by $(BW_{\text{human}}/BW_{\text{mouse}})^{0.25} = 7$.

^eDichloromethane tumor risk factor (extra risk per unit internal dose) derived by dividing the BMR (0.1) by the mouse BMDL₁₀ and by the allometric-scaled human BMDL₁₀ for the scaling = 1.0 and allometric-scaled risk factors, respectively.

Linear extrapolation from the internal human BMDL₁₀ values (0.1/BMDL₁₀) was used to derive oral risk factors for liver tumors based on tumor responses in male mice. Proposed key events for dichloromethane carcinogenesis are discussed in Sections 4.7 and 5.4.1.1. The linear low-dose extrapolation approach for agents with a mutagenic mode of action was selected.

Application of allometric scaling factor. As discussed in Section 4.7 and summarized in 5.4.1.2, several lines of evidence point to the involvement of the GST metabolic pathway in the carcinogenic response seen in dichloromethane. The role of specific metabolites has not been firmly established, however. S-(chloromethyl)-glutathione is an intermediate to the production of formaldehyde through this pathway (Hashmi et al., 1994). Formation of the free hydrogen ion is also hypothesized, although no direct evidence supporting this has been presented. The pattern of HPRT gene mutations seen in CHO cells incubated with GST-complete mouse liver cytosol preparations suggest that S-(chloromethyl)glutathione, rather than formaldehyde, is responsible for the mutagenic effects associated with dichloromethane (Graves et al., 1996). DNA reaction products (e.g., DNA adducts) produced by S-(chloromethyl)glutathione have not been quantified,

possibly due to potential instability of these compounds (Watanabe et al., 2007; Hashmi et al., 1994).

The question of the role of specific metabolites and particularly how these metabolites are transformed or removed is a key question affecting the choice of a scaling factor to be used in conjunction with the internal dose metric based on rate of GST metabolism. If the key metabolite is established and is known to be sufficiently reactive to not spread in systemic circulation, then it can be assumed that: (1) the level of reactivity and rate of clearance (i.e., disappearance due to local reactivity) for this metabolite per volume tissue is equal in rodents and humans, and (2) risk is proportional to the long-term daily average concentration of the metabolite. Under these assumptions, rodent internal BMDL₁₀ values based on tissue-specific dichloromethane metabolism require no allometric scaling to account for toxicodynamic differences and predict the corresponding level of human risk as a function of the metric (i.e., the scaling factor in Figure 5-13 was equal to 1.0). (A single metabolite is referenced, but the same argument holds in general for more than one metabolite). Under this scenario and assumptions, humans and rodents with the same long-term daily average metabolite formation per volume tissue (e.g., equal internal BMDL₁₀) should both experience the same long-term average concentration of the metabolite when the metabolite is highly reactive and hence experience the same extra risk.

Although the evidence points to a specific metabolic pathway and to site-specific actions resulting from a reactive metabolite that does not escape the tissue in which it is formed, some assumptions remain concerning this hypothesis. Specifically, the active metabolite(s) have not been established, and data pertaining to the reactivity or clearance rate of these metabolite(s) are lacking. Quantitative measurements of adducts of interest or of the half life of relevant compounds in humans and in mice are not available. To address the uncertainties in the available data, it may be appropriate to use a scaling factor that addresses the possibility that the rate of clearance for the metabolite is limited by processes that are known to scale allometrically, such as blood perfusion, enzyme activity, or availability of reaction cofactors that is limited by overall metabolism. This case would result in use of a mouse:human dose-rate scaling factor of $(BW_{\text{human}}/BW_{\text{mouse}})^{0.25} = 7$ to adjust the mouse-based BMDL₁₀ values downward. Using this internal dose metric (liver-specific metabolism with allometric scaling), equivalent rodent and human internal BMDL₁₀ values result in a human liver tumor risk factor (0.1/BMDL₁₀) that is assumed equal to that for the mouse, given a 70-year lifetime exposure.

Another alternative that can be used is based on an allometrically-scaled whole-body metabolism metric. In this case, less weight is given to the evidence of site-specificity, as this metric allows for systemic circulation of the relevant metabolites.

The cancer toxicity values derived using each of these metrics and scaling factors (i.e., liver-specific metabolism with and without allometric-scaling and the whole-body metabolism metric) are presented in the following tables. Considering the lack of data pertaining to

clearance rates or the actual AUC of the active carcinogenic metabolite(s) in mice and humans, the OSF recommended by EPA is based on the allometrically-scaled tissue-specific GST metabolism rate dose metric.

Calculation of OSFs. The human PBPK model adapted from David et al. (2006) (see Appendix B), using Monte Carlo sampling techniques, was used to calculate distributions of human internal dose metrics of daily mass of dichloromethane metabolized via the liver-specific GST pathway per unit volume of liver resulting from a long-term average daily drinking water dose of 1 mg/kg dichloromethane. In another analysis of whole body metabolism, a dose metric based on the total metabolites formed in liver and lungs via GST metabolism per BW was used. The human model used parameter values derived from Monte Carlo sampling of probability distributions for each parameter, including MCMC-derived distributions for the metabolic parameters (David et al., 2006). The drinking water exposures comprised six discrete drinking-water episodes for specified times and percentage of total daily intake (Reitz et al., 1997) (Appendix B).

The distribution of cancer OSFs shown in Table 5-13 was derived by multiplying the human oral liver tumor risk factors by the respective distributions of human average daily internal doses resulting from chronic, unit oral exposures of 1 mg/kg-day dichloromethane. Because adjustments for interindividual variability are not generally used or recommended in cancer risk analysis, the mean slope factor was selected as the recommended value to be used in deterministic risk assessments; other values at the upper end of the distribution are also presented.

Table 5-13. Cancer OSFs for dichloromethane based on PBPK model-derived internal liver doses in B6C3F₁ mice exposed via drinking water for 2 years, based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype

Internal dose metric and scaling factor ^a	Population genotype ^b	Human tumor risk factor ^c	Distribution of human internal dichloromethane doses from 1 mg/kg-d exposure ^d			Resulting candidate human OSF ^e (mg/kg-d) ⁻¹		
			Mean	95 th percentile	99 th percentile	Mean	95 th percentile	99 th percentile
Liver-specific, allometric-scaled	GST-T1 ^{+/+}	1.77×10^{-2}	0.94×10^{-1}	2.98×10^{-1}	5.43×10^{-1}	1.7×10^{-3}	5.3×10^{-3}	9.6×10^{-3}
	Mixed	1.77×10^{-2}	0.53×10^{-1}	1.96×10^{-1}	3.78×10^{-1}	9.4×10^{-4}	3.5×10^{-3}	6.7×10^{-3}
Liver-specific, scaling = 1.0	GST-T1 ^{+/+}	2.53×10^{-3}	0.94×10^{-1}	2.98×10^{-1}	5.43×10^{-1}	2.4×10^{-4}	7.5×10^{-4}	1.4×10^{-3}
	Mixed	2.53×10^{-3}	0.53×10^{-1}	1.96×10^{-1}	3.78×10^{-1}	1.3×10^{-4}	5.0×10^{-4}	9.6×10^{-4}
Whole-body, allometric-scaled	GST-T1 ^{+/+}	4.24×10^{-1}	2.20×10^{-3}	7.20×10^{-3}	1.30×10^{-2}	9.3×10^{-4}	3.1×10^{-3}	5.5×10^{-3}
	Mixed	4.24×10^{-1}	1.27×10^{-3}	4.66×10^{-3}	9.41×10^{-3}	5.4×10^{-4}	2.0×10^{-3}	4.0×10^{-3}

^aLiver specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue per d; Whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

^bGST-T1^{+/+} = homozygous, full enzyme activity; mixed = population reflecting estimated frequency of genotypes in current U.S. population: 20% GST-T^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^cDichloromethane tumor risk factor (extra risk per unit internal dose per d) derived by dividing the BMR (0.1) by the allometric-scaled human BMDL₁₀ and the mouse BMDL₁₀ for the allometric-scaled and scaling = 1.0 risk factors, respectively (from Table 5-12).

^dMean, 95th, and 99th percentile of the human PBPK model-derived probability distribution of daily average internal dichloromethane dose resulting from chronic oral exposure of 1 mg/kg-d.

^eDerived by multiplying the dichloromethane tumor risk factor by the PBPK model-derived probabilistic internal doses from daily exposure to 1 mg/kg-d.

Consideration of Sensitive Human Subpopulations. An important issue in the derivation process used by EPA, pertaining to the use of the human PBPK model, stems from the assumption regarding the population for which the derivation should be applied. The inclusion of the GST-T1 null subpopulation in effect dilutes the risk that would be experienced by those who carry a GST-T1 allele by averaging in nonresponders (i.e., the GST-T1^{-/-} genotype). Thus, the cancer OSF was derived specifically for carriers of the GST-T1 homozygous positive (+/+) genotype, the population that would be expected to be most sensitive to the carcinogenic effects of dichloromethane given the GST-related dose metric under consideration. In addition, cancer values derived for a population reflecting the estimated frequency of GST-T1 genotypes in the current U.S. population (20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+}, i.e., the “mixed” population) are also presented. All simulations also included a distribution of CYP activity based on data from Lipscomb et al. (2003).

5.4.1.5. Oral Cancer Slope Factor

The recommended cancer OSF for dichloromethane is 2×10^{-3} (mg/kg-day)⁻¹ (rounded from 1.7×10^{-3}) and is based on liver tumor responses in male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years (Serota et al., 1986b; Hazleton Laboratories, 1983). The OSF was derived by using a tissue-specific GST metabolism dose metric with allometric scaling for the population that is presumed to have the greatest sensitivity (the GST-T1^{+/+} genotype). The application of ADAFs to the cancer OSF is recommended and is described in Section 5.4.4.

5.4.1.6. Alternative Derivation Based on Route-to-Route Extrapolation

For comparison, alternative cancer OSFs were derived via route-to-route extrapolations from the data for liver tumors in male and female B6C3F₁ mice exposed by inhalation for 2 years (Mennear et al., 1988; NTP, 1986). This derivation, shown in Table 5-14, uses the cancer IUR derived in Section 5.4.2.4 (see Table 5-19 for these IUR values) and the distribution of human internal dichloromethane exposures from 1 mg/kg-day exposure using the tissue-specific GST metabolism dose metric (mg dichloromethane metabolized via the GST pathway per liter liver per day). The cancer OSFs based on the route-to-route extrapolations from liver tumors in mice exposed by inhalation (Table 5-14) are about one order of magnitude lower than those based on the liver tumor responses in mice exposed via drinking water.

Table 5-14. Alternative route-to-route cancer OSFs for dichloromethane extrapolated from male B6C3F₁ mouse inhalation liver tumor incidence data using a tissue-specific GST metabolism dose metric, by population genotype

Internal dose metric and scaling factor	Population genotype ^a	Human tumor risk factor ^b	Distribution of human internal dichloromethane doses from 1 mg/kg-d exposure ^c			Resulting candidate human OSF ^d (mg/kg-d) ⁻¹		
			Mean	95 th percentile	99 th percentile	Mean	95 th percentile	99 th percentile
Liver-specific, allometric-scaled	GST-T1 ^{+/+}	1.29 × 10 ⁻³	0.94 × 10 ⁻¹	2.98 × 10 ⁻¹	5.43 × 10 ⁻¹	1.2 × 10 ⁻⁴	3.8 × 10 ⁻⁴	7.0 × 10 ⁻⁴
	Mixed	1.29 × 10 ⁻³	0.53 × 10 ⁻¹	1.96 × 10 ⁻¹	3.78 × 10 ⁻¹	6.8 × 10 ⁻⁵	2.5 × 10 ⁻⁴	4.9 × 10 ⁻⁴
Liver-specific, scaling = 1.0	GST-T1 ^{+/+}	1.84 × 10 ⁻⁴	0.94 × 10 ⁻¹	2.98 × 10 ⁻¹	5.43 × 10 ⁻¹	1.7 × 10 ⁻⁵	5.5 × 10 ⁻⁵	1.0 × 10 ⁻⁴
	Mixed	1.84 × 10 ⁻⁴	0.53 × 10 ⁻¹	1.96 × 10 ⁻¹	3.78 × 10 ⁻¹	9.7 × 10 ⁻⁶	3.6 × 10 ⁻⁵	6.9 × 10 ⁻⁵
Whole-body metabolism	GST-T1 ^{+/+}	3.03 × 10 ⁻²	2.20 × 10 ⁻³	7.20 × 10 ⁻³	1.30 × 10 ⁻²	6.7 × 10 ⁻⁵	2.2 × 10 ⁻⁴	3.9 × 10 ⁻⁴
	Mixed	3.03 × 10 ⁻²	1.27 × 10 ⁻³	4.66 × 10 ⁻³	9.41 × 10 ⁻³	3.9 × 10 ⁻⁵	1.4 × 10 ⁻⁴	2.9 × 10 ⁻⁴

^aGST-T1^{+/+} = homozygous, full enzyme activity; mixed = population reflecting estimated frequency of genotypes in current U.S. population: 20% GST-T^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^bDichloromethane tumor risk factor (extra risk per milligram dichloromethane metabolized via GST pathway per liter tissue per d) derived by dividing the BMR (0.1) by the allometric-scaled human BMDL₁₀ and the mouse BMDL₁₀ for the allometric-scaled and scaling = 1.0 risk factors, respectively (from IUR data, Table 5-19).

^cMean, 95th, and 99th percentile of the human PBPK model-derived probability distribution of daily average internal dichloromethane dose (mg dichloromethane metabolized via GST pathway per liter tissue per d) resulting from chronic oral exposure of 1 mg/kg-d.

^dDerived by multiplying the dichloromethane tumor risk factor by the PBPK model-derived probabilistic internal doses from daily exposure to 1 mg/kg-d.

5.4.1.7. Alternative Based On Administered Dose

One comparison that can be made is with an alternative OSF based on liver tumors in mice, using the external concentrations of dichloromethane in the mouse as converted to human equivalent doses and then applying this by using BMD modeling to obtain the BMDL₁₀ and resulting oral cancer risk. Mouse bioassay exposures were adjusted to human equivalent doses as follows:

$$\text{human equivalent dose} = (\text{nominal daily intake/BW scaling factor}) \times \text{daily exposure adjustment factor}$$

where BW scaling factor = $(\text{BW}_{\text{human}}/\text{BW}_{\text{mouse}})^{0.25} = 7$
 and
 daily exposure adjustment factor = 5/7

The human equivalent doses for the 0, 60, 125, 185, and 250 mg/kg-day dose groups used in the liver tumor analysis (Table 5-11) (from Serota et al. [1986b]) were 0, 6.12, 12.75, 18.87, and 25.51 mg/kg-day, respectively. The BMD modeling and OSF derived from these values are shown in Table 5-15. The resulting OSF based on the liver tumors in the mouse is approximately one order of magnitude higher than the current recommended value obtained by using the mouse and human PBPK models.

Table 5-15. Cancer OSF based on a human BMDL₁₀ using administered dose for liver tumors in male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years

Sex, tumor type	BMDS model ^a	χ^2 goodness of fit p-value	Human BMD ₁₀ ^c	Human BMDL ₁₀ ^c	Cancer OSF ^d (mg/kg-d) ⁻¹
Male, liver	MS (0,1)	0.55	19.4	10.4	1.0×10^{-2}

^aThe multistage (MS) model in EPA BMDS version 2.0 was fit to the mouse liver tumor data shown in Table 5-11. The human equivalent doses for the 0, 60, 125, 185, and 250 mg/kg-d dose groups used in the liver tumor analysis were 0, 6.12, 12.75, 18.87, and 25.51 mg/kg-d, respectively, based on application of BW scaling factor = $(\text{BW}_{\text{human}}/\text{BW}_{\text{mouse}})^{0.25} = 7$ and adjusting for daily exposure by multiplying by 5/7 d. Numbers in parentheses indicate: (1) the number of dose groups dropped in order to obtain an adequate fit, starting with the highest dose group, and (2) the degree polynomial of the model.

^cBMD₁₀ and BMDL₁₀ refer to the BMD-model-predicted human equivalent dose (mg/kg-d) and its 95% lower confidence limit, associated with a 10% extra risk for the incidence of tumors.

^dCancer OSF (risk per mg/kg-d) = 0.1/human BMDL₁₀.

The administered dose methodology can be considered equivalent to using a single-compartment, whole-body model of dichloromethane where the internal dose metric is the AUC of dichloromethane itself, and clearance of dichloromethane scales from mice to humans as $\text{BW}^{0.75}$. The estimates based on the PBPK model, in contrast, use the rate of metabolism of dichloromethane (GST) as the metric. Another difference is that the administered dose

methodology does not account in any way for the GST polymorphism and so might be considered as representing the general/mixed-GST-genotype population rather than the +/+ subpopulation.

5.4.1.8. Previous IRIS Assessment: Cancer OSF

The previous IRIS assessment derived a cancer OSF of 7.5×10^{-3} (mg/kg-day)⁻¹ by the application of the multistage model to combined incidence of hepatocellular adenomas, carcinomas from two studies. These were the 2-year drinking water study of dichloromethane in B6C3F₁ mice by the Hazleton Laboratories (1983) and the 2-year inhalation study of dichloromethane in B6C3F₁ mice by NTP (1986). The slope factor was the arithmetic mean of two candidate slope factors, 1.2×10^{-2} (mg/kg-day)⁻¹ (Hazleton Laboratories, 1983) and 2.6×10^{-3} (mg/kg-day)⁻¹ (NTP, 1986). Since the NTP (1986) animal data were from inhalation exposures, the estimated inhaled doses were calculated for mice and humans (assuming near complete uptake into lung tissues and blood) and converted to administered doses in units of mg/kg-day. Assumed inhalation rates of 0.0407 and 20 m³/day were used for mice and humans, respectively. No adjustments were made for species differences in metabolism or toxicokinetics.

5.4.1.9. Comparison of Cancer OSFs Using Different Methodologies

Cancer OSFs derived using different dose metrics and assumptions are summarized in Table 5-16. The recommended OSF of 2×10^{-3} per mg/kg-day (rounded to one significant digit) is based on a tissue-specific GST-internal dose metric with allometric scaling (=7) because of some uncertainty regarding the rate of clearance of the relevant metabolite(s) formed via the GST pathway. The value derived specifically for the GST-T1^{+/+} population is recommended to provide protection for the population that is hypothesized to be most sensitive to the carcinogenic effect. The values based on the GST-T1^{+/+} group are approximately twofold higher than those for the full population for the dose metrics used in this assessment (Table 5-16). Within a genotype population, the values of the OSF among most of the various dose metrics vary by about one to two orders of magnitude.

Table 5-16. Comparison of OSFs derived using various assumptions and metrics, based on tumors in male mice

Population ^a	Dose metric	Species, sex	Tumor	Scaling factor	Mean OSF (mg/kg-d) ⁻¹	Source (table)
GST-T1 ^{+/+} ^b	Tissue-specific GST-metabolism rate	Mouse, male	Liver	7.0	1.7 × 10⁻³	Table 5-13
	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	2.4 × 10 ⁻⁴	Table 5-13
	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	9.3 × 10 ⁻⁴	Table 5-13
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	7.0	1.2 × 10 ⁻⁴	Table 5-14
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	1.0	1.7 × 10 ⁻⁵	Table 5-14
	Route-to-route extrapolation, whole-body metabolism	Mouse, male	Liver	7.0	6.7 × 10 ⁻⁵	Table 5-14
Mixed	Tissue-specific GST-metabolism rate ^b	Mouse, male	Liver	7.0	9.4 × 10 ⁻⁴	Table 5-13
	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	1.3 × 10 ⁻⁴	Table 5-13
	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	5.4 × 10 ⁻⁴	Table 5-13
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	7.0	6.8 × 10 ⁻⁵	Table 5-14
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	1.0	9.7 × 10 ⁻⁶	Table 5-14
	Route-to-route extrapolation, whole-body metabolism	Mouse, male	Liver	7.0	3.9 × 10 ⁻⁵	Table 5-14
	Applied dose (human equivalent dose)	Mouse, male	Liver		1.0 × 10 ⁻²	Table 5-15
	1995 IRIS assessment	Mouse, male	Liver		7.5 × 10 ⁻³	

^aGST-T1^{+/+} = homozygous, full enzyme activity; Mixed = genotypes based on a population reflecting the estimated frequency of genotypes in the current U.S. population: 20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^bBolded value is the basis for the recommended OSF of 2 × 10⁻³ per mg/kg-d.

5.4.2. Cancer IUR

5.4.2.1. *Choice of Study/Data—with Rationale and Justification*

As discussed in Section 4.7, results from several cohort mortality studies of workers repeatedly exposed to dichloromethane and several case-control studies provide some supporting evidence of carcinogenicity in humans, specifically with respect to liver and brain cancer. However, the epidemiologic studies do not provide adequate data to estimate exposure-response relationships for dichloromethane exposure and these cancers.

Results from several bioassays provide sufficient evidence of the carcinogenicity of dichloromethane in mice and rats exposed by inhalation, as well as adequate data to describe dose-response relationships. As discussed in Section 4.7.2, repeated inhalation exposure to concentrations of 2,000 or 4,000 ppm dichloromethane produced increased incidences of lung and liver tumors in male and female B6C3F₁ mice (Maronpot et al., 1995; Foley et al., 1993; Kari et al., 1993; Mennear et al., 1988; NTP, 1986). A weaker trend ($p = 0.08$) was seen with respect to liver tumor incidence (described as neoplastic nodules or hepatic carcinomas) in female rats, but this trend was not seen when limited to hepatic carcinomas (NTP, 1986). A statistically significant increased incidence of brain tumors has not been observed in any of the animal cancer bioassays, but a 2-year study using relatively low exposure levels (0, 50, 200, and 500 ppm) in Sprague-Dawley rats observed a total of six astrocytoma or glioma (mixed glial cell) tumors (combining males and females) in the exposed groups (Nitschke et al., 1988a). These tumors are exceedingly rare in rats, and there are few examples of statistically significant trends in animal bioassays (Sills et al., 1999). Male and female F344 rats exposed by inhalation to 2,000 or 4,000 ppm showed significantly increased incidences of benign mammary tumors (adenomas or fibroadenomas) and the male rats also exhibited a low rate of sarcoma or fibrosarcoma in mammary gland or subcutaneous tissue around the mammary gland (NTP, 1986).

The NTP inhalation study in B6C3F₁ mice (NTP, 1986) was used to derive an IUR for dichloromethane because of the completeness of the data, adequate sample size, and clear dose response with respect to liver and lung tumors. The liver tumor incidence in male mice increased from 44% in controls to 66% in the highest dose group; in females, the incidence of this tumor rose from 6 to 83%. For lung tumors, the incidence rose from 10 to 80% in males and from 6 to 85% in females. Compelling evidence exists for the role of GST-mediated metabolism of dichloromethane in carcinogenicity in mice (Section 4.7.3), and both mice and humans possess this metabolic pathway. Modeling intake, metabolism, and elimination of dichloromethane in mice and humans is feasible. Thus, it is reasonable to apply the best available PBPK models to estimate equivalent internal doses in mice and humans.

The mammary tumor data from the NTP (1986) study was also used to derive a comparative IUR. However, the toxicokinetic or mechanistic events that might lead to mammary gland tumor development in rats are unknown, and so a clear choice of the optimal

internal dose metric could not be made. Thus, this derivation is based on the average daily AUC for dichloromethane in blood. The role of CYP- or GST-mediated metabolism in the mammary gland is uncertain, although both GST-T1 (Lehmann and Wagner, 2008) and CYP2E1 (El-Rayes et al., 2003; Hellmold et al., 1998) expressions have been detected in human mammary tissue. It is also possible that some metabolites enter systemic circulation from the liver and lung where they are formed.

The female rat liver cancer data from the NTP (1986) inhalation study was not used to derive an IUR because the trend was weaker than that seen in the mouse (incidence increased from 4% in controls to 10% in the highest dose group, trend $p = 0.08$), and because the effect categorization included neoplastic nodule or hepatocellular carcinoma. The brain tumor data seen in the Nitschke et al. (1988a) study in Sprague-Dawley rats were not used to develop an IUR because of the low incidence of this rare tumor (a total of four astrocytoma or glioma tumors in exposed males and two in exposed females). The mechanistic issues with respect to mammary tumors and health effects issues with respect to brain tumors represent data gaps in the understanding of the health effects of dichloromethane and relevance of the rat data to humans.

5.4.2.2. Derivation of the Cancer IUR

The derivation of the IUR parallels the process described in Section 5.4.1.2 for the cancer OSF. Since modeling metabolism and elimination kinetics of dichloromethane in mice and humans is feasible, it is reasonable to apply the best available PBPK models to determine equivalent target organ doses in mice and humans. Although the GST metabolic pathway takes on a greater role as the CYP pathway is saturated, both the GST and CYP pathways are operating even at low exposures. The PBPK model incorporates the metabolic shift and expected nonlinearity (GST dose attenuation with low exposures) in the exposure-dose relationship across exposure levels.

5.4.2.3. Dose-Response Data

Data for liver and lung tumors in male and female B6C3F₁ mice following exposure to airborne dichloromethane were used to develop IURs for dichloromethane (Mennear et al., 1988; NTP, 1986). As discussed in Section 5.4.1.8, the liver tumor dose-response data were also the basis of an OSF derived by route-to-route extrapolation using the PBPK models to compare with an OSF based on liver tumor data in mice exposed to dichloromethane in drinking water (Serota et al., 1986b). In the NTP (1986) study, significant increases in incidence of liver and lung adenomas and carcinomas were observed in both sexes of B6C3F₁ mice exposed 6 hours/day, 5 days/week for 2 years (Table 5-17). Since significant decreases in survival were observed in the treated groups of both sexes, the at-risk study populations (represented by the denominators in the incidence data) were determined by excluding all animals dying prior 52 weeks.

Table 5-17. Incidence data for liver and lung tumors and internal doses based on GST metabolism dose metrics in male and female B6C3F₁ mice exposed to dichloromethane via inhalation for 2 years

Sex, tumor type	BW (g)	External dichloromethane concentration (ppm)	Mouse tumor incidence	Mouse internal tissue dose ^a	Mouse whole body metabolism dose ^b
Male, liver ^c	–	0	22/50 (44%) ^d	0	0
	34.0	2,000	24/47 (51%)	2,363.7	100.2
	32.0	4,000	33/47 (70%)	4,972.2	210.7
Male, lung ^e	–	0	5/50 (10%) ^d	0	0
	34.0	2,000	27/47 (55%)	475.0	100.2
	32.0	4,000	40/47 (85%)	992.2	210.7
Female, liver ^c	–	0	3/47 (6%) ^d	0	0
	30.0	2,000	16/46 (35%)	2,453.2	104.0
	29.0	4,000	40/46 (87%)	5,120.0	217.0
Female, lung ^e	–	0	3/45 (6%) ^d	0	0
	30.0	2,000	30/46 (65%)	493.0	104.0
	29.0	4,000	41/46 (89%)	1,021.8	217.0

^aFor liver tumors: mg dichloromethane metabolized via GST pathway/L liver tissue/d from 6 hrs/d, 5 d/wk exposure; for lung tumors: mg dichloromethane metabolized via GST pathway/L lung tissue/d from 6 hrs/d, 5 d/wk exposure.

^bBased on the sum of dichloromethane metabolized via the GST pathway in the lung plus the liver, normalized to total BW (i.e., [lung GST metabolism (mg/d) + liver GST metabolism (mg/d)]/kg BW). Units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

^cHepatocellular carcinoma or adenoma. Mice dying prior to 52 wks were excluded from the denominators.

^dStatistically significant increasing trend (by incidental and life-table tests; $p \leq 0.01$).

^eBronchoalveolar carcinoma or adenoma. Mice dying prior to 52 wks were excluded from the denominators.


Sources: Mennear et al. (1988); NTP (1986).

5.4.2.4. Dose Conversion and Extrapolation Methods: Cancer IUR


Dose conversion. The PBPK model of Marino et al. (2006) for dichloromethane in the mouse was used to simulate inhalation exposures of 6 hours/day, 5 days/week (Mennear et al., 1988; NTP, 1986) and to calculate long-term daily average internal doses. Study-, group-, and sex-specific mean BWs were used. Based on evidence that metabolites of dichloromethane produced via the GST pathway are primarily responsible for dichloromethane carcinogenicity in mouse liver and lung (summarized in Section 4.7.3) and the assumption that these metabolites are sufficiently reactive that they do not have substantial distribution outside these tissues, the recommended selected internal dose metrics for liver tumors and lung tumors were long-term average daily mass of dichloromethane metabolized via the GST pathway per unit volume of liver and lung, respectively (Table 5-17). Figure 5-15 shows the comparison between inhalation external and internal doses in the liver and lung using this dose metric for the mouse and for the

human. A whole-body metabolism metric was also examined; however, this metric would be more relevant under a scenario of slowly cleared metabolites that undergo general circulation.

A.



B.



Average daily doses were calculated from simulated mouse exposures of 6 hours/day, 5 days/week, while simulated human exposures were continuous. The GST metabolism rate in each simulated human population was obtained by generating 1,000 random samples from each population (ages 0.5–80 years, males and females) for each exposure level and calculating the average GST metabolic rate for each sample.

Figure 5-15. PBPK model-derived internal doses (mg dichloromethane metabolized via the GST pathways per liter tissue per day) for liver (A) and lung (B) in mice and humans and their associated external exposures (ppm) used for the derivation of cancer IURs.

Dose-response modeling and extrapolation. The multistage dose-response model was fit to the mouse tumor incidence and PBPK model-derived internal dose data to derive mouse internal BMD₁₀ and BMDL₁₀ values associated with 10% extra risk (Table 5-18). Different polynomial models and models dropping dose groups starting with the highest dose group were compared based on adequacy of model fit as assessed by overall χ^2 goodness of fit (p -value > 0.10)) and examination of residuals at the 0 dose exposure (controls) and in the region of the BMR (U.S. EPA, 2000c). Appendix E-2 provides details of the BMD modeling results for the male. The mouse liver and lung tumor risk factors (extra risk per unit internal dose) were calculated by dividing 0.1 by the mouse BMDL₁₀ for liver and lung tumors, respectively.

Table 5-18. BMD modeling results and tumor risk factors associated with 10% extra risk for liver and lung tumors in male and female B6C3F₁ mice exposed by inhalation to dichloromethane for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics

Internal dose metric ^a		BMDS model ^b	χ^2 goodness of fit <i>p</i> -value	Mouse BMD ₁₀ ^c	Mouse BMDL ₁₀ ^c	Allometric-scaled human BMDL ₁₀ ^d	Tumor risk factor ^e	
							Scaling = 1.0	Allometric-scaled
Tissue-specific	Male, liver	MS (0,1)	0.40	913.9	544.4	77.8	1.84×10^{-4}	1.29×10^{-3}
	Male, lung	MS (0,1)	0.64	61.7	48.6	7.0	2.06×10^{-3}	1.44×10^{-2}
	Female, liver	MS (0,2)	0.53	1,224.1	659.7	94.2	1.52×10^{-4}	1.06×10^{-3}
	Female, lung	MS (0,1)	0.87	51.2	40.7	5.8	2.46×10^{-3}	1.72×10^{-2}
Whole body	Male, liver	MS (0,1)	0.40	38.7	23.1	3.3	-	3.03×10^{-2}
	Male, lung	MS (0,1)	0.66	13.1	10.3	1.5	-	6.80×10^{-2}
	Female, liver	MS (0,2)	0.53	51.9	28.0	4.0	-	2.50×10^{-2}
	Female, lung	MS (0,1)	0.88	10.8	8.6	1.2	-	8.14×10^{-2}

^aTissue-specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue (liver or lung) per d; Whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d).

^bThe multistage (MS) model in EPA BMDS version 2.0 was fit to the mouse dose-response data shown in Table 5-17 using internal dose metrics calculated with the mouse PBPK model. Numbers in parentheses indicate: (1) the number of dose groups dropped in order to obtain an adequate fit, and (2) the degree polynomial of the model.

^cBMD₁₀ and BMDL₁₀ refer to the BMD-model-predicted mouse internal dose and its 95% lower confidence limit, associated with a 10% extra risk for the incidence of tumors.

^dMouse BMDL₁₀ divided by $(BW_{\text{human}}/BW_{\text{mouse}})^{0.25} = 7$.

^eDichloromethane tumor risk factor (extra risk per unit internal dose) derived by dividing the BMR (0.1) by the mouse BMDL₁₀ and by the allometric-scaled human BMDL₁₀, for the scaling = 1.0 and allometric-scaled risk factors, respectively.

Linear extrapolation from the internal BMDL₁₀ (0.1/BMDL₁₀) was used to derive inhalation risk factors for lung and liver tumors in male and female mice (Table 5-18). As discussed in Section 4.7, the linear low-dose extrapolation approach is applied for agents with a mutagenic mode of action.

Application of allometric scaling factor. As discussed in Section 5.4.1.4, the choice of a scaling factor is based on the question of the role of specific metabolites and particularly how these metabolites are transformed or removed. If the key metabolite is established and is known to be sufficiently reactive to not spread in systemic circulation, then it can be assumed that: (1) the level of reactivity and rate of clearance (i.e., disappearance due to local reactivity) for this metabolite per volume tissue is equal in rodents and humans, and (2) risk is proportional to the long-term daily average concentration of the metabolite. Under these assumptions, rodent internal BMDL₁₀ values based on tissue-specific dichloromethane metabolism require no allometric scaling to account for toxicodynamic differences and predict the corresponding level of human risk as a function of the metric (i.e., the scaling factor in Figure 5-13 was equal to 1.0). (A single metabolite is referenced, but the same argument holds in general for more than one metabolite). Under this scenario and assumptions, humans and rodents with the same long-term daily average metabolite formation per volume tissue (e.g., equal internal BMDL₁₀) should both experience the same long-term average concentration of the metabolite when the metabolite is highly reactive and, hence, experience the same extra risk. However, some uncertainties remain concerning the hypothesized role of a highly reactive metabolite in the carcinogenic effects seen with dichloromethane. The active metabolite(s) have not been established, and data pertaining to the reactivity or removal (clearance) rate of these metabolite(s) are lacking. For example, quantitative measurements of adducts of interest or of the half life of relevant compounds in humans and in mice are not available. To address these uncertainties, use of a scaling factor that addresses the possibility that the rate of clearance for the metabolite is limited by processes that scale allometrically, such as blood perfusion, reaction cofactor supply (e.g., antioxidant supply), or enzyme activity, may be appropriate. This case would result in use of a mouse:human dose-rate scaling factor of $(BW_{\text{human}}/BW_{\text{mouse}})^{0.25} = 7$ to adjust the mouse-based BMDL₁₀ values downward. Using this internal dose metric (liver-specific metabolism with allometric scaling), equivalent rodent and human internal BMDL₁₀ values result in a human liver tumor risk factor (0.1/BMDL₁₀) that is assumed equal to that for the mouse, given a 70-year lifetime exposure. Another alternative that can be used is based on an allometrically-scaled, whole-body metabolism metric. In this case, less weight is given to the evidence of site-specificity of the effects. As with the OSF derivations, the cancer toxicity values derived using each of these metrics and scaling factors (i.e., liver-specific metabolism with and without allometric-scaling and the whole-body metabolism metric) are presented in the following tables. Considering the lack of data pertaining to clearance rates or the actual AUC of the active carcinogenic

metabolite(s) in mice and humans, the IURs recommended by EPA are based on the allometrically-scaled tissue-specific GST metabolism rate dose metric.

Calculation of IURs. A probabilistic PBPK model for dichloromethane in humans, adapted from David et al. (2006) (see Appendix B), was used with Monte Carlo sampling to calculate distributions of internal lung, liver, or blood doses associated with chronic unit inhalation ($1 \mu\text{g}/\text{m}^3$) exposures. The data on which the model is based indicate that the relationship between exposure and internal dose is linear at low doses. Parameters in the human PBPK model developed by David et al. (2006) are distributions that incorporate information about dichloromethane toxicokinetic variability and uncertainty among humans. Monte Carlo sampling was performed in which each human model parameter was defined by a value randomly drawn from each respective parameter distribution. The model was then executed by using the external unit exposure as input, and the resulting human equivalent internal dose was recorded. This process was repeated for 10,000 iterations to generate a distribution of human internal doses.

The resulting distribution of IURs shown in Table 5-19 was derived by multiplying the human internal dose tumor risk factor (in units of reciprocal internal dose) by the respective distributions of human average daily internal dose resulting from a chronic unit inhalation exposure of $1 \mu\text{g}/\text{m}^3$ dichloromethane. Table 5-19 presents the analysis using the male data. Analyses based on the female data produced very similar results, and are summarized in Appendix F. Because adjustments for interindividual variability are not generally used or recommended in cancer risk analysis, the mean slope factor was selected as the recommended value to be used in deterministic risk assessments; other values at the upper end of the distribution are also presented. As with the cancer OSF derivation, the cancer IUR is derived for a population composed entirely of carriers of the GST-T1 homozygous positive genotype (the group that would be expected to be most sensitive to the carcinogenic effects of dichloromethane), and a population reflecting the estimated frequency of GST-T1 genotypes in the current U.S. population (20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+}, the “mixed” population). All simulations also included a distribution of CYP activity, based on data from Lipscomb et al. (2003).

Table 5-19. IURs for dichloromethane based on PBPK model-derived internal liver and lung doses in B6C3F₁ male mice exposed via inhalation for 2 years, based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype

Internal dose metric and scaling factor ^a	Population genotype ^b	Tumor type	Human tumor risk factor ^c	Distribution of human internal dichloromethane doses from 1 µg/m ³ exposure ^d			Resulting candidate human IUR ^e (µg/m ³) ⁻¹		
				Mean	95 th percentile	99 th percentile	Mean	95 th percentile	99 th percentile
Tissue-specific, allometric-scaled	GST-T1 ^{+/+}	Liver	1.29 × 10 ⁻³	6.61 × 10 ⁻⁶	2.21 × 10 ⁻⁵	4.47 × 10 ⁻⁵	8.5 × 10 ⁻⁹	2.8 × 10 ⁻⁸	5.8 × 10 ⁻⁸
	GST-T1 ^{+/+}	Lung	1.44 × 10 ⁻²	3.89 × 10 ⁻⁷	1.24 × 10 ⁻⁶	2.42 × 10 ⁻⁶	5.6 × 10 ⁻⁹	1.8 × 10 ⁻⁸	3.5 × 10 ⁻⁸
	Mixed	Liver	1.29 × 10 ⁻³	3.71 × 10 ⁻⁶	1.43 × 10 ⁻⁵	3.03 × 10 ⁻⁵	4.8 × 10 ⁻⁹	1.8 × 10 ⁻⁸	3.9 × 10 ⁻⁸
	Mixed	Lung	1.44 × 10 ⁻²	2.20 × 10 ⁻⁷	8.06 × 10 ⁻⁷	1.69 × 10 ⁻⁶	3.2 × 10 ⁻⁹	1.2 × 10 ⁻⁸	2.4 × 10 ⁻⁸
Tissue-specific, scaling = 1.0	GST-T1 ^{+/+}	Liver	1.84 × 10 ⁻⁴	6.61 × 10 ⁻⁶	2.21 × 10 ⁻⁵	4.47 × 10 ⁻⁵	1.2 × 10 ⁻⁹	4.1 × 10 ⁻⁹	8.2 × 10 ⁻⁹
	GST-T1 ^{+/+}	Lung	2.06 × 10 ⁻³	3.89 × 10 ⁻⁷	1.24 × 10 ⁻⁶	2.42 × 10 ⁻⁶	8.0 × 10 ⁻¹⁰	2.6 × 10 ⁻⁹	5.0 × 10 ⁻⁹
	Mixed	Liver	1.84 × 10 ⁻⁴	3.71 × 10 ⁻⁶	1.43 × 10 ⁻⁵	3.03 × 10 ⁻⁵	6.8 × 10 ⁻¹⁰	2.6 × 10 ⁻⁹	5.6 × 10 ⁻⁹
	Mixed	Lung	2.06 × 10 ⁻³	2.20 × 10 ⁻⁷	8.06 × 10 ⁻⁷	1.69 × 10 ⁻⁶	4.5 × 10 ⁻¹⁰	1.7 × 10 ⁻⁹	3.5 × 10 ⁻⁹
Whole-body, allometric-scaled	GST-T1 ^{+/+}	Liver	3.03 × 10 ⁻²	1.80 × 10 ⁻⁷	6.38 × 10 ⁻⁷	1.41 × 10 ⁻⁶	5.5 × 10 ⁻⁹	1.9 × 10 ⁻⁸	4.3 × 10 ⁻⁸
	GST-T1 ^{+/+}	Lung	6.80 × 10 ⁻²	1.80 × 10 ⁻⁷	6.38 × 10 ⁻⁷	1.41 × 10 ⁻⁶	1.2 × 10 ⁻⁸	4.3 × 10 ⁻⁸	9.6 × 10 ⁻⁸
	Mixed	Liver	3.03 × 10 ⁻²	1.01 × 10 ⁻⁷	4.00 × 10 ⁻⁷	9.43 × 10 ⁻⁷	3.1 × 10 ⁻⁹	1.2 × 10 ⁻⁸	2.9 × 10 ⁻⁸
	Mixed	Lung	6.80 × 10 ⁻²	1.01 × 10 ⁻⁷	4.00 × 10 ⁻⁷	9.43 × 10 ⁻⁷	6.9 × 10 ⁻⁹	2.7 × 10 ⁻⁸	6.4 × 10 ⁻⁸

^aTissue specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue (liver or lung, respectively, for liver and lung tumors) per d; whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

^bGST-T1^{+/+} = homozygous, full enzyme activity; mixed = population reflecting estimated frequency of genotypes in current U.S. population: 20% GST-T^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^cDichloromethane tumor risk factor (extra risk per unit internal dose) derived by dividing the BMR (0.1) by the allometric-scaled human BMDL₁₀ or by the mouse BMDL₁₀ (from Table 5-18) for the allometric-scaled and scaling = 1.0 risk factors, respectively.

^dMean, 95th, and 99th percentile of the human PBPK model-derived probability distribution of daily average internal dichloromethane dose resulting from chronic exposure to 1 µg/m³ (0.00029 ppm).

^eDerived by multiplying the dichloromethane tumor risk factor by the PBPK model-derived probabilistic internal doses from daily exposure to 1 µg/m³.

5.4.2.5. *Cancer IUR*

The recommended cancer IURs are $9 \times 10^{-9} (\mu\text{g}/\text{m}^3)^{-1}$ and $6 \times 10^{-9} (\mu\text{g}/\text{m}^3)^{-1}$ for the development of liver and lung cancer, respectively, based on the mean for the GST-T1^{+/+} population (the group with the greatest presumed sensitivity). These values are based on male B6C3F₁ mice, using a tissue-specific GST metabolism dose metric with allometric scaling (Table 5-19). Risk estimates were similar to the values based on female mice in the NTP (1986) inhalation study: $7 \times 10^{-9} (\mu\text{g}/\text{m}^3)^{-1}$ and $7 \times 10^{-9} (\mu\text{g}/\text{m}^3)^{-1}$ for the development of liver and lung cancer, respectively, in the GST-T1^{+/+} population (see Appendix F).

Consideration of combined risk (summing risk across tumors). With two significant tumor sites, focusing on the more sensitive response may underestimate the overall cancer risk associated with exposure to this chemical. Following the recommendations of the NRC (1994) and the *Guidelines for Carcinogen Risk Assessment* (U.S. EPA, 2005a), an upper bound on total risk was estimated in order to gain some understanding of the total risk from multiple tumor sites in the selected data set. Note that this estimate of overall risk describes the risk of developing either tumor type, not just the risk of developing both simultaneously.

NRC (1994) stated that an approach based on counts of animals with one or more tumors (or tumor-bearing animals) would tend to underestimate overall risk when tumor types occur independently and that an approach based on combining the risk estimates from each separate tumor type should be used. For dichloromethane, there is no reason to expect that the occurrence of one tumor type depends on the incidence of the other, given the association of different dose metrics with each tumor response. Therefore, it appears reasonable to assume that the two tumor types occur independently. However, simply summing upper limit risks may result in an overestimation of overall of combined risk because of the statistical issues with respect to summing variances of distributions. An additional challenge results from the use of different internal dose metrics for different tumors, as is the case with the dose metrics based on tissue-specific metabolism. Statistical methods based on a common metric cannot be used with the tissue-specific metabolism metric used in these derivations.

An alternative approach is to derive an upper bound on the combined risk estimates by summing central tendency risks and calculating a pooled SD by using BMD₁₀ and BMDL₁₀ values for liver and lung tumors. The SD associated with the IUR for each tumor site is calculated as the difference between 95th percentiles of the distribution for upper bound and maximum likelihood estimate IURs (based on either female or male mouse tumor risk factors), divided by 1.645 (the relevant *t* statistic, assuming normal distributions of summed quantities). Variances for each tumor site are the squares of the SDs. Pooled variance and SD are defined as the sum of variances for lung and liver tumors and the square root of that sum, respectively. Finally, the upper bound on the combined lung and liver cancer risk is determined by multiplying the cumulative SD by 1.645 and adding it to the summed central tendency IURs. The

calculations of these upper bound estimates for combined liver and lung tumor risks are shown in Table 5-20.

Table 5-20. Upper bound estimates of combined human IURs for liver and lung tumors resulting from lifetime exposure to 1 µg/m³ dichloromethane based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype

Internal dose metric and scaling factor ^a	Population genotype ^b	Tumor site	Upper bound IUR ^c	Central tendency IUR ^d	Variance of tissue-specific tumor risk ^e	Combined tumor risk SD ^f	Upper bound on combined tumor risk ^g (µg/m ³) ⁻¹
Tissue-specific, allometric-scaled	GST-T1 ^{+/+}	Liver	8.5×10^{-9}	5.1×10^{-9}	4.36×10^{-18}		
		Lung	5.6×10^{-9}	4.4×10^{-9}	5.18×10^{-19}		
		Liver or lung		9.5×10^{-9}		2.2×10^{-9}	1.3×10^{-8}
	Mixed	Liver	4.8×10^{-9}	2.8×10^{-9}	1.37×10^{-18}		
		Lung	3.2×10^{-9}	2.5×10^{-9}	1.66×10^{-19}		
		Liver or lung		5.3×10^{-9}		1.2×10^{-9}	7.4×10^{-9}
Tissue-specific, scaling = 1.0	GST-T1 ^{+/+}	Liver	1.2×10^{-9}	7.2×10^{-10}	8.91×10^{-20}		
		Lung	8.0×10^{-10}	6.3×10^{-10}	1.07×10^{-20}		
		Liver or lung		1.4×10^{-9}		3.2×10^{-10}	1.9×10^{-9}
	Mixed	Liver	6.8×10^{-10}	4.1×10^{-10}	2.81×10^{-20}		
		Lung	4.5×10^{-10}	3.6×10^{-10}	3.41×10^{-21}		
		Liver or lung		7.6×10^{-10}		1.7×10^{-10}	1.1×10^{-9}
Whole-body, allometric-scaled	GST-T1 ^{+/+}	Liver	5.5×10^{-9}	3.3×10^{-9}	1.79×10^{-18}		
		Lung	1.2×10^{-8}	9.6×10^{-9}	2.55×10^{-18}		
		Liver or lung		1.3×10^{-8}		2.1×10^{-9}	1.6×10^{-8}
	Mixed	Liver	3.1×10^{-9}	1.8×10^{-9}	5.62×10^{-19}		
		Lung	6.9×10^{-9}	5.4×10^{-9}	8.03×10^{-19}		
		Liver or lung		7.2×10^{-9}		1.2×10^{-9}	9.2×10^{-9}

^aTissue specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue (liver or lung, respectively, for liver and lung tumors) per d; whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

^bGST-T1^{+/+} = homozygous, full enzyme activity); mixed = population reflecting estimated frequency of genotypes in current U.S. population: 20% GST-T^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^cEstimated at the human equivalent BMDL₁₀ (0.1/BMDL₁₀) (see Table 5-18).

^dEstimated at the human equivalent BMD₁₀ (0.1/BMD) (see Table 5-18).

^eCalculated as the square of the difference of the upper bound and central tendency IURs divided by the *t* statistic, 1.645.

^fCalculated as the square root of the sum of the variances for liver and lung tumors.

^gCalculated as the product of the cumulative tumor risk SD and the *t* statistic, 1.645, added to the sum of central tendency IURs.

Using this approach and the male mouse-derived risk factors, the combined human equivalent IUR values for both tumor types is $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$ (rounded from 1.3×10^{-8}) in the most sensitive (GST-T1^{+/+}) population. This is the recommended inhalation cancer unit risk value to be used in deterministic risk assessments for chronic exposure to dichloromethane. The corresponding value for a population with the frequency distribution of GST-T1 genotypes currently found in the U.S. population is $7 \times 10^{-9} (\mu\text{g}/\text{m}^3)^{-1}$.

5.4.2.6. Comparative Derivation Based on Rat Mammary Tumor Data

Mammary gland tumor data from male and female F344 rats following an inhalation exposure to dichloromethane were considered in development of a comparative IUR for dichloromethane (Mennear et al., 1988; NTP, 1986). In both the male and female rats, there were significant increases in the incidence of adenomas, fibroadenomas, or fibromas in or near the mammary gland. These were characterized as benign tumors in the NTP report (NTP, 1986). Increased numbers of benign mammary tumors per animal in exposed groups were also seen in two studies of Sprague-Dawley rats (Nitschke et al., 1988a; Burek et al., 1984). A gavage study in Sprague-Dawley rats reported an increased incidence of malignant mammary tumors, mainly adenocarcinomas (8, 6, and 18% in the control, 100, and 500 mg/kg dose groups, respectively), but the increase was not statistically significant. Data were not provided to allow an analysis that accounts for differing mortality rates (Maltoni et al., 1988). There are considerably more uncertainties regarding the interpretation of these data with respect to carcinogenic risk compared with the data pertaining to liver and lung tumors. The trends were driven in large part by benign tumors; adenocarcinomas and carcinomas were seen only in the females with incidences of 1, 2, 2, and 0 in the 0, 1,000, 2,000, and 4,000 ppm exposure groups, respectively. There are little data to guide the choice of relevant dose metric, and the genotoxicity and mechanistic studies have not included mammary tissue. For these reasons, the analysis and the calculation of the comparative IUR based on rat mammary tumor data are presented in Appendix G. The IUR based on the female rat data was $1 \times 10^{-7} (\mu\text{g}/\text{m}^3)^{-1}$.

5.4.2.7. Alternative Based on Administered Concentration

Another comparison that can be made is with an alternative IUR based on liver and lung tumors in mice using the external concentrations of dichloromethane in the mouse studies as converted to HECs, and then applying this using BMD modeling to obtain the BMDL₁₀ and resulting IUR. Mouse bioassay exposures were adjusted to HECs as follows:

- Adjusting to continuous exposure: $\text{External concentration}_{\text{ADJ}} = \text{External concentration} \times (6 \text{ hours}/24 \text{ hours}) \times (5 \text{ days}/7 \text{ days});$
- Concentrations in $\text{mg}/\text{m}^3 = \text{concentrations in ppm} \times 84.93/24.45;$ and

- $[H_{b/g}]_A/[H_{b/g}]_H$ = the ratio of blood:gas (air) partition coefficients in animals and humans. Because the partition coefficient for mice (23.0) is higher than for humans (9.7), a value of 1.0 was used, as per U.S. EPA (1994b) guidance.

Thus, $HECs = \text{External concentration}_{ADJ} \times [H_{b/g}]_A/[H_{b/g}]_H = \text{External concentration}_{ADJ} \times 1$.

The HECs (mg/m^3) for the 0, 2,000, and 4,000 ppm exposure groups were 0, 1,241, and 2,481 mg/m^3 , respectively. The BMD modeling and IURs derived from these values, in conjunction with the liver and lung tumor data from Table 5-17 (NTP, 1986), are shown in Table 5-21. The resulting IURs based on the liver or lung tumors in the mouse are approximately one order of magnitude higher than the currently recommended value obtained by using the mouse and human PBPK models.

Table 5-21. Inhalation units risks based on human BMDL₁₀ values using administered concentration for liver and lung tumors in B6C3F₁ mice exposed by inhalation to dichloromethane for 2 years

Sex, tumor type	BMS model ^a	χ^2 goodness of fit p-value	BMD ₁₀ ^b	BMDL ₁₀ ^b	Inhalation unit risk ^c ($\mu\text{g}/\text{m}^3$) ⁻¹
Male, liver	MS (0,1)	0.37	463.89	276.15	3.6×10^{-7}
Male, lung	MS (0,1)	0.54	157.23	124.10	8.1×10^{-7}
Female, liver	MS (0,2)	0.38	601.84	342.83	2.9×10^{-7}
Female, lung	MS (0,1)	0.77	126.40	100.61	9.9×10^{-7}

^aThe multistage (MS) model in EPA BMDS version 2.0 was fit to each of the four sets of mouse dose-response data shown in Table 5-17. The HEC used in these models for the 0, 2,000, and 4,000 ppm exposure groups were 0, 1,241, and 2,481 mg/m^3 , respectively. Numbers in parentheses indicate: (1) the number of dose groups dropped in order to obtain an adequate fit, and (2) the lowest degree polynomial of the model showing an adequate fit.

^bBMD₁₀ and BMDL₁₀ refer to the BMD-model-predicted HECs (mg dichloromethane per cubic meter), and its 95% lower confidence limit associated with a 10% extra risk for the incidence of tumors.

^cIUR ($\text{risk}/\mu\text{g}\cdot\text{m}^3$) = 0.1/human BMDL₁₀.

Sources: Mennear et al. (1988); NTP (1986).

The difference between the administered concentration methodology and PBPK-based approaches depends on two key differences: the use of a dichloromethane-metabolite dose-metric rather than dichloromethane AUC, and the fact that the rate of dichloromethane conversion to that metabolite is estimated in humans by using human data rather than default allometric scaling ($BW^{0.75}$). In addition, the administered concentration methodology does not account in any way for the GST polymorphism and so might be considered as representing the general/mixed-GST-genotype population rather than the +/+ subpopulation.

5.4.2.8. Previous IRIS Assessment: Cancer IUR

The IUR in the previous IRIS assessment was determined from the combined incidence of liver and lung adenomas and carcinomas in B6C3F₁ mice exposed to dichloromethane for 2 years by NTP (1986). A value of $4.7 \times 10^{-7} (\mu\text{g}/\text{m}^3)^{-1}$ was derived by the application of a modified version of the PBPK model of Andersen et al. (1987), which incorporated the pharmacokinetics and metabolism of dichloromethane. Internal dose estimates based on dichloromethane metabolism via the GST pathway were used and corrected for differences in interspecies sensitivity by applying to the human risks an interspecies scaling factor of 12.7, which was based on dose equivalence adjusted to BW to the $2/3$ power (Rhomberg, 1995; U.S. EPA, 1987a).

5.4.2.9. Comparison of Cancer IUR Using Different Methodologies

In this assessment, cancer IURs derived by using different dose metrics and assumptions were examined, as summarized in Table 5-22. The recommended IUR value of $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$ is based on a tissue-specific, GST-internal dose metric with allometric scaling because of the evidence for the involvement of highly reactive metabolites formed via the GST pathway. The value derived specifically for the GST-T1^{+/+} population is recommended to provide protection for the population that is hypothesized to be most sensitive to the carcinogenic effect. The values based on the GST-T1^{+/+} group are approximately two to fivefold higher than those for the full population for all dose metrics used in this assessment. Within a genotype population, the values of the IUR among the various dose metrics vary by about one to two orders of magnitude.

Table 5-22. Comparison of IURs derived by using various assumptions and metrics

Population ^a	Dose metric	Species, sex	Tumor type	Scaling factor	IUR ^b ($\mu\text{g}/\text{m}^3$) ⁻¹	Source (Table)
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate^c	Mouse, male	Liver and lung	7.0	1.3×10^{-8}	Table 5-20
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Liver	7.0	8.5×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Lung	7.0	5.6×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Liver and lung	1.0	1.9×10^{-9}	Table 5-20
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	1.2×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Lung	1.0	8.0×10^{-10}	Table 5-19
GST-T1 ^{+/+}	Whole-body GST metabolism rate	Mouse, male	Liver and lung	7.0	1.6×10^{-8}	Table 5-20
GST-T1 ^{+/+}	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	5.5×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Whole-body GST metabolism rate	Mouse, male	Lung	7.0	1.2×10^{-8}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver and lung	7.0	7.4×10^{-9}	Table 5-20
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver	7.0	4.8×10^{-9}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Lung	7.0	3.2×10^{-9}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver and lung	1.0	1.1×10^{-9}	Table 5-20
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	6.8×10^{-10}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Lung	1.0	4.5×10^{-10}	Table 5-19
Mixed	Whole-body GST metabolism rate	Mouse, male	Liver and lung	7.0	9.2×10^{-9}	Table 5-20
Mixed	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	3.1×10^{-9}	Table 5-19
Mixed	Whole-body GST metabolism rate	Mouse, male	Lung	7.0	6.9×10^{-9}	Table 5-19
	Administered concentration (HEC)	Mouse, male	Liver		3.6×10^{-7}	Table 5-21
	Administered concentration (HEC)	Mouse, male	Lung		8.1×10^{-7}	Table 5-21
	1995 IRIS assessment ^c	Mouse, male	Liver and lung	12.7	4.7×10^{-7}	

^aGST-T1^{+/+} = homozygous, full enzyme activity; mixed = genotypes based on a population reflecting the estimated frequency of genotypes in the current U.S. population: 20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^bBased on value of the derived distributions.

^cBolded value is the basis for the recommended IUR of $1 \times 10^{-8} \mu\text{g}/\text{m}^3$ per mg/kg-d.

5.4.3. Differences Between Current Assessment and Previous IRIS PBPK-based Assessment

To better understand the changes in assessment risk predictions between previous EPA evaluations and the current assessment, the differences in PBPK model parameters for the B6C3F₁ mouse were evaluated. Values that differed significantly between the model version used previously and that of Marino et al. (2006), along with derived group parameters that lend further insight, are shown in Table 5-23.

Table 5-23. Comparison of key B6C3F₁ mouse parameters differing between prior and current PBPK model application

Parameter ^a	Marino et al. (2006); mean values as applied (posterior)	U.S. EPA (1988b, 1987a, b)
<i>Partition coefficients</i>		
PB blood/air	23	8.29
PF fat/blood	5.1	14.5
PF·PB (fat/blood)·(blood/air) = fat/air	117.3	120.2
PL liver/blood	1.6	1.71
PL·PB (liver/blood)·(blood/air) = liver/air	36.8	14.2
PLu lung (tissue)/blood	0.46	1.71
PLu·PB (lung/blood)·(blood/air) = lung/air	10.6	14.2
PR rapidly perfused/blood	0.52	1.71
PR·PB rapidly perfused/air	12.0	14.2
PS slowly perfused/blood	0.44	0.96
PS·PB slowly perfused/air	10.1	7.96
<i>Flow rates</i>		
QCC cardiac output (L/hr/kg ^{0.74})	24.2	14.3
VPR ventilation:perfusion ratio	1.45	1.0
<i>Metabolism parameters</i>		
V _{maxC} maximum CYP metabolic rate (mg/hr/kg ^{0.7})	9.27	11.1
K _m CYP affinity (mg/L)	0.574	0.396
V _{maxC} /K _m (L/hr/kg ^{0.7})	16.1	28
A1 ratio of lung V _{maxC} to liver V _{maxC}	0.207	0.416
Total lung + liver V _{maxC} /K _m	19.5	39.7
k _{fC} first-order GST metabolic rate constant (kg ^{0.3} /hr)	1.41	1.46
A2 ratio of lung k _{fC} to liver k _{fC}	0.196	0.137
Total lung + liver k _{fC}	1.69	1.66

^aParameters not listed differed by <10% between versions. See Table 3-5 and associated text for details.

While a number of the tissue:blood partition coefficients in Table 5-23 differ significantly between the two models (e.g., PF, PLu, and PR), the corresponding tissue:air coefficients (e.g. the products PF·PB, PLu·PB, and PR·PB) generally do not. Since the latter tend to determine the long-term equilibration between the tissue (tissue group) and air, the differences in the tissue:blood coefficients are not expected to significantly impact long-term risk predictions. Thus, the partition coefficients that most significantly differ (the blood:air and liver:air partition coefficients) are, respectively, 2.8- and 2.6-fold higher in the current version. The increased PB results in a tendency for simulated blood concentrations to rise more quickly and reach higher

values with other parameters being equal. The significantly increased QCC and VPR contribute even more to this difference, resulting in an even faster rise to steady state during inhalation exposure simulations but also more rapid delivery to the liver (decreasing blood-flow limitation of hepatic metabolism) and more rapid exhalation. The increased liver:air partition coefficient leads to higher predicted liver concentrations (again, other parameters being equal) and hence, higher rates of metabolism.

For metabolism, a much reduced oxidative metabolism is seen, which at low doses depends on $V_{\max C}/K_m$. The revised hepatic metabolism is over 40% lower and the total of lung + liver metabolism is 50% lower than previously used. This lower rate of metabolism means that far less of parent dichloromethane will be removed through metabolism and hence, predicted blood concentrations will be higher still relative to the impact of changes in partition coefficient, QCC, and VPR, as noted above.

The result of having higher predicted blood and liver dichloromethane concentrations is that, while the GSH-pathway metabolic constant, k_{TC} , is virtually the same for the mouse liver in both cases, the much higher concentration of dichloromethane available will lead to a much higher predicted rate of metabolism via this pathway. For the lung, since the A2 is 43% higher in the model of Marino et al. (2006), the relative increase will be even greater.

Because the revised rate of GST metabolism in mice was estimated by using data with CYP2E1 inhibited by a suicide inhibitor, there is considerable confidence in the relative rate of metabolism by these two pathways and the GST pathway in particular. The partition coefficients used by Marino et al. (2006) are as measured by Clewell et al. (1993) and expected to be at least as reliable as those used in the 1995 assessment. Considering that the revised PBPK model does an excellent job of reproducing closed-chamber gas uptake data that were not available for calibration of the 1987 model, as well as blood concentrations after intravenous injection, there is fairly high confidence in its predictions.

The net result of these model changes is that, under mouse bioassay conditions, the predicted dose metrics for liver and lung cancer (i.e., GST-mediated metabolism) are higher than those obtained with the previous model, resulting in a lower risk estimated per unit of dose.

The other piece of the PBPK-based risk estimation is the human model. In updating the parameter estimates for the human model (see Appendix B for details), the oxidative metabolism $V_{\max C}/K_m$ approximately doubled, which leads to lower predicted blood concentrations of dichloromethane available for metabolism by GST. In addition, the liver GST was reduced by almost 60%, and the lung:liver GST ratio decreased by almost fivefold, for a net change in lung GST of over 90%. Given the larger human data set available to David et al. (2006) and the sophisticated Bayesian analysis used to recalibrate the model, the expectation is that these values are more reliable than the values used in the 1995 IRIS assessment.

Since actual rates of metabolism at a given exposure level also depend on respiration rate and blood flows, these changes in metabolic parameters do not completely determine the relative

(predicted) dosimetry. But the difference in cancer risk predictions between the current and previous assessments is primarily explained by the overall prediction of higher GST-mediated dosimetry in the mouse (at bioassay conditions) and lower human GST metabolism (due in part to greater predicted clearance of dichloromethane by oxidative metabolism). In addition to these changes in PBPK parameters, the reduction of scaling factor from 12.7 to 7 is a significant factor in the overall change from the previous assessments.

5.4.4. Application of Age-Dependent Adjustment Factors (ADAFs)

The available dichloromethane studies do not include an evaluation of early-life susceptibility to dichloromethane cancer risk. In the absence of this type of data, and if a chemical follows a mutagenic mode of action for carcinogenicity like dichloromethane, the *Supplemental Guidance for Assessing Susceptibility from Early-Life Exposure to Carcinogens* (U.S. EPA, 2005b) recommends that ADAFs be applied to the cancer values. Since the OSF of $1 \times 10^{-3} \text{ (mg/kg-day)}^{-1}$ and the IUR of $1 \times 10^{-8} \text{ (}\mu\text{g/m}^3\text{)}^{-1}$ were calculated from adult dichloromethane exposures, early-life cancer susceptibility has not been accounted for in these values, and ADAFs need to be applied in combination with exposure information when estimating cancer risks that include early-life exposures. Sample calculations that incorporate ADAFs into the cancer risks are presented in subsequent sections. Additional examples of evaluations of cancer risks incorporating early-life exposure are provided in Section 6 of the *Supplemental Guidance* (U.S. EPA, 2005b).

In the *Supplemental Guidance* (U.S. EPA, 2005b), ADAFs are established for three age groups. An ADAF of 10 is applied for age groups <2 years, 3 is applied for ages 2–<16 years, and 1 is applied for ≥ 16 years (U.S. EPA, 2005b). The 10- and 3-fold adjustments in cancer values are combined with age-specific exposure estimates when early-life exposure considerations need to be included in cancer risk estimates. The most current information on usage of ADAFs can be found at <http://www.epa.gov/cancerguidelines>. For estimation of risk, the *Supplemental Guidance* (U.S. EPA, 2005b) recommends obtaining and using age-specific values for exposure and cancer potency. In the absence of age-specific cancer potency values, as is true for dichloromethane, age-specific cancer values are estimated by using the appropriate ADAFs. Using this process, a cancer risk is derived for each age group. The risks are summed across the age groups to get the total cancer risk for the age-exposure period of interest.

5.4.4.1. Application of ADAFs in Oral Exposure Scenarios

Sample calculations incorporating the use of ADAFs are presented for three exposure duration scenarios. These scenarios include full lifetime exposure (assuming a 70-year lifespan), and two 30-year exposures at ages 0–30 and ages 20–50. A constant dichloromethane exposure of 1 mg/kg-day was assumed for each scenario.

Table 5-24 lists the four factors (ADAFs, OSF, assumed exposure, and duration adjustment) that are needed to calculate the partial cancer risk based on the early age-specific group. The partial cancer risk for each age group is the product of the four factors in columns 2–5. Therefore, the partial cancer risk following daily dichloromethane oral exposure in the age group 0 to <2 years is the product of the values in columns 2–5 or $10 \times (2 \times 10^{-3}) \times 1 \times 2/70 = 5.7 \times 10^{-4}$. The partial risks that are listed in the last column of Table 5-24 are added together to get the total risk. Thus, a 70-year (lifetime) risk estimate for continuous exposure to 1 mg/kg-day dichloromethane is 3.3×10^{-3} per mg/kg-day, which is adjusted for early-life susceptibility and assumes a 70-year lifetime and constant exposure across age groups.

Table 5-24. Application of ADAFs to dichloromethane cancer risk following a lifetime (70-year) oral exposure

Age group (yrs)	ADAF	Unit risk (per mg/kg-d)	Exposure concentration (mg/kg-d)	Duration adjustment	Partial risk
0–<2	10	2×10^{-3}	1	2 yrs/ 70 yrs	5.7×10^{-4}
2–<16	3	2×10^{-3}	1	14 yrs/ 70 yrs	1.2×10^{-3}
≥16	1	2×10^{-3}	1	54 yrs/ 70 yrs	1.5×10^{-3}
Total risk					3.3×10^{-3}

In calculating the cancer risk for a 30-year constant exposure to dichloromethane at an exposure level of 1 mg/kg-day from ages 0–30, the duration adjustments would be 2/70, 14/70, and 14/70, and the partial risks for the three age groups would be 5.7×10^{-4} , 1.2×10^{-3} , and 4.0×10^{-4} , which would result in a total risk estimate of 2.2×10^{-3} .

In calculating the cancer risk for a 30-year constant exposure to dichloromethane at an exposure level of 1 mg/kg-day from ages 20–50, the duration adjustments would be 0/70, 0/70, and 30/70. The partial risks for the three groups are 0, 0, and 8.6×10^{-4} , which would result in a total risk estimate of 8.6×10^{-4} .

5.4.4.2. Application of ADAFs in Inhalation Exposure Scenarios

Sample calculations incorporating the use of ADAFs are presented for three exposure duration scenarios involving inhalation exposure. These scenarios include full lifetime exposure (assuming a 70-year lifespan) and two 30-year exposures from ages 0–30 and ages 20–50. A constant dichloromethane inhalation exposure of $1 \mu\text{g}/\text{m}^3$ was assumed for each scenario.

Similar to the oral exposure scenarios presented in Section 5.4.4.1, Table 5-25 lists the four factors (ADAFs, unit risk, assumed exposure, and duration adjustment) that are needed to calculate the partial cancer risk based on the early age-specific group. The partial cancer risk for each age group is the product of the four factors in columns 2–5. Therefore, the partial cancer

risk following daily dichloromethane inhalation exposure in the age group 0 to <2 years is the product of the values in columns 2–5 or $10 \times (1 \times 10^{-8}) \times 1 \times 2/70 = 2.9 \times 10^{-9}$. The partial risks that are listed in the last column of Table 5-25 are added together to get the total risk. Thus, a 70-year (lifetime) risk estimate for continuous exposure to $1 \mu\text{g}/\text{m}^3$ dichloromethane is 1.8×10^{-8} per $\mu\text{g}/\text{m}^3$, which is adjusted for early-life susceptibility and assumes a 70-year lifetime and constant exposure across age groups.

Table 5-25. Application of ADAFs to dichloromethane cancer risk following a lifetime (70-year) inhalation exposure

Age group (yrs)	ADAF	Unit risk (per mg/kg-d)	Exposure concentration (mg/kg-d)	Duration adjustment	Partial risk
0–<2	10	1×10^{-8}	1	2 yrs/ 70 yrs	2.9×10^{-9}
2–<16	3	1×10^{-8}	1	14 yrs/ 70 yrs	6.0×10^{-9}
≥ 16	1	1×10^{-8}	1	54 yrs/ 70 yrs	7.7×10^{-9}
Total risk					1.7×10^{-8}

In calculating the cancer risk for a 30-year constant exposure to dichloromethane at a level of $1 \mu\text{g}/\text{m}^3$ from ages 0–30, the duration adjustments would be 2/70, 14/70, and 14/70, and the partial risks for the three age groups are 2.9×10^{-9} , 6.0×10^{-9} , and 2.0×10^{-9} . These partial risks result in a total risk estimate of 1.1×10^{-8} .

In calculating the cancer risk for a 30-year constant exposure to dichloromethane at a level of $1 \mu\text{g}/\text{m}^3$ from ages 20–50, the duration adjustments would be 0/70, 0/70, and 30/70, and the partial risks for the three age groups are 0, 0, and 4.3×10^{-9} , resulting in a total risk estimate of 4.3×10^{-9} .

5.4.5. Uncertainties in Cancer Risk Values

The derivation of cancer risk values often involves a number of uncertainties in the extrapolation of dose-response data in animals to cancer risks in human populations. Several types of uncertainty have been quantitatively integrated into the derivation of the recommended OSFs and IURs for dichloromethane, while others are qualitatively considered. Table 5-26 and the ensuing discussion summarize the principal uncertainties identified, their possible effects on the cancer risk values, and decisions made in the derivations.

Table 5-26. Summary of uncertainty in the derivation of cancer risk values for dichloromethane

Consideration and Impact On Cancer Risk Value	Decision	Justification and Discussion
Selection of data set (Selection of an alternative data set could change the recommended cancer risk values.)	Select NTP (1986) as principal inhalation study and Serota et al. (1986b) as principal oral (drinking water) study to derive cancer risks for humans	NTP (1986) inhalation mouse bioassay provides the strongest cancer responses (liver and lung tumors) and the best dose-response data in the animal database. The oral mouse study (Serota et al. (1986b; Hazleton Laboratories, 1983) provide an adequate basis for dose-response modeling
Selection of target organ (Selection of a target organ could change the recommended cancer risk values.)	Liver, and for inhalation, lung selected as target organ. Cancer risk values based on mammary gland tumors in rats also examined; potential brain cancer risk was identified as a data gap.	The evidence for mammary gland tumors from dichloromethane exposure is less consistent than evidence for liver and lung tumors. Inhalation cancer risk values based on mammary tumors in rats are about one order of magnitude higher than risk values based on liver or lung tumors in mice, but
Selection of extrapolation approach (Selection of extrapolation approach could change the recommended cancer risk values.)	Oral data used for OSF and inhalation data used for IUR. Oral cancer risk values based on route-to-route extrapolation from inhalation study also examined.	Derivation from oral exposure is preferred to route-to-route extrapolation when adequate oral data are available. Oral cancer risk values based on route-to-route extrapolation from the NTP (1986) inhalation mouse study were about one order of magnitude lower than values based on oral exposure study
Selection of dose metric (Selection of dose metric could change the recommended cancer risk values.)	Use tissue-specific GST-metabolism dose metric. Cancer risk estimates based on alternative (whole-body) metrics also examined.	Contribution of CYP pathway to cancer risk unknown, but strong evidence of GSTrole in carcinogenesis supports focus on this pathway. Values based on tissue-specific GSTmetabolism recommended based on evidence of site locality of effects.
Dose-response modeling (Human risk values could increase or decrease, depending on fits of alternative models)	Use multistage dose-response model to derive a BMD	The multistage model has biological support and is the model most consistently used in EPA cancer assessments.
Low-dose extrapolation (Human risk values would be expected to decrease with the application of nonlinear tumor responses in low-dose regions of dose-response curves.)	Use linear extrapolation of risk in low-dose region	PBPK model incorporates the metabolic shift and expected nonlinearity (GST dose attenuation) in the exposure-dose relationship across exposure levels. Linear low-dose extrapolation for agents with a mutagenic mode of action is supported.
Interspecies extrapolation of dosimetry and risk (Alternative values for PBPK model parameters and cross-species scaling factor could increase or decrease human cancer risk values.)	Use PBPK model and allometric scaling factor for the primary dose metric	Use of rodent and human PBPK models reduced uncertainty due to interspecies differences in toxicokinetics. Examination of impact of different values for key parameters in human model, and sensitivity analysis of rodent PBPK model parameters identified influential metabolic parameters for which limited experimental data exist

(Table 5-26 continues on next page)

Table 5-26. Summary of uncertainty in the derivation of cancer risk values for dichloromethane

Consideration and Impact On Cancer Risk Value	Decision	Justification and Discussion
Sensitive subpopulations (Differences in CYP and GST metabolic rates could change cancer risk values.)	Risk estimates generated for presumed most sensitive (GST-T1 ^{+/+}) genotype; CYP variability incorporated into PBPK model	No data are available to determine the range of human toxicodynamic variability or sensitivity, including whether children are more sensitive than adults.

Data selections for derivation of IUR and OSF. The database of animal bioassays identifies the liver and lung as the most sensitive target organs for dichloromethane-induced tumor development. These effects demonstrate a dose-response relationship in mice exposed orally (liver, males only) or by inhalation (liver and lung, males and females). The liver cancer effects seen in the oral exposure study (Serota et al., 1986b; Hazleton Laboratories, 1983) were not strong (increasing from approximately 20% in controls to 30% in the highest 3 dose groups), but rather can be characterized as by marginally increased (trend test $p = 0.058$) combined hepatocellular adenomas and carcinomas and by statistically significantly increased ($p < 0.05$) hepatocellular adenomas and carcinomas at dose levels of 125, 185, and 250 mg/kg-day. These data are considered adequate for dose-response modeling. Although EPA's interpretation of these data differs from that of the study authors, the reasons for this difference were described in Section 4.2.1.2.2. In addition, as shown in Table 4-43, the lower incidence of liver tumors induced by the highest doses used in the oral exposure study (Serota et al., 1986b; Hazleton Laboratories, 1983) compared with the higher incidence induced by inhalation exposure to 2,000 ppm (NTP, 1986) is consistent with the predicted lower liver dose of GST metabolites (and hence lower probability of DNA modification) with oral exposure.

Statistically significant increases in benign mammary gland tumors were observed in one study of F344 rats exposed by inhalation to 2,000 or 4,000 ppm (Mennear et al., 1988; NTP, 1986), and evidence for a tumorigenic mammary gland response in Sprague-Dawley rats was limited to increased numbers of benign mammary tumors per animal at levels of 50–500 ppm (Nitschke et al., 1988a) or 500–3,500 ppm (Burek et al., 1984). A gavage study in female Sprague-Dawley rats reported an increased incidence of malignant mammary tumors, mainly adenocarcinomas (8, 6, and 18% in the 0, 100, and 500 mg/kg dose groups, respectively), but the increase was not statistically significant. Data were not provided to allow an analysis that accounts for differing mortality rates (Maltoni et al., 1988). The toxicokinetic or mechanistic events that might lead to mammary gland tumor development in rats are unknown, although CYP2E1 (El-Rayes et al., 2003; Hellmold et al., 1998) and GST-T1 expression have been detected in human mammary tissue (Lehmann and Wagner, 2008).

Rare CNS tumors were observed in one study in rats spanning a relatively low range of exposures (0–500 ppm). These cancers were not seen in two other studies (NTP, 1986; Burek et

al., 1984) in rats, both involving higher doses (1,000–4,000 ppm), or in a similar high-dose study (NTP, 1986) in mice. The relative rarity of the tumors precludes the use of the low-dose exposure study in a quantitative dose-response assessment.

The *in vivo* genotoxicity and mechanistic data in rodents provide a detailed sequence of steps from generation of reactive metabolites to mutagenic effects, such as DNA-protein cross-links and DNA strand breaks. Further, the toxicokinetic pathways implicated in production of the putative carcinogenic metabolites in animals also exist in humans. Thus, there is high confidence that the dose-response data for liver and lung cancer in mice represents the best data currently available for derivation of human cancer risks. A more complete understanding of the carcinogenic potential of dichloromethane would be achieved by addressing data gaps identified with respect to issues regarding potential risk and mechanisms relating to brain cancer and mammary tumors.

Target organ. The liver and lung tumor incidence from chronic exposure bioassays provide clear evidence of the carcinogenic potential of dichloromethane exposure. The bioassays are supported by a substantial database of genotoxicity and mechanistic studies (summarized in Section 4.5). The evidence for mammary gland tumors from dichloromethane exposure is based primarily on observations of benign tumors in rats with inhalation exposure (NTP, 1986). Derivation of cancer potency values based on these data are presented in Appendix G. The potential brain cancer risk, suggested by a limited number of these relatively rare tumors in both animal and human studies, is identified as a data gap which would benefit from additional research.

Extrapolation approach. A route-to-route extrapolation from the NTP (1986) inhalation mouse bioassay was used to develop an oral cancer slope value for purposes of comparison. Although the exposure-response effect seen in the oral exposure study is not strong, the direct derivation from oral exposure is preferred to the route-to-route extrapolation when adequate oral data are available.

The comparison of the OSF derived from the oral exposure data and from the route-to-route extrapolation from the inhalation data provides a crude measure of the uncertainty in recommending a human OSF based on the available rodent bioassay data. The cancer OSF based on route-to-route extrapolations from liver tumors in mice exposed by inhalation are about an order of magnitude lower than those based on the liver tumor responses in mice exposed via drinking water. This difference may be explained, at least partially, by the heterogeneity of hepatic cell types within the sinusoid, resulting in regio-specific toxicity. Oral exposure may result in a higher internal exposure of hepatocytes in the periportal region (particularly those lining the portal vein, through which all gastrointestinal-absorbed dichloromethane passes) than in the centrilobular region (SRC, 1989). Further, the metabolic capacity of hepatic cells is also

regio-specific, with higher CYP activity found in the centrilobular region compared to the periportal region. Thus, liver perfusion via the systemic arterial circulation or portal drainage of the gastrointestinal tract, through which inhaled dichloromethane would be introduced, may influence regio-specific hepatotoxicity, resulting in the route-of-exposure effects on toxicity. The available PBPK models do not have the capability to predict regio-specific disposition of dichloromethane in the liver.

Dose metric. There is considerable data supporting the role of GST-related metabolism of dichloromethane in carcinogenicity, as described in Sections 4.5.1 and 4.7. Pretreatment of mice with buthionine sulphoximine, a GSH depletor, caused a decrease to levels seen in controls in the amount of DNA damage detected immediately after in vivo exposure in liver and lung tissue (Graves et al., 1995). Although the results of Landi et al. (2003) indicate that GST activity is not needed for the observation of DNA damage by the comet assay from some trihalomethanes (e.g., bromodichloromethane), the results for dichloromethane were much weaker and of uncertain significance.

Dose-response modeling. The multistage model is used because of its biological relevance and because of the lack of information supporting a biologically-based or other particular model instead of the multistage model. The multistage model is the most commonly used model for cancer derivations and its use maintains comparability with existing assessments.



Because of the adequacy of the fit of the multistage model to the data, little modeling uncertainty would be expected to be introduced by the choice of this model. Application of the multistage model allowed for estimation of a POD in the lower region of exposure for observable cancer effects.

For human oral exposure, ingestion is assumed to occur as six discrete boluses during the course of the day: 25% of the daily dose consumed at 7 am, noon, and 6 pm; 10% at 10 am and 3 pm; and 5% at 10 pm. When exposure occurs as a bolus, the short-term (peak) concentration of dichloromethane will be higher, leading to a higher degree of CYP saturation and hence a higher fraction metabolized by GST, as compared to more continuous exposure such as occurs by inhalation. Thus if actual ingestion is in fewer/larger boluses than those assumed, the cancer risk will be somewhat under-predicted. On the other hand if ingestion is in more/smaller boluses, the opposite will occur. However, when ingestion is fairly small, such that the peak concentration is well below the saturation constant (K_m) for CYP, the difference in metabolic outcome will be negligible. The pattern used here assumes in effect that the amount of food and liquid ingested is divided neatly into meals and snacks or breaks as indicated *and* that the concentration of dichloromethane in the food and beverages ingested is constant. Thus if one meal or drink happens to include the bulk of that ingested for a day, total ingestion will be more like a single daily bolus. But to the extent that people sip beverages or ingest foods over longer periods of

time, actual ingestion will be more continual. Given that both of these are likely to occur to some extent, the population ingestion pattern is expected to be a distribution that includes the one used for simulation purposes. While it cannot be said that this pattern is an exact average, given that the differences in saturation at low total exposure levels will be small, it is considered sufficiently representative of the population and the uncertainty resulting from inexact knowledge of actual ingestion is unlikely to be significant.

Low-dose extrapolation. The mode of action is a key consideration in determining how risks should be estimated for low-dose exposure. The *in vitro* and *in vivo* genotoxicity data suggest that mutagenicity is the most plausible mode of action, although key mutagenic events in the development of liver or lung tumors have not been identified. Because it was concluded that dichloromethane acts through a mutagenic mode of action, a linear-low-dose extrapolation approach was used to estimate OSFs and IURs.

While the rate equation for GST metabolism in the PBPK model is first-order, consistent with metabolism increasing in direct proportion to the concentration of dichloromethane, because the GST and CYP pathways compete for dichloromethane and the CYP kinetics are nonlinear (saturable), the interaction of the two pathways within the whole-body system results in a nonlinear exposure-dose relationship for both pathways. These nonlinearities are demonstrated for a simulated group of 30-year-old women (population mean kinetics for continuous inhalation exposure) in Figure 5-16. The upper panel (A) of Figure 5-16 provides a full-scale plot of CYP liver metabolism (mg/L liver/d) up to 2000 ppm exposure while the lower panel expands the CYP metabolism curve up to 400 ppm exposure (with CYP metabolism still included, but not far off the the y-axis with that scale). Because both CYP and GST metabolism are linear at very low concentrations (below 10-30 ppm), the exposure-response relationship at low exposures is linear for both pathway metrics, initially increasing from zero dose at zero concentration in direct proportion to the exposure level until CYP saturation begins. As CYP becomes saturated, starting around 50 ppm and reaching half saturation at around 200 ppm, a lower fraction of dichloromethane is eliminated by CYP metabolism. As a lower fraction is metabolized by CYP the blood concentration increases faster than directly proportional to exposure concentration with the result that GST metabolism also increases faster than direct proportionality: the upward curvature seen in the lower panel (B) of Figure 5-16. While GST metabolism remains less than CYP over the entire exposure range shown here for humans, in mice, where the GST pathway has relatively higher activity compared to CYP, GST metabolism increases above CYP metabolism in the range of bioassay exposures. This transition from CYP-dominated (or vastly-dominated) clearance at low exposures to a higher fraction of GST metabolism at high exposures has at times been referred to as a “switch,” but as shown in Figure 5-16 the transition is smooth and continuous, and there is some GST metabolism at all exposure levels with the exposure-response approaching linearity, without a threshold, at low exposures.



The curves represent average results for a simulated population of 1000 women with the GST-T1 +/+ genotype. A) Relationships scaled to show full range of CYP metabolism up to 2,000 ppm inhalation exposure. B) Relationships scaled to show low-dose linearity (below 50 ppm) and curvature (transition) in GST metabolism (above 50 ppm), as CYP metabolism saturates.

Figure 5-16. PBPK-model-predicted exposure-response relationships for hepatic CYP and GST metabolism for continuous inhalation exposure to dichloromethane in 30-year-old GST +/+ women.

One other important note is that in calculating IUR for humans the relationship between external exposure and internal dose was determined using the PBPK model at a very low level of exposure (i.e., $1 \mu\text{g}/\text{m}^3$ or 0.00029 ppm), where the relationship is effectively linear:

the difference between the actual exposure-dose curve and a straight line is less than 1%. However as one goes to higher concentrations the relationship becomes significantly nonlinear, and hence application of the cancer toxicity values (IUR) will not accurately represent the risk. Because GST metabolism increases faster than proportional to exposure level with concentration, in fact the IUR will under-predict risk at those higher exposure levels. Analysis of the PBPK model versus the low-exposure linear estimate shows that the extent of nonlinearity is less than 20% for oral exposures at low doses and for inhalation exposures at less than 30 ppm (100 $\mu\text{g}/\text{m}^3$). The dose used for calculating the internal dose:exposure ratio for oral exposures, 1 mg/kg-d, was above the transition to nonlinear dosimetry, but only to a small extent. For oral exposures the linear approximation used differed from the full model by less than 30% for exposures less than 2 mg/kg-d, but at doses below 1 mg/kg-d the error would be in the direction of an over-prediction of risk (i.e., actual cancer risks may be somewhat lower, but no more than 1.3-fold lower), than indicated by the linear model.

Interspecies extrapolation of dosimetry and risk. Target organ dosimetry for neoplastic mouse responses and estimation of equivalent internal human doses were accomplished using PBPK models for dichloromethane in mice and humans. Uncertainty in the ability of the PBPK models to estimate animal and human internal doses from lifetime bioassay low-level environmental exposures may affect the confidence in the cancer risk extrapolated from animal data. Uncertainties in the mouse and human model parameter values were integrated quantitatively into parameter estimation by utilizing hierarchical Bayesian methods to calibrate the models at the population level (David et al., 2006; Marino et al., 2006). The use of Monte Carlo sampling to define human model parameter distributions allowed for derivation of human distributions of dosimetry and cancer risk, providing for bounds on the recommended risk values.

A detailed discussion of PBPK model structure (CYP rate equation) and parameter uncertainties is provided in Sections 3.5.2 and 3.5.5, respectively. While the structure and equations used in the existing model have been described in multiple peer-reviewed publications over the past two decades, there are discrepancies between dichloromethane kinetics observed in vitro and the model parameters obtained from in vivo data, and the model poorly fits some of the in vivo data (e.g., fraction of dose exhaled as CO at higher exposure levels in mice). The discrepancies are significant enough that simply re-estimating model parameters is unlikely to resolve them, but based on a limited analysis, it appears that an alternative (dual-binding-site) CYP metabolic equation (Korzekwa et al., 1998) may provide the resolution. At present, the suggestion of this alternate equation is a hypothesis which should be tested experimentally. Further, integration of the alternate rate equation into the PBPK modeling and then quantitative risk assessment will likely require several years of further research and, hence, is beyond the scope of the current assessment. Since the GST activity in the current model is within a factor of

three of that measured in vitro (when both are evaluated per gram of liver), the impact of that model uncertainty is also expected to be no more than a factor of three.

Also as detailed in Section 3.5.2, the results of David et al. (2006) for the GST-T1 activity parameter, k_{fC} , for the combined human data set appear to be discrepant from the range of results for each of the individual data sets. Therefore, sensitivity to the human PBPK parameter distributions was evaluated by rescaling the parameters to the mean values obtained by David et al. (2006) for a specific data set (DiVincenzo and Kaplan, 1981) for which the GST activity was intermediate among those obtained across individual data sets. When this was done, the upper bound estimates on GST dosimetry (for low, fixed inhalation or oral exposures) in the GST-T1 +/+ subpopulation increased by over an order of magnitude, as did the estimate of the mean activity for an inhalation exposure, although the estimated mean GST activity for an oral exposure only increased about twofold. Thus, while correspondence of the in vivo GST activity with that measured in vitro suggests a lower degree of quantitative uncertainty, it is possible that revision of the PBPK model could have a larger impact. The ultimate impact will depend on how revisions effect model predictions for both the animal and the human. If the predicted GST metabolism per unit exposure increases in both mice and humans by a similar factor, there will be little impact on the risk estimate. But if the GST activity predicted in the mouse is decreased by a factor of 3, while that in the human is increased by a factor of 3, for example, then the net impact would be an increase of ninefold in human risk estimates.

Sensitivity analysis of the mouse PBPK parameters. The mouse and rat PBPK models were utilized deterministically; i.e., the single-value parameter estimates for the rat PBPK model were used for rat dosimetry simulations, and the mean parameter estimates from the Bayesian analysis of Marino et al. (2006) were used for the mouse dosimetry simulations. To assess the effect of using point estimates of parameter values for calculation of rodent dosimetry, a sensitivity analysis was performed to identify model parameters most influential on the predictions of dose metrics used to estimate oral and inhalation cancer risks. As was described in the RfD and RfC sensitivity analysis calculation, this procedure used a univariate analysis in which the value of an individual model parameter was perturbed by an amount (Δ) in the forward and reverse direction (i.e., an increase and decrease from the nominal value), and the change in the output variable was determined. Results are for the effects of a perturbation of $\pm 1\%$ from the nominal value of each parameter on the output values at the end of a minimum of 10,000 simulated hours. This time was chosen to achieve a stable daily value of the dose metric, given that the simulated bioassay exposures did not include weekend exposures. The exposure conditions represented the lowest bioassay exposure resulting in significant increases in the critical effect. For inhalation exposures in mice, the PB, followed closely by the first-order GST-mediated metabolism rate (k_{fC}), had the greatest impact on the dose metric for liver cancer (mg dichloromethane metabolized via GST pathway per liter liver per day) (Figure 5-17). For

drinking water exposures in mice, the k_{fC} , followed by the CYP-mediated maximum reaction velocity (V_{maxC}), affected the liver cancer dose metric to the greatest extent (Figure 5-18). For mice inhaling dichloromethane, the lung cancer dose metric (mg dichloromethane metabolized via GST pathways per liter lung per day), like the liver cancer metric, was highly affected by the k_{fC} and the PB (Figure 5-19). However, since GST-mediated lung metabolism is calculated as a constant fraction of the liver metabolism rate ($A2 \times k_{fC}$), the lung cancer dose metric was most sensitive to the proportional yield of liver GST-mediated metabolic activity attributed to the lung. The PB was experimentally determined, lending high confidence to its value. Values for the three metabolic parameters were determined by computational optimization against data sets not directly measuring dichloromethane or its metabolites in the target/metabolizing tissues. It is uncertain how alternative values for these three parameters would affect the estimation of animal $BMDL_{10}$ values and, ultimately, the OSFs and IURs.




Figure 5-17. Sensitivity coefficients for long-term mass GST-mediated metabolites per liver volume from a long-term average daily inhalation concentration of 2,000 ppm in mice.




Figure 5-18. Sensitivity coefficients for long-term mass GST-mediated metabolites per liver volume from a long-term average daily drinking water concentration of 500 mg/L in mice.




Figure 5-19. Sensitivity coefficients for long-term mass GST-mediated metabolites per lung volume from a long-term average daily inhalation concentration of 500 ppm in mice.

There is uncertainty as to whether the reactivity of the toxic dichloromethane metabolites is sufficiently high enough to preclude systemic distribution. Therefore, alternative derivations of cancer risk values were performed under the assumption that high reactivity leads to complete clearance from the tissue in which the active metabolite is formed (scaling factor = 1.0). The difference in scaling factor (7.0 for allometric scaling versus 1.0) results in a sevenfold decrease in estimated cancer toxicity values. Using a whole-body GST metabolism dose metric, the resulting OSF and IUR for liver cancer was approximately fivefold lower than when tissue-specific dose metrics were used (Table 5-16 and Table 5-22); however, the IURs for lung cancer and for the combined liver and lung cancer risk were higher with the whole-body compared with the tissue-specific metric (Table 5-22). This difference reflects the lower metabolism that occurs in human versus mouse lung (relative to total); lung-specific metabolism is lower in humans than mice, so the predicted risk in the lung is lower when based on that metabolism versus when whole-body metabolism is used. The mechanistic data support the hypothesis that reactive metabolites produced in the target tissues do not distribute significantly beyond those tissues and cause deleterious effects in the metabolizing tissues soon after generation. Thus, there is less uncertainty in the cancer risk values derived by using a tissue-specific GST metabolism dose metric compared with those derived using a whole-body GST metabolism dose metric.

Sensitive human populations. Possible sensitive populations include persons with altered CYP (e.g., obese individuals, alcoholics, diabetics, and the very young) and GST (e.g., GST-T1 homozygous conjugators) metabolic capacity. The PBPK model includes an estimate of the

variability of CYP metabolism (sixfold variation) within the general population but does not specifically address what could be greater variation in these other groups. However, the known polymorphisms for GST-T1 expression were integrated into the human model. The distributions of human IUR values (from which the recommended [i.e., mean] values were taken) show that the 99th percentiles are approximately seven- and sixfold higher than means for liver and lung cancer, respectively. For the distribution of OSFs, the 99th percentile is approximately twofold higher than the mean for liver cancer.

To further characterize the potential sensitivity of specific subpopulations, internal dose distributions for oral exposure to 1 mg/kg-day or inhalation exposure to 1 mg/m³ were estimated for 1-year-old children and 70-year-old men and women to compare with the broader population results used to estimate cancer risks above. Since the recommended cancer risk estimate is based on the GST-T1^{+/+} subpopulation, this analysis was also restricted to that subpopulation so that only the factors of age and gender would differ. The impact of considering other GST-T1 groups can be seen where risk estimates for the GST-T1^{+/-} and entire population mix are given above. Specification of age- and gender-specific parameters are as described in Appendix B. This sensitivity analysis is qualitatively similar to that described previously for the noncancer assessments of dichloromethane, where the variability in human equivalent dose and HEC values was estimated.

For this analysis, however, consideration of exclusively GST-T1^{+/+} individuals will clearly narrow any estimate of variability. This analysis will also differ from that for noncancer effects in that the inverse of the former relationship is being considered (i.e., the variation in a specific internal dose for a fixed exposure is being computed, whereas for the human equivalent dose and HEC, the variability in exposure levels corresponding to a fixed internal dose are estimated). The results of this analysis are shown in Figure 5-20 and Table 5-27 for oral exposures and in Figure 5-20 and Table 5-28 for inhalation exposures.




Figure 5-20. Histograms for a liver-specific dose of GST metabolism (mg GST metabolites per liter liver per day) for the general population (0.5- to 80-year-old males and females), and specific age/gender groups within the population of GST-T1^{+/+} genotypes, given a daily oral dose-rate of 1 mg/kg-day dichloromethane.

Table 5-27. Statistical characteristics of human internal doses for 1 mg/kg-day oral exposures in specific populations

Population	Internal dose (mg/L liver per d) ^a		
	Mean	95 th percentile	99 th percentile
All ages ^b	9.43×10^{-2}	2.98×10^{-1}	5.43×10^{-1}
1-yr-old children	7.82×10^{-2}	2.41×10^{-1}	4.00×10^{-1}
70-yr-old men	9.71×10^{-2}	2.99×10^{-1}	5.51×10^{-1}
70-yr-old women	1.01×10^{-1}	5.33×10^{-1}	9.84×10^{-1}

^aLiver-specific GST-T1 metabolism in GST-T1^{+/+} individuals exposed orally to 1 mg/kg-d dichloromethane.

^b0.5- to 80-yr-old males and females.




Figure 5-21. Histograms for liver-specific dose of GST metabolism (mg GST metabolites per liter liver per day) for the general population (0.5- to 80-year-old males and females), and specific age/gender groups within the population of GST-T1^{+/+} genotypes, given a continuous inhalation exposure to 1 mg/m³ dichloromethane.

Table 5-28. Statistical characteristics of human internal doses for 1 mg/m³ inhalation exposures in specific subpopulations

Population	Internal dose (mg/L liver per d) ^a		
	Mean	95 th percentile	99 th percentile
All ages ^b	6.61×10^{-6}	2.21×10^{-5}	4.47×10^{-5}
1-yr-old children	1.65×10^{-5}	5.11×10^{-5}	9.04×10^{-5}
70-yr-old men	5.09×10^{-6}	1.68×10^{-5}	3.12×10^{-5}
70-yr-old women	4.14×10^{-6}	1.37×10^{-5}	2.56×10^{-5}

^aLiver-specific GST-T1 metabolism in GST-T1^{+/+} individuals exposed continuously by inhalation to 1 mg/m³ dichloromethane.

^b0.5- to 80-yr-old males and females.

For the oral exposure analysis, the distribution of internal doses shows little variation among the different age/gender groups (Figure 5-21, Table 5-27). The cancer analysis is based on a very low internal dose where little enzymatic saturation is expected to occur, allowing for efficient first-pass metabolism, which is independent of differences in respiration; differences will be more significant at the higher doses analyzed for the noncancer human equivalent applied

dose. Thus, the consideration of only GST metabolism and the narrower range of metabolic rate for that pathway in the +/+ population at low oral exposure rates results in minimal age/gender sensitivity differences (the 7-year-old female is only 5% more sensitive from pharmacokinetic factors than the general population).

For inhalation, an internal liver GST dose (mean value) about 2.5 times higher in the child than the general population is predicted due to the higher inhalation rate. The results for the liver GST dose for inhalation (Figure 5-21 and Table 5-28) indicate that the 70-year-old male and female populations are only slightly shifted from the general population, while the population for the 1-year-old child is a distinct upper tail of the general distribution.

6. MAJOR CONCLUSIONS IN THE CHARACTERIZATION OF HAZARD AND DOSE RESPONSE

6.1. HUMAN HAZARD POTENTIAL

Dichloromethane (CASRN 75-09-2), also known as methylene chloride, is a colorless liquid with a penetrating, ether-like odor. It is produced by the direct reaction of methane with chlorine at either high temperatures or low temperatures under catalytic or photolytic conditions. The principal uses for dichloromethane have been in paint strippers and removers, as a propellant in aerosols, in the manufacture of drugs, pharmaceuticals, film coatings, electronics, and polyurethane foam, and as a metal-cleaning solvent.

Dichloromethane is rapidly absorbed through both oral administration and inhalation exposure with a steady-state saturation occurring with inhalation. Results from studies of animals show that following absorption, dichloromethane is rapidly distributed throughout the body and has been detected in all tissues that have been evaluated. Metabolism of dichloromethane involves two primary pathways. Dichloromethane is metabolized to CO in a CYP-dependent oxidative pathway (CYP2E1) that is predominant at low exposure levels. The other major pathway for dichloromethane metabolism involves the conjugation of dichloromethane to GSH, catalyzed by GST (GST-T1). This results in the formation of a GSH conjugate that is eventually metabolized to CO₂. The conjugation of dichloromethane to GSH results in the formation of two reactive intermediates that have been hypothesized to be involved in dichloromethane carcinogenicity, S-(chloromethyl)glutathione and formaldehyde. Formation of formaldehyde leads to several covalent modifications of cellular macromolecules, including DNA-protein cross-links (Casanova et al., 1996) and RNA-formaldehyde adducts (Casanova et al., 1997). Evidence is also available that S-(chloromethyl)glutathione can result in both DNA SSBs and DNA mutations, presumably through DNA alkylation (Green, 1997; Graves and Green, 1996; Graves et al., 1996, 1994a; Hashmi et al., 1994). However, DNA reaction products (e.g., DNA adducts) produced by S-(chloromethyl)glutathione have not been found in vivo, possibly due to potential instability of these compounds or due to the limited doses used in attempts to detect them (Watanabe et al., 2007; Hashmi et al., 1994). DNA adducts, however, have been observed in in vitro studies in which calf thymus DNA was incubated with dichloromethane and GST or was incubated with S-(1-acetoxymethyl)glutathione, a compound structurally similar to S-(chloromethyl)glutathione (Marsch et al., 2004; Kayser and Vuilleumier, 2001).

Information on noncancer effects in humans exposed orally to dichloromethane are restricted to case reports of neurological impairment (general CNS depression), liver and kidney effects (as severe as organ failure), and gastrointestinal irritation in individuals who ingested amounts ranging from about 25 to 300 mL (Chang et al., 1999; Hughes and Tracey, 1993). The

animal toxicity database identifies hepatic effects (hepatic vacuolation, liver foci) as the critical dose-dependent noncancer endpoint associated with oral exposure to dichloromethane. The most frequently observed liver effect was hepatocyte vacuolation, seen with drinking water exposure (90 days) in F344 rats at ≥ 166 mg/kg-day and B6C3F₁ mice at 586 mg/kg-day (Kirschman et al., 1986) and with gavage exposure (14 days) in CD-1 mice at 333 mg/kg-day (Condie et al., 1983). Hepatocyte degeneration or necrosis was observed in female F344 rats exposed in drinking water for 90 days to 1,469 mg/kg-day (Kirschman et al., 1986) and in female F344 rats exposed by gavage for 14 days to 337 mg/kg-day (Berman et al., 1995). In the chronic-duration (104-week) study, liver effects (areas of foci alteration) were observed in F344 rats exposed to drinking water doses between 50 and 250 mg/kg-day (Serota et al., 1986a). In the reproductive oral administration studies, no significant effect on reproductive function or parameters was observed in rats up to 225 mg/kg-day (General Electric Company, 1976) or in mice up to 500 mg/kg-day (Raje et al., 1988). The NOAEL and LOAEL for altered neurological functions in female F344 rats were 101 and 337 mg/kg-day (as reported by Moser et al., 1995).

Acute inhalation exposure of humans to dichloromethane has been associated with cardiovascular impairments due to decreased oxygen availability from COHb formation and neurological impairment from interaction of dichloromethane with nervous system membranes (Bos et al., 2006; ACGIH, 2001; ATSDR, 2000; Cherry et al., 1983; Putz et al., 1979; Gamberale et al., 1975; Winneke, 1974). Relatively little is known about the long-term neurological effects of chronic exposures, although there are studies that provide some evidence of an increased prevalence of neurological symptoms among workers with average exposures of 75–100 ppm (Cherry et al., 1981) and long-term effects on some neurological measures (i.e., possible detriments in attention and reaction time in complex tasks) in retired workers whose past exposures were in the 100–200 ppm range (Lash et al., 1991). These studies are limited by the relatively small sample sizes and low power for detecting statistically significant results for these endpoints.

Following repeated inhalation to dichloromethane, the liver is the most sensitive target for noncancer toxicity in rats and mice. Lifetime exposure was associated with hepatocyte vacuolation and necrosis in F344 rats exposed to 1,000 ppm 6 hours/day (Mennear et al., 1988; NTP, 1986), hepatocyte vacuolation in Sprague-Dawley rats exposed to 500 ppm 6 hours/day (Nitschke et al., 1988a; Burek et al., 1984), and hepatocyte degeneration in B6C3F₁ mice exposed to 2,000 ppm 6 hours/day (lower concentrations were not tested in mice) (Mennear et al., 1988; NTP, 1986). Other effects observed include renal tubular degenerations in F344 rats and B6C3F₁ mice at 2,000 ppm, testicular atrophy in B6C3F₁ mice at 4,000 ppm, and ovarian atrophy in B6C3F₁ mice at 2,000 ppm.

Other studies with inhalation exposure to dichloromethane revealed no significant effects on reproductive performance in rats (up to 1,500 ppm) (Nitschke et al., 1988b), although some evidence of a decrease in fertility index was seen in male mice exposed to 150 and 200 ppm

(Raje et al., 1988), and no adverse effects on fetal development of mice or rats exposed up to 1,250 ppm were seen by Schwetz et al. (1975). Decreases in fetal BW and changes in behavioral habituation were observed in Long-Evans rats exposed to 4,500 ppm during the gestational period (Bornschein et al., 1980; Hardin and Manson, 1980). Exposure-related noncancer effects on the lungs consisted of foreign-body pneumonia in rats exposed to 8,400 ppm 6 hours/day for 13 weeks (NTP, 1986), Clara cell vacuolation in mice exposed to 4,000 ppm 6 hours/day for 13 weeks (Foster et al., 1992), and pulmonary congestion in guinea pigs exposed to 5,000 ppm 7 hours/day for 6 months (Heppel et al., 1944). Several neurological mediated parameters including decreased activity (Kjellstrand et al., 1985; Weinstein et al., 1972; Heppel and Neal, 1944), impairment of learning and memory (Alexeef and Kilgore, 1983), and changes in responses to sensory stimuli (Rebert et al., 1989) are reported from acute and short-term dichloromethane exposure. Evidence of a localized immunosuppressive effect in the lung resulting from inhalation dichloromethane exposure was seen in an acute exposure (3 hours, 100 ppm) study in CD-1 mice (Aranyi et al., 1986).

Numerous in vitro studies have demonstrated mutagenic and genotoxic effects associated with dichloromethane exposure. For example, bacterial assays, yeast, and fungi provide evidence that the mutagenic action of dichloromethane in bacterial systems is enhanced by metabolic activation (e.g., Dillon et al., 1992; Jongen et al., 1982; Gocke et al., 1981). Positive results from assays of DNA damage with in vitro mammalian systems provide support that dichloromethane genotoxicity is linked to metabolism by GST enzymes (Graves et al., 1996, 1995, 1994b). Consistent evidence for several genotoxic endpoints in target tissues (liver and lung) in mice following in vivo exposure to dichloromethane provides supporting evidence that GST-pathway metabolites are key actors in the mutagenic and carcinogenic mode of action for dichloromethane. Pretreatment of mice with buthionine sulphoximine, a GSH depletor, caused a decrease to levels seen in controls in the amount of DNA damage detected immediately after in vivo exposure in liver and lung tissue, indicating GSH involvement in the genotoxic process (Graves et al., 1995). DNA damage (detected by the comet assay) was also reported in liver and lung tissues from male CD-1 mice sacrificed 24 hours after administration of a single oral dose of 1,720 mg/kg of dichloromethane (Sasaki et al., 1998). In this study, DNA damage in lung and liver was not detected 3 hours after dose administration, and no DNA damage occurred at either time point in several other tissues in which a carcinogenic response was not seen in chronic animal cancer bioassays (e.g., stomach, kidney, bone marrow). The weight of evidence from these studies suggests that dichloromethane is carcinogenic by a mutagenic mode of action.

Dichloromethane is “likely to be carcinogenic in humans” under the *Guidelines for Carcinogen Risk Assessment* (U.S. EPA, 2005a). Results from 2-year bioassays provide adequate evidence of the carcinogenicity of dichloromethane in mice and rats exposed by inhalation, as well as adequate data to describe dose-response relationships. Oral exposure to dichloromethane produced statistically significant increases in hepatocellular adenomas and

carcinomas in male B6C3F₁ mice (Serota et al., 1986b; Hazleton Laboratories, 1983). Inhalation exposure to concentrations of 2,000 or 4,000 ppm dichloromethane produced increased incidences of lung and liver tumors in B6C3F₁ mice (Maronpot et al., 1995; Foley et al., 1993; Kari et al., 1993; Mennear et al., 1988; NTP, 1986). Significantly increased incidences of benign mammary tumors (adenomas or fibroadenomas) were observed in male and female F344/N rats exposed by inhalation to 2,000 or 4,000 ppm (Mennear et al., 1988; NTP, 1986). A statistically significant increased incidence of brain or CNS tumors has not been observed in any of the animal cancer bioassays, but a 2-year study using relatively low exposure levels (0, 50, 200, and 500 ppm) in Sprague-Dawley rats observed a total of six astrocytoma or glioma (mixed glial cell) tumors in the exposed groups (Nitschke et al., 1988a). These tumors are exceedingly rare in rats, and there are few examples of statistically significant trends in animal bioassays (Sills et al., 1999).

6.2. DOSE RESPONSE

6.2.1. Oral RfD

The available oral toxicity data for animals identify hepatic effects (hepatic vacuolation, liver foci) as the most sensitive noncancer endpoint associated with chronic oral exposure to dichloromethane. The 104-week drinking-water study in F344 rats (Serota et al., 1986a) was selected as the principal study for RfD derivation because the study provided a sensitive endpoint (liver foci) and used lower doses in comparison to other chronic oral administration studies. In this study, four doses (6, 52, 125, and 235 mg/kg-day in males; 6, 58, 136, and 263 mg/kg-day in females) were used. A NOAEL of 6 mg/kg-day in males and females and a LOAEL of 52 (male) and 58 (female) mg/kg-day for alterations of liver foci was identified.

An RfD of 7×10^{-3} mg/kg-day is recommended for use in humans. The RfD derivation process involved first fitting all available dichotomous models in BMDS version 2.0 to the incidence data for male and female rats; the male data were used because a greater sensitivity was seen in males compared with females in this study. A dose metric of average daily mass of dichloromethane metabolized via the CYP pathway per unit volume of liver was derived from a EPA-modified rat PBPK model (see Appendix C). This metric was chosen because there are no data to support the role of a specific metabolite in the development of the noncancer liver lesions seen in oral and inhalation exposure studies and the CYP-metabolism dose metric was determined to be most consistent with the data. Then, the BMDL₁₀ for liver lesions was derived based on the best fitting model (in terms of the value of the AIC and examination of model fit and residuals). Because the metric is a rate of metabolism rather than the concentration of putative toxic metabolites and the clearance of these metabolites may be slower per volume tissue in the human compared with the rat, this rodent internal dose metric for noncancer effects was adjusted by dividing by a pharmacokinetic allometric $BW^{0.75}$ scaling factor (operationalized as $[BW_{\text{human}}/BW_{\text{rat}}]^{0.25} \approx 4.09$) to obtain a human equivalent internal BMDL₁₀. This BMDL₁₀

was then converted to the human equivalent dose by using a human PBPK model (adapted from David et al., 2006; see Appendix B) that utilizes Monte Carlo sampling techniques to provide a distribution of human equivalent doses. The first percentile of the distribution of human equivalent doses was chosen to include the most sensitive population while staying within bounds of what is considered computationally stable. The first percentile human equivalent dose was used as a POD and was divided by a composite UF of 30 ($3 [10^{0.5}]$ to account for uncertainty about interspecies toxicodynamic equivalence, $3 [10^{0.5}]$ to account for uncertainty about toxicodynamic variability in humans, and $3 [10^{0.5}]$ for database deficiencies) to arrive at an RfD of 7×10^{-3} mg/kg-day.

Use of the mean value (3.99×10^{-1} mg/kg-day) of the human equivalent dose distribution instead of the 1st percentile, with an additional UF of 3 ($10^{0.5}$) to account for human toxicokinetic variability, would yield a candidate RfD of 4×10^{-3} , which is relatively similar to the recommended RfD of 7×10^{-3} .

Confidence in the principal study, Serota et al. (1986a), is high. The 2-year drinking water study in rats is a well-conducted, peer-reviewed study that used four dose groups plus a control. Confidence in the oral database is medium-high. The oral database includes a 2-year drinking water study in rats (Serota et al., 1986a) and mice (Serota et al., 1986b) as well as a supporting subchronic exposure study (Kirschman et al., 1986) that reports similar liver effects to those observed in the chronic oral exposure studies. The toxicity of orally-administered dichloromethane has also been investigated in an oral administration immunotoxicity study (Warbrick et al., 2003), a one-generation oral reproductive toxicity study (General Electric Company, 1976), and an orally dosed developmental toxicity study (Narotsky and Kavlock, 1995). Several studies have also evaluated neurotoxicity associated with oral exposure to dichloromethane. The oral database lacks a two-generation reproductive study and a developmental neurotoxicity study; neurodevelopmental outcomes are relevant endpoints given the increase in blood CO (a known developmental neurotoxicant) that occurs through the CYP2E1 metabolic pathway for dichloromethane after oral and inhalation exposures. Overall confidence in the RfD is high.

6.2.2. Inhalation RfC

The liver is the most sensitive target for noncancer toxicity in rats and mice following repeated inhalation exposure to dichloromethane. Liver lesions (specifically, hepatic vacuolation) in rats are the critical noncancer effect from chronic dichloromethane inhalation in animals. Inhalation bioassays with Sprague-Dawley rats identified the lowest inhalation LOAEL for liver lesions in the database: 500 ppm (6 hours/day, 5 days/week for 2 years) (Nitschke et al., 1988a; Burek et al., 1984). Nitschke et al. (1988a) identified a NOAEL of 200 ppm for hepatocyte vacuolation in female rats. Because the Nitschke et al. (1988a) study more

adequately covers the range spanning the BMR compared with the study by Burek et al. (1984), the former study was selected as the principal study for derivation of a chronic inhalation RfC.

An RfC of 0.2 mg/m³ is derived based on the observed critical effect in the principal study. As was described above for the RfD, the RfC derivation process was based on a dose metric of average daily mass of dichloromethane metabolized via the CYP pathway per unit volume of liver. This metric was derived from a EPA-modified rat PBPK model (see Appendix C). Then, the BMDL₁₀ risk for liver lesions was derived based on the best fitting model in terms of the value of the AIC and examination of model fit and residuals. Because the metric is a rate of metabolism rather than the concentration of putative toxic metabolites and the clearance of these metabolites may be slower per volume tissue in the human compared with the rat, this rodent internal dose metric for noncancer effects was adjusted by dividing by a pharmacokinetic allometric BW^{0.75} scaling factor (operationalized as [BW_{human}/BW_{rat}]^{0.25} ≈ 4.09) to obtain a human-equivalent internal BMDL₁₀. This BMDL₁₀ was then converted to the HEC by using a human PBPK model (adapted from David et al., 2006; see Appendix B) that utilizes Monte Carlo sampling techniques to provide a distribution of HECs.

The first percentile HEC was used as a POD. This percentile was chosen because it included the most sensitive population while staying within bounds of what is considered computationally stable. This POD was divided by a composite UF of 100 (3 [10^{0.5}] to account for uncertainty about interspecies toxicodynamic equivalence, 3 [10^{0.5}] to account for uncertainty about toxicodynamic variability in humans, and 10 for database deficiencies) to arrive at an RfC of 0.2 mg/m³.

Use of the mean value (47.24 mg/m³) of the HEC distribution instead of the 1st percentile with an additional UF of 3 (10^{0.5}) to account for human toxicokinetic variability would yield a candidate RfC identical to the recommended value of 0.2 mg/m³. In addition, two comparison values derived from occupational studies produced values of 3.5 mg/m³ (Cherry et al., 1983) and 0.55 mg/m³ (Lash et al., 1991). The animal-derived candidate RfC is preferable to the human-derived candidate RfC because of the uncertainties about the exposure durations for the workers in the Cherry et al. (1983) study and uncertainties regarding the exposures and effect sizes in Lash et al. (1991), and because the RfC based on the rat data is more health protective.

Confidence in the principal study, Nitschke et al. (1988a), is high. The 2-year inhalation study in mice is a well-conducted, peer-reviewed study that used three concentration groups plus a control. Confidence in the inhalation database is medium-low. The inhalation database includes several well-conducted chronic inhalation studies that consistently identified the liver as the most sensitive noncancer target organ in rats (Nitschke et al., 1988a; NTP, 1986; Burek et al., 1984). A two-generation reproductive toxicity study (Nitschke et al., 1988b), developmental studies at relatively high exposures (≥1,250 ppm), several neurotoxicity studies, and an immunotoxicity study have been conducted in animals following inhalational exposures to dichloromethane. The results from the single dose developmental toxicity study in rats

(Bornschein et al., 1980; Hardin and Manson, 1980), in addition to the known increase in CO, the placental transfer of dichloromethane, and the relatively high activity of CYP2E1 in the brain compared to the liver of the developing human fetus (Hines, 2007; Johnsrud et al., 2003; Brzezinski et al., 1999), raise uncertainty regarding possible neurodevelopmental toxicity from gestational exposure to inhaled dichloromethane. An acute, 3-hour exposure to 100 ppm dichloromethane demonstrated evidence of immunosuppression in CD-1 mice (Aranyi et al., 1986). This study used a functional immune assay that is relevant to humans (i.e., increased risk of Streptococcal pneumonia-related mortality and decreased clearance of Klebsiella bacteria). Chronic and/or repeated exposure studies evaluating functional immunity are not available and represent a data gap. The inhalation database lacks adequate developmental neurotoxicity and immunotoxicity studies at chronic low exposures. Overall confidence in the RfC is medium.

6.2.3. Uncertainties in RfD and RfC Values

One data uncertainty identified is the potential for neurodevelopmental effects. Animal bioassays have not identified gross or microscopic effects on neural tissues from long-term exposures or single (Schwetz et al., 1975) or multigenerational (Nitschke et al., 1988b) developmental toxicity studies. However, behavioral changes were observed in pups born to rats exposed to high levels (4,500 ppm) of dichloromethane (Bornschein et al., 1980; Hardin and Manson, 1980); 4,500 ppm was the only dose used in this study. Thus, uncertainty exists as to the development of neurological effects from lower gestational exposures in animals or in humans. Immunotoxicity data revealed an additional area of data uncertainty specifically with respect to inhalation exposure. Data from Aranyi et al. (1986) demonstrated evidence of immunosuppression following a single 100 ppm dichloromethane exposure for 3 hours in CD-1 mice. The weight of evidence for nonneoplastic effects in humans and animals suggests that the development of liver lesions is the most sensitive effect, with a UF applied because of the lack of reproductive and neurodevelopmental studies for the RfD and, for the RfC, the uncertainty regarding the lack of neurodevelopmental, developmental, and immune system toxicity studies at low exposures.

The extrapolation of internal dichloromethane dosimetry from rat liver responses to human risk was accomplished by using PBPK models for dichloromethane in rats and humans. Uncertainties in rat and human dosimetry used for RfD and RfC derivation can arise from uncertainties in the PBPK models to accurately simulate the toxicokinetics of dichloromethane for animals under bioassay conditions and humans experiencing relatively low, chronic environmental exposures. Specific uncertainties identified with the PBPK models used here are described in detail in Sections 3.5.2 and 3.5.5. In brief, there is both a structural uncertainty in that the equation used to describe the CYP2E1-mediated metabolism that could be the source of discrepancies between the model and some of the data (both in vitro and in vivo) and a parametric uncertainty in the GST-T1 parameter, k_{fC} , evident from an inconsistency in the values

obtained by David et al. (2006). For noncancer reference values, the impact of these uncertainties appears to be moderate, no more than 30%. Further, the dose metric used in the models is the rate of metabolism to a putative toxic metabolite rather than the concentration (average or area under the concentration curve of the metabolite), so the model specifically fails to account for rodent-human differences in clearance or removal of the toxic metabolite. A scaling factor based on BW ratios was used to account for this difference.

Uncertainties in the human population model parameters and variability in that population were quantitatively accounted for by utilizing hierarchical Bayesian calibration methods during model development (David et al., 2006), as modified here by EPA. The rat model was modified, recalibrated, and utilized in a deterministic manner (Appendix C). Data were not available to perform a hierarchical Bayesian calibration in the rat, but uncertainties in the rat model predictions were assessed qualitatively. For both oral and inhalation exposures, the liver volume, followed closely by the volume of slowly perfused tissues, had the greatest impact on the internal dose of mg dichloromethane metabolized via CYP pathway per liter tissue per day. This was due to the fact that the dose metric is a tissue-specific measure, the majority of CYP metabolism is attributed to the liver, and changes in liver volume have a greater impact on the total CYP metabolism than either of the individual V_{\max} values. There is high confidence in the values used for volume of liver and slowly perfused tissues in the rat, as these are well studied (Brown et al., 1997). Therefore, except as described in the preceding paragraph, the uncertainties associated with use of the rat PBPK model should not markedly affect the values of the RfD and RfC.

An additional uncertainty inherent in this process, however, is the lack of knowledge concerning the most relevant dose metric (e.g., a specific metabolite) within the context of the development of the noncancer liver effects. This basic research question represents a data gap, and this uncertainty is not addressed quantitatively or qualitatively in the assessment.

The effect of dichloromethane on human populations that are sensitive due to pharmacokinetic differences was addressed quantitatively by using a human probabilistic PBPK model to generate distributions of human exposures likely to occur given a specified internal BMDL₁₀. The model and resulting distributions take into account the known differences in human physiology and metabolic capability with regard to dichloromethane dosimetry. The first percentile values of the distributions of human equivalent doses (Table 5-3) and HECs (Table 5-7) served as points of departure for candidate RfDs and RfCs, respectively, to protect toxicokinetically sensitive individuals. No data are available regarding toxicodynamic differences within a human population. Therefore, a UF of 3 for possible differences in human toxicodynamic responses is intended to be protective for sensitive individuals.

6.2.4. Oral Cancer Slope Factor

The recommended cancer OSF for dichloromethane is 2×10^{-3} (mg/kg-day)⁻¹, which is based on liver tumor responses in male B6C3F₁ mice exposed to dichloromethane in drinking water for 2 years (Serota et al., 1986b; Hazleton Laboratories, 1983). This value was derived by using a tissue-specific GST metabolism dose metric with allometric scaling to account for uncertainty regarding the reactivity and clearance of the metabolite(s) involved in the carcinogenic response.

There was only one adequate oral exposure cancer bioassay evaluating the carcinogenic potential of orally administered dichloromethane in F344 rats and B6C3F₁ mice (Serota et al., 1986a, b; Hazleton Laboratories, 1983). Significant increases in incidence of liver adenomas and carcinomas were observed in male but not female B6C3F₁ mice (female data were not presented in the summary reports) (Serota et al., 1986b; Hazleton Laboratories, 1983). The study authors concluded that in the male bioassay, there was no dose-related trend and that there were no significant differences comparing the individual dose groups with the combined control group, and that the observed incidences were “within the normal fluctuation of this type of tumor incidence.” (The trend *p*-value and pairwise test *p*-values were not given in the Serota et al. [1986b] paper but can be found in the full report [Hazleton Laboratories, 1983]). Although Serota et al. (1986b) state that a two-tailed significance level of *p* = 0.05 was used for all tests, Hazleton Laboratories (1983) indicated that a correction factor for multiple comparisons was used specifically for the liver cancer data, reducing the nominal *p*-value from 0.05 to 0.0125; none of these individual group comparisons are statistically significant when a *p*-value of 0.0125 is used.

Based on the Hazleton Laboratories (1983) statistical analysis, EPA concluded that dichloromethane induced a carcinogenic response in male B6C3F₁ mice as evidenced by a marginally increased trend test (*p* = 0.058) for combined hepatocellular adenomas and carcinomas, and by small but statistically significant (*p* < 0.05) increases in hepatocellular adenomas and carcinomas at dose levels of 125 (*p* = 0.023), 185 (*p* = 0.019), and 250 mg/kg-day (*p* = 0.036). EPA did not consider the use of a multiple comparisons correction factor for the evaluation of the liver tumor data (a primary a priori hypothesis) to be warranted.

In addition, the incidence in the control groups (19%) was almost identical to the mean seen in the historical controls from this laboratory (17.8% based on 354 male B6C3F₁ mice), so there is no indication that the observed trend is being driven by an artificially low rate in controls, no indication that the experimental conditions resulted in a systematic increase in the incidence of hepatocellular adenomas and carcinomas, and it is unlikely that the pattern of incidence rates observed in this study (increased incidence in all four dose groups, with three of these increases significant at a *p*-value < 0.05) reflect normal fluctuations in the incidence of these tumors. In F344 rats (Serota et al., 1986a), no increased incidence of liver tumors was seen in male rats, and the pattern in female rats was characterized by a jagged stepped pattern of

increasing incidence of hepatocellular carcinoma or neoplastic nodules. However, the potential malignant characterization of the nodules was not described, and the data for hepatocellular carcinomas are much more limited. The derivation of the oral cancer slope factor is based on the male mice data because of their greater sensitivity to liver cancer compared with female mice and to male and female rats.

A modified mouse PBPK model of Marino et al. (2006) was used to approximate the internal dose of daily dichloromethane (mg) metabolized via the GST pathway per unit volume of liver from the daily oral administered doses. This approach was taken based on evidence that GST-pathway metabolites produced from dichloromethane are primarily responsible for dichloromethane carcinogenicity in mouse liver. The multistage dose-response model (BMDS version 2.0) was used to fit the mouse liver tumor incidence and PBPK model-derived internal dose data and to derive a mouse internal BMD and BMDL₁₀. Because the metric is a rate of metabolism rather than the concentration of putative toxic metabolites and data pertaining to the reactivity or clearance rate of the relevant metabolite(s) are lacking, the human BMDL₁₀ was derived by multiplying the mouse BMDL₁₀ by a BW^{0.75} allometric scaling factor (operationalized as $[BW_{\text{human}}/BW_{\text{mouse}}]^{0.25} \approx 7$) to account for the potential slower clearance per volume tissue in the human compared with the mouse. Linear extrapolation from the internal human BMDL₁₀ ($0.1/\text{BMDL}_{10}$) was used to derive oral risk factors for liver tumors based on tumor responses in male mice. The linear low-dose extrapolation approach for agents with a mutagenic mode of action was selected because GST-metabolism of dichloromethane is expected to occur at and below exposures producing the mouse BMDL₁₀, even though CYP2E1 metabolism is expected to be unsaturated and to represent the predominant metabolic pathway in the liver. Currently, there are no data from chronic oral cancer bioassays in mice providing support for a nonlinear dose-response relationship.

Probability distributions of human oral cancer slope factors were derived by using a human PBPK model (adapted from David et al. [2006]; see Appendix B). The cancer reference values (OSF and IUR) were derived for a sensitive population: a population composed entirely of carriers of the GST-T1^{+/+} homozygous genotype (that is, the group that would be expected to be most sensitive to the carcinogenic effects of dichloromethane). In addition, cancer values derived for a population reflecting the estimated frequency of GST-T1 genotypes in the current U.S. population (20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+}) were presented. All simulations also included a distribution of CYP activity based on data from Lipscomb et al. (2003). The mean OSF based on liver tumors in mice exposed to dichloromethane in drinking water, $2 \times 10^{-3} \text{ (mg/kg-day)}^{-1}$, based on what is assumed to be the most sensitive of the populations (the GST-T1^{+/+} group) is the recommended OSF to be used in deterministic risk assessments for chronic oral exposures to dichloromethane.

An OSF derived from the liver tumor data in the Serota et al. (1986b) study using administered dose dosimetry rather than PBPK modeling is approximately one order of

magnitude higher than the current recommended value of 2×10^{-3} (per mg/kg-day). There is approximately one to two orders of magnitude difference among the values based on different dose metrics, scaling factors, and populations (Table 6-1).

Table 6-1. Comparison of OSFs derived by using various assumptions and metrics, based on liver tumors in male mice

Population ^a	Dose metric	Species, sex	Tumor	Scaling factor	Mean OSF (mg/kg-d) ⁻¹	Source (table)
GST-T1^{+/+}	Tissue-specific GST-metabolism rate^b	Mouse, male	Liver	7.0	1.7 × 10⁻³	Table 5-13
	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	2.4 × 10 ⁻⁴	Table 5-13
	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	9.3 × 10 ⁻⁴	Table 5-13
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	7.0	1.2 × 10 ⁻⁴	Table 5-14
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	1.0	1.7 × 10 ⁻⁵	Table 5-14
	Route-to-route extrapolation, whole-body metabolism	Mouse, male	Liver	7.0	6.7 × 10 ⁻⁵	Table 5-14
Mixed	Tissue-specific GST-metabolism rate ^b	Mouse, male	Liver	7.0	9.4 × 10 ⁻⁴	Table 5-13
	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	1.3 × 10 ⁻⁴	Table 5-13
	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	5.4 × 10 ⁻⁴	Table 5-13
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	7.0	6.8 × 10 ⁻⁵	Table 5-14
	Route-to-route extrapolation, tissue-specific metabolism	Mouse, male	Liver	1.0	9.7 × 10 ⁻⁶	Table 5-14
	Route-to-route extrapolation, whole-body metabolism	Mouse, male	Liver	7.0	3.9 × 10 ⁻⁵	Table 5-14
	Applied dose (human equivalent dose)	Mouse, male	Liver		1.0 × 10 ⁻²	Table 5-15
	1995 IRIS assessment	Mouse, male	Liver		7.5 × 10 ⁻³	

^aGST-T1^{+/+} = homozygous, full enzyme activity; Mixed = genotypes based on a population reflecting the estimated frequency of genotypes in the current U.S. population: 20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^bBolded value is the basis for the recommended OSF of 2 × 10⁻³ per mg/kg-d.

The recommended OSF of 2×10^{-3} (per mg/kg-day) is based on a tissue-specific GST internal dose metric with allometric scaling. Although the involvement of the GST pathway in carcinogenic response has been established, some uncertainty remains as to the metabolite(s) involved and the rate at which those metabolites are cleared. The value derived specifically for the GST-T1^{+/+} population is recommended to provide protection for the population that is hypothesized to be most sensitive to the carcinogenic effect. Application of ADAFs to the cancer OSF is recommended in combination with appropriate exposure data when assessing risks associated with early-life exposure (see Section 5.4.4 for more details).

6.2.5. Cancer IUR

The recommended cancer IUR is 1×10^{-8} ($\mu\text{g}/\text{m}^3$)⁻¹ for the development of liver and lung cancers based on data from male B6C3F₁ mice using a tissue-specific GST metabolism dose metric. Data for liver and lung tumors in male and female B6C3F₁ mice following exposure to airborne dichloromethane were used to develop IURs for dichloromethane (Mennear et al., 1988; NTP, 1986). This study was selected as the principal study to derive an IUR for dichloromethane because of the completeness of the data, adequate sample size, and clear dose response. In the NTP (1986) study, significant increases in incidence of liver and lung adenomas and carcinomas were observed in both sexes of B6C3F₁ mice exposed 6 hours/day, 5 days/week for 2 years.

The PBPK model of Marino et al. (2006) for dichloromethane in the mouse was used to calculate long-term daily average internal liver doses. The selected internal dose metrics for liver tumors and lung tumors were long-term average daily mass of dichloromethane metabolized via the GST pathway per unit volume of liver and lung, respectively. This approach was taken based on evidence that GST-pathway metabolites produced from dichloromethane are primarily responsible for dichloromethane carcinogenicity in mouse liver. The multistage dose-response model (BMDS version 2.0) was used to fit the mouse liver tumor incidence and PBPK model-derived internal dose data and to derive a mouse internal BMD and BMDL₁₀. Because the metric is a rate of metabolism rather than the concentration of putative toxic metabolites and data pertaining to the reactivity or clearance rate of the relevant metabolite(s) are lacking, the human BMDL₁₀ was derived by multiplying the mouse BMDL₁₀ by a $\text{BW}^{0.75}$ allometric scaling factor (operationalized as $[\text{BW}_{\text{human}}/\text{BW}_{\text{mouse}}]^{0.25} \approx 7$) to account for the potential slower clearance per volume tissue in the human compared with the mouse. A linear extrapolation approach using the internal human BMDL₁₀ for liver and lung tumors was used to calculate human tumor risk factors by dividing the BMR of 0.1 by the human BMDL for each tumor type. Currently, there are no data from chronic inhalation cancer bioassays in mice or rats providing support for a nonlinear dose-response relationship.

The human PBPK model (adapted from David et al. [2006]; see Appendix B) provided distributions of human internal dose metrics of daily mass of dichloromethane metabolized via

the GST pathway per unit volume of liver and lung resulting from chronic inhalation exposure to a unit concentration of $1 \mu\text{g}/\text{m}^3$ dichloromethane (0.00029 ppm). As with the OSF, the cancer IUR was derived for a sensitive population: a population composed entirely of carriers of the GST-T1 homozygous positive genotype (that is, the group that would be expected to be most sensitive to the carcinogenic effects of dichloromethane). In addition, cancer values derived for a population reflecting the estimated frequency of GST-T1 genotypes in the current U.S. population (20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+}) were also presented. The distributions of IURs for liver or lung tumors were generated by multiplying the human tumor risk factor for each tumor type and sex by the distribution of internal doses from chronic exposure to $1 \mu\text{g}/\text{m}^3$ dichloromethane. A procedure to combine risks for liver and lung tumors using different dose metrics for the different tumors (i.e., liver-specific and lung-specific metabolism for liver and lung tumors, respectively) was used to derive the recommended IUR of $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$ based on what is assumed to be the most sensitive of the populations, the GST-T1^{+/+} group.

The current recommended IUR value of $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$ is approximately 47-fold lower than the previous IRIS value of $4.7 \times 10^{-7} (\mu\text{g}/\text{m}^3)^{-1}$. An IUR derived from the liver tumor data of the NTP (1986) study using applied concentration dosimetry rather than PBPK modeling, $3.6 \times 10^{-7} (\mu\text{g}/\text{m}^3)^{-1}$, is approximately one order of magnitude higher than the currently recommended value of $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$ (Table 6-2). There is approximately one to two orders of magnitude difference among the values based on different dose metrics, scaling factors, and populations.

The recommended IUR value of $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$ is based on a tissue-specific GST-internal dose metric with allometric scaling. Although the involvement of the GST pathway in carcinogenic response has been established, some uncertainty remains as to the metabolite(s) involved and the rate at which those metabolites are cleared. The value derived specifically for the GST-T1^{+/+} population is recommended to provide protection for the population that is hypothesized to be most sensitive to the carcinogenic effect. Application of ADAFs to the cancer IUR is recommended when assessing risks associated with early-life exposure (see Section 5.4.4 for more details).

Table 6-2. Comparison of IURs derived by using various assumptions and metrics

Population ^a	Dose metric	Species, sex	Tumor type	Scaling factor	IUR ^b ($\mu\text{g}/\text{m}^3$) ⁻¹	Source (table)
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate^c	Mouse, male	Liver and lung	7.0	1.3×10^{-8}	Table 5-20
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Liver	7.0	8.5×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Lung	7.0	5.6×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Liver and lung	1.0	1.9×10^{-9}	Table 5-20
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	1.2×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Tissue-specific GST-metabolism rate	Mouse, male	Lung	1.0	8.0×10^{-10}	Table 5-19
GST-T1 ^{+/+}	Whole-body GST metabolism rate	Mouse, male	Liver and lung	7.0	1.6×10^{-8}	Table 5-20
GST-T1 ^{+/+}	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	5.5×10^{-9}	Table 5-19
GST-T1 ^{+/+}	Whole-body GST metabolism rate	Mouse, male	Lung	7.0	1.2×10^{-8}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver and lung	7.0	7.4×10^{-9}	Table 5-20
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver	7.0	4.8×10^{-9}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Lung	7.0	3.2×10^{-9}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver and lung	1.0	1.1×10^{-9}	Table 5-20
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Liver	1.0	6.8×10^{-10}	Table 5-19
Mixed	Tissue-specific GST-metabolism rate	Mouse, male	Lung	1.0	4.5×10^{-10}	Table 5-19
Mixed	Whole-body GST metabolism rate	Mouse, male	Liver and lung	7.0	9.2×10^{-9}	Table 5-20
Mixed	Whole-body GST metabolism rate	Mouse, male	Liver	7.0	3.1×10^{-9}	Table 5-19
Mixed	Whole-body GST metabolism rate	Mouse, male	Lung	7.0	6.9×10^{-9}	Table 5-19
	Administered concentration (HEC)	Mouse, male	Liver		3.6×10^{-7}	Table 5-21
	Administered concentration (HEC)	Mouse, male	Lung		8.1×10^{-7}	Table 5-21
	1995 IRIS assessment ^c	Mouse, male	Liver and lung	12.7	4.7×10^{-7}	

^aGST-T1^{+/+} = homozygous, full enzyme activity; Mixed = genotypes based on a population reflecting the estimated frequency of genotypes in the current U.S. population: 20% GST-T1^{-/-}, 48% GST-T1^{+/-}, and 32% GST-T1^{+/+} (Haber et al., 2002).

^bBased on mean value of the derived distributions.

^cBolded value is the basis for the recommended IUR of 1×10^{-8} $\mu\text{g}/\text{m}^3$ per mg/kg-d.

6.2.6. Uncertainties in Cancer Risk Values

The database of animal bioassays identifies the liver and lung as the most sensitive target organs for dichloromethane-induced tumor development, and there is high confidence that the dose-response data for liver and lung cancer in mice represent the best available data for derivation of human cancer risks. A dose-response relationship was seen with respect to liver cancer in mice exposed orally and with respect to liver and lung cancer in mice exposed by inhalation. Statistically significant increases in benign mammary gland tumors were observed in one study of F344 rats exposed by inhalation to 2,000 or 4,000 ppm (Mennear et al., 1988; NTP, 1986); evidence for a tumorigenic mammary gland response in Sprague-Dawley rats was limited to increased numbers of benign mammary tumors per animal at levels of 50–500 ppm (Nitschke et al., 1988a) or 500–3,500 ppm (Burek et al., 1984). A gavage study in female Sprague-Dawley rats reported an increased incidence of malignant mammary tumors, mainly adenocarcinomas (8, 6, and 18% in the control, 100, and 500 mg/kg dose groups, respectively), but the increase was not statistically significant. Data were not provided to allow an analysis that accounts for differing mortality rates (Maltoni et al., 1988). The toxicokinetic or mechanistic events that might lead to mammary gland tumor development in rats are unknown, although CYP2E1 (El-Rayes et al., 2003; Hellmold et al., 1998) and GST-T1 expression have been detected in human mammary tissue (Lehmann and Wagner, 2008). Rare CNS tumors were observed in one study in rats spanning a relatively low range of exposures (0–500 ppm) (Nitschke et al., 1988a). These cancers were not seen in two other studies in rats, both involving higher doses (1,000–4,000 ppm) (NTP, 1986; Burek et al., 1984), or in a similar high-dose study in mice (NTP, 1986). The relative rarity of the tumors precludes the use of the low-dose exposure study (Nitschke et al., 1988a) in a quantitative dose-response assessment. The available epidemiologic studies provide some evidence of an association between dichloromethane and brain cancer. The available epidemiologic studies do not provide an adequate basis for the evaluation of the role of dichloromethane in breast cancer because there are currently no cohort studies with adequate statistical power and no case-control studies with adequate exposure methodology to examine this relationship.

There is uncertainty as to whether the reactivity of the toxic dichloromethane metabolites is sufficiently high enough to preclude systemic distribution. Therefore, alternative derivations of cancer risk values were performed under the assumption that high reactivity leads to complete clearance from the tissue in which the active metabolite is formed (scaling factor = 1.0). The difference in scaling factor (7.0 for allometric scaling versus 1.0) results in a sevenfold decrease in estimated cancer toxicity values. Using a whole-body GST metabolism dose metric, the resulting OSF and IUR for liver cancer was approximately fivefold lower than when tissue-specific dose metrics were used; however, the IURs for lung cancer and for the combined liver and lung cancer risk were higher with the whole-body compared with the tissue-specific metric

(Table 6-1 and Table 6-2). This difference reflects the lower metabolism that occurs in human versus mouse lung (relative to total): lung-specific metabolism is lower in humans than mice, so the predicted risk in the lung is lower when based on that metabolism versus whole-body metabolism. The mechanistic data support the hypothesis that reactive metabolites produced in the target tissues do not distribute significantly beyond those tissues and cause deleterious effects in the metabolizing tissues soon after generation. Thus, there is less uncertainty in the cancer risk values derived by using a tissue-specific GST metabolism dose metric compared with those derived using a whole-body GST metabolism dose metric.

Uncertainty in the ability of the PBPK models to estimate animal and human internal doses from lifetime bioassay low-level environmental exposures may affect the confidence in the cancer risk extrapolated from animal data. Uncertainties in the mouse and human model parameter values were integrated quantitatively into parameter estimation by utilizing hierarchical Bayesian methods to calibrate the models at the population level (David et al., 2006; Marino et al., 2006). However, with the subsequent deterministic application of the mouse model (using only the mean value for each parameter distribution), the information contained in the mouse parameter uncertainties reported by Marino et al. (2006) is not integrated into the final risk estimates described here.

The use of Monte Carlo sampling to define human model parameter distributions allowed for derivation of human distributions of dosimetry and cancer risk, providing for bounds on the recommended risk values. A sensitivity analysis was performed to identify model parameters most influential on the predictions of dose metrics used to estimate oral and inhalation cancer risks. For inhalation exposures in mice, the PB, followed closely by the first-order GST-mediated metabolism rate, had the greatest impact on the dose metric for liver cancer (mg dichloromethane metabolized via GST pathway per liter liver per day). For drinking water exposures in mice, the first-order GST-mediated metabolism rate, followed by the CYP-mediated maximum reaction velocity ($V_{\max C}$) affected the liver cancer dose metric to the greatest extent. For mice inhaling dichloromethane, the lung cancer dose metric (mg dichloromethane metabolized via GST pathways per liter lung per day), like the liver cancer metric, was highly affected by the first-order GST-mediated metabolism rate and the PB. However, the lung cancer dose metric was most sensitive to the proportional yield of liver GST-mediated metabolic activity attributed to the lung. The PB was experimentally determined, lending high confidence to its value. In contrast, values for the three metabolic parameters were determined by computational optimization against data sets not directly measuring dichloromethane or its metabolites in the target/metabolizing tissues. It is uncertain how alternative values for these three parameters would affect the estimation of animal BMDL₁₀ values and, ultimately, IURs and OSFs. In addition, specific uncertainty remains concerning the human PBPK parameter distributions (see discussion on k_{fC} in Section 3.5.5).

Finally, while the structure and equations used in the existing model have been described extensively in peer-reviewed publications, uncertainty remains concerning the model structure. Specifically an alternative (dual-binding-site) CYP metabolic rate equation (Korzekwa et al., 1998) for dichloromethane may better describe CYP2E1-mediated GST metabolism. However, this hypothesis requires further testing in the laboratory and integration of the alternate rate equation into the PBPK modeling.

7. REFERENCES

- ACGIH (American Conference of Governmental Industrial Hygienists). (2001) Dichloromethane. In: Documentation of the threshold limit values and biological exposure indices. 7th edition. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- Alexeef, GV; Kilgore, WW. (1983) Learning impairment in mice following acute exposure to dichloromethane and carbon tetrachloride. *J Toxicol Environ Health* 11:569–581.
- Allen, J; Kligerman, A; Campbell, J; et al. (1990) Cytogenetic analyses of mice exposed to dichloromethane. *Environ Mol Mutagen* 15:221–228.
- Anders, MW; Sunram, JM. (1982) Transplacental passage of dichloromethane and carbon monoxide. *Toxicol Lett* 12:231–234.
- Andersen, ME; Clewell, HJ, III; Gargas, ML; et al. (1987) Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol Appl Pharmacol* 87:185–205.
- Andersen, ME; Clewell, HJ, III; Gargas, ML; et al. (1991) Physiologically based pharmacokinetic modeling with dichloromethane, its metabolite, carbon monoxide, and blood carboxyhemoglobin in rats and humans. *Toxicol Appl Pharmacol* 108:14–27.
- Andrae, U; Wolff, T. (1983) Dichloromethane is not genotoxic in isolated rat hepatocytes. *Arch Toxicol* 52:287–290.
- Angelo, MJ; Pritchard, AB; Hawkins, DR; et al. (1986a) The pharmacokinetics of dichloromethane. I. Disposition in B6C3F₁ mice following intravenous and oral administration. *Food Chem Toxicol* 24(9):965–974.
- Angelo, MJ; Pritchard, AB; Hawkins, DR; et al. (1986b) The pharmacokinetics of dichloromethane. II. Disposition in Fischer 344 rats following intravenous and oral administration. *Food Chem Toxicol* 24(9):975–980.
- Anthony, T. (1979) Methylene chloride. In: Kirk, RE; Othmer, DF; eds. *Kirk-Othmer encyclopedia of chemical technology*. 3rd edition. New York, NY: John Wiley & Sons, pp. 686–693.
- Anundi, H; Lind, ML; Friis, L; et al. (1993) High exposures to organic solvents among graffiti removers. *Int Arch Occup Environ Health* 65:247–251.
- Aranyi, C; O'Shea, WJ; Graham, JA; et al. (1986) The effects of inhalation of organic chemical air contaminants on murine lung host defenses. *Fundam Appl Toxicol* 6:713–720.
- Arcus-Arth, A; Blaisdell, RJ. (2007) Statistical distributions of daily breathing rates for narrow age groups of infants and children. *Risk Anal* 27:97–110.
- Astrand, I; Ovrum, P; Carlsson, A. (1975) Exposure to methylene chloride. I. Its concentration in alveolar air and blood during rest and exercise and its metabolism. *Scand J Work Environ Health* 1(2):78–94.
- Atkinson, R. (1989) Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. *J Phys Chem Ref Data* 1:63.
- ATSDR (Agency for Toxic Substances and Disease Registry). (2000) Toxicological profile for methylene chloride. Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA. Available online at <http://www.atsdr.cdc.gov/toxpro2.html> (accessed March 9, 2010).
- Bakinson, MA; Jones, RD. (1985) Gassings due to methylene chloride, xylene, toluene, and styrene reported to Her Majesty's factory inspectorate 1961–80. *Br J Ind Med* 42:184–190.

- Bell, BP; Franks, P; Hildreth, N; et al. (1991) Methylene chloride exposure and birth weight in Monroe County, New York. *Environ Res* 55:31–39.
- Berman, E; Schlicht, M; Moser, VC; et al. (1995) A multidisciplinary approach to toxicological screening: I. Systemic toxicity. *J Toxicol Environ Health* 45:27–143.
- Bernauer, U; Vieth, B; Ellrich, R; et al. (2000) CYP2E1 expression in bone marrow and its intra- and interspecies variability: approaches for a more reliable extrapolation from one species to another in the risk assessment of chemicals. *Arch Toxicol* 73:618–624.
- Blair, A; Hartge, P; Stewart, P; et al. (1998) Mortality and cancer incidence of aircraft maintenance workers exposed to trichloroethylene and other organic solvents and chemicals: extended follow up. *Occup Environ Med* 55:161–171.
- Blocki, FA; Logan, MSP; Baoli, C; et al. (1994) Reaction of rat liver glutathione s-transferase and bacterial dichloromethane dehalogenase with dihalomethanes. *J Biol Chem* 269(12):8826–8830.
- Bogaards, JJP; van Ommen, B; van Bladeren, PJ. (1993) Interindividual differences in the in vitro conjugation of methylene chloride with glutathione by cytosolic glutathione S-transferase in 22 human liver samples. *Biochem Pharmacol* 45(10):2166–2169.
- Bornschein, RL; Hastings, L; Manson, JM. (1980) Behavioral toxicity in the offspring of rats following maternal exposure to dichloromethane. *Toxicol Appl Pharmacol* 52:29–37.
- Bos, PMJ; Zeilmaier, MJ; van Eijkeren, JCH. (2006) Application of physiologically based pharmacokinetic modeling in setting acute exposure guideline levels for methylene chloride. *Toxicol Sci* 91(2):576–585.
- Boublik, T; Fried, V; Hala, E. (1984) The vapor pressures of pure substances: selected values of the temperature dependence of the vapour pressures of some pure substances in the normal and low pressure region. Vol. 17. 2nd revised edition. Amsterdam, Netherlands: Elsevier, p. 42.
- Bowen, SE; Batis, JC; Paez-Martinez, N; et al. (2006) The last decade of solvent research in animal models of abuse: mechanistic and behavioral studies. *Neurotoxicol Teratol* 28(6):636–647.
- Breslow, NE; Day, NE. (1987) Statistical methods in cancer research. Volume II. The design and analysis of cohort studies. *IARC Sci Publ* 82:1–406.
- Briving, C; Hamberg, A; Kjellstrand, P; et al. (1986) Chronic effects of dichloromethane on amino acids, glutathione and phosphoethanolamine in gerbil brain. *Scand J Work Environ Health* 12:216–220.
- Brown, RP; Delp, MD; Lindstedt, SL; et al. (1997) Physiological parameter values for physiologically-based pharmacokinetic models. *Toxicol Ind Health* 13(4):407–484.
- Brown-Woodman, PDC; Hayes, LC; Huq, F; et al. (1998) In vitro assessment of the effect of halogenated hydrocarbons: chloroform, dichloromethane, and dibromoethane on embryonic development of the rat. *Teratology* 57:321–333.
- Bruhn, C; Bröckmoller, J; Kerb, R; et al. (1998) Concordance between enzyme activity and genotype of glutathione S-transferase theta (GST-T1). *Biochem Pharmacol* 56:1189–1193.
- Brzezinski, MR; Boutelet-Bochan, H; Person, RE; et al. (1999) Catalytic activity and quantitation of cytochrome P-450 2E1 in prenatal human brain. *J Pharmacol Exp Ther* 289:1648–1653.
- Burek, JD; Nitschke, KD; Bell, TJ; et al. (1984) Methylene chloride: a two-year inhalation toxicity and oncogenicity study in rats and hamsters. *Fundam Appl Toxicol* 4:30–47.
- Cagiano, R; Ancona, D; Cassano, T; et al. (1998) Effects of prenatal exposure to low concentrations of carbon monoxide on sexual behaviour and mesolimbic dopaminergic function in rat offspring. *Br J Pharmacol* 125(4):909–915.

- Callen, DF; Wolf, CR; Philpot, RM. (1980) Cytochrome P-450 mediated genetic activity and cytotoxicity of seven halogenated aliphatic hydrocarbons in *Saccharomyces cerevisiae*. *Mutat Res* 77:55–63.
- Candrian, U; You, M; Goodrow, T; et al. (1991) Activation of protooncogenes in spontaneously occurring non-liver tumors from C57BL/6 x C3H F1 mice. *Cancer Res* 51:1148–1153.
- Cantor, KP; Stewart, PA; Brinton, LA; et al. (1995) Occupational exposures and female breast cancer mortality in the United States. *J Occup Environ Med* 37:336–348.
- Carlsson, A; Hultengren, M. (1975) Exposure to methylene chloride. III. Metabolism of ¹⁴C-labelled methylene chloride in rat. *Scand J Work Environ Health* 1:104–108.
- Casanova, M; Deyo, DF; Heck, Hd'A. (1992) Dichloromethane (methylene chloride): metabolism to formaldehyde and formation of DNA-protein cross links in B6C3F₁ mice and Syrian golden hamsters. *Toxicol Appl Pharmacol* 114:162–165.
- Casanova, M; Conolly, RB; Heck, Hd'A. (1996) DNA-protein cross-links (DPX) and cell proliferation in B6C3F₁ mice but not Syrian golden hamsters exposed to dichloromethane: pharmacokinetics and risk assessment with DPX as dosimeter. *Fundam Appl Toxicol* 31:103–106.
- Casanova, M; Bell, DA; Heck, Hd'A. (1997) Dichloromethane metabolism to formaldehyde and reaction of formaldehyde with nucleic acids in hepatocytes of rodents and humans with and without glutathione S-transferase T1 and M1 genes. *Fundam Appl Toxicol* 37:168–180.
- Chang, Y; Yang, CC; Deng, JF; et al. (1999) Diverse manifestations of oral methylene dichloride poisoning: report of 6 cases. *Clin Toxicol* 37(4):497–504.
- Cherry, N; Venables, H; Waldron, HA; et al. (1981) Some observations on workers exposed to methylene chloride. *Br J Ind Med* 38:351–355.
- Cherry, N; Venables, H; Waldron, HA. (1983) The acute behavioural effects of solvent exposure. *J Soc Occup Med* 33:13–18.
- Clewell, HJ, III; Gearhart, JM; Andersen, ME. (1993) Analysis of the metabolism of methylene chloride in the B6C3F₁ mouse and its implications for human carcinogenic risk. Submitted to Occupational Safety and Health Administration, U.S. Department of Labor, Washington, DC. Docket # H-071, Exhibit #96.
- Clewell, HJ; Gentry, PR; Covington, TR; et al. (2004) Evaluation of the potential impact of age- and gender-specific pharmacokinetic differences on tissue dosimetry. *Toxicol Sci* 79:381–393.
- CMR (Chemical Marketing Reporter). (1973) Chemical profile on methylene chloride. *Chemical Marketing Reporter*, October 22, 1973.
- CMR (Chemical Marketing Reporter). (1979) Chemical profile on methylene chloride. *Chemical Marketing Reporter*, August 6, 1979.
- CMR (Chemical Marketing Reporter). (1982) Chemical profile on methylene chloride. *Chemical Marketing Reporter*, July 12, 1982.
- CMR (Chemical Marketing Reporter). (1997) Chemical profile on methylene chloride. *Chemical Marketing Reporter*, November 24, 1997.
- CMR (Chemical Marketing Reporter). (2000) Chemical profile on methylene chloride. *Chemical Marketing Reporter*, October 9, 2000.
- Cocco, P; Heineman, EF; Dosemeci, M. (1999) Occupational risk factors for cancer of the central nervous system (CNS) among US women. *Am J Ind Med* 36:70–74.

- Cohen J. (1987) Statistical power analysis for the behavioral sciences. Hillsdale, NJ: Lawrence Erlbaum Associates, p. 36–37.
- Condie, LW; Smallwood, CL; Laurie, RD. (1983) Comparative renal and hepatotoxicity of halomethanes: bromodichloromethane, bromoform, chloroform, dibromochloromethane, and methylene chloride. *Drug Chem Toxicol* 6(6):563–578.
- Crebelli, R; Benigni, R; Franekic, J; et al. (1988) Induction of chromosome malsegregation by halogenated organic solvents in *Aspergillus nidulans*: unspecific or specific mechanism? *Mutat Res* 201:401–411.
- Crebelli, R; Carere, A; Leopardi, P; et al. (1999) Evaluation of 10 aliphatic halogenated hydrocarbons in the mouse bone marrow micronucleus test. *Mutagenesis* 14(2):207–215.
- Dankovic, DA; Bailer, AJ. (1994) The impact of exercise and intersubject variability on dose estimates for dichloromethane derived from a physiologically based pharmacokinetic model. *Fundam Appl Toxicol* 22:20–25.
- David, RM; Clewell, HJ; Gentry, PR; et al. (2006) Revised assessment of cancer risk to dichloromethane II. Application of probabilistic methods to cancer risk determinations. *Regul Toxicol Pharmacol* 45:55–65.
- De Salvia, MA; Cagiano, R; Carratù, MR; et al. (1995) Irreversible impairment of active avoidance behavior in rats prenatally exposed to mild concentrations of carbon monoxide. *Psychopharmacology (Berl)* 122(1):66–71.
- Dearfield, KL; Moore, MM. (2005) Use of genetic toxicology information for risk assessment. *Environ Mol Mutagen* 46(4):236–245.
- DeMarini, DM; Shelton, ML; Warren, SH; et al. (1997) Glutathione S-transferase-mediated induction of GC→AT transitions by halomethanes in Salmonella. *Environ Mol Mutagen* 30:440–447.
- Devereux, TR; Foley, JF; Maronpot, RR; et al. (1993) *Ras* proto-oncogene activation in liver and lung tumors from B6C3F₁ mice exposed chronically to methylene chloride. *Carcinogenesis* 14(5):795–801.
- Dillon, D; Edwards, I; Combes, R; et al. (1992) The role of glutathione in the bacterial mutagenicity of vapour phase dichloromethane. *Environ Mol Mutagen* 20:211–217.
- Dinse GE; Lagakos SW. (1982) Nonparametric estimation of lifetime and disease onset distributions from incomplete observations. *Biometrics* 38(4):921–932.
- DiVincenzo, GD; Kaplan, CJ. (1981) Uptake, metabolism, and elimination of methylene chloride vapor by humans. *Toxicol Appl Pharmacol* 59:130–140.
- DiVincenzo, GD; Yanno, FJ; Astill, BD. (1971) The gas chromatographic analysis of methylene chloride in breath, blood, and urine. *Am Ind Hyg Assoc J* 32:387–391.
- DiVincenzo, GD; Yanno, FJ; Astill, BD. (1972) Human and canine exposures to methylene chloride vapor. *Am Ind Hyg Assoc J* 33:125–134.
- Doherty, AT; Ellard, S; Parry, EM; et al. (1996) An investigation into the activation and deactivation of chlorinated hydrocarbons to genotoxins in metabolically competent human cells. *Mutagenesis* 11(3):247–274.
- Dosemeci, M; Cocco, P; Chow, WH. (1999) Gender differences in risk of renal cell carcinoma and occupational exposures to chlorinated aliphatic hydrocarbons. *Am J Ind Med* 36:54–59.
- Dumas, S; Parent, ME; Siemiatycki, J. (2000) Rectal cancer and occupational risk factors: a hypothesis-generating, exposure-based case-control study. *Int J Cancer* 87:874–879.
- Eastmond, DA; Hartwig, A; Anderson, D; et al. (2009) Mutagenicity testing for chemical risk assessment: update of WHO/IPCS harmonized scheme. *Mutagenesis* 24:341–349.

- Ehrlich, R. (1980) Interaction between environmental pollutants and respiratory infections. *Environ Health Perspect* 35:89–100
- El-Masri, HA; Bell, DA; Portier, CJ. (1999) Effects of glutathione transferase theta polymorphism on the risk estimates of dichloromethane to humans. *Toxicol Appl Pharmacol* 158:221–230.
- El-Rayes, BF; Ali, S; Heilbrun, LK; et al. (2003) Cytochrome P450 and glutathione transferase expression in human breast cancer. *Clin Cancer Res* 9(5):1705–1709.
- Engström, J; Bjurström, R. (1977) Exposure to methylene chloride. Content in subcutaneous adipose tissue. *Scand J Work Environ Health* 3:215–224.
- Estill, CF; Spencer, AB. (1996) Case study: control of methylene chloride exposures during furniture stripping. *Am Ind Hyg Assoc J* 57:43–49.
- Evans, MV; Caldwell, JC. (2010) Evaluation of two different metabolic hypotheses for dichloromethane toxicity using physiologically based pharmacokinetic modeling for in vivo inhalation gas uptake data exposure in female B6C3F1 mice. *Toxicol Appl Pharmacol* Feb 10. [Epub ahead of print].
- Fechner, G; Ortmann, C; Du Chesne, A; et al. (2001) Fatal intoxication due to excessive dichloromethane inhalation. *Forensic Sci Int* 122:69–72.
- Fechter, LD. (1987) Neurotoxicity of prenatal carbon monoxide exposure. *Res Rep Health Eff Inst* 12:3–22.
- Fisher, JW; Whittaker, TA; Taylor, DH; et al. (1989) Physiologically based pharmacokinetic modeling of the pregnant rat: a multiroute exposure model for trichloroethylene and its metabolite, trichloroacetic acid. *Toxicol Appl Pharmacol* 99(3):395–414.
- Fodor, GG; Prajsnar, D; Schlipkoter, H. (1973) Endogenous CO formation by incorporated halogenated hydrocarbons of the methane series. *Staub Reinhalt Luft* 33:260–261.
- Foley, JF; Tuck, PD; Ton, T; et al. (1993) Inhalation exposure to a hepatocarcinogenic concentration of methylene chloride does not induce sustained replicative DNA synthesis in hepatocytes of female B6C3F₁ mice. *Carcinogenesis* 14(5):811–817.
- Forster, HV; Graff, S; Hake, CL; et al. (1974) Pulmonary-hematologic studies on humans during exposure to methylene chloride. Prepared by the Department of Environmental Medicine, Medical College of Wisconsin, Milwaukee, WI, for the National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, Cincinnati, OH. Available from the National Technical Information Service, Springfield, VA; PB82-151697.
- Foster, JR; Green, T; Smith, LL; et al. (1992) Methylene chloride-an inhalation study to investigate pathological and biochemical events occurring in the lungs of mice over an exposure period of 90 days. *Fundam Appl Toxicol* 18:376–388.
- Foster, JR; Green, T; Smith, LL; et al. (1994) Methylene chloride: an inhalation study to investigate toxicity in the mouse lung using morphological, biochemical, and Clara cell culture techniques. *Toxicology* 91:221–234.
- Friedlander, BR; Hearne, T; Hall, S. (1978) Epidemiologic investigation of employees chronically exposed to methylene chloride. *J Occup Med* 20:657–666.
- Fujimoto, K; Arakawa, S; Watanabe, T; et al. (2007) Generation and functional characterization of mice with a disrupted glutathione S-transferase, theta 1 gene. *Drug Metab Dispos* 35:2196–2202.
- Fuxe, K; Andersson, K; Hansson, T; et al. (1984) Central catecholamine neurons and exposure to dichloromethane. Selective changes in amine levels and turnover in tel- and diencephalic and Na nerve terminal systems and in the secretion of anterior pituitary hormone in the male rat. *Toxicol* 29:293–305.

- Gamberale, F; Annwall, G; Hultengren, M (1975) Exposure to methylene chloride. II. Psychological function. *Scand J Work Environ Health* 1:95–103.
- Gargas, ML; Clewell, HJ; Andersen, ME. (1986) Metabolism of inhaled dihalomethanes in vivo: differentiation of kinetic constants for two independent pathways. *Toxicol Appl Pharmacol* 82:211–223.
- Garrett, NE; Lewtas, J. (1983) Cellular toxicity in Chinese hamster ovary cells culture. I. Analysis of cytotoxicity endpoints for twenty-nine priority pollutants. *Environ Res* 32:455–465.
- Garte, S; Gaspari, L; Alexandrie, AK; et al. (2001) Metabolic gene polymorphism frequencies in control populations. *Cancer Epidemiol Biomarkers Prev* 10:1239–1248.
- General Electric Company. (1976) Dichloromethane and ninety day oral toxicity study in rats. Prepared by the International Research and Development Corporation, Mattawan, MI for the Plastics Tech Department, General Electric Co., Pittsfield, MA. Submitted under TSCA Section 8D; EPA Document No. 86-878210707; NTIS No. OTS0205887.
- Gibbs, GW. (1992) The mortality of workers employed at a cellulose acetate and triacetate fibers plant in Cumberland, Maryland: a “1970” cohort followed 1970–1989 [final report]. Prepared by Safety Health Environment International Consultants Corporation, Winterburn, Alberta, Canada, for the Hoechst Celanese Corporation, Somerville, NJ.
- Gibbs, GW; Amsel, J; Soden, K. (1996) A cohort mortality study of cellulose triacetate-fiber workers exposed to methylene chloride. *J Occup Environ Med* 38(7):693–697.
- Giustino, A; Cagiano, R; Carratù, MR; et al. (1999) Prenatal exposure to low concentrations of carbon monoxide alters habituation and non-spatial working memory in rat offspring. *Brain Res* 844(1–2):201–205.
- Glatzel, W; Tietze, K; Gutewort, R; et al. (1987) Interaction of dichloromethane and ethanol in rats: toxicokinetics and nerve conduction velocity. *Alcoholism: Clin Exp Res* 11:450–455.
- Gocke, E; King, MT; Eckhardt, K; et al. (1981) Mutagenicity of cosmetic ingredients licensed by the European communities. *Mutat Res* 90:91–109.
- Gomez, MR. (1996) Exposure determinants needed to improve the assessment of exposure. *Am J Ind Med* 29:569–570.
- Gomez, MR; Cocco, P; Dosemeci, M; et al. (1994) Occupational exposure to chlorinated aliphatic hydrocarbons: job exposure matrix. *Am J Ind Med* 26:171–183.
- Goodman, DG; Maronpot, RR; Newbene, PM, et al. (1994) Proliferative and selected other lesions in the liver of rats. G1-5. In: *Guides for toxicologic pathology. STP/ARP/AFIP*, Washington DC:1–24.
- Goulle, JP; Lacroix, C; Vaz, E; et al. (1999) Fatal case of dichloromethane poisoning. *J Anal Toxicol* 23:380–383.
- Graves, RJ; Green, T. (1996) Mouse liver glutathione S-transferase mediated metabolism of methylene chloride to a mutagen in the CHO/HPRT assay. *Mutat Res* 367:143–150.
- Graves, RJ; Callander, RD; Green, T. (1994a) The role of formaldehyde and S-chloromethylglutathione in the bacterial mutagenicity of methylene chloride. *Mutat Res* 320:235–243.
- Graves, RJ; Coutts, C; Eyton-Jones, H; et al. (1994b) Relationship between hepatic DNA damage and methylene chloride-induced hepatocarcinogenicity in B6C3F₁ mice. *Carcinogenesis* 15(5):991–996.
- Graves, RJ; Coutts, C; Green, T. (1995) Methylene chloride-induced DNA damage: an interspecies comparison. *Carcinogenesis* 16(8):1919–1926.

- Graves, RJ; Trueman, P; Jones, S; et al. (1996) DNA sequence analysis of methylene chloride-induced HPRT mutations in Chinese hamster ovary cells: comparison with the mutation spectrum obtained for 1,2-dibromoethane and formaldehyde. *Mutagenesis* 11(3):229–233.
- Green, T. (1983) The metabolic activation of dichloromethane and chloroform in a bacterial mutation assay using *Salmonella typhimurium*. *Mutat Res* 118:277–288.
- Green, T. (1989) A biological data base for methylene chloride risk assessment. In: Travis, CC; ed. *Biologically based methods for cancer risk assessment*. New York, NY: Plenum Press, p. 289–300.
- Green, T. (1997) Methylene chloride induced mouse liver and lung tumours: an overview of the role of mechanistic studies in human safety assessment. *Hum Exp Toxicol* 16:3–13.
- Guengerich, FP. (1997) Mechanisms of mutagenicity of DNA adducts derived from alkyl and vinyl halides. *Jpn J Toxicol Environ Health* 43:69–82.
- Haber, LT; Maier, A; Gentry, PR; et al. (2002) Genetic polymorphisms in assessing interindividual variability in delivered dose. *Reg Toxicol Pharmacol* 35:177–197.
- Hall, AH; Rumack, BH. (1990) Methylene chloride exposure in furniture-stripping shops: ventilation and respirator use practices. *J Occup Med* 32:33–37.
- Hallier, E; Schröder, KR; Asmuth, K; et al. (1994) Metabolism of dichloromethane (methylene chloride) to formaldehyde in human erythrocytes: influence of polymorphism of glutathione transferase theta (GST T1-1). *Arch Toxicol* 68:423–427.
- Hansch, C; Leo, A; Hoekman, D. (1995) Exploring QSAR. Hydrophobic, electronic, and steric constants. ACS professional reference book. Washington, DC: American Chemical Society; p. 3.
- Hardie, DWF. (1964) Methylene chloride. In: Kirk-Othmer encyclopedia of chemical technology. 2nd edition. New York, NY: John Wiley & Sons; pp. 111–119.
- Hardin, BD; Manson, J. (1980) Absence of dichloromethane teratogenicity with inhalation exposure in rats. *Toxicol Appl Pharmacol* 52:22–28.
- Hashmi, M; Dechert, S; Dekant, W; et al. (1994) Bioactivation of [¹³C]dichloromethane in mouse, rat, and human liver cytosol: ¹³C nuclear magnetic resonance spectroscopic studies. *Chem Res Toxicol* 7:291–296.
- Haufroid, V; Ligoocka, D; Buyschaert, M; et al. (2003) Cytochrome P4502E1 (CYP2E1) expression in peripheral blood lymphocytes: evaluation in hepatitis C and diabetes. *Eur J Clin Pharmacol* 59:29–33.
- Haun, CC; Harris, ES; Darmer, KI, Jr. (1971) Continuous animal exposure to methylene chloride. In: Proceedings of the 2nd conference of environmental toxicology; August 31–September 2; Fairborn, OH; Paper No. 10; AMRL-TR-71-120. Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base, OH, pp. 125–135. Available from the National Technical Information Service, Springfield, VA; AD751432 (individual paper); AD746660 (entire proceedings).
- Haun, CC; Vernot, EH; Darmer, KI, Jr; et al. (1972) Continuous animal exposure to low levels of dichloromethane. Proceedings of the 3rd conference of environmental toxicology; October 25–27; Fairborn, OH; Paper No. 12; AMRL-TR-72-120. Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base, OH, pp. 199–208. Available from the National Technical Information Service, Springfield, VA; AD773766.
- Hazleton Laboratories. (1983) 24-Month oncogenicity study of methylene chloride in mice [final report]. Performed by Hazleton Laboratories America, Inc., Vienna, VA for the National Coffee Association, New York, NY (Project No. 2112-106).
- Hearne, FT; Pifer, JW. (1999) Mortality study of two overlapping cohorts of photographic film base manufacturing employees exposed to methylene chloride. *J Occup Environ Med* 41(12):1154–1169.

Hearne, FT; Grose, F; Pifer, JW; et al. (1987) Methylene chloride mortality study: dose-response characterization and animal model comparison. *J Occup Med* 29(3):217–228.

Hearne, FT; Pifer, JW; Grose, F. (1990) Absence of adverse mortality effects in workers exposed to methylene chloride: an update. *J Occup Med* 32(3):234–240.

Hegi, ME; Söderkvist, P; Foley, JF; et al. (1993) Characterization of p53 mutations in methylene chloride-induced lung tumors from B6C3F₁ mice. *Carcinogenesis* 14(5):803–810.

Heineman, EF; Cocco, P; Gomez, MR; et al. (1994) Occupational exposure to chlorinated aliphatic hydrocarbons and risk of astrocytic brain cancer. *Am J Ind Med* 26:155–169.

Heineman, EF; Gomez, MR; Dosemeci, M; et al. (1996) Methylene chloride and brain cancer: interpreting a new study in light of existing literature. *Am J Ind Med* 30:506–507.

Hellmold, H; Rylander, T; Magnusson, M; et al. (1998) Characterization of cytochrome P450 enzymes in human breast tissue from reduction mammoplasties. *J Clin Endocrinol Metab* 83(3):886–895.

Heppel, LA; Neal, PA. (1944) Toxicology of dichloromethane (methylene chloride). II. Its effect upon running activity in the male rat. *J Ind Hyg Toxicol* 26(1):17–21.

Heppel, LA; Neal, PA; Perrin, TL; et al. (1944) Toxicology of dichloromethane (methylene chloride). *J Ind Hyg Toxicol* 26:8–16.

Herr, DW; Boyes, WK. (1997) A comparison of the acute neuroactive effects of dichloromethane, 1,3-dichloropropane, and 1,2-dichlorobenzene on rat flash evokes potentials (FEPs). *Fundam Appl Toxicol* 35:31–48.

Hines, RN. (2007) Ontogeny of human hepatic cytochromes P450. *J Biochem Mol Toxicol* 21:169–175.

Holbrook, MT. (2003) Methylene chloride. In: Kirk-Othmer encyclopedia of chemical technology. New York, NY: John Wiley & Sons. Available online at http://www.mrw.interscience.wiley.com/kirk/kirk_search_fs.html (accessed March 9, 2010). (subscription required).

Horvath, AL. (1982) Halogenated hydrocarbons: solubility-miscibility with water. New York, NY: Marcel Dekker, Inc., p. 543.

Hu, Y; Kabbler, SL; Tennant, AH et al. (2006) Induction of DNA-protein crosslinks by dichloromethane in a V79 cell line transfected with the murine glutathione-S-transferase theta 1 gene. *Mutation Res* 607:231–239.

Hughes, NJ; Tracey, JA. (1993) A case of methylene chloride (nitromors) poisoning, effects on carboxyhaemoglobin levels. *Hum Exp Toxicol* 12:159–160.

IARC (International Agency for Research on Cancer). (1999) Dichloromethane. IARC monographs on the evaluation of carcinogenic risk of chemicals to humans. Vol. 71. Re-evaluation of some organic chemicals, hydrazine and hydrogen peroxide. Lyon, France: International Agency for Research on Cancer, pp. 251–315. Available online at <http://inchem.org/documents/iarc/vol71/004-dichloromethane.html> (accessed March 9, 2010).

Infante-Rivard, C; Siemiatycki, J; Lakhani, R; et al. (2005) Maternal exposure to occupational solvents and childhood leukemia. *Environ Health Perspect* 113:787–792.

Ingelman-Sundberg, M. (2004) Human drug metabolising cytochrome P450 enzymes: properties and polymorphisms. *Naunyn-Schmiedeberg's Arch Pharmacol* 369:89–104.

Jacobovich, RM; Landau, D; Dayan, YB; et al. (2005) Facial nerve palsy after acute exposure to dichloromethane. *Am J Ind Med* 48:389–392.

Johnsrud, EK; Koukouritaki, SB; Divakaran, K; et al. (2003) Human hepatic CYP2E1 expression during development. *J Pharmacol Exp Ther* 307:402–407.

- Jongen, WMF; Alink, GM; Koeman, JH. (1978) Mutagenic effect of dichloromethane on *Salmonella typhimurium*. *Mutat Res* 56:245–248.
- Jongen, WMF; Lohman, PHM; Kottenhagen, MJ; et al. (1981) Mutagenicity testing of dichloromethane in short-term mammalian test systems. *Mutat Res* 81:203–213.
- Jongen, WMF; Harmsen, EGM; Alink, GM; et al. (1982) The effect of glutathione conjugation and microsomal oxidation on the mutagenicity of dichloromethane in *S. typhimurium*. *Mutat Res* 95:183–189.
- Jonsson, F; Johanson, G. (2001) A Bayesian analysis of the influence of GST-T1 polymorphism on the cancer risk estimate for dichloromethane. *Toxicol Appl Pharmacol* 174:99–112.
- Jonsson, F; Johanson, G. (2003) The Bayesian population approach to physiological toxicokinetic-toxicodynamic models—an example using the MCSim software. *Toxicol Lett* 138:143–150.
- Jonsson, F; Bois, F; Johanson, G. (2001) Physiologically based pharmacokinetic modeling of inhalation exposure of humans to dichloromethane during moderate to heavy exercise. *Toxicol Sci* 59:209–218.
- Juronen, E; Tasa, G; Uusküla, M; et al. (1996) Purification, characterization and tissue distribution of human class theta glutathione S-transferase T1-1. *Biochem Mol Biol Int* 39:21–9
- Kanada, M; Miyagawa, M; Sato, M; et al. (1994) Neurochemical profile of effects of 28 neurotoxic chemicals on the central nervous system in rats (1). Effects of oral administration on brain contents of biogenic amines and metabolites. *Ind Health* 32:145–164.
- Kanno, J; Foley, JF; Kari, F; et al. (1993) Effect of methylene chloride inhalation of replicative DNA synthesis in the lungs of female B6C3F₁ mice. *Environ Health Perspect* 101(Suppl. 5):271–276.
- Kari, FW; Foley, JF; Seilkop, SK; et al. (1993) Effect of varying exposure regimens on methylene chloride-induced lung and liver tumors in female B6C3F₁ mice. *Carcinogenesis* 14(5):819–826.
- Karlsson, JE; Rosengren, LE; Kjellstrand, P; et al. (1987) Effects of low-dose inhalation of three chlorinated aliphatic organic solvents on deoxyribonucleic acid in gerbil brain. *Scand J Work Environ Health* 13:453–458.
- Kayser, MF; Vuilleumier, S. (2001) Dehalogenation of dichloromethane by dichloromethane dehalogenase/glutathione s-transferase leads to formation of DNA adducts. *J Bacteriol* 183:5209–5212.
- Kelly, M. (1988) Case reports of individuals with oligospermia and methylene chloride exposures. *Reprod Toxicol* 2:13–17.
- Kernan, GJ; Ji, BT; Dosemeci, M; et al. (1999) Occupational risk factors for pancreatic cancer: a case-control study based on death certificates from 24 US states. *Am J Ind Med* 36:260–270.
- Kim, SK; Kim YC. (1996) Effect of a single administration of benzene, toluene or *m*-xylene on carboxyhaemoglobin elevation and metabolism of dichloromethane in rats. *J Appl Toxicol* 16:437–444.
- Kim, RB; O’Shea, D; Wilkinson, GR. (1995) Interindividual variability of chlorzoxazone 6-hydroxylation in men and women and its relationship to CYP2E1 genetic polymorphisms. *Clin Pharmacol Ther* 57(6):645–655.
- Kim, NY; Park, SW; Suh, JK. (1996) Two fatal cases of dichloromethane or chloroform poisoning. *J Forensic Sci* 41:527–529.
- Kirschman, JC; Brown, NM; Coots, RH; et al. (1986) Review of investigations of dichloromethane metabolism and subchronic oral toxicity as the basis for the design of chronic oral studies in rats and mice. *Food Chem Toxicol* 24(9):943–949.
- Kitchin, KT; Brown, JL. (1989) Biochemical effects of three carcinogenic chlorinated methanes in rat liver. *Teratog Carcinog Mutagen* 9:61–69.

- Kjellstrand, P; Holmquist, B; Jonsson, I; et al. (1985) Effects of organic solvents on motor activity in mice. *Toxicol* 35:35–46.
- Kolodner, K; Cameron, L; Gittlesohn, A; et al. (1990) Morbidity study of occupational exposure to methylene chloride using a computerized surveillance system. Center of Occupational Health and Environmental Health, Johns Hopkins School of Hygiene and Public Health, Baltimore, MD. Submitted to TSCA under Section 8D; EPA Document No. 86-900000421; NTIS No. OTS0522984.
- Korzekwa, KR; Krishnamachary, N; Shou, M; et al. (1998) Evaluation of atypical cytochrome P450 kinetics with two-substrate models: evidence that multiple substrates can simultaneously bind to cytochrome P450 active sites. *Biochemistry* 37:4137–4147.
- Kramers, PGN; Mout, HCA; Bissumbhar, B; et al. (1991) Inhalation exposure in *Drosophila* mutagenesis assays: experiments with aliphatic halogenated hydrocarbons, with emphasis on the genetic activity profile of 1,2-dichloroethane. *Mutat Res* 252:17–33.
- Landi, S; Naccarati, A; Ross, MK; et al. (2003) Induction of DNA strand breaks by trihalomethanes in primary human lung epithelial cells. *Mutat Res* 538:41–50.
- Landry, TD; Ramsey, JC; McKenna, MJ. (1983) Pulmonary physiology and inhalation dosimetry in rats: development of a method and two examples. *Toxicol Appl Pharmacol* 71:72–83.
- Lanes, SF; Cohen, A; Rothman, KJ; et al. (1990) Mortality of cellulose fiber production workers. *Scand J Work Environ Health* 16:247–251.
- Lanes, SF; Rothman, KJ; Dreyer, NA; et al. (1993) Mortality update of cellulose fiber production workers. *Scand J Work Environ Health* 19:426–428.
- Lash, AA; Becker, CE; So, Y; et al. (1991) Neurotoxic effects of methylene chloride: are they long lasting in humans? *Br J Ind Med* 48:418–426.
- Lefevre, PA; Ashby, J. (1989) Evaluation of dichloromethane as an inducer of DNA synthesis in the B6C3F₁ mouse liver. *Carcinogenesis* 10(6):1067–1072.
- Lehmann, L; Wagner, J. (2008) Gene expression of 17beta-estradiol-metabolizing isozymes: comparison of normal human mammary gland to normal human liver and to cultured human breast adenocarcinoma cells. *Adv Exp Med Biol* 617:617–624.
- Lehnebach, A; Kuhn, C; Pankow, D. (1995) Dichloromethane as an indicator of cytochrome c oxidase in different tissues of rats. *Arch Toxicol* 69:180–184.
- Leighton, DT, Jr; Calo, JM. (1981) Distribution coefficients of chlorinated hydrocarbons in dilute air-water systems for groundwater contamination applications. *J Chem Eng* 26:382–385.
- Leikin, JB; Kaufman, D; Lipscomb, JW; et al. (1990) Methylene chloride: report of five exposures and two deaths. *Am J Emerg Med* 8:534–537.
- Leitao, MM, Jr; White, P; Cracchiolo, B. (2008) Cervical cancer in patients infected with the human immunodeficiency virus. *Cancer* 112:2683–2689.
- Leuschner, F; Neumann, B; Hübscher, F. (1984) Report on subacute toxicological studies with dichloromethane in rats and dogs by inhalation. *Arzneim Forsch/Drug Res* 34(12):1772–1774.
- Lewis, RJ, Sr; ed. (1997) *Hawley's condensed chemical dictionary*. 13th edition. New York, NY: John Wiley & Sons, Inc., p. 736.
- Lide, DR; ed. (2000) *CRC handbook of chemistry and physics*. 81st edition. Boca Raton, FL: CRC Press, pp. 3–206.

Lipscomb, JC; Garrett, CM; Snawder, JE. (1997) Cytochrome P450-dependent metabolism of trichloroethylene: interindividual differences in humans. *Toxicol Appl Pharmacol* 142:311–318.

Lipscomb, JC; Teuschler, LK; Swartout, J; et al. (2003) The impact of cytochrome P450 2E1-dependent metabolic variance on a risk-relevant pharmacokinetic outcome in humans. *Risk Anal* 23(6):1221–1238.

Lorenz, J; Glatt, HR; Fleischmann, R; et al. (1984) Drug metabolism in man and its relationship to that in three rodent species: monooxygenase, epoxide hydrolase, and glutathione S-transferase activities in subcellular fractions of lung and liver. *Biochem Med* 32(1):43–56

Lowenfels, AB; Maisonneuve, P. (2005) Risk factors for pancreatic cancer. *J Cell Biochem* 95:649–656.

Lucas, D; Ferrara, R; Gonzalez, E; et al. (1999) Chlorzoxazone, a selective probe for phenotyping CYP2E1 in humans. *Pharmacogenetics* 9:377–388.

Lucas, D; Ferrara, R; Gonzales, E; et al. (2001) Cytochrome CYP2E1 phenotyping and genotyping in the evaluation of health risks from exposure to polluted environments. *Toxicol Lett* 124:71–81.

Madle, S; Dean, SW; Andrae, U; et al. (1994) Recommendations for the performance of UDS tests in vitro and in vivo. *Mutat Res* 312:263–285.

Mahmud, M; Kales, SN. (1999) Methylene chloride poisoning in a cabinet worker. *Environ Health Perspect* 107:769–772.

Mainwaring, GW; Williams, SM; Foster, JR; et al. (1996) The distribution of Theta-class glutathione S-transferase in the liver and lung of mouse, rat and human. *Biochem J* 318:297–303.

Maltoni, C; Cotti, G; Perino, G. (1988) Long-term carcinogenicity bioassays on methylene chloride administered by ingestion to Sprague-Dawley rats and Swiss mice and by inhalation to Sprague-Dawley rats. *Ann NY Acad Sci* 534:352–366.

Manno, M; Ruge, M; Cocheo, V. (1992) Double fatal inhalation of dichloromethane. *Hum Exp Toxicol* 11:540–545.

Marino, DJ; Clewell, HJ; Gentry, PR; et al. (2006) Revised assessment of cancer risk to dichloromethane: part I Bayesian PBPK and dose-response modeling in mice. *Regul Toxicol Pharmacol* 45:44–54.

Maronpot, RR; Devereux, TR; Hegi, M; et al. (1995) Hepatic and pulmonary carcinogenicity of methylene chloride in mice: a search for mechanisms. *Toxicology* 102:73–81.

Marriott, HM; Dockrell, DH. (2007) The role of the macrophage in lung disease mediated by bacteria. *Exp Lung Res* 33:493–505.

Marsch, GA; Mundkowski, R; Morris, BKJ; et al. (2001) Characterization of nucleoside and DNA adducts formed by s-(1-acetoxymethyl)glutathione and implications for dihalomethane-glutathione conjugates. *Chem Res Toxicol* 14:600–608.

Marsch, GA; Botta, S; Martin, MV; et al. (2004) Formation and mass spectrometric analysis of DNA and nucleoside adducts by S-(1-acetoxymethyl)glutathione and by glutathione S-transferase-mediated activation of dihalomethanes. *Chem Res Toxicol* 17:45–54.

Mathews, JM; Raymer, JH; Etheridge, AS; et al. (1997) Do endogenous volatile organic chemicals measured in breath reflect and maintain CYP2E1 levels in vivo? *Toxicol Appl Pharmacol* 146:255–260.

Mattsson, JL; Albee, RR; Eisenbrandt, DL. (1990) Neurotoxicologic evaluation of rats after 13 weeks of inhalation exposure to dichloromethane or carbon monoxide. *Pharmacol Biochem Behav* 36:671–681.

McKenna, MJ; Zempel, JA. (1981) The dose-dependent metabolism of [¹⁴C]methylene chloride following oral administration to rats. *Food Cosmet Toxicol* 19:73–78.

- McKenna, MJ; Zempel, JA; Braun, WH. (1982) The pharmacokinetics of inhaled methylene chloride in rats. *Toxicol Appl Pharmacol* 65:1–10.
- Mendoza-Cantú, A; Castorena-Torres, F; Bermudez, M; et al. (2004) Genotype and allele frequencies of polymorphic cytochromes P450 CYP1A2 and CYP2E1 in Mexicans. *Cell Biochem Funct* 22(1):29–34.
- Menear, JH; McConnell, EE; Huff, JE; et al. (1988) Inhalation and carcinogenesis studies of methylene chloride (dichloromethane) in F344/n rats and B6C3F₁ mice. *Ann NY Acad Sci* 534:343–351.
- Meyer, DJ; Coles, B; Pemble, SE; et al. (1991) Theta, a new class of glutathione transferases purified from rat and man. *Biochem J* 274:409–414.
- Miksys, S; Tyndale, RF. (2004) The unique regulation of brain cytochrome P450 2 (CYP2) family enzymes by drugs and genetics. *Drug Metab Rev* 36(2):313–333.
- Mirsalis, JC; Tyson, CK; Steinmetz, KL; et al. (1989) Measurement of unscheduled DNA synthesis and s-phase synthesis in rodent hepatocytes following in vivo treatment: testing of 24 compounds. *Environ Mol Mutagen* 14:155–164.
- Moser, VC; Cheek, BM; MacPhail, RC. (1995) A multidisciplinary approach to toxicological screening: III. Neurobehavioral toxicity. *J Toxicol Environ Health* 45:173–210.
- Narotsky, MG; Kavlock, RJ. (1995) A multidisciplinary approach to toxicological screening: II. Developmental toxicity. *J Toxicol Environ Health* 45:145–171.
- Nelson, HH; Wiencke, JK; Christiani, DC; et al. (1995) Ethnic differences in the prevalence of the homozygous deleted genotype of glutathione S-transferase theta. *Carcinogenesis* 16(5):1243–1245.
- NIOSH (National Institute of Occupational Safety and Health). (1986) Methylene chloride. Current intelligence bulletin 46. National Institute of Occupational Safety and Health, Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, Cincinnati, OH; DHHS (NIOSH) Publication No. 86-114. Available online at: http://www.cdc.gov/niosh/86114_46.html (accessed June 29, 2006).
- Nishimura, M; Yaguti, H; Yoshitsugu, H. (2003) Tissue distribution of mRNA expression of human cytochrome P450 isoforms assessed by high-sensitivity real-time reverse transcription PCR. *Yakugaku Zasshi* 123(5):369–375.
- Nitschke, KD; Burek, JD; Bell, TJ; et al. (1988a) Methylene chloride: a 2-year inhalation toxicity and oncogenicity study in rats. *Fundam Appl Toxicol* 11:48–59.
- Nitschke, KD; Eisenbrandt, DL; Lomax, LG; et al. (1988b) Methylene chloride: two-generation inhalation reproductive study in rats. *Fundam Appl Toxicol* 11:60–67.
- NLM (National Library of Medicine). (2003) Dichloromethane. HSDB (Hazardous Substances Data Bank). National Library of Medicine, National Institutes of Health, U.S. Department of Health and Human Services, Bethesda, Maryland. Available online at <http://toxnet.nlm.nih.gov/> (accessed March 9, 2010).
- Norman, WC, III; Boggs, P. (1996) Flawed estimates of methylene chloride exposures. *Am J Ind Med* 30:504–509.
- NRC (National Research Council). (1983) Risk assessment in the federal government: managing the process. Washington, DC: National Academy Press.
- NRC (National Research Council). (1994) Science and judgment in risk assessment. Washington, DC: National Academy Press.
- NTP (National Toxicology Program). (1986) Toxicology and carcinogenesis studies of dichloromethane (methylene chloride) (CAS No. 75-09-2) in F344/N rats and B6C3F₁ mice (inhalation studies). Public Health Service, U.S. Department of Health and Human Services; NTP TR 306. Available from the National Institute of Environmental

Health Sciences, Research Triangle Park, NC. Available online at http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr306.pdf (accessed March 9, 2010).

Oda, Y; Yamazaki, H; Thier, R; et al. (1996) A new *Salmonella typhimurium* NM5004 strain expressing rat glutathione S-transferase 5-5: use in detection of genotoxicity of dihaloalkanes using an SOS/*umu* test system. *Carcinogenesis* 17:297–302.

Oggenovski, VM; Marder, W; Somers EC; et al. (2004) Increased incidence of cervical intraepithelial neoplasia in women with systemic lupus erythematosus treated with intravenous cyclophosphamide. *J Rheumatol* 31:1763–1767.

Oh, SJ; Kim, SK; Kim, YC. (2002) Role of glutathione in metabolic degradation of dichloromethane in rats. *Toxicol Lett* 129:107–114.

Ojajärvi, A; Partanen, T; Ahlbom, A; et al. (2001) Risk of pancreatic cancer in workers exposed to chlorinated hydrocarbon solvents and related compounds: a meta-analysis. *Am J Epidemiol* 153:841–850.

O’Neil, MJ; Smith, A; Heckelman, PE; et al. (2001) Methylene chloride. In: *The Merck index: an encyclopedia of chemicals, drugs, and biologicals*. Whitehouse Station, NJ: Merck & Co., Inc., p. 1082.

Osterman-Golkar, S; Hussain, S; Walles, S; et al. (1983) Chemical reactivity and mutagenicity of some dihalomethanes. *Chem Biol Interact* 46:121–130.

OSHA (Occupational Safety and Health Administration). (1997) Occupational exposure to methylene chloride. *Federal Register* 62(7):1494–1611.

Ott, MG; Skory, LK; Holder, BB; et al. (1983a) Health evaluation of employees occupationally exposed to 7methylene chloride: general study design and environmental considerations. *Scand J Work Environ Health* 9(Suppl. 1):1–7.

Ott, MG; Skory, LK; Holder, BB; et al. (1983b) Health evaluation of employees occupationally exposed to methylene chloride: mortality. *Scand J Work Environ Health* 9(Suppl. 1):8–16.

Ott, MG; Skory, LK; Holder, BB; et al. (1983c) Health evaluation of employees occupationally exposed to methylene chloride: clinical laboratory evaluation. *Scand J Work Environ Health* 9(Suppl. 1):17–25.

Ott, MG; Skory, LK; Holder, BB; et al. (1983d) Health evaluation of employees occupationally exposed to methylene chloride: twenty-four hour electrocardiographic monitoring. *Scand J Work Environ Health* 9(Suppl. 1):26–30.

Ott, MG; Skory, LK; Holder, BB; et al. (1983e) Health evaluation of employees occupationally exposed to methylene chloride: metabolism data and oxygen half-saturation pressures. *Scand J Work Environ Health* 9(Suppl. 1):31–38.

Ott, MG; Carlo, GL; Steinberg, S; et al. (1985) Mortality among employees engaged in chemical manufacturing and related activities. *Am J Epidemiol* 122:311–322.

Pankow, D. (1988) Enhancement of dichloromethane-induced carboxyhemoglobinemia by isoniazid pretreatment. *Biomed Biochem Act* 3:293–295.

Pankow, D; Hoffmann, P. (1989) Dichloromethane metabolism to carbon monoxide can be induced by isoniazid, acetone and fasting. *Arch Toxicol Suppl* 13:302–303.

Pankow, D; Jagielki, S. (1993) Effect of methanol or modifications of the hepatic glutathione concentration on the metabolism of dichloromethane to carbon monoxide in rats. *Hum Exp Toxicol* 12:227–231.

Pankow, D; Kretschmer, S; Weise, M. (1991a) Effect of pyrazole on dichloromethane metabolism to carbon monoxide. *Arch Toxicol Suppl* 14:246–248.

- Pankow, D; Matschiner, F; Weigmann, H. (1991b) Influence of aromatic hydrocarbons on the metabolism of dichloromethane to carbon monoxide in rats. *Toxicol* 68:89–100.
- Pegram, RA; Andersen, ME; Warren, SH; et al. (1997) Glutathione S-transferase-mediated mutagenicity of trihalomethanes in *Salmonella typhimurium*: contrasting results with bromodichloromethane and chloroform. *Toxicol Appl Pharmacol* 144:183–188.
- Perocco, P; Prodi, G. (1981) DNA damage by haloalkanes in human lymphocytes cultured in vitro. *Cancer Lett* 13:213–218.
- Portier, K; Tolson, JK; Roberts, SM. (2007) Body weight distributions for risk assessment. *Risk Anal* 27(1):11–26.
- Putz, VR; Johnson BL; Setzer, JV. (1979) A comparative study of the effects of carbon monoxide and methylene chloride on human performance. *J Environ Pathol Toxicol* 2:97–112.
- Quondamatteo, F; Schulz, TG; Bunzel, N; et al. (1998) Immunohistochemical localization of glutathione S-transferase-T1 in murine kidney, liver, and lung. *Histochem Cell Biol* 110:417–423.
- Raimondi, S; Paracchini, V; Autrup, H; et al. (2006) Human genome epidemiology (HuGE) review. Meta- and pooled analysis of GST-T1 and lung cancer: a HuGE-GSEC review. *Am J Epidemiol* 164(11):1027–1042.
- Raje, R; Basso, M; Tolen, T; et al. (1988) Evaluation of in vivo mutagenicity of low-dose methylene chloride in mice. *J Am Coll Toxicol* 7(5):699–703.
- Ramsey, JR; Andersen, ME. (1984) A physiologically-based description of the inhalation pharmacokinetics of styrene in rats and humans. *Toxicol Appl Pharmacol* 73:159–175.
- Raphael, M; Nadiras, P; Flacke-Vordos, N. (2002) Acute methylene chloride intoxication—a case report on domestic poisoning. *Eur J Emerg Med* 9:57–59.
- Rebert, CS; Matteucci, MJ; Pryor, GT. (1989) Acute effects of inhaled dichloromethane on the EEG and sensory-evoked potentials of Fischer-344 rats. *Pharmacol Biochem Behav* 34:619–629.
- Reitz, RH. (1991) Estimating the risk of human cancer associated with exposure to methylene chloride. *Ann Ist Super Sanita* 27(4):609–614.
- Reitz, RH; McDougal, JN; Himmelstein, MW; et al. (1988a) In vitro metabolism of methylene chloride in human and animal tissues: use in physiologically based pharmacokinetic models. *Toxicol Appl Pharmacol* 97:230–246.
- Reitz, RH; Mendrala, AL; Park, CN; et al. (1988b) Incorporation of in vitro enzyme data into the physiologically-based pharmacokinetic (PB-PK) model for methylene chloride: implications for risk assessment. *Toxicol Lett* 43:97–116.
- Reitz, RH; Mendrala, AL; Guengerich, FP. (1989) In vitro metabolism of methylene chloride in human and animal tissues: use in physiologically based pharmacokinetic models. *Toxicol Appl Pharmacol* 97:230–246.
- Reitz, RH; Hays, SM; Gargas, ML. (1997) Addressing priority data needs for methylene chloride with physiologically based pharmacokinetic modeling. Prepared for the Agency for Toxic Substances and Disease Registry, Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA, on behalf of the Halogenated Solvents Industry Alliance (HSIA), Arlington, VA.
- Rhomberg, L. (1995) Use of quantitative modeling in methylene chloride risk assessment. *Toxicology* 102:95–114.
- Riley, EC; Fassett, DW; Sutton, WL. (1966) Methylene chloride vapor in expired air of human subjects. *Am Ind Hyg Assoc J* 27:341–348.
- Rioux, JP; Myers, RA. (1988) Methylene chloride poisoning: a paradigmatic review. *J Emerg Med* 6:227–238.

- Rodkey, FL; Collison, HA. (1977) Effect of dihalogenated methanes on the in vivo production of carbon monoxide and methane by rats. *Toxicol Appl Pharmacol* 40:39–47.
- Rodriguez-Arnaiz, R. (1998) Biotransformation of several structurally related 2B compounds to reactive metabolites in the somatic *w/w+* assay of *Drosophila melanogaster*. *Environ Mol Mutagen* 31(4):390–401.
- Roldán-Arjona, T; Pueyo, C. (1993) Mutagenic and lethal effects of halogenated methanes in the Ara test of *Salmonella typhimurium*: quantitative relationship with chemical reactivity. *Mutagenesis* 8(2):127–131.
- Rosengren, LE; Kjellstrand, P; Aurell, A; et al. (1986) Irreversible effects of dichloromethane on the brain after long term exposure: a quantitative study of DNA and the glial cell marker proteins S-100 and GFA. *Br J Ind Med* 43:291–299.
- Roth, RP; Drew, RT; Lo, RJ; et al. (1975) Dichloromethane inhalation, carboxyhemoglobin concentrations, and drug metabolizing enzymes in rabbits. *Toxicol Appl Pharmacol* 33:427–437.
- Rothman, KJ; Greenland, S. (1998) Precision and validity in epidemiologic studies. In: *Modern epidemiology*. 2nd edition. Philadelphia, PA: Lippincott-Raven Publishers, pp. 115–134.
- Sakai, T; Morita, Y; Wakui, C. (2002) Biological monitoring of workers exposed to dichloromethane using head-space gas chromatography. *J Chromat B* 778:245–250.
- Sasaki, YF; Saga, A; Akasaka, M; et al. (1998) Detection of in vivo genotoxicity of haloalkanes and haloalkenes carcinogenic to rodents by the alkaline single cell gel electrophoresis (comet) assay in multiple mouse organs. *Mutat Res* 419:13–20.
- Savolainen, H; Pfäffli, P; Tengén M; et al. (1977) Biochemical and behavioural effects of inhalation exposure to tetrachloroethylene and dichloromethane. *J Neuropathol Exp Neurol* 36:941–949.
- Savolainen, H; Kurppa, K; Pfäffli, P; et al. (1981) Dose-related effects of dichloromethane in rat brain in short-term inhalations exposure. *Chem-Biol Interact* 34:315–322.
- Schwetz, BA; Leong, BKJ; Gehring, PJ. (1975) The effect of maternally inhaled trichloroethylene, perchloroethylene, methyl chloroform, and methylene chloride on embryonal and fetal development in mice and rats. *Toxicol Appl Pharmacol* 32:84–96.
- Searles, J; McPhail, HA. (1949) Methylene chloride, CH₂Cl₂. In: Kirk, RE; Othmer, DF; eds. *Encyclopedia of chemical technology*. New York, NY: Interscience Encyclopedia, Inc., pp. 747–751.
- Selgrade, MK; Gilmour, MI. (2006) Immunotoxicology of inhaled compounds—assessing risks of local immune suppression and hypersensitivity. *J Toxicol Environ Health A* 69:827–844.
- Serota, DG; Thakur, AK; Ulland, BM; et al. (1986a) A two-year drinking water study of dichloromethane in rodents. I. Rats. *Food Chem Toxicol* 24(9):951–958.
- Serota, DG; Thakur, AK; Ulland, BM; et al. (1986b) A two-year drinking water study of dichloromethane in rodents. II. Mice. *Food Chem Toxicol* 24(9):959–963.
- Sheldon, T; Richardson, CR; Elliott, BM. (1987) Inactivity of methylene chloride in the mouse bone marrow micronucleus assay. *Mutagenesis* 2(1):57–59.
- Sherratt, PJ; Pulford, DJ; Harrison, DJ; et al. (1997) Evidence that human class Theta glutathione S-transferase T1-1 can catalyse the activation of dichloromethane, a liver and lung carcinogen in the mouse. *Biochem J* 326:837–846.
- Sherratt, PJ; Williams, S; Foster, J; et al. (2002) Direct comparison of the nature of mouse and human GST T1-1 and the implications on dichloromethane carcinogenicity. *Toxicol Appl Pharmacol* 179:89–97.

Shimada, T; Yamazaki, H; Mimura, M; et al. (1994) Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther* 270(1):414–423.

Shusterman, D; Quinlan, P; Lowengart, R; et al. (1990) Methylene chloride intoxication in a furniture refinisher. A comparison of exposure estimates utilizing workplace air sampling and blood carboxyhemoglobin measurements. *J Occup Med* 32:451–454.

Sills, RC; Hailey, JR; Neal, J; et al. (1999) Examination of low-incidence brain tumor responses in F344 rats following chemical exposures in National Toxicology Program carcinogenicity studies. *Toxicol Pathol* 27(5):589–599.

Simula, TP; Glancey, MJ; Wolf, CR. (1993) Human glutathione S-transferase-expressing *Salmonella typhimurium* tester strains to study the activation/detoxification of mutagenic compounds: studies with halogenated compounds, aromatic amines and aflatoxin B1. *Carcinogenesis* 14(7):1371–1376.

Soden, KJ. (1993) An evaluation of chronic methylene chloride exposure. *J Occup Med* 35(3):282–286.

Soden, KJ; Marras, G; Amsel, J. (1996) Carboxyhemoglobin levels in methylene chloride-exposed employees. *J Occup Environ Med* 38(4):367–371.

Spirtas, R; Stewart, PA; Lee, JS; et al. (1991) Retrospective cohort mortality study of workers at an aircraft maintenance facility. I. Epidemiological results. *Br J Ind Med* 48:515–530.

SRC (Syracuse Research Corporation). (1989) A review of in vitro test methodology for assessment of hepatotoxicity with a view to application to chemical mixtures. Prepared for U.S. EPA, Environmental Criteria and Assessment Office, U.S. Environmental Protection Agency; SRC TR-89-205.

Stephens, EA; Taylor, JA; Kaplan, N; et al. (1994) Ethnic variation in the CYP2E1 gene: polymorphism analysis of 695 African-Americans, European-Americans and Taiwanese. *Pharmacogenetics* 4:185–192.

Stewart, RD; Fisher, TN; Hosko, MJ; et al. (1972a) Experimental human exposure to methylene chloride. *Arch Environ Health* 25:342–348.

Stewart, RD; Fisher, TN; Hosko, MJ; et al. (1972b) Carboxyhemoglobin elevation after exposure to dichloromethane. *Science* 176:295–296.

Stewart, PA; Lee, JS; Marano, DE; et al. (1991) Retrospective cohort mortality study of workers at an aircraft maintenance facility. II Exposures and their assessment. *Br J Ind Med* 48:531–537.

Stott, WT; McKenna, MJ. (1984) The comparative absorption and excretion of chemical vapors by the upper, lower, and intact respiratory tract of rats. *Fundam Appl Toxicol* 4:594–602.

Strange, RC; Howie, AF; Hume, R; et al. (1989) The development expression of alpha-, mu- and pi-class glutathione S-transferases in human liver. *Biochim Biophys Acta* 993(2–3):186–190.

Sweeney, LM; Gargas, ML; Strother, DE; et al. (2003) Physiologically based pharmacokinetic model parameter estimation and sensitivity and variability analyses for acrylonitrile disposition in humans. *Toxicol Sci* 71:27–40.

Sweeney, LM; Kirman, CR; Morgott, DA; et al. (2004) Estimation of interindividual variation in oxidative metabolism of dichloromethane in human volunteers. *Toxicol Lett* 154:201–216.

Takehita, H; Mogi, K; Yasuda, T; et al. (2000) Postmortem absorption of dichloromethane: a case study and animal experiments. *Int J Legal Med* 114:96–100.

Taskinen, H; Lindbohm, ML; Hemminki, K. (1986) Spontaneous abortions among women working in the pharmaceutical industry. *Br J Ind Med* 43:199–205.

- Tay, P; Tan, KT; Sam, CT. (1995) Fatal gassing due to methylene chloride—a case report. *Singapore Med J* 36:444–445.
- Teschke, K; Olshan, AF; Daniels, JL; et al. (2002) Occupational exposure assessment in case-control studies: opportunities for improvement. *Occup Environ Med* 59:575–593.
- Thier, R; Taylor, JB; Pemble, SE; et al. (1993) Expression of mammalian glutathione S-transferase 5-5 in *Salmonella typhimurium* TA1535 leads to base-pair mutations upon exposure to dihalomethanes. *Proc Natl Acad Sci USA* 90:8576–8580.
- Thier, R; Wiebel, FA; Hinkel, A; et al. (1998) Species differences in the glutathione transferase GST-T1-1 activity towards the model substrates methyl chloride and dichloromethane in the liver and kidney. *Arch Toxicol* 72:622–629.
- Thilagar, AK; Kumaroo, V. (1983) Induction of chromosome damage by methylene chloride in CHO cells. *Mutat Res* 116:361–367.
- Thilagar, AK; Back, AM; Kirby, PE; et al. (1984) Evaluation of dichloromethane in short term in vitro genetic toxicity assays. *Environ Mutagen* 6:418–419.
- Thomas, AA; Pinkerton, MK; Warden, JA. (1972) Effects of low level dichloromethane exposure on the spontaneous activity of mice. Proceedings of the 3rd conference of environmental toxicology; October 25–27; Fairborn, OH; Paper No. 14; AMRL-TR-72-130. Aerospace Medical Research Laboratory, Wright-Patterson Air Force Base, OH; pp. 223–238. Available from the National Technical Information Service, Springfield, VA; AD773766.
- Tomenson, JA; Bonner, SM; Heijne, CG; et al. (1997) Mortality of workers exposed to methylene chloride employed at a plant producing cellulose triacetate film base. *Occup Environ Med* 54:470–476.
- Treluyer, JM; Cheron, G; Sonnier M; et al. (1996) Cytochrome P-450 expression in sudden infant death syndrome. *Biochem Pharmacol* 52:497–504.
- Trueman, RW; Ashby, J. (1987) Lack of UDS activity in the livers of mice and rats exposed to dichloromethane. *Environ Mol Mutagen* 10:189–195.
- U.S. Coast Guard. (1999) Dichloromethane. CHRIS: hazardous chemical data. U.S. Coast Guard, Department of Transportation, Washington, DC. Available online at <http://www.chrismanual.com/findform.htm>.
- U.S. EPA (Environmental Protection Agency). (1986a) Guidelines for the health risk assessment of chemical mixtures. *Federal Register* 51(185):34014–34025.
- U.S. EPA (Environmental Protection Agency). (1986b) Guidelines for mutagenicity risk assessment. *Federal Register* 51(185):34006–34012. Available online at <http://www.epa.gov/iris/backgrd.html> (accessed March 9, 2010).
- U.S. EPA (Environmental Protection Agency). (1987a) Update to the health assessment document and addendum for dichloromethane (methylene chloride): pharmacokinetics, mechanism of action and epidemiology [review draft]. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/8-87/030A. Available from the National Technical Information Service, Springfield, VA, PB87228565, and online at <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=30001GFH.txt> (accessed March 9, 2010).
- U.S. EPA (Environmental Protection Agency). (1987b) Technical analysis of new methods and data regarding dichloromethane hazard assessments [review draft]. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/8-87/029A. Available from the National Technical Information Service, Springfield, VA, PB87-228557, and online at <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=30001GAN.txt> (accessed March 9, 2010).
- U.S. EPA (Environmental Protection Agency). (1988a) Recommendations for and documentation of biological values for use in risk assessment. Environmental Criteria and Assessment Office, Office of Health and Environmental Assessment, Cincinnati, OH; EPA/600/6-87/008. Available from the National Technical Information

Service, Springfield, VA, PB88-179874/AS, and online at <http://cfpub.epa.gov/ncea/cfm/recorddisplay.cfm?deid=34855> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (1988b) The impact of pharmacokinetics on the risk assessment of dichloromethane. Office of Health and Environmental Assessment, Office of Research and Development, Washington, DC; EPA/600/D-88/219. Available from National Technical Information Service, Springfield, VA; PB89-173249.

U.S. EPA (Environmental Protection Agency). (1991) Guidelines for developmental toxicity risk assessment. Federal Register 56(234):63798–63826. Available online at <http://www.epa.gov/iris/backgrd.html> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (1994a) Interim policy for particle size and limit concentration issues in inhalation toxicity: notice of availability. Federal Register 59(206):53799. Available online at <http://www.epa.gov/EPA-PEST/1994/October/Day-26/pr-11.html> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (1994b) Methods for derivation of inhalation reference concentrations and application of inhalation dosimetry. Environmental Criteria and Assessment Office, Office of Health and Environmental Assessment, Cincinnati, OH; EPA/600/8-90/066F. Available from the National Technical Information Service, Springfield, VA, PB2000-500023, and online at <http://cfpub.epa.gov/ncea/raf/recorddisplay.cfm?deid=71993> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (1995) Use of the benchmark dose approach in health risk assessment. Risk Assessment Forum, Washington, DC; EPA/630/R-94/007. Available from the National Technical Information Service, Springfield, VA, PB95-213765, and online at http://cfpub.epa.gov/ncea/raf/raf_pubtitles.cfm?detype=document&excCol=archive (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (1996) Guidelines for reproductive toxicity risk assessment. Federal Register 61(212):56274–56322. Available online at <http://www.epa.gov/iris/backgrd.html> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (1998) Guidelines for neurotoxicity risk assessment. Federal Register 63(93):26926–26954. Available online at <http://www.epa.gov/iris/backgrd.html> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2000a) Science policy council handbook: risk characterization. Office of Science Policy, Office of Research and Development, Washington, DC; EPA/100-B-00-002. Available online at <http://www.epa.gov/OSA/spc/pdfs/prhandbk.pdf> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2000b) Benchmark dose technical guidance document [external review draft]. Risk Assessment Forum, Washington, DC; EPA/630/R-00/001. Available online at <http://www.epa.gov/iris/backgrd.html> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2000c) Supplementary guidance for conducting health risk assessment of chemical mixtures. Risk Assessment Forum, Washington, DC; EPA/630/R-00/002. Available online at http://cfpub.epa.gov/ncea/raf/chem_mix.cfm (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2000d) Toxicological review of vinyl chloride. Integrated Risk Information System (IRIS), National Center for Environmental Assessment, Washington, DC; EPA/635/R-00/004. Available online at <http://www.epa.gov/iris> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2002) A review of the reference dose concentration and reference concentration processes. Risk Assessment Forum, Washington, DC; EPA/630/P-02/002F. Available online at http://cfpub.epa.gov/ncea/raf/raf_pubtitles.cfm?detype=document&excCol=archive (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2005a) Guidelines for carcinogen risk assessment. Federal Register 70(66):17765–18717. Available online at <http://www.epa.gov/cancerguidelines> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2005b) Supplemental guidance for assessing susceptibility from early-life exposure to carcinogens. Risk Assessment Forum, Washington, DC; EPA/630/R-03/003F. Available online at <http://www.epa.gov/cancerguidelines> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2006a) Science policy council handbook: peer review. 3rd edition. Office of Science Policy, Office of Research and Development, Washington, DC; EPA/100/B-06/002. Available online at <http://www.epa.gov/OSA/spc/2peerrev.htm> (accessed March 9, 2010).

U.S. EPA (Environmental Protection Agency). (2006b) A framework for assessing health risk of environmental exposures to children. National Center for Environmental Assessment, Washington, DC; EPA/600/R-05/093F. Available online at <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=158363> (accessed March 9, 2010).

Vieira, I; Sonnier, M; Cresteil, T. (1996) Developmental expression of CYP2E1 in human liver. Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 238:476–483.

Warbrick, EV; Kilgour, JD; Dearman, RJ; et al. (2003) Inhalation exposure to methylene chloride does not induce systemic immunotoxicity in rats. *J Toxicol Environ Health A* 66:1207–1219.

Warholm, M; Alexandrie, AK; Högberg, J; et al. (1994) Polymorphic distribution of glutathione transferase activity with methyly chloride in human blood. *Pharmacogenetics* 4:307–311.

Watanabe, K; Guengerich, FP. (2006) Limited reactivity of formyl chloride with glutathione and relevance to metabolism and toxicity of dichloromethane. *Chem Res Toxicol* 19(8):1091–1096.

Watanabe, K; Liberman, RG; Skipper, PL; et al. (2007) Analysis of DNA adducts formed in vivo in rats and mice from 1,2-dibromoethane, 1,2-dichloroethane, dibromomethane, and dichloromethane using HPLC/accelerator mass spectrometry and relevance to risk estimates. *Chem Res Toxicol* 20:1594–600.

Weinstein, RS; Boyd, DD; Back, KC. (1972) Effects of continuous inhalation of dichloromethane in the mouse: morphologic and functional observations. *Toxicol Appl Pharmacol* 23:660–679.

Wells, VE; Schrader, SM; McCamon, CS; et al. (1989) Letter to the editor. *Reprod Toxicol* 3:281–282.

Westbrook-Collins, B; Allen, JW; Sharief, Y; et al. (1990) Further evidence that dichloromethane does not induce chromosome damage. *J Appl Toxicol* 10(2):79–81.

Winneke, G. (1974) Behavioral effects of methylene chloride and carbon monoxide as assessed by sensory and psychomotor performance. In: Xintaras, C; Johnson, BL; de Groot, I; eds. *Behavioral toxicology: early detection of occupational hazards*. Washington, DC: National Institute for Occupational Safety and Health, Center for Disease Control, Public Health Service, U.S. Department of Health, Education and Welfare; pp. 130–144.

Wirkner, K; Damme, B; Peolchen, W; et al. (1997) Effect of long-term ethanol pretreatment on the metabolism of dichloromethane to carbon monoxide in rats. *Toxicol Appl Pharmacol* 143:83–88.

Withey, JR; Karpinski, K. (1985) The fetal distribution of some aliphatic chlorinated hydrocarbons in the rat after vapor phase exposure. *Biol Res Pregnancy Perinatol* 6(2):79–88.

Zarrabeitia, MT; Ortega, C; Altuzarra, E; et al. (2001) Accidental dichloromethane fatality: a case report. *J Forensic Sci* 46:726–727

Zeiger, E. (1990) Mutagenicity of 42 chemicals in Salmonella. *Environ Mol Mutagen* 16:32–54.

Zielenska, M; Ahmed, A; Pienkowska, M; et al. (1993) Mutational specificities of environmental carcinogens in the *lacI* gene of *Escherichia coli*. VI. Analysis of methylene chloride-induced mutational distribution in Uvr+ and UvrB- strains. *Carcinogenesis* 14(5):789–794.

**APPENDIX A: SUMMARY OF EXTERNAL PEER REVIEW AND PUBLIC
COMMENTS AND DISPOSITION**

[page intentionally left blank]

APPENDIX B: HUMAN PBPK DICHLOROMETHANE MODEL

B.1. HUMAN MODEL DESCRIPTION

The basic model structure used by David et al. (2006) was that of Andersen et al. (1987) with the addition of the CO submodel of Andersen et al. (1991), refinements from the Marino et al. (2006) mouse model, and an inclusion of CYP metabolism in rapidly perfused tissue (see Figure B-1).




Figure B-1. Schematic of the David et al. (2006) PBPK model for dichloromethane in the human.

In order to incorporate known variability in human physiology and metabolism of dichloromethane into internal dosimetry, Monte Carlo analysis of the human model was performed to derive probability distributions of internal dose, as reported in David et al. (2006), but with changes in some of the key distributions as described below. The shape of the resulting dose distribution can be used to quantify the variability and uncertainty in internal dose with respect to variability in human physiology and variability and uncertainty in dichloromethane metabolism. The human model was run repeatedly using a random sample of each parameter from its respective parameter distribution in each iteration. Internal doses predicted for all iterations collectively defined a distribution for internal dose. The Monte Carlo analysis was run for 10- or 20,000 iterations. Repeated Monte Carlo analyses (at 10,000 iterations each) yielded 99th percentile values of internal dose in the liver or lung that differed by <2%. Normal or log-normal distributions of physiological and metabolic parameter and partition coefficient values were described by a mean, SD, and in most cases upper and lower truncation bounds. Physiological parameter and partition coefficient values were initially taken from the literature as described in David et al. (2006) and their distributions were assumed to be true variability (physiological parameters) or a level of uncertainty and variability (partition coefficients), neither of which could be meaningfully informed by the dichloromethane pharmacokinetic data.

Hence, the distributions for physiological parameters and partition coefficients were not updated in the Bayesian analysis of David et al. (2006).

The first-order oral absorption rate constant, $k_a = 5.0/\text{hour}$, was used in conjunction with human drinking water exposures simulated as six discrete drinking water episodes for specified times (25, 10, 25, 10, 25, and 5% of total daily intake at hours 0, 3, 5, 8, 11, and 15 of each day) (Reitz et al., 1997). Metabolic parameter distributions were derived from multiple human data sets by using MCMC calibration, also described in David et al. (2006).

The variability of genotypic expression of GST-T1 activity (the mechanism for GST-mediated metabolism of dichloromethane) was simulated as a uniform discrete distribution of the three GST-T1 genotypes (+/+, +/-, -/-) with varying activities in the liver and lung. The genotype frequency was based on data from Haber et al. (2002), with a frequency of genotypes of 32, 48, and 20 in the +/+, +/-, and -/- groups, respectively. GST activities measured by Warholm et al. (1994) for the three genotypes in a group of 208 healthy male and female subjects from Sweden were scaled by David et al. (2006) to obtain distributions of k_{fC} for each genotype that, when weighted by estimated frequencies of the genotypes in U.S. populations, would result in a distribution of k_{fC} activities with a mean equal to $0.852/\text{hour}\cdot\text{kg}^{0.3}$, which is the mean estimate of the population-mean value of k_{fC} obtained from the Bayesian analysis. The resulting distributions of internal lung and liver dose in human populations would have a theoretical probability of 20% for zero exposure to GST-mediated metabolites, and hence zero cancer risk for that 20% of the population. The final parameter distributions used by David et al. (2006) are summarized in Table B-1.

Table B-1. Parameter distributions used in human Monte Carlo analysis for dichloromethane by David et al. (2006)

Parameter		Distribution		Source
		Mean (arithmetic)	SD	
BW	Body weight (kg)	70.0	21.0	Humans ^a
QCC	Cardiac output (L/hr·kg ^{0.74})	16.5	1.49	Humans ^a
VPR	Ventilation:perfusion ratio	1.45	0.203	Humans ^a
QFC	Fat	0.05	0.0150	Humans ^a
QLC	Liver	0.26	0.0910	Humans ^a
QRC	Rapidly perfused tissues	0.50	0.10	Humans ^a
QSC	Slow perfused tissues	0.19	0.0285	Humans ^a
Tissue volumes (fraction BW)				
VFC	Fat	0.19	0.0570	Humans ^a
VLC	Liver	0.026	0.00130	Humans ^a
VLuC	Lung	0.0115	0.00161	Humans ^a
VRC	Rapidly perfused tissues	0.064	0.00640	Humans ^a
VSC	Slowly perfused tissues (muscle)	0.63	0.189	Humans ^a

Table B-1. Parameter distributions used in human Monte Carlo analysis for dichloromethane by David et al. (2006)

Parameter		Distribution		Source
		Mean (arithmetic)	SD	
Partition coefficients				
PB	Blood:air	9.7	0.970	Humans ^b
PF	Fat:blood	12.4	3.72	Rats ^b
PL	Liver:blood	1.46	0.292	Rats ^b
PLu	Lung:arterial blood	1.46	0.292	Rats ^b
PR	Rapidly perfused tissue:blood	1.46	0.292	Rats ^b
PS	Slowly perfused tissue (muscle:blood)	0.82	0.164	Rats ^b
Metabolism parameters				
V _{maxC}	Maximum metabolism rate (mg/hr·kg ^{0.7})	9.34	1.73	Calibration ^c
K _m	Affinity (mg/L)	0.433	0.146	Calibration ^c
A1	Ratio of lung V _{max} to liver V _{max}	0.000993	0.000396	Calibration ^c
A2	Ratio of lung KF to liver KF	0.0102	0.00739	Calibration ^c
FracR	Fractional CYP2E1 capacity in rapidly perfused tissue	0.0193	0.0152	Calibration ^c
First order metabolism rate (/hr·kg ^{0.3})				
k _{fC}	Homozygous (-/-)	0	0	Calibration ^c
	Heterozygous (+/-)	0.676	0.123	Calibration ^c
	Homozygous (+/+)	1.31	0.167	Calibration ^c
Oral absorption				
k _a	First-order oral absorption rate constant (/hr)	5.0		Reitz et al. (1997) (point estimate)

^aU.S. EPA (2000d) human PBPK model used for vinyl chloride.

^bAndersen et al. (1987). Blood:air partition measured by using human samples; other partition coefficients based on estimates from tissue measures in rats.

^cBayesian calibration based on five data sets (see text for description); posterior distributions presented in this table.

Source: David et al. (2006).

B.2. REVISIONS TO PARAMETER DISTRIBUTIONS OF DAVID ET AL. (2006)

An evaluation of the David et al. (2006) model and parameterization was undertaken, focusing on the adequacy of the characterization of parameter distributions in the full human population. EPA's conclusion is that the reported distributions for physiological parameters in particular, but also key metabolic parameters, only represented a narrow set of adults (with the exception of BW) or failed to include the parameter uncertainty from the Bayesian analysis. Therefore, supplemental data sources were chosen to define a number of the physiological parameter distributions in a way that should fully characterize the variability in the human population for individuals between 6 months and 80 years of age. Since many physiological parameters vary with age and gender, a structured approach will be used where an individual's age and then gender may be selected from the overall population distribution for these

characteristics (the male:female ratio in the population declines with age, for example). Dosimetry simulations can then be run for each such individual to obtain an overall population distribution of internal doses. Thus, each dosimetry distribution represents a “snapshot” of the dosimetry in a given individual at a given age and age- and gender-appropriate sampled body composition (fraction of BW in each tissue group). Finally, the sampling for two key metabolic parameters representing metabolism by CYP2E1 (i.e., $V_{\max C}$) and GST-T1 (i.e., k_{fC}) was adjusted to explicitly account for both the interindividual variability and the parameter-value uncertainty among humans.

In estimating the human equivalent exposure levels for noncancer endpoints, which are presumed to be relatively short-term effects possibly occurring from exposures of several weeks or months, using the distribution of such dosimetry “snapshots” should provide precisely the correct distribution to estimate overall population risk. For estimating cancer risk where risk is due to the cumulative exposure over months or years, however, the ideal approach would be to simulate the time-course of internal doses in a given individual tracked over a lifetime or significant portion thereof. But doing so would require estimating time-courses for each physiological and metabolic parameter in the individual over that time-period, a task which would be far more complicated than the structured “snapshot” approach used here. For example, while the CYP2E1 activity in an individual at age 12 is probably predictive of the activity in that individual at age 70 (e.g., someone who has above-average CYP2E1 activity when younger may well continue to be above average throughout his or her lifetime), we simply do not have the information or model structure to predict the time-dependences. Further, we know, for example, that some individuals who are lean in their youth may become obese by middle-age, while others (through lifestyle-changes) change in the opposite direction; and these changes may be reversed by the time the individual reaches 70 or 80 years of age.

Therefore a “life-course” dosimetry for specific individuals has not been calculated. For calculating HECs for noncancer effects, however, this means that the exposure level is identified such that 99% of the population at a given time is predicted to have an internal dose at or below a POD (defined as an internal dose level). It then seems highly likely that if one were to track individual exposure over time, one would also find that this equivalent exposure keeps 99% of the population from exceeding the POD. In short, if 99% of individuals’ snapshot-internal doses are below the POD (i.e., 99% of the internal doses in a cross-section of society on a given day), it can be anticipated that no more than 1% of all people will exceed that POD at any point in their life-times, even though the model simulations did not specifically track the changes in internal dose with age. For cancer risk, it can likewise be assumed that the average internal dose per unit exposure in the population as a whole (or the GST-T1 +/- portion of the population) at a given point in time is a good estimate of the average one would estimate if the internal dose was tracked over the lifetime in the same population—where the distribution of physiological and

metabolic characteristics at a given age is the same as used to estimate the average internal dose distribution for a cross-section of society.

B.3. CYP2E1 AND GST-T1

This evaluation incorporated additional data concerning the variability in CYP2E1 activity among humans, based on Lipscomb et al. (2003). The Lipscomb et al. (2003) study was based on in vitro analysis of liver samples from 75 human tissue donors (activity towards trichloroethylene and measurements of protein content) to estimate a distribution of activity in the population. The distribution used by David et al. (2006) for hepatic CYP2E1 activity for dichloromethane ($V_{\max C}$) was a truncated log-normal distribution with GM = 9.34 mg/hour-kg^{0.7}, GSD = 1.14, lower bound = 6.33 (68% of mean), and upper bound = 13.8 (218% of mean). However, Lipscomb et al. (2003; Table IV), analyzing data from a larger set of human tissue donors, derived an ultimate distribution for CYP2E1 activity with TCE (in units of pmol oxidized/minute/g liver) with GSD = 1.7274, 5th percentile = 40.7% of the mean, and 95th percentile = 245.8% of the mean. These data support a wider distribution in CYP2E1 activity than had been used in the David et al. (2006) model, with approximately a sixfold range between the upper and lower bounds in Lipscomb et al. (2003) and a twofold range in David et al. (2006). Since the distribution for $V_{\max C}$ (CYP2E1) parameterized by the posterior-distribution parameters in Table 4 of David et al. (2006) represents the population mean and uncertainty in that mean, that uncertainty is not reduced or replaced by the knowledge of variability gained from the data of Lipscomb et al. (2003). Therefore, in EPA's Monte-Carlo simulations, a two-dimensional sampling process was used for $V_{\max C}$. First the population-mean value of $V_{\max C}$, $V_{\max C, \text{mean}}$, was sampled from the range of uncertainty represented by a log-normal distribution with GM = 9.34 mg/hour-kg^{0.7} and GSD = 1.14 mg/hour-kg^{0.7} (values converted from linear-space mean/SD of 9.42 and 1.23 mg/hour-kg^{0.7} reported by David et al. [2006]) and upper/lower bounds of 7.20 and 12.11 mg/hour-kg^{0.7}, respectively (± 2 SD in log-space). After obtaining the sample population mean ($V_{\max C, \text{mean}}$), an individual $V_{\max C}$ value was then obtained by sampling from the log-normal distribution with that mean but GSD = 1.73 as obtained from the data of Lipscomb et al. (2003). Further, since even the data available to Lipscomb et al. (2003) were limited, and the log-normal distribution is naturally bounded to be greater than zero, a nontruncated distribution was used for this second dimension (step) of parameter sampling.

For GST-T1-mediated metabolism characterized by the rate coefficient, k_{TC} , David et al. (2006) replaced their estimates of population mean, uncertainty, and variability (the latter is not reported by David et al. [2006] but would have been estimated along with the uncertainty as part of the Bayesian analysis) with a measure of population variability alone. The population variability was obtained by using the known distribution of GST-T1 genotypes in the U.S. population (from Haber et al., 2002) and the genotype-specific activity distributions from Warholm et al. (1994), scaled to have the same mean value as the overall mean estimate of the

population mean obtained by David et al. (2006): $0.852 \text{ kg}^{0.3}/\text{hour}$. This treatment, however, fails to incorporate the uncertainty in the population mean characterized by the CV for k_{fC} in Table 4 of David et al. (2006), which is 0.711. Therefore, like $V_{\max C}$, EPA chose to use a two-dimensional sampling technique for k_{fC} . First, $k_{fC, \text{mean}}$ is sampled from a log-normal distribution with GM = $0.6944 \text{ kg}^{0.3}/\text{hour}$ and GSD = $1.896 \text{ kg}^{0.3}/\text{hour}$ (converted from the linear-space mean and CV of $0.852 \text{ kg}^{0.3}/\text{hour}$ and 0.711, respectively) with upper/lower bounds of $0.193 \text{ kg}^{0.3}/\text{hour}$ and $2.50 \text{ kg}^{0.3}/\text{hour}$, respectively (± 2 SD in log-space). After obtaining the sample population mean ($k_{fC, \text{mean}}$), an individual's genotype was sampled from the discrete incidence distribution (32% chance to be GST-T1 +/+, 48% chance to be +/-, and 20% chance to be -/-; Haber et al., 2002). Given those genotype frequencies, the interindividual variability was then characterized by rescaling the activity distributions from Warholm et al. (1994), but with the upper and lower bounds set to zero and mean + 5 SDs, respectively. David et al. (2006) also used zero for the lower bounds, but set upper bounds to mean + 3 SDs. However, in keeping with the decision to use an unbounded [log-normal] distribution for CYP2E1, the GST-T1 upper bounds were set at 5 SDs above the mean to assure characterization of the upper end of the human distribution. This rescaling and choice of bounds yields:

$$k_{fC} = \begin{cases} 0 & \text{for GST-T1 -/-,} \\ k_{fC, \text{mean}} \times N(0.8929, 0.1622 \mid 0 \leq x \leq 1.704) & \text{for GST-T1 +/-,} \\ k_{fC, \text{mean}} \times N(1.786, 0.2276 \mid 0 \leq x \leq 2.924) & \text{for GST-T1 +/+,} \end{cases}$$

where $N(\mu, \sigma \mid \text{LB} \leq x \leq \text{UB})$ is the truncated-normal distribution with mean = μ and SD = σ , bounded between LB and UB. To be clear, when the GST-T1 genotype is sampled as indicated above, the mean value for the tri-model distribution defined here for k_{fC} is $k_{fC, \text{mean}}$, the mean for the GST-T1 +/- subpopulation is one-half that for the GST-T1 +++ subpopulation, and the CV for each subpopulation is the same as used by David et al. (2006) (from the results of Warholm et al., 1994).


To assure that the age-dependence of CYP2E1 is properly characterized, particularly for children, the data of Johnsrud et al. (2003)¹⁰ was analyzed. This study measures CYP2E1 activity and other parameters from individuals up to 18 years of age. For a significant subset of the individuals in that study, values were available for the liver CYP2E1 content (activity/mg microsomal protein), liver weight, and BW. For individuals 14–18 years old there appeared no significant trend in the CYP2E1 activity, and the average BW was 69.6 kg, essentially identical to the value of 70 kg used as a representative adult. Therefore, the data from the 14- to 18-year-old individuals were assumed to represent adult values, and hence, an evaluation of the change

¹⁰Individual data supplied by the corresponding author D. Gail McCarver, to Paul Schlosser, U.S. EPA.

versus adult could be made by normalizing to these data. In particular, assuming that the microsomal content per gram of liver is constant across the age-range considered here (6 months–18 years), the total activity for an individual relative to that in an adult could be estimated as:

$$\frac{V_{\max}(\text{individual})}{V_{\max}(14-18)} = \frac{\text{CYP2E1}(\text{individual}) \cdot \text{LW}(\text{individual})}{[\text{CYP2E1}(14-18) \cdot \text{LW}(14-18)]}$$

where $V_{\max}(\text{individual})$, $\text{CYP2E1}(\text{individual})$, and $\text{LW}(\text{individual})$ are the individual's total activity, activity (per mg microsomal protein), and liver weight, respectively, while $V_{\max}(14-18)$, $\text{CYP2E1}(14-18)$, and $\text{LW}(14-18)$ are the respective average values for individuals 14–18 years old. (If not normalized, $V_{\max} = \text{CYP2E1} \cdot \text{msp} \cdot \text{LW}$, where msp is the microsomal content [mg/kg liver], but if msp is the same in all individuals, it drops from the equation when dividing by the 14–18-year-old average.) These normalized activities are plotted against the relative BW, $\text{BW}(\text{individual})/\text{BW}(14-18)$ in Figure B-2.



Source: Johnsrud et al. (2003).

Figure B-2. Total CYP2E1 activity (V_{\max}) normalized to the average total activity in 14–18 year-old individuals ($V_{\max}[14-18]$) plotted against normalized BW for individuals ranging from 6 months to 18 years of age.


The data in Figure B-2 are compared to two model predictions: the allometric-based prediction used by David et al. (2006) that V_{\max} will scale as $\text{BW}^{0.7}$, and an alternate scaling obtained by fitting to these data, $\text{BW}^{0.88}$. Both alternatives do a fairly good job of representing

the average trend in the data, but the scaling by $BW^{0.7}$ tends to under-predict the data in the range of 0.4–0.8 for $BW/BW(14-18)$. Therefore, this fitted coefficient will be used when estimating the total activity for individuals <18 years of age. However, since these data indicate a slower trend (rise in V_{max} with BW) for normalized $BW \sim 0.7$ and above, and there are no data to indicate that total activity continues to increase so rapidly in adults over 70 kg, the coefficient will be kept at 0.7 for individuals >18 years of age.

Using the allometric coefficient of 0.88, the normalized V_{maxC} values were computed in the same group of individuals as:

$$V_{maxC}(\text{individual})/V_{maxC}(14-18) = [V_{max}(\text{individual})/BW(\text{individual})^{0.88}]/[V_{max}(14-18)/BW(14-18)^{0.88}].$$

To test the revised allometric function versus the data, these values were then plotted against individual age, and a linear regression was performed, as shown in Figure B-3. While the slight trend in the regression indicates that not all of the age-dependence may be captured by this function, the low R^2 and small value of the slope indicate that the observation is not statistically significant and that further attempts to explicitly account for age dependence would lead to minimal improvement. This representation of the data also clearly shows that the overall variability in the scaled activity (V_{maxC}) is fairly constant across ages: approximately sixfold, ranging from ~0.3 to 1.8. This is the same range observed in adults by Lipscomb et al. (2003), as noted above. Thus, these data support the use of a constant variability (GSD) in simulating population variability in CYP2E1 activity for children above 6 months of age, as well as adults.



Source: Johnsrud et al. (2003).

Figure B-3. Body-weight scaled CYP2E1 activity (V_{maxC}) normalized to the average scaled activity in 14–18 year-old individuals ($V_{maxC}[14-18]$) plotted against age individuals ranging from 6 months to 18 years of age.

Note that scaling CYP2E1 activity by $BW^{0.88}$ for children and by $BW^{0.7}$ for adults, rather than per liver weight (which is expected to scale as $BW^{1.0}$ with only a 5% CV in liver fraction), leads to a lower range in CYP2E1 than scaling by BW^1 would indicate. In particular, the distribution predicted by the model for total CYP2E1 activity (mg/hour, upper/lower bound, given BW between 7 and 130 kg) will be about twofold less than if one assumed the activity varied as liver weight and simply multiplied the two sets of upper/lower bounds (BW and activity/g liver). However, given the database for this analysis from adults and children, the resulting distribution is expected to provide a good prediction of variability in the overall population.

Unfortunately, there is not a rich data set for the age-dependence of GST-T1 such as is available for CYP2E1. Strange et al. (1989) examined the developmental patterns for two other GST classes, mu and pi, and found that the child:adult activity for mu followed a similar pattern as CYP2E1, increasing from a low level near birth over time, but generally being higher than the ratio for CYP2E1 in a given age-range. GST-pi, however, was expressed at 21 times adult values in children up to 1 year of age, then declined with age, so it is difficult to draw specific conclusions regarding age-dependent variation in GST-T1 from these data (i.e., a quantitative analysis of the activity data for other GST classes will not be used to estimate the variation in GST-T1).

As is, the model uses a first-order rate constant for GST-T1 which is scaled as $BW^{-0.3}$. While the more recent “standard” for scaling of first-order constants is as $BW^{-0.25}$, the difference between these two is small and $BW^{-0.3}$ was used in the Bayesian model calibration. Because first-order constants are multiplied by tissue volumes in calculating total metabolism rates, and tissue volume scales approximately as BW^1 , scaling the GST-T1 constant by $BW^{-0.3}$ is equivalent to scaling a V_{\max} by $BW^{0.7}$. Even though the results cited above for GST-pi have the opposite trend, data on CYP2E1 activity discussed above and the trend in GST-mu activity are both at least qualitatively consistent with this scaling. Therefore this scaling is assumed to appropriately account for age-dependent changes in GST-T1 activity for individuals over 6 months of age via the explicit dependence on BW.

B.4. ANALYSIS OF HUMAN PHYSIOLOGICAL DISTRIBUTIONS FOR PBPK MODELING

While the BW distribution in the David et al. (2006) PBPK model used ranges from 7 to 130 kg, thus covering 6-month-old children to obese adults, there are age-dependent changes and gender-dependent differences in ventilation rates and body fat that are not explicitly included. To more accurately reflect the distribution of physiological parameters in the entire population, the unstructured distributions of David et al. (2006) for certain primary or key parameters were replaced with distributions based on available information that specifically accounts for the population distributions of age and gender and the age- and gender- specific distributions or functions for BW, QCC, alveolar ventilation, body fat (fraction), and liver fraction. In the following, $v \sim U[0,1]$ indicates that v is a random sample from the uniform distribution from 0 to 1.

B.4.1. Age

U.S. Census Bureau statistics¹¹ for 6 months to 80 years of age were normalized (population for 6 months to 1 year assumed to be one-half of 0- to 1-year population), and the resulting quantiles were plotted against the corresponding ages with a polynomial function fit, as shown in Figure B-4. A sample individual’s age can be determined by using the polynomial, given $v \sim U[0,1]$. (Alternately, age can be specified.)

¹¹ Available at <http://factfinder.census.gov/servlet/DatasetMainPageServlet>; use "enter a table number" or "list all tables" to select tables QT-P1 and QT-P2 for the (entire) United States.




Figure B-4. U.S. age distribution, 6 months to 80 years (values from U.S. Census Bureau).

B.4.2. Gender

U.S. Census Bureau statistics for fraction of males versus age in 5-year intervals were plotted and an empirical function was fit, as shown in Figure B-5. Given the individual's age, the gender is randomly selected as male if $v \sim U[0,1]$ is less than or equal to the polynomial and, otherwise female. (Alternately, it can be specified.)




Figure B-5. U.S. age-specific gender distribution (values from U.S. Census Bureau).

B.4.3. BW

Portier et al. (2007) reported statistical mean and SD for BW as a function of age and gender based on the NHANES IV. Portier et al. (2007) also described empirical functions fit to these results; however, those functions were found to be exquisitely sensitive to the fitted parameters to the point that entering the functions by using the four significant figures for parameters given by Portier et al. (2007) gave results that significantly deviated from the results as shown by the authors in plots at higher age ranges. Therefore, a somewhat different functional form was chosen. In particular for BW mean and SD for each gender, a function of the form was fitted:

$$\text{value} = \begin{cases} \exp[\text{poly}_1((\text{age}_c - \text{age})/10)], & \text{age} \leq \text{age}_c \\ \exp[\text{poly}_2((\text{age}_c - \text{age})/10)], & \text{age} \geq \text{age}_c \end{cases}$$

where $\text{poly}_i(x) = a \cdot x + b_i \cdot x^2 + c_i \cdot x^3$, $i = 1$ and 2 , age is in years, age_c is a “cut” age, dividing the early-age function from the later age function, and poly_i ($i = 1$ or 2). Because there is no explicit constant term in poly_i and the linear coefficient, a , is common to the two functions, the functions will automatically satisfy the condition of being equal and of having equal slope (first derivative) at $\text{age} = \text{age}_c$, so the overall function will be smooth and continuous.

The functions were fitted to the summarized data based on minimizing a weighted sum of square errors, $\text{error} = \sum n_{\text{age}} \times [\text{data}(\text{age}) - \text{function}(\text{age})]^2$, where n_{age} was the number of observations for the age. The resulting parameter values are listed in Table B-2, and the curve fits are shown in Figure B-6. Given the age and gender from Sections B.4.1 and B.4.2, the BW is then randomly selected from a normal distribution, with the resulting mean and SD truncated at the 1st and 99th percentile.

Table B-2. Parameters for BW distributions as functions of age and gender

Parameter	Male BW (kg)		Female BW (kg)	
	Polynomial parameters for mean ^a	Polynomial parameters for SD ^a	Polynomial parameters for mean ^a	Polynomial parameters for SD ^a
age_c	21	16	16	13
a	4.406	2.87	4.146	2.574
b_1	-0.0285	0.06	-0.147	-0.358
c_1	-0.729	-2.56	-1.36	-2.55
b_2	0.115	0.96	0.44	1.16
b_2	0.0048	0.0448	-0.0278	-0.0861

$${}^a\text{Mean or SD} = \begin{cases} \exp[\text{poly}_1((age_c - age)/10)], & age \leq age_c \\ \exp[\text{poly}_2((age_c - age)/10)], & age \geq age_c \end{cases}$$

where $\text{poly}_i(x) = a \cdot x + b_i \cdot x^2 + c_i \cdot x^3$, $i = 1$ and 2




Figure B-6. Function fits to age-dependent data for BW mean and SDs for males and females in the United States (values from Portier et al. [2007]).

An example output BW distribution, from a Monte Carlo simulation for ages 0.5–80 years, both genders, is shown in Figure B-7. The range, 6.6–131.4 kg, is only slightly larger than that set by David et al. (2006) (i.e., 7–130 kg). But the bimodal form is an unexpected but reproducible result, presumably occurring because the fraction of a given life span spent at intermediate BW values is smaller, as evidenced by the most rapid growth rate occurring between ~7 and 18 years of age (Figure B-6).




Figure B-7. Example BW histogram from Monte Carlo simulation for 0.5- to 80-year-old males and females in the United States (simulated n = 10,000).

B.4.4. Alveolar Ventilation

Clewell et al. (2004) tabulated values for the alveolar ventilation constant QAlvC (L/hour·kg^{0.75}) for males and females at different ages (QAlvC is multiplied by BW^{0.75} to obtain the total rate). Smooth functions of age were fitted to those results to use as age- and gender-specific mean values, shown in Figure B-8. Arcus-Arth and Blaisdell (2007) reported GSD values for respiration rates for 0–18 years of age; a smooth function was fit to those results and the value at 18 years was assumed to apply for all adults greater than age 18 (Figure B-9). An individual's QAlvC was then selected from a log-normal distribution with the resulting mean and GSD (given age and gender) truncated between the 5th and 95th percentile.




Figure B-8. Mean value respiration rates for males and females as a function of age (values from Clewell et al. [2004]).




Figure B-9. GSDs for respiration rates for males and females as a function of age (values from Arcus-Arth and Blaisdell [2007]).

B.4.5. QCC

Clewell et al. (2004) provide a function that will be taken to represent the mean for the QCC, $QCC_{\text{mean}} = 56.906 \times (1.0 - e^{-0.681 \times \exp[0.0454 \times QAlvC]}) - 29.747$. However, using this function alone will tightly link cardiac flow and ventilation rates, rather than using a distribution in the VPR ($VPR = QAlv/QC = QAlvC/QCC$) as was done by David et al. (2006). Since a distribution is already defined for QAlvC, above, a VPR subject to variability will be estimated but renormalized to match the ratio of QAlvC and QCC_{mean} as defined above. In effect, this means $QCC_{\text{sample}} = QCC_{\text{mean}} \times VPR_{\text{mean}}/VPR_{\text{sample}}$ will be chosen, where VPR_{mean} and VPR_{sample} are the mean and a random sample from the distribution defined by David et al. (2006). Clewell et al.

(2004) suggest scaling QCC and alveolar ventilation as $BW^{0.75}$, while David et al. (2006) used $BW^{0.74}$. While virtually identical, it is noted that the implementation here uses $BW^{0.75}$.

B.4.6. Fat Fraction

Tabulated values from Clewell et al. (2004) show indistinguishable values for the fraction of BW as fat (VFC) for males and females up to 7 years of age, after which they diverge. Polynomial functions were fit separately for 0–7 years, 7–20 years, and 20–80 years, with the latter two ranges being gender-specific, as shown in Figure B-10. The ratio of the resulting gender- and age-specific value to the VFC_{mean} from David et al. (2006) was then used to scale the bounded normal distribution as specified by David et al. (2006) for the selected age and gender.




Figure B-10. Fraction body fat (VFC) over various age ranges in males and females (data from Clewell et al. [2004]).

B.4.7. Liver Fraction

Tabulated values from Clewell et al. (2004) showed an interesting age dependence for the liver fraction of BW, VLC, as shown in Figure B-11. While female VLC values were somewhat higher than males between 18 and 40 years, and the difference may be statistically significant, the overall variation is not large; the two were indistinguishable during other age-ranges, and model predictions are not overly sensitive to VLC. Therefore, only the male data were used. Smooth functions were fit separately for 0–18 years and 18–80 years and used to scale the distribution from David et al. (2006) for the given age, as was done for body fat.




Figure B-11. Fraction liver (VLC) as a function of age (data from Clewell et al. [2004]).

B.4.8. Tissue Volume Normalization

While not explicitly stated by David et al. (2006), total tissue volume must remain roughly the same as a fraction of BW. While this fraction could also change with age, gender, and other characteristics, it was assumed that any change in it would be modest and not significantly affect model predictions, given the fairly broad distribution implemented for total BW. This normalization was applied irrespective of those factors. Therefore, after drawing sample values of the tissue volume fractions from each of their respective distributions, the fractions were then normalized to a total fraction of 0.9215, which is the sum of the mean values for the fractions for the distributions as described by David et al. (2006). The remaining body mass is taken to be bone, teeth, hair, nails, and any other minimally or nonperfused components.

B.5. SUMMARY OF REVISED HUMAN PBPK MODEL

The resulting set of parameter distribution characteristics, including those used as defined by David et al. (2006), are described in Table B-3. The metabolic parameter statistics reported in Table 4 of David et al. (2006) are summary statistics of the converged parameter chains obtained in that analysis for the population mean of each parameter. As such, those statistics (means and

CVs) are assumed to represent the most likely value of the mean and the degree of uncertainty in that mean. However, for the metabolic parameters other than $V_{\max C}$ (representing CYP2E1 activity) and k_{fC} (GST-T1 activity), EPA considers it reasonable to assume negligible variability among the population compared to the estimated uncertainties. So while EPA's objective is to account for both variability and parameter uncertainty in the population, the statistics for those other parameters (K_m , A1, A2, and FracR) were used as is to define population distributions. For $V_{\max C}$ and k_{fC} , two-dimensional sampling routines were used, as described in detail above, to explicitly account for both uncertainty and the known high degree of interindividual variability. Distributions for a number of the physiological parameters, which are assumed to represent a well known degree of variability, were also revised from those used by David et al. (2006).

Table B-3. Parameter distributions for the human PBPK model for dichloromethane used by EPA

Parameter		Distribution					Section or source
		Shape	(Geometric mean ^a)	SD/GSD ^a	Lower bound	Upper bound	
BW	Body weight (kg)	Normal	$f(\text{age, gender})$		1 st %tile	99 th %tile	B-4.3; NHANES IV
Flow rates							
QAlvC	Alveolar ventilation (L/hr/kg ^{0.75})	Normal	$f(\text{age, gender})$	$f(\text{age})$	5 th %tile	95 th %tile	B-4.4; mean: Clewell et al. (2004); SD: Arcus-Arth and Blaisdell (2007)
vprv	Variability in ventilation/perfusion ratio	Log-normal	1.00	0.203	0.69	1.42	VPR/VPR _{mean} of David et al. (2006)
QCC	Cardiac output (L/hr/kg ^{0.75})	QCC _{mean} = $f(\text{QAlvC})$		QCC = QCC _{mean} /vprv			B-4.5; Clewell et al. (2004) (mean)
Fractional flow rates (fraction of cardiac output)							
QFC	Fat	Normal	0.05	0.0150	0.0050	0.0950	David et al. (2006); <i>after sampling from these distributions, normalize:</i> $Q_i = \frac{Q_C \cdot Q_i C}{\sum Q_j C}$
QLC	Liver	Normal	0.26	0.0910	0.010	0.533	
QRC	Rapidly perfused tissues	Normal	0.50	0.10	0.20	0.80	
QSC	Slowly perfused tissues	Normal	0.19	0.0285	0.105	0.276	
Tissue volumes (fraction BW)							
VFC	Fat	Normal	$f(\text{age, gender})$	0.3·mean	0.1·mean	1.9·mean	Fat mean: B-4.6; (Clewell et al., 2004); Liver mean: B-4.7; (Clewell et al., 2004); otherwise David et al. (2006); <i>after sampling from these distributions, normalize:</i> $V_i = \frac{0.9215 \cdot BW \cdot V_i C}{\sum V_j C}$
VLC	Liver	Normal	$f(\text{age})$	0.05·mean	0.85·mean	1.15·mean	
VLuC	Lung	Normal	0.0115	0.00161	0.00667	0.0163	
VRC	Rapidly perfused tissues	Normal	0.064	0.00640	0.0448	0.0832	
VSC	Slowly perfused tissues	Normal	0.63	0.189	0.431	0.829	
Partition coefficients							
PB	Blood/air	Log-normal	9.7	1.1	7.16	13.0	GM & GSD values listed here, converted from arithmetic mean and SD values of David et al. (2006)
PF	Fat/blood	Log-normal	11.9	1.34	4.92	28.7	
PL, PLu, & PR	Liver/blood, lung/arterial blood, and rapidly perfused tissue/blood	Log-normal	1.43	1.22	0.790	2.59	
PS	Slowly perfused tissue (muscle)/blood	Log-normal	0.80	1.22	0.444	1.46	

Table B-3. Parameter distributions for the human PBPK model for dichloromethane used by EPA

Parameter	Distribution					Section or source	
	Shape	(Geometric) mean ^a	SD/GSD ^a	Lower bound	Upper bound		
Metabolism parameters (based on Monte Carte calibration from five human data sets)							
$V_{\max C, \text{mean}} / V_{\max C}$	Population mean / individual maximum metabolism rate (mg/hr/kg ^{X_{vmax}})	Log-normal	9.34	1.14	7.20	12.11	B-3; mean: David et al. (2006); Individual GSD: Lipscomb et al. (2003); X _{vmax} = 0.88 for age <18. X _{vmax} = 0.70 for age ≥18.
		Log-normal	$V_{\max C, \text{mean}}$	1.73	(none)	(none)	
K_m	Affinity (mg/L)	Log-normal	0.41	1.39	0.154	1.10	
A1	Ratio of lung V_{\max} to liver V_{\max}	Log-normal	0.00092	1.47	0.000291	0.00292	
A2	Ratio of lung KF to liver KF	Log-normal	0.0083	1.92	0.00116	0.0580	GM & GSD values listed here, converted from arithmetic mean and SD values of David et al. (2006)
FracR	Fractional MFO capacity in rapidly perfused tissue	Log-normal	0.0152	2.0	0.00190	0.122	
First order metabolism rate (/hr/kg ^{0.3})							
$k_{fC, \text{mean}}$	Population average	Log-normal	0.6944	1.896	0.1932	2.496	Adapted from David et al. (2006); $k_{fC, \text{mean}}$ is first sampled, then the relative individual value, $k_{fC}/k_{fC, \text{mean}}$, given the genotype; k_{fC} is then the product.
$k_{fC} / k_{fC, \text{mean}}$	Homozygous (-/-)	N/A	$k_{fC} = 0$	-	-	-	
	Heterozygous (+/-)	Normal	0.8929	0.1622	0	1.704	
	Homozygous (+/+)	Normal	1.786	0.2276	0	2.924	

^aArithmetic mean and SD listed for normal distributions; GM and GSD listed for log-normal distributions.

The Monte-Carlo sampling approach used effectively assumes that all the parameters are distributed independently, ignoring the covariance that was likely represented in the actual posterior chains. This approach will tend to overestimate the overall range of parametric variability and uncertainty and, hence, distribution of dose metrics in the population compared to what one would obtain if the covariance were explicitly included. Thus, if the covariance (i.e., the variance-covariance matrix) for the set of parameters had been reported by David et al. (2006) it could have been used to narrow the predicted distribution of internal doses or equivalent applied doses, but lacking such information, the approach used will not underestimate risk or overestimate lower bounds on human equivalent exposure levels. However, this source of overestimation in variability and uncertainty is probably offset to some extent by the fact that the analysis leaves out the degree of interindividual variability for K_m , A1, A2, and FracR that should have also been estimated by the Bayesian analysis.

APPENDIX C. RAT DICHLOROMETHANE PBPK MODELS

The critical studies and data chosen for derivation of a recommended RfD and RfC were based on nonneoplastic liver lesions in rats inhaling dichloromethane for 2 years (Nitschke et al., 1988a; Serota et al., 1986a). For this reason, a PBPK model for inhaled and orally absorbed dichloromethane in rats was needed to provide estimates of internal dosimetry for dose-response modeling and to extrapolate internal liver doses from rats to humans. Several deterministic PBPK rat models have been reported in the scientific literature (Sweeney et al., 2004; Andersen et al., 1991; Reitz, 1991; Reitz et al., 1988a, b; EPA 1988b, 1987a, b; Andersen et al., 1987; Gargas et al., 1986). Unlike in the mouse study (Marino et al., 2006), however, no probabilistic models are available in which the uncertainty in model parameters was reduced by utilizing multiple data sets for parameter estimation. Rat data were not available that would allow for MCMC calibration of individual metabolic parameters for the CYP or GST pathways. For example, the MCMC calibration of the mouse model (Marino et al., 2006) relied on dichloroethylene inhalation data (not available for the rat) to specifically estimate GST metabolism in isolation of CYP metabolism, thereby improving the estimate of metabolic flux through the competing pathways. Thus, the selected model includes parameter values estimated by deterministic methods only. In order to use the latest data for dichloromethane toxicokinetics in rats, an assessment was conducted of multiple rat models (or modified versions of those models) to select the most appropriate model for use in the derivation of the RfD and RfC.

C.1. METHODS OF ANALYSIS


C.1.1. Selection of Evaluation Data Sets and PBPK Models

Published studies of dichloromethane metabolism and toxicokinetics in rats were reviewed to identify data sets for use in model evaluation and possible calibration. Toxicokinetic data were available for:

- blood levels of dichloromethane, the percent saturation of hemoglobin as COHb, and expired dichloromethane and CO following intravenous injection (Angelo et al., 1986b);
- dichloromethane air concentrations in closed chamber experiments (Gargas et al., 1986);
- dichloromethane and %COHb blood levels during and after a 4-hour open chamber (constant concentration) inhalation exposure (Andersen et. al., 1991, 1987);
- %COHb levels from 30 minutes to 12 hours postexposure following a single oral dose of 526 mg/kg (6.2 mmol/kg in Oleum pedum tauri vehicle) (Pankow et al., 1991a); and

- Cumulative dichloromethane (mg) expired up to 96 hours following gavage doses of 250, 500, 1,000, or 2,000 mg/kg in corn oil or water (Kirschman et al., 1986).

Four variations on the PBPK models of Andersen et al. (1991, 1987) (Figure C-1) were assessed for the ability to predict these data. In each model, metabolism involves two competing pathways: the GST pathway, described with a linear first-order kinetic model, and the CYP pathway, described with a saturable Michaelis-Menten kinetic model.



Models A, C, and D also included metabolism in the lung tissue compartment via GST and CYP.

Figure C-1. Schematic of the Andersen et al. (1991) PBPK model (model B) for dichloromethane in the rat.

Model A is a hybrid of Andersen et al. (1991) and Andersen et al. (1987) in that it included both CO production resulting from CYP-mediated metabolism (Andersen et al., 1987) and lung metabolism of the parent compound (Andersen et al., 1991). Andersen et al. (1987) based the lung-to-liver metabolism ratios (A1 and A2 for CYP- and GST-mediated metabolism, respectively) on the reaction rates reported by Lorenz et al. (1984). However, for model A the Bayesian-calibrated values of A1 and A2 were used for mice (Marino et al., 2006), which were identified using in vivo data that distinguished CYP and GST metabolism. Specifically, EPA sought to determine if use of these mouse-calibrated ratios would improve the model fit to the rat data. With A1 and A2 fixed, the model was fit to data by adjusting the parameters for CYP and GST metabolism in the liver: $V_{\max C}$ (allometrically-scaled maximum rate of CYP metabolism),

K_m (CYP metabolic saturation constant), and k_{fC} (allometrically-scaled first-order GST rate constant).

Model B is the model of Andersen et al. (1991) using the same published values for V_{maxC} , K_m , and k_{fC} , but lung metabolism is nullified.

Model C is the model of Andersen et al. (1991) with lung metabolism defined as 4 and 14% of CYP- and GST-mediated liver metabolite production, respectively. The lung-to-liver metabolic ratios (A1 [CYP] and A2 [GST]) were estimated from in vitro metabolic activity measured in lung and liver cytosolic (GST) and microsomal (CYP) tissue fractions (Reitz et al., 1989). V_{maxC} , K_m , and k_{fC} were unchanged from Andersen et al. (1991).

Model D is Model C with V_{maxC} , K_m , and k_{fC} re-optimized against: the inhalation gas uptake data of Gargas et al. (1986; closed-chamber gas-uptake data); the inhalation data of Andersen et al. (1987; blood dichloromethane levels); and the 24-hour cumulative exhalation data of Angelo et al. (1986b; exhaled breath dichloromethane and CO levels, oral exposures). The latter data, which were insensitive to the oral absorption constant (because absorption and excretion are essentially complete by 24 hours), were used to inform the balance between GST-mediated and CYP-mediated metabolism.

C.1.2. Analysis

Models A–D were implemented using the acslXtreme simulation software (version 2.4, The Aegis Technologies Group, Huntsville, AL). Parameter estimation was performed using the Nelder-Mead algorithm as implemented in the acslXtreme package.

After completing the calibration for model D as described above, the first-order oral absorption constant, k_a , was then estimated by optimizing it, with all other parameters held constant, to the oral pharmacokinetic data of Angelo et al. (1986b): blood and liver dichloromethane levels, expired dichloromethane levels (all time-points), and expired CO levels (24-hour cumulative). Only the 24-hour cumulative CO exhalation data were used to inform the estimate k_a (earlier time-points not used) for reasons explained below. Two other data sets of gavage exposures (Pankow et al., 1991a; Kirschman et al., 1986) were also used in the evaluation of k_a .

In order to obtain reasonable fits, the assumed error in observations of blood and liver dichloromethane was modeled as absolute (e.g., unrelated to the magnitude of the response variable). If the assumed error for these variables was modeled as relative, then the fit to a few low blood concentration data points disproportionately impacted the outcome, resulting in poor fits to the higher blood concentration data. The parameter values for each of the four models are given in Table C-1. Differences between observed values and predictions of uptake, blood and liver concentrations, and expiration of dichloromethane from Models A–D were assessed visually.

Table C-1. Parameter values used in rat PBPK models

Parameter	Model A	Model B	Model C	Model D
Flow rates				
QCC (L/hr·kg ^{0.74})	15.9	Same as Model A		
VPR	0.94			
Fractional flow rates (percent of QCC)				
Fat	9	Same as Model A		
Liver	20			
Rapidly perfused tissues	56			
Slowly perfused tissues	15			
Tissue volumes (percent BW)				
Fat	7	Same as Model A		
Liver	4			
Lung (scaled as BW ^{0.99})	1.15			
Rapidly perfused tissues	5			
Slowly perfused tissues	75			
Partition coefficients				
Blood:air	19.4	Same as Model A		
Fat:blood	6.19			
Liver/blood	0.732			
Lung/arterial blood	0.46			
Rapidly perfused tissue/blood	0.732			
Slowly perfused tissue (muscle)/blood	0.408			
Metabolism and absorption parameters				
Maximum metabolism rate, V _{maxC} (mg/hr·kg ^{0.7})	6.21	4.0	4.0	3.93
Affinity, K _m (mg/L)	0.23	0.4	0.4	0.524
Ratio of lung V _{max} to liver V _{max} , A1	0.21	Not applicable	0.04	0.04
Ratio of lung KF to liver KF, A2	0.20	Not applicable	0.14	0.14
1 st order metabolism rate (liver), k _{FC} (1/hr·kg ^{0.3})	2.89	2.0	2.0	2.46
First-order oral absorption rate constant, k _a (1/hr)	5.0	5.0	4.0	1.80

C.2. RESULTS


Four versions of the PBPK model for dichloromethane (Models A–D) were evaluated for goodness of fit to a common set of pharmacokinetic data. The k_a was then calibrated using oral pharmacokinetic data for the model that was deemed best.

C.2.1. Evaluation of Model Structure for Description of Carboxyhemoglobin Levels

The model of Andersen et al. (1991) calculated the amount of COHb by assuming instantaneous equilibration between the free CO and hemoglobin-bound CO in the blood. However, when Angelo et al. (1986b) observed the kinetics of dichloromethane in blood and exhaled dichloromethane and CO after oral administration, they found that blood dichloromethane peaked rapidly in <30 minutes. The rate of exhalation of dichloromethane,

shown by the rate of rise in the exhaled dichloromethane data in Figure C-2A where the plateau in exhaled dichloromethane levels from 6 to 24 hours, indicates that blood dichloromethane levels had dropped to nearly 0 by 6 hours. Exhaled CO, however, shows a more gradual rise than dichloromethane, continuing between 6 and 24 hours. Pankow et al. (1991a) measured COHb in blood after dichloromethane exposure, albeit at a higher dose (526 mg/kg versus 200 mg/kg by Angelo et al. [1986b]), and found that blood COHb levels did not peak until 6 hours after dosing (Figure C-2B), while Angelo et al. (1986b) found that blood dichloromethane had declined to ~4% of the peak level (10-minute concentration) by 6 hours. Some delay between peak dichloromethane and peak (cumulative) CO levels is predicted by the published (1991) and modified (Models A and D) Andersen et al. (1991) models because CO is produced from oxidative dichloromethane metabolism. However, we found that none of the PBPK models could simultaneously describe both the very short-time peak in blood dichloromethane with the concurrent rapid rise in exhaled dichloromethane levels (Figure C-2A) and the late peak blood CO (Figure C-2B) and slower rate of CO exhalation (Figure C-2A). The final models were able to adequately predict dichloromethane exhalation data and dichloromethane blood concentration data but overestimated CO exhalation data (these findings are illustrated for Model D later in the results section).

EPA hypothesized that the models' inability to simultaneously fit these data might specifically be due to the assumption of instantaneous equilibrium between free CO and COHb in blood. The model structure (set of equations) characterizes the kinetics of free CO and COHb as being at instantaneous equilibrium with one another so that the kinetics of COHb are completely determined by the predicted rate of appearance and elimination of CO. The rate of production of CO is in turn determined by the rate of metabolism of dichloromethane and the rate of elimination of CO is determined by physiological parameters. Therefore, EPA considered that the only way to alter the predicted time-course of COHb with this model structure and these assumptions is to alter the rate of dichloromethane metabolism, which would degrade the fit to the dichloromethane data, or to alter physiological parameters which are well established (based on direct measurements). If this hypothesis is correct, or there is some other model-structure error involved, then adjusting physiological parameters in an attempt to fit the CO exhalation would only cover up the error and reduce the model's reliability in predicting other aspects of dosimetry, particularly the pharmacokinetics of dichloromethane and its rates of metabolism which are critical to the assessment. In summary, it appears that given the model structure and assumptions, one cannot explain the difference between the exhalation time-courses for dichloromethane and CO/COHb shown in Figure C-2 while retaining realistic physiological parameters; that difference might be explained if the release of CO from COHb is treated as rate-limiting, in which case the two would not be in rapid equilibration.



Points are data collected on 1st, 7th, or 14th days of exposure. Lines are plotted through average of days 1, 7, and 14 data. (Arrows indicate corresponding y-axis).

Figure C-2. A: Observations of exhaled [¹⁴C]-labelled dichloromethane (left y-axis) and CO (right y-axis) after a bolus oral dose of 200 mg/kg [¹⁴C]-dichloromethane in rats (data of Angelo et al., 1986b). B: Blood COHb (percent of total hemoglobin) from a single gavage dose of 526 mg/kg dichloromethane in rats.

While the analysis of the CO and COHb data above led us to the conclusion that the purpose of the model for chronic-exposure risk assessment was best served by not using some of those data (i.e., the short-term dichloromethane and CO exhalation data), the combination of dichloromethane concentration data in various compartments as described below (primarily blood and air) and the more limited long-term CO exhalation data were considered adequate and necessary to estimate model parameters for dichloromethane. Some comparisons of model

predictions to the CO/COHb data are shown, however, to both further illustrate the discrepancy in kinetics and the agreement in mass balance. If the model was to be used for evaluation of the risk of acute carbon-monoxide exposure, then it would be necessary to demonstrate adequate reproduction of the COHb data in particular.

C.2.2. Evaluation of Prediction of Uptake, Blood and Liver Concentrations, and Expiration of Dichloromethane

One data set used for model calibration is the closed chamber experiments of Gargas et al. (1986) (100–3,000 ppm dichloromethane). Because actual chamber concentrations can differ from target concentrations, a simple exponential function was fit to the first four data points at each concentration and used to extrapolate back to an actual initial concentration. The initial concentrations so obtained and used in the modeling were 107, 498, 1,028, and 3,206 ppm versus the target concentrations of 100, 500, 1,000, and 3,000 ppm. Predictions from models B, C, and D all fit the observed chamber air concentrations reasonably well and noticeably better than model A, but model D is clearly superior to model C and is slightly but noticeably better than model B (Figure C-3; comparing model B and D fits to the 1,000 and 3,000 ppm data). Model A did not simulate the observed break between linear metabolism at 100 ppm and partially saturated metabolism at 300 ppm, ostensibly because of the higher value for maximal rate of CYP metabolism, $V_{\max C}$ (6.21 mg/hour/kg^{0.7} in model A), which prevented saturation of oxidative metabolism at these exposure levels. Use of $V_{\max C} \sim 4$ mg/hour/kg^{0.7} in models B, C, and D (Table C-1) provides a much better fit to the metabolic saturation pattern across all exposure levels.





Figure C-3. Observations of Gargas et al. (1986; data) and predictions (models A–D) for respiratory uptake by three rats of 100–3,000 ppm dichloromethane in a 9-L closed chamber.

Model simulations were next compared to measurements of blood dichloromethane concentration after intravenous injections of 10 or 50 mg/kg (Angelo et al., 1986b). Intravenous data are considered particularly useful for evaluating metabolic rate constants because the kinetics and complexity of uptake by other routes of exposure have been bypassed. All four models matched the observed blood dichloromethane levels fairly well following intravenous injections of 10 or 50 mg/kg (Figure C-4): simulations are within a factor of 3 of the data (often closer) and the slope of the model curve after the initial distribution phase (~5 minutes) appears close to that of the data. However, each model over-predicted the observed data to some extent; the 50 mg/kg dose over-predicts all of the observed data points, whereas the 10 mg/kg simulation matches the first three data points well. Because the y-axis in Figure C-4 is plotted on a log scale, the discrepancy between model simulations and the 50 mg/kg data appears similar to the discrepancy vs. the 10 mg/kg data at later time points, but in fact is much larger in an absolute sense.




Legend for models B and C is the same as for model A. Data are for dichloromethane in blood following 10 and 50 mg/kg intravenous injection in rats. Model predictions with doses at 5 and 25 mg/kg, are shown for comparison as thinner lines for model D.

Figure C-4. Observations (points) of Angelo et al. (1986b) and predictions (curves, denoted “sim” in legend) for models A–D.

The most likely explanation for the over-prediction of all four models vs. the experimental observations in Figure C-4 is some shortcoming in model structure or parameter specification (e.g., the model lacks an explicit blood-volume compartment, which could significantly impact predictions for this particular route of administration). The discrepancy appears to exist primarily in the initial distribution phase, as the slope of the simulated clearance curve closely matches that of the data beyond 10 minutes (0.17 hours). The error was further evaluated by running simulations with doses resulting in blood dichloromethane levels that more closely matched the data at the end of the initial distribution phase. These simulations can indicate whether the error is just in the initial distribution phase (primarily impacted by the tissue-compartment volumes and partition coefficients) or is also in the rate of clearance by

metabolism and other mechanisms (which determine the slope of the curve at later times). When the simulated administered doses were decreased 50% (i.e., to 5 and 25 mg/kg), the model predictions under-predicted the early time-point data but closely matched the data from 10 minutes on, as shown in the model D panel of Figure C-4 (thin simulation curves). This result gives further confidence in the model's ability to predict dosimetry over longer periods of time, if not immediately following intravenous exposures, which is appropriate since we are concerned with correctly predicting dosimetry from chronic exposures.


Predictions of percent dichloromethane dose expired as dichloromethane were also similar between all models simulating intravenous injections of 10 or 50 mg/kg (Figure C-5) but with notable differences. Model D predictions for the 10 and 50 mg/kg exposure were higher than both the data and the predictions of the other models, but all models over-predict the amount exhaled from the 50 mg/kg dose. As with the simulations of blood concentrations (Figure C-4), the fit of model D to the percent dichloromethane exhaled data is significantly improved if the simulated doses are reduced by 50% (model D panel of Figure C-5), suggesting that the primary error is in the model's ability to describe the initial distribution rather than longer-term clearance.



Legend for models B and C is the same as for model A. Data are percent of dichloromethane dose expired as dichloromethane. Model predictions with doses at 5 and 25 mg/kg, are shown for comparison as thinner lines for model D.

Figure C-5. Observations of Angelo et al. (1986b) and predictions (curves, denoted “sim” in legend) for models A–D following 10 and 50 mg/kg intravenous injection in rats.

Simulations were then run to match the open-chamber inhalation data of Andersen et al. (1987), as shown in Figure C-6. Despite reasonably good fits for dichloromethane blood concentrations during 4-hour inhalation phases of exposure to 200 or 1,000 ppm, none of the models fit the postexposure clearance phase of the inhalation-exposure blood dichloromethane data very well, especially the 1,000 ppm data (Figure C-6). Model C describes the 200 ppm data quite well, but the improvement versus models B and D is not sufficient to offset the lack-of-fit by model C to the closed-chamber uptake data. Models B, C, and D all over-predict the blood concentration for 200 ppm exposure during the 4-hour exposure period.



Model prediction for a 150 ppm exposure (thin dashed line) also shown for comparison.

Figure C-6. Observations of Andersen et al. (1987; data points) and simulations for models A–D (curves, denoted “sim” in legend) for dichloromethane in rat blood from inhalation of 200 and 1,000 ppm dichloromethane for 4 hours.

As with the intravenous exposures where the initial distribution phase was over-predicted, EPA examined whether adjusting the uptake portion of the model simulation would also improve the model’s fit to the later data. Running a simulation with the exposure concentration set to 150 ppm (Figure C-6, model D, thin dashed lines) gives much better agreement with both the inhalation phase (time ≤ 4 hours) and clearance phase (time > 4 hours) 200 ppm data. It seems unlikely that such a large error in the exposure concentration would have occurred, but the simulation result suggests that there was some alteration in uptake conditions. For example, rodents are capable of reducing their respiration rate when exposed to irritant gases, thereby reducing the rate of inhalation. In the absence of plethysmography and animal

handling data for the Andersen et al. (1987) experiments, alternate simulations were performed with the QCC and alveolar ventilation reduced by 38% (the two rates are held in constant proportion), which provided a better fit to the data up to 4 hours (during exposure; quite similar to the result shown in Figure C-6 for model D with 150 ppm), while the clearance after 4 hours was slower than indicated by the experimental observations (not shown). It is also possible that the rats reduced respiration and QCC due to the exposure and resumed breathing at a normal rate once the exposure ended. Experiments in which respiration rates are monitored could be conducted to test this hypothetical explanation. However, none of the attempts to simulate reduced QCC and respiration during exposure, with or without a return to normal rates at 4 hours, matched the data as well as the 150 ppm exposure with default cardiac and respiration rates throughout (shown in Figure C-6).

The results with the 150 ppm simulated exposure show that if the inhalation portion of the blood dichloromethane concentration curve for the 200 ppm exposure is correctly matched, the clearance phase simulation (determined in large part by the metabolic parameters) then matches those data well with QCC and respiration at their default rates. However, since the model simulation already matches the uptake portion of the data for the 1,000 ppm exposure, it appears that the rate of metabolic (or other) clearance predicted by the model is slower than exhibited in that experiment. Thus, the model appears to over-estimate the degree of metabolic saturation (hence under-predicting metabolism) at that higher exposure level.

C.2.3. Evaluation of Relative Flux of CYP and GST Metabolism of Dichloromethane

The relative flux of dichloromethane metabolism through the CYP or GST pathways during simulated chronic, medium-, and high-level inhalation exposures of rats to dichloromethane is shown in Table C-2. Specifically, models B and D were considered as the best models with zero or nonzero lung metabolism, respectively, so the impact of this structural difference could be evaluated. Chronic exposures of 200, 1,000, 2,000, and 4,000 ppm, reflecting exposures used by Andersen et al. (1991) and NTP (1986) were simulated.

Table C-2. Effect of PBPK model configuration on predicted dichloromethane metabolite production in the liver of (male) rats from inhalation exposures^a

Model configuration	Exposure (ppm)	CYP-mediated metabolite production (mg/L liver/d)	GST-mediated metabolite production (mg/L liver/d)	Total metabolite production (mg/L liver/d)	CYP:GST metabolite production ratio
B	200	584	58	642	10
D	200	559	72	631	7.8
B	1,000	813	549	1,361	1.5
D	1,000	785	646	1,431	1.2
B	2,000	830	1,259	2,089	0.66
D	2,000	806	1,502	2,308	0.54
B	4,000	892	2,610	3,502	0.34
D	4,000	866	3,111	3,977	0.28

^aInhalation exposures of 200 or 1,000 ppm for 4 hr/d (Andersen et al., 1991) or 2,000 or 4,000 ppm dichloromethane for 6 hr/d, 5 d/wk for 2 yrs (NTP, 1986).

Only a modest difference was seen in predicted dosimetry between models B and D. Model B predicted 15–20% less GST metabolite production and 3–4% more CYP metabolite production in the liver than model D, depending on the exposure level (Table C-2). Total metabolite production was predicted to be 2% higher by model B versus model D at 200 ppm, but 12% lower by model B versus model D at 4,000 ppm (Table 2). This occurs because model D has a higher GST-rate constant (k_{TC}) than model B (2.46 versus 2.0/hour/kg^{0.3}; Table C-1), as well as lung metabolism that is absent in model B, both of which reduce the amount of dichloromethane available for CYP metabolism. For both models, CYP-mediated metabolism dominated during the 200 ppm exposure, yielding 8- to 10-fold the metabolic production of GST. However, GST-mediated metabolism dominated at 2,000 and 4,000 ppm, with CYP accounting for roughly 1/3 of total metabolite production during the 4,000 ppm exposure.

To elucidate the exposure ranges where internal doses are linear and where CYP metabolism saturates, weekly average CYP and GST liver metabolic rates were simulated for models B and D over a wide span of inhalation concentrations, given a regimen of 6 hours/day, 5 days/week, with results shown in Figure C-7. Metabolite production is predicted to be approximately linear up to ~100 ppm for both models. CYP metabolism becomes mostly saturated between 100 and 1,000 ppm but is not fully saturated even at 2,000 ppm. CYP metabolism is not predicted to be fully saturated by 2,000 ppm because exposures are only 6 hours/day, and blood concentrations fall quickly after each exposure. Since dichloromethane blood levels still increase with exposure level during that exposure-off period (because they are higher at the end of the exposure-on period), there is also increased metabolite production with exposure level during these exposure-off periods. However, the increase in metabolism per

exposure-ppm is much less in this high-concentration range than occurs in the lower-concentration range because blood concentrations are falling throughout the exposure-off period. Depending on the model, GST metabolism is predicted to become dominant above 1,100–1,300 ppm.




Figure C-7. Simulation results using models B and D for weekly average metabolic rates by the GST and CYP pathways for 6 hours/day, 5 days/week inhalation exposures.

Thus, models B and D have been shown to perform similarly in matching the inhalation and intravenous data (see above), with model D fitting the closed chamber inhalation data slightly better (Figure C-3), and in predicting liver metabolism (Table C-2). Since model D is mechanistically superior in that it explicitly describes metabolism in the lung, model D is considered the better of the two models for the purpose of predicting dichloromethane dosimetry in rats. (As discussed above, model D is also clearly superior to models A and C.)

Finally, as an absolute measure of model D's ability to predict the correct proportion of metabolism by the CYP vs. GST pathway, model D predictions of exhaled CO are compared to model predictions in Figure C-8, panel C, below. Since CO is produced by the CYP pathway and not the GST pathway, discrepancy between model predictions and these data indicates the extent to which the fraction of metabolism by each pathway is in error. (The data are from Angelo et al. [1986b] used [¹⁴C]-labeled dichloromethane and measured the amount of ¹⁴CO, so the data should have no interference from other sources of CO.) Given the observations about CO kinetics and model structure discussed in Section C.2.1, this comparison focuses on the data

at 24 hours. At 50 mg/kg, the model predicts 15.0% of the dose exhaled as CO vs. an observed mean of 16.2% (a relative error of only 7%), while at 200 mg/kg, the model predicts 5.5% exhaled as CO vs. an observed mean of 7.2% (a relative error of 23%). Model predictions for the GST pathway are 10% of the dose at 50 mg/kg and 13% of the dose at 200 mg/kg; so if one assumes that the absolute error in amount exhaled of CO comes from an over-prediction of GST metabolism, then the error in GST metabolism would be about 12% at 50 mg/kg and 13% at 200 mg/kg (i.e., GST metabolism would have to be decreased by 12–13% from its current total amount with the corresponding mass shifted to CYP to exactly match the CO data). Overall, this agreement with the data appears quite good and does not indicate a strong systematic error in the proportion of metabolism attributed to the CYP vs. GST pathways.





Figure C-8. Observations of Angelo et al. (1986b; data) and model D predictions for: (A) percent dose expired as dichloromethane; (B) blood dichloromethane; (C) percent expired as CO; and (D) liver DCM in rats given a single dichloromethane gavage dose of 50 and 200 mg/kg, using a numerically fitted k_a of 1.8 hours⁻¹ (see Section C.2.4).

Considering the overall quality of model D compared to the other three model versions evaluated and the agreement with total mass balance indicated by the comparison to the CO exhalation data, only model D was considered in the subsequent parameter estimation and evaluation of oral dosimetry.

C.2.4. Evaluation of Model Predictions of Oral Absorption of Dichloromethane

For model D, the k_a was numerically fit to the data for blood and liver dichloromethane levels and total expired dichloromethane and CO levels, as measured by Angelo et al. (1986b) for rats exposed to gavage doses of 50 and 200 mg/kg dichloromethane. Resulting fits to data are shown in Figure C-8. (Note that for the percentage expired as CO, panel C, the percentage exhaled at 200 ppm (closed circles) is lower than the percent exhaled at 50 ppm (open squares), so the percent declines as concentration increases, balancing the increased percentage exhaled as dichloromethane). The optimized value of $k_a = 1.8/\text{hour}$ resulted primarily from fitting to dichloromethane data, as it was optimized against three data sets (blood, liver, and exhaled dichloromethane) and only one CO data set. With $k_a = 1.8/\text{hour}$, model simulations fit the blood and liver dichloromethane levels adequately (Figure C-8, panels B and D), but the predicted peak in blood concentration (panel B) clearly occurs later than indicated by the data, and the predicted liver concentrations (panel D) are considerably higher than observed. The model fit to the observed dichloromethane expiration data is excellent (panel A), and model fit to the percent expired as CO is also very good (panel C).

Pankow et al. (1991a) measured blood COHb following a gavage dose of 525 mg/kg, data which might also be used to estimate k_a . When k_a was re-optimized against those data (other parameters as in model D), the resulting value was 0.47/hour. Model simulations with both $k_a = 0.47/\text{hour}$ and the previously-estimated value of 1.8/hour are compared to the COHb data of Pankow et al. (1991a) in Figure C-9. While the lower value of k_a (0.47/hour) provides a better fit to the COHb data than simulations with $k_a = 1.8/\text{hour}$, the model still does not capture the more gradual rise to a peak and subsequent decline in the data. Moreover, with $k_a = 0.47/\text{hour}$, the peak blood and liver concentrations of dichloromethane after gavage are predicted to occur at just beyond 1 hour after exposure (not shown), which is completely inconsistent with the data of Angelo et al. (1986b). Therefore, as discussed previously, the inability to describe the kinetics of COHb (with $k_a = 1.8/\text{hour}$) may be due to the assumption of a rapid equilibrium between free and hemoglobin-bound CO in the blood: the more gradual rise and fall shown by the data in Figure C-9 could be better predicted if the binding and release steps were treated as rate-limiting. The difference in kinetics might also occur because the CO model does not include body/tissue compartments other than blood and so does not account for diffusion (of free CO) into and out of the rest of the body.



Model simulations performed with model D (heavy line, $k_a = 1.8/\text{hour}$) or with an alternate value of $k_a = 0.47/\text{hour}$, fit to these data.

Figure C-9. Model predictions of blood COHb (percent of total hemoglobin) from a single gavage dose of 526 mg/kg dichloromethane in rats, compared to the data of Pankow et al. (1991a).

Another factor in attempting to simulate the CO data along with other data sets is the difference in vehicle: Pankow et al. (1991a) used *Oleum pedum tauri* while Angelo et al. (1986b) used water with PEG. The kinetics of orally administered dichloromethane with the water/PEG vehicle are shown in Figure C-8, with the blood concentration already clearly falling from the peak at 30 minutes (panel B) and most of the dichloromethane exhalation being complete by 4 hours (panel A). If such dichloromethane kinetics had occurred in the experiments of Pankow et al. (1991a), it seems unlikely that COHb levels would continue to rise between 4 and 6 hours. In short, the late peak of COHb observed by Pankow et al. (1991a) strongly indicates that absorption was slower (a lower value of k_a applies) and, hence, the dichloromethane peak was also later in those experiments than observed by Angelo et al. (1986b); such a shift might be due to the use of a more lipophilic vehicle by Pankow et al. (1991a). But the failure to describe the COHb kinetics when the value of k_a was reduced, as shown in Figure C-9, indicates that changes in absorption rate due to vehicle do not completely explain discrepancies between model predictions and the observed data.

Model D simulations of total expired dichloromethane for oral doses with $k_a = 0.47$ or $1.8/\text{hour}$ are compared to the data of Kirschman et al. (1986) in Table C-3. Agreement with the data for water vehicle at 250 and 500 mg/kg for the model with $k_a = 1.8/\text{hour}$ is quite good, especially considering that in rats at 500 mg/kg, Kirschman et al. (1986) only accounted for 74% of the dose in all excreta (breath, urine, and feces). The lack of mass balance by Kirschman et al. (1986) (normal for such studies) could result from binding of dichloromethane metabolites to

tissue macromolecules, a mechanism not included in the model, as well as incomplete quantitation. Using $k_a = 0.47/\text{hour}$ obtained by fitting the oil-vehicle data of Pankow et al. (1991a), the model predicts the qualitative trend of less dichloromethane exhaled than with the water vehicle after administration of a 250 mg/kg dose, but not the extent of reduction observed for the amount exhaled (Table C-3). Since the model assumes that 100% of an oral dose is absorbed, this discrepancy may result if a significant fraction of the amount dosed in corn oil is excreted in the feces. (Only 0.3 and 0.05% was detected in the feces of mice after oral doses of 100 and 500 mg/kg dichloromethane in a water vehicle, respectively, by Kirschman et al. [1986].) Thus, the degree to which model simulations do match these data provides reasonably good confirmation of the model predictions, given the shortcomings noted. While the two oral datasets used result in quite different estimates of k_a , the difference appears to be at least partly due to the administration vehicle, with the lower $k_a = 0.47/\text{hour}$ obtained for an oil vehicle and a larger $k_a = 1.8/\text{hour}$ for water as the vehicle.

Table C-3. Observations and predictions of total expired dichloromethane resulting from gavage doses in rats^a

Dose (mg/kg)	Total dichloromethane exhaled (mg/kg)			
	Observations		Predictions	
	Corn oil vehicle	Water vehicle	$k_a = 0.47 \text{ hr}^{-1}$	$k_a = 1.8 \text{ hr}^{-1}$
250	91.7	166.3	171.4	193.4
500	Not reported	391.4	376.9	403.6
1,000	457.0	Not reported	795.5	826.8
2,000	1,314.5	Not reported	1,640.2	1,676.2

^aThe y-axis of Figure C-2 of Kirschman et al. (1986) appears to be mislabeled as mg instead of mg/kg; mg would result in unrealistic values. Assuming this is supposed to be mg/kg, and assuming average value of 250 g for a F344 rat, an expiration of 1,300 mg/kg given a dose of 2,000 mg/kg corresponds to 65% of the administered dose. This value is consistent with their observation in mice where 55% was observed expired as dichloromethane from a dose of 500 mg/kg, with the percentage expired increasing with dose.

Source: Kirschman et al. (1986).

C.3. MODEL OPTION SUMMARY

Our evaluation indicates that model D, a rat PBPK model represented by the basic model structure of Andersen et al. (1991), with the inclusion of lung dichloromethane metabolism via CYP (4% of liver metabolite production) and GST (14% of liver metabolite production) pathways (estimated from Reitz et al., 1989) and with liver metabolic parameters re-calibrated against data of Gargas et al. (1986), provides biological realism (including lung metabolism) and the best overall model agreement using a single set of model parameters, with the available rat data sets. The resulting model parameters are clearly dependent on the inclusion of lung

metabolism since the metabolic parameters for the liver ($V_{\max C}$, K_m , and k_{fC}) were re-optimized based on the presence of lung metabolism.

Our evaluation also revealed some deficiencies of the available models. The on-off rate of COHb binding or transport of CO between the blood and various tissues may be significant, rate-limiting steps in determining the kinetics of CO, which the existing model does not include. Explicitly including either of these mechanisms could allow the model to describe a slower rise to peak COHb levels and slower rate of exhalation of CO without compromising the model's ability to describe the observed, more rapid dichloromethane kinetics. The existing model structure cannot accurately describe the COHb and some of the CO exhalation data while also providing a good match to the dichloromethane data.

Reitz et al. (1997) employed a k_a value of 5.0 hours⁻¹ for dichloromethane in deriving acute- and intermediate-duration oral minimal risk levels. These investigators cited previous work in which a k_a value of 5.0/hour resulted in reasonable kinetic predictions for bolus or drinking water exposures of 1,1,1-trichloroethane in rats (Reitz et al., 1997), while a value of 5.4/hour provided good agreement between toxicokinetic observations and predictions for bolus doses of trichloroethylene in rats (Fisher et al., 1989). Our analysis indicates that k_a depends on vehicle (i.e., there is slower absorption from oil vehicles than water). Since a primary intention for the model is to predict dosimetry during bioassays conducted in drinking water, the k_a estimate from aqueous vehicle exposures (see Figure C-8) is considered most relevant. The value obtained here for aqueous vehicle, 1.8/hour, is driven by the observed rate of dichloromethane exhalation (Figure C-8A) and the peak height for dichloromethane concentration in blood (Figure C-8B) and liver (Figure C-8D). It can be noted that the timing of the blood dichloromethane peak predicted in Figure C-8B is later than the actual peak; this timing would be better matched with a larger value of k_a . However, increasing k_a would also increase the predicted peak height (that for liver is already over-predicted, Figure C-8D) and result in a more rapid rise in predicted exhaled dichloromethane than measured (Figure C-8A). So different components or aspects of the data set indicate different values for k_a , and the value of k_a obtained here can be considered an average among those which would best represent each data set.

In summary, we have examined four PBPK model structures and parameter sets, models A–D, to describe dichloromethane dosimetry in rats as candidates for use in risk assessment where the purpose is to estimate internal doses of dichloromethane that occurred during various bioassays. In comparing model predictions to a variety of data, one can say that while all of the four models do a fairly good job of fitting some of the data, none of the models fits all of the data very well, and there are some data for which none of the models provides a particularly good fit. Nevertheless there are some clear distinctions in model goodness of fit, and we conclude that model D, which includes lung metabolism via the CYP and GST pathways with primary metabolic parameters re-estimated for the liver (given defined lung:liver ratios), is the

best of the four models. This evaluation indicates that the existing model structure for CO and COHb does not adequately describe the corresponding data and that attempts to specifically use those data in setting key parameters could compromise the accuracy of those estimates. However, model D, the best of the models evaluated, should be adequate to predict rat internal dosimetry (dichloromethane blood concentrations or rates of metabolism) under bioassay exposure conditions.

**APPENDIX D. SUMMARY OF BENCHMARK DOSE (BMD) MODELING OF
NONCANCER ENDPOINTS**

D.1. ORAL RfD: BMD MODELING OF LIVER LESION INCIDENCE DATA FOR RATS EXPOSED TO DICHLOROMETHANE IN DRINKING WATER FOR 2 YEARS (SEROTA ET AL., 1986a)

BMD and BMDL refer to the model-predicted dose (and its lower 95% confidence limit) associated with 10% extra risk for the incidence of liver foci/areas of cellular alteration in male and female F344 rats given dichloromethane in drinking water for 2 years (Serota et al., 1986a) (Table D-1).

Table D-1. Incidence data for liver lesions and internal liver doses based on various metrics in male and female F344 rats exposed to dichloromethane in drinking water for 2 years (Serota et al., 1986a)

Sex	Nominal (actual) daily intake (mg/kg-d)	Rat liver lesion incidence ^a	Rat internal liver dose ^b			
			CYP	GST	GST and CYP	Parent AUC
Male (BW = 380 g)	0 (0)	52/76 (68%)	0	0	0	0
	5 (6)	22/34 (65%)	133.9	2.1	136.1	0.47
	50 (52)	35/38 (92%) ^c	872.7	58.8	931.4	13.1
	125 (125)	34/35 (97%) ^c	1,433.1	236.0	1,669.1	52.6
	250 (235)	40/41 (98%) ^c	1,868.6	561.5	2,430.0	125.0
Female (BW = 229 g)	0 (0)	34/67 (51%)	0	0	0	0
	5 (6)	12/29 (41%)	134.5	2.1	136.6	0.4
	50 (58)	30/41 (73%) ^c	977.8	66.0	1,043.8	12.6
	125 (136)	34/38 (89%) ^c	1,577.0	258.7	1,835.7	49.5
	250 (263)	31/34 (91%) ^c	2,070.0	642.4	2,712.3	122.9

^aLiver foci/areas of cellular alteration; number affected divided by total sample size.

^bInternal doses were estimated using a rat PBPK model from simulations of actual daily doses reported by the study authors. CYP dose is in units of mg dichloromethane metabolized via CYP pathway/L tissue/d; GST dose is in units of mg dichloromethane metabolized via GST pathway/L tissue/d; GST and CYP dose is in units of mg dichloromethane metabolized via CYP and GST pathways/L tissue/d; and Parent AUC dose is in units of mg dichloromethane × hrs/L tissue.

^cSignificantly ($p < 0.05$) different from control with Fisher's exact test.

Source: Serota et al. (1986a).

All available dichotomous models in the BMDS (version 2.0) were fit to male and female rat internal tissue doses of dichloromethane metabolized by the CYP pathway and incidences for animals with these liver lesions observed at the time of death (Table D-2). (The quantal model is identical to the one-stage multistage model and so is not included in this set of models). The male rats exhibited a greater sensitivity compared to the female rats (based on lower BMDL₁₀

values for all of the models), and thus, the male data are used as the basis for the RfD derivation. The logistic model was the best fitting model for the male incidence data based on AIC value among models with adequate fit (U.S. EPA, 2000b). (If more than one model shares the lowest AIC value, BMDL₁₀ values from these models may be averaged to obtain a POD. However, this average is not a well-defined lower bound, and should be referred to only as averages of BMDL₁₀s. U.S. EPA does not support averaging BMDLs in situations in which AIC values are similar, but not identical, because the level of statistical confidence is lost and because there is no consensus regarding a cut-off between similar and dissimilar AIC values.) Results for this model are presented below.

Table D-2. BMD modeling results for incidence of liver lesions in male and female F344 rats exposed to dichloromethane in drinking water for 2 years, based on liver-specific CYP metabolism dose metric (mg dichloromethane metabolism via CYP pathway per liter liver tissue per day)


Sex and model ^a	BMD ₁₀	BMDL ₁₀	χ^2 goodness of fit p-value	AIC
Males				
Gamma ^a	151.73	48.93	0.62	185.33
Logistic^b	85.17	61.78	0.75	183.61
Log-logistic ^a	213.73	37.06	0.83	184.79
Multistage (1) ^a	68.62	47.58	0.71	183.74
Probit	98.87	75.49	0.69	183.81
Log-probit ^a	197.65	77.56	0.81	184.84
Weibull ^a	117.29	48.39	0.57	185.49
Females				
Gamma ^a	336.38	98.70	0.52	233.07
Logistic	169.77	134.87	0.59	231.70
Log-logistic ^a	404.87	101.15	0.60	232.80
Multistage (1) ^a	123.59	91.46	0.47	232.32
Probit	179.59	146.27	0.59	231.70
Log-probit ^a	400.95	173.57	0.60	232.80
Weibull ^a	283.24	97.31	0.47	233.27

^aThese models in U.S. EPA BMDS version 2.0 were fit to the rat dose-response data shown in Table 5-1 by using internal dose metrics calculated with the rat PBPK model. Details of the models are as follows: Gamma and Weibull models restrict power ≥ 1 ; Log-logistic and Log-probit models restrict to slope > 1 , multistage model restrict betas ≥ 0 ; lowest degree polynomial with an adequate fit is reported (degree of polynomial noted in parentheses).

^bBolded model is the best-fitting model in the most sensitive sex (males), which is used in the RfD derivation.

Source: Serota et al. (1986a).

Logistic Model, Male Rats (Serota et al., 1986a), CYP Metabolism Metric



11:43 02/25 2009

Figure D-1. Predicted (logistic model) and observed incidence of noncancer liver lesions in male F344 rats exposed to dichloromethane in drinking water for 2 years (Serota et al., 1986a).

```

=====
      Logistic Model. (Version: 2.12; Date: 05/16/2008)
      Input Data File:
C:\USEPA\IRIS\DCM\noncancer\dataset1_liverlesions\male\2LogSerlog.(d)
      Gnuplot Plotting File:
C:\USEPA\IRIS\DCM\noncancer\dataset1_liverlesions\male\2LogSerlog.plt
                                          Wed Feb 25 11:43:00 2009
=====
  
```

BMDS Model Run

~~~~~  
 The form of the probability function is:

$$P[\text{response}] = 1/[1+\text{EXP}(-\text{intercept}-\text{slope}*\text{dose})]$$

Dependent variable = incidence  
 Independent variable = CYP  
 Slope parameter is not restricted

Total number of observations = 5  
 Total number of records with missing values = 0  
 Maximum number of iterations = 250  
 Relative Function Convergence has been set to: 1e-008  
 Parameter Convergence has been set to: 1e-008

Default Initial Parameter Values  
background = 0 Specified  
intercept = 0.683407  
slope = 0.00155073

Asymptotic Correlation Matrix of Parameter Estimates

( \*\*\* The model parameter(s) -background  
have been estimated at a boundary point, or have been specified by  
the user, and do not appear in the correlation matrix )

|           | intercept | slope |
|-----------|-----------|-------|
| intercept | 1         | -0.46 |
| slope     | -0.46     | 1     |

Parameter Estimates

| Variable  | Estimate   | Std. Err.   | 95.0% Wald Confidence Interval |                   |
|-----------|------------|-------------|--------------------------------|-------------------|
|           |            |             | Lower Conf. Limit              | Upper Conf. Limit |
| intercept | 0.669705   | 0.208716    | 0.260629                       | 1.07878           |
| slope     | 0.00182324 | 0.000412451 | 0.00101485                     | 0.00263163        |

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance | Test d.f. | P-value |
|---------------|-----------------|-----------|----------|-----------|---------|
| Full model    | -89.2097        | 5         |          |           |         |
| Fitted model  | -89.8065        | 2         | 1.19363  | 3         | 0.7545  |
| Reduced model | -106.616        | 1         | 34.8133  | 4         | <.0001  |

**AIC: 183.613**

Goodness of Fit

| Dose      | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|-----------|------------|----------|----------|------|-----------------|
| 0.0000    | 0.6614     | 50.269   | 52.000   | 76   | 0.420           |
| 133.9000  | 0.7138     | 24.269   | 22.000   | 34   | -0.861          |
| 872.7000  | 0.9056     | 34.412   | 35.000   | 38   | 0.326           |
| 1433.1000 | 0.9638     | 33.734   | 34.000   | 35   | 0.241           |
| 1868.6000 | 0.9833     | 40.316   | 40.000   | 41   | -0.385          |

**Chi^2 = 1.23      d.f. = 3      P-value = 0.7458**

Benchmark Dose Computation

Specified effect = 0.1  
Risk Type = Extra risk  
Confidence level = 0.95  
**BMD = 85.1668**  
**BMDL = 61.7795**



**D.2. INHALATION RfC: BMD MODELING OF LIVER LESION INCIDENCE DATA FOR RATS EXPOSED TO DICHLOROMETHANE VIA INHALATION FOR 2 YEARS (NITSCHKE ET AL., 1988a)**

BMD and BMDL refer to the model-predicted dose (and its lower 95% confidence limit) associated with 10% extra risk for the incidence of hepatic vacuolation in female F344 rats exposed to dichloromethane via inhalation for 2 years (Nitschke et al., 1988a) (Table D-3).

**Table D-3. Incidence data for liver lesions (hepatic vacuolation) and internal liver doses based on various metrics in female Sprague-Dawley rats exposed to dichloromethane via inhalation for 2 years (Nitschke et al., 1988a)**

| Sex                 | Exposure (ppm) | Liver lesion incidence <sup>a</sup> | Rat internal liver dose <sup>b</sup>                                                       |       |             |            |
|---------------------|----------------|-------------------------------------|--------------------------------------------------------------------------------------------|-------|-------------|------------|
|                     |                |                                     | CYP                                                                                        | GST   | GST and CYP | Parent AUC |
| Male                | 0              | 22/70 (31)                          | Not modeled because results from male rats were not provided for the 50 and 200 ppm groups |       |             |            |
|                     | 50             | Not reported                        | Not modeled because results for middle two doses were not reported                         |       |             |            |
|                     | 200            | Not reported                        |                                                                                            |       |             |            |
|                     | 500            | 28/70 (40)                          |                                                                                            |       |             |            |
| Female (BW = 229 g) | 0              | 41/70 (59%)                         | 0                                                                                          | 0     | 0           | 0          |
|                     | 50             | 42/70 (60%)                         | 280.3                                                                                      | 6.3   | 286.6       | 1.2        |
|                     | 200            | 41/70 (58%)                         | 656.5                                                                                      | 93.2  | 749.7       | 17.8       |
|                     | 500            | 53/70 (76%) <sup>c</sup>            | 772.6                                                                                      | 359.0 | 1,131.6     | 68.7       |

<sup>a</sup>Number affected divided by total sample size.

<sup>b</sup>Internal doses were estimated using a rat PBPK model using exposures reported by study authors (50 ppm = 174 mg/m<sup>3</sup>, 200 ppm = 695 mg/m<sup>3</sup>, and 500 ppm = 1,737 mg/m<sup>3</sup>) and are weighted-average daily values for 1 wk of exposure at 6 hrs/d, 5 d/wk. CYP dose is in units of mg dichloromethane metabolized via CYP pathway/L tissue/d; GST dose is in units of mg dichloromethane metabolized via GST pathway/L tissue/d; GST and CYP dose is in units of mg dichloromethane metabolized via CYP and GST pathways/L tissue/d; and Parent AUC dose is in units of mg dichloromethane × hrs/L tissue.

<sup>c</sup>Significantly (*p* < 0.05) different from control with Fisher's exact test.

Source: Nitschke et al. (1988a).

All available dichotomous models in the BMDS (version 2.0) were fit to male and female rat internal tissue doses of dichloromethane metabolized by the CYP pathway and incidences for animals with these liver lesions observed at the time of death (Table D-4). (The quantal model is identical to the one-stage multistage model and so is not included in this set of models). The log-probit model was the best fitting model for the female incidence data based on AIC value among models with adequate fit (U.S. EPA, 2000c). (If more than one model shares the lowest AIC value, BMDL<sub>10</sub> values from these models may be averaged to obtain a POD. However, this average is not a well-defined lower bound, and should be referred to only as averages of BMDL<sub>10</sub>s. U.S. EPA does not support averaging BMDLs in situations in which AIC values are

similar, but not identical, because the level of statistical confidence is lost and because there is no consensus regarding a cut-off between similar and dissimilar AIC values. Results for this model are presented below.)

**Table D-4. BMD modeling results for incidence of liver lesions in female Sprague-Dawley rats exposed to dichloromethane by inhalation for 2 years, based on liver specific CYP metabolism metric (mg dichloromethane metabolized via CYP pathway per liter liver tissue per day)**


| Model <sup>a</sup>              | BMD <sub>10</sub> | BMDL <sub>10</sub> | $\chi^2$<br>goodness of fit<br>p-value | AIC           |
|---------------------------------|-------------------|--------------------|----------------------------------------|---------------|
| Gamma <sup>a</sup>              | 614.27            | 225.96             | 0.48                                   | 367.22        |
| Logistic                        | 274.58            | 150.43             | 0.14                                   | 369.77        |
| Log-logistic <sup>a</sup>       | 697.90            | 499.42             | 0.94                                   | 365.90        |
| Multistage (3) <sup>a</sup>     | 506.94            | 153.13             | 0.25                                   | 368.53        |
| Probit                          | 275.49            | 152.52             | 0.14                                   | 369.75        |
| <b>Log-probit<sup>a,b</sup></b> | <b>728.96</b>     | <b>523.94</b>      | <b>0.98</b>                            | <b>365.82</b> |
| Weibull <sup>a</sup>            | 706.45            | 487.45             | 0.95                                   | 365.87        |

<sup>a</sup>These models in U.S. EPA BMDS version 2.0 were fit to the rat dose-response data shown in Table 5-5 by using internal dose metrics calculated with the rat PBPK model. Gamma and Weibull models restrict power  $\geq 1$ ; Log-logistic and Log-probit models restrict to slope  $> 1$ , multistage model restrict betas  $\geq 0$ ; lowest degree polynomial with an adequate fit reported (degree of polynomial in parentheses).

<sup>b</sup>Bolded model is the best-fitting model in the most sensitive sex (females), which is used in the RfC derivation.

Source: Nitschke et al. (1988a).

**Log Probit Model, Female Rats (Nitschke et al., 1988a), CYP Metabolism Metric**



16:07 02/25 2009

**Figure D-2. Predicted (log-probit model) and observed incidence of noncancer liver lesions in female Sprague-Dawley rats inhaling dichloromethane for 2 years (Nitschke 1988a).**

```

=====
      Probit Model. (Version: 3.1; Date: 05/16/2008)
      Input Data File:
C:\USEPA\BMDS21Beta\Data\dichloromethane\Dataset2\CYP\4LogCYPlog.(d)
      Gnuplot Plotting File:
C:\USEPA\BMDS21Beta\Data\dichloromethane\Dataset2\CYP\4LogCYPlog.plt
                                          Wed Feb 25 16:07:52 2009
=====

BMDS Model Run
~~~~~
The form of the probability function is:

P[response] = Background
 + (1-Background) * CumNorm(Intercept+Slope*Log(Dose)),

where CumNorm(.) is the cumulative normal distribution function

Dependent variable = Incidence
Independent variable = Dose
Slope parameter is restricted as slope >= 1

Total number of observations = 4
Total number of records with missing values = 0

```

Maximum number of iterations = 250  
 Relative Function Convergence has been set to: 1e-008  
 Parameter Convergence has been set to: 1e-008  
 User has chosen the log transformed model

Default Initial (and Specified) Parameter Values  
 background = 0.585714  
 intercept = -7.69845  
 slope = 1

Asymptotic Correlation Matrix of Parameter Estimates

( \*\*\* The model parameter(s) -slope  
 have been estimated at a boundary point, or have been specified by  
 the user,  
 and do not appear in the correlation matrix )

|            | background | intercept |
|------------|------------|-----------|
| background | 1          | -0.37     |
| intercept  | -0.37      | 1         |

Parameter Estimates

| Variable   | Estimate | Std. Err. | 95.0% Wald Confidence Interval |                   |
|------------|----------|-----------|--------------------------------|-------------------|
|            |          |           | Lower Conf. Limit              | Upper Conf. Limit |
| background | 0.590379 | 0.0339871 | 0.523765                       | 0.656992          |
| intercept  | -119.931 | 0.346784  | -120.61                        | -119.251          |
| slope      | 18       | NA        |                                |                   |

NA - Indicates that this parameter has hit a bound implied by some inequality constraint and thus has no standard error.

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance  | Test d.f. | P-value |
|---------------|-----------------|-----------|-----------|-----------|---------|
| Full model    | -180.889        | 4         |           |           |         |
| Fitted model  | -180.909        | 2         | 0.0403244 | 2         | 0.98    |
| Reduced model | -184.186        | 1         | 6.5937    | 3         | 0.08604 |

**AIC: 365.818**

Goodness of Fit

| Dose     | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|----------|------------|----------|----------|------|-----------------|
| 0.0000   | 0.5904     | 41.326   | 41.000   | 70   | -0.079          |
| 280.3400 | 0.5904     | 41.326   | 42.000   | 70   | 0.164           |
| 656.4800 | 0.5907     | 41.349   | 41.000   | 70   | -0.085          |
| 772.5900 | 0.7571     | 52.998   | 53.000   | 70   | 0.001           |

**Chi^2 = 0.04      d.f. = 2      P-value = 0.9801**

Benchmark Dose Computation

Specified effect = 0.1  
 Risk Type = Extra risk  
 Confidence level = 0.95  
**BMD = 728.956**  
**BMDL = 523.944**

## **APPENDIX E: SUMMARY OF BENCHMARK DOSE (BMD) MODELING OF CANCER ENDPOINTS**

### **E.1. ORAL CANCER SLOPE FACTORS: BMD MODELING OF LIVER TUMOR INCIDENCE DATA FOR MICE EXPOSED TO DICHLOROMETHANE IN DRINKING WATER FOR 2 YEARS (SEROTA ET AL., 1986b; HAZLETON LABORATORIES, 1983)**

BMD<sub>10</sub> and BMDL<sub>10</sub> refer to the model-predicted dose (and its lower 95% confidence limit) associated with 10% extra risk for the incidence of hepatocellular adenoma and carcinoma in male mice given dichloromethane in drinking water for 2 years (Serota et al., 1986b; Hazleton Laboratories, 1983) (Table E-1). Multistage models were fit to male mouse internal tissue doses of dichloromethane metabolized by the GST pathway and incidences for animals with liver tumors observed at the time of death. Different polynomial models and models dropping dose groups starting with the highest dose group were compared based on adequacy of model fit as assessed by overall  $\chi^2$  goodness of fit ( $p$ -value > 0.10) and examination of residuals at the 0 dose exposure (controls) and in the region of the BMR. The predicted BMD<sub>10</sub> and BMDL<sub>10</sub> for the incidence data are 73.0 and 39.6 mg dichloromethane metabolized via GST pathways per liter tissue per day, respectively, for the internal liver metabolism metric, and 3.05 and 1.65 mg dichloromethane metabolized via GST pathway in lung and liver/kg-day, respectively, for the whole body metabolism metric (Table E-2).

**Table E-1. Incidence data for liver tumors and internal liver doses, based on GST metabolism dose metrics, in male B6C3F<sub>1</sub> mice exposed to dichloromethane in drinking water for 2 years**

| Sex                      | Nominal (actual) daily intake (mg/kg-d) | Mouse liver tumor incidence <sup>a</sup> | Mouse internal liver metabolism dose <sup>b</sup> | Mouse whole body metabolism dose <sup>c</sup> |
|--------------------------|-----------------------------------------|------------------------------------------|---------------------------------------------------|-----------------------------------------------|
| Male<br>(BW =<br>37.3 g) | 0 (0)                                   | 24/125 (19%)                             | 0                                                 | 0                                             |
|                          | 60 (61)                                 | 51/199 (26%)                             | 17.5                                              | 0.73                                          |
|                          | 125 (124)                               | 30/99 (30%)                              | 63.3                                              | 2.65                                          |
|                          | 185 (177)                               | 31/98 (32%)                              | 112.0                                             | 4.68                                          |
|                          | 250 (234)                               | 35/123 (28%)                             | 169.5                                             | 7.1                                           |

<sup>a</sup>Hepatocellular carcinoma or adenoma combined. Mice dying prior to 52 wks were excluded from the denominators. Cochran-Armitage trend *p*-value = 0.058. *P*-values for comparisons with the control group were 0.071, 0.023, 0.019, and 0.036 in the 60, 125, 185, and 250 mg/kg-d groups, respectively, based on statistical analyses reported by Hazleton Laboratories (1983).

<sup>b</sup>mg dichloromethane metabolized via GST pathway/L liver/d. Internal doses were estimated from simulations of actual daily doses reported by the study authors.

<sup>c</sup>Based on the sum of dichloromethane metabolized via the GST pathway in the lung plus the liver, normalized to total BW (i.e., [lung GST metabolism (mg/d) + liver GST metabolism (mg/d)]/kg BW). Units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

Sources: Serota et al. (1986b); Hazleton Laboratories (1983).

**Table E-2. BMD modeling results and tumor risk factors for internal dose metric associated with 10% extra risk for liver tumors in male B6C3F<sub>1</sub> mice exposed to dichloromethane in drinking water for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics**

| Internal dose metric <sup>a</sup> | BMDS model <sup>b</sup> | $\chi^2$ goodness of fit <i>p</i> -value | Mouse BMD <sub>10</sub> <sup>c</sup> | Mouse BMDL <sub>10</sub> <sup>c</sup> | Allometric-scaled human BMDL <sub>10</sub> <sup>d</sup> | Tumor risk factor <sup>e</sup> |                       |
|-----------------------------------|-------------------------|------------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------------------------|--------------------------------|-----------------------|
|                                   |                         |                                          |                                      |                                       |                                                         | Scaling = 1.0                  | Allometric-scaled     |
| Liver-specific                    | MS (1,1)                | 0.56                                     | 73.0                                 | 39.6                                  | 5.66                                                    | $2.53 \times 10^{-3}$          | $1.77 \times 10^{-2}$ |
| Whole-body                        | MS (1,1)                | 0.56                                     | 3.05                                 | 1.65                                  | 0.24                                                    | –                              | $4.24 \times 10^{-1}$ |

<sup>a</sup>Liver specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue per d; Whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

<sup>b</sup>The multistage (MS) model in U.S. EPA BMDS version 2.0 was fit to the mouse dose-response data shown in Table 5-11 using internal dose metrics calculated with the mouse PBPK model. Numbers in parentheses indicate (1) the number of dose groups dropped in order to obtain an adequate fit, and (2) the degree polynomial of the model.

<sup>c</sup>BMD<sub>10</sub> and BMDL<sub>10</sub> refer to the BMD-model-predicted mouse internal and its 95% lower confidence limit, associated with a 10% extra risk for the incidence of tumors.

<sup>d</sup>Mouse BMDL<sub>10</sub> divided by  $(BW_{\text{human}}/BW_{\text{mouse}})^{0.25} = 7$ .


<sup>e</sup>Dichloromethane tumor risk factor (extra risk per unit internal dose) derived by dividing the BMR (0.1) by the mouse BMDL<sub>10</sub> and by the allometric-scaled human BMDL<sub>10</sub>, for the scaling = 1.0 and allometric-scaled risk factors, respectively.

Modeling results are presented in the subsequent sections for the tissue-specific liver-metabolism metric (Section E.1.1) and the whole-body metabolism metric (Section E.1.2).

**E.1.1. Modeling Results for the Internal Liver Metabolism Metric**

**Serota et al. (1986b), Hazleton Laboratories, 1983: Internal liver dose-response, highest dose dropped**

1-degree polynomial



**Figure E-1. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F<sub>1</sub> mice exposed to dichloromethane in drinking water for 2 years, using liver-specific metabolism dose metric (Serota et al., 1986b; Hazleton Laboratories, 1983).**

```

=====
Multistage Cancer Model. (Version: 1.7; Date: 05/16/2008)
Input Data File: C:\USEPA\IRIS\DCM\Serota\highdosedropped\1MulSerMS_.(d)
Gnuplot Plotting File:
C:\USEPA\IRIS\DCM\Serota\highdosedropped\1MulSerMS_.plt
Thu Feb 19 13:46:49 2009
=====
BMDS Model Run
=====
The form of the probability function is:
P[response] = background + (1-background)*[1-EXP(
 -beta1*dose^1)]

```

The parameter betas are restricted to be positive  
 Dependent variable = incidence  
 Independent variable = dose

Total number of observations = 4  
 Total number of records with missing values = 0  
 Total number of parameters in model = 2  
 Total number of specified parameters = 0  
 Degree of polynomial = 1

Maximum number of iterations = 250  
 Relative Function Convergence has been set to: 2.22045e-016  
 Parameter Convergence has been set to: 1.49012e-008

\*\*\*\* We are sorry but Relative Function and Parameter Convergence \*\*\*\*  
 \*\*\*\* are currently unavailable in this model. Please keep checking \*\*\*\*  
 \*\*\*\* the web sight for model updates which will eventually \*\*\*\*  
 \*\*\*\* incorporate these convergence criterion. Default values used. \*\*\*\*

Default Initial Parameter Values  
 Background = 0.218634  
 Beta(1) = 0.00136788

Asymptotic Correlation Matrix of Parameter Estimates

|            | Background | Beta(1) |
|------------|------------|---------|
| Background | 1          | -0.7    |
| Beta(1)    | -0.7       | 1       |

Parameter Estimates

| Variable   | Estimate   | Std. Err. | 95.0% Wald Confidence Interval |                   |
|------------|------------|-----------|--------------------------------|-------------------|
|            |            |           | Lower Conf. Limit              | Upper Conf. Limit |
| Background | 0.218642   | *         | *                              | *                 |
| Beta(1)    | 0.00144288 | *         | *                              | *                 |

\* - Indicates that this value is not calculated.

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance | Test d.f. | P-value |
|---------------|-----------------|-----------|----------|-----------|---------|
| Full model    | -296.282        | 4         |          |           |         |
| Fitted model  | -296.87         | 2         | 1.1754   | 2         | 0.5556  |
| Reduced model | -299.126        | 1         | 5.68747  | 3         | 0.1278  |

**AIC: 597.74**

Goodness of Fit

| Dose     | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|----------|------------|----------|----------|------|-----------------|
| 0.0000   | 0.2186     | 27.330   | 24.000   | 125  | -0.721          |
| 17.5000  | 0.2381     | 47.387   | 51.000   | 199  | 0.601           |
| 63.3000  | 0.2868     | 28.398   | 30.000   | 99   | 0.356           |
| 112.0000 | 0.3352     | 32.853   | 31.000   | 98   | -0.397          |

**Chi^2 = 1.16      d.f. = 2      P-value = 0.5585**

Benchmark Dose Computation  
 Specified effect = 0.1  
 Risk Type = Extra risk  
 Confidence level = 0.95  
**BMD = 73.0211**



**BMDL = 39.6034**

**BMDU = 335.18**


Taken together, (39.6034, 335.18 ) is a 90 % two-sided confidence interval for the BMD

**Multistage Cancer Slope Factor = 0.00252504**

## E.1.2. Modeling Results for the Whole Body Metabolism Metric

Serota et al. (1986b), Hazleton Laboratories, 1983: Internal whole-body metabolism dose-response in mice, highest dose dropped

1-degree polynomial



**Figure E-2. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F<sub>1</sub> mice exposed to dichloromethane in drinking water for 2 years, using whole-body metabolism dose metric (Serota et al., 1986b; Hazleton Laboratories, 1983).**

```

=====
Multistage Cancer Model. (Version: 1.7; Date: 05/16/2008)
Input Data File: C:\USEPA\IRIS\DCM\Serota\highdosedropped\1MulSerMS_.(d)
Gnuplot Plotting File:
C:\USEPA\IRIS\DCM\Serota\highdosedropped\1MulSerMS_.plt
Sat Feb 21 17:33:49 2009
=====

```

BMDS Model Run

The form of the probability function is:

$$P[\text{response}] = \text{background} + (1 - \text{background}) * [1 - \text{EXP}(-\text{beta}1 * \text{dose}^1)]$$

The parameter betas are restricted to be positive  
 Dependent variable = incidence  
 Independent variable = dose

Total number of observations = 4  
 Total number of records with missing values = 0  
 Total number of parameters in model = 2  
 Total number of specified parameters = 0  
 Degree of polynomial = 1  
 Maximum number of iterations = 250  
 Relative Function Convergence has been set to: 2.22045e-016  
 Parameter Convergence has been set to: 1.49012e-008

\*\*\*\* We are sorry but Relative Function and Parameter Convergence \*\*\*\*  
 \*\*\*\* are currently unavailable in this model. Please keep checking \*\*\*\*  
 \*\*\*\* the web sight for model updates which will eventually \*\*\*\*  
 \*\*\*\* incorporate these convergence criterion. Default values used. \*\*\*\*

Default Initial Parameter Values  
 Background = 0.218649  
 Beta(1) = 0.032703

Asymptotic Correlation Matrix of Parameter Estimates

|            | Background | Beta(1) |
|------------|------------|---------|
| Background | 1          | -0.7    |
| Beta(1)    | -0.7       | 1       |

Parameter Estimates

| Variable   | Estimate  | Std. Err. | 95.0% Wald Confidence Interval |                   |
|------------|-----------|-----------|--------------------------------|-------------------|
|            |           |           | Lower Conf. Limit              | Upper Conf. Limit |
| Background | 0.218662  | *         | *                              | *                 |
| Beta(1)    | 0.0344939 | *         | *                              | *                 |

\* - Indicates that this value is not calculated.

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance | Test d.f. | P-value |
|---------------|-----------------|-----------|----------|-----------|---------|
| Full model    | -296.282        | 4         |          |           |         |
| Fitted model  | -296.871        | 2         | 1.17643  | 2         | 0.5553  |
| Reduced model | -299.126        | 1         | 5.68747  | 3         | 0.1278  |
| <b>AIC:</b>   | <b>597.741</b>  |           |          |           |         |

Goodness of Fit

| Dose   | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|--------|------------|----------|----------|------|-----------------|
| 0.0000 | 0.2187     | 27.333   | 24.000   | 125  | -0.721          |
| 0.7310 | 0.2381     | 47.385   | 51.000   | 199  | 0.602           |
| 2.6470 | 0.2868     | 28.397   | 30.000   | 99   | 0.356           |
| 4.6840 | 0.3352     | 32.853   | 31.000   | 98   | -0.396          |

**Chi^2 = 1.17      d.f. = 2      P-value = 0.5582**

Benchmark Dose Computation  
 Specified effect = 0.1  
 Risk Type = Extra risk  
 Confidence level = 0.95

**BMD = 3.05447**

**BMDL = 1.65649**

BMDU = 14.0263

Taken together, (1.65649, 14.0263) is a 90 % two-sided confidence interval for the BMD

**Multistage Cancer Slope Factor = 0.0603686**

**E.2. CANCER IUR: BMD MODELING OF LIVER AND LUNG TUMOR INCIDENCE DATA FOR MALE MICE EXPOSED TO DICHLOROMETHANE VIA INHALATION FOR 2 YEARS (MENNEAR ET AL., 1988; NTP, 1986)**

BMD<sub>10</sub> and BMDL<sub>10</sub> refer to the model-predicted dose (and its lower 95% confidence limit) associated with 10% extra risk for the combined incidence of adenoma and carcinoma of the liver or lung of male B6C3F<sub>1</sub> mice inhaling dichloromethane for 2 years (Mennear et al., 1988; NTP, 1986) (Table E-3).

**Table E-3. Incidence data for liver and lung tumors and internal doses based on GST metabolism dose metrics in male B6C3F<sub>1</sub> mice exposed to dichloromethane via inhalation for 2 years**

| Sex, tumor type          | BW (g) | External dichloromethane concentration (ppm) | Mouse tumor incidence    | Mouse internal tissue dose <sup>a</sup> | Mouse whole body metabolism dose <sup>b</sup> |
|--------------------------|--------|----------------------------------------------|--------------------------|-----------------------------------------|-----------------------------------------------|
| Male, liver <sup>c</sup> | –      | 0                                            | 22/50 (44%) <sup>d</sup> | 0                                       | 0                                             |
|                          | 34.0   | 2,000                                        | 24/47 (51%)              | 2,363.7                                 | 100.2                                         |
|                          | 32.0   | 4,000                                        | 33/47 (70%)              | 4,972.2                                 | 210.7                                         |
| Male, lung <sup>e</sup>  | –      | 0                                            | 5/50 (10%) <sup>d</sup>  | 0                                       | 0                                             |
|                          | 34.0   | 2,000                                        | 27/47 (55%)              | 475.0                                   | 100.2                                         |
|                          | 32.0   | 4,000                                        | 40/47 (85%)              | 992.2                                   | 210.7                                         |

<sup>a</sup>For liver tumors: mg dichloromethane metabolized via GST pathway/L liver tissue/d from 6 hrs/d, 5 d/wk exposure; for lung tumors: mg dichloromethane metabolized via GST pathway/L lung tissue/d from 6 hrs/d, 5 d/wk exposure.

<sup>b</sup>Based on the sum of dichloromethane metabolized via the GST pathway in the lung plus the liver, normalized to total BW (i.e., [lung GST metabolism (mg/d) + liver GST metabolism (mg/d)]/kg BW). Units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

<sup>c</sup>Hepatocellular carcinoma or adenoma. Mice dying prior to 52 wks were excluded from the denominators.

<sup>d</sup>Statistically significant increasing trend (by incidental and life-table tests;  $p \leq 0.01$ ).

<sup>e</sup>Bronchoalveolar carcinoma or adenoma. Mice dying prior to 52 wks were excluded from the denominators.

Sources: Mennear et al. (1988); NTP (1986).

Multistage models were fit to male mouse internal tissue doses of dichloromethane metabolized by the GST pathway and incidences for animals with liver tumors observed at the time of death. The predicted BMD<sub>10</sub> and BMDL<sub>10</sub> for the liver tumor incidence data are 913.9 and 544.4 mg dichloromethane metabolized via GST pathways per liter liver per day, respectively, for the internal liver metabolism metric, and 38.7 and 23.1 mg dichloromethane metabolized via GST pathway in lung and liver/kg-day, respectively, for the whole body metabolism metric (Table E-4). For lung tumors, the BMD<sub>10</sub> and BMDL<sub>10</sub> are 61.7 and 48.7 mg dichloromethane metabolized via GST pathway per liter tissue per day, respectively, for the lung-specific metric, and 13.1 and 10.3 mg dichloromethane metabolized via GST pathway in lung and liver/kg-day, respectively, for the whole body metabolism metric.

**Table E-4. BMD modeling results and tumor risk factors associated with 10% extra risk for liver and lung tumors in male B6C3F<sub>1</sub> mice exposed by inhalation to dichloromethane for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics**

| Internal dose metric <sup>a</sup> |             | BMDS model <sup>b</sup> | $\chi^2$ goodness of fit <i>p</i> -value | Mouse BMD <sub>10</sub> <sup>c</sup> | Mouse BMDL <sub>10</sub> <sup>c</sup> | Allometric-scaled human BMDL <sub>10</sub> <sup>d</sup> | Tumor risk factor <sup>e</sup> |                       |
|-----------------------------------|-------------|-------------------------|------------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------------------------|--------------------------------|-----------------------|
|                                   |             |                         |                                          |                                      |                                       |                                                         | Scaling = 1.0                  | Allometric-scaled     |
| Tissue-specific                   | Male, liver | MS (0,1)                | 0.40                                     | 913.9                                | 544.4                                 | 77.8                                                    | $1.84 \times 10^{-4}$          | $1.29 \times 10^{-3}$ |
|                                   | Male, lung  | MS (0,1)                | 0.64                                     | 61.7                                 | 48.6                                  | 7.0                                                     | $2.06 \times 10^{-3}$          | $1.44 \times 10^{-2}$ |
| Whole body                        | Male, liver | MS (0,1)                | 0.40                                     | 38.7                                 | 23.1                                  | 3.3                                                     | –                              | $3.03 \times 10^{-2}$ |
|                                   | Male, lung  | MS (0,1)                | 0.66                                     | 13.1                                 | 10.3                                  | 1.5                                                     | –                              | $6.80 \times 10^{-2}$ |

<sup>a</sup>Tissue specific dose units = mg dichloromethane metabolized via GST pathway per liter (liver or lung) tissue per d; whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d).

<sup>b</sup>The multistage (MS) model in EPA BMDS version 2.0 was fit to the mouse dose-response data shown in Table 5-17 using internal dose metrics calculated with the mouse PBPK model. Numbers in parentheses indicate: (1) the number of dose groups dropped in order to obtain an adequate fit, and (2) the degree polynomial of the model.

<sup>c</sup>BMD<sub>10</sub> and BMDL<sub>10</sub> refer to the BMD-model-predicted mouse internal dose and its 95% lower confidence limit, associated with a 10% extra risk for the incidence of tumors.


<sup>d</sup>Mouse BMDL<sub>10</sub> divided by  $(BW_{\text{human}}/BW_{\text{mouse}})^{0.25} = 7$ .

<sup>e</sup>Dichloromethane tumor risk factor (extra risk per unit internal dose) derived by dividing the BMR (0.1) by the mouse BMDL<sub>10</sub> and by the allometric-scaled human BMDL<sub>10</sub>, for the scaling = 1.0 and allometric-scaled risk factors, respectively.

Modeling results are presented in the subsequent sections for the tissue-specific liver-metabolism metric for liver tumors (Section E.2.1), tissue-specific lung metabolism metric for lung tumors (section E-2.2), and the whole-body metabolism metric for liver tumors (Section E.2.3) and lung tumors (Section E.2.4).

**E.2.1. Modeling Results for the Internal Liver Metabolism Metric, Liver Tumors. Mennear et al. (1988); NTP (1986): Internal Liver Dose-Response for Liver Tumors in Male Mice**

1-degree polynomial



**Figure E-3. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F<sub>1</sub> mice exposed by inhalation to dichloromethane for 2 years, using liver-specific metabolism dose metric (Mennear et al., 1988; NTP, 1986).**

```

=====
Multistage Cancer Model. (Version: 1.7; Date: 05/16/2008)
Input Data File: C:\USEPA\IRIS\DCM\NTP\lung\male\lMulNTPMS_.(d)
Gnuplot Plotting File: C:\USEPA\IRIS\DCM\NTP\lung\male\lMulNTPMS_.plt
Thu Feb 19 16:08:19 2009
=====

```

BMDS Model Run

The form of the probability function is:

$$P[\text{response}] = \text{background} + (1-\text{background}) * [1 - \text{EXP}(-\text{beta}1 * \text{dose}^1)]$$

The parameter betas are restricted to be positive

Dependent variable = incidence

Independent variable = dose

Total number of observations = 3  
 Total number of records with missing values = 0  
 Total number of parameters in model = 2  
 Total number of specified parameters = 0  
 Degree of polynomial = 1  
 Maximum number of iterations = 250  
 Relative Function Convergence has been set to: 2.22045e-016  
 Parameter Convergence has been set to: 1.49012e-008

\*\*\*\* We are sorry but Relative Function and Parameter Convergence \*\*\*\*  
 \*\*\*\* are currently unavailable in this model. Please keep checking \*\*\*\*  
 \*\*\*\* the web sight for model updates which will eventually \*\*\*\*  
 \*\*\*\* incorporate these convergence criterion. Default values used. \*\*\*\*

Default Initial Parameter Values

Background = 0.406706  
 Beta(1) = 0.00012805

Asymptotic Correlation Matrix of Parameter Estimates

|            | Background | Beta(1) |
|------------|------------|---------|
| Background | 1          | -0.69   |
| Beta(1)    | -0.69      | 1       |

Parameter Estimates

| Variable   | Estimate    | Std. Err. | 95.0% Wald Confidence Interval |                   |
|------------|-------------|-----------|--------------------------------|-------------------|
|            |             |           | Lower Conf. Limit              | Upper Conf. Limit |
| Background | 0.421771    | *         | *                              | *                 |
| Beta(1)    | 0.000115283 | *         | *                              | *                 |

\* - Indicates that this value is not calculated.

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance | Test d.f. | P-value |
|---------------|-----------------|-----------|----------|-----------|---------|
| Full model    | -95.4892        | 3         |          |           |         |
| Fitted model  | -95.8368        | 2         | 0.695297 | 1         | 0.4044  |
| Reduced model | -99.1316        | 1         | 7.28482  | 2         | 0.02619 |

**AIC: 195.674**

Goodness of Fit

| Dose      | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|-----------|------------|----------|----------|------|-----------------|
| 0.0000    | 0.4218     | 21.089   | 22.000   | 50   | 0.261           |
| 2363.7000 | 0.5597     | 26.305   | 24.000   | 47   | -0.677          |
| 4972.2000 | 0.6740     | 31.680   | 33.000   | 47   | 0.411           |

Chi^2 = 0.70      d.f. = 1      P-value = 0.4042



Benchmark Dose Computation  
Specified effect = 0.1

Risk Type = Extra risk

Confidence level = 0.95

**BMD = 913.932**

**BMDL = 544.35**


BMDU = 2569.01

Taken together, (544.35 , 2569.01) is a 90 % two-sided confidence interval for the BMD

**Multistage Cancer Slope Factor = 0.000183705**

**E.2.2. Modeling Results for the Internal Lung Metabolism Metric, Lung Tumors.  
Mennear et al. (1988); NTP (1986): Internal Lung Dose-Response for Lung Tumors in  
Male Mice**

1-degree polynomial



**Figure E-4. Predicted and observed incidence of animals with carcinoma or adenoma in the lung of male B6C3F<sub>1</sub> mice exposed by inhalation to dichloromethane for 2 years, using liver-specific metabolism dose metric (Mennear et al., 1988; NTP, 1986).**

```

=====
Multistage Cancer Model. (Version: 1.7; Date: 05/16/2008)
Input Data File: C:\USEPA\IRIS\DCM\NTP\lung\male\1MulNTPMS_.d)
Gnuplot Plotting File: C:\USEPA\IRIS\DCM\NTP\lung\male\1MulNTPMS_.plt
Thu Feb 19 21:20:36 2009
=====
BMDS Model Run
=====

```

The form of the probability function is:

$$P[\text{response}] = \text{background} + (1 - \text{background}) * [1 - \text{EXP}(-\text{beta}1 * \text{dose}^1)]$$

The parameter betas are restricted to be positive

Dependent variable = incidence  
Independent variable = dose

Total number of observations = 3  
Total number of records with missing values = 0  
Total number of parameters in model = 2  
Total number of specified parameters = 0  
Degree of polynomial = 1

Maximum number of iterations = 250  
Relative Function Convergence has been set to: 2.22045e-016  
Parameter Convergence has been set to: 1.49012e-008

\*\*\*\* We are sorry but Relative Function and Parameter Convergence \*\*\*\*  
\*\*\*\* are currently unavailable in this model. Please keep checking \*\*\*\*  
\*\*\*\* the web sight for model updates which will eventually \*\*\*\*  
\*\*\*\* incorporate these convergence criterion. Default values used. \*\*\*\*

Default Initial Parameter Values

Background = 0.0642604  
Beta(1) = 0.00181622

Asymptotic Correlation Matrix of Parameter Estimates

|            | Background | Beta(1) |
|------------|------------|---------|
| Background | 1          | -0.56   |
| Beta(1)    | -0.56      | 1       |

Parameter Estimates

| Variable   | Estimate   | Std. Err. | 95.0% Wald Confidence Interval |                   |  |
|------------|------------|-----------|--------------------------------|-------------------|--|
|            |            |           | Lower Conf. Limit              | Upper Conf. Limit |  |
| Background | 0.0980033  | *         | *                              | *                 |  |
| Beta(1)    | 0.00170868 | *         | *                              | *                 |  |

\* - Indicates that this value is not calculated.

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance | Test d.f. | P-value |
|---------------|-----------------|-----------|----------|-----------|---------|
| Full model    | -68.0892        | 3         |          |           |         |
| Fitted model  | -68.199         | 2         | 0.219579 | 1         | 0.6394  |
| Reduced model | -99.8132        | 1         | 63.4479  | 2         | <.0001  |

**AIC: 140.398**

Goodness of Fit

| Dose     | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|----------|------------|----------|----------|------|-----------------|
| 0.0000   | 0.0980     | 4.900    | 5.000    | 50   | 0.047           |
| 475.0000 | 0.5994     | 28.171   | 27.000   | 47   | -0.349          |
| 992.2000 | 0.8345     | 39.219   | 40.000   | 47   | 0.306           |

**Chi^2 = 0.22      d.f. = 1      P-value = 0.6408**

Benchmark Dose Computation

Specified effect = 0.1  
Risk Type = Extra risk

Confidence level = 0.95

**BMD = 61.6618**

**BMDL = 48.628**


BMDU = 80.2137

Taken together, (48.628 , 80.2137) is a 90 % two-sided confidence interval for the BMD

Multistage Cancer Slope Factor = 0.00205643

### E.2.3. Modeling Results for the Whole Body Metabolism Metric, Liver Tumors. Mennear et al. (1988); NTP (1986): Internal Whole-Body Metabolism Dose-Response for Liver Tumors in Male Mice

1-degree polynomial



17:59 02/21 2009

**Figure E-5. Predicted and observed incidence of animals with hepatocellular carcinoma or adenoma in male B6C3F<sub>1</sub> mice exposed by inhalation to dichloromethane for 2 years, using whole-body metabolism dose metric (Mennear et al., 1988; NTP, 1986).**

```
=====
Multistage Cancer Model. (Version: 1.7; Date: 05/16/2008)
Input Data File: C:\USEPA\IRIS\DCM\NTP\liver\male\1MulNTPMS_.(d)
Gnuplot Plotting File: C:\USEPA\IRIS\DCM\NTP\liver\male\1MulNTPMS_.plt
Sat Feb 21 17:59:59 2009
=====
BMDS Model Run
```

~~~~~  
The form of the probability function is:

$$P[\text{response}] = \text{background} + (1-\text{background}) * [1 - \text{EXP}(-\text{beta1} * \text{dose}^1)]$$

The parameter betas are restricted to be positive

Dependent variable = incidence  
Independent variable = dose

Total number of observations = 3  
Total number of records with missing values = 0  
Total number of parameters in model = 2  
Total number of specified parameters = 0  
Degree of polynomial = 1  
Maximum number of iterations = 250  
Relative Function Convergence has been set to: 2.22045e-016  
Parameter Convergence has been set to: 1.49012e-008

\*\*\*\* We are sorry but Relative Function and Parameter Convergence \*\*\*\*  
\*\*\*\* are currently unavailable in this model. Please keep checking \*\*\*\*  
\*\*\*\* the web sight for model updates which will eventually \*\*\*\*  
\*\*\*\* incorporate these convergence criterion. Default values used. \*\*\*\*

Default Initial Parameter Values

Background = 0.406695  
Beta(1) = 0.00302163

Asymptotic Correlation Matrix of Parameter Estimates

|            | Background | Beta(1) |
|------------|------------|---------|
| Background | 1          | -0.69   |
| Beta(1)    | -0.69      | 1       |

Parameter Estimates

| Variable   | Estimate   | Std. Err. | 95.0% Wald Confidence Interval |                   |
|------------|------------|-----------|--------------------------------|-------------------|
|            |            |           | Lower Conf. Limit              | Upper Conf. Limit |
| Background | 0.421768   | *         | *                              | *                 |
| Beta(1)    | 0.00272018 | *         | *                              | *                 |

\* - Indicates that this value is not calculated.

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance | Test d.f. | P-value |
|---------------|-----------------|-----------|----------|-----------|---------|
| Full model    | -95.4892        | 3         |          |           |         |
| Fitted model  | -95.8372        | 2         | 0.696122 | 1         | 0.4041  |
| Reduced model | -99.1316        | 1         | 7.28482  | 2         | 0.02619 |

**AIC: 195.674**

Goodness of Fit

| Dose     | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|----------|------------|----------|----------|------|-----------------|
| 0.0000   | 0.4218     | 21.088   | 22.000   | 50   | 0.261           |
| 100.2000 | 0.5597     | 26.307   | 24.000   | 47   | -0.678          |
| 210.7000 | 0.6740     | 31.679   | 33.000   | 47   | 0.411           |

**Chi^2 = 0.70      d.f. = 1      P-value = 0.4039**

Benchmark Dose Computation

Specified effect = 0.1

Risk Type = Extra risk

Confidence level = 0.95

**BMD = 38.733**

**BMDL = 23.0698**


BMDU = 108.885

Taken together, (23.0698, 108.885) is a 90 % two-sided confidence interval for the BMD

**Multistage Cancer Slope Factor = 0.00433467**

**E.2.4. Modeling Results for the Whole Body Metabolism Metric, Lung Tumors. Mennear et al. (1988); NTP (1986): Internal Whole-Body Metabolism Dose-Response for Lung Tumors in Male Mice**

1-degree polynomial



**Figure E-6. Predicted and observed incidence of animals with carcinoma or adenoma in the lung of male B6C3F<sub>1</sub> mice exposed by inhalation to dichloromethane for 2 years, using whole-body metabolism dose metric (Mennear et al., 1988; NTP, 1986).**

```

=====
Multistage Cancer Model. (Version: 1.7; Date: 05/16/2008)
Input Data File: C:\USEPA\IRIS\DCM\NTP\lung\male\lMulNTPMS_.(d)
Gnuplot Plotting File: C:\USEPA\IRIS\DCM\NTP\lung\male\lMulNTPMS_.plt
Sat Feb 21 21:17:36 2009
=====
BMDS Model Run
~~~~~
The form of the probability function is:
P[response] = background + (1-background)*[1-EXP(
                -beta1*dose^1)]

The parameter betas are restricted to be positive
    
```

Dependent variable = incidence  
 Independent variable = dose

Total number of observations = 3  
 Total number of records with missing values = 0  
 Total number of parameters in model = 2  
 Total number of specified parameters = 0  
 Degree of polynomial = 1  
 Maximum number of iterations = 250  
 Relative Function Convergence has been set to: 2.22045e-016  
 Parameter Convergence has been set to: 1.49012e-008

\*\*\*\* We are sorry but Relative Function and Parameter Convergence \*\*\*\*  
 \*\*\*\* are currently unavailable in this model. Please keep checking \*\*\*\*  
 \*\*\*\* the web sight for model updates which will eventually \*\*\*\*  
 \*\*\*\* incorporate these convergence criterion. Default values used. \*\*\*\*

Default Initial Parameter Values

Background = 0.0659119  
 Beta(1) = 0.00855407

Asymptotic Correlation Matrix of Parameter Estimates

|            | Background | Beta(1) |
|------------|------------|---------|
| Background | 1          | -0.56   |
| Beta(1)    | -0.56      | 1       |

Parameter Estimates

| Variable   | Estimate   | Std. Err. | 95.0% Wald Confidence Interval |                   |
|------------|------------|-----------|--------------------------------|-------------------|
|            |            |           | Lower Conf. Limit              | Upper Conf. Limit |
| Background | 0.0980803  | *         | *                              | *                 |
| Beta(1)    | 0.00807004 | *         | *                              | *                 |

\* - Indicates that this value is not calculated.

Analysis of Deviance Table

| Model         | Log(likelihood) | # Param's | Deviance | Test d.f. | P-value |
|---------------|-----------------|-----------|----------|-----------|---------|
| Full model    | -68.0892        | 3         |          |           |         |
| Fitted model  | -68.1887        | 2         | 0.198975 | 1         | 0.6555  |
| Reduced model | -99.8132        | 1         | 63.4479  | 2         | <.0001  |

**AIC: 140.377**

Goodness of Fit

| Dose     | Est._Prob. | Expected | Observed | Size | Scaled Residual |
|----------|------------|----------|----------|------|-----------------|
| 0.0000   | 0.0981     | 4.904    | 5.000    | 50   | 0.046           |
| 100.2000 | 0.5982     | 28.116   | 27.000   | 47   | -0.332          |
| 210.7000 | 0.8353     | 39.259   | 40.000   | 47   | 0.291           |

**Chi^2 = 0.20      d.f. = 1      P-value = 0.6569**

Benchmark Dose Computation

Specified effect = 0.1

Risk Type = Extra risk

Confidence level = 0.95

**BMD = 13.0558**

**BMDL = 10.2947**



BMDU = 16.9865

Taken together, (10.2947, 16.9865) is a 90 % two-sided confidence interval for the BMD

**Multistage Cancer Slope Factor = 0.00971371**

## APPENDIX F. COMPARATIVE CANCER IUR BASED ON FEMALE MICE DATA

Using the male B6C3F<sub>1</sub> mouse data from a 2-year inhalation exposure study (Mennear et al., 1988; NTP, 1986), the recommended cancer IURs are  $7 \times 10^{-9} (\mu\text{g}/\text{m}^3)^{-1}$  and  $5 \times 10^{-9} (\mu\text{g}/\text{m}^3)^{-1}$  for the development of liver and lung cancer, respectively, based on the mean for the GST-T1<sup>+/+</sup> population. These values were derived using a tissue-specific GST metabolism dose metric with allometric scaling. The combined human equivalent IUR values for both tumor types is  $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$ . As described in detail below, the resulting combined human equivalent IUR values for both tumors did not differ appreciably by gender.

BMD<sub>10</sub> and BMDL<sub>10</sub> refer to the model-predicted dose (and its lower 95% confidence limit) associated with 10% extra risk for the combined incidence of adenoma and carcinoma of the liver or lung of female B6C3F<sub>1</sub> mice inhaling dichloromethane for 2 years (Mennear et al., 1988; NTP, 1986) (Table F-1).

**Table F-1. Incidence data for liver and lung tumors and internal doses based on GST metabolism dose metrics in female B6C3F<sub>1</sub> mice exposed to dichloromethane via inhalation for 2 years**

| Sex, tumor type            | BW (g) | External dichloromethane concentration (ppm) | Mouse tumor incidence  | Mouse internal tissue dose <sup>a</sup> | Mouse whole body metabolism dose <sup>b</sup> |
|----------------------------|--------|----------------------------------------------|------------------------|-----------------------------------------|-----------------------------------------------|
| Female, liver <sup>c</sup> | –      | 0                                            | 3/47 (6%) <sup>d</sup> | 0                                       | 0                                             |
|                            | 30.0   | 2,000                                        | 16/46 (35%)            | 2,453.2                                 | 104.0                                         |
|                            | 29.0   | 4,000                                        | 40/46 (87%)            | 5,120.0                                 | 217.0                                         |
| Female, lung <sup>e</sup>  | –      | 0                                            | 3/45 (6%) <sup>d</sup> | 0                                       | 0                                             |
|                            | 30.0   | 2,000                                        | 30/46 (65%)            | 493.0                                   | 104.0                                         |
|                            | 29.0   | 4,000                                        | 41/46 (89%)            | 1,021.8                                 | 217.0                                         |

<sup>a</sup>For liver tumors: mg dichloromethane metabolized via GST pathway/L liver tissue/d from 6 hrs/d, 5 d/wk exposure; for lung tumors: mg dichloromethane metabolized via GST pathway/L lung tissue/d from 6 hrs/day, 5 d/wk exposure.

<sup>b</sup>Based on the sum of dichloromethane metabolized via the GST pathway in the lung plus the liver, normalized to total BW (i.e., [lung GST metabolism (mg/d) + liver GST metabolism (mg/d)]/kg BW). Units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

<sup>c</sup>Hepatocellular carcinoma or adenoma. Mice dying prior to 52 wks were excluded from the denominators.

<sup>d</sup>Statistically significant increasing trend (by incidental and life-table tests;  $p \leq 0.01$ ).

<sup>e</sup>Bronchoalveolar carcinoma or adenoma. Mice dying prior to 52 wks were excluded from the denominators.

Sources: Mennear et al. (1988); NTP (1986).

Multistage models were fit to the female mouse internal tissue doses of dichloromethane metabolized by the GST pathway and incidences for animals with liver tumors observed at the

time of death. The predicted  $BMD_{10}$  and  $BMDL_{10}$  for the liver and lung tumor incidence data are shown in Table F-2.

**Table F-2. BMD modeling results and tumor risk factors associated with 10% extra risk for liver and lung tumors in female B6C3F<sub>1</sub> mice exposed by inhalation to dichloromethane for 2 years, based on liver-specific GST metabolism and whole body GST metabolism dose metrics**

| Internal dose metric <sup>a</sup> |               | BMDS model <sup>b</sup> | $\chi^2$ goodness of fit <i>p</i> -value | Mouse BMD <sub>10</sub> <sup>c</sup> | Mouse BMDL <sub>10</sub> <sup>c</sup> | Allometric-scaled human BMDL <sub>10</sub> <sup>d</sup> | Tumor risk factor <sup>e</sup> |                       |
|-----------------------------------|---------------|-------------------------|------------------------------------------|--------------------------------------|---------------------------------------|---------------------------------------------------------|--------------------------------|-----------------------|
|                                   |               |                         |                                          |                                      |                                       |                                                         | Scaling = 1.0                  | Allometric-scaled     |
| Liver-specific                    | Female, liver | MS (0,2)                | 0.53                                     | 1,224.1                              | 659.7                                 | 94.2                                                    | $1.52 \times 10^{-4}$          | $1.06 \times 10^{-3}$ |
|                                   | Female, lung  | MS (0,1)                | 0.87                                     | 51.2                                 | 40.7                                  | 5.8                                                     | $2.46 \times 10^{-3}$          | $1.72 \times 10^{-2}$ |
| Whole body                        | Female, liver | MS (0,2)                | 0.53                                     | 51.9                                 | 28.0                                  | 4.0                                                     | –                              | $2.50 \times 10^{-2}$ |
|                                   | Female, lung  | MS (0,1)                | 0.88                                     | 10.8                                 | 8.6                                   | 1.2                                                     | –                              | $8.14 \times 10^{-2}$ |

<sup>a</sup>Liver specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue per d; whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

<sup>b</sup>The multistage (MS) model in EPA BMDS version 2.0 was fit to the mouse dose-response data shown in Table 5-17 using internal dose metrics calculated with the mouse PBPK model. Numbers in parentheses indicate: (1) the number of dose groups dropped in order to obtain an adequate fit, and (2) the degree polynomial of the model.

<sup>c</sup>BMD<sub>10</sub> and BMDL<sub>10</sub> refer to the BMD-model-predicted mouse internal dose and its 95% lower confidence limit, associated with a 10% extra risk for the incidence of tumors.

<sup>d</sup>Mouse BMDL<sub>10</sub> divided by  $(BW_{\text{human}}/BW_{\text{mouse}})^{0.25} = 7$ .

<sup>e</sup>Dichloromethane tumor risk factor (extra risk per unit internal dose) derived by dividing the BMR (0.1) by the mouse BMDL<sub>10</sub> and by the allometric-scaled human BMDL<sub>10</sub>, for the scaling = 1.0 and allometric-scaled risk factors, respectively.

A probabilistic PBPK model for dichloromethane in humans adapted from David et al. (2006) (see Appendix B) was used with Monte Carlo sampling to calculate distributions of internal lung, liver, or blood doses associated with chronic unit inhalation ( $1 \mu\text{g}/\text{m}^3$ ) exposures. The model was then executed by using the external unit exposure as input, and the resulting human equivalent internal dose was recorded. This process was repeated for 10,000 iterations to generate a distribution of human internal doses. The resulting distribution of IURs shown in Table F-3 was derived by multiplying the human internal dose tumor risk factor (in units of reciprocal internal dose) by the respective distributions of human average daily internal dose resulting from a chronic unit inhalation exposure of  $1 \mu\text{g}/\text{m}^3$  dichloromethane. Risk estimates were slightly higher for liver tumors and essentially equivalent for lung tumors in males compared to females.

**Table F-3. IURs for dichloromethane based on PBPK model-derived internal liver and lung doses in B6C3F<sub>1</sub> female mice exposed via inhalation for 2 years, based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype**

| Internal dose metric and scaling factor <sup>a</sup> | Population genotype <sup>b</sup> | Tumor type | Human tumor risk factor <sup>c</sup> | Distribution of human internal dichloromethane doses from 1 µg/m <sup>3</sup> exposure <sup>d</sup> |                             |                             | Resulting candidate human IUR <sup>e</sup> (µg/m <sup>3</sup> ) <sup>-1</sup> |                             |                             |
|------------------------------------------------------|----------------------------------|------------|--------------------------------------|-----------------------------------------------------------------------------------------------------|-----------------------------|-----------------------------|-------------------------------------------------------------------------------|-----------------------------|-----------------------------|
|                                                      |                                  |            |                                      | Mean                                                                                                | 95 <sup>th</sup> percentile | 99 <sup>th</sup> percentile | Mean                                                                          | 95 <sup>th</sup> percentile | 99 <sup>th</sup> percentile |
| Tissue-specific, allometric-scaled                   | GST-T1 <sup>+/+</sup>            | Liver      | 1.06 × 10 <sup>-3</sup>              | 6.61 × 10 <sup>-6</sup>                                                                             | 2.21 × 10 <sup>-5</sup>     | 4.47 × 10 <sup>-5</sup>     | 7.0 × 10 <sup>-9</sup>                                                        | 2.4 × 10 <sup>-8</sup>      | 4.7 × 10 <sup>-8</sup>      |
|                                                      | GST-T1 <sup>+/+</sup>            | Lung       | 1.72 × 10 <sup>-2</sup>              | 3.89 × 10 <sup>-7</sup>                                                                             | 1.24 × 10 <sup>-6</sup>     | 2.42 × 10 <sup>-6</sup>     | 6.7 × 10 <sup>-9</sup>                                                        | 2.1 × 10 <sup>-8</sup>      | 4.2 × 10 <sup>-8</sup>      |
|                                                      | Mixed                            | Liver      | 1.06 × 10 <sup>-3</sup>              | 3.71 × 10 <sup>-6</sup>                                                                             | 1.43 × 10 <sup>-5</sup>     | 3.03 × 10 <sup>-5</sup>     | 3.9 × 10 <sup>-9</sup>                                                        | 1.5 × 10 <sup>-8</sup>      | 3.2 × 10 <sup>-8</sup>      |
|                                                      | Mixed                            | Lung       | 1.72 × 10 <sup>-2</sup>              | 2.20 × 10 <sup>-7</sup>                                                                             | 8.06 × 10 <sup>-7</sup>     | 1.69 × 10 <sup>-6</sup>     | 3.8 × 10 <sup>-9</sup>                                                        | 1.4 × 10 <sup>-8</sup>      | 2.9 × 10 <sup>-8</sup>      |
| Tissue-specific, scaling = 1.0                       | GST-T1 <sup>+/+</sup>            | Liver      | 1.52 × 10 <sup>-4</sup>              | 6.61 × 10 <sup>-6</sup>                                                                             | 2.21 × 10 <sup>-5</sup>     | 4.47 × 10 <sup>-5</sup>     | 1.0 × 10 <sup>-9</sup>                                                        | 3.4 × 10 <sup>-9</sup>      | 6.8 × 10 <sup>-9</sup>      |
|                                                      | GST-T1 <sup>+/+</sup>            | Lung       | 2.46 × 10 <sup>-3</sup>              | 3.89 × 10 <sup>-7</sup>                                                                             | 1.24 × 10 <sup>-6</sup>     | 2.42 × 10 <sup>-6</sup>     | 9.6 × 10 <sup>-10</sup>                                                       | 3.1 × 10 <sup>-9</sup>      | 6.0 × 10 <sup>-9</sup>      |
|                                                      | Mixed                            | Liver      | 1.52 × 10 <sup>-4</sup>              | 3.71 × 10 <sup>-6</sup>                                                                             | 1.43 × 10 <sup>-5</sup>     | 3.03 × 10 <sup>-5</sup>     | 5.6 × 10 <sup>-10</sup>                                                       | 2.2 × 10 <sup>-9</sup>      | 4.6 × 10 <sup>-9</sup>      |
|                                                      | Mixed                            | Lung       | 2.46 × 10 <sup>-3</sup>              | 2.20 × 10 <sup>-7</sup>                                                                             | 8.06 × 10 <sup>-7</sup>     | 1.69 × 10 <sup>-6</sup>     | 5.4 × 10 <sup>-10</sup>                                                       | 2.0 × 10 <sup>-9</sup>      | 4.2 × 10 <sup>-9</sup>      |
| Whole-body, allometric-scaled                        | GST-T1 <sup>+/+</sup>            | Liver      | 2.50 × 10 <sup>-2</sup>              | 1.80 × 10 <sup>-7</sup>                                                                             | 6.38 × 10 <sup>-7</sup>     | 1.41 × 10 <sup>-6</sup>     | 4.5 × 10 <sup>-9</sup>                                                        | 1.6 × 10 <sup>-8</sup>      | 3.5 × 10 <sup>-8</sup>      |
|                                                      | GST-T1 <sup>+/+</sup>            | Lung       | 8.14 × 10 <sup>-2</sup>              | 1.80 × 10 <sup>-7</sup>                                                                             | 6.38 × 10 <sup>-7</sup>     | 1.41 × 10 <sup>-6</sup>     | 1.5 × 10 <sup>-8</sup>                                                        | 5.2 × 10 <sup>-8</sup>      | 1.2 × 10 <sup>-7</sup>      |
|                                                      | Mixed                            | Liver      | 2.50 × 10 <sup>-2</sup>              | 1.01 × 10 <sup>-7</sup>                                                                             | 4.00 × 10 <sup>-7</sup>     | 9.43 × 10 <sup>-7</sup>     | 2.5 × 10 <sup>-9</sup>                                                        | 1.0 × 10 <sup>-8</sup>      | 2.4 × 10 <sup>-8</sup>      |
|                                                      | Mixed                            | Lung       | 8.14 × 10 <sup>-2</sup>              | 1.01 × 10 <sup>-7</sup>                                                                             | 4.00 × 10 <sup>-7</sup>     | 9.43 × 10 <sup>-7</sup>     | 8.2 × 10 <sup>-9</sup>                                                        | 3.3 × 10 <sup>-8</sup>      | 7.7 × 10 <sup>-8</sup>      |

<sup>a</sup>Tissue specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue (liver or lung, respectively, for liver and lung tumors) per d; whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

<sup>b</sup>GST-T1<sup>+/+</sup> = homozygous, full enzyme activity; mixed = population reflecting estimated frequency of genotypes in current U.S. population: 20% GST-T<sup>-/-</sup>, 48% GST-T1<sup>+/-</sup>, and 32% GST-T1<sup>+/+</sup> (Haber et al., 2002).

<sup>c</sup>Dichloromethane tumor risk factor (extra risk per unit internal dose) derived by dividing the BMR (0.1) by the allometric-scaled human BMDL<sub>10</sub> or by the mouse BMDL<sub>10</sub> (from Table 5-18) for the allometric-scaled and scaling = 1.0 risk factors, respectively.

<sup>d</sup>Mean, 95<sup>th</sup>, and 99<sup>th</sup> percentile of the human PBPK model-derived probability distribution of daily average internal dichloromethane dose resulting from chronic exposure to 1 µg/m<sup>3</sup> (0.00029 ppm).

<sup>e</sup>Derived by multiplying the dichloromethane tumor risk factor by the PBPK model-derived probabilistic internal doses from daily exposure to 1 µg/m<sup>3</sup>.

For the female mouse, the combined human equivalent IUR values for both tumor types is  $1 \times 10^{-8} (\mu\text{g}/\text{m}^3)^{-1}$  in the most sensitive (GST-T1<sup>+/+</sup>) population (Table F-4), which is the same value that was obtained using the male mouse data.

**Table F-4. Upper bound estimates of combined human IURs for liver and lung tumors resulting from lifetime exposure to 1 µg/m<sup>3</sup> dichloromethane based on liver-specific GST metabolism and whole body metabolism dose metrics, by population genotype, using female mouse data for derivation of risk factors**

| Internal dose metric and scaling factor <sup>a</sup> | Population genotype <sup>b</sup> | Tumor site    | Upper bound IUR <sup>c</sup> | Central tendency IUR <sup>d</sup> | Variance of tissue-specific tumor risk <sup>e</sup> | Combined tumor risk SD <sup>f</sup> | Upper bound on combined tumor risk <sup>g</sup> (µg/m <sup>3</sup> ) <sup>-1</sup> |
|------------------------------------------------------|----------------------------------|---------------|------------------------------|-----------------------------------|-----------------------------------------------------|-------------------------------------|------------------------------------------------------------------------------------|
| Tissue-specific, allometric-scaled                   | GST-T1 <sup>+/+</sup>            | Liver         | $7.0 \times 10^{-9}$         | $3.8 \times 10^{-9}$              | $3.87 \times 10^{-18}$                              |                                     |                                                                                    |
|                                                      |                                  | Lung          | $6.7 \times 10^{-9}$         | $5.3 \times 10^{-9}$              | $7.12 \times 10^{-19}$                              |                                     |                                                                                    |
|                                                      |                                  | Liver or lung |                              | $9.1 \times 10^{-9}$              |                                                     | $2.1 \times 10^{-9}$                | $1.3 \times 10^{-8}$                                                               |
|                                                      | Mixed                            | Liver         | $3.9 \times 10^{-9}$         | $2.1 \times 10^{-9}$              | $1.22 \times 10^{-18}$                              |                                     |                                                                                    |
|                                                      |                                  | Lung          | $3.8 \times 10^{-9}$         | $3.0 \times 10^{-9}$              | $2.28 \times 10^{-19}$                              |                                     |                                                                                    |
|                                                      |                                  | Liver or lung |                              | $5.2 \times 10^{-9}$              |                                                     | $1.2 \times 10^{-9}$                | $7.1 \times 10^{-9}$                                                               |
| Tissue-specific, scaling = 1.0                       | GST-T1 <sup>+/+</sup>            | Liver         | $1.0 \times 10^{-9}$         | $5.4 \times 10^{-10}$             | $7.89 \times 10^{-20}$                              |                                     |                                                                                    |
|                                                      |                                  | Lung          | $9.6 \times 10^{-10}$        | $7.6 \times 10^{-10}$             | $1.42 \times 10^{-21}$                              |                                     |                                                                                    |
|                                                      |                                  | Liver or lung |                              | $1.3 \times 10^{-9}$              |                                                     | $3.1 \times 10^{-10}$               | $1.8 \times 10^{-9}$                                                               |
|                                                      | Mixed                            | Liver         | $5.6 \times 10^{-10}$        | $3.0 \times 10^{-10}$             | $2.48 \times 10^{-20}$                              |                                     |                                                                                    |
|                                                      |                                  | Lung          | $5.4 \times 10^{-10}$        | $4.3 \times 10^{-10}$             | $4.54 \times 10^{-21}$                              |                                     |                                                                                    |
|                                                      |                                  | Liver or lung |                              | $7.3 \times 10^{-10}$             |                                                     | $1.7 \times 10^{-10}$               | $1.0 \times 10^{-9}$                                                               |
| Whole-body, allometric-scaled                        | GST-T1 <sup>+/+</sup>            | Liver         | $4.5 \times 10^{-9}$         | $2.4 \times 10^{-9}$              | $1.59 \times 10^{-18}$                              |                                     |                                                                                    |
|                                                      |                                  | Lung          | $1.5 \times 10^{-8}$         | $1.2 \times 10^{-8}$              | $4.11 \times 10^{-18}$                              |                                     |                                                                                    |
|                                                      |                                  | Liver or lung |                              | $1.4 \times 10^{-8}$              |                                                     | $2.4 \times 10^{-9}$                | $1.8 \times 10^{-8}$                                                               |
|                                                      | Mixed                            | Liver         | $2.5 \times 10^{-9}$         | $1.4 \times 10^{-9}$              | $5.00 \times 10^{-19}$                              |                                     |                                                                                    |
|                                                      |                                  | Lung          | $8.2 \times 10^{-9}$         | $6.7 \times 10^{-9}$              | $1.29 \times 10^{-18}$                              |                                     |                                                                                    |
|                                                      |                                  | Liver or lung |                              | $7.9 \times 10^{-9}$              |                                                     | $1.3 \times 10^{-9}$                | $1.0 \times 10^{-8}$                                                               |

<sup>a</sup>Tissue specific dose units = mg dichloromethane metabolized via GST pathway per liter tissue (liver or lung, respectively, for liver and lung tumors) per d; whole-body dose units = mg dichloromethane metabolized via GST pathway in lung and liver/kg-d.

<sup>b</sup>GST-T1<sup>+/+</sup> = homozygous, full enzyme activity); mixed = population reflecting estimated frequency of genotypes in current U.S. population: 20% GST-T<sup>-/-</sup>, 48% GST-T1<sup>+/+</sup>, and 32% GST-T1<sup>+/+</sup> (Haber et al., 2002).

<sup>c</sup>Estimated at the human equivalent BMDL<sub>10</sub> (0.1/BMDL<sub>10</sub>) (see Table F-2).

<sup>d</sup>Estimated at the human equivalent BMD<sub>10</sub> (0.1/BMD) (see Table F-2).

<sup>e</sup>Calculated as the square of the difference of the upper bound and central tendency IURs divided by the *t* statistic, 1.645.

<sup>f</sup>Calculated as the square root of the sum of the variances for liver and lung tumors.

<sup>g</sup>Calculated as the product of the cumulative tumor risk SD and the *t* statistic, 1.645, added to the sum of central tendency IURs.



## APPENDIX G. COMPARATIVE CANCER IUR BASED ON BENIGN MAMMARY GLAND TUMORS IN RATS

Data for mammary gland tumors in male and female F344 rats following exposure to airborne dichloromethane were used to develop a comparative IUR for dichloromethane (Mennear et al., 1988; NTP, 1986) (Table G-1). Significantly increased incidences of mammary gland or subcutaneous tissue adenoma, fibroadenomas, or fibromas were observed in male rats at 4,000 ppm, while mammary gland adenomas or fibroadenomas were increased in female rats exposed 6 hours/day, 5 days/week for 2 years at concentrations  $\geq 1,000$  ppm. Significant decreases in survival were observed in the treated groups of both sexes. The at-risk study populations (represented by the denominators in the incidence data) were determined by excluding all animals dying prior to 52 weeks.

**Table G-1. Incidence data for mammary gland tumors and internal doses based on different dose metrics in male and female F344 rats exposed to dichloromethane via inhalation for 2 years**

| Sex    | BW (g) | External dichloromethane concentration (ppm) | Rat tumor incidence <sup>a</sup> | Rat internal dose, AUC in blood <sup>b</sup> |
|--------|--------|----------------------------------------------|----------------------------------|----------------------------------------------|
| Male   | –      | 0                                            | 1/50 (2%) <sup>c</sup>           | 0                                            |
|        | 390.5  | 1,000                                        | 1/50 (2%)                        | 93.3                                         |
|        | 385.2  | 2,000                                        | 4/50 (8%)                        | 196.4                                        |
|        | 384.8  | 4,000                                        | 9/50 (18%)                       | 403.5                                        |
| Female | –      | 0                                            | 6/49 (12%) <sup>c</sup>          | 0                                            |
|        | 245.5  | 1,000                                        | 13/50 (26%)                      | 93.3                                         |
|        | 244.3  | 2,000                                        | 14/50 (28%)                      | 196.2                                        |
|        | 242.2  | 4,000                                        | 23/50 (46%)                      | 403.0                                        |

<sup>a</sup>Male tumors include mammary gland or subcutaneous tissue adenoma, fibroadenomas, or fibroma. Female tumors include mammary gland adenoma or fibroadenomas. Rats dying prior to 52 wks were excluded from the denominators.


<sup>b</sup>Average daily AUC for dichloromethane in slowly perfused tissue (mg  $\times$  hr/L) (see text for rationale for using this dose metric).

<sup>c</sup>Statistically significant increasing trend ( $p \leq 0.01$ ).

Sources: Mennear et al. (1988); NTP (1986).

The rat PBPK model of Andersen et al. (1991) (modified as described in Appendix D) was used to simulate inhalation exposures of 6 hours/day, 5 days/week and calculate long-term daily average internal doses for the 2-year bioassays (Mennear et al., 1988; NTP, 1986). Study-, group-, and sex-specific mean BWs for rats were used. The modified PBPK model did not include a compartment for the mammary gland nor did it account for metabolism of dichloromethane occurring in the mammary gland. The selected internal dose metric for

mammary gland tumors was the average daily AUC for dichloromethane in slowly perfused tissue; the mammary gland and the regions around it consist primarily of fatty tissue, which is slowly perfused tissue. The role of CYP- or GST-mediated metabolism in the mammary gland is uncertain. GST-T1 (Lehmann and Wagner, 2008) and CYP2E1 (El-Rayes et al., 2003; Hellmold et al., 1998) expression has been detected in human mammary tissue, and it is also possible that some metabolites enter systemic circulation from the liver and lung where they are formed. Figure G-1 shows the comparison between inhalation external and internal doses in the liver and lung, respectively, using this dose metric for the rat and the human.



Average daily doses were calculated from simulated rat exposures of 6 hours/day, 5 days/week, while simulated human exposures were continuous. The GST metabolism rate in the human population for the mixed GST-T1 group (+/+, +/-, and -/-) in the current U.S. population was estimated as the mean of a simulated sample of 3,000 individuals at each exposure concentration, based on GST-T1 polymorphism data from Haber et al. (2002). The results for the GST-T1 +/- and +/+ subpopulations were then calculated as the means of the subsets of the mixed population sample with the respective genotypes.

**Figure G-1. PBPK model-derived internal doses (daily average AUC for dichloromethane in blood) in rats and humans and their associated external exposures (ppm) used for the derivation of cancer IURs based on mammary tumors in rats.**

The multistage model was fit to the rat mammary gland tumor incidence and PBPK model-derived internal dose data to derive rat internal BMD<sub>10</sub> and BMDL<sub>10</sub> values associated with 10% extra risk (Table G-2).

**Table G-2. BMD modeling results associated with 10% extra risk for mammary gland tumors in F344 rats exposed by inhalation to dichloromethane for 2 years based on AUC for dichloromethane in slowly perfused tissue**

| Sex    | Tumor type                                                             | BMDS model <sup>a</sup> | $\chi^2$ goodness of fit <i>p</i> -value | Rat BMD <sub>10</sub> <sup>c</sup> | Rat BMDL <sub>10</sub> <sup>c</sup> | Tumor risk factor <sup>d</sup> |
|--------|------------------------------------------------------------------------|-------------------------|------------------------------------------|------------------------------------|-------------------------------------|--------------------------------|
| Male   | Mammary gland or subcutaneous tissue adenoma, fibroadenoma, or fibroma | MS (0,1)                | 0.53                                     | 275.3                              | 172.3                               | $5.80 \times 10^{-4}$          |
| Female | Mammary gland adenoma or fibroadenoma                                  | MS (0,1)                | 0.71                                     | 91.0                               | 61.5                                | $1.63 \times 10^{-3}$          |

<sup>a</sup>The multistage (MS) model in EPA BMDS version 2.0 was fit to each of the two sets of rat dose-response data shown in Table G-1 using internal dose metrics calculated with the rat PBPK model. Numbers in parentheses indicate: (1) the number of dose groups dropped in order to obtain an adequate fit, and (2) the degree polynomial of the model.

<sup>c</sup>BMD<sub>10</sub> and BMDL<sub>10</sub> refer to the BMD-model-predicted rat internal dose (average daily AUC for dichloromethane in slowly perfused tissue [mg × hr/L]) and its 95% lower confidence limit associated with a 10% extra risk for the incidence of tumors.

<sup>d</sup>Dichloromethane tumor risk factor (extra risk per average daily AUC for dichloromethane in slowly perfused tissue [mg × hr/L]) was derived by dividing the BMR (0.1) by the rat BMDL<sub>10</sub>. The rat BMDL<sub>10</sub> is assumed to be equivalent to human BMDL<sub>10</sub>; humans exposed to the same average daily AUC for dichloromethane in slowly perfused tissue as rats will have the same risks for mammary tumors.

Rat mammary gland tumor risk factors (extra risk per unit internal dose) were calculated by dividing 0.1 by the rat internal BMDL<sub>10</sub>. Because this risk factor is based on the internal concentration of dichloromethane rather than a rate of reaction, it is assumed that the human risk factor is equal to that of the rat (i.e., that humans exposed for a 70-year lifetime to the same weekly average AUC of dichloromethane will have the same risk as rats exposed for 2 years). The human PBPK model (David et al., 2006) was used to calculate a distribution of human average daily AUCs for dichloromethane in slowly perfused tissue resulting from chronic inhalation exposure to a unit concentration of 1 µg/m<sup>3</sup> dichloromethane (0.00029 ppm). The distribution of IURs for mammary gland tumors was generated by multiplying the human tumor risk factor for each sex by the distribution of internal doses from chronic human exposure to 1 µg/m<sup>3</sup> dichloromethane. Because this analysis is not based on the assumption that either metabolic pathway is or is not influencing the cancer risk, this distribution was derived by using weights reflecting the estimated frequency of GST-T1 genotypes in the current U.S. population (20% GST-T1<sup>-/-</sup>, 48% GST-T1<sup>+/-</sup>, and 32% GST-T1<sup>+/+</sup>). As shown in Table G-3, the mean human IUR based on mammary gland tumors in rats is  $4 \times 10^{-8}$  and  $1 \times 10^{-7}$  (µg/m<sup>3</sup>)<sup>-1</sup> based on

male and female rat-derived risk factors, respectively. Identical values were obtained using slowly perfused tissue as the internal dose metric.

**Table G-3. IURs for dichloromethane based on benign mammary tumors and PBPK model-derived internal doses in F344N rats exposed via inhalation for 2 years based on AUC for dichloromethane in slowly perfused tissue dose metric**

| Sex, tumor type | Human tumor risk factor <sup>a</sup> | Distribution of human internal dichloromethane doses from 1 µg/m <sup>3</sup> exposure <sup>b</sup> |                             |                             | Resulting candidate human IUR <sup>c</sup> |                             |                             |
|-----------------|--------------------------------------|-----------------------------------------------------------------------------------------------------|-----------------------------|-----------------------------|--------------------------------------------|-----------------------------|-----------------------------|
|                 |                                      | Mean                                                                                                | 95 <sup>th</sup> percentile | 99 <sup>th</sup> percentile | Mean                                       | 95 <sup>th</sup> percentile | 99 <sup>th</sup> percentile |
| Male            | $5.80 \times 10^{-4}$                | $6.81 \times 10^{-5}$                                                                               | $1.07 \times 10^{-4}$       | $1.33 \times 10^{-4}$       | $3.95 \times 10^{-8}$                      | $6.19 \times 10^{-8}$       | $7.69 \times 10^{-8}$       |
| Female          | $1.63 \times 10^{-3}$                | $6.81 \times 10^{-5}$                                                                               | $1.07 \times 10^{-4}$       | $1.33 \times 10^{-4}$       | $1.11 \times 10^{-7}$                      | $1.74 \times 10^{-7}$       | $2.16 \times 10^{-7}$       |

<sup>a</sup>Dichloromethane tumor risk factor (extra risk per average daily AUC for dichloromethane in slowly perfused tissue [mg × hr/L]) was derived by dividing the BMR (0.1) by the rat BMDL<sub>10</sub> (from Table G-2). The rat BMDL<sub>10</sub> is assumed to be equivalent to human BMDL<sub>10</sub>; humans exposed to the same average daily AUC for dichloromethane in slowly perfused tissue as rats will have the same risks for mammary tumors.

<sup>b</sup>Mean, 95<sup>th</sup>, and 99<sup>th</sup> percentile of the human PBPK model-derived probability distribution of daily average internal dichloromethane dose (average daily AUC for dichloromethane in slowly perfused tissue [mg × hr/L]) resulting from chronic inhalation exposure to a unit concentration of 1 µg/m<sup>3</sup> (0.00029 ppm), based on a distribution of GST-T1 genotypes that reflects the frequency distribution in the current U.S. population (Haber et al., 2002).

<sup>c</sup>Mean, 95<sup>th</sup>, and 99<sup>th</sup> percentile of a distribution of human IURs (extra risk per µg/m<sup>3</sup>) derived by multiplying the dichloromethane tumor risk factor by the PBPK model-derived probabilistic distribution of human internal dichloromethane doses from unit dichloromethane inhalation exposure.

## APPENDIX H: SOURCE CODE AND COMMAND FILES FOR DICHLOROMETHANE PBPK MODELS

The following is a copy of the primary acslXtreme code (.csl file for implementation under acslXtreme v. 2.4.2.1) used for the dichloromethane simulations. Portions of the code which had been commented out (i.e., unused code) were deleted for brevity. Note that most parameters are set in the subsequent script/m-files, used to specify simulations and call the .csl code.

While the code indicates modifications by G Diamond and M Lumpkin (employees of the EPA contractor, SRC, Inc.), these are changes from the version of the code as received by them to bring it into alignment with the version as described in the publications of Marino et al. (2006) and David et al. (2006) (changes noted by comments). Other changes allow for the calculation of various metrics to improve time-efficiency for computational convergence; for example, calculating an average AUC over only the last week of simulated exposure requires a much shorter overall simulated time for the calculation to give the steady-state or very-long-term average, when the approach to steady-state or repeating periodic solution occurs over the first days or weeks of exposure. Thus, the code exactly replicates the published models when the published parameter values are used.

PROGRAM: DCM\_2010\_EPA.csl

! Code from Reitz et al. 1997, Addressing Data Needs for Methylene Chloride with  
! Physiologically Based Pharmacokinetic Modeling (Appendix I)  
! Translated by GDiamond (05/2004); DCM.CSL.Reitz.  
! Revision date 18-Dec-96 by RHR (Peak moved to dynamic)  
! Modified by GDiamond (08/2004) based on Andersen et al. 1987 (TAP 87:185-205)  
! and 1991 (TAP 108:14-27):  
! Deleted brain compartment  
! Added lung compartment, with lung metabolism  
! Adjusted metabolism parameter values to match Andersen et al. 1987  
! Adjusted physiological parameters to match Andersen et al. 1987  
! Modified by M Lumpkin (11/2005) (MHL) to include extrahepatic MFO metabolism and  
! CO kinetics in blood  
! Additional comments and changes by Paul Schlosser (PS), U.S. EPA (7/2008 - 2/2010)  
! Removed unused (legacy) code bits, added comments, PS 9/2009

INITIAL

! Simulation, T=hour  
NSTP=1000           ! Initial integration cycle length at CINT/1000  
! MERROR ALU=0.00001   ! Error tolerance for Gear

CONSTANT POINTS=96.0 ! Number of points in plot  
 CINTERVAL CINT=0.1 ! Communication interval  
 CONSTANT TEND=25.0 ! Termination time (hr)  
 TSTOP=TEND  
 ! Initial values for 250 g rat  
  
 ! Body masses and fractional volumes  
 CONSTANT BW=0.25 ! Body weight (kg) MHL  
 CONSTANT VBL2C=0.059 ! Fractional volume of blood MHL  
 CONSTANT VFC=0.07 ! Fractional volume fat  
 CONSTANT VLC=0.04 ! Fractional volume of liver  
 CONSTANT VRC=0.05 ! Fractional volume of rapidly-perfused tissues  
 CONSTANT VSC=0.75 ! Fractional volume of slowly-perfused tissues \*\*  
 CONSTANT VLUC=0.0115 ! Allometric scaling factor for lung volume \*\*  
  
 ! Tissue masses (kg)  
 VTOT=VFC+VLC+(VLUC/(BW\*\*0.01))+VRC+VSC  
 ! Total volume fractions constrained to sum to 0.9215 (7.85% carcass) MHL  
 VBL2=VBL2C\*BW ! Blood  
 VF=VFC\*BW\*0.9215/VTOT ! Fat  
 VL=VLC\*BW\*0.9215/VTOT ! Liver  
 VR=VRC\*BW\*0.9215/VTOT ! Rapidly-perfused tissue  
 VS=VSC\*BW\*0.9215/VTOT ! Slowly-perfused tissue  
 VLU=VLUC\*(BW\*\*0.99)\*0.9215/VTOT ! Lung  
  
 ! Flow constants and fractions  
 CONSTANT VPR=0.42 ! Ventilation/perfusion ratio (for QP calc) MHL  
 CONSTANT DSPC=0.15 ! Fractional lung dead space MHL  
 CONSTANT QCC=15.0 ! Allometric constant for cardiac output  
 CONSTANT QCCR=10.0 ! For 'resting' period PMS 8/4/09  
 CONSTANT QCSW=0 ! Set to 1 to use QCCR (resting cardiac rate) up to TCHNG  
 CONSTANT QLC=0.20 ! Fractional flow to liver  
 CONSTANT QFC=0.09 ! Fractional flow to fat  
 CONSTANT QSC=0.15 ! Fractional flow to slowly-perfused tissue  
 CONSTANT QRC=0.56 ! Fractional flow to rapidly-perfused tissue \*\* MHL  
  
 ! Flow rates (L/hr)  
  
 IF (QCSW) THEN  
 QC=QCCR\*BW\*\*0.74 ! Cardiac output if resting, PMS 8/4/09  
 ELSE

$QC=QCC*BW**0.74$  ! Cardiac output if not resting, PMS 8/4/09  
 ENDIF  
 $QP=QC*VPR$  ! Alveolar ventilation rate MHL  
 $QPDP=QP/DL/PAIR$  ! MHL  
 $QTOT=QFC+QLC+QRC+QSC$  ! MHL  
 $QL=QLC*QC/QTOT$  ! Liver MHL  
 $QF=QFC*QC/QTOT$  ! Fat MHL  
 $QR=QRC*QC/QTOT$  ! Rapidly-perfused tissue MHL  
 $QS=QSC*QC/QTOT$  ! Slowly-perfused MHL  
  
 ! Partition coefficients  
 $CONSTANT PL=0.732$  ! Liver:blood  
 $CONSTANT PF=6.19$  ! Fat:blood  
 $CONSTANT PS=0.408$  ! Muscle:blood  
 $CONSTANT PR=0.732$  ! Liver:blood  
 $CONSTANT PLU=0.732$  ! Lung:blood \*\*  
 $CONSTANT PB=19.4$  ! Blood:air  
  
 ! Metabolism  
 $CONSTANT VMAXC=4.0$  ! Allometric scaling constant for VMAX  
 $CONSTANT KM=0.4$  ! Michaelis constant for MFO pathway (mg/L)  
 $CONSTANT FRACR=0.1$  ! Oxidative metab in rapidly perfused MHL  
 $CONSTANT KFC=2.56$  ! Allometric scaling constant for KF  
 $CONSTANT MOLWT=85.0$  ! Molecular weight of DCM  
 $CONSTANT MWCO=28.0$  ! Molecular weight of CO MHL  
 $CONSTANT A1=0.416$  ! Ratio of specific activities of MFO, lung/liver MHL  
 $CONSTANT A2=0.137$  ! Ratio of specific activities of GST, lung/liver MHL  
  
 $VMAX=VMAXC*BW**0.7$  ! Maximum rate of MFO pathway (mg/hr)  
 $VMAXR=VMAX*FRACR$  ! MHL  
 $KF=KFC/(BW**0.3)$  !First-order rate constant for GSH pathway  
 $CONSTANT AFFG=0.0$  ! 1.57e-4 ! Affinity constant (1/Km) for GST pathway  
 ! Used to test impact of low affinity / slight saturation PMS 12/09  
 ! CO Submodel params MHL  
 $CONSTANT DLC=0.060$  ! MHL  
 $CONSTANT RENCOC=0.035$  ! Rate of endogenous CO production MHL  
 $CONSTANT ABCOC=0.117$  ! Conc of background CO (mg/kg) MHL  
 $CONSTANT HBTOT=10.0$  ! Conc of hemoglobin (mmoles/L) MHL  
 $CONSTANT P1=0.80$  ! CO yield factor MHL  
 $CONSTANT F1=1.21$  ! CO elimination factor MHL  
 $CONSTANT M=150.0$  ! Haldane coefficient MHL



CONSTANT COINH=2.2 ! Background CO inhalation concentration (ppm) MHL  
 CONSTANT O2=0.13 !4/16/32 mmoles/L MHL  
 CONSTANT PAIR=713.0 ! Pressure of air (mm Hg) MHL  
 CONSTANT RHO=1102.0 ! Density of CO (mg/L) MHL  
 CONSTANT SOL=0.03 ! mg/L/mmHG, MHL  
 DL=DLC\*BW\*\*0.92 ! MHL  
 RENCO=RENCOC\*BW\*\*0.7 ! MHL  
 ABCO=ABCOC\*BW ! MHL  
 Mmm=1.1325\*M ! MHL  
 L12D=1.0/P1 ! MHL  
 LCO=28.0/MOLWT ! MHL  
 L12=LCO\*P1 ! MHL

! Exposure schedule

CONSTANT CONC=0.0 ! Inhaled concentration (ppm)  
 CONSTANT TCHNG=6.0 ! Exposure pulse 1 width (hr)  
 CONSTANT TDUR=24.0 ! Exposure duration (hr)  
 CONSTANT TCHNG2=120.0 ! Exposure pulse 2 width (hr)  
 CONSTANT TDUR2=168.0 ! Exposure duration 2 (hr)  
 CIZONE = 1.0 ! Start with inhalation on  
 PEAK = 0.0 ! Zero peak concentration in brain  
 CONSTANT IVDOSE=0.0 ! IV infusion (mg/kg) MHL  
 CONSTANT TIV=0.0028 ! IV dosing time (hrs, or 10 seconds) MHL  
 CONSTANT VCHC=10.0 ! Uncorrected rat chamber volume (L) MHL  
 CONSTANT CC=0 ! Flag for closed chamber MHL  
 CONSTANT NCH=5 ! Number of animals in chamber MHL  
 CONSTANT KL=0.0 ! MHL  
 VCH=VCHC-(NCH\*BW)

! Oral Gavage Dosing added by MHL

CONSTANT BOLUS=0.0  
 TOTALBOLUS=BOLUS\*BW  
 TOTALDOSE=(IVDOSE+BOLUS)\*BW  
 IVORBOL=(TOTALDOSE.GT.0.0)  
 CONSTANT DRCONC=0.0 ! User-specified concentration in drinking water (mg/L)  
 CONSTANT FIXDRDOSE=0.0! Set to constant DW dose in mg/kg-day  
 ! Assume 70 kg human drinks 2 L/day, calculate rodent allometrically  
 DRVOL=0.102\*BW\*\*0.7 ! Daily water intake, based on body weight  
 DRDOSE=DRVOL\*DRCONC + FIXDRDOSE\*BW ! Total dose from water (mg (in a day))  
 DDOSE=DRDOSE/BW ! Daily dose (mg/kg/day)  
 NEWDAY=0.0 ! To reset area-under-curve values each 24 hours

```

PEAKPCTHBCO=0.0
CONSTANT KA=5.0          ! Rate constant for absorption

! Periodic drinking water intake schedule
! Assume T=0 is 7 AM (T+4=11 AM)
INTEGER I
I=2
DIMENSION DRTIME(32)      ! Store drinking times in array
DIMENSION DRPCT(32)      ! Store drinking percentages array
CONSTANT DRTIME=32*0.0, DRPCT=32*0.0
      GASD=MOLWT/24450.0

IF (DRDOSE.GT.0.0) SCHEDULE drink.AT.DRTIME(2)      ! Assume DRTIME(1)=0 ...
      ! and initial drink applied here.
CUMORALDOSE=DRPCT(1)*DRDOSE      !To calculate input to GI (mg)

! Switching 'lastday' and 'lastwk' to be defined by discrete blocks, below, PS 3/2009
lastday=0.0
IF (TSTOP.GT.24) SCHEDULE lday .AT. TSTOP-24
lastwk=0.0
if (TSTOP.GT.168) SCHEDULE lwk .AT. TSTOP-168

CIDAY=1.0
CIWK=1.0
SCHEDULE idayoff .AT.TCHNG
SCHEDULE idayon .AT.TDUR
SCHEDULE iwkoff.AT.TCHNG2
SCHEDULE iwkon .AT.TDUR2
!zzstopflag = .FALSE.
END      ! End of INITIAL section of program

DYNAMIC
      DISCRETE idayoff
            CIDAY=0.0
            QC=QCC*BW**0.74
            SCHEDULE idayoff .AT. (T+TDUR)
      END
      DISCRETE idayon
            CIDAY=1.0
            IF (QCSW) THEN
                  QC=QCCR*BW**0.74

```

```

ENDIF
SCHEDULE idayon .AT. (T+TDUR)
END
DISCRETE iwkoff
    CIWK=0.0
    SCHEDULE iwkoff .AT. (T+TDUR2)
END
DISCRETE iwkon
    CIWK=1.0
    SCHEDULE iwkon .AT. (T+TDUR2)
END
CIZONE = CIDAY*CIWK

DISCRETE lday      ! Value -> 1 for last 24 h, PS 3/2009
    lastday=1.0
end      ! of lday
DISCRETE lwk      ! Value -> 1 for last 168 h, PS 3/2009
    lastwk=1.0
end      ! of lwk

DISCRETE drink      ! Loop for drinking water intake schedule
    CUMORALDOSE = CUMORALDOSE + DRPCT(I)*DRDOSE
    I=I+1
    IF (I .EQ. 32) THEN
        I=1
        NEWDAY=NEWDAY + 24.0
    ENDIF
    SCHEDULE drink.AT.(NEWDAY+DRTIME(I))
End      ! Drink

```

#### DERIVATIVE

ALGORITHM IALG=2 ! Gear for stiff systems

```

! Following are daily averages over final week of simulations, PS 8/2008
WAVGLIVCYPDOSE=integ(lastwk*RAM1L,0.0)/(7.0*VL)
WAVGLIVGSTDOSE=integ(lastwk*RAM2L,0.0)/(7.0*VL)
WAVGLUNGGSTDOSE=integ(lastwk*RAM2LU,0.0)/(7.0*VLU)
WAVGLUNG CYPDOSE=integ(lastwk*RAM1LU,0.0)/(7.0*VLU)
WAVGWBDYGSTDOSE=integ(lastwk*(RAM2L+RAM2LU),0.0)/(7.0*BW)
WAVGWBDYCYPDOSE=integ(lastwk*(RAM1L+RAM1LU),0.0)/(7.0*BW)
WAVGAUCV=integ(lastwk*CV,0.0)/7.0

```

WAVGAUCS=integ(lastwk\*CS,0.0)/7.0

WAVGAUCL=integ(lastwk\*CL,0.0)/7.0

! Following are daily averages calculated from final day's dose-rate, PS 8/2008

LDAYLIVCYPDOSE=integ(lastday\*RAM1L,0.0)/VL

LDAYLIVGSTDOSE=integ(lastday\*RAM2L,0.0)/VL

LDAYLUNGGSTDOSE=integ(lastday\*RAM2LU,0.0)/VLU

LDAYWBDYGSTDOSE=integ(lastday\*(RAM2L+RAM2LU),0.0)/BW

LDAYWBDYCYPDOSE=integ(lastday\*(RAM1L+RAM1LU),0.0)/BW

LDAYLIVAUC=integ(lastday\*CL,0.0)

LDAYAUCV=integ(lastday\*CV,0.0)

LDAYAUCS=integ(lastday\*CS,0.0)

! GI compartment for drinking water inputs

RSTOM=-KA\*STOM

STOM=INTEG(RSTOM,TOTALBOLUS)+CUMORALDOSE

R= 0.21\*pulse(0.0,1.0,0.015)\*pulse(0.0,24.0,12.0)/(12\*0.015) + &

0.79\*pulse(0.0,0.6,0.015)\*pulse(12.0,24.0,12.0)/(20\*0.015)

test=integ(rdriink,0.0)

RIV=IVDOSE\*BW/TIV ! IV dose rate (mg/hr) MHL

blip=STEP(tiv)

IV=RIV\*(1.0-blip)

! Chamber calcs MHL

RACH=CC\*NCH\*QP\*((CA1/PB)-CCH)-(KL\*ACH) ! MHL

ACH=INTEG(RACH,CONC\*GASD\*VCH) ! MHL added initial condition

CCH=CC\*ACH/VCH ! MHL

CCHPPM=CCH/GASD ! MHL original

! If/else statements removed by use of multipliers; PMS 7-23-08

CI=(1.0-CC)\*CONC\*CIZONE\*GASD ! Convert to mg/L

DIDOSE=24.0\*QP\*INTEG(CI,0.0)/(BW\*(T+1.0E-8)) ! Daily inhaled dose (mg/kg-d)

! CA1=Arterial blood concentration from gas exchange region

! to lung tissue compartment (mg DCM/L)

CA1=(QP\*CI+QP\*CCH+QC\*CV)/((QP/PB)+QC) !\*\*

! AX=Amount eliminated by exhalation (mg DCM)

CX=CA1/PB ! Concentration in air leaving gas exchange region

RAX=QP\*CX

AX=INTEG(RAX, 0.0)

! Concentration DCM in exhaled air

CEX1=0.7\*CX + 0.3\*CI

! Assumes mixing with 30% of inhaled that only goes to deadspace

RAEX1=CEX1\*QP ! MHL

AEX1=INTEG(RAEX1,0.0) ! MHL

PCTIVEXH=100.0\*IVORBOL\*AX/(TOTALDOSE + 1.0E-8)

! % IV/BOLUS dose exhaled as DCM, MHL

! Amount in lung tissue (mg DCM)

RAM1LU=A1\*VMAX\*CA\*(VLU/VL)/(KM+CA)

AM1LU=INTEG(RAM1LU, 0.0)

RAM2LU=A2\*KF\*CA\*VLU/(1.0+AFFG\*CA)

RALU=QC\*(CA1-CA)-RAM1LU-RAM2LU

ALU=INTEG(RALU, 0.0)

CLU=ALU/VLU

AUCLU=INTEG(CLU,0.0)

CA=CLU/PLU ! Amount in arterial blood to body

! AF=Amount in fat (mg DCM)

RAF=QF\*(CA-CVF)

AF=INTEG(RAF, 0.0)

CF=AF/VF

CVF=CF/PF

!AL=Amount in liver (mg DCM)

RAL=QL\*(CA-CVL)-RAM1L-RAM2L+KA\*STOM

RAM1L=VMAX\*CVL/(KM+CVL) !\*\*

RAM2L=KF\*CVL\*VL/(1.0+AFFG\*CVL) !\*\*

AL=INTEG(RAL, 0.0)

CL=AL/VL

CVL=CL/PL

AUCL=INTEG(CL, 0.0) !\*\*

!AS= Amount in slowly-perfused tissues

RAS=QS\*(CA-CVS) !-RAMS ! VMAXS=0, so not needed, PS 7/2008

AS=INTEG(RAS, 0.0)

CS=AS/VS

CVS=CS/PS

!AR=Amount in rapidly-perfused tissues  
 RAMR=VMAXR\*CVR/(KM+CVR) ! MHL  
 AMR=INTEG(RAMR,0.0) ! Total metabolized (MFO) in RP, MHL  
 RAR=QR\*(CA-CVR)-RAMR  
 AR=INTEG(RAR, 0.0)  
 CR=AR/VR  
 CVR=CR/PR

! AM1=Amount metabolized in MFO pathway (mg DCM)  
 RAM1=RAM1LU+RAM1L+Ramr !+ams ! AMS = 0, term not needed, PS 7/2008  
 TAM1=INTEG(RAM1,0.0)  
 ! DDAM2=Amount metabolized in GSH pathway (mg DCM)  
 RAM2=RAM2LU+RAM2L  
 TAM2=INTEG(RAM2,0.0)  
 ! Total rate/amount metabolized (MFO plus GSH pathways)  
 RAMTOT=RAM1+RAM2 !\*\*  
 AMTOT=INTEG(RAMTOT, 0.0)

! CV=Average venous blood concentration (mg DCM/L)  
 CV=(QF\*CVF+QL\*CVL+QS\*CVS+QR\*CVR+IV)/QC !MHL added RIV  
 CVP=0.23\*CV ! Venous plasma conc  
 ABL=(CA+CV)\*VBL2  
 AUCBL=INTEG(ABL, 0.0)  
 AUCV=INTEG(CV, 0.0) ! GLD

! CO appearance in and elimination from blood, MHL  
 ACO=ABL2/(VBL2\*MWCO) !MHL  
 BO2=O2/Mmm ! MHL  
 CHBT=HBTOT ! MHL  
 HBCO=((ACO+BO2+CHBT)-SQRT((ACO+BO2+CHBT)\*(ACO+BO2+CHBT)-4.0\*ACO\*CHBT))/2.0  
 ! Conc of HB-bound CO (mmol/L) MHL  
 COFREE=ACO-HBCO ! Conc of free CO in blood (mmol/L) MHL  
 PICO=COINH/1402.5 ! CO concentration in inhaled air (mg/L) MHL

PCTHBCO=100.0\*HBCO/HBTOT ! Percent carboxyhemoglobin in blood  
 PEAKPCTHBCO = MAX(PEAKPCTHBCO,PCTHBCO) ! Capture peak CO concentration

! COE = Amount of expired CO, MHL  
 PCCO=COFREE\*MWCO/SOL  
 PACO=(PCCO+PICO\*QPDP)/(1.0+QPDP)

RCOE=DL\*RHO\*(PCCO-PACO)\*F1  
 COE=INTEG(RCOE,0.0) ! Expired amount of CO (mg) MHL  
 COEC=0.7\*COE/QP ! Conc expired CO (mg/L)  
 ! Multiplier changed from 2/3 to 0.7 in above to be consistent with...  
 ! other expired concentration calculations, PS 7/2008.  
 COECPPM=COEC\*24450.0/MWCO  
 PCTIVCOEXH=100.0\*IVORBOL\*COE/(TOTALDOSE\*MWCO/MOLWT + 1.0E-8) ! % dose exhaled  
 as CO MHL

! ABL2 = Amount of CO in blood (mg) MHL  
 RFROMDCM=(RAMIL+RAM1LU+RAMR)\*L12 ! RAMS left out since not used, =0  
 ! Inputs calculated for mass balance  
 AENCO = INTEG(RENCO, 0.0) ! Amount produced endogenously (mg)  
 AFROMDCM = INTEG(RFROMDCM, 0.0) ! Amount produced from metabolism of DCM  
 RABL2=RFROMDCM+RENCO-RCOE  
 ABL2=INTEG(RABL2,0.0) ! MHL  
 CABL2=ABL2/VBL2 ! MHL

! AI=Total mass input (mg DCM)  
 RAI=(QP\*(CI+CCH))+RIV !MHL  
 AI=INTEG(RAI, 0.0)

! TMASS=Mass balance (mg DCM)  
 MASSBAL=MASSIN-STOM-AF-AL-ALU-AS-AR-ABL-MASSOUT  
 MASSIN=AI+CUMORALDOSE + TOTALBOLUS  
 MASSOUT=AX+AMTOT

! Mass Balance for CO  
 MASSBALCO = 100\*(-ABL2+AENCO-COE+AFROMDCM)/(AENCO+AFROMDCM+1e-10)

! IF (T .GE. TSTOP) zzstopflag = .TRUE. ! Termination condition for model run  
 END ! End of DERIVATIVE section of program  
 TERMT(T.GE.TEND)  
 END ! End of Dynamic section of program

TERMINAL

AVGT= 24.0/(T+1.0e-8)  
 METDOS = AVGT\*RAMTOT/VL  
 LDAYREFDOSE = LDAYLIVCYPDOSE + LDAYLIVGSTDOSE ! MHL  
 WAVGREFDOSE = WAVGLIVCYPDOSE + WAVGLIVGSTDOSE ! PS, 8/2008  
 LDAYWBDYCYPGSTDOSE = LDAYWBDYCYPDOSE + LDAYWBDYGSTDOSE

```

METABRATIO = RAM1/(RAM2+1.0E-12)
DDOSECHECK = CUMORALDOSE*avgt/BW ! Should equal DDOSE
END ! End of TERMINAL section of program
END ! End of PROGRAM

```

## Computational Files (.m files)

The following are files (functions) created as .m files to support the Monte Carlo statistical sampling and subsequent calculations for the DCM analysis.

### Utility files used for all human simulations

---

```

function v=normbnd(mu, sigma, lo, up)
% -----
% FILE: normbnd.m
%
% NORMBND - Returns a random sample from a truncated normal distribution with
%     mean = mu, standard deviation = sigma, lower bound = lo, & upper bound = up.
% Modified by Paul Schlosser, U.S. EPA, 7/2008
% -----
    if (lo >= up)
        v = NaN; disp('Lower bound must be less than upper bound.')
        return;
    end
    FL = normcdf(lo, mu, sigma); % normal probability of lower bound
    FU = normcdf(up, mu, sigma); % normal probability of upper bound
    p = rand*(FU - FL) + FL; % rand = uniform random(0,1)
    v = norminv(p, mu, sigma);
end

```

---

```

function v=lnormbnd(mu, sigma, lo, up)
% -----
% FILE: lnormbnd.m
%
% LNORMBND - Returns a truncated LOGnormal distribution with distribution with
%     mean = mu, standard deviation = sigma, lower bound = lo, & upper bound = up.
% Modified by Paul Schlosser, U.S. EPA, 7/2008
% -----
v = exp(normbnd(log(mu), log(sigma), log(lo), log(up)));
end

```

---

```

function v=prctile(y,x)
% -----
% FILE: prctile.m
% Computes percentiles of vector y at percentile values x
% If x = [50 90 95], that's 50th, 90th, 95th
% Created by Paul Schlosser, U.S. EPA, July, 2008
% -----
if any((x<0) | (x>100))
    disp('Error! All percentiles must be between 0 and 100!');
end

```



```

    return
end
dy = sort(y);
ps = 100*((1:length(dy))-0.5)/length(dy);
v = x;
for id=1:length(x)
    if x(id)<ps(1)
        v(id)=dy(1);
    elseif x(id)>max(ps)
        v(id)=max(dy);
    else
        v(id)=interp1(ps,dy,x(id));
    end
end
end

```

---

```

% -----
% File clearT.m
% Creates list of variable names for tracking, rnames, and
% performs other initializations for human Monte Carlo chains.
% Created by Paul Schlosser, U.S. EPA, 7/2008
% -----
rnames=["age";"BW";"DRVOL";"DRCONC";"DDOSE";"CONC";"VMAXC";"KM";"FRACR";"KFC";
"A1";"A2";"BW";"VLC";"VFC";"VRC";"VLUC";"VSC";"QAlvC";"QCC";"QLC";"QFC";"QSC";
"QRC";"PL";"PLU";"PF";"PS";"PR";"PB";"GSTGT";"CV";"LDAYLIVGSTDOSE";
"LDAYLIVCYPDOSE";"LDAYLUNGGSTDOSE";"LDAYWBDYGSTDOSE";"LDAYREFDOSE";
"LDAYAUCV";"LDAYAUCS"];
sr='['; rns=length(rnames);
for ir=1:(rns-1)
    sr=[sr,ctot(rnames(ir)),', '];
end
sr=[sr,ctot(rnames(rns)),','];
% save current model values before perturbations
save @file='dcm_human.sav'
runo=[];
% prepare time history values.
prepare @clear T CV
WESITG=0; WEDITG=0; CONC=0.0; IVDOSE=0.0; DRCONC=0.0; FIXDRDOSE=0.0;
IVORBOL=0.0;
DRPCT = [0.25, 0.1, 0.25, 0.1, 0.25, 0.05];
DRTIME= [0.0, 3.0, 5.0, 8.0, 11.0, 15.0];
TEND=600.; TCHNG=2000.0; TCHNG2=2000.0; TDUR=2000.0; TDUR2=2000.0;
start @NoCallback

exist popn;    % Test to see if 'popn' defined.
if ~ans % If not...
    popn="mix"; % Mix of GST types.
end
exist agem;    % Check if agem defined ...
if ~ans % If not...
    agem=0;
end
exist gendm;   % Check if gendm defined ...
if ~ans % If not...

```

```
    gendm="both"; % males and females
end
```

---

```
function nn=findnames(nm,rnames)
```

```
% -----
% File findnames.m
%
% Function to find indices of specific names in list nm
% within the larger list (of all variables), rnames.
% Created by Paul Schlosser, U.S. EPA, 7/2008
% -----
    nn=[];
    for n=1:length(nm)
        n1=find(rnames==nm(n));
        if isempty(n1)
            disp(['Error, 'ctot(nm(n)),' not in saved variable list.']); return
        end
        nn=[nn,n1];
    end
end
```

```
% File human_par1.m
%
% Code for selecting human model parameters from MC distribution for
% dichlormethane PBPK model from probability density functions as
% described by David et al (2006), with *most of* the revisions as
% described in the U.S. EPA IRIS Toxicological Review for Dichloromethane,
% Appendix B.
% ** This code uses "one-dimensional" sampling distributions for CYP
% (VMAXC) and GSTT-1 (KFC). The sampling of KFC is as described by David
% et al (2006). The sampling for VMAXC uses the mean value of David et
% al (2006) but substitutes a larger geometric standard deviation and
% uses log-normal distribution without bounds. Further details below.
% Genotypic distribution of GSTT-1 activity taken from Tables 8 and 9
% of Environ report for Eastman Kodak.
% Arithmetic means and SDs for lognormal distributions converted
% by ML to geometric mean/SD to match acsl m-file lnormbnd.m values.
%
% Programmed by Michael Lumpkin (ML)
% Syracuse Research Corporation, 11/2005
% Modified by Paul Schlosser (PS), U.S. EPA 7/2008 and 1/2010
%
% Gender parameter 'gendm' can be "both" (default), "male", or "female".
% Age parameter 'agem' can be 0 (default) to simulate the full range
% from 0.5-80 years, or any value in that range for a specific age.
% Specific parameters, as noted below, set for those life-stages.
% -----
% choose uniform discrete distribution for GSTT-1 genotypes.
if popn=="++"
    GSTGT = 0; % for ++ only
elseif popn=="+-"
    GSTGT = 1; % for +/- only
```

```

elseif popn=="--"
    GSTGT = 2; % for -/- only
else
    r=rand(1)*gsmult; GSTGT=(r>0.32)+(r>0.8); %For general/mixed population
    popn = "mix of +/+, +/-, and -/-";
end

% selection of GSTT1 activity distribution based on ethnic distribution
% upper bounds changed from mean + 3*SD to mean + 5*SD, P.S., 2/2009
if (GSTGT == 0)
    KFC = (5.87/0.852)*normbnd(1.31, 0.167, 0.0, 2.145); %
elseif (GSTGT == 1)
    KFC = (5.87/0.852)*normbnd(0.676, 0.123, 0.0, 1.291); %
else
    KFC = 0.0;
end
AFFG=0.0; % 1.57e-4 % Affinity constant (1/Km) for GSTT-T1 pathway
% Allows for some saturation of the GSTT-1 pathway if set > 0, P.S., 1/2010

KA=5.0; %From Reitz et al (1997); this parameter has no variability data and is
% described as point estimate
VMAXC=lognorm(9.34, 1.73); %lnormbnd(9.34, 1.73, 3.8, 23.0);
%geometric mean and GSD, converted from arith mean/SD of 9.42 and 1.23
% VMAXC switched to unbounded distribution, PS, 2/2009
KM=lnormbnd(0.410, 1.39, 0.154, 1.10);
%geometric mean and SD, converted from arith mean/SD of 0.433 and 1.46
FRACR=lnormbnd(0.0152, 2.0, 0.0019, 0.122);
%geometric mean and SD, converted from arith mean/SD of 0.0193 and 0.0152
A1=lnormbnd(0.00092, 1.47, 0.000291, 0.00292);
%geometric mean & SD, converted from arith mean/SD of 0.000993 & 0.000396
A2=lnormbnd(0.0083, 1.92, 0.00116, 0.0580);
%geometric mean and SD, converted from arith mean/SD of 0.0102 and 0.00739

age=agem;
if age==0 % agem=0 => age = random from population, otherwise leave as set
    p=rand;
    age = 165.86*p^4 - 253.19*p^3 + 113.27*p^2 + 53.356*p + 0.5;
end
VMP = 0.7 + 0.18*(age<18);
% power for scaling Vmax is 0.88 if age < 18, otherwise 0.7
% Change in VMP introduced 2/2009, PS

gend=gendm;
if ~((gendm=="male")|(gendm=="female")) % if not defined as one
    gend="female";
    if rand()<(0.513*((125.3 - age)^4)/(33.7^4 + (125.36 - age)^4));
        gend="male";
    end
end

if mod(RUNN,50)==0
    disp(['GSTGT group: ',ctot(popn),'. GSTGT = ',num2str(GSTGT),'.']);
    disp(['Simulation for ',num2str(age), ' year old ',ctot(gend),'; gender mix: ',ctot(gendm)]);
    disp(' ');
end

```

```

end

% Human physiologic parameters

if gend=="female"
    aged=(16-age)/10;
    bwmean = 4.146 - 0.147*aged - 1.36*aged^2 + 0.44*aged^3;
    if aged<0
        bwmean = 4.146 - 0.147*aged - 0.0278*aged^2 - 0.00095*aged^3;
    end
    aged=(13-age)/10;
    bwSD = 2.574 - 0.358*aged - 2.55*aged^2 + 1.16*aged^3;
    if aged<0
        bwSD = 2.574 - 0.358*aged - 0.0861*aged^2 - 0.00469*aged^3;
    end
else % males
    aged=(21-age)/10;
    bwmean = 4.406 - 0.0285*aged - 0.729*aged^2 + 0.115*aged^3;
    if aged<0
        bwmean = 4.406 - 0.0285*aged + 0.0048*aged^2 + 0.0018*aged^3;
    end
    aged=(16-age)/10;
    bwSD = 2.87 + 0.06*aged - 2.56*aged^2 + 0.96*aged^3;
    if aged<0
        bwSD = 2.87 + 0.06*aged + 0.0448*aged^2 + 0.0067*aged^3;
    end
end
end
BW=norminv((0.01+0.98*rand),exp(bwmean),exp(bwSD));

QAlvmean = 13.6 + 13.3*exp(-0.05*age);
if gend=="female"
    QAlvmean = 10.7 + 22.1*exp(-0.08*age);
end
Qgsd = -0.1948*(age/10)^3 + 0.6095*(age/10)^2 - 0.3978*(age/10) + 1.4261;
if age>16.81
    Qgsd=1.554;
end
QAlvC = QAlvmean*exp(norminv((0.05+0.9*rand),0,log(Qgsd)));

QCCmean = 56.906*(1.0 - exp(-0.681*exp(0.0454*QAlvC))) - 29.747;
QCC = QCCmean/lnormbnd(1.0, 0.203, 0.69, 1.42);
VPR = QAlvC/QCC;

vlm=-0.0036*(age/10)^2 - 0.0051*(age/10) + 0.0395;
if age>17
    vlm=-0.0004*(age/10)^2 + 0.0034*(age/10) + 0.0169;
end
VLC=vlm*normbnd(1.0, 0.05, 0.85, 1.15);

vfm=0.1612*(age/10)^3 + 0.0846*(age/10)^2 - 0.3083*(age/10) + 0.2709;
if ((age>7)&(age<=20))
    vfm=-0.0458*(age/10)^2 + 0.2082*(age/10) + 0.0274;
    if gend=="male"
        vfm= -0.0057*(age/10)^2 + 0.0293*(age/10) + 0.1303;
    end
end

```

```

        end
elseif age>20
    vfm=-0.0024*(age/10)^3 + 0.0355*(age/10)^2 - 0.115*(age/10) + 0.3678;
    if gend=="male"
        vfm= -0.0015*(age/10)^2 + 0.0384*(age/10) + 0.0908;
    end
end
VFC=vfm*normbnd(1.0, 0.3, 0.1, 1.9);

VRC=normbnd(0.064, 0.0064, 0.0448, 0.0832);
VLUC=normbnd(0.0115, 0.00161, 0.00667, 0.0163);
VSC=normbnd(0.63, 0.189, 0.431, 0.829);
vtr=0.9215/(VFC+VLC+VLUC+VRC+VSC);
VFC = VFC*vtr; VLC=VLC*vtr; VLUC=VLUC*vtr; VRC=VRC*vtr; VSC=VSC*vtr;
VBL2C=0.059;

DSPC=0.15;
QLC=normbnd(0.26, 0.0910, 0.010, 0.533);
QFC=normbnd(0.05, 0.0150, 0.0050, 0.0950);
QSC=normbnd(0.19, 0.0285, 0.105 ,0.276);
QRC=normbnd(0.50 ,0.10, 0.20, 0.80);

% Human partition coefficients for DCM
PL=lnormbnd(1.43, 1.22, 0.790, 2.59);
    %geometric mean and SD, converted from arith mean/SD of 1.46 and 0.292
PLU=lnormbnd(1.43, 1.22, 0.790, 2.59);
    %geometric mean and SD, converted from arith mean/SD of 1.46 and 0.292
PF=lnormbnd(11.9, 1.34, 4.92, 28.7);
    %geometric mean and SD, converted from arith mean/SD of 12.4 and 3.72
PS=lnormbnd(0.80, 1.22, 0.444, 1.46);
    %geometric mean and SD, converted from arith mean/SD of 0.82 and 1.64
PR=lnormbnd(1.43, 1.22, 0.790, 2.59);
    %geometric mean and SD, converted from arith mean/SD of 1.46 and 0.292
PB=lnormbnd(9.7, 1.10, 7.16, 13.0);
    %geometric mean and SD, converted from arith mean/SD of 9.7 and 0.97



---


% File human_par2.m
%
% Code for selecting human model parameters from MC distribution for
% dichlormethane PBPK model from probability density functions as
% described by David et al (2006), with revisions as described in the
% U.S. EPA IRIS Toxicological Review for Dichloromethane, Appendix B.
%
% ** This code uses "two-dimensional" sampling distributions for CYP
% (VMAXC) and GSTT-1 (KFC), as described in the current assessment.
% Also see comments further details below.
%
% Genotypic distribution of GSTT-1 activity taken from Tables 8 and 9
% of Environ report for Eastman Kodak.
% Arithmetic means and SDs for lognormal distributions converted
% by ML to geometric mean/SD to match acsl m-file lnormbnd.m values.
%
% Programmed by Michael Lumpkin (ML)

```

```

% Syracuse Research Corporation, 11/2005
% Modified by Paul Schlosser (PS), U.S. EPA 7/2008 and 1/2010
%
% Gender parameter 'gendm' can be "both" (default), "male", or "female".
% Age parameter 'agem' can be 0 (default) to simulate the full range
% from 0.5-80 years, or any value in that range for a specific age.
% Specific parameters, as noted below, set for those life-stages.
% -----
% 'par2'
% choose uniform discrete distribution for GSTT-1 genotypes.
if popn=="++"
    GSTGT = 0; % for ++ only
elseif popn=="+-"
    GSTGT = 1; % for +/- only
elseif popn=="--"
    GSTGT = 2; % for -/- only
else
    r=rand(1)*gsmult; GSTGT=(r>0.32)+(r>0.8); %For general/mixed population
    popn = "mix of +/+, +/-, and -/-";
end

% selection of GSTT1 activity distribution based on ethnic distribution
% upper bounds changed from mean + 3*SD to mean + 5*SD, P.S., 2/2009
% From Table 4 of David et al. (2006) for kfc, mu = 0.852, CV = 0.711
% Lognorm tranform: m = mu/sqrt(CV^2 + 1) = 0.6944
% and s = exp(sqrt(ln(CV^2 + 1))) = 1.896
% kfc=lnormbnd(m,s,m/(s^2),m*(s^2))/((0.48/2)+0.32);
% The last term (divisor) accounts for relative weighting and activity
% of the three genotypes.
kfc=lnormbnd(0.6944,1.896,0.1932,2.496);
if (GSTGT == 0)
    KFC = kfc*normbnd(1.786, 0.2276, 0.0, 2.924); % Relative activity in ++ pop'n
    %=kfc*normbnd(1, 0.167/1.31, 0.0, 1.0+5*0.167/1.31)/0.56;
    % 0.167 and 1.31 are s.d and mean, respectively, for ++ from Table 2 of David et al.
    (2006)
elseif (GSTGT == 1)
    KFC = kfc*normbnd(0.8929, 0.1622, 0.0, 1.704); % Relative activity in +/- pop'n
    %=kfc*normbnd(1, 0.123/0.676, 0.0, 1.0+5*0.123/0.676);
    % 0.123 and 0.676 are s.d and mean, respectively, for +/- from Table 2 of David et al.
    (2006)
else
    KFC = 0.0;
end
AFFG=0.0; % 1.57e-4 % Affinity constant (1/Km) for GSTT-T1 pathway
% Allows for some saturation of the GSTT-1 pathway if set > 0, P.S., 1/2010

KA=5.0; %from Reitz et al (1997); this parameter has no variability data and is described as point
estimate
vmaxcm=lnormbnd(9.34,1.14,7.20,12.11);
%geometric mean and GSD, converted from arith mean/SD of 9.42 and 1.23
% lower/upper bounds = GM/(GSD^2) and GM*(GSD^2); i.e., +/- 2 SD in log-space
% bound calculations done with GM and GSD *not rounded* to 2 decimal places
VMAXC=lognorm(vmaxcm, 1.73); %lnormbnd(9.34, 1.73, 3.8, 23.0);
% VMAXC switched to unbounded distribution, PS, 2/2009

```

```

KM=lnormbnd(0.410, 1.39, 0.154, 1.10);           %geometric mean and SD, converted from arith
mean/SD of 0.433 and 1.46
FRACR=lnormbnd(0.0152, 2.0, 0.0019, 0.122);     %geometric mean and SD, converted from arith
mean/SD of 0.0193 and 0.0152
A1=lnormbnd(0.00092, 1.47, 0.000291, 0.00292);   %geometric mean & SD, converted
from arith mean/SD of 0.000993 & 0.000396
A2=lnormbnd(0.0083, 1.92, 0.00116, 0.0580);     %geometric mean and SD, converted from arith
mean/SD of 0.0102 and 0.00739

age=agem;
if age==0      % agem=0 => age = random from population, otherwise leave
    p=rand;
    age = 165.86*p^4 - 253.19*p^3 + 113.27*p^2 + 53.356*p + 0.5;
end
VMP = 0.7 + 0.18*(age<18);    % power for scaling Vmax is 0.88 if age < 18, otherwise 0.7
% Change in VMP introduced 2/2009, PS

gend=gendm;
if ~((gendm=="male")|(gendm=="female"))    % if not defined as one
    gend="female";
    if rand()<(0.513*((125.3 - age)^4)/(33.7^4 + (125.36 - age)^4));
        gend="male";
    end
end

if mod(RUNN,50)==0
    disp(['GSTGT group: ',ctot(popn),'. GSTGT = ',num2str(GSTGT),'.']);
    disp(['Simulation for ',num2str(age),' year old ',ctot(gend),'; gender mix: ',ctot(gendm)]);
    disp(' ');
end
    % Human physiologic parameters

if gend=="female"
    aged=(16-age)/10;
    bwmean = 4.146 - 0.147*aged - 1.36*aged^2 + 0.44*aged^3;
    if aged<0
        bwmean = 4.146 - 0.147*aged - 0.0278*aged^2 - 0.00095*aged^3;
    end
    aged=(13-age)/10;
    bwSD = 2.574 - 0.358*aged - 2.55*aged^2 + 1.16*aged^3;
    if aged<0
        bwSD = 2.574 - 0.358*aged - 0.0861*aged^2 - 0.00469*aged^3;
    end
else % males
    aged=(21-age)/10;
    bwmean = 4.406 - 0.0285*aged - 0.729*aged^2 + 0.115*aged^3;
    if aged<0
        bwmean = 4.406 - 0.0285*aged + 0.0048*aged^2 + 0.0018*aged^3;
    end
    aged=(16-age)/10;
    bwSD = 2.87 + 0.06*aged - 2.56*aged^2 + 0.96*aged^3;
    if aged<0
        bwSD = 2.87 + 0.06*aged + 0.0448*aged^2 + 0.0067*aged^3;
    end
end

```

```

end
BW=norminv((0.01+0.98*rand),exp(bwmean),exp(bwSD));

QAlvmean = 13.6 + 13.3*exp(-0.05*age);
if gend=="female"
    QAlvmean = 10.7 + 22.1*exp(-0.08*age);
end
Qgsd = -0.1948*(age/10)^3 + 0.6095*(age/10)^2 - 0.3978*(age/10) + 1.4261;
if age>16.81
    Qgsd=1.554;
end
QAlvC = QAlvmean*exp(norminv((0.05+0.9*rand),0,log(Qgsd)));

QCCmean = 56.906*(1.0 - exp(-0.681*exp(0.0454*QAlvC))) - 29.747;
QCC = QCCmean/lnormbnd(1.0, 0.203, 0.69, 1.42);
VPR = QAlvC/QCC;

vlm=-0.0036*(age/10)^2 - 0.0051*(age/10) + 0.0395;
if age>17
    vlm=-0.0004*(age/10)^2 + 0.0034*(age/10) + 0.0169;
end
VLC=vlm*normbnd(1.0, 0.05, 0.85, 1.15);

vfm=0.1612*(age/10)^3 + 0.0846*(age/10)^2 - 0.3083*(age/10) + 0.2709;
if ((age>7)&(age<=20))
    vfm=-0.0458*(age/10)^2 + 0.2082*(age/10) + 0.0274;
    if gend=="male"
        vfm= -0.0057*(age/10)^2 + 0.0293*(age/10) + 0.1303;
    end
elseif age>20
    vfm=-0.0024*(age/10)^3 + 0.0355*(age/10)^2 - 0.115*(age/10) + 0.3678;
    if gend=="male"
        vfm= -0.0015*(age/10)^2 + 0.0384*(age/10) + 0.0908;
    end
end
VFC=vfm*normbnd(1.0, 0.3, 0.1, 1.9);

VRC=normbnd(0.064, 0.0064, 0.0448, 0.0832);
VLUC=normbnd(0.0115, 0.00161, 0.00667, 0.0163);
VSC=normbnd(0.63, 0.189, 0.431, 0.829);
vtr=0.9215/(VFC+VLC+VLUC+VRC+VSC);
VFC = VFC*vtr; VLC=VLC*vtr; VLUC=VLUC*vtr; VRC=VRC*vtr; VSC=VSC*vtr;
VBL2C=0.059;

DSPC=0.15;
QLC=normbnd(0.26, 0.0910, 0.010, 0.533);
QFC=normbnd(0.05, 0.0150, 0.0050, 0.0950);
QSC=normbnd(0.19, 0.0285, 0.105 ,0.276);
QRC=normbnd(0.50 ,0.10, 0.20, 0.80);

PL=lnormbnd(1.43, 1.22, 0.790, 2.59);
    %geometric mean and SD, converted from arith mean/SD of 1.46 and 0.292
PLU=lnormbnd(1.43, 1.22, 0.790, 2.59);
    %geometric mean and SD, converted from arith mean/SD of 1.46 and 0.292

```



```

PF=lnormbnd(11.9, 1.34, 4.92, 28.7);
    %geometric mean and SD, converted from arith mean/SD of 12.4 and 3.72
PS=lnormbnd(0.80, 1.22, 0.444, 1.46);
    %geometric mean and SD, converted from arith mean/SD of 0.82 and 1.64
PR=lnormbnd(1.43, 1.22, 0.790, 2.59);
    %geometric mean and SD, converted from arith mean/SD of 1.46 and 0.292
PB=lnormbnd(9.7, 1.10, 7.16, 13.0);
    %geometric mean and SD, converted from arith mean/SD of 9.7 and 0.97

```

---

```

% -----
% File: finish.m
% Programmed by Paul Schlosser, U.S. EPA, 8/2008, rev. 1/2010
% Performs final analysis on saved results in runo from simulations.
% -----
disp([num2str(size(runo,1)), ' simulations completed.']);
contsim=0; p=ctot(popn); p=p(1:min([3,length(p)]));
eval(['save runo @file=',astp,num2str(agem),p,'_',ctot(gendm), ...
    '_ ',model(7:10),'.csv @format=ascii @separator=comma']);
eval(['save @file=',astp,num2str(agem),p,'_',ctot(gendm),'_',model(7:10),'.mat']);
disp(['GSTT-1 group: ',ctot(popn)]);
aget=['a ',num2str(agem),' year-old;'];
if agem==0
    aget='0.5-80 years of age;';
end
gendt=[ctot(gendm),' population'];
if gendm=="both"
    gendt='males and females';
end
disp(['Simulation for ',aget,gendt,']);
disp(['Metric =',dtxt,']);
res=[];
pcs=num2str(percs(1));
for n=2:length(percs)
    pcs=[pcs,' ',num2str(percs(n))];
end
for n=1:length(nm)
    disp([ctot(nm(n)), ' mean, median, percentiles = ',pcs]);
    r=[mean(runo(:,nn(n))),median(runo(:,nn(n))),prctile(runo(:,nn(n)),percs/gsmult)]*mult;
    disp(r); disp(' '); res=[res;r];
end

```

---

### Files used specifically for cancer analysis

---

```

% -----
% File: straightsims.m
% Created by Paul Schlosser, U.S. EPA, 8/2008
% Runs Monte Carlo (MC) simulations withOUT search (for human equivalent exposures)
% Requires set exposure, NRUNS and rnames (list of variables to save)
% PBPK parameters set using MC selection by file human_pars.m.
% -----
exist contsim; % Test to see if contsim defined.

```

```

if ~ans % If not...
    contsim=0; % Not a continuation.
end
if contsim==0 % When starting a new set of simulation;
    % set contsim=1 if continuing an interrupted chain.
    runo=[]; rns=length(rnames); ns=1;
else
    ns=RUNN;
end
for RUNN = ns:NRUNS
    if mod(RUNN,50)==0
        disp(['Remaining runs = ',num2str(NRUNS-RUNN)])
    end
    human_pars; start @NoCallback
    runo=[runo;eval(sr)];
end

% -----
% File: Human drinking water MCA OSF.m
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 10/2007
% Modified by Paul Schlosser, U.S. EPA 8/2008, rev. 1/2010
%
% This script file sets up the control parameters to simulate human unit
% drinking water exposure (1 mg/kg/day) and calls straightsim which
% generates a Monte-Carlo chain for the internal doses (identified
% in text array nn) to be used in calculating oral slope factors (OSFs).
% -----
% Test to see if contsim defined
exist contsim
if ~ans % If not...
    contsim=0; % Not a continuation.
end
if contsim==0 % When starting a new set of simulation;
    % set contsim=1 if continuing an interrupted chain.
    clearT
    nm=["LDAYLIVGSTDOSE";"LDAYWBDYGSTDOSE"]; nn=findnames(nm,rnames);
    if (length(nn) ~= length(nm))
        disp("Not all names in list nm.")
        return
    end
    TEND=95.0; NRUNS=10000; CINT=1.0; % Total iterations for Monte Carlo analysis
    FIXDRDOSE=1.0; % Drinking fixed mg/kg-day
    gendm="female"; % Gender mix; choose "male", "female", or "both"
    agem=70; % Age "mix"; if 0 draws from distribution (0.5-80 years)
    % otherwise agem value is used exclusively
    popn="++"; % GSTGT "++", "+-", or "mix" of +/+, +/-, and -/-
    model='human_par2';
    % Choose model between 'human_par1' (original) and 'human_par2' (+ uncertainty)
    astp='OSF_age'; percs=[95 99]; dtxt='n/a'; gsmult=1.0;
end
straightsim; mult=1.0/DDOSE; finish

```

---

```

% -----
% File: Human inhalation MCA IUR.m
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 11/2007
% Modified by Paul Schlosser, 7/2008, rev. 1/2010
%
% This script file sets up the control parameters to simulate human unit inhalation exposures
% (1 mcg/m3) and calls straightsim which generates a Monte-Carlo chain for the internal
% doses (identified in text array nm) to be used in calculating inhalation unit risks (IURs)
% -----
exist contsim; % Test to see if contsim defined.
if ~ans % If not...
    contsim=0; % Not a continuation.
end
if contsim==0 % When starting a new set of simulation;
    % set contsim=1 if continuing an interrupted chain.
    clearT; CINT=0.1;
    TEND=24*4; CONC=1.0e-6/GASD; CC=0; % Daily dose-rates now calculated from
    % final day of simulations, & only need to go to 60 hr to reach SS
    % CONC in ppm; 0.00029 ppm = 1 ug/m3 DCM = 1e-3 ug/L = 1e-6 mg/L
    nm=["LDAYLIVGSTDOSE";"LDAYLUNGGSTDOSE";"LDAYWBDYGSTDOSE";
    "CV";"LDAYAUCS"];
    nn=findnames(nm,rnames);
    if (length(nn) ~= length(nm))
        disp("Not all names in list nm.")
        return
    end
    NRUNS=10000; %Total iterations for Monte Carlo analysis
    gendm="female"; % Gender mix; choose "male", "female", or "both"
    agem=70; % Age "mix"; if 0 draws from distribution (0.5-80 years)
    % otherwise agem value is used exclusively
    popn="++"; % GSTGT "++", "+-", or "mix" of +/+, +/-, and -/-
    gsmult = 1.0; % Multiplies draw for GST individual selection, divides the percentiles at end.
    astp='IUR_age'; mult= 1.0e-6/(CONC*GASD); percs=[95 99]; dtxt=' n/a';
    model='human_par2';
    % Choose model between 'human_par1' (original) and 'human_par2' (+ uncertainty)
end
straightsim; finish

```

---

Files used specifically for non-cancer analysis

---

```
% -----
% File: searchsim_refdose.m
% Created by Paul Schlosser, U.S. EPA, 8/2008
% Runs Monte Carlo (MC) simulations WITH search for human equivalent exposures, using variable
% LDAYREFDOSE ('reference dose based on last day of simulated exposure -- presumed "periodicity")
% Requires exposure variable named = expnm = "DRCONC" or "CONC",
% target value = heqt, relative tolerance = hetol, initial set
% exposure (CONC or DRCONC value >0), NRUNS, and rnames (list of
% variables to save)
% PBPK parameters set using MC selection by file human_pars.m.
% -----
% set contsim=1 if continuing an interrupted chain.
if contsim==0 % When starting a new set of simulations
    if ~(expnm=="CONC" | expnm=="DRCONC")
        disp('Variable expnm must be "CONC" or "DRCONC", in double quotes.');
```

```
        return
    end
    if expnm=="CONC" % atxt used as variable command below
        DRCONC=0.0; atxt='CONC = c1'; c2=CONC;
    else
        CONC=0.0; atxt='DRCONC = c1'; c2=DRCONC;
    end
    runo=[]; mns=length(rnames); ns=1; c1s=['c1 = c2*heqt/',dtxt];
    v1s=['v1 = ',dtxt]; v2s=['v2 = ',dtxt];
else % Continuing a set of simulations
    ns=RUNN;
end
for RUNN = ns:NRUNS
    disp(['Remaining runs = ',num2str(NRUNS-RUNN)])
    human_pars
    % Following block calculates daily dose resulting in specified internal dose
    start @NoCallback
    nstep=1; eval(v2s); eval(c1s); eval(atxt);
    start @NoCallback
    eval(v1s); het1=hetol;
    while (abs((v1/heqt)-1)>hetol); % Specify corresponding dose metric
        cn = abs((heqt-v2)*(c1-c2)/(v1-v2) + c2); % Linear interpolation to heqt
        c2=c1; v2=v1; c1=cn; eval(atxt); % Assign values
        start @NoCallback
        eval(v1s);
        nstep=nstep+1;
        if nstep==20
            hetol=hetol*10; nstep=1;
            disp('WARNING! hetol increased to allow convergence!')
        end
    end
    hetol=het1; runo=[runo;eval(sr)];
end
% -----
```

```

% File: human drinking water MCA RfD.m
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 11/2005
% Modified by Paul Schlosser, U.S. EPA 8/2008 and 1/2010
%
% This script file sets up the control parameters to simulate human
% drinking water exposure for a given internal dose (heqt) and calls
% searchsim_refdose which generates a Monte-Carlo chain for the human
% equivalent administered daily dose (HEAD; mg/kg/d).
% -----
exist contsim;
if ~ans
    contsim=0;
end
if contsim==0 % When starting a new set of simulation;
    % set contsim=1 if continuing an interrupted chain.
    clearT
    nm=["DDOSE"]; nn=findnames(nm,rnames);
    if (length(nn) ~= length(nm))
        disp("Not all names in list nm.")
        return
    end
    hetol=1.0e-4; %relative tolerance
    expnm="DRCONC"; DRCONC=9.0; % 35 mg/L is 1 mg/kg/day for 70 kg human drinking 2L
    TEND=95.0; NRUNS=10000; % Total iterations for Monte Carlo analysis
    gendm="female"; % Gender mix; choose "male", "female", or "both"
    agem=70; % Age "mix"; if 0 draws from distribution (0.5-80 years)
    % otherwise agem value is used exclusively
    popn="--"; % GSTGT "++", "+-", "--", or "mix" of +/+, +/-, and -/-
    % dtxt = dose metric to use
    % heqt = target for HEQ search; set value for specific dose metric
    dtxt='LDAYREFDOSE'; dt1='W';
    dtxt='LDAYLIVGSTDOSE'; dt1='G';
    dtxt='LDAYLIVAUC'; dt1='A'; heqt=0.0562;
    dtxt='LDAYLIVCYPDOSE'; dt1='Y'; heqt=15.11;
    model='human_par2';
    % Choose model between 'human_par1' (original) and 'human_par2' (+ uncertainty)
    astp=['RfD_',dt1,'_',num2str(heqt),'_age']; percs=[5 1]; mult=1; gsmult=1;
end
searchsim_refdose; finish

```

---

```

% -----
% File: Human inhalation MCA RfC.m
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 11/2005
% Modified by Paul Schlosser, U.S. EPA, 7/2008 and 1/2010
%
% This script file sets up the control parameters to simulate human
% inhalation exposure for a given internal dose (heqt) and calls
% searchsim_refdose which generates a Monte-Carlo chain for the human
% Equivalent concentration (HEC; mg/m^3)
% -----
exist contsim;
if ~ans
    contsim=0;
end
if contsim==0 % When starting a new set of simulation;
    % set contsim=1 if continuing an interrupted chain.
    use clearT
    nm=["CONC"]; nn=findnames(nm,rnames);
    if (length(nn) ~= length(nm))
        disp("Not all names in list nm.")
        return
    end
    TEND=96; NRUNS=10000; % Total iterations for Monte Carlo analysis
    expnm="CONC"; CONC=67; % CONC in ppm; 0.29 ppm = 1 mg/m3 DCM
    hetol=1.0e-5; %relative tolerance
    gendm="both"; % Gender mix; choose "male", "female", or "both"
    agem=1; % Age "mix"; if 0 draws from distribution (0.5-80 years)
        % otherwise agem value is used exclusively
    popn="--"; % GSTGT "++", "+-", or "mix" of +/+, +/-, and -/-
    gsmult = 1.0; % Multiplies draw for GST individual selection, divides the percentiles at end.
    % dtxt = dose metric to use
    % heqt = target for HEQ search; set value for specific dose metric
    % The last statement below is the metric and target value used.
    % Move desired one to end and save this file before running.
    dtxt='LDAYLIVAUC'; dt1='A'; heqt=0.0562;
    dtxt='LDAYLIVGSTDOSE'; dt1='G'; heqt=10.97;
    dtxt='LDAYREFDOSE'; dt1='W'; heqt=76.71;
    dtxt='LDAYLIVCYPDOSE'; dt1='Y'; heqt=128.1;
    model='human_par2';
        % Choose between 'human_par1' (original) and 'human_par2' (+ uncertainty)
    astp=['RfC_',num2str(heqt),'_age']; mult=GASD*1000; percs=[5 1];
end
searchsim_refdose; finish

```

---

## Files used for mouse dose analyses

---

```
% -----
% File: mouse_set.m
% Programmed by Paul Schlosser, U.S. EPA, 7/2008
% Clears previous variables and sets parameters for mouse simulations.
% -----
% prepare time history values.
prepare @clear @all
CINT=0.1; CONC=0.0; IVDOSE=0.0; BOLUS=0.0; DRCONC=0.0; FIXDRDOSE=0.0;
WESITG=0; WEDITG=0; CC=0;
% from Reitz et al. 1997, Table 1
DRPCT = [0.233, 0.1, 0.1, 0.1, 0.233, 0.234];
DRTIME= [0.0, 4.0, 8.0, 12.0, 16.0, 20.0];
TEND=336;    % 2 weeks
TCHNG=6;    % daily exposure duration = 6 hours/day
TCHNG2=120; % weekly dose width = 5 days/week = 120 hours
TDUR=24;    % daily dose period = 24 hours
TDUR2=168;  % weekly exposure period = 7 days = 168 hours

% U.S. EPA (1988) reference value for B6C3F1 mice: males=0.0373, females=0.0353
BW=0.0373;
% Mouse uptake & metabolism parameters (defined by MCMC calibration)
KA=5.0; VMAXC=9.27; KM=0.574; FRACR=0.0; KFC=1.41; AFFG=0.0; A1=0.207; A2=0.196;
% Mouse physiologic parameters (from prior distributions or MCMC
% calibration)
VLC=0.04; VFC=0.04; VRC=0.05; VLUC=0.0115; VBL2C=0.059; VSC=0.78;
QCC=24.2; VPR=1.45; DSPC=0.15; QLC=0.24; QFC=0.05; QSC=0.19; QRC=0.52;
% Mouse partition coefficients for DCM
PL=1.6; PLU=0.46; PF=5.1; PS=0.44; PR=0.52; PB=23.0;
start @NoCallback

% -----
% File: Mouse drinking water National Coffee 1983.m
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 01/2007
% Modified by Paul Schlosser, U.S. EPA, 8/2008
%
% This run time file sets species-specific constants and exposure
% values for National Coffee Association study of DCM in drinking water
%   in the mouse; parameterized to run with DCM.07.rev3.csl.
% -----
use mouse_set
BW=0.0373; TEND=4*7*24; runo=[];

% Mouse exposure parameters: DDOSE males: 61, 124, 177, or 244 mg/kg-day
for FIXDRDOSE=[61, 124, 177, 244]
    start @NoCallback;
    runo=[runo;[BW,DDOSE,WAVGLIVGSTDOSE,WAVGLUNGGSTDOSE,
WAVGWBDYGSTDOSE]];
end
runo
```

```
save runo @file='mouseOSFdrink.csv' @format=ascii @separator=comma
```

---

```
% -----  
% File: Mouse Inhalation NTP 1986.m  
%  
% Programmed by Michael Lumpkin  
% Syracuse Research Corporation, 11/2007  
% Modified by Paul Schlosser, U.S. EPA, 7/2008  
%  
% This run time file sets species-specific constants and exposure  
% values for NTP (1986) inhalation exposures of DCM in the mouse.  
% -----  
use mouse_set  
CINT=0.01;  
% Kodak value; study average male: 2k=0.034, 4k=0.032; female 2k=0.030, 4k=0.029  
TEND=5*7*24; Bws=[0.034, 0.032; 0.030,0.029]; runo=[]; Cs=[2000, 4000];  
for s=[1 2]  
    for c=[1,2];  
        BW=Bws(s,c); CONC=Cs(c);  
        start @NoCallback  
runo=[runo;[BW,CONC,WAVGLIVGSTDOSE,WAVGLUNGGSTDOSE,WAVGWBDYGSTDOSE]];  
    end  
end  
runo  
%% % Saving last simulation results to file  
run=[_t_cv_cv1_cvf_cvs_cvr_cabl2_wavglivgstdose_wavglunggstdose_ddose];  
save run @file='RUNOUT_NTP_inhal.csv' @format=ascii @separator=comma
```

---

### Files used for rat dose analyses

```
% -----  
% File: rat_set_D.m  
% Programmed by Paul Schlosser, U.S. EPA, 9/2009  
% Clears previous variables and sets parameters for rat simulations.  
% Rat model "D".  
% -----  
% prepare time history values.  
prepare @clear @all  
WESITG=0; WEDITG=0;  
% Rat exposure controls  
CINT=0.01; CONC=0.0; IVDOSE=0.0; DRCONC=0.0; FIXDRDOSE=0.0; BOLUS=0; CC=0;  
DRPCT = [0.233, 0.1, 0.1, 0.1, 0.233, 0.234];  
DRTIME = [0.0, 4.0, 8.0, 12.0, 16.0, 20.0];  
TEND=6; % 7 days  
TCHNG=6; % daily exposure = 6 hours  
TCHNG2=120; % weekly dose width = 5 days/week = 120 hours  
TDUR=24; % daily dose period = 24 hours  
TDUR2=168; % weekly exposure period = 7 days = 168 hours  
  
% Rat UPTAKE & metabolism parameters  
KA=5.0; FRACR=0.0;  
VMAXC=3.66; KM=0.59; KFC=2.97; AFFG=0.0; % MLE fitted
```



A1=0.04; A2=0.14; % from Reitz et al. 1989  
% Rat physiologic parameters (from Andersen et al 1991)  
BW=0.380; % males: 0.380 kg, females: 0.229 kg, from U.S. EPA (1988)  
VLC=0.04; VFC=0.07; VRC=0.05; VLUC=0.0115; VBL2C=0.059; VSC=0.75;  
QCC=15.9; VPR=0.94; DSPC=0.15; QLC=0.20; QFC=0.09; QSC=0.15; QRC=0.56;  
%Rat partition coefficients for DCM (Andersen et al 1991)  
PL=0.732; PLU=0.46; PF=6.19; PS=0.408; PR=0.732; PB=19.4;  
%CO submodel parameters  
DLC=0.060; RENCOC=0.035; ABCOC=0.117; HBTOT=10.0; P1=0.80; F1=1.21;  
M=197.0; COINH=2.2; O2=0.13; PAIR=713.0; RHO=1102.0; SOL=0.03;  
start @NoCallBack  
%RENCOC=0.0;  
VMAXC = 3.923; KM = 0.524; KFC = 2.46; KA=1.8;  
use allfit\_rat-params  
use KAFit-params  
RENCOC=0.0; ABCO=0.0; COINH=0.0;

---

% -----  
% File: rat\_set\_A.m  
% Programmed by Paul Schlosser, U.S. EPA, 9/2009  
% Clears previous variables and sets parameters for rat simulations.  
% Rat model "A".  
% -----  
use rat\_set\_D  
VMAXC = 6.21; KM = 0.23;  
A1 = 0.21; A2 = 0.20;  
KFC = 2.89; KA = 5.0;

---

% -----  
% File: rat\_set\_B.m  
% Programmed by Paul Schlosser, U.S. EPA, 9/2009  
% Clears previous variables and sets parameters for rat simulations.  
% Rat model "B".  
% -----  
use rat\_set\_D  
VMAXC = 4.0; KM = 0.4;  
A1 = 0.0; A2 = 0.0;  
KFC = 2.0; KA = 5.0;

---

% -----  
% File: rat\_set\_C.m  
% Programmed by Paul Schlosser, U.S. EPA, 9/2009  
% Clears previous variables and sets parameters for rat simulations.  
% Rat model "C".  
% -----  
use rat\_set\_D  
VMAXC = 4.0; KM = 0.4;  
A1 = 0.4; A2 = 0.14;  
KFC = 2.0; KA = 4.0;

---

% -----  
% File: Rat inhalation Nitschke 1988a.M  
%

```

% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 10/2007
% Modified by Paul Schlosser, U.S. EPA, 7/2008
%
% This run time file sets species-specific constants values for modified
% Andersen et al (1987, 1991) simulating Nitschke et al. (1988a) exposures
% of DCM in the male and female rats.
% -----
use rat_set_D
TEND=5*7*24; runo=[]; BW=0.229; TCHNG=6; TCHNG2=120;
% males:0.380 kg, females:0.229 kg, from U.S. EPA (1988)
for CONC=[50, 200, 500]
    start @NoCallback
    runo=[runo;[BW,CONC,DIDOSE,WAVGREFDOSE,WAVGLIVGSTDOSE,
        WAVGLIVCYPDOSE,WAVGDAILYAUCL]];
end
runo
save runo @file='RUNOUT_rat_inhal_Nitschke.csv' @format=ascii @separator=comma

```

---

```

% -----
% File: Rat inhalation Burek et al. (1984).m
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 10/2007
% Modified by Paul Schlosser, U.S. EPA, 7/2008
%
% This run time file sets species-specific constants values for modified Andersen et al
% (1987, 1991) simulating Burek et al. (1984) exposures of DCM in male & female rats
% -----
use rat_set_D
TEND=5*7*24; TCHNG=6; TCHNG2=120; runo=[];
for BW=[0.523, 0.338]
    for CONC=[500, 1500, 3500]
        start @NoCallback
        runo=[runo;[BW,DIDOSE,CONC,WAVGREFDOSE,WAVGLIVGSTDOSE,
            WAVGLIVCYPDOSE,WAVGDAILYAUCL]];
    end
end
runo
save runo @file='RUNOUT_rat_inhal_Burek.csv' @format=ascii @separator=comma

```

---

```

% -----
% File: Rat drinking water Serota 1986a.m
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 1/2007
% Modified by Paul Schlosser, U.S. EPA, 7/2008
%
% This run time file sets species-specific constants values for modified Andersen et al
% (1987, 1991) simulating Serota et al. (1986a) exposures of DCM to male & female rats
% -----
use rat_set_D
%DRCONC values from Serota et al. (1986a)

```

```

% males:0, 44.00, 381.36, 916.74, 1723.47
% females: 0, 37.80, 365.41, 856.82, 1656.94
TEND=7*24; runo=[]; BW=0.380; % Males
for FIXDRDOSE=[6, 52, 125, 235]
    start @NoCallback
    runo=[runo;[BW,DDOSE,LDAYREFDOSE,LDAYLIVGSTDOSE,LDAYLIVCYPDOSE,
        LDAYLIVAUC]];
end
BW=0.229; % Females
for FIXDRDOSE=[6, 58, 136, 263]
    start @NoCallback
    runo=[runo;[BW,DDOSE,LDAYREFDOSE,LDAYLIVGSTDOSE,LDAYLIVCYPDOSE,
        LDAYLIVAUC]];
end
runo
save runo @File="Serota_rat_DW.csv" @Format=ascii @Separator=comma

```

---

```

% -----
% File: Rat inhalation NTP 1986.M
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 11/2007
% Modified by Paul Schlosser, U.S. EPA, 7/2008
%
% This run time file sets species-specific constants values for modified Andersen et al
% (1987, 1991) simulating NTP (1986) exposures of DCM to male & female rats
% -----
use rat_set_D
CINT=0.1;
bws=[0.3905, 0.3852, 0.3848, 0.2455, 0.2443, 0.2422];
TEND=6*24*7; concs=[1 2 4 1 2 4]*1000; runo=[];
for n=1:length(bws)
    BW=bws(n); CONC=concs(n); start @NoCallback
    runo=[runo;[BW,CONC,DIDOSE,WAVGLIVGSTDOSE,WAVGAUCV,WAVGAUCS]];
end
runo
save runo @File="RUNOUT_rat_inhal_NTP.csv" @Format=ascii @Separator=comma

```

---

```

% -----
% File: Figure 5_3.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Creates dichloromethane exposure-dose relationship for humans
% vs. the rat shown in Figure 5-3 of the IRIS assessment
% -----
use rat_set_D
ods=[10:10:100,120,140,170,200,220,250];
TEND=7*24; ratr=[]; BW=0.380; % Male rats
for FIXDRDOSE=ods
    start @NoCallback
    ratr=[ratr;[LDAYLIVGSTDOSE,LDAYLIVCYPDOSE,LDAYREFDOSE]];
    disp(['Rat, oral dose = ',num2str(FIXDRDOSE),', livGSTdose =
',num2str(LDAYLIVGSTDOSE),'.']);
end

```

```

end

clearT; TEND=95.0; humr=[]; contsim=0;
NRUNS=1000; % NRUNS = Total iterations for Monte Carlo analysis
gendm="both"; % Gender mix; choose "male", "female", or "both"
agem=0; % Age "mix"; if 0 draws from distribution (0.5-80 years)
% otherwise agem value is used exclusively
popn="mix"; gsmult=1; % GSTGT "++", "+-", or "mix" of +/+, +/-, and -/-
model='human_par2'; % Choose between 'human_par1' (original) and 'human_par2' (+
uncertainty)
nm=["LDAYLIVGSTDOSE";"LDAYLIVCYPDOSE";"LDAYREFDOSE"];
nn=findnames(nm,rnames);
    if (length(nn) ~= length(nm))
        disp("Not all names in list nm.")
        return
    end
for FIXDRDOSE=ods
    rres=[FIXDRDOSE]; straight sims;
    for n=1:length(nm)
        rres=[rres,[mean(runo(:,nn(n))),prctile(runo(:,nn(n)),[5 95])]];
    end
    disp(['Human, dose = ',num2str(FIXDRDOSE),', internal doses = ...']); rres
    humr=[humr;rres];
end
save @File="Fig5_3.mat"
plot(ods,ratr(:,1),ods,humr(:,2),ods,humr(:,4),'Fig5_3a.aps');
plot(ods,ratr(:,2),ods,humr(:,5),ods,humr(:,6),ods,humr(:,7),'Fig5_3b.aps');
plot(ods,ratr(:,3),ods,humr(:,8),ods,humr(:,9),ods,humr(:,10),'Fig5_3c.aps');

% -----
% File: Figure 5_7.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Creates dichloromethane exposure-dose relationship for humans
% vs. the rat shown in Figure 5-7 of the IRIS assessment
% -----
use rat_set_D
ccs=[10:10:60,80,100,150,200:100:600,800,1000]; %,1500,2000:1000:5000];
TEND=2*7*24; ratr=[]; BW=0.229; TCHNG=6; TCHNG2=120;
for CONC=ccs
    start @NoCallback
        ratr=[ratr;[CONC,WAVGLIVGSTDOSE,WAVGLIVCYPDOSE,WAVGREFDOSE]];
    disp(['Rat, conc = ',num2str(CONC),', livGSTdose = ',num2str(WAVGLIVGSTDOSE),'.']);
end
save ratr @file='Fig5_7-rat.csv' @format=ascii @separator=comma
clearT; TEND=95.0; huml=[]; contsim=0;
NRUNS=1000; % NRUNS = Total iterations for Monte Carlo analysis
gendm="both"; % Gender mix; choose "male", "female", or "both"
agem=0; % Age "mix"; if 0 draws from distribution (0.5-80 years)
% otherwise agem value is used exclusively

```

```

popn="mix"; gsmult=1; % GSTGT "++", "+-", or "mix" of +/+, +/-, and -/-
model='human_par2';
    % Choose between 'human_par1' (original) and 'human_par2' (+ uncertainty)
nm=["LDAYLIVGSTDOSE","LDAYLIVCYPDOSE","LDAYREFDOSE";
    "LDAYLUNGGSTDOSE"];
nn=findnames(nm,rnames);
    if (length(nn) ~= length(nm))
        disp("Not all names in list nm.")
        return
    end
for CONC=ccs
    rres=[CONC]; straightsim;
    for n=1:length(nm)
        rres=[rres,[mean(runo(:,nn(n))),prctile(runo(:,nn(n)),[5 95])]];
    end
    disp(['Human, conc = ',num2str(CONC),', internal doses = ...']); rres
    hum1=[hum1; rres];
end
save hum1 @file='Fig5_7-human.csv' @format=ascii @separator=comma
save @File="Fig5_7.mat"
plot(ccs,ratr(:,2),ccs,hum1(:,2),ccs,hum1(:,4),'Fig5_7a.aps')
plot(ccs,ratr(:,3),ccs,hum1(:,5),ccs,hum1(:,6),ccs,hum1(:,7),'Fig5_7b.aps')
plot(ccs,ratr(:,4),ccs,hum1(:,8),ccs,hum1(:,9),ccs,hum1(:,10),'Fig5_7c.aps')

```

---

```

% -----
% File: Figure 5_9n10_rat_sense.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Performs sensitivity analysis for rat oral (Figure 5-9) and inhalation
% (Figure 5-10) dichloromethane exposures, and writes results to file
% 'ratsense_Figs59n10.csv' for plotting in Excel
% -----
use rat_set_D

FIXDRDOSE=10; TEND=7*24; ratr=[]; BW=0.380; % Male rats
start @nocallback
r0=[LDAYLIVGSTDOSE,LDAYLIVCYPDOSE,LDAYREFDOSE];
pars=["QCC","VPR","VLC","VSC","PB","VMAXC","KA","A2","KFC"];
for pt=pars
    p0=eval(pt); setbase(pt,1.01*p0); start @nocallback
    r=[LDAYLIVGSTDOSE,LDAYLIVCYPDOSE,LDAYREFDOSE];
    setbase(pt,0.99*p0); start @nocallback
    r= 50*(r-[LDAYLIVGSTDOSE,LDAYLIVCYPDOSE,LDAYREFDOSE])./r0;
    setbase(pt,p0);
    ratr=[ratr;r];
end

FIXDRDOSE=0; CONC=500; TEND=5*7*24; BW=0.229; TCHMG=6; TCHNG2=120;
start @nocallback
r0=[WAVGLIVGSTDOSE,WAVGLIVCYPDOSE,WAVGREFDOSE];

```

```

for pt=pars
    p0=eval(pt); setbase(pt,1.01*p0); start @nocallback
    r=[WAVGLIVGSTDOSE,WAVGLIVCYPDOSE,WAVGREFDOSE];
    setbase(pt,0.99*p0); start @nocallback
    r= 50*(r-[WAVGLIVGSTDOSE,WAVGLIVCYPDOSE,WAVGREFDOSE])./r0;
    setbase(pt,p0);
    ratr=[ratr;r];
end

ratr
save ratr @file="ratsense_Figs59n10.csv" @format=ascii @separator=comma

% -----
% File: Figure 5_14.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Creates dichloromethane exposure-dose relationship for humans vs. the mouse
% shown in Figure 5-14 of the IRIS assessment; results are written to file
% "Fig5-14_human.csv" for plotting in Excel
% -----
use mouse_set
ods=[10:10:100,120,140,170,200,220,250];
TEND=7*24; mour=[]; BW=0.033; TCHNG=2000; TCHNG2=2000;
model='human_par2';
for FIXDRDOSE=ods
    start @NoCallBack
    mour=[mour;WAVGLIVGSTDOSE];
    disp(['Mouse, dose = ',num2str(FIXDRDOSE),',
        livGSTdose = ',num2str(WAVGLIVGSTDOSE),'.']);
end
save mour @file="Fig5-14_mouse.csv" @format=ascii @separator=comma
clearT; TEND=95.0; hum2=[]; conssim=0;
NRUNS=1000; % NRUNS = Total iterations for Monte Carlo analysis
gendm="both"; % Gender mix; choose "male", "female", or "both"
agem=0; % Age "mix"; if 0 draws from distribution (0.5-80 years)
% otherwise agem value is used exclusively
gsmult=1; nm=["LDAYLIVGSTDOSE"]; nn=findnames(nm,rnames);
if isempty(nn)
    disp("Variable name not in list nm.")
    return
end
for FIXDRDOSE=ods
    rres=[FIXDRDOSE]
    for popn=["mix","+","++"] % GSTGT "+", "+-", or "mix" of +/+, +/-, and -/-
        strightsims;
        for n=1:length(nm)
            rres=[rres,[mean(runo(:,nn(n))),prctile(runo(:,nn(n)),[5 95])]]
        end
        disp(['Human',ctot(popn),', dose = ',num2str(FIXDRDOSE),', internal doses = ...']); rres
    end
end

```

```

        hum2=[hum2; rres];
end
save hum2 @file="Fig5-14_human.csv" @format=ascii @separator=comma
save @file="Fig5_14.mat"
plot(ods,mour,ods,hum2(:,2),ods,hum2(:,4), ods,hum2(:,5),ods,hum2(:,6),ods,hum2(:,7), ...
      ods,hum2(:,8),ods,hum2(:,9),ods,hum2(:,10),'Fig5_14.aps')

```

---

```

% -----
% File: Figure 5_15.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Creates dichlromethane exposure-dose relationship for humans vs. the
% mouse shown in Figure 5-15 (panels A and B) of the IRIS assessment;
% results are written to file "Fig5-15_human.csv" for plotting in Excel
% -----
use mouse_set
ccs=[10:10:100,120,150,200,230,300,330,400,430,500:100:1000,1300,2000:1000:5000];
TEND=5*7*24; mour=[]; BW=0.031; TCHNG=6; TCHNG2=120;
for CONC=ccs
    start @NoCallBack
        mour=[mour;[CONC,WAVGLIVGSTDOSE,WAVGLUNGGSTDOSE]];
    disp(['Mouse, conc = ',num2str(CONC),', livGSTdose = ',num2str(WAVGLIVGSTDOSE),'.']);
end

clearT; TEND=95.0; hum3=[]; contsim=0;
NRUNS=1000;      % NRUNS = Total iterations for Monte Carlo analysis
gendm="both";    % Gender mix; choose "male", "female", or "both"
agem=0;         % Age "mix": draws from distribution (0.5-80 years)
model='human_par2'; gsmult=1;
nm=["LDAYLIVGSTDOSE";"LDAYLUNGGSTDOSE"]; nn=findnames(nm,rnames);
    if (length(nn) ~= length(nm))
        disp("Not all names in list nm.")
        return
    end
for CONC=ccs
    rres=[CONC];
    for popn=["+","++"] % GSTGT "++", "+-", or "mix" of +/+, +/-, and -/-
        strightsims;
        for n=nn
            rres=[rres,[mean(runo(:,n)),prctile(runo(:,n),[5 95])]];
        end
        if popn=="mix"
            for ij=[0,1]
                runo2=runo(runo(:,31)==ij,:);
                for n=1:length(nm)
                    rres=[rres,[mean(runo2(:,nn(n))),prctile(runo2(:,nn(n)),[5 95])]];
                end
            end
        end
    end
    disp(['Human, conc = ',num2str(CONC),', internal doses = ...']); rres

```

```

        end
        hum3=[hum3;rres];
end
save hum3 @file="Fig5-15_human.csv" @format=ascii @separator=comma
save @File="Fig5_15.mat"
% Liver dose plot...
% plot(ccs,mour(:,2),hum1(:,1),hum1(:,2),hum1(:,1),hum1(:,4), ...
%     ccs,hum3(:,2),ccs,hum3(:,3),ccs,hum3(:,4), ...
%     ccs,hum3(:,8),ccs,hum3(:,9),ccs,hum3(:,10),'Fig4_3.aps')
plot(ccs,mour(:,2),hum1(:,1),hum1(:,2),ccs,hum3(:,2),ccs,hum3(:,8),'Fig5_15a.aps')
% Lung dose plot...
% plot(ccs,mour(:,1),hum1(:,1),hum1(:,11),hum1(:,1),hum1(:,13), ...
%     ccs,hum3(:,5),ccs,hum3(:,6),ccs,hum3(:,7), ...
%     ccs,hum3(:,11),ccs,hum3(:,12),ccs,hum3(:,13),'Fig4_4.aps')
plot(ccs,mour(:,3),hum1(:,1),hum1(:,11),ccs,hum3(:,5),ccs,hum3(:,11),'Fig5_15b.aps')

% -----
% File: Fig5_16to18_mouse_sense.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Runs sensitivity analyses for dichloromethane exposures in mice,
% producing results plotted in Figures 5-18 to 5-18 of the IRIS assessment.
% Values saved to file "mousesense_Figs5_16to18.csv" for plotting in Excel.
% -----
use mouse_set
CINT=0.001; TEND=(3*7*24)-CINT;
CONC=2000; TCHNG=6; TCHNG2=120; mour=[]; BW=0.031; % average mouse
start @nocallback
r0=WAVGLIVGSTDOSE;
pars=["QCC","VPR","VLC","VSC","PB","VMAXC","KA","A2","KFC"];
for pt=pars
    p0=eval(pt);
    setbase(pt,1.01*p0); start @nocallback
    r=WAVGLIVGSTDOSE;
    setbase(pt,0.99*p0); start @nocallback
    r= 50*(r-WAVGLIVGSTDOSE)/r0;
    setbase(pt,p0);
    mour=[mour,r];
end

DRCONC=500; CONC=0; start @nocallback
r0=LDAYLIVGSTDOSE; m5=[];
for pt=pars
    p0=eval(pt); setbase(pt,1.01*p0); start @nocallback
    r=LDAYLIVGSTDOSE;
    setbase(pt,0.99*p0); start @nocallback
    r= 50*(r-LDAYLIVGSTDOSE)/r0;
    setbase(pt,p0);
    m5=[m5,r];
end

```



```

mour=[mour;m5];

CONC=500; DRCONC=0; start @nocallback
r0=WAVGLUNGGSTDOSE; m5=[];
for pt=pars
    p0=eval(pt); setbase(pt,1.01*p0); start @nocallback
    r=WAVGLUNGGSTDOSE;
    setbase(pt,0.99*p0); start @nocallback
    r= 50*(r-WAVGLUNGGSTDOSE)/r0;
    setbase(pt,p0);
    m5=[m5,r];
end
mour=[mour;m5]
save mour @file="mousesense_Figs5_16to18.csv" @format=ascii @separator=comma

```

---

```

% -----
% File: FigC3 rat inhal Gargas 86.m
% Figure C-3 (creates 4 sub-plots)
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 10/2007
% Modified by Paul Schlosser, U.S. EPA, 9/2009
%
% This run time file sets species-specific constants values for modified
% Andersen et al (1987, 1991) simulating Burek et al. (1984) exposures
% of dichloromethane in the male and female rats
% -----
dch1=[0.103, 93.73; 0.2178, 81.12; 0.3707, 66.26; 0.5543, 52.58; 0.7227, 41.73;
      0.8834, 34.085; 1.044, 27.84; 1.22, 22.095; 1.3807, 18.312; 1.55, 14.532;
      1.717, 12.58; 1.893, 10.28; 2.061, 8.3948];

dch5=[0.1015, 448.9; 0.216, 400.0; 0.3843, 326.7; 0.5522, 282.85; 0.7128, 234.4;
      0.8806, 208.9; 1.056, 183.5; 1.231, 163.5; 1.384, 145.7; 1.5514, 133.7;
      1.719, 119.14; 1.887, 106.2; 2.0395, 95.99; 2.192, 85.5385; 2.360, 75.13;
      2.5276, 65.99; 2.688, 57.96; 2.8484, 49.45; 3.024, 40.99; 3.1996, 36.00;
      3.360, 31.165; 3.5356, 25.46; 3.719, 21.10; 3.864, 18.00; 4.025, 15.585;
      4.193, 13.297; 4.353, 11.51];

dch10=[0.0973, 927.1; 0.2195, 826.0; 0.3957, 627.6; 0.5556, 575.7; 0.7235, 513.1;
       0.8836, 457.2; 1.059, 419.5; 1.226, 401.9; 1.3936, 379.6; 1.561, 348.2;
       1.721, 333.66; 1.873, 329.1; 2.0556, 301.9; 2.200, 285.1; 2.3754, 265.4;
       2.5426, 258.0; 2.695, 236.7; 2.87, 226.8; 3.030, 211.1; 3.1975, 193.66;
       3.365, 182.9; 3.532, 175.24; 3.7076, 156.175; 3.875, 145.4; 4.035, 135.3;
       4.195, 124.1; 4.340, 112.2; 4.515, 101.5; 4.683, 89.12];

dch30=[0.1058, 2958.6; 0.2288, 2280; 0.3819, 1862.5; 0.5419, 1684; 0.7095, 1545;
       0.8848, 1397; 1.067, 1338.6; 1.227, 1264; 1.387, 1229; 1.5615, 1195;
       1.7136, 1178.5; 1.881, 1129; 2.056, 1051; 2.208, 1052; 2.383, 1008;
       2.5424, 994.2; 2.6945, 966.46; 2.869, 939.6; 3.0366, 913.45; 3.196, 914.2;

```

```
3.356, 863.3; 3.531, 851.6; 3.713, 816.1; 3.880, 805.0; 4.032, 794.0;
4.192, 760.75; 4.352, 750.4; 4.5114, 708.6; 4.686, 709.2; 4.853, 699.6;
5.0055, 670.3; 5.173, 642.26; 5.34, 633.53];
```

```
for model=['A' 'B' 'C' 'D']
    eval(['use rat_set_',model]);
    BW=0.225; TEND=6; TCHNG=7; CC=1; VCHC=9; NCH=3; CINT=0.05;
    prepare @clear T CCHPPM
    res=[];
    for CONC=[107, 498, 1028, 3206]
        start @nocallback
        res=[res,_t,_cchppm];
    end
    eval(['save res @file=gargas-inh-sim-',model,'.csv @format=ascii @separator=comma'])
    plot(dch1(:,1),dch1(:,2),dch5(:,1),dch5(:,2),dch10(:,1),dch10(:,2), ...
        dch30(:,1),dch30(:,2),res(:,1),res(:,2),res(:,3),res(:,4), ...
        res(:,5),res(:,6),res(:,7),res(:,8),'gargas86',model,'.aps'])
end
```

```
dch1=[dch1;dch5;dch10;dch30];
save dch1 @file='gargas-inh-dat.csv' @format=ascii @separator=comma
```

---

```
% -----
% File: FigC4n5 Angelo_IV_comp.m
% Figures C-4 and C-5 (creates up to 4 sub-plots for each)
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 10/2007
% Modified by Paul Schlosser, U.S. EPA, 9/2009
%
% This run time file sets species-specific constants values for the
% modified Andersen et al (1987, 1991) PBPK model for dichloromethane,
% simulating Angelo et al. (1986b) IV exposures of DCM in rats
% -----
d10bl=[2, 17.624; 5, 8.6635; 10, 3.533; 15, 2.858; 20, 1.7125;
30, 0.826; 40, 0.6192];
d50bl=[2, 62.686; 5, 34.105; 10, 16.56; 15, 12.33; 20, 12.80;
30, 10.25; 40, 5.561];
d10ex=[0.3333, 32.245; 0.6667, 38.56; 1, 41.396; 4, 44.42];
d50ex=[0.3333, 35.38; 0.6667, 45.95; 1, 51.38; 4, 57.59];

for model='D' %['A' 'B' 'C' 'D'] % Use set of letters for models to test
    eval(['use rat_set_',model]);
    prepare @clear T CV PCTIVEXH
    TEND=10; CINT=0.01;
    IVDOSE=10
start @nocallback
    res=[_t,_cv,_pctivexh]; r2=[_t*60,_cv,_t,_pctivexh];
    IVDOSE=5; start @nocallback
```

```

res=[res,_t,_cv,_pctivexh]; r2=[r2,_t*60,_cv,_t,_pctivexh];
IVDOSE=25; start @nocallback
res=[res,_t,_cv,_pctivexh]; r2=[r2,_t*60,_cv,_t,_pctivexh];
IVDOSE=50; start @nocallback
res=[res,_t,_cv,_pctivexh];r2=[r2,_t*60,_cv,_t,_pctivexh];
plot(d10bl(:,1),d10bl(:,2),d50bl(:,1),d50bl(:,2), ...
      res(:,1)*60,res(:,2),_t*60,_cv,res(:,4)*60,res(:,5), ...
      res(:,7)*60,res(:,8),['angivbl',model,'aps'])
plot(d10ex(:,1),d10ex(:,2),d50ex(:,1),d50ex(:,2), ...
      res(:,1),res(:,3),_t,_pctivexh,res(:,4),res(:,6), ...
      res(:,7),res(:,9),['angivex',model,'aps'])
eval(['save r2 @file=Angelo_IV_sims_',model,'.csv @format=ascii @separator=comma'])
end

```

---

```

% -----
% File: FigC6 rat_inhal_Andersen.m
% Figures C-6 (creates 4 sub-plots)
%
% Programmed by Michael Lumpkin
% Syracuse Research Corporation, 10/2007
% Modified by Paul Schlosser, U.S. EPA, 9/2009
%
% This run time file sets species-specific constants values for the
% modified Andersen et al (1987, 1991) PBPK model for dichloromethane,
% simulating Andersen et al. (1987) inhalation exposures of DCM in rats
% -----
dan2=[1, 2.6634; 2, 4.290; 3, 4.9915; 4, 4.464; 4.25, 1.3334; 4.5, 0.5513;
      4.75, 0.4234; 5, 0.2618; 5.25, 0.1206; 5.5, 0.1049];
dan10=[1, 29.34; 2, 49.50; 3, 47.10; 4, 50.72; 4.25, 14.92; 4.5, 9.225;
      4.75, 6.260; 5, 3.811; 5.25, 1.3085; 5.5, 0.6515; 5.75, 0.6025];
can2=[0.9942, 3.967; 1.027, 4.121; 1.039, 3.798; 2.045, 7.151;
      2.0565, 6.598; 2.057, 5.7836; 3.053, 8.4295; 3.076, 7.493;
      3.086, 8.7215; 4.073, 9.094; 4.074, 8.003; 4.084, 8.940; 4.5545, 5.302;
      4.597, 6.853; 4.608, 7.1145; 5.078, 3.5525; 5.079, 2.723; 5.0795, 2.124;
      5.302, 1.464; 5.313, 1.971; 5.325, 1.142];
can10=[1.006, 4.004; 1.018, 4.305; 1.021, 5.114; 2.0494, 9.257;
      2.061, 9.488; 2.068, 7.9395; 3.047, 10.42; 3.0535, 8.777; 3.055, 9.170;
      4.030, 6.449; 4.030, 10.31; 4.036, 12.11; 5.0275, 7.448; 5.061, 7.656;
      5.067, 9.598; 5.5285, 4.978; 5.540, 5.001; 5.552, 9.115; 6.028, 1.999;
      6.030, 6.460; 6.049, 1.653; 6.510, 0.4305; 6.525, 2.072; 6.539, 2.973];
for model=['A' 'B' 'C' 'D']
eval(['use rat_set_',model]);
QCSW=0;
prepare @clear T CV PCTHBCO
TEND=6; CINT=0.01; TCHNG=4.0;
CONC=200; start @nocallback
res=[_t,_cv,_pcthbco];
if model=='D'
CONC=150; start @nocallback

```

```

        res=[res,_cv];
        %use rat_set_D
        %QCSW=0;
    end

    CONC=750; CINT=0.01; TCHNG=4.0; start @nocallback
    res2=[_t,_cv]; res=[res,res2];
    TEND=7; CONC=1014; start @nocallback
    plot(dan2(:,1),dan2(:,2),dan10(:,1),dan10(:,2), ...
        res(:,1),res(:,2),res2(:,1),res2(:,2),res(:,1),res(:,4),['andinhbl',model,'.aps'])
    %plot(can2(:,1),can2(:,2),can10(:,1),can10(:,2), ...
    %    res(:,1),res(:,3),_t,_pcthbco,['andinhhb',model,'.aps'])
eval(['save res @file=Andersen87_inh_sims_',model,'.csv @format=ascii @separator=comma'])
end

```

---

```

% -----
% File: FigC7 rat inhal exp_dose.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Creates exposure-dose relationship for rat inhalation exposures to
% dichloromethane, for models B and D, as shown in Figure C-7 of the
% IRIS assessment
% -----

```

```

use rat_set_D
prepare @clear T CV PCTHBCO

bwm=mean([0.3905, 0.3852, 0.3848, 0.2455, 0.2443, 0.2422]);
res=[0,0,0,0,0];
for C=[10:10:100, 100:20:300, 400:100:1000, 1200:200:2000]
    use rat_set_B
    CONC=C; TEND=6*24*7; CINT=0.1; TCHNG=6.0; BW=bwm;
    start @nocallback
    r=[CONC,WAVGLIVGSTDOSE,WAVGLIVCYPDOSE];
    use rat_set_D
    CONC=C; TEND=6*24*7; CINT=0.1; TCHNG=6.0; BW=bwm;
    start @nocallback
    res=[res;[r,WAVGLIVGSTDOSE,WAVGLIVCYPDOSE]];
end
plot(res(:,1),res(:,2),res(:,1),res(:,3), ...
    res(:,1),res(:,4),res(:,1),res(:,5),'ratlinear.aps')

```

---

```

% -----
% File: FigC8n9 rat_oral_fits.m
% Programmed by Paul Schlosser, U.S. EPA, 9/2009
% Creates plots shown in Figure C-8 (4 panels) and Figure C-9 of the
% dichloromethane IRIS assessment. Figure C-8 shows model D fits
% to data of Angelo et al (1986b) for 50 and 200 mg/kg bolus oral
% exposures. Figure C-9 is fit of model D with KA = 0.47 or = 1.8.
% -----
use rat_set_D

```

```

use Kfit-params
KA = 1.8; % Comment this line out to use one above, if alternate fitted
% value is available
prepare @clear T CV CL PCTIVEXH PCTIVCOEXH PCTHBCO ACO BO2 CHBT HBCO
TEND=25; BW=0.25; CINT=0.05;
RENCOC=0.0; ABCO=0.0; COINH=0.0;

% Angelo et al. (1986b) rat timecourse data from oral dosing of 50 or 200 mg/kg
% time (hr), blood DCM, liver DCM, % exhaled DCM, % exh CO
load @file=angelo_86_50mg.csv @Format=Ascii; da50=angelo_86_50mg; % 50 mg/kg
load @file=angelo_86_200mg.csv @Format=Ascii; da200=angelo_86_200mg; % 200 mg/kg

% Pankow et al. '91; COHb following a single gavage dose of
% 6.2 mmol/kg DCM (526 mg/kg) in 259-gram male Wistar rats
load @file=Pankow_91_526mgHb.csv @Format=Ascii; dpank=Pankow_91_526mgHb;

BOLUS=50; start @nocallback
ra50=[_t, _cv, _cl, _pctivexh, _pctivcoexh];

BOLUS=200; start @nocallback
ra200=[_t, _cv, _cl, _pctivexh, _pctivcoexh];

plot(da50(:,1),da50(:,2),ra50(:,1),ra50(:,2),da200(:,1),da200(:,2),ra200(:,1),ra200(:,2),'angbl.aps')
plot(da50(:,1),da50(:,3),ra50(:,1),ra50(:,3),da200(:,1),da200(:,3),ra200(:,1),ra200(:,3),'angli.aps')
plot(da50(:,1),da50(:,4),ra50(:,1),ra50(:,4),da200(:,1),da200(:,4),ra200(:,1),ra200(:,4),
'angexd.aps')
plot(da50(:,1),da50(:,5),ra50(:,1),ra50(:,5),da200(:,1),da200(:,5),ra200(:,1),ra200(:,5),
'angexc.aps')

TEND=12.5; BOLUS=526; start @nocallback
ps=[_t,_pcthbco];
kasave=KA;
KA=0.47; start @nocallback
KA=kasave;
plot(dpank(:,1),dpank(:,2),ps(:,1),ps(:,2),_t,_pcthbco,'pankhh.aps')
ra50=[ra50,ra200];
save ra50 @file=Angelo-oral-sim.csv @format=ascii @separator=comma
da50=[da50;da200];
save da50 @file=Angelo-oral-dat.csv @format=ascii @separator=comma

```