



# TOXICOLOGICAL REVIEW

OF

## Trichloroethylene

(CAS No. 79-01-6)

**In Support of Summary Information on the  
Integrated Risk Information System (IRIS)**

*June 2011*

### NOTICE

This document is a Final Agency/Interagency **Review draft**. This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by U.S. EPA. It does not represent and should not be construed to represent any Agency determination or policy. It is being circulated for review of its technical accuracy and science policy implications.

U.S. Environmental Protection Agency  
Washington, DC

## **DISCLAIMER**

This document is a preliminary draft for review purposes only. This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by EPA. It does not represent and should not be construed to represent any Agency determination or policy. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

## GUIDE TO READERS OF THIS DOCUMENT

**Due to the length of the TCE toxicological review, it is recommended that Chapters 1 and 6 be read prior to Chapters 2–5.**

Chapter 1 is the standard introduction to an IRIS Toxicological Review, describing the purpose of the assessment and the guidelines used in its development.

Chapter 2 is an exposure characterization that summarizes information about TCE sources, releases, media levels and exposure pathways for the general population (occupational exposure is also discussed to a lesser extent).

Chapter 3 describes the toxicokinetics and physiologically based pharmacokinetic (PBPK) modeling of TCE and metabolites (PBPK modeling details are in Appendix A).

Chapter 4 is the hazard characterization of TCE. Section 4.1 summarizes the evaluation of epidemiologic studies of cancer and TCE (qualitative details in Appendix B; meta-analyses in Appendix C). Each of the Sections 4.2–4.9 provides self-contained summary and syntheses of the epidemiologic and laboratory studies on TCE and metabolites, organized by tissue/type of effects, in the following order: genetic toxicity, central nervous system (CNS), kidney, liver, immune system, respiratory tract, reproduction and development, and other cancers. Additional details are provided in Appendix D for CNS effects and Appendix E for liver effects. Section 4.10 summarizes the available data on susceptible lifestages and populations. Section 4.11 describes the overall hazard characterization, including the weight of evidence for noncancer effects and for carcinogenicity.

Chapter 5 is the dose-response assessment of TCE. Section 5.1 describes the dose-response analyses for noncancer effects, and Section 5.2 describes the dose-response analyses for cancer. Additional computational details are described in Appendix F for noncancer dose-response analyses, Appendix G for cancer dose-response analyses based on rodent bioassays, and Appendix H for cancer dose-response analyses based on human epidemiologic data.

Chapter 6 is the summary of the major conclusions in the characterization of TCE hazard and dose response.

**CONTENTS of TOXICOLOGICAL REVIEW for TRICHLOROETHYLENE  
(CAS No. 79-01-6)**

<b>TOXICOLOGICAL REVIEW OF Trichloroethylene</b> .....	i
<b>DISCLAIMER</b> .....	ii
<b>GUIDE TO READERS OF THIS DOCUMENT</b> .....	iii
<b>CONTENTS of TOXICOLOGICAL REVIEW for TRICHLOROETHYLENE</b> .....	iv
<b>LIST OF TABLES</b> .....	xv
<b>LIST OF FIGURES</b> .....	xxv
<b>LIST OF ABBREVIATIONS AND ACRONYMS</b> .....	xxxii
<b>FOREWORD</b> .....	xxxviii
<b>AUTHORS, CONTRIBUTORS, AND REVIEWERS</b> .....	xxxix
<b>EXECUTIVE SUMMARY</b> .....	xliv
<b>1. INTRODUCTION</b> .....	1
<b>2. EXPOSURE CHARACTERIZATION</b> .....	1
1     2.1. ENVIRONMENTAL SOURCES.....	2
2     2.2. ENVIRONMENTAL FATE.....	7
2.2.1. Fate in Terrestrial Environments .....	7
2.2.2. Fate in the Atmosphere .....	7
2.2.3. Fate in Aquatic Environments .....	8
3     2.3. EXPOSURE CONCENTRATIONS.....	8
2.3.1. Outdoor Air—Measured Levels .....	8
2.3.2. Outdoor Air—Modeled Levels.....	9
2.3.3. Indoor Air.....	12

*This document is a draft for review purposes only and does not constitute Agency policy.*

	2.3.4. Water.....	15
	2.3.5. Other Media .....	17
	2.3.6. Biological Monitoring.....	19
1	2.4. EXPOSURE PATHWAYS AND LEVELS .....	19
	2.4.1. General Population.....	19
2	2.4.1.1.1. Inhalation .....	20
3	2.4.1.1.2. Ingestion.....	21
4	2.4.1.1.3. Dermal.....	22
5	2.4.1.1.4. Exposure to Trichloroethylene (TCE) Related Compounds.....	23
	2.4.2. Potentially Highly Exposed Populations .....	24
6	2.4.2.1.1. Occupational Exposure .....	24
7	2.4.2.1.2. Consumer Exposure .....	26
	2.4.3. Exposure Standards.....	26
8	2.5. EXPOSURE SUMMARY .....	27
	<b>3. TOXICOKINETICS .....</b>	<b>1</b>
9	3.1. ABSORPTION .....	2
	3.1.1. Oral .....	2
	3.1.2. Inhalation .....	4
	3.1.3. Dermal.....	11
10	3.2. DISTRIBUTION AND BODY BURDEN .....	13
11	3.3. METABOLISM .....	24
	3.3.1. Introduction.....	24
	3.3.2. Extent of Metabolism.....	25
	3.3.3. Pathways of Metabolism.....	28
12	3.3.3.1.1. Cytochrome P450-Dependent Oxidation.....	28
13	3.3.3.1.3. Formation of chloral hydrate (CH), trichloroethanol (TCOH)	
14	and trichloroacetic acid (TCA) .....	33
15	3.3.3.1.8. Glutathione (GSH) Conjugation Pathway .....	46
16	3.3.3.1.17. Relative Roles of the Cytochrome P450 (CYP) and	
17	Glutathione (GSH) Pathways .....	64
18	3.4. TRICHLOROETHYLENE (TCE) EXCRETION .....	68
	3.4.1. Exhaled Air .....	68
	3.4.2. Urine .....	71
	3.4.3. Feces .....	73
19	3.5. PHYSIOLOGICALLY BASED PHARMACOKINETIC (PBPK)	
20	MODELING OF TRICHLOROETHYLENE (TCE) AND ITS	
21	METABOLITES.....	74
	3.5.1. Introduction.....	74
	3.5.2. Previous Physiologically Based Pharmacokinetic (PBPK) Modeling of	
	Trichloroethylene (TCE) for Risk Assessment Application.....	74

*This document is a draft for review purposes only and does not constitute Agency policy.*

	3.5.3. Development and Evaluation of an Interim “Harmonized” Trichloroethylene (TCE) Physiologically Based Pharmacokinetic (PBPK) Model .....	76
1	3.5.4. Physiologically Based Pharmacokinetic (PBPK) Model for Trichloroethylene (TCE) and Metabolites Used for This Assessment .....	77
2	3.5.4.1.1. Introduction .....	77
3	3.5.4.1.2. Updated Physiologically Based Pharmacokinetic (PBPK) Model Structure .....	80
4	3.5.4.1.3. Specification of Baseline Physiologically Based Pharmacokinetic (PBPK) Model Parameter .....	84
5	3.5.4.1.4. Dose-Metric Predictions .....	85
6	3.5.5. Bayesian Estimation of Physiologically Based Pharmacokinetic (PBPK) Model Parameters, and Their Uncertainty and Variability .....	86
7	3.5.5.1.1. Updated Pharmacokinetic Database .....	86
8	3.5.5.1.2. Updated Hierarchical Population Statistical Model and Prior Distributions .....	94
9	3.5.5.1.3. Use of Interspecies Scaling to Update Prior Distributions in the Absence of Other Data .....	96
10	3.5.5.1.4. Implementation .....	100
11	3.5.6. Evaluation of Updated Physiologically Based Pharmacokinetic (PBPK) Model .....	100
12	3.5.6.1.1. Convergence .....	100
13	3.5.6.1.2. Evaluation of Posterior Parameter Distributions .....	103
14	3.5.6.1.3. Comparison of Model Predictions With Data .....	126
15	3.5.6.1.7. Sensitivity Analysis With Respect to Calibration Data .....	150
16	3.5.6.1.8. Summary Evaluation of Updated Physiologically Based Pharmacokinetic (PBPK) Model .....	157
17	3.5.7. Physiologically Based Pharmacokinetic (PBPK) Model Dose-Metric Predictions .....	157
18	3.5.7.1.1. Characterization of Uncertainty and Variability .....	157
19	3.5.7.1.2. Local Sensitivity Analysis With Respect to Dose-Metric Predictions .....	158
20	3.5.7.1.3. Implications for the Population Pharmacokinetics of Trichloroethylene (TCE) .....	174
21	3.5.7.1.6. Key Limitations and Potential Implications of Violating Key Assumptions .....	189
22	3.5.7.1.7. Overall Evaluation of Physiologically Based Pharmacokinetic (PBPK) Model-Based Internal Dose Predictions .....	190
23		
24		
25		
26		
27		
28		
	<b>4. HAZARD CHARACTERIZATION .....</b>	<b>1</b>
29	4.1. EPIDEMIOLOGIC STUDIES ON CANCER AND TRICHLOROETHYLENE (TCE)—METHODODOLOGICAL OVERVIEW .....	1
30		
31	4.2. GENETIC TOXICITY .....	33
	4.2.1. Trichloroethylene (TCE) .....	34

*This document is a draft for review purposes only and does not constitute Agency policy.*

1	4.2.1.1.1. DNA Binding Studies .....	34
2	4.2.1.1.2. Bacterial Systems—Gene Mutations .....	36
3	4.2.1.1.3. Fungal and Yeast Systems—Gene Mutations, Conversions	
4	and Recombination .....	40
5	4.2.1.1.4. Mammalian Systems Including Human Studies .....	44
6	4.2.1.1.14. Summary .....	57
	4.2.2. Trichloroacetic Acid (TCA).....	58
7	4.2.2.1.1. Bacterial Systems—Gene Mutations .....	59
8	4.2.2.1.2. Mammalian Systems .....	60
9	4.2.2.1.7. Summary .....	67
	4.2.3. Dichloroacetic Acid (DCA) .....	67
10	4.2.3.1.1. Bacterial and Fungal Systems—Gene Mutations .....	67
11	4.2.3.1.2. Mammalian Systems .....	72
12	4.2.3.1.6. Summary .....	74
	4.2.4. Chloral Hydrate.....	74
13	4.2.4.1.1. DNA Binding Studies .....	75
14	4.2.4.1.2. Bacterial and Fungal Systems—Gene Mutations .....	82
15	4.2.4.1.3. Mammalian Systems .....	83
16	4.2.4.1.9. Summary .....	86
	4.2.5. Dichlorovinyl Cysteine (DCVC) and S-Dichlorovinyl Glutathione	
	(DCVG).....	87
	4.2.6. Trichloroethanol (TCOH) .....	93
	4.2.7. Synthesis and Overall Summary .....	94
17	4.3. CENTRAL NERVOUS SYSTEM (CNS) TOXICITY .....	98
	4.3.1. Alterations in Nerve Conduction .....	99
18	4.3.1.1.1. Trigeminal Nerve Function: Human Studies .....	99
19	4.3.1.1.2. Nerve Conduction Velocity—Human Studies .....	106
20	4.3.1.1.3. Trigeminal Nerve Function: Laboratory Animal Studies .....	106
21	4.3.1.1.4. Discussion and Conclusions: Trichloroethylene (TCE)-	
22	Induced Trigeminal Nerve Impairment .....	109
	4.3.2. Auditory Effects.....	110
23	4.3.2.1.1. Auditory Function: Human Studies .....	110
24	4.3.2.1.2. Auditory Function: Laboratory Animal Studies .....	114
25	4.3.2.1.3. Summary and Conclusion of Auditory Effects .....	119
	4.3.3. Vestibular Function.....	121
26	4.3.3.1.1. Vestibular Function: Human Studies .....	121
27	4.3.3.1.2. Vestibular Function: Laboratory Animal Data .....	121
28	4.3.3.1.3. Summary and Conclusions for the Vestibular Function	
29	Studies.....	123
	4.3.4. Visual Effects.....	124
30	4.3.4.1.1. Visual Effects: Human Studies .....	124
31	4.3.4.1.2. Visual Effects: Laboratory Animal Data .....	127
32	4.3.4.1.3. Summary and Conclusion of Visual Effects .....	129
	4.3.5. Cognitive Function.....	131
33	4.3.5.1.1. Cognitive Effects: Human Studies.....	131
34	4.3.5.1.2. Cognitive Effects: Laboratory Animal Studies.....	132

1	4.3.5.1.3. Summary and Conclusions of Cognitive Function Studies .....	133
	4.3.6. Psychomotor Effects .....	138
2	4.3.6.1.1. Psychomotor Effects: Human Studies .....	139
3	4.3.6.1.4. Psychomotor Effects: Laboratory Animal Data.....	143
4	4.3.6.1.8. Summary and Conclusions for Psychomotor Effects .....	149
	4.3.7. Mood Effects and Sleep Disorders .....	150
5	4.3.7.1.1. Effects on Mood: Human Studies.....	150
6	4.3.7.1.2. Effects on Mood: Laboratory Animal Findings.....	150
7	4.3.7.1.3. Sleep Disturbances.....	151
	4.3.8. Developmental Neurotoxicity.....	152
8	4.3.8.1.1. Human Studies.....	152
9	4.3.8.1.2. Animal Studies.....	153
10	4.3.8.1.3. Summary and Conclusions for the Developmental	
11	Neurotoxicity Studies .....	158
	4.3.9. Mechanistic Studies of Trichloroethylene (TCE) Neurotoxicity.....	159
12	4.3.9.1.1. Dopamine Neuron Disruption.....	159
13	4.3.9.1.5. Neurochemical and Molecular Changes .....	161
	4.3.10. Potential Mechanisms for Trichloroethylene (TCE)-Mediated	
	Neurotoxicity .....	164
	4.3.11. Overall Summary and Conclusions—Weight of Evidence .....	167
14	4.4. KIDNEY TOXICITY AND CANCER .....	172
	4.4.1. Human Studies of Kidney.....	172
15	4.4.1.1.1. Nonspecific Markers of Nephrotoxicity .....	172
16	4.4.1.1.2. End-Stage Renal Disease .....	179
	4.4.2. Human Studies of Kidney Cancer.....	179
17	4.4.2.1.1. Studies of Job Titles and Occupations with Historical	
18	Trichloroethylene (TCE) Usage.....	188
19	4.4.2.1.2. Cohort and Case-Controls Studies of Trichloroethylene	
20	(TCE) Exposure .....	192
21	4.4.2.1.4. Examination of Possible Confounding Factors.....	197
22	4.4.2.1.5. Susceptible Populations—Kidney Cancer and	
23	Trichloroethylene (TCE) Exposure .....	202
24	4.4.2.1.6. Meta-Analysis for Kidney Cancer .....	205
	4.4.3. Human Studies of Somatic Mutation of von Hippel-Lindau (VHL)	
	Gene .....	211
	4.4.4. Kidney Noncancer Toxicity in Laboratory Animals .....	217
	4.4.5. Kidney Cancer in Laboratory Animals.....	228
25	4.4.5.1.1. Inhalation Studies of Trichloroethylene (TCE) .....	228
26	4.4.5.1.2. Gavage and Drinking Water Studies of Trichloroethylene	
27	(TCE) .....	231
28	4.4.5.1.3. Conclusions: Kidney Cancer in Laboratory Animals .....	233
	4.4.6. Role of Metabolism in Trichloroethylene (TCE) Kidney Toxicity .....	233
29	4.4.6.1.1. In Vivo Studies of the Kidney Toxicity of Trichloroethylene	
30	(TCE) Metabolites .....	234
31	4.4.6.1.4. In Vitro Studies of Kidney Toxicity of Trichloroethylene	
32	(TCE) and Metabolites.....	243



1	4.4.6.1.5. Conclusions as to the Active Agents of Trichloroethylene	
2	(TCE)-Induced Nephrotoxicity.....	245
3	4.4.7. Mode(s) of Action for Kidney Carcinogenicity.....	246
4	4.4.7.1.1. Hypothesized Mode of Action: Mutagenicity.....	246
5	4.4.7.1.3. Hypothesized Mode of Action: Cytotoxicity and	
6	Regenerative Proliferation.....	253
7	4.4.7.1.5. Additional Hypothesized Modes of Action with Limited	
8	Evidence or Inadequate Experimental Support.....	256
9	4.4.7.1.9. Conclusions About the Hypothesized Modes of Action.....	258
10	4.4.8. Summary: Trichloroethylene (TCE) Kidney Toxicity, Carcinogenicity,	
11	and Mode-of-Action.....	261
12	4.5. LIVER TOXICITY AND CANCER.....	264
13	4.5.1. Liver Noncancer Toxicity in Humans.....	264
14	4.5.2. Liver Cancer in Humans.....	269
15	4.5.3. Experimental Studies of Trichloroethylene (TCE) in Rodents—	
16	Introduction.....	287
17	4.5.4. Trichloroethylene (TCE)-Induced Liver Noncancer Effects.....	290
18	4.5.4.1.1. Liver Weight.....	290
19	4.5.4.1.2. Cytotoxicity and Histopathology.....	301
20	4.5.4.1.3. Measures of DNA Synthesis, Cellular Proliferation, and	
21	Apoptosis.....	308
22	4.5.4.1.4. Peroxisomal Proliferation and Related Effects.....	311
23	4.5.4.1.5. Oxidative Stress.....	313
24	4.5.4.1.6. Bile Production.....	315
25	4.5.4.1.7. Summary: Trichloroethylene (TCE)-Induced Noncancer	
26	Effects in Laboratory Animals.....	316
27	4.5.5. Trichloroethylene (TCE)-Induced Liver Cancer in Laboratory Animals.....	317
28	4.5.5.1.1. Negative or Inconclusive Studies of Mice and Rats.....	317
29	4.5.5.1.2. Positive Trichloroethylene (TCE) Studies of Mice.....	327
30	4.5.5.1.3. Summary: Trichloroethylene (TCE)-Induced Cancer in	
31	Laboratory Animals.....	329
32	4.5.6. Role of Metabolism in Liver Toxicity and Cancer.....	329
33	4.5.6.1.1. Pharmacokinetics of Chloral Hydrate (CH), Trichloroacetic	
34	Acid (TCA), and Dichloroacetic Acid (DCA) From	
35	Trichloroethylene (TCE) Exposure.....	330
36	4.5.6.1.2. Comparisons Between Trichloroethylene (TCE) and	
	Trichloroacetic Acid (TCA), Dichloroacetic Acid (DCA),	
	and Chloral Hydrate (CH) Noncancer Effects.....	331
	4.5.6.1.10. Comparisons of Trichloroethylene (TCE)-Induced	
	Carcinogenic Responses With Trichloroacetic Acid (TCA),	
	Dichloroacetic Acid (DCA), and Chloral Hydrate (CH)	
	Studies.....	352
	4.5.6.1.15. Conclusions Regarding the Role of Trichloroacetic Acid	
	(TCA), Dichloroacetic Acid (DCA), and Chloral Hydrate	
	(CH) in Trichloroethylene (TCE)-Induced Effects in the	
	Liver.....	380

	4.5.7. Mode of Action (MOA) for Trichloroethylene (TCE) Liver Carcinogenicity .....	380
1	4.5.7.1.1. Mutagenicity .....	381
2	4.5.7.1.3. Peroxisome Proliferator Activated Receptor Alpha (PPAR $\alpha$ )	
3	Receptor Activation .....	382
4	4.5.7.1.6. Additional Proposed Hypotheses and Key Events with	
5	Limited Evidence or Inadequate Experimental Support .....	392
6	4.5.7.1.15. Mode of Action (MOA) Conclusions .....	402
7	4.6. IMMUNOTOXICITY AND CANCERS OF THE IMMUNE SYSTEM .....	412
	4.6.1. Human Studies .....	399
8	4.6.1.1.1. Noncancer Immune-Related Effects .....	399
9	4.6.1.1.6. Cancers of the Immune System, Including Childhood	
10	Leukemia .....	413
	4.6.2. Animal Studies .....	453
11	4.6.2.1.1. Immunosuppression .....	453
12	4.6.2.1.5. Hypersensitivity .....	463
13	4.6.2.1.6. Autoimmunity .....	467
14	4.6.2.1.7. Cancers of the Immune System .....	481
	4.6.3. Summary .....	484
15	4.6.3.1.1. Noncancer Effects .....	484
16	4.6.3.1.2. Cancer .....	486
17	4.7. RESPIRATORY TRACT TOXICITY AND CANCER .....	488
	4.7.1. Epidemiologic Evidence .....	488
18	4.7.1.1.1. Chronic Effects: Inhalation .....	488
19	4.7.1.1.2. Cancer .....	489
	4.7.2. Laboratory Animal Studies .....	501
20	4.7.2.1.1. Respiratory Tract Animal Toxicity .....	501
21	4.7.2.1.3. Respiratory Tract Cancer .....	510
	4.7.3. Role of Metabolism in Pulmonary Toxicity .....	515
	4.7.4. Mode of Action for Pulmonary Carcinogenicity .....	520
22	4.7.4.1.1. Mutagenicity via Oxidative Metabolism .....	520
23	4.7.4.1.3. Cytotoxicity Leading to Increased Cell Proliferation .....	522
24	4.7.4.1.5. Additional Hypothesized Modes of Action with Limited	
25	Evidence or Inadequate Experimental Support .....	524
26	4.7.4.1.7. Conclusions About the Hypothesized Modes of Action .....	525
	4.7.5. Summary and Conclusions .....	527
27	4.8. REPRODUCTIVE AND DEVELOPMENTAL TOXICITY .....	529
	4.8.1. Reproductive Toxicity .....	529
28	4.8.1.1.1. Human Reproductive Outcome Data .....	530
29	4.8.1.1.6. Animal Reproductive Toxicity Studies .....	538
30	4.8.1.1.9. Discussion/Synthesis of Noncancer Reproductive Toxicity	
31	Findings .....	553
	4.8.2. Cancers of the Reproductive System .....	561
32	4.8.2.1.1. Human Data .....	561

1	4.8.2.1.5. Animal Studies.....	578
2	4.8.2.1.6. Mode of Action for Testicular Tumors.....	581
	4.8.3. Developmental Toxicity.....	582
3	4.8.3.1.1. Human Developmental Data.....	582
4	4.8.3.1.5. Animal Developmental Toxicology Studies.....	608
5	4.8.3.1.9. Discussion/Synthesis of Developmental Data.....	635
6	4.9. OTHER SITE-SPECIFIC CANCERS.....	653
	4.9.1. Esophageal Cancer.....	653
	4.9.2. Bladder Cancer.....	664
	4.9.3. Central Nervous System and Brain Cancers.....	671
7	4.10. SUSCEPTIBLE LIFESTAGES AND POPULATIONS.....	678
	4.10.1. Lifestages.....	678
8	4.10.1.1.1. Early Lifestages.....	679
9	4.10.1.1.5. Later Lifestages.....	690
	4.10.2. Other Susceptibility Factors.....	691
10	4.10.2.1.1. Gender.....	691
11	4.10.2.1.4. Genetic Variability.....	698
12	4.10.2.1.8. Race/Ethnicity.....	699
13	4.10.2.1.9. Preexisting Health Status.....	700
14	4.10.2.1.13. Lifestyle Factors and Nutrition Status.....	701
15	4.10.2.1.19. Mixtures.....	704
	4.10.3. Uncertainty of Database and Research Needs for Susceptible Populations.....	705
16	4.11. HAZARD CHARACTERIZATION.....	706
	4.11.1. Characterization of Noncancer Effects.....	706
17	4.11.1.1.1. Neurotoxicity.....	706
18	4.11.1.1.2. Kidney Toxicity.....	711
19	4.11.1.1.3. Liver Toxicity.....	712
20	4.11.1.1.4. Immunotoxicity.....	714
21	4.11.1.1.5. Respiratory Tract Toxicity.....	715
22	4.11.1.1.6. Reproductive Toxicity.....	716
23	4.11.1.1.7. Developmental Toxicity.....	717
	4.11.2. Characterization of Carcinogenicity.....	721
24	4.11.2.1.1. Summary Evaluation of Epidemiologic Evidence of Trichloroethylene (TCE) and Cancer.....	722
25		
26	4.11.2.1.12. Summary of Evidence for Trichloroethylene (TCE) Carcinogenicity in Rodents.....	731
27		
28	4.11.2.1.13. Summary of Additional Evidence on Biological Plausibility.....	734
29		
	4.11.3. Characterization of Factors Impacting Susceptibility.....	739
	<b>5. DOSE-RESPONSE ASSESSMENT.....</b>	<b>1</b>
30	5.1. DOSE-RESPONSE ANALYSES FOR NONCANCER ENDPOINTS.....	1

*This document is a draft for review purposes only and does not constitute Agency policy.*

	5.1.1. Modeling Approaches and Uncertainty Factors for Developing Candidate Reference Values Based on Applied Dose .....	3
	5.1.2. Candidate Critical Effects by Effect Domain .....	7
1	5.1.2.1.1. Candidate Critical Neurological Effects on the Basis of Applied Dose .....	8
2		
3	5.1.2.1.2. Candidate Critical Kidney Effects on the Basis of Applied Dose .....	16
4		
5	5.1.2.1.3. Candidate Critical Liver Effects on the Basis of Applied Dose .....	22
6		
7	5.1.2.1.4. Candidate Critical Body Weight Effects on the Basis of Applied Dose .....	23
8		
9	5.1.2.1.5. Candidate Critical Immunological Effects on the Basis of Applied Dose .....	24
10		
11	5.1.2.1.6. Candidate Critical Respiratory Tract Effects on the Basis of Applied Dose .....	29
12		
13	5.1.2.1.7. Candidate Critical Reproductive Effects on the Basis of Applied Dose .....	30
14		
15	5.1.2.1.10. Candidate Critical Developmental Effects on the Basis of Applied Dose .....	42
16		
17	5.1.2.1.11. Summary of cRfCs, cRfDs, and Candidate Critical Effects .....	54
	5.1.3. Application of Physiologically Based Pharmacokinetic (PBPK) Model to Inter- and Intraspecies Extrapolation for Candidate Critical Effects.....	54
18	5.1.3.1.1. Selection of Dose-metrics for Different Endpoints .....	58
19	5.1.3.1.7. Methods for Inter- and Intraspecies Extrapolation Using Internal Doses .....	66
20		
21	5.1.3.1.8. Results and Discussion of p-RfCs and p-RfDs for Candidate Critical Effects .....	85
22		
	5.1.4. Uncertainties in cRfCs and cRfDs .....	86
23	5.1.4.1.1. Qualitative Uncertainties .....	86
24	5.1.4.1.2. Quantitative Uncertainty Analysis of Physiologically Based Pharmacokinetic (PBPK) Model-Based Dose-metrics for Lowest-Observed-Adverse-Effect Level (LOAEL) or No-Observed-Adverse-Effect Level (NOAEL)-Based Points of Departure (PODs).....	89
25		
26		
27		
28		
	5.1.5. Summary of Noncancer Reference Values .....	101
29	5.1.5.1.1. Preferred Candidate Reference Values (cRfCs, cRfD, p-cRfCs and p-cRfDs) for Candidate Critical Effects .....	101
30		
31	5.1.5.1.2. Reference Concentration.....	108
32	5.1.5.1.3. Reference Dose .....	111
	5.2. DOSE-RESPONSE ANALYSIS FOR CANCER ENDPOINTS.....	115
33		
	5.2.1. Dose-Response Analyses: Rodent Bioassays .....	115
34	5.2.1.1.1. Rodent Dose-Response Analyses: Studies and Modeling Approaches .....	116
35		
36	5.2.1.1.2. Rodent Dose-Response Analyses: Dosimetry .....	125
37	5.2.1.1.5. Rodent Dose-Response Analyses: Results.....	134

1	5.2.1.1.6. Uncertainties in Dose-Response Analyses of Rodent	
2	Bioassays .....	146
	5.2.2. Dose-Response Analyses: Human Epidemiologic Data .....	154
3	5.2.2.1.1. Inhalation Unit Risk Estimate for Renal Cell Carcinoma	
4	Derived from Charbotel et al. (2006) Data .....	158
5	5.2.2.1.6. Adjustment of the Inhalation Unit Risk Estimate for Multiple	
6	Sites.....	167
7	5.2.2.1.7. Route-to-Route Extrapolation Using Physiologically Based	
8	Pharmacokinetic (PBPK) Model .....	171
	5.2.3. Summary of Unit Risk Estimates.....	177
9	5.2.3.1.1. Inhalation Unit Risk Estimate.....	177
10	5.2.3.1.2. Oral Slope Factor Estimate .....	178
11	5.2.3.1.3. Application of Age-Dependent Adjustment Factors.....	179
12	5.3. KEY RESEARCH NEEDS FOR TCE DOSE-RESPONSE ANALYSES .....	189
	<b>6. MAJOR CONCLUSIONS IN THE CHARACTERIZATION OF HAZARD AND</b>	
	<b>DOSE RESPONSE.....</b>	<b>1</b>
13	6.1. HUMAN HAZARD POTENTIAL.....	1
	6.1.1. Exposure (see Section 2).....	1
	6.1.2. Toxicokinetics and Physiologically Based Pharmacokinetic Modeling	
	(see Section 3 and Appendix A) .....	2
	6.1.3. Noncancer Toxicity.....	4
14	6.1.3.1.1. Neurological Effects (see Sections 4.3 and 4.11.1.1 and	
15	Appendix D) .....	4
16	6.1.3.1.2. Kidney Effects (see Sections 4.4.1, 4.4.4, 4.4.6, and	
17	4.11.1.2).....	5
18	6.1.3.1.3. Liver Effects (see Sections 4.5.1, 4.5.3, 4.5.4, 4.5.6, and	
19	4.11.1.3, and Appendix E).....	6
20	6.1.3.1.4. Immunological Effects (see Sections 4.6.1.1, 4.6.2, and	
21	4.11.1.4).....	7
22	6.1.3.1.5. Respiratory Tract Effects (see Sections 4.7.1.1, 4.7.2.1,	
23	4.7.3, and 4.11.1.5) .....	8
24	6.1.3.1.6. Reproductive Effects (see Sections 4.8.1 and 4.11.1.6).....	9
25	6.1.3.1.7. Developmental Effects (see Sections 4.8.3 and 4.11.1.7).....	10
	6.1.4. Carcinogenicity (see Sections 4.1, 4.2, 4.4.2, 4.4.5, 4.4.7, 4.5.2, 4.5.5,	
	4.5.6, 4.5.7, 4.6.1.2, 4.6.2.4, 4.7.1.2, 4.7.2.2, 4.7.4, 4.8.2, 4.9, and	
	4.11.2, and Appendices B and C) .....	12
	6.1.5. Susceptibility (see Sections 4.10 and 4.11.3) .....	17
26	6.2. DOSE-RESPONSE ASSESSMENT .....	19
	6.2.1. Noncancer Effects (see Section 5.1).....	19
27	6.2.1.1.1. Background and Methods .....	19
28	6.2.1.1.2. Uncertainties and Application of Uncertainty Factors (UFs)	
29	(see Section 5.1.1 and 5.1.4).....	20

1	6.2.1.1.3. Candidate Critical Effects and Reference Values (see	
2	Sections 5.1.2 and 5.1.3).....	24
3	6.2.1.1.11. Noncancer Reference Values (see Section 5.1.5).....	30
4	6.2.2. Cancer (see Section 5.2).....	33
5	6.2.2.1.1. Background and Methods (rodent: see Section 5.2.1.1;	
6	human: see Section 5.2.2.1).....	33
7	6.2.2.1.2. Inhalation Unit Risk Estimate (rodent: see Section 5.2.1.3;	
8	human: see Section 5.2.2.1 and 5.2.2.2).....	34
9	6.2.2.1.3. Oral Slope Factor Estimate (rodent: see Section 5.2.1.3;	
10	human: see Section 5.2.2.3).....	36
11	6.2.2.1.4. Uncertainties in Cancer Dose-Response Assessment.....	38
12	6.2.2.1.7. Application of Age-Dependent Adjustment Factors (see	
13	Section 5.2.3.3).....	43
14	6.3. OVERALL CHARACTERIZATION OF TCE HAZARD AND DOSE	
	RESPONSE.....	44
	7. References.....	1
	APPENDIX A: PBPK MODELING OF TCE AND METABOLITES–DETAILED	
	METHODS AND RESULTS.....	A-1
	APPENDIX B: SYSTEMATIC REVIEW OF EPIDEMIOLOGIC STUDIES ON	
	CANCER AND TRICHLOROETHYLENE (TCE) EXPOSURE.....	B-1
	APPENDIX C: META-ANALYSIS OF CANCER RESULTS FROM	
	EPIDEMIOLOGICAL STUDIES.....	C-1
	APPENDIX D: NEUROLOGICAL EFFECTS OF TRICHLOROETHYLENE.....	D-1
	APPENDIX E: ANALYSIS OF LIVER AND COEXPOSURE ISSUES FOR	
	THE TCE TOXICOLOGICAL REVIEW.....	E-1
	APPENDIX F: TCE NONCANCER DOSE-RESPONSE ANALYSES.....	F-1
	APPENDIX G: TCE CANCER DOSE-RESPONSE ANALYSES WITH RODENT	
	CANCER BIOASSAY DATA.....	G-1
	APPENDIX H: LIFETABLE ANALYSIS AND WEIGHTED LINEAR REGRESSION	
	BASED ON RESULTS FROM CHARBOTEL ET AL.....	H-1
	APPENDIX I: EPA RESPONSE TO MAJOR PEER REVIEW AND PUBLIC	
	COMMENTS.....	I-1

## LIST OF TABLES

1	Table 2-1. TCE metabolites and related parent compounds <sup>a</sup> .....	1
2	Table 2-2. Chemical properties of TCE.....	2
3	Table 2-3. Properties and uses of TCE related compounds.....	3
4	Table 2-4. TRI releases of TCE (pounds/year).....	5
5	Table 2-5. Concentrations of trichloroethylene in ambient air.....	10
6	Table 2-6. TCE ambient air monitoring data ( $\mu\text{g}/\text{m}^3$ ).....	11
7	Table 2-7. Mean TCE air levels across monitors by land setting and use (1985–1998).....	11
8	Table 2-8. Concentrations of trichloroethylene in water based on pre-1990 studies.....	15
9	Table 2-9. Levels in food.....	18
10	Table 2-10. TCE levels in whole blood by population percentile.....	19
11	Table 2-11. Modeled 1999 annual exposure concentrations ( $\mu\text{g}/\text{m}^3$ ) for trichloroethylene.....	20
12	Table 2-12. Preliminary estimates of TCE intake from food ingestion.....	22
13	Table 2-13. Preliminary intake estimates of TCE and TCE-related chemicals.....	23
14	Table 2-14. Years of solvent use in industrial degreasing and cleaning operations.....	25
15	Table 2-15. TCE standards.....	27
16	Table 3-1. Blood:air PC values for humans.....	6
17	Table 3-2. Blood:air PC values for rats and mice.....	7
18	Table 3-3. Air and blood concentrations during exposure to TCE in humans (Astrand and	
19	Ovrum, 1976).....	9
20	Table 3-4. Retention of inhaled TCE vapor in humans (Jakubowski and Wieczorek, 1988).....	10
21	Table 3-5. Uptake of TCE in human volunteers following 4 hour exposure to 70 ppm (Monster	
22	et al., 1979).....	10
23	Table 3-6. Concentrations of TCE in maternal and fetal blood at birth.....	15
24	Table 3-7. Distribution of TCE to rat tissues <sup>a</sup> following inhalation exposure (Savolainen et al.,	
25	1977).....	17
26	Table 3-8. Tissue:blood partition coefficient values for TCE.....	18
27	Table 3-9. Age-dependence of tissue:air partition coefficients in rats.....	21
28	Table 3-10. Predicted maximal concentrations of TCE in rat blood following a 6-hour inhalation	
29	exposure (Rodriguez et al., 2007).....	21
30	Table 3-11. Tissue distribution of TCE metabolites following inhalation exposure.....	22
31	Table 3-12. Binding of $^{14}\text{C}$ from [ $^{14}\text{C}$ ]TCE in rat liver and kidney at 72 hours after oral	
32	administration of 200 mg/kg [ $^{14}\text{C}$ ]TCE (Dekant et al., 1986b).....	24
33	Table 3-13. In vitro TCE oxidative metabolism in hepatocytes and microsomal fractions.....	30

1	Table 3-14. In vitro kinetics of trichloroethanol and trichloroacetic acid formation from chloral hydrate in rat, mouse, and human liver homogenates.....	36
2		
3	Table 3-15. In vitro kinetics of DCA metabolism in hepatic cytosol of mice, rats, and humans	38
4	Table 3-16. TCOH and TCA formed from CH in vitro in lysed whole blood of rats and mice or fractionated blood of humans (nmoles formed in 400 $\mu$ L samples over 30 minutes) .....	40
5		
6	Table 3-17. Reported TCA plasma binding parameters .....	41
7	Table 3-18. Partition coefficients for TCE oxidative metabolites .....	41
8	Table 3-19. Urinary excretion of trichloroacetic acid by various species exposed to trichloroethylene [based on data reviewed in (Fisher et al., 1991)].....	44
9		
10	Table 3-20. P450 isoform kinetics for metabolism of TCE to CH in human, rat, and mouse recombinant P450s.....	45
11		
12	Table 3-21. P450 isoform activities in human liver microsomes exhibiting different affinities for TCE.....	45
13		
14	Table 3-22. Comparison of peak blood concentrations in humans exposed to 100 ppm (537 $\text{mg}/\text{m}^3$ ) TCE for 4 hours (Fisher et al., 1998; Lash et al., 1999a).....	50
15		
16	Table 3-23. GSH conjugation of TCE (at 1–2 mM) in liver and kidney cellular fractions in humans, male F344 rats, and male B6C3F1 mice from Lash laboratory .....	51
17		
18	Table 3-24. Kinetics of TCE metabolism via GSH conjugation in male F344 rat kidney and human liver and kidney cellular and subcellular fractions from Lash laboratory .....	52
19		
20	Table 3-25. GSH conjugation of TCE (at 1.4–4 mM) in liver and kidney cellular fractions in humans, male F344 rats, and male B6C3F1 mice from Green and Dekant laboratories.....	53
21		
22	Table 3-26. GGT activity in liver and kidney subcellular fractions of mice, rats, and humans ..	61
23	Table 3-27. Multispecies comparison of whole-organ activity levels of GGT and dipeptidase..	62
24	Table 3-28. Comparison of hepatic in vitro oxidation and conjugation of TCE .....	67
25	Table 3-29. Estimates of DCVG in blood relative to inhaled TCE dose in humans exposed to 50 and 100 ppm (269 and 537 $\text{mg}/\text{m}^3$ ) (Fisher et al., 1998; Lash et al., 1999b) .....	68
26		
27	Table 3-30. Concentrations of TCE in expired breath from inhalation-exposed humans (Astrand and Ovrum, 1976).....	71
28		
29	Table 3-31. Conclusions from evaluation of Hack et al. (2006), and implications for PBPK model development.....	78
30		
31	Table 3-32. Discussion of changes to the Hack et al. (2006) PBPK model implemented for this assessment.....	83
32		
33	Table 3-33. PBPK model-based dose-metrics .....	87
34	Table 3-34. Rodent studies with pharmacokinetic data considered for analysis .....	88
35	Table 3-35. Human studies with pharmacokinetic data considered for analysis.....	92



1	Table 3-36. Parameters for which scaling from mouse to rat, or from mouse and rat to human,	
2	was used to update the prior distributions.....	99
3	Table 3-37. Prior and posterior uncertainty and variability in mouse PBPK model parameters	106
4	Table 3-38. Prior and posterior uncertainty and variability in rat PBPK model parameters .....	111
5	Table 3-39. Prior and posterior uncertainty and variability in human PBPK model parameters	
6	.....	116
7	Table 3-40. Confidence interval (CI) widths (ratio of 97.5% to 2.5% estimates) and fold-shift in	
8	median estimate for the PBPK model population median parameters, sorted in order of	
9	decreasing CI width. Shifts in the median estimate greater than threefold are in bold to denote	
10	larger shifts between the prior and posterior distributions.....	121
11	Table 3-41. Estimates of the residual-error.....	128
12	Table 3-42. Summary comparison of updated PBPK model predictions and in vivo data in mice	
13	.....	132
14	Table 3-43. Summary comparison of updated PBPK model predictions and in vivo data used for	
15	“calibration” in rats.....	138
16	Table 3-44. Summary comparison of updated PBPK model predictions and in vivo data used for	
17	“out-of-sample” evaluation in rats.....	141
18	Table 3-45. Summary comparison of updated PBPK model predictions and in vivo data used for	
19	“calibration” in humans .....	147
20	Table 3-46. Summary comparison of updated PBPK model predictions and in vivo data used for	
21	“out-of-sample” evaluation in humans .....	149
22	Table 3-47. Summary of scaling parameters ordered by fraction of calibration data of moderate	
23	or high sensitivity.....	155
24	Table 3-48. Posterior predictions for representative internal doses: mouse .....	169
25	Table 3-49. Posterior predictions for representative internal doses: rat .....	170
26	Table 3-50. Posterior predictions for representative internal doses: human.....	171
27	Table 3-51. Degree of variance in dose-metric predictions due to incomplete convergence	
28	(columns 2–4), combined uncertainty and population variability (columns 5–7), uncertainty in	
29	particular human population percentiles (columns 8–10), model fits to in vivo data (column 11).	
30	The GSD is the geometric standard deviation, which is a “fold-change” from the central	
31	tendency.....	192
32	Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies	
33	assessing cancer and TCE exposure .....	2
34	Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure.....	9
35	Table 4-3. Geographic-based studies assessing cancer and TCE exposure.....	22

1	Table 4-4. Standards of epidemiologic study design and analysis use for identifying cancer	
2	hazard and TCE exposure .....	25
3	Table 4-5. Summary of criteria for meta-analysis study selection .....	29
4	Table 4-6. TCE genotoxicity: bacterial assays .....	38
5	Table 4-7. TCE genotoxicity: fungal and yeast systems.....	42
6	Table 4-8. TCE genotoxicity: mammalian systems—gene mutations and chromosome	
7	aberrations.....	46
8	Table 4-9. TCE genotoxicity: mammalian systems—micronucleus, sister chromatic exchanges	
9	.....	50
10	Table 4-10. TCE genotoxicity: mammalian systems—unscheduled DNA synthesis, DNA strand	
11	breaks/protein crosslinks, cell transformation .....	55
12	Table 4-11. Genotoxicity of trichloroacetic acid—bacterial systems.....	59
13	Table 4-12. TCA Genotoxicity—mammalian systems (both in vitro and in vivo).....	62
14	Table 4-13. Genotoxicity of dichloroacetic acid (bacterial systems) .....	69
15	Table 4-14. Genotoxicity of dichloroacetic acid—mammalian systems.....	70
16	Table 4-15. Chloral hydrate genotoxicity: bacterial, yeast and fungal systems .....	76
17	Table 4-16.. Chloral hydrate genotoxicity: mammalian systems—all genetic endpoints, in vitro	
18	.....	78
19	Table 4-17.. Chloral hydrate genotoxicity: mammalian systems—all genetic damage, in vivo .	80
20	Table 4-18. TCE GSH conjugation metabolites genotoxicity .....	89
21	Table 4-19. Genotoxicity of trichloroethanol .....	93
22	Table 4-20. Summary of human trigeminal nerve and nerve conduction velocity studies.....	100
23	Table 4-21. Summary of animal trigeminal nerve studies.....	108
24	Table 4-22. Summary of human auditory function studies.....	112
25	Table 4-23. Summary of animal auditory function studies.....	117
26	Table 4-26. Summary of animal visual system studies.....	129
27	Table 4-27. Summary of human cognition effect studies .....	134
28	Table 4-28. Summary of animal cognition effect studies .....	137
29	Table 4-29. Summary of human choice reaction time studies.....	141
30	Table 4-30. Summary of animal psychomotor function and reaction time studies .....	145
31	Table 4-31. Summary of animal locomotor activity studies.....	147
32	Table 4-33. Summary of human developmental neurotoxicity associated with TCE exposures	
33	.....	152
34	Table 4-34. Summary of mammalian in vivo developmental neurotoxicity studies—oral	
35	exposures.....	154
36	Table 4-35. Summary of animal dopamine neuronal studies .....	160

1	Table 4-36. Summary of neurophysiological, neurochemical, and neuropathological effects with	
2	TCE exposure.....	161
3	Table 4-37. Summary of in vitro ion channel effects with TCE exposure .....	165
4	Table 4-38. Summary of human kidney toxicity studies .....	174
5	Table 4-39. Summary of human studies on TCE exposure and kidney cancer .....	181
6	Table 4-40. Summary of case-control studies on kidney cancer and occupation or job title ....	189
7	Table 4-41. Summary of lung and kidney cancer risks in active smokers (from IARC, 2004b)	
8	.....	200
9	Table 4-42. Summary of human studies on somatic mutations of the VHL gene <sup>a</sup> .....	214
10	Table 4-43. Inhalation studies of kidney noncancer toxicity in laboratory animals .....	219
11	Table 4-44. Oral and i.p. studies of kidney noncancer toxicity in laboratory animals .....	220
12	Table 4-46. Summary of renal toxicity and tumor findings in gavage studies of trichloroethylene	
13	by NCI (1976).....	224
14	Table 4-47. Summary of renal toxicity findings in gavage studies of trichloroethylene by	
15	Maltoni et al. (1988) .....	225
16	Table 4-48. Summary of renal toxicity and tumor incidence in gavage studies of	
17	trichloroethylene by NTP (1988).....	226
18	Table 4-49. Summary of renal toxicity and tumor findings in inhalation studies of	
19	trichloroethylene by Maltoni et al. (1988) <sup>a</sup> .....	227
20	Table 4-50. Summary of renal tumor findings in inhalation studies of trichloroethylene by	
21	Henschler et al. (1980) <sup>a</sup> and Fukuda et al. (1983) <sup>b</sup> .....	230
22	Table 4-51. Summary of renal tumor findings in gavage studies of trichloroethylene by	
23	Henschler et al. (1984) <sup>a</sup> and Van Duuren et al. (1979) <sup>b</sup> .....	232
24	Table 4-52. Laboratory animal studies of kidney noncancer toxicity of TCE metabolites .....	236
25	Table 4-53. Summary of histological changes in renal proximal tubular cells induced by chronic	
26	exposure to TCE, DCVC, and TCOH.....	238
27	Table 4-54. Summary of major mode of action conclusions for TCE kidney carcinogenesis ..	248
28	Table 4-55. Summary of human liver toxicity studies.....	265
29	Table 4-56. Selected results from epidemiologic studies of TCE exposure and cirrhosis .....	271
30	Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer ...	274
31	Table 4-58. Oral studies of TCE-induced liver effects in mice and rats.....	292
32	Table 4-59. Inhalation and i.p. studies of TCE-induced liver effects in mice and rats.....	295
33	Table 4-60. Summary of liver tumor findings in gavage studies of trichloroethylene by NTP	
34	(1990) <sup>a</sup> .....	319
35	Table 4-61. Summary of liver tumor findings in gavage studies of trichloroethylene by NCI	
36	(1976).....	320

1	Table 4-62. Summary of liver tumor incidence in gavage studies of trichloroethylene by NTP	
2	(1988).....	321
3	Table 4-63. Summary of liver tumor findings in inhalation studies of trichloroethylene by	
4	Maltoni et al. (1988) <sup>a</sup> .....	323
5	Table 4-64. Summary of liver tumor findings in inhalation studies of trichloroethylene by	
6	Henschler et al. (1980) <sup>a</sup> and Fukuda et al. (1983) .....	324
7	Table 4-65. Summary of liver tumor findings in gavage studies of trichloroethylene by	
8	Henschler et al. (1984) <sup>a</sup> .....	325
9	Table 4-66. Potency indicators for mouse hepatocarcinogenicity and in vitro transactivation of	
10	mouse PPAR $\alpha$ for four PPAR $\alpha$ agonists .....	390
11	Table 4-67. Potency indicators for rat hepatocarcinogenicity and common short-term markers of	
12	PPAR $\alpha$ activation for four PPAR $\alpha$ agonists .....	391
13	Table 4-68. Summary of mode of action conclusions for TCE-induced liver carcinogenesis ..	403
14	Table 4-69. Studies of immune parameters (IgE antibodies and cytokines) and trichloroethylene	
15	in humans .....	401
16	Table 4-70. Case-control studies of autoimmune diseases with measures of trichloroethylene	
17	exposure .....	411
18	Table 4-71. Incidence cohort studies of TCE exposure and lymphopoietic and hematopoietic	
19	cancer risk .....	418
20	Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and	
21	hematopoietic cancer risk .....	422
22	Table 4-73. Case-control studies of TCE exposure and lymphopoietic cancer, leukemia or	
23	multiple myeloma .....	433
24	Table 4-74. Geographic-based studies of TCE and non-Hodgkin lymphoma or leukemia in	
25	adults .....	438
26	Table 4-75. Selected results from epidemiologic studies of TCE exposure and childhood	
27	leukemia .....	443
28	Table 4-76. Summary of TCE immunosuppression studies .....	455
29	Table 4-77. Summary of TCE hypersensitivity studies .....	465
30	Table 4-78. Summary of autoimmune-related studies of TCE and metabolites in mice and rats	
31	(by sex, strain, and route of exposure) <sup>a</sup> .....	469
32	Table 4-79. Malignant lymphomas incidence in mice exposed to TCE in gavage and inhalation	
33	exposure studies .....	482
34	Table 4-80. Leukemia incidence in rats exposed to TCE in gavage and inhalation exposure	
35	studies .....	483
36	Table 4-81. Selected results from epidemiologic studies of TCE exposure and lung cancer ....	491

1	Table 4-82. Selected results from epidemiologic studies of TCE exposure and laryngeal cancer	
2	.....	498
3	Table 4-83. Animal toxicity studies of trichloroethylene .....	503
4	Table 4-84. Animal carcinogenicity studies of trichloroethylene.....	512
5	Table 4-85. Human reproductive effects .....	531
6	Table 4-86. Summary of mammalian in vivo reproductive toxicity studies—inhaleation exposures	
7	.....	539
8	Table 4-87. Summary of mammalian in vivo reproductive toxicity studies—oral exposures ..	541
9	Table 4-88. Summary of adverse female reproductive outcomes associated with TCE exposures	
10	.....	553
11	Table 4-89. Summary of adverse male reproductive outcomes associated with TCE exposures	
12	.....	555
13	Table 4-90. Summary of human studies on TCE exposure and prostate cancer .....	562
14	Table 4-91. Summary of human studies on TCE exposure and breast cancer .....	567
15	Table 4-92. Summary of human studies on TCE exposure and cervical cancer .....	571
16	Table 4-93. Histopathology findings in reproductive organs .....	580
17	Table 4-94. Testicular tumors in male rats exposed to TCE, adjusted for reduced survival <sup>a</sup> ....	581
18	Table 4-95. Developmental studies in humans.....	584
19	Table 4-96. Summary of mammalian in vivo developmental toxicity studies—inhaleation	
20	exposures.....	608
21	Table 4-97. Ocular defects observed (Narotsky et al., 1995) .....	610
22	Table 4-98. Summary of mammalian in vivo developmental toxicity studies—oral exposures	612
23	Table 4-99. Types of congenital cardiac defects observed in TCE-exposed fetuses (Dawson et	
24	al., 1993, Table 3).....	621
25	Table 4-100. Types of heart malformations per 100 fetuses (Johnson et al., 2003, Table 2, p.	
26	290).....	622
27	Table 4-101. Congenital cardiac malformations (Johnson et al., 1998a, Table 2, p. 997) .....	625
28	Table 4-102. Summary of adverse fetal and early neonatal outcomes associated with TCE	
29	exposures.....	636
30	Table 4-103. Summary of studies that identified cardiac malformations associated with TCE	
31	exposures.....	637
32	Table 4-104. Events in cardiac valve formation in mammals and birds <sup>a</sup> .....	641
33	Table 4-105. Summary of other structural developmental outcomes associated with TCE	
34	exposures.....	645
35	Table 4-106. Summary of developmental neurotoxicity associated with TCE exposures .....	647
36	Table 4-107. Summary of developmental immunotoxicity associated with TCE exposures ....	650

1	Table 4-108. Summary of childhood cancers associated with TCE exposures .....	652
2	Table 4-109. Selected observations from case-control studies of TCE exposure and esophageal	
3	cancer .....	655
4	Table 4-110. Summary of human studies on TCE exposure and esophageal cancer .....	658
5	Table 4-111. Summary of human studies on TCE exposure and bladder cancer .....	666
6	Table 4-112. Summary of human studies on TCE exposure and brain cancer .....	673
7	Table 4-113. Estimated lifestage-specific daily doses for TCE in water <sup>a</sup> .....	682
8	Table. 5-1 Summary of studies of neurological effects suitable for dose-response assessment....	9
9	Table 5-2. Neurological effects in studies suitable for dose-response assessment, and	
10	corresponding cRfCs and cRfDs.....	11
11	Table 5-3. Summary of studies of kidney, liver, and body weight effects suitable for	
12	dose-response assessment .....	17
13	Table 5-4. Kidney, liver, and body weight effects in studies suitable for dose-response	
14	assessment, and corresponding cRfCs and cRfDs .....	19
15	Table 5-5. Summary of studies of immunological effects suitable for dose-response assessment	
16	.....	26
17	Table 5-6. Immunological effects in studies suitable for dose-response assessment, and	
18	corresponding cRfCs and cRfDs.....	28
19	Table 5-7. Summary of studies of reproductive effects suitable for dose-response assessment .	32
20	Table 5-8. Reproductive effects in studies suitable for dose-response assessment, and	
21	corresponding cRfCs and cRfDs.....	37
22	Table 5-9. Summary of studies of developmental effects suitable for dose-response assessment	
23	.....	45
24	Table 5-10. Developmental effects in studies suitable for dose-response assessment, and	
25	corresponding cRfCs and cRfDs.....	49
26	Table 5-11. Ranges of cRfCs based on applied dose for various noncancer effects associated	
27	with inhalation TCE exposure .....	56
28	Table 5-12. Ranges of cRfDs based on applied dose for various noncancer effects associated	
29	with oral TCE exposure .....	57
30	Table 5-13. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on	
31	PBPK modeled internal dose-metrics) for candidate critical neurological effects .....	71
32	Table 5-14. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on	
33	PBPK modeled internal dose-metrics) for candidate critical kidney effects .....	73
34	Table 5-15. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on	
35	PBPK modeled internal dose-metrics) for candidate critical liver effects.....	76

1	Table 5-16. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on	
2	PBPK modeled internal dose-metrics) for candidate critical immunological effects.....	77
3	Table 5-17. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on	
4	PBPK modeled internal dose-metrics) for candidate critical reproductive effects.....	79
5	Table 5-18. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on	
6	PBPK modeled internal dose-metrics) for candidate critical developmental effects .....	82
7	Table 5-19. Comparison of “sensitive individual” HECs or HEDs for neurological effects based	
8	on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the	
9	NOAEL or LOAEL.....	93
10	Table 5-20. Comparison of “sensitive individual” HECs or HEDs for kidney and liver effects	
11	based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at	
12	the NOAEL or LOAEL.....	94
13	Table 5-21. Comparison of “sensitive individual” HECs or HEDs for immunological effects	
14	based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at	
15	the NOAEL or LOAEL.....	96
16	Table 5-22. Comparison of “sensitive individual” HECs or HEDs for reproductive effects based	
17	on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the	
18	NOAEL or LOAEL.....	97
19	Table 5-23. Comparison of “sensitive individual” HECs or HEDs for developmental effects	
20	based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at	
21	the NOAEL or LOAEL.....	99
22	Table 5-24. Lowest p-cRfCs or cRfCs for different effect domains.....	103
23	Table 5-25. Lowest p-cRfDs or cRfDs for different effect domains .....	105
24	Table 5-26. Lowest p-cRfCs for candidate critical effects for different types of effect based on	
25	primary dose-metric .....	107
26	Table 5-27. Lowest p-cRfDs for candidate critical effects for different types of effect based on	
27	primary dose-metric .....	107
28	Table 5-28. Summary of critical studies, effects, PODs, and UFs used to derive the RfC .....	109
29	Table 5-29. Summary of supporting studies, effects, PODs, and UFs for the RfC .....	109
30	Table 5-30. Summary of critical studies, effects, PODs, and UFs used to derive the RfD .....	112
31	Table 5-31. Summary of supporting studies, effects, PODs, and UFs for the RfD.....	114
32	Table 5-32. Inhalation bioassays.....	116
33	Table 5-33. Oral bioassays.....	118
34	Table 5-34. Specific dose-response analyses performed and dose-metrics used.....	122
35	Table 5-35. Mean PBPK model predictions for weekly internal dose in humans exposed	
36	continuously to low levels of TCE via inhalation (ppm) or orally (mg/kg/day) .....	133

1	Table 5-36. Summary of PODs and unit risk estimates for each sex/species/bioassay/tumor type	
2	(inhalation).....	135
3	Table 5-37. Summary of PODs and slope factor estimates for each sex/species/bioassay/tumor	
4	type (oral).....	137
5	Table 5-38. Comparison of survival-adjusted results for 3 oral male rat data sets <sup>a</sup> .....	141
6	Table 5-39. Inhalation: most sensitive bioassay for each sex/species combination <sup>a</sup> .....	145
7	Table 5-40. Oral: most sensitive bioassay for each sex/species combination <sup>a</sup> .....	145
8	Table 5-41. Summary of PBPK model-based uncertainty analysis of unit risk estimates for each	
9	sex/species/bioassay/tumor type (inhalation) .....	155
10	Table 5-42. Summary of PBPK model-based uncertainty analysis of slope factor estimates for	
11	each sex/species/bioassay/tumor type (oral).....	156
12	Table 5-43. Results from Charbotel et al. (2006) on relationship between TCE exposure and	
13	RCC.....	160
14	Table 5-44. Extra risk estimates for RCC incidence from various levels of lifetime exposure to	
15	TCE, using linear cumulative exposure model.....	162
16	Table 5-45. EC <sub>01</sub> , LEC <sub>01</sub> , and unit risk estimates for RCC incidence, using linear cumulative	
17	exposure model.....	163
18	Table 5-46. Relative contributions to extra risk for cancer incidence from TCE exposure for	
19	multiple tumor types .....	171
20	Table 5-47. Route-to-route extrapolation of site-specific inhalation unit risks to oral slope	
21	factors.....	176
22	Table 5-48. Sample calculation for total lifetime cancer risk based on the kidney unit risk	
23	estimate, potential risk for NHL and liver cancer, and potential increased early-life susceptibility,	
24	assuming a constant lifetime exposure to 1 µg/m <sup>3</sup> of TCE in air .....	183
25	Table 5-49. Sample calculation for total lifetime cancer risk based on the kidney cancer slope	
26	factor estimate, potential risk for NHL and liver cancer, and potential increased early-life	
27	susceptibility, assuming a constant lifetime exposure to 1 µg/L of TCE in drinking water .....	187

## LIST OF FIGURES

28	Figure 2-1. Molecular structure of TCE. ....	2
29	Figure 2-2. Source contribution to TCE emissions.....	6
30	Figure 2-3. Annual emissions of TCE. ....	6
31	Figure 2-4. Modeled ambient air concentrations of TCE. ....	13
32	Figure 3-1. Gas uptake data from closed-chamber exposure of rats to TCE. Symbols represent	
33	measured chamber concentrations. Source: Simmons et al. (2002). ....	12



1 Figure 3-2. Disposition of [<sup>14</sup>C]TCE administered by oral gavage in mice (Dekant et al., 1984;  
2 Dekant et al., 1986b; Green and Prout, 1985; Prout et al., 1985). ..... 26  
3 Figure 3-3. Disposition of [<sup>14</sup>C]TCE administered by oral gavage in rats (Dekant et al., 1984;  
4 Dekant et al., 1986b; Green and Prout, 1985; Prout et al., 1985). ..... 27  
5 Figure 3-4. Scheme for the oxidative metabolism of TCE. .... 29  
6 Figure 3-5. Scheme for GSH-dependent metabolism of TCE. .... 48  
7 Figure 3-6. Interorgan TCE transport and metabolism via the GSH pathway. See Figure 3-5 for  
8 enzymes involved in metabolic steps. Source: Lash et al.(2000a; 2000b); NRC (2006). ..... 58  
9 Figure 3-7. Overall structure of PBPK model for TCE and metabolites used in this assessment.  
10 Boxes with underlined labels are additions or modifications of the Hack et al. (2006) model,  
11 which are discussed in Table 3-32. .... 81  
12 Figure 3-8. Schematic of how posterior predictions were generated for comparison with  
13 experimental data. Two sets of posterior predictions were generated: population predictions  
14 (diagonal hashing) and subject-specific predictions (vertical hashing). (Same as Figure A-2 in  
15 Appendix A)..... 127  
16 Figure 3-9. Comparison of mouse data and PBPK model predictions from a random posterior  
17 sample. Each panel shows results for a different measurement. The solid line represents  
18 prediction = data, and the grey dotted lines show prediction = data × GSD<sub>err</sub> and data ÷ GSD<sub>err</sub>,  
19 where GSD<sub>err</sub> is the median estimate of the residual-error GSD shown in Table 3-41. .... 130  
20 Figure 3-10. Comparison of rat data and PBPK model predictions from a random posterior  
21 sample. Each panel shows results for a different measurement. The solid line represents  
22 prediction = data, and the grey lines show prediction = data × GSD<sub>err</sub> and data ÷ GSD<sub>err</sub>, where  
23 GSD<sub>err</sub> is the lowest (dotted) and highest (dashed) median estimate of the residual-error GSD  
24 shown in Table 3-41..... 134  
25 Figure 3-11. Comparison of urinary excretion data for NAcDCVC and predictions from the  
26 Hack et al. (2006) and the updated PBPK models. Data are from Bernauer et al. (1996) for (A  
27 and B) rats or (C and D) humans exposed for 6 hour to 40 (○), 80 (Δ), or 160 (+) ppm in air  
28 (thick horizontal line denotes the exposure period). Predictions from Hack et al. (2006) and the  
29 corresponding data (A and C) are only for the 1,2 isomer, whereas those from the updated model  
30 (B and D) are for both isomers combined. Parameter values used for each prediction are a  
31 random sample from the subject-specific parameters from the rat and human MCMC chains (the  
32 last iteration of the first chain was used in each case). Note that in the Hack et al. (2006) model,  
33 each dose group had different model parameters, whereas in the updated model, all dose groups  
34 are required to have the same model parameters. See files linked to Appendix A for comparisons  
35 with the full distribution of predictions. .... 143

1 Figure 3-12. Comparison of human data and PBPK model predictions from a random posterior  
2 sample. Each panel shows results for a different measurement. The solid line represents  
3 prediction = data, and the grey lines show prediction = data  $\times$  GSD<sub>err</sub> and data  $\div$  GSD<sub>err</sub>, where  
4 GSD<sub>err</sub> is the lowest (dotted) and highest (dashed) median estimate of the residual-error GSD  
5 shown in Table 3-41..... 145

6 Figure 3-13. Comparison of DCVG concentrations in human blood and predictions from the  
7 updated model. Data are mean concentrations for males ( $\Delta$ ) and females ( $\circ$ ) reported in Lash  
8 et al. (1999a) for humans exposed for 4 hours to 100 ppm TCE in air (thick horizontal line  
9 denotes the exposure period). Data for oxidative metabolites from the same individuals were  
10 reported in Fisher et al. (1998) but could not be matched with the individual DCVG data (Lash  
11 2007, personal communication). The vertical error bars are standard errors of the mean as  
12 reported in Lash et al. (1999a) ( $n = 8$ , so standard deviation is 80.5-fold larger). Lines are PBPK  
13 model predictions for individual male (solid) and female (dashed) subjects. Parameter values  
14 used for each prediction are a random sample from the individual-specific parameters from the  
15 human MCMC chains (the last iteration of the 1<sup>st</sup> chain was used). See files linked to Appendix  
16 A for comparisons with the full distribution of predictions..... 151

17 Figure 3-14. Sensitivity analysis results: Number of mouse calibration data points with SC in  
18 various categories for each scaling parameter. .... 152

19 Figure 3-15. Sensitivity analysis results: Number of rat calibration data points with SC in  
20 various categories for each scaling parameter. .... 153

21 Figure 3-16. Sensitivity analysis results: Number of human calibration data points with SC in  
22 various categories for each scaling parameter. .... 154

23 Figure 3-17. PBPK model predictions for the fraction of intake that is metabolized under  
24 continuous inhalation (A) and oral (B) exposure conditions in mice (white), rats (diagonal  
25 hashing), and humans (horizontal hashing). Bars and thin error bars represent the median  
26 estimate and 95% CI for a random subject, and reflect combined uncertainty and variability.  
27 Circles and thick error bars represent the median estimate and 95% CI for the population mean,  
28 and reflect uncertainty only. .... 160

29 Figure 3-18. PBPK model predictions for the fraction of intake that is metabolized by oxidation  
30 (in the liver and lung) under continuous inhalation (A) and oral (B) exposure conditions in mice  
31 (white), rats (diagonal hashing), and humans (horizontal hashing). Bars and thin error bars  
32 represent the median estimate and 95% CI for a random subject, and reflect combined  
33 uncertainty and variability. Circles and thick error bars represent the median estimate and 95%  
34 CI for the population mean, and reflect uncertainty only. .... 161

35 Figure 3-19. PBPK model predictions for the fraction of intake that is metabolized by GSH  
36 conjugation (in the liver and kidney) under continuous inhalation (A) and oral (B) exposure

1 conditions in mice (dotted line), rats (dashed line), and humans (solid line). X-values are slightly  
2 offset for clarity. Open circles (connected by lines) and thin error bars represent the median  
3 estimate and 95% CI for a random subject, and reflect combined uncertainty and variability.  
4 Filled circles and thick error bars represent the median estimate and 95% CI for the population  
5 mean, and reflect uncertainty only..... 162

6 Figure 3-20. PBPK model predictions for the fraction of intake that is bioactivated DCVC in the  
7 kidney under continuous inhalation (A) and oral (B) exposure conditions in rats (dashed line) and  
8 humans (solid line). X-values are slightly offset for clarity. Open circles (connected by lines)  
9 and thin error bars represent the median estimate and 95% CI for a random subject, and reflect  
10 combined uncertainty and variability. Filled circles and thick error bars represent the median  
11 estimate and 95% CI for the population mean, and reflect uncertainty only..... 163

12 Figure 3-21. PBPK model predictions for fraction of intake that is oxidized in the respiratory  
13 tract under continuous inhalation (A) and oral (B) exposure conditions in mice (dotted line), rats  
14 (dashed line), and humans (solid line). X-values are slightly offset for clarity. Open circles  
15 (connected by lines) and thin error bars represent the median estimate and 95% CI for a random  
16 subject, and reflect combined uncertainty and variability. Filled circles and thick error bars  
17 represent the median estimate and 95% CI for the population mean, and reflect uncertainty only.  
18 ..... 164

19 Figure 3-22. PBPK model predictions for the fraction of intake that is “untracked” oxidation of  
20 TCE in the liver under continuous inhalation (A) and oral (B) exposure conditions in mice  
21 (dotted line), rats (dashed line), and humans (solid line) X-values are slightly offset for clarity.  
22 Open circles (connected by lines) and thin error bars represent the median estimate and 95% CI  
23 for a random subject, and reflect combined uncertainty and variability. Filled circles and thick  
24 error bars represent the median estimate and 95% CI for the population mean, and reflect  
25 uncertainty only. .... 165

26 Figure 3-23. PBPK model predictions for the weekly AUC of TCE in venous blood  
27 (mg-hour/L-week) per unit exposure (ppm or mg/kg-day) under continuous inhalation (A) and  
28 oral (B) exposure conditions in mice (dotted line), rats (dashed line), and humans (solid line).  
29 X-values are slightly offset for clarity. Open circles (connected by lines) and thin error bars  
30 represent the median estimate and 95% CI for a random subject, and reflect combined  
31 uncertainty and variability. Filled circles and thick error bars represent the median estimate and  
32 95% CI for the population mean, and reflect uncertainty only..... 166

33 Figure 3-24. PBPK model predictions for the weekly AUC of TCOH in blood  
34 (mg-hour/L-week) per unit exposure (ppm or mg/kg-day) under continuous inhalation (A) and  
35 oral (B) exposure conditions in mice (dotted line), rats (dashed line), and humans (solid line).  
36 X-values are slightly offset for clarity. Open circles (connected by lines) and thin error bars

1	represent the median estimate and 95% CI for a random subject, and reflect combined	
2	uncertainty and variability. Filled circles and thick error bars represent the median estimate and	
3	95% CI for the population mean, and reflect uncertainty only.....	167
4	Figure 3-25. PBPK model predictions for the weekly AUC of TCA in the liver	
5	(mg-hour/L-week) per unit exposure (ppm or mg/kg-day) under continuous inhalation (A) and	
6	oral (B) exposure conditions in mice (dotted line), rats (dashed line), and humans (solid line).	
7	X-values are slightly offset for clarity. Open circles (connected by lines) and thin error bars	
8	represent the median estimate and 95% CI for a random subject, and reflect combined	
9	uncertainty and variability. Filled circles and thick error bars represent the median estimate and	
10	95% CI for the population mean, and reflect uncertainty only.....	168
11	Figure 3-26. Sensitivity analysis results: SC for mouse scaling parameters with respect to	
12	dose-metrics following 100 ppm (light bars) and 600 ppm (dark bars), 7 h/day, 5 day/wk	
13	inhalation exposures.....	176
14	Figure 3-27. Sensitivity analysis results: SC for mouse scaling parameters with respect to	
15	dose-metrics following 300 mg/kg-day (light bars) and 1,000 mg/kg-day (dark bars), 5 day/wk	
16	oral gavage exposures.....	177
17	Figure 3-28. Sensitivity analysis results: SC for rat scaling parameters with respect to	
18	dose-metrics following 100 ppm (light bars) and 600 ppm (dark bars), 7 h/day, 5 day/wk	
19	inhalation exposures.....	178
20	Figure 3-29. Sensitivity analysis results: SC for rat scaling parameters with respect to	
21	dose-metrics following 300 mg/kg-day (light bars) and 1,000 mg/kg-day (dark bars), 5 day/wk	
22	oral gavage exposures.....	179
23	Figure 3-30. Sensitivity analysis results: SC for female (light bars) and male (dark bars) human	
24	scaling parameters with respect to dose-metrics following 0.001 ppm continuous inhalation	
25	exposures.....	180
26	Figure 3-31. Sensitivity analysis results: SC for female (light bars) and male (dark bars) human	
27	scaling parameters with respect to dose-metrics following 0.001 mg/kg-day continuous oral	
28	exposures.....	181
29	Figure 4-1. Meta-analysis of kidney cancer and overall TCE exposure (the summary estimate is	
30	in the bottom row, represented by the diamond). Random effects model; fixed effect model	
31	same. Symbol sizes reflect relative weights of the studies.....	207
32	Figure 4-2. Meta-analysis of kidney cancer and TCE exposure—highest exposure groups. With	
33	assumed null RR estimates for Antilla, Axelson, and Hansen (see Appendix C text). Random	
34	effects model; fixed effect model same. The summary estimate is in the bottom row, represented	
35	by the diamond. Symbol sizes reflect relative weights of the studies. ....	209

1 Figure 4-3. Relative risk estimates of liver and biliary tract cancer and overall TCE exposure.  
2 Random effects model; fixed effect model same. The summary estimate is in the bottom row,  
3 represented by the diamond. Symbol sizes reflect relative weights of the studies. .... 284  
4 Figure 4-4. Meta-analysis of liver cancer and TCE exposure—highest exposure groups, with  
5 assumed null RR estimates for Hansen and Zhao (see Appendix C text). Random effects model;  
6 fixed effect model same. The summary estimate is in the bottom row, represented by the  
7 diamond. Symbol sizes reflect relative weights of the studies. .... 285  
8 Figure 4-5. Comparison of average fold-changes in relative liver weight to control and exposure  
9 concentrations of 2 g/L or less in drinking water for TCA and DCA in male B6C3F1 mice for  
10 14–30 days (Carter et al., 1995; DeAngelo et al., 1989; 2008; Kato-Weinstein et al., 2001;  
11 Parrish et al., 1996; Sanchez and Bull, 1990). .... 334  
12 Figure 4-6. Comparisons of fold-changes in average relative liver weight and gavage dose of  
13 (top panel) male B6C3F1 mice for 10–28 days of exposure (Dees and Travis, 1993; Elcombe et  
14 al., 1985; Goldsworthy and Popp, 1987; Merrick et al., 1989) and (bottom panel) in male  
15 B6C3F1 and Swiss mice. .... 336  
16 Figure 4-7. Comparison of fold-changes in relative liver weight for data sets in male B6C3F1,  
17 Swiss, and NRMI mice between TCE studies Kjellstrand et al., 1983a (Buben and O'Flaherty,  
18 1985; Goel et al., 1992; Merrick et al., 1989) [duration 28–42 days]) and studies of direct oral  
19 TCA administration to B6C3F1 mice (DeAngelo et al., 1989; DeAngelo et al., 2008; Kato-  
20 Weinstein et al., 2001; Parrish et al., 1996)[duration 14–28 days]). Abscissa for TCE studies  
21 consists of the median estimates of the internal dose of TCA predicted from metabolism of TCE  
22 using the PBPK model described in Section 3.5 of the TCE risk assessment. Lines show linear  
23 regression with intercept fixed at unity. All data were reported fold-change in mean liver  
24 weight/body weight ratios, except for Kjellstrand et al. (1983b), with were the fold-change in the  
25 ratio of mean liver weight to mean body weight. In addition, in Kjellstrand et al. (1983b), some  
26 systemic toxicity as evidence by decreased total body weight was reported in the highest-dose  
27 group. .... 338  
28 Figure 4-8. Comparison of hepatomegaly as a function of AUC of TCA in liver, using values for  
29 the TCA drinking water fractional absorption (Fabs). Fold-changes in relative liver weight for  
30 data sets in male B6C3F1, Swiss, and NRMI mice between TCE studies Kjellstrand et al., 1983a  
31 (Buben and O'Flaherty, 1985; Goel et al., 1992; Merrick et al., 1989) [duration 28–42 days] and  
32 studies of direct oral TCA administration to B6C3F1 mice (DeAngelo et al., 1989; DeAngelo et  
33 al., 2008; Kato-Weinstein et al., 2001; Parrish et al., 1996) Green, 2003b [duration 14–28 days].  
34 Linear regressions were compared using ANOVA to assess whether the TCE studies were  
35 consistent with the TCA studies, using TCA as the dose-metric. For each analysis of drinking  
36 water fraction absorption, ANOVA  $p$ -values were  $<10^{-4}$  when comparing the assumption that all

1 the data had a common slope with the assumption that TCE and TCA data had different slopes.  
2 ..... 340

3 Figure 4-9. Fold-changes in relative liver weight for data sets in male B6C3F1, Swiss, and  
4 NRM1 mice reported by TCE studies of duration 28–42 days (Kjellstrand et al., 1983a(Buben  
5 and O'Flaherty, 1985; Goel et al., 1992; Merrick et al., 1989) using internal dose-metrics  
6 predicted by the PBPK model described in Section 3.5: (A) dose-metric is the median estimate of  
7 the daily AUC of TCE in blood, (B) dose-metric is the median estimate of the total daily rate of  
8 TCE oxidation. Lines show linear regression. Use of liver oxidative metabolism as a  
9 dose-metric gives results qualitatively similar to (B), with  $R^2 = 0.86$ ..... 341

10 Figure 4-10. Dose-response relationship, expressed as (A) percentage incidence and  
11 (B) fold-increase over controls, for TCE hepatocarcinogenicity in NCI (1976). For comparison,  
12 incidences of carcinomas for NTP (1990), Anna et al. (1994), and Bull et al. (2002) are included,  
13 but without connecting lines since they are not appropriate for assessing the shape of the  
14 dose-response relationship..... 357

15 Figure 4-11. Dose-response relationship, expressed as (A) incidence and (B) fold-increase over  
16 controls, for TCE hepatocarcinogenicity in Maltoni et al. (1986). Note that the BT306  
17 experiment reported excessive mortality due to fighting, and so the paradigm was repeated in  
18 experiment BT306bis using mice from a different source..... 358

19 Figure 4-12. Dose-response data for hepatocellular carcinomas (HCC) (A) incidence and  
20 (B) multiplicity, induced by DCA from DeAngelo et al. (1999). Drinking water concentrations  
21 were 0, 0.05, 0.5, 1, 2, and 3.5 g/L, from which daily average doses were calculated using  
22 observed water consumption in the study..... 359

23 Figure 4-13. Reported incidences of hepatocellular carcinomas (HCC) and hepatocellular  
24 adenomas plus carcinomas (HCA + HCC) in various studies in B6C3F1 mice (DeAngelo et al.,  
25 2008; Pereira, 1996). Combined HCA + HCC were not reported in Pereira (1996)..... 361

26 Figure 4-14. Reported incidence of hepatocellular carcinomas induced by DCA and TCA in  
27 104-week studies (DeAngelo et al., 2008; DeAngelo et al., 1999). Only carcinomas were  
28 reported in DeAngelo et al. (1999), so combined adenomas and carcinomas could not be  
29 compared..... 363

30 Figure 4-15. Effects of dietary control on the dose-response curves for changes in liver tumor  
31 incidences induced by CH in diet (Leakey et al., 2003a). ..... 368

32 Figure 5-1. Flow-chart of the process used to derive the RfD and RfC for noncancer effects..... 2

33 Figure 5-2. Flow-chart for dose-response analyses of rodent noncancer effects using PBPK  
34 model-based dose-metrics. Square nodes indicate point values, circle nodes indicate  
35 distributions, and the inverted triangle indicates a (deterministic) functional relationship. .... 68

1 Figure 5-3. Schematic of combined interspecies, intraspecies, and route-to-route extrapolation  
2 from a rodent study LOAEL or NOAEL. In the case where BMD modeling is performed, the  
3 applied dose values are replaced by the corresponding median internal dose estimate, and the  
4 idPOD is the modeled BMDL in internal dose units. .... 69  
5 Figure 5-4. Flow-chart for uncertainty analysis of HECs and HEDs derived using PBPK  
6 model-based dose-metrics. Square nodes indicate point values, circle nodes indicate  
7 distributions, and the inverted triangle indicates a (deterministic) functional relationship. .... 92  
8 Figure 5-5. Flow-chart for dose-response analyses of rodent bioassays using PBPK model-based  
9 dose-metrics. Square nodes indicate point values, circular nodes indicate distributions, and the  
10 inverted triangles indicate a (deterministic) functional relationship. .... 132  
11 Figure 5-6. Flow-chart for uncertainty analysis of dose-response analyses of rodent bioassays  
12 using PBPK model-based dose-metrics. Square nodes indicate point values, circular nodes  
13 indicate distributions, and the inverted triangles indicate a (deterministic) functional relationship.  
14 ..... 153  
15 Figure 5-7. Flow-chart for route-to-route extrapolation of human site-specific cancer inhalation  
16 unit risks to oral slope factors. Square nodes indicate point values, circle nodes indicate  
17 distributions, and the inverted triangle indicates a (deterministic) functional relationship. .... 175

## LIST OF ABBREVIATIONS AND ACRONYMS

[ <sup>14</sup> C]TCE	[ <sup>14</sup> C]-radio labeled TCE
1,2-DCVC	S-(1,2-dichlorovinyl)-L-cysteine
17-β-HSD	17-β-hydroxy steroid dehydrogenase
8-OHdG	8-hydroxy-2' deoxyguanosine
ACO	acyl CoA oxidase
ADAF	age-dependent adjustment factor
ADME	absorption, distribution, metabolism, and excretion
AIC	Akaike Information Criteria
ALL	acute lymphoblastic leukemia
ALP	alkaline phosphatase
ALT	alanine aminotransferase
ANA	antinuclear antibodies
ANCA	antineutrophil-cytoplasmic antibody
AOAA	a beta-lyase inhibitor
ASD	autism spectrum disorder
ASPEN	Assessment System for Population Exposure Nationwide
AST	aspartate aminotransferase
ATF-2	activating transcription factor 2
ATSDR	Agency for Toxic Substances and Disease Registry
AUC	area-under-the-curve
AV	atrioventricular
AVC	atrioventricular canal
AZ DHS	Arizona Department of Health Services
BAER	brainstem auditory-evoked response
BAL	bronchoalveolar lavage
BMD	benchmark dose
BMDL	benchmark dose lower bound
BMDs	BenchMark Dose Software
BMI	body mass index
BMR	benchmark response
BUN	blood urea nitrogen
BW	body weight
CA DHS	California Department of Health Services
CH	chloral hydrate
CI	confidence interval
CLL	chronic lymphocytic leukemia
CNS	central nervous system
CO <sub>2</sub>	carbon dioxide
CoA	coenzyme A
cRfC	candidate RfC
cRfD	candidate RfD
CRT	choice reaction time
CYP	cytochrome P450

*This document is a draft for review purposes only and does not constitute Agency policy.*



## LIST OF ABBREVIATIONS AND ACRONYMS (continued)

DAL	dichloroacetyl lysine
DASO <sub>2</sub>	diallyl sulfone
DBP	dibutyl phthalate
DCA	dichloroacetic acid
DCAA	dichloroacetic anhydride
DCAC	dichloroacetyl chloride
DCE	dichloroethane
DCVC	dichlorovinyl cysteine
DCVG	S-dichlorovinyl glutathione
DEHP	di(2-ethylhexyl) phthalate
DHEAS	dehydroepiandrosterone sulphate
DNP	dinitrophenol
DPM	disintegrations per minute
dsDNA	double-stranded DNA
EC <sub>x</sub>	concentration of the chemical at which x% of the maximal effect is produced
EEG	electroencephalogram
EPA	U.S. Environmental Protection Agency
ERG	electroretinogram
FAA	fumarylacetoacetate
FDVE	fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether
FMO	flavin mono-oxygenase
FOB	functional observational battery
FSH	follicle-stimulating hormone
G6PDH	glucose 6-p dehydrogenase
GA	glomerular antigen
GABA	gamma-amino butyric acid
G-CSF	granulocyte colony stimulating factor
GD	gestation day
GFR	glomerular filtration rate
GGT	γ-glutamyl transpeptidase or γ-transpeptidase
GI	gastro-intestinal
GIS	geographic information system
GSH	glutathione
GSSG	oxidized GSH
GST	glutathione-S-transferase
GT	glutamyl transferase
H&E	hematoxylin and eosin
H <sub>2</sub> O	water
HCC	hepatocellular carcinoma
hCG	human chorionic gonadotropin
HCl	hydrochloric acid
HDL-C	high density lipoprotein-cholesterol
HEC	human equivalent concentration

*This document is a draft for review purposes only and does not constitute Agency policy.*

## LIST OF ABBREVIATIONS AND ACRONYMS (continued)

HED	human equivalent dose
HgCl <sub>2</sub>	mercuric chloride
HH	Hamberger and Hamilton
HPLC	high-performance liquid chromatography
HPT	hypothalamic-pituitary-testis
i.a.	intra-arterial
i.p.	intraperitoneal
i.v.	intravenous
IARC	International Agency for Research on Cancer
ICC	intrahepatic cholangiocarcinoma
ICD	International Classification of Disease
ICRP	The International Commission on Radiological Protection
idPOD	internal dose points of departure
IDR	incidence density ratio
IFN	interferon
IgE	immunoglobulin E
IGF-II	insulin-like growth factor-II (gene)
IL	interleukin
IPCS	International Programme on Chemical Safety
IUGR	intrauterine growth restriction
JEM	job-exposure matrix
JTEM	job-task-exposure matrix
LC	lethal concentration
LCL	lower confidence limit
LDH	lactate dehydrogenase
LEC <sub>x</sub>	lowest effective concentration corresponding to an extra risk of x%
LH	luteinizing hormone
lnPBC	blood-air partition coefficient
lnQCC	cardiac output
lnVMAXC	VMAX for oxidation
lnVPRC	ventilation-perfusion ratio
LOAEL	lowest observed adverse effect level
LOH	loss of heterozygosity
LORR	loss of righting reflex
MA	maleylacetone
MA DPH	Massachusetts Department of Public Health
MAA	maleylacetoacetate
MCA	monochloroacetic acid
MCMC	Markov chain Monte Carlo
MCP	methylclofenapate
MDA	malondialdehyde
MLE	maximum likelihood estimate
MNU	methyl nitrosourea

*This document is a draft for review purposes only and does not constitute Agency policy.*

## LIST OF ABBREVIATIONS AND ACRONYMS (continued)

MOA	mode of action
MS	
MSW	multistage Weibull
NAcDCVC	N-acetyl-S-(1,2-dichlorovinyl)-L-cysteine
NADH	nicotinamide adenine dinucleotide
NAG	N-acetyl- $\beta$ -D-glucosaminidase
NAT	N-acetyl transferase
NCI	National Cancer Institute
NF	
NHL	non-Hodgkin lymphoma
NK	natural killer
NOAEL	no-observed-adverse-effect level
NOEC	no-observed-effect concentration
NOEL	no-observed-effect level
NPMC	nonpurified rat peritoneal mast cells
NRC	National Research Council
NSATA	National-Scale Air Toxics Assessment
NTP	National Toxicology Program
NYS DOH	New York State Department of Health
ODE	ordinary differential equation
OECD	Organization for Economic Co-operation and Development
OFT	outflow tract
OP	oscillatory potential
OR	odds ratio
PAS	periodic acid-Schiff
PBPK	physiologically based pharmacokinetics
PC	partition coefficient
PCEs	polychromatic erythrocytes
PCNA	proliferating cell nuclear antigen
PCO	palmitoyl-CoA oxidase
PCR	polymerase chain reaction
p-cRfC	PBPK model-based candidate RfCs
p-cRfD	PBPK model-based candidate RfDs
PEG 400	polyethylene glycol 400
PFC	plaque-forming cell
PFU	plaque-forming units
PMR	proportionate mortality ratio
PND	postnatal day
PO <sub>2</sub>	partial pressure oxygen
POD	point of departure
PPAR	peroxisome proliferator activated receptor
QC	quality control
RBL-2H3	rat basophilic leukemia

*This document is a draft for review purposes only and does not constitute Agency policy.*

## LIST OF ABBREVIATIONS AND ACRONYMS (continued)

RCC	renal cell carcinoma
RfC	inhalation reference concentration
RfD	oral reference dose
ROS	reactive oxygen species
RR	relative risk
RRm	summary RR
RT	reaction time
S9	metabolic activation system
SBA	serum bile acids
SC	sensitivity coefficient
SCEs	sister chromatid exchanges
S-D	Sprague-Dawley
SD	standard deviation
SDH	sorbitol dehydrogenase
SE	standard error
SEER	Surveillance, Epidemiology, and End Results
SES	socioeconomic status
SGA	small for gestational age
SHBG	sex-hormone binding globulin
SIR	standardized incidence ratio
SMR	standardized mortality ratio
SNP	single nucleotide polymorphism
SRBC	sheep red blood cells
SRT	simple reaction time
SSB	single-strand breaks
SSCP	single strand conformation polymorphism
ssDNA	single-stranded DNA
TaClo	tetrahydro-beta-carbolines
TBARS	thiobarbiturate acid-reactive substances
TCA	trichloroacetic acid
TCAA	trichloroacetaldehyde
TCAH	trichloroacetaldehyde hydrate
TCE	trichloroethylene
TCOG	trichloroethanol-glucuronide conjugate
TCOH	trichloroethanol
ThX	T-helper Type X
TNF	tumor necrosis factor
TNFAI	wild- type TNF- $\alpha$ (G $\rightarrow$ A substitution at position -238)
TNFAII	TNF- $\alpha$ (G $\rightarrow$ A substitution at position -308)
TRI	Toxics Release Inventory
TSEP	trigeminal somatosensory evoked potential
TTC	total trichloro compounds
TWA	time-weighted average

## LIST OF ABBREVIATIONS AND ACRONYMS (continued)

U.S. EPA	U.S. Environmental Protection Agency
UCL	upper confidence limit
UDS	unscheduled DNA synthesis
UF	uncertainty factor
USGS	United States Geological Survey
U-TCA	urinary-TCA
U-TTC	urinary total trichloro-compounds
VEGF	vascular endothelial growth factor
VEP	visual evoked potential
VHL	von Hippel-Lindau
VLivC	liver volume
VOC	volatile organic compound
VSCC	voltage sensitive calcium channel
W	wakefulness
WHO	World Health Organization
YFF	fluorescent Y-bodies

## FOREWORD

The purpose of this Toxicological Review is to provide scientific support and rationale for the hazard and dose-response assessment in IRIS pertaining to chronic exposure to **trichloroethylene**. It is not intended to be a comprehensive treatise on the chemical or toxicological nature of **trichloroethylene**.

The intent of Chapter 6, *Major Conclusions in the Characterization of Hazard and Dose Response*, is to present the major conclusions reached in the derivation of the reference dose, reference concentration and cancer assessment, where applicable, and to characterize the overall confidence in the quantitative and qualitative aspects of hazard and dose response by addressing the quality of the data and related uncertainties. The discussion is intended to convey the limitations of the assessment and to aid and guide the risk assessor in the ensuing steps of the risk assessment process.

For other general information about this assessment or other questions relating to IRIS, the reader is referred to EPA's IRIS Hotline at (202) 566-1676 (phone), (202) 566-1749 (fax), or [hotline.iris@epa.gov](mailto:hotline.iris@epa.gov) (email address).

## AUTHORS, CONTRIBUTORS, AND REVIEWERS

### **CHEMICAL MANAGER**

Weihsueh A. Chiu  
National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

### **AUTHORS**

Ambuja Bale  
National Center for Environmental Assessment—Immediate Office  
U.S. Environmental Protection Agency  
Washington, DC

Stanley Barone  
National Center for Environmental Assessment—Immediate Office  
U.S. Environmental Protection Agency  
Washington, DC

Rebecca Brown  
National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Jane C. Caldwell  
National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Chao Chen  
National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Weihsueh A. Chiu  
National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Glinda Cooper  
National Center for Environmental Assessment—Immediate Office  
U.S. Environmental Protection Agency

Washington, DC

**AUTHORS, CONTRIBUTORS, AND REVIEWERS (continued)**

Ghazi Dannan

National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Marina Evans

National Health and Environmental Effects Research Laboratory  
(on detail to National Center for Environmental Assessment—Washington Office)  
U.S. Environmental Protection Agency  
Research Triangle Park, NC

John Fox

National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Kathryn Z. Guyton

National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Maureen R. Gwinn

National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Jennifer Jinot

National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Nagalakshmi Keshava

National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

John Lipscomb

National Center for Environmental Assessment—Cincinnati Office  
U.S. Environmental Protection Agency  
Cincinnati, OH



## **AUTHORS, CONTRIBUTORS, AND REVIEWERS (continued)**

Susan Makris  
National Center for Environmental Assessment—Washington Office  
U.S. Environmental Protection Agency  
Washington, DC

Miles Okino  
National Exposure Research Laboratory—Las Vegas Office  
U.S. Environmental Protection Agency  
Las Vegas, NV

Fred Power  
National Exposure Research Laboratory—Las Vegas Office  
U.S. Environmental Protection Agency  
Las Vegas, NV

John Schaum  
National Center for Environmental Assessment—Washington Office  
Office of Research and Development  
Washington, DC

Cheryl Siegel Scott  
National Center for Environmental Assessment—Washington Office  
Office of Research and Development  
Washington, DC

### **REVIEWERS**

This document has been reviewed by U.S. EPA scientists, reviewers from other Federal agencies and White House offices, and the public, and peer reviewed by independent scientists external to U.S. EPA. A summary and U.S. EPA's disposition of the comments received from the independent external peer reviewers and from the public is included in Appendix I.

### **INTERNAL EPA REVIEWERS**

Daniel Axelrad  
National Center for Environmental Economics

Robert Benson  
U.S. EPA Region 8

**AUTHORS, CONTRIBUTORS, AND REVIEWERS (continued)**

Ted Birner  
National Center for Environmental Assessment—Immediate Office

Nancy Chiu  
Office of Water

David Farrar  
National Center for Environmental Assessment—Cincinnati Office

Lynn Flowers  
National Center for Environmental Assessment—Immediate Office

Stiven Foster  
Office of Solid Waste and Emergency Response

Susan Griffin  
U.S. EPA Region 8

Samantha Jones  
National Center for Environmental Assessment—Immediate Office

Leonid Kopylev  
National Center for Environmental Assessment—Washington Office

Allan Marcus  
National Center for Environmental Assessment—Immediate Office

Gregory Miller  
Office of Children's Health Protection and Environmental Education

Deirdre Murphy  
Office of Air Quality Planning and Standards

Marian Olsen  
U.S. EPA Region 2

Peter Preuss  
National Center for Environmental Assessment—Immediate Office

Kathleen Raffaele  
National Center for Environmental Assessment—Washington Office

## AUTHORS, CONTRIBUTORS, AND REVIEWERS (continued)

William Sette  
Office of Solid Waste and Emergency Response

Bob Sonawane  
National Center for Environmental Assessment—Washington Office

Suryanarayana Vulimiri  
National Center for Environmental Assessment—Washington Office

Nina Ching Y. Wang  
National Center for Environmental Assessment—Cincinnati Office

Paul White  
National Center for Environmental Assessment—Washington Office

Marcia Bailey  
U.S. EPA Region 10

### ACKNOWLEDGMENTS

Drafts of Section 3.3 (TCE metabolism) were prepared for the U.S. EPA by Syracuse Research Corporation under contract. Additional support, including literature searches and retrievals and drafts of Appendix D were prepared for the U.S. EPA by the Oak Ridge Institute for Science and Education (ORISE) through interagency agreement number DW-89939822-01-0 with the U.S. Department of Energy (DOE). ORISE is managed by Oak Ridge Associated Universities under a contract with DOE. The PBPK modeling sections of this report are dedicated to the memory of Fred Power (1938–2007). His keen analytical mind will be greatly missed, but his gentle heart and big smile will be missed even more.

Additionally, we gratefully acknowledge Terri Konoza and Ellen Lorang of NCEA for their management of the document production and reference/citation management processes, and the following individuals for their support of these processes:

IntelliTech Systems, Inc.	ECFlex, Inc.
Chris Broyles	Heidi Glick
Stacey Lewis	Debbie Kleiser
Kathleen Secor	Crystal Lewis
Linda Tackett	Lana Wood

## EXECUTIVE SUMMARY

1  
2  
3  
4 There is substantial potential for human exposure to trichloroethylene (TCE), as it has a  
5 widespread presence in ambient air, indoor air, soil, and groundwater. At the same time, humans  
6 are likely to be exposed to a variety of compounds that are either metabolites of TCE or which  
7 have common metabolites or targets of toxicity. Once exposed, humans, as well as laboratory  
8 animal species, rapidly absorb TCE, which is then distributed to tissues via systemic circulation,  
9 extensively metabolized, and then excreted primarily in breath as unchanged TCE or carbon  
10 dioxide, or in urine as metabolites.

11 Based on the available human epidemiologic data and experimental and mechanistic  
12 studies, it is concluded that TCE poses a potential human health hazard for noncancer toxicity to  
13 the central nervous system, the kidney, the liver, the immune system, the male reproductive  
14 system, and the developing fetus. The evidence is more limited for TCE toxicity to the  
15 respiratory tract and female reproductive system. Following U.S. Environmental Protection  
16 Agency (U.S. EPA, 2005c) *Guidelines for Carcinogen Risk Assessment*, TCE is characterized as  
17 “*carcinogenic in humans by all routes of exposure.*” This conclusion is based on convincing  
18 evidence of a causal association between TCE exposure in humans and kidney cancer. The  
19 human evidence of carcinogenicity from epidemiologic studies of TCE exposure is compelling  
20 for non-Hodgkin Lymphoma but less convincing than for kidney cancer, and more limited for  
21 liver and biliary tract cancer. Less human evidence is found for an association between TCE  
22 exposure and other types of cancer, including bladder, esophageal, prostate, cervical, breast, and  
23 childhood leukemia, breast. Further support for the characterization of TCE as “*carcinogenic in*  
24 *humans by all routes of exposure*” is derived from positive results in multiple rodent cancer  
25 bioassays in rats and mice of both sexes, similar toxicokinetics between rodents and humans,  
26 mechanistic data supporting a mutagenic mode of action (MOA) for kidney tumors, and the lack  
27 of mechanistic data supporting the conclusion that any of the MOA(s) for TCE-induced rodent  
28 tumors are irrelevant to humans.

29 As TCE toxicity and carcinogenicity are generally associated with TCE metabolism,  
30 susceptibility to TCE health effects may be modulated by factors affecting toxicokinetics,  
31 including lifestage, gender, genetic polymorphisms, race/ethnicity, preexisting health status,  
32 lifestyle, and nutrition status. In addition, while these some of these factors are known risk  
33 factors for effects associated with TCE exposure, it is not known how TCE interacts with known  
34 risk factors for human diseases.

35 For noncancer effects, the most sensitive types of effects, based either on human  
36 equivalent concentrations/doses or on candidate inhalation reference concentrations (RfCs)/oral

1 reference doses (RfDs), appear to be developmental, kidney, and immunological (adult and  
2 developmental) effects. The neurological and reproductive effects appear to be about an order of  
3 magnitude less sensitive, with liver effects another two orders of magnitude less sensitive. The  
4 RfC estimate of **0.0004 ppm** (0.4 ppb or 2  $\mu\text{g}/\text{m}^3$ ) is based on route-to-route extrapolated results  
5 from oral studies for the critical effects of heart malformations (rats) and immunotoxicity (mice).  
6 This RfC value is further supported by route-to-route extrapolated results from an oral study of  
7 toxic nephropathy (rats). Similarly, the RfD estimate for noncancer effects of **0.0005 mg/kg-day**  
8 is based on the critical effects of heart malformations (rats), adult immunological effects (mice),  
9 and developmental immunotoxicity (mice), all from oral studies. This RfD value is further  
10 supported by results from an oral study for the effect of toxic nephropathy (rats) and route-to-  
11 route extrapolated results from an inhalation study for the effect of increased kidney weight  
12 (rats). There is high confidence in these noncancer reference values, as they are supported by  
13 moderate- to high-confidence estimates for multiple effects from multiple studies.

14 For cancer, the inhalation unit risk is  **$2 \times 10^{-2}$  per ppm** [ **$4 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$** ], based on  
15 human kidney cancer risks reported by Charbotel et al. (2006) and adjusted, using human  
16 epidemiologic data, for potential risk for tumors at multiple sites. The oral unit risk for cancer is  
17  **$5 \times 10^{-2}$  per mg/kg-day**, resulting from physiologically based pharmacokinetic model-based  
18 route-to-route extrapolation of the inhalation unit risk estimate based on the human kidney  
19 cancer risks reported in Charbotel et al. (2006) and adjusted, using human epidemiologic data,  
20 for potential risk for tumors at multiple sites. There is high confidence in these unit risks for  
21 cancer, as they are based on good quality human data, as well as being similar to unit risk  
22 estimates based on multiple rodent bioassays. There is both sufficient weight of evidence to  
23 conclude that TCE operates through a mutagenic MOA for kidney tumors and a lack of TCE-  
24 specific quantitative data on early-life susceptibility. Generally, the application of age-  
25 dependent adjustment factors (ADAFs) is recommended when assessing cancer risks for a  
26 carcinogen with a mutagenic MOA. However, because the ADAF adjustment applies only to the  
27 kidney cancer component of the total risk estimate, it is likely to have a minimal impact on the  
28 total cancer risk except when exposures are primarily during early life.

## 1. INTRODUCTION

1 This document presents background information and justification for the Integrated Risk  
2 Information System (IRIS) Summary of the hazard and dose-response assessment of  
3 **trichloroethylene**. IRIS Summaries may include oral reference dose (RfD) and inhalation  
4 reference concentration (RfC) values for chronic and other exposure durations, and a  
5 carcinogenicity assessment.

6 The RfD and RfC, if derived, provide quantitative information for use in risk assessments  
7 for health effects known or assumed to be produced through a nonlinear (presumed threshold)  
8 mode of action. The RfD (expressed in units of mg/kg/d) is defined as an estimate (with  
9 uncertainty spanning perhaps an order of magnitude) of a daily exposure to the human  
10 population (including sensitive subgroups) that is likely to be without an appreciable risk of  
11 deleterious effects during a lifetime. The inhalation RfC (expressed in units of ppm or  $\mu\text{g}/\text{m}^3$ ) is  
12 analogous to the oral RfD, but provides a continuous inhalation exposure estimate. The  
13 inhalation RfC considers toxic effects for both the respiratory system (portal-of-entry) and for  
14 effects peripheral to the respiratory system (extrarrespiratory or systemic effects). Reference  
15 values are generally derived for chronic exposures (up to a lifetime), but may also be derived for  
16 acute ( $\leq 24$  hours), short-term ( $>24$  hours up to 30 days), and subchronic ( $>30$  days up to 10% of  
17 lifetime) exposure durations, all of which are derived based on an assumption of continuous  
18 exposure throughout the duration specified. Unless specified otherwise, the RfD and RfC are  
19 derived for chronic exposure duration.

20 The carcinogenicity assessment provides information on the carcinogenic hazard  
21 potential of the substance in question and quantitative estimates of risk from oral and inhalation  
22 exposure may be derived. The information includes a weight-of-evidence judgment of the  
23 likelihood that the agent is a human carcinogen and the conditions under which the carcinogenic  
24 effects may be expressed. Quantitative risk estimates may be derived from the application of a  
25 low-dose extrapolation procedure. If derived, the oral slope factor is a plausible upper bound on  
26 the estimate of risk per mg/kg/d of oral exposure. Similarly, an inhalation unit risk is a plausible  
27 upper bound on the estimate of risk per ppm or  $\mu\text{g}/\text{m}^3$  in air breathed.

28 Development of these hazard identification and dose-response assessments for  
29 **trichloroethylene** has followed the general guidelines for risk assessment as set forth by the  
30 National Research Council (1983). U.S. EPA Guidelines and Risk Assessment Forum Technical  
31 Panel Reports that may have been used in the development of this assessment include the  
32 following: EPA Guidelines and Risk Assessment Forum technical panel reports that may have  
33 been used in the development of this assessment include the following: *Guidelines for the Health*  
34 *Risk Assessment of Chemical Mixtures* (U.S. EPA, 1986a), *Guidelines for Mutagenicity Risk*

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 *Assessment* (U.S. EPA, 1986b), *Recommendations for and Documentation of Biological Values*  
2 *for Use in Risk Assessment* (U.S. EPA, 1988), *Guidelines for Developmental Toxicity Risk*  
3 *Assessment* (U.S. EPA, 1991), *Interim Policy for Particle Size and Limit Concentration Issues in*  
4 *Inhalation Toxicity* (U.S. EPA, 1994a), *Methods for Derivation of Inhalation Reference*  
5 *Concentrations and Application of Inhalation Dosimetry* (U.S. EPA, 1994b), *Use of the*  
6 *Benchmark Dose Approach in Health Risk Assessment* (U.S. EPA, 1995), *Guidelines for*  
7 *Reproductive Toxicity Risk Assessment* (U.S. EPA, 1996), *Guidelines for Neurotoxicity Risk*  
8 *Assessment* (U.S. EPA, 1998), *Science Policy Council Handbook: Risk Characterization*  
9 (U.S. EPA, 2000a), *Benchmark Dose Technical Guidance Document* (U.S. EPA, 2000b),  
10 *Supplementary Guidance for Conducting Health Risk Assessment of Chemical Mixtures*  
11 (U.S. EPA, 2000c), *A Review of the Reference Dose and Reference Concentration Processes*  
12 (U.S. EPA, 2002), *Guidelines for Carcinogen Risk Assessment* (U.S. EPA, 2005a), *Supplemental*  
13 *Guidance for Assessing Susceptibility from Early-Life Exposure to Carcinogens* (U.S. EPA,  
14 2005b), *Science Policy Council Handbook: Peer Review* (U.S. EPA, 2006a), and *A Framework*  
15 *for Assessing Health Risks of Environmental Exposures to Children* (U.S. EPA, 2006b).

16         The literature search strategy employed for this compound was based on the Chemical  
17 Abstracts Service Registry Number and at least one common name. Any pertinent scientific  
18 information submitted by the public to the IRIS Submission Desk was also considered in the  
19 development of this document. The relevant literature was reviewed through December, 2010.  
20 It should be noted that references have been added to the Toxicological Review after the external  
21 peer review in response to peer reviewer's comments and for the sake of completeness. These  
22 references have not changed the overall qualitative and quantitative conclusions.

1  
2  
3

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 DRAFT—DO NOT CITE OR QUOTE



## 2. EXPOSURE CHARACTERIZATION

1           The purpose of this exposure characterization is to summarize information about  
 2 trichloroethylene (TCE) sources, releases, media levels, and exposure pathways for the general  
 3 population (occupational exposure is also discussed to a lesser extent). It is not meant as a  
 4 substitute for a detailed exposure assessment for a particular risk assessment application. While  
 5 this section primarily addresses TCE, it also includes some information on a number of related  
 6 compounds. These related compounds include metabolites of TCE and other parent compounds  
 7 that produce similar metabolites as shown in Table 2-1. The first column in this table lists the  
 8 principal TCE metabolites in humans (trichloroethanol, trichloroethanol-glucuronide and  
 9 trichloroacetic acid) as well as a number of minor ones (ATSDR, 1997b). The subsequent  
 10 columns list parent compounds that can produce some of the same metabolites. The metabolic  
 11 reaction pathways are much more complicated than implied here and it should be understood that  
 12 this table is intended only to provide a general understanding of which parent compounds lead to  
 13 which TCE metabolites. Exposure to the TCE-related compounds can alter or enhance TCE's  
 14 metabolism and toxicity by generating higher internal metabolite concentrations than would  
 15 result from TCE exposure by itself. This characterization is based largely on earlier work by Wu  
 16 and Schaum (2000, 2001), but also provides updates in a number of areas.

**Table 2-1. TCE metabolites and related parent compounds<sup>a</sup>**

TCE metabolites	Parent compounds				
	Tetrachloro-ethylene	1,1-Dichloro-ethane	1,1,1-Tri-chloroethane	1,1,1,2-Tetra-chloroethane	1,2-Dichloro-ethylene
Oxalic acid				X	X
Chloral	X				
Chloral hydrate	X				
Monochloroacetic acid	X	X	X	X	X
Dichloroacetic acid	X	X		X	
Trichloroacetic acid	X		X	X	
Trichloroethanol	X		X	X	
Trichloroethanol-glucuronide	X		X	X	

17 <sup>a</sup> X indicates that the parent compound can produce the corresponding metabolite (Hazardous Substances Data  
 18 Bank, <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>).

## 2.1. ENVIRONMENTAL SOURCES

TCE is a stable, colorless liquid with a chloroform-like odor and chemical formula  $C_2Cl_3H$  as diagrammed in Figure 2-1 (Lewis, 2001). Its chemical properties are listed in Table 2-2.




Figure 2-1. Molecular structure of TCE.

Table 2-2. Chemical properties of TCE

Property	Value	Reference
Molecular weight	131.39	Lide et al. (1998)
Boiling point	87.2°C	Lide et al. (1998)
Melting point	-84.7°C	Lide et al. (1998)
Density	1.4642 at 20°C	Budavari (1996)
Solubility	1,280 mg/L water at 25°C	Horvath et al. (1999)
Vapor pressure	69.8 mmHG @ 25°C	Boublik et al.(1984)
Vapor density	4.53 (air = 1)	Budavari (1996)
Henry's Law Constant	$9.85 \times 10^{-3}$ atm-cu m/mol @ 25°C	Leighton and Calo (1981)
Octanol/water partition coefficient	$\log K_{ow} = 2.61$	Hansch et al. (1995)
Air concentration conversion	1 ppb = $5.38 \mu\text{g}/\text{m}^3$	HSDB (2002)

Trichloroethylene has been produced commercially since the 1920s in many countries by chlorination of ethylene or acetylene. Its use in vapor degreasing began in the 1920s. In the 1930s, it was introduced for use in dry cleaning. This use was largely discontinued in the 1950s and was replaced with tetrachloroethylene (ATSDR, 1997b). More recently, 80–90% of trichloroethylene production worldwide is used for degreasing metals (IARC, 1995a). It is also used in adhesives, paint-stripping formulations, paints, lacquers, and varnishes (SRI

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 International, 1992). A number of past uses in cosmetics, drugs, foods, and pesticides have now  
 2 been discontinued including use as an extractant for spice oleoresins, natural fats and oils, hops,  
 3 and decaffeination of coffee (IARC, 1995a), and as a carrier solvent for the active ingredients of  
 4 insecticides and fungicides, and for spotting fluids (ATSDR, 1997b; WHO, 1985). The  
 5 production of TCE in the United States peaked in 1970 at 280 million kg (616 million pounds)  
 6 and declined to 60 million kg (132 million pounds) in 1998 (USGS, 2006). In 1996, the United  
 7 States imported 4.5 million kg (10 million pounds) and exported 29.5 million kg (65 million  
 8 pounds) ("Chemical Profile: Trichloroethylene," 1997). Table 2-3 summarizes the basic  
 9 properties and principal uses of the TCE related compounds.

10  
 11 **Table 2-3. Properties and uses of TCE related compounds**

	<b>Water solubility (mg/L)</b>	<b>Vapor pressure (mmHG)</b>	<b>Uses</b>	<b>Sources</b>
Tetrachloroethylene	150	18.5 @25°C	Dry cleaning, degreasing, solvent	a
1,1,1-Trichloroethane	4,400	124 @25°C	Solvents, degreasing	a
1,2-Dichloroethylene	3,000–6,000	273–395 @30°C	Solvents, chemical intermediates	a
1,1,1,2-Tetrachloroethane	1,100	14 @25°C	Solvents, but currently not produced in United States	a,b
1,1-Dichloroethane	5,500	234 @25°C	Solvents, chemical intermediates	a
Chloral	High	35 @20°C	Herbicide production	a
Chloral hydrate	High	NA	Pharmaceutical production	a
Monochloroacetic acid	High	1 @43°C	Pharmaceutical production	a
Dichloroacetic acid	High	<1 @20°C	Pharmaceuticals, not widely used	a
Trichloroacetic acid	High	1 @50°C	Herbicide production	a
Oxalic acid	220,000	0.54 @105°C	Scouring/cleaning agent, degreasing	b
Dichlorovinyl	Not	Not available	Not available	

*This document is a draft for review purposes only and does not constitute Agency policy.*

cysteine	available			
Trichloroethanol	Low	NA	Anesthetics and chemical intermediate	<sup>c</sup>

1 <sup>a</sup> Wu and Schaum (2001).

2 <sup>b</sup> HSDB (2002).

3 <sup>c</sup> Lewis (2001).

1 Releases of TCE from nonanthropogenic activities are negligible (HSDB, 2002). Most of the  
 2 TCE used in the United States is released to the atmosphere, primarily from vapor degreasing  
 3 operations (ATSDR, 1997b). Releases to air also occur at treatment and disposal facilities, water  
 4 treatment facilities, and landfills (ATSDR, 1997b). TCE has also been detected in stack  
 5 emissions from municipal and hazardous waste incineration (ATSDR, 1997b). TCE is on the list  
 6 for reporting to U.S. Environmental Protection Agency (EPA)'s Toxics Release Inventory (TRI).  
 7 Reported releases into air predominate over other types and have declined over the period  
 8 1994–2004 (see Table 2-4).


9  
 10 **Table 2-4. TRI releases of TCE (pounds/year)**

Year	On-site fugitive air	On-site stack air	Total on-site air emissions	On-site surface water discharges	Total on-site underground injection	Total on-site releases to land	Total off-site disposal or other releases	Total on- and off-site disposal or other releases
1994	15,018,818	15,929,943	30,948,761	1,671	288	4,070	96,312	31,051,102
1995	12,498,086	13,784,853	26,282,939	1,477	550	3,577	74,145	26,362,688
1996	10,891,223	10,995,228	21,886,451	541	1,291	9,740	89,527	21,987,550
1997	9,276,150	8,947,909	18,224,059	568	986	3,975	182,423	18,412,011
1998	6,769,810	6,504,289	13,274,099	882	593	800	136,766	13,413,140
1999	5,861,635	4,784,057	10,645,692	1,034	0	148,867	192,385	10,987,978
2000	5,485,493	4,375,516	9,861,009	593	47,877	9,607	171,952	10,091,038
2001	4,968,282	3,453,451	8,421,733	406	98,220	12,609	133,531	8,666,499
2002	4,761,104	3,436,289	8,197,393	579	140,190	230	139,398	8,477,790
2003	3,963,054	3,121,718	7,084,772	595	90,971	150,642	66,894	7,393,873
2004	3,040,460	3,144,980	6,185,440	216	123,637	2	71,780	6,381,075
2005	2,733,983	2,893,168	5,627,152	533	86,817	4,711	60,074	5,779,287
2006	2,816,241	2,795,184	5,611,425	482	0	77,339	90,758	5,780,004

12  
 13 Source: EPA TRI Explorer, <http://www.epa.gov/triexplorer/trends.htm>.


14  
 15  
 16 Under the National-Scale Air Toxics Assessment (NSATA) program, EPA has developed  
 17 an emissions inventory for TCE (U.S. EPA, 2006b). The inventory includes sources in the  
 18 United States plus the Commonwealth of Puerto Rico and the U.S. Virgin Islands. The types of  
 19 emission sources in the inventory include large facilities, such as waste incinerators and factories  
 20 and smaller sources, such as dry cleaners and small manufacturers. Figures 2-2 and 2-3

*This document is a draft for review purposes only and does not constitute Agency policy.*



1  
2

**Figure 2-2. Source contribution to TCE emissions.**



3  
4

**Figure 2-3. Annual emissions of TCE.**

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 show the results of the 1999 emissions inventory for TCE. Figure 2-2 shows the percent  
2 contribution to total emissions by source category. A variety of sources have TCE emissions  
3 with the largest ones identified as halogenated solvent cleaners and metal parts and products.  
4 Figure 2-3 shows a national map of the emission density (tons/sq mi-yr) for TCE. This map  
5 shows the highest densities in the far west and northeastern regions of the United States.  
6 Emissions range from 0–4.12 tons/sq mi-yr.

7

## 2.2. ENVIRONMENTAL FATE

### 2.2.1. Fate in Terrestrial Environments

8 The dominant fate of trichloroethylene released to surface soils is volatilization. Because  
9 of its moderate water solubility, trichloroethylene introduced into soil (e.g., landfills) also has the  
10 potential to migrate through the soil into groundwater. The relatively frequent detection of  
11 trichloroethylene in groundwater confirms this. Biodegradation in soil and groundwater may  
12 occur at a relatively slow rate (half-lives on the order of months to years) (Howard et al., 1991).  
13

### 2.2.2. Fate in the Atmosphere

14 In the atmosphere, trichloroethylene is expected to be present primarily in the vapor  
15 phase, rather than sorbed to particulate, because of its high vapor pressure. Some removal by  
16 scavenging during wet precipitation is expected because of its moderate water solubility. The  
17 major degradation process affecting vapor phase trichloroethylene is photo-oxidation by  
18 hydroxyl radicals. Photolysis in the atmosphere proceeds very slowly, if at all.  
19 Trichloroethylene does not absorb ultraviolet light at wavelengths of less than 290 nm and thus  
20 will not directly photolyze. Based on measured rate data for the vapor phase photo-oxidation  
21 reaction with hydroxyl radicals, the estimated half-life of trichloroethylene in the atmosphere is  
22 on the order of 1–11 days with production of phosgene, dichloroacetyl chloride, and formyl  
23 chloride. Under smog conditions, degradation is more rapid (half-life on the order of hours)  
24 (Howard et al., 1991; HSDB, 2002).

25

### 2.2.3. Fate in Aquatic Environments

1           The dominant fate of trichloroethylene released to surface waters is volatilization  
2 (predicted half-life of minutes to hours). Bioconcentration, biodegradation, and sorption to  
3 sediments and suspended solids are not thought to be significant (HSDB, 2002).  
4 Trichloroethylene is not hydrolyzed under normal environmental conditions. However, slow  
5 photo-oxidation in water (half-life of 10.7 months) has been reported (Howard et al., 1991;  
6 HSDB, 2002).

## 2.3. EXPOSURE CONCENTRATIONS

7           TCE levels in the various environmental media result from the releases and fate processes  
8 discussed in Sections 2.1 and 2.2. No statistically based national sampling programs have been  
9 conducted that would allow estimates of true national means for any environmental medium. A  
10 substantial amount of air and groundwater data, however, has been collected as well as some  
11 data in other media, as described below.  
12

### 2.3.1. Outdoor Air—Measured Levels

13           TCE has been detected in the air throughout the United States. According to ATSDR  
14 (1997b), atmospheric levels are highest in areas concentrated with industry and population, and  
15 lower in remote and rural regions. Table 2-5 shows levels of TCE measured in the ambient air at  
16 a variety of locations in the United States.

17           More recent ambient air measurement data for TCE were obtained from EPA's Air  
18 Quality System database at the AirData Web site: <http://www.epa.gov/air/data/index.html>  
19 (2007). These data were collected from a variety of sources including state and local  
20 environmental agencies. The data are not from a statistically based survey and cannot be  
21 assumed to provide nationally representative values. The most recent data (2006) come from  
22 258 monitors located in 37 states. The means for these monitors range from 0.03–7.73  $\mu\text{g}/\text{m}^3$   
23 and have an overall average of 0.23  $\mu\text{g}/\text{m}^3$ . Table 2-6 summarizes the data for the years  
24 1999–2006. The data suggest that levels have remained fairly constant since 1999 at about  
25 0.3  $\mu\text{g}/\text{m}^3$ . Table 2-7 shows the monitoring data organized by land setting (rural, suburban, or  
26 urban) and land use (agricultural, commercial, forest, industrial, mobile, and residential). Urban  
27 air levels are almost four times higher than rural areas. Among the land use categories, TCE  
28 levels are highest in commercial/industrial areas and lowest in forest areas.  
29



### 2.3.2. Outdoor Air—Modeled Levels

1 Under the National-Scale Air Toxics Assessment program, EPA has compiled emissions  
2 data and modeled air concentrations/exposures for the Criteria Pollutants and Hazardous Air  
3 Pollutants (U.S. EPA, 2006b). The results of the 1999 emissions inventory for TCE were  
4 discussed earlier and results presented in Figures 2-2 and 2-3. A computer simulation model  
5 known as the Assessment System for Population Exposure Nationwide (ASPEN) is used to  
6 estimate toxic air pollutant concentrations (<http://www.epa.gov/ttnatw01/nata/aspn.html>). This  
7 model is based on the EPA's Industrial Source Complex Long Term model which simulates the  
8 behavior of the pollutants after they are emitted into the atmosphere. ASPEN uses estimates of  
9 toxic air pollutant emissions and meteorological data from National Weather Service Stations to  
10 estimate

1 **Table 2-5. Concentrations of trichloroethylene in ambient air**

2

Area	Year	Concentration ( $\mu\text{g}/\text{m}^3$ )	
		Mean	Range
<b><i>Rural</i></b>			
Whiteface Mountain, NY <sup>a</sup>	1974	0.5	<0.3–1.9
Badger Pass, CA <sup>a</sup>	1977	0.06	0.005–0.09
Reese River, NV <sup>a</sup>	1977	0.06	0.005–0.09
Jetmar, KS <sup>a</sup>	1978	0.07	0.04–0.11
All rural sites	1974–1978		0.005–1.9
<b><i>Urban and Suburban</i></b>			
New Jersey <sup>a</sup>	1973–79	9.1	ND–97
New York City, NY <sup>a</sup>	1974	3.8	0.6–5.9
Los Angeles, CA <sup>a</sup>	1976	1.7	0.14–9.5
Lake Charles, LA <sup>a</sup>	1976–78	8.6	0.4–11.3
Phoenix, AZ <sup>a</sup>	1979	2.6	0.06–16.7
Denver, CO <sup>a</sup>	1980	1.07	0.15–2.2
St. Louis, MO <sup>a</sup>	1980	0.6	0.1–1.3
Portland, OR <sup>a</sup>	1984	1.5	0.6–3.9
Philadelphia, PA <sup>a</sup>	1983–1984	1.9	1.6–2.1
Southeast Chicago, IL <sup>b</sup>	1986–1990	1.0	
East St. Louis, IL <sup>b</sup>	1986–1990	2.1	
District of Columbia <sup>c</sup>	1990–1991	1.94	1–16.65
Urban Chicago, IL <sup>d</sup>	pre–1993	0.82–1.16	
Suburban Chicago, IL <sup>d</sup>	pre–1993	0.52	
300 cities in 42 states <sup>e</sup>	pre–1986	2.65	
Several Canadian Cities <sup>f</sup>	1990	0.28	
Several United States Cities <sup>f</sup>	1990	6.0	
Phoenix, AZ <sup>g</sup>	1994–1996	0.29	0–1.53
Tucson, AZ <sup>g</sup>	1994–1996	0.23	0–1.47
All urban/suburban sites	1973–1996		0–97

3

4 <sup>a</sup> IARC (1995a).

5 <sup>b</sup> Sweet (1992).

6 <sup>c</sup> Hendler (1992).

7 <sup>d</sup> Scheff (1993).

8 <sup>e</sup> Shah (1988).

9 <sup>f</sup> Bunce (1994).

10 <sup>g</sup> Zielinska-Psuja (1998).

11

12 ND = nondetect

13

**Table 2-6. TCE ambient air monitoring data ( $\mu\text{g}/\text{m}^3$ )**

Year	Number of monitors	Number of states	Mean	Standard deviation	Median	Range
1999	162	20	0.30	0.53	0.16	0.01–4.38
2000	187	28	0.34	0.75	0.16	0.01–7.39
2001	204	31	0.25	0.92	0.13	0.01–12.90
2002	259	41	0.37	1.26	0.13	0.01–18.44
2003	248	41	0.35	0.64	0.16	0.02–6.92
2004	256	37	0.32	0.75	0.13	0.00–5.78
2005	313	38	0.43	1.05	0.14	0.00–6.64
2006	258	37	0.23	0.55	0.13	0.03–7.73

Source: EPA’s Air Quality System database at the AirData Web site:  
<http://www.epa.gov/air/data/index.html>.

**Table 2-7. Mean TCE air levels across monitors by land setting and use (1985–1998)**

	Rural	Suburban	Urban	Agricultural	Commercial	Forest	Industrial	Mobile	Residential
Mean concentration ( $\mu\text{g}/\text{m}^3$ )	0.42	1.26	1.61	1.08	1.84	0.1	1.54	1.5	0.89
<i>n</i>	93	500	558	31	430	17	186	39	450

Source: EPA’s Air Quality System database at the AirData Web site:  
<http://www.epa.gov/air/data/index.html>.

air toxics concentrations nationwide. The ASPEN model takes into account important determinants of pollutant concentrations, such as

- rate of release;
- location of release;
- the height from which the pollutants are released;

*This document is a draft for review purposes only and does not constitute Agency policy.*

- 1 • wind speeds and directions from the meteorological stations nearest to the release;
- 2 • breakdown of the pollutants in the atmosphere after being released (i.e., reactive decay);
- 3 • settling of pollutants out of the atmosphere (i.e., deposition); and
- 4 • transformation of one pollutant into another (i.e., secondary formation).


5  
6  
7 The model estimates toxic air pollutant concentrations for every census tract in the  
8 continental United States, the Commonwealth of Puerto Rico and the U.S. Virgin Islands.  
9 Census tracts are land areas defined by the U.S. Bureau of the Census and typically contain about  
10 4,000 residents each. Census tracts are usually smaller than 2 square miles in size in cities but  
11 much larger in rural areas.

12 Figure 2-4 shows the results of the 1999 ambient air concentration modeling for TCE.  
13 The county median air levels range from 0–3.79  $\mu\text{g}/\text{m}^3$  and an overall median of 0.054  $\mu\text{g}/\text{m}^3$ .  
14 They have a pattern similar to the emission densities shown in Figure 2-3. These NSATA  
15 modeled levels appear lower than the monitoring results presented above. For example, the 1999  
16 air monitoring data (see Table 2-6) indicates a median outdoor air level of 0.16  $\mu\text{g}/\text{m}^3$  which is  
17 about three times as high as the modeled 1999 county median (0.054  $\mu\text{g}/\text{m}^3$ ). However, it should  
18 be understood that the results from these two efforts are not perfectly comparable. The modeled  
19 value is a median of county levels for the entire United States which includes many rural areas.  
20 The monitors cover many fewer areas ( $n = 162$  for 1999) and most are in nonrural locations. A  
21 better analysis is provided by EPA (2006b) which presents a comparison of modeling results  
22 from NSATA to measured values at the same locations. For 1999, it was found that  
23 formaldehyde levels were underestimated at 79% of the sites ( $n = 92$ ). Thus, while the NSATA  
24 modeling results are useful for understanding geographic distributions, they may frequently  
25 underestimate ambient levels.  
26

### 27 **2.3.3. Indoor Air**

28 TCE can be released to indoor air from use of consumer products that contain it (i.e.,  
29 adhesives and tapes), vapor intrusion (migration of volatile chemicals from the subsurface into  
30 overlying buildings) and volatilization from the water supply. Where such sources are present, it  
31 is likely that indoor levels will be higher than outdoor levels. A number of studies have  
32 measured indoor levels of TCE:  
33

- The 1987 EPA Total Exposure Assessment Methodology study (U.S. EPA, 1987) showed that the ratio of indoor to outdoor TCE concentrations for residences in Greensboro, NC, was about 5:1.



**Figure 2-4. Modeled ambient air concentrations of TCE.**

- In two homes using well water with TCE levels averaging 22–128  $\mu\text{g/L}$ , the TCE levels in bathroom air ranged from  $<500$ – $40,000 \mu\text{g/m}^3$  when the shower ran less than 30 minutes (Andelman, 1985).
- Shah and Singh (1988) report an average indoor level of  $7.2 \mu\text{g/m}^3$  based on over 2,000 measurements made in residences and workplaces during 1981–1984 from various locations across the United States.
- Hers et al. (2001) provides a summary of indoor air TCE measurements at locations in United States, Canada, and Europe with a range of  $<1$ – $165 \mu\text{g/m}^3$ .
- Sapkota et al. (2005) measured TCE levels inside and outside of the Baltimore Harbor Tunnel toll booths during the summer of 2001. Mean TCE levels were  $3.11 \mu\text{g/m}^3$  indoors and  $0.08 \mu\text{g/m}^3$  outdoors based on measurements on 7 days. The authors

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 speculated that indoor sources, possibly dry cleaning residues on uniforms, were the  
2 primary source of the indoor TCE.

- 3 • Sexton et al. (2005) measured TCE levels inside and outside residences in  
4 Minneapolis/St. Paul metropolitan area. Two day samples were collected over  
5 three seasons in 1999. Mean TCE levels were  $0.5 \mu\text{g}/\text{m}^3$  indoors ( $n = 292$ ),  $0.2 \mu\text{g}/\text{m}^3$   
6 outdoors ( $n = 132$ ) and  $1.0 \mu\text{g}/\text{m}^3$  based on personal sampling ( $n = 288$ ).
- 7 • Zhu et al. (2005) measured TCE levels inside and outside of residences in Ottawa,  
8 Canada. Seventy-five homes were randomly selected and measurements were made  
9 during the winter of 2002/2003. TCE was above detection limits in the indoor air of  
10 33% of the residences and in the outdoor air of 19% of the residences. The mean levels  
11 were  $0.06 \mu\text{g}/\text{m}^3$  indoors and  $0.08 \mu\text{g}/\text{m}^3$  outdoors. Given the high frequency of  
12 nondetects, a more meaningful comparison can be made on basis of the 75<sup>th</sup> percentiles:  
13  $0.08 \mu\text{g}/\text{m}^3$  indoors and  $0.01 \mu\text{g}/\text{m}^3$  outdoors.

14  
15  
16 TCE levels measured indoors have been directly linked to vapor intrusion at two sites in New  
17 York:

- 18  
19  
20 • TCE vapor intrusion has occurred in buildings/residences near a former Smith Corona  
21 manufacturing facility located in Cortlandville, NY. An extensive sampling program  
22 conducted in 2006 has detected TCE in groundwater ( $1\text{--}13 \mu\text{g}/\text{L}$ ), soil gas ( $6\text{--}97 \mu\text{g}/\text{m}^3$ ),  
23 subslab gas ( $2\text{--}1,600 \mu\text{g}/\text{m}^3$ ), and indoor air ( $1\text{--}17 \mu\text{g}/\text{m}^3$ ) (NYSDEC, 2006a).
- 24 • Evidence of vapor intrusion of TCE has also been reported in buildings and residences in  
25 Endicott, NY. Sampling in 2003 showed total volatile organic compounds (VOCs) in  
26 soil gas exceeding  $10,000 \mu\text{g}/\text{m}^3$  in some areas. Indoor air sampling detected TCE levels  
27 ranging from  $1\text{--}140 \mu\text{g}/\text{m}^3$  (NYSDEC, 2006b).

28  
29  
30 Little et al. (1992) developed attenuation coefficients relating contaminants in soil gas  
31 (assumed to be in chemical equilibrium with the groundwater) to possible indoor levels as a  
32 result of vapor intrusion. On this basis they estimated that TCE groundwater levels of  $540 \mu\text{g}/\text{L}$ ,  
33 (a high contamination level) could produce indoor air levels of  $5\text{--}500 \mu\text{g}/\text{m}^3$ . Vapor intrusion is  
34 likely to be a significant source only in situations where residences are located near soils or  
35 groundwater with high contamination levels. EPA (2002b) recommends considering vapor  
36 intrusion when volatiles are suspected to be present in groundwater or soil at a depth of  
37  $<100$  feet. Hers et al. (2001) concluded that the contribution of VOCs from subsurface sources  
38 relative to indoor sources is small for most chemicals and sites.

### 2.3.4. Water

A number of early (pre-1990) studies measured TCE levels in natural water bodies (levels in drinking water are discussed later in this section) as summarized in Table 2-8.

**Table 2-8. Concentrations of trichloroethylene in water based on pre-1990 studies**

Water type	Location	Year	Mean (µg/L)	Median (µg/L)	Range (µg/L)	Number of samples	Ref.
Industrial effluent	U.S.	83		0.5		NR	IARC (1995a)
Surface waters	U.S.	83		0.1		NR	IARC (1995a)
Rainwater	Portland, OR	84	0.006		0.002–0.02	NR	Ligocki et al. (1985)
Groundwater	MN	83			0.2–144	NR	Sabel and Clark (1984)
	NJ	76			≤1,530	NR	Burmester et al. (1982)
	NY	80			≤3,800	NR	Burmester et al. (1982)
	PA	80			≤27,300	NR	Burmester et al. (1982)
	MA	76			≤900	NR	Burmester et al. (1982)
	AZ				8.9–29	NR	IARC (1995a)
Drinking water	U.S.	76			0.2–49		IARC (1995a)
	U.S.	77			0–53		IARC (1995a)
	U.S.	78			0.5–210		IARC (1995a)
	MA	84			max. 267		IARC (1995a)
	NJ	84	23.4		max. 67	1130	Cohn et al. (1994a)
	CA	85			8–12	486	EPA, (1987)
	CA	84	66			486	EPA, (1987)
	NC	84	5			48	EPA, (1987)
ND	84	5			48	EPA, (1987)	

NR = Not Reported.

According to IARC (1995a), the reported median concentrations of TCE in 1983–1984 were 0.5 µg/L in industrial effluents and 0.1 µg/L in ambient water. Results from an analysis of the EPA STORET Data Base (1980–1982) showed that TCE was detected in 28% of 9,295 surface water reporting stations nationwide (ATSDR, 1997b). A more recent search of the STORET database for TCE measurements nationwide during 2008 in streams, rivers and lakes indicated three detects (0.03–0.04 µg/L) out of 150 samples (STORET Database, <http://www.epa.gov/storet/dbtop.html>).

ATSDR (1997b) has reported that TCE is the most frequently reported organic contaminant in groundwater and the one present in the highest concentration in a summary of ground water analyses reported in 1982. It has been estimated that between 9 and 34% of the drinking water supply sources tested in the United States may have some trichloroethylene contamination. This estimate is based on available Federal and State surveys (ATSDR, 1997b).

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Squillace et al. (2004) reported TCE levels in shallow groundwater based on data from  
2 the National Water Quality Assessment Program managed by USGS. Samples from 518 wells  
3 were collected from 1996–2002. All wells were located in residential or commercial areas and  
4 had a median depth of 10 m. The authors reported that approximately 8.3% of the well levels  
5 were above the detection limit (level not specified), 2.3% were above 0.1 µg/L and 1.7% were  
6 above 0.2 µg/L.

7 As part of the Agency’s first Six-Year Review, EPA obtained analytical results for over  
8 200,000 monitoring samples reported at 23,035 public water systems (PWS) in 16 states (U.S.  
9 EPA, 2003a). Approximately 2.6% of the systems had at least one sample exceed a minimum  
10 reporting level of 0.5 µg/L; almost 0.65% had at least one sample that exceeds the maximum  
11 contaminant level (MCL) of 5 µg/L. Based on average system concentrations estimated by EPA,  
12 54 systems (0.23%) had an average concentration that exceeded the MCL. EPA’s statistical  
13 analysis to extrapolate the sample result to all systems regulated for TCE resulted in an estimate  
14 of 154 systems with average TCE concentrations that exceed the MCL.

15 TCE concentrations in ground water have been measured extensively in California. The  
16 data were derived from a survey of water utilities with more than 200 service connections. The  
17 survey was conducted by the California Department of Health Services (CDHS, 1986). From  
18 January 1984 through December 1985, untreated water from wells in 819 water systems were  
19 sampled for organic chemical contamination. The water systems use a total of 5,550 wells,  
20 2,947 of which were sampled. TCE was found in 187 wells at concentrations up to 440 µg/L,  
21 with a median concentration among the detects of 3.0 µg/L. Generally, the wells with the highest  
22 concentrations were found in the heavily urbanized areas of the state. Los Angeles County  
23 registered the greatest number of contaminated wells (149).

24 A second California study collected data on TCE levels in public drinking water  
25 (Williams et al., 2002). The data were obtained from the CA DHS. The data spanned the years  
26 1995–2001 and the number of samples for each year ranged from 3,447–4,226. The percent of  
27 sources that were above the detection limit ranged from 9.6–11.7 per year (detection limits not  
28 specified). The annual average detected concentrations ranged from 14.2–21.6 µg/L. Although  
29 not reported, the overall average concentration of the samples (assuming an average of 20 µg/L  
30 among the samples above the detection limit, 10% detection rate and 0 for the nondetects) would  
31 be about 2 µg/L.

32 The USGS (2006) conducted a national assessment of 55 VOCs, including  
33 trichloroethylene, in ground water. A total of 3,500 water samples were collected during  
34 1985–2001. Samples were collected at the well head prior to any form of treatment. The types  
35 of wells sampled included 2,400 domestic wells and 1,100 public wells. Almost 20% of the



1 samples contained one or more of the VOCs above the assessment level of 0.2 µg/L. The  
2 detection frequency increased to over 50% when a subset of samples was analyzed with a low  
3 level method that had an assessment level of 0.02 µg/L. The largest detection frequencies were  
4 observed in California, Nevada, Florida, the New England States, and Mid-Atlantic states. The  
5 most frequently detected VOCs (>1% of samples) include TCE, tetrachloroethylene,  
6 1,1,1-trichloroethane (methyl chloroform), 1,2 dichloroethylene, and 1,1-dichloroethane.  
7 Findings specific to TCE include the following:

- 8
- 9
- 10 • Detection frequency was 2.6% at 0.2 µg/L and was 3.8% at 0.02 µg/L.
- 11 • The median concentration was 0.15 µg/L with a range of 0.02–100 µg/L.
- 12 • The number of samples exceeding the MCL (5 µg/L) was six at domestic wells and nine  
13 at public wells.
- 14
- 15

16 USGS (2006) also reported that four solvents (TCE, tetrachloroethylene,  
17 1,1,1-trichloroethane and methylene chloride) occurred together in 5% of the samples. The most  
18 frequently occurring two-solvent mixture was TCE and tetrachloroethylene. The report stated  
19 that the most likely reason for this co-occurrence is the reductive dechlorination of  
20 tetrachloroethylene to TCE.  
21

### 22 **2.3.5. Other Media**

23 Levels of TCE were found in the sediment and marine animal tissue collected in  
24 1980–1981 near the discharge zone of a Los Angeles County waste treatment plant.  
25 Concentrations were 17 µg/L in the effluent, <0.5 µg/kg in dry weight in sediment, and  
26 0.3–7 µg/kg wet weight in various marine animal tissue (IARC, 1995a). TCE has also been  
27 found in a variety of foods. U.S. Food and Drug Administration (FDA) has limits on TCE use as  
28 a food additive in decaffeinated coffee and extract spice oleoresins (see Table 2-15). Table 2-9  
29 summarizes data from two sources:

- 30
- 31 • IARC (1995a) reports average concentrations of TCE in limited food samples collected in  
32 the United States.
- 33 • Jones and Smith (2003) measured VOC levels in over 70 foods collected from 1996–  
34 2000 as part of the FDA’s Total Diet Program. All foods were collected directly from

1 supermarkets. Analysis was done on foods in a ready-to-eat form. Sample sizes for most  
 2 foods were in the 2–5 range.

3  
 4 **Table 2-9. Levels in food**  
 5

IARC (1995a)	Fleming-Jones and Smith (2003)
Cheese 3.8 µg/kg Butter and Margarine 73.6 µg/kg	Cheese 2–3 µg/kg Butter 7–9 µg/kg Margarine 2–21 µg/kg Cheese Pizza 2 µg/kg
Peanut Butter 0.5 µg/kg	Nuts 2–5 µg/kg Peanut Butter 4–70 µg/kg
	Ground Beef 3–6 µg/kg Beef Frankfurters 2–105 µg/kg Hamburger 5–9 µg/kg Cheeseburger 7 µg/kg Chicken Nuggets 2–5 µg/kg Bologna 2–20 µg/kg Pepperoni Pizza 2 µg/kg
	Banana 2 µg/kg Avocado 2–75 µg/kg Orange 2 µg/kg
	Chocolate Cake 3–57 µg/kg Blueberry Muffin 3–4 µg/kg Sweet Roll 3 µg/kg Chocolate Chip Cookies 2–4 µg/kg Apple Pie 2–4 µg/kg Doughnuts 3 µg/kg
	Tuna 9–11 µg/kg
Cereals 3 µg/kg Grain-based Foods 0.9 µg/kg	Cereal 3 µg/kg
	Popcorn 4–8 µg/kg French Fries 3 µg/kg Potato Chips 4–140 µg/kg Coleslaw 3 µg/kg

6  
 7

### 2.3.6. Biological Monitoring

1 Biological monitoring studies have detected TCE in human blood and urine in the United  
2 States and other countries such as Croatia, China, Switzerland, and Germany (IARC, 1995a).  
3 Concentrations of TCE in persons exposed through occupational degreasing operations were  
4 most likely to have detectable levels (IARC, 1995a). In 1982, eight of eight human breastmilk  
5 samples from four United States urban areas had detectable levels of TCE. The levels of TCE  
6 detected, however, are not specified (ATSDR, 1997b; HSDB, 2002).

7 The Third National Health and Nutrition Examination Survey (NHANES III) examined  
8 TCE concentrations in blood in 677 nonoccupationally exposed individuals. The individuals  
9 were drawn from the general U.S. population and selected on the basis of age, race, gender and  
10 region of residence (Ashley et al., 1994; IARC, 1995a). The samples were collected during  
11 1988–1994. TCE levels in whole blood were below the detection limit of 0.01 µg/L for about  
12 90% of the people sampled (see Table 2-10). Assuming that nondetects equal half of the  
13 detection limit, the mean concentration was about 0.017 µg/L.

14  
15 **Table 2-10. TCE levels in whole blood by population percentile**  
16

Percentiles	10	20	30	40	50	60	70	80	90
Concentration (µg/L)	ND	ND	ND	ND	ND	ND	ND	ND	0.012

17  
18 ND = Nondetect, i.e., below detection limit of 0.01 µg/L.  
19 Data from IARC (1995a) and Ashley et al. (1994).  
20  
21

## 2.4. EXPOSURE PATHWAYS AND LEVELS

### 2.4.1. General Population

22 Because of the pervasiveness of TCE in the environment, most people are likely to have  
23 some exposure via one or more of the following pathways: ingestion of drinking water,  
24 inhalation of outdoor/indoor air, or ingestion of food (ATSDR, 1997b). As noted earlier, the  
25 NHANES survey suggests that about 10% of the population has detectable levels of TCE in  
26 blood. Each pathway is discussed below.  
27

#### 2.4.1.1.1. Inhalation

As discussed earlier, EPA has estimated emissions and modeled air concentrations for the Criteria Pollutants and Hazardous Air Pollutants under the National-Scale Air Toxics Assessment program (U.S. EPA, 2006b). This program has also estimated inhalation exposures on a nationwide basis. The exposure estimates are based on the modeled concentrations from outdoor sources and human activity patterns (Rosenbaum, 2005). Table 2-11 shows the 1999 results for TCE.

These modeled inhalation exposures would have a geographic distribution similar to that of the modeled air concentrations as shown in Figure 2-4. Table 2-11 indicates that TCE inhalation exposures in urban areas are generally about twice as high as rural areas. While these modeling results are useful for understanding the geographic distribution of exposures, they

**Table 2-11. Modeled 1999 annual exposure concentrations ( $\mu\text{g}/\text{m}^3$ ) for trichloroethylene**

Percentile	Exposure concentration ( $\mu\text{g}/\text{m}^3$ )		
	Rural areas	Urban areas	Nationwide
5	0.030	0.048	0.038
10	0.034	0.054	0.043
25	0.038	0.065	0.056
50	0.044	0.086	0.076
75	0.053	0.122	0.113
90	0.070	0.189	0.172
95	0.097	0.295	0.262
Mean	0.058	0.130	0.116

Percentiles and mean are based on census tract values.

Source: <http://www.epa.gov/ttn/atw/nata/ted/exporisk.html#indb>.

appear to underestimate actual exposures. This is based on the fact that, as discussed earlier, the modeled ambient air levels are generally lower than measured values. Also, the modeled exposures do not consider indoor sources. Indoor sources of TCE make the indoor levels higher than ambient levels. This is particularly important to consider since people spend about 90% of their time indoors (U.S. EPA, 1997a). A number of measurement studies were presented earlier that showed higher TCE levels indoors than outdoors. Sexton et al. (2005) measured TCE levels in Minneapolis/St. Paul area and found means of  $0.5 \mu\text{g}/\text{m}^3$  indoors ( $n = 292$ ) and  $1.0 \mu\text{g}/\text{m}^3$

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 based on personal sampling ( $n = 288$ ). Using  $1.0 \mu\text{g}/\text{m}^3$  and an average adult inhalation rate of  
2  $13 \text{ m}^3$  air/day (U.S. EPA, 1997a) yields an estimated intake of  $13 \mu\text{g}/\text{day}$ . This is consistent with  
3 ATSDR (1997b), which reports an average daily air intake for the general population of  
4  $11\text{--}33 \mu\text{g}/\text{day}$ .

#### 2.4.1.1.2. Ingestion

6 The median value from the nationwide survey of domestic and public wells by USGS for  
7 1985–2001 is  $0.15 \mu\text{g}/\text{L}$ . This value was selected for exposure estimation purposes because it  
8 was the most current and most representative of the national population. Using this value and an  
9 average adult water consumption rate of  $1.4 \text{ L}/\text{d}$  yields an estimated intake of  $0.2 \mu\text{g}/\text{day}$ . [This is  
10 from U.S. EPA (1997a), but note that U.S. EPA (2004) indicates a mean per capita daily average  
11 total water ingestion from all sources of  $1.233 \text{ L}$ ]. This is lower than the ATSDR (1997b)  
12 estimate water intake for the general population of  $2\text{--}20 \mu\text{g}/\text{day}$ . The use of the USGS survey to  
13 represent drinking water is uncertain in two ways. First, the USGS survey measured only  
14 groundwater and some drinking water supplies use surface water. Second, the USGS measured  
15 TCE levels at the well head, not the drinking water tap. Further discussion about the possible  
16 extent and magnitude of TCE exposure via drinking water is presented below.

17 According to ATSDR (1997b), TCE is the most frequently reported organic contaminant  
18 in ground water (1997b), and between 9 and 34% of the drinking water supply sources tested in  
19 the United States may have some TCE contamination. Approximately 90% of the 155,000  
20 public drinking water systems<sup>1</sup> in the United States are ground water systems. The drinking  
21 water standard for TCE only applies to community water systems (CWSs) and approximately  
22 78% of the 51,972 CWSs in the United States are ground water systems (U.S. EPA, 2008c).  
23 Although commonly detected in water supplies, the levels are generally low because, as  
24 discussed earlier, MCL violations for TCE in public water supplies are relatively rare for any  
25 extended period (U.S. EPA, 1998). The USGS (2006) survey found that the number of samples  
26 exceeding the MCL ( $5 \mu\text{g}/\text{L}$ ) was six at domestic wells ( $n = 2,400$ ) and nine at public wells ( $n =$   
27  $1,100$ ). Private wells, however, are often not closely monitored and if located near TCE  
28 disposal/contamination sites where leaching occurs, may have undetected contamination levels.  
29 About 10% of Americans (27 million people) obtain water from sources other than public water  
30 systems, primarily private wells (U.S. EPA, 1995). TCE is a common contaminant at Superfund

---

<sup>1</sup> PWSs are defined as systems which provide water for human consumption through pipes or other constructed conveyances to at least 15 service connections or serves an average of at least 25 people for at least 60 days a year. EPA further specifies three types of PWSs, including CWS—a PWS that supplies water to the same population year-round.

1 sites. It has been identified in at least 861 of the 1,428 hazardous waste sites proposed for  
 2 inclusion on the EPA National Priorities List (NPL) (ATSDR, 1997b). Studies have shown that  
 3 many people live near these sites: 41 million people live less than 4 miles from one or more of  
 4 the nation's NPL sites, and on average 3,325 people live within 1 mile of any given NPL site  
 5 (ATSDR, 1996a).

6 Table 2-12 presents preliminary estimates of TCE intake from food. They are based on  
 7 average adult food ingestion rates and food data from Table 2-9. This approach suggests a total  
 8 ingestion intake of about 5 µg/d. It is important to consider this estimate as preliminary because  
 9 it is derived by applying data from very limited food samples to broad classes of food.

10  
 11 **Table 2-12. Preliminary estimates of TCE intake from food ingestion**

	Consumption rate (g/kg-day)	Consumption rate (g/day)	Concentration in food (µg/kg)	Intake (µg/day)
Fruit	3.4	238	2	0.48
Vegetables	4.3	301	3	0.90
Fish		20	10	0.20
Meat	2.1	147	5	0.73
Dairy products	8	560	3	1.68
Grains	4.1	287	3	0.86
Sweets	0.5	35	3	0.10
Total				4.96

13  
 14 <sup>a</sup> Consumption rates are per capita averages from EPA (1997a).

15 <sup>b</sup> Consumption rates in g/d assume 70 kg body weight.

16  
 17  
 18 **2.4.1.1.3. Dermal**

19 TCE in bathing water and consumer products can result in dermal exposure. A modeling  
 20 study has suggested that a significant fraction of the total dose associated with exposure to  
 21 volatile organics in drinking water results from dermal absorption (Brown et al., 1984). EPA  
 22 (2004) used a prediction model based on octanol-water partitioning and molecular weight to  
 23 derive a dermal permeability coefficient for TCE in water of 0.012 cm/hour. EPA used this  
 24 value to compute the dermally absorbed dose from a 35 minute shower and compared it to the  
 dose from drinking 2 L of water at the same concentration. This comparison indicated that the

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 dermal dose would be 17% of the oral dose. Much higher dermal permeabilities were reported  
 2 by Nakai et al. (1999) based on human skin in vitro testing. For dilute aqueous solutions of  
 3 TCE, they measured a permeability coefficient of 0.12 cm/hour (26°C). Nakai et al. (1999) also  
 4 measured a permeability coefficient of 0.018 cm/hour for tetrachloroethylene in water. Poet  
 5 et al. (2000) measured dermal absorption of TCE in humans from both water and soil matrices.  
 6 The absorbed dose was estimated by applying a physiologically based pharmacokinetic model to  
 7 TCE levels in breath. The permeability coefficient was estimated to be 0.015 cm/hour for TCE  
 8 in water and 0.007 cm/hour for TCE in soil (Poet et al., 2000).

9

#### 2.4.1.1.4. Exposure to Trichloroethylene (TCE) Related Compounds

10 Table 2-13 presents adult exposure estimates that have been reported for the TCE related  
 11 compounds. This table was originally compiled by Wu and Schaum (2001). The exposure/dose  
 12 estimates are taken directly from the listed sources or derived based on monitoring data  
 13 presented in the source documents. They are considered “preliminary” because they are  
 14 generally based on very limited monitoring data. These preliminary estimates suggest that  
 15 exposures to most of the TCE related compounds are comparable to or greater than TCE itself.

16  
 17 **Table 2-13. Preliminary intake estimates of TCE and TCE-related chemicals**

Chemical	Population	Media	Range of estimated adult exposures (µg/day)	Range of adult doses (mg/kg-day)	Data sources <sup>a</sup>
Trichloroethylene	General	Air	11–33	1.57E-04–4.71E-04	ATSDR (1997b)
	General	Water	2–20 <sup>b</sup>	2.86E-05–2.86E-04	ATSDR (1997b)
	Occupational	Air	2,232–9,489	3.19E-02–1.36E-01	ATSDR (1997b)
Tetrachloroethylene	General	Air	80–200	1.14E-03–2.86E-03	ATSDR (1997a)
	General	Water	0.1–0.2	1.43E-06–2.86E-06	ATSDR (1997a)
	Occupational	Air	5,897–219,685	8.43E-02–3.14	ATSDR (1997a)
1,1,1-Trichloroethane	General	Air	10.8–108	1.54E-04–1.54E-03	ATSDR (1995)
	General	Water	0.38–4.2	5.5E-06–6.00E-05	ATSDR (1995)
1,2-Dichloroethylene	General	Air	1–6	1.43E-05–8.57E-05	ATSDR (1996b)
	General	Water	2.2	3.14E -05	ATSDR (1996b)
Cis-1,2-Dichloroethylene	General	Air	5.4	7.71E -05	HSDB (1996)
	General	Water	0.5–5.4	7.14E-06–7.71E-05	HSDB (1996)
1,1,1,2-Tetrachloroethane	General	Air	142	2.03E -03	HSDB (2002)

*This document is a draft for review purposes only and does not constitute Agency policy.*

1,1-Dichloroethane	General	Air	4	5.71E -05	ATSDR (1990)
	General	Water	2.47–469.38	3.53E-05–6.71E-03	ATSDR (1990)
Chloral	General	Water	0.02–36.4	2.86E-07–5.20E-04	HSDB (1996)
Monochloroacetic acid	General	Water	2–2.4	2.86E-05–3.43E-05	EPA (1994b)
Dichloroacetic acid	General	Water	10–266	1.43E-04–3.80E-03	IARC (1995a)
Trichloroacetic acid	General	Water	8.56–322	1.22E-03–4.60E-03	IARC (1995a)

<sup>a</sup> Originally compiled in Wu and Schaum (2001).

<sup>b</sup> New data from USGS (2006) suggests much lower water intakes, i.e., 0.2 µg/d.

## 2.4.2. Potentially Highly Exposed Populations

Some members of the general population may have elevated TCE exposures. ATSDR (1997b) has reported that TCE exposures may be elevated for people living near waste facilities where TCE may be released, residents of some urban or industrialized areas, people exposed at work (discussed further below) and individuals using certain products (also discussed further below). Because TCE has been detected in breast milk samples of the general population, infants who ingest breast milk may be exposed, as well. Increased TCE exposure is also a possible concern for bottle-fed infants because they ingest more water on a bodyweight basis than adults (the average water ingestion rate for adults is 21 mL/kg-day and for infants under one year old it is 44 mL/kg-day) (U.S. EPA, 1997a). Also, because TCE can be present in soil, children may be exposed through activities such as playing in or ingesting soil.

### 2.4.2.1.1. Occupational Exposure

Occupational exposure to TCE in the United States has been identified in various degreasing operations, silk screening, taxidermy, and electronics cleaning (IARC, 1995a). The major use of trichloroethylene is for metal cleaning or degreasing (IARC, 1995a). Degreasing is used to remove oils, greases, waxes, tars, and moisture before galvanizing, electroplating, painting, anodizing, and coating. The five primary industries using TCE degreasing are furniture and fixtures; electronic and electric equipment; transport equipment; fabricated metal products; and miscellaneous manufacturing industries (IARC, 1995a). Additionally, TCE is used in the manufacture of plastics, appliances, jewelry, plumbing fixtures, automobile, textiles, paper, and glass (IARC, 1995a).

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 Table 2-14 lists the primary types of industrial degreasing procedures and the years that  
 2 the associated solvents were used. Vapor degreasing has the highest potential for exposure  
 3 because vapors can escape into the work place. Hot dip tanks, where trichloroethylene is heated  
 4 to close to its boiling point of 87°C, are also major sources of vapor that can create exposures as  
 5 high as vapor degreasers. Cold dip tanks have a lower exposure potential, but they have a large  
 6 surface area which enhances volatilization. Small bench-top cleaning operations with a rag or  
 7 brush and open bucket have the lowest exposure potential. In combination with the vapor  
 8 source, the size and ventilation of the workroom are the main determinants of exposure intensity  
 9 (NRC, 2006).

10 Occupational exposure to TCE has been assessed in a number of epidemiologic and  
 11 industrial hygiene studies. Bakke et al. (2007) estimated that the arithmetic mean of TCE  
 12 occupational exposures across all industries and decades (mostly 1950s, 1970s, and 1980s)  
 13 was 38.2 ppm (210 mg/m<sup>3</sup>). They also reported that the highest personal and area air levels were

14 **Table 2-14. Years of solvent use in industrial degreasing and cleaning**  
 15 **operations**  
 16

Years	Vapor degreasers	Cold dip tanks	Rag or brush and bucket on bench top
~1934–1954	Trichloroethylene (poorly controlled)	Stoddard solvent <sup>a</sup>	Stoddard solvent (general use), alcohols (electronics shop), carbon tetrachloride (instrument shop).
~1955–1968	Trichloroethylene (poorly controlled, tightened in 1960s)	Trichloroethylene (replaced some Stoddard solvent)	Stoddard solvent, trichloroethylene (replaced some Stoddard solvent), perchloroethylene, 1,1,1-trichloroethane (replaced carbon tetrachloride, alcohols, ketones).
~1969–1978	Trichloroethylene, (better controlled)	Trichloroethylene, Stoddard solvent	Trichloroethylene, perchloroethylene, 1,1,1-trichloroethane, alcohols, ketones, Stoddard solvent.
~1979–1990s	1,1,1-Trichloroethane (replaced trichloroethylene)	1,1,1-Trichloroethane (replaced trichloroethylene), Stoddard solvent	1,1,1-Trichloroethane, perchloroethylene, alcohols, ketones, Stoddard solvent.

17 <sup>a</sup> A mixture of straight and branched chain paraffins (48%), naphthenes (38%), and aromatic hydrocarbons (14%).  
 18 Source: Stewart and Dosemeci (2005) and Bakke et al. (2007).  
 19  
 20  
 21

22 found in vapor degreasing operations (arithmetic mean of 44.6 ppm or 240 mg/m<sup>3</sup>). Hein et al.  
 23 (2010) developed and evaluated statistical models to estimate the intensity of occupational

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 exposure to trichloroethylene (and other solvents) using a database of air measurement data and  
2 associated exposure determinants. The measurement database was compiled from the published  
3 literature and National Institute for Occupational Safety and Health (NIOSH) reports from 1940–  
4 1998 ( $n = 484$ ) and were split between personal (47%) and area (53%) measurements. The  
5 predicted arithmetic mean exposure intensity levels for the evaluated exposure scenarios ranged  
6 from 0.21–3,700 ppm (1.1–20,000  $\text{mg}/\text{m}^3$ ) with a median of 30 ppm (160  $\text{mg}/\text{m}^3$ ). Landrigan  
7 et al. (1987) used air and biomonitoring techniques to quantify the exposure of degreasing  
8 workers who worked around a heated, open bath of TRI. Exposures were found to be between  
9 22 and 66 ppm (117–357  $\text{mg}/\text{m}^3$ ) on average, with short-term peaks between 76 and 370 ppm  
10 (413–2,000  $\text{mg}/\text{m}^3$ ). High peak exposures have also been reported for cardboard workers who  
11 were involved with degreasing using a heated and open process (Henschler et al., 1995).  
12 Lacking industrial hygiene data and making some assumptions about plant environment and TCE  
13 usage, Cherrie et al. (2001) estimated that cardboard workers at a plant in Germany had peak  
14 exposures in the range of 200–4,000 ppm (1,100–22,000  $\text{mg}/\text{m}^3$ ) and long-term average  
15 exposures of 10–225 ppm (54–1,200  $\text{mg}/\text{m}^3$ ). ATSDR (1997b) reports that the majority of  
16 published worker exposure data show time-weighted average concentrations ranging from  
17 <50 ppm–100 ppm (<270–540  $\text{mg}/\text{m}^3$ ). NIOSH conducted a survey of various industries from  
18 1981–1983 and estimated that approximately 401,000 U.S. employees in 23,225 plants in the  
19 United States were potentially exposed to TCE during this timeframe (ATSDR, 1997b; IARC,  
20 1995a). Occupational exposure to TCE has likely declined since the 1950s and 1960s due to  
21 decreased usage, better release controls, and improvements in worker protection. Reductions in  
22 TCE use are illustrated in Table 2-14, which shows that by about 1980, common degreasing  
23 operations had substituted other solvents for TCE.  
24

#### 2.4.2.1.2. Consumer Exposure

25 Consumer products reported to contain TCE include wood stains, varnishes, and finishes;  
26 lubricants; adhesives; typewriter correction fluids; paint removers; and cleaners (ATSDR,  
27 1997b). Use of TCE has been discontinued in some consumer products (i.e., as an inhalation  
28 anesthetic, fumigant, and an extractant for decaffeinating coffee) (ATSDR, 1997b).  
29

#### 2.4.3. Exposure Standards

30 Table 2-15 summarizes the federal regulations limiting TCE exposure.  
31

*This document is a draft for review purposes only and does not constitute Agency policy.*

## 2.5. EXPOSURE SUMMARY

1 TCE is a volatile compound with moderate water solubility. Most TCE produced today  
 2 is used for metal degreasing. The highest environmental releases are to the air. Ambient air  
 3 monitoring data suggests that levels have remained fairly constant since 1999 at about 0.3 µg/m<sup>3</sup>.  
 4 Indoor levels are commonly three or more times higher than outdoor levels due to releases from  
 5 building materials and consumer products. TCE is among the most common groundwater  
 6 contaminants and the median level based on a large survey by USGS for 1985–2001 is  
 7 0.15 µg/L. It has also been detected in a wide variety of foods in the 1–100 µg/kg range. None  
 8 of the environmental sampling has been done using statistically based national surveys.  
 9 However, a substantial amount of air and groundwater data have been collected allowing  
 10 reasonably well supported estimates of typical daily intakes by the general population:  
 11 inhalation—13 µg/day and water ingestion—0.2 µg/day. The limited food data suggests an  
 12 intake of about 5 µg/day, but this must be considered preliminary.

13 Much higher exposures have occurred to various occupational groups. For example, past  
 14 studies of aircraft workers have shown short term peak exposures in the hundreds of ppm

15 **Table 2-15. TCE standards**

Standard	Value	Reference
OSHA Permissible Exposure Limit: Table Z-2 8-h time-weighted average.	100 ppm (538 mg/m <sup>3</sup> )	29 CFR 1910.1000 (7/1/2000)
OSHA Permissible Exposure Limit: Table Z-2 Acceptable ceiling concentration (this cannot be exceeded for any time period during an 8-h shift except as allowed in the maximum peak standard below).	200 ppm (1,076 mg/m <sup>3</sup> )	29 CFR 1910.1000 (7/1/2000)
OSHA Permissible Exposure Limit: Table Z-2 Acceptable maximum peak above the acceptable ceiling concentration for an 8-h shift. Maximum Duration: 5 minutes in any 2 h.	300 ppm (1,614 mg/m <sup>3</sup> )	29 CFR 1910.1000 (7/1/2000)
MCL under the Safe Drinking Water Act.	5 ppb (5 µg/L)	40 CFR 141.161
FDA Tolerances for decaffeinated ground coffee decaffeinated soluble (instant) coffee extract spice oleoresins.	25 ppm (25 µg/g) 10 ppm (10 µg/g) 30 ppm (30 µg/g)	21 CFR 173.290 (4/1/2000)

18  
 19 OSHA = Occupational Safety and Health Administration.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2  
3 (>540,000  $\mu\text{g}/\text{m}^3$ ) and long term exposures in the low tens of ppm (>54,000  $\mu\text{g}/\text{m}^3$ ).  
4 Occupational exposures have likely decreased in recent years due to better release controls and  
5 improvements in worker protection.

6 Preliminary exposure estimates were presented for a variety of TCE related compounds  
7 which include metabolites of TCE and other parent compounds that produce similar metabolites.  
8 Exposure to the TCE related compounds can alter or enhance TCE's metabolism and toxicity by  
9 generating higher internal metabolite concentrations than would result from TCE exposure by  
10 itself. The preliminary estimates suggest that exposures to most of the TCE related compounds  
11 are comparable to or greater than TCE itself.

1

### 3. TOXICOKINETICS

1  
2  
3 Trichloroethylene (TCE) is a lipophilic compound that readily crosses biological  
4 membranes. Exposures may occur via the oral, dermal, and inhalation route, with evidence for  
5 systemic availability from each route. TCE is rapidly and nearly completely absorbed from the  
6 gut following oral administration, and studies with animals indicate that exposure vehicle may  
7 impact the time-course of absorption: oily vehicles may delay absorption whereas aqueous  
8 vehicles result in a more rapid increase in blood concentrations.

9 Following absorption to the systemic circulation, TCE distributes from blood to solid  
10 tissues by each organ's solubility. This process is mainly determined by the blood:tissue  
11 partition coefficients (PCs), which are largely established by tissue lipid content. Adipose  
12 partitioning is high, adipose tissue may serve as a reservoir for TCE, and accumulation into  
13 adipose tissue may prolong internal exposures. TCE attains high concentrations relative to blood  
14 in the brain, kidney, and liver—all of which are important target organs of toxicity. TCE is  
15 cleared via metabolism mainly in three organs: the kidney, liver, and lungs.

16 The metabolism of TCE is an important determinant of its toxicity. Metabolites are  
17 generally thought to be responsible for toxicity—especially for the liver and kidney. Initially,  
18 TCE may be oxidized via cytochrome P450 (CYP) xenobiotic metabolizing isozymes or  
19 conjugated with glutathione (GSH) by glutathione S-transferase enzymes. While CYP2E1 is  
20 generally accepted to be the CYP form most responsible for TCE oxidation at low  
21 concentrations, others forms may also contribute, though their contributions may be more  
22 important at higher, rather than lower, environmentally-relevant exposures.

23 Once absorbed, TCE is excreted primarily either in breath as unchanged TCE or carbon  
24 dioxide (CO<sub>2</sub>), or in urine as metabolites. Minor routes of elimination include excretion of  
25 metabolites in saliva, sweat, and feces. Following oral administration or upon cessation of  
26 inhalation exposure, exhalation of unmetabolized TCE is a major elimination pathway. Initially,  
27 elimination of TCE upon cessation of inhalation exposure demonstrates a steep  
28 concentration-time profile: TCE is rapidly eliminated in the minutes and hours postexposure, and  
29 then the rate of elimination via exhalation decreases. Following oral or inhalation exposure,  
30 urinary elimination of parent TCE is minimal, with urinary elimination of the metabolites  
31 trichloroacetic acid (TCA) and trichloroethanol (TCOH) accounting for the bulk of the absorbed  
32 dose of TCE.

1 Sections 3.1–3.4 below describe the absorption, distribution, metabolism, and excretion  
2 (ADME) of TCE and its metabolites in greater detail. Section 3.5 then discusses physiologically  
3 based pharmacokinetic(PBPK) modeling of TCE and its metabolites.

### 3.1. ABSORPTION

4 Trichloroethylene is a low-molecular-weight lipophilic solvent; these properties explain  
5 its rapid transfer from environmental media into the systemic circulation after exposure. As  
6 discussed below, it is readily absorbed into the bloodstream following exposure via oral  
7 ingestion and inhalation, with more limited data indicating dermal penetration.  
8

#### 3.1.1. Oral

9 Available reports on human exposure to TCE via the oral route are largely restricted to  
10 case reports of occupational or intentional (suicidal) ingestions and suggest significant gastric  
11 absorption (e.g., Brüning et al., 1998; Perbellini et al., 1991; Yoshida et al., 1996). Clinical  
12 symptoms attributable to TCE or metabolites were observed in these individuals within a few  
13 hours of ingestion (such as lack of consciousness), indicating absorption of TCE. In addition,  
14 TCE and metabolites were measured in blood or urine at the earliest times possible after  
15 ingestion, typically upon hospital admission, while urinary excretion of TCE metabolites was  
16 followed for several days following exposure. Therefore, based on these reports, it is likely that  
17 TCE is readily absorbed in the gastrointestinal tract; however, the degree of absorption cannot be  
18 confidently quantified because the ingested amounts are not known.

19 Experimental evidence in mice and rats supports rapid and extensive absorption of TCE,  
20 although variables such as stomach contents, vehicle, and dose may affect the degree of gastric  
21 absorption. D’Souza et al. (1985) reported on bioavailability and blood kinetics in fasted and  
22 nonfasted male Sprague-Dawley rats following intragastric administration of TCE at 5–25 mg/kg  
23 in 50% polyethylene glycol (PEG 400) in water. TCE rapidly appeared in peripheral blood (at  
24 the initial 0.5 minutes sampling) of fasted and nonfasted rats with peak levels being attained  
25 shortly thereafter (6–10 minutes), suggesting that absorption is not diffusion limited, especially  
26 in fasted animals. The presence of food in the gastro-intestinal (GI) tract, however, seems to  
27 influence TCE absorption based on findings in the nonfasted animals of lesser bioavailability  
28 (60–80% vs. 90% in fasted rats), smaller peak blood levels (two–threefold lower than nonfasted  
29 animals), and a somewhat longer terminal half-life ( $t_{1/2}$ ) (174 vs. 112 minutes in fasted rats).

30 Studies by Prout et al. (1985) and Dekant et al. (1986b) have shown that up to 98% of  
31 administered radiolabel was found in expired air and urine of rats and mice following gavage

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 administration of [<sup>14</sup>C]-radio labeled TCE ([<sup>14</sup>C]TCE). Prout et al. (1985) and Green and Prout  
2 (1985) compared the degree of absorption, metabolites, and routes of elimination among  
3 two strains each of male rats (Osborne-Mendel and Park Wistar) and male mice (B6C3F1 and  
4 Swiss-Webster) following a single oral administration of 10, 500, or 1,000 [<sup>14</sup>C]TCE. Additional  
5 dose groups of Osborne-Mendel male rats and B6C3F1 male mice also received a single oral  
6 dose of 2,000 mg/kg [<sup>14</sup>C]TCE. At the lowest dose of 10 mg/kg, there were no major differences  
7 between rats and mice in routes of excretion with most of the administered radiolabel (nearly  
8 60–70%) being in the urine. At this dose, the expired air from all groups contained 1–4% of  
9 unchanged TCE and 9–14% CO<sub>2</sub>. Fecal elimination of the radiolabel ranged from 8.3% in  
10 Osborne-Mendel rats to 24.1% in Park Wistar rats. However, at doses between 500 and  
11 2,000 mg/kg, the rat progressively excreted a higher proportion of the radiolabel as unchanged  
12 TCE in expired air such that 78% of the administered high dose was found in expired air (as  
13 unchanged TCE) while only 13% was excreted in the urine.

14 Following exposure to a chemical by the oral route, distribution is determined by delivery  
15 to the first organ encountered in the circulatory pathway—the liver (i.e., the first-pass effect),  
16 where metabolism and elimination may limit the proportion that may reach extrahepatic organs.  
17 Lee et al. (1996) evaluated the efficiency and dose-dependency of presystemic elimination of  
18 TCE in male Sprague-Dawley rats following administration into the carotid artery, jugular vein,  
19 hepatic portal vein, or the stomach of TCE (0.17, 0.33, 0.71, 2, 8, 16, or 64 mg/kg) in a  
20 5% aqueous Alkamus emulsion (polyethoxylated vegetable oil) in 0.9% saline. The first-pass  
21 elimination, decreased from 57.5 to <1% with increasing dose (0.17–16 mg/kg) which implied  
22 that hepatic TCE metabolism may be saturated at doses above 16 mg/kg in the male rat. At  
23 doses of 16 mg/kg or higher, hepatic first-pass elimination was almost nonexistent indicating  
24 that, at relatively large doses, virtually all of TCE passes through the liver without being  
25 extracted (Lee et al., 1996). In addition to the hepatic first-pass elimination findings, pulmonary  
26 extraction, which was relatively constant (at nearly 5–8% of dose) over the dose range, also  
27 played a role in eliminating TCE.

28 In addition, oral absorption appears to be affected by both dose and vehicle used. The  
29 majority of oral TCE studies have used either aqueous solution or corn oil as the dosing vehicle.  
30 Most studies that relied on an aqueous vehicle delivered TCE as an emulsified suspension in  
31 Tween 80<sup>®</sup> or PEG 400 in order to circumvent the water solubility problems. Lee et al. (2000a;  
32 2000b) used Alkamuls (a polyethoxylated vegetable oil emulsion) to prepare a 5% aqueous  
33 emulsion of TCE that was administered by gavage to male Sprague-Dawley rats. The findings  
34 confirmed rapid TCE absorption but reported decreasing absorption rate constants (i.e., slower  
35 absorption) with increasing gavage dose (2–432 mg/kg). The time to reach blood peak

1 concentrations increased with dose and ranged between 2 and 26 minutes postdosing. Other  
2 pharmacokinetics data, including area under the blood concentration time curve (AUC) and  
3 prolonged elevation of blood TCE levels at the high doses, indicated prolonged GI absorption  
4 and delayed elimination due to metabolic saturation occurring at the higher TCE doses.

5 A study by Withey et al. (1983) evaluated the effect of dosing TCE with corn oil versus  
6 pure water as a vehicle by administering four volatile organic compounds separately in each  
7 dosing vehicle to male Wistar rats. Based on its limited solubility in pure water, the dose for  
8 TCE was selected at 18 mg/kg (administered in 5 mL/kg). Times to peak in blood reported for  
9 TCE averaged 5.6 minutes when water was used. In comparison, the time to peak in blood was  
10 much longer (approximately 100 minutes) when the oil vehicle was used and the peaks were  
11 smaller, below the level of detection, and not reportable.

12 Time-course studies reporting times to peak in blood or other tissues have been  
13 performed using both vehicles (D'Souza et al., 1985; Dekant et al., 1984; Green and Prout, 1985;  
14 Larson and Bull, 1992a; 1992b; Withey et al., 1983). Related data for other solvents (Chieco et  
15 al., 1981; Dix et al., 1997; Kim et al., 1990b; Lilly et al., 1994) confirmed differences in TCE  
16 absorption and peak height between the two administered vehicles. One study has also evaluated  
17 the absorption of TCE from soil in rats (Kadry et al., 1991) and reported absorption within  
18 16 hours for clay and 24 hours for sandy soil. In summary, these studies confirm that TCE is  
19 relatively quickly absorbed from the stomach, and that absorption is dependent on vehicle used.  
20

### 3.1.2. Inhalation

21 Trichloroethylene is a lipophilic volatile compound that is readily absorbed from inspired  
22 air. Uptake from inhalation is rapid and the absorbed dose is proportional to exposure  
23 concentration and duration, and pulmonary ventilation rate. Distribution into the body via  
24 arterial blood leaving the lungs is determined by the net dose absorbed and eliminated by  
25 metabolism in the lungs. Metabolic clearance in the lungs will be further discussed in  
26 Section 3.3, below. In addition to metabolism, solubility in blood is the major determinant of the  
27 TCE concentration in blood entering the heart and being distributed to the each body organ via  
28 the arterial blood. The measure of TCE solubility in each organ is the partition coefficient, or the  
29 concentration ratio between both organ phases of interest. The blood-to-air partition coefficient  
30 quantifies the resulting concentration in blood leaving the lungs at equilibrium with alveolar air.  
31 The value of the blood-to-air partition coefficient is used in PBPK modeling (see Section 3.5).  
32 The blood-to-air partition has been measured in vitro using the same principles in different



1 studies and found to range between 8.1–11.7 in humans and somewhat higher values in mice and  
2 rats (13.3–25.8) (see Tables 3-1–3-2, and references therein).

3 TCE enters the human body quickly by inhalation, and, at high concentrations, it may  
4 lead to death (Coopman et al., 2003), narcosis, unconsciousness, and

5

1  
2

**Table 3-1. Blood:air PC values for humans**

<b>Blood:air partition coefficient</b>	<b>Reference/notes</b>
8.1 ± 1.8	Fiserova-Bergerova et al. (1984); mean ± SD (SD converted from SE based on <i>n</i> = 5)
8.11	Gargas et al. (1989); ( <i>n</i> = 3–15)
9.13 ± 1.73 [6.47–11]	Fisher et al. (1998); mean ± SD [range] of females ( <i>n</i> = 6)
9.5	Sato and Nakajima (1979); ( <i>n</i> = 1)
9.77	Koizumi (1989)
9.92	Sato et al. (1977); ( <i>n</i> = 1)
11.15 ± 0.74 [10.1–12.1]	Fisher et al. (1998); mean ± SD [range] of males ( <i>n</i> = 7)
11.2 ± 1.8 [7.9–15]	Mahle et al. (2007); mean ± SD; 20 male pediatric patients aged 3–7 years (range; USAF, 2004)
11.0 ± 1.6 [6.6–13.5]	Mahle et al. (2007); mean ± SD; 18 female pediatric patients aged 3–17years (range; USAF, 2004)
11.7 ± 1.9 [6.7–16.8]	Mahle et al. (2007); mean ± SD; 32 male patients aged 23–82 years (range; USAF, 2004)
10.6 ± 2.3 [3–14.4]	Mahle et al., (2007); mean ± SD; 27 female patients aged 23–82 years (range; USAF, 2004)

3  
4  
5

SD = standard deviation, SE = standard error.

1  
2

**Table 3-2. Blood:air PC values for rats and mice**

<b>Blood:air partition coefficient</b>	<b>Reference/notes</b>
Rat	
15 ± 0.5	Fisher et al. (1998); mean ± SD (SD converted from SE based on <i>n</i> = 3)
17.5	Rodriguez et al. (2007)
20.5 ± 2.4	Barton et al. (1995); mean ± SD (SD converted from SE based on <i>n</i> = 4)
20.69 ± 3.3	Simmons et al. (2002); mean ± SD ( <i>n</i> = 7–10)
21.9	Gargas et al. (1989) ( <i>n</i> = 3–15)
25.8	Koizumi (1989) (pooled <i>n</i> = 3)
25.82 ± 1.7	Sato et al. (1977); mean ± SD ( <i>n</i> = 5)
13.3 ± 0.8 [11.6–15]	Mahle et al. (2007); mean ± SD; 10 PND 10 male rat pups (range; USAF, 2004)
13.4 ± 1.8 [11.8–17.2]	Mahle et al. (2007); mean ± SD; 10 PND 10 female rat pups (range; USAF, 2004)
17.5 ± 3.6 [11.7–23.1]	Mahle et al. (2007); mean ± SD; 9 adult male rats (range; USAF, 2004)
21.8 ± 1.9 [16.9–23.5]	Mahle et al. (2007); mean ± SD; 11 aged male rats (range; USAF, 2004)
Mouse	
13.4	Fisher et al. (1991); male
14.3	Fisher et al. (1991); female
15.91	Abbas and Fisher (1997)

3  
4  
5  
6  
7  
8  
9  
10  
11  
12

SD = standard deviation, SE = standard error, PND = postnatal day.

acute kidney damage (Carrieri et al., 2007). Controlled exposure studies in humans have shown absorption of TCE to approach a steady state within a few hours after the start of inhalation exposure (Fernandez et al., 1977; Monster et al., 1976; Vesterberg and Astrand, 1976; Vesterberg et al., 1976). Several studies have calculated the net dose absorbed by measuring the difference between the inhaled concentration and the exhaled air concentration. Soucek and Vlachova (1960) reported between 58–70% absorption of the amount inhaled for 5-hour

1 exposures between 93–158 ppm. Bartonicek (1962) obtained an average retention value of 58%  
2 after 5 hours of exposure to 186 ppm. Monster et al. (1976) also took into account minute  
3 ventilation measured for each exposure, and calculated between 37–49% absorption in subjects  
4 exposed to 70 and 140 ppm. The impact of exercise, the increase in workload, and its effect on  
5 breathing has also been measured in controlled inhalation exposures. Astrand and Ovrum (1976)  
6 reported 50–58% uptake at rest and 25–46% uptake during exercise from exposure at 100 or  
7 200 ppm (540 or 1,080 mg/m<sup>3</sup>, respectively) of TCE for 30 minutes (see Table 3-3). These  
8 authors also monitored heart rate and pulmonary ventilation. In contrast, Jakubowski and  
9 Wieczorek (1988) calculated about 40% retention in their human volunteers exposed to TCE at 9  
10 ppm (mean inspired concentration of 48–49 mg/m<sup>3</sup>) for 2 hours at rest, with no change in  
11 retention during increase in workload due to exercise (see Table 3-4).

12 Environmental or occupational settings may results from a pattern of repeated exposure  
13 to TCE. Monster et al. (1979) reported 70-ppm TCE exposures in volunteers for 4 hours for  
14 5 consecutive days, averaging a total uptake of 450 mg per 4 hours exposure (see Table 3-5). In  
15 dry-cleaning workers, Skender et al. (1991) reported initial blood concentrations of 0.38 µmol/L,  
16 increasing to 3.4 µmol/L 2 days after. Results of these studies support rapid absorption of TCE  
17 via inhalation.

18 Direct measurement of retention after inhalation exposure in rodents is more difficult  
19 because exhaled breath concentrations are challenging to obtain. The only available data are  
20 from Dallas et al. (1991), who designed a nose-only exposure system for rats using a facemask  
21 equipped with one-way breathing valves to obtain measurements of TCE in inspired and exhaled  
22 air. In addition, indwelling carotid artery cannulae were surgically implanted to facilitate the  
23 simultaneous collection of blood. After a 1-hour acclimatization period, rats were exposed to  
24 50- or 500-ppm TCE for 2 hours and the time course of TCE in blood and expired air was  
25 measured during and for 3 hours following exposure. When air concentration data were  
26 analyzed to reveal absorbed dose (minute volume multiplied by the concentration difference  
27 between inspired and exhaled breath), it was demonstrated that the fractional absorption of either  
28 concentration was more than 90% during the initial 5 minutes of exposure. Fractional absorption  
29 then decreased to 69 and 71% for the 50 and 500-ppm groups during the second hour of  
30 exposure. Cumulative uptake appeared linear with respect to time over the 2-hour exposure,  
31 resulting in absorbed doses of 8.4 mg/kg and 73.3 mg/kg in rats exposed to 50 and 500 ppm,  
32 respectively. Given the 10-fold difference in inspired concentration and the 8.7-fold difference  
33 in uptake, the authors interpreted this information to indicate that metabolic saturation occurred  
34 at some concentration below 500 ppm. In comparing the absorbed doses to those developed for

1  
2  
3

**Table 3-3. Air and blood concentrations during exposure to TCE in humans (Astrand and Ovrum, 1976)**

TCE conc. (mg/m <sup>3</sup> )	Work load (watt)	Exposure series	TCE concentration in			Uptake as % of amount available	Amount taken up (mg)
			Alveolar air (mg/m <sup>3</sup> )	Arterial blood (mg/kg)	Venous blood (mg/kg)		
540	0	I	124 ± 9	1.1 ± 0.1	0.6 ± 0.1	53 ± 2	79 ± 4
540	0	II	127 ± 11	1.3 ± 0.1	0.5 ± 0.1	52 ± 2	81 ± 7
540	50	I	245 ± 12	2.7 ± 0.2	1.7 ± 0.4	40 ± 2	160 ± 5
540	50	II	218 ± 7	2.8 ± 0.1	1.8 ± 0.3	46 ± 1	179 ± 2
540	50	II	234 ± 12	3.1 ± 0.3	2.2 ± 0.4	39 ± 2	157 ± 2
540	50	II	244 ± 16	3.3 ± 0.3	2.2 ± 0.4	37 ± 2	147 ± 9
1,080	0	I	280 ± 18	2.6 ± 0.0	1.4 ± 0.3	50 ± 2	156 ± 9
1,080	0	III	212 ± 7	2.1 ± 0.2	1.2 ± 0.1	58 ± 2	186 ± 7
1,080	50	I	459 ± 44	6.0 ± 0.2	3.3 ± 0.8	45 ± 2	702 ± 31
1,080	50	III	407 ± 30	5.2 ± 0.5	2.9 ± 0.7	51 ± 3	378 ± 18
1,080	100	III	542 ± 33	7.5 ± 0.7	4.8 ± 1.1	36 ± 3	418 ± 39
1,080	150	III	651 ± 53	9.0 ± 1.0	7.4 ± 1.1	25 ± 5	419 ± 84

4  
5  
6  
7  
8

Series I consisted of 30-minute exposure periods of rest, rest, 50W and 50W; Series II consisted of 30-minute exposure periods of rest, 50W, 50W, 50W; Series III consisted of 30-minute exposure periods of rest, 50W, 100W, 150W.

1 **Table 3-4. Retention of inhaled TCE vapor in humans (Jakubowski and**  
 2 **Wieczorek, 1988)**  
 3

<b>Workload</b>	<b>Inspired concentration (mg/m<sup>3</sup>)</b>	<b>Pulmonary ventilation (m<sup>3</sup>/hour)</b>	<b>Retention</b>	<b>Uptake (mg/hour)</b>
Rest	48 ± 3 <sup>a</sup>	0.65 ± 0.07	0.40 ± 0.05	12 ± 1.1
25 W	49 ± 1.3	1.30 ± 0.14	0.40 ± 0.05	25 ± 2.9
50 W	49 ± 1.6	1.53 ± 0.13	0.42 ± 0.06	31 ± 2.8
75 W	48 ± 1.9	1.87 ± 0.14	0.41 ± 0.06	37 ± 4.8

4  
 5 <sup>a</sup> Mean ± standard deviation, n = 6 adult males.  
 6

7 W = watts.  
 8  
 9

10 **Table 3-5. Uptake of TCE in human volunteers following 4 hour exposure to**  
 11 **70 ppm (Monster et al., 1979)**  
 12

	<b>BW (kg)</b>	<b>MV (L/min)</b>	<b>% Retained</b>	<b>Uptake (mg/day)</b>	<b>Uptake (mg/kg-day)</b>
A	80	9.8 ± 0.4	45 ± 0.8	404 ± 23	5.1
B	82	12.0 ± 0.7	44 ± 0.9	485 ± 35	5.9
C	82	10.9 ± 0.8	49 ± 1.2	493 ± 28	6.0
D	67	11.8 ± 0.8	35 ± 2.6	385 ± 38	5.7
E	90	11.0 ± 0.7	46 ± 1.1	481 ± 25	5.3
Mean					5.6 ± 0.4

13  
 14 MV = minute-volume.  
 15 BW = body weight.  
 16  
 17

18 the 70-ppm-exposed human (see Monster et al., 1979), Dallas et al. (1991) concluded that on a  
 19 systemic dose (mg/kg) basis, rats receive a much higher TCE dose from a given  
 20 inhalation exposure than do humans. In particular, using the results cited above, the absorption  
 21 per ppm-hour was 0.084 and 0.073 mg/kg-ppm-hour at 50 and 500 ppm in rats (Dallas et al.,  
 22 1991) and 0.019 mg/kg-ppm-hour at 70 ppm in humans (Monster et al., 1979)—a difference of  
 23 around fourfold. However, rats have about a 10-fold higher alveolar ventilation rate per unit  
 24 body weight (BW) than humans (Brown et al., 1997), which more than accounts for the observed  
 25 increase in absorption.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Other experiments, such as closed-chamber gas uptake experiments or blood  
2 concentration measurements following open-chamber (fixed concentration) experiments,  
3 measure absorption indirectly but are consistent with significant retention. Closed-chamber  
4 gas-uptake methods (Gargas et al., 1988) place laboratory animals or in vitro preparations into  
5 sealed systems in which a known amount of TCE is injected to produce a predetermined  
6 chamber concentration. As the animal retains a quantity of TCE inside its body, due to  
7 metabolism, the closed-chamber concentration decreases with time when compared to the start of  
8 exposure. Many different studies have made use of this technique in both rats and mice to  
9 calculate total TCE metabolism (i.e., Andersen et al., 1987a; Fisher et al., 1991; Simmons et al.,  
10 2002). This inhalation technique is combined with PBPK modeling to calculate metabolic  
11 parameters, and the results of these studies are consistent with rapid absorption of TCE via the  
12 respiratory tract. Figure 3-1 shows an example from Simmons et al. (2002), in Long Evans rats,  
13 that demonstrates an immediate decline in chamber concentrations of TCE indicating absorption,  
14 with multiple initial concentrations needed for each metabolic calculation. At concentrations  
15 below metabolic saturation, a secondary phase of uptake appears, after 1 hour from starting the  
16 exposure, indicative of metabolism. At concentrations above 1,000 ppm, metabolism appears  
17 saturated, with time course curves having a flat phase after absorption. At intermediate  
18 concentrations, between 100–1,000 ppm, the secondary phase of uptake appears after  
19 distribution as continued decreases in chamber concentration as metabolism proceeds. Using a  
20 combination of experiments that include both metabolic linear decline and saturation obtained by  
21 using different initial concentrations, both components of metabolism can be estimated from the  
22 gas uptake curves, as shown in Figure 3-1.


23 Several other studies in humans and rodents have measured blood concentrations of TCE  
24 or metabolites and urinary excretion of metabolites during and after inhalation exposure (e.g.,  
25 Filser and Bolt, 1979; 1991; Fisher et al., 1998; 1990). While qualitatively indicative of  
26 absorption, blood concentrations are also determined by metabolism, distribution, and excretion,  
27 so comparisons between species may reflect similarities or differences in any of the absorption,  
28 distribution, metabolism, and excretion processes.

29

### 3.1.3. Dermal

30 Skin membrane is believed to present a diffusional barrier for entrance of the chemical into the  
31 body, and TCE absorption can be quantified using a permeability rate or permeability constant,  
32 though not all studies performed such a calculation. Absorption through the skin has been shown

1 to be rapid by both vapor and liquid TCE contact with the skin. Human dermal absorption of  
2 TCE vapors was investigated by Kezic et al. (2000). Human volunteers were exposed to  
3



4  
5 **Figure 3-1. Gas uptake data from closed-chamber exposure of rats to TCE.**  
6 Symbols represent measured chamber concentrations. Source: Simmons et al.  
7 (2002).  
8  
9

10  $3.18 \times 10^4$  ppm around each enclosed arm for 20 minutes. Adsorption was found to be rapid  
11 (within 5 minutes), reaching a peak in exhaled breath around 30 minutes, with a calculated  
12 dermal penetration rate averaging 0.049 cm/hour for TCE vapors.

13 With respect to dermal penetration of liquid TCE, Nakai et al. (1999) used surgically  
14 removed skin samples exposed to TCE in aqueous solution in a chamber designed to measure the  
15 difference between incoming and outgoing [ $^{14}\text{C}$ ]TCE. The in vitro permeability constant  
16 calculated by these researchers averaged 0.12 cm/hour. In vivo, Sato and Nakajima (1978)  
17 exposed adult male volunteers dermally to liquid TCE for 30 minutes, with exhaled TCE  
18 appearing at the initial sampling time of 5 minutes after start of exposure, with a maximum  
19 observed at 15 minutes. In Kezic et al. (2001), human volunteers were exposed dermally for  
20 3 minutes to neat liquid TCE, with TCE detected in exhaled breath at the first sampling point of



1 3 minutes, and maximal concentrations observed at 5 minutes. Skin irritancy was reported in all  
2 subjects, which may have increased absorption. A dermal flux of  $430 \pm 295$  (mean  $\pm$  standard  
3 error [SE]) nmol/cm<sup>2</sup>/minute was reported in these subjects, suggesting high interindividual  
4 variability.

5 Another species where dermal absorption for TCE has been reported is in guinea pigs.  
6 Jakobson et al. (1982) applied liquid TCE to the shaved backs of guinea pigs and reported peak  
7 blood TCE levels at 20 minutes after initiation of exposure. Bogen et al. (1992) estimated  
8 permeability constants for dermal absorption of TCE in hairless guinea pigs between  
9 0.16–0.47 mL/cm<sup>2</sup>/hour across a range of concentrations (19–100,000 ppm).

10

### 3.2. DISTRIBUTION AND BODY BURDEN

11 TCE crosses biological membranes and quickly results in rapid systemic distribution to  
12 tissues—regardless of the route of exposure. In humans, in vivo studies of tissue distribution are  
13 limited to tissues taken from autopsies following accidental poisonings or from surgical patients  
14 exposed environmentally, so the level of exposure is typically unknown. Tissue levels reported  
15 after autopsy show wide systemic distribution across all tested tissues, including the brain,  
16 muscle, heart, kidney, lung, and liver (Coopman et al., 2003; De Baere et al., 1997; Dehon et al.,  
17 2000; Ford et al., 1995). However, the reported levels themselves are difficult to interpret  
18 because of the high exposures and differences in sampling protocols. In addition, human  
19 populations exposed environmentally show detectable levels of TCE across different tissues,  
20 including the liver, brain, kidney, and adipose tissues (Kroneld, 1989; McConnell et al., 1975;  
21 Pellizzari et al., 1982).

22 In addition, TCE vapors have been shown to cross the human placenta during childbirth  
23 (Laham, 1970), with experiments in rats confirming this finding (Withey and Karpinski, 1985).  
24 In particular, Laham (1970) reported determinations of TCE concentrations in maternal and fetal  
25 blood following administration of TCE vapors (concentration unreported) intermittently and at  
26 birth (see Table 3-6). TCE was present in all samples of fetal blood, with ratios of  
27 concentrations in fetal:maternal blood ranging from approximately 0.5 to approximately 2. The  
28 concentration ratio was less than 1.0 in six pairs, greater than one in three pairs, and  
29 approximately one in one pair; in general, higher ratios were observed at maternal concentrations  
30 below 2.25 mg/100 mL. Because no details of exposure concentration, duration, or time  
31 postexposure were given for samples taken, these results are not suitable for use in PBPK  
32 modeling, but they do demonstrate the placental transfer of TCE in humans. Withey and  
33 Karpinski (1985) exposed pregnant rats to TCE vapors (302, 1,040, 1,559, or 2,088 ppm for

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 5 hours) on gestation Day 17 and concentrations of TCE in maternal and fetal blood were  
2 determined. At all concentrations, TCE concentration in fetal blood was approximately one-third  
3 the concentration in corresponding maternal blood. Maternal blood concentrations approximated  
4

1 Table 3-6. Concentrations of TCE in maternal and fetal blood at birth  
2

TCE concentration in blood (mg/100 mL)		Ratio of concentrations fetal:maternal
Maternal	Fetal	
4.6	2.4	0.52
3.8	2.2	0.58
8	5	0.63
5.4	3.6	0.67
7.6	5.2	0.68
3.8	3.3	0.87
2	1.9	0.95
2.25	3	1.33
0.67	1	1.49
1.05	2	1.90

3  
4 Source: Laham (1970).  
5  
6

7 15, 60, 80, and 110 µg/gram blood. When the position along the uterine horn was examined,  
8 TCE concentrations in fetal blood decreased toward the tip of the uterine horn. TCE appears to  
9 also distribute to mammary tissues and is excreted in milk. Pellizzari et al. (1982) conducted a  
10 survey of environmental contaminants in human milk using samples from cities in the  
11 northeastern region of the United States and one in the southern region. No details of times  
12 postpartum, milk lipid content, or TCE concentration in milk or blood are reported, but TCE was  
13 detected in 8 milk samples taken from 42 lactating women. Fisher et al. (1990) exposed lactating  
14 rats to 600-ppm TCE for 4 hours and collected milk immediately following the cessation of  
15 exposure. TCE was clearly detectable in milk, and, from a visual interpretation of the graphic  
16 display of their results, concentrations of TCE in milk approximated 110 µg/mL milk.

17 In rodents, detailed tissue distribution experiments have been performed using different  
18 routes of administration (Abbas and Fisher, 1997; Greenberg et al., 1999; Keys et al., 2003;  
19 Pfaffenberger et al., 1980; Savolainen et al., 1977; Simmons et al., 2002). Savolainen et al.  
20 (1977) exposed adult male rats to 200-ppm TCE for 6 hours/day for a total of 5 days.  
21 Concentrations of TCE in the blood, brain, liver, lung, and perirenal fat were measured 17 hours  
22 after cessation of exposure on the fourth day and after 2, 3, 4, and 6 hours of exposure on the

1 fifth day (see Table 3-7). TCE appeared to be rapidly absorbed into blood and distributed to  
2 brain, liver, lungs, and perirenal fat. TCE concentrations in these tissues reached near-maximal  
3 values within 2 hours of initiation of exposure on the fifth day. Pfaffenberger et al. (1980) dosed  
4 rats by gavage with 1 or 10 mg TCE/kg/day in corn oil for 25 days to evaluate the distribution  
5 from serum to adipose tissue. During the exposure period, concentrations of TCE in serum were  
6 below the limit of detection (1 µg/L) and were 280 and 20,000 ng per gram of fat in the 1 and  
7 10 mg/day dose groups, respectively. Abbas and Fisher (1997) and Greenberg et al. (1999)  
8 measured tissue concentrations in the liver, lung, kidney, and fat of mice administered TCE by  
9 gavage (300–2,000 mg/kg) and by inhalation exposure (100 or 600 ppm for 4 hours). In a study  
10 to investigate the effects of TCE on neurological function, Simmons et al. (2002) conducted  
11 pharmacokinetic experiments in rats exposed to 200, 2,000, or 4,000 ppm TCE vapors for 1 hour.  
12 Time-course data were collected on blood, liver, brain, and fat. The data were used to develop a  
13 PBPK model to explore the relationship between internal dose and neurological effect. Keys  
14 et al. (2003), exposed groups of rats to TCE vapors of 50 or 500 ppm for 2 hours and sacrificed  
15 at different time points during exposure. In addition to inhalation, this study also includes oral  
16 gavage and intra-arterial dosing, with the following time course measured: liver, fat, muscle,  
17 blood, GI, brain, kidney, heart, lung, and spleen. These pharmacokinetic data were presented  
18 with an updated PBPK model for all routes.

19 Besides the route of administration, another important factor contributing to body  
20 distribution is the individual solubility of the chemical in each organ, as measured by a partition  
21 coefficient. For volatile compounds, partition coefficients are measured in vitro using the vial  
22 equilibration technique to determine the ratio of concentrations between organ and air at  
23 equilibrium. Table 3-8 reports values developed by several investigators from mouse, rat, and  
24 human tissues. In humans, partition coefficients in the following tissues have been measured:  
25 brain, fat, kidney, liver, lung, and muscle; but the organ having the highest TCE partition  
26 coefficient is fat (63–70), while the lowest is the lung (0.5–1.7). The adipose tissue also has the  
27 highest measured value in rodents, and is one of the considerations needed to be accounted for  
28 when extrapolating across species. However, the rat adipose partition coefficient value is  
29 smaller (23–36), when compared to humans, that is, TCE is less lipophilic in rats than humans.  
30 For the mouse, the measured fat partition coefficient averages 36, ranging between rats and  
31 humans. The value of the partition coefficient plays a role in distribution for each organ and is  
32 computationally described in computer simulations using a PBPK model. Due to its high  
33 lipophilicity in fat, as compared to blood, the adipose tissue behaves as a storage compartment

1  
2  
3  
  
4  
5  
6  
7  
8  
9

**Table 3-7. Distribution of TCE to rat tissues<sup>a</sup> following inhalation exposure (Savolainen et al., 1977)**

Exposure on 5 <sup>th</sup> day	Tissue (concentration in nmol/gram tissue)					
	Cerebrum	Cerebellum	Lung	Liver	Perirenal fat	Blood
0 <sup>b</sup>	0	0	0.08	0.04	0.23 ± 0.09	0.35 ± 0.1
2	9.9 ± 2.7	11.7 ± 4.2	4.9 ± 0.3	3.6	65.9 ± 1.2	7.5 ± 1.6
3	7.3 ± 2.2	8.8 ± 2.1	5.5 ± 1.4	5.5 ± 1.7	69.3 ± 3.3	6.6 ± 0.9
4	7.2 ± 1.7	7.6 ± 0.5	5.8 ± 1.1	2.5 ± 1.4	69.5 ± 6.3	6.0 ± 0.2
6	7.4 ± 2.1	9.5 ± 2.5	5.6 ± 0.5	2.4 ± 0.2	75.4 ± 14.9	6.8 ± 1.2

<sup>a</sup>Data presented as mean of two determinations ± range.  
<sup>b</sup>Sample taken 17 hours following cessation of exposure on Day 4.

1  
2

**Table 3-8. Tissue:blood partition coefficient values for TCE**

Species / tissue	TCE partition coefficient		References
	Tissue:blood	Tissue:air	
Human			
Brain	2.62	21.2	Fiserova-Bergerova et al. (1984)
Fat	63.8–70.2	583–674.4	Sato et al. (1977); Fiserova-Bergerova et al. (1984); Fisher et al. (1998)
Kidney	1.3–1.8	12–14.7	Fiserova-Bergerova et al. (1984); Fisher et al. (1998)
Liver	3.6–5.9	29.4–54	Fiserova-Bergerova et al. (1984); Fisher et al. (1998)
Lung	0.48–1.7	4.4–13.6	Fiserova-Bergerova et al. (1984); Fisher et al. (1998)
Muscle	1.7–2.4	15.3–19.2	Fiserova-Bergerova et al. (1984); Fisher et al. (1998)
Rat			
Brain	0.71–1.29	14.6–33.3	Sato et al. (1977); Simmons et al. (2002); Rodriguez et al. (2007)
Fat	22.7–36.1	447–661	Gargas et al. (1989); Sato et al. (1977); Simmons et al. (2002); Rodriguez et al. (2007); Fisher et al. (1989); Koizumi (1989); Barton et al. (1995)
Heart	1.1	28.4	Sato et al. (1977)
Kidney	1.0–1.55	17.7–40	Sato et al., (1977); Barton et al., (1995); Rodriguez et al., (2007)
Liver	1.03–2.43	20.5–62.7	Gargas et al. (1989); Sato et al. (1977); Simmons et al. (2002); Rodriguez et al. (2007); Fisher et al. (1989); Koizumi, (1989); Barton et al. (1995)
Lung	1.03	26.6	Sato et al. (1977)
Muscle	0.46–0.84	6.9–21.6	Gargas et al. (1989); Sato et al. (1977); Simmons et al. (2002); Rodriguez et al. (2007); Fisher et al. (1989); Koizumi, (1989); Barton et al. (1995)
Spleen	1.15	29.7	Sato et al. (1977)
Testis	0.71	18.3	Sato et al. (1977)
Milk	7.10	N.R.	Fisher et al. (1990)
Mouse			
Fat	36.4	578.8	Abbas and Fisher (1997)

*This document is a draft for review purposes only and does not constitute Agency policy.*

Kidney	2.1	32.9	Abbas and Fisher (1997)
Liver	1.62	23.2	Fisher et al. (1991)
Lung	2.6	41.5	Abbas and Fisher (1997)
Muscle	2.36	37.5	Abbas and Fisher (1997)

1 N.R. = not reported.

2

1 for this chemical, affecting the slower component of the chemical's distribution. For example  
2 Monster et al. (1979) reported that, following repeated inhalation exposures to TCE, TCE  
3 concentrations in expired breath postexposure were highest for the subject with the greatest  
4 amount of adipose tissue (adipose tissue mass ranged 3.5-fold among subjects). The intersubject  
5 range in TCE concentration in exhaled breath increased from approximately twofold at 20 hours  
6 to approximately 10-fold 140 hours postexposure. Notably, they reported that this difference  
7 was not due to differences in uptake, as body weight and lean body mass were most closely  
8 associated with TCE retention. Thus, adipose tissue may play an important role in postexposure  
9 distribution, but does not affect its rapid absorption.

10 Mahle et al. (2007) reported age-dependent differences in partition coefficients in rats,  
11 (see Table 3-9) that can have implications as to life-stage-dependent differences in tissue TCE  
12 distribution. To investigate the potential impact of these differences, Rodriguez et al. (2007)  
13 developed models for the postnatal Day 10 rat pup; the adult and the aged rat, including  
14 age-specific tissue volumes and blood flows; and age-scaled metabolic constants. The models  
15 predict similar uptake profiles for the adult and the aged rat during a 6-hour exposure to  
16 500 ppm; uptake by the postnatal day (PND) 10 rat was higher (see Table 3-10). The effect was  
17 heavily dependent on age-dependent changes in anatomical and physiological parameters  
18 (alveolar ventilation rates and metabolic rates); age-dependent differences in partition coefficient  
19 values had minimal impact on predicted differences in uptake.

20 Finally, TCE binding to tissues or cellular components within tissues can affect overall  
21 pharmacokinetics. The binding of a chemical to plasma proteins, for example, affects the  
22 availability of the chemical to other organs and the calculation of the total half-life. However,  
23 most studies have evaluated binding using [<sup>14</sup>C]TCE, from which one cannot distinguish  
24 covalent binding of TCE from that of TCE metabolites. Nonetheless, several studies have  
25 demonstrated binding of TCE-derived radiolabel to cellular components (Mazzullo et al., 1992;  
26 Moslen et al., 1977). Bolt and Filser (1977) examined the total amount irreversibly bound to  
27 tissues following 9-, 100-, and 1,000-ppm exposures via inhalation in closed-chambers. The  
28 largest percent of in vivo radioactivity taken up occurred in the liver; albumin is the protein  
29 favored for binding (see Table 3-11). Banerjee and van Duuren (1978) evaluated the in vitro  
30 binding of TCE to microsomal proteins from the liver, lung, kidney, and stomachs in rats and  
31 mice. In both rats and mice, radioactivity was similar in stomach and lung, but about 30% lower  
32 in kidney and liver.

33 Based on studies of the effects of metabolizing enzyme induction on binding, there is  
34 some evidence that a major contributor to the observed binding is from TCE metabolites rather  
35 than from TCE itself. Dekant et al. (1986b) studied the effect of enzyme modulation on the



**Table 3-9. Age-dependence of tissue:air partition coefficients in rats**

Age	Liver	Kidney	Fat	Muscle	Brain
PND10 male	22.1 ± 2.3	15.2 ± 1.3	398.7 ± 89.2	43.9 ± 11.0	11.0 ± 0.6
PND10 female	21.2 ± 1.7	15.0 ± 1.1	424.5 ± 67.5	48.6 ± 17.3	11.6 ± 1.2
Adult male	20.5 ± 4.0	17.6 ± 3.9 <sup>a</sup>	631.4 ± 43.1 <sup>a</sup>	12.6 ± 4.3	17.4 ± 2.6
Aged male	34.8 ± 8.7 <sup>a,b</sup>	19.9 ± 3.4 <sup>a</sup>	757.5 ± 48.3 <sup>a,b</sup>	26.4 ± 10.3 <sup>a,b</sup>	25.0 ± 2.0 <sup>a,b</sup>

<sup>a</sup> Statistically significant ( $p \leq 0.05$ ) difference between either the adult or aged partition coefficient and the PND10 male partition coefficient.

<sup>b</sup> Statistically significant ( $p \leq 0.05$ ) difference between aged and adult partition coefficient.

Data are mean ± standard deviation;  $n = 10$ , adult male and pooled male and female litters; 11, aged males. Source: Mahle et al. (2007).

**Table 3-10. Predicted maximal concentrations of TCE in rat blood following a 6-hour inhalation exposure (Rodriguez et al., 2007)**

Age	Exposure concentration					
	50 ppm			500 ppm		
	Predicted peak concentration (mg/L) in: <sup>a</sup>		Predicted time to reach 90% of steady state (hour) <sup>b</sup>	Predicted peak concentration (mg/L) in: <sup>a</sup>		Predicted time to reach 90% of steady state (hour) <sup>b</sup>
	Venous blood	Brain		Venous blood	Brain	
PND 10	3.0	2.6	4.1	33	28	4.2
Adult	0.8	1.0	3.5	22	23	11.9
Aged	0.8	1.2	6.7	21	26	23.3

<sup>a</sup> During a 6 hour exposure.

<sup>b</sup> Under continuous exposure.

1 **Table 3-11. Tissue distribution of TCE metabolites following inhalation**  
 2 **exposure**  
 3

Tissue <sup>a</sup>	Percent of radioactivity taken up/g tissue					
	TCE = 9 ppm, n = 4		TCE = 100 ppm, n = 4		TCE = 1,000 ppm, n = 3	
	Total metabolites	Irreversibly bound	Total metabolites	Irreversibly bound	Total metabolites	Irreversibly bound
Lung	0.23 ± 0.026	0.06 ± 0.002	0.24 ± 0.025	0.06 ± 0.006	0.22 ± 0.055	0.1 ± 0.003
Liver	0.77 ± 0.059	0.28 ± 0.027	0.68 ± 0.073	0.27 ± 0.019	0.88 ± 0.046	0.48 ± 0.020
Spleen	0.14 ± 0.015	0.05 ± 0.002	0.15 ± 0.001	0.05 ± 0.004	0.15 ± 0.006	0.08 ± 0.003
Kidney	0.37 ± 0.005	0.09 ± 0.007	0.40 ± 0.029	0.09 ± 0.007	0.39 ± 0.045	0.14 ± 0.016
Small intestine	0.41 ± 0.058	0.05 ± 0.010	0.38 ± 0.062	0.07 ± 0.008	0.28 ± 0.015	0.09 ± 0.015
Muscle	0.11 ± 0.005	0.014 ± 0.001	0.11 ± 0.013	0.012 ± 0.001	0.10 ± 0.011	0.027 ± 0.003

4  
 5 <sup>a</sup> Male Wistar rats, 250 g.

6  
 7 n = number of animals.

8 Values shown are means ± standard deviation.

9 Source: Bolt and Filser (1977).

10  
 11  
 12 binding of radiolabel from [<sup>14</sup>C]TCE by comparing tissue binding after administration of  
 13 200 mg/kg via oral gavage in corn oil between control (naïve) rats and rats pretreated with  
 14 phenobarbital (a known inducer of CYP2B family) or Aroclor 1254 (a known inducer of both  
 15 CYP1A and CYP2B families of isoenzymes) (see Table 3-12). The results indicate that  
 16 induction of total cytochromes P-450 content by three- to fourfold resulted in nearly 10-fold  
 17 increase in radioactivity (disintegrations per minute; [DPM]) bound in liver and kidney. By  
 18 contrast, Mazzullo et al. (1992) reported that, phenobarbital pretreatment did not result in  
 19 consistent or marked alterations of in vivo binding of radiolabel to DNA, RNA, or protein in rats  
 20 and mice at 22 hours after an intraperitoneal (i.p.) injection of [<sup>14</sup>C]TCE. On the other hand, in  
 21 vitro experiments by Mazzullo et al. (1992) reported reduction of TCE-radiolabel binding to calf  
 22 thymus DNA with introduction of a CYP inhibitor into incubations containing rat liver

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 microsomal protein. Moreover, increase/decrease of glutathione levels in incubations containing  
2 lung cytosolic protein led to a parallel increase/decrease in TCE-radiolabel binding to calf  
3 thymus DNA.

4

5

1 **Table 3-12. Binding of <sup>14</sup>C from [<sup>14</sup>C]TCE in rat liver and kidney at 72 hours**  
 2 **after oral administration of 200 mg/kg [<sup>14</sup>C]TCE (Dekant et al., 1986b)**  
 3

Tissue	DPM/gram tissue		
	Untreated	Phenobarbital	Arochlor 1254
Liver	850 ± 100	9,300 ± 1,100	8,700 ± 1,000
Kidney	680 ± 100	5,700 ± 900	7,300 ± 800

4  
5  
**3.3. METABOLISM**

6 This section focuses on both in vivo and in vitro studies of the biotransformation of  
 7 trichloroethylene, identifying metabolites that are deemed significant for assessing toxicity and  
 8 carcinogenicity. In addition, metabolism studies may be used to evaluate the flux of parent  
 9 compound through the known metabolic pathways. Sex-, species-, and interindividual  
 10 differences in the metabolism of TCE are discussed, as are factors that possibly contribute to this  
 11 variability. Additional discussion of variability and susceptibility is presented in Section 4.10.  
 12

**3.3.1. Introduction**

13 The metabolism of TCE has been studied mostly in mice, rats, and humans and has been  
 14 extensively reviewed (IARC, 1995d; Lash et al., 2000a; U.S. EPA, 1985). It is now well  
 15 accepted that TCE is metabolized in laboratory animals and in humans through at least  
 16 two distinct pathways: (1) oxidative metabolism via the cytochrome P450 mixed-function  
 17 oxidase system and (2) GSH conjugation followed by subsequent further biotransformation and  
 18 processing, either through the cysteine conjugate beta lyase pathway or by other enzymes (Lash  
 19 et al., 2000b). While the flux through the conjugative pathway is less, quantitatively, than the  
 20 flux through oxidation (Bloemen et al., 2001), GSH conjugation is an important route  
 21 toxicologically, giving rise to relatively potent toxic biotransformation products (Elfarra et al.,  
 22 1987; Elfarra et al., 1986).


23 Information about metabolism is important because, as discussed extensively in  
 24 Chapter 4, certain metabolites are thought to cause one or more of the same acute and chronic  
 25 toxic effects, including carcinogenicity, as TCE. Thus, in many of these cases, the toxicity of  
 26 TCE is generally considered to reside primarily in its metabolites rather than in the parent  
 27 compound itself.

*This document is a draft for review purposes only and does not constitute Agency policy.*

### 3.3.2. Extent of Metabolism


2 TCE is extensively metabolized in animals and humans. The most comprehensive  
3 mass-balance studies are in mice and rats (Dekant et al., 1984, 1986a; Dekant et al., 1986b;  
4 Green and Prout, 1985; Prout et al., 1985) in which [<sup>14</sup>C]TCE is administered by oral gavage at  
5 doses of 2 –2,000 mg/kg, the data from which are summarized in Figure 3-2 and Figure 3-3. In  
6 both mice and rats, regardless of sex and strain, there is a general trend of increasing exhalation  
7 of unchanged TCE with dose, suggesting saturation of a metabolic pathway. The increase is  
8 smaller in mice (from 1–6% to 10–18%) than in rats (from 1–3% to 43–78%), suggesting  
9 greater overall metabolic capacity in mice. The dose at which apparent saturation occurs appears  
10 to be more sex- or strain-dependent in mice than in rats. In particular, the marked increase in  
11 exhaled TCE occurred between 20 and 200 mg/kg in female NMRI mice, between 500 and  
12 1,000 mg/kg in B6C3F1 mice, and between 10 and 500 mg/kg in male Swiss-Webster mice.  
13 However, because only one study is available in each strain, interlot or interindividual variability  
14 might also contribute to the observed differences. In rats, all three strains tested showed marked  
15 increase in unchanged TCE exhaled between 20 and 200 mg/kg or 10 and 500 mg/kg.  
16 Recovered urine, the other major source of excretion, had mainly TCA, trichloroethanol, and  
17 trichloroethanol-glucuronide conjugate (TCOG), but revealed no detectable TCE. The source of  
18 radioactivity in feces was not analyzed, but it is presumed not to include substantial TCE given  
19 the complete absorption expected from the corn oil vehicle. Therefore, at all doses tested in  
20 mice, and at doses <200 mg/kg in rats, the majority of orally administered TCE is metabolized.  
21 Pretreatment of rats with P450 inducers prior to a 200 mg/kg dose did not change the pattern of  
22 recovery, but it did increase the amount recovered in urine by 10–15%, with a corresponding  
23 decrease in the amount of exhaled unchanged TCE (Dekant et al., 1986b).

24 Comprehensive mass balance studies are not available in humans, but several studies  
25 have measured or estimated recovery of TCE in exhaled breath and/or TCA and TCOH in urine  
26 following controlled inhalation exposures to TCE (Monster et al., 1976; Opdam, 1989b; Soucek  
27 and Vlachova, 1960). Opdam (1989b) only measured exhaled breath, and estimated that, on  
28 average, 15–20% of TCE uptake (retained dose) was exhaled after exposure to 5.8–38 ppm for  
29 29–62 minutes. Soucek and Vlachova (1960) and Bartonicek (1962) did not measure exhaled  
30 breath but did report 69–73% of the retained dose excreted in urine as TCA and TCOH  
31 following exposure to 93–194 ppm (500–1,043 mg/m<sup>3</sup>) for 5 hours. Soucek and Vlachova  
32 (1960) additionally reported 4% of the retained dose excreted in urine as monochloroacetic acid  
33 (MCA). Monster et al. (1976) reported that an average of 10% of the retained TCE dose was



1 eliminated unchanged following 6 hour exposures to 70–140 ppm (376–752 mg/m<sup>3</sup>) TCE, along  
 2 with an average of 57% of the retained dose excreted in urine as TCA and free or conjugated  
 3

4  
 5 **Figure 3-2. Disposition of [<sup>14</sup>C]TCE administered by oral gavage in mice**  
 6 **(Dekant et al., 1984; Dekant et al., 1986b; Green and Prout, 1985; Prout et**  
 7 **al., 1985).**  
 8  
 9



1  
2  
3  
4  
5  
6

**Figure 3-3. Disposition of [<sup>14</sup>C]TCE administered by oral gavage in rats (Dekant et al., 1984; Dekant et al., 1986b; Green and Prout, 1985; Prout et al., 1985).**

1 TCOH. The differences among these studies may reflect a combination of interindividual  
2 variability and errors due to the difficulty in precisely estimating dose in inhalation studies, but  
3 in all cases less than 20% of the retained dose was exhaled unchanged and greater than 50% was  
4 excreted in urine as TCA and TCOH. Therefore, it is clear that TCE is extensively metabolized  
5 in humans. No saturation was evident in any of these human recovery studies at the exposure  
6 levels tested.

7

### 3.3.3. Pathways of Metabolism

8 As mentioned in Section 3.3.1, TCE metabolism in animals and humans has been  
9 observed to occur via two major pathways: P450-mediated oxidation and GSH conjugation.  
10 Products of the initial oxidation or conjugation step are further metabolized to a number of other  
11 metabolites. For P450 oxidation, all steps of metabolism occur primarily in the liver, although  
12 limited oxidation of TCE has been observed in the lungs of mice, as discussed below. The GSH  
13 conjugation pathway also begins predominantly in the liver, but toxicologically significant  
14 metabolic steps occur extrahepatically—particularly in the kidney (Lash et al., 1999a; 2006;  
15 Lash et al., 1998b; Lash et al., 1995). The mass-balance studies cited above found that at  
16 exposures below the onset of saturation, >50% of TCE intake is excreted in urine as oxidative  
17 metabolites (primarily as TCA and TCOH), so TCE oxidation is generally greater than TCE  
18 conjugation. This is discussed in detail in Section 3.3.3.3.


19

#### 3.3.3.1.1. Cytochrome P450-Dependent Oxidation

20 Oxidative metabolism by the cytochrome P450, or CYP-dependent, pathway is  
21 quantitatively the major route of TCE biotransformation (IARC, 1995d; 2000a; Lash et al.,  
22 2000b; U.S. EPA, 1985). The pathway is operative in humans and rodents and leads to several  
23 metabolic products, some of which are known to cause toxicity and carcinogenicity (IARC,  
24 1995d; U.S. EPA, 1985). Although several of the metabolites in this pathway have been clearly  
25 identified, others are speculative or questionable. Figure 3-4 depicts the overall scheme of TCE  
26 P450 metabolism.

27 In brief, TCE oxidation via P450, primarily CYP2E1 (Guengerich and Shimada, 1991),  
28 yields an oxygenated TCE-P450 intermediate. The TCE-P450 complex is a transition state that  
29 goes on to form chloral or TCE oxide. In the presence of water, chloral rapidly equilibrates with  
30 chloral hydrate (CH), which undergoes reduction and oxidation by alcohol dehydrogenase and





1 aldehyde dehydrogenase or aldehyde oxidase to form TCOH and TCA, respectively (Dekant et  
 2 al., 1986b; Green and Prout, 1985; Miller and Guengerich, 1983). TCE oxide can rearrange to  
 3

4 **Figure 3-4. Scheme for the oxidative metabolism of TCE.**

5  
 6 Adapted from: Clewell et al. (2000); Cummings et al. (2001); Forkert et al. (2006); Lash et al.  
 7 (2000b); Lash et al. (2000a); Tong et al. (1998).  
 8  
 9  
 10

1  
2  
3

**Table 3-13. In vitro TCE oxidative metabolism in hepatocytes and microsomal fractions**

In vitro system	K <sub>M</sub>	V <sub>MAX</sub>	1,000 × V <sub>MAX</sub> /K <sub>M</sub>	Source
	μM in medium	nmol TCE oxidized/min/mg MSP <sup>a</sup> or 10 <sup>6</sup> hepatocytes		
Human hepatocytes	210 ± 159 (45–403)	0.268 ± 0.215 (0.101–0.691)	2.45 ± 2.28 (0.46–5.57)	Lipscomb et al. (1998a)
Human liver microsomal protein	16.7 ± 2.45 (13.3–19.7)	1.246 ± 0.805 (0.490–3.309)	74.1 ± 44.1 (38.9–176)	Lipscomb et al. (1997) (Low K <sub>M</sub> )
	30.9 ± 3.3 (27.0–36.3)	1.442 ± 0.464 (0.890–2.353)	47.0 ± 16.0 (30.1–81.4)	Lipscomb et al. (1997) (Mid K <sub>M</sub> )
	51.1 ± 3.77 (46.7–55.7)	2.773 ± 0.577 (2.078–3.455)	54.9 ± 14.1 (37.3–69.1)	Lipscomb et al. (1997) (High K <sub>M</sub> )
	24.6	1.44	58.5	Lipscomb et al. (1998b) (pooled)
	12 ± 3 (9–14)	0.52 ± 0.17 (0.37–0.79)	48 ± 23 (26–79)	Elfarra et al. (1998) (males, high affinity)
	26 ± 17 (13–45)	0.33 ± 0.15 (0.19–0.48)	15 ± 10 (11–29)	Elfarra et al. (1998) (females, high affinity)
Rat liver microsomal protein	55.5	4.826	87.0	Lipscomb et al. (1998b) (pooled)
	72 ± 82	0.96 ± 0.65	24 ± 21	Elfarra et al. (1998) (males, high affinity)
	42 ± 21	2.91 ± 0.71	80 ± 34	Elfarra et al. (1998) (females, high affinity)
Rat kidney microsomal protein	940	0.154	0.164	Cummings et al. (2001)
Mouse liver microsomal protein	35.4	5.425	153	Lipscomb et al. (1998b) (pooled)
	378 ± 414	8.6 ± 4.5	42 ± 29	Elfarra et al. (1998) (males)
	161 ± 29	26.06 ± 7.29	163 ± 37	Elfarra et al. (1998) (females)

4  
5  
6

<sup>a</sup> MSP = Microsomal protein.

1 Notes: Results presented as mean  $\pm$  standard deviation (minimum–maximum).  $K_M$  for human hepatocytes  
2 converted from ppm in headspace to  $\mu\text{M}$  in medium using reported hepatocyte:air partition coefficient (Lipscomb et  
3 al., 1998a).  
4

1 dichloroacetyl chloride. Table 3-13 summarizes available in vitro measurements of TCE  
2 oxidation, as assessed by the formation of CH, TCOH, and TCA. Glucuronidation of TCOH  
3 forms TCOG, which is readily excreted in urine. Alternatively, TCOG can be excreted in bile  
4 and passed to the small intestine where it is hydrolyzed back to TCOH and reabsorbed (Bull,  
5 2000). TCA is poorly metabolized but may undergo dechlorination to form dichloroacetic  
6 acid(DCA). However, TCA is predominantly excreted in urine, albeit at a relatively slow rate as  
7 compared to TCOG. Like the TCE-P450 complex, TCE oxide also seems to be a transient  
8 metabolite. Recent data suggest that it is transformed to dichloroacetyl chloride, which  
9 subsequently decomposes to form DCA (Cai and Guengerich, 1999). As shown in Figure 3-4,  
10 several other metabolites, including oxalic acid and *N*-(hydroxyacetyl) aminoethanol, may form  
11 from the TCE oxide or the TCE-O-P450 intermediate and have been detected in the urine of  
12 rodents and humans following TCE exposure. Pulmonary excretion of CO<sub>2</sub> has been identified  
13 in exhaled breath from rodents exposed to <sup>14</sup>C-labeled TCE and is thought to arise from  
14 metabolism of DCA. The following sections provide details as to pathways of TCE oxidation,  
15 including discussion of inter- and intraspecies differences in metabolism.  
16

#### **3.3.3.1.2. Formation of trichloroethylene oxide**

17 In previous studies of halogenated alkene metabolism, the initial step was the generation  
18 of a reactive epoxide (Anders and Jakobson, 1985). Early studies in anesthetized human patients  
19 (Powell, 1945), dogs (Butler, 1949), and later reviews (e.g., Goeptar et al., 1995) suggest that the  
20 TCE epoxide may be the initial reaction product of TCE oxidation.

21 Epoxides can form acyl chlorides or aldehydes, which can then form aldehydes,  
22 carboxylic acids, or alcohols, respectively. Thus, earlier studies suggesting the appearance of  
23 CH, TCA, and TCOH as the primary metabolites of TCE were considered consistent with the  
24 oxidation of TCE to an epoxide intermediate (Butler, 1949; Powell, 1945). Following in vivo  
25 exposures to 1,1-dichloroethylene, a halocarbon very similar in structure to TCE, mouse liver  
26 cytosol and microsomes and lung Clara cells exhibited extensive P450-mediated epoxide  
27 formation (Dowsley et al., 1996; Forkert, 1999a, b; Forkert et al., 1999) Indeed, TCE oxide  
28 inhibits purified CYP2E1 activity (Cai and Guengerich, 2001a) similarly to TCE inhibition of  
29 CYP2E1 in human liver microsomes (Lipscomb et al., 1997).

30 Conversely, cases have been made against TCE oxide as an obligate intermediate to the  
31 formation of chloral. Using liver microsomes and reconstituted P450 systems (Miller and  
32 Guengerich, 1982, 1983) or isolated rat hepatocytes (Miller and Guengerich, 1983), it has been  
33 suggested that chlorine migration and generation of a TCE-O-P450 complex (via the heme

1 oxygen) would better explain the observed destruction of the P450 heme, an outcome not likely  
2 to be epoxide-mediated. Miller and Guengerich (1982) found CYP2E1 to generate an epoxide  
3 but argued that the subsequent production of chloral was not likely related to the epoxide. Green  
4 and Prout (1985) argued against epoxide (free form) formation in vivo in mice and rats,  
5 suggesting that the expected predominant metabolites would be carbon monoxide, CO<sub>2</sub>, MCA,  
6 and DCA, rather than the observed predominant appearance of TCA, TCOH, and TCOG.

7 It appears likely that both a TCE-O-P450 complex and a TCE oxide are formed, resulting  
8 in both CH and dichloroacetyl chloride, respectively, though it appears that the former  
9 predominates. In particular, it has been shown that dichloroacetyl chloride can be generated  
10 from TCE oxide, dichloroacetyl chloride can be trapped with lysine (Cai and Guengerich, 1999),  
11 and that dichloroacetyl-lysine adducts are formed in vivo (Forkert et al., 2006). Together, these  
12 data strongly suggest TCE oxide as an intermediate metabolite, albeit short-lived, from TCE  
13 oxidation in vivo.  
14

### **3.3.3.1.3. Formation of chloral hydrate (CH), trichloroethanol (TCOH) and trichloroacetic acid (TCA)**

15 CH (in equilibrium with chloral) is a major oxidative metabolite produced from TCE as  
16 has been shown in numerous in vitro systems, including human liver microsomes and purified  
17 P450 CYP2E1 (Guengerich et al., 1991) as well as recombinant rat, mouse, and human P450s  
18 including CYP2E1 (Forkert et al., 2005). However, in rats and humans, in vivo circulating CH is  
19 generally absent from blood following TCE exposure. In mice, CH is detectable in blood and  
20 tissues but is rapidly cleared from systemic circulation (Abbas and Fisher, 1997). The low  
21 systemic levels of CH are because of its rapid transformation to other metabolites.

22 CH is further metabolized predominantly to TCOH (Sellers et al., 1972; Shultz and  
23 Weiner, 1979) and/or CYP2E1 (Ni et al., 1996). The role for alcohol dehydrogenase was  
24 suggested by the observation that ethanol inhibited CH reduction to TCOH (Larson and Bull,  
25 1989; Muller et al., 1975; Sellers et al., 1972). For instance, Sellers et al. (1972) reported that  
26 coexposure of humans, to ethanol and CH resulted in a higher percentage of urinary TCOH (24%  
27 of CH metabolites) compared to TCA (19%). When ethanol was absent, 10 and 11% of CH was  
28 metabolized to TCOH and TCA, respectively. However, because ethanol can be oxidized by  
29 both alcohol dehydrogenase and CYP2E1, there is some ambiguity as to whether these  
30 observations involve competition with one or the other of these enzymes. For instance, Ni et al.  
31 (1996) reported that CYP2E1 expression was necessary for metabolism of CH to mutagenic  
32 metabolites in a human lymphoblastoid cell line, suggesting a role for CYP2E1. Furthermore, Ni

1 et al. (1996) reported that cotreatment of mice with CH and pyrazole, a specific CYP2E1  
2 inducer, resulted in enhanced liver microsomal lipid peroxidation, while treatment with DPEA,  
3 an inhibitor of CYP2E1, suppressed lipid peroxidation, suggesting CYP2E1 as a primary enzyme  
4 for CH metabolism in this system. Lipscomb et al. (1996) suggested that two enzymes are likely  
5 responsible for CH reduction to TCOH based on observation of bi-phasic metabolism for this  
6 pathway in mouse liver microsomes. This behavior has also been observed in mouse liver  
7 cytosol, but was not observed in rat or human liver microsomes. Moreover, CH metabolism to  
8 TCOH increased significantly both in the presence of nicotinamide adenine dinucleotide  
9 (NADH) in the 700× g supernatant of mouse, rat, and human liver homogenate as well as with  
10 the addition of NADPH in human samples, suggesting two enzymes may be involved (Lipscomb  
11 et al., 1996).

12 TCOH formed from CH is available for oxidation to TCA (see below) or glucuronidation  
13 via uridine 5'-diphospho-glucuronyltransferase to TCOG, which is excreted in urine or in bile  
14 (Stenner et al., 1997). Biliary TCOG is hydrolyzed in the gut and available for reabsorption to  
15 the liver as TCOH, where it can be glucuronidated again or metabolized to TCA. This  
16 enterohepatic circulation appears to play a significant role in the generation of TCA from TCOH  
17 and in the observed lengthy residence time of this metabolite, compared to TCE. Using jugular-,  
18 duodenal-, and bile duct-cannulated rats, Stenner et al. (1997) showed that enterohepatic  
19 circulation of TCOH from the gut back to the liver and subsequent oxidation to TCA was  
20 responsible for 76% of TCA measured in the systemic blood.

21 Oxidation of CH and TCOH to TCA has been demonstrated in vivo in mice (Dekant et  
22 al., 1986b; Green and Prout, 1985; Larson and Bull, 1992a), rats (Dekant et al., 1986b; Green  
23 and Prout, 1985; Larson and Bull, 1992a; Pravecsek et al., 1996; Stenner et al., 1997; Templin et  
24 al., 1995b), dogs (Templin et al., 1995b), and humans (Sellers et al., 1978). Urinary metabolite  
25 data in mice and rats exposed to 200 mg/kg TCE (Dekant et al., 1986b; Larson and Bull, 1992a);  
26 and humans following oral CH exposure (Sellers et al., 1978) show greater TCOH production  
27 relative to TCA production. However, because of the much longer urinary half-life in humans of  
28 TCA relative to TCOH, the total amount of TCA excreted may be similar to TCOH (Fisher et al.,  
29 1998; Monster et al., 1976). This is thought to be primarily due to conversion of TCOH to TCA,  
30 either directly or via "back-conversion" of TCOH to CH, rather than due to the initial formation  
31 of TCA from CH (Owens and Marshall, 1955).

32 In vitro data are also consistent with CH oxidation to TCA being much less than CH  
33 reduction to TCOH. For instance, Lipscomb et al. (1996) reported 1,832-fold differences in  $K_M$   
34 values and 10–195-fold differences in clearance efficiency ( $V_{MAX}/K_M$ ) for TCOH and TCA in all  
35 three species (see Table 3-14). Clearance efficiency of CH to TCA in mice is very similar to

1 humans but is 13-fold higher than rats. Interestingly, Bronley-DeLancey et al. (2006) recently  
2 reported that similar amounts of TCOH and TCA were generated from CH using cryopreserved  
3 human hepatocytes. However, the intersample variation was extremely high, with measured  
4  $V_{MAX}$  ranging from eightfold greater TCOH to fivefold greater TCA and clearance ( $V_{MAX}/K_M$ )  
5 ranging from 13-fold greater TCOH to 17-fold greater TCA. Moreover, because a comparison  
6 with fresh hepatocytes or microsomal protein was not made, it is not clear to what extent these  
7 differences are due to population heterogeneity or experimental procedures.

8 The metabolism of CH to TCA and TCOH involves several enzymes including CYP2E1,  
9 alcohol dehydrogenase, and aldehyde dehydrogenase enzymes (Guengerich et al., 1991; Miller  
10 and Guengerich, 1983; Ni et al., 1996; Shultz and Weiner, 1979; Wang et al., 1993). Because  
11 these enzymes have preferred cofactors (NADPH, NADH, and NAD<sup>+</sup>), cellular cofactor ratio  
12 and redox status of the liver may have an impact on the preferred pathway (Kawamoto et al.,  
13 1988a; Lipscomb et al., 1996).

#### 14 **3.3.3.1.4. Formation of dichloroacetic acid (DCA) and other products**

15 As discussed above, DCA could hypothetically be formed via multiple pathways. The  
16 work reviewed by Guengerich (2004) has suggested that one source of DCA may be through a  
17 TCE oxide intermediary. Miller and Guengerich (1983) report evidence of formation of the  
18 epoxide, and Cai and Guengerich (1999) report that a significant amount (about 35%) of DCA is  
19 formed from aqueous decomposition of TCE oxide via hydrolysis in an almost pH-independent  
20 manner. Because this reaction forming DCA from TCE oxide is a chemical process rather than a  
21 process mediated by enzymes, and because evidence suggests that some epoxide was formed  
22 from TCE oxidation, Guengerich (2004) notes that DCA would be an expected product of TCE  
23 oxidation (see also Yoshioka et al., 2002). Alternatively, dechlorination of TCA and oxidation  
24 of TCOH have been proposed as sources of DCA (Lash et al., 2000a). Merdink et al. (2000)  
25 investigated dechlorination of TCA and reported trapping a DCA radical with the spin-trapping  
26 agent phenyl-tert-butyl nitroxide, identified by gas chromatography/mass spectroscopy, in both a  
27 chemical Fenton system and rodent microsomal incubations with TCA as substrate.  
28 Dose-dependent catalysis of TCA to DCA was observed in cultured microflora from B6C3F1  
29 mice (Moghaddam et al., 1996). However, while antibiotic-treated mice lost the ability to  
30 produce DCA in the gut, plasma DCA levels were unaffected by antibiotic treatment, suggesting  
31 that the primary site of murine DCA production is

**Table 3-14. In vitro kinetics of trichloroethanol and trichloroacetic acid formation from chloral hydrate in rat, mouse, and human liver homogenates**

Species	TCOH			TCA		
	$K_m^a$	$V_{MAX}^b$	$V_{MAX}/K_m^c$	$K_m^a$	$V_{MAX}^b$	$V_{MAX}/K_m^c$
Rat	0.52	24.3	46.7	16.4	4	0.24
Mouse <sup>d</sup>	0.19	11.3	59.5	3.5	10.6	3.0
High affinity	0.12	6.3	52.5	na <sup>e</sup>	na	na
Low affinity	0.51	6.1	12.0	na	na	na
Human	1.34	34.7	25.9	23.9	65.2	2.7

<sup>a</sup>  $K_m$  presented as mM CH in solution.

<sup>b</sup>  $V_{MAX}$  presented as nmoles/mg supernatant protein/min.

<sup>c</sup> Clearance efficiency represented by  $V_{MAX}/K_m$ .

<sup>d</sup> Mouse kinetic parameters derived for observations over the entire range of CH exposure as well as discrete, bi-phasic regions for CH concentrations below (high affinity) and above (low affinity) 1.0 mM.

<sup>e</sup> na = not applicable.

Source: Lipscomb et al. (1996).

other than the gut (Moghaddam et al., 1997).

However, direct evidence for DCA formation from TCE exposure remains equivocal. In vitro studies in human and animal systems have demonstrated very little DCA production in the liver (James et al., 1997). In vivo, DCA was detected in the blood of mice (Larson and Bull, 1992a; Templin et al., 1993) and humans (Fisher et al., 1998) and in the urine of rats and mice (Larson and Bull, 1992b) exposed to TCE by aqueous oral gavage. However, the use of strong acids in the analytical methodology produces *ex vivo* conversion of TCA to DCA in mouse blood (Ketcha et al., 1996). This method may have resulted in the appearance of DCA as an artifact in human plasma (Fisher et al., 1998) and mouse blood in vivo (Templin et al., 1995b). Evidence for the artifact is suggested by DCA AUCs that were larger than would be expected from the available TCA (Templin et al., 1995b). After the discovery of these analytical issues, Merdink et al. (1998) reevaluated the formation of DCA from TCE, TCOH, and TCA in mice, with particular focus on the hypothesis that DCA is formed from dechlorination of TCA. They were unable to detect blood DCA in naive mice after administration of TCE, TCOH, or TCA. Low levels of DCA were detected in the blood of children administered therapeutic doses of CH (Henderson et al., 1997), suggesting TCA or TCOH as the source of DCA. Oral TCE exposure in rats and dogs failed to produce detectable levels of DCA (Templin et al., 1995b).

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 Another difficulty in assessing the formation of DCA is its rapid metabolism at low  
2 exposure levels. Degradation of DCA is mediated by glutathione-S-transferase (GST)-zeta  
3 (Saghir and Schultz, 2002; Tong et al., 1998), apparently occurring primarily in the hepatic  
4 cytosol. DCA metabolism results in suicide inhibition of the enzyme, evidenced by decreased  
5 DCA metabolism in DCA-treated animals (Gonzalez-Leon et al., 1999) and humans (Shroads et  
6 al., 2008) and loss of DCA metabolic activity and enzymatic protein in liver samples from  
7 treated animals (Schultz et al., 2002). This effect has been noted in young mice exposed to DCA  
8 in drinking water at doses approximating 120 mg/kg-day (Schultz et al., 2002). The  
9 experimental data and pharmacokinetic model simulations of several investigators (Jia et al.,  
10 2006; Keys et al., 2004; Li et al., 2008; Merdink et al., 1998; Shroads et al., 2008) suggest that  
11 several factors prevent the accumulation of measurable amounts of DCA: (1) its formation as a  
12 short-lived intermediate metabolite, and (2) its rapid elimination relative to its formation from  
13 TCA. While DCA elimination rates appear approximately one order of magnitude higher in rats  
14 and mice than in humans (James et al., 1997) (see Table 3-15), they still may be rapid enough so  
15 that even if DCA were formed in humans, it would be metabolized too quickly to appear in  
16 detectable quantities in blood.

17 A number of other metabolites, such as oxalic acid, MCA, glycolic acid, and glyoxylic  
18 acid, are formed from DCA (Lash et al., 2000a; Saghir and Schultz, 2002). Unlike other  
19 oxidative metabolites of TCE, DCA appears to be metabolized primarily via hepatic cytosolic  
20 proteins. Since P450 activity resides almost exclusively in the microsomal and mitochondrial  
21 cell fractions, DCA metabolism appears to be independent of P450. Rodent microsomal and  
22 mitochondrial metabolism of DCA was measured to be  $\leq 10\%$  of cytosolic metabolism  
23 (Lipscomb et al., 1995). DCA in the liver cytosol from rats and humans is transformed to  
24 glyoxylic acid via a GSH-dependent pathway (James et al., 1997). In rats, the  $K_M$  for GSH was  
25 0.075 mM with a  $V_{MAX}$  for glyoxylic acid formation of 1.7 nmol/mg protein/minute. While this  
26 pathway may not involve GST (as evidenced by very low GST activity in this study), Tong et al.  
27 (1998) showed GST-zeta, purified from rat liver, to be involved in metabolizing DCA to  
28 glyoxylic acid, with a  $V_{MAX}$  of 1,334 nmol/mg protein/minute and  $K_M$  of 71.4  $\mu$ M for glyoxylic  
29 acid formation and a GSH  $K_M$  of 59  $\mu$ M.

30

**Table 3-15. In vitro kinetics of DCA metabolism in hepatic cytosol of mice, rats, and humans**

Species	$V_{MAX}$ (nmol/min/mg protein)	$K_M$ ( $\mu$ M)	$V_{MAX}/K_M$
Mouse	13.1	350	37.4
Rat	11.6	280	41.4
Human	0.37	71	5.2

Source: James et al. (1997).

### 3.3.3.1.5. Tissue distribution of oxidative metabolism and metabolites

Oxidative metabolism of TCE, irrespective of the route of administration, occurs predominantly in the liver, but TCE metabolism via the P450 (CYP) system also occurs at other sites because CYP isoforms are present to some degree in most tissues of the body. For example, both the lung and kidneys exhibit cytochrome P450 enzyme activities (Cummings et al., 2001; Forkert et al., 2005; 1997a; Green et al., 1997b). Green et al. (1997b) detected TCE oxidation to chloral in microsomal fractions of whole-lung homogenates from mice, rats, and humans, with the activity in mice the greatest and in humans the least. The rates were slower than in the liver (which also has a higher microsomal protein content as well as greater tissue mass) by 1.8-, 10-, and >10-fold in mice, rats, and humans, respectively. While qualitatively informative, these rates were determined at a single concentration of about 1 mM TCE. A full kinetic analysis was not performed, so clearance and maximal rates of metabolism could not be determined. With the kidney, Cummings et al. (2001) performed a full kinetic analysis using kidney microsomes, and found clearance rates ( $V_{MAX}/K_M$ ) for oxidation were more than 100-fold smaller than average rates that were found in the liver (see Table 3-13). In human kidney microsomes, Amet et al.(1997) reported that CYP2E1 activity was weak and near detection limits, with no CYP2E1 detectable using immunoblot analysis. Cummings and Lash (2000) reported detecting oxidation of TCE in only one of four kidney microsome samples, and only at the highest tested concentration of 2 mM, with a rate of 0.13 nmol/minute/mg protein. This rate contrasts with the  $V_{MAX}$  values for human liver microsomal protein of 0.19–3.5 nmol/minute/mg protein reported in various experiments (see Table 3-13, above). Extrahepatic oxidation of TCE may play an important role for generation of toxic metabolites in situ. The roles of local

1 metabolism in kidney and lung toxicity are discussed in detail in Sections 4.4 and 4.7,  
2 respectively.

3         With respect to further metabolism beyond oxidation of TCE, CH has been shown to be  
4 metabolized to TCA and TCOH in lysed whole blood of mice and rats and fractionated human  
5 blood (Lipscomb et al., 1996) (see Table 3-16). TCOH production is similar in mice and rats and  
6 is approximately twofold higher in rodents than in human blood. However, TCA formation in  
7 human blood is two- or threefold higher than in mouse or rat blood, respectively. In human  
8 blood, TCA is formed only in the erythrocytes. TCOH formation occurs in both plasma and  
9 erythrocytes, but fourfold more TCOH is found in plasma than in an equal volume of packed  
10 erythrocytes. While blood metabolism of CH may contribute further to its low circulating levels  
11 in vivo., the metabolic capacity of blood (and kidney) may be substantially lower than liver.  
12 Regardless, any CH reaching the blood may be rapidly metabolized to TCA and TCOH. DCA  
13 and TCA are known to bind to plasma proteins. Schultz et al. (1999) measured DCA binding in  
14 rats at a single concentration of about 100  $\mu\text{M}$  and found a binding fraction of less than 10%.  
15 However, these data are not greatly informative for TCE exposure in which DCA levels are  
16 significantly lower, and limitation to a single concentration precludes fitting to Templin et al.  
17 (1993, 1995a; 1995b), Schultz et al. (1999), Lumpkin et al. (2003), and Yu et al. (2003) all  
18 measured TCA binding in various species and at various concentration ranges. Of these,  
19 Templin et al. (1995a; 1995b) and Lumpkin et al. (2003) measured levels in humans, mice, and  
20 rats. Lumpkin et al. (2003) studied the widest concentration range, spanning reported TCA  
21 plasma concentrations from experimental studies. Table 3-17 shows derived binding parameters.  
22 However, these data are not entirely consistent among researchers; two- to fivefold differences in  
23  $B_{\text{MAX}}$  and  $K_d$  are noted in some cases, although some differences existed in the rodent strains and  
24 experimental protocols used. In general, however, at lower concentrations, the bound fraction  
25 appears greater.

26

1 **Table 3-16. TCOH and TCA formed from CH in vitro in lysed whole blood**  
 2 **of rats and mice or fractionated blood of humans (nmoles formed in 400  $\mu$ L**  
 3 **samples over 30 minutes)**  
 4

	Rat	Mouse	Human	
			Erythrocytes	Plasma
TCOH	45.4 $\pm$ 4.9	46.7 $\pm$ 1.0	15.7 $\pm$ 1.4	4.48 $\pm$ 0.2
TCA	0.14 $\pm$ 0.2	0.21 $\pm$ 0.3	0.42 $\pm$ 0.0	not detected

5  
 6 Source: Lipscomb et al. (1996).  
 7  
 8

9 in humans than in rats and mice. Typical human TCE exposures, even in controlled experiments  
 10 with volunteers, lead to TCA blood concentrations well below the reported  $K_d$  (see Table 3-17,  
 11 below), so the TCA binding fraction should be relatively constant. However, in rats and mice,  
 12 experimental exposures may lead to peak concentrations similar to, or above, the reported  $K_d$   
 13 (e.g., Templin et al., 1993; Yu et al., 2000), meaning that the bound fraction should temporarily  
 14 decrease following such exposures.

15 Limited data are available on tissue:blood partitioning of the oxidative metabolites CH,  
 16 TCA, TCOH and DCA, as shown in Table 3-18. As these chemicals are all water soluble and  
 17 not lipophilic, it is not surprising that their partition coefficients are close to one (within about  
 18 twofold). It should be noted that the TCA tissue:blood partition coefficients reported in  
 19 Table 3-18 were measured at concentrations 1.6–3.3 M, over 1,000-fold higher than the reported  
 20  $K_d$ . Therefore, these partition coefficients should reflect the equilibrium between tissue and free  
 21 blood concentrations. In addition, only one in vitro measurement has been reported of  
 22 blood:plasma concentration ratios for TCA: Schultz et al. (1999) reported a value of 0.76 in rats.  
 23

### 3.3.3.1.6. Species-, sex-, and age-dependent differences of oxidative metabolism

24 The ability to describe species- and sex-dependent variations in TCE metabolism is  
 25 important for species extrapolation of bioassay data and identification of human populations that  
 26 are particularly susceptible to TCE toxicity. In particular, information on the variation in the  
 27 initial oxidative step of CH formation from TCE is desirable, because this is the rate-limiting  
 28 step in the eventual formation and distribution of the putative toxic metabolites TCA and DCA  
 29 (Lipscomb et al., 1997).

1 Inter- and intraspecies differences in TCE oxidation have been investigated in vitro using  
 2 cellular or subcellular fractions, primarily of the liver. The available in vitro metabolism data on  
 3 TCE oxidation in the liver (see Table 3-13) show substantial inter and intraspecies variability.

4 **Table 3-17. Reported TCA plasma binding parameters**

	A	B <sub>MAX</sub> (μM)	K <sub>d</sub> (μM)	A+ B <sub>MAX</sub> /K <sub>d</sub>	Concentration range (μM bound+free)
Human					
Templin et al. (1995b)	–	1,020	190	5.37	3–1,224
Lumpkin et al. (2003)	–	708.9	174.6	4.06	0.06–3,065
Rat					
Templin et al. (1995b)	–	540	400	1.35	3–1,224
Yu et al. (2000)	0.602	312	136	2.90	3.8–1,530
Lumpkin et al. (2003)	–	283.3	383.6	0.739	0.06–3,065
Mouse					
Templin et al. (1993)	–	310	248	1.25	3–1,224
Lumpkin et al. (2003)	–	28.7	46.1	0.623	0.06–1,226

6  
 7 Notes: Binding parameters based on the equation  $C_{\text{bound}} = A \times C_{\text{free}} + B_{\text{MAX}} \times C_{\text{free}} / (K_d + C_{\text{free}})$ , where  $C_{\text{bound}}$  is the  
 8 bound concentration,  $C_{\text{free}}$  is the free concentration, and  $A = 0$  for Templin et al. (1993; 1995b) and Lumpkin et al.  
 9 (2003). The quantity  $A + B_{\text{MAX}}/K_d$  is the ratio of bound-to-free at low concentrations.

10 **Table 3-18. Partition coefficients for TCE oxidative metabolites**

Species/tissue	Tissue: blood partition coefficient			
	CH	TCA	TCOH	DCA
<b>Human<sup>a</sup></b>				
Kidney	–	0.66	2.15	-
Liver	–	0.66	0.59	-
Lung	–	0.47	0.66	-
Muscle	–	0.52	0.91	-
<b>Mouse<sup>b</sup></b>				
Kidney	0.98	0.74	1.02	0.74
Liver	1.42	1.18	1.3	1.08
Lung	1.65	0.54	0.78	1.23
Muscle	1.35	0.88	1.11	0.37

11 *This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2           <sup>a</sup> Fisher et al. (1998).

3           <sup>b</sup> Abbas and Fisher (1997).

4           Note: TCA and TCOH partition coefficients have not been reported for rats.  
5

6 Across species, microsomal data show that mice apparently have greater capacity ( $V_{MAX}$ ) than  
7 rat or humans, but the variability within species can be 2- to 10-fold. Part of the explanation may  
8 be related to CYP2E1 content. Although liver P450 content is similar across species, mice and  
9 rats exhibit higher levels of CYP2E1 content (0.85 and 0.89 nmol/mg protein, respectively)  
10 (Davis et al., 2002; Nakajima et al., 1993) than humans (approximately 0.25–0.30 nmol/mg  
11 protein) (Davis et al., 2002; Elfarra et al., 1998). Thus, the data suggest that rodents would have  
12 a higher capacity than humans to metabolize TCE, but this is difficult to verify in vivo because  
13 very high exposure concentrations in humans would be necessary to assess the maximum  
14 capacity of TCE oxidation.

15           With respect to the  $K_M$  of liver microsomal TCE oxidative metabolism, where  $K_M$  is  
16 indicative of affinity (the lower the numerical value of  $K_M$ , the higher the affinity), the trend  
17 appears to be mice and rats have higher  $K_M$  values (i.e., lower affinity) than humans, but with  
18 substantial overlap due to interindividual variability. Note that, as shown in Table 3-13, the  
19 ranking of rat and mouse liver microsomal  $K_M$  values between the two reports by Lipscomb et al.  
20 (1998b) and Elfarra et al. (1998) is not consistent. However, both studies clearly show that  $K_M$  is  
21 the lowest (i.e., affinity is highest) in humans. Because clearance at lower concentrations is  
22 determined by the ratio  $V_{MAX}$  to  $K_M$ , the lower apparent  $K_M$  in humans may partially offset the  
23 lower human  $V_{MAX}$ , and lead to similar oxidative clearances in the liver at environmentally  
24 relevant doses. However, differences in activity measured in vitro may not translate into in vivo  
25 differences in metabolite production, as the rate of metabolism in vivo depends also on the rate  
26 of delivery to the tissue via blood flow (Lipscomb et al., 2003). The interaction of enzyme  
27 activity and blood flow is best investigated using PBPK models and is discussed, along with  
28 descriptions of in vivo data, in Section 3.5.

29           Data on sex- and age-dependence in oxidative TCE metabolism are limited but suggest  
30 relatively modest differences in humans and animals. In an extensive evaluation of  
31 CYP-dependent activities in human liver microsomal protein and cryopreserved hepatocytes,  
32 Parkinson et al. (2004) identified no age or gender-related differences in CYP2E1 activity. In  
33 liver microsomes from 23 humans, the  $K_M$  values for females was lower than males, but  $V_{MAX}$   
34 values were very similar (Lipscomb et al., 1997). Appearance of total trichloro compounds  
35 (TTC) in urine following intraperitoneal dosing with TCE was 28% higher in female rats than in  
36 males (Verma and Rana, 2003). The oxidation of TCE in male and female rat liver microsomes  
37 was not significantly different; however, pregnancy resulted in a decrease of 27–39% in the rate

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 of CH production in treated microsomes from females (Nakajima et al., 1992b). Formation of  
2 CH in liver microsomes in the presence of 0.2 or 5.9 mM TCE exhibited some dependency on  
3 age of rats, with formation rates in both sexes of 1.1–1.7 nmol/mg protein/minute in 3-week-old  
4 animals and 0.5–1.0 nmol/mg protein/minute in 18-week-old animals (Nakajima et al., 1992b).

5 Fisher et al. (1991) reviewed data available at that time on urinary metabolites to  
6 characterize species differences in the amount of urinary metabolism accounted for by TCA (see  
7 Table 3-19). They concluded that TCA seemed to represent a higher percentage of urinary  
8 metabolites in primates than in other mammalian species, indicating a greater proportion of  
9 oxidation leading ultimately to TCA relative to TCOG.

10

### 3.3.3.1.7. Cytochrome P450 (CYP) isoforms and genetic polymorphisms

11 A number of studies have identified multiple P450 isozymes as having a role in the oxidative  
12 metabolism of TCE. These isozymes include CYP2E1 (Guengerich et al., 1991; Guengerich and  
13 Shimada, 1991; Nakajima et al., 1988; Nakajima et al., 1992a; Nakajima et al., 1990), CYP3A4  
14 (Shimada et al., 1994), CYP1A1/2, CYP2C11/6 (Nakajima et al., 1992a, 1993), CYP2F, and  
15 CYP2B1 (Forkert et al., 2005). Recent studies in CYP2E1-knockout mice have shown that in  
16 the absence of CYP2E1, mice still have substantial capacity for TCE oxidation (Forkert et al.,  
17 2006; Kim and Ghanayem, 2006). However, CYP2E1 appears to be the predominant (i.e.,  
18 higher affinity) isoform involved in oxidizing TCE (Forkert et al., 2005; Guengerich et al., 1991;  
19 Guengerich and Shimada, 1991; Nakajima et al., 1992a). In rat liver, CYP2E1 catalyzed TCE  
20 oxidation more than CYP2C11/6 (Nakajima et al., 1992a). In rat recombinant-derived P450s,  
21 the CYP2E1 had a lower  $K_M$  (higher affinity) and higher  $V_{MAX}/K_M$  ratio (intrinsic clearance)  
22 than CYP2B1 or CYP2F4 (Forkert et al., 2005). Interestingly, there was substantial differences  
23 in  $K_M$  between rat and human CYP2E1s and between rat CYP2F4 and mouse CYP2F2,  
24 suggesting that species-specific isoforms have different kinetic behavior (see Table 3-20).

25 The presence of multiple P450 isoforms in human populations affects the variability in  
26 individuals' ability to metabolize TCE. Studies using microsomes from human liver or from  
27 human lymphoblastoid cell lines expressing CYP2E1, CYP1A1, CYP1A2, or CYP3A4 have  
28 shown that CYP2E1 is responsible for greater than 60% of oxidative TCE metabolism  
29 (Lipscomb et al., 1997). Similarities between metabolism of chlorzoxazone (a CYP2E1  
30 substrate) in liver microsomes from 28 individuals (Peter et al., 1990) and TCE metabolism  
31 helped identify CYP2E1 as the predominant (high affinity) isoform for TCE oxidation.  
32 Additionally, Lash et al. (2000a) suggested that, at concentrations above the  $K_M$  value for  
33 CYP2E1, CYP1A2 and CYP2A4 may also metabolize TCE in humans; however, their

1 **Table 3-19. Urinary excretion of trichloroacetic acid by various species**  
 2 **exposed to trichloroethylene [based on data reviewed in (Fisher et al., 1991)]**  
 3

Species	Percentage of urinary excretion of TCA		Dose route	TCE dose (mg TCE/kg)	References, comments
	Male	Female			
Baboon <sup>a,c</sup>	16	—	Intramuscular injection	50	Mueller et al. (1982)
Chimpanzee <sup>a</sup>	24	22	Intramuscular injection	50	Mueller et al. (1982)
Monkey, Rhesus <sup>a,c</sup>	19	—	Intramuscular injection	50	Mueller et al. (1982)
Mice, NMRI <sup>b</sup>	—	8–20	Oral intubation	2–200	Dekant et al. (1986b)
Mice, B6C3F1 <sup>a</sup>	7–12	—	Oral intubation	10–2,000	Green and Prout (1985)
Rabbit, Japanese White <sup>a,c</sup>	0.5	—	Intraperitoneal injection	200	Nomiyama and Nomiyama (1979)
Rat, Wistar <sup>b</sup>	—	14–17	Oral intubation	2–200	Dekant et al. (1986b)
Rat, Osborne-Mendel <sup>a</sup>	6–7	—	Oral intubation	10–2,000	Green and Prout (1985)
Rat, Holtzman <sup>a</sup>	7	—	Intraperitoneal injection	10 mg TCE/rat	Nomiyama and Nomiyama (1979)

4  
 5 <sup>a</sup> Percentage urinary excretion determined from accumulated amounts of TCOH and TCA in urine 3–6 days  
 6 postexposure.

7 <sup>b</sup> Percentage urinary excretion determined from accumulated amounts of TCOH, dichloroacetic acid, oxalic acid,  
 8 and *N*-(hydroxyacetyl)aminoethanol in urine 3 days postexposure.

9 <sup>c</sup> Sex is not specified.

10  
 11 Note: The human data tabulated in Fisher et al. (1991) from Nomiyama and Nomiyama (1971) were not included  
 12 here because they were relative to urinary excretion of total trichloro-compounds—not as fraction of intake as was  
 13 the case for the other data included here.  
 14



**Table 3-20. P450 isoform kinetics for metabolism of TCE to CH in human, rat, and mouse recombinant P450s**

<b>Experiment</b>	<b>K<sub>M</sub> μM</b>	<b>V<sub>MAX</sub> pmol/min/pmol P450</b>	<b>V<sub>MAX</sub>/K<sub>M</sub></b>
Human rCYP2E1	196 ± 40	4 ± 0.2	0.02
Rat rCYP2E1	14 ± 3	11 ± 0.3	0.79
Rat rCYP2B1	131 ± 36	9 ± 0.5	0.07
Rat rCYP2F4	64 ± 9	17 ± 0.5	0.27
Mouse rCYP2F2	114 ± 17	13 ± 0.4	0.11

Source: Forkert et al. (2005)

contribution to the overall TCE metabolism was considered low compared to that of CYP2E1. Given the difference in expression of known TCE-metabolizing P450 isoforms (see Table 3-21) and the variability in P450-mediated TCE oxidation (Lipscomb et al., 1997), significant variability may exist in individual human susceptibility to TCE toxicity. Differences in content and/or intrinsic catalytic properties (K<sub>M</sub>, V<sub>MAX</sub>) of specific enzymes among species, strains, and individuals may play an important role in the observed differences in TCE metabolism and resulting toxicities. Lipscomb et al. (1997) reported observing three statistically distinct groups of K<sub>M</sub> values for TCE oxidation using human microsomes. The mean ± standard deviation [SD] (μM TCE) for each of the three groups was 16.7 ± 2.5 (n = 10), 30.9 ± 3.3 (n = 9), and 51.1 ± 3.8 (n = 4). Within each group, there were no significant differences in sex or ethnicity. However, the overall observed K<sub>M</sub> values in female microsomes (21.9 ± 3.5 μM, n = 10) were significantly lower than males (33.1 ± 3.5 μM, n = 13). Interestingly, in human liver microsomes, different groups of individuals with different affinities for TCE oxidation appeared to also have different activities for other substrates not only with respect to CYP2E1 but also CYP1A2 (Lash et al., 2000a) (see Table 3-21). Genetic polymorphisms in humans have been identified in the CYP isozymes thought to be responsible for TCE metabolism (Pastino et al., 2000), but no data exist correlating these polymorphisms with enzyme activity. It is relevant to note that repeat polymorphism (Hu et al., 1999) or polymorphism in the regulatory sequence (McCarver et al., 1998) were not involved in the constitutive expression of human CYP2E1; however, it is unknown if these types of polymorphisms may play a role in the inducibility of the respective gene.

**Table 3-21. P450 isoform activities in human liver microsomes exhibiting different affinities for TCE**

*This document is a draft for review purposes only and does not constitute Agency policy.*

1

Affinity group	CYP isoform activity (pmol/min/mg protein)		
	CYP2E1	CYP1A2	CYP3A4
Low $K_M$	520 $\pm$ 295	241 $\pm$ 146	2.7 $\pm$ 2.7
Mid $K_M$	820 $\pm$ 372	545 $\pm$ 200	2.9 $\pm$ 2.8
High $K_M$	1,317 $\pm$ 592	806 $\pm$ 442	1.8 $\pm$ 1.1

2

3 Activities of CYP1A2, CYP2E1, and CYP3A4 were measured with phenacetin, chlorzoxazone, and testosterone as  
4 substrates, respectively. Data are means  $\pm$  standard deviation from 10, 9, and four samples for the low-, mid-, and  
5 high- $K_M$  groups, respectively. Only CYP3A4 activities are not significantly different ( $p < 0.05$ ) from one another  
6 by Kruskal-Wallis one-way analysis of variance.

7 Source: Lash et al. (2000a).

8

9

10 Individual susceptibilities to TCE toxicity may also result from variations in enzyme  
11 content, either at baseline or due to enzyme induction/inhibition, which can lead to alterations in  
12 the amounts of metabolites formed. Certain physiological and pathological conditions or  
13 exposure to other chemicals (e.g., ethanol and acetaminophen) can induce, inhibit, or compete  
14 for enzymatic activity. Given the well established (or characterized) role of the liver to  
15 oxidatively metabolize TCE (by CYP2E1), increasing the CYP2E1 content or activity (e.g., by  
16 enzyme induction) may not result in further increases in TCE oxidation. Indeed, Kaneko et al.  
17 (1994a) reported that enzyme induction by ethanol consumption in humans increased TCE  
18 metabolism only at high concentrations (500 ppm, 2,687 mg/m<sup>3</sup>) in inspired air. However, other  
19 interactions between ethanol and the enzymes that oxidatively metabolize TCE metabolites can  
20 result in altered metabolic fate of TCE metabolites. In addition, enzyme inhibition or  
21 competition can decrease TCE oxidation and subsequently alter the TCE toxic response via, for  
22 instance, increasing the proportion undergoing GSH conjugation Lash et al. (2000a). TCE itself  
23 is a competitive inhibitor of CYP2E1 activity (Lipscomb et al., 1997), as shown by reduced  
24 *p*-nitrophenol hydroxylase activity in human liver microsomes, and so may alter the toxicity of  
25 other chemicals metabolized through that pathway. On the other hand, suicidal CYP heme  
26 destruction by the TCE-oxygenated CYP intermediate has also been shown (Miller and  
27 Guengerich, 1983).

28

### 3.3.3.1.8. Glutathione (GSH) Conjugation Pathway

29 Historically, the conjugative metabolic pathways have been associated with xenobiotic  
30 detoxification. This is true for GSH conjugation of many compounds. However, several

*This document is a draft for review purposes only and does not constitute Agency policy.*


1 halogenated alkanes and alkenes, including TCE, are bioactivated to cytotoxic metabolites by the  
2 GSH conjugate processing pathway (mercapturic acid) pathways (Elfarra et al., 1987; Elfarra et  
3 al., 1986). In the case of TCE, production of reactive species several steps downstream from the  
4 initial GSH conjugation is believed to cause cytotoxicity and carcinogenicity, particularly in the  
5 kidney. Since the GSH conjugation pathway is in competition with the P450 oxidative pathway  
6 for TCE biotransformation, it is important to understand the role of various factors in  
7 determining the flux of TCE through each pathway. Figure 3-5 depicts the present  
8 understanding of TCE metabolism via GSH conjugation.  
9

### 3.3.3.1.9. Formation of S-(1,2-dichlorovinyl)glutathione or S-(2,2-dichlorovinyl)glutathione (DCVG)

10 The conjugation of TCE to GSH produces S-(1,2-dichlorovinyl)glutathione or its isomer  
11 S-(2,2-dichlorovinyl)glutathione (DCVG). There is some uncertainty as to which GST isoforms  
12 mediate TCE conjugation. Lash and colleagues studied TCE conjugation in renal tissue  
13 preparations, isolated renal tubule cells from male F344 rats and purified GST alpha-class  
14 isoforms 1-1, 1-2 and 2-2 (Cummings et al., 2000a; Cummings and Lash, 2000; Lash et al.,  
15 2000b). The results demonstrated high conjugative activity in renal cortex and in proximal  
16 tubule cells. Although the isoforms studied had similar  $V_{MAX}$  values, the  $K_M$  value for GST 2-2  
17 was significantly lower than the other forms, indicating that this form will catalyze TCE  
18 conjugation at lower (more physiologically relevant) substrate concentrations. In contrast, using  
19 purified rat and human enzymes, Hissink et al. (2002) reported in vitro activity for DCVG  
20 formation only for mu- and pi-class GST isoforms, and none towards alpha-class isoforms;  
21 however, the rat mu-class GST 3-3 was several folds more active than the human mu-class  
22 GST M1-1. Although GSTs are present in tissues throughout the body, the majority of TCE

23 GSH conjugation is thought to occur in the liver (Lash et al., 2000a). Using in vitro  
24 studies with renal preparations, it has been demonstrated that GST catalyzed conjugation of TCE  
25 is increased following the inhibition of CYP-mediated oxidation (Cummings and Lash, 2000).

26 In F344 rats, following gavage doses of 263–1,971 mg/kg TCE in 2 mL corn oil, DCVG  
27 was observed in the liver and kidney of females only, in blood of both sexes (Lash et al., 2006),  
28 and in bile of males (Dekant, 1990). The data from Lash et al. (2006) are difficult to interpret  
29 because the time courses seem extremely erratic, even for the oxidative metabolites TCOH and  
30 TCA. Moreover, a comparison of blood levels of TCA and TCOH with other studies  
31



1  
2  
3 **Figure 3-5. Scheme for GSH-dependent metabolism of TCE.**

4  
5 Adapted from: Lash et al. (2000a); Cummings and Lash (2000); NRC (2006).

6  
7  
8 in rats at similar doses reveals differences of over 1,000-fold in reported concentrations. For  
9 instance, at the lowest dose of 263 mg/kg, the peak blood levels of TCE and TCA in male F344  
10 rats were 10.5 and 1.6 µg/L, respectively (Lash et al., 2006). By contrast, Larson and Bull  
11 (1992a) reported peak blood TCE and TCA levels in male Sprague-Dawley rats over 1,000-fold  
12 higher—around 10 and 13 mg/L, respectively—following oral doses of 197 mg/kg as a  
13 suspension in 1% aqueous Tween 80<sup>®</sup>. The results of Larson and Bull (1992a) are similar to

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Lee et al. (2000b), who reported peak blood TCE levels of 20–50 mg/L after male  
2 Sprague-Dawley rats received oral doses of 144–432 mg/kg in a 5% aqueous Alkamus emulsion  
3 (polyethoxylated vegetable oil), and to Stenner et al. (1997), who reported peak blood levels of  
4 TCA in male F344 rats of about 5 mg/L at a slightly lower TCE oral dose of 100 mg/kg  
5 administered to fasted animals in 2% Tween 80<sup>®</sup>. Thus, while useful qualitatively as an  
6 indicator of the presence of DCVG in rats, the quantitative reliability of reported concentrations,  
7 for metabolites of either oxidation or GSH conjugation, may be questionable.

8 In humans, DCVG was readily detected at in human blood following onset of a 4-hour  
9 TCE inhalation exposure to 50 or 100 ppm (269 or 537 mg/m<sup>3</sup>) (Lash et al., 1999a). At 50 ppm,  
10 peak blood levels ranged from 2.5–30 μM, while at 100 ppm, the mean ( $\pm$  SE,  $n = 8$ ) peak blood  
11 levels were  $46.1 \pm 14.2$  μM in males and  $13.4 \pm 6.6$  μM in females. Although on average, male  
12 subjects had threefold higher peak blood levels of DCVG than females, DCVG blood levels in  
13 half of the male subjects were similar to or lower than those of female subjects. This suggests a  
14 polymorphism in GSH conjugation of TCE rather than a true gender difference (Lash et al.,  
15 1999a) as also has been indicated by Hissink et al. (2002) for the human mu-class GST M1-1  
16 enzyme. Interestingly, as shown in Table 3-22, the peak blood levels of DCVG are similar on a  
17 molar basis to peak levels of TCE, TCA, and TCOH in the same subjects, as reported in  
18 Fisher et al. (1998).

19 Tables 3-23–3-25 summarize DCVG formation from TCE conjugation from in vitro  
20 studies of liver and kidney cellular and subcellular fractions in mouse, rat, and human  
21 (tissue-distribution and species-and gender-differences in DCVG formation are discussed  
22 below). As shown by these tables, different investigators have reported considerably different  
23 rates for TCE conjugation in human liver and kidney cell fractions. For instance, values in  
24 Table 3-23 from Lash et al. (1999a) are between two and five orders of magnitude higher than  
25 those reported by Green et al. (1997a) or Dekant et al. (1990) (see Table 3-25). In addition,  
26 Green et al. (1997a) and Dekant et al. (1990) reported a difference in the relative importance of  
27 rat liver cytosol and rat liver microsomes for GSH conjugation, with Green et al. (1997a)  
28 reporting activity in the cytosol and none in the microsomes and Dekant et al. (1990) reporting  
29 the opposite.

30 The reasons for such discrepancies are unclear, but they may be related to different  
31 analytical methods (Lash et al., 2000a). In particular, Lash et al. (1999a) employed the “Reed  
32 method,” which used ion-exchange high-performance liquid chromatography (HPLC) of  
33 derivatized analytes. This HPLC method is characterized by variability and an overall decline in  
34 retention times over the life of the HPLC column due to derivatization of amine groups on the  
35

1  
2  
3

**Table 3-22. Comparison of peak blood concentrations in humans exposed to 100 ppm (537 mg/m<sup>3</sup>) TCE for 4 hours (Fisher et al., 1998; Lash et al., 1999a)**

Chemical species	Peak blood concentration (mean $\pm$ SD, $\mu$ M)	
	Males	Females
TCE	23 $\pm$ 11	14 $\pm$ 4.7
TCA	56 $\pm$ 9.8	59 $\pm$ 12
TCOH	21 $\pm$ 5.0	15 $\pm$ 5.6
DCVG	46.1 $\pm$ 14.2	13.4 $\pm$ 6.6

4  
5  
6

1  
2  
3  
4

**Table 3-23. GSH conjugation of TCE (at 1–2 mM) in liver and kidney cellular fractions in humans, male F344 rats, and male B6C3F1 mice from Lash laboratory**

Species and cellular/subcellular fraction (TCE concentration)	DCVG formation (nmol/hour/mg protein or 10 <sup>6</sup> cells)	
	Male	Female
Human		
Hepatocytes (0.9 mM) [pooled]	11 ± 3	
Liver cytosol (1 mM) [individual samples]	156 ± 16	174 ± 13
Liver cytosol (2 mM) [pooled]	346	
Liver microsomes (1 mM) [individual samples]	108 ± 24	83 ± 11
Liver microsomes (1 mM) [pooled]	146	
Kidney cytosol (2 mM) [pooled]	42	
Kidney microsomes (1 mM) [pooled]	320	
Rat		
Liver cytosol (2 mM)	7.30 ± 2.8	4.86 ± 0.14
Liver microsomes (2 mM)	10.3 ± 2.8	7.24 ± 0.24
Kidney cortical cells (2 mM)	0.48 ± 0.02	0.65 ± 0.15
Kidney cytosol (2 mM)	0.45 ± 0.22	0.32 ± 0.02
Kidney microsomes (2 mM)	not detected	0.61 ± 0.06
Mouse		
Liver cytosol (2 mM)	24.5 ± 2.4	21.7 ± 0.9
Liver microsomes (2 mM)	40.0 ± 3.1	25.6 ± 0.8
Kidney cytosol (2 mM)	5.6 ± 0.24	3.7 ± 0.48
Kidney microsomes (2 mM)	5.47 ± 1.41	16.7 ± 4.7

5  
6  
7  
8

Mean ± SE. Source: Lash et al. (1999b; 1998a; 1995); Cummings and Lash (2000); Cummings et al. (2000a).

1 **Table 3-24. Kinetics of TCE metabolism via GSH conjugation in male F344**  
 2 **rat kidney and human liver and kidney cellular and subcellular fractions**  
 3 **from Lash laboratory**  
 4

Tissue and cellular fraction	$K_M$ ( $\mu\text{M TCE}$ )	$V_{MAX}$ (nmol DCVG/min/mg protein or $10^6$ hepatocytes)	$1,000 \times$ $V_{MAX}/K_M$
Rat			
Kidney proximal tubular cells: low affinity	2,910	0.65	0.22
Kidney proximal tubular cells: high affinity	460	0.47	1.0
Human			
Liver hepatocytes <sup>a</sup>	37~106	0.16~0.26	2.4~4.5
Liver cytosol: low affinity	333	8.77	2.6
Liver cytosol: high affinity	22.7	4.27	190
Liver microsomes: low affinity	250	3.1	12
Liver microsomes: high affinity	29.4	1.42	48
Kidney proximal tubular cells: low affinity	29,400	1.35	0.046
Kidney proximal tubular cells: high affinity	580	0.11	0.19
Kidney cytosol	26.3	0.81	31
Kidney microsomes	167	6.29	38

5  
 6 <sup>a</sup> Kinetic analyses of first six–nine (out of 10) data points from Figure 1 from Lash et al. (1999a) using  
 7 Lineweaver-Burk or Eadie-Hofstee plots and linear regression ( $R^2 = 0.50\sim 0.95$ ). Regression with best  $R^2$  used  
 8 first 6 data points and Eadie-Hofstee plot, with resulting  $K_M$  and  $V_{MAX}$  of 106 and 0.26, respectively.  
 9

10 Source: Lash et al. (1999a); Cummings and Lash (2000); Cummings et al. (2000a).  
 11



1  
2  
3  
4

**Table 3-25. GSH conjugation of TCE (at 1.4–4 mM) in liver and kidney cellular fractions in humans, male F344 rats, and male B6C3F1 mice from Green and Dekant laboratories**

Species and cellular/subcellular fraction (TCE concentration)	DCVG formation (nmol/hour/mg protein) [substrate concentration in mM]	
	Dekant et al. (1990)	Green et al. (1997a)
Human		
Liver cytosol	-	0.00019 ± 0.00014
Liver microsomes	-	n.d.
Kidney cytosol	-	n.d.
Kidney microsomes	-	n.d.
Rat		
Liver cytosol	<0.002	0.00162 ± 0.00002
Liver microsomes	0.002	n.d.
Kidney cytosol	-	n.d.
Kidney microsomes	-	n.d.
Mouse		
Liver cytosol	-	0.0025
Liver microsomes	-	n.d.
Kidney cytosol	-	n.d.
Kidney microsomes	-	n.d.

5  
6  
7  
8  
9

n.d. = not determined  
where available, mean ± SD. Source: Dekant et al. (1990), Green et al. (1997a)

1 column (Lash et al., 1999b). Although data are limited, the GSH pathway metabolite levels  
2 reported by methods that utilize <sup>14</sup>C TCE and radiochemical detection followed by mass  
3 spectrometry identification of the metabolites are lower. In particular, Green et al. (1997a) and  
4 Dekant et al. (1990) both used HPLC with radiochemical detection. Peak identity was confirmed  
5 by Green et al. (1997a) using liquid chromatography/mass spectrometry (LC/MS) and by  
6 GC/MS following hydrolysis by Dekant et al. (1990). In addition, studies using HPLC-MS/MS  
7 techniques with stable isotope-labeled DCVG and dichlorovinyl cysteine (DCVC) standards  
8 have also been used to detect GSH pathway metabolite levels Kim et al. (2009). Based on the in  
9 vitro work presented in Table 3-23 using the “Reed method,” one would expect mouse serum  
10 DCVG levels to be ~4-6 times lower than humans. However, using the HPLC-MS/MS  
11 technique of Kim et al. (2009), the peak DCVG serum levels are ~1,000 times lower in mouse  
12 serum than determined by Lash et al. (1999a) in human serum. Although advances in LC  
13 technology, and differences in exposure routes (inhalation versus oral, with different first pass),  
14 exposure doses, and the degree of competition with TCE oxidation (greater in mouse than in  
15 human) should be considered, this much-larger-than-expected difference is consistent with the  
16 suggestion that the “Reed method” provides an overestimation of DCVG levels in humans. This  
17 could occur if the “Reed method” identifies nonspecific derivatives as DCVG or other GSH  
18 pathway metabolites. However, the degree of overestimation is unclear, and differing results in  
19 humans may be attributable to true interindividual variation (especially since GSTs are known to  
20 be polymorphic). However, overall, there remains significant uncertainty in the quantitative  
21 estimation of DCVG formation from TCE both in vivo and in vitro.  
22

#### **3.3.3.1.10. Formation of S-(1,2-dichlorovinyl) cysteine or S-(2,2-dichlorovinyl) cysteine (DCVC)**

23 The cysteine conjugate, isomers S-(1,2-dichlorovinyl) cysteine (1,2-DCVC) or  
24 S-(2,2-dichlorovinyl) cysteine (2,2-DCVC), is formed from DCVG in a two-step sequence.  
25 DCVG is first converted to the cysteinylglycine conjugate  
26 S-(1,2-dichlorovinyl)-L-cysteinylglycine or its isomer S-(2,2-dichlorovinyl)-L-cysteinylglycine  
27 by  $\gamma$ -glutamyl transpeptidase (GGT) in the renal brush border (Elfarra and Anders, 1984; Lash et  
28 al., 1988).

29 Cysteinylglycine dipeptidases in the renal brush border and basolateral membrane  
30 convert DCVG to DCVC via glycine cleavage (Goeptar et al., 1995; Lash et al., 1995). This  
31 reaction can also occur in the bile or gut, as DCVG excreted into the bile is converted to DCVC  
32 and reabsorbed into the liver where it may undergo further acetylation.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1

#### **3.3.3.1.11. Formation of N-Acetyl-S-(1,2-dichlorovinyl)-L-cysteine or N-Acetyl-S-(2,2-dichlorovinyl)-L-cysteine (NAcDCVC)**

2 N-acetylation of DCVC can either occur in the kidney, as demonstrated in rat kidney  
3 microsomes (Duffel and Jakoby, 1982), or in the liver (Birner et al., 1997). Subsequent release  
4 of DCVC from the liver to blood may result in distribution to the kidney resulting in increased  
5 internal kidney exposure to the acetylated metabolite over and above what the kidney already is  
6 capable of generating. In the kidney, N-Acetyl-S-(1,2-dichlorovinyl)-L-cysteine or N-Acetyl-S-  
7 (2,2-dichlorovinyl)-L-cysteine (NAcDCVC) may undergo deacetylation, which is considered a  
8 rate-limiting-step in the production of proximal tubule damage (Wolfgang et al., 1989b; Zhang  
9 and Stevens, 1989). As a polar mercapturate, NAcDCVC may be excreted in the urine as  
10 evidenced by findings in mice (Birner et al., 1993), rats (Bernauer et al., 1996; Commandeur and  
11 Vermeulen, 1990), and humans who were exposed to TCE (Bernauer et al., 1996; Birner et al.,  
12 1993), suggesting a common glutathione-mediated metabolic pathway for DCVC among species.  
13

#### **3.3.3.1.12. Beta lyase metabolism of S-(1,2-dichlorovinyl) cysteine (DCVC)**

14 The enzyme cysteine conjugate  $\beta$ -lyase catalyzes the breakdown of DCVC to reactive  
15 nephrotoxic metabolites (Goepfert et al., 1995). This reaction involves removal of pyruvate and  
16 ammonia and production of S-(1,2-dichlorovinyl) thiol (DCVT), an unstable intermediate, which  
17 rearranges to other reactive alkylation metabolites that form covalent bonds with cellular  
18 nucleophiles (Dekant et al., 1988; Goepfert et al., 1995). The rearrangement of DCVT to  
19 enethiols and their acetylating agents has been described in trapping experiments (Dekant et al.,  
20 1988) and proposed to be responsible for nucleophilic adduction and toxicity in the kidney. The  
21 quantification of acid-labile adducts was proposed as a metric for TCE flux through the GSH  
22 pathway. However, the presence of analytical artifacts precluded such analysis. In fact,  
23 measurement of acid-labile adduct products resulted in higher values in mice than in rats (Eyre et  
24 al., 1995a, b).

25 DCVC metabolism to reactive species via a  $\beta$ -lyase pathway has been observed in vitro  
26 by Green et al. (1997a), who reported greater  $\beta$ -lyase activity in rats than in mice or humans.  
27 However, in vitro DCVC metabolism by the competing enzyme *N*-acetyl transferase was also  
28 reported to be greater in rats than mice and humans. In vivo,  $\beta$ -lyase activity in humans and rats  
29 (reaction rates were not reported) was demonstrated using a surrogate substrate,  
30 2-(fluoromethoxy)-1,1,3,3,3-pentafluoro-1-propene (Iyer et al., 1998).  $\beta$ -lyase -mediated

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 reactive adducts have been described in several extrarenal tissues, including rat and human liver  
2 and intestinal microflora (Dohn and Anders, 1982; Larsen and Stevens, 1986; Stevens, 1985;  
3 Stevens and Jakoby, 1983; Tateishi et al., 1978; Tomisawa et al., 1986; Tomisawa et al.,  
4 1984) and rat brain (Alberati-Giani et al., 1995; Malherbe et al., 1995).

5 In the kidneys, glutamine transaminase K appears to be primarily responsible for  $\beta$ -lyase  
6 metabolism of DCVC (Jones et al., 1988; Lash et al., 1986; Lash et al., 1990b; Perry et al., 1993;  
7 Stevens et al., 1988; Stevens et al., 1986).  $\beta$ -lyase transformation of DCVC appears to be  
8 regulated by 2-keto acids. DCVC toxicity in isolated rat proximal tubular cells was significantly  
9 increased with the addition of  $\alpha$ -keto- $\gamma$ -methiolbutyrate or phenylpyruvate (Elfarra et al., 1986).  
10 The presence of  $\alpha$ -keto acid cofactors is necessary to convert the inactive form of the  $\beta$ -lyase  
11 enzyme (containing pyridoxamine phosphate) to the active form (containing pyridoxal  
12 phosphate) (Goeptar et al., 1995).

13 Both low- and high-molecular-weight enzymes with  $\beta$ -lyase activities have been  
14 identified in rat kidney cytosol and mitochondria (Abraham et al., 1995a; Abraham et al., 1995b;  
15 Lash et al., 1986; Stevens et al., 1988). While glutamine transaminase K and  
16 kynureninase-associated  $\beta$ -lyase activities have been identified in rat liver (Alberati-Giani et al.,  
17 1995; Stevens, 1985), they are quite low compared to renal glutamine transaminase K activity  
18 and do not result in hepatotoxicity in DCVG- or DCVC-treated rats (Elfarra and Anders, 1984).  
19 Similar isoforms of  $\beta$ -lyase have also been reported in mitochondrial fractions of brain tissue  
20 (Cooper, 2004).

21 The kidney enzyme L- $\alpha$ -hydroxy (L-amino) acid oxidase is capable of forming an  
22 iminium intermediate and keto acid analogues (pyruvate or  
23 S-(1,2-dichlorovinyl)-2-oxo-3-mercaptpropionate) of DCVC, which decomposes to  
24 dichlorovinylthiol (Lash et al., 1990a; Stevens et al., 1989). In rat kidney homogenates, this  
25 enzyme activity resulted in as much as 35% of GSH pathway-mediated bioactivation. However,  
26 this enzyme is not present in humans, an important consideration for extrapolation of renal  
27 effects across species.

28

### **3.3.3.1.13. Sulfoxidation of S-(1,2-dichlorovinyl) cysteine (DCVC) and N-Acetyl-S-(1,2-dichlorovinyl)-L-cysteine (NAcDCVC)**

29 A second pathway for bioactivation of TCE S-conjugates involves sulfoxidation of either  
30 the cysteine or mercapturic acid conjugates (Lash et al., 2003; Park et al., 1992; Sausen and  
31 Elfarra, 1990) (Birner et al., 1998; Krause et al., 2003; Lash et al., 1994; Werner et al., 1995a,  
32 1996; Werner et al., 1995b). Sulfoxidation of DCVC was mediated mainly by flavin


1 monooxygenase 3 (FMO3), rather than CYP, in rabbit liver microsomes (Ripp et al., 1997) and  
2 human liver microsomes (Krause et al., 2003). Krause et al., (2003) also reported DCVC  
3 sulfoxidation by human cDNA-expressed FMO3, as well as detection of FMO3 protein in human  
4 kidney samples. While Krause et al. (2003) were not able to detect sulfoxidation in human  
5 kidney microsomes, the authors noted FMO3 expression in the kidney was lower and more  
6 variable than that in the liver. However, sulfoxidation products in tissues or urine have not been  
7 reported in vivo.

8 Sulfoxidation of NAcDCVC, by contrast, was found to be catalyzed predominantly, if not  
9 exclusively, by CYP3A enzymes (Werner et al., 1996), whose expressions are highly  
10 polymorphic in humans. Sulfoxidation of other haloalkyl mercapturic acid conjugates has also  
11 been shown to be catalyzed by CYP3A (Altuntas et al., 2004; Werner et al., 1995a; Werner et al.,  
12 1995b). While Lash et al. (2000a) suggested that this pathway would be quantitatively minor  
13 because of the relatively low CYP3A levels in the kidney, no direct data exist to establish the  
14 relative toxicological importance of this pathway relative to bioactivation of DCVC by  $\beta$ -lyase or  
15 FMO3. However, the contribution of CYP3A in S-conjugate sulfoxidation to nephrotoxicity in  
16 vivo was recently demonstrated by Sheffels et al. (2004) with  
17 fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether (FDVE). In particular, in vivo  
18 production and urinary excretion of FDVE-mercapturic acid sulfoxide metabolites were  
19 unambiguously established by mass spectrometry, and CYP inducers/inhibitors  
20 increased/decreased nephrotoxicity in vivo while having no effect on urinary excretion of  
21 metabolites produced through  $\beta$ -lyase (Sheffels et al., 2004). These data suggest that, by  
22 analogy, sulfoxidation of NAcDCVC may be an important bioactivating pathway.

23

#### **3.3.3.1.14. Tissue distribution of glutathione (GSH) metabolism**

24 The sites of enzymatic metabolism of TCE to the various GSH pathway-mediated  
25 metabolites are significant in determining target tissue toxicity along this pathway. Figure 3-6  
26 presents a schematic of interorgan transport and metabolism of TCE along the glutathione  
27 pathway. TCE is taken up either by the liver or kidney and conjugated to DCVG. The primary  
28 factors affecting TCE flux via this pathway include high hepatic GST activity, efficient transport  
29 of DCVG from the liver to the plasma or bile, high renal brush border and low hepatic GGT  
30 activities, and the capability for GSH conjugate uptake into the renal basolateral membranes with  
31 limited or no uptake into liver cell plasma membranes.



1 As discussed previously, GST activity is present in many different cell types. However,  
 2 the liver is the major tissue for GSH conjugation. GST activities in rat and mouse cytosolic  
 3 fractions were measured using 1-chloro-2,4-dinitrobenzene, a GST substrate that is nonspecific

4  
 5 **Figure 3-6. Interorgan TCE transport and metabolism via the GSH pathway.** See  
 6 Figure 3-5 for enzymes involved in metabolic steps. Source: Lash et al.(2000a; 2000b);  
 7 NRC (2006).  
 8

9  
 10 for particular isoforms (Lash et al., 1998b). Specific activities (normalized for protein content)  
 11 in whole kidney cytosol were slightly less than those in the liver (0.64 compared to 0.52 mU/mg  
 12 protein for males and females). However, the much larger mass of the liver compared to the  
 13 kidney indicates that far more total GST activity resides in the liver. This is consistent with in  
 14 vitro data on TCE conjugation to DCVG, discussed previously (see Tables 3-23 and 3-24). For

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 instance, in humans, rats, and mice, liver cytosol exhibits greater DCVG production than kidney  
2 cytosol. Distinct high- and low-affinity metabolic profiles were observed in the liver but not in  
3 the kidney (see Table 3-24). In microsomes, human liver and kidney had similar rates of DCVG  
4 production, while for rats and mice, the production in the liver was substantially greater.  
5 According to studies by Lash et al.(1998a; 1998b), the activity of GGT, the first step in the  
6 conversion of DCVG to DCVC, is much higher in the kidney than the liver of mice, rats, and  
7 humans, with most of the activity being concentrated in the microsomal, rather than the  
8 cytosolic, fraction of the cell (see Table 3-26). In rats, this activity is quite high in the kidney but  
9 is below the level of detection in the liver while the relative kidney to liver levels in humans and  
10 mice were higher by 18- and up to 2,300-fold, respectively. Similar qualitative findings were  
11 also reported in another study (Hinchman and Ballatori, 1990) when total organ GGT levels were  
12 compared in several species (see Table 3-27). Cysteinylglycine dipeptidase was also  
13 preferentially higher in the kidney than the liver of all tested species although the interorgan  
14 differences in this activity (one–ninefold) seemed to be less dramatic than for GGT (see  
15 Table 3-27). High levels of both GGT and dipeptidases have also been reported in the small  
16 intestine of rat (Kozak and Tate, 1982) and mouse (Habib et al., 1996), as well as GGT in the  
17 human jejunum (Fairman et al., 1977). No specific human intestinal cysteinylglycine  
18 dipeptidase has been identified; however, a related enzyme (EC 3.4.13.11) from human kidney  
19 microsomes has been purified and studied (Adachi et al., 1989) while several human intestinal  
20 dipeptidases have been characterized including a membrane dipeptidase (EC 3.4.13.19) which  
21 has a wide dipeptide substrate specificity including cysteinylglycine (Hooper et al., 1994; Ristoff  
22 and Larsson, 2007).

23

#### **3.3.3.1.15. Sex- and species-dependent differences in glutathione (GSH) metabolism**

24 Diverse sex and species differences appear to exist in TCE metabolism via the  
25 glutathione pathway. In rodents, rates of TCE conjugation to GSH in male rats and mice are  
26 higher than females (see Table 3-23). Verma and Rana (2003) reported twofold higher GST  
27 activity values in liver cytosol of female rats, compared to males, given 15 intraperitoneal  
28 injections of TCE over 30 days period. This effect may be due to sex-dependent variation in  
29 induction, as GST activities in male and female controls were similar. DCVG formation rates by  
30 liver and kidney subcellular fractions were much higher in both sexes of mice than in rats and,  
31 except for mouse kidney microsomes, the rates were generally higher in males than in females of  
32 the same species (see Table 3-23).

1            In terms of species differences, comparisons at 1–2 mM TCE concentrations (see  
2 Table 3-23) suggest that, in liver and kidney cytosol, the greatest DCVG production rate was in  
3 humans, followed by mice and then rats. However, different investigators have reported  
4 considerably different rates for TCE conjugation in human liver and kidney cell fractions. For  
5 instance,



1  
2  
3

**Table 3-26. GGT activity in liver and kidney subcellular fractions of mice, rats, and humans**

Species	Sex	Tissue	Cellular fraction	Activity (mU/mg)
Mouse	Male	Liver	Cytosol	0.07 ± 0.04
			Microsomes	0.05 ± 0.04
		Kidney	Cytosol	1.63 ± 0.85
			Microsomes	92.6 ± 15.6
	Female	Liver	Cytosol	0.10 ± 0.10
			Microsomes	0.03 ± 0.03
		Kidney	Cytosol	0.79 ± 0.79
			Microsomes	69.3 ± 14.0
Rat	Male	Liver	Cytosol	<0.02
			Microsomes	<0.02
		Kidney	Cytosol	<0.02
			Microsomes	1,570 ± 100
	Female	Liver	Cytosol	<0.02
			Microsomes	<0.02
		Kidney	Cytosol	<0.02
			Microsomes	1,840 ± 40
Human	Male	Liver	Cytosol	8.89 ± 3.58
			Microsomes	29
		Kidney	Cytosol	13.2 ± 1.0
			Microsomes	960 ± 77

4  
5  
6  
7  
8  
9

Source: Lash et al. (1999b; 1998a)

1 **Table 3-27. Multispecies comparison of whole-organ activity levels of GGT**  
 2 **and dipeptidase**  
 3

Species	Whole organ enzyme activity (μmol substrate/organ)			
	Kidney		Liver	
	GGT	Dipeptidase	GGT	Dipeptidase
Rat	1,010 ± 41	20.2 ± 1.1	7.1 ± 1.4	6.1 ± 0.4
Mouse	60.0 ± 4.2	3.0 ± 0.3	0.47 ± 0.05	1.7 ± 0.2
Rabbit	1,119 ± 186	112 ± 17	71.0 ± 9.1	12.6 ± 1.0
Guinea pig	148 ± 13	77 ± 10	46.5 ± 4.2	13.2 ± 1.5
Pig	3,800 ± 769	2,428 ± 203	1,600 ± 255	2,178 ± 490
Macaque	988	136	181	71

4  
 5 Source: Hinchman and Ballatori (1990).  
 6  
 7

8 values in Table 3-23 from Lash et al. (1999a) are between two and five orders of magnitude  
 9 higher than those reported by Green et al. (1997a). (The rates of DCVG formation by liver  
 10 cytosol from male F344 rat, male B6C3F1 mouse, and human were 1.62, 2.5, and  
 11 0.19 pmol/minute/mg protein, respectively, while there was no measurable activity in liver  
 12 microsomes or subcellular kidney fractions (Green et al., 1997a). The reasons for such  
 13 discrepancies are unclear but may be related to different analytical methods employed such as  
 14 detection of radiolabeled substrate vs. derivatized analytes (Lash et al., 2000a).

15 Expression of GGT activity does not appear to be influenced by sex (see Table 3-26); but  
 16 species differences in kidney GGT activity are notable with rat subcellular fractions exhibiting  
 17 the highest levels and mice and humans exhibiting about 4–6% and 50%, respectively, of rat  
 18 levels (Lash et al., 1999b; Lash et al., 1998a). Table 3-27 shows measures of whole-organ GGT  
 19 and dipeptidase activities in rats, mice, guinea pigs, rabbits, pigs, and monkeys. These data show  
 20 that the whole kidney possesses higher activities than liver for these enzymes, despite the  
 21 relatively larger mass of the liver.

22 As discussed above, the three potential bioactivating pathways subsequent to the  
 23 formation of DCVC are catalyzed by β-lyase, FMO3 or CYP3A. Lash et al. (2000a) compared  
 24 in vitro β-lyase activities and kinetic constants (when available) for kidney of rats, mice, and  
 25 humans. They reported that variability of these values spans up to two orders of magnitude  
 26 depending on substrate, analytical method used, and research group. Measurements of rat,

1 mouse, and human  $\beta$ -lyase activities collected by the same researchers following  
2 tetrachloroethylene exposure (Green et al., 1990) resulted in higher  $K_M$  and lower  $V_{MAX}$  values  
3 for mice and humans than rats. Further, female rats exhibited higher  $K_M$  and lower  $V_{MAX}$  values  
4 than males.

5 With respect to FMO3, Ripp et al. (1999) found that this enzyme appeared catalytically  
6 similar across multiple species, including humans, rats, dogs, and rabbits, with respect to several  
7 substrates, including DCVC, but that there were species differences in expression. Specifically,  
8 in male liver microsomes, rabbits had threefold higher methionine S-oxidase activity than mice  
9 and dogs had 1.5-fold higher activity than humans and rats. Species differences were also noted  
10 in male and female kidney microsomes; rats exhibited two- to sixfold higher methionine  
11 S-oxidase activity than the other species. Krause et al. (2003) detected DCVC sulfoxidation in  
12 incubations with human liver microsomes but did not in an incubation with a single sample of  
13 human kidney microsomes. However, FMO3 expression in the 26 human kidney samples was  
14 found to be highly variable, with a range of five–sixfold (Krause et al., 2003).

15 No data on species differences in CYP3A-mediated sulfoxidation of NAcDCVC are  
16 available. However, Altuntas et al. (2004) examined sulfoxidation of cysteine and mercapturic  
17 acid conjugates of FDVE (fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether) in rat and  
18 human liver and kidney microsomes. They reported that the formation of sulfoxides from the  
19 mercapturates *N*-Ac-FFVC and (*Z*)-*N*-Ac-FFVC (FFVC is  
20 (*E,Z*)-S-(1-fluoro-2-fluoromethoxy-2-(trifluoromethyl)vinyl-L-cysteine) were greatest in rat liver  
21 microsomes, and 2- to 30-fold higher than in human liver microsomes (which had high  
22 variability). Sulfoxidation of *N*-Ac-FFVC could not be detected in either rat or human kidney  
23 microsomes, but sulfoxidation of (*Z*)-*N*-Ac-FFVC was detected in both rat and human kidney  
24 microsomes at rates comparable to human liver microsomes. Using human- and rat-expressed  
25 CYP3A, Altuntas et al. (2004) reported that rates of sulfoxidation of (*Z*)-*N*-Ac-FFVC were  
26 comparable in human CYP3A4 and rat CYP3A1 and CYP3A2, but that only rat CYP3A1 and  
27 A2 catalyzed sulfoxidation of *N*-Ac-FFVC. As the presence or absence of the species  
28 differences in mercapturate sulfoxidation appears to be highly chemical-specific, no clear  
29 inferences can be made as to whether species differences exist for sulfoxidation of NAcDCVC

30 Also relevant to assess the flux through the various pathways are the rates of  
31 *N*-acetylation and de-acetylation of DCVC. This is demonstrated by the results of Elfarra and  
32 Hwang (1990) using S-(2-benzothiazolyl)-L-cysteine as a marker for  $\beta$ -lyase metabolism in rats,  
33 mice, hamsters, and guinea pigs. Guinea pigs exhibited about twofold greater flux through the  
34  $\beta$ -lyase pathway, but this was not attributable to higher  $\beta$ -lyase activity. Rather, guinea pigs  
35 have relatively low *N*-acetylation and high deacetylation activities, leading to a high level of

1 substrate recirculation (Lau et al., 1995). Thus, a high *N*-deacetylase:*N*-acetylase activity ratio  
2 may favor DCVC recirculation and subsequent metabolism to reactive species. In human,  
3 Wistar rat, Fischer rat, and mouse cytosol, deacetylation rates for NAcDCVC varied less than  
4 threefold (0.35, 0.41, 0.61, and 0.94 nmol DCVC formed/minute/mg protein in humans, rats, and  
5 mice) (Birner et al., 1993). However, similar experiments have not been carried out for  
6 *N*-acetylation of DCVC, so the balance between its *N*-acetylation and de-acetylation has not been  
7 established.

#### **3.3.3.1.16. Human variability and susceptibility in glutathione (GSH) conjugation**

8 Knowledge of human variability in metabolizing TCE through the glutathione pathway is  
9 limited to in vitro comparisons of variance in GST activity rates. Unlike CYP-mediated  
10 oxidation, quantitative differences in the polymorphic distribution or activity levels of GST  
11 isoforms in humans are not presently known. However, the available data (Lash et al., 1999a;  
12 Lash et al., 1999b) do suggest that significant variation in GST-mediated conjugation of TCE  
13 exists in humans. In particular, at a single substrate concentration of 1 mM, the rate of GSH  
14 conjugation of TCE in human liver cytosol from 9 male and 11 females spanned a range of  
15 2.4-fold (34.7–83.6 nmol DCVG formed/20-minute/mg protein) (Lash et al., 1999b). In liver  
16 microsomes from 5 males and 15-females, the variation in activity was 6.5-fold (9.9–64.6 nmol  
17 DCVG formed/20 minute/mg protein). No sex-dependent variation was identified. Despite  
18 being less pronounced than the known variability in human CYP-mediated oxidation, the impact  
19 on risk assessment of the variability in GSH conjugation to TCE is currently unknown especially  
20 in the absence of data on variability for *N*-acetylation and bioactivation via  $\beta$ -lyase, FMO3, or  
21 CYP3A in the human kidney.

22

#### **3.3.3.1.17. Relative Roles of the Cytochrome P450 (CYP) and Glutathione (GSH) Pathways**

23 In vivo mass balance studies in rats and mice, discussed above, have shown  
24 unequivocally that in these species, CYP oxidation of TCE predominates over GSH conjugation.  
25 In these species, at doses from 2–2,000 mg/kg of [<sup>14</sup>C]TCE, the sum of radioactivity in exhaled  
26 TCE, urine, and exhaled CO<sub>2</sub> constitutes 69–94% of the dose, with the vast majority of the  
27 radioactivity in urine (95–99%) attributable to oxidative metabolites (Dekant et al., 1984; Dekant  
28 et al., 1986b; Green and Prout, 1985; Prout et al., 1985). The rest of the radioactivity was found  
29 mostly in feces and the carcass. More rigorous quantitative limits on the amount of GSH  
30 conjugation based on in vivo data such as these can be obtained using PBPK models, discussed  
31 in Section 3.5.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Comprehensive mass-balance studies are unavailable in humans. DCVG and DCVC in  
2 urine have not been detected in any species, while the amount of urinary NAcDCVC from  
3 human exposures is either below detection limits or very small from a total mass balance point of  
4 view (Bernauer et al., 1996; Birner et al., 1993; Bloemen et al., 2001; Lash et al., 1999b). For  
5 instance, the ratio of primary oxidative metabolites (TCA + TCOH) to NAcDCVC in urine of  
6 rats and humans exposed to 40–160 ppm (215–860 mg/m<sup>3</sup>) TCE heavily favored oxidation,  
7 resulting in ratios of 986–2,562:1 in rats and 3,292–7,163:1 in humans (Bernauer et al., 1996).  
8 Bloemen et al. (2001) reported that at most 0.05% of an inhaled TCE dose would be excreted as  
9 NAcDCVC, and concluded that this suggested TCE metabolism by GSH conjugation was of  
10 minor importance. While it is a useful biomarker of exposure and an indicator of GSH  
11 conjugation, NAcDCVC may capture only a small fraction of TCE flux through the GSH  
12 conjugation pathway due to the dominance of bioactivating pathways (Lash et al., 2000a).

13 A number of lines of evidence suggest that the amount of TCE conjugation to GSH in  
14 humans, while likely smaller than the amount of oxidation, may be much more substantial than  
15 analysis of urinary mercapturates would suggest. In Table 3-28, in vitro estimates of the V<sub>MAX</sub>,  
16 K<sub>M</sub>, and clearance (V<sub>MAX</sub>/K<sub>M</sub>) for hepatic oxidation and conjugation of TCE are compared in a  
17 manner that accounts for differences in cytosolic and microsomal partitioning and protein  
18 content. Surprisingly, the range of in vitro kinetic estimates for oxidation and conjugation of  
19 TCE substantially overlap, suggesting similar flux through each pathway, though with high  
20 interindividual variation. The microsomal and cytosolic protein measurements of GSH  
21 conjugation should be caveated by the observation by Lash et al. (1999a) that GSH conjugation  
22 of TCE was inhibited by ~50% in the presence of oxidation. Note that this comparison cannot be  
23 made in rats and mice because in vitro kinetic parameters for GSH conjugation in the liver are  
24 not available in those species (only activity at 1 or 2 mM have been measured).  
25 Furthermore, as shown earlier in Table 3-22, the human in vivo data of Lash et al. (1999a) show  
26 blood concentrations of DCVG similar, on a molar basis, to that of TCE, TCA, or TCOH,  
27 suggesting substantial conjugation of TCE. In addition, these data give a lower limit as to the  
28 amount of TCE conjugated. In particular, by multiplying the peak blood concentration of DCVG  
29 by the blood volume, a minimum amount of DCVG in the body at that time can be derived (i.e.,  
30 assuming the minimal empirical distribution volume equal to the blood volume). As shown in  
31 Table 3-29, this lower limit amounts to about 0.4–3.7% of the inhaled TCE dose. Since this is  
32 the minimum amount of DCVG in the body at a single time point, the total amount of DCVG  
33 formed is likely to be substantially greater owing to possible distribution outside of the blood as  
34 well as the metabolism and/or excretion of DCVG. Lash et al. (1999b) found levels of urinary  
35 mercapturates were near or below the level of detection of 0.19 µM, results that are consistent

1 with those of Bloemen et al. (2001), who reported urinary concentrations below 0.04  $\mu\text{M}$  at two-  
2 to fourfold lower cumulative exposures. Taken together, these results confirm the suggestion by  
3 Lash et al. (2000a) that NAcDCVC is a poor quantitative marker for the flux through the GSH  
4 pathway.

5         However, as discussed in Section 3.3.3.2.1, data from other laboratories have reported  
6 substantially lower amounts of GSH conjugation in vitro. The reasons for such discrepancies are  
7 unclear, but they may be related to different analytical methods (Lash et al., 2000a). More recent  
8 in vivo data from Kim et al. (2009) in mice reported  $\sim 1000$  times lower DCVG in mouse serum  
9 as compared to the levels of DCVG reported by Lash et al. (1999a) in human blood. These data  
10 are consistent with the suggestion that the “Reed method” employed by Lash et al. (1999a)  
11 overestimated DCVG levels in humans. However, the degree of overestimation is unclear, as is  
12 the degree to which differences may be attributable to true inter-species or inter-individual  
13 variability.

14         In summary, TCE oxidation is likely to be greater quantitatively than conjugation with  
15 GSH in mice, rats, and humans. Some evidence suggests that the flux through the GSH pathway,  
16 particularly in humans, may be greater by an order of magnitude or more than the  $<0.1\%$   
17 typically excreted of NAcDCVC in urine. This is evidenced both by a direct comparison of in  
18 vitro rates of oxidation and conjugation, as well as by in vivo data on the amount of DCVG in  
19 blood. PBPK models can be used to more quantitatively synthesize these data and put more  
20 rigorous limits on the relative amounts of TCE oxidation and conjugation with GSH. Such  
21 analyses are discussed in Section 3.5. However, these data are not consistent with studies in  
22 other laboratories using different analytical methods, which report two to five orders of  
23 magnitude lower estimates of GSH conjugation. Because the reason for these differences have  
24 not been fully determined, substantial uncertainty remains in the degree of GSH conjugation,  
25 particularly in humans.

Table 3-28. Comparison of hepatic in vitro oxidation and conjugation of TCE

Cellular or subcellular fraction	V <sub>MAX</sub> (nmol TCE metabolized/min/g tissue)		K <sub>M</sub> (μM in blood)		V <sub>MAX</sub> /K <sub>M</sub> (mL/min/g tissue)	
	Oxidation	GSH conjugation	Oxidation	GSH conjugation	Oxidation	GSH conjugation
Hepatocytes	10.0–68.4	16~25	22.1–198	16~47	0.087–1.12	0.55~1.0
Liver microsomes	6.1–111	45	2.66–11.1 <sup>a</sup>	5.9 <sup>a</sup>	1.71–28.2 <sup>a</sup>	7.6 <sup>a</sup>
			71.0–297 <sup>b</sup>	157 <sup>b</sup>	0.064–1.06 <sup>b</sup>	0.29 <sup>b</sup>
Liver cytosol	–	380	–	4.5 <sup>a</sup>	–	84 <sup>a</sup>
	–		–	22.7 <sup>b</sup>	–	16.7 <sup>b</sup>

Note: When biphasic metabolism was reported, only high affinity pathway is shown here.

Conversion assumptions for VMAX:

Hepatocellularity of 99 million cells/g liver (Barter et al., 2007);

Liver microsomal protein content of 32 mg protein/g tissue (Barter et al., 2007); and

Liver cytosolic protein content of 89 mg protein/g tissue (based on rats: (Prasanna et al., 1989; van Bree et al., 1990).

Conversion assumptions for K<sub>M</sub>:

For hepatocytes, K<sub>M</sub> in headspace converted to K<sub>M</sub> in blood using blood:air partition coefficient of 9.5 (reported range of measured values 6.5–12.1, Table 3-1);

For microsomal protein, option (a) assumes K<sub>M</sub> in medium is equal to K<sub>M</sub> in tissue, and converts to K<sub>M</sub> in blood by using a liver:blood partition coefficient of 5 (reported ranges of measured values 3.6–5.9, Table 3-8), and option (b) converts K<sub>M</sub> in medium to K<sub>M</sub> in air using the measured microsomal protein:air partition coefficient of 1.78 (Lipscomb et al., 1997), and then converts to K<sub>M</sub> in blood by using the blood:air partition coefficient of 9.5; and

For cytosolic protein, option (a) assumes K<sub>M</sub> in medium is equal to K<sub>M</sub> in tissue, and converts to K<sub>M</sub> in blood by using a liver:blood partition coefficient of 5 (reported ranges of measured values 3.6–5.9, Table 3-8), and option (b) assumes K<sub>M</sub> in medium is equal to K<sub>M</sub> in blood, so no conversion is necessary.

1 **Table 3-29. Estimates of DCVG in blood relative to inhaled TCE dose in**  
 2 **humans exposed to 50 and 100 ppm (269 and 537 mg/m<sup>3</sup>) (Fisher et al., 1998;**  
 3 **Lash et al., 1999b)**  
 4

Sex exposure	Estimated inhaled TCE dose (mmol) <sup>a</sup>	Estimated peak amount of DCVG in blood (mmol) <sup>b</sup>
Males		
50 ppm × 4 h	3.53	0.11 ± 0.08
100 ppm × 4 h	7.07	0.26 ± 0.08
Females		
50 ppm × 4 h	2.36	0.010 ± 0
100 ppm × 4 h	4.71	0.055 ± 0.027

5  
 6 <sup>a</sup> Inhaled dose estimated by (50 or 100 ppm)/(24,450 ppm/mM) × (240 min) × Q<sub>p</sub>, where alveolar ventilation rate Q<sub>p</sub>  
 7 is 7.2 L/min for males and 4.8 L/min for females. Q<sub>p</sub> is calculated as (V<sub>T</sub> - V<sub>D</sub>) × f<sub>R</sub> with the following  
 8 respiratory parameters: tidal volume V<sub>T</sub> (0.75 L for males, 0.46 L for females), dead space V<sub>D</sub> (0.15 L for males,  
 9 0.12 L for females), and respiration frequency f<sub>R</sub> (12 min<sup>-1</sup> for males, 14 min<sup>-1</sup> for females) (assumed sitting,  
 10 awake from The International Commission on Radiological Protection (ICRP, 2003).

11 <sup>b</sup> Peak amount of DCVG in blood estimated by multiplying the peak blood concentration by the estimated blood  
 12 volume: 5.6 L in males and 4.1 L in females (ICRP, 2003).  
 13  
 14  
 15

### 3.4. TRICHLOROETHYLENE (TCE) EXCRETION

16 This section discusses the major routes of excretion of TCE and its metabolites in exhaled  
 17 air, urine, and feces. Unmetabolized TCE is eliminated primarily via exhaled air. As discussed  
 18 in Section 3.3, the majority of TCE absorbed into the body is eliminated by metabolism. With  
 19 the exception of CO<sub>2</sub>, which is eliminated solely via exhalation, most TCE metabolites have low  
 20 volatility and, therefore, are excreted primarily in urine and feces. Though trace amounts of TCE  
 21 metabolites have also been detected in sweat and saliva (Bartonicek, 1962), these excretion  
 22 routes are likely to be relatively minor.  
 23

#### 3.4.1. Exhaled Air

24 In humans, pulmonary elimination of unchanged trichloroethylene and other volatile  
 25 compounds is related to ventilation rate, cardiac output, and the solubility of the compound in  
 26 blood and tissue, which contribute to final exhaled air concentration of TCE. In their study of

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 the impact of workload on TCE absorption and elimination, Astrand and Ovrum (1976)  
2 characterized the postexposure elimination of TCE in expired breath. TCE exposure (540 or  
3 1,080 mg/m<sup>3</sup>; 100 or 200 ppm) was for a total of 2 hours, at workloads from 0–150 Watts.  
4 Elimination profiles were roughly equivalent among groups, demonstrating a rapid decline in  
5 TCE concentrations in expired breath postexposure (see Table 3-30).

6 The lung clearance of TCE represents the volume of air from which all TCE can be  
7 removed per unit time, and is a measure of the rate of excretion via the lungs. Monster et al.  
8 (1976) reported lung clearances ranging from 3.8–4.9 L/minute in four adults exposed at rest to  
9 70 ppm and 140 ppm of trichloroethylene for 4 hours. Pulmonary ventilation rates in these  
10 individuals at rest ranged from 7.7–12.3 L/minute. During exercise, when ventilation rates  
11 increased to 29–30 L/minute, lung clearance was correspondingly higher, 7.7–12.3 L/minute.  
12 Under single and repeated exposure conditions, Monster et al. (1979; 1976) reported from  
13 7–17% of absorbed TCE excreted in exhaled breath. Pulmonary elimination of unchanged  
14 trichloroethylene at the end of exposure is a first-order diffusion process across the lungs from  
15 blood into alveolar air, and it can be thought of as the reversed equivalent of its uptake from the  
16 lungs. Exhaled pulmonary excretion occurs in several distinct (delayed) phases corresponding to  
17 release from different tissue groups, at different times. Sato et al. (1977) detected 3 first-order  
18 phases of pulmonary excretion in the first 10 hours after exposure to 100 ppm for 4 hours, with  
19 fitted half-times of pulmonary elimination of 0.04 hour, 0.67 hour, and 5.6 hours, respectively.  
20 Opdam (1989a) sampled alveolar air up to 20–310 hours after 29–62 minute exposures to 6–38  
21 ppm, and reported terminal half-lives of 8–44 hours at rest. Chiu et al. (2007) sampled alveolar  
22 air up to 100 hours after 6-hour exposures to 1 ppm and reported terminal half-lives of 14–23  
23 hours. The long terminal half-time of TCE pulmonary excretion indicates that a considerable  
24 time is necessary to completely eliminate the compound, primarily due to the high partitioning to  
25 adipose tissues (see Section 3.2).

26 As discussed above, several studies (Dekant et al., 1984; Green and Prout, 1985; Prout et  
27 al., 1985) have investigated the disposition of [<sup>14</sup>C]TCE in rats and mice following gavage  
28 administrations (see Section 3.3.2). These studies have reported CO<sub>2</sub> as an exhalation excretion  
29 product in addition to unchanged TCE. With low doses, the amount of TCE excreted unchanged  
30 in exhaled breath is relatively low. With increasing dose in rats, a disproportionately increased  
31 amount of radiolabel is expired as unchanged TCE. This may indicate saturation of metabolic  
32 activities in rats at doses 200 mg/kg and above, which is perhaps only minimally apparent in the  
33 data from mice. In addition, exhaled air TCE concentration has been measured after constant  
34 inhalation exposure for 2 hours to 50 or 500 ppm in rats (Dallas et al., 1991), and after dermal

- 1 exposure in rats and humans (Poet et al., 2000). Exhaled TCE data from rodents and humans
- 2 have been integrated into the PBPK model presented in Section 3.5.

1 **Table 3-30. Concentrations of TCE in expired breath from**  
 2 **inhalation-exposed humans (Astrand and Ovrum, 1976)**  
 3

Time postexposure	Alveolar air		
	I <sup>a</sup>	II	III
0	459 ± 44	244 ± 16	651 ± 53
30	70 ± 5	51 ± 3	105 ± 18
60	40 ± 4	28 ± 2	69 ± 8
90	35 ± 9	21 ± 1	55 ± 2
120	31 ± 8	16 ± 1	45 ± 1
300	8 ± 1	9 ± 2	14 ± 2
420	5 ± 0.5	4 ± 0.5	8 ± 1.3
19 hours	2 ± 0.3	2 ± 0.2	4 ± 0.5

4 <sup>a</sup> Roman numerals refer to groups assigned different workloads.

5 Concentrations are in mg/m<sup>3</sup> for expired air.  
 6  
 7  
 8  
 9

10 Finally, TCOH is also excreted in exhaled breath, though at a rate about 10,000-fold  
 11 lower than unmetabolized TCE (Monster, 1979; Monster et al., 1976).  
 12

### 3.4.2. Urine

13 Urinary excretion after TCE exposure consists predominantly of the metabolites TCA  
 14 and TCOH, with minor contributions from other oxidative metabolites and GSH conjugates.  
 15 Measurements of unchanged TCE in urine have been at or below detection limits (e.g., Chiu et  
 16 al., 2007; Fisher et al., 1998). The recovery of urinary oxidative metabolites in mice, rats, and  
 17 humans was addressed earlier (see Section 3.3.2) and will not be discussed here. Because of  
 18 their relatively long elimination half-life, urinary oxidative metabolites have been used as an  
 19 occupational biomarker of TCE exposure for many decades (Carrieri et al., 2007; Ikeda and  
 20 Imamura, 1973). Ikeda and Imamura (1973) measured total trichloro compounds TCOH and  
 21 TCA, in urine over 3 consecutive postexposure days for 4 exposure groups totaling 24 adult  
 22 males and one exposure group comprising 6 adult females. The elimination half-life for TTC  
 23 ranged 26.1–48.8 hours in males and was 50.7 hours in females. The elimination half-life for  
 24 TCOH was 15.3 hours in the only group of males studied and was 42.7 hours in females. The

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 elimination half-life for TCA was 39.7 hours in the only group of males studied and was 57.6  
2 hours in females. These authors compared their results to previously published elimination  
3 half-lives for TTC, TCOH, and TCA. Following experimental exposures of groups of two–five  
4 adults, elimination half-lives ranged from 31–50 hours for TTC; 19–29 hours for TCOH; and  
5 36–55 hours for TCA (Bartonicek, 1962; Nomiya and Nomiya, 1971; Ogata et al., 1971;  
6 Stewart et al., 1970). The urinary elimination half-life of TCE metabolites in a subject who  
7 worked with and was addicted to sniffing TCE for 6–8 years approximated 49.7 hours for  
8 TCOH, 72.6 hours for TCA, and 72.6 hours for TTC (Ikeda et al., 1971).

9         The quantitative relationship between urinary concentrations of oxidative metabolites and  
10 exposure in an occupational setting was investigated by Ikeda (1977). This study examined the  
11 urinary elimination of TCE and metabolites in urine of 51 workers from 10 workshops. The  
12 concentration of TCA and TCOH in urine demonstrated a marked concentration-dependence,  
13 with concentrations of TCOH being approximately twice as high as those for TCA. Urinary  
14 half-life values were calculated for six males and six females from five workshops; males were  
15 intermittently exposed to 200 ppm and females were intermittently exposed to 50 ppm  
16 (269 mg/m<sup>3</sup>). Urinary elimination half-lives for TTC, TCOH and TCA were 26.1, 15.3, and  
17 39.7 hours; and 50.7, 42.7 and 57.6 hours in males and females, respectively, which were similar  
18 to the range of values previously reported. These authors estimated that urinary elimination of  
19 parent TCE during exposure might account for one-third of the systemically absorbed dose.  
20 Importantly, urinary TCA exhibited marked saturation at exposures higher than 50 ppm.  
21 Because neither TTC nor urinary TCOH (in the form of the glucuronide TCOG) showed such an  
22 effect, this saturation cannot be due to TCE oxidation itself, but must rather be from one of the  
23 metabolic processes forming TCA from TCOH. Unfortunately, since biological monitoring  
24 programs usually measure only urinary TCA, rather than TTC, urinary TCA levels above around  
25 150 mg/L cannot distinguish between exposures at 50 ppm and at much higher concentrations.

26         It is interesting to attempt to extrapolate on a cumulative exposure basis the Ikeda (1977)  
27 results for urinary metabolites obtained after occupational exposures at 50 ppm to the controlled  
28 exposure study by Chiu et al. (2007) at 1.2 ppm for 6 hours (the only controlled exposure study  
29 for which urinary concentrations, rather than only cumulative excretion, are available). Ikeda  
30 (1977) reported that measurements were made during the second half of the week, so one can  
31 postulate a cumulative exposure duration of 20–40 hours. At 50 ppm, Ikeda (1977) report a  
32 urinary TCOH concentration of about 290 mg/L, so that per ppm-hour, the expected urinary  
33 concentration would be  $290 / (50 \times 20 \sim 40) = 0.145 \sim 0.29$  mg/L-ppm-hour. The cumulative  
34 exposure in Chiu et al. (2007) is  $1.2 \times 6 = 7.2$  ppm-hour, so the expected urinary TCOH  
35 concentration would be  $7.2 \times (0.145 \sim 0.29) = 1.0 \sim 2.1$  mg/L. This estimate is somewhat

1 surprisingly consistent with the actual measurements of Chiu et al. (2007) during the first day  
2 postexposure, which ranged from 0.8~1.2 mg/L TCOH in urine.

3 On the other hand, extrapolation of TCA concentrations was less consistent. At 50 ppm,  
4 Ikeda (1977) report a urinary TCA concentration of about 140 mg/L, so that per ppm-hour, the  
5 expected urinary concentration would be  $140/(50 \times 20 \sim 40) = 0.07 \sim 0.14$  mg/L-ppm-hour. The  
6 cumulative exposure in Chiu et al. (2007) is  $1.2 \times 6 = 7.2$  ppm-hour, so the expected urinary  
7 TCA concentration would be  $7.2 \times (0.07 \sim 0.14) = 0.5 \sim 1.0$  mg/L, whereas Chiu et al. (2007)  
8 reported urinary TCA concentrations on the first day after exposure of 0.03~0.12 mg/L.  
9 However, as noted in Chiu et al. (2007), relative urinary excretion of TCA was 3- to 10-fold  
10 lower in Chiu et al. (2007) than other studies at exposures of 50~140 ppm, which may explain  
11 part of the discrepancies. However, this may be due in part to saturation of many urinary TCA  
12 measurements, and, furthermore, interindividual variance, observed to be substantial in Fisher  
13 et al.(1998), cannot be ruled out.

14 Urinary elimination kinetics have been reported to be much faster in rodents than in  
15 humans. For instance, adult rats were exposed to 50, 100, or 250 ppm (269, 537, or  
16  $1,344 \text{ mg/m}^3$ ) via inhalation for 8 hours or were administered an i.p. injection (1.47 g/kg) and the  
17 urinary elimination of total trichloro compounds was followed for several days (Ikeda and  
18 Imamura, 1973). These authors calculated urinary elimination half-lives of 14.3–15.6 hours for  
19 female rats and 15.5–16.6 hours for male rats; the route of administration did not appear to  
20 influence half-life value. In other rodent experiments using orally administered radiolabeled  
21 TCE, urinary elimination was complete within one or two days after exposure (Dekant et al.,  
22 1984; Green and Prout, 1985; Prout et al., 1985).

### 3.4.3. Feces

23 Fecal elimination accounts for a small percentage of TCE as shown by limited  
24 information in the available literature. Bartonicek (1962) exposed 7 human volunteers to  
25 1.042 mg TCE/L air for 5 hours and examined TCOH and TCA in feces on the 3<sup>rd</sup> and 7<sup>th</sup> day  
26 following exposure. The mean amount of TCE retained during exposure was 1,107 mg,  
27 representing 51–64% (mean 58%) of administered dose. On the third day following TCE  
28 exposure, TCOH and TCA in feces demonstrated mean concentrations of 17.1 and  
29 18.5 mg/100 grams feces, similar to concentrations in urine. However, because of the 10-fold  
30 smaller daily rate of excretion of feces relative to urine, this indicates fecal excretion of these  
31 metabolites is much less significant than urinary excretion. Neither TCOH nor TCA was  
32 detected in feces on the seventh day following exposure.

1            In rats and mice, total radioactivity has been used to measure excretion in feces after oral  
2 gavage TCE administration in corn oil, but since the radiolabel was not characterized it is not  
3 possible to determine whether the radiolabel in feces represented unabsorbed parent compound,  
4 excreted parent compound, and/or excreted metabolites. Dekant et al. (1984) reported mice  
5 eliminated 5% of the total administered TCE, while rats eliminated 2% after oral gavage.  
6 Dekant et al. (1986b) reported a dose response related increase in fecal elimination with dose,  
7 ranging between 0.8–1.9% in rats and 1.6–5% in mice after oral gavage in corn oil. Due to the  
8 relevant role of CYP2E1 in the metabolism of TCE (see Section 3.3.3.1.6), Kim and Ghanayem  
9 (2006) compared fecal elimination in both wild-type and CYP2E1 knockout mice and reported  
10 fecal elimination ranging between 4.1–5.2% in wild-type and 2.1–3.8% in knockout mice  
11 exposed by oral gavage in aqueous solution.  
12

### **3.5. PHYSIOLOGICALLY BASED PHARMACOKINETIC (PBPK) MODELING OF TRICHLOROETHYLENE (TCE) AND ITS METABOLITES**

#### **3.5.1. Introduction**

13            PBPK models are extremely useful tools for quantifying the relationship between  
14 external measures of exposure and internal measures of toxicologically relevant dose. In  
15 particular, for the purposes of this assessment, PBPK models are evaluated for the following:  
16 (1) providing additional quantitative insights into the ADME of TCE and metabolites described  
17 in the sections above; (2) cross-species pharmacokinetic extrapolation of rodent studies of both  
18 cancer and noncancer effects, (3) exposure-route extrapolation; and (4) characterization of  
19 human pharmacokinetic variability. The following sections first describe and evaluate previous  
20 and current TCE PBPK modeling efforts, then discuss the insights into ADME (1, above), and  
21 finally present conclusions as to the utility of the model to predict internal doses for use in  
22 dose-response assessment (2–4, above).  
23

#### **3.5.2. Previous Physiologically Based Pharmacokinetic (PBPK) Modeling of Trichloroethylene (TCE) for Risk Assessment Application**

24            TCE has an extensive number of both in vivo pharmacokinetic and PBPK modeling  
25 studies (see Chiu et al., 2006, supplementary material, for a review). Models previously  
26 developed for occupational or industrial hygiene applications are not discussed here but are  
27 reviewed briefly in Clewell et al. (2000). Models designed for risk assessment applications have

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 focused on descriptions of TCE and its major oxidative metabolites TCA, TCOH, and TCOG.  
2 Most of these models were extensions of the “first generation” of models developed by Fisher  
3 and coworkers (Allen and Fisher, 1993; Fisher et al., 1991) in rats, mice, and humans. These  
4 models, in turn, are based on a Ramsey and Andersen (1984) structure with flow-limited tissue  
5 compartments and equilibrium gas exchange, saturable Michaelis-Menten kinetics for oxidative  
6 metabolism, and lumped volumes for the major circulating oxidative metabolites TCA and  
7 TCOH. Fisher and coworkers updated their models with new in vivo and in vitro experiments  
8 performed in mice (Abbas and Fisher, 1997; Greenberg et al., 1999) and human volunteers  
9 (Fisher et al., 1998) and summarized their findings in Fisher (2000). Clewell et al. (2000) added  
10 enterohepatic recirculation of TCOG and pathways for local oxidative metabolism in the lung  
11 and GST metabolism in the liver. While Clewell et al. (2000) does not include the updated  
12 Fisher data, they have used a wider set of in vivo and in vitro mouse, rat, and human data than  
13 previous models. Finally, Bois (2000a, b) performed reestimations of PBPK model parameters  
14 for the Fisher and Clewell models using a Bayesian population approach (Gelman et al., 1996,  
15 and discussed further below).

16 As discussed in Rhomberg (2000), the choice as to whether to use the Fisher, Clewell,  
17 and Bois models for cross-species extrapolation of rodent cancer bioassays led to quantitative  
18 results that differed by as much as an order of magnitude. There are a number of differences in  
19 modeling approaches that can explain their differing results. First, the Clewell et al. (2000)  
20 model differed structurally in its use of single-compartment volume-of-distribution models for  
21 metabolites as opposed to the Fisher (2000) models’ use of multiple physiologic compartments.  
22 Also, the Clewell et al. (2000) model, but not the Fisher models, includes enterohepatic  
23 recirculation of TCOH/TCOG (although reabsorption was set to zero in some cases). In addition  
24 to structural differences in the models, the input parameter values for these various models were  
25 calibrated using different subsets of the overall in vivo database (see Chiu et al., 2006,  
26 supplementary material, for a review). The Clewell et al. (2000) model is based primarily on a  
27 variety of data published before 1995; the Fisher (2000) models were based primarily on new  
28 studies conducted by Fisher and coworkers (after 1997); and the Bois (2000a, b) reestimations of  
29 the parameters for the Clewell et al. (2000) and Fisher (2000) models used slightly different data  
30 sets than the original authors. The Bois (2000a, b) reanalyses also led to somewhat different  
31 parameter estimates than the original authors, both because of the different data sets used as well  
32 as because the methodology used by Bois allowed many more parameters to be estimated  
33 simultaneously than were estimated in the original analyses.

34 Given all these methodological differences, it is not altogether surprising that the  
35 different models led to different quantitative results. Even among the Fisher models themselves,

1 Fisher (2000) noted some inconsistencies, including differing estimates for metabolic parameters  
2 between mouse gavage and inhalation experiments. These authors included possible  
3 explanations for these inconsistencies: the impact of corn oil vehicle use during gavage (Staats et  
4 al., 1991) and the impact of a decrease in ventilation rate in mice due to sensory irritation during  
5 the inhalation of solvents [e.g., Stadler and Kennedy (1996)].

6 As discussed in a report by the National Research Council (NRC, 2006), several  
7 additional PBPK models relevant to TCE pharmacokinetics have been published since 2000 and  
8 are reviewed briefly here. Poet et al. (2000) incorporated dermal exposure to TCE in PBPK  
9 models in rats and humans, and published in vivo data in both species from dermal exposure  
10 (Poet et al., 2000; Thrall and Poet, 2000). Albanese et al. (2002) published a series of models  
11 with more complex descriptions of TCE distribution in adipose tissue but did not show  
12 comparisons with experimental data. Simmons et al. (2002) developed a PBPK model for TCE  
13 in the Long-Evans rat that focused on neurotoxicity endpoints and compared model predictions  
14 with experimentally determined TCE concentrations in several tissues—including the brain.  
15 Keys et al. (2003) investigated the lumping and un lumping of various tissue compartments in a  
16 series of PBPK models in the rat and compared model predictions with TCE tissue  
17 concentrations in a multitude of tissues. Although none of these TCE models included  
18 metabolite descriptions, the experimental data was available for either model or evaluation.  
19 Finally, Keys et al. (2004) developed a model for DCA in the mouse that included a description  
20 of suicide inhibition of GST-zeta, but this model was not been linked to TCE.

### 3.5.3. Development and Evaluation of an Interim “Harmonized” Trichloroethylene (TCE) Physiologically Based Pharmacokinetic (PBPK) Model

22 Throughout 2004, EPA and the U.S. Air Force jointly sponsored an integration of the  
23 Fisher, Clewell, and Bois modeling efforts (Hack et al., 2006). In brief, a single interim PBPK  
24 model structure combining features from both the Fisher and Clewell models was developed and  
25 used for all three species of interest (mice, rats, and humans). An effort was made to combine  
26 structures in as simple a manner as possible; the evaluation of most alternative structures was left  
27 for future work. The one level of increased complexity introduced was inclusion of species- and  
28 dose-dependent TCA plasma binding, although only a single in vitro study of Lumpkin et al.  
29 (2003) was used as parameter inputs. As part of this joint effort, a hierarchical Bayesian  
30 population analysis using Markov chain Monte Carlo (MCMC) sampling (similar to the Bois  
31 [2000a, b] analyses) was performed on the revised model with a cross-section of the combined  
32 database of kinetic data to provide estimates of parameter uncertainty and variability (Hack et al.,



1 2006). Particular attention was given to using data from each of the different efforts, but owing  
2 to time and resource constraints, a combined analysis of all data was not performed. The results  
3 from this effort suggested that a single model structure could provide reasonable fits to a variety  
4 of data evaluated for TCE and its major oxidative metabolites TCA, TCOH, and TCOG.  
5 However, in many cases, different parameter values—particularly for metabolism—were  
6 required for different studies, indicating significant interindividual or interexperimental  
7 variability. In addition, these authors concluded that dosimetry of DCA, conjugative  
8 metabolites, and metabolism in the lung remained highly uncertain (Hack et al., 2006).

9 Subsequently, EPA conducted a detailed evaluation of the Hack et al. (2006) model that  
10 included (1) additional model runs to improve convergence; (2) evaluation of posterior  
11 distributions for population parameters; and (3) comparison of model predictions both with the  
12 data used in the Hack et al. (2006) analysis as well as with additional data sets identified in the  
13 literature. Appendix A provides the details and conclusions of this evaluation, briefly  
14 summarized in Table 3-31, along with their pharmacokinetic implications.

### 3.5.4. Physiologically Based Pharmacokinetic (PBPK) Model for Trichloroethylene (TCE) and Metabolites Used for This Assessment

#### 3.5.4.1.1. Introduction

16 Based on the recommendations of the NRC (2006) as well as additional analysis and  
17 evaluation of the Hack et al. (2006) PBPK model, an updated PBPK model for TCE and  
18 metabolites was developed for use in this risk assessment. The updated model is reported in  
19 Evans et al. (2009) and Chiu et al. (2009), and the discussion below provides some details in  
20 additional to the information in the published articles.

21 This updated model included modification of some aspects of the Hack et al. (2006)  
22 PBPK model structure, incorporation of additional in vitro and in vivo data for estimating model  
23 parameters, and an updated hierarchical Bayesian population analysis of PBPK model  
24 uncertainty and variability. In the subsections below, the updated PBPK model and baseline

**Table 3-31. Conclusions from evaluation of Hack et al. (2006), and implications for PBPK model development**

Conclusion from evaluation of Hack et al. (2006) model	Implications for PBPK model parameters, structure, or data
<p>For some model parameters, posterior distributions were somewhat inconsistent with the prior distributions.</p> <ul style="list-style-type: none"> <li>• For parameters with strongly informative priors (e.g., tissue volumes and flows), this may indicate errors in the model.</li> <li>• For many parameters, the prior distributions were based on visual fits to the same data. If the posteriors are inconsistent, then that means the priors were “inappropriately” informative, and, thus, the same data was used twice.</li> </ul>	<p>Reevaluation of all prior distributions</p> <ul style="list-style-type: none"> <li>• Update priors for parameters with independent data (physiological parameters, partition coefficients, in vitro metabolism), looking across all available data sets.</li> <li>• For priors without independent data (e.g., many metabolism parameters), use less informative priors (e.g., log-uniform distributions with wide bounds) so as prevent bias.</li> </ul> <p>Evaluate modifications to the model structure, as discussed below.</p>
<p>A number of data sets involve TCE (i.a., portal vein), TCA (oral, i.v.), and TCOH (oral, i.v.) dosing routes that are not currently in the model, but could be useful for calibration.</p>	<ul style="list-style-type: none"> <li>• Additional dosing routes can be added easily.</li> </ul>
<p>TCE concentrations in blood, air, and tissues well-predicted only in rats, not in mice and humans. Specifically:</p> <ul style="list-style-type: none"> <li>• In mice, the oral uptake model could not account for the time-course of several data sets. Blood TCE concentrations after inhalation consistently over-predicted.</li> <li>• In rats, tissue concentrations measured in data not used for calibration were accurately predicted.</li> <li>• In humans, blood and air TCE concentrations were consistently over-predicted in the majority of (but not all) data sets.</li> </ul>	<ul style="list-style-type: none"> <li>• In mice, uptake from the stomach compartment (currently zero), but previously included in Abbas and Fisher (1997), may improve the model fit.</li> <li>• In mice and humans, additional extrahepatic metabolism, either presystemic (e.g., in the lung) or postsystemic (e.g., in the kidney) and/or a wash-in/wash-out effect may improve the model fit.</li> </ul>
<p>Total metabolism appears well-predicted in rats and mice based on closed-chamber data, but required significantly different <math>V_{MAX}</math> values between dose groups. Total recovery in humans (60–70%) is less than the model would predict. In all three species, the ultimate disposition of metabolism is uncertain. In particular, there are uncertainties in attributing the “missing” metabolism to</p> <ul style="list-style-type: none"> <li>• GSH pathway (e.g., urinary mercapturates may only capture a fraction of the total flux; moreover, in Bernauer et al. (1996), excretion was still on-going at end of collection period; model does not accurately depict time-course of mercapturate excretion).</li> <li>• Other hepatic oxidation (currently attributed to DCA).</li> <li>• Extrahepatic systemic metabolism (e.g., kidney).</li> <li>• Presystemic metabolism in the lung.</li> <li>• Additional metabolism of TCOH or TCA (see below).</li> </ul>	<ul style="list-style-type: none"> <li>• Calibration of GSH pathway may be improved by utilizing in vitro data on liver and kidney GSH metabolism, adding a DCVG compartment to improve the prediction of the time-course for mercapturate excretion, and/or using the Lash et al. (1999b) blood DCVG in humans (necessitating the addition of a DCVG compartment).</li> <li>• Presystemic lung metabolism can only be evaluated if added to the model (in vitro data exists to estimate the <math>V_{MAX}</math> for such metabolism). In addition, a wash-in/wash-out effect [e.g., suggested by Greenberg et al. (1999)] can be evaluated using a continuous breathing model that separately tracks inhaled and exhaled air, with adsorption/desorption in the respiratory tract.</li> <li>• Additional elimination pathways for TCOH and TCA can be added for evaluation.</li> </ul>

**Table 3-31. Conclusions from evaluation of Hack et al. (2006), and implications for PBPK model development (continued)**

Conclusion from evaluation of Hack et al. (2006) model	Implications for PBPK model parameters, structure, or data
<p>TCA blood/plasma concentrations well predicted following TCE exposures in all species. However, there may be inaccuracies in the total flux of TCA production, as well as its disposition.</p> <ul style="list-style-type: none"> <li>• In TCA dosing studies, the majority (&gt;50%), but substantially &lt;100%, was recovered in urine, suggesting significant metabolism of TCA. Although urinary TCA was well predicted in mice and humans (but not in rats), if TCA metabolism is significant, then this means that the current model underestimates the flux of TCE metabolism to TCA.</li> <li>• An improved TCOH/TCOG model may also provide better estimates of TCA kinetics (see below).</li> </ul> <p>TCOH/TCOG concentrations and excretion were inconsistently predicted, particularly after TCOH dosing.</p> <ul style="list-style-type: none"> <li>• In mice and rats, first-order clearance for TCOH glucuronidation was predicted to be greater than hepatic blood flow, which is consistent with a first pass effect that is not currently accounted for.</li> <li>• In humans, the estimated clearance rate for TCOH glucuronidation was substantially smaller than hepatic blood flow. However, the presence of substantial TCOG in blood (as opposed to free TCOH) in the Chiu et al. (2007) data are consistent with greater glucuronidation than predicted by the model.</li> <li>• In TCOH dosing studies, substantially &lt;100% was recovered in urine as TCOG and TCA, suggesting another metabolism or elimination pathway.</li> </ul>	<ul style="list-style-type: none"> <li>• Additional elimination pathways for TCOH and TCA can be added for evaluation.</li> <li>• The addition of a liver compartment for TCOH and TCOG would permit hepatic first-pass effects to be accounted for, as appears necessary for mice and rats.</li> </ul>

i.a. = intra-arterial, i.v. = intravenous.

1 parameter values are described, as well as the approach and results of the analysis of PBPK  
2 model uncertainty and variability. Appendix A provides more detailed descriptions of the model  
3 and parameters, including background on hierarchical Bayesian analyses, model equations,  
4 statistical distributions for parameter uncertainty and variability, data sources for these parameter  
5 values, and the PBPK model code. Additional computer codes containing input files to the  
6 MCSim program are available electronically.

7


### 3.5.4.1.2. Updated Physiologically Based Pharmacokinetic (PBPK) Model Structure

8 The updated TCE PBPK model is illustrated in Figure 3-7, with detailed descriptions of  
9 the model structure, equations, and parameters found in Appendix A (see Section A.4), and the  
10 major changes from the Hack et al. (2006) model described here. The TCE submodel was  
11 augmented by the addition of kidney and venous blood compartments, and an updated  
12 respiratory tract model that included both metabolism and the possibility of local storage in the  
13 respiratory tissue. In particular, in the updated lung, separate processes describing inhalation and  
14 exhalation allowed for adsorption and desorption from tracheobronchial epithelium  
15 (wash-in/wash-out), with the possibility of local metabolism as well. In addition, conjugative  
16 metabolism in the kidney was added, motivated by the in vitro data on TCE conjugation  
17 described in Sections 3.3.3.2–3.3.3.3. With respect to oxidation, a portion of the lung  
18 metabolism was assumed to produce systemically available oxidative metabolites, including  
19 TCOH and TCA, with the remaining fraction assumed to be locally cleared. This is clearly a  
20 lumping of a multistep process, but the lack of data precludes the development of a more  
21 sequential model. TCE oxidation in the kidney was not included because it was not likely to  
22 constitute a substantial flux of total TCE oxidation given the much lower CYP activity in the  
23 kidney relative to the liver (Cummings and Lash, 2000; Cummings et al., 1999) and the greater  
24 tissue mass of the liver.<sup>2</sup> In addition, liver compartments were added to the TCOH and TCOG

---

2 The extraction ratio for kidney oxidation is likely to be very low, as shown by the following calculation in rats and humans. In rats, the in vitro kidney oxidative clearance ( $V_{MAX}/K_M$ ) rate (see Table 3-13, converting units) is  $1.64 \times 10^{-7}$  L/min/mg microsomal protein. Converting units using 16 mg microsomal protein to g tissue (Bong et al., 1985) gives a clearance rate per unit tissue mass of  $2.6 \times 10^{-6}$  L/min/g kidney. This is more than a 1000-fold smaller than the kidney specific blood flow rate of  $6.3 \times 10^{-3}$  L/min/g kidney (Brown et al., 1997). In humans, an in vitro clearance rate of  $6.5 \times 10^{-8}$  L/min/mg microsomal protein is derived from the only detectable in vitro oxidation rate from Cummings and Lash (2000) of 0.13 nmol/minute/mg protein at 2 mM. Using the same conversion from microsomal protein to tissue mass gives a clearance rate of  $1.0 \times 10^{-6}$  L/min/g kidney, more than 1000-fold smaller than the kidney specific blood flow of  $3.25 \times 10^{-3}$  L/min/g kidney (Brown et al., 1997). No data on kidney metabolism are available in mice, but the results are likely to be similar. Therefore, even accounting for uncertainties of up to an order of magnitude in the in vitro-to-in vivo conversion, kidney oxidation should contribute negligibly to total metabolism of TCE.

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 submodels to account properly for first-pass hepatic metabolism, which is important for  
 2 consistency across

3 **Figure 3-7. Overall structure of PBPK model for TCE and metabolites used**  
 4 **in this assessment.** Boxes with underlined labels are additions or modifications  
 5 of the Hack et al. (2006) model, which are discussed in Table 3-32.

*This document is a draft for review purposes only and does not constitute Agency policy.*



1  
2  
3

**Table 3-32. Discussion of changes to the Hack et al. (2006) PBPK model implemented for this assessment**

<b>Change to Hack et al. (2006) PBPK model</b>	<b>Discussion</b>
TCE respiratory tract compartments and metabolism	<p>In vitro data indicate that the lung (at least in the mouse) has a significant capacity for oxidizing TCE. However, in the Hack et al. (2006) model, respiratory metabolism was blood flow-limited. The model structure used was inconsistent with other PBPK models in which the same mechanism for respiratory metabolism is assumed [e.g., styrene, Sarangapani et al. (2003)]. In these models, the main source of exposure in the respiratory tract tissue is from the respiratory lumen—not from the tracheobronchial blood flow. In addition, a wash-in/wash-out effect has also been postulated. The current structure, which invokes a “continuous breathing” model with separate “inhaled” and “exhaled” respiratory lumens, can accommodate both respiratory metabolism due to exposure from the respiratory lumen as well as a wash-in/wash-out effect in which there is temporary storage in the respiratory tract tissue.</p> <p>Moreover, preliminary analyses indicated that these changes to the model structure allowed for a substantially better fit to mouse closed-chamber data under the requirement that all the dose levels are modeled using the same set of parameters.</p>
TCE kidney compartment	<p>In vitro data indicate that the kidney has a significant capacity for conjugating TCE with GSH.</p>
TCE venous blood compartment	<p>Many PBPK models have used a separate blood compartment. It was believed to be potentially important and feasible to implement here because (1) TCE blood concentrations were often not well predicted by the Hack et al. (2006) model; (2) the TCA submodel has a plasma compartment, which is a fraction of the blood volume based on the blood volume; (3) adequate independent information on blood volume is available; and (4) the updated model was to include the intravenous route of exposure.</p>
TCOH and TCOG liver compartments	<p>In mice and rats, the Hack et al. (2006) model estimated a rate of TCOH glucuronidation that exceeded hepatic blood flow (all glucuronidation is assumed to occur in the liver), which indicated a significant first-pass effect. Therefore, a separate liver compartment is necessary to account properly for hepatic first-pass.</p>
TCOH and TCA “other” elimination pathways	<p>Mass-balance studies with TCOH and TCA dosing indicated that, although the majority of TCOH and TCA are excreted in urine, the amount is still substantially less than 100%. Therefore, additional elimination of TCOH and TCA must exist and should be accounted for.</p>

*This document is a draft for review purposes only and does not constitute Agency policy.*

DCVG compartment (human model only)	Blood DCVG data in humans exist as part of the Fisher et al. (1998) experiments, reported in Lash et al. (1999b), and a DCVG compartment is necessary in order to utilize those data.
-------------------------------------	---

1  
2 routes of exposure. Furthermore, additional clearance pathways of TCOH and TCA were added  
3 to their respective submodels. With respect to TCE conjugation, in humans, an additional  
4 DCVG compartment was added between TCE conjugation and production of DCVC. In  
5 addition, it should be noted that the urinary clearance of DCVC represents a lumping of  
6 *N*-acetylation of DCVC, deacetylation of NAcDCVC, and urinary excretion NAcDCVC, and that  
7 the bioactivation of DCVC represents a lumping of thiol production from DCVC by beta-lyase,  
8 sulfoxidation of DCVC by FMO3, and sulfoxidation of NAcDCVC by CYP3A. Such lumping  
9 was used because these processes are not individually identifiable given the available data.  
10

**3.5.4.1.3. Specification of Baseline Physiologically Based Pharmacokinetic (PBPK) Model Parameter**

11 Point estimates for PBPK model parameters (“baseline values”), used as central estimates  
12 in the prior distributions for population mean parameters in the hierarchical Bayesian statistical  
13 model (see Appendix A), were developed using standard methodologies to ensure biological  
14 plausibility, and were a refinement of those used in Hack et al. (2006). Because the Bayesian  
15 parameter estimation methodology utilizes the majority of the useable *in vivo* data on TCE  
16 pharmacokinetics, all baseline parameter estimates were based solely on measurements  
17 independent of the *in vivo* data. This avoids using the same data in both the prior and the  
18 likelihood. These parameters were, in turn, given truncated normal or lognormal distributions  
19 for the uncertainty in the population mean. If no independent data were available, as is the case  
20 for many “downstream” metabolism parameters, then no baseline value was specified, and a  
21 noninformative prior was used. Section 3.5.5.4, below, discusses the updating of these  
22 noninformative priors using interspecies scaling.

23 In keeping with standard practice, many of the PBPK model parameters were “scaled” by  
24 body or organ weights, cardiac output, or allometrically by an assumed (fixed) power of body  
25 weight. Metabolic capacity and cardiac output were scaled by the <sup>3</sup>/<sub>4</sub> power of body weight and  
26 rate coefficients were scaled by the—<sup>1</sup>/<sub>4</sub> power of body weight, in keeping with general  
27 expectations as to the relationship between metabolic rates and body size (U.S. EPA, 1992; West  
28 et al., 2002). So as to ensure a consistent model structure across species as well as improve the  
29 performance of the MCMC algorithm, parameters were further scaled to the baseline



1 point-estimates where available, as was done by Hack et al. (2006). For example, to obtain the  
2 actual liver volume (VLivC) in liters, a point estimate is first obtained by multiplying the fixed,  
3 species-specific baseline point estimate for the fractional liver volume by a fixed body weight  
4 (measured or species-specific default) with density of 1 kg per liter assumed to convert from kg  
5 to liters. Then, any deviation from this point estimate is represented by multiplying by a separate  
6 “scaled” parameter VLivC that has a value of one if there is no deviation from the point estimate.  
7 These “scaled” parameters are those estimated by the MCMC algorithm, and for which  
8 population means and variances are estimated.

9 Baseline physiological parameters were reestimated based on the updated tissue lumping  
10 (e.g., separate blood and kidney compartments) using the standard references International  
11 Commission on Radiological Protection (ICRP, 2003) and Brown et al. (1997). For a few of  
12 these parameters, such as hematocrit and respiratory tract volumes in rodents, additional  
13 published sources were used as available, but no attempt was made to compile a comprehensive  
14 review of available measurements. In addition, a few parameters, such as the slowly perfused  
15 volume, were calculated rather than sampled in order to preserve total mass or flow balances.

16 For chemical-specific distribution and metabolism parameters, in vitro data from various  
17 sources were used. Where multiple measurements had been made, as was the case for many  
18 partition coefficients, TCA plasma protein binding parameters, and TCE metabolism, different  
19 results were pooled together, with their uncertainty reflected appropriately in the prior  
20 distribution. Such in vitro measurements were available for most chemical partition coefficients,  
21 except for those for TCOG (TCOH used as a proxy) and DCVG. There were also such data to  
22 develop baseline values for the oxidative metabolism of TCE in the liver ( $V_{MAX}$  and  $K_M$ ), the  
23 relative split in TCE oxidation between formation of TCA and TCOH, and the  $V_{MAX}$  for TCE  
24 oxidation in the lung. For GSH conjugation, the geometric means of the in vitro data from Lash  
25 et al. (1999a) and Green et al. (1997a) were used as central estimates, with a wide enough  
26 uncertainty range to encompass both (widely disparate) estimates. Thus, the prior distribution  
27 for these parameters was only mildly informative, and the results are primarily determined by the  
28 available in vivo data. All other metabolism parameters were not given baseline values and  
29 needed to be estimated from the in vivo data.

#### 3.5.4.1.4. Dose-Metric Predictions

31 The purpose of this PBPK model is to make predictions of internal dose in rodents used in  
32 toxicity studies or in humans in the general population, and not in the groups or individuals for  
33 which pharmacokinetic data exist. Therefore, to evaluate its predictive utility for risk

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 assessment, a number of dose-metrics were selected for simulation in a “generic” mouse, rat, or  
2 human, summarized in Table 3-33. The parent dose-metric was AUC in blood. TCE  
3 metabolism dose-metrics (i.e., related to the amount metabolized) included both total  
4 metabolism, metabolism splits between oxidation versus conjugation, oxidation in the liver  
5 versus the lung, the amount of oxidation in the liver to products *other* than TCOH and TCA, and  
6 the amount of TCA produced. These metabolism rate dose-metrics are scaled by body weight in  
7 the case of TCA produced, by the metabolizing tissue volume and by body weight to the  
8  $\frac{3}{4}$  power in the cases of the lung and “other” oxidation in the liver, and by body weight to the  
9  $\frac{3}{4}$  power only in other cases. With respect to the oxidative metabolites, liver concentrations of  
10 TCA and blood concentrations of free TCOH were used. With respect to conjugative  
11 metabolites, the dose-metrics considered were total GSH metabolism scaled by body weight to  
12 the  $\frac{3}{4}$  power, and the amount of DCVC bioactivated (rather than excreted in urine) per unit body  
13 weight to the  $\frac{3}{4}$  power and per unit kidney mass.

14 All dose-metrics are converted to daily or weekly averages based on simulations lasting  
15 10 weeks for rats and mice and 100 weeks for humans. These simulation times were the shortest  
16 for which additional simulation length did not add substantially to the average (i.e., less than a  
17 few percent change with a doubling of simulation time).

18

### **3.5.5. Bayesian Estimation of Physiologically Based Pharmacokinetic (PBPK) Model Parameters, and Their Uncertainty and Variability**

#### **3.5.5.1.1. Updated Pharmacokinetic Database**

19 An extensive search was made for data not previously considered in the PBPK modeling  
20 of TCE and metabolites, with a few studies identified or published subsequent to the review by  
21 Chiu et al. (2006b). The studies considered for analysis are listed in Tables 3-34–3-35, along  
22 with an indication of whether and how they were used.<sup>3</sup>

23 The least amount of data was available for mice, so an effort was made to include as  
24 many studies as feasible for use in calibrating the PBPK model parameters. Exceptions include  
25 mouse studies with CH or DCA dosing, since those metabolites are not included in the PBPK  
26 model. In addition, the Birner et al. (1993) data only reported urine concentrations, not the  
27 amount excreted in urine. Because there is uncertainty as to total volume of urine excreted, and  
28 over what time period, these data were not used. Moreover, many other studies had urinary

---

<sup>3</sup> Additional in vivo data on TCE or metabolites published after the PBPK modeling was completed ([Sweeney et al., 2009](#)) ([Kim et al., 2009](#); [Liu et al., 2009](#)) were evaluated separately, and discussed in Appendix A.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 excretion data, so this exclusion should have minimal impact. Several data sets not included by  
 2 Hack et al. (2006) were used here. Of particular importance was the inclusion of TCA and  
 3 TCOH dosing data from Abbas et al. (1997), Green and Prout (1985), Larson and Bull (1992a),  
 4 and Templin et al. (1993). A substantial amount of data is available in rats, so some data that  
 5 appeared to be redundant were excluded from the calibration set and saved for comparison with  
 6 posterior predictions (a “validation” set). In particular, those used for “validation” are one  
 7 closed-chamber experiment (Andersen et al., 1987b), several data sets with only TCE blood data  
 8 (D'Souza et al., 1985; Jakobson et al., 1986; Lee et al., 1996), and selected time courses from  
 9 Fisher et al. (1991) and Lee et al. (2000a; 2000b), and one unpublished data set (Bruckner et al.,  
 10 unpublished). The

11 **Table 3-33. PBPK model-based dose-metrics**

Abbreviation	Description
ABioactDCVCBW3 4	Amount of DCVC bioactivated in the kidney (mg) per unit body weight <sup>3/4</sup> (kg <sup>3/4</sup> )
ABioactDCVCKid	Amount of DCVC bioactivated in the kidney (mg) per unit kidney mass (kg)
AMetGSHBW34	Amount of TCE conjugated with GSH (mg) per unit body weight <sup>3/4</sup> (kg <sup>3/4</sup> )
AMetLiv1BW34	Amount of TCE oxidized in the liver per unit body weight <sup>3/4</sup> (kg <sup>3/4</sup> )
AMetLivOtherBW3 4	Amount of TCE oxidized to metabolites other than TCA and TCOH in the liver (mg) per unit body weight <sup>3/4</sup> (kg <sup>3/4</sup> )
AMetLivOtherLiv	Amount of TCE oxidized to metabolites other than TCA and TCOH in the liver (mg) per unit liver mass (kg)
AMetLngBW34	Amount of TCE oxidized in the respiratory tract (mg) per unit body weight <sup>3/4</sup> (kg <sup>3/4</sup> )
AMetLngResp	Amount of TCE oxidized in the respiratory tract (mg) per unit respiratory tract tissue mass (kg)
AUCCBld	Area under the curve of the venous blood concentration of TCE (mg-hour/L)
AUCCTCOH	Area under the curve of the blood concentration of TCOH (mg-hour/L)
AUCLivTCA	Area under the curve of the liver concentration of TCA (mg-hour/L)
TotMetabBW34	Total amount of TCE metabolized (mg) per unit body weight <sup>3/4</sup> (kg <sup>3/4</sup> )
TotOxMetabBW34	Total amount of TCE oxidized (mg) per unit body weight <sup>3/4</sup> (kg <sup>3/4</sup> )
TotTCAInBW	Total amount of TCA produced (mg) per unit body weight (kg)

13  
14

**Table 3-34. Rodent studies with pharmacokinetic data considered for analysis**

Reference	Species (strain)	Sex	TCE exposures	Other exposures	Calibration	Validation	Not used	Comments
Mouse studies								
Abbas et al. (1996)	Mouse (B6C3F1)	M	--	CH i.v.			√	CH not in model.
Abbas and Fisher (1997)	Mouse (B6C3F1)	M	Oral (corn oil)	--	√ <sup>a</sup>			
Abbas et al. (1997)	Mouse (B6C3F1)	M	--	TCOH, TCA i.v.	√			
Barton et al. (1999)	Mouse (B6C3F1)	M	--	DCA i.v. and oral (aqueous)			√	DCA not in model.
Birner et al. (1993)	Mouse (NMRI)	M+F	Gavage	--			√	Only urine concentrations available, not amount.
Fisher and Allen, (1993)	Mouse (B6C3F1)	M+F	Gavage (corn oil)	--	√			
Fisher et al. (1991)	Mouse (B6C3F1)	M+F	Inhalation	--	√ <sup>a</sup>			
Green and Prout (1985)	Mouse (B6C3F1)	M	Gavage (corn oil)	TCA i.v.	√			
Greenberg et al. (1999)	Mouse (B6C3F1)	M	Inhalation	--	√ <sup>a</sup>			
Larson and Bull (1992a)	Mouse (B6C3F1)	M	--	DCA, TCA oral (aqueous)	√			Only data on TCA dosing was used, since DCA is not in the model.
Larson and Bull (1992b)	Mouse (B6C3F1)	M	Oral (aqueous)	--	√			
Merdink et al. (1998)	Mouse (B6C3F1)	M	i.v.	CH i.v.	√			Only data on TCE dosing was used, since CH is not in the model.

This document is a draft for review purposes only and does not constitute Agency policy.

**Table 3-34. Rodent studies with pharmacokinetic data considered for analysis (continued)**

Reference	Species (strain)	Sex	TCE exposures	Other exposures	Calibration	Validation	Not used	Comments
Prout et al. (1985)	Mouse (B6C3F1, Swiss)	M	Gavage (corn oil)	--	√ <sup>a</sup>			
Templin et al. (1993)	Mouse (B6C3F1)	M	Oral (aqueous)	TCA oral	√ <sup>a</sup>			
Rat studies								
Andersen et al. (1997)	Rat (F344)	M	Inhalation	--		√ <sup>a</sup>		
Barton et al. (1995)	Rat (S-D)	M	Inhalation	--			√	Initial chamber concentrations unavailable, so not used.
Bernauer et al. (1996)	Rat (Wistar)	M	Inhalation	--	√ <sup>a</sup>			
Birner et al. (1993)	Rat (Wistar, F344)	M+F	Gavage (ns)	--			√	Only urine concentrations available, not amount.
Birner et al. (1997)	Rat (Wistar)	M+F	--	DCVC i.v.			√	Single dose, route does not recapitulate how DCVC is formed from TCE, excreted NAcDCVC ~100-fold greater than that from relevant TCE exposures (Bernauer et al., 1996).
Bruckner et al. unpublished	Rat (S-D)	M	Inhalation	--		√		Not published, so not used for calibration. Similar to Keys et al. (2003) data.
Dallas et al. (1991)	Rat (S-D)	M	Inhalation	--	√			
D'Souza et al. (1985)	Rat (S-D)	M	i.v., oral (aqueous)	--			√	Only TCE blood measurements, and ≥10-fold greater than other similar studies.
Fisher et al. (1989)	Rat (F344)	F	Inhalation	--	√			

Table 3-34. Rodent studies with pharmacokinetic data considered for analysis (continued)

Reference	Species (strain)	Sex	TCE exposures	Other exposures	Calibration	Validation	Not used	Comments
Fisher et al. (1991)	Rat (F344)	M+F	Inhalation	--	√ <sup>a</sup>	√		Experiment with blood only data not used for calibration.
Green and Prout (1985)	Rat (Osborne-Mendel)	M	Gavage (corn oil)	TCA gavage (aqueous)	√			
Hissink et al. (2002)	Rat (Wistar)	M	Gavage (corn oil), i.v.	--	√			
Jakobson et al. (1986)	Rat (S-D)	F	Inhalation	Various pretreatments (oral)		√		Pretreatments not included. Only blood TCE data available.
Kaneko et al. (1994a)	Rat (Wistar)	M	Inhalation	Ethanol pretreatment (oral)	√			Pretreatments not included.
Keys et al. (2003)	Rat (S-D)	M	Inhalation, oral (aqueous), i.a.	--	√			
Kimmerle and Eben (1973a)	Rat (Wistar)	M	Inhalation	--	√			
Larson and Bull (1992a)	Rat (F344)	M	--	DCA, TCA oral (aqueous)	√			Only TCA dosing data used, since DCA is not in the model.
Larson and Bull (1992b)	Rat (S-D)	M	Oral (aqueous)	--	√ <sup>a</sup>			
Lash et al. (2006)	Rat (F344)	M+F	Gavage (corn oil)	--			√	Highly inconsistent with other studies.
Lee et al. (1996)	Rat (S-D)	M	Arterial, venous, portal, stomach injections	--		√		Only blood TCE data available.
Lee et al. (2000a; 2000b)	Rat (S-D)	M	Stomach injection, i.v., p.v.	p-nitrophenol pretreatment (i.a.)	√	√		Pretreatments not included. Only experiments with blood and liver data used for calibration.

**Table 3-34. Rodent studies with pharmacokinetic data considered for analysis (continued)**

Reference	Species (strain)	Sex	TCE exposures	Other exposures	Calibration	Validation	Not used	Comments
Merdink et al. (1999)	Rat (F344)	M	--	CH, TCOH i.v.	√			TCOH dosing used; CH not in model.
Poet et al. (2000)	Rat (F344)	M	Dermal	--			√	Dermal exposure not in model.
Prout et al. (1985)	Rat (Osborne-Mendel, Wistar)	M	Gavage (corn oil)	--	√ <sup>a</sup>			
Saghir et al. (2002)	Rat (F344)	M	--	DCA i.v., oral (aqueous)			√	DCA not in model.
Simmons et al. (2002)	Rat (Long-Evans)	M	Inhalation	--	√			
Stenner et al. (1997)	Rat (F344)	M	intraduodenal	TCOH, TCA i.v.	√			
Templin et al. (1995b)	Rat (F344)	M	Oral (aqueous)	--	√ <sup>a</sup>			
Thrall et al. (2000)	Rat (F344)	M	i.v., i.p.	with toluene			√	Only exhaled breath data available from i.v. study. i.p. dosing not in model.
Yu et al. (2000)	Rat (F344)	M	--	TCA i.v.	√			

<sup>a</sup> Part or all of the data in the study was used for calibration in Hack et al. (2006).

i.a. = intra-arterial, i.p. = intraperitoneal, i.v. = intravenous, p.v. = intraperivenous.

**Table 3-35. Human studies with pharmacokinetic data considered for analysis**

Reference	Species (number of individuals)	Sex	TCE exposures	Other exposures	Calibration	Validation	Not used	Comments
Bartonicek (1962)	Human ( <i>n</i> = 8)	M+F	Inhalation	--		√		Sparse data, so not included for calibration to conserve computational resources.
Bernauer et al. (1996)	Human	M	Inhalation	--	√ <sup>a</sup>			Grouped data, but unique in that includes NAcDCVC urine data.
Bloemen et al. (2001)	Human ( <i>n</i> = 4)	M	Inhalation	--		√		Sparse data, so not included for calibration to conserve computational resources.
Chiu et al. (2007)	Human ( <i>n</i> = 6)	M	Inhalation	--	√			
Ertle et al. (1972)	Human	M	Inhalation	CH oral			√	Very similar to Muller data.
Fernandez et al. (1977)	Human	M	Inhalation	--		√		
Fisher et al. (1998)	Human ( <i>n</i> = 17)	M+F	Inhalation	--	√ <sup>a</sup>			
Kimmerle and Eben (1973b)	Human ( <i>n</i> = 12)	M+F	Inhalation	--	√			
Lapare et al. (1995)	Human ( <i>n</i> = 4)	M+F	Inhalation	--		√ <sup>b</sup>		Complex exposure patterns, and only grouped data available for urine, so used for validation.
Lash et al. (1999b)	Human	M+F	Inhalation	--	√			Grouped only, but unique in that DCVG blood data available (same individuals as Fisher et al. (1998)].
Monster et al. (1976)	Human ( <i>n</i> = 4)	M	Inhalation	--	√ <sup>b</sup>			Experiments with exercise not included.
Monster et al. (1979)	Human	M	Inhalation	--		√ <sup>a</sup>		Grouped data only.
Muller et al. (1972)	Human	ns	Inhalation	--			√	Same data also included in Muller et al. (1975).



**Table 3-35. Human studies with pharmacokinetic data considered for analysis (continued)**

Reference	Species (number of individuals)	Sex	TCE exposures	Other exposures	Calibration	Validation	Not used	Comments
Muller et al. (1974)	Human	M	Inhalation	CH, TCA, TCOH oral	√	√ <sup>a</sup>		TCA and TCOH dosing data used for calibration, since it is rare to have metabolite dosing data. TCE dosing data used for validation, since only grouped data available. CH not in model.
Muller et al. (1975)	Human	M	Inhalation	Ethanol oral		√ <sup>a</sup>		Grouped data only.
Paykoc et al. (1945)	Human ( <i>n</i> = 3)	ns	--	TCA i.v.	√			
Poet et al. (2000)	Human	M+F	Dermal	--				Dermal exposure not in model.
Sato et al. (1977)	Human	M	Inhalation	--		√		
Stewart et al. (1970)	Human	ns	Inhalation	--		√ <sup>a</sup>		
Treibig et al. (1976)	Human	ns	Inhalation	--		√ <sup>a</sup>		
Vesterberg and Astrand (1976)	Human	M	Inhalation	--			√	All experiments included exercise, so were not included.

<sup>a</sup> Part or all of the data in the study was used for calibration in Hack et al. (2006).

<sup>b</sup> Grouped data from this study was used for calibration in Hack et al. (2006), but individual data was used here.

1 Andersen et al. (1987b) data were selected randomly from the available closed-chamber data,  
2 while the other data sets were selected because they were unpublished or because they were  
3 more limited in scope (e.g., TCE blood only) and so were not as efficient for use in the  
4 computationally-intensive calibration stage. As with the mouse analyses, TCA and TCOH  
5 dosing data were incorporated to better calibrate those pathways.

6 The human pharmacokinetic database of controlled exposure studies is extensive but also  
7 more complicated. For the majority of the studies, only grouped or aggregated data were  
8 available, and most of those data were saved for “validation” since there remained a large  
9 number of studies for which individual data were available. However, some data that may be  
10 uniquely informative are only available in grouped form, in particular DCVG blood  
11 concentrations, NAcDCVC urinary excretion, and data from TCA and TCOH dosing. While  
12 there are analytic uncertainties as to the DCVG blood measurements, discussed above in Section  
13 3.3.3.2.1, they were nonetheless included here because they are the only in vivo data available on  
14 this measurement in humans. The uncertainty associated with their use is discussed below (see  
15 Section 3.5.7.3.2).

16 In addition, several human data sets, while having individual data, involved sparse  
17 collection at only one or a few time points per exposure (Bartonicek, 1962; Bloemen et al., 2001)  
18 and were subsequently excluded to conserve computational resources. Lapare et al. (1995),  
19 which involved multiple, complex exposure patterns over the course of a month and was missing  
20 the individual urine data, was also excluded due to the relatively low amount of data given the  
21 large computational effort required to simulate it. Several studies also investigated the effects of  
22 exercise during exposure on human TCE toxicokinetics. The additional parameters in a model  
23 including exercise would include those for characterizing the changes in cardiac output, alveolar  
24 ventilation, and regional blood flow as well as their interindividual variability, and would have  
25 further increased the computational burden. Therefore, it was decided that such data would be  
26 excluded from this analysis. Even with these exclusions, data on a total of 42 individuals, some  
27 involving multiple exposures, were included in the calibration.  
28

### 3.5.5.1.2. Updated Hierarchical Population Statistical Model and Prior Distributions

29 While the individual animals of a common strain and sex within a study are likely to vary  
30 to some extent, this variability was not included as part of the hierarchical population model for  
31 several reasons. First, generally, only aggregated pharmacokinetic data (arithmetic mean and  
32 standard deviation or standard error) are available from rodent studies. While methods exist for  
33 addressing between-animal variability with aggregated data (e.g., Chiu and Bois, 2007), they

1 require a higher level of computational intensity. Second, dose-response data are generally also  
2 only separated by sex and strain, and otherwise aggregated. Thus, in analyzing dose-response  
3 data (see Chapter 5), one usually has no choice but to treat all the animals in a particular study of  
4 a particular strain and sex as identical units. In the Hack et al. (2006) model, each simulation  
5 was treated as a separate observational unit, so different dosing levels within the same study  
6 were treated separately and assigned different PBPK model parameters. However, the animals  
7 within a study are generally inbred and kept under similarly controlled conditions, whereas  
8 animals in different studies—even if of the same strain and sex—likely have differences in  
9 genetic lineage, diet, and handling. Thus, animals *within* a study are likely to be much more  
10 homogeneous than animals *between* studies. As a consequence, in the revised model, for  
11 rodents, different animals of the same sex and strain in the same study (or series of studies  
12 conducted simultaneously) were treated as identical, and grouped together as a single “subject.”  
13 Thus, the predictions from the population model in rodents simulate “average” pharmacokinetics  
14 for a particular “lot” of rodents of a particular species, strain, and sex. Between-animal  
15 variability is not explicitly modeled, but it is incorporated in a “residual” error term as part of the  
16 likelihood function (see Appendix A, Section A.4.3.4). Therefore, a high degree of within-study  
17 variability would be reflected in a high posterior value in the variance of the residual-error.

18 In humans, however, interindividual variability is of interest, and, furthermore,  
19 substantial individual data are available in humans. However, in some studies, the same  
20 individual was exposed more than once, so those data should be grouped together (in the Hack  
21 et al. [2006] model, they were treated as different “individuals”). Because the primary interest  
22 here is chronic exposure, and because it would add substantially to the computational burden,  
23 interoccasion variability—changes in pharmacokinetic parameters in a single individual over  
24 time—is not addressed. Therefore, each individual is considered a single “subject,” and the  
25 predictions from the population model in humans are the “average” across different occasions for  
26 a particular individual (adult). Between-occasion variability is not explicitly modeled, but it is  
27 incorporated in a “residual” error term as part of the likelihood function (see Appendix A,  
28 Section A.4.3.4). Therefore, a high degree of between-occasion variability would be reflected in  
29 a high posterior value in the variance of the residual-error.

30 As discussed in Section 3.3.3.1, sex and (in rodents) strain differences in oxidative  
31 metabolism were modest or minimal. While some sex-differences have been noted in GSH  
32 metabolism (see Sections 3.3.3.2.7–3.3.3.2.8), almost all of the available *in vivo* data is in males,  
33 making it more difficult to statistically characterize that difference with PBPK modeling.  
34 Therefore, within a species, different sexes and (in rodents) strains were considered to be drawn

1 from a single, species-level population. For humans, each individual was considered to be drawn  
2 from a single (adult) human population.

3 Thus, from here forward, the term “subject” will be used to refer to both a particular “lot”  
4 of a particular rodents’ species, strain, and sex for, and a particular human individual. The term  
5 “population” will, therefore, refer to the collection of rodent “lots” of the same species and the  
6 collection of human individuals.

7 Figure A-1 in Appendix A illustrates the hierarchical structure. Informative prior  
8 distributions reflecting the uncertainty in the population mean and variance, detailed in  
9 Appendix A, were updated from those used in Hack et al. (2006) based on an extensive analysis  
10 of the available literature. The population variability of the scaling parameter across subjects is  
11 assumed to be distributed as a truncated normal distribution, a standard assumption in the  
12 absence of specific data suggesting otherwise. Because of the truncation of extreme values, the  
13 sensitivity to this choice is expected to be small as long as the true underlying distribution is  
14 uni-modal and symmetric. In addition, most scaling parameters, being strictly positive in their  
15 original units, were log-transformed—so these parameters have lognormal distributions in their  
16 original units. The uncertainty distribution for the population parameters was assumed to be a  
17 truncated normal distribution for population mean parameters and an inverse gamma distribution  
18 for population variance parameters—both standard choices in hierarchical models.  
19 Section 3.5.5.3, next, discusses specification of prior distributions in the case where no data  
20 independent of the calibration data exist.

### 3.5.5.1.3. Use of Interspecies Scaling to Update Prior Distributions in the Absence of Other Data

22 For many metabolic parameters, little or no in vitro or other prior information is available  
23 to develop prior distributions. Initially, for such parameters, noninformative priors in the form of  
24 log-uniform distributions with a range spanning at least  $10^4$  were specified. However, in the  
25 time available for analysis (up to about 100,000 iterations), only for the mouse did all these  
26 parameters achieve adequate convergence. This suggests that some of these parameters are  
27 poorly identified for the rat and human. Additional preliminary runs indicated replacing the  
28 log-uniform priors with lognormal priors and/or requiring more consistency between species  
29 could improve identifiability sufficiently for adequate convergence. However, an objective  
30 method of “centering” the lognormal distributions that did not rely on the in vivo data (e.g., via  
31 visual fitting or limited optimization) being calibrated against was necessary in order to  
32 minimize potential bias.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1           Therefore, the approach taken was to consider three species sequentially, from mouse to  
2 rat to human, and to use interspecies scaling to update the prior distributions across species. This  
3 sequence was chosen because the models are essentially “nested” in this order, the rat model  
4 adds to the mouse model the “downstream” GSH conjugation pathways, and the human model  
5 adds to the rat model the intermediary DCVG compartment. Therefore, for those parameters  
6 with little or no independent data *only*, the mouse posteriors were used to update the rat priors,  
7 and both the mouse and rat posteriors were used to update the human priors. Table 3-36 contains  
8 a list of the parameters for which this scaling was used to update prior distributions. The scaling  
9 relationship is defined by the “scaled parameters” listed in Appendix A (see Section A.4.1,  
10 Table A-4), and generally follows standard practice. For instance,  $V_{MAX}$  and clearance rates  
11 scale by body weight to the  $3/4$  power, whereas  $K_M$  values are assumed to not scale, and rate  
12 constants (inverse time units) scale by body weight to the  $-1/4$  power.

13           The scaling model is given explicitly as follows. If  $\theta_i$  are the “scaled” parameters  
14 (usually also natural-log-transformed) that are actually estimated, and  $A$  is the “universal”  
15 (species-independent) parameter, then  $\theta_i = A + \varepsilon_i$ , where  $\varepsilon_i$  is the species-specific “departure”  
16 from the scaling relationship, assumed to be normally distributed with variance  $\sigma_\varepsilon^2$ . Therefore,  
17 the mouse model gives an initial estimate of “ $A$ ,” which is used to update the priordistribution  
18 for  $\theta_r = A + \varepsilon_r$  in the rat. The rat and mouse together then give a “better” estimate of  $A$ , which is  
19 used to update the prior distribution for  $\theta_h = A + \varepsilon_h$  in the human, with the assumed distribution  
20 for  $\varepsilon_h$ . The mathematical details are given in Appendix A, but three key points in this model are  
21 worth noting here:

- 22
- 23
- 24       • It is known that interspecies scaling is not an exact relationship, and that, therefore, in  
25 any *particular* case it may either over- or underestimate. Therefore, the variance in the  
26 new priors reflect a combination of (1) the uncertainty in the “previous” species’  
27 posteriors as well as (2) a “prediction error” that is distributed lognormally withgeometric  
28 standard deviation (GSD) of 3.16-fold, so that the 95% confidence range about the  
29 central estimate spans 100-fold. This choice was dictated partially by practicality, as  
30 larger values of the GSD used in preliminary runs did not lead to adequate convergence  
31 within the time available for analysis.
- 32       • The rat posterior is a product of its prior (which is based on the mouse posterior) and its  
33 likelihood. Therefore, using the rat and mouse posteriors together to update the human  
34 priors would use the mouse posterior “twice.” Therefore, the rat posterior is  
35 disaggregated into its prior and its likelihood using a lognormal approximation (since the  
36 prior is lognormal), and only the (approximate) likelihood is used along with the mouse  
37 posterior to develop the human prior.
- 38       • The model transfers the marginal distributions for each parameter across species, so  
39 correlations between parameters are not retained. This is a restriction on the software

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 used for conducting MCMC analyses. However, assuming independence will lead to a  
2 “broader” joint distribution, given the same marginal distributions. Therefore, this  
3 assumption tends to reduce the weight of the interspecies scaling as compared to the  
4 species-specific calibration data.

**Table 3-36. Parameters for which scaling from mouse to rat, or from mouse and rat to human, was used to update the prior distributions**

Parameter with no or highly uncertain a priori data	Mouse → Rat	Rat → Human	Mouse+ Rat → Human	Comments
Respiratory lumen→tissue diffusion flow rate	√		√	No a priori information
TCOG body/blood partition coefficient	√		√	Prior centered on TCOH data, but highly uncertain
TCOG liver/body partition coefficient	√		√	Prior centered on TCOH data, but highly uncertain
Fraction of hepatic TCE oxidation not to TCA+TCOH	√		√	No a priori information
V <sub>MAX</sub> for hepatic TCE GSH conjugation	√			Rat data on at 1 and 2 mM. Human data at more concentrations, so V <sub>MAX</sub> and K <sub>M</sub> can be estimated
K <sub>M</sub> for hepatic TCE GSH conjugation	√			
V <sub>MAX</sub> for renal TCE GSH conjugation	√			Rat data on at 1 and 2 mM. Human data at more concentrations, so V <sub>MAX</sub> and K <sub>M</sub> can be estimated
K <sub>M</sub> for renal TCE GSH conjugation	√			
V <sub>MAX</sub> for Tracheo-bronchial TCE oxidation	√		√	Prior based on activity at a single concentration
K <sub>M</sub> for Tracheo-bronchial TCE oxidation	√		√	No a priori information
Fraction of respiratory oxidation entering systemic circulation	√		√	No a priori information
V <sub>MAX</sub> for hepatic TCOH→TCA	√		√	No a priori information
K <sub>M</sub> for hepatic TCOH→TCA	√		√	No a priori information
V <sub>MAX</sub> for hepatic TCOH→TCOG	√		√	No a priori information
K <sub>M</sub> for hepatic TCOH→TCOG	√		√	No a priori information
Rate constant for hepatic TCOH→other	√		√	No a priori information
Rate constant for TCA plasma→urine	√		√	Prior centered at GFR, but highly uncertain
Rate constant for hepatic TCA→other	√		√	No a priori information
Rate constant for TCOG liver→bile	√		√	No a priori information
Lumped rate constant for TCOG bile→TCOH liver	√		√	No a priori information
Rate constant for TCOG→urine	√		√	Prior centered at GFR, but highly uncertain
Lumped rate constant for DCVC→Urinary NAcDCVC		√		Not included in mouse model
Rate constant for DCVC bioactivation		√		Not included in mouse model

GFR = glomerular filtration rate.  
See Appendix A, Table A-4 for scaling relationships.

1 To summarize, in order to improve rate of the convergence of the MCMC analyses in rats  
2 and humans, a sequential approach was used for fitting scaling parameters without strong prior  
3 species-specific information. In particular, an additional assumption was made that *across*  
4 *species*, these scaling parameters were, in absence of other information, expected to have a  
5 common underlying value. These assumptions are generally based on allometric scaling  
6 principles—with partition coefficients and concentrations scaling directly and rate constants  
7 scaling by  $BW^{-1/4}$  (so clearances and maximum metabolic capacities would scale by  $BW^{3/4}$ ).  
8 These assumptions are used consistently throughout the parameter calibration process.  
9 Therefore, after running the mouse model, the posterior distribution for these parameters was  
10 used, with an additional error term, as priors for the rat model. Subsequently, after the mouse  
11 and rat model were run, their posterior distributions were combined, with an additional error  
12 term, to use as priors for the human model. With this methodology for updating the prior  
13 distributions, adequate convergence was achieved for the rat and human after  
14 110,000~140,000 iterations (discussed further below).  
15

#### **3.5.5.1.4. Implementation**

16 The PBPK model was coded in for use in the MCSim software (version 5.0.0), which was  
17 developed particularly for implementing MCMC simulations. As a quality control (QC) check,  
18 results were checked against the original Hack et al. (2006) model, with the original structures  
19 restored and parameter values made equivalent, and the results were within the error tolerances  
20 of the ordinary differential equation (ODE) solver after correcting an error in the Hack et al.  
21 (2006) model for calculating the TCA liver plasma flow. In addition, the model was translated to  
22 MatLab (version 7.2.0.232) with simulation results checked and found to be within the error  
23 tolerances of the ODE solver used (“ode15s”). Mass balances were also checked using the  
24 baseline parameters, as well as parameters from preliminary MCMC simulations, and found to  
25 be within the error tolerances of the ODE solver. Appendix A contains the MCSim model code.  
26

### **3.5.6. Evaluation of Updated Physiologically Based Pharmacokinetic (PBPK) Model**

#### **3.5.6.1.1. Convergence**

27 As in previous similar analyses (Bois, 2000a, b; David et al., 2006; Gelman et al., 1996;  
28 Hack et al., 2006), the potential scale reduction factor “*R*” is used to determine whether different

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 independent MCMC chains have converged to a common distribution. The  $R$  diagnostic is  
2 calculated for each parameter in the model, and represents the factor by which the standard  
3 deviation or other measure of scale of the posterior distribution (such as a confidence interval  
4 [CI]) may potentially be reduced with additional samples (Gelman et al., 2004). This  
5 convergence diagnostic declines to 1 as the number of simulation iterations approaches infinity,  
6 so values close to 1 indicate approximate convergence, with values of 1.1 and below commonly  
7 considered adequate (Gelman et al., 2004). However, as an additional diagnostic, the  
8 convergence of model dose-metric predictions was also assessed. Specifically, dose-metrics for  
9 a number of generic exposure scenarios similar to those used in long-term bioassays were  
10 generated, and their natural log (due to their approximate lognormal posterior distributions) was  
11 assessed for convergence using the potential scale reduction factor “ $R$ .” This is akin to the idea  
12 of utilizing sensitivity analysis so that effort is concentrated on calibrating the most sensitive  
13 parameters for the purpose of interest. In addition, predictions of interest which do not  
14 adequately converge can be flagged as such, so that the statistical uncertainty associated with the  
15 limited sample size can be considered.

16 The mouse model had the most rapid reduction in potential scale reduction factors.  
17 Initially, four chains of 42,500 iterations each were run, with the first 12,500 discarded as  
18 “burn-in” iterations. The initial decision for determining “burn-in” was determined by visual  
19 inspection. At this point, evaluating the 30,000 remaining iterations, all the population  
20 parameters except for the  $V_{MAX}$  for DCVG formation had  $R < 1.2$ , with only the first-order  
21 clearance rate for DCVG formation and the  $V_{MAX}$  and  $K_M$  for TCOH glucuronidation having  
22  $R > 1.1$ . For the samples used for inference, all of these initial iterations were treated as  
23 “burn-in” iterations, and each chain was then restarted and run for an additional  
24 68,700–71,400 iterations (chains were terminated at the same time, so the number of iterations  
25 per chains was slightly different). For these iterations, all values of  $R$  were  $< 1.03$ . Dose-metric  
26 predictions calculated for exposure scenarios 10–600 ppm either continuously or 7 hour/day,  
27 5 day/week and 10–3,000 mg/kg-day either continuously or by gavage 5 day/week. These  
28 predictions were all adequately converged, with all values of  $R < 1.03$ .

29 As discussed above, for parameters with little or no a priori information, the posterior  
30 distributions from the mouse model were used to update prior distributions for the rat model,  
31 accounting for both the uncertainty reflected in the mouse posteriors as well as the uncertainty in  
32 interspecies extrapolation. Four chains were run to 111,960–128,000 iterations each (chains  
33 were terminated at the same time and run on computers with slightly different processing speeds,  
34 so the number of iterations per chains was slightly different). As is standard, about the  
35 first “half” of the chains—i.e., the first 64,000 iterations—were discarded as “burn-in” iterations,

1 and the remaining iterations were used for inferences. For these remaining iterations, the  
2 diagnostic  $R$  was  $<1.1$  for all population parameters except the fraction of oxidation not  
3 producing TCA or TCOH ( $R = 1.44$  for population mean,  $R = 1.35$  for population variance), the  
4  $K_M$  for TCOH  $\rightarrow$  TCA ( $R = 1.19$  for population mean), the  $V_{MAX}$  and  $K_M$  for TCOH  
5 glucuronidation ( $R = 1.23$  and  $1.12$ , respectively for population mean, and  $R = 1.13$  for both  
6 population variances), and the rate of “other” metabolism of TCOH ( $R = 1.29$  for population  
7 mean and  $R = 1.18$  for population variance). Due to resource constraints, chains needed to be  
8 stopped at this point. However, these are similar to the degree of convergence reported in Hack  
9 et al. (2006). Dose-metric predictions calculated for two inhalation exposure scenarios  
10 (10–600 ppm continuously or 7 hours/day, 5 day/week) and two oral exposure scenarios  
11 (10–3,000 mg/kg-day continuously or by gavage 5 day/week).

12 All dose-metric predictions had  $R < 1.04$ , except for the amount of “other” oxidative  
13 metabolism (i.e., not producing TCA or TCOH), which had  $R = 1.12$ – $1.16$ , depending on the  
14 exposure scenario. The poorer convergence of this dose-metric is expected given that a key  
15 determining parameter, the fraction of oxidation not producing TCA or TCOH, had the poorest  
16 convergence among the population parameters.

17 For the human model, a set of four chains was run for 74,160–84,690 iterations using  
18 “preliminary” updated prior distributions based on the mouse posteriors and preliminary runs of  
19 the rat model. Once the rat chains were completed, final updated prior distributions were  
20 calculated and the last iteration of the preliminary runs were used as starting points for the final  
21 runs. The center of the final updated priors shifted by less than 25% of the standard deviation of  
22 either the preliminary or revised priors, so that the revised median was between the  
23 40<sup>th</sup> percentile and 60<sup>th</sup> percentile of the preliminary median, and vice versa. The standard  
24 deviations changed by less than 5%. Therefore, the use of the preliminary chains as a starting  
25 point should introduce no bias, as long as an appropriate burn-in period is used for the final runs.

26 The final chains were run for an additional 59,140–61,780 iterations, at which point, due  
27 to resource constraints, chains needed to be stopped. After the first 20,000 iterations, visual  
28 inspection revealed the chains were no longer dependent on the starting point. These iterations  
29 were therefore discarded as “burn-in” iterations, and for the remaining ~40,000 iterations used  
30 for inferences. All population mean parameters had  $R < 1.1$  except for the respiratory tract  
31 diffusion constant ( $R = 1.20$ ), the liver:blood partition coefficient for TCOG ( $R = 1.23$ ), the rate  
32 of TCE clearance in the kidney producing DCVG ( $R = 1.20$ ), and the rate of elimination of  
33 TCOG in bile ( $R = 1.46$ ). All population variances also had  $R < 1.1$  except for the variance for  
34 the fraction of oxidation not producing TCOH or TCA ( $R = 1.10$ ). Dose-metric predictions were

1 assessed for continuous exposure scenarios at 1–60 ppm in air or 1–300 mg/kg-day orally.  
2 These predictions were all adequately converged with all values of  $R < 1.02$ .  
3

### 3.5.6.1.2. Evaluation of Posterior Parameter Distributions

4 Posterior distributions of the population parameters need to be checked as to whether  
5 they appear reasonable given the prior distributions. Inconsistency between the prior and  
6 posterior distributions may indicate insufficiently broad (i.e., due to overconfidence) or  
7 otherwise incorrectly specified priors, a misspecification of the model structure (e.g., leading to  
8 pathological parameter estimates), or an error in the data. As was done with the evaluation of  
9 Hack et al. (2006) in Appendix A, parameters were flagged if the interquartile regions of their  
10 prior and posterior distributions did not overlap.

11 Appendix A contains detailed tables of the “sampled” parameters, and their prior and  
12 posterior distributions. Because these parameters are generally scaled one or more times to  
13 obtain a physically meaningful parameter, they are difficult to interpret. Therefore, in  
14 Tables 3-37–3-39, the prior and posterior population distributions for the PBPK model  
15 parameters obtained *after* scaling are summarized. Since it is desirable to characterize the  
16 contributions from both uncertainty in population parameters and variability within the  
17 population, the following procedure is adopted. First, 500 sets of population parameters (i.e.,  
18 population mean and variance for each scaling parameter) are either generated from the prior  
19 distributions via Monte Carlo or extracted from the posterior MCMC samples—these represent  
20 the uncertainty in the population parameters. To minimize autocorrelation, for the posteriors, the  
21 samples were obtained by “thinning” the chains to the appropriate degree. From each of these  
22 sets of population parameters, 100 sets of “subject”-level parameters were generated by Monte  
23 Carlo—each of these represents the population variability, given a *particular* set of population  
24 parameters. Thus, a total of 50,000 subjects, representing 100 (variability) each for 500 different  
25 populations (uncertainty), were generated. For each of the 500 populations, the scaling  
26 parameters are converted to PBPK model parameters, and the population median and GSD is  
27 calculated—representing the central tendency and variability for that population. Then, the  
28 median and the 95% CIs for the population median and GSD are calculated, and presented in the  
29 tables that follow. Thus, these tables summarize separately the uncertainty in population  
30 distribution as well as the variability in the population, while also accounting for correlations  
31 among the population-level parameters. Finally, Table 3-40 shows the change in the CI in the  
32 population median for the PBPK model parameters between the prior and posterior distributions,

1 as well as the shift in the central estimate (median) of the population median PBPK model  
2 parameter.

3 The prior and posterior distributions for most physiological parameters were similar. The  
4 posterior distribution was substantially narrower (i.e., less uncertainty) than the prior distribution  
5 only in the case of the diffusion rate from the respiratory lumen to the respiratory tissue, which  
6 also was to be expected given the very wide, noninformative prior for that parameter.

7 For distribution parameters, there were only relatively minor changes between prior and  
8 posterior distributions for TCE and TCOH partition coefficients. The posterior distributions for  
9 several TCA partition coefficients and plasma binding parameters were substantially narrower  
10 than their corresponding priors, but the central estimates were similar, meaning that values at the  
11 high and low extremes were not likely. For TCOG as well, partition coefficient posterior  
12 distributions were substantially narrower, which was expected given the greater uncertainty in  
13 the prior distributions (TCOH partition coefficients were used as a proxy).

14 Again, posterior distributions indicated that the high and low extremes were not likely.  
15 Finally, posterior distribution for the distribution volume for DCVG was substantially narrower  
16 than the prior distribution, which only provided a lower bound given by the blood volume. In  
17 this case, the upper bounds were substantially lower in the posterior.

18 Posterior distributions for oral absorption parameters in mice and rats (there were no oral  
19 studies in humans) were also informed by the data, as reflected in their being substantially more  
20 narrow than the corresponding priors. Finally, with a few exceptions, TCE and metabolite  
21 kinetic parameters showed substantially narrower posterior distributions than prior distributions,  
22 indicating that they were fairly well specified by the in vivo data. The exceptions were the  $V_{MAX}$   
23 for hepatic oxidation in humans (for which there was substantial in vitro data) and the  $V_{MAX}$  for  
24 respiratory metabolism in mice and rats (although the posterior distribution for the  $K_M$  for this  
25 pathway was substantially narrower than the corresponding prior).

26 However, for some parameters, the posterior distributions in the population medians had  
27 CIs greater than 100-fold. In mice, the absorption parameter for TCA still had posterior CI of  
28 400-fold, reflecting the fact that the absorption rate is poorly estimated from the few available  
29 studies with TCA dosing. In addition, mouse metabolism parameters for GSH conjugation have  
30 posterior CIs greater than 10,000-fold, reflecting the lack of any direct data on GSH conjugation  
31 in mice. In rats, two parameters related to TCOH and TCOG had CIs between 100- and  
32 1,000-fold, reflecting the poor identifiability of these parameters given the available data. In  
33 humans, only the oral absorption parameters for TCA and TCOH had CIs greater than 100-fold,  
34 reflecting the fact that the absorption rate is poorly estimated from the few available studies with  
35 TCOH and TCA dosing.

1           In terms of general consistency between prior and posterior distributions, in most cases,  
2 the central estimate of the population median shifted by less than threefold. In almost all the  
3 cases that the shift was greater (see bold entries in Table 3-40), the prior distribution had a wide  
4 distribution, with CI greater (sometimes substantially greater) than 100-fold. The only exception

**Table 3-37. Prior and posterior uncertainty and variability in mouse PBPK model parameters**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Cardiac output (L/hour)	QC	0.84 (0.59, 1.2)	1 (0.79, 1.3)	1.17 (1.1, 1.4)	1.35 (1.15, 1.54)
Alveolar ventilation (L/hour)	QP	2.1 (1.3, 3.5)	2.1 (1.5, 2.7)	1.27 (1.17, 1.54)	1.45 (1.28, 1.66)
Scaled fat blood flow	QFatC	0.07 (0.03, 0.11)	0.072 (0.044, 0.1)	1.65 (1.22, 2.03)	1.64 (1.3, 1.99)
Scaled gut blood flow	QGutC	0.14 (0.11, 0.17)	0.16 (0.14, 0.17)	1.15 (1.09, 1.19)	1.12 (1.07, 1.19)
Scaled liver blood flow	QLivC	0.02 (0.016, 0.024)	0.021 (0.017, 0.024)	1.15 (1.09, 1.19)	1.15 (1.09, 1.19)
Scaled slowly perfused blood flow	QSlwC	0.22 (0.14, 0.29)	0.21 (0.15, 0.28)	1.3 (1.15, 1.38)	1.3 (1.17, 1.39)
Scaled rapidly perfused blood flow	QRapC	0.46 (0.37, 0.56)	0.45 (0.37, 0.52)	1.15 (1.11, 1.2)	1.17 (1.12, 1.2)
Scaled kidney blood flow	QKidC	0.092 (0.054, 0.13)	0.091 (0.064, 0.12)	1.34 (1.14, 1.45)	1.34 (1.18, 1.44)
Respiratory lumen:tissue diffusive clearance rate (L/hour)	DResp	0.017 (3.2e-05, 15)	2.5 (1.4, 5.1)	1.37 (1.25, 1.62)	1.53 (1.37, 1.73)
Fat fractional compartment volume	VFatC	0.071 (0.032, 0.11)	0.089 (0.061, 0.11)	1.59 (1.19, 1.93)	1.4 (1.19, 1.78)
Gut fractional compartment volume	VGutC	0.049 (0.041, 0.057)	0.048 (0.042, 0.055)	1.11 (1.07, 1.14)	1.11 (1.08, 1.14)
Liver fractional compartment volume	VLivC	0.054 (0.038, 0.071)	0.047 (0.037, 0.06)	1.22 (1.12, 1.29)	1.23 (1.17, 1.3)
Rapidly perfused fractional compartment volume	VRapC	0.1 (0.087, 0.11)	0.099 (0.09, 0.11)	1.08 (1.05, 1.11)	1.09 (1.06, 1.11)
Fractional volume of respiratory lumen	VRespLumC	0.0047 (0.004, 0.0053)	0.0047 (0.0041, 0.0052)	1.09 (1.06, 1.12)	1.09 (1.07, 1.12)
Fractional volume of respiratory tissue	VRespEffC	7e-04 (6e-04, 0.00079)	7e-04 (0.00062, 0.00078)	1.09 (1.06, 1.12)	1.1 (1.07, 1.12)
Kidney fractional compartment volume	VKidC	0.017 (0.015, 0.019)	0.017 (0.015, 0.019)	1.08 (1.05, 1.11)	1.09 (1.06, 1.11)
Blood fractional compartment volume	VBldC	0.049 (0.042, 0.056)	0.048 (0.043, 0.054)	1.1 (1.06, 1.13)	1.1 (1.08, 1.13)

**Table 3-37. Prior and posterior uncertainty and variability in mouse PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Slowly perfused fractional compartment volume	VSlwC	0.55 (0.5, 0.59)	0.54 (0.51, 0.57)	1.05 (1.04, 1.07)	1.05 (1.04, 1.07)
Plasma fractional compartment volume	VPlasC	0.026 (0.016, 0.036)	0.022 (0.016, 0.029)	1.24 (1.15, 1.35)	1.27 (1.19, 1.36)
TCA body fractional compartment volume [not incl. blood+liver]	VBodC	0.79 (0.77, 0.8)	0.79 (0.78, 0.81)	1.01 (1.01, 1.02)	1.01 (1.01, 1.02)
TCOH/G body fractional compartment volume [not incl. liver]	VBodTCOHC	0.84 (0.82, 0.85)	0.84 (0.83, 0.85)	1.01 (1.01, 1.02)	1.01 (1.01, 1.02)
TCE blood:air partition coefficient	PB	15 (10, 23)	14 (11, 17)	1.22 (1.12, 1.42)	1.44 (1.28, 1.53)
TCE fat:blood partition coefficient	PFat	36 (21, 62)	36 (26, 49)	1.26 (1.14, 1.52)	1.32 (1.16, 1.56)
TCE gut:blood partition coefficient	PGut	1.9 (0.89, 3.8)	1.5 (0.94, 2.6)	1.36 (1.2, 1.75)	1.36 (1.2, 1.79)
TCE liver:blood partition coefficient	PLiv	1.7 (0.89, 3.5)	2.2 (1.3, 3.3)	1.37 (1.2, 1.75)	1.39 (1.21, 1.84)
TCE rapidly perfused:blood partition coefficient	PRap	1.8 (0.98, 3.7)	1.8 (1.1, 3)	1.37 (1.2, 1.76)	1.37 (1.2, 1.77)
TCE respiratory tissue:air partition coefficient	PResp	2.7 (1.2, 5)	2.5 (1.5, 4.2)	1.36 (1.19, 1.78)	1.37 (1.19, 1.74)
TCE kidney:blood partition coefficient	PKid	2.2 (0.96, 4.6)	2.6 (1.7, 4)	1.36 (1.2, 1.77)	1.51 (1.25, 1.88)
TCE slowly perfused:blood partition coefficient	PSlw	2.4 (1.2, 4.9)	2.2 (1.4, 3.5)	1.38 (1.2, 1.78)	1.39 (1.21, 1.8)
TCA blood:plasma concentration ratio	TCAPlas	0.76 (0.4, 16)	1.1 (0.75, 1.8)	1.21 (1.09, 1.58)	1.23 (1.1, 1.73)

**Table 3-37. Prior and posterior uncertainty and variability in mouse PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Free TCA body:blood plasma partition coefficient	PBodTCA	0.77 (0.27, 17)	0.87 (0.59, 1.5)	1.41 (1.23, 1.8)	1.39 (1.24, 1.9)
Free TCA liver:blood plasma partition coefficient	PLivTCA	1.1 (0.36, 21)	1.1 (0.64, 1.9)	1.41 (1.23, 1.8)	1.4 (1.24, 1.87)
Protein:TCA dissociation constant (µmole/L)	kDissoc	100 (13, 790)	130 (24, 520)	2.44 (1.73, 5.42)	2.64 (1.75, 5.45)
Maximum binding concentration (µmole/L)	B <sub>MAX</sub>	87 (9.6, 790)	140 (28, 690)	2.72 (1.92, 5.78)	2.88 (1.93, 5.89)
TCOH body:blood partition coefficient	PBodTCOH	1.1 (0.61, 2.1)	0.89 (0.65, 1.3)	1.29 (1.16, 1.66)	1.31 (1.17, 1.61)
TCOH liver:body partition coefficient	PLivTCOH	1.3 (0.73, 2.3)	1.9 (1.2, 2.6)	1.3 (1.16, 1.61)	1.35 (1.18, 1.68)
TCOG body:blood partition coefficient	PBodTCOG	0.95 (0.016, 77)	0.48 (0.18, 1.1)	1.36 (1.19, 2.05)	1.41 (1.22, 2.19)
TCOG liver:body partition coefficient	PLivTCOG	1.3 (0.019, 92)	1.3 (0.64, 2.6)	1.36 (1.18, 2.13)	1.56 (1.28, 2.52)
DCVG effective volume of distribution	VDCVG	0.033 (0.0015, 15)	0.027 (0.0016, 4.1)	1.28 (1.08, 1.97)	1.31 (1.1, 2.19)
TCE stomach absorption coefficient (/hour)	kAS	1.7 (0.0049, 450)	1.7 (0.37, 13)	4.74 (2.29, 23.4)	4.28 (2.39, 13.4)
TCE stomach-duodenum transfer coefficient (/hour)	kTSD	1.4 (0.043, 51)	4.5 (0.51, 26)	3.84 (2.09, 10.6)	4.79 (2.53, 10.9)
TCE duodenum absorption coefficient (/hour)	kAD	1.2 (0.0024, 200)	0.27 (0.067, 1.6)	4.33 (2.14, 26)	4.17 (2.34, 14.4)
TCA stomach absorption coefficient (/hour)	kASTCA	0.63 (0.0027, 240)	4 (0.2, 74)	4.26 (2.27, 23.4)	5.15 (2.56, 22)
V <sub>MAX</sub> for hepatic TCE oxidation (mg/hour)	V <sub>MAX</sub>	3.9 (1.4, 15)	2.5 (1.6, 4.2)	2.02 (1.56, 2.85)	1.86 (1.59, 2.47)
K <sub>M</sub> for hepatic TCE oxidation (mg/L)	K <sub>M</sub>	34 (1.6, 620)	2.7 (1.4, 8)	1.25 (1.15, 1.61)	2.08 (1.48, 3.49)



**Table 3-37. Prior and posterior uncertainty and variability in mouse PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Fraction of hepatic TCE oxidation not to TCA+TCOH	FracOther	0.43 (0.0018, 1)	0.023 (0.0037, 0.15)	1.23 (1, 2.13)	1.49 (1.25, 2.83)
Fraction of hepatic TCE oxidation to TCA	FracTCA	0.086 (0.00022, 0.66)	0.13 (0.084, 0.21)	1.48 (1.12, 2.56)	1.4 (1.21, 1.96)
V <sub>MAX</sub> for hepatic TCE GSH conjugation (mg/hour)	V <sub>MAX</sub> DCVG	3.7 (0.0071, 2800)	0.6 (0.01, 480)	1.55 (1.33, 2.52)	1.61 (1.37, 2.91)
K <sub>M</sub> for hepatic TCE GSH conjugation (mg/L)	K <sub>M</sub> DCVG	250 (0.0029, 6500000)	2200 (0.17, 2300000)	1.81 (1.47, 3.62)	1.93 (1.49, 3.68)
V <sub>MAX</sub> for renal TCE GSH conjugation (mg/hour)	V <sub>MAX</sub> KidDCVG	0.34 (0.00051, 180)	0.027 (0.0012, 13)	1.49 (1.26, 2.49)	1.54 (1.28, 2.72)
K <sub>M</sub> for renal TCE GSH conjugation (mg/L)	K <sub>M</sub> KidDCVG	150 (0.0053, 6200000)	160 (0.078, 280000)	1.79 (1.43, 3.45)	1.91 (1.5, 3.91)
V <sub>MAX</sub> for tracheo-bronchial TCE oxidation (mg/hour)	V <sub>MAX</sub> Clara	0.24 (0.03, 3.9)	0.42 (0.1, 1.5)	2.32 (1.74, 3.66)	4.13 (2.27, 6.79)
K <sub>M</sub> for tracheo-bronchial TCE oxidation (mg/L)	K <sub>M</sub> Clara	1.5 (0.0018, 630)	0.011 (0.0024, 0.09)	1.47 (1.25, 2.58)	1.63 (1.28, 5.02)
Fraction of respiratory metabolism to systemic circ.	FracLungSys	0.34 (0.0016, 1)	0.78 (0.18, 0.99)	1.24 (1, 2.1)	1.11 (1, 1.72)
V <sub>MAX</sub> for hepatic TCOH→TCA (mg/hour)	V <sub>MAX</sub> TCOH	0.064 (1.4e-05, 380)	0.12 (0.048, 0.28)	1.5 (1.24, 2.61)	1.6 (1.28, 2.92)
K <sub>M</sub> for hepatic TCOH→TCA (mg/L)	K <sub>M</sub> TCOH	1.4 (0.00018, 5300)	0.92 (0.26, 2.7)	1.48 (1.24, 2.41)	1.49 (1.26, 2.4)
V <sub>MAX</sub> for hepatic TCOH→TCOG (mg/hour)	V <sub>MAX</sub> Gluc	0.11 (1.3e-05, 310)	4.6 (1.9, 16)	1.48 (1.26, 2.53)	1.47 (1.26, 2.14)
K <sub>M</sub> for hepatic TCOH→TCOG (mg/L)	K <sub>M</sub> Gluc	1.8 (0.0018, 610)	30 (5.3, 130)	1.48 (1.25, 2.48)	1.8 (1.3, 4.72)
Rate constant for hepatic TCOH→other (/hour)	kMetTCOH	0.19 (3.9e-05, 1400)	8.8 (1.9, 23)	1.47 (1.25, 2.36)	1.54 (1.26, 2.92)

**Table 3-37. Prior and posterior uncertainty and variability in mouse PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Rate constant for TCA plasma→urine (/hour)	kUrnTCA	32 (0.38, 1700)	3.2 (1.2, 7.1)	1.57 (1.34, 2.61)	1.84 (1.44, 2.94)
Rate constant for hepatic TCA→other (/hour)	kMetTCA	0.12 (4e-04, 130)	1.5 (0.63, 2.9)	1.48 (1.25, 2.32)	1.51 (1.26, 2.27)
Rate constant for TCOG liver→bile (/hour)	kBile	0.3 (4e-04, 160)	2.4 (0.74, 8.4)	1.48 (1.24, 2.29)	1.51 (1.26, 2.39)
Lumped rate constant for TCOG bile→TCOH liver (/hour)	kEHR	0.21 (0.00036, 150)	0.039 (0.0026, 0.11)	1.47 (1.23, 2.29)	1.53 (1.28, 2.94)
Rate constant for TCOG→urine (/hour)	kUrnTCOG	1 (0.00015, 6200)	12 (2.6, 77)	1.71 (1.4, 3.13)	3.44 (1.89, 9.49)
Rate constant for hepatic DCVG→DCVC (/hour)	kDCVG	0.24 (4e-04, 160)	0.81 (0.0033, 46)	1.48 (1.25, 2.39)	1.52 (1.25, 2.5)
Lumped rate constant for DCVC→urinary NAcDCVC (/hour)	kNAT	0.29 (4e-04, 160)	0.37 (0.0021, 34)	1.5 (1.25, 2.49)	1.53 (1.25, 2.77)
Rate constant for DCVC bioactivation (/hour)	kKidBioact	0.18 (4e-04, 150)	0.23 (0.0024, 33)	1.48 (1.25, 2.51)	1.53 (1.25, 3.03)

**Table 3-38. Prior and posterior uncertainty and variability in rat PBPK model parameters**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Cardiac output (L/hour)	QC	5.3 (4.2, 6.9)	6.1 (5.2, 7.4)	1.12 (1.07, 1.28)	1.26 (1.12, 1.36)
Alveolar ventilation (L/hour)	QP	10 (5.1, 18)	7.5 (5.8, 10)	1.32 (1.18, 1.71)	1.52 (1.33, 1.84)
Scaled fat blood flow	QFatC	0.071 (0.032, 0.11)	0.081 (0.06, 0.1)	1.66 (1.21, 2.02)	1.5 (1.3, 1.86)
Scaled gut blood flow	QGutC	0.15 (0.12, 0.18)	0.17 (0.15, 0.19)	1.15 (1.09, 1.19)	1.13 (1.08, 1.18)
Scaled liver blood flow	QLivC	0.021 (0.017, 0.026)	0.022 (0.018, 0.025)	1.15 (1.09, 1.2)	1.15 (1.1, 1.19)
Scaled slowly perfused blood flow	QSlwC	0.33 (0.21, 0.46)	0.31 (0.23, 0.4)	1.31 (1.15, 1.4)	1.32 (1.22, 1.41)
Scaled rapidly perfused blood flow	QRapC	0.28 (0.15, 0.42)	0.28 (0.18, 0.36)	1.38 (0.0777, 1.72)	1.42 (0.0856, 1.75)
Scaled kidney blood flow	QKidC	0.14 (0.12, 0.16)	0.14 (0.12, 0.16)	1.11 (1.07, 1.14)	1.11 (1.08, 1.14)
Respiratory lumen:tissue diffusive clearance rate (L/hour)	DResp	9.9 (0.48, 85)	21 (9.5, 46)	1.41 (1.26, 1.77)	1.59 (1.41, 1.9)
Fat fractional compartment volume	VFatC	0.069 (0.031, 0.11)	0.069 (0.046, 0.091)	1.61 (1.2, 1.93)	1.59 (1.34, 1.88)
Gut fractional compartment volume	VGutC	0.032 (0.027, 0.037)	0.032 (0.028, 0.036)	1.11 (1.07, 1.14)	1.11 (1.08, 1.14)
Liver fractional compartment volume	VLivC	0.034 (0.026, 0.042)	0.033 (0.028, 0.039)	1.16 (1.09, 1.21)	1.17 (1.12, 1.21)
Rapidly perfused fractional compartment volume	VRapC	0.087 (0.076, 0.1)	0.088 (0.079, 0.097)	1.1 (1.06, 1.13)	1.1 (1.07, 1.13)
Fractional volume of respiratory lumen	VRespLumC	0.0046 (0.0037, 0.0057)	0.0047 (0.0039, 0.0055)	1.16 (1.1, 1.21)	1.16 (1.11, 1.21)
Fractional volume of respiratory tissue	VRespEffC	5e-04 (0.00039, 0.00061)	5e-04 (0.00041, 0.00058)	1.16 (1.09, 1.21)	1.16 (1.11, 1.2)
Kidney fractional compartment volume	VKidC	0.0069 (0.0056, 0.0082)	0.007 (0.006, 0.008)	1.13 (1.08, 1.17)	1.13 (1.09, 1.17)

**Table 3-38 Prior and posterior uncertainty and variability in rat PBPK model parameters (contuined)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Blood fractional compartment volume	VBldC	0.073 (0.063, 0.085)	0.074 (0.066, 0.082)	1.1 (1.06, 1.13)	1.1 (1.07, 1.13)
Slowly perfused fractional compartment volume	VSlwC	0.6 (0.55, 0.63)	0.6 (0.57, 0.62)	1.05 (1.04, 1.06)	1.05 (1.04, 1.06)
Plasma fractional compartment volume	VPlasC	0.039 (0.025, 0.054)	0.04 (0.032, 0.049)	1.24 (1.15, 1.35)	1.22 (1.16, 1.33)
TCA body fractional compartment volume [not incl. blood+liver]	VBodC	0.79 (0.78, 0.81)	0.79 (0.78, 0.8)	1.01 (1.01, 1.01)	1.01 (1.01, 1.01)
TCOH/G body fractional compartment volume [not incl. liver]	VBodTCOHC	0.87 (0.86, 0.87)	0.87 (0.86, 0.87)	1.01 (1, 1.01)	1.01 (1, 1.01)
TCE blood:air partition coefficient	PB	22 (14, 33)	19 (16, 24)	1.26 (1.19, 1.35)	1.3 (1.22, 1.38)
TCE fat:blood partition coefficient	PFat	27 (16, 46)	31 (24, 42)	1.32 (1.22, 1.44)	1.32 (1.23, 1.43)
TCE gut:blood partition coefficient	PGut	1.3 (0.69, 3)	1.1 (0.79, 1.7)	1.36 (1.21, 1.79)	1.36 (1.2, 1.68)
TCE liver:blood partition coefficient	PLiv	1.5 (1.2, 1.9)	1.6 (1.3, 1.8)	1.15 (1.11, 1.2)	1.15 (1.11, 1.2)
TCE rapidly perfused:blood partition coefficient	PRap	1.3 (0.66, 2.7)	1.3 (0.82, 2.1)	1.35 (1.18, 1.82)	1.37 (1.2, 1.76)
TCE respiratory tissue:air partition coefficient	PResp	0.97 (0.48, 2.1)	1 (0.62, 1.6)	1.37 (1.19, 1.77)	1.36 (1.19, 1.78)
TCE kidney:blood partition coefficient	PKid	1.3 (0.77, 2.2)	1.2 (0.9, 1.7)	1.31 (1.19, 1.5)	1.3 (1.2, 1.45)
TCE slowly perfused:blood partition coefficient	PSlw	0.57 (0.35, 0.97)	0.73 (0.54, 0.97)	1.32 (1.23, 1.43)	1.33 (1.25, 1.46)
TCA blood:plasma concentration ratio	TCAPlas	0.78 (0.6, 0.96)	0.78 (0.71, 0.86)	1.12 (1.06, 1.22)	1.11 (1.07, 1.17)

**Table 3-38 Prior and posterior uncertainty and variability in rat PBPK model parameters (contuined)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Free TCA body:blood plasma partition coefficient	PBodTCA	0.7 (0.18, 2.2)	0.76 (0.46, 1.3)	1.72 (1.39, 2.81)	1.65 (1.4, 2.19)
Free TCA liver:blood plasma partition coefficient	PLivTCA	0.84 (0.25, 3.3)	1.1 (0.61, 2.1)	1.71 (1.39, 2.78)	1.66 (1.38, 2.37)
Protein:TCA dissociation constant ( $\mu\text{mole/L}$ )	kDissoc	270 (95, 790)	280 (140, 530)	1.62 (1.31, 2.43)	1.6 (1.31, 2.31)
Maximum binding concentration ( $\mu\text{mole/L}$ )	B <sub>MAX</sub>	320 (80, 1300)	320 (130, 750)	1.89 (1.5, 2.64)	1.84 (1.49, 2.57)
TCOH body:blood partition coefficient	PBodTCOH	1 (0.33, 4)	1.1 (0.51, 2.1)	1.71 (1.37, 2.69)	1.76 (1.38, 2.45)
TCOH liver:body partition coefficient	PLivTCOH	1.3 (0.39, 4.5)	1.2 (0.59, 2.8)	1.71 (1.37, 2.8)	1.78 (1.37, 2.75)
TCOG body:blood partition coefficient	PBodTCOG	0.48 (0.021, 14)	1.6 (0.091, 16)	1.39 (1.2, 1.97)	1.42 (1.21, 2.52)
TCOG liver:body partition coefficient	PLivTCOG	1.3 (0.078, 39)	10 (2.7, 41)	1.4 (1.2, 2.14)	1.42 (1.21, 2.3)
DCVG effective volume of distribution	VDCVG	0.27 (0.27, 0.27)	0.27 (0.27, 0.27)	1 (1, 1)	1 (1, 1)
TCE stomach absorption coefficient (/hour)	kAS	0.73 (0.0044, 400)	2.5 (0.32, 19)	4.16 (2.21, 20)	9.3 (4.07, 31.1)
TCE stomach-duodenum transfer coefficient (/hour)	kTSD	1.4 (0.04, 45)	3.2 (0.31, 19)	3.92 (2.13, 10.4)	5.54 (2.77, 10.7)
TCE duodenum absorption coefficient (/hour)	kAD	0.96 (0.0023, 260)	0.17 (0.038, 1)	4.17 (2.15, 20.8)	4.07 (2.51, 11.9)
TCA stomach absorption coefficient (/hour)	kASTCA	0.83 (0.0024, 240)	1.4 (0.13, 13)	4.15 (2.2, 18.7)	4.21 (2.4, 11.4)
V <sub>MAX</sub> for hepatic TCE oxidation (mg/hour)	V <sub>MAX</sub>	5.8 (2, 19)	5.3 (3.9, 7.7)	1.97 (1.54, 2.92)	1.69 (1.47, 2.15)
K <sub>M</sub> for hepatic TCE oxidation (mg/L)	K <sub>M</sub>	18 (1.9, 240)	0.74 (0.54, 1.4)	2.76 (1.89, 6.46)	1.84 (1.51, 2.7)
Fraction of hepatic TCE	FracOther	0.027 (0.0018, 0.59)	0.29 (0.047, 0.56)	1.42 (1.15, 2.33)	2.15 (1.32, 5.06)

**Table 3-38 Prior and posterior uncertainty and variability in rat PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
oxidation not to TCA+TCOH					
Fraction of hepatic TCE oxidation to TCA	FracTCA	0.2 (0.027, 0.76)	0.046 (0.023, 0.087)	1.35 (1.11, 2.14)	1.84 (1.36, 2.8)
V <sub>MAX</sub> for hepatic TCE GSH conjugation (mg/hour)	V <sub>MAX</sub> DCVG	2 (0.015, 1100)	5.8 (0.16, 340)	1.52 (1.3, 2.67)	1.57 (1.32, 2.93)
K <sub>M</sub> for hepatic TCE GSH conjugation (mg/L)	K <sub>M</sub> DCVG	1500 (1.2, 1800000)	6300 (120, 720000)	1.83 (1.45, 3.15)	1.88 (1.48, 3.49)
V <sub>MAX</sub> for renal TCE GSH conjugation (mg/hour)	V <sub>MAX</sub> KidDCVG	0.038 (0.00027, 13)	0.0024 (5e-04, 0.014)	1.52 (1.3, 2.81)	1.56 (1.29, 2.72)
K <sub>M</sub> for renal TCE GSH conjugation (mg/L)	K <sub>M</sub> KidDCVG	470 (0.47, 530000)	0.25 (0.038, 2.2)	1.84 (1.47, 4.27)	1.93 (1.49, 3.57)
V <sub>MAX</sub> for tracheo-bronchial TCE oxidation (mg/hour)	V <sub>MAX</sub> Clara	0.2 (0.0077, 2.4)	0.17 (0.042, 0.69)	2.26 (1.71, 3.3)	4.35 (1.99, 6.7)
K <sub>M</sub> for tracheo-bronchial TCE oxidation (mg/L)	K <sub>M</sub> Clara	0.016 (0.0014, 0.58)	0.025 (0.005, 0.15)	1.47 (1.26, 2.39)	1.65 (1.28, 10.5)
Fraction of respiratory metabolism to systemic circ.	FracLungSys	0.82 (0.027, 1)	0.73 (0.06, 0.98)	1.09 (1, 1.71)	1.13 (1.01, 1.86)
V <sub>MAX</sub> for hepatic TCOH→TCA (mg/hour)	V <sub>MAX</sub> TCOH	0.75 (0.037, 20)	0.71 (0.27, 2.2)	1.51 (1.25, 2.64)	1.68 (1.3, 3.23)
K <sub>M</sub> for hepatic TCOH→TCA (mg/L)	K <sub>M</sub> TCOH	1 (0.029, 23)	19 (3.6, 94)	1.52 (1.26, 2.7)	1.72 (1.26, 3.93)
V <sub>MAX</sub> for hepatic TCOH→TCOG (mg/hour)	V <sub>MAX</sub> Gluc	27 (0.83, 620)	11 (4.1, 32)	1.5 (1.25, 2.59)	2.3 (1.41, 5.19)
K <sub>M</sub> for hepatic TCOH→TCOG (mg/L)	K <sub>M</sub> Gluc	31 (1, 570)	6.3 (1.2, 20)	1.5 (1.25, 2.74)	2.04 (1.3, 8.4)
Rate constant for hepatic TCOH→other (/hour)	kMetTCOH	4.2 (0.17, 150)	3 (0.57, 15)	1.49 (1.27, 2.67)	1.72 (1.3, 8.31)
Rate constant for TCA plasma→urine (/hour)	kUrnTCA	1.9 (0.21, 47)	0.92 (0.51, 1.7)	1.56 (1.33, 2.81)	1.58 (1.36, 2.25)

**Table 3-38 Prior and posterior uncertainty and variability in rat PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Rate constant for hepatic TCA→other (/hour)	kMetTCA	0.76 (0.037, 19)	0.47 (0.17, 1.2)	1.5 (1.26, 2.74)	1.52 (1.27, 2.45)
Rate constant for TCOG liver→bile (/hour)	kBile	1.4 (0.052, 31)	14 (2.7, 39)	1.5 (1.25, 2.8)	1.63 (1.29, 4.1)
Lumped rate constant for TCOG bile→TCOH liver (/hour)	kEHR	0.013 (0.00055, 0.64)	1.7 (0.34, 7.4)	1.5 (1.25, 2.49)	1.67 (1.26, 5.91)
Rate constant for TCOG→urine (/hour)	kUrnTCOG	11 (0.063, 1000)	12 (0.45, 370)	1.74 (1.42, 2.99)	1.86 (1.43, 3.54)
Rate constant for hepatic DCVG→DCVC (/hour)	kDCVG	30000 (30000, 30000)	30000 (30000, 30000)	1 (1, 1)	1 (1, 1)
Lumped rate constant for DCVC→urinary NAcDCVC (/hour)	kNAT	0.15 (0.00024, 84)	0.0029 (0.00066, 0.015)	1.49 (1.24, 2.8)	1.54 (1.26, 2.45)
Rate constant for DCVC bioactivation (/hour)	kKidBioact	0.12 (0.00023, 83)	0.0092 (0.0012, 0.043)	1.48 (1.24, 2.68)	1.52 (1.25, 2.5)

**Table 3-39. Prior and posterior uncertainty and variability in human PBPK model parameters**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Cardiac output (L/hour)	QC	390 (280, 560)	330 (280, 390)	1.17 (1.1, 1.39)	1.39 (1.26, 1.54)
Alveolar ventilation (L/hour)	QP	380 (220, 640)	440 (360, 530)	1.27 (1.17, 1.52)	1.58 (1.44, 1.73)
Scaled fat blood flow	QFatC	0.051 (0.021, 0.078)	0.043 (0.033, 0.055)	1.64 (1.23, 2)	1.92 (1.72, 2.09)
Scaled gut blood flow	QGutC	0.19 (0.15, 0.23)	0.16 (0.14, 0.18)	1.16 (1.1, 1.21)	1.16 (1.12, 1.2)
Scaled liver blood flow	QLivC	0.063 (0.029, 0.099)	0.039 (0.026, 0.055)	1.62 (1.22, 1.92)	1.8 (1.62, 1.98)
Scaled slowly perfused blood flow	QSlwC	0.22 (0.13, 0.3)	0.17 (0.14, 0.21)	1.34 (1.18, 1.45)	1.39 (1.31, 1.46)
Scaled rapidly perfused blood flow	QRapC	0.29 (0.18, 0.4)	0.39 (0.34, 0.43)	1.31 (1.14, 1.57)	1.22 (1.16, 1.3)
Scaled kidney blood flow	QKidC	0.19 (0.16, 0.22)	0.19 (0.18, 0.21)	1.1 (1.07, 1.13)	1.1 (1.07, 1.12)
Respiratory lumen:tissue diffusive clearance rate (L/hour)	DResp	560 (44, 3300)	270 (130, 470)	1.37 (1.25, 1.61)	1.71 (1.52, 2.35)
Fat fractional compartment volume	VFatC	0.19 (0.088, 0.31)	0.16 (0.12, 0.21)	1.66 (1.23, 1.93)	1.65 (1.4, 1.9)
Gut fractional compartment volume	VGutC	0.02 (0.018, 0.022)	0.02 (0.019, 0.021)	1.07 (1.04, 1.08)	1.06 (1.05, 1.08)
Liver fractional compartment volume	VLivC	0.026 (0.018, 0.032)	0.026 (0.022, 0.03)	1.21 (1.12, 1.28)	1.2 (1.13, 1.26)
Rapidly perfused fractional compartment volume	VRapC	0.087 (0.079, 0.096)	0.088 (0.083, 0.093)	1.07 (1.05, 1.09)	1.06 (1.05, 1.08)
Fractional volume of respiratory lumen	VRespLumC	0.0024 (0.0018, 0.003)	0.0024 (0.0021, 0.0027)	1.18 (1.1, 1.23)	1.17 (1.12, 1.22)
Fractional volume of respiratory tissue	VRespEffC	0.00018 (0.00014, 0.00022)	0.00018 (0.00015, 0.00021)	1.18 (1.1, 1.24)	1.17 (1.13, 1.23)
Kidney fractional compartment volume	VKidC	0.0043 (0.0034, 0.0052)	0.0043 (0.0038, 0.0048)	1.15 (1.09, 1.19)	1.14 (1.1, 1.19)



**Table 3-39 Prior and posterior uncertainty and variability in human PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Blood fractional compartment volume	VBldC	0.077 (0.066, 0.088)	0.078 (0.072, 0.084)	1.1 (1.06, 1.13)	1.1 (1.07, 1.13)
Slowly perfused fractional compartment volume	VSlwC	0.45 (0.33, 0.55)	0.48 (0.43, 0.52)	1.18 (1.1, 1.24)	1.16 (1.12, 1.22)
Plasma fractional compartment volume	VPlasC	0.044 (0.037, 0.051)	0.044 (0.04, 0.048)	1.11 (1.08, 1.14)	1.11 (1.08, 1.14)
TCA body fractional compartment volume [not incl. blood+liver]	VBodC	0.75 (0.74, 0.77)	0.75 (0.74, 0.76)	1.01 (1.01, 1.01)	1.01 (1.01, 1.01)
TCOH/G body fractional compartment volume [not incl. liver]	VBodTCOHC	0.83 (0.82, 0.84)	0.83 (0.83, 0.83)	1.01 (1, 1.01)	1.01 (1, 1.01)
TCE blood:air partition coefficient	PB	9.6 (6.5, 13)	9.2 (8.2, 10)	1.18 (1.13, 1.26)	1.21 (1.16, 1.28)
TCE fat:blood partition coefficient	PFat	68 (46, 98)	57 (49, 66)	1.18 (1.11, 1.33)	1.18 (1.11, 1.3)
TCE gut:blood partition coefficient	PGut	2.6 (1.3, 5.3)	2.9 (1.9, 4.1)	1.37 (1.2, 1.78)	1.41 (1.21, 1.77)
TCE liver:blood partition coefficient	PLiv	4 (1.9, 8.5)	4.1 (2.7, 5.9)	1.37 (1.22, 1.81)	1.33 (1.19, 1.6)
TCE rapidly perfused:blood partition coefficient	PRap	2.6 (1.2, 5.7)	2.4 (1.8, 3.2)	1.37 (1.21, 1.78)	1.5 (1.25, 1.87)
TCE respiratory tissue:air partition coefficient	PResp	1.3 (0.65, 2.7)	1.3 (0.9, 1.9)	1.36 (1.19, 1.81)	1.32 (1.2, 1.56)
TCE kidney:blood partition coefficient	PKid	1.6 (1.1, 2.3)	1.6 (1.3, 1.9)	1.17 (1.1, 1.33)	1.15 (1.09, 1.25)
TCE slowly perfused:blood partition coefficient	PSlw	2.1 (1.2, 3.5)	2.3 (1.9, 2.8)	1.28 (1.14, 1.53)	1.51 (1.36, 1.66)
TCA blood:plasma concentration ratio	TCAPlas	0.78 (0.55, 15)	0.65 (0.6, 0.77)	1.08 (1.03, 1.53)	1.52 (1.23, 2.03)

**Table 3-39 Prior and posterior uncertainty and variability in human PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
Free TCA body:blood plasma partition coefficient	PBodTCA	0.45 (0.19, 8.1)	0.44 (0.33, 0.55)	1.36 (1.19, 1.75)	1.67 (1.38, 2.2)
Free TCA liver:blood plasma partition coefficient	PLivTCA	0.59 (0.24, 10)	0.55 (0.39, 0.77)	1.36 (1.18, 1.76)	1.65 (1.37, 2.16)
Protein:TCA dissociation constant (μmole/L)	kDissoc	180 (160, 200)	180 (170, 190)	1.05 (1.03, 1.09)	1.04 (1.03, 1.07)
Maximum binding concentration (μmole/L)	B <sub>MAX</sub>	830 (600, 1100)	740 (630, 880)	1.17 (1.1, 1.3)	1.16 (1.1, 1.28)
TCOH body:blood partition coefficient	PBodTCOH	0.89 (0.51, 1.7)	1.5 (1.3, 1.7)	1.29 (1.16, 1.64)	1.34 (1.25, 1.47)
TCOH liver:body partition coefficient	PLivTCOH	0.58 (0.32, 1.1)	0.63 (0.45, 0.87)	1.29 (1.16, 1.65)	1.29 (1.17, 1.5)
TCOG body:blood partition coefficient	PBodTCOG	0.67 (0.036, 16)	0.72 (0.3, 1.8)	1.38 (1.2, 2.42)	7.83 (4.86, 12.6)
TCOG liver:body partition coefficient	PLivTCOG	1.8 (0.11, 28)	3.1 (0.87, 8.1)	1.38 (1.19, 2.04)	4.94 (2.73, 8.58)
DCVG effective volume of distribution	VDCVG	73 (5.2, 36000)	6.1 (5.4, 7.3)	1.27 (1.08, 1.95)	1.1 (1.07, 1.16)
TCE stomach absorption coefficient (/hour)	kAS	1.4 (1.4, 1.4)	1.4 (1.4, 1.4)	1 (1, 1)	1 (1, 1)
TCE stomach-duodenum transfer coefficient (/hour)	kTSD	1.4 (1.4, 1.4)	1.4 (1.4, 1.4)	1 (1, 1)	1 (1, 1)
TCE duodenum absorption coefficient (/hour)	kAD	0.75 (0.75, 0.75)	0.75 (0.75, 0.75)	1 (1, 1)	1 (1, 1)
TCA stomach absorption coefficient (/hour)	kASTCA	0.58 (0.0022, 210)	3 (0.061, 180)	4.26 (2.13, 17.6)	5.16 (2.57, 22.3)
TCOH stomach absorption coefficient (/hour)	kASTCOH	0.49 (0.0024, 210)	7.6 (0.11, 150)	4.19 (2.22, 21.5)	5.02 (2.44, 18.5)
V <sub>MAX</sub> for hepatic TCE	V <sub>MAX</sub>	430 (130, 1500)	190 (130, 290)	1.98 (1.69, 2.31)	2.02 (1.77, 2.38)

**Table 3-39 Prior and posterior uncertainty and variability in human PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
oxidation (mg/hour)					
K <sub>M</sub> for hepatic TCE oxidation (mg/L)	K <sub>M</sub>	3.7 (0.22, 63)	0.18 (0.078, 0.4)	2.74 (2.1, 5.62)	4.02 (2.9, 5.64)
Fraction of hepatic TCE oxidation not to TCA+TCOH	FracOther	0.12 (0.0066, 0.7)	0.11 (0.024, 0.23)	1.4 (1.11, 2.38)	2.71 (1.37, 5.33)
Fraction of hepatic TCE oxidation to TCA	FracTCA	0.19 (0.036, 0.56)	0.035 (0.024, 0.05)	2.55 (1.51, 3.96)	2.25 (1.89, 2.87)
V <sub>MAX</sub> for hepatic TCE GSH conjugation (mg/hour)	V <sub>MAX</sub> DCVG	100 (0.0057, 690000)	340 (110, 1100)	1.91 (1.55, 3.76)	6.18 (3.35, 11.3)
K <sub>M</sub> for hepatic TCE GSH conjugation (mg/L)	K <sub>M</sub> DCVG	3.1 (0.21, 42)	3.6 (1.2, 11)	1.52 (1.26, 2.91)	4.2 (2.48, 8.01)
V <sub>MAX</sub> for renal TCE GSH conjugation (mg/hour)	V <sub>MAX</sub> KidDCVG	220 (0.028, 6700000)	2.1 (0.17, 9.3)	1.86 (1.51, 3.33)	4.02 (1.57, 33.9)
K <sub>M</sub> for renal TCE GSH conjugation (mg/L)	K <sub>M</sub> KidDCVG	2.7 (0.14, 41)	0.76 (0.29, 5.8)	1.5 (1.27, 2.56)	1.49 (1.27, 2.32)
V <sub>MAX</sub> for tracheo-bronchial TCE oxidation (mg/hour)	V <sub>MAX</sub> Clara	25 (1, 260)	18 (3.8, 41)	2.25 (1.85, 3.25)	2.9 (2.12, 6.49)
K <sub>M</sub> for tracheo-bronchial TCE oxidation (mg/L)	K <sub>M</sub> Clara	0.019 (0.0017, 0.5)	0.31 (0.057, 1.4)	1.48 (1.25, 2.39)	10.8 (1.99, 37.6)
Fraction of respiratory metabolism to systemic circ.	FracLungSys	0.75 (0.051, 0.99)	0.96 (0.86, 0.99)	1.12 (1, 1.75)	1.02 (1, 1.1)
V <sub>MAX</sub> for hepatic TCOH→TCA (mg/hour)	V <sub>MAX</sub> TCOH	42 (0.77, 2200)	9.2 (5.5, 20)	1.83 (1.46, 3.43)	3.15 (2.3, 5.44)
K <sub>M</sub> for hepatic TCOH→TCA (mg/L)	K <sub>M</sub> TCOH	5 (0.23, 81)	2.2 (1.3, 4.5)	1.49 (1.25, 2.57)	2.58 (1.75, 4.5)
V <sub>MAX</sub> for hepatic TCOH→TCOG (mg/hour)	V <sub>MAX</sub> Gluc	720 (12, 50000)	900 (340, 2000)	1.83 (1.48, 3.5)	2.29 (1.84, 4.57)
K <sub>M</sub> for hepatic	K <sub>M</sub> Gluc	10 (0.53, 190)	130 (47, 290)	1.5 (1.25, 2.6)	1.58 (1.26, 3.69)

**Table 3-39 Prior and posterior uncertainty and variability in human PBPK model parameters (continued)**

<b>Parameter description</b>	<b>PBPK Parameter</b>	<b>Prior Population Median: Median (2.5%, 97.5%)</b>	<b>Posterior Population Median: Median (2.5%, 97.5%)</b>	<b>Prior Population GSD: Median (2.5%, 97.5%)</b>	<b>Posterior Population GSD: Median (2.5%, 97.5%)</b>
TCOH→TCOG (mg/L)					
Rate constant for hepatic TCOH→other (/hour)	kMetTCOH	0.83 (0.035, 10)	0.25 (0.042, 0.7)	1.5 (1.26, 3)	5.13 (2.72, 16.7)
Rate constant for TCA plasma→urine (/hour)	kUrnTCA	0.26 (0.038, 4)	0.11 (0.083, 0.15)	1.48 (1.29, 2.29)	1.86 (1.58, 2.28)
Rate constant for hepatic TCA→other (/hour)	kMetTCA	0.19 (0.01, 2.6)	0.096 (0.038, 0.19)	1.48 (1.26, 2.57)	2.52 (1.79, 4.34)
Rate constant for TCOG liver→bile (/hour)	kBile	1.2 (0.059, 16)	2.5 (1.1, 6.9)	1.47 (1.25, 2.75)	1.56 (1.27, 3.21)
Lumped rate constant for TCOG bile→TCOH liver (/hour)	kEHR	0.074 (0.004, 1.4)	0.053 (0.033, 0.087)	1.52 (1.26, 2.64)	1.72 (1.35, 2.51)
Rate constant for TCOG→urine (/hour)	kUrnTCOG	2.9 (0.061, 260)	2.4 (0.83, 7)	1.75 (1.4, 3.31)	18.7 (11.6, 31.8)
Rate constant for hepatic DCVG→DCVC (/hour)	kDCVG	0.044 (6.3e-05, 22)	2.5 (1.9, 3.4)	1.48 (1.25, 2.83)	1.51 (1.3, 1.86)
Lumped rate constant for DCVC→urinary NAcDCVC (/hour)	kNAT	0.00085 (5.5e-05, 0.041)	1e-04 (4.7e-05, 7e-04)	1.51 (1.25, 2.34)	1.47 (1.24, 2.48)
Rate constant for DCVC bioactivation (/hour)	kKidBioact	0.0022 (9.5e-05, 0.079)	0.023 (0.0062, 0.061)	1.51 (1.25, 2.57)	1.52 (1.25, 2.69)

**Table 3-40. Confidence interval (CI) widths (ratio of 97.5% to 2.5% estimates) and fold-shift in median estimate for the PBPK model population median parameters, sorted in order of decreasing CI width. Shifts in the median estimate greater than threefold are in bold to denote larger shifts between the prior and posterior distributions**

Mouse				Rat				Human			
PBPK Parameter	Width of CI on population median		Fold-shift in pop. median	PBPK Parameter	Width of CI on population median		Fold-shift in pop. median	PBPK Parameter	Width of CI on population median		Fold-shift in pop. median
	Prior	Posterior			Prior	Posterior			Prior	Posterior	
K <sub>M</sub> DCVG	2230000000	13400000	× <b>8.8</b>	K <sub>M</sub> DCVG	1500000	5800	× <b>4.29</b>	kASTCA	94300	3040	× <b>5.18</b>
K <sub>M</sub> KidDCVG	1170000000	3540000	×1.05	V <sub>MAX</sub> DCVG	71100	2130	×2.86	kASTCOH	85900	1420	× <b>15.6</b>
V <sub>MAX</sub> DCVG	400000	46200	÷ <b>6.18</b>	kUrnTCOG	16700	822	×1.04	V <sub>MAX</sub> KidDCVG	236000000	55.1	÷ <b>105</b>
V <sub>MAX</sub> KidDCVG	357000	11000	÷ <b>12.8</b>	PBodTCOG	666	172	× <b>3.43</b>	K <sub>M</sub> Clara	289	23.9	× <b>16.2</b>
kASTCA	89300	374	× <b>6.3</b>	kASTCA	98200	95.7	×1.69	K <sub>M</sub> KidDCVG	287	20	÷ <b>3.48</b>
kTSD	1190	51.1	× <b>3.26</b>	kTSD	1130	61.8	×2.29	kMetTCOH	289	16.6	÷ <b>3.28</b>
kEHR	412000	42.1	÷ <b>5.43</b>	kAS	91000	60.2	× <b>3.41</b>	kNAT	756	15.1	÷ <b>8.14</b>
FracOther	567	39.5	÷ <b>18.5</b>	K <sub>M</sub> KidDCVG	1130000	58.6	÷ <b>1880</b>	V <sub>MAX</sub> Clara	255	10.6	÷1.41
K <sub>M</sub> Clara	351000	37.5	÷ <b>134</b>	kKidBioact	366000	35.6	÷ <b>13.3</b>	kKidBioact	833	9.91	× <b>10.5</b>
kAS	91900	35.9	×1	K <sub>M</sub> Clara	406	29.9	×1.53	V <sub>MAX</sub> DCVG	122000000	9.78	× <b>3.29</b>
kUrnTCOG	40500000	29.9	× <b>11.8</b>	V <sub>MAX</sub> KidDCVG	48500	27.5	÷ <b>15.6</b>	FracOther	106	9.75	÷1.09
B <sub>MAX</sub>	81.8	24.4	×1.66	kMetTCOH	891	26.4	÷1.41	PLivTCOG	253	9.32	×1.77
K <sub>M</sub> Gluc	344000	24.3	× <b>16.3</b>	kAD	115000	26.3	÷ <b>5.53</b>	K <sub>M</sub> DCVG	198	9.13	×1.18
kAD	84900	23.8	÷ <b>4.53</b>	K <sub>M</sub> TCOH	781	26	× <b>18.7</b>	kUrnTCOG	4290	8.5	÷1.19
kDissoc	60.3	21.8	×1.33	kNAT	351000	22.7	÷ <b>50.2</b>	kBile	274	6.54	×2.01
V <sub>MAX</sub> Clara	131	15	×1.75	kEHR	1160	21.9	× <b>134</b>	K <sub>M</sub> Gluc	365	6.07	× <b>13.4</b>
kMetTCOH	35500000	12.1	× <b>47.4</b>	K <sub>M</sub> Gluc	562	17.1	÷ <b>4.98</b>	PBodTCOG	454	5.85	×1.08
kBile	390000	11.3	× <b>8.23</b>	V <sub>MAX</sub> Clara	305	16.5	÷1.21	V <sub>MAX</sub> Gluc	4330	5.71	×1.25
K <sub>M</sub> TCOH	29600000	10.5	÷1.47	FracLungSys	36.7	16.3	÷1.12	K <sub>M</sub>	288	5.1	÷ <b>20.5</b>
V <sub>MAX</sub> Gluc	23600000	8.28	× <b>41.1</b>	PLivTCOG	501	14.8	× <b>8.07</b>	kMetTCA	248	4.89	÷1.94

**Table 3-40. Confidence interval (CI) widths (ratio of 97.5% to 2.5% estimates) and fold-shift in median estimate for the PBPK model population median parameters, sorted in order of decreasing CI width (continued)**

Mouse				Rat				Human			
PBPK Parameter	Width of CI on population median		Fold-shift in pop. median	PBPK Parameter	Width of CI on population median		Fold-shift in pop. median	PBPK Parameter	Width of CI on population median		Fold-shift in pop. median
	Prior	Posterior			Prior	Posterior			Prior	Posterior	
PBodTCOG	4770	6.27	÷1.95	kBile	588	14.8	× <b>9.67</b>	DResp	74.3	3.71	÷2.06
V <sub>MAX</sub> TCOH	27100000	5.78	×1.8	FracOther	331	11.9	× <b>10.7</b>	V <sub>MAX</sub> TCOH	2900	3.62	÷ <b>4.56</b>
K <sub>M</sub>	386	5.76	÷ <b>12.5</b>	V <sub>MAX</sub> TCOH	550	8.25	÷1.06	K <sub>M</sub> TCOH	359	3.48	÷2.33
kUrnTCA	4540	5.76	÷ <b>10.2</b>	V <sub>MAX</sub> Gluc	740	7.79	÷2.4	kEHR	339	2.62	÷1.39
FracLungSys	608	5.55	×2.27	kMetTCA	507	6.93	÷1.61	V <sub>MAX</sub>	11.5	2.27	÷2.33
kMetTCA	316000	4.59	× <b>12</b>	B <sub>MAX</sub>	16.2	5.79	×1	PResp	4.1	2.16	÷1.01
PLivTCOG	4860	3.99	×1.04	DResp	180	4.81	×2.12	PLiv	4.44	2.14	×1.02
DResp	475000	3.64	× <b>147</b>	PLivTCOH	11.5	4.7	÷1.09	QLivC	3.46	2.11	÷1.62
PLivTCA	58.3	2.88	×1	PBodTCOH	12.1	4.03	×1.03	PGut	4.21	2.1	×1.11
PResp	4	2.85	÷1.07	kDissoc	8.38	3.85	×1.04	FracTCA	15.5	2.06	÷ <b>5.37</b>
PRap	3.78	2.79	÷1.03	FracTCA	28.1	3.85	÷ <b>4.27</b>	PLivTCA	42.6	1.98	÷1.07
PGut	4.33	2.77	÷1.25	PLivTCA	13.3	3.49	×1.37	PLivTCOH	3.52	1.93	×1.08
V <sub>MAX</sub>	10.7	2.67	÷1.58	kUrnTCA	219	3.28	÷2	kDCVG	344000	1.8	× <b>55.7</b>
PBodTCA	62.6	2.55	×1.14	PBodTCA	12	2.8	×1.09	kUrnTCA	105	1.79	÷2.32
PSlw	4.04	2.54	÷1.06	PResp	4.32	2.6	×1.04	VFatC	3.49	1.76	÷1.21
PLiv	3.87	2.5	×1.26	K <sub>M</sub>	123	2.56	÷ <b>24</b>	PRap	4.66	1.74	÷1.09
FracTCA	3060	2.49	×1.49	PRap	4.01	2.53	÷1.01	QFatC	3.7	1.7	÷1.19
TCAPlas	40.6	2.38	×1.46	PGut	4.35	2.16	÷1.17	PBodTCA	42.9	1.7	÷1.04
PKid	4.78	2.37	×1.2	V <sub>MAX</sub>	9.5	1.98	÷1.11	PSlw	2.9	1.5	×1.11
QFatC	3.62	2.26	×1.02	QRapC	2.77	1.97	÷1	PKid	2.05	1.49	÷1.01
PLivTCOH	3.19	2.13	×1.48	VFatC	3.58	1.96	÷1	QP	2.97	1.48	×1.16
PBodTCOH	3.41	2.01	÷1.27	PKid	2.89	1.85	÷1.11	QSlwC	2.25	1.48	÷1.26
QKidC	2.39	1.91	÷1.01	QP	3.59	1.79	÷1.38	QC	2.04	1.39	÷1.19
PFat	3.01	1.89	÷1.01	PSlw	2.76	1.79	×1.28	B <sub>MAX</sub>	1.92	1.38	÷1.12
QSlwC	2.04	1.88	÷1.02	PFat	2.91	1.77	×1.16	VLivC	1.79	1.36	×1.01
VPlasC	2.18	1.87	÷1.17	QSlwC	2.19	1.69	÷1.06	PFat	2.13	1.34	÷1.2

**Table 3-40. Confidence interval (CI) widths (ratio of 97.5% to 2.5% estimates) and fold-shift in median estimate for the PBPK model population median parameters, sorted in order of decreasing CI width (continued)**

Mouse				Rat				Human			
PBPK Parameter	Width of CI on population median		Fold-shift in pop. median	PBPK Parameter	Width of CI on population median		Fold-shift in pop. median	PBPK Parameter	Width of CI on population median		Fold-shift in pop. median
	Prior	Posterior			Prior	Posterior			Prior	Posterior	
VFatC	3.49	1.83	×1.25	QFatC	3.47	1.66	×1.14	VDCVG	6820	1.34	÷12
QP	2.75	1.82	÷1.02	VPlasC	2.17	1.55	×1.03	VRespEffC	1.66	1.33	÷1.02
VLivC	1.85	1.6	÷1.16	PB	2.37	1.51	÷1.15	PBodTCOH	3.32	1.32	×1.68
QC	2.1	1.59	×1.2	QC	1.64	1.43	×1.15	VRespLumC	1.65	1.31	÷1
PB	2.3	1.54	÷1.07	VRespEffC	1.56	1.43	÷1	TCAPlas	26.9	1.29	÷1.21
QLivC	1.55	1.42	×1.02	VRespLumC	1.56	1.41	×1	VKidC	1.54	1.28	÷1.01
QRapC	1.51	1.41	÷1.03	VLivC	1.57	1.4	÷1.05	PB	2.04	1.28	÷1.04
VGutC	1.38	1.3	÷1.01	PLiv	1.67	1.37	×1.05	QRapC	2.22	1.25	×1.34
VBldC	1.34	1.27	÷1.02	QLivC	1.53	1.34	×1.04	QGutC	1.59	1.23	÷1.19
VRespLumC	1.32	1.26	÷1.01	VKidC	1.47	1.33	×1.01	VSlwC	1.66	1.21	×1.07
VRespEffC	1.31	1.26	÷1	QKidC	1.39	1.28	×1	VPlasC	1.39	1.2	×1.01
QGutC	1.52	1.24	×1.15	VGutC	1.38	1.28	÷1.01	QKidC	1.36	1.17	÷1
VKidC	1.29	1.24	÷1	VBldC	1.34	1.25	×1.01	VBldC	1.34	1.17	×1.02
VRapC	1.3	1.23	÷1.01	VRapC	1.34	1.23	×1	FracLungSys	19.4	1.14	×1.29
VSlwC	1.19	1.11	÷1.01	QGutC	1.53	1.22	×1.14	VRapC	1.22	1.12	×1
VBodC	1.05	1.03	×1.01	TCAPlas	1.6	1.21	÷1.01	kDissoc	1.23	1.12	÷1.01
VBodTCOHC	1.04	1.03	×1.01	VSlwC	1.15	1.09	×1	VGutC	1.22	1.11	×1.01
				VBodC	1.04	1.03	×1	VBodC	1.04	1.02	÷1
				VBodTCOHC	1.02	1.01	×1	VBodTCOHC	1.02	1.01	÷1

1 was the fraction of TCE oxidation directly producing TCA, which shifted by fourfold in rats and  
2 fivefold in mice, with prior CIs of 28-fold and 16-fold, respectively. These shifts are still  
3 relatively modest in comparison to the prior CI, and moreover, the posterior CI is quite narrow  
4 (fourfold in rats, twofold in humans), suggesting that the parameter is well identified by the in  
5 vivo data.

6 In addition, in only a few cases did the interquartile regions of the prior and posterior  
7 distributions not overlap. In most of these cases, including the diffusion rate from respiratory  
8 lumen to tissue, the  $K_{MS}$  for renal TCE GSH conjugation and respiratory TCE oxidation, and  
9 several metabolite kinetic parameters, the prior distributions themselves were noninformative.  
10 For a noninformative prior, the lack of overlap would only be an issue if the posterior  
11 distributions were affected by the truncation limit, which was not the case here. The only other  
12 parameter for which there was a lack of interquartile overlap between the prior and posterior  
13 distribution was the  $K_M$  for hepatic TCE oxidation in mice and in rats, though the prior and  
14 posterior 95% CIs did overlap within each species. As discussed Section 3.3, there is some  
15 uncertainty in the extrapolation of in vitro  $K_M$  values to in vivo values (within the same species).  
16 In addition, in mice, it has been known for some time that  $K_M$  values appear to be discordant  
17 among different studies (Abbas and Fisher, 1997; Fisher et al., 1991; Greenberg et al., 1999).

18 In terms of estimates of population variability, for the vast majority of parameters, the  
19 posterior estimate of the population GSD was either twofold or less, indicating modest  
20 variability. In some cases, while the posterior population GSD was greater than twofold, it was  
21 similar to the prior estimate of the population GSD, indicating limited additional informative  
22 data on variability. This was the case for oral absorption parameters, which are expected to be  
23 highly variable because the current model lumps parameters for different oral dosing vehicles  
24 together, and a relatively wide prior distribution was given. In addition, in some cases this was  
25 due to in vitro data showing a higher degree of variability. Examples of this include TCA  
26 plasma binding parameters in the mouse, and the  $V_{MAX}$  for hepatic oxidation and the fraction of  
27 oxidation to TCA in humans. In a few other cases, the in vivo data appeared to indicate greater  
28 than twofold variability between subjects, and these are discussed in more detail below.

29 In the mouse, the two parameters for which this is the case are the  $V_{MAX}$  for respiratory  
30 tract oxidation and the urinary excretion rate for TCOG. In the first case, the variability is driven  
31 by the need for a higher respiratory tract  $V_{MAX}$  for males in the Fisher et al. (1991) study as  
32 compared to other studies. In the second case, it is driven by the relatively low estimate of  
33 urinary excretion of TCOG in the Abbas and Fisher (1997), Abbas et al. (1997), and Greenberg  
34 et al. (1999) studies as compared with the relatively high estimate Green and Prout (1985) and  
35 Prout et al. (1985).



1           In the rat, the two parameters for which the in vivo data suggest greater than twofold  
2 variability are the fraction of oxidation to not producing TCA or TCOH, and the  $V_{MAX}$  for  
3 respiratory tract oxidation. In the first case, this is driven by three studies that appeared to  
4 require greater (Bernauer et al., 1996; Kimmerle and Eben, 1973a) or lower (Hissink et al., 2002)  
5 estimates for this parameter as compared with the other studies. Nonetheless, the degree of  
6 variability is not much greater than twofold, with a central estimate population GSD of 2.15-fold.  
7 In the case of the  $V_{MAX}$  for respiratory tract oxidation, two studies appeared to require higher  
8 (Fisher et al., 1989) or lower (Simmons et al., 2002) values for this parameter as compared with  
9 the other studies.

10           In humans, as would be expected, more parameters appeared to exhibit greater than  
11 twofold variability. In terms of distribution, the partition coefficients for TCOG had rather large  
12 posterior estimates for the population GSD of eightfold for the body and fivefold for the liver. In  
13 terms of the body, a few of the subjects in Fisher et al. (1998) and all the subjects in Monster  
14 et al. (1976) appeared to require much higher partition coefficients for TCOG. For the liver, the  
15 variability did not have a discernable trend across studies. In addition, almost all the metabolism  
16 and clearance parameters had posterior estimates for population variability of greater than a  
17 twofold GSD. The largest of these was the urinary excretion rate for TCOG, with a GSD of  
18 19-fold. In this case, the variability was driven by individuals in the Chiu et al. (2007) 1 ppm  
19 study, who were predicted to have much lower rate of urinary excretion as compared to that  
20 estimated in the other, higher exposure studies.

21           In sum, the Bayesian analysis of the updated PBPK model and data exhibited no major  
22 inconsistencies in prior and posterior parameter distributions. The most significant issue in terms  
23 of population central estimates was the  $K_M$  for hepatic oxidative metabolism, for which the  
24 posterior estimates were low compared to, albeit somewhat uncertain, in vitro estimates, and it  
25 could be argued that a wider prior distribution would have been better. However, the central  
26 estimates were not at or near the truncation boundary, so it is unlikely that wider priors would  
27 change the results substantially. In terms of population variability, in rodents, the estimates of  
28 variability were generally modest, which is consistent with more homogeneous and controlled  
29 experimental subjects and conditions, whereas the estimates of human population variability  
30 were greater—particularly for metabolism and clearance. Overall, there were no indications  
31 based on this evaluation of prior and posterior distributions either that prior distributions were  
32 overly restrictive or that model specification errors led to pathological parameter estimates.

33


### 3.5.6.1.3. Comparison of Model Predictions With Data

1 Comparisons of model predictions and data for each species are discussed in the  
2 sub-sections below. First, as an overall summary, for each species and each output  
3 measurement, the data and predictions generated from a random sample of the MCMC chain are  
4 scatter-plotted to show the general degree of consistency between data and predictions. Next, as  
5 with the Hack et al. (2006) model, the sampled subject-specific parameters were used to generate  
6 predictions for comparison to the calibration data (see Figure 3-8). Thus, the predictions for a  
7 particular data set are conditioned on the posterior parameter distributions for same data set.  
8 Because these parameters were “optimized” for each experiment, these subject-specific  
9 predictions should be accurate by design—and, on the whole, were so. In addition, the  
10 “residual-error” estimate for each measurement (see Table 3-41) provides some quantitative  
11 measure of the degree to which there were deviations due to intrastudy variability and model  
12 misspecification, including any difficulties fitting multiple dose levels in the same study using  
13 the same model parameters.

14 Next, only samples of the population parameters (means and variances) were used, and  
15 new subjects were sampled from appropriate distribution using these population means and  
16 variances (see Figure 3-8). That is, the predictions were only conditioned on the  
17 population-level parameters distributions, representing an “average” over all the data sets, and  
18 not on the specific predictions for that data set. These “new” subjects then represent the  
19 predicted population distribution, incorporating variability in the population as well as  
20 uncertainty in the population means and variances. Because of the limited amount of mouse  
21 data, all available data for that species were utilized for calibration, and there were no data  
22 available for “out-of-sample” evaluation (often referred to as “validation data,” but this term is  
23 not used here due to ambiguities as to its definition). In rats, several studies that contained  
24 primarily blood TCE data, which were abundant, were used for out-of-sample evaluation. In  
25 humans, there were substantial individual and aggregated (mean of individuals in a study) data  
26 that was available for out-of-sample evaluation, as computational intensity limited the number of  
27 individuals that could be used in the MCMC-based calibration.

### 3.5.6.1.4. Mouse model and data

29 Each panel of Figure 3-9 shows a scatter plot of the calibration data and a random posterior  
30 prediction for each of the measured endpoint. The endpoint abbreviations are listed in  
31 Table 3-41, as are the implied GSDs for the “residual” errors, which include intrastudy  
32 variability, interindividual variability, and measurement and model errors. The residual-error



1 GSDs are also shown as grey dotted lines in Figure 3-9. Table 3-42 provides an evaluation of  
 2 the predictions of the mouse model for each data set, with figures showing individual  
 3 time-course data and predictions in Appendix A.

4  
 5  
 6  
 7  
 8  
 9  
 10  
 11

**Figure 3-8. Schematic of how posterior predictions were generated for comparison with experimental data.** Two sets of posterior predictions were generated: population predictions (diagonal hashing) and subject-specific predictions (vertical hashing). (Same as Figure A-2 in Appendix A)

1  
2

**Table 3-41. Estimates of the residual-error**

Measurement abbreviation	Measurement description	GSD for "residual" error (median estimate)		
		Mouse	Rat	Human
RetDose	Retained TCE dose (mg)	-	-	1.13
CAIvPPM	TCE concentration in alveolar air (ppm)	-	-	1.44~1.83
CIInhPPM	TCE concentration in closed-chamber (ppm)	1.18	1.11~1.12	-
CMixExh	TCE concentration in mixed exhaled air (mg/L)	-	1.5	-
CArt	TCE concentration in arterial blood (mg/L)	-	1.17~1.52	-
CVen	TCE concentration in venous blood (mg/L)	<b>2.68</b>	1.22~ <b>4.46</b>	1.62~ <b>2.95</b>
CBldMix	TCE concentration in mixed arterial and venous blood (mg/L)	1.61	1.5	-
CFat	TCE concentration in fat (mg/L)	<b>2.49</b>	1.85~ <b>2.66</b>	-
CGut	TCE concentration in gut (mg/L)	-	1.86	-
CKid	TCE concentration in kidney (mg/L)	<b>2.23</b>	1.47	-
CLiv	TCE concentration in liver (mg/L)	1.71	1.67~1.78	-
CMus	TCE concentration in muscle (mg/L)	-	1.65	-
AExhpost	Amount of TCE exhaled postexposure (mg)	1.23	1.12~1.17	-
CPlasTCA	TCA concentration in plasma (mg/L)	1.40	1.13~1.21	1.12~1.17
CBldTCA	TCA concentration in blood (mg/L)	1.49	1.13~1.59	1.12~1.49
CLivTCA	TCA concentration in liver (mg/L)	1.34	1.67	-
AUrnTCA	Cumulative amount of TCA excreted in urine (mg)	1.34	1.18~1.95	1.11~1.54
AUrnTCA_collect	Cumulative amount of TCA collected in urine (noncontinuous sampling) (mg)	-	-	<b>2~2.79</b>
CTCOH	Free TCOH concentration in blood (mg/L)	1.54	1.14~1.64	1.14~ <b>2.1</b>
CLivTCOH	Free TCOH concentration in liver (mg/L)	1.59	-	-
TotCTCOH	Total TCOH concentration in blood (mg/L)	1.85	1.49	1.2~1.69
ABileTCOG	Cumulative amount of bound TCOH excreted in bile (mg)	-	<b>2.13</b>	-
CTCOG	Bound TCOH concentration in blood	-	<b>2.76</b>	-
CTCOGTCOH	Bound TCOH concentration in blood in free TCOH equivalents	1.49	-	-
CLivTCOGTCOH	Bound TCOH concentration in liver in free TCOH equivalents (mg/L)	1.63	-	-
AUrnTCOGTCOH	Cumulative amount of total TCOH excreted in urine (mg)	1.26	1.12~ <b>2.27</b>	1.11~1.13
AUrnTCOGTCOH_collect	Cumulative amount of total TCOH collected in urine (noncontinuous sampling) (mg)	-	-	1.3~1.63
AUrnTCTotMole	Cumulative amount of TCA+total TCOH excreted in urine (mmol)	-	1.12~1.54	-
CDCVGMol	DCVG concentration in blood (mmol/L)	-	-	1.53
AUrnNDCVC	Cumulative amount of NAcDCVC excreted in urine (mg)	-	1.17	1.17


3  
4

Values higher than twofold are in bold.


1 In terms of total metabolism, closed-chamber data (see Figure 3-9, panel A) were fit  
2 accurately with the updated model, with a small residual-error GSD of 1.18. While the previous  
3 analyses of Hack et al. (2006) allowed for each chamber experiment to be fit with different  
4 parameters, the current analysis made the more restrictive assumption that all experiments in a  
5 single study utilize the same parameters. Furthermore, the accuracy of closed-chamber  
6 predictions did not require the very high values for cardiac output that were used by Fisher et al.  
7 (1991), confirming the suggestion (discussed in Appendix A) that additional respiratory  
8 metabolism would resolve this discrepancy. The accurate model means that uncertainty with  
9 respect to possible wash-in/wash-out, respiratory metabolism, and extrahepatic metabolism could  
10 be well characterized. For instance, the absence of in vivo data on GSH metabolism in mice  
11 means that this pathway remains relatively uncertain; however, the current model should be  
12 reliable for estimating lower and upper bounds on the GSH pathway flux.

13 In terms of the parent compound TCE (see Figure 3-9, panels B-G), the parent PBPK  
14 model (for TCE) appears to now be robust, with the exception of the remaining over-prediction  
15 of TCE in blood following inhalation exposure. As expected, the venous-blood TCE  
16 concentration had the largest residual-error, with a GSD of 2.7, reflecting largely the difficulty in  
17 fitting TCE blood levels following inhalation exposure. In addition, the fat and kidney TCE  
18 concentrations also are somewhat uncertain, with a GSD for the residual-error of 2.5 and 2.2,  
19 respectively. These tissues were only measured in two studies, Abbas and Fisher (1997) and  
20 Greenberg et al. (1999), and the residual-error reflects the difficulties in simultaneously fitting  
21 the model to the different dose levels with the same parameters. Residual-error GSDs for other  
22 TCE measurements were less than twofold. Thus, most of the problems previously encountered  
23 with the Abbas and Fisher (1997) gavage data were solved by allowing absorption from both the  
24 stomach and duodenal compartments. Notably, the addition of possible wash-in/wash-out,  
25 respiratory metabolism, and extrahepatic metabolism (i.e., kidney GSH conjugation) was  
26 insufficient to remove the long-standing discrepancy of PBPK models over-predicting TCE  
27 blood levels from mouse inhalation exposures, suggesting another source of model or  
28 experimental error is the cause. However, the availability of tissue concentration levels of TCE  
29 somewhat ameliorates this limitation.

30 In terms of TCA and TCOH, the overall mass balance and metabolic disposition to these  
31 metabolites also appeared to be robust, as urinary excretion following dosing with TCE, TCOH,  
32 as well as TCA could be modeled accurately (see Figure 3-9, panels K and Q). The residual  
33 GSDs for the urinary excretions are small: 1.34 for TCA and 1.26 for total TCOH. In addition,  
34 the blood and tissue concentrations were also accurately predicted (see Figure 3-9, panels H-J,  
35 L-P). All the residual GSDs were less than twofold, with those for TCA measurements less  
36 than 1.5-fold. This improvement over the Hack et al. (2006) model was likely due in part to the



**Figure 3-9. Comparison of mouse data and PBPK model predictions from a random posterior sample.** Each panel shows results for a different measurement. The solid line represents prediction = data, and the grey dotted lines show prediction = data  $\times$   $GSD_{err}$  and data  $\div$   $GSD_{err}$ , where  $GSD_{err}$  is the median estimate of the residual-error GSD shown in Table 3-41.



1

2

3

4

5 **Figure 3-9 (continued). Comparison of mouse data and PBPK model**

6 **predictions from a random posterior sample.** Each panel shows results for a

7 different measurement. The solid line represents prediction = data, and the grey

8 dotted lines show prediction = data × GSD<sub>err</sub> and data ÷ GSD<sub>err</sub>, where GSD<sub>err</sub> is

9 the median estimate of the residual-error GSD shown in Table 3-41.

10

11

1  
2  
3

**Table 3-42. Summary comparison of updated PBPK model predictions and in vivo data in mice**

Study	Exposure(s)	Discussion
Abbas and Fisher (1997)	TCE gavage (corn oil)	<p>Generally, model predictions were quite good, especially with respect to tissue concentrations of TCE, TCA, and TCOH. There were some discrepancies in TCA and TCOG urinary excretion and TCA and TCOG concentrations in blood due to the requirement (unlike in Hack et al., 2006) that all experiments in the same study utilize the same parameters. Thus, for instance, TCOG urinary excretion was accurately predicted at 300 mg/kg, underpredicted at 600 mg/kg, over-predicted at 1,200 mg/kg, and underpredicted again at 2,000 mg/kg, suggesting significant intraexperimental variability (not addressed in the model).</p> <p>Population predictions were quite good, with the almost all of the data within the 95% CI of the predictions, and most within the interquartile region.</p>
Abbas et al. (1997)	TCOH, TCA i.v.	<p>Both subject-specific and population predictions were quite good. Urinary excretion, which was over-predicted by the Hack et al. (2006) model, was accurately predicted due to the allowance of additional “untracked” clearance. In the case of population predictions, almost all of the data were within the 95% CI of the predictions, and most within the interquartile region.</p>
Fisher and Allen (1993)	TCE gavage (corn oil)	<p>Both subject-specific and population predictions were quite good. Some discrepancies in the time-course of TCE blood concentrations were evidence across doses in the subject-specific predictions, but not in the population predictions, suggesting significant intrasubject variability (not addressed in the model).</p>
Fisher et al. (1991)	TCE inhalation	<p>Blood TCE levels during and following inhalation exposures were still over-predicted at the higher doses. However, there was the stringent requirement (absent in Hack et al., 2006) that the model utilize the same parameters for all doses and in both the closed and open-chamber experiments. Moreover, the Hack et al. (2006) model required significant differences in the parameters for the different closed-chamber experiments, while predictions here were accurate utilizing the same parameters across different initial concentrations. These conclusions were the same for subject-specific and population predictions (e.g., TCE blood levels remained over-predicted in the later case).</p>
Green and Prout (1985)	TCE gavage (corn oil)	<p>Both subject-specific and population predictions were adequate, though the data collection was sparse. In the case of population predictions, almost all of the data were within the 95% CI of the predictions, and about half within the interquartile region.</p>
Greenberg et al. (1999)	TCE inhalation	<p>Model predictions were quite good across a wide variety of measures that included tissue concentrations of TCE, TCA, and TCOH. However, as with the Hack et al. (2006) predictions, TCE blood levels were over-predicted by up to twofold. Population predictions were quite good, with the exception of TCE blood levels. Almost all of the other data was within the 95% CI of the predictions, and most within the interquartile region.</p>

4



**Table 3-42. Summary comparison of updated PBPK model predictions and in vivo data in mice (continued)**

Study	Exposure(s)	Discussion
Larson and Bull (1992b)	TCE gavage (aqueous)	Both subject-specific and population predictions were quite good, though the data collection was somewhat sparse. In the case of population predictions, all of the data were within the 95% CI of the predictions.
Larson and Bull (1992a)	TCA gavage (aqueous)	Both subject-specific and population predictions were quite good. In the case of population predictions, most of the data were within the interquartile region.
Merdink et al. (1998)	TCE i.v.	Both subject-specific and population predictions were quite good, though the data collection was somewhat sparse. In the case of population predictions, all of the data were within the 95% CI of the predictions.
Prout et al. (1985)	TCE gavage (corn oil)	Both subject-specific and population predictions were adequate, though there was substantial scatter in the data due to the use of single animals at each data point.
Templin et al. (1993)	TCE gavage (aqueous)	Both subject-specific and population predictions were quite good. With respect to population predictions, almost all of the other data was within the 95% CI of the predictions, and most within the interquartile region.


1  
2  
3 addition of nonurinary clearance (“untracked” metabolism) of TCA and TCOH. Also, the  
4 addition of a liver compartment for TCOH and TCOG, so that first-pass metabolism could be  
5 properly accounted for, was essential for accurate simulation of the metabolite pharmacokinetics  
6 both from intravenous (i.v.) dosing of TCOH and from exposure to TCE.  
7

### 3.5.6.1.5. Rat model and data


8 Each panel of Figure 3-10 shows a scatter plot of the calibration data and a random  
9 posterior prediction for each of the measured endpoint. The endpoint abbreviations are listed in  
10 Table 3-41, as are the implied GSDs for the “residual” errors, which include intrastudy  
11 variability, interindividual variability, and measurement and model errors. The residual-error  
12 GSDs are also shown as grey dashed or dotted lines in Figure 3-10. A summary evaluation of  
13 the predictions of the rat model as compared to the data are provided in Tables 3-43 and 3-44,  
14 with figures showing individual time-course data and predictions in Appendix A.

15 Similar to previous analyses (Hack et al., 2006), the TCE submodel for the rat appears to  
16 be robust, accurately predicting blood and tissue concentrations (see Figure 3-10, panels A-K),  
17 with residual-error GSDs generally less than twofold. The only exceptions are the predictions of  
18 venous blood from Kimmerle and Eben (1973a), which have residual-error GSDs greater than  
19 fourfold, and and predictions of fat concentrations from Simmons et al. (2002), with

1 residual-error GSD of 2.7-fold. For Kimmerle and Eben (1973b), the inaccuracy was primarily  
2 at the 3,000-ppm exposure, which might reflect other factors related to the high exposure. For



3  
4  
5  
6  
7 **Figure 3-10. Comparison of rat data and PBPK model predictions from a**  
8 **random posterior sample.** Each panel shows results for a different  
9 measurement. The solid line represents prediction = data, and the grey lines show  
10 prediction = data × GSD<sub>err</sub> and data ÷ GSD<sub>err</sub>, where GSD<sub>err</sub> is the lowest (dotted)  
11 and highest (dashed) median estimate of the residual-error GSD shown in  
12 Table 3-41.  
13



1

2

3

4

5

6

7


8

9

10

11

**Figure 3-10 (continued). Comparison of rat data and PBPK model predictions from a random posterior sample.** Each panel shows results for a different measurement. The solid line represents prediction=data, and the grey lines show prediction = data  $\times$  GSD<sub>err</sub> and data  $\div$  GSD<sub>err</sub>, where GSD<sub>err</sub> is the lowest (dotted) and highest (dashed) median estimate of the residual-error GSD shown in Table 3-41.



1

2

3

4

5

6

7

8

9

10

**Figure 3-10 (continued). Comparison of rat data and PBPK model predictions from a random posterior sample.** Each panel shows results for a different measurement. The solid line represents prediction=data, and the grey lines show prediction = data  $\times$  GSD<sub>err</sub> and data  $\div$  GSD<sub>err</sub>, where GSD<sub>err</sub> is the lowest (dotted) and highest (dashed) median estimate of the residual-error GSD shown in Table 3-41.

1 Simmons et al. (2002), the high residual-error appears to reflect scatter due to intra-study  
2 variability. Unlike in the mouse, some data consisting of TCE blood and tissue concentrations  
3 were used for “out-of-sample evaluation” (sometimes loosely termed “validation”). These data  
4 were generally well simulated (see Table 3-44); most of the data within the 95% CI of posterior  
5 predictions. This provides additional confidence in the predictions for the parent compound.  
6 In terms of TCA and TCOH, as with the mouse, the overall mass balance and metabolic  
7 disposition to these metabolites also appeared to be robust: urinary excretion following dosing  
8 with TCE, TCOH, as well as TCA, could be modeled accurately (see Figure 3-10 panels O, T,  
9 and U), with the residual-errors also indicating good predictions in most cases. Residual-error  
10 for these measurements was larger for Green and Prout (1985), Prout et al. (1985), and Stenner  
11 et al. (1997), ranging from GSD of 1.8–2.3, reflecting largely intra-study variability.  
12 Residual-errors for the other studies had GSDs of 1.1–1.5. This improvement over the Hack  
13 et al. (2006) model was likely due in part to the addition of nonurinary clearance (“untracked”  
14 metabolism) of TCA and TCOH. In addition, the addition of a liver compartment for TCOH and  
15 TCOG, so that first-pass metabolism could be properly accounted for, was essential for accurate  
16 simulation of the metabolite pharmacokinetics both from i.v. dosing of TCOH and from TCE  
17 exposure. Blood and plasma concentrations of TCA and free or total TCOH were also fairly  
18 well simulated (see Figure 3-10, panels L, M, P, Q, and S), with GSD for the residual-error of  
19 1.1–1.6. A bit more discrepancy (residual-error GSD of 1.7) was evident with TCA liver  
20 concentrations (see Figure 3-10, panel N). However, TCA liver concentrations were only  
21 available in one study (Yu et al., 2000), and the data show a change in the ratio of liver to blood  
22 concentrations at the last time point, which may be the source of the added residual-error.  
23 Predictions of biliary excretion of TCOG in bile-cannulated rats (see Figure 3-10, panel R), from  
24 Green and Prout (1985), and TCOG in blood (see Figure 3-10, panel S), from Stenner et al.  
25 (1997), were less accurate, with residual-error GSD > 2. However, the biliary excretion data  
26 consisted of a single measurement, and the amount of free TCOH in the same experiment from  
27 Stenner et al. (1997) was accurately predicted.

28 In terms of total metabolism, as with the mouse, closed-chamber data (see Figure 3-10,  
29 panel A) were fit accurately with the updated model (residual-error GSD of about 1.1). In  
30 addition, the data on NAcDCVC urinary excretion was well predicted (see Figure 3-10, panel V),  
31 with residual-error GSD of 1.18. In particular, the fact that excretion was still ongoing at the end  
32 of the experiment was accurately predicted (see Figure 3-11, panels A and B). Thus, there is  
33 greater confidence in the estimate of the flux through the GSH pathway than there was from the  
34 Hack et al. (2006) model. However, the overall flux is still estimated indirectly, and there  
35 remains some ambiguity as to the relative contributions of respiratory wash-in/wash-out,  
36 respiratory metabolism, extrahepatic metabolism, DCVC bioactivation versus *N*-acetylation, and  
37

1  
2  
3

**Table 3-43. Summary comparison of updated PBPK model predictions and in vivo data used for “calibration” in rats**

Study	Exposure(s)	Discussion
Bernauer et al. (1996)	TCE inhalation	<p>Posterior fits to these data were adequate, especially with the requirement that all predictions for dose levels utilize the same PBPK model parameters. Predictions of TCOG and TCA urinary excretion was relatively accurate, though the time-course of TCA excretion seemed to proceed more slowly with increasing dose, an aspect not captured in by model. Importantly, unlike the Hack et al. (2006) results, the time-course of NAcDCVC excretion was quite well simulated, with the excretion rate remaining nonnegligible at the last time point (48 hour). It is likely that the addition of the DCVG submodel between TCE and DCVC, along with prior distributions that accurately reflected the lack of reliable independent (e.g., in vitro) data on bioactivation, allowed for the better fit.</p>
Dallas et al. (1991)	TCE inhalation	<p>These data, consisting of arterial blood and exhaled breath concentrations of TCE, were accurately predicted by the model using both subject-specific and population sampled parameters. In the case of population predictions, most of the data were within the 95% CI of the predictions.</p>
Fisher et al. (1989)	TCE inhalation	<p>These data, consisting of closed-chamber TCE concentrations, were accurately simulated by the model using both subject-specific and population sampled parameters. In the case of population predictions, most of the data were within the 95% CI of the predictions.</p>
Fisher et al. (1991)	TCE inhalation	<p>These data, consisting of TCE blood, and TCA blood and urine time-courses, were accurately simulated by the model using both subject-specific and population sampled parameters. In the case of population predictions, most of the data were within the 95% CI of the predictions.</p>
Green and Prout (1985)	TCE gavage (corn oil) TCA i.v. TCA gavage (aqueous)	<p>For TCE treatment, these data, consisting of one time point each in urine for TCA, TCA +TCOG, and TCOG, were accurately simulated by both subject-specific and population predictions.</p> <p>For TCA i.v. treatment, the single datum of urinary TCA+TCOG at 24 hour was at the lower 95% CI in the subject-specific simulations, but accurately predicted with the population sampled parameters, suggesting intrastudy variability is adequately accounted for by population variability.</p> <p>For TCA gavage treatment, the single datum of urinary TCA+TCOG at 24 hour was accurately simulated by both subject-specific and population predictions.</p>
Hissink et al. (2002)	TCE gavage (corn oil) TCE i.v.	<p>These data, consisting of TCE blood, and TCA+TCOG urinary excretion time-courses, were accurately simulated by the model using subject-specific parameters. In the case of population predictions, TCA+TCOH urinary excretion appeared to be somewhat under-predicted.</p>
Kaneko et al. (1994b)	TCE inhalation	<p>These data, consisting of TCE blood and TCA and TCOG urinary excretion time-courses, were accurately predicted by the model using both subject-specific and population sampled parameters. In the case of population predictions, TCA+TCOH urinary excretion appeared to be somewhat underpredicted, However, all of the data were within the 95% CI of the predictions.</p>

4

**Table 3-43. Summary comparison of updated PBPK model predictions and in vivo data used for “calibration” in rats (continued)**

Study	Exposure(s)	Discussion
Keys et al. (2003)	TCE inhalation, gavage (aqueous), i.a.	These data, consisting of TCE blood, gut, kidney, liver, muscle and fat concentration time-courses, were accurately predicted by the model using both subject-specific and population sampled parameters. In the case of population predictions, most of the data were within the 95% CI of the predictions.
Kimmerle and Eben (1973a)	TCE inhalation	Some inaccuracies were noted in subject-specific predictions, particularly with TCA and TCOG urinary excretion, TCE exhalation postexposure, and TCE venous blood concentrations. In the case of TCA excretion, the rate was underpredicted at the lowest dose (49 mg/kg) and over-predicted at 330 ppm. In terms of TCOG urinary excretion, the rate was over-predicted at 175 ppm and underpredicted at 330 ppm. Similarly for TCE exhaled postexposure, there was some over-prediction at 175 ppm and some underprediction at 300 ppm. Finally, venous blood concentrations were over-predicted at 3,000 ppm. However, for population predictions, most of the data were within with 95% confidence region.
Larson and Bull (1992b)	TCA gavage (aqueous)	These data, consisting of TCA plasma time-courses, were accurately predicted by the model using both subject-specific and population sampled parameters. In the case of population predictions, all of the data were within the 95% CI of the predictions.
Larson and Bull (1992a)	TCE gavage (aqueous)	These data, consisting of TCE, TCA, and TCOH in blood, were accurately predicted by the model using both subject-specific and population sampled parameters. In the case of population predictions, all of the data were within the 95% CI of the predictions.
Lee et al. (2000a)	TCE i.v., p.v.	These data, consisting of TCE concentration time course in mixed arterial and venous blood and liver, were predicted using both the subject specific and population predictions. In both cases, most of the data were within the 95% CI of the predictions.
Merdink et al. (1999)	TCOH i.v.	TCOH blood concentrations were accurately predicted using subject-specific parameters. However, population-based parameters seemed to lead to some under-prediction, though most of the data were within the 95% CI of the predictions.
Prout et al. (1985)	TCE gavage (corn oil)	Most of these data were accurately predicted using both subject-specific and population-sampled parameters. However, at the highest two doses (1,000 and 2,000 mg/kg), there were some discrepancies in the (very sparsely collected) urinary excretion measurements. In particular, using subject-specific parameters, TCA+TCOH urinary excretion was under-predicted at 1,000 mg/kg and over-predicted at 2,000 mg/kg. Using population-sampled parameters, this excretion was underpredicted in both cases, though not entirely outside of the 95% CI.
Simmons et al. (2002)	TCE inhalation	Most of these data were accurately predicted using both subject-specific and population-sampled parameters. In the open-chamber experiments, there was some scatter in the data that did not seem to be accounted for in the model. The closed-chamber data were accurately fit.

**Table 3-43. Summary comparison of updated PBPK model predictions and in vivo data used for “calibration” in rats (continued)**

Study	Exposure(s)	Discussion
Stenner et al. (1997)	TCE intraduodenal TCOH i.v. TCOH i.v., bile-cannulated	These data, consisting of TCA and TCOH in blood and TCA and TCOG in urine, were generally accurately predicted by the model using both subject-specific and population sampled parameters. However, using subject-specific parameters, the amount of TCOG in urine was over-predicted for 100 TCOH mg/kg i.v. dosing, though total TCOH in blood was accurately simulated. In addition, in bile-cannulated rats, the TCOG excretions at 5 and 20 mg/kg i.v. were underpredicted, while the amount at 100 mg/kg was accurately predicted. On the other hand, in the case of population predictions, all of the data were within the 95% CI of the predictions, and mostly within the interquartile region, even for TCOG urinary excretion. This suggests that intrastudy variability may be a source of the poor fit in using the subject-specific parameters.
Templin et al. (1995b)	TCE oral (aqueous)	These data, consisting of TCE, TCA, and TCOH in blood, were accurately predicted by the model using both subject-specific and population sampled parameters. In the case of population predictions, all of the data were within the 95% CI of the predictions.
Yu et al. (2000)	TCA i.v.	These data, consisting of TCA in blood, liver, plasma, and urine, were generally accurately predicted by the model using both subject-specific and population sampled parameters. The only notable discrepancy was at the highest dose of 50 mg/kg, in which the rate of urinary excretion from 0–6 hour appeared to more rapid than the model predicted. However, all of the data were within the 95% CI of the predictions based on population-sampled parameters.

1  
2 i.a. = intra-arterial, i.v. = intravenous, p.v. = intraperivenous.  
3



**Table 3-44. Summary comparison of updated PBPK model predictions and in vivo data used for “out-of-sample” evaluation in rats**

Study	Exposure(s)	Discussion
Andersen et al. (1987a)	TCE inhalation	These closed-chamber data were well within the 95% CI of the predictions based on population-sampled parameters.
Bruckner et al. unpublished	TCE inhalation	These data on TCE in blood, liver, kidney, fat, muscle, gut, and venous blood, were generally accurately predicted based on population-sampled parameters. The only notable exception was TCE in the kidney during the exposure period at the 500 ppm level, which were somewhat under-predicted (though levels postexposure were accurately predicted).
Fisher et al. (1991)	TCE inhalation	These data on TCE in blood were well within the 95% CI of the predictions based on population-sampled parameters.
Jakobson et al. (1986)	TCE inhalation	These data on TCE in arterial blood were well within the 95% CI of the predictions based on population-sampled parameters.
Lee et al. (1996)	TCE i.a., i.v., p.v., gavage	Except at some very early time-points (<0.5 hour), these data on TCE in blood were well within the 95% CI of the predictions based on population-sampled parameters.
Lee et al. (2000a; 2000b)	TCE gavage	These data on TCE in blood were well within the 95% CI of the predictions based on population-sampled parameters.


i.a. = intra-arterial, i.v. = intravenous, p.v. = intraperivenous.

oxidation in the liver producing something other than TCOH or TCA. Therefore, there remains a large range of possible values for the flux through the GSH conjugation and other indirectly estimated pathways that are nonetheless consistent with all the available in vivo data. The use of noninformative priors for the metabolism parameters for which there were no in vitro data means that a fuller characterization of the uncertainty in these various metabolic pathways could be achieved. Thus, the model should be reliable for estimating lower and upper bounds on several of these pathways.

### 3.5.6.1.6. Human model and data

Each panel of Figure 3-12 shows a scatter plot of the calibration data and a random posterior prediction for each of the measured endpoint. The endpoint abbreviations are listed in Table 3-41, as are the implied GSDs for the “residual” errors, which include intrastudy variability, interindividual variability, and measurement and model errors. The residual-error GSDs are also shown as grey dashed or dotted lines in Figure 3-12. Table 3-45–3-46 provide a summary evaluation of the predictions of the model as compared to the human data, with figures showing individual time-course data and predictions in Appendix A.

1  
2




**Figure 3-11. Comparison of urinary excretion data for NAcDCVC and predictions from the Hack et al. (2006) and the updated PBPk models.** Data are from Bernauer et al. (1996) for (A and B) rats or (C and D) humans exposed for 6 hour to 40 ( $\circ$ ), 80 ( $\Delta$ ), or 160 (+) ppm in air (thick horizontal line denotes the exposure period). Predictions from Hack et al. (2006) and the corresponding data (A and C) are only for the 1,2 isomer, whereas those from the updated model (B and D) are for both isomers combined. Parameter values used for each prediction are a random sample from the subject-specific parameters from the rat and human MCMC chains (the last iteration of the first chain was used in each case). Note that in the Hack et al. (2006) model, each dose group had different model parameters, whereas in the updated model, all dose groups are required to have the same model parameters. See files linked to Appendix A for comparisons with the full distribution of predictions.

1 With respect to the TCE submodel, retained dose, blood and exhaled air measurements  
2 (see Figure 3-12, panels A-C) appeared more robust than previously found from the Hack et al.  
3 (2006) model. TCE blood concentrations from most studies were well predicted, with  
4 residual-error GSD in most studies < twofold. However, those from Chiu et al. (2007) were  
5 consistently over-predicted (i.e., data < 0.1 mg/L in Figure 3-12, panel C), with residual-error  
6 GSD of almost threefold, and a few of those from Fisher et al. (1989) were consistently  
7 underpredicted. Alveolar breath concentrations and retained dose of TCE were well predicted  
8 (residual-error GSD < 1.5-fold) from all studies except Fisher et al. (1998), which had a  
9 residual-error GSD of 1.8-fold. However, the discrepancy in alveolar breath appeared smaller  
10 than that originally reported by Fisher et al. (1998) for their PBPK model. In addition, the  
11 majority of the “out-of-sample” evaluation data consisted of TCE in blood or breath, and were  
12 generally well predicted (see Table 3-46), lending confidence to the model predictions for the  
13 parent compound.

14 In terms of TCA and TCOH, as with the mouse and rat, the overall mass balance and  
15 metabolic disposition to these metabolites also appeared to be robust, as urinary excretion  
16 following TCE exposure could be modeled accurately (see Figure 3-12, panels F, G, J, and K).  
17 In most cases, the residual-error GSD was less than twofold. However, TCA urinary data from  
18 Chiu et al. (2007) (panel G in Figure 3-12) indicated greater interoccasion variability, reflected in  
19 the residual-error GSD of 2.8. In this study, the same individual exposed to the same  
20 concentration on different occasions sometimes had substantial differences in urinary excretion.  
21 In addition, many TCA urine measurements in this study were saturated, and had to be omitted,  
22 and the fact that the remaining data were sparse and possibly censored may have contributed to  
23 the greater intrastudy variability. Blood and plasma concentrations of TCA and free TCOH (see  
24 Figure 3-12, panels D, E, and H) were fairly well simulated, with GSD for the residual-error of  
25 1.1–1.4, though total TCOH in blood (see Figure 3-12, panel I) had slightly greater residual-error  
26 with GSD of about 1.6. This partially reflects the “sharper” peak concentrations of total TCOH  
27 in the Chiu et al. (2007) data relative to the model predictions. In addition, TCA and TCOH  
28 blood and urine data were available from several studies for “out-of-sample” evaluation and  
29 were generally well predicted by the model (see Table 3-46), lending further confidence to the  
30 model predictions for these metabolites.

31 In terms of total metabolism, no closed-chamber data exist in humans, but, as discussed  
32 above, alveolar breath concentrations and retained dose (see Figure 3-12, panels A and B) were  
33 generally well simulated, suggesting that total metabolism may be fairly robust. In addition, as  
34 with the rat, the data on NAcDCVC urinary excretion was well predicted (see Figure 3-11,  
35 Figure 3-12 panel M), with residual-error GSD of 1.12). In particular the model accurately  
36 predicted the fact that excretion was still ongoing at the end of the experiment (48 hrs after the

1






2

RetDose Data

CAIvPPM Data

CVen Data






CPlasTCA Data

CBidTCA Data

AUrnTCA Data

3



AUrnTCA\_collect Data

CTCOH Data

TotCTCOH Data

4

5

6

7

8


9

10

11

12

**Figure 3-12. Comparison of human data and PBPK model predictions from a random posterior sample.** Each panel shows results for a different measurement. The solid line represents prediction = data, and the grey lines show prediction = data  $\times$   $GSD_{err}$  and data  $\div$   $GSD_{err}$ , where  $GSD_{err}$  is the lowest (dotted) and highest (dashed) median estimate of the residual-error GSD shown in Table 3-41.



1

2

3

4

5

6

7

8

9

10

11

12

**Figure 3-12 (continued). Comparison of rat data and PBPK model predictions from a random posterior sample.** Each panel shows results for a different measurement. The solid line represents prediction = data, and the grey lines show prediction = data  $\times$  GSD<sub>err</sub> and data  $\div$  GSD<sub>err</sub>, where GSD<sub>err</sub> is the lowest (dotted) and highest (dashed) median estimate of the residual-error GSD shown in Table 3-41.

1  
2  
3

**Table 3-45. Summary comparison of updated PBPK model predictions and in vivo data used for “calibration” in humans**

Reference	Exposure(s)	Discussion
Bernauer et al. (1996)	TCE inhalation	<p>These data, consisting of TCA, TCOG and NAcDCVC excreted in urine, were accurately predicted by the model using both individual-specific and population sampled parameters. The posterior NAcDCVC predictions were an important improvement over the predictions of Hack et al. (2006), which predicted much more rapid excretion than observed. The fit improvement is probably a result of the addition of the DCVG submodel between TCE and DCVC, along with the broader priors on DCVC excretion and bioactivation. Interestingly, in terms of population predictions, the NAcDCVC excretion data from this study were on the low end, though still within the 95% CI.</p>
Chiu et al. (2007)	TCE inhalation	<p>Overall, posterior predictions were quite accurate across most of the individuals and exposure occasions. TCE alveolar breath concentrations were well simulated for both individual-specific and population-generated simulations, though there was substantial scatter (intraoccasion variability). However, TCE blood concentrations were consistently over-predicted in most of the experiments, both using individual-specific and population-generated parameters. This was not unexpected, as Chiu et al. (2007) noted the TCE blood measurements to be lower by about twofold relative to previously published studies. As discussed in Chiu et al. (2007) wash-in/wash-out and extrahepatic (including respiratory) metabolism were not expected to be able to account for the difference, and indeed all these processes were added to the current model without substantially improving the discrepancy.</p> <p>With respect to metabolite data, TCA and total TCOH in blood were relatively accurately predicted. There was individual experimental variability observed for both TCA and TCOH in blood at six hours (end of exposure). The population-generated simulations over-predicted TCA in blood, while they were accurate in predicting blood TCOH. Predictions of free TCOH in blood also showed over-prediction for individual experiments, with variability at the end of exposure timepoint. However, TCOH fits were improved for the population-generated simulations. TCA and TCOG urinary excretion was generally well simulated, with simulations slightly under- or over-predicting the individual experimental data in some cases.</p>
Fisher et al. (1998)	TCE inhalation	<p>The majority of the predictions for these data were quite accurate. Interestingly, in contrast to the predictions for Chiu et al. (2007), TCE blood levels were somewhat underpredicted in a few cases, both from using individual-specific and population-generated predictions. These two results together suggest some unaccounted-for study-to-study variance, though interindividual variability cannot be discounted as the data from Chiu et al. (2007) were from individuals in the Netherlands and that from Fisher et al. (1998) were from individuals in the United States. As reported by Fisher et al. (1998), TCE in alveolar air was somewhat over-predicted in several cases, however, the discrepancies seemed smaller than originally reported for the Fisher et al. model.</p>

4

**Table 3-45. Summary comparison of updated PBPK model predictions and in vivo data used for “calibration” in humans (continued)**

Reference	Exposure(s)	Discussion
Fisher et al. (1998) (continued)	TCE inhalation (continued)	<p>With respect to metabolite data, TCOH and TCA in blood and TCOG and TCA in urine were generally well predicted, though data for some individuals appeared to exhibit inter- and/or intraoccasion variability. For example, in one case in which the same individual (female) was exposed to both 50 and 100 ppm, the TCOH blood data was over-predicted at the higher one exposure. In addition, in one individual, initial individual-specific simulations for TCA in urine were underpredicted but shifted to over-predictions towards the end of the simulations. The population-generated results over-predicted TCA in urine for the same individual. Given the results from Chiu et al. (2007) interoccasion variability is likely to be the cause, though some dose-related effect cannot be ruled out.</p> <p>Finally, DCVG data was well predicted in light of the high variability in the data and availability of only grouped data or data from multiple individual who cannot be matched to the appropriate TCE and oxidative metabolite data set. In all cases, the basic shape (plateau and then sharp decline) and order of magnitude of the time-course were well predicted. Furthermore, the range of the data was well-captured by the 95% CI of the population-generated predictions.</p>
Kimmerle and Eben (1973b)	TCE inhalation	These data were well fit by the model, using either individual-specific or population-generated parameters.
Monster et al. (1976)	TCE inhalation	The data simulated in this case were exhaled alveolar TCE, TCE in venous blood, TCA in blood, TCA in urine, and TCOG in urine. Both using individual-specific and population-generated simulations, all fits are within the 95% CI. The one exception was the retained dose for a male exposed to 65 ppm, which was outside the 95% CI for the population-generated results.
Muller et al. (1974)	TCA,  TCOH oral	<p>The data measured after oral TCA was timecourse TCA measured in plasma and urine. Individual-specific predictions were accurate, but both data sets were over-predicted in the population-generated simulations.</p> <p>The data measured after oral TCOH was timecourse TCOH in blood, TCOG in urine, TCA in plasma, and TCA in urine. Individual-specific predictions were accurate, but the population-generated simulations over-predicted TCOH in blood and TCOG in urine. The population-based TCA predictions were accurate.</p> <p>These results indicate that “unusual” parameter values were necessary in the individual-specific simulations to give accurate predictions.</p>
Paykoc et al. (1945)	TCA i.v.	These data were well fit by the model, using either individual-specific or population-generated parameters.

1  
2



**Table 3-46. Summary comparison of updated PBPK model predictions and in vivo data used for “out-of-sample” evaluation in humans**

Reference	Exposure(s)	Discussion
Bartonicek (1962)	TCE inhalation	While these data were mostly within the 95% CI of the predictions, they tended to be at the high end for all the individuals in the study.
Bloemen et al. (2001)	TCE inhalation	These data were all well within the 95% CI of the predictions.
Fernandez et al. (1977)	TCE inhalation	These data were all well within the 95% CI of the predictions.
Lapare et al. (1995)	TCE inhalation	These data were all well within the 95% CI of the predictions.
Monster et al. (1979)	TCE inhalation	These data were all well within the 95% CI of the predictions.
Muller et al. (1974; 1975)	TCE inhalation	Except for TCE in alveolar air, which was over-predicted during exposure, these data were all well within the 95% CI of the predictions.
Sato et al. (1977)	TCE inhalation	These data were all well within the 95% CI of the predictions.
Stewart et al. (1970)	TCE inhalation	These data were all well within the 95% CI of the predictions.
Triebig et al. (1976)	TCE inhalation	Except for TCE in alveolar air, these data were all well within the 95% CI of the predictions.

end of exposure). Thus, there is greater confidence in the estimate of the flux through this part of the GSH pathway than there was from the Hack et al. (2006) model, in which excretion was completed within the first few hours after exposure (see Figure 3-11, panels C and D).

If only urinary NAcDCVC data were available, as is the case for the rat, the overall GSH conjugation flux would still be estimated indirectly, and there would remain some ambiguity as to the relative contributions of respiratory wash-in/wash-out, respiratory metabolism, extrahepatic metabolism, DCVC bioactivation versus *N*-acetylation, and oxidation in the liver producing something other than TCOH or TCA. However, unlike in the rat, the blood DCVG data, while highly variable, nonetheless provide substantial constraints (at least a strong lower bound) on the flux of GSH conjugation, and is well fit by the model (see Figure 3-12, panel L, and Figure 3-13). Importantly, the high residual-error GSD for blood DCVG reflects the fact that only grouped or unmatched individual data were available, so in this case, the residual-error includes interindividual variability, which is not included in the other residual-error estimates. However, as discussed above in Section 3.3.3.2.1, there are uncertainties as to the accuracy of analytical method used by Lash et al. (1999a) in the measurement of DCVG in blood. Because these data are so determinative of the overall GSH conjugation flux, these analytical uncertainties are important to consider in the overall evaluation of the PBPK model predictions (see below, section 3.5.7).

For the other indirectly estimated pathways, there remain a large range of possible values that are nonetheless consistent with all the available in vivo data. The use of noninformative

1 priors for the metabolism parameters for which there were no in vitro data means that a fuller  
2 characterization of the uncertainty in these various metabolic pathways could be achieved. Thus,  
3 as with the rat, the model should be reliable for estimating lower and upper bounds on several of  
4 these pathways.  
5

### 3.5.6.1.7. Sensitivity Analysis With Respect to Calibration Data


6 To assess the informativeness of the calibration data to the parameters, local sensitivity  
7 analysis is performed with respect to the calibration data points. For each scaling parameter, the  
8 central difference is used to estimate the partial derivatives by centering on the sample mean of  
9 its estimated population mean, and then increasing and decreasing by 5%. The relative change in  
10 the model output  $f(\theta)$  is used to estimate a local sensitivity coefficient (SC) as follows:

$$11 \quad \quad \quad SC = 10 \times \{f(\theta_+) - f(\theta_-)\} / [\frac{1}{2} \times \{f(\theta_+) + f(\theta_-)\}]. \quad \quad \quad \text{Eq. 3-1}$$

12  
13  
14  
15  
16 Here,  $f(\theta)$  is one of the model predictions of the calibration data,  $\theta_{\pm}$  is the maximum likelihood  
17 estimate (MLE) or baseline value of  $\pm 5\%$ . For log-transformed parameters, 0.05 was added or  
18 subtracted from the baseline value, whereas for untransformed parameters, the baseline value  
19 was multiplied by 1.05 or 0.95. The resulting values of SC are binned into five categories  
20 according to their sensitivity coefficient: negligible ( $|SC| \leq 0.01$ ) very low ( $0.01 < |SC| \leq 0.1$ ),  
21 low ( $0.1 < |SC| \leq 0.5$ ), medium ( $0.5 < |SC| \leq 1.0$ ), and high ( $|SC| > 1.0$ ).

22 Note that local sensitivity analyses as typically performed in deterministic PBPK  
23 modeling can only inform the “primary” effects of parameter uncertainties—i.e., the direct  
24 change on the quantity of interest due to change in a parameter. They cannot address the  
25 *propagation* of uncertainties, such as those that can arise due to parameter correlations in the  
26 parameter fitting process. Those can only be addressed in a global sensitivity analysis, which is  
27 left for future research.


28 The results of local sensitivity analyses are shown in Figures 3-14–3-16. For each  
29 parameter, the number of data points (out of the entire calibration set) which have sensitivity  
30 coefficients in the various categories are shown graphically. As summarized in Table 3-47, most



1 of the parameters have at least some calibration data to which they are at least moderately  
 2 sensitive ( $|SC| > 0.5$ ). Across species, the cardiac output (lnQCC), ventilation-perfusion ratio


3  
 4  
 5  
 6  
 7  
 8  
 9  
 10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19

**Figure 3-13. Comparison of DCVG concentrations in human blood and predictions from the updated model.** Data are mean concentrations for males ( $\Delta$ ) and females ( $\circ$ ) reported in Lash et al. (1999a) for humans exposed for 4 hours to 100 ppm TCE in air (thick horizontal line denotes the exposure period). Data for oxidative metabolites from the same individuals were reported in Fisher et al. (1998) but could not be matched with the individual DCVG data (Lash 2007, personal communication). The vertical error bars are standard errors of the mean as reported in Lash et al. (1999a) ( $n = 8$ , so standard deviation is 80.5-fold larger). Lines are PBPK model predictions for individual male (solid) and female (dashed) subjects. Parameter values used for each prediction are a random sample from the individual-specific parameters from the human MCMC chains (the last iteration of the 1<sup>st</sup> chain was used). See files linked to Appendix A for comparisons with the full distribution of predictions.




1  
2  
3  
4  
5  
6  
7  
8

**Figure 3-14. Sensitivity analysis results: Number of mouse calibration data points with SC in various categories for each scaling parameter.**



1  
2  
3  
4  
5

**Figure 3-15. Sensitivity analysis results: Number of rat calibration data points with SC in various categories for each scaling parameter.**



1  
2  
3  
4  
5  
6

**Figure 3-16. Sensitivity analysis results: Number of human calibration data points with SC in various categories for each scaling parameter.**

1  
2  
3

**Table 3-47. Summary of scaling parameters ordered by fraction of calibration data of moderate or high sensitivity**

Mouse		Rat		Human	
Parameter	Fraction with  SC >0.5	Parameter	Fraction with  SC >0.5	Parameter	Fraction with  SC >0.5
lnV <sub>MAX</sub> C	0.4405	VLivC	0.4213	lnQCC	0.4159
VLivC	0.428	lnQCC	0.4182	lnVPRC	0.3777
lnPBC	0.3233	lnVPRC	0.4158	lnCITCOHC	0.2871
lnQCC	0.2454	lnV <sub>MAX</sub> C	0.3984	QGutC	0.2137
lnkAD	0.1675	lnPBC	0.2893	lnClGlucC	0.186
lnPBodTCAC	0.1642	VFatC	0.1455	lnkUrnTCAC	0.1789
lnVPRC	0.1575	QFatC	0.1273	FracPlasC	0.1553
lnFracTCAC	0.1323	lnPBodTCAC	0.1162	lnPBodTCOHC	0.1486
lnV <sub>MAX</sub> GlucC	0.1147	lnPFatC	0.1154	lnV <sub>MAX</sub> C	0.1358
lnPFatC	0.093	lnV <sub>MAX</sub> GlucC	0.1083	lnPBC	0.1269
lnPLivTCAC	0.0896	QGutC	0.0885	VLivC	0.1225
lnkAS	0.0863	lnkUrnTCAC	0.0696	lnPBodTCAC	0.12
VFatC	0.0762	lnPSlwC	0.0664	lnBMaxkDC	0.0897
lnKMGluc	0.0762	lnFracTCAC	0.064	VBldC	0.0586
lnkMetTCAC	0.0762	lnKMGluc	0.0625	lnkDCVGC	0.0515
lnkUrnTCAC	0.0754	lnkBileC	0.0538	lnPLivTCOGC	0.0446
lnKMC	0.0653	lnPLivTCOGC	0.0514	lnCIDCVGC	0.0435
lnkUrnTCOGC	0.0544	lnPLivC	0.0482	lnkBileC	0.0422
lnV <sub>MAX</sub> LungLivC	0.0511	lnkAD	0.0474	QFatC	0.0401
lnkTSD	0.0469	lnKMC	0.0427	lnPSlwC	0.0372
QGutC	0.0452	lnV <sub>MAX</sub> TCOHC	0.0427	QSlwC	0.0345
QFatC	0.0402	lnPKidC	0.0324	lnKMTCOH	0.0305
lnPLivC	0.0402	lnPGutC	0.03	lnPFatC	0.0292
lnPLivTCOHC	0.0377	lnFracOtherC	0.03	lnCIC	0.0288
lnPKidC	0.0352	lnPLivTCAC	0.0292	lnkUrnTCOGC	0.0282
lnPLivTCOGC	0.0352	lnBMaxkDC	0.0285	lnPRBCPlasTCAC	0.0147
lnPRBCPlasTCAC	0.031	lnkMetTCAC	0.0213	lnPLivTCAC	0.0135
lnV <sub>MAX</sub> TCOHC	0.0235	lnV <sub>MAX</sub> LungLivC	0.0182	lnkMetTCAC	0.013
lnPBodTCOHC	0.0201	lnKMTCOH	0.0182	lnFracTCAC	0.0103
lnPSlwC	0.0134	lnkAS	0.0158	lnPBodTCOGC	0.0095
lnBMaxkDC	0.0134	lnPBodTCOHC	0.015	VRapC	0.0063
lnDRespC	0.0109	FracPlasC	0.0126	VKidC	0.0057
lnkBileC	0.0084	lnkTSD	0.0103	lnClKidDCVGC	0.0057
FracPlasC	0.0059	VKidC	0.0095	lnkNATC	0.0057
lnPBodTCOGC	0.005	lnV <sub>MAX</sub> KidDCVGC	0.0095	lnPRapC	0.005
VGutC	0.0025	lnkNATC	0.0095	lnPLivTCOHC	0.005
lnPGutC	0.0025	lnDRespC	0.0063	lnkMetTCOHC	0.005
lnKMTCOH	0.0017	QSlwC	0.0055	lnFracOtherC	0.0046
lnkMetTCOHC	0.0017	lnPLivTCOHC	0.0016	VFatC	0.0036
lnkEHRC	0.0017	lnkASTCA	0.0016	lnkEHRC	0.0036
QKidC	0.0008	lnkMetTCOHC	0.0016	lnDRespC	0.0011
VKidC	0.0008	VGutC	0.0008	lnKMDCVGC	0.0011
		lnPRBCPlasTCAC	0.0008	lnkKidBioactC	0.0002
		lnkUrnTCOGC	0.0008		

4  
5  
6  
7

Parameters not shown have no data with |SC| > 0.5

1 (lnVPRC), blood-air partition coefficient (lnPBC),  $V_{MAX}$  for oxidation (ln $V_{MAX}C$ ), and  $V_{Liv}C$   
2 are consistently among the most sensitive parameters, with >10% of the calibration data  
3 exhibiting  $|SC| > 0.5$  to these parameters. Note that the reason the liver volume is sensitive is that  
4 it is used to scale the capacity or clearance rate for oxidation.

5 For scaling parameters for which all the calibration data are negligibly sensitive  
6 ( $|SC| < 0.01$ ), it is important that they either have informative prior data or are unimportant for  
7 dose-metric predictions. For mice, these parameters are the volumes of the respiratory lumen  
8 and tissue ( $V_{RespLum}C$ ,  $V_{RespEff}C$ ), the partition coefficient for the respiratory tissue  
9 (ln $P_{Resp}C$ ), and the  $V_{MAX}$  values for GSH conjugation in the liver and kidney. For the  
10 respiratory tract parameters, there are prior data to identify the parameters. Moreover, none of  
11 the dose-metric predictions are sensitive to these parameters (see Section 3.5.7.2, below). For  
12 GSH conjugation, it should be noted that for the clearance in the liver and lung ( $V_{MAX}/K_M$ ), some  
13 data are available with sensitivity  $0.01 < |SC| < 0.1$ . The data are not at all informative as to the  
14 maximum capacity for GSH conjugation.

15 For rats, all the scaling parameters have at least one calibration data point with  
16  $|SC| > 0.01$ . However, for the volumes of the respiratory lumen and tissue ( $V_{RespLum}C$ ,  
17  $V_{RespEff}C$ ), the partition coefficient for the respiratory tissue (ln $P_{Resp}C$ ), and the  $V_{MAX}$  values  
18 for GSH conjugation in the liver, these consist of only one or two data points. As with  
19 mice, there are prior data to help identify the respiratory tract parameters. Moreover, none of the  
20 dose-metric predictions are sensitive to the respiratory tract parameters (see Section 3.5.7.2,  
21 below). The data are not very informative as to maximum capacity for GSH conjugation in the  
22 liver. However, there are some data that have low or moderate informativeness ( $0.1 < |SC| < 1$ )  
23 as to the maximum capacity for GSH conjugation in the kidney, and clearance via GSH  
24 conjugation ( $V_{MAX}/K_M$ ) in the liver and kidney, which have much greater impact on the  
25 dose-metric predictions than the maximum capacity in the liver (see Section 3.5.7.2, below).

26 For humans, all the scaling parameters have at least one calibration data point with  
27  $|SC| > 0.01$ . However, for the volumes of the respiratory lumen and tissue ( $V_{RespLum}C$ ,  
28  $V_{RespEff}C$ ), the partition coefficient for the respiratory tissue (ln $P_{Resp}C$ ), and the oral  
29 absorption rate for TCA, these consist of only one or two data points. As with mice and rats,  
30 there are prior data to help identify the respiratory tract parameters. Moreover, none of the  
31 dose-metric predictions are sensitive to the respiratory or TCA oral absorption parameters (see  
32 Section 3.5.7.2, below).



1           Therefore, the local sensitivity analysis with respect to calibration data confirms that  
2 most of the scaling parameters are informed by at least some of the calibration data. In addition,  
3 the parameters for which the calibration data have very little or negligible sensitivity are either  
4 informed by prior data or have little impact on dose-metric predictions.  
5

### **3.5.6.1.8. Summary Evaluation of Updated Physiologically Based Pharmacokinetic (PBPK) Model**

6           Overall, the updated PBPK model, utilizing parameters consistent with the available  
7 physiological and in vitro data from published literature, provides reasonable fits to an extremely  
8 large database of in vivo pharmacokinetic data in mice, rats, and humans. Posterior parameter  
9 distributions were obtained by MCMC sampling using a hierarchical Bayesian population  
10 statistical model and a large fraction of this in vivo database. Convergence of the MCMC  
11 samples for model parameters was good for mice, and adequate for rats and humans. Evaluation  
12 of posterior parameter distributions suggest reasonable results in light of prior expectations and  
13 the nature of the available calibration data. In addition, in rats and humans, the model produced  
14 predictions that are consistent with in vivo data from many studies not used for calibration  
15 (insufficient studies were available in mice for such “out of sample” evaluation). Finally, the  
16 local sensitivity analysis with respect to calibration data confirms that most of the scaling  
17 parameters are informed by at least some of the calibration data, and those that were not either  
18 were informed by prior data or would not have great impact on dose-metric predictions.  
19

### **3.5.7. Physiologically Based Pharmacokinetic (PBPK) Model Dose-Metric Predictions**

#### **3.5.7.1.1. Characterization of Uncertainty and Variability**

20           Since it is desirable to characterize the contributions from both uncertainty in population  
21 parameters and variability within the population, so the following procedure is adopted. First,  
22 500 sets of population parameters (i.e., population mean and variance for each parameter) are  
23 extracted from the posterior MCMC samples—these represent the uncertainty in the population  
24 parameters. To minimize autocorrelation, they were obtained by “thinning” the chains to the  
25 appropriate degree. From each of these sets of population parameters, 100 subject-specific  
26 parameters were generated by Monte Carlo—each of these represents the population variability,  
27 given a *particular* set of population parameters. Thus a total of 50,000 subjects, representing  
28 100 (variability) each for 500 different populations (uncertainty), were generated.

1 Each set was run for a variety of generic exposure scenarios. The combined distribution  
2 of all 50,000 individuals reflects both uncertainty and variability—i.e., the case in which one is  
3 trying to predict the dosimetry for a single random subject. In addition, for each dose-metric, the  
4 mean predicted internal dose was calculated from each of the 500 sets of 100 individuals,  
5 resulting in a distribution for the uncertainty in the population mean. Comparing the combined  
6 uncertainty and variability distribution with the uncertainty distribution in the population mean  
7 gives a sense of how much of the overall variation is due to uncertainty versus variability.

8 Figures 3-17–3-25 show the results of these simulations for a number of representative  
9 dose-metrics across species continuously exposed via inhalation or orally. For display purposes,  
10 dose-metrics have been scaled by total intake (resulting in a predicted “fraction” metabolized) or  
11 exposure level (resulting in an internal dose per ppm for inhalation or per mg/kg-day for oral  
12 exposures). In these figures, the thin error bars represent the 95% CI for overall uncertainty and  
13 variability, and the thick error bars represent the 95% CI for the uncertainty in the population  
14 mean. The interpretation of these figures is that if the thick error bars are much smaller (or  
15 greater) than the thin error bars, then variability (or uncertainty) contributes the most to overall  
16 uncertainty and variability.

17 For application to human health risk assessment, the uncertainty in and variability among  
18 rodent internal dose estimates both contribute to uncertainty in human risk estimates. Therefore,  
19 it is appropriate to combine uncertainty and variability when applying rodent dose-metric  
20 predictions to quantitative risk assessment. The median and 95% CI for each dose-metric at  
21 some representative exposures in rodents are given in Tables 3-48–3-49, and the CI in these  
22 tables includes both uncertainty in the population mean and variance as well as variability in the  
23 population. On the other hand, for use in predicting human risk, it is often necessary to separate,  
24 to the extent possible, interindividual variability from uncertainty, and this disaggregation is  
25 summarized in Table 3-50.


#### 3.5.7.1.2. Local Sensitivity Analysis With Respect to Dose-Metric Predictions

27 To assess the parameter sensitivity of dose-metric predictions, a local sensitivity analysis  
28 is performed. The representative exposure scenarios in Tables 3-48–3-50 are used, but with  
29 metabolic flux dose-metrics converted to “fraction of intake” (i.e., amount metabolized through a  
30 pathway divided by total dose). Each parameter is centered on the sample mean of its estimated  
31 population mean, and then increased and decreased by 5%. The relative change in the model  
32 output  $f(\theta)$  is used to estimate a local SC as follows:


1  
2

$$SC = 10 \times \{f(\theta_+) - f(\theta_-)\} / [\frac{1}{2} \times \{f(\theta_+) + f(\theta_-)\}].$$

Eq. 3-2




1  
2 **Figure 3-17. PBPK model predictions for the fraction of intake that is**  
3 **metabolized under continuous inhalation (A) and oral (B) exposure**  
4 **conditions in mice (white), rats (diagonal hashing), and humans (horizontal**  
5 **hashing).** Bars and thin error bars represent the median estimate and 95% CI for  
6 a random subject, and reflect combined uncertainty and variability. Circles and  
7 thick error bars represent the median estimate and 95% CI for the population  
8 mean, and reflect uncertainty only.  
9




1  
 2 **Figure 3-18. PBPK model predictions for the fraction of intake that is**  
 3 **metabolized by oxidation (in the liver and lung) under continuous inhalation**  
 4 **(A) and oral (B) exposure conditions in mice (white), rats (diagonal hashing),**  
 5 **and humans (horizontal hashing).** Bars and thin error bars represent the median  
 6 estimate and 95% CI for a random subject, and reflect combined uncertainty and  
 7 variability. Circles and thick error bars represent the median estimate and 95% CI  
 8 for the population mean, and reflect uncertainty only.


9  
 10  
 11  
 12



1  
2 **Figure 3-19. PBPK model predictions for the fraction of intake that is**  
3 **metabolized by GSH conjugation (in the liver and kidney) under continuous**  
4 **inhalation (A) and oral (B) exposure conditions in mice (dotted line), rats**  
5 **(dashed line), and humans (solid line).** X-values are slightly offset for clarity.  
6 Open circles (connected by lines) and thin error bars represent the median  
7 estimate and 95% CI for a random subject, and reflect combined uncertainty and  
8 variability. Filled circles and thick error bars represent the median estimate and  
9 95% CI for the population mean, and reflect uncertainty only.




1  
 2 **Figure 3-20. PBPK model predictions for the fraction of intake that is**  
 3 **bioactivated DCVC in the kidney under continuous inhalation (A) and oral**  
 4 **(B) exposure conditions in rats (dashed line) and humans (solid line).**  
 5 X-values are slightly offset for clarity. Open circles (connected by lines) and thin  
 6 error bars represent the median estimate and 95% CI for a random subject, and  
 7 reflect combined uncertainty and variability. Filled circles and thick error bars  
 8 represent the median estimate and 95% CI for the population mean, and reflect  
 9 uncertainty only.




1  
 2 **Figure 3-21. PBPK model predictions for fraction of intake that is oxidized**  
 3 **in the respiratory tract under continuous inhalation (A) and oral (B)**  
 4 **exposure conditions in mice (dotted line), rats (dashed line), and humans**  
 5 **(solid line).** X-values are slightly offset for clarity. Open circles (connected by  
 6 lines) and thin error bars represent the median estimate and 95% CI for a random  
 7 subject, and reflect combined uncertainty and variability. Filled circles and thick  
 8 error bars represent the median estimate and 95% CI for the population mean, and  
 9 reflect uncertainty only.






1  
2  
3  
4  
5  
6  
7  
8  
9  
10


**Figure 3-22. PBPK model predictions for the fraction of intake that is “untracked” oxidation of TCE in the liver under continuous inhalation (A) and oral (B) exposure conditions in mice (dotted line), rats (dashed line), and humans (solid line)** X-values are slightly offset for clarity. Open circles (connected by lines) and thin error bars represent the median estimate and 95% CI for a random subject, and reflect combined uncertainty and variability. Filled circles and thick error bars represent the median estimate and 95% CI for the population mean, and reflect uncertainty only.



1  
2 **Figure 3-23. PBPK model predictions for the weekly AUC of TCE in venous**  
3 **blood (mg-hour/L-week) per unit exposure (ppm or mg/kg-day) under**  
4 **continuous inhalation (A) and oral (B) exposure conditions in mice (dotted**  
5 **line), rats (dashed line), and humans (solid line). X-values are slightly offset**  
6 **for clarity. Open circles (connected by lines) and thin error bars represent the**  
7 **median estimate and 95% CI for a random subject, and reflect combined**  
8 **uncertainty and variability. Filled circles and thick error bars represent the**  
9 **median estimate and 95% CI for the population mean, and reflect uncertainty**  
10 **only.**  
11



1  
2 **Figure 3-24. PBPK model predictions for the weekly AUC of TCOH in blood**  
3 **(mg-hour/L-week) per unit exposure (ppm or mg/kg-day) under continuous**  
4 **inhalation (A) and oral (B) exposure conditions in mice (dotted line), rats**  
5 **(dashed line), and humans (solid line). X-values are slightly offset for clarity.**  
6 Open circles (connected by lines) and thin error bars represent the median  
7 estimate and 95% CI for a random subject, and reflect combined uncertainty and  
8 variability. Filled circles and thick error bars represent the median estimate and  
9 95% CI for the population mean, and reflect uncertainty only.  
10



1  
2 **Figure 3-25. PBPK model predictions for the weekly AUC of TCA in the**  
3 **liver (mg-hour/L-week) per unit exposure (ppm or mg/kg-day) under**  
4 **continuous inhalation (A) and oral (B) exposure conditions in mice (dotted**  
5 **line), rats (dashed line), and humans (solid line).** X-values are slightly offset  
6 for clarity. Open circles (connected by lines) and thin error bars represent the  
7 median estimate and 95% CI for a random subject, and reflect combined  
8 uncertainty and variability. Filled circles and thick error bars represent the  
9 median estimate and 95% CI for the population mean, and reflect uncertainty  
10 only.  
11

**Table 3-48. Posterior predictions for representative internal doses: mouse**

Dose-metric	Posterior predictions for mouse dose-metrics: median (2.5%, 97.5%)				Units
	100 ppm, 7 hour/day, 5 day/wk	600 ppm, 7 hour/day, 5 day/wk	300 mg/kg-day, 5 day/wk	1,000 mg/kg-day, 5 day/wk	
ABioactDCVCBW34	0.304 (0.000534, 12.4)	2.35 (0.00603, 37)	0.676 (0.00193, 18.4)	2.81 (0.0086, 42.4)	mg/wk-kg <sup>3/4</sup>
ABioactDCVCKid	43.7 (0.0774, 1780)	336 (0.801, 5,240)	96.8 (0.281, 2,550)	393 (1.23, 6,170)	mg/wk-kg tissue
AMetGSHBW34	0.684 (0.0307, 17.6)	5.15 (0.285, 44.9)	1.66 (0.0718, 24.5)	6.37 (0.567, 49.4)	mg/wk-kg <sup>3/4</sup>
AMetLiv1BW34	170 (61.2, 403)	878 (342, 2,030)	400 (125, 610)	874 (233, 1,960)	mg/wk-kg <sup>3/4</sup>
AMetLivOtherBW34	3.81 (0.372, 38.4)	20 (1.86, 192)	8.38 (0.773, 80.1)	20 (1.55, 202)	mg/wk-kg <sup>3/4</sup>
AMetLivOtherLiv	196 (19, 2,070)	1,030 (96.5, 10,100)	437 (39.5, 4,180)	1,020 (82.1, 10,400)	mg/wk-kg tissue
AMetLngBW34	187 (7.75, 692)	263 (10.9, 2,240)	38.5 (3.49, 147)	127 (8.59, 484)	mg/wk-kg <sup>3/4</sup>
AMetLngResp	638,000 (26,500, 2,510,000)	918,000 (36,800, 7,980,000)	134,000 (12,500, 514,000)	433,000 (30,200, 1,690,000)	mg/wk-kg tissue
AUCBld	96.9 (45, 211)	822 (356, 2,040)	110 (6.95, 411)	592 (56, 1,910)	mg-hour/L-wk
AUCCTCOH	87.9 (9.9, 590)	480 (42.1, 4,140)	132 (14.4, 670)	389 (34, 2,600)	mg-hour/L-wk
AUCLivTCA	1,880 (444, 7,190)	5,070 (1,310, 18,600)	2,260 (520, 8,750)	4,660 (939, 18,900)	mg-hour/L-wk
TotMetabBW34	377 (140, 917)	1,260 (475, 3,480)	472 (165, 617)	1,110 (303, 2,010)	mg/wk-kg <sup>3/4</sup>
TotOxMetabBW34	375 (139, 916)	1,250 (451, 3,450)	465 (161, 616)	1,100 (294, 2,010)	mg/wk-kg <sup>3/4</sup>
TotTCAInBW	272 (88.9, 734)	729 (267, 1,950)	334 (106, 875)	694 (185, 1,910)	mg/wk-kg

Note: Mouse body weight is assumed to be 0.03 kg. Predictions are weekly averages over 10 weeks of the specified exposure protocol. CI reflects both uncertainties in population parameters (mean, variance) as well as population variability.

**Table 3-49. Posterior predictions for representative internal doses: rat**

Dose-metric	Posterior predictions for rat dose-metrics: median (2.5%,97.5%)				Units
	100 ppm, 7 hour/day, 5 day/wk	600 ppm, 7 hour/day, 5 day/wk	300 mg/kg-day, 5 day/wk	1,000 mg/kg-day, 5 day/wk	
ABioactDCVCBW34	0.341 (0.0306, 2.71)	2.3 (0.175, 22.6)	2.15 (0.17, 20.2)	8.89 (0.711, 84.1)	mg/wk-kg <sup>3/4</sup>
ABioactDCVCKid	67.8 (6.03, 513)	450 (35.4, 4,350)	420 (31.6, 3,890)	1,720 (134, 15,800)	mg/wk-kg tissue
AMetGSHBW34	0.331 (0.0626, 2.16)	2.27 (0.315, 19.3)	2.13 (0.293, 16)	8.84 (1.35, 69.3)	mg/wk-kg <sup>3/4</sup>
AMetLiv1BW34	176 (81.1, 344)	623 (271, 1,270)	539 (176, 1,060)	951 (273, 2,780)	mg/wk-kg <sup>3/4</sup>
AMetLivOtherBW34	45.5 (2.52, 203)	160 (7.84, 749)	134 (6.83, 659)	238 (11.3, 1390)	mg/wk-kg <sup>3/4</sup>
AMetLivOtherLiv	1,870 (92.1, 8,670)	6,660 (313, 31,200)	5,490 (280, 27,400)	9,900 (492, 59,600)	mg/wk-kg tissue
AMetLngBW34	15 (0.529, 173)	24.5 (0.819, 227)	15.1 (0.527, 115)	32.1 (1.01, 311)	mg/wk-kg <sup>3/4</sup>
AMetLngResp	41,900 (1,460, 496,000)	67,900 (2,350, 677,000)	40,800 (1,500, 325,000)	85,700 (2,660, 877,000)	mg/wk-kg tissue
AUCCBid	86.7 (39.2, 242)	1,160 (349, 2,450)	670 (47.8, 1,850)	3,340 (828, 8,430)	mg-hour/L-wk
AUCCTCOH	83.6 (1.94, 1,560)	446 (6, 10,900)	304 (4.71, 7,590)	685 (8.14, 32,500)	mg-hour/L-wk
AUCLivTCA	587 (53.7, 4,740)	2,030 (186, 13,400)	1,730 (124, 11,800)	3,130 (200, 21,000)	mg-hour/L-wk
TotMetabBW34	206 (103, 414)	682 (288, 1,430)	572 (199, 1,080)	1,030 (302, 2,920)	mg/wk-kg <sup>3/4</sup>
TotOxMetabBW34	206 (103, 414)	677 (285, 1,430)	568 (191, 1,080)	1,010 (286, 2,910)	mg/wk-kg <sup>3/4</sup>
TotTCAInBW	31.7 (3.92, 174)	110 (13.8, 490)	90.1 (10.4, 417)	164 (17.3, 800)	mg/wk-kg

Note: Rat body weight is assumed to be 0.3 kg. Predictions are weekly averages over 10 weeks of the specified exposure protocol. CI reflects both uncertainties in population parameters (mean, variance) as well as population variability.

**Table 3-50. Posterior predictions for representative internal doses: human**

Dose-metric	Posterior predictions for human dose-metrics: 2.5% population: median (2.5%, 97.5%) 50% population: median (2.5%, 97.5%) 97.5% population: median (2.5%, 97.5%)			
	Female 0.001 ppm continuous	Male 0.001 ppm continuous	Female 0.001 mg/kg-day continuous	Male 0.001 mg/kg-day continuous
ABioactDCVCBW34	0.000256 (6.97e-5, 0.000872) 0.00203 (0.00087, 0.00408) 0.0119 (0.00713, 0.0177)	0.000254 (6.94e-5, 0.000879) 0.00202 (0.000859, 0.00413) 0.012 (0.00699, 0.0182)	0.000197 (6.13e-5, 0.000502) 0.00262 (0.0012, 0.00539) 0.021 (0.0118, 0.0266)	0.0002 (6.24e-5, 0.000505) 0.00271 (0.00125, 0.00559) 0.022 (0.0124, 0.0277)
ABioactDCVCKid	0.02 (0.00549, 0.0709) 0.16 (0.0671, 0.324) 0.95 (0.56, 1.45)	0.0207 (0.00558, 0.0743) 0.163 (0.0679, 0.342) 0.979 (0.563, 1.51)	0.0152 (0.0048, 0.0384) 0.207 (0.0957, 0.43) 1.68 (0.956, 2.26)	0.016 (0.00493, 0.0407) 0.22 (0.102, 0.459) 1.81 (1.03, 2.43)
AMetGSHBW34	0.000159 (4.38e-05, 0.000539) 0.00126 (0.000536, 0.00253) 0.00736 (0.00442, 0.011)	0.000157 (4.37e-05, 0.00054) 0.00125 (0.000528, 0.00254) 0.00736 (0.00434, 0.0112)	0.000121 (3.82e-05, 0.000316) 0.00161 (0.000748, 0.00331) 0.013 (0.00725, 0.0164)	0.000123 (3.82e-05, 0.000323) 0.00167 (0.000777, 0.00343) 0.0136 (0.00759, 0.0171)
AMetLiv1BW34	0.00161 (0.000619, 0.00303) 0.00637 (0.00501, 0.00799) 0.0157 (0.0118, 0.0206)	0.00157 (0.000608, 0.00292) 0.00619 (0.00484, 0.00779) 0.0152 (0.0115, 0.02)	0.00465 (0.00169, 0.0107) 0.0172 (0.0153, 0.0183) 0.0192 (0.019, 0.0193)	0.00498 (0.00184, 0.0112) 0.018 (0.0161, 0.0191) 0.02 (0.0198, 0.0201)
AMetLivOtherBW34	4.98e-5 (8.59e-6, 0.000222) 0.000671 (0.000134, 0.00159) 0.00507 (0.00055, 0.00905)	4.87e-5 (8.33e-6, 0.000214) 0.000652 (0.000129, 0.00153) 0.00491 (0.000531, 0.00885)	0.000143 (2.35e-5, 0.000681) 0.00166 (0.00035, 0.00365) 0.00993 (0.00109, 0.0153)	0.00015 (2.49e-5, 0.000713) 0.00173 (0.000365, 0.00382) 0.0103 (0.00113, 0.0159)
AMetLivOtherLiv	0.000748 (0.000138, 0.00335) 0.0104 (0.00225, 0.0237) 0.0805 (0.00871, 0.147)	0.00065 (0.000119, 0.00288) 0.00898 (0.00193, 0.0203) 0.0691 (0.00751, 0.127)	0.00214 (0.000354, 0.00979) 0.0253 (0.00564, 0.0543) 0.157 (0.0188, 0.251)	0.00197 (0.00033, 0.00907) 0.0234 (0.00526, 0.0503) 0.146 (0.0173, 0.232)
AMetLngBW34	6.9e-6 (6.13e-7, 7.99e-5) 0.00122 (0.000309, 0.0032) 0.0123 (0.00563, 0.0197)	7.25e-6 (6.44e-7, 8.39e-5) 0.00127 (0.000325, 0.00329) 0.0124 (0.00582, 0.0199)	7.54e-8 (6.59e-9, 7.85e-7) 1.51e-5 (3.44e-6, 4.6e-5) 0.000396 (0.000104, 0.00097)	7.05e-8 (6.1e-9, 7.25e-7) 1.39e-5 (3.21e-6, 4.24e-5) 0.000366 (9.54e-5, 0.000906)

**Table 3-50. Posterior predictions for representative internal doses: human (continued)**

Dose-metric	Posterior predictions for human dose-metrics: 2.5% population: median (2.5%, 97.5%) 50% population: median (2.5%, 97.5%) 97.5% population: median (2.5%, 97.5%)			
	Female 0.001 ppm continuous	Male 0.001 ppm continuous	Female 0.001 mg/kg-day continuous	Male 0.001 mg/kg-day continuous
AMetLngResp	0.0144 (0.00116, 0.155) 2.44 (0.613, 6.71) 25.8 (12.4, 42.3)	0.0146 (0.00118, 0.157) 2.44 (0.621, 6.65) 25.3 (12.2, 41.2)	0.00015 (1.27e-05, 0.00153) 0.0313 (0.00725, 0.0963) 0.813 (0.216, 2.13)	0.000134 (1.15e-05, 0.00137) 0.0279 (0.00644, 0.086) 0.716 (0.189, 1.9)
AUCCBld	0.00151 (0.00122, 0.00186) 0.00285 (0.00252, 0.00315) 0.00444 (0.00404, 0.00496)	0.00158 (0.00127, 0.00191) 0.00295 (0.00262, 0.00326) 0.00456 (0.00416, 0.00507)	4.33e-05 (3.3e-05, 6.23e-05) 0.000229 (0.000122, 0.000436) 0.00167 (0.000766, 0.00324)	3.84e-05 (2.89e-05, 5.61e-05) 0.000204 (0.000109, 0.000391) 0.00153 (0.000693, 0.00303)
AUCCTCOH	0.00313 (0.00135, 0.00547) 0.0181 (0.0135, 0.0241) 0.082 (0.0586, 0.118)	0.00305 (0.00134, 0.00532) 0.0179 (0.0133, 0.0238) 0.0812 (0.0585, 0.117)	0.00584 (0.00205, 0.0122) 0.0333 (0.025, 0.0423) 0.115 (0.0872, 0.163)	0.00615 (0.00213, 0.0127) 0.035 (0.0264, 0.0445) 0.122 (0.0919, 0.172)
AUCLivTCA	0.0152 (0.00668, 0.0284) 0.126 (0.0784, 0.194) 0.754 (0.441, 1.38)	0.0137 (0.00598, 0.0258) 0.114 (0.0704, 0.177) 0.699 (0.408, 1.3)	0.029 (0.0116, 0.0524) 0.227 (0.138, 0.343) 1.11 (0.661, 1.87)	0.0279 (0.0114, 0.0501) 0.219 (0.133, 0.33) 1.09 (0.64, 1.88)
TotMetabBW34	0.0049 (0.00383, 0.00595) 0.0107 (0.00893, 0.0129) 0.0246 (0.0185, 0.0326)	0.00482 (0.0038, 0.00585) 0.0105 (0.00877, 0.0127) 0.0244 (0.0183, 0.0324)	0.0163 (0.0136, 0.0181) 0.0191 (0.0188, 0.0193) 0.0194 (0.0194, 0.0194)	0.0173 (0.0147, 0.019) 0.0199 (0.0196, 0.0201) 0.0202 (0.0202, 0.0202)
TotOxMetabBW34	0.00273 (0.00143, 0.00422) 0.00871 (0.0069, 0.0111) 0.0224 (0.0158, 0.0309)	0.00269 (0.00143, 0.00415) 0.00857 (0.00675, 0.011) 0.0222 (0.0155, 0.0308)	0.0049 (0.00183, 0.0108) 0.0173 (0.0154, 0.0183) 0.0192 (0.019, 0.0193)	0.00516 (0.00194, 0.0114) 0.018 (0.0161, 0.0191) 0.02 (0.0198, 0.0201)



**Table 3-50. Posterior predictions for representative internal doses: human (continued)**

<b>Dose-metric</b>	<b>Posterior predictions for human dose-metrics: 2.5% population: median (2.5%, 97.5%) 50% population: median (2.5%, 97.5%) 97.5% population: median (2.5%, 97.5%)</b>			
	<b>Female 0.001 ppm continuous</b>	<b>Male 0.001 ppm continuous</b>	<b>Female 0.001 mg/kg-day continuous</b>	<b>Male 0.001 mg/kg-day continuous</b>
TotTCAInBW	0.000259 (0.000121, 0.000422) 0.00154 (0.00114, 0.00202) 0.00525 (0.00399, 0.00745)	0.000246 (0.000114, 0.000397) 0.00146 (0.00109, 0.00193) 0.00499 (0.0038, 0.0071)	0.000501 (0.000189, 0.000882) 0.00286 (0.00222, 0.00357) 0.00659 (0.00579, 0.00724)	0.000506 (0.000192, 0.00089) 0.00289 (0.00222, 0.0036) 0.00662 (0.00581, 0.00726)

Note: Human body weight is assumed to be 70 kg for males, 60 kg for females. Predictions are weekly averages over 100 weeks of continuous exposure (dose-metric units same as previous tables). Each row represents a different population percentile (2.5, 50, and 97.5%), and the CI in each entry reflects uncertainty in population parameters (mean, variance).

1 Here,  $f(\theta)$  is one of dose-metric predictions,  $\theta_{\pm}$  is the MLE or baseline value of  $\pm 5\%$ . For  
2 log-transformed parameters, 0.05 was added or subtracted from the baseline value, whereas for  
3 untransformed parameters, the baseline value was multiplied by 1.05 or 0.95.

4 Note that local sensitivity analyses as typically performed in deterministic PBPK  
5 modeling can only inform the “primary” effects of parameter uncertainties— i.e., the direct  
6 change on the quantity of interest due to change in a parameter. They cannot address the  
7 *propagation* of uncertainties through an analysis, such as those that can arise due to parameter  
8 correlations in the parameter fitting process. Those can only be addressed in a global sensitivity  
9 analysis, which is left for future research.

10 The results of local sensitivity analyses are shown in Figures 3-26–3-31. As expected,  
11 each dose-metric is sensitive to a only a small fraction of the scaling parameters. Many of these  
12 are well-specified a priori, either due to their being physiological parameters or partition  
13 coefficients that can be measured in vitro. The remaining sensitive parameters are generally  
14 related to metabolism or clearance.

### 3.5.7.1.3. Implications for the Population Pharmacokinetics of Trichloroethylene (TCE)


#### 3.5.7.1.4. Results

16 The overall uncertainty and variability in key toxicokinetic predictions, as a function of  
17 dose and species, is shown in Figures 3-17–3-25. As expected, TCE that is inhaled or ingested is  
18 substantially metabolized in all species, predominantly by oxidation (see Figures 3-17–3-18). At  
19 higher exposures, metabolism becomes saturated and the fraction metabolized declines. Mice on  
20 average have a greater capacity to oxidize TCE than rats or humans, and this is reflected in the  
21 predictions at the two highest levels for each route. The uncertainty in the predictions for the  
22 population means for total and oxidative metabolism is relatively modest, therefore, the wide CI  
23 for combined uncertainty and variability largely reflects intersubject variability. Of particular  
24 note is the high variability in oxidative metabolism at low doses in humans, with the 95% CI  
25 spanning from 0.1–0.7 for inhalation and 0.2–1.0 for ingestion.


26 Predictions of GSH conjugation and renal bioactivation of DCVC are highly uncertain in  
27 rodents, spanning more than 1,000-fold in mice and 100-fold in rats (see Figures 3-19–3-20). In  
28 both mice and rats, the uncertainty in the population mean virtually overlaps with the combined  
29 uncertainty and variability. The uncertainty in mice reflects the lack of GSH-conjugate specific  
30 data in that species, and is, therefore, based on overall mass balance only. The somewhat smaller  
31 uncertainty in rats reflects the fact that, in addition to overall mass balance, urinary NAcDCVC

*This document is a draft for review purposes only and does not constitute Agency policy.*


- 1 excretion data are available in that species. However, while the lower bound of GSH
- 2 conjugation is informed by NAcDCVC excretion data, the upper bound for GSH conjugation and




**Figure 3-26. Sensitivity analysis results: SC for mouse scaling parameters with respect to dose-metrics following 100 ppm (light bars) and 600 ppm (dark bars), 7 h/day, 5 day/wk inhalation exposures.**




**Figure 3-27. Sensitivity analysis results: SC for mouse scaling parameters with respect to dose-metrics following 300 mg/kg-day (light bars) and 1,000 mg/kg-day (dark bars), 5 day/wk oral gavage exposures.**



**Figure 3-28. Sensitivity analysis results: SC for rat scaling parameters with respect to dose-metrics following 100 ppm (light bars) and 600 ppm (dark bars), 7 h/day, 5 day/wk inhalation exposures.**



**Figure 3-29. Sensitivity analysis results: SC for rat scaling parameters with respect to dose-metrics following 300 mg/kg-day (light bars) and 1,000 mg/kg-day (dark bars), 5 day/wk oral gavage exposures.**



**Figure 3-30. Sensitivity analysis results: SC for female (light bars) and male (dark bars) human scaling parameters with respect to dose-metrics following 0.001 ppm continuous inhalation exposures.**






Figure 3-31. Sensitivity analysis results: SC for female (light bars) and male (dark bars) human scaling parameters with respect to dose-metrics following 0.001 mg/kg-day continuous oral exposures.

1 the amount of DCVC bioactivation are still indirectly estimated from data on other clearance  
2 pathways. In humans, however, overall GSH conjugation is strongly constrained by the  
3 blood concentrations of DCVG from Lash et al. (1999a), with 95% CIs on the population mean  
4 spanning only threefold or so. DCVC bioactivation is still indirectly estimated, derived from the  
5 difference between overall GSH conjugation flux and NAcDCVC excretion data from Bernauer  
6 et al. (1996). However, substantial variability is predicted (reflecting variability in the  
7 measurements of Last et al. (1999a), for the error bars for the population mean are substantially  
8 smaller than that for overall uncertainty and variability. Of particular note is the prediction of  
9 one or two orders of magnitude more GSH conjugation and DCVC bioactivation, on average, in  
10 humans than in rats, although importantly, the 95% CIs for the predicted population means do  
11 overlap. However, as discussed above in Section 3.3.3.2.1, there are uncertainties as to the  
12 accuracy of analytical method used by Lash et al. (1999a) in the measurement of DCVG in  
13 blood. Because these data are so influential, the analytical uncertainties contribute substantially  
14 to the overall uncertainty in the estimates of the overall GSH conjugation flux, and may be  
15 greater than the statistical uncertainties calculated using the model.

16 Predictions for respiratory tract oxidative metabolism were, as expected, greatest in mice,  
17 followed by rats and then humans (see Figure 3-21). In addition, due to the “presystemic” nature  
18 of the respiratory tract metabolism model as well as the hepatic first-pass effect, substantially  
19 more metabolism was predicted from inhalation exposures as compared to oral exposures.  
20 Interestingly, the population means appeared to be fairly well constrained despite the lack of  
21 direct data, suggesting that overall mass balance is an important constraint for the presystemic  
22 respiratory tract metabolism modeled here.

23 Some constraints were also placed on “other” hepatic oxidation—i.e., through a pathway  
24 that does not result in chloral formation and subsequent formation of TCA and TCOH (see  
25 Figure 3-22). The 95% CI for overall uncertainty and variability spanned about 100-fold, a large  
26 fraction of that due to uncertainty in the population mean. Interestingly, a higher rate per kg  
27 tissue was predicted for rats than for mice or humans, although importantly, the 95% CIs for the  
28 population means overlap among all three species.

29 The AUC of TCE in blood (see Figure 3-23) showed the expected nonlinear behavior  
30 with increasing dose, with the nonlinearity more pronounced with oral exposure, as would be  
31 expected by hepatic first-pass. Notably, the predicted AUC of TCE in blood from inhalation  
32 exposures corresponds closely with cross-species ppm-equivalence, as is often assumed. For low  
33 oral exposures ( $\leq 1$  mg/kg-day), cross-species mg/kg-day equivalence appears to be fairly  
34 accurate (within twofold), implying the usual assumption of  $\text{mg/kg}^{3/4}$ -day equivalence would be  
35 somewhat less accurate, at least for humans. Interestingly, the AUC of TCOH in blood (see

1 Figure 3-24) was relatively constant with dose, reflecting the parallel saturation of both TCE  
2 oxidation and TCOH glucuronidation. In fact, in humans, the mean AUC for TCOH in blood  
3 increases up to 100 ppm or 100 mg/kg-day, due to saturation of TCOH glucuronidation, before  
4 decreasing at 1,000 ppm or 1,000 mg/kg-day, due to saturation of TCE oxidation.

5 The predictions for the AUC for TCA in the liver showed some interesting features (see  
6 Figure 3-25). The predictions for all three species with within an order of magnitude of each  
7 other, with a relatively modest uncertainty in the population mean (reflecting the substantial  
8 amount of data on TCA). The shape of the curves, however, differs substantially, with humans  
9 showing saturation at much lower doses than rodents, especially for oral exposures. In fact, the  
10 ratio between the liver TCA AUC and the rate of TCA production, although differing between  
11 species, is relatively constant as a function of dose within species (not shown). Therefore, the  
12 shape of the curves largely reflect saturation in the production of TCA from TCOH, *not* in the  
13 oxidation of TCE itself, for which saturation is predicted at higher doses, particularly via the oral  
14 route (see Figure 3-18). In addition, while for the same exposure (ppm or mg/kg-day TCE) more  
15 TCA (on a mg/kg-day basis) is produced in mice relative to rats and humans, humans and rats  
16 have longer TCA half-lives even though plasma protein binding of TCA is on average greater.

### 3.5.7.1.5. Discussion

18 This analysis substantially informs four of the major areas of pharmacokinetic  
19 uncertainty previously identified in numerous reports (reviewed in Chiu et al., 2006b): GSH  
20 conjugation pathway, respiratory tract metabolism, alternative pathways of TCE oxidation  
21 including DCA formation, and the impact of plasma binding on TCA kinetics particularly in the  
22 liver. In addition, the analysis helps identify data that have the potential to further reduce the  
23 uncertainties in TCE toxicokinetics and risk assessment.

24 With respect to the first, previous estimates of the degree of TCE GSH conjugation and  
25 subsequent bioactivation of DCVC in humans were based on urinary excretion data alone  
26 (Bernauer et al., 1996; Birner et al., 1993). For instance, Bloemen et al. (2001) concluded that  
27 due to the low yield of identified urinary metabolites through this pathway (<0.05% as compared  
28 to 20–30% in urinary metabolites of TCE oxidation), GSH conjugation of TCE is likely of minor  
29 importance. However, as noted by Lash et al. (2000a; 2000b), urinary excretion is a poor  
30 quantitative marker of flux through the GSH pathway because it only accounts for the portion  
31 detoxified, and not the portion bioactivated (a limitation acknowledged by Bloemen et al., 2001).

32 A reexamination of the available in vitro data on GSH conjugation by Chiu et al.(2006b)  
33 suggested that the difference in flux between TCE oxidation and GSH conjugation may not be as

1 large as suggested by urinary excretion data. For example, the formation rate of DCVG from  
2 TCE in freshly isolated hepatocytes was similar in order of magnitude to the rate measured for  
3 oxidative metabolites (Lash et al., 1999b; Lipscomb et al., 1998a). A closer examination of the  
4 only other available human in vivo data on GSH conjugation, the DCVG blood levels reported in  
5 Lash et al. (Lash et al., 1998a) also suggests a substantially greater flux through this pathway  
6 than inferred from urinary data. In particular, the peak DCVG blood levels reported in this study  
7 were comparable on a molar basis to peak blood levels of TCOH, the major oxidative metabolite,  
8 in the same subjects, as previously reported by Fisher et al. (1998). A lower bound estimate of  
9 the GSH conjugation flux can be derived as follows. The reported mean peak blood DCVG  
10 concentrations of 46  $\mu\text{M}$  in males exposed to 100 ppm TCE for 4 hours (Lash et al., 1999a),  
11 multiplied by a typical blood volume of 5 L (ICRP, 2003), yields a peak amount of DCVG in  
12 blood of 0.23 mmoles. In comparison, the retained dose from 100 ppm exposure for 4 hours is  
13 4.4 mmol, assuming retention of about 50% (Monster et al., 1976) and minute-volume of  
14 9 L/minute (ICRP, 2003). Thus, in these subjects, about 5% of the retained dose is present in  
15 blood as DCVG at the time of peak blood concentration. This is a strong lower bound on the  
16 total fraction of retained TCE undergoing GSH conjugation because DCVG clearance is ongoing  
17 at the time of peak concentration, and DCVG may be distributed to tissues other than blood. It  
18 should be reiterated that only grouped DCVG blood data were available for PBPK model-based  
19 analysis; however, this should only result in an underestimation of the degree of *variation* in  
20 GSH conjugation. Finally, this hypothesis of a significant flux through the human GSH  
21 conjugation pathway is consistent with the limited available total recovery data in humans in  
22 which only 60–70% of the TCE dose is recovered as TCE in breath and excreted urinary  
23 metabolites (reviewed in Chiu et al., 2007).

24 Thus, there is already substantial qualitative and semi-quantitative evidence to suggest a  
25 substantially greater flux through the GSH conjugation pathway than previously estimated based  
26 on urinary excretion data alone. The scientific utility of applying a combination of PBPK  
27 modeling and Bayesian statistical methods to this question comes from being able to  
28 systematically integrate these different types of data—in vitro and in vivo, direct (blood DCVG)  
29 and indirect (total recovery, urinary excretion)—and quantitatively assess their consistency and  
30 implications. For example, the in vitro data discussed above on GSH conjugation were used for  
31 developing prior distributions for GSH conjugation rates, and were not used in previous PBPK  
32 models for TCE. Then, both the direct and indirect in vivo data were used to the extent possible  
33 either in the Bayesian calibration or model evaluation steps.

34 However, this evidence—both qualitative and quantitative—is highly dependent on the  
35 reliability of the human DCVG measurements, both in vitro and in vivo, from Lash et al. (Lash et

1 al., 1999a; Lash et al., 1999b). In vitro, Green et al. (1997a) reported much lower rates of  
2 DCVG formation in humans using a different analytical method. Similarly, the rates of in vitro  
3 DCVG formation in rats have uneven consistency among studies. In male rat liver cytosol,  
4 Green et al. (1997a) reported a rate of 0.54 pmol/min-mg, consistent with the <2 pmol/min-mg  
5 reported by Dekant et al. (1990), but much less than the 121 pmol/min-mg reported by Lash et al.  
6 (1999b). However, in microsomes, Green et al. (1997a) reported no enzymatic formation,  
7 whereas Dekant et al. (1990) reported a higher rate (i.e., 2 pmol/min-mg) and Lash et al. (1999b)  
8 reported a much higher rate (i.e., 171 pmol/min-mg). Differing results in humans may be  
9 attributable to true interindividual variation (especially since GSTs are known to be  
10 polymorphic). However, this may be less plausible for rats, suggesting that significant  
11 uncertainties remain in the quantitative estimation of GSH conjugation flux.

12 Several other aspects of the predictions related to GSH conjugation of TCE are worthy of  
13 note. Predictions for rats and mice remain more uncertain due to their having less direct  
14 toxicokinetic data, but are better constrained by total recovery studies. For instance, the total  
15 recovery of 60-70% of dose in exhaled breath and oxidative metabolites in human studies is  
16 substantially less than the >90% reported in rodent studies (also noted by Goepfert et al., 1995).  
17 In addition, it has been suggested that “saturation” of the oxidative pathway for volatiles in  
18 general, and TCE in particular, may lead to marked increases in flux through the GSH  
19 conjugation pathway (Goepfert et al., 1995; 2004a; Slikker et al., 2004b), but the PBPK model  
20 predicts only a modest, at most ~twofold, change in flux. This is because there is evidence that  
21 both pathways are saturable in the liver for this substrate at similar exposures and because GSH  
22 conjugation also occurs in the kidney. Therefore, the available data are not consistent with  
23 toxicokinetics alone causing substantially nonlinearities in TCE kidney toxicity or cancer, or in  
24 any other effects associated with GSH conjugation of TCE.

25 Finally, the present analysis suggests a number of areas where additional data can further  
26 reduce uncertainty in and better characterize the TCE GSH conjugation pathway. The Bayesian  
27 analysis predicts a relatively low distribution volume for DCVG in humans, a hypothesis that  
28 could be tested experimentally. In addition, in vivo measurements of DCVG in blood via a  
29 different, validated analytical method, in humans with known exposures to TCE, would be  
30 highly influential in either corroborating the DCVG blood levels reported in Lash et al. (1999a)  
31 or providing evidence that those reported DCVG blood levels are too high due to analytical  
32 issues. Moreover, it would be useful in such studies to be able to match individuals with respect  
33 to toxicokinetic data on oxidative and GSH conjugation metabolites so as to better characterize  
34 variability. A consistent picture as to which GST isozymes are involved in TCE GSH  
35 conjugation, along with data on variability in isozyme polymorphisms and activity levels, can

1 further inform the extent of human variability. In rodents, more direct data on GSH metabolites,  
2 such as reliably-determined DCVG blood concentrations, preferably coupled with simultaneous  
3 data on oxidative metabolites, would greatly enhance the assessment of GSH conjugation flux in  
4 laboratory animals. Given the large apparent variability in humans, data on interstrain variability  
5 in rodents may also be useful.

6 With respect to oxidative metabolism, as expected, the liver is the major site of oxidative  
7 metabolism in all three species, especially after oral exposure, where >85% of total metabolism  
8 is oxidation in the liver in all three species. However, after inhalation exposure, the model  
9 predicts a greater proportion of metabolism via the respiratory tract than previous models for  
10 TCE. This is primarily because previous models for TCE respiratory tract metabolism (Clewell  
11 et al., 2000; Hack et al., 2006) were essentially flow-limited—i.e., the amount of respiratory tract  
12 metabolism (particularly in mice) was determined primarily by the (relatively small) blood flow  
13 to the tracheobronchial region. However, the respiratory tract structure used in the present model  
14 is more biologically plausible, is more consistent with that of other volatile organics metabolized  
15 in the respiratory tract (e.g., styrene), and leads to a substantially better fit to closed-chamber  
16 data in mice.

17 Consistent with the qualitative suggestions from in vitro data, the analysis here predicts  
18 that mice have a greater rate of respiratory tract oxidative metabolism as compared to rats and  
19 humans. However, the predicted difference of 50-fold or so on average between mice and  
20 humans is not as great as the 600-fold suggested by previous reports (Green, 2000; Green et al.,  
21 1997b; NRC, 2006). The suggested factor of 600-fold was based on multiplying the Green et al.  
22 (1997b) data on TCE oxidation in lung microsomes from rats versus mice (23-fold lower) by a  
23 factor for the total CYP content of human lung compared to rat lung (27-fold lower) (incorrectly  
24 cited as being from Raunio et al., 1998; Wheeler and Guenther, 1990). However, because of  
25 the isozyme-specificity of TCE oxidation, and the differing proportions of different isozymes  
26 across species, total CYP content may not be the best measure of interspecies differences in TCE  
27 respiratory tract oxidative metabolism. Wheeler et al. (1992) reported that CYP2E1 content of  
28 human lung microsomes is about 10-fold lower than that of human liver microsomes. Given that  
29 Green et al. (1997b) report that TCE oxidation by human liver microsomes is about threefold  
30 lower than that in mouse lung microsomes, this suggests that the mouse-to-human comparison  
31 TCE oxidation in lung microsomes would be about 30-fold. Moreover, the predicted amount of  
32 metabolism corresponds to about the detection limit reported by Green et al. (1997b) in their  
33 experiments with human lung microsomes, suggesting overall consistency in the various results.  
34 Therefore, the 50-fold factor predicted by our analysis is biologically plausible given the

1 available in vitro data. More direct in vivo measures of respiratory tract metabolism would be  
2 especially beneficial to reduce its uncertainty as well as better characterize its human variability.

3 TCA dosimetry is another uncertainty that was addressed in this analysis. In particular,  
4 the predicted interspecies differences in liver TCA AUC are modest, with a range of 10-fold or  
5 so across species, due to the combined effects of interspecies differences in the yield of TCA  
6 from TCE, plasma protein binding, and elimination half-life. This result is in contrast to  
7 previous analyses which did not include TCA protein binding (Clewell et al., 2000; Fisher,  
8 2000), which predicted significantly more than an order of magnitude difference in TCA AUC  
9 across species. In addition, in order to be consistent with available data, the model requires some  
10 metabolism or other clearance of TCA in addition to urinary excretion. That urinary excretion  
11 does not represent 100% of TCA clearance is evident empirically, as urinary recovery after TCA  
12 dosing is not complete even in rodents (Abbas et al., 1997; Yu et al., 2000). Additional  
13 investigation into possible mechanisms, including metabolism to DCA or enterohepatic  
14 recirculation with fecal excretion, would be beneficial to provide a stronger biological basis for  
15 this empirical finding.

16 With respect to “untracked” oxidative metabolism, this pathway appears to be a relatively  
17 small contribution to total oxidative metabolism. While it is tempting to use this pathway as a  
18 surrogate for DCA production through from the TCE epoxide (Cai and Guengerich, 1999), one  
19 should be reminded that DCA may be formed through multiple pathways (see Section 3.3).  
20 Therefore, this pathway at best represents a lower bound on DCA production. In addition, better  
21 quantitative markers of oxidative metabolism through the TCE epoxide pathway (e.g.,  
22 dichloroacetyl lysine protein adducts, as reported in (e.g., dichloroacetyl lysine protein adducts,  
23 as reported in Forkert et al., 2006) are needed in order to more confidently characterize its flux.

24 In a situation such as TCE in which there is large database of studies coupled with  
25 complex toxicokinetics, the Bayesian approach provides a systematic method of simultaneously  
26 estimating model parameters and characterizing their uncertainty and variability. While such an  
27 approach is not necessarily needed for all applications, such as route-to-route extrapolation (Chiu  
28 and White, 2006), as discussed in Barton et al. (2007), characterization of uncertainty and  
29 variability is increasingly recognized as important for risk assessment while representing a  
30 continuing challenge for both PBPK modelers and users. If there is sufficient reason to  
31 characterize uncertainty and variability in a highly transparent and objective manner, there is no  
32 reason why our approach could not be applied to other chemicals. However, such an endeavor is  
33 clearly not trivial, though the high level of effort for TCE is partially due to the complexity of its  
34 metabolism and the extent of its toxicokinetic database.

1           It is notable that, with experience, the methodology for the Bayesian approach to PBPK  
2 modeling of TCE has evolved significantly from that of Bois (2000a, b), to Hack et al. (2006), to  
3 the present analysis. Part of this evolution has been a more refined specification of the problem  
4 being addressed, showing the importance of “problem formulation” in risk assessment  
5 applications of PBPK modeling. The particular hierarchical population model for each species  
6 was specified based on the intended use of the model predictions, so that relevant data can be  
7 selected for analysis (e.g., excluding most grouped human data in favor of individual human  
8 data) and data can be appropriately grouped (e.g., in rodent data, grouping by sex and strain  
9 within a particular study). Thus, the predictions from the population model in rodents are the  
10 “average” for a particular “lot” of rodents of a particular species, strain, and sex. This is in  
11 contrast to the Hack et al. (2006) model, in which each dose group was treated as a separate  
12 subject. As discussed above, this previous population model structure led to the unlikely result  
13 that different dose groups within a closed-chamber study had significantly different  $V_{MAX}$  values.  
14 In humans, however, interindividual variability is of interest, and furthermore, substantial  
15 individual data are available in humans. Hack et al. (2006) mixed individual- and group-level  
16 data, depending on the availability from the published study, but this approach likely  
17 underestimates population variability due to group means being treated as individuals. In  
18 addition, in some studies, the same individual was exposed more than once, and in Hack et al.  
19 (2006), these were treated as different “individuals.” In this case, actual interindividual  
20 variability may be either over- or underestimated, depending on the degree of interoccasion  
21 variability. While it is technically feasible to include interoccasion variability, it would have  
22 added substantially to the computational burden and reduced parameter identifiability. In  
23 addition, a primary interest for this risk assessment is chronic exposure, so the predictions from  
24 the population model in humans are the “average” across different occasions for a particular  
25 individual (adult).

26           The second aspect of this evolution is the drive towards increased objectivity and  
27 transparency. For instance, available information, or the lack thereof, is formally codified and  
28 explicit either in prior distributions or in the data used to generate posterior distributions, and not  
29 both. Methods at minimizing subjectivity (and hence improving reproducibility) in parameter  
30 estimation include: (1) clear separation between the in vitro or physiologic data used to develop  
31 prior distributions and the in vivo data used to generate posterior distributions; (2) use of  
32 noninformative distributions, first updated using a probabilistic model of interspecies-scaling  
33 that allows for prediction error, for parameters lacking in prior information; and (3) use of a  
34 more comprehensive database of physiologic data, in vitro measurements, and in vivo data for  
35 parameter calibration or for out-of-sample evaluation (“validation”). These measures increase



1 the confidence that the approach employed also provides adequate characterization of the  
2 uncertainty in metabolic pathways for which available data was sparse or relatively indirect, such  
3 as GSH conjugation in rodents and respiratory tract metabolism. Moreover, this approach yields  
4 more confident insights into what additional data can reduce these uncertainties than approaches  
5 that rely on more subjective methods.  
6

### 3.5.7.1.6. Key Limitations and Potential Implications of Violating Key Assumptions

7 Like all analyses, this one has a number of limitations and opportunities for refinement,  
8 both biological and statistical. Of course, the modeling results are highly dependent on the  
9 assumed PBPK model structure. However, most of the elements of the model structure are well  
10 established for volatile, lipophilic chemicals such as TCE, and, thus, these assumptions are  
11 unlikely to introduce much bias or inaccuracy. In terms of the statistical model, a key  
12 assumption is the choice of prior and population distributions, —particularly the choice of  
13 unimodal distributions for population variability. While reasonable as a first approximation,  
14 especially without data to suggest otherwise, this assumption may introduce inaccuracies in the  
15 predictions of population variability. For example, if there were an underlying bimodal  
16 distribution, then fitting using a unimodal population distribution would lead to a high estimate  
17 for the variance, and potentially overestimate the degree of population variability. In some cases  
18 in the human model where larger population variance distributions are estimated, this may be the  
19 underlying cause. However, only in the case of GSH conjugation in humans do the larger  
20 estimates of population variability impact the dose-metric predictions used in the dose-response  
21 assessment, so the impact of this assumption is limited for this assessment.

22 In addition, certain sources of variability, such as between-animal variability in rodents  
23 and between-occasion variability in humans were not included in the hierarchical model, but  
24 were aggregated with other sources of variability in a “residual” error term. Based on the  
25 posterior predictions, it does not appear that this assumption has introduced significant bias in  
26 the estimates because the residuals between predictions and data do not overall appear  
27 systematically high or low. However, this could be verified by addressing between-animal  
28 variability in rodents (requiring a more rigorous treatment of aggregated data, e.g., Chiu and  
29 Bois, 2007) and incorporation of interoccasion variability in humans (e.g., Bernillon and Bois,  
30 2000).

31 Some key potential refinements are as follows. First would be the inclusion of a CH  
32 submodel, so that pharmacokinetic data, such as that recently published by Merdink et al. (2008),  
33 could be incorporated. In addition, the current analysis is still dependent on a model structure

1 substantially informed by deterministic analyses that test alternative model structures (Evans  
2 et al., submitted), as probabilistic methods for discrimination or selection among complex,  
3 nonlinear models such as that for TCE toxicokinetics have not yet been widely accepted.  
4 Therefore, additional refinement of the respiratory tract model may be possible, though more  
5 direct in vivo data would likely be necessary to strongly discriminating among models. In terms  
6 of validation, application of more sophisticated methods such as cross-validation, may be useful  
7 in further assessing the robustness of the modeling. Finally, additional model changes that may  
8 be of utility to risk assessment, such as development of models for different lifestages (including  
9 childhood and pregnancy), would likely require additional in vivo or in vitro data, particularly as  
10 to metabolism, to ensure model identifiability.

11

### **3.5.7.1.7. Overall Evaluation of Physiologically Based Pharmacokinetic (PBPK) Model-Based Internal Dose Predictions**

12 The utility of the PBPK model developed here for making predictions of internal dose  
13 can be evaluated based on four different components: (1) the degree to which the simulations  
14 have converged to the true posterior distribution; (2) the degree of overall uncertainty and  
15 variability; (3) for humans, the degree of uncertainty in the population; and (4) the degree to  
16 which the model predictions are consistent with in vivo data that are informative to a particular  
17 dose-metric. Table 3-51 summarizes these considerations for each dose-metric prediction. Note  
18 that this evaluation does not consider in any way the extent to which a dose-metric may be the  
19 appropriate choice for a particular toxic endpoint.

20 Overall, the least uncertain dose-metrics are the fluxes of total metabolism  
21 (TotMetabBW34), total oxidative metabolism (TotOxMetabBW34), and hepatic oxidation  
22 (AMetLiv1BW34). These all have excellent posterior convergence ( $R$  diagnostic  $\leq 1.01$ ),  
23 relatively low uncertainty and variability (GSD  $< 2$ ), and relatively low uncertainty in human  
24 population variability (GSD for population percentiles  $< 2$ ). In addition, the PBPK model  
25 predictions compare well with the available in vivo pharmacokinetic data.

26 Predictions for TCE in blood (AUCCBld) are somewhat more uncertain. Although  
27 convergence was excellent across species ( $R \leq 1.01$ ), overall uncertainty and variability was  
28 about threefold. In humans, the uncertainty in human population variability was relatively low  
29 (GSD for population percentiles  $< 1.5$ ). TCE blood level predictions were somewhat high in  
30 comparison to the Chiu et al. (2006a) study at 1 ppm, though the predictions were better for most  
31 of the other studies at higher exposure levels. In mice, TCE blood levels were somewhat  
32 over-predicted in open-chamber inhalation studies. In both mice and rats, there were some cases

1 in which fits were inconsistent across dose groups if the same parameters were used across dose  
2 groups, indicating unaccounted-for dose-related effects or intrastudy variability. However, in  
3 both rats and humans, TCE blood (humans and rats) and tissue (rats only) concentrations from

**Table 3-51. Degree of variance in dose-metric predictions due to incomplete convergence (columns 2–4), combined uncertainty and population variability (columns 5–7), uncertainty in particular human population percentiles (columns 8–10), model fits to in vivo data (column 11). The GSD is the geometric standard deviation, which is a “fold-change” from the central tendency.**

Dose-metric abbreviation	Convergence: R for generic scenarios			GSD for combined uncertainty and variability			GSD for uncertainty in human population percentiles			Comments regarding model fits to in vivo data
	Mouse	Rat	Human	Mouse	Rat	Human	1~5%	25~75%	95~99%	
ABioactDCVCBW34, ABioactDCVCKid	–	≤1.016	≤1.015	–	≤3.92	≤3.77	≤2.08	≤1.64	≤1.30	Good fits to urinary NAcDCVC, and blood DCVG.
AMetGSHBW34	≤1.011	≤1.024	≤1.015	≤9.09	≤3.28	≤3.73	≤2.08	≤1.64	≤1.29	Good fits to urinary NAcDCVC, and blood DCVG.
AMetLiv1BW34	≤1.000	≤1.003	≤1.004	≤2.02	≤1.84	≤1.97	≤1.82	≤1.16	≤1.16	Good fits to oxidative metabolites.
AMetLivOtherBW34, AMetLivOtherLiv	≤1.004	≤1.151	≤1.012	≤3.65	≤3.36	≤3.97	≤2.63	≤1.92	≤2.05	No direct in vivo data.
AMetLngBW34, AMetLngResp	≤1.001	≤1.003	≤1.002	≤4.65	≤4.91	≤10.4	≤4.02	≤2.34	≤1.83	No direct in vivo data, but good fits to closed-chamber.
AUCCBld	≤1.001	≤1.004	≤1.005	≤3.04	≤3.16	≤3.32	≤1.20	≤1.43	≤1.49	Generally good fits, but poor fit to a few mouse and human studies
AUCCTCOH	≤1.001	≤1.029	≤1.002	≤3.35	≤8.78	≤5.84	≤1.73	≤1.20	≤1.23	Good fits across all three species.
AUCLivTCA	≤1.000	≤1.005	≤1.002	≤2.29	≤3.18	≤2.90	≤1.65	≤1.30	≤1.40	Good fits to rodent data.
TotMetabBW34	≤1.001	≤1.004	≤1.004	≤1.92	≤1.82	≤1.81	≤1.13	≤1.12	≤1.18	Good fits to closed-chamber.
TotOxMetabBW34	≤1.001	≤1.003	≤1.004	≤1.94	≤1.85	≤1.96	≤1.77	≤1.15	≤1.20	Good fits to closed-chamber and oxidative metabolites.
TotTCAInBW	≤1.002	≤1.002	≤1.001	≤1.96	≤2.69	≤2.30	≤1.68	≤1.19	≤1.19	Good fits to TCA data.

1 studies not used for calibration (i.e., saved for “out-of-sample” evaluation/“validation”) were  
2 well simulated, adding confidence to the parent compound dose-metric predictions.

3 For the TCA dose-metric predictions (TotTCAInBW, AUCLivTCA) convergence in all  
4 three species was excellent ( $R \leq 1.01$ ). Overall uncertainty and variability was intermediate  
5 between dose-metrics for metabolism and that for TCE in blood, with GSD of about two to  
6 threefold. Uncertainty in human population percentiles was relatively low (GSD of 1.2–1.7).  
7 While liver TCA levels were generally well fit, the data was relatively sparse. Plasma and blood  
8 TCA levels were generally well fit, though in mice, there were again some cases in which fits  
9 were inconsistent across dose groups if the same parameters were used across dose groups,  
10 indicating unaccounted-for dose-related effects or intrastudy variability. In humans, the accurate  
11 predictions for TCA blood and urine concentrations from studies used for “out of sample”  
12 evaluation lends further confidence to dose-metrics involving TCA.

13 The evaluation of TCOH in blood followed a similar pattern. Convergence in all three  
14 species was good, though the rat model had slightly worse convergence ( $R \sim 1.03$ ) than the  
15 mouse and humans ( $R \leq 1.01$ ). In mice, overall uncertainty and variability was slightly more  
16 than for TCE in blood. There much higher overall uncertainty and variability in the rat  
17 predictions (GSD of almost 9) that likely reflects true interstudy variability. The  
18 population-generated predictions for TCOH and TCOG in blood and urine were quite wide, with  
19 some in vivo data both at the upper and lower ends of the range of predictions. In humans, the  
20 overall uncertainty and variability was intermediate between mice and rats (GSD = 5.8). As with  
21 the rats, this likely reflects true population heterogeneity, as the uncertainty in human population  
22 percentiles was relatively low (GSD of around 1.2~1.7-fold). For all three species, fits to in vivo  
23 data are generally good. In mice, however, there were again some cases in which fits were  
24 inconsistent across dose groups if the same parameters were used across dose groups, indicating  
25 unaccounted-for dose-related effects or intrastudy variability. In humans, the accurate  
26 predictions for TCOH blood and urine concentrations from studies used for “out of sample”  
27 evaluation lends further confidence to those dose-metrics involving TCOH.

28 GSH metabolism dose-metrics (ABioactDCVCBW34, ABioactDCVCKid,  
29 AMetGSHBW34) had the greatest overall uncertainty in mice but was fairly well characterized  
30 in rats and humans. In mice, there was no in vivo data informing this pathway except for the  
31 indirect constraint of overall mass balance. So although convergence was adequate ( $R < 1.02$ ),  
32 the uncertainty/variability was very large, with a GSD of ninefold for the overall flux (the  
33 amount of bioactivation was not characterized because there are no data constraining  
34 downstream GSH pathways). For rats, there were additional constraints from (well-fit) urinary  
35 NAcDCVC data, which reduced the overall uncertainty and variability substantially (GSD <

1 fourfold). In humans, in addition to urinary NAcDCVC data, DCVG blood concentration data  
2 was available, though only at the group level. These data, both of which were well fit, in  
3 addition to the greater amount of in vitro metabolism data, allowed for the flux through the GSH  
4 pathway and the rate of DCVC bioactivation to be fairly well constrained, with overall  
5 uncertainty and variability having GSD < fourfold, and uncertainty in population percentiles no  
6 more than about twofold. However, these predictions may need to be interpreted with caution,  
7 given potential analytical issues with quantifying DCVG either in vitro or in vivo (see  
8 Section 3.3.3.2). Thus, the substantial inconsistencies across studies and methods in the  
9 quantification of DCVG following TCE exposure suggest lower confidence in the accuracy of  
10 these predictions.

11 The final two dose-metrics, respiratory metabolism (AMetLngBW34, AMetLngResp)  
12 and “other” oxidative metabolism (AMetLivOtherBW34, AMetLivOtherLiv), also lacked direct  
13 in vivo data and were predicted largely on the basis of mass balance and physiological  
14 constraints. Respiratory metabolism had good convergence ( $R < 1.01$ ), helped by the availability  
15 of closed-chamber data in rodents. In rats and mice, overall uncertainty and variability was  
16 rather uncertain (GSD of four~fivefold), but the overall uncertainty and variability was much  
17 greater in humans, with a GSD of about 10-fold. This largely reflects the significant variability  
18 across individuals as well as substantial uncertainty in the low population percentiles (GSD of  
19 fourfold). However, the middle (i.e., “typical” individuals) and upper percentiles (i.e., the  
20 individuals at highest risk) are fairly well constrained with a GSD of around twofold. For the  
21 “other” oxidative metabolism dose-metric, convergence was good in mice and humans  
22 ( $R < 1.02$ ), but less than ideal in rats ( $R \sim 1.15$ ). In rodents, the overall uncertainty and  
23 variability were moderate, with a GSD around 3.5-fold, slightly higher than that for TCE in  
24 blood. The overall uncertainty and variability in this metric in humans had a GSD of about  
25 fourfold, slightly higher than for GSH conjugation metrics. However, uncertainty in the middle  
26 and upper population percentiles had GSDs of only about twofold, similar to that for respiratory  
27 metabolism.

28 Overall, as shown in Table 3-51, the updated PBPK model appears to be most reliable for  
29 the fluxes of total, oxidative, and hepatic oxidative metabolism. In addition, dose-metrics related  
30 to blood levels of TCE and oxidative metabolites TCOH and TCA had only modest uncertainty.  
31 In the case of TCE in blood, for some data sets, model predictions over-predicted the in vivo  
32 data, and, in the case of TCOH in rats, substantial interstudy variability was evident. For GSH  
33 metabolism, dose-metric predictions for rats and humans had only slightly greater uncertainty  
34 than the TCE and metabolism metrics. Predictions for mice were much more uncertain,  
35 reflecting the lack of GSD-specific in vivo data. Finally, for “other” oxidative metabolism and

1 respiratory oxidative metabolism, predictions also had somewhat more uncertainty than the TCE  
2 and metabolism metrics, though uncertainty in middle and upper human population percentiles  
3 was modest.  
4

## 4. HAZARD CHARACTERIZATION

1  
2 This section presents the hazard characterization of trichloroethylene (TCE) health  
3 effects. Because of the number of studies and their relevance to multiple endpoints, the  
4 evaluation of epidemiologic studies of cancer and TCE is summarized in Section 4.1  
5 (endpoint-specific results are presented in subsequent sections). Genotoxicity data are discussed  
6 in Section 4.2. Due to the large number of endpoints and studies in the toxicity database,  
7 subsequent sections (see Sections 4.3–4.10) are organized by tissue/organ system. Each section  
8 is further organized by noncancer and cancer endpoints, discussing data from human  
9 epidemiologic and laboratory experimental studies. In cases where there is adequate  
10 information, the role of metabolism in toxicity, comparisons of toxicity between TCE and its  
11 metabolites, and carcinogenic mode of action (MOA) are also discussed. Finally, Section 4.11  
12 summarizes the overall hazard characterization and the weight of evidence for noncancer and  
13 carcinogenic effects.  
14

### 4.1. EPIDEMIOLOGIC STUDIES ON CANCER AND TRICHLOROETHYLENE (TCE)—METHODOLOGICAL OVERVIEW

15 This brief overview of the epidemiologic studies on cancer and TCE below provides  
16 background to the discussion contained in Sections 4.4–4.10. Over 50 epidemiologic studies on  
17 cancer and TCE exposure (see Tables 4-1 through 4-3) were examined to assess their ability to  
18 inform weight-of-evidence evaluation, i.e., to inform the cancer hazard from TCE exposure,  
19 according to 15 standards of study design (see Table 4-4), conduct, and analysis. The analysis of  
20 epidemiologic studies on cancer and TCE serves to document essential design features, exposure  
21 assessment approaches, statistical analyses, and potential sources of confounding and bias. This  
22 analysis, furthermore, supports the discussion of site-specific cancer observations in  
23 Sections 4.4–4.9. In those sections, study findings are presented with an assessment and  
24 discussion of their observations according to a study’s weight of evidence and the potential for  
25 alternative explanations, including bias and confounding. Tables containing observed findings  
26 for site-specific cancers are also found in Sections 4.4–4.9. Full details of the weight-of  
27 evidence-review to identify a cancer hazard and study selections for meta-analysis may be found  
28 in Appendix B.



**Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies assessing cancer and TCE exposure**

Reference	Description	Study group (N) comparison group (N)	Exposure assessment and other information
<i>Aircraft and aerospace workers</i>			
Radican et al. (2008); Blair et al. (1998)	Civilian aircraft-maintenance workers with at least 1 yr in 1952–1956 at Hill Air Force Base, UT. VS to 1990 (Blair et al., 1998) or 2000 (Radican et al., 2008); cancer incidence 1973–1990 (Blair et al., 1998)	14,457 (7,204 ever exposed to TCE). Incidence (Blair et al., 1998) and mortality rates (Blair et al., 1998; Radican et al., 2008) of nonchemical exposed subjects.	Most subjects ( $n = 10,718$ ) with potential exposure to 1–25 solvents. Cumulative TCE assigned to individual subjects using JEM. Exposure-response patterns assessed using cumulative exposure, continuous or intermittent exposures, and peak exposure. TCE replaced in 1968 with 1,1,1-trichloroethane and was discontinued in 1978 in vapor degreasing activities. <b>Median TCE exposures were about 10 ppm for rag and bucket; 100–200 ppm for vapor degreasing.</b> Poisson regression analyses controlled for age, calendar time, sex (Blair et al., 1998) or Cox proportional hazard model for age and race.
Krishnadasan et al. (2007)	Nested case-control study within a cohort of 7,618 workers employed for between 1950 and 1992, or who had started employment before 1980 at Boeing/Rockwell/Rocketdyne [Santa Susana Field Laboratory, the UCLA cohort of (Morgenstern et al., 1997)]. Cancer incidence 1988–1999.	326 prostate cancer cases, 1,805 controls. Response rate: Cases, 69%; Controls, 60%.	JEM for TCE, hydrazine, PAHs, benzene, mineral oil constructed from company records, walk-through, or interviews. Lifestyle factors obtained from living subjects through mail and telephone surveys. Conditional logistic regression controlled for cohort, age at diagnosis, physical activity, SES and other occupational exposure (benzene, PAHs, mineral oil, hydrazine).

**Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies assessing cancer and TCE exposure (continued)**

Reference	Description	Study group (N) comparison group (N)	Exposure assessment and other information
Ritz et al. (1999); Zhao et al. (2005)	Aerospace workers with $\geq 2$ yr of employment at Rockwell/Rocketdyne (now Boeing) and who worked at Santa Susana Field Laboratory, Ventura, CA, from 1950–1993 [the UCLA cohort of (Morgenstern et al., 1997)]. Cancer mortality as of December 31, 2001. Cancer incidence 1988–2000 for subjects alive as of 1988.	6,044 (2,689 with high cumulative exposure to TCE). Mortality rates of subjects in lowest TCE exposure category. 5,049 (2,227 with high cumulative exposure to TCE). Incidence rates of subjects in lowest TCE exposure category.	JEM for TCE, hydrazine, PAHs, mineral oil, and benzene. IH ranked each job title ranked for presumptive TCE exposure as high (3), medium (2), low (1), or no (0) exposure for 3 three time periods (1951–1969, 1970–1979, 1980–1989). Cumulative TCE score: low (up to 3), medium (over 3 up to 12), high (over 12) assigned to individual subjects using JEM. Cox proportional hazard, controlled for time, since 1 <sup>st</sup> employment, SES, age at diagnosis and hydrazine.
Boice et al. (2006b)	Aerospace workers with $\geq 6$ mo employment at Rockwell/Rocketdyne (Santa Susana Field Laboratory and nearby facilities) from 1948–1999 [IEI cohort, (IEI, 2005)]. VS to 1999.	41,351, 1,642 male hourly test stand mechanics (1,111 with potential TCE exposure). Mortality rates of U.S. population and California population. Internal referent groups including male hourly nonadministrative Rocketdyne workers; male hourly, nonadministrative SSFL workers; and test stand mechanics with no potential exposure to TCE.	Potential TCE exposure assigned to test stands workers only whose tasks included the cleaning or flushing of rocket engines (engine flush) ( $n = 639$ ) or for general utility cleaning ( $n = 472$ ); potential for exposure to large quantities of TCE was much greater during engine flush than when TCE used as a utility solvent. JEM for TCE and hydrazine without semiquantitative intensity estimates. Exposure to other solvents not evaluated due to low potential for confounding (few exposed, low exposure intensity, or not carcinogenic). Exposure metrics included employment duration, employment decade, yr worked with potential TCE exposure, and yr worked with potential TCE exposure via engine cleaning, weighted by number of tests. Lifetable (SMR); Cox proportional hazard controlling for birth yr, hire yr, and hydrazine exposure.

**Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies assessing cancer and TCE exposure (continued)**

Reference	Description	Study group (N) comparison group (N)	Exposure assessment and other information
Boice et al. (1999)	Aircraft-manufacturing workers with at least 1 yr $\geq$ 1960 at Lockheed Martin (Burbank, CA). VS to 1996.	77,965 (2,267 with potential routine TCE exposures and 3,016 with routine or intermittent TCE exposure). Mortality rates of U.S. population (routine TCE exposed subjects) and nonexposed internal referents (routine and intermittent TCE exposed subjects).	12% with potential routine mixed solvent exposure and 30% with route or intermittent solvent exposure. JEM for potential TCE exposure on (1) routine basis or (2) intermittent or routine basis without semiquantitative intensity estimate. Exposure-response patterns assessed by any exposure or duration of exposure and internal control group. Vapor degreasing with TCE before 1966 and PCE, afterwards. Lifetable analyses; Poisson regression analysis adjusting for birth date, starting employment date, finishing employment date, sex and race.
Morgan et al. (1998)	Aerospace workers with $\geq$ 6 mo 1950–1985 at Hughes (Tucson, AZ). VS to 1993.	20,508 (4,733 with TCE exposures). Mortality rates of U.S. population for overall TCE exposure; mortality rates of all-other cohort subjects (internal referents).	TCE exposure intensity assigned using JEM. Exposure-response patterns assessed using cumulative exposure (low vs. high) and job with highest TCE exposure rating (peak, medium/high exposure vs. no/low exposure). <b>“High exposure” job classification defined as &gt;50 ppm.</b> Vapor degreasing with TCE 1952–1977, but limited IH data <1975. Limited IH data before 1975 and medium/low rankings likely misclassified given temporal changes in exposure intensity not fully considered (NRC, 2006).
Costa et al. (1989)	Aircraft manufacturing workers employed 1954–1981 at plant in Italy. VS to 1981.	8,626 subjects Mortality rates of the Italian population.	No exposure assessment to TCE and job titles grouped into one of four categories: blue- and white-collar workers, technical staff, and administrative clerks. Lifetable (SMR).
Garabrant et al. (1988)	Aircraft manufacturing workers $\geq$ 4 yr employment and who had worked at least 1 d at San Diego, CA, plant 1958–1982. VS to 1982.	14,067 Mortality rates of U.S. population.	TCE exposure assessment for 70 of 14,067 subjects; 14 cases of esophageal cancer and 56 matched controls. For these 70 subjects, company work records identified 37% with job title with potential TCE exposure without quantitative estimates. Lifetable (SMR).

**Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies assessing cancer and TCE exposure (continued)**

Reference	Description	Study group (N) comparison group (N)	Exposure assessment and other information
<i>Cohorts Identified From Biological Monitoring (U-TCA)</i>			
Hansen et al. (2001)	Workers biological monitored using U-TCA and air-TCE, 1947–1989. Cancer incidence from 1964–1996.	803 total. Cancer incidence rates of the Danish population.	712 with U-TCA, 89 with air-TCE measurement records, two with records of both types. U-TCA from 1947–1989; air TCE measurements from 1974. Historic median exposures estimated from the U-TCA concentrations were: 9 ppm for 1947–1964, 5 ppm for 1965–1973, 4 ppm for 1974–1979, and 0.7 ppm for 1980–1989. Air TCE measurements from 1974 onward were 19 ppm (mean) and 5 ppm (median). <b>Overall, median TCE exposure to cohort as extrapolated from air TCE and U-TCA measurements was 4 ppm (arithmetic mean, 12 ppm).</b> Exposure metrics: yr 1 <sup>st</sup> employed, employment duration, mean exposure, cumulative exposure. Exposure metrics: employment duration, average TCE intensity, cumulative TCE, period 1 <sup>st</sup> employment. Lifetable analysis (SIR).
Anttila et al. (1995)	Workers biological monitored using U-TCA, 1965–1982. VS 1965–1991 and cancer incidence 1967–1992.	3,974 total (3,089 with U-TCA measurements). Mortality and cancer incidence rates of the Finnish population.	Median U-TCA, 63 µmol/L for females and 48 µmol/L for males; mean U-TCA was 100 µmol/L. Average 2.5 U-TCA measurements per individual. Using the Ikeda et al. (1972) relationship for TCE exposure to U-TCA, <b>TCE exposures were roughly 4 ppm (median) and 6 ppm (mean).</b> Exposure metrics: yr since 1 <sup>st</sup> measurement. Lifetable analysis (SMR, SIR).
Axelsson et al. (1994)	Workers biological monitored using U-TCA, 1955–1975. VS to 1986 and cancer incidence 1958–1987.	1,421 males. Mortality and cancer incidence rates of Swedish male population.	Biological monitoring for U-TCA from 1955 and 1975. <b>Roughly ¾ of cohort had U-TCA concentrations equivalent to &lt;20 ppm TCE.</b> Exposure metrics: duration exposure, mean U-TCA. Lifetable analysis (SMR, SIR).
<i>Other Cohorts</i>			
Clapp and Hoffman (2008)	Deaths between 1969–2001 among employees ≥5 yr employment duration at an IBM facility (Endicott, NY).	360 deaths. Proportion of deaths among New York residents during 1979–1998.	No exposure assessment to TCE. PMR analysis.

**Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies assessing cancer and TCE exposure (continued)**

Reference	Description	Study group (N) comparison group (N)	Exposure assessment and other information
Sung et al. (2007; 2008)	Female workers 1 <sup>st</sup> employed 1973–1997 at an electronics (RCA) manufacturing factory (Taoyuan, Taiwan). Cancer incidence 1979–2001 (Sung et al., 2007). Childhood leukemia 1979–2001 among first born of female subjects in Sung et al. (2007; 2008)	63,982 females and 40,647 females with 1 <sup>st</sup> live born offspring. Cancer incidence rates of Taiwan population (Sung et al., 2007). Childhood leukemia incidence rates of first born live births of Taiwan population (Sung et al., 2007).	No exposure assessment. Chlorinated solvents including TCE and PCE found in soil and groundwater at factory site. Company records indicated TCE not used 1975–1991 and PCE 1975–1991 and PCE after 1981. No information for other time periods. Exposure-response using employment duration. Lifetable analysis (SMR, SIR) (Chang et al., 2003; Chang et al., 2005; Sung et al., 2007) or Poisson regression adjusting for maternal age, education, sex, and birth yr (Sung et al., 2008).
Chang et al. (2005),	Male and female workers employed 1978–1997 at electronics factory as studied by Sung et al. (2007). VS from 1985–1997 and cancer incidence 1979–1997.	86,868 total. Incidence (Chang et al., 2005) mortality (Chang et al., 2003) rates Taiwan population.	
ATSDR (2004)	Workers 1952–1980 at the View-Master factory (Beaverton, OR).	616 deaths 1989–2001. Proportion of deaths between 1989–2001 in Oregon population.	No exposure information on individual subjects. TCE and other VOCs detected in well water at the time of the plant closure in 1998 were TCE, 1,220–1,670 µg/L; 1,1-DCE, up to 33 µg/L; and, PCE up to 56 µg/L. PMR analysis.
Raaschou-Nielsen et al. (2003)	Blue-collar workers employed >1968 at 347 Danish TCE-using companies. Cancer incidence through 1997.	40,049 total (14,360 with presumably higher level exposure to TCE). Cancer incidence rates of the Danish population.	Employers had documented TCE usage. Blue-collar vs. white-collar workers and companies with <200 workers were variables identified as increasing the likelihood for TCE exposure. Subjects from iron and metal, electronics, painting, printing, chemical, and dry cleaning industries. <b>Median exposures to trichloroethylene were 40–60 ppm for the yr before 1970, 10–20 ppm for 1970–1979, and approximately 4 ppm for 1980–1989.</b> Exposure metrics: employment duration, yr 1 <sup>st</sup> employed, and # employees in company. Lifetable (SIR).

**Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies assessing cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Description</b>	<b>Study group (N) comparison group (N)</b>	<b>Exposure assessment and other information</b>
Ritz (1999a)	Male uranium-processing plant workers $\geq 3$ mo employment 1951–1972 at DOE facility (Fernald, OH). VS 1951–1989, cancer.	3,814 white males monitored for radiation (2,971 with potential TCE exposure). Mortality rates of the U.S. population; non-TCE exposed internal controls for TCE exposure-response analyses.	JEM for TCE, cutting fluids, kerosene, and radiation generated by employees and industrial hygienists. Subjects assigned potential TCE according to intensity: light (2,792 subjects), moderate (179 subjects), heavy (no subjects). Lifetable (SMR) and conditional logistic regression adjusted for pay status, date first hire, radiation.
Henschler et al. (1995)	Male workers $\geq 1$ yr 1956–1975 at cardboard factory (Arnsberg region, Germany). VS to 1992.	169 exposed; 190 unexposed. Mortality rates from German Democratic Republic (broad categories) or renal cell carcinoma incidence rates from Danish population, German Democratic, or non-TCE exposed subjects.	Walk-through surveys and employee interviews used to identify work areas with TCE exposure. TCE exposure assigned to renal cancer cases using workman's compensation files. Lifetable (SMR, SIR) or Mantel-Haenszel.
Greenland et al. (1994)	Cancer deaths, 1969–1984, among pensioned workers employed <1984 at GE transformer manufacturing plant (Pittsfield, MA), and who had job history record; controls were noncancer deaths among pensioned workers.	512 cases, 1,202 controls. Response rate: Cases, 69%; Controls, 60%.	Industrial hygienist assessment from interviews and position descriptions. TCE (no/any exposure) assigned to individual subjects using JEM. Logistic regression.
Sinks et al. (1992)	Workers employed 1957–1980 at a paperboard container manufacturing and printing plant (Newnan, GA). VS to 1988. Kidney and bladder cancer incidence through 1990.	2,050 total. Mortality rates of the U.S. population, bladder and kidney cancer incidence rates from the Atlanta-SEER registry for the yr 1973–1977.	No exposure assessment to TCE; analyses of all plant employees including white- and blue-collar employees. Assignment of work department in case-control study based upon work history; Material Safety Data Sheets identified chemical usage by department. Lifetable (SMR, SIR) or conditional logistic regression adjusted for hire date and age at hire, and using 5- and 10-yr lagged employment duration.

**Table 4-1. Description of epidemiologic cohort and proportionate mortality ratio (PMR) studies assessing cancer and TCE exposure (continued)**

Reference	Description	Study group (N) comparison group (N)	Exposure assessment and other information
Blair et al. (1989)	Workers employed 1942–1970 in U.S. Coast. VS to 1980.	3,781 males of whom 1,767 were marine inspectors (48%). Mortality rates of the U.S. population. Mortality rates of marine inspectors also compared to that of noninspectors.	No exposure assessment to TCE. Marine inspectors worked in confined spaces and had exposure potential to multiple chemicals. TCE was identified as one of 10 potential chemical exposures. Lifetable (SMR) and directly adjusted relative risks.
Shannon et al. (1988)	Workers employed $\geq 6$ mo at GE lamp manufacturing plant, 1960–1975. Cancer incidence from 1964–1982.	1,870 males and females, 249 (13%) in coiling and wire-drawing area. Cancer incidence rates from Ontario Cancer Registry.	No exposure assessment to TCE. Workers in CWD had potential exposure to many chemicals including metals and solvents. A 1955-dated engineering instruction sheet identified trichloroethylene used as degreasing solvent in CWD. Lifetable (SMR).
Shindell and Ulrich (1985)	Workers employed $\geq 3$ mo at a TCE manufacturing plant 1957–1983. VS to 1983.	2,646 males and females. Mortality rates of the U.S. population.	No exposure assessment to TCE; job titles categorized as either white- or blue-collar. Lifetable analysis (SMR).
Wilcosky et al. (1984)	Respiratory, stomach, prostate, lymphosarcoma, and lymphatic leukemia cancer deaths 1964–1972 among 6,678 active and retired production workers at a rubber plant (Akron, OH); controls were a 20% age-stratified random sample of the cohort.	183 cases (101 respiratory, 33 prostate, 30 stomach, 9 lymphosarcoma and 10 lymphatic leukemia cancer deaths).	JEM without quantitative intensity estimates for 20 exposures including TCE. Exposure metric: ever held job with potential TCE exposure.

1, 1- DCE = dichloroethylene, CWD = coiling and wire drawing; DOE = U.S. Department of Energy, GE = General Electric, IBM = International Business Machines Corporation, IEI = International Epidemiology Institute, IH = industrial hygienist, JEM = job-exposure matrix, NRC = National Research Council, PAH = polycyclic aromatic hydrocarbon, PCE = perchloroethylene, PMR = proportionate mortality ratio, SES = socioeconomic status, SIR = standardized incidence ratio, SMR = standardized mortality ratio, SSFL = Santa Susanna Field Laboratory, U-TCA = urinary trichloroacetic acid, UCLA = University of California, Los Angeles, VOCs = volatile organic compounds, VS = vital status.

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
<i>Bladder</i>			
Pesch et al. (2000a)	Histologically confirmed urothelial cancer (bladder, ureter, renal pelvis) cases from German hospitals (5 regions) in 1991–1995; controls randomly selected from residency registries matched on region, sex, and age.	1,035 cases. 4,298 controls. Cases, 84%; Controls, 71%.	Occupational history using job title or self-reported exposure. JEM and JTEM to assign exposure potential to metals and solvents (chlorinated solvents, TCE, PCE). Lifetime exposure to TCE exposure examined as 30 <sup>th</sup> , 60 <sup>th</sup> , and 90 <sup>th</sup> percentiles (medium, high, and substantial) of exposed control exposure index. Duration used to examine occupational title and job task duties and defined as 30 <sup>th</sup> , 60 <sup>th</sup> , and 90 <sup>th</sup> percentiles (medium, long, and very long) of exposed control durations. Logistic regression with covariates for age, study center, and smoking.
Siemiatycki et al. (1994) Siemiatycki (1991)	Male bladder cancer cases, age 35–75 yr, diagnosed in 16 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	484 cases. 533 population controls; 740 other cancer controls. Cases, 78%; Controls, 72%.	JEM to assign 294 exposures including TCE on semiquantitative scales categorized as any or substantial exposure. Other exposure metrics included exposure duration in occupation or job title. Logistic regression adjusted for age, ethnic origin, socioeconomic status, smoking, coffee consumption, and respondent status (occupation or job title) or Mantel-Haenszel stratified on age, income, index for cigarette smoking, coffee consumption, and respondent status (TCE).
<i>Brain</i>			
DeRoos et al. (2001a); Olshan et al. (1999)	Neuroblastoma cases in children of <19 yr selected from Children’s Cancer Group and Pediatric Oncology Group with diagnosis in 1992–1994; population controls (RDD) matched to control on birth date.	504 cases. 504 controls. Cases, 73% Controls, 74%.	Telephone interview with parent using questionnaire to assess parental occupation and self-reported exposure history and judgment-based attribution of exposure to chemical classes (halogenated solvents) and specific solvents (TCE). Exposure metric was any potential exposure. Logistic regression with covariate for child’s age and material race, age, and education.



**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Heineman et al. (1994)	White, male cases, age $\geq$ 30 yr, identified from death certificates in 1978–1981; controls identified from death certificates and matched for age, yr of death and study area.	300 cases. 386 controls. Cases, 74%; Controls, 63%.	In-person interview with next-of-kin; questionnaire assessing lifetime occupational history using job title and JEM of Gomez et al. (1994). Cumulative exposure metric (low, medium or and high) based on weighted probability and duration. Logistic regression with covariates for age and study area.
<i>Colon and Rectum</i>			
Goldberg et al. (2001); Siemietycki (1991)	Male colon cancer cases, 35–75 yr, from 16 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	497 cases. 533 population controls and 740 cancer controls. Cases, 82%; Controls, 72%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (294 exposures on semiquantitative scales); potential TCE exposure defined as any or substantial exposure. Logistic regression adjusted for age, ethnic origin, birthplace, education, income, parent's occupation, smoking, alcohol consumption, tea consumption, respondent status, heating source socioeconomic status, smoking, coffee consumption, and respondent status (occupation, some chemical agents) or Mantel-Haenszel stratified on age, income, index for cigarette smoking, coffee consumption, and respondent status (TCE).
Dumas et al. (2000); Simeitycki (1991)	Male rectal cancer cases, age 35–75 yr, diagnosed in 16 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	292 cases. 533 population controls and 740 other cancer controls. Cases, 78%; Controls, 72%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (294 exposures on semiquantitative scales); potential TCE exposure defined as any or substantial exposure. Logistic regression adjusted for age, education, respondent status, cigarette smoking, beer consumption and body mass index (TCE) or Mantel-Haenszel stratified on age, income, index for cigarette smoking, coffee consumption, ethnic origin, and beer consumption (TCE).

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Fredriksson et al. (1989)	Colon cancer cases aged 30–75 yr identified through the Swedish Cancer Registry among patients diagnosed in 1980–1983; population-based controls were frequency-matched on age and sex and were randomly selected from a population register.	329 cases. 658 controls. Not available.	Mailed questionnaire assessing occupational history with telephone interview follow-up. Self-reported exposure to TCE defined as any exposure. Mantel-Haenszel stratified on age, sex, and physical activity.
<i>Esophagus</i>			
Parent et al. (2000b); Siemiatycki (1991)	Male esophageal cancer cases, 35–75 yr, diagnosed in 19 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	292 cases. 533 population controls; 740 subjects with other cancers. Cases, 78%; controls, 72%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (294 exposures on semiquantitative scales); potential TCE exposure defined as any or substantial exposure. Logistic regression adjusted for age, education, respondent status, cigarette smoking, beer consumption and body mass index (solvents) or Mantel-Haenszel stratified on age, income, index for cigarette smoking, coffee consumption, ethnic origin, and beer consumption (TCE).

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

Reference	Population	Study group (N) comparison group (N) response rates	Exposure assessment and other information
<i>Lymphoma</i>			
Purdue et al. (2011)	Cases aged 20–74 with histologically-confirmed NHL (B-cell diffuse and follicular, T-cell, lymphoreticular) without HIV in 1998–2000 and identified from four SEER areas (Los Angeles County and Detroit metropolitan area, random sample; Seattle_Puget Sound and Iowa, all consecutive cases); population controls aged 20–74 with no previous diagnosis of HIV infection or NHL, identified through (1) if >65 yr of age, RDD, or (2) if ≥65 yr, identified from Medicare eligibility files and stratified on geographic area, age, and race.	1,321 cases. 1,057 controls. Cases, 76%; Controls, 78%.	In-person interview using questionnaire or computer-assisted personal interview questionnaire specific for jobs held for >1 yr since the age of 16 yr, hobbies, and medical and family history. For occupational history, 32 job- or industry-specific interview modules asked for detailed information on individual jobs and focused on solvents exposure, including TCE, assessment by expert industrial hygienist blinded to case and control status by levels of probability, frequency, and intensity. Exposure metric of overall exposure, average weekly exposure, yr exposed, average exposure intensity, and cumulative exposure. Logistic regression adjusted for sex, age, race, education and SEER site.
Gold et al.	Cases aged 35–74 with histologically-confirmed multiple myeloma in 2000–2002 and identified from Seer areas (Detroit, Seattle-Puget Sound); population controls.	181 cases. 481 controls. Cases, 71%; Controls, 52%.	In-person interview using computer-assisted personal interview questionnaire for jobs held ≥1 yr since 1941 (cases) or 1946 (controls) and since age 18 yr. For occupational history, 20 occupations, job- or industry-specific interview modules asked for detailed information on individual jobs held at least 2 yr and focused on solvents exposure, including TCE, assessment by expert industrial hygienist blinded to case and control status by levels of probability, duration and cumulative exposure. Logistic regression adjusted for sex, age, race, education and SEER site.

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Cocco et al. (2010)	Histologically or cytologically confirmed cases aged $\geq 17$ yr with lymphoma (B-cell, T-cell, CLL, multiple myeloma, Hodgkin) in 1998–2004 and residents of referral areas from seven European countries (Czech Republic, Finland, France, Germany, Ireland, Italy, and Spain); hospital (4 participating countries) or population controls (all others); controls from (1) Germany and Italy selected by RDD from general population and matched (individually in German and group-based in Italy) to cases by sex, age and residence area, and, (2) for all other countries, matched hospital controls with diagnoses other than cancer, infectious diseases and immunodeficient diseases (individually in Czech Republic group-based in all other countries).	2,348 cases. 2,462 controls. Cases, 88%; controls, 81% hospital and 52% population.	In-person interviews using same structured questionnaire translated to the local language for information on sociodemographic factors, lifestyle, health history and all full-time job held $\geq 1$ yr. Assessment by industrial hygienists in each participating center to 43 agents, including TCE, by confidence, exposure intensity, and exposure frequency. Exposure metric of overall TCE exposure and cumulative TCE exposure for subjects assessed with high degree of confidence. Logistic regression adjusted for age, gender, education and study center.
German centers: Seidler et al. (2007); Mester et al. (2006); Becker et al. (2004)	NHL and Hodgkin's disease cases aged 18–80 yr identified through all hospitals and ambulatory physicians in six regions of Germany between 1998 and 2003; population controls were identified from population registers and matched on age, sex, and region.	710 cases. 710 controls. Cases, 87%; Controls, 44%.	n-person interview using questionnaire assessing personal characteristics, lifestyle, medical history, UV light exposure, and occupational history of all obs held for $\geq 1$ yr. Exposure of a prior interest were assessed using job ask-specific supplementary questionnaires. JEM used to assign cumulative quantitative TCE exposure metric, categorized according to the distribution among the control persons (50 <sup>th</sup> and 90 <sup>th</sup> percentile of the exposed controls). Conditional logistic regression adjusted for age, sex, region, smoking and alcohol consumption.

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Wang et al. (2009)	Cases among females aged 21 and 84 yr with NHL in 1996–2000 and identified from Connecticut Cancer Registry; population-based female controls (1) if <65 yr of age, having Connecticut address stratified by 5-yr age groups identified from random digit dialing or (2) ≥65 yr of age, by random selection from Centers for Medicare and Medicaid Service files.	601 cases. 717 controls. Cases, 72%; Controls, 69% (<65 yr), 47% (≥65 yr)	In-person interview using questionnaire assessment specific for jobs held for >1 yr. Intensity and probability of exposure to broad category of organic solvents and to individual solvents, including TCE, estimated using JEM (Dosemeci et al., 1994; Gomez et al., 1994) and assigned blinded. Exposure metric of any exposure, exposure intensity (low, medium/high), and exposure probability (low, medium/high) Logistic regression adjusted for age, family history of hematopoietic cancer, alcohol consumption and race.
Costantini et al. (2008); Miligi et al. (2006)	Cases aged 20–74 with NHL, including CLL, all forms of leukemia, or MM in 1991–1993 and identified through surveys of hospital and pathology departments in study areas and in specialized hematology centers in eight areas in Italy; population-based controls stratified by 5-yr age groups and by sex selected through random sampling of demographic or of National Health Service files.	1,428 NHL + CLL, 586 Leukemia, 263, MM. 1,278 controls (leukemia analysis). 1,100 controls (MM analysis). Cases, 83%; Controls, 73%.	In-person interview primarily at interviewee's home (not blinded) using questionnaire assessing specific jobs, extra occupational exposure to solvents and pesticides, residential history, and medical history. Occupational exposure assessed by job-specific or industry-specific questionnaires. JEM used to assign TCE exposure and assessed using intensity (2 categories) and exposure duration (2 categories). All NHL diagnoses and 20% sample of all cases confirmed by panel of three pathologists. Logistic regression with covariates for sex, age, region, and education. Logistic regression for specific NHL included an additional covariate for smoking.

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Persson and Fredriksson (1999); combined analysis of NHL cases in Persson et al. (1989; 1993)	Histologically confirmed cases of B-cell NHL, age 20–79 yr, identified in two hospitals in Sweden: Orebro in 1964–1986 (Persson et al., 1989) and in Linkoping between 1975–1984 (Persson et al., 1993); controls were identified from previous studies and were randomly selected from population registers.	NHL cases, 199. 479 controls. Cases, 96% (Oreboro), 90% (Linkoping); controls, not reported.	Mailed questionnaire to assess self reported occupational exposures to TCE and other solvents. Unadjusted Mantel-Haenszel chi-square.
Nordstrom et al. (1998)	Histologically-confirmed cases in males of hairy-cell leukemia reported to Swedish Cancer Registry in 1987–1992 (includes one case later identified with an incorrect diagnosis date); population-based controls identified from the National Population Registry and matched (1:4 ratio) to cases for age and county.	111 cases. 400 controls. Cases, 91%; Controls, 83%.	Mailed questionnaire to assess self reported working history, specific exposure, and leisure time activities. Univariate analysis for chemical-specific exposures (any TCE exposure).
Fritschi and Siemiatycki (1996b); Siemiatycki, (1991)	Male NHL cases, age 35–75 yr, diagnosed in 16 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	215 cases. 533 population controls (Group 1) and 1,900 subjects with other cancers (Group 2). Cases, 83%; Controls, 71%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (294 exposures on semiquantitative scales). Exposure metric defined as any or substantial exposure. Logistic regression adjusted for age, proxy status, income, and ethnicity (solvents) or Mantel-Haenszel stratified by age, body mass index, and cigarette smoking (TCE).

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Hardell et al. (1994; 1981)	Histologically-confirmed cases of NHL in males, age 25–85 yr, admitted to Swedish (Umea) hospital between 1974–1978; living controls (1:2 ratio) from the National Population Register, matched to living cases on sex, age, and place of residence; deceased controls from the National Registry for Causes of Death, matched (1:2 ratio) to dead cases on sex, age, place of residence, and yr of death.	105 cases. 335 controls . Response rate not available.	Self-administered questionnaire assessing self-reported solvent exposure; phone follow-up with subject, if necessary. Unadjusted Mantel-Haenszel chi-square.
Persson et al. (1989; 1993)	Histologically confirmed cases of Hodgkin's disease, age 20–80 yr, identified in two hospitals in Sweden: Oreboro in 1964–1986 (Persson et al., 1989) and in Linkoping between 1975–1984 (Persson et al., 1993); controls randomly selected from population registers.	54 cases (1989 study); 31 cases (1993 study). 275 controls (1989 study); 204 controls (1993 study). Response rate not available.	Mailed questionnaire to assess self reported occupational exposures to TCE and other solvents. Logistic regression with adjustment for age and other exposure; unadjusted Mantel-Haenszel chi-square.
<i>Childhood Leukemia</i>			
Shu et al. (2004; 1999)	Childhood leukemia cases, <15 yr, diagnosed between 1989 and 1993 by a Children's Cancer Group member or affiliated institute; population controls (random digit dialing), matched for age, race, and telephone area code and exchange.	1,842 cases. 1,986 controls. Cases, 92%; controls, 77%.	Telephone interview with mother, and whenever available, fathers using questionnaire to assess occupation using job-industry title and self-reported exposure history. Questionnaire included questions specific for solvent, degreaser or cleaning agent exposures. Logistic regression with adjustment for maternal or paternal education, race, and family income. Analyses of paternal exposure also included age and sex of the index child.

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Costas et al. (2002); MDPH (1997a, b)	Childhood leukemia (<19 yr age) diagnosed in 1969–1989 and who were resident of Woburn, MA; controls randomly selected from Woburn public School records, matched for age.	19 cases. 37 controls . Cases, 91%; Controls, not available.	Questionnaire administered to parents separately assessing demographic and lifestyle characteristics, medical history information, environmental and occupational exposure and use of public drinking water in the home. Hydraulic mixing model used to infer delivery of TCE and other solvents water to residence. Logistic regression with composite covariate, a weighted variable of individual covariates.
McKinney et al. (1991)	Incident childhood leukemia and non-Hodgkin lymphoma cases, 1974–1988, ages not identified, from three geographical areas in England; controls randomly selected from children of residents in the three areas and matched for sex and birth health district.	109 cases. 206 controls. Cases, 72%; Controls, 77%.	In-person interview with questionnaire with mother to assess maternal occupational exposure history, and with father and mother, as surrogate, to assess paternal occupational exposure history. No information provided in paper whether interviewer was blinded as to case and control status. Matched pair design using logistic regression for univariate and multivariate analysis.
Lowengart et al. (1987)	Childhood leukemia cases aged ≤10 yr and identified from the Los Angeles (CA) Cancer Surveillance Program in 1980–1984; controls selected from RDD or from friends of cases and matched on age, sex, and race.	123 cases. 123 controls. Cases, 79%; Controls, not available.	Telephone interview with questionnaire to assess parental occupational and self-reported exposure history. Matched (discordant) pair analysis.



**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
<i>Melanoma</i>			
Fritschi and Siemiatycki (1996b); Siemiatycki (1991)	Male melanoma cases, age 35–75 yr, diagnosed in 16 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	103 cases. 533 population controls and 533 other cancer controls. Cases, 78%; Controls, 72%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (294 exposures on semiquantitative scales); potential TCE exposure defined as any or substantial exposure. Logistic regression adjusted for age, education, and ethnic origin (TCE) or Mantel-Haenszel stratified on age, income, index for cigarette smoking, and ethnic origin (TCE).
<i>Pancreas</i>			
Kernan et al. (1999)	Pancreatic cancer deaths from 1984–1993 in 24 U.S. states; age-, sex-, race-, and state-matched noncancer deaths, excluding other pancreatic diseases and pancreatitis, controls.	63,097 cases. 252,386 population controls. Response rates not identified.	Exposure surrogate assigned for 111 chlorinated hydrocarbons, including TCE, and two broad chemical categories using usual occupation on death certificate and job-exposure-matrix of Gomez et al. (1994). Race and sex-specific mortality odds ratios from logistic regression analysis adjusted for age, marital status, metropolitan area, and residential status.
<i>Prostate</i>			
Aronson et al. (1996); Siemiatycki (1991)	Male prostate cancer cases, age 35–75 yr, diagnosed in 16 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	449 cases. 533 population controls (Group 1) and other cancer cases from same study (Group 2). Cases, 81%; Controls, 72%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (294 exposures on semiquantitative scales). Logistic regression adjusted for age, ethnic origin, socioeconomic status, Quetlet, and respondent status (occupation) or Mantel-Haenszel stratified on age, income, index for cigarette smoking, ethnic origin, and respondent status (TCE).

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
<i>Renal Cell</i>			
Moore et al. (2010)	Cases aged 20–74 yr from four European countries (Russia, Romania, Poland, Czech Republic) with histologically confirmed kidney cancer in 1999–2003; hospital controls with diagnoses unrelated to smoking or genitourinary disorders in 1998–2003 and frequency-matched by sex, age and study center.	1,097 cases (825 renal cell carcinomas). 1,184 controls. Cases, 90–99%; Controls, 90.3–96%.	In-person interview using questionnaire for information on lifestyle habits, smoking, anthropometric measures, personal and family medical history and occupational history. Specialized job-specific questionnaire for specific jobs or industries of interest focused on solvents exposure, including TCE, with exposure assignment by expert blinded to case and control status by frequency, intensity and confidence of TCE exposure. Exposure metric of overall exposure, duration (total h, yr) and cumulative exposure. Logistic regression adjusted for sex, age, and study center. BMI, hypertension, smoking, residence location also included in initial models but did not alter odds ratios by >10%.
Charbotel et al. (2006; 2009)	Cases from Arve Valley region in France identified from local urologists files and from area teaching hospitals; age- and sex-matched controls chosen from file of same urologist as who treated case or recruited among the patients of the case's general practitioner.	87 cases. 316 controls. Cases, 74%; controls, 78%.	Telephone interview with case or control, or, if deceased, with next-of-kin (22% cases, 2% controls). Questionnaire assessing occupational history, particularly, employment in the screw cutting jobs, and medical history. Semiquantitative TCE exposure assigned to subjects using a task/TCE-Exposure Matrix designed using information obtained from questionnaires and routine atmospheric monitoring of work shops or biological monitoring (U-TCA) of workers carried out since the 1960s. Cumulative exposure, cumulative exposure with peaks, and TWA. Conditional logistic regression with covariates for tobacco smoking and body mass index.
Brüning et al. (2003)	Histologically-confirmed cases 1992–2000 from German hospitals (Arnsberg); hospital controls (urology department) serving area, and local geriatric department, for older controls, matched by sex and age.	134 cases. 401 controls. Cases, 83%; Controls, not available.	In-person interviews with case or next-of-kin; questionnaire assessing occupational history using job title. Exposure metrics included longest job held, JEM of Pannett et al. (1985) to assign cumulative exposure to TCE and PCE, and exposure duration. Logistic regression with covariates for age, sex, and smoking.

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
Pesch et al. (2000b)	Histologically-confirmed cases from German hospitals (5 regions) in 1991–1995; controls randomly selected from residency registries matched on region, sex, and age.	935 cases. 4,298 controls. Cases, 88%; Controls, 71%.	In-person interview with case or next-of-kin; questionnaire assessing occupational history using job title (JEM approach), self-reported exposure, or job task (JTEM approach) to assign TCE and other exposures. Logistic regression with covariates for age, study center, and smoking.
Parent et al. (2000a); Siemiatycki (1991)	Male renal cell carcinoma cases, age 35–75 yr, diagnosed in 16 large Montreal-area hospitals in 1979–1985 and histologically confirmed; controls identified concurrently at 18 other cancer sites; age-matched, population-based controls identified from electoral lists and RDD.	142 cases. 533 population controls (Group 1) and other cancer controls (excluding lung and bladder cancers) (Group 2). Cases, 82%; Controls, 71%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (about 300 exposures on semiquantitative scales); TCE defined as any or substantial exposure. Mantel-Haenszel stratified by age, body mass index, and cigarette smoking (TCE) or logistic regression adjusted for respondent status, age, smoking, and body mass index (occupation, job title).
Dosemeci et al. (1999)	Histologically-confirmed cases, 1988–1990, white males and females, 20–85 yr, from Minnesota Cancer Registry; controls stratified for age and sex using RDD, 21–64 yr, or from HCFA records, 64–85 yr.	438 cases. 687 controls. Cases, 87%; Controls, 86%.	In-person interviews with case or next-of-kin; questionnaire assessing occupational history of TCE using job title and JEM of Gomez et al. (1994). Exposure metric was any TCE exposure. Logistic regression with covariates for age, smoking, hypertension, and body mass index.
Vamvakas et al. (1998)	Cases who underwent nephrectomy in 1987–1992 in a hospital in Arnsberg region of Germany; controls selected accident wards from nearby hospital in 1992.	58 cases. 84 controls. Cases, 83%; Controls, 75%.	In-person interview with case or next-of-kin; questionnaire assessing occupational history using job title or self-reported exposure to assign TCE and PCE exposure. Logistic regression with covariates for age, smoking, body mass index, hypertension, and diuretic intake.

**Table 4-2. Case-control epidemiologic studies examining cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Population</b>	<b>Study group (N) comparison group (N) response rates</b>	<b>Exposure assessment and other information</b>
<i>Multiple or Other Sites</i>			
Lee et al., (2003)	Liver, lung, stomach, colorectal cancer deaths in males and females between 1966–1997 from two villages in Taiwan; controls were cardiovascular and cerebral-vascular disease deaths from same underlying area as cases.	53 liver, 39 stomach, 26 colorectal, 41 lung cancer cases. 286 controls. Response rate not reported.	Residence as recorded on death certificate. Mantel-Haenszel stratified by age, sex, and time period.
Siemiatycki (1991)	Male cancer cases, 1979–1985, 35–75 yr, diagnosed in 16 Montreal-area hospitals, histologically confirmed; cancer controls identified concurrently; age-matched, population-based controls identified from electoral lists and RDD.	857 lung and 117 pancreatic cancer cases. 533 population controls (Group 1) and other cancer cases from same study (Group 2). Cases, 79% (lung), 71% (pancreas); Controls, 72%.	In-person interviews (direct or proxy) with segments on work histories (job titles and self-reported exposures); analyzed and coded by a team of chemists and industrial hygienists (294 exposures on semiquantitative scales); TCE defined as any or substantial exposure. Mantel-Haenszel stratified on age, income, index for cigarette smoking, ethnic origin, and respondent status (lung cancer) and age, income, index for cigarette smoking, and respondent status (pancreatic cancer).

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

BMI = body mass index, CLL = chronic lymphocytic leukemia, HCFA = Health Care Financing Administration, JEM = job-exposure matrix, JTEM = job-task-exposure matrix, MM = multiple myeloma, NCI = National Cancer Institute, NHL = non-Hodgkin lymphoma, PCE = perchloroethylene, RDD = random digit dialing, TWA = time-weighted average, U-TCA = urinary trichloroacetic acid, UV = ultra-violet.

**Table 4-3. Geographic-based studies assessing cancer and TCE exposure**

Reference	Description	Analysis approach	Exposure assessment
<i>Broome County, NY Studies</i>			
ATSDR (2006a) (2008)	Total, 22 site-specific, and childhood cancer incidence from 1980–2001 among residents in two areas in Endicott, NY.	SIR among all subjects (ATSDR, 2006a) or among white subjects only (ATSDR, 2008) with expected numbers of cancers derived using age-specific cancer incidence rates for New York State, excluding New York City. Limited assessment of smoking and occupation using medical and other records in lung and kidney cancer subjects (ATSDR, 2008).	Two study areas, Eastern and Western study areas, identified based on potential for soil vapor intrusion exposures as defined by the extent of likely soil vapor contamination. Contour lines of modeled VOC soil vapor contamination levels based on exposure model using GIS mapping and soil vapor sampling results taken in 2003. The study areas were defined by 2000 Census block boundaries to conform to model predicted areas of soil vapor contamination. TCE was the most commonly found contaminant in indoor air in Eastern study area at levels ranging from 0.18–140 $\mu\text{g}/\text{m}^3$ , with tetrachloroethylene, cis-1,2-dichloroethene, 1,1,1-trichloroethane, 1,1-dichloroethylene, 1,1-dichloroethane, and Freon 113 detected at lower levels. PCE was most common contaminant in indoor air in Western study area with other VOCs detected at lower levels.
<i>Maricopa County, AZ Studies</i>			
Aickin et al. (1992); Aickin (2004)	Cancer deaths, including leukemia, 1966–1986, and childhood ( $\leq 19$ yr old) leukemia incident cases (1965–1986), Maricopa County, AZ.	Standardized mortality rate ratio from Poisson regression modeling. Childhood leukemia incidence data evaluated using Bayes methods and Poisson regression modeling.	Location of residency in Maricopa County, AZ, at the time of death as surrogate for exposure. Some analyses examined residency in West Central Phoenix and cancer. Exposure information is limited to TCE concentration in two drinking water wells in 1982.
<i>Pima County, AZ Studies</i>			
ADHS (1990, 1995)	Cancer incidence in children ( $\leq 19$ yr old) and testicular cancer in 1970–1986 and 1987–1991, Pima County, AZ.	Standardized incidence RR from Poisson regression modeling using method of Aickin et al. (Aickin et al., 1992). Analysis compares incidence in Tucson Airport Area to rate for rest of Pima County.	Location of residency in Pima, County, AZ, at the time of diagnosis or death as surrogate for exposure. Exposure information is limited to monitoring since 1981 and include VOCs in soil gas samples (TCE, PCE, 1,1-dichloroethylene, 1,1,1-trichloroacetic acid); PCBs in soil samples, and TCE in municipal water supply wells.

**Table 4-3. Geographic-based studies assessing cancer and TCE exposure (continued)**

Reference	Description	Analysis approach	Exposure assessment
<i>Other</i>			
Coyle et al. (2005)	Incident breast cancer cases among men and women, 1995–2000, reported to Texas Cancer Registry.	Correlation study using rank order statistics of mean average annual breast cancer rate among women and men and atmospheric release of 12 hazardous air pollutants.	Reporting to EPA Toxic Release Inventory the number of pounds released for 12 hazardous air pollutants, (carbon tetrachloride, formaldehyde, methylene chloride, styrene, tetrachloroethylene, trichloroethylene, arsenic, cadmium, chromium, cobalt, copper, and nickel).
Morgan and Cassady (2002)	Incident cancer cases, 1988–1989, among residents of 13 census tracts in Redlands area, San Bernardino County, CA.	SIR for all cancer sites and 16 site-specific cancers; expected numbers using incidence rates of site-specific cancer of a four-county region between 1988–1992.	TCE and perchlorate detected in some county wells; no information on location of wells to residents, distribution of contaminated water, or TCE exposure potential to individual residents in studied census tracts.
Vartiainen et al. (1993)	Total cancer and site-specific cancer cases (lymphoma sites and liver) from 1953–1991 in two Finnish municipalities.	SIR with expected number of cancers and site-specific cancers derived from incidence of the Finnish population.	Monitoring data from 1992 indicated presence of TCE, tetrachloroethylene and 1,1,1,-trichloroethane in drinking water supplies in largest towns in municipalities. Residence in town used to infer exposure to TCE.
Cohn et al. (1994b); Fagliano et al. (1990)	Incident leukemia and NHL cases, 1979–1987, from 75 municipalities and identified from the New Jersey State Cancer Registry. Histological type classified using WHO scheme and the classification of NIH Working Formulation Group for grading NHL.	Logistic regression modeling adjusted for age.	Monitoring data from 1984–1985 on TCE, THM, and VOCs concentrations in public water supplies, and historical monitoring data conducted in 1978–1984.
Mallin (1990)	Incident bladder cancer cases and deaths, 1978–1985, among residents of nine NW Illinois counties.	SIR and SMR by county of residence and zip code; expected numbers of bladder cancers using age-race-sex specific incidence rates from SEER or bladder cancer mortality rates of the U.S. population from 1978–1985.	Exposure data are lacking for the study population with the exception of noting one of two zip code areas with observed elevated bladder cancer rates also had groundwater supplies contaminated with TCE, PCE and other solvents.

**Table 4-3. Geographic-based studies assessing cancer and TCE exposure (continued)**

<b>Reference</b>	<b>Description</b>	<b>Analysis approach</b>	<b>Exposure assessment</b>
Isacson et al. (1985)	Incident bladder, breast, prostate, colon, lung and rectal cancer cases reported to Iowa cancer registry between 1969–1981.	Age-adjusted site-specific cancer incidence in Iowa towns with populations of 1,000–10,000 and who were serviced by a public drinking water supply.	Monitoring data of drinking water at treatment plant in each Iowa municipality with populations of 1,000–10,000 used to infer TCE and other volatile organic compound concentrations in finished drinking water supplies.

EPA = U.S. Environmental Protection Agency, GIS = geographic information system, NHL = non-Hodgkin lymphoma, NIH = National Institutes of Health, NW = Northwestern, PCB = polychlorinated biphenyl, PCE = perchloroethylene, -, SEER = Surveillance, Epidemiology, and End Results, SIR = standardized incidence ratio, SMR = standardized mortality ratio, VOCs = volatile organic compounds, WHO = World Health Organization.

**Table 4-4. Standards of epidemiologic study design and analysis use for identifying cancer hazard and TCE exposure**

<p><b>Category A: Study Design</b></p>
<p><b>Clear articulation of study objectives or hypothesis.</b> The ideal is a clearly stated hypothesis or study objectives and the study is designed to achieve the identified objectives.</p>
<p><b>Selection and characterization in cohort studies of exposure and control groups and of cases and controls (case-control studies) is adequate.</b> The ideal is for selection of cohort and referents from the same underlying population and differences between these groups to be due to TCE exposure or level of TCE exposure and not to physiological, health status, or lifestyle factors. Controls or referents are assumed to lack or to have background exposure to TCE. These factors may lead to a downward bias including one of which is known as “healthy worker bias,” often introduced in analyses when mortality or incidence rates from a large population such as the U.S. population are used to derive expected numbers of events. The ideal in case-control studies is cases and controls are derived from the same population and are representative of all cases and controls in that population. Any differences between controls and cases are due to exposure to TCE itself and not to confounding factors related to both TCE exposure and disease. Additionally, the ideal is for controls to be free of any disease related to TCE exposure. In this latter case, potential bias is toward the null hypothesis.</p>
<p><b>Category B: Endpoint Measured</b></p>
<p><b>Levels of health outcome assessed.</b> Three levels of health outcomes are considered in assessing the human health risks associated with exposure to TCE: biomarkers of effects and susceptibility, morbidity, and mortality. Both morbidity as enumerated by incidence and mortality as identified from death certificates are useful indicators in risk assessment for hazard identification. The ideal is for accurate and predictive indicator of disease. Incidence rates are generally considered to provide an accurate indication of disease in a population and cancer incidence is generally enumerated with a high degree of accuracy in cancer registries. Death certifications are readily available and have complete national coverage but diagnostic accuracy is reduced and can vary by specific diagnosis. Furthermore, diagnostic inaccuracies can contribute to death certificates as a poor surrogate for disease incidence. Incidence, when obtained from population-based cancer registries, is preferred for identifying cancer hazards.</p>
<p><b>Changes in diagnostic coding systems for lymphoma, particularly non-Hodgkin lymphoma.</b> Classification of lymphomas today is based on morphologic, immunophenotypic, genotypic, and clinical features using the World Health Organization (WHO) classification, introduced in 2001, and incorporation of WHO terminology into International Classification of Disease (ICD)-0-3. ICD Versions 7 and earlier had rubrics for general types of lymphatic and hematopoietic cancer, but no categories for distinguishing specific types of cancers, such as acute leukemia. Epidemiologic studies based on causes of deaths as coded using these older ICD classifications typically grouped together lymphatic neoplasms instead of examining individual types of cancer or specific cell types. Before the use of immunophenotyping, these grouping of ambiguous diseases such as non-Hodgkin lymphoma and Hodgkin’s lymphoma may be have misclassified. With the introduction of ICD-10 in 1990, lymphatic tumors coding, starting in 1994 with the introduction of the Revised European-American Lymphoma classification, the basis of the current WHO classification, was more similar to that presently used. Misclassification of specific types of cancer, if unrelated to exposure, would have attenuated estimate of relative risk and reduced statistical power to detect associations. When the outcome was mortality, rather than incidence, misclassification would be greater because of the errors in the coding of underlying causes of death on death certificates (IOM, 2003). Older studies that combined all lymphatic and hematopoietic neoplasms must be interpreted with care.</p>



**Table 4-4. Standards of epidemiologic study design and analysis use for identifying cancer hazard and TCE exposure (continued)**

<p><b>Category C: TCE-Exposure Criteria</b></p> <p><b>Adequate characterization of exposure.</b> The ideal is for TCE exposure potential known for each subject and quantitative assessment [job-exposure-matrix approach] of TCE exposure assessment for each subject as a function of job title, yr exposed, duration, and intensity. The assessment approach is accurate for assigning TCE intensity [TCE concentration or a time-weighted-average] to individual study subjects and estimates of TCE intensity are validated using monitoring data from the time period. For the purpose of this report, the objective for cohort and case-controls studies is to differentiate TCE-exposed subjects from subjects with little or no TCE exposure. A variety of dose-metrics may be used to quantify or classify exposures for an epidemiologic study. They include precise summaries of quantitative exposure, concentrations of biomarkers, cumulative exposure, and simple qualitative assessments of whether exposure occurred (yes or no). Each method has implicit assumptions and potential problems that may lead to misclassification. Studies in which it was unclear that the study population was actually exposed to TCE are excluded from analysis.</p>
<p><b>Category D: Follow-up (Cohort)</b></p> <p>Loss to follow-up. The ideal is complete follow-up of all subjects; however, this is not achievable in practice, but it seems reasonable to expect loss to follow-up not to exceed 10%. The bias from loss to follow-up is indeterminate. Random loss may have less effect than if subjects who are not followed have some significant characteristics in common.</p> <p>Follow-up period allows full latency period for over 50% of the cohort. The ideal to follow all study subjects until death. Short of the ideal, a sufficient follow-up period to allow for cancer induction period or latency over 15 or 20 yr is desired for a large percentage of cohort subjects.</p>
<p><b>Category E: Interview Type (Case-control)</b></p> <p><b>Interview approach.</b> The ideal interviewing technique is face-to-face by trained interviewers with more than 90% of interviews with cases and control subjects conducted face-to-face. The effect on the quality of information from other types of data collection is unclear, but telephone interviews and mail-in questionnaires probably increase the rate of misclassification of subject information. The bias is toward the null hypothesis if the proportion of interview by type is the same for case and control, and of indeterminate direction otherwise.</p> <p><b>Blinded interviewer.</b> The ideal is for the interviewer to be unaware whether the subject is among the cases or controls and the subject to be unaware of the purpose and intended use of the information collected. Although desirable for case-control studies, blinding is usually not possible to fully accomplish because subject responses during the interview provide clues as to subject status. The potential for bias from face-to-face interviews is probably less than with mail-in interviews. Some studies have assigned exposure status in a blinded manner using a job-exposure matrix and information collected in the unblinded interview. The potential for bias in this situation is probably less with this approach than for nonblinded assignment of exposure status.</p>

**Table 4-4. Standards of epidemiologic study design and analysis use for identifying cancer hazard and TCE exposure (continued)**

<p><b>Category F: Proxy Respondents</b></p> <p><b>Proxy respondents.</b> The ideal is for data to be supplied by the subject because the subject generally would be expected to be the most reliable source; less than 10% of either total cases or total controls for case-control studies. A subject may be either deceased or too ill to participate, however, making the use of proxy responses unavoidable if those subjects are to be included in the study. The direction and magnitude of bias from use of proxies is unclear, and may be inconsistent across studies.</p>
<p><b>Category G: Sample Size</b></p> <p>The ideal is for the sample size is large enough to provide sufficient statistical power to ensure that any elevation of effect in the exposure group, if present, would be found, and to ensure that the confidence bounds placed on relative risk estimates can be well characterized.</p>
<p><b>Category H: Analysis Issues</b></p> <p><b>Control for potentially confounding factors of importance in analysis.</b> The ideal in cohort studies is to derive expected numbers of cases based on age-sex- and time-specific cancer rates in the referent population and in case-control studies by matching on age and sex in the design and then adjusting for age in the analysis of data. Age and sex are likely correlated with exposure and are also risk factors for cancer development. Similarly, other factors such as cigarette smoking and alcohol consumption are risk factors for several site-specific cancers reported as associative with TCE exposure. To be a confounder of TCE, exposure to the other factor must be correlated, and the association of the factor with the site-specific cancer must be causal. The expect effect from controlling for confounders is to move the estimated relative risk estimate closer to the true value.</p> <p><b>Statistical methods are appropriate.</b> The ideal is that conclusions are drawn from the application of statistical methods that are appropriate to the problem and accurately interpreted.</p> <p><b>Evaluation of exposure-response.</b> The ideal is an examination of a linear exposure-response as assessed with a quantitative exposure metric such as cumulative exposure. Some studies, absent quantitative exposure metrics, examine exposure response relationships using a semiquantitative exposure metric or by duration of exposure. A positive dose-response relationship is usually more convincing of an association as causal than a simple excess of disease using TCE dose-metric. However, a number of reasons have been identified for a lack of linear exposure-response finding and the failure to find such a relationship mean little from an etiological viewpoint.</p> <p><b>Documentation of results.</b> The ideal is for analysis observations to be completely and clearly documented and discussed in the published paper, or provided in supplementary materials accompanying publication.</p>

1 Twenty-six of the studies identified in a systematic review were selected for inclusion in  
2 the meta-analysis through use of the following meta-analysis inclusion criteria: (1) cohort or  
3 case-control designs; (2) evaluation of incidence or mortality; (3) adequate selection in cohort  
4 studies of exposure and control groups and of cases and controls in case-control studies; (4) TCE  
5 exposure potential inferred to each subject and quantitative assessment of TCE exposure  
6 assessment for each subject by reference to industrial hygiene records indicating a high  
7 probability of TCE use, individual biomarkers, job exposure matrices, water distribution models,  
8 or obtained from subjects using questionnaire (case-control studies); and (5) relative risk  
9 estimates for kidney cancer, liver cancer, or non-Hodgkin lymphoma (NHL) adjusted, at  
10 minimum, for possible confounding of age, sex, and race (see Table 4-5). This evaluation is  
11 summarized below, separately for cohort and case-control studies. Appendix C contains a full  
12 discussion of the meta-analysis, its analytical methodology, including sensitivity analyses, and  
13 findings. The meta-analysis focuses on kidney cancer, liver cancer, and NHL, as most studies  
14 reported relative risks for these sites. Fewer numbers of studies reported relative risks for other  
15 site-specific cancers and TCE exposure and examination of these site-specific cancers and TCE  
16 exposure using meta-analysis was not attempted.

17 The cohort studies (Anttila et al., 1995; ATSDR, 2004; Axelson et al., 1994; Blair et al.,  
18 1989; Blair et al., 1998; Boice et al., 1999; Boice et al., 2006b; Chang et al., 2003; Chang et al.,  
19 2005; Clapp and Hoffman, 2008; Costa et al., 1989; Garabrant et al., 1988; Greenland et al.,  
20 1994; Hansen et al., 2001; Henschler et al., 1995; Krishnadasan et al., 2007; Morgan et al., 1998;  
21 Raaschou-Nielsen et al., 2003; Radican et al., 2008; Ritz, 1999a; Shannon et al., 1988; Shindell  
22 and Ulrich, 1985; Sinks et al., 1992; Sung et al., 2007; Sung et al., 2008; Wilcosky et al., 1984;  
23 Zhao et al., 2005) (see Table 4-1), with data on the incidence or morality of site-specific cancer  
24 in relation to TCE exposure, range in size 803 (Hansen et al., 2001) to 86,868 (Chang et al.,  
25 2003; Chang et al., 2005), and were conducted in Denmark, Sweden, Finland, Germany, Taiwan,  
26 and the United States (see Table 4-1). Three case-control studies nested within cohorts  
27 (Greenland et al., 1994; Krishnadasan et al., 2007; Wilcosky et al., 1984) are considered as  
28 cohort studies because the summary risk estimate from a nested case-control study, the odds  
29 ratio, was estimated from incidence density sampling. This is considered an unbiased estimate of  
30 the hazard ratio, similar to a relative risk estimate from a cohort study, if, as is the case for these  
31 studies, controls are selected from the same source population as the cases, the sampling rate is  
32 independent of exposure status, and the selection probability is proportional to time-at-risk  
33 (Rothman et al., 2008). Cohort and nested case-control study designs are analytical  
34 epidemiologic studies and are generally relied on for identifying a causal association between  
35 human exposure and adverse health effects (U.S. EPA, 2005c).

36

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-5. Summary of criteria for meta-analysis study selection**

Decision outcome	Studies	Primary reason(s)
Studies recommended for meta-analysis:		
	Axelson et al. (1994); Greenland et al. (1994); Hardell et al. (1994); Siemiatycki (1991); Anttila et al. (1995); Morgan et al. (1998); Nordstrom et al. (1998); Boice et al. (1999); Boice et al. (2006b); Dosemeci et al. (1999); Persson and Fredriksson (1999); Pesch et al. (2000b); Hansen et al. (2001); Brüning et al. (2003); Raaschou-Nielsen et al. (2003); Zhao et al. (2005); Miligi et al. (2006); Seidler et al. (2007); Charbotel et al. (2006; 2009); Blair et al. (1998); its follow-up Radican et al. (2008); Wang et al. (2009); Cocco et al. (2010); Moore et al. (2010); Purdue et al. (2011)	Analytical study designs of cohort or case-control; evaluation of incidence or mortality; adequate selection in cohort studies of exposure and control groups and of cases and controls in case-control studies; TCE exposure potential inferred to each subject and quantitative assessment of TCE exposure assessment for each subject by reference to industrial hygiene records indicating a high probability of TCE use, individual biomarkers, job exposure matrices, water distribution models, or obtained from subjects using questionnaire (case-control studies); relative risk estimates for kidney cancer, liver cancer, or NHL adjusted, at minimum, for possible confounding of relevant risk factors, e.g., age, sex, and race).
Studies not recommended for meta-analysis:		
	Clapp and Hoffman (2008); ATSDR (2004; Cohn et al., 1994b)	Weakness with respect to analytical study design (i.e., geographic-based, ecological or proportional mortality ratio design).
	Garabrant et al.(1988); Isacson et al.(1985); Shindell and Ulrich (1985); Wilcosky et al. (1984); Shannon et al. (1988); Blair et al. (1989); Costa et al. (1989); (ADHS, 1990, 1995); Mallin (1990); Aickin et al. (1992); Sinks et al. (1992); Vartiainen et al. (1993); Morgan and Cassady (2002); Lee et al. (2003); Aickin (2004); Chang et al. (Chang et al., 2003; 2005); Coyle et al. (2005); ATSDR (2006a); ATSDR (2008); Sung et al. (2007; 2008)	TCE exposure potential not assigned to individual subjects using job exposure matrix, individual biomarkers, water distribution models, or industrial hygiene data from other process indicating a high probability of TCE use (cohort studies).
	Lowengart et al. (1987); Fredriksson et al. (1989); McKinney et al. (1991); Heineman et al. (1994); Siemiatycki et al. (1994); Aronson et al. (1996); Fritschi and Siemiatycki (1996b); Dumas et al. (2000); Kernan et al. (1999); Shu et al. (2004; 1999); Parent et al. (2000b); Pesch et al. (2000a); DeRoos et al. (2001a); Goldberg et al. (2001); Costas et al. (2002); Krishnadasan et al. (2007) Costantini et al. (2008); Gold et al.	Cancer incidence or mortality reported for cancers other than kidney, liver, or NHL.
	Ritz (1999a)	Subjects monitored for radiation exposure with likelihood for potential confounding; cancer mortality and TCE exposure not reported for kidney cancer and all hemato- and lymphopoietic cancer reported as broad category.

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-5. Summary of criteria for meta-analysis study selection (continued)**

Decision outcome	Studies	Primary reason(s)
Studies not recommended for meta-analysis: (continued)		
	Henschler et al. (1995)	Incomplete identification of cohort and index kidney cancer cases included in case series.

1  
2 NHL = non-Hodgkin lymphoma.

3  
4  
5 While all of these cohort studies are considered in the overall weight of evidence, eleven  
6 of them met all meta-analysis inclusion criteria: the cohorts of Blair et al. (1998) and its  
7 follow-up by Radican et al. (2008); Morgan et al. (1998), Boice et al. (1999; Boice et al., 2006b),  
8 and Zhao et al.(2005), of aerospace workers or aircraft mechanics; and Axelson et al. (1994),  
9 Anttila et al. (1995), Hansen et al. (2001), and Raaschou-Nielsen et al. (2003) of Nordic workers  
10 in multiple industries with TCE exposure; and Greenland et al. (1994) of electrical  
11 manufacturing workers. Subjects or cases and controls in these studies are considered to  
12 sufficiently represent the underlying population, and the bias associated with selection of referent  
13 populations is considered minimal. The exposure-assessment approaches included detailed  
14 job-exposure matrix, biomonitoring data, or use of industrial hygiene data on TCE exposure  
15 patterns and factors that affect such exposure, with high probability of TCE exposure potential to  
16 individual subjects. The statistical analyses methods were appropriate and well documented, the  
17 measured endpoint was an accurate indicator of disease, and the follow-up was sufficient for  
18 cancer latency. These studies are also considered as strong studies for identifying kidney, liver  
19 and NHL cancer hazard. The remaining cohort studies less satisfactorily meet identified criteria  
20 or standards of epidemiologic design and analysis, having deficiencies in multiple criteria  
21 (ATSDR, 2004; Chang et al., 2003; Chang et al., 2005; Clapp and Hoffman, 2008; Costa et al.,  
22 1989; Garabrant et al., 1988; Henschler et al., 1995; Ritz, 1999a; Shindell and Ulrich, 1985;  
23 Sinks et al., 1992; Sung et al., 2007; Sung et al., 2008; Wilcosky et al., 1984). Krishnandansen  
24 et al. (2007), who reported on prostate cancer, met four of the five meta-analysis inclusion  
25 criteria except that for reporting a relative risk estimate cancer of the kidney, liver or NHL, the  
26 site-specific cancers examined using meta-analysis.

27 The case-control studies on TCE exposure are of several site-specific cancers, including  
28 bladder (Pesch et al., 2000a; Siemiatycki, 1991; Siemiatycki et al., 1994); brain (De Roos et al.,  
29 2001b; Heineman et al., 1994); childhood lymphoma or leukemia (Costas et al., 2002;  
30 Lowengart et al., 1987; McKinney et al., 1991; Shu et al., 2004; Shu et al., 1999); colon cancer  
31 (Goldberg et al., 2001; Siemiatycki, 1991); esophageal cancer (Parent et al., 2000b; Siemiatycki,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 1991); liver cancer (Lee et al., 2003); lung (Siemiatycki, 1991); adult lymphoma or leukemia  
2 (Hardell et al., 1994) [non-Hodgkin lymphoma (NHL), Hodgkin lymphoma]; (Fritschi and  
3 Siemiatycki, 1996a; Siemiatycki, 1991) [NHL]; (Nordström et al., 1998) [hairy cell leukemia];  
4 (Persson and Fredrikson, 1999) [NHL]; (Miligi et al., 2006) [NHL and chronic lymphocytic  
5 leukemia (CLL)]; (Seidler et al., 2007) [NHL, Hodgkin lymphoma and subjects included in  
6 (Cocco et al., 2010; Costantini et al., 2008) [leukemia types, CLL included with NHL in (Miligi  
7 et al., 2006; Wang et al., 2009) [NHL]; (Cocco et al., 2010) [B-cell including CLL and multiple  
8 myeloma, T-cell and Hodgkin lymphomas]; (Purdue et al., 2011) [NHL]; Gold et al. [multiple  
9 myeloma]); melanoma (Fritschi and Siemiatycki, 1996b; Siemiatycki, 1991); rectal cancer  
10 (Dumas et al., 2000; Siemiatycki, 1991); renal cell carcinoma, a form of kidney cancer (Brüning  
11 et al., 2003; Charbotel et al., 2006; Dosemeci et al., 1999; Moore et al., 2010; Parent et al.,  
12 2000a; Pesch et al., 2000b; Siemiatycki, 1991; Vamvakas et al., 1998); pancreatic cancer  
13 (Siemiatycki, 1991), and prostate cancer (Aronson et al., 1996; Siemiatycki, 1991) (see  
14 Table 4-2). No case-control studies of reproductive cancers (breast or cervix) and TCE exposure  
15 were found in the peer-reviewed literature.

16 While all of these case-control studies are considered in the overall weight of evidence,  
17 fifteen of them met the meta-analysis inclusion criteria identified in Section B.2.9 (Brüning et al.,  
18 2003; Charbotel et al., 2006; Charbotel et al., 2009; Cocco et al., 2010; Dosemeci et al., 1999;  
19 Hardell et al., 1994; Miligi et al., 2006; Moore et al., 2010; Nordström et al., 1998; Persson and  
20 Fredrikson, 1999; Pesch et al., 2000b; Purdue et al., 2011; Seidler et al., 2007; Siemiatycki,  
21 1991; Wang et al., 2009). They were of analytical study design, cases and controls were  
22 considered to represent underlying populations and selected with minimal potential for bias;  
23 exposure assessment approaches included assignment of TCE exposure potential to individual  
24 subjects using information obtained from face-to-face, mailed, or telephone interviews; analyses  
25 methods were appropriate, well-documented, included adjustment for potential confounding  
26 exposures, with relative risk estimates and associated confidence intervals reported for kidney  
27 cancer, liver cancer or NHL.

28 These studies were also considered, to varying degrees, as strong studies for weight-of  
29 evidence characterization of hazard. Both Brüning et al. (2003) and Charbotel et al. (2006;  
30 2009) had a priori hypotheses for examining renal cell carcinoma and TCE exposure. Strengths  
31 of both studies are in their examination of populations with potential for high exposure intensity  
32 and in areas with high frequency of TCE usage and their assessment of TCE potential. An  
33 important feature of the exposure assessment approach of Charbotel et al. (2006) is their use of a  
34 large number of studies on biological monitoring of workers in the screw-cutting industry a  
35 predominant industry with documented TCE exposures as support. Other studies were either

1 large multiple-center studies (Cocco et al., 2010; Miligi et al., 2006; Moore et al., 2010; Pesch et  
2 al., 2000a; Pesch et al., 2000b; Purdue et al., 2011; Wang et al., 2009) or reporting from one  
3 location of a larger international study (Dosemeci et al., 1999; Seidler et al., 2007). Cocco et al.  
4 (2010) includes subjects in Seidler et al. (2007) and is preferred because of the larger number of  
5 subjects from four other European countries. In contrast to Brüning et al. (2003) and Charbotel  
6 et al. (2006; 2009), two studies conducted in geographical areas with widespread TCE usage and  
7 potential for exposure to higher intensity, in these other studies, a lower exposure prevalence to  
8 TCE is found (any TCE exposure: 15% of cases (Dosemeci et al., 1999); 6% of cases (Miligi et  
9 al., 2006); 13% of cases (Wang et al., 2009); 4% of cases (Cocco et al., 2010) and most subjects  
10 were identified as exposed to TCE probably had minimal contact (3% of cases with  
11 moderate/high TCE exposure (Miligi et al., 2006); 2% of cases with high intensity, but of low  
12 probability TCE exposure (Wang et al., 2009). This pattern of lower exposure prevalence and  
13 intensity is common to community-based population case-control studies (Teschke et al., 2002).

14 Fourteen case-control studies did not meet specific meta-analysis inclusion criterion  
15 (Aronson et al., 1996; Costas et al., 2002; Dumas et al., 2000; Fritschi and Siemiatycki, 1996b;  
16 Gold et al., In Press; Goldberg et al., 2001; Kernan et al., 1999; Lee et al., 2003; Parent et al.,  
17 2000b; Pesch et al., 2000a; Shu et al., 2004; Shu et al., 1999; Siemiatycki et al., 1994; Vamvakas  
18 et al., 1998). Twelve studies reported relative risk estimates for site-specific cancers other than  
19 kidney, liver, and NHL (Aronson et al., 1996; Costas et al., 2002; Dumas et al., 2000; Fritschi  
20 and Siemiatycki, 1996b; Gold et al., In Press; Goldberg et al., 2001; Kernan et al., 1999; Parent  
21 et al., 2000b; Pesch et al., 2000a; Shu et al., 2004; Shu et al., 1999; Siemiatycki et al., 1994).  
22 Vamvakas et al. (1998) has been subject of considerable controversy (Bloemen and Tomenson,  
23 1995; Cherrie et al., 2001; Green and Lash, 1999; Mandel, 2001; McLaughlin and Blot, 1997;  
24 Swaen, 1995) with questions raised on potential for selection bias related to the study's controls.  
25 This study was deficient in the criterion for adequacy of case and control selection. Brüning  
26 et al. (2003), a study from the same region as Vamvakas et al. (1998), is considered a stronger  
27 study for identifying cancer hazard since it addresses many of the deficiencies of Vamvakas et al.  
28 (1998). Lee et al. (2003) in their study of hepatocellular cancer assigns one level of exposure to  
29 all subjects in a geographic area, and inherent measurement error and misclassification bias  
30 because not all subjects are exposed uniformly. Additionally, statistical analyses in this study  
31 did not control for hepatitis viral infection, a known risk factor for hepatocellular cancer and of  
32 high prevalence in the study area.

33 The geographic-based studies (ADHS, 1990, 1995; Aickin, 2004; Aickin et al., 1992;  
34 ATSDR, 2006a, 2008; Cohn et al., 1994b; Isacson et al., 1985; Mallin, 1990; Morgan and  
35 Cassady, 2002; Vartiainen et al., 1993) with data on cancer incidence are correlation studies to

1 examine cancer outcomes of residents in communities with TCE and other chemicals detected in  
2 groundwater wells or in municipal drinking water supplies (see Table 4-3). These studies did not  
3 meet all five meta-analysis inclusion criteria. The geographic-base studies are not of analytical  
4 designs such as cohort and case-control designs. Another deficiency in all studies is their low  
5 level of detail to individual subjects for TCE. One level of exposure to all subjects in a  
6 geographic area is assigned without consideration of water distribution networks, which may  
7 influence TCE concentrations delivered to a home, or a subject's ingestion rate to estimate TCE  
8 exposure to individual study subjects. Some inherent measurement error and misclassification  
9 bias is likely in these studies because not all subjects are exposed uniformly. Additionally, in  
10 contrast to case-control studies, the geographic-based studies, including the Agency for Toxic  
11 Substances and Disease Registry (ATSDR, 2008), had limited accounting for other potential risk  
12 factors. These studies are of low sensitivity for weight-of-evidence characterization of hazard  
13 compared to other cohort and case-control studies.  
14

#### 4.2. GENETIC TOXICITY

15 This section discusses the genotoxic potential of TCE and its metabolites. A summary is  
16 provided at the end of each section for TCE or its metabolite for their mutagenic potential in  
17 addition to an overall synthesis summary at the end of the genotoxicity section. The liver and  
18 kidney are subjects of study for the genotoxic potential of TCE and its metabolites, and are  
19 discussed more in-depth in sections 4.4.3, 4.4.7, 4.5.6.2.7, 4.5.7, E.2.3, and E.2.4.

20 The application of genotoxicity data to predict potential carcinogenicity is based on the  
21 principle that genetic alterations are found in all cancers. Genotoxicity is the ability of chemicals  
22 to alter the genetic material in a manner that permits changes to be transmitted during cell  
23 division. Although most tests for mutagenicity detect changes in DNA or chromosomes, some  
24 specific modifications of the epigenome including proteins associated with DNA or RNA, can  
25 also cause transmissible changes. Changes that occur due to the modifications in the epigenome  
26 are discussed in endpoint-specific Sections 4.3–4.9 as well as Sections E.3.1–E.3.4.

27 Genetic alterations can occur through a variety of mechanisms including gene mutations,  
28 insertions, deletions, translocations, or amplification; evidence of mutagenesis provides  
29 mechanistic support for the inference of potential for carcinogenicity in humans.

30 Evaluation of genotoxicity data entails a weight-of-evidence approach that includes  
31 consideration of the various types of genetic damage that can occur. In acknowledging that  
32 genotoxicity tests are by design complementary evaluations of different mechanisms of  
33 genotoxicity, a recent International Programme on Chemical Safety (IPCS) publication

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 (Eastmond et al., 2009) notes that “multiple negative results may not be sufficient to remove  
2 concern for mutagenicity raised by a clear positive result in a single mutagenicity assay.” These  
3 considerations inform the present approach. In addition, consistent with U.S. Environmental  
4 Protection Agency’s (EPA) *Guidelines on Carcinogenic Risk Assessment and Supplemental*  
5 *Guidance for Assessing Susceptibility from Early-Life Exposure to Carcinogens* (U.S. EPA,  
6 2005b, d), the approach does not address relative potency (e.g., among TCE metabolites, or of  
7 such metabolites with other known genotoxic carcinogens) *per se*, nor does it consider  
8 quantitative issues related to the probable production of these metabolites in vivo. Instead, the  
9 analysis of genetic toxicity data presented here focuses on the identification of a genotoxic  
10 hazard of these metabolites; a quantitative analysis of TCE metabolism to reactive intermediates,  
11 via physiologically based pharmacokinetic (PBPK) modeling, is presented in Section 3.5.

12 TCE and its known metabolites trichloroacetic acid (TCA), dichloroacetic acid (DCA),  
13 chloral hydrate (CH), trichloroethanol (TCOH), S-(1,2-dichlorovinyl)-L-cysteine (DCVC) and  
14 S-dichlorovinyl glutathione (DCVG) have been studied to varying degrees for their genotoxic  
15 potential. The following section summarizes available data on genotoxicity for both TCE and its  
16 metabolites for each potential genotoxic endpoints, when available, in different organisms.

17

#### 4.2.1. Trichloroethylene (TCE)

##### 4.2.1.1.1. DNA Binding Studies

18 Covalent binding of TCE to DNA and protein in cell-free systems has been studied by  
19 several investigators. Incubation of [<sup>14</sup>C]-radio labeled TCE ([<sup>14</sup>C]TCE) with salmon sperm  
20 DNA in the presence of microsomal preparations from B6C3F1 mice resulted in dose-related  
21 covalent binding of TCE to DNA. The binding was enhanced when the microsomes were taken  
22 from mice pretreated with phenobarbital, which induces cytochrome (CYP) P450 enzymes,  
23 suggesting the binding may be related to an oxidative metabolite, or when  
24 1,2-epoxy-3,3,3-trichloropropane, an inhibitor of epoxide hydrolase, was added to the  
25 incubations (Banerjee and Van Duuren, 1978). In addition, covalent binding of [<sup>14</sup>C]TCE with  
26 microsomal proteins was detected after incubation with microsomal preparations from mouse  
27 lung, liver, stomach and kidney, and rat liver (Banerjee and Van Duuren, 1978). Furthermore,  
28 incubation of [<sup>14</sup>C]TCE with calf thymus DNA in the presence of hepatic microsomes from  
29 phenobarbital-pretreated rats yielded significant covalent binding (DiRenzo et al., 1982).

30 A number of studies have also examined the role TCE metabolism in covalent binding to  
31 DNA and proteins. Miller and Guengerich (1983) used liver microsomes from control,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 b-naphthoflavone- and phenobarbital-induced B6C3F1 mice, Osborne-Mendel rats, and human  
2 liver microsomes. Significant covalent binding of TCE metabolites to calf thymus DNA and  
3 proteins was observed in all experiments. Phenobarbital treatment increased the formation of  
4 chloral and TCE oxide formation, DNA and protein adducts. In contrast, b-naphthoflavone  
5 treatment did not induce the formation of any microsomal metabolite suggesting that the forms  
6 of CYP induced by phenobarbital are primarily involved in TCE metabolism while the  
7 b-naphthoflavone-inducible forms of CYP have only a minor role in TCE metabolism. TCE  
8 metabolism (based on TCE-epoxide and DNA-adduct formation) was 2.5–3-fold higher in  
9 mouse than in rat microsomes due to differences in rates and clearance of metabolism (discussed  
10 in Section 3.3.3.1). The levels of DNA and protein adducts formed in human liver microsomal  
11 system approximated those observed in liver microsomes prepared from untreated rats. It was  
12 also shown that whole hepatocytes of both untreated mice and phenobarbital-induced rats and  
13 mice could activate TCE into metabolites able to covalently bind extracellular DNA. A study by  
14 Cai and Guengerich (2001b) postulate TCE oxide (an intermediate in the oxidative metabolism  
15 of TCE in rat and mouse liver microsomes) is responsible for the covalent binding of TCE with  
16 protein, and to a lesser extent, DNA. Mass spectrometry was used to analyze the reaction of  
17 TCE oxide (synthesized by m-chloroperbenzoic acid treatment of TCE) with nucleosides,  
18 oligonucleotides and protein to understand the transient nature of the inhibition of enzymes in the  
19 context of adduct formation. Protein amino acid adducts were observed during the reaction of  
20 TCE oxide with the model peptides. The majority of these adducts were unstable under  
21 physiological conditions. Results using other peptides also indicate that adducts formed from the  
22 reaction of TCE oxide with macromolecules and their biological effects are likely to be relatively  
23 short-lived.

24 Studies have been conducted using in vitro and in vivo systems to understand the DNA  
25 and protein binding capacity of TCE. In a study in male mice, after repeated intraperitoneal (i.p.)  
26 injections of [<sup>14</sup>C]TCE, radioactivity was detected in the DNA and RNA of all organs studied  
27 (kidney, liver, lung, spleen, pancreas, brain, and testis) (Bergman, 1983). However, in vivo  
28 labeling was shown to be due to metabolic incorporation of C1 fragments, particularly in guanine  
29 and adenine, rather than to DNA-adduct formation. In another study (Stott et al., 1982),  
30 following i.p. injection of [<sup>14</sup>C]TCE in male Sprague-Dawley rats (10–100 mg/kg) and B6C3F1  
31 mice (10–250 mg/kg), high liver protein labeling was observed while very low DNA labeling  
32 was detected. Stott et al. (1982) also observed very low levels of DNA binding ( $0.62 \pm 0.43$   
33 alkylation/ $10^6$  nucleotides) in mice administered 1,200 mg/kg of TCE. In addition, a  
34 dose-dependent binding of TCE to hepatic DNA and protein at low doses in mice was  
35 demonstrated by Kautiainen et al. (1997). In their dose-response study (doses between 2 µg/kg

1 and 200 mg/kg body weight [BW]), the highest level of protein binding (2.4 ng/g protein) was  
2 observed 1 hour after the treatment followed by a rapid decline, indicating pronounced instability  
3 of the adducts and/or rapid turnover of liver proteins. Highest binding of DNA (120 pg/g DNA)  
4 was found between 24 and 72 hours following treatment. Dose-response curves were linear for  
5 both protein and DNA binding. In this study, the data suggest that TCE does bind to DNA and  
6 proteins in a dose-dependent fashion, however, the type and structure of adducts were not  
7 determined.

8 Mazzullo et al. (1992) reported that TCE was covalently bound in vivo to DNA, RNA  
9 and proteins of rat and mouse organs 22 hours after i.p. injection. Labeling of proteins from  
10 various organs of both species was higher than that of DNA. Bioactivation of TCE to its  
11 intermediates using various microsomal fractions was dependent on CYP enzyme induction and  
12 the capacity of these intermediates to bind to DNA. It appeared that mouse lung microsomes  
13 were more efficient in forming the intermediates than rat lung microsomes, although no other  
14 species specific differences were found (Mazzullo et al., 1992) This also supports the results  
15 described by Miller and Guengerich (1983). The authors suggest some binding ability of TCE to  
16 interact covalently with DNA (Mazzullo et al., 1992).

17 In summary, studies report that TCE exposure in vivo can lead to binding to nucleic acids  
18 and proteins, and some authors have suggested that such binding is likely due to conversion to  
19 one or more reactive metabolites.  
20

#### 4.2.1.1.2. Bacterial Systems—Gene Mutations

21 Gene mutation studies (Ames assay) in various *Salmonella typhimurium* (*S. typhimurium*)  
22 strains of bacteria exposed to TCE both in the presence and absence of stabilizing agent have  
23 been conducted by different laboratories (Baden et al., 1979; Crebelli et al., 1982; Henschler et  
24 al., 1977; McGregor et al., 1989; Mortelmans et al., 1986; Shimada et al., 1985; Simmon et al.,  
25 1977; Waskell, 1978) (see Table 4-6). It should be noted that these studies have tested TCE  
26 samples of different purities using various experimental protocols. In all in vitro assays,  
27 volatilization is a concern when TCE is directly administered.

28 Waskell (1978) studied the mutagenicity of several anesthetics and their metabolites.  
29 Included in their study was TCE (and its metabolites) using the Ames assay. The study was  
30 conducted both in the presence and absence of a metabolic activation system, S9, and caution  
31 was exercised to perform the experiment under proper conditions (incubation of reaction mixture  
32 in sealed dessicator vials). This study was performed in both TA98 and TA100 *S. typhimurium*  
33 strains at a dose range of 0.5–10% between 4 and 48 hours. No change in revertant colonies was

1 observed in any of the doses or time courses tested. No information either on the presence or  
2 absence of stabilizers in TCE obtained commercially nor its effect on cytotoxicity was provided  
3 in the study.

**Table 4-6. TCE genotoxicity: bacterial assays**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<i>S. typhimurium</i> (TA100)	0.1–10 µL (epoxide-free)	–	–	Plate incorporation assay	Henschler et al. (1977)
<i>S. typhimurium</i> (TA1535, TA100)	1–2.5% (epoxide-free)	+ (TA100) – (TA1535)			Simmon et al. (1977)
<i>S. typhimurium</i> (TA98, TA100)	0.5–10%	–	–	The study was conducted in sealed dessicator vials	Waskell (1978)
<i>S. typhimurium</i> (TA100, TA1535)	1–3% (epoxide-free)	+ (TA100) ± (TA1535)	–		Baden et al. (1979)
<i>S. typhimurium</i> (TA100)	5–20% (v/v)	–	–	Negative under normal conditions, but twofold increase in mutations in a preincubation assay	Bartsch et al. (1979)
	0.33–1.33% (epoxide-free)	+	–		Crebelli et al. (1982)
<i>S. typhimurium</i> (TA1535, TA100)	1–5% (higher and lower purity)	– (higher purity) + (lower purity)	–	Extensive cytotoxicity	Shimada et al. (1985)
<i>S. typhimurium</i> (TA98, TA100, TA1535, TA1537, TA97)	10–1,000 µL/plate	–	–	Preincubation protocol	Mortelmans et al. (1986)
<i>S. typhimurium</i> (TA98, TA100, TA1535)	≤10,000 µg/plate (unstabilized)	–	ND	Vapor assay	McGregor et al. (1989)
	≤10,000 µg/plate (oxirane-stabilized)	+	+	Vapor assay	McGregor et al. (1989)
<i>S. typhimurium</i>	≤10,000 µg/plate (epoxybutane stabilized)	ND	+	Preincubation assay	McGregor et al. (1989)
	≤10,000 µg/plate (epichlorohydrin stabilized)	ND	+	Vapor assay	McGregor et al. (1989)
<i>S. typhimurium</i> (YG7108)	1.000–3.000 µg/plate	ND	+	Microcolony assay/revertants	Emmert et al. (2006)
<i>E. coli</i> (K12)	0.9 mM (analytical grade)	+	–	Revertants at arg56 but not nad113 or other loci	Greim et al. (1975)

ND = not determined.

1 In other studies highly purified, epoxide free TCE samples were not mutagenic in  
2 experiments with and without exogenous metabolic activation by S9 in *S. typhimurium* strain  
3 TA100 using the plate incorporation assay (Henschler et al., 1977). Furthermore, no mutagenic  
4 activity was found in several other strains including TA1535, TA1537, TA97, TA98, and  
5 TA100 using the preincubation protocol (Mortelmans et al., 1986). Simmon et al. (1977)  
6 observed a less than twofold but reproducible and dose-related increase in *his* + revertants in  
7 plates inoculated with *S. typhimurium* TA100 and exposed to a purified, epoxide-free TCE  
8 sample. The authors observed no mutagenic response in strain TA1535 with S9 mix and in  
9 either TA1535 or TA100 without rat or mouse liver S9. Similar results were obtained by Baden  
10 et al. (1979), Bartsch et al. (1979) and Crebelli et al. (1982). In all these studies purified,  
11 epoxide-free TCE samples induced slight but reproducible and dose-related increases in  
12 *his* + revertants in *S. typhimurium* TA100 only in the presence of S9. No mutagenic activity was  
13 detected without exogenous metabolic activation or when liver S9 from naïve rats, mice, and  
14 hamsters (Crebelli et al., 1982) was used for activation. Therefore, a number of these studies  
15 showed positive results in TA100 with metabolic activation, but not in other strains or without  
16 metabolic activation.

17 Shimada et al. (1985) tested a low-stabilized, highly purified TCE sample in an Ames  
18 reversion test, modified to use vapor exposure, in *S. typhimurium* TA1535 and TA100. No  
19 mutagenic activity was observed—either in the presence or absence of S9 mix. However, at the  
20 same concentrations (1, 2.5, and 5%), a sample of lower purity, containing undefined stabilizers,  
21 was directly mutagenic in TA100 (>fivefold) and TA1535 (>38-fold) at 5% concentration  
22 regardless of the presence of S9. It should be noted that the doses used in this study resulted in  
23 extensive killing of bacterial population, particularly at 5% concentration, more than 95%  
24 toxicity was observed.

25 A series of studies evaluating TCE (with and without stabilizers) was conducted by  
26 McGregor et al. (1989). The authors tested high purity and oxirane-stabilized TCE samples for  
27 their mutagenic potential in *S. typhimurium* strains TA1535, TA98, and TA100. Preincubation  
28 protocol was used to test stabilized TCE (up to 10,000 µg/plate). Mutagenic response was not  
29 observed either in the presence or absence of metabolic activation. When TCE was tested in a  
30 vapor delivery system without the oxirane stabilizers, the authors did not observe any mutagenic  
31 activity. However, TA1535 and TA100 produced a mutagenic response both in the presence and  
32 absence of S9 when exposed to TCE containing 0.5–0.6% 1,2-epoxybutane. Furthermore,  
33 exposure to epichlorohydrin also increased the frequency of mutants.

34 Emmert et al. (2006) used a CYP2E1-competent bacterial strain (*S. typhimurium*  
35 containing YG7108pin3ERb<sub>5</sub> plasmid) in their experiments. TCE was among several other

1 compounds investigated and was tested at concentrations of 1,000–3,000 µg/plate. TCE induced  
2 toxicity and microcolonies at or above 1,000 µg per plate. A study on *Escherichia coli* (*E. coli*)  
3 K12 strain was conducted by Greim et al. (1975) using analytical-grade TCE samples.  
4 Revertants were scored at two loci: *arg56*, sensitive to base-pair substitution and *nad113*, reverted  
5 by frameshift mutagens. In addition, forward mutations to 5-methyltryptophan resistance and  
6 galactose fermentation were selected. Approximately twofold increase in *arg* + colonies was  
7 observed. No change in other sites was observed. No definitive conclusion can be drawn from  
8 this study due to lack of information on reproducibility and dose-response. .

9 In addition to the above studies, the ability of TCE to induce gene mutations in bacterial  
10 strains has been reviewed and summarized by several authors (Clewell and Andersen, 2004;  
11 Crebelli and Carere, 1989; Douglas et al., 1999; Fahrig et al., 1995; Moore and Harrington-  
12 Brock, 2000). In summary, TCE, in its pure form as a parent compound is unlikely to induce  
13 point mutations in most bacterial strains. It is possible that some mutations observed in response  
14 to exposure to technical grade TCE may be contributed by the contaminants/impurities such as  
15 1,2-epoxybutane and epichlorohydrin, which are known bacterial mutagens. However, several  
16 studies of TCE reported low, but positive responses in the TA100 strain in the presence of S9  
17 metabolic activation, even when genotoxic stabilizers were not present.

#### 4.2.1.1.3. Fungal and Yeast Systems—Gene Mutations, Conversions and Recombination

19 Gene mutations, conversions, and recombinations have been studied to identify the effect  
20 of TCE in fungi and yeast systems (see Table 4-7).

21 Crebelli et al. (1985) studied the mutagenicity of TCE in *Aspergillus nidulans*  
22 (*A. nidulans*) both for gene mutations and mitotic segregation. No increase in mutation  
23 frequency was observed when *A. nidulans* was plated on selective medium and then exposed to  
24 TCE vapors. A small but statistically significant increase in mutations was observed when  
25 conidia of cultures were grown in the presence of TCE vapors and then plated on selective  
26 media. Since TCE required actively growing cells to exerts its genotoxic activity and previous  
27 studies (Bignami et al., 1980) have shown activity in the induction of *methG1* suppressors by  
28 trichloroethanol and chloral hydrate, it is possible that endogenous metabolic conversion of TCE  
29 into trichloroethanol or chloral hydrate may have been responsible for the positive response.

30 To understand the cytochrome P450 mediated genotoxic activity of TCE, Callen et al.  
31 (1980) conducted a study in two yeast strains (D7 and D4) CYP. The D7 strain in it log-phase  
32 had a CYP concentration up to five times higher than a similar cell suspension of D4 strain. Two

1 different concentrations (15 and 22 mM) at two different time points (1 and 4 hours) were  
2 studied. A significant increase in frequencies of mitotic gene conversion and recombination was



**Table 4-7. TCE genotoxicity: fungal and yeast systems**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<b>Gene conversions</b>					
<i>S. cerevisiae</i> D7 and D4	15 and 22 mM; 1 and 4 h	ND	+ at 1 h, D7 strain; – at 4 h, both D7 and D4	gene conversion; CYP content fivefold greater in D7 strain; high cytotoxicity at 22 mM	Callen et al. (1980)
<i>S. cerevisiae</i> D7	11.1, 16.6, and 22.2 mM	–	–	both stationary and log phase/production of phototropic colonies	Koch et al. (1988)
<i>S. pombe</i>	0.2–200 mM (“pure” and technical grade)	–	–	forward mutation, different experiments with different doses and time	Rossi et al. (1983)
<i>S. cerevisiae</i> D7		+	–		Bronzetti et al. (1980)
<i>A. nidulans</i>		no data	+	forward mutation	Crebelli et al. (1985)
<b>Recombination</b>					
<i>S. cerevisiae</i>		+	–	gene conversion	Bronzetti et al. (1980)
<i>S. cerevisiae</i> D7 and D4	15 and 22 mM; 1 and 4 h	ND	+		Callen et al. (1980)
<i>A. nidulans</i>		ND	+	gene cross over	Crebelli et al. (1985)
<b>Mitotic aneuploidy</b>					
<i>S. cerevisiae</i> D61.M	5.5, 11.1, and 16.6 mM	+	+	loss of dominant color homolog	Koch et al. (1988)

ND = not determined .

1 observed at 15 mM concentrations at 1-hour exposure period in the D7 strain, however, the  
2 22 mM concentration was highly cytotoxic (only 0.3% of the total number of colonies survived).  
3 No changes were seen in D4 strain, suggesting that metabolic activation via CYP played an  
4 important role in both genotoxicity and cytotoxicity. However, marginal or no genotoxic activity  
5 was observed when incubation of cells and test compounds were continued for 4 hours in either  
6 strain, possibly because of increased cytotoxicity, or a destruction of the metabolic system.

7 Koch et al. (1988) studied the genotoxic effects of chlorinated ethylenes including TCE  
8 in various yeast *Saccharomyces cerevisiae* strains. Strain D7 was tested (11.1, 16.6, and 22.2  
9 mM TCE) both in stationary-phase cells without S9, stationary-phase cells with S9 and  
10 logarithmic-phase cells using different concentrations. No significant change in mitotic gene  
11 conversion or reverse mutation was observed in either absence or presence of S9. In addition,  
12 there was a considerable increase in the induction of mitotic aneuploidy in Strain D61.M, though  
13 no statistical analysis was performed.

14 Rossi et al. (1983) studied the effect of TCE on yeast species *S. pombe* both using in vitro  
15 and host mediated mutagenicity studies and the effect of two stabilizers, epichlorohydrin and  
16 1,2-epoxybutane that are contained in the technical grade of TCE. The main goal of this study  
17 was to evaluate genotoxic activity of TCE samples of different purity and if the effect is due to  
18 the additives present in the TCE or TCE itself. Forward mutations at five loci (*ade 1, 3, 4, 5, 9*)  
19 of the adenine pathway in the yeast, strain P1 was evaluated. The stationary-phase cells were  
20 exposed to 25 mM concentration of TCE for 2, 4, and 8 hours in the presence and absence of S9.  
21 No change in mutation frequency was observed either in pure-grade samples or technical-grade  
22 samples either in the presence or absence of S9 at any of the time-points tested. Interestingly,  
23 this suggests that the stabilizers used in technical-grade TCE are not genotoxic in yeast. In a  
24 follow-up experiment, the same authors studied the effect of different concentrations (0.22, 2.2  
25 and 22.0 mM) in a host mediated assay using liver microsome preparations obtained from  
26 untreated mice, from phenobarbital-pretreated and naphthoflavone-pretreated mice and rats,  
27 which also suggested that stabilizers were not genotoxic in yeast. This experiment is described  
28 in more detail in Section 4.2.1.4.1.

29 Furthermore, TCE was tested for its ability to induce both point mutation and mitotic  
30 gene conversion in diploid strain of yeast *S. cerevisiae* (strain D7) both with and without a  
31 mammalian microsomal activation system. In a suspension test with D7, TCE was active only  
32 with microsomal activation (Bronzetti et al., 1980).

33 These studies are consistent with those of bacterial systems in indicating that pure TCE as  
34 a parent compound is not likely to cause mutations, gene conversions, or recombinations in  
35 fungal or yeast systems. In addition, the data suggest that contaminants used as stabilizers in

1 technical grade TCE are not genotoxic in these systems, and that the observed genotoxic activity  
2 in these systems is predominantly mediated by TCE metabolites.  
3

#### 4.2.1.1.4. Mammalian Systems Including Human Studies

##### 4.2.1.1.5. Gene mutations (bacterial, fungal, or yeast with a mammalian host)

4 Very few studies have been conducted to identify the effect of TCE, particularly on gene  
5 (point) mutations using mammalian systems (see Table 4-8). An overall summary of different  
6 endpoints using mammalian systems will be provided at the end of this section. In order to  
7 assess the potential mutagenicity of TCE and its possible contaminants, Rossi et al. (1983)  
8 performed genotoxicity tests using two different host mediated assays with pure- and  
9 technical-grade TCE. Male mice were administered with one dose of 2 g/kg of pure or technical  
10 grade TCE by gavage. Following the dosing, for the intraperitoneal host-mediated assay, yeast  
11 cell suspensions ( $2 \times 10^9$  cells/mL) were inoculated into the peritoneal cavity of the animals.  
12 Following 16 hours, animals were sacrificed and yeast cells were recovered to detect the  
13 induction of forward mutations at five loci (*ade 1, 2, 4, 5, 9*) of the adenine pathway. A second  
14 host-mediated assay was performed by exposing the animals to 2 g/kg of pure or technical grade  
15 TCE and inoculating the cells into the blood system. Yeast cells were recovered from livers  
16 following 4h of exposure. Forward mutations in the five loci (*ade 1,2,4,5,9*) were not observed  
17 in host-mediated assay either with pure or technical-grade TCE. Genotoxic activity was not  
18 detected when the mutagenic epoxide stabilizers were tested for mutagenicity independently or  
19 in combination. To confirm the sensitivity of the assay, the authors tested a positive  
20 control—*N*-nitroso-dimethyl-nitrosamine (1 mg/kg) and found a mutation frequency of more  
21 than 20 times the spontaneous level. The authors suggest that the negative result could have  
22 been due to an inadequate incubation time of the sample with the yeast cells.

23 Male and female transgenic *lac Z* mice were exposed by inhalation to an actual  
24 concentrations of 0, 203, 1,153, and 3,141 ppm TCE, 6 hours/day for 12 days (Douglas et al.,  
25 1999). Following 14 and 60 days of last exposure, animals were sacrificed and the mutation  
26 frequencies were determined in various organs such as bone marrow, kidney, spleen, liver, lung,  
27 and testicular germ cells. No statistically significant increases in base-changes or small-deletions  
28 were observed at any of the doses tested in male or female lung, liver, bone marrow, spleen, and  
29 kidney, or in male testicular germ cells when the animals were sampled 60 days after exposure.  
30 In addition, statistically significantly increased gene mutations were not observed in the lungs at  
31 14 days after the end of exposure (Douglas et al., 1999). The authors acknowledge that *lacZ*

*This document is a draft for review purposes only and does not constitute Agency policy.*

- 1 bacteriophage transgenic assay does not detect large deletions. The authors also acknowledge
- 2 that their hypothesis does not readily explain the increases in small deletions and base-change

**Table 4-8. TCE genotoxicity: mammalian systems—gene mutations and chromosome aberrations**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<b>Gene mutations (forward mutations)</b>					
<i>Schizosaccharomyces pombe</i>	2 g/kg, 4 and 16 h	ND	–	Host-mediated: intravenous and intraperitoneal injections of yeast cells	Rossi et al. (1983)
<b>Gene mutations (mutations frequency)</b>					
lac Z transgenic mice	0, 203, 1,153, or 3,141 ppm	No base changes or small deletions	No base changes or small deletions	Lung, liver, bone marrow, spleen, kidney, testicular germ cells used	Douglas et al. (1999)
<b>Chromosomal aberrations<sup>a</sup></b>					
CHO	745–14,900 µg/mL	ND	–	8–14 h	Galloway et al. (1987)
	499–14,900 µg/mL	–	ND	2 h exposure	Galloway et al. (1987)
C57BL/6J mice	5, 50, 500, or 5,000 ppm (6 h)	–	NA	Splenocytes	Kligerman et al. (1994)
S-D rats	5, 50, 500, or 5,000 ppm (6 h, single and 4-d exposure)	–	NA	Peripheral blood lymphocytes	Kligerman et al. (1994)

<sup>a</sup> It should be noted that results of most chromosomal aberration assays report the combined incidence of multiple effects, including chromatid breaks, isochromatid or chromosome breaks, chromatid exchanges, dicentric chromosomes, ring chromosomes, and other aberrations.

ND = not determined, NA = not applicable.

1 mutations found in the *von Hippel-Lindau* tumor suppressor gene in renal cell carcinomas of the  
2 TCE-exposed population. DCA, a TCE metabolite has been shown to increase *lacI* mutations in  
3 transgenic mouse liver, however, only after 60-weeks-of-exposure to high concentration  
4 (>1,000 ppm) in drinking water (Leavitt et al., 1997). DCA induced relatively small increase in  
5 *lac I* mutations when the animals were exposed for 60 weeks, a significantly longer duration than  
6 the TCE exposure in the Douglas et al. (1999) study (<2 weeks). Because a relatively small  
7 fraction of TCE is metabolized to DCA (see Section 3.3), the mutagenic effect of DCA is  
8 unlikely to have been detected in the experiments in Douglas et al. (1999). Glutathione (GSH)  
9 conjugation, which leads to the production of genotoxic metabolites (see Section 4.2.5),  
10 constitutes a relatively small (and relatively uncertain) portion of TCE metabolism in mice, with  
11 little data on the extent of renal DCVC bioactivation versus detoxification in mice (see  
12 Sections 3.3 and 3.5). In addition, statistically significantly increased kidney tumors have not  
13 been reported in mice with TCE treatment, and the increased incidence of kidney tumors in rats,  
14 while considered biologically significant, are quite low and not always statistically significant  
15 (see Section 4.4). Therefore, although Douglas et al. (1999) did not detect increased mutations  
16 in the kidney, these results are not highly informative as to the role of mutagenicity in  
17 TCE-induced kidney tumors, given the uncertainties in the production in genotoxic GSH  
18 conjugation metabolites in mice and the low carcinogenic potency of TCE for kidney tumors in  
19 rodents relative to what is detectable in experimental bioassays.

20

#### 4.2.1.1.6. *von Hippel-Lindau* (VHL) gene mutations

21 Studies have been conducted to determine the role of *VHL* gene mutations in renal cell  
22 carcinoma, with and without TCE exposure, and are summarized here. Most of these studies are  
23 epidemiologic, comparing *VHL* mutation frequencies of TCE-exposed to nonexposed cases from  
24 renal cell carcinoma case-control studies, or to background mutation rates among other renal cell  
25 carcinoma case series (described in Section 4.4.3). Inactivation of the *VHL* gene through  
26 mutations, loss of heterozygosity and imprinting has been observed in about 70% of renal clear  
27 cell carcinomas (Alimov et al., 2000; Kenck et al., 1996). Recent studies have also examined the  
28 role of other genes or pathways in renal cell carcinoma subtypes, including c-myc activation and  
29 vascular endothelial growth factor (VEGF) (Furge et al., 2007; Toma et al., 2008).

30 Several studies have examined the role of *VHL* gene inactivation in renal cell carcinoma,  
31 including a recent study that measured not only mutations but also promoter hypermethylation  
32 (Nickerson et al., 2008). This study focused on kidney cancer regardless of cause, and found that  
33 91% of cc-renal cell carcinoma (RCC) exhibited alterations of the *VHL* gene, suggesting a role

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 for *VHL* mutations as an early event in cc-RCC. A recent analysis of current epidemiological  
2 studies of renal cell cancer suggests *VHL* gene alterations as a marker of cc-RCC, but that  
3 limitations of previous studies may make the results difficult to interpret (Chow and Devesa,  
4 2008). Conflicting results have been reported in epidemiological studies of *VHL* mutations in  
5 TCE-exposed cases and are described in detail in Section 4.5.2. Both Brüning et al. (1997b) and  
6 Brauch et al. (1999; 2004) associated increased *VHL* mutation frequency in TCE-exposed renal  
7 cell carcinoma cases. The two other available studies of Schraml et al. (1999) and  
8 Charbotel et al. (2007) because of their limitations and lower mutation detection rate in the case  
9 of Charbotel et al. (2007) neither add nor detract to the conclusions from the earlier studies.  
10 Additional discussion of these data are in Section 4.4.3.

11 Limited animal studies have examined the role of TCE and *VHL* mutations, although  
12 Mally et al. (2006) have recently conducted both in vitro and in vivo studies using the Eker rat  
13 model (see Section 4.4.6.1.1). The Eker rat model (*Tsc-2<sup>±</sup>*) is at increased risk for the  
14 development of spontaneous renal cell carcinoma and as such has been used to understand the  
15 mechanisms of renal carcinogenesis (Stemmer et al., 2007; Wolf et al., 2000). One study has  
16 demonstrated similar pathway activation in Eker rats as that seen in humans with *VHL* mutations  
17 leading to renal cell carcinoma, suggesting *Tsc-2* inactivation is analogous to inactivation of *VHL*  
18 in human renal cell carcinoma (Liu et al., 2003). In Mally et al. (2006), male rats carrying the  
19 Eker mutation were exposed to TCE (0, 100, 250, 500, or 1,000 mg/kg BW by gavage, 5 days a  
20 week) for 13 weeks to determine the renal effects (additional data from this study on in vitro  
21 DCVC exposure are discussed below, Section 4.2.5). A significant increase in labeling index in  
22 kidney tubule cells was observed, however, no enhancement of preneoplastic lesions or tumor  
23 incidence was found in Eker rat kidneys compared to controls. In addition, no *VHL* gene  
24 mutations in exons 1–3 were detected in tumors obtained from either control or TCE-exposed  
25 Eker rats. Although no other published studies have directly examined *VHL* mutations following  
26 exposure to TCE, two studies performed mutational analysis of archived formalin-fixed paraffin  
27 embedded tissues from renal carcinomas from previous rat studies. These carcinomas were  
28 induced by the genotoxic carcinogens potassium bromate (Shiao et al., 2002) or  
29 *N*-nitrosodimethylamine (Shiao et al., 1998). Limited mutations in the *VHL* gene were observed  
30 in all samples, but, in both studies, these were found only in the clear cell renal carcinomas.  
31 Limitations of these two studies include the small number of total samples analyzed, as well as  
32 potential technical issues with DNA extraction from archival samples (see Section 4.4.3).  
33 However, analyses of *VHL* mutations in rats may not be informative as to the potential  
34 genotoxicity of TCE in humans because the *VHL* gene may not be the target for  
35 nephrocarcinogenesis in rats to the extent that it appears to be in humans.

#### 4.2.1.1.7. Chromosomal aberrations

1 A few studies were conducted to investigate the ability of TCE to induce chromosomal  
2 aberrations in mammalian systems (see Table 4-8). Galloway et al. (1987) studied the effect of  
3 TCE on chromosome aberrations in Chinese hamster ovary cells. When the cells were exposed  
4 to TCE (499–14,900 µg/mL) for 2 hours with metabolic activation, S9, no chromosomal  
5 aberrations were observed. Furthermore, without metabolic activation, no changes in  
6 chromosomal aberrations were found when the cells were exposed to TCE concentrations of  
7 745–14,900 µg/mL for 8–14 hours. It should be noted that in this study, liquid incubation  
8 method was used and the experiment was part of a larger study to understand the genotoxic  
9 potential of 108 chemicals.

10 Three inhalation studies in mice and rats examined if TCE could induce cytogenetic  
11 damage (Kligerman et al., 1994). In the first two studies, CD rats or C57Bl/6 mice, were  
12 exposed to 0-, 5-, 500-, or 5,000-ppm TCE for 6 hours. Peripheral blood lymphocytes in rats and  
13 splenocytes in mice were analyzed for induction of chromosomal aberrations, sister chromatid  
14 exchanges, and micronucleus formation. The results of micronucleus and sister chromatid  
15 exchanges will be discussed in the next sections (see Sections 4.2.1.4.4 and 4.2.1.4.5). No  
16 significant increase in chromosomal aberrations was observed in binucleated peripheral blood  
17 lymphocytes. In the third study, the authors exposed the same strain of rats for 6 hours/day over  
18 4 consecutive days. No statistically significant concentration-related increases in chromosomal  
19 aberrations were observed. The limited results of the above studies have not reported TCE to  
20 cause chromosomal aberrations either in in vitro or in vivo mammalian systems.

#### 4.2.1.1.8. Micronucleus induction

22 The appearance of micronuclei is another endpoint that can demonstrate the genotoxic  
23 effect of a chemical. Several studies have been conducted to identify if TCE can cause  
24 micronucleus formation (see Table 4-9).

25 Wang et al. (2001) investigated micronucleus formation by TCE administered as a vapor  
26 in CHO-K1 cells in vitro. Cells were grown in culture media with an inner Petri dish containing  
27 TCE that would evaporate into the media containing cells. The concentration of TCE in cultured  
28 medium was determined by gas chromatography. The actual concentration of TCE ranged from  
29 0.8 and 1.4 ppm after a 24-hour treatment. A significant dose-dependent increase in micronuclei  
30 formation was observed. A dose-dependent decrease in cell growth and cell number was also  
31 observed. The authors did not test if the micronuclei formed was due to direct damage to the  
32 DNA or spindle formation.

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-9. TCE genotoxicity: mammalian systems—micronucleus, sister chromatic exchanges**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<b>Micronucleus</b>					
Human hepatoma HepG2 cells	0.5–4 mM, 24 h	NA	+		Hu et al. (2008)
Primary cultures of human and rat kidney cells	1.0, 2.0, or 4.0 mM	NA	+	Dose-dependent significant increase	Robbiano et al. (2004)
<b>Sprague-Dawley rats</b>	3,591 mg/kg	+	–		Robbiano et al. (2004)
CHO-K1 cells	0.8–1.4 ppm		+	Dose-dependent significant increase	Wang et al. (2001)
Male CD-1 mice	457 mg/kg	+	NA	Bone marrow, correlated with TCOH in urine	Hrelia et al. (1994)
C56BL/6J mice	5, 50, 500, or 5,000 ppm	–	NA	Splenocytes	Kligerman et al. (1994)
S-D rats	5, 50, 500, or 5,000 ppm	+	NA	Dose dependent; peripheral blood lymphocytes	Kligerman et al. (1994)
<b>Sister chromatid exchanges</b>					
CHO	0.17%	–	ND	1 h (vapor)	White et al. (1979)
	17.9–700 µg/mL	ND	+	25 h (liquid)	Galloway et al. (1987)
	49.7–14,900 µg/mL	+	ND	2 h	Galloway et al. (1987)
Human lymphocytes	178 µg/mL	ND	+		Gu et al. (1981a; 1981b)
S-D rats	5, 50, 500, or 5,000 ppm	–	NA	Peripheral blood lymphocytes	Kligerman et al. (1994)
Peripheral blood lymphocytes from humans occupationally exposed	Occupational exposure	–	NA		Nagaya et al., 1989a
C57BL/6J mice	5, 50, 500, or 5,000 ppm	–	NA	Splenocytes	Kligerman et al. (1994)

ND = not determined, NA = not applicable.

1 Robbiano et al. (2004) conducted an in vitro study on DNA damage and micronuclei  
2 formation in rat and human kidney cells exposed to six carcinogenic chemicals including TCE.  
3 The authors examined for the ability of TCE to induce DNA fragmentation and formation of  
4 micronuclei in primary cultures of rat and human kidney cells derived from kidney cancer  
5 patients with 1–4 mM TCE concentrations. A significant dose-dependent increase in the  
6 frequency of micronuclei was obtained in primary kidney cells from both male rats and human of  
7 both genders. The authors acknowledge that the significance of the results should be considered  
8 in light of the limitations including (1) examination of TCE on cells from only three rats,  
9 (2) considerable variation in the frequency of DNA lesions induced in the cells, and (3) the  
10 possibility that kidney cells derived from kidney cancer patients may be more sensitive to  
11 DNA-damaging activity due to a more marked expression of enzymes involved in the metabolic  
12 activation of kidney procarcinogens and suppression of DNA repair processes. Never the less,  
13 this study is important and provides information of the possible genotoxic effects of TCE.

14 In the same study, Robbiano et al. (2004) administered rats a single oral dose of TCE  
15 (3,591 mg/kg) corresponding to  $\frac{1}{2}$  LD<sub>50</sub>, which had been pre-exposed to folic acid for 48 hours  
16 and the rats were euthanized 48 hours later following exposure to TCE. The frequency of  
17 binucleated cells was taken as an index of kidney cell proliferation. A statistically significant  
18 increase in the average frequency of micronucleus was observed.

19 Hu et al. (2008) studied the effect of TCE on micronuclei frequencies using human  
20 hepatoma HepG2 cells. The cells were exposed to 0.5, 1, 2, and 4 mM TCE for 24 hours. TCE  
21 caused a significant increase in micronuclei frequencies at all concentrations tested. It is  
22 important to note that similar concentrations were used in Robbiano et al. (2004).

23 As described in the chromosomal aberration section (see Section 4.2.1.4.3), inhalation  
24 studies were performed using male C57BL/6 mice and CD rats (Kligerman et al., 1994) to  
25 determine if TCE could induce micronuclei. In the first and second study, rats or mice  
26 respectively, were exposed to 0-, 5-, 500-, or 5,000-ppm TCE for 6 hours. Peripheral blood  
27 lymphocytes in rats and splenocytes in mice were cultured and analyzed for induction of  
28 micronuclei formation. Bone marrow polychromatic erythrocytes (PCEs) were also analyzed for  
29 micronuclei. TCE caused a statistically significant increase in micronuclei formation at all  
30 concentrations in rat bone marrow PCEs but not in mice. The authors note that TCE was  
31 significantly cytotoxic at the highest concentration tested as determined by significant  
32 concentration-related decrease in the ratio of PCEs/normochromatic erythrocytes. In the  
33 third study, to confirm the results of the first study, the authors exposed rats to one dose of  
34 5,000 ppm for 6 hours. A statistical increase in bone marrow micronuclei-PCEs was observed  
35 confirming the results of the first study.

1 Hrelia et al. (1994) treated male CD-1 mice with TCE (457 mg/kg BW; i.p.) for 30 hours.  
2 Bone marrow cells were harvested for determination of micronuclei frequencies in PCEs. An  
3 increase in micronuclei frequency at 30 hours after treatment was observed. Linear regression  
4 analysis showed that micronuclei frequency induced by TCE correlated with trichloroethanol  
5 concentrations in urine, a marker of TCE oxidative metabolism (Hrelia et al., 1994).

6 In summary, based on the results of the above studies, TCE is capable of inducing  
7 micronuclei in different in vitro and in vivo systems tested. Specific methods were not used that  
8 could definitively identify the mechanism of micronuclei formation. These are important  
9 findings that indicate TCE has genotoxic potential as measured by the micronucleus formation.  
10

#### 4.2.1.1.9. Sister chromatid exchanges (SCEs)

11 Studies have been conducted to understand the ability of TCE to induce sister chromatid  
12 exchanges (SCEs) both in vitro and in vivo systems (see Table 4-9). White et al. (1979)  
13 evaluated the possible induction of SCE in CHO using a vapor exposure procedure by exposing  
14 the cells to TCE (0.17%) for 1 hour in the presence of S9 metabolic activation. No change in  
15 SCE frequencies were observed between the control and the treatment group. However, in  
16 another study by Galloway et al. (1987) a dose-related increase in SCE frequency in repeated  
17 experiments both with and without metabolic activation was observed. It should be noted that in  
18 this study, liquid incubation was used, and the exposure times were 25 hours without metabolic  
19 activation at a concentration between 17.9–700 µg/mL and 2 hours in the presence of S9 at a  
20 concentration of 49.7–14,900 µg/mL. Due to the difference in the dose, length of exposure and  
21 treatment protocol (vapor exposure vs. liquid incubation), no direct comparison can be made. It  
22 should also be noted that inadequacy of dose selection and the absence of positive control in the  
23 White et al. (1979) makes it difficult to interpret the study. In another study (Gu et al.,  
24 1981a)sss, a small but positive response was observed in assays with peripheral lymphocytes.

25 No statistically significant increase in SCEs was found when male C57Bl/6 mice or CD  
26 rats were exposed to TCE at concentrations of 5,500, or 5,000 ppm for 6 hours (Kligerman et al.,  
27 1994). Furthermore, in another study by Nagaya et al. (1989a), lymphocytes of TCE-exposed  
28 workers ( $n = 22$ ) and matched controls ( $n = 22$ ) were analyzed for SCEs. The workers had  
29 constantly used TCE in their jobs although the exact exposure was not provided. The duration of  
30 their employment ranged from 0.7–34 years, averaging about 10 years. It should be noted that  
31 there were both smokers and nonsmokers among the exposed population. If a subject had not  
32 smoked for at least 2 years before the samples were taken, then they were considered as  
33 nonsmokers. There were eight nonsmokers in the group. If they were classified as smokers,

1 then they smoked between 10–50 cigarettes per day. No significant increase in mean SCE  
2 frequencies were found in exposed population compared to controls, though the study is  
3 relatively small.

4 In summary, induction of SCEs have been reported in several, though not all, paradigms  
5 of TCE exposure, consistent with the structural damage to DNA/chromosomes indicated by  
6 excess micronuclei formation.

7

#### 4.2.1.1.10. **Unscheduled DNA synthesis**

8 In vitro studies are briefly described here, with additional discussion of effects related to  
9 TCE-induced unscheduled DNA synthesis in the context of the liver in Section E.2.4.1. Perocco  
10 and Prodi (1981) studied unscheduled DNA synthesis in human lymphocytes cultured in vitro  
11 (see Table 4-10). Three doses of TCE (2.5, 5.0, and 10  $\mu\text{L}/\text{mL}$ ) were used as final  
12 concentrations with and without S9. The results indicate that there was an increase in  
13 unscheduled DNA synthesis (UDS) only in the presence of S9, and in addition, the increase was  
14 maximal at the TCE concentration of 5  $\mu\text{L}/\text{mL}$ . Three chlorinated ethane and ethylene solvent  
15 products were examined for their genotoxicity in hepatocyte primary culture DNA repair assays  
16 using vapor phase exposures. Rat hepatocytes primary cultures were initiated and exposed to  
17 low-stabilized or standard stabilized TCE (0.1–2.5%) for 3 or 18 hours. Unscheduled DNA  
18 synthesis or DNA repair was not observed using either low or standard stabilized TCE, even at  
19 vapor phase doses up to those that produced extensive cell killing after 3 or 18 hour exposure  
20 (Shimada et al., 1985). Costa and Ivanetich (1984) examined the ability of TCE to induce  
21 unscheduled DNA synthesis hepatocytes isolated from phenobarbital treated rats. The UDS was  
22 assessed only at the highest concentration that is tolerated by the hepatocytes (2.8 mM TCE).

23 These results indicate that TCE stimulated unscheduled DNA synthesis in isolated rodent  
24 hepatocytes, and, importantly, in human lymphocytes in vitro.

25

#### 4.2.1.1.11. **DNA strand breaks**

26 DNA damage in response to TCE exposure was studied using comet assay in human  
27 hepatoma HepG2 cells (Hu et al., 2008; see Table 4-10). The cells were exposed to 0.5, 1, 2, and  
28 4 mM for 24 hours. TCE increased the DNA migration in a significant dose-dependent manner  
29 at all tested concentrations suggesting TCE caused DNA strand breaks and chromosome damage.

30 TCE (4–10 mmol/kg BW) were given to male mice by i.p. injection. The induction of  
31 single-strand breaks (SSB) in DNA of liver, kidney, and lung was studied by the DNA

*This document is a draft for review purposes only and does not constitute Agency policy.*

- 1 unwinding technique. There was a linear increase in the level of single strand breaks in kidney
- 2 and liver DNA but not in lung DNA 1 hour after administration (Wallis, 1986).

**Table 4-10. TCE genotoxicity: mammalian systems—unscheduled DNA synthesis, DNA strand breaks/protein crosslinks, cell transformation**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<b>Unscheduled DNA synthesis</b>					
Rat primary hepatocytes		ND	–		Shimada et al. (1985)
Human lymphocytes	2.5, 5, 10 µL/mL	±	–	Increase was only in certain doses and maximum at 5 µL/mL conc.	Perocco and Prodi (1981)
Phenobarbital induced rat hepatocytes	2.8 mM	ND	+		Costa and Ivanetich (1984)
<b>DNA strand breaks/protein crosslinks</b>					
Primary rat kidney cells	0.5, 1.0, 2.0, 4.0 mM	NA	+	Dose-dependent significant increase	Robbiano et al. (2004)
Primary cultures of human kidney cells	1.0, 2.0, 4.0 mM	ND	+	Dose-dependent significant increase	Robbiano et al. (2004)
Sprague-Dawley rats	3,591 mg/kg	+	NA	Single oral administration	Robbiano et al. (2004)
Sprague-Dawley rats	500, 1,000, and 2,000 ppm	–	NA	Comet assay	Clay (2008)
<b>Cell transformation</b>					
BALB/c 3T3 mouse cells	4, 20, 100, 250 µg/mL	NA	+	Weakly positive compared to other halogenated compounds tested in the same experiment	Tu et al. (1985)
Rat embryo cells		NA	+		Price et al. (1978)
Syrian hamster embryo cells	5, 10, 25 µg/mL	NA	–		Amacher and Zelljadt (1983)

ND = not determined, NA = not applicable.

1 Robbiano et al. (2004) conducted an in vitro study on DNA damage in rat and human  
2 kidney cells exposed to six carcinogenic chemicals including TCE in the comet assay. The  
3 authors examined the ability of TCE to induce DNA fragmentation in primary cultures of rat and  
4 human kidney cells with 1–4 mM TCE concentrations. TCE was dissolved in ethanol with a  
5 maximum concentration of 0.3% and the rat cultures were exposed to 20 hours. Primary human  
6 kidney cells were isolated from fragments of kidney discarded during the course of surgery for  
7 carcinoma of both male and female donors with an average age of 64.2 years and were also  
8 exposed to 20 hours. Significant dose-dependent increases in the ratio of treated/control tail  
9 length (average 4–7  $\mu$ M compared to control) was observed as measured by comet assay in  
10 primary kidney cells from both male rats and human of both genders.

11 Clay et al. (2008) studied the DNA damage inducing capacity of TCE using the comet  
12 assay in rat kidney proximal tubules. Rats were exposed by inhalation to a range of TCE  
13 concentrations (500, 1,000, or 2,000 ppm) for 6 hours/day for 5 days. TCE did not induce DNA  
14 damage (as measured by tail length and percentage tail DNA and tail movement) in rat kidney  
15 proximal tubules in any of the doses tested possibly due to study limitations (small number of  
16 animals tested [ $n = 5$ ] and limited exposure time [6 hours/day for only 5 days]). These results  
17 are in contrast to the findings of Robbiano et al. (2004) which showed DNA damage and  
18 increased micronuclei in the rat kidney 20 hours following a single dose (3,591 mg/kg BW) of  
19 TCE. The DNA damage reported by comet assay is consistent with results for other markers  
20 of chromosomal damage or DNA structural damage such as excess micronuclei formation and  
21 SCE induced by TCE exposure.

22

#### 4.2.1.1.12. DNA damage related to oxidative stress, polymorphisms

23 A detailed description of studies related to lipid peroxidation of TCE is presented in  
24 conjunction with discussion of liver toxicity (see Section 4.5, E.2.4.3, and E.3). A recent study  
25 reported on genetic polymorphism in solvent exposed population (Kumar et al., 2009). Normal  
26 ( $n = 220$ ) and solvent-exposed ( $n = 97$ ) population was genotyped for CYP1A1, GSTM1,  
27 GSTT1 and GSTP1 polymorphisms. No exposure related differences were observed. In  
28 addition, the authors also examined TCE-exposed lymphocytes for the presence of chromosomal  
29 aberrations and micronucleus at concentration of 2, 4 or 6 mM TCE. No significant changes in  
30 any of the parameters were observed.

31

#### 4.2.1.1.13. Cell transformation

1           In vitro cell transformation using BALB/c-3T3 cells was conducted using TCE with  
2 concentrations varying from 0–250 µg/mL in liquid phase exposed for 72 hours (see Table 4-10).  
3 The cytotoxicity of TCE at the concentration tested in the transformation assay was determined  
4 by counting cells from duplicate plates of each test conditions at the end of the treatment period.  
5 A dose-dependent increase in Type III foci was observed although no statistical analysis was  
6 conducted (Tu et al., 1985). In another study by Amacher and Zelljadt (1983), Syrian hamster  
7 embryo cells were exposed to 5, 10, or 25 µg/mL of TCE. In this experiment, two different  
8 serums (horse serum and fetal bovine serum) were also tested to understand the importance of  
9 serum quality in the transformation assay. Preliminary toxicity assay was performed to select  
10 dose levels which had 50–90% cell survival. One week after dosing, the cell colonies were fixed  
11 and counted for variability determination and examination of individual colonies for the evidence  
12 of morphological transformation. No significant change in morphological cell transformation  
13 was obtained. Furthermore, no significant changes were seen in transformed colonies when  
14 tested in different serum. However, these studies are of limited use for determining the  
15 genotoxic potential of TCE because they did not examine the foci for mutations, for instance in  
16 oncogenes or tumor suppressor genes.  
17

#### 4.2.1.1.14. Summary

18           Evidence from a number of different analyses and a number of different laboratories  
19 using a fairly complete array of endpoints suggests that TCE, following metabolism, has the  
20 potential to be genotoxic. A series of carefully controlled studies evaluating TCE itself (without  
21 mutagenic stabilizers and without metabolic activation) found it to be incapable of inducing gene  
22 mutations in most standard mutation bacterial assays (Baden et al., 1979; Bartsch et al., 1979;  
23 Crebelli et al., 1982; Henschler et al., 1977; Mortelmans et al., 1986; Shimada et al., 1985;  
24 Simmon et al., 1977; Waskell, 1978). Therefore, it appears that it is unlikely that TCE is a  
25 direct-acting mutagen, though TCE has shown potential to affect DNA and chromosomal  
26 structure. Low, but positive responses were observed in the TA100 strain in the presence of S9  
27 metabolic activation, even when genotoxic stabilizers were not present, suggesting metabolites  
28 of TCE are genotoxic. TCE is also positive in some but not all fungal and yeast systems (Callen  
29 et al., 1980; Crebelli et al., 1985; Koch et al., 1988; Rossi et al., 1983). Data from human  
30 epidemiological studies support the possible mutagenic effect of TCE leading to *VHL* gene  
31 damage and subsequent occurrence of renal cell carcinoma. Association of increased *VHL*



1 mutation frequency in TCE-exposed renal cell carcinoma cases has been observed (Brauch et al.,  
2 1999; Brauch et al., 2004; Brüning et al., 1997b).

3 TCE can lead to binding to nucleic acids and proteins (Bergman, 1983; DiRenzo et al.,  
4 1982; Kautiainen et al., 1997; Mazzullo et al., 1992; Miller and Guengerich, 1983), and such  
5 binding appears to be due to conversion to one or more reactive metabolites. For instance,  
6 increased binding was observed in samples bioactivated with mouse and rat microsomal fractions  
7 (Banerjee and Van Duuren, 1978; DiRenzo et al., 1982; Mazzullo et al., 1992; Miller and  
8 Guengerich, 1983). DNA binding is consistent with the ability to induce DNA and chromosomal  
9 perturbations. Several studies report the induction of micronuclei in vitro and in vivo from TCE  
10 exposure (Hrelia et al., 1994; Hu et al., 2008; Kligerman et al., 1994; Robbiano et al., 2004;  
11 Wang et al., 2001). Reports of SCE induction in some studies are consistent with DNA effects,  
12 but require further study (Gu et al., 1981a; Gu et al., 1981b; Kligerman et al., 1994; Nagaya et  
13 al., 1989a; White et al., 1979).

14 Overall, evidence from a number of different analyses and a number of different  
15 laboratories using various genetic endpoints indicates that TCE has a potential to induce damage  
16 to the structure of the chromosome in a number of targets but has a more limited ability to induce  
17 mutation in bacterial systems.

18 Below, the genotoxicity data for TCE metabolites TCA, DCA, TCOH, chloral hydrate,  
19 DCVC, and DCVG are briefly reviewed. The contributions of these data are twofold. First, to  
20 the extent that these metabolites may be formed in the in vitro and in vivo test systems for TCE,  
21 they provide insight into what agent or agents may contribute to the limited activity observed  
22 with TCE in these genotoxicity assays. Second, because the in vitro systems do not necessarily  
23 fully recapitulate in vivo metabolism, the genotoxicity of the known in vivo metabolites  
24 themselves provide data as to whether one may expect genotoxicity to contribute to the toxicity  
25 of TCE following in vivo exposure.

#### 26 **4.2.2. Trichloroacetic Acid (TCA)**

27 The TCE metabolite, TCA, has been studied using a variety of genotoxicity assay for its  
28 genotoxic potential (see International Agency for Research on Cancer [IARC, 2004] for  
29 additional information). Evaluation of in vitro studies of TCA must consider toxicity and  
30 acidification of medium resulting in precipitation of proteins, as TCA is commonly used as a  
31 reagent to precipitate proteins.

#### 4.2.2.1.1. Bacterial Systems—Gene Mutations

TCA has been evaluated in a number of in vitro test systems including the bacterial assays (Ames) using different *S. typhimurium* strains such as TA98, TA100, TA104, TA1535, and RSJ100 (see Table 4-11). The majority of these studies did not report positive findings for genotoxicity Shirasu et al., (1976); Nestmann et al., (1980) Moriya et al., (1983) (DeMarini et al., 1994; Kargalioglu et al., 2002; Nelson et al., 2001; Rapson et al., 1980; Waskell, 1978). Waskell (1978) studied the effect of TCA (0.45 mg/plate) on bacterial strains TA98 and TA100 both in

**Table 4-11. Genotoxicity of trichloroacetic acid—bacterial systems**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
λ Prophage induction, <i>E. coli</i> WP2s	10,000	-	-	DeMarini et al. (1994)
SOS chromotest, <i>Escherichia coli</i> PQ37	10,000	-	-	Giller et al. (1997)
<i>S. typhimurium</i> TA1535, 1536, 1537, 1538, reverse mutation	20 µg/plate	NT	-	Shirasu et al., (1976)
<i>S. typhimurium</i> TA100, 98, reverse mutation	450 µg/plate	-	-	Waskell (1978)
<i>S. typhimurium</i> TA100, 1535, reverse mutation	4,000 µg/plate	-	-	Nestmann et al., (1980)
<i>S. typhimurium</i> TA1537, 1538, 98, reverse mutation	2,000 µg/plate	-	-	Nestmann et al., (1980)
<i>S. typhimurium</i> TA100, reverse mutation	520 µg/plate	NT	-	Rapson et al. (1980)
<i>S. typhimurium</i> TA100, 98, reverse mutation	5,000 µg/plate	-	-	Moriya et al., (1983)
<i>S. typhimurium</i> TA100, reverse mutation	600 ppm	-	-	DeMarini et al. (1994)
<i>S. typhimurium</i> TA100, reverse mutation, liquid medium	1,750	+	+	Giller et al. (1997)
<i>S. typhimurium</i> TA104, reverse mutation, microsuspension	250 µg/plate	-	-	Nelson et al. (2001)
<i>S. typhimurium</i> TA100, RSJ100, reverse mutation	16,300	-	-	Kargalioglu et al. (2002)
<i>S. typhimurium</i> TA98, reverse mutation	13,100	-	-	Kargalioglu et al. (2002)
<i>S. typhimurium</i> TA1535, SOS DNA repair		+	-	Ono et al. (1991)

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose; doses are in µg/mL for in vitro tests unless specified.

<sup>b</sup> Results: +, positive; -, negative; NT, not tested.

Table adapted from IARC monograph (2004a) and modified/updated for newer references.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2  
3 the presence and absence of S9. The author did not find any revertants at the maximum nontoxic  
4 dose tested. Following exposure to TCA, Rapson et al. (1980) reported no change in mutagenic  
5 activity in strain TA100 in the absence of S9. DeMarini et al. (1994) performed different studies  
6 to evaluate the genotoxicity of TCA, including the Microscreen prophage-induction assay (TCA  
7 concentrations 0–10 mg/mL) and use of the *S. typhimurium* TA100 strain using bag vaporization  
8 technique (TCA concentrations 0–100 ppm), neither of which yielded positive results.  
9 Nelson et al. (2001) reported no positive findings with TCA using a *S. typhimurium*  
10 microsuspension bioassay (*S. typhimurium* strain TA104) following incubation of TCA for  
11 various lengths of time, with or without rat cecal microbiota. Similarly, no activity was observed  
12 in a study conducted by Kargalioglu et al. (2002) where *S. typhimurium* strains TA98, TA100,  
13 and RSJ100 were exposed to TCA (0.1–100 mM) either in the presence or absence of S9  
14 (Kargalioglu et al., 2002).

15 TCA was also negative in other bacterial systems. The SOS chromotest (which measures  
16 DNA damage and induction of the SOS repair system) in *E. coli* PQ37, ± S9 (Giller et al., 1997)  
17 evaluated the genotoxic activity of TCA ranging from 10–10,000 µg/mL and did not find any  
18 response. Similarly, TCA was not genotoxic in the Microscreen prophage-induction assay in *E.*  
19 *coli* with TCA concentrations ranging from 0–10,000 µg/mL, with and without S9 activation  
20 (DeMarini et al., 1994).

21 However, TCA induced a small increase in SOS DNA repair (an inducible error-prone  
22 repair system) in *S. typhimurium* strain TA1535 in the presence of S9 (Ono et al., 1991).  
23 Furthermore, Giller et al. (1997) reported that TCA demonstrated genotoxic activity in an Ames  
24 fluctuation test in *S. typhimurium* TA100 in the absence of S9 at noncytotoxic concentrations  
25 ranging from 1,750–2,250 µg/mL. The addition of S9 decreased the genotoxic response, with  
26 effects observed at 3,000–7,500 µg/mL. Cytotoxic concentrations in the Ames fluctuation assay  
27 were 2,500 and 10,000 µg/mL without and with microsomal activation, respectively.  
28

#### 4.2.2.1.2. Mammalian Systems

##### 4.2.2.1.3. Gene mutations

29 The mutagenicity of TCA has also been tested in cultured mammalian cells (see  
30 Table 4-12). Harrington-Brock et al. (1998) examined the potential of TCA to induce mutations  
31 in L5178Y/TK<sup>±</sup>-3.7.2C mouse lymphoma cells. In this study, mouse lymphoma cells were  
32 incubated in culture medium treated with TCA concentrations up to 2,150 µg/mL in the presence

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 of S9 metabolic activation and up to 3,400  $\mu\text{g}/\text{mL}$  in the absence of S9 mixture. In the presence  
2 of S9, a doubling of mutant frequency was seen at concentrations of 2,250  $\mu\text{g}/\text{mL}$  and higher,  
3 including several concentrations with survival  $>10\%$ . In the absence of S9, TCA increased the  
4 mutant frequency by twofold or greater only at concentrations of 2,000  $\mu\text{g}/\text{mL}$  or higher. These  
5 results were obtained at  $\leq 11\%$  survival rates. The authors noted that the mutants included both  
6 large-colony and small-colony mutants. The small-colony mutants are indicative of  
7 chromosomal damage. It should be noted that no rigorous statistical evaluation was conducted  
8 on these data. Cytotoxic and genotoxic effects of TCA were tested in a microplate-based  
9 cytotoxicity test and a HGPRT gene mutation assay using Chinese hamster ovary K1 cells,  
10 respectively (Zhang et al., 2010). TCA was the least cytotoxic when compared to six other

**Table 4-12. TCA Genotoxicity—mammalian systems (both in vitro and in vivo)**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
Gene mutation, mouse lymphoma L5178Y/TK ± cells, in vitro	3,000	(+)	?	Harrington-Brock et al. (1998)
Gene mutation, Chinese hamster ovary cells in vitro, HGPRT gene mutation assay	10,000 µM	NT	-	Zhang et al. (2010)
DNA strand breaks, B6C3F1 mouse and Fischer 344 rat hepatocytes, in vitro	1,630	NT	-	Chang et al. (1992)
DNA strand breaks, human CCRF-CEM lymphoblastic cells, in vitro	1,630	NT	-	Chang et al. (1992)
DNA damage, Chinese hamster ovary cells, in vitro, comet assay	3 mM	NT	-	Plewa et al. (2002)
DNA strand breaks, B6C3F1 mouse liver, in vivo	1.0, oral, × 1	+		Nelson and Bull (1988)
DNA strand breaks, B6C3F1 mouse liver, in vivo	500, oral, × 1	+		Nelson et al. (1989)
DNA strand breaks, B6C3F1 mouse liver, in vivo	500, oral, 10 repeats	-		Nelson et al. (1989)
DNA strand breaks, B6C3F1 mouse liver and epithelial cells from stomach and duodenum, in vivo	1,630, oral, × 1	-		Chang et al. (1992)
DNA strand breaks, male B6C3F1 mice, in vivo	500 (neutralized)	-		Styles et al. (1991)
Micronucleus formation, Swiss mice, in vivo	125, i.p., × 2	+		Bhunya and Behera (1987)
Micronucleus formation, female C57BL/6JfBL10/Alpk mouse bone-marrow erythrocytes, in vivo	1,300, i.p., × 2	-		Mackay et al. (1995)
Micronucleus formation, male C57BL/6JfBL10/Alpk mouse bone-marrow erythrocytes, in vivo	1,080, i.p., × 2	-		Mackay et al. (1995)
Micronucleus formation, <i>Pleurodeles waltl</i> newt larvae peripheral erythrocytes, in vivo	80	+		Giller et al. (1997)
Chromosomal aberrations, Swiss mouse bone-marrow cells in vivo	125, i.p., × 1	+		Bhunya and Behera (1987)
	100, i.p., × 5	+		Bhunya and Behera (1987)
	500, oral, × 1	+		Bhunya and Behera (1987)
Chromosomal aberrations, chicken <i>Gallus domesticus</i> bone marrow, in vivo	200, i.p., × 1	+		Bhunya and Jena (1996)

5/12/11

*This document is a draft for review purposes only and does not constitute Agency policy.*

63

**DRAFT—DO NOT CITE OR QUOTE**

**Table 4-12. TCA Genotoxicity—mammalian systems (both in vitro and in vivo) (continued)**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
Chromosomal aberrations, human lymphocytes, in vitro	5,000, (neutralized)	-		Mackay et al. (1995)
Sperm morphology, Swiss mice, in vivo	125, i.p., × 5	+		Bhunya and Behera (1987)

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose; doses are in µg/mL for in vitro tests; mg/kg for in vivo tests unless specified.

<sup>b</sup> Results: + = positive; (+) = weakly positive; - = negative; NT = not tested; ? = inconclusive.

Table adapted from IARC monograph (2004a) and modified/updated for newer references.

1 haloacetic acids. TCA at concentrations 0, 200, 1,000, 5,000 and 10,000  $\mu\text{M}$  induced a visible  
2 increase in mutant frequency but did not show any statistically significant increase at any of the  
3 doses tested.  
4

#### 4.2.2.1.4. Chromosomal aberrations

5 Mackay et al. (1995) investigated the ability of TCA to induce chromosomal damage in  
6 an in vitro chromosomal aberration assay using cultured human cells. The authors treated the  
7 cells with TCA as free acid, both in the presence and absence of metabolic activation. TCA  
8 induced chromosomal damage in cultured human peripheral lymphocytes at concentrations  
9 (2,000 and 3,500  $\mu\text{g}/\text{mL}$ ) that significantly reduced the pH of the medium. However, exposure  
10 of cells to neutralized TCA did not have any effect even at a cytotoxic concentration of  
11 5,000  $\mu\text{g}/\text{mL}$ . It is possible that the reduced pH was responsible for the TCA-induced  
12 clastogenicity in this study. To further evaluate the role of pH changes in the induction of  
13 chromosome damage, the authors isolated liver-cell nuclei from B6C3F1 mice and suspended in  
14 a buffer at various pH levels. The cells were stained with chromatin-reactive (fluorescein  
15 isothiocyanate) and DNA-reactive (propidium iodide) fluorescent dyes. A decrease in chromatin  
16 staining intensity was observed with the decrease in pH, suggesting that pH changes,  
17 independent of TCA exposure, can alter chromatin conformation. It was concluded by the  
18 authors that TCA-induced pH changes are likely to be responsible for the chromosomal damage  
19 induced by un-neutralized TCA. In another in vitro study, Plewa et al. (2002) evaluated the  
20 induction of DNA strand breaks induced by TCA (1–25 mM) in CHO cells and did not observe  
21 any genotoxicity.  
22

#### 4.2.2.1.5. Micronucleus

23 Relative genotoxicity of TCA was tested in a mouse in vivo system (see Table 4-12)  
24 using three different cytogenetic assay (bone marrow chromosomal aberrations, micronucleus  
25 and sperm-head abnormalities) (Bhunya and Behera, 1987) and for chromosomal aberrations in  
26 chicken (Bhunya and Jena, 1996). TCA induced a variety of anomalies including micronucleus  
27 in the bone marrow of mice and chicken. A small increase in the frequency of micronucleated  
28 erythrocytes at 80  $\mu\text{g}/\text{mL}$  in a newt (*Pleurodeles waltl* larvae) micronucleus test was observed in  
29 response to TCA exposure (Giller et al., 1997). Mackay et al. (1995) investigated the ability of  
30 TCA to induce chromosomal DNA damage in the in vivo bone-marrow micronucleus assay in  
31 mice. C57BL mice were given TCA intraperitoneally at doses of 0, 337, 675, or

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 1,080 mg/kg-day for males and 0, 405, 810, or 1,300 mg/kg-day for females for two consecutive  
2 days, and bone-marrow samples were collected 6 and 24 hours after the last dose. The  
3 administered doses represented 25, 50, and 80% of the median lethal dose, respectively. No  
4 treatment-related increase in micronucleated polychromatic erythrocytes was observed.  
5

#### 4.2.2.1.6. Other DNA damage studies

6 DNA unwinding assays have been used as indicators of single strand breaks and are  
7 discussed in detail in Section E.2.3. Studies were conducted on the ability of TCA to induce  
8 single-strand breaks (Chang et al., 1992; Nelson and Bull, 1988; Nelson et al., 1989; Styles et al.,  
9 1991; see Table 4-12). Nelson and Bull (1988) evaluated the ability of TCA and other  
10 compounds to induce single-strand DNA breaks in vivo in Sprague-Dawley rats and B6C3F1  
11 mice. Single oral doses were administered to three groups of three animals, with an additional  
12 group as a vehicle control. Animals were sacrificed after 4 hours, and 10% liver suspensions  
13 were analyzed for single-strand DNA breaks by the alkaline unwinding assay. Dose-dependent  
14 increases in single-strand DNA breaks were induced in both rats and mice, with mice being more  
15 susceptible than rats. The lowest dose of TCA that produced significant SSBs was 0.6 mmol/kg  
16 (98 mg/kg) in rats but 0.006 mmol/kg (0.98 mg/kg) in mice.

17 However, in a follow-up study, Nelson et al. (1989) male B6C3F1 mice were treated with  
18 500 mg/kg TCA, and single strand breaks in whole liver homogenate were examined, and no  
19 significant differences from controls were reported. Moreover, in the experiments in the same  
20 study with DCA, increased single strand breaks were reported, but with no dose-response  
21 between 10 and 500 mg/kg, raising concerns about the reliability of the DNA unwinding assay  
22 used in these studies. For further details, see Section E.2.3. In an additional follow-up  
23 experiment with a similar experimental paradigm, Styles et al. (1991) tested TCA for its ability  
24 to induce strand breaks in male B6C3F1 mice in the presence and absence of liver growth  
25 induction. The test animals were given one, two, or three daily doses of neutralized TCA  
26 (500 mg/kg) by gavage and killed 1 hour after the final dose. Additional mice were given a  
27 single 500-mg/kg gavage dose and sacrificed 24 hours after treatment. Liver nuclei DNA were  
28 isolated, and the induction of single strand breaks was evaluated using the alkaline unwinding  
29 assay. Exposure to TCA did not induce strand breaks under the conditions tested in this assay.  
30 In a study by Chang et al. (1992), administration of single oral doses of TCA (1–10 mmol/kg) to  
31 B6C3F1 mice did not induce DNA strand breaks in a dose-related manner as determined by the  
32 alkaline unwinding assay. No genotoxic activity (evidence for strand breakage) was detected in  
33 F344 rats administered by gavage up to 5 mmol/kg (817 mg/kg).

1 In summary, although Nelson and Bull (1988) report effects on DNA unwinding for TCE  
2 and its metabolites with DCA having the highest activity and TCA the lowest, Nelson et al.  
3 (1989), using the same assay, reported no effect for TCA and the same effect at 10 and  
4 500 mg/kg for DCA in mice. Moreover, Styles et al. (1991) did not find a positive result for  
5 TCA using the same paradigm as Nelson and Bull (1988) and Nelson et al. (1989). Furthermore,  
6 Chang et al. (1992) also did not find increased single strand breaks for TCA exposure in rats.  
7 (see Section E.2.4.3).  
8

#### 4.2.2.1.7. Summary

9 In summary, TCA has been studied using a variety of genotoxicity assays, including the  
10 recommended battery. No mutagenicity was reported in *S. typhimurium* strains in the presence  
11 or absence of metabolic activation or in an alternative protocol using a closed system, except in  
12 one study on strain TA100 using a modified protocol in liquid medium. This is largely  
13 consistent with the results from TCE, which was negative in most bacterial systems except some  
14 studies with the TA100 strain. Mutagenicity in mouse lymphoma cells was only induced at  
15 cytotoxic concentrations. Measures of DNA-repair responses in bacterial systems have been  
16 inconclusive, with induction of DNA repair reported in *S. typhimurium* but not in *E. coli*.  
17 TCA-induced clastogenicity may be secondary to pH changes and not a direct effect of TCA.  
18

#### 4.2.3. Dichloroacetic Acid (DCA)

19 DCA is another metabolite of TCE that has been studied using a variety of genotoxicity  
20 assay for its genotoxic potential (see Tables 4-13 and 4-14; see IARC (2004a) for additional  
21 information).  
22

##### 4.2.3.1.1. Bacterial and Fungal Systems—Gene Mutations

23 Studies were conducted to evaluate mutagenicity of DCA in different *S. typhimurium* and  
24 *E. coli* strains (DeMarini et al., 1994; Fox et al., 1996a; Fox et al., 1996b; Giller et al., 1997;  
25 Herbert et al., 1980; Kargalioglu et al., 2002; Nelson et al., 2001; Waskell, 1978). DCA was  
26 mutagenic in three strains of *S. typhimurium*: strain TA100 in three of five studies, strain RSJ100  
27 in a single study, and strain TA98 in two of three studies. DCA failed to induce point mutations  
28 in other strains of *S. typhimurium* (TA104, TA1535, TA1537, and TA1538) or in *E. coli* strain

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 WP2uvrA. In one study, DCA caused a weak induction of SOS repair in *E. coli* strain PQ37  
2 (Giller et al., 1997).

3 DeMarini et al. (1994), in the same study as described in the TCA section of this section,  
4 also studied DCA as one of their compounds for analysis. In the prophage-induction assay using  
5 *E. coli*, DCA, in the presence of S9, was genotoxic producing 6.6–7.2 plaque-forming units  
6 (PFU)/mM and slightly less than threefold increase in PFU/plate in the absence of S9. In the  
7 second set of studies, which involved the evaluation of DCA at concentrations of 0–600 ppm for

**Table 4-13. Genotoxicity of dichloroacetic acid (bacterial systems)**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
$\lambda$ Prophage induction, <i>E. coli</i> WP2s	2,500	+	–	DeMarini et al. (1994)
SOS chromotest, <i>E. coli</i> PQ37	500	–	(+)	Giller et al. (1997)
<i>S. typhimurium</i> , DNA repair-deficient strains TS24, TA2322, TA1950	31,000	–	–	Waskell (1978)
<i>S. typhimurium</i> TA100, TA1535, TA1537, TA1538, reverse mutation		–	–	Herbert et al. (1980)
<i>S. typhimurium</i> TA100, reverse mutation	50	+	+	DeMarini et al. (1994)
<i>S. typhimurium</i> TA100,TA1535, TA1537, TA98, reverse mutation	5,000	–	–	Fox et al., (1996b)
<i>S. typhimurium</i> TA100, reverse mutation, liquid medium	100	+	+	Giller et al. (1997)
<i>S. typhimurium</i> RSJ100, reverse mutation	1,935	–	+	Kargalioglu et al. (2002)
<i>S. typhimurium</i> TA104, reverse mutation, microsuspension	150 $\mu$ g/plate	–	–	Nelson et al. (2001)
<i>S. typhimurium</i> TA98, reverse mutation	10 $\mu$ g/plate	(+)	–	Herbert et al. (1980)
	5,160	–	+	Kargalioglu et al. (2002)
<i>S. typhimurium</i> TA100, reverse mutation	1,935	+	+	Kargalioglu et al. (2002)
<i>E. coli</i> WP2uvrA, reverse mutation	5,000	–	–	Fox et al., (1996b)

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose; doses are in  $\mu$ g/mL for in vitro tests unless specified.

<sup>b</sup> Results: + = positive; (+) = weakly positive; – = negative.

Table adapted from IARC monograph (2004a) and modified/updated for newer references.

**Table 4-14. Genotoxicity of dichloroacetic acid—mammalian systems**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
Gene mutation, mouse lymphoma cell line L5178Y/TK ± in vitro	5,000	–	–	Fox et al., (1996b)
Gene mutation, mouse lymphoma cell line L5178Y/TK ± –3.7.2C in vitro	400	NT	+	Harrington-Brock et al. (1998)
Gene mutation, Chinese hamster ovary cells in vitro, HGPRT gene mutation assay	1,000 µM	NT	+	Zhang et al. (2010)
DNA strand breaks and alkali-labile damage, Chinese hamster ovary cells in vitro (single-cell gel electrophoresis assay)	3,225 µg/mL	NT	–	Plewa et al. (2002)
DNA strand breaks, B6C3F1 mouse hepatocytes in vitro	2,580	NT	–	Chang et al. (1992)
DNA strand breaks, Fischer 344 rat hepatocytes in vitro	1,290	NT	–	Chang et al. (1992)
Micronucleus formation, mouse lymphoma L5178Y/TK ± –3.7.2C cell line in vitro	800	NT	–	Harrington-Brock et al. (1998)
Chromosomal aberrations, Chinese hamster ovary in vitro	5,000	–	–	Fox et al., (1996b)
Chromosomal aberrations, mouse lymphoma L5178Y/Tk ± –3.7.2C cell line in vitro	600	NT	+	Harrington-Brock et al. (1998)
Aneuploidy, mouse lymphoma L5178Y/Tk ± –3.7.2C cell line in vitro	800	NT	–	Harrington-Brock et al. (1998)
DNA strand breaks, human CCRF-CEM lymphoblastoid cells in vitro	1,290	NT	–	Chang et al. (1992)
DNA strand breaks, male B6C3F1 mouse liver in vivo	13, oral, × 1	+		Nelson and Bull (1988)
DNA strand breaks, male B6C3F1 mouse liver in vivo	10, oral, × 1	+		Nelson et al. (1989)
DNA strand breaks, male B6C3F1 mouse liver in vivo	1,290, oral, × 1	–		Chang et al. (1992)
DNA strand breaks, male B6C3F1 mouse splenocytes in vivo	1,290, oral, × 1	–		Chang et al. (1992)
DNA strand breaks, male B6C3F1 mouse epithelial cells from stomach and duodenum in vivo	1,290, oral, × 1	–		Chang et al. (1992)
DNA strand breaks, male B6C3F1 mouse liver in vivo	5,000, dw, × 7–14 d	–		Chang et al. (1992)

DNA strand breaks, alkali-labile sites, cross linking, male B6C3F1 mouse blood leukocytes in vivo (single-cell gel electrophoresis assay)	3,500, dw, × 28 d	+	Fuscoe et al. (1996)
---	-------------------	---	----------------------

**Table 4-14. Genotoxicity of Dichloroacetic Acid—mammalian systems (continued)**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
DNA strand breaks, male Sprague-Dawley rat liver in vivo	30, oral, × 1	+		Nelson and Bull (1988)
DNA strand breaks, male Fischer 344 rat liver in vivo	645, oral, × 1	–		Chang et al. (1992)
DNA strand breaks, male Fischer 344 rat liver in vivo	2,000, dw, × 30 wk	–		Chang et al. (1992)
Gene mutation, lacI transgenic male B6C3F1 mouse liver assay in vivo	1,000, dw, × 60 wk	+		Leavitt et al. (1997)
Micronucleus formation, male B6C3F1 mouse peripheral erythrocytes in vivo	3,500, dw, × 9 d	+		Fuscoe et al. (1996)
Micronucleus formation, male B6C3F1 mouse peripheral erythrocytes in vivo	3,500, dw, × 28 d	–		Fuscoe et al. (1996)
Micronucleus formation, male B6C3F1 mouse peripheral erythrocytes in vivo	3,500, dw, × 10 wk	+		Fuscoe et al. (1996)
Micronucleus formation, male and female Crl:CD (S-D) BR rat bone-marrow erythrocytes in vivo	1,100, i.v., × 3	–		Fox et al., (1996b)
Micronucleus formation, Pleurodeles waltl newt larvae peripheral erythrocytes in vivo	80 d	–		Giller et al. (1997)

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose; doses are in µg/mL for in vitro tests; mg/kg for in vivo tests unless specified; dw = drinking-water (in mg/L); d = day; wk = week; i.v. = intravenous.

<sup>b</sup> Results: + = positive; - = negative; NT = not tested.

Table adapted from IARC monograph (2004a) and modified/updated for newer references.

1 mutagenicity in *S. typhimurium* TA100 strain, DCA was mutagenic both in the presence and  
2 absence of S9, producing three–five times increases in the revertants/plate compared to the  
3 background. The lowest effective concentration for DCA without S9 was 100 ppm and 50 ppm  
4 in the presence of S9. In the third and most important study, mutation spectra of DCA were  
5 determined at the base-substitution allele *hisG46* of *S. typhimurium* TA100. DCA-induced  
6 revertants were chosen for further molecular analysis at concentrations that produced mutant  
7 yields that were two–fivefold greater than the background. The mutation spectra of DCA were  
8 significantly different from the background mutation spectrum. Thus, despite the modest  
9 increase in the mutant yields (three–five times) produced by DCA, the mutation spectra confirm  
10 that DCA is mutagenic. DCA primarily induced GC-AT transitions.

11 Kargalioglu et al. (2002) analyzed the cytotoxicity and mutagenicity of the drinking  
12 water disinfection by-products including DCA in *S. typhimurium* strains TA98, TA100, and  
13 RSJ100 ± S9. DCA was mutagenic in this test although the response was low when compared to  
14 other disinfection by-products tested in strain TA100. This study was also summarized in a  
15 review by Plewa et al. (2002). Nelson et al. (2001) investigated the mutagenicity of DCA using  
16 a *S. typhimurium* microsuspension bioassay following incubation of DCA for various lengths of  
17 time, with or without rat cecal microbiota. No mutagenic activity was detected for DCA with  
18 *S. typhimurium* strain TA104.

19 Although limited data, it appears that DCA has mutagenic activity in the *S. typhimurium*  
20 strains, particularly TA100.

21

#### 4.2.3.1.2. Mammalian Systems

#### 4.2.3.1.3. Gene mutations

22 The mutagenicity of DCA has been tested in mammalian systems, particularly, mouse  
23 lymphoma cell lines in vitro (Fox et al., 1996b; Harrington-Brock et al., 1998); and *lacI*  
24 transgenic mice in vivo (Leavitt et al., 1997). Harrington-Brock et al. (1998) evaluated DCA for  
25 its mutagenic activity in L5178Y/Tk ± (-) 3.7.2C mouse lymphoma cells. A dose-related  
26 increase in mutation (and cytotoxic) frequency was observed at concentrations between 100 and  
27 800 µg/mL. Most mutagenic activity of DCA at the *Tk* locus was due to the production of  
28 small-colony *Tk* mutants (indicating chromosomal mutations). Different pH levels were tested  
29 in induction of mutant frequencies and it was determined that the mutagenic effect observed was  
30 due to the chemical and not pH effects.

1 Mutation frequencies were studied in male transgenic B6C3F1 mice harboring the  
2 bacterial *lacI* gene administered DCA at either 1.0 or 3.5 g/L in drinking water (Leavitt et al.,  
3 1997). No significant difference in mutant frequency was observed after 4 or 10 weeks of  
4 treatment in both the doses tested as compared to control. However, at 60 weeks, mice treated  
5 with 1.0 g/L DCA showed a slight increase (1.3-fold) in the mutant frequency over the control,  
6 but mice treated with 3.5 g/L DCA had a 2.3-fold increase in the mutant frequency. Mutational  
7 spectra analysis revealed that ~33% had G:C-A:T transitions and 21% had G:C-T:A  
8 transversions and this mutation spectra was different than that was seen in the untreated animals,  
9 indicating that the mutations were likely induced by the DCA treatment. The authors conclude  
10 that these results are consistent with the previous observation that the proportion of mutations at  
11 T:A sites in codon 61 of the H-ras gene was increased in DCA-induced liver tumors in B6C3F1  
12 mice (Leavitt et al., 1997).

13 Zhang et al. (2010) tested the cytotoxic and genotoxic effects of DCA in a  
14 microplate-based cytotoxicity test and HGPRT gene mutation assay using Chinese hamster ovary  
15 K1 cells, respectively. The concentrations at which these tests were conducted was 0, 200,  
16 1,000, 5,000 and 10,000  $\mu\text{M}$ . Two parameters were used to indicate chronic cytotoxicity: the  
17 lowest cytotoxic concentration and the percent C1/2 value. The lowest cytotoxic concentration  
18 for DCA was  $2.87 \times 10^{-3}\text{M}$ . Statistically significant increase in HGPRT mutant frequency was  
19 observed at concentration 1,000  $\mu\text{M}$  and above.

20

#### 4.2.3.1.4. Chromosomal aberrations and micronucleus

21 Harrington-Brock et al. (1998) evaluated DCA for its potential to induce chromosomal  
22 aberrations in DCA-treated (0, 600, and 800  $\mu\text{g/mL}$ ) mouse lymphoma cells. A clearly positive  
23 induction of aberrations was observed at both concentrations tested. No significant increase in  
24 micronucleus was observed in DCA-treated (0, 600, and 800  $\mu\text{g/mL}$ ) mouse lymphoma cells  
25 (Harrington-Brock et al., 1998). However, no chromosomal aberrations were found in Chinese  
26 hamster ovary cells exposed to DCA (Fox et al., 1996b)

27 Fuscoe et al. (1996) investigated in vivo genotoxic potential of DCA in bone marrow and  
28 blood leukocytes using the peripheral-blood-erythrocyte micronucleus assay (to detect  
29 chromosome breakage and/or malsegregation) and the alkaline single cell gel electrophoresis  
30 (comet) assay, respectively. Mice were exposed to DCA in drinking water, available *ad libitum*,  
31 for up to 31 weeks. A statistically significant dose-related increase in the frequency of  
32 micronucleated PCEs was observed following subchronic exposure to DCA for 9 days.  
33 Similarly, a significant increased was also observed when exposed for  $\geq 10$  weeks particularly at

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 the highest dose of DCA tested (3.5 g/L). DNA cross-linking was observed in blood leukocytes  
2 in mice exposed to 3.5 g/L DCA for 28 days. These data provide evidence that DCA may have  
3 some potential to induce chromosome damage when animals were exposed to concentrations  
4 similar to those used in the rodent bioassay.

#### 4.2.3.1.5. Other DNA damage studies

5 Nelson and Bull (1988) and Nelson et al. (1989) have been described above in  
6 Section 4.2.2.4 and E.2.3, with positive results for DNA unwinding for DCA, though Nelson  
7 et al. (1989) reported the same response at 10 and 500 mg/kg in mice, raising concerns about the  
8 reliability of the assay in these studies. Chang et al. (1992) conducted both in vitro and in vivo  
9 studies to determine the ability of DCA to cause DNA damage. Primary rat (Fischer 344)  
10 hepatocytes and primary mouse hepatocytes treated with DCA for 4 hours did not induce  
11 DNA single strand breaks as detected by alkaline DNA unwinding assay. No DNA strand breaks  
12 were observed in human CCRF-CEM lymphoblastoid cells in vitro exposed to DCA. Similarly,  
13 analysis of the DNA single strand breaks in mice killed 1 hour after a single dose of 1, 5 or  
14 10 mM/kg DCA did not cause DNA damage. None of the Fischer 344 rats killed 4 hours after a  
15 single gavage treatment (1–10 mM/kg) produced any detectable DNA damage.

16

#### 4.2.3.1.6. Summary

17 In summary, DCA has been studied using a variety but limited number of genotoxicity  
18 assays. Within the available data, DCA has been demonstrated to be mutagenic in the  
19 *S. typhimurium* assay, particularly in strain TA100, the in vitro mouse lymphoma assay and in  
20 vivo cytogenetic and gene mutation assays. DCA can cause DNA strand breaks in mouse and rat  
21 liver cells following in vivo administration by gavage.

22

#### 4.2.4. Chloral Hydrate

23 Chloral hydrate has been evaluated for its genotoxic potential using a variety of  
24 genotoxicity assays (see Tables 4-15, 4-16, and 4-17). These data are particularly important  
25 because it is known that a large flux of TCE metabolism leads to chloral hydrate as an  
26 intermediate, so a comparison of their genotoxicity profiles is likely to be highly informative.

27

#### 4.2.4.1.1. DNA Binding Studies

1 Limited analysis has been performed examining DNA binding potential of chloral  
2 hydrate (Keller and Hd'A, 1988; Ni et al., 1995; Von Tungeln et al., 2002). Keller and Heck  
3 (1988) conducted both in vitro and in vivo experiments using B6C3F1 mouse strain. The mice  
4 were pretreated with 1,500 mg/kg TCE for 10 days and then given 800 mg/kg [<sup>14</sup>C] chloral. No  
5 detectable covalent binding of [<sup>14</sup>C] to DNA in the liver was observed. Another study with in  
6 vivo exposures to nonradioactive chloral hydrate at a concentration of 1,000 and 2,000 nmol in  
7 mice B6C3F1 demonstrated an increase in malondialdehyde-derived and 8-oxo-2'-  
8 deoxyguanosine adducts in liver DNA (Von Tungeln et al., 2002). Ni et al. (1995) observed

**Table 4-15. Chloral hydrate genotoxicity: bacterial, yeast and fungal systems**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
SOS chromotest, <i>Escherichia coli</i> PQ37	10,000	-	-	Giller et al. (1995)
<i>S. typhimurium</i> TA100, TA1535, TA98, reverse mutation	10,000	-	-	Waskell (1978)
<i>S. typhimurium</i> TA100, TA1537, TA1538, TA98, reverse mutation	1,000	+	+	Haworth et al. (1983)
<i>S. typhimurium</i> TA100, reverse mutation	5,000 µg/plate	-	-	Leuschner and Leuschner (1991)
<i>S. typhimurium</i> TA100, reverse mutation	2,000 µg/plate	+	+	Ni et al. (1994)
<i>S. typhimurium</i> TA100, reverse mutation, liquid medium	300	+	-	Giller et al. (1995)
<i>S. typhimurium</i> TA100, TA104, reverse mutation	1,000 µg/plate	+	+	Beland (1999)
<i>S. typhimurium</i> TA104, reverse mutation	1,000 µg/plate	+	+	Ni et al. (1994)
<i>S. typhimurium</i> TA1535, reverse mutation	1,850	-	-	Leuschner and Leuschner (1991)
<i>S. typhimurium</i> TA1535, TA1537 reverse mutation	6,667	-	-	Haworth et al. (1983)
<i>S. typhimurium</i> TA1535, reverse mutation	10,000	-	-	Beland (1999)
<i>S. typhimurium</i> TA98, reverse mutation	7,500	-	-	Haworth et al. (1983)
<i>S. typhimurium</i> TA98, reverse mutation	10,000 µg/plate	-	+	Beland (1999)
<i>A.nidulans</i> , diploid strain 35X17, mitotic cross-overs	1,650	NT	-	Crebelli et al. (1985)
<i>A. nidulans</i> , diploid strain 30, mitotic cross-overs	6,600	NT	-	Kafer (1986)
<i>A. nidulans</i> , diploid strain NH, mitotic cross-overs	1,000	NT	-	Kappas (1989)
<i>A. nidulans</i> , diploid strain P1, mitotic cross-overs	990	NT	-	Crebelli et al. (1991)
<i>A. nidulans</i> , diploid strain 35X17, nondisjunctions	825	NT	+	Crebelli et al. (1985)
<i>A. nidulans</i> , diploid strain 30, aneuploidy	825	NT	+	Kafer (1986)
<i>A. nidulans</i> , haploid conidia, aneuploidy, polyploidy	1,650	NT	+	Kafer (1986)
<i>A. nidulans</i> , diploid strain NH, nondisjunctions	450	NT	+	Kappas (1989)

**Table 4-15. Chloral hydrate genotoxicity: bacterial, yeast and fungal systems (continued)**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
<i>A. nidulans</i> , diploid strain P1, nondisjunctions	660	NT	+	Crebelli et al. (1991)
<i>A. nidulans</i> , haploid strain 35, hyperploidy	2,640	NT	+	Crebelli et al. (1991)
<i>S. cerevisiae</i> , meiotic recombination	3,300	NT	?	Sora and Agostini Carbone (1987)
<i>S. cerevisiae</i> , disomy in meiosis	2,500	NT	+	Sora and Agostini Carbone (1987)
<i>S. cerevisiae</i> , disomy in meiosis	3,300	NT	+	Sora and Agostini Carbone (1987)
<i>S. cerevisiae</i> , D61.M, mitotic chr. malsegregation	1,000	NT	+	Albertini, (1990)
<i>Drosophila melanogaster</i> , somatic mutation wing spot test	825		+	Zordan et al. (1994)
<i>Drosophila melanogaster</i> , induction of sex-linked lethal mutation	37.2 feed		?	Beland (1999)
<i>Drosophila melanogaster</i> , induction of sex-linked lethal mutation	67.5 inj		-	Beland (1999)

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose; doses are in µg/mL for in vitro tests; inj = injection.

<sup>b</sup> Results: + = positive; - = negative; NT = not tested; ? = inconclusive.

Table adapted from IARC monograph (2004a) and modified/updated for newer references.

**Table 4-16.. Chloral hydrate genotoxicity: mammalian systems—all genetic endpoints, in vitro**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
DNA-protein cross-links, rat nuclei in vitro	41,250	NT	-	Keller and Heck (1988)
DNA single-strand breaks, rat primary hepatocytes in vitro	1,650	NT	-	Chang et al. (1992)
Gene mutation, mouse lymphoma L5178Y/TK <sup>±</sup> , in vitro	1,000		(+)	Harrington-Brock et al. (1998)
Sister chromatid exchange, CHO cells, in vitro	100	+	+	Beland (1999)
Micronucleus formation, (kinetochore-positive), Chinese hamster C1 cells, in vitro	165	NT	+	Degrassi and Tanzarella (1988)
Micronucleus formation, (kinetochore-negative), Chinese hamster C1 cells, in vitro	250	NT	-	Degrassi and Tanzarella (1988)
Micronucleus formation, (kinetochore-positive), Chinese hamster LUC2 cells, in vitro	400	NT	+	Parry et al., (1990)
Micronucleus formation, (kinetochore-positive), Chinese hamster LUC2 cells, in vitro	400	NT	+	Lynch and Parry (1993)
Micronucleus formation, Chinese hamster V79 cells, in vitro	316	NT	+	Seelbach et al. (1993)
Micronucleus formation, mouse lymphoma L5178Y/TK <sup>±</sup> , in vitro	1,300	NT	-	Harrington-Brock et al. (1998)
Micronucleus formation, mouse lymphoma L5178Y/TK <sup>±</sup> , in vitro	500	NT	+	Nesslany and Marzin (1999)
Chromosomal aberrations, Chinese Hamster CHED cells, in vitro	20	NT	+	Furnus et al. (1990)
Chromosomal aberrations, Chinese Hamster ovary cells, in vitro	1,000	+	+	Beland (1999)
Chromosomal aberrations, mouse lymphoma L5178Y/TK <sup>±</sup> cells line, in vitro	1,250	NT	(+)	Harrington-Brock et al. (1998)
Aneuploidy, Chinese hamster CHED cells, in vitro	10	NT	+	Furnus et al. (1990)
Aneuploidy, primary Chinese hamster embryonic cells, in vitro	250	NT	+	Natarajan et al. (1993)
Aneuploidy, Chinese hamster LUC2p4 cells, in vitro	250	NT	+	Warr et al. (1993)
Aneuploidy, mouse lymphoma L5178Y/TK <sup>±</sup> , in vitro	1,300	NT	-	Harrington-Brock et al. (1998)
Tetraploidy and endoreduplication, Chinese hamster LUC2p4cells, in vitro	500	NT	+	Warr et al. (1993)
Cell transformation, Syrian hamster embryo cells (24-h treatment)	350	NT	+	Gibson et al. (1995)
Cell transformation, Syrian hamster dermal cell line (24-h treatment)	50	NT	+	Parry et al. (1996)
DNA single-strand breaks, human lymphoblastoid cells, in vitro	1,650	NT	-	Chang et al. (1992)

**Table 4-16. Chloral hydrate genotoxicity: mammalian systems—all genetic endpoints, in vitro (continued)**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
Gene mutation, <i>tk</i> and <i>hprt</i> locus, human lymphoblastoid	1,000	NT	+	Beland (1999)
Sister chromatid exchanges, human lymphocytes, in vitro	54	NT	(+)	Gu et al. (1981a)
Micronucleus formation, human lymphocytes, in vitro	100	-	+	Van Hummelen and Kirsch-Volders (1992)
Micronucleus formation, human lymphoblastoid AHH-1 cell line, in vitro	100	NT	+	Parry et al. (1996)
Micronucleus formation, human lymphoblastoid MCL-5 cell line, in vitro	500	NT	-	Parry et al. (1996)
Micronucleus formation (kinetochore-positive), human diploid LEO fibroblasts, in vitro	120	NT	+	Bonatti et al. (1992)
Aneuploidy (double Y induction), human lymphocytes, in vitro	250	NT	+	Vagnarelli et al. (1990)
Aneuploidy (hyperdiploidy and hypodiploidy), human lymphocytes in vitro	50	NT	+	Sbrana et al. (1993)
Polyploidy, human lymphocytes, in vitro	137	NT	+	Sbrana et al. (1993)
C-Mitosis, human lymphocytes, in vitro	75	NT	+	Sbrana et al. (1993)

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose; doses are in µg/mL for in vitro tests.

<sup>b</sup> Results: + = positive; (+) = weakly positive in an inadequate study; - = negative; NT = not tested.

Table adapted from IARC monograph (2004a) and modified/updated for newer references.

**Table 4-17.. Chloral hydrate genotoxicity: mammalian systems—all genetic damage, in vivo**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>	Reference
DNA single-strand breaks, male Sprague-Dawley rat liver	300, oral	+	Nelson and Bull (1988)
DNA single-strand breaks, male Fischer 344 rat liver	1,650, oral	-	Chang et al. (1992)
DNA single-strand breaks, male B6C3F1 mouse liver	100, oral	+	Nelson and Bull (1988)
DNA single-strand breaks, male B6C3F1 mouse liver	825, oral	-	Chang et al. (1992)
Micronucleus formation, male and female NMRI mice, bone-marrow erythrocytes	500, i.p.	-	Leuschner and Leuschner (1991)
Micronucleus formation, BALB/c mouse spermatids	83, i.p.	-	Russo and Levis, 1992
Micronucleus formation, male BALB/c mouse bone-marrow erythrocytes and early spermatids	83, i.p.	+	Russo and Levis, 1992
Micronucleus formation, male BALB/c mouse bone-marrow erythrocytes	200, i.p.	+	Russo et al. (1992)
Micronucleus formation, male F1 mouse bone-marrow erythrocytes	400, i.p.	-	Leopardi et al. (1993)
Micronucleus formation, C57B1 mouse spermatids	41, i.p.	+	Allen et al., 1994
Micronucleus formation, male Swiss CD-1 mouse bone-marrow erythrocytes	200, i.p.	+	Marrazzini et al., (1994)
Micronucleus formation, B6C3F1 mouse spermatids after spermatogonial stem-cell treatment	165, i.p.	+	Nutley et al. (1996)
Micronucleus formation, B6C3F1 mouse spermatids after meiotic cell treatment	413, i.p.	-	Nutley et al. (1996)
Micronucleus formation, male F1, BALB/c mouse peripheral-blood erythrocytes	200, i.p.	—	Grawe et al. (1997)
Micronucleus formation, male B6C3F1 mouse bone-marrow erythrocytes	500, i.p., × 3	+	Beland (1999)
Micronucleus formation, infants, peripheral lymphocytes	50, oral	+	Ikbal et al. (2004)
Chromosomal aberrations, male and female F1 mouse bone marrow cells	600, i.p.	-	Xu and Alder (1990)
Chromosomal aberrations, male and female Sprague-Dawley rat bone-marrow cells	1,000, oral	—	Leuschner and Leuschner (1991)
Chromosomal aberrations, BALB/c mouse spermatogonia treated	83, i.p.	-	Russo and Levis, (1992b)
Chromosomal aberrations, F1 mouse secondary spermatocytes	82.7, i.p.	+	Russo et al. (1984)
Chromosomal aberrations, male Swiss CD-1 mouse bone-marrow erythrocytes	400, i.p.	—	Marrazzini et al. (1994)
Chromosomal aberrations, ICR mouse oocytes	600, i.p.	-	Mailhes et al. (1993)

**Table 4-17. Chloral hydrate genotoxicity: mammalian systems—all genetic damage, in vivo (continued)**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>	Reference
Micronucleus formation, infants, peripheral lymphocytes	50, oral	+	Ikbal et al. (2004)
Polyploidy, male and female F1, mouse bone-marrow cells	600, i.p.	-	Xu and Alder (1990)
Aneuploidy F1 mouse secondary spermatocytes	200, i.p.	+	Miller and Adler (1992)
Aneuploidy, male F1 mouse secondary spermatocytes	400, i.p.	-	Leopardi et al. (1993)
Hyperploidy, male Swiss CD-1 mouse bone-marrow erythrocytes	200, i.p.	+	Marrazzini et al., 1994

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose; doses are in mg/kg BW for in vivo tests, i.p. = intraperitoneally.

<sup>b</sup> Results: + = positive; - = negative.

Table adapted from IARC monograph (2004a) and modified/updated for newer references.



1 malondialdehyde adducts in calf thymus DNA when exposed to chloral hydrate and microsomes  
2 from male B6C3F1 mouse liver.

3 Keller and Heck (1988) investigated the potential of chloral to form DNA-protein  
4 cross-links in rat liver nuclei using concentrations 25, 100, or 250 mM. No statistically  
5 significant increase in DNA-protein cross-links was observed. DNA and RNA isolated from the  
6 [<sup>14</sup>C] chloral-treated nuclei did not have any detectable [<sup>14</sup>C] bound. However, the proteins from  
7 chloral-treated nuclei did have a concentration-related binding of [<sup>14</sup>C].  
8

#### 4.2.4.1.2. Bacterial and Fungal Systems—Gene Mutations

9 Chloral hydrate induced gene mutations in *S. typhimurium* TA100 and TA104 strains, but  
10 not in most other strains assayed. Four of six studies of chloral hydrate exposure in  
11 *S. typhimurium* TA100 and two of two studies in *S. typhimurium* TA104 were positive for  
12 revertants (Beland, 1999; Giller et al., 1995; Haworth et al., 1983; Ni et al., 1994). Waskell  
13 (1978) studied the effect of chloral hydrate along with TCE and its other metabolites. Chloral  
14 hydrate was tested at different doses (1.0–13 mg/plate) in different *S. typhimurium* strains  
15 (TA98, TA100, TA1535) for gene mutations using Ames assay. No revertant colonies were  
16 observed in strains TA98 or TA1535 both in the presence and absence of S9 mix. Similar results  
17 were obtained by Leuschner and Leuschner (1991). However, in TA100, a dose-dependent  
18 statistically significant increase in revertant colonies was obtained both in the presence and  
19 absence of S9. It should be noted that chloral hydrate that was purchased from Sigma was  
20 recrystallized from one to six times from chloroform and the authors describe this as crude  
21 chloral hydrate. However, this positive result is consistent with other studies in this strain as  
22 noted above. Furthermore, Giller et al. (1995) studied chloral hydrate genotoxicity in  
23 three short-term tests. Chloral-induced mutations in strain TA100 of *S. typhimurium* (fluctuation  
24 test). Similar results were obtained by Haworth et al. (1983). These are consistent with several  
25 studies of TCE, in which low, but positive responses were observed in the TA100 strain in the  
26 presence of S9 metabolic activation, even when genotoxic stabilizers were not present.

27 A significant increase in mitotic segregation was observed in *Aspergillus nidulans* when  
28 exposed to 5 and 10 mM chloral hydrate (Crebelli et al., 1985). Studies of mitotic crossing-over  
29 in *Aspergillus nidulans* have been negative while these same studies were positive for  
30 aneuploidy (Crebelli et al., 1985, 1991; Kafer, 1986; Kappas, 1989).

31 Two studies were conducted in *Saccharomyces cerevisiae* to understand the  
32 chromosomal malsegregation as a result of exposure to chloral hydrate (Albertini, 1990; Sora  
33 and Agostini Carbone, 1987). Chloral hydrate (1–25 mM) was dissolved in sporulation medium

1 and the frequencies of various meiotic events such as recombination, disomy were analyzed.  
2 Chloral hydrate inhibited sporulation as a function of dose and increased diploid and disomic  
3 clones . Chloral hydrate was also tested for mitotic chromosome malsegregation using  
4 *Saccharomyces cerevisiae* D61.M (Albertini, 1990). The tester strain was exposed to a dose  
5 range of 1–8 mg/mL. An increase in the frequency of chromosomal malsegregation was  
6 observed as a result of exposure to chloral hydrate.

7 Limited analysis of chloral hydrate mutagenicity has been performed in *Drosophila*  
8 (Beland, 1999; Zordan et al., 1994). Of these two studies, chloral hydrate was positive in the  
9 somatic mutation wing spot test (Zordan et al., 1994), equivocal in the induction of sex-linked  
10 lethal mutation when in feed but negative when exposed via injection (Beland, 1999).  
11

#### 4.2.4.1.3. Mammalian Systems

##### 4.2.4.1.4. Gene mutations

12 Harrington-Brock et al. (1998) noted that chloral hydrate-induced concentration related  
13 cytotoxicity in TK± mouse lymphoma cell lines without S9 activation. A nonstatistical increase  
14 in mutant frequency was observed in cells treated with chloral hydrate. The mutants were  
15 primarily small colony TK mutants, indicating that most chloral hydrate-induced mutants  
16 resulted from chromosomal mutations rather than point mutations. It should be noted that in  
17 most concentrations tested (350–1,600 µg/mL), cytotoxicity was observed. Percentage cell  
18 survival ranged from 96–4%.  
19

##### 4.2.4.1.5. Micronucleus

20 Micronuclei induction following exposure to chloral hydrate is positive in most test  
21 systems in both in vitro and in vivo assays, although some negative tests do also exist (Allen et  
22 al., 1994; Harrington-Brock et al., 1998; Marrazzini et al., 1994) (Beland, 1999; Bonatti et al.,  
23 1992; Degrassi and Tanzarella, 1988; Giller et al., 1995; Grawé et al., 1997; Ikbali et al., 2004;  
24 Leopardi et al., 1993; Leuschner and Leuschner, 1991; Lynch and Parry, 1993; Nesslany and  
25 Marzin, 1999; Nutley et al., 1996; Parry et al., 1996; Russo and Levis, 1992a; Russo and Levis,  
26 1992b; Russo et al., 1992; Seelbach et al., 1993; Van Hummelen and Kirsch-Volders, 1992).  
27 Some studies have attempted to make inferences regarding aneuploidy induction or  
28 clastogenicity as an effect of chloral hydrate. Aneuploidy results from defects in chromosome

1 segregation during mitosis and is a common cytogenetic feature of cancer cells (see  
2 Section E.3.1.5).

3 Giller et al. (1995) studied chloral hydrate genotoxicity in three short-term tests. Chloral  
4 hydrate caused a significant increase in the frequency of micronucleated erythrocytes following  
5 in vivo exposure of the amphibian *Pleurodeles waltl* newt larvae.

6 Chloral hydrate induced aneuploidy in vitro in multiple Chinese hamster cell lines  
7 (Furnus et al., 1990; Natarajan et al., 1993; Warr et al., 1993) and human lymphocytes (Sbrana et  
8 al., 1993; Vagnarelli et al., 1990) but not mouse lymphoma cells Harrington-Brock et al. (1998).  
9 In vivo studies performed in various mouse strains led to increased aneuploidy in spermatocytes  
10 (Liang and Pacchierotti, 1988; Miller and Adler, 1992; Russo et al., 1984) but not oocytes  
11 (Mailhes et al., (1993)) or bone marrow cells (Leopardi et al., 1993; Xu and Adler, 1990).

12 The potential of chloral hydrate to induce aneuploidy in mammalian germ cells has been  
13 of particular interest since Russo et al. (1984) first demonstrated that chloral hydrate treatment of  
14 male mice results in significant increase in frequencies of hyperploidy in metaphase II cells.  
15 This hyperploidy was thought to have arisen from chromosomal nondisjunction in  
16 premeiotic/meiotic cell division and may be a consequence of chloral hydrate interfering with  
17 spindle formation [reviewed by Russo et al. (1984) and Liang and Brinkley (1985)]. Chloral  
18 hydrate also causes meiotic delay, which may be associated with aneuploidy (Miller and Adler,  
19 1992). Chloral hydrate has been shown to induce micronuclei but not structural chromosomal  
20 aberrations in mouse bone-marrow cells. Micronuclei induced by nonclastogenic agents are  
21 generally believed to represent intact chromosomes that failed to segregate into either  
22 daughter-cell nucleus at cell division (Russo et al., 1992; Xu and Adler, 1990). Furthermore,  
23 chloral hydrate-induced micronuclei in mouse bone-marrow cells (Russo et al., 1992) and in  
24 cultured mammalian cells (Bonatti et al., 1992; Degrassi and Tanzarella, 1988) have shown to be  
25 predominantly kinetochore-positive in composition upon analysis with immunofluorescent  
26 methods. The presence of a kinetochore in a micronucleus is considered evidence that the  
27 micronucleus contains a whole chromosome lost at cell division (Degrassi and Tanzarella, 1988;  
28 Eastmond and Tucker, 1989; Hennig et al., 1988). Therefore, both TCE and chloral hydrate  
29 appear to increase the frequency of micronuclei.

30 Allen et al. (1994) treated male C57B1/6J mice were given a single intraperitoneal  
31 injection of 0, 41, 83, or 165 mg/kg chloral hydrate. Spermatids were harvested at 22 hours, 11,  
32 13.5, and 49 days following exposure (Allen et al., 1994). Harvested spermatids were processed  
33 to identify both kinetochore-positive micronucleus (aneugen) and kinetochore-negative  
34 micronucleus (clastogen). All chloral hydrate doses administered 49 days prior to cell harvest  
35 were associated with significantly increased frequencies of kinetochore-negative micronuclei in

1 spermatids, however, dose dependence was not observed. This study is in contrast with other  
2 studies (Bonatti et al., 1992; Degrassi and Tanzarella, 1988) who demonstrated predominantly  
3 kinetochore-positive micronucleus.

4 The ability of chloral hydrate to induce aneuploidy and polyploidy was tested in human  
5 lymphocyte cultures established from blood samples obtained from two healthy nonsmoking  
6 donors (Sbrana et al., 1993). Cells were exposed for 72 and 96 hours at doses between 50 and  
7 250 µg/mL. No increase in percentage hyperdiploid, tetraploid, or endoreduplicated cells were  
8 observed when cells were exposed to 72 hours at any doses tested. However, at 96 hours of  
9 exposure, significant increase in hyperdiploid was observed at one dose (150 µg/mL) and was  
10 not dose dependent. Significant increase in tetraploid was observed at dose 137 mg/mL, again,  
11 no dose dependence was observed.

12 Ikbal et al. (2004) assessed the genotoxic effects in cultured peripheral blood  
13 lymphocytes of 18 infants (age range of 31–55 days) before and after administration of a single  
14 dose of chloral hydrate (50 mg/kg of body weight) for sedation before a hearing test for  
15 micronucleus frequency. A significant increase in micronuclei frequency was observed after  
16 administration of chloral hydrate.

#### 17 18 **4.2.4.1.6. Chromosomal aberrations**

19 Several studies have included chromosomal aberration analysis in both in vitro and in  
20 vivo systems exposed to chloral hydrate and have resulted in positive in in vitro  
21 studies—although not all studies had statistically significant increase Harrington-Brock et al.  
22 (1998) (Beland, 1999; Furnus et al., 1990).

23 Analysis of chloral hydrate treated mouse lymphoma cell lines for chromosomal  
24 aberrations resulted in a nonsignificant increase in chromosomal aberrations Harrington-Brock  
25 et al. (1998). However, it should be noted that the concentrations tested (1,250 and  
26 1,300 µg/mL) were cytotoxic (with a cell survival of 11 and 7%, respectively). Chinese hamster  
27 embryo cells were also exposed to 0.001, 0.002, and 0.003% chloral hydrate for 1.5 hours  
28 (Furnus et al., 1990). A nonstatistically significant increase in frequency of chromosomal  
29 aberrations was observed only 0.002 and 0.003% concentrations, with the increase not  
30 dose-dependent. In this study, it should be noted that the cells were only exposed for 1.5 hours  
31 to chloral hydrate and cells were allowed to grow for 48 hours (two cell cycles) to obtain similar  
32 mitotic index before analyzing for chromosomal aberrations. No information on cytotoxicity  
33 was provided except that higher doses decreased the frequency of mitotic cells at the time of  
fixation.

1 In vivo chromosome aberration studies have mostly reported negative or null results  
2 (Leuschner and Leuschner, 1991; Liang and Pacchierotti, 1988; Mailhes et al., 1993; Russo and  
3 Levis, 1992a; Russo and Levis, 1992b; Xu and Adler, 1990) with the exception of one study  
4 (Russo et al., 1984) in an F1 cross of mouse strain between C57B1/Cne × C3H/Cne.

#### 4.2.4.1.7. Sister chromatid exchanges (SCEs)

5 SCEs were assessed by Ikbal et al. (2004) in cultured peripheral blood lymphocytes of  
6 18 infants (age range of 31–55 days) before and after administration of a single dose of chloral  
7 hydrate (50 mg/kg of body weight) for sedation before a hearing test. The authors report a  
8 significant increase in the mean number of SCEs, from before administration ( $7.03 \pm 0.18$   
9 SCEs/cell) and after administration ( $7.90 \pm 0.19$  SCEs/cell) , with each of the 18 individuals  
10 showing an increase with treatment. Micronuclei were also significantly increased. SCEs were  
11 also assessed by Gu et al. (1981a) in human lymphocytes exposed in vitro with inconclusive  
12 results, although positive results were observed by Beland (1999) in Chinese hamster ovary cells  
13 exposed in vitro with and without an exogenous metabolic system.

#### 4.2.4.1.8. Cell transformation

15 Chloral hydrate was positive in the two studies designed to measure cellular  
16 transformation (Gibson et al., 1995; Parry et al., 1996). Both studies exposed Syrian hamster  
17 cells (embryo and dermal) to chloral hydrate and induced cellular transformation.

#### 4.2.4.1.9. Summary

19 Chloral hydrate has been reported to induce micronuclei formation, aneuploidy, and  
20 mutations in multiple in vitro systems and in vivo. In vivo studies have limited results to an  
21 increased micronuclei formation mainly in mouse spermatocytes. CH is positive to in some  
22 studies in in vitro genotoxicity assays that detect point mutations, micronuclei induction,  
23 chromosomal aberrations, and/or aneuploidy. The in vivo data exhibit mixed results (Allen et  
24 al., 1994) (Adler, 1993; Leuschner and Beuscher, 1998; Mailhes et al., 1993; Nutley et al., 1996;  
25 Russo et al., 1992; Xu and Adler, 1990). Most of the positive studies show that chloral hydrate  
26 induces aneuploidy. Based on the existing array of data, CH has the potential to be genotoxic,  
27 particularly when aneuploidy is considered in the weight of evidence for genotoxic potential.  
28 Some have suggested that chloral hydrate may act through a mechanism of spindle poisoning and

1 resulting in numerical changes in the chromosomes, but some data also suggest induction of  
2 chromosomal aberrations. These results are consistent with TCE, albeit there are more limited  
3 data on TCE for these genotoxic endpoints.  
4

#### 4.2.5. Dichlorovinyl Cysteine (DCVC) and S-Dichlorovinyl Glutathione (DCVG)

5 DCVC and DCVG have been studied for their genotoxic potential; however, since there  
6 is limited number of studies to evaluate them based on each endpoint, particularly in mammalian  
7 systems, the following section has been combined to include all the available studies for different  
8 endpoints of genotoxicity. Study details can be found in Table 4-18.

9 DCVC and DCVG, cysteine intermediates of TCE formed by the glutathione-  
10 S-transferase (GST) pathway, are capable of inducing point mutations as evidenced by the fact  
11 that they are positive in the Ames assay. Dekant et al. (1986c) demonstrated mutagenicity of  
12 DCVC in *S. typhimurium* strains (TA100, TA2638, and TA98) using the Ames assay in the  
13 absence of S9. The effects were decreased with the addition of a beta-lyase inhibitor  
14 aminooxyacetic acid, suggesting that bioactivation by this enzyme plays a role in genotoxicity.  
15 Vamvakas et al. (1987) tested *N*-acetyl-S-(1,2-dichlorovinyl)-L-cysteine (NAcDCVC) for  
16 mutagenicity following addition of rat kidney cytosol and found genotoxic activity.  
17 Furthermore, Vamvakas (1988b), in another experiment, investigated the mutagenicity of DCVG  
18 and DCVC in *S. typhimurium* strain TA2638, using kidney subcellular fractions for metabolic  
19 activation and AOAA (a beta-lyase inhibitor) to inhibit genotoxicity. DCVG and DCVC both  
20 exhibited direct-acting mutagenicity, with kidney mitochondria, cytosol, or microsomes  
21 enhancing the effects for both compounds and AOAA diminishing, but not abolishing the  
22 effects. Importantly, addition of liver subcellular fractions did not enhance the mutagenicity of  
23 DCVG, consistent with in situ metabolism playing a significant role in the genotoxicity of these  
24 compounds in the kidney.

25 While additional data are not available on DCVG or NAcDCVC, the genotoxicity of  
26 DCVC is further supported by the predominantly positive results in other available in vitro and  
27 in vivo assays. Jaffe et al. (1985) reported DNA strand breaks due to DCVC administered in  
28 vivo, in isolated perfused kidneys, and in isolated proximal tubules of albino male rabbits.  
29 Vamvakas et al. (1989) reported dose-dependent increases in unscheduled DNA synthesis in  
30 LLC-PK1 cell clones at concentrations without evidence of cytotoxicity. In addition,  
31 Vamvakas et al. (1996) reported that 7-week DCVC exposure to LLC-PK1 cell clones at  
32 noncytotoxic concentrations induces morphological and biochemical de-differentiation that  
33 persists for at least 30 passages after removal of the compound. This study also reported

1 increased expression of the proto-oncogene *c-fos* in the cells in this system. In a Syrian hamster  
2 embryo fibroblast system, DCVC did not induce micronuclei, but demonstrated an unscheduled  
3 DNA synthesis response (Vamvakas et al., 1988a).

4 Two more recent studies are discussed in more detail. Mally et al. (2006) isolated  
5 primary rat kidney epithelial cells from *Tsc-2<sup>E<sup>k</sup>/+</sup>* (Eker) rats, and reported increased  
6 transformation when exposed to 10 μM DCVC, similar to that of the genotoxic renal carcinogens  
7 *N*-methyl-*N*'-nitro-*N*-nitrosoguanidine (Horesovsky et al., 1994). The frequency was variable  
8 but consistently higher than background. No loss-of-heterozygosity (LOH) of the *Tsc-2* gene

**Table 4-18. TCE GSH conjugation metabolites genotoxicity**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<b>Gene mutations (Ames test)</b>					
<i>S. typhimurium</i> , TA100, 2638, 98	0.1–0.5 nmol	ND	+	DCVC was mutagenic in all three strains of <i>S. typhimurium</i> without the addition of mammalian subcellular fractions.	Dekant et al., (1986c)
<i>S. typhimurium</i> , TA2638	50–300 nmol	+	+	Increase in number of revertants in DCVC alone at low doses; further increase in revertants was observed in the presence of microsomal fractions. Toxicity as indicated by decreased revertants per plate were seen at higher doses.	Vamvakas et al. (1988b)
<b>Mutation analysis</b>					
In vitro—rat kidney epithelial cells, LOH in <i>Tsc</i> gene	10 µM	NA	-	Only 1/9 transformed cells showed LOH.	Mally et al. (2006)
In vitro—rat kidney epithelial cells, <i>VHL</i> gene (exons 1–3)	10 µM	NA	-	No mutations in <i>VHL</i> gene. <u>Note:</u> <i>VHL</i> is not a target gene in rodent models of chemical-induced or spontaneous renal carcinogenesis.	Mally et al. (2006)
<b>Unscheduled DNA synthesis</b>					
Porcine kidney tubular epithelial cell line (LLC-PK1)	2.5 µM–5, 10, 15, 24 h; 2.5–100 µM	NA	+	Dose-dependent in UDS up to 24 h tested at 2.5 µM. Also, there was a dose dependent increase at lower conc. Higher concentrations were cytotoxic as determined by LDH release from the cells.	Vamvakas et al. (1989)
Syrian hamster embryo fibroblasts		NA	+	Increase in UDS in treatment groups.	Vamvakas et al. (1988a)



**Table 4-18. TCE GSH conjugation metabolites genotoxicity (continued)**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<b>DNA strand breaks</b>					
Male rabbit renal tissue (perfused kidneys and proximal tubules)	0–100 mg/kg or 10 µM to 10 mM	ND	+	Dose dependent increase in strand breaks in both i.v. and i.p. injections (i.v. injections were done only for 10 and 20 mg/kg) were observed. Perfusion of rabbit kidney (45 min exposure) and proximal tubules (30 min exposure) experiment resulted in a dose dependent difference in the amount of single strand breaks.	Jaffe et.al. (1985)
Primary kidney cells from both male rats and human	1–4 mM; 20 h exposure	NA	+	Statistically significant increase in all doses (1, 2, or 4 mM) both in rats and human cells.	Robbiano et al. (2004)
In vivo—male Sprague-Dawley rats exposed to TCE or DCVC—comet assay	TCE: 500–2,000 ppm, inhalation, 6 h/d, 5 d DCVC: 1 or 10 mg/kg, single oral dose for 16 h	+ (DCVC) – (TCE)	NA	No significant increase in tail length in any of the TCE exposed groups. In Expt. 1. 2 h exposure—1 or 10 mg to DCVC resulted in significant increase with no dose-response, but not at 16 h. In Expt. 2 ND for 1 mg, significant increase at 10 mg.	Clay (2008)
<b>Micronucleus</b>					
Syrian hamster embryo fibroblasts		NA	-	No micronucleus formation.	Vamvakas et al. (1988a)
Primary kidney cells from both male rats and human	1–4 mM; 20 h exposure	NA	+	Statistically significant increase in all doses (1, 2, and 4 mM) both in rats and human cells.	Robbiano et al. (2004)
Male Sprague-Dawley rats; proximal tubule cells (in vivo)	4 mM/kg TCE exposure, single dose	NA	+	Statistically significant increase in the average frequency of micronucleated kidney cells was observed.	Robbiano et al. (1998)

**Table 4-18. TCE GSH conjugation metabolites genotoxicity (continued)**

Test system/endpoint	Doses tested	With activation	Without activation	Comments	References
<b>Cell transformation</b>					
Kidney tubular epithelial cell line (LLC-PK1)	1 or 5 $\mu$ M; 7 wk	NA	+	Induced morphological cell transformation at both concentrations tested. Furthermore, cells maintained both biochemical and morphological alterations remained stable for 30 passages.	Vamvakas et al. (1996)
Rat kidney epithelial cells (in vitro)	10 $\mu$ M; 24 h exposure, 7 wk post incubation	NA	+	Cell transformation was higher than control, however, cell survival percentage ranged from 39–64% indicating cytotoxicity.	Mally et al. (2006)
<b>Gene expression</b>					
Kidney tubular epithelial cell line (LLC-PK1)	1 or 5 $\mu$ M clones, 30, 60, 90 min	NA	+	Increased <i>c-fos</i> expression in 1 and 5 $\mu$ M exposed clones at three different times tested.	Vamvakas et al. (1996)
Kidney tubular epithelial cell line (LLC-PK1)		NA	+	Expression of <i>c-fos</i> and <i>c-myc</i> increased in a time-dependent manner.	Vamvakas et al. (1993)

i.v. = intravenous, LDH = lactate dehydrogenase, ND = not determined, NA = not applicable.

1 was reported either in these DCVC transformants or in renal tumors (which were not increased in  
2 incidence) from TCE-treated Eker rats, which Mally et al. (2006) suggested support a  
3 nongenotoxic mechanism because a substantial fraction of spontaneous renal tumors in Eker rats  
4 showed LOH at this locus (Kubo et al., 1994; Yeung et al., 1995) and because LOH was  
5 exhibited both in vitro and in vivo with 2,3,4-tris(glutathion-S-yl)-hydroquinone treatment in  
6 Eker rats (Yoon et al., 2001). However, 2,3,4-tris(glutathion-S-yl)-hydroquinone is not  
7 genotoxic in standard mutagenicity assays (Yoon et al., 2001), and Kubo et al. (1994) also  
8 reported that none of renal tumors induced by the genotoxic carcinogen *N*-ethyl-*N*-nitrosourea  
9 showed LOH. Therefore, the lack of LOH at the *Tsc-2* locus induced by DCVC in vitro, or TCE  
10 in vivo, reported by Mally et al. (2006) is actually more similar to the response from the  
11 genotoxic carcinogen *N*-ethyl-*N*-nitrosourea than the nongenotoxic carcinogen  
12 2,3,4-tris(glutathion-S-yl)-hydroquinone. Therefore, these data do not substantially contradict  
13 the body of evidence on DCVC genotoxicity.

14 Finally, Clay (2008) evaluated the genotoxicity of DCVC in vivo using the comet assay  
15 to assess DNA breakage in the proximal tubules of rat kidneys. Rats were exposed orally to a  
16 single dose of DCVC (1 or 10 mg/kg). The animals were sacrificed either 2 or 16 hours after  
17 dosing and samples prepared for detecting the DNA damage. DCVC (1 and 10 mg/kg) induced  
18 no significant DNA damage in rat kidney proximal tubules at the 16-hour sampling time or after  
19 1 mg/kg DCVC at the 2-hour sampling time. While Clay et al. (2008) concluded that these data  
20 were insufficient to indicate a positive response in this assay, the study did report a statistically  
21 significant increase in percentage tail DNA 2 hours after treatment with 10 mg/kg DCVC,  
22 despite the small number of animals at each dose ( $n = 5$ ) and sampling time. Therefore, these  
23 data do not substantially contradict the body of evidence on DCVC genotoxicity.

24 Overall, DCVC, and to a lesser degree DCVG and NAcDCVC, have demonstrated  
25 genotoxicity based on consistent results in a number of available studies. While some recent  
26 studies (Clay, 2008; Mally et al., 2006) have reported a lack of positive responses in some in  
27 vivo measures of genotoxicity with DCVC treatment, due to a number of limitations discussed  
28 above, these studies do not substantially contradict the body of evidence on DCVC genotoxicity.  
29 It is known that these metabolites are formed in vivo following TCE exposure, specifically in the  
30 kidney, so they have the potential to contribute to the genotoxicity of TCE, especially in that  
31 tissue. Moreover, DCVC and DCVG genotoxic responses were enhanced when metabolic  
32 activation using *kidney* subcellular fractions was used (Vamvakas et al., 1988b). Finally, the  
33 lack of similar responses in in vitro genotoxicity assays with TCE, even with metabolic  
34 activation, is likely the result of the small yield (if any) of DCVC under in vitro conditions, since  
35 in vivo, DCVC is likely formed predominantly in situ in the kidney while S9 fractions are

1 typically derived from the liver. This hypothesis could be tested in experiments in which TCE is  
 2 incubated with subcellular fractions from the kidney, or from both the kidney and the liver (for  
 3 enhanced GSH conjugation).  
 4

**4.2.6. Trichloroethanol (TCOH)**

5 Limited studies are available on the effect of TCOH on genotoxicity (see Table 4-19).  
 6 TCOH is negative in the *S. typhimurium* assay using the TA100 strain (Bignami et al., 1980;  
 7 DeMarini et al., 1994; Waskell, 1978). A study by Beland (1999) using *S. typhimurium* strain  
 8 TA104 did not induce reverse mutations without exogenous metabolic activation, however did  
 9 increase mutant frequency in the presence of exogenous metabolic activation at a dose above  
 10 2,500 µg/plate. TCOH has not been evaluated in the other recommended screening assays.  
 11 Therefore, the database is limited for the determination of TCOH genotoxicity.  
 12  
 13

**Table 4-19. Genotoxicity of trichloroethanol**

Test system/endpoint	Doses (LED or HID) <sup>a</sup>	Results <sup>b</sup>		Reference
		With activation	Without activation	
<i>S. typhimurium</i> TA100, 98, reverse mutation	7,500 µg/plate	-	-	Waskell (1978)
<i>S. typhimurium</i> TA100, reverse mutation	0.5 µg/cm <sup>3</sup> vapor	-	-	DeMarini et al. (1994)
<i>S. typhimurium</i> TA104, reverse mutation	2,500 µg/plate	+	-	Beland (1999)
<i>S. typhimurium</i> TA100, 1535 reverse mutation	NA	-	-	Bignami et al. (1980)
Sister chromatid exchanges	NA	NA	+	Gu et al., 1981a

<sup>a</sup> LED = lowest effective dose; HID = highest ineffective dose.

<sup>b</sup> Results: + = positive; - = negative; NA = doses not available, results based on the abstract.

16  
 17  
 18  
 19  
 20

#### 4.2.7. Synthesis and Overall Summary

1 Trichloroethylene and its metabolites (TCA, DCA, CH, DCVC, DCVG, and TCOH) have  
2 been evaluated to varying degrees for their genotoxic activity in several of in vitro systems such  
3 as bacteria, yeast, and mammalian cells and, also, in in vivo systems.

4 There are several challenges in interpreting the genotoxicity results obtained from TCE  
5 exposure. For example, some studies in bacteria should be interpreted with caution if conducted  
6 using technical grade TCE since it may contain known bacterial mutagens in trace amounts as  
7 stabilizers (e.g., 1,2-epoxybutane and epichlorohydrin). Because of the volatile nature of TCE,  
8 there could be false negative results if proper precautions are not taken to limit evaporation, such  
9 as the use of a closed sealed system. The adequacy of the enzyme-mediated activation of TCE in  
10 vitro tests is another consideration. For example, it is not clear if standard S9 fractions can  
11 adequately recapitulate the complex in vivo metabolism of TCE to reactive intermediates, which  
12 in some cases entails multiple sequential steps involving multiple enzyme systems (e.g., CYP,  
13 GST, etc.) and interorgan processing (as is described in more detail in Section 3.3). In addition,  
14 the relative potency of the metabolites in vitro may not necessarily inform their relative  
15 contribution to the overall mechanistic effects of the parent compound, TCE. Furthermore,  
16 although different assays provided data relevant to different types of genotoxic endpoints, not all  
17 effects that are relevant for carcinogenesis are encompassed. The standard battery of prokaryotic  
18 as well as mammalian genotoxicity test protocols typically specify the inclusion of significantly  
19 cytotoxic concentrations of the test compound.

20 With respect to potency, several TCE studies have been conducted along with numerous  
21 other chlorinated compounds and the results interpreted as a comparison of the group of  
22 compounds tested (relative potency). However, for the purposes of hazard characterization, such  
23 comparisons are not informative—particularly if they are not necessarily correlated with in vivo  
24 carcinogenic potency. Also, differentiating the effects of TCE with respect to its potency can be  
25 influenced by many factors such as the type of cells, their differing metabolic capacities,  
26 sensitivity of the assay, need for greater concentration to show any effect, interpretation of data  
27 when the effects are marginal, and gradation of severity of effects.

28 Also, type of samples used, methodology used for the isolation of genetic material, and  
29 duration of exposure can particularly influence the results of several studies. This is particularly  
30 true for human epidemiological studies. For example, while some studies use tissues obtained  
31 directly from the patients others use formalin fixed tissues sections to isolate DNA for mutation  
32 detection. Type of fixing solution, fixation time, and period of storage of the tissue blocks often  
33 affect the quality of DNA. Formic acid contained in the formalin solution or picric acid  
34 contained in Bouin's solution is known to degrade nucleic acids resulting in either low yield or

1 poor quality of DNA. In addition, during collection of tumor tissues, contamination of  
2 neighboring normal tissue can easily occur if proper care is not exercised. This could lead to the  
3 ‘dilution effect’ of the results, i.e., because of the presence of some normal tissue; frequency of  
4 mutations detected in the tumor tissue can be lower than expected. Due to some of these  
5 technical difficulties in obtaining proper material (DNA) for the detection of mutation, the results  
6 of these studies should be interpreted cautiously.

7 The following synthesis, summary, and conclusions focus on the available studies that  
8 may provide some insight into the potential genotoxicity of TCE considering the above  
9 challenges when interpreting the mutagenicity data for TCE.

10 Overall, evidence from a number of different analyses and a number of different  
11 laboratories using a fairly complete array of endpoints suggests that TCE, following metabolism,  
12 has the potential to be genotoxic. TCE has a limited ability to induce mutation in bacterial  
13 systems, but greater evidence of potential to bind or to induce damage in the structure of DNA or  
14 the chromosome in a number of targets. A series of carefully controlled studies evaluating TCE  
15 itself (without mutagenic stabilizers and without metabolic activation) found it to be incapable of  
16 inducing gene mutations in most standard mutation bacterial assays (Baden et al., 1979; Bartsch  
17 et al., 1979; Crebelli et al., 1982; Henschler et al., 1977; Mortelmans et al., 1986; Shimada et al.,  
18 1985; Simmon et al., 1977; Waskell, 1978). Therefore, it appears that it is unlikely that TCE is a  
19 direct-acting mutagen, though TCE has shown potential to affect DNA and chromosomal  
20 structure. TCE is also positive in some but not all fungal and yeast systems (Callen et al., 1980;  
21 Crebelli et al., 1985; Koch et al., 1988; Rossi et al., 1983). Data from human epidemiological  
22 studies support the possible mutagenic effect of TCE leading to *VHL* gene damage and  
23 subsequent occurrence of renal cell carcinoma. Association of increased *VHL* mutation  
24 frequency in TCE-exposed renal cell carcinoma cases has been observed (Brauch et al., 1999;  
25 Brauch et al., 2004; Brüning et al., 1997b).

26 TCE can lead to binding to nucleic acids and proteins (Bergman, 1983; DiRenzo et al.,  
27 1982; Kautiainen et al., 1997; Mazzullo et al., 1992; Miller and Guengerich, 1983), and such  
28 binding appears to be due to conversion to one or more reactive metabolites. For instance,  
29 increased binding was observed in samples bioactivated with mouse and rat microsomal fractions  
30 (Banerjee and Van Duuren, 1978; DiRenzo et al., 1982; Mazzullo et al., 1992; Miller and  
31 Guengerich, 1983). DNA binding is consistent with the ability to induce DNA and chromosomal  
32 perturbations. Several studies report the induction of micronuclei in vitro and in vivo from TCE  
33 exposure (Hrelia et al., 1994; Hu et al., 2008; Kligerman et al., 1994; Robbiano et al., 2004;  
34 Wang et al., 2001). Reports of SCE induction in some studies are consistent with DNA effects,

1 but require further study (Gu et al., 1981a; Gu et al., 1981b; Kligerman et al., 1994; Nagaya et  
2 al., 1989a; White et al., 1979).

3 TCA, an oxidative metabolite of TCE, exhibits little, if any genotoxic activity in vitro.  
4 TCA did not induce mutations in *S. typhimurium* strains in the absence of metabolic activation or  
5 in an alternative protocol using a closed system (DeMarini et al., 1994; Giller et al., 1997;  
6 Kargalioglu et al., 2002; Nelson et al., 2001; Rapson et al., 1980; Waskell, 1978) but a  
7 mutagenic response was induced in TA100 in the Ames fluctuation test (Giller et al., 1997).  
8 However, in vitro experiments with TCA should be interpreted with caution if steps have not  
9 been taken to neutralize pH changes caused by the compound (Mackay et al., 1995). Measures  
10 of DNA-repair responses in bacterial systems have shown induction of DNA repair reported in *S.*  
11 *typhimurium* but not in *E. coli*. Mutagenicity in mouse lymphoma cells was only induced at  
12 cytotoxic concentrations (Harrington-Brock et al., 1998). TCA was positive in some  
13 genotoxicity studies in vivo mouse, newt, and chick test systems (Bhunya and Behera, 1987;  
14 Bhunya and Jena, 1996; Birner et al., 1994; Giller et al., 1997). DNA unwinding assays have  
15 either shown TCA to be much less potent than DCA (Nelson and Bull, 1988) or negative (Nelson  
16 et al., 1989; Styles et al., 1991). Due to limitations in the genotoxicity database, the possible  
17 contribution of TCA to TCE genotoxicity is unclear.

18 DCA, a chloroacid metabolite of TCE, has also been studied using different types of  
19 genotoxicity assays. Although limited studies are conducted for different genetic endpoints,  
20 DCA has been demonstrated to be mutagenic in the *S. typhimurium* assays, in vitro (DeMarini et  
21 al., 1994; Kargalioglu et al., 2002; Plewa et al., 2002) in some strains, mouse lymphoma assay,  
22 (Harrington-Brock et al., 1998) in vivo cytogenetic tests (Fusco et al., 1996; Leavitt et al.,  
23 1997), the micronucleus induction test, the Big Blue mouse system, and other tests (Bignami et  
24 al., 1980; Chang et al., 1992; DeMarini et al., 1994; Fuscoe et al., 1996; Harrington-Brock et al.,  
25 1998; Leavitt et al., 1997; Nelson and Bull, 1988; Nelson et al., 1989). DCA can cause DNA  
26 strand breaks in mouse and rat liver cells following in vivo mice and rats (Fusco et al., 1996).  
27 Because of uncertainties as to the extent of DCA formed from TCE exposure, inferences as to  
28 the possible contribution from DCA genotoxicity to TCE toxicity are difficult to make.

29 Chloral hydrate is mutagenic in the standard battery of screening assays. Effects include  
30 positive results in bacterial mutation tests for point mutations and in the mouse lymphoma assay  
31 for mutagenicity at the Tk locus (Haworth et al., 1983). In vitro tests showed that CH also  
32 induced micronuclei and aneuploidy in human peripheral blood lymphocytes and Chinese  
33 hamster pulmonary cell lines. Micronuclei were also induced in Chinese hamster embryonic  
34 fibroblasts. Several studies demonstrate that chloral hydrate induces aneuploidy (loss or gain of  
35 whole chromosomes) in both mitotic and meiotic cells, including yeast (Gualandi, 1987; Kafer,

1 1986; Singh and Sinha, 1976, 1979; Sora and Agostini Carbone, 1987), cultured mammalian  
2 somatic cells (Degrassi and Tanzarella, 1988), and spermatocytes of mice (Liang and  
3 Pacchierotti, 1988; Russo et al., 1984). Chloral hydrate was negative for sex-linked recessive  
4 lethal mutations in *Drosophila* (Yoon et al., 1985). It induces SSB in hepatic DNA of mice and  
5 rats (Nelson and Bull, 1988) and mitotic gene conversion in yeast (Bronzetti et al., 1984).  
6 Schatten and Chakrabarti (1998) showed that chloral hydrate affects centrosome structure, which  
7 results in the inability to reform normal microtubule formations and causes abnormal fertilization  
8 and mitosis of sea urchin embryos. Based on the existing array of data, CH has the potential to  
9 be genotoxic, particularly when aneuploidy is considered in the weight of evidence for genotoxic  
10 potential. Chloral hydrate appears to act through a mechanism of spindle poisoning and resulting  
11 in numerical changes in the chromosomes. These results are consistent with TCE, albeit there  
12 are limited data on TCE for these genotoxic endpoints.

13 DCVC, and to a lesser degree DCVG, has demonstrated bacterial mutagenicity based on  
14 consistent results in a number of available studies (Dekant et al., 1986c; Vamvakas et al., 1987;  
15 Vamvakas et al., 1988b). DCVC has demonstrated a strong, direct-acting mutagenicity both  
16 with and without the presence of mammalian activation enzymes. It is known that these  
17 metabolites are formed in vivo following TCE exposure, so they have the potential to contribute  
18 to the genotoxicity of TCE. The lack of similar response in bacterial assays with TCE is likely  
19 the result of the small yield (if any) of DCVC under in vitro conditions, since in vivo, DCVC is  
20 likely formed predominantly in situ in the kidney (S9 fractions are typically derived from the  
21 liver). DCVC and DCVG have not been evaluated extensively in other genotoxicity assays, but  
22 the available in vitro and in vivo data are predominantly positive. For instance, several studies  
23 have reported the DCVC can induce primary DNA damage in mammalian cells in vitro and in  
24 vivo (Clay, 2008; Jaffe et al., 1985; Vamvakas et al., 1989). Long-term exposure to DCVC  
25 induced de-differentiation of cells (Vamvakas et al., 1996). It has been shown to induce  
26 expression of the protooncogene *c-fos* (Vamvakas et al., 1996) and cause cell transformation in  
27 rat kidney cells (Mally et al., 2006). In LLC-PK1 cell clones, DCVC was reported to induce  
28 unscheduled DNA synthesis, but not micronuclei (Vamvakas et al., 1988a). Finally, DCVC  
29 induced transformation in kidney epithelial cells isolated from Eker rats carrying the  
30 heterozygous *Tsc-2* mutations (Mally et al., 2006). Moreover, the lack of LOH at the *Tsc-2* locus  
31 observed in exposed cells does not constitute negative evidence of DCVC genotoxicity, as none  
32 of renal tumors induced in Eker rats by the genotoxic carcinogen *N*-ethyl-*N*-nitrosourea showed  
33 LOH (Kubo et al., 1994).

34 In support of the importance of metabolism, there is some concordance between effects  
35 observed from TCE and those from several metabolites. For instance, both TCE and chloral



1 hydrate have been shown to induce micronucleus in mammalian systems, but chromosome  
2 aberrations have been more consistently observed with chloral hydrate than with TCE. The role  
3 of TCA in TCE genotoxicity is less clear, as there is less concordance between the results from  
4 these two compounds. Finally, several other TCE metabolites show at least some genotoxic  
5 activity, with the strongest data from DCA, DCVG, and DCVC. While quantitatively smaller in  
6 terms of flux as compared to TCA and TCOH (for which there is almost no genotoxicity data),  
7 these metabolites may still be toxicologically important.

8 Thus, uncertainties with regard to the characterization of TCE genotoxicity remain,  
9 particularly because not all TCE metabolites have been sufficiently tested in the standard  
10 genotoxicity screening battery to derive a comprehensive conclusion. However, the metabolites  
11 that have been tested particularly DCVC have predominantly resulted in positive data although  
12 to a lesser extent in DCVG and NAcDCVC, supporting the conclusion that these compounds are  
13 genotoxic, particularly in the kidney, where in situ metabolism produces and/or bioactivates  
14 these TCE metabolites.

#### 15

### 16 **4.3. CENTRAL NERVOUS SYSTEM (CNS) TOXICITY**

17 TCE exposure results in central nervous system (CNS) effects in both humans and  
18 animals that can result from acute, subchronic, or chronic exposure. There are studies indicating  
19 that TCE exposure results in CNS tumors and this discussion can be found in Section 4.9. The  
20 studies discussed in this section focus on the most critical neurological effects that were  
21 extracted from the neurotoxicological literature. Although there are several studies and reports  
22 that have evaluated TCE as an anesthetic, those studies were not included in this section because  
23 of the high exposure levels in comparison to the selected critical neurological effects described  
24 below. The critical neurological effects are nerve conduction changes, sensory effects, cognitive  
25 deficits, changes in psychomotor function, and changes in mood and sleep behaviors. The  
26 selection criteria that were used to determine study importance included study design and  
27 validity, pervasiveness of neurological effect, and for animal studies, the relevance of these  
28 reported outcomes in humans. More detailed information on human and animal neurological  
29 studies with TCE can be found in Appendix D.

### **4.3.1. Alterations in Nerve Conduction**

#### **4.3.1.1.1. Trigeminal Nerve Function: Human Studies**

1           A number of human studies have been conducted that examined the effects of  
2 occupational or drinking water exposures to TCE on trigeminal nerve function (see Table 4-20).  
3 Many studies reported that humans exposed to TCE present trigeminal nerve function  
4 abnormalities as measured by blink reflex and masseter reflex test measurements (Feldman et al.,  
5 1988; Feldman et al., 1992; Kilburn, 2002b; Kilburn and Warshaw, 1993)Ruitjen et al., 1991.  
6 The blink and masseter reflexes are mediated primarily by the trigeminal nerve and changes in  
7 measurement suggest impairment in nerve conduction. Other studies measured the trigeminal  
8

1  
2  
3

**Table 4-20. Summary of human trigeminal nerve and nerve conduction velocity studies**

Reference	Subjects	Exposure	Effect
Barret et al. (1982)	11 workers with chronic TCE exposure. Controls: 20 unexposed subjects.	Presence of TCE and TCA found through urinalysis. Atmospheric TCE concentrations and duration of exposure not reported in paper.	Following stimulation of the trigeminal nerve, significantly higher voltage stimuli was required to obtain a normal response and there was a significant increase in latency for response and decreased response amplitude.
Barret et al. (1984)	188 factory workers. No unexposed controls; lowest exposure group used as comparison.	>150 ppm; $n = 54$ < 150 ppm; $n = 134$ . 7 h/d for 7 yr.	Trigeminal nerve and optic nerve impairment, asthenia and dizziness were significantly increased with exposure.
Barret et al. (1987)	104 degreaser machine operators. Controls: 52 unexposed subjects Mean age 41.6 yr.	Mean duration, 8.2 yr, average daily exposure 7 h/d. Average TCOH range = 162–245 mg/g creatinine. Average TCA range = 93–131 mg/g creatinine.	Evoked trigeminal responses were measured following stimulation of the nerve and revealed increased latency to respond, amplitude or both and correlated with length of exposure ( $p < 0.01$ ) and with age ( $p < 0.05$ ), but not concentration.
El-Ghawabi et al. (1973)	30 money printing shop workers. Controls: 20 nonexposed males . 10 workers exposed to inks not containing TCE.	Mean TCE air concentrations ranged from 41 ppm to 163 ppm. Exposure durations:  Less than 1 yr: $n = 3$ 1 yr: $n = 1$ 2 yr: $n = 2$ 3 yr: $n = 11$ 4 yr: $n = 4$ 5 yr or greater: $n = 9$	No effect on trigeminal nerve function was noted.
Feldman et al. (1988)	21 Woburn, MA residents. 27 controls.	TCE maximum reported concentration in well water was 267 ppb; other solvents also present.  Exposure duration ranged from 1–12 yr.	Measurement of the blink reflex as mediated by the trigeminal nerve resulted in significant increases in the latency of reflex components ( $p < 0.001$ ).

Feldman et al. (1992)	18 workers. 30 controls.	<p>TCE exposure categories of “extensive”, “occasional,” and “chemical other than TCE”.</p> <p>“Extensive” = chronically exposed (<math>\geq 1</math> yr) to TCE for 5 d/wk and &gt;50% workday.</p> <p>“Occupational” = chronically exposed to TCE for 1–3 d/wk and &gt;50% workday.</p>	The blink reflex as mediated by the trigeminal was measured. The “extensive” group revealed latencies greater than three SD above the nonexposed group mean on blink reflex components.
-----------------------	-----------------------------	---	---

1

**Table 4-20. Summary of human trigeminal nerve and nerve conduction velocity studies (continued)**

Reference	Subjects	Exposure	Effect
Kilburn and Warshaw (1993)	160 residents living in Southwest Tucson with TCE, other solvents, and chromium in groundwater. Control: 113 histology technicians from a previous study (Kilburn et al., 1987; Kilburn and Warshaw, 1992).	>500 ppb of TCE in well water before 1981 and 25–100 ppb afterwards. Duration ranged from 1–25 yr.	Significant impairments in sway speed with eyes open and closed and blink reflex latency (R-1) which suggests trigeminal nerve impairment.
Kilburn (2002a)	236 residents near a microchip plant in Phoenix, AZ. Controls: 161 regional referents from Wickenburg, AZ and 67 referents in northeastern Phoenix.	<0.2–10,000 ppb of TCE, <0.2–260,000 ppb TCA, <0.2–6,900 ppb 1,1-DCE, <0.2–1,600 1,2-DCE, <0.2–23,000 ppb PCE, <0.02–330 ppb VC in well water. Exposure duration ranged from 2–37 yr.	Trigeminal nerve impairment as measured by the blink reflex test; both right and left blink reflex latencies (R-1) were prolonged. Exposed group mean 14.2 + 2.1 ms (right) or 13.9 + 2.1 ms (left) vs. referent group mean of 13.4 + 2.1 ms (right) or 13.5 + 2.1ms (left), <i>p</i> = 0.0001 (right) and 0.008 (left).
Mhiri et al. (2004)	<b>23 phosphate industry workers.</b> <b>Controls: 23 unexposed workers.</b>	Exposure ranged from 50–150 ppm, for 6 h/d for at least 2 yr. Mean urinary trichloroethanol and trichloroacetic acid levels were 79.3 ± 42 and 32.6 ± 22 mg/g creatinine.	<b>TSEPs were recorded. Increase in the TSEP latency was observed in 15 out of 23 (65%) workers.</b>

**Table 4-20. Summary of human trigeminal nerve and nerve conduction velocity studies (continued)**

Reference	Subjects	Exposure	Effect
Rasmussen et al. (1993d)	96 Danish metal degreasers. Age range: 19–68. No unexposed controls; low exposure group used as comparison.	<p>Average exposure duration: 7.1 yr.); range of full-time degreasing: 1 mo to 36 yr. Exposure to TCE or to CFC113 .</p> <p>1) Low exposure: <math>n = 19</math>, average full-time exposure 0.5 yr.</p> <p>2) Medium exposure: <math>n = 36</math>, average full-time exposure 2.1 yr.</p> <p>3) High exposure: <math>n = 41</math>, average full-time exposure 11 yr. TCA in high exposure group = 7.7 mg/L (max = 26.1 mg/L).</p>	No statistically significant trend on trigeminal nerve function, although some individuals had abnormal function.
Ruijten et al. (1991)	31 male printing workers. Mean age 44 yr; mean duration 16 yr. Controls: 28 unexposed; mean age 45 yr.	<p>Mean cumulative exposure = 704 ppm × yr (SD 583, range: 160–2,150 ppm × yr.</p> <p>Mean, 17 ppm at time of study; historic TCE levels from 1976–1981, mean of 35 ppm.</p> <p>Mean duration of 16 yr.</p>	<b>Measurement of trigeminal nerve function by using the blink reflex resulted in no abnormal findings. Increased latency in the masseter reflex is indicative of trigeminal nerve impairment.</b>

**Table 4-20. Summary of human trigeminal nerve and nerve conduction velocity studies (continued)**

Reference	Subjects	Exposure	Effect
Triebig et al. (1982)	24 workers (20 males, 4 females) occupationally exposed—ages 17–56. Controls: 144 individuals to establish normal nerve conduction parameters. Matched group: 24 unexposed workers (20 males, 4 females).	Exposure duration of 1 mo to 258 mo (mean 83 mo). Air exposures were between 5–70 ppm.	No statistically significant difference in nerve conduction velocities between the exposed and unexposed groups.
Triebig et al. (1983)	66 workers occupationally exposed. Control: 66 workers not exposed to solvents.	Subjects were exposed to a mixture of solvents, including TCE.	Exposure-response relationship observed between length of solvent exposure and statistically significant reduction in mean sensory ulnar nerve conduction velocities.

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

DCE = dichloroethylene, PCE = perchloroethylene, SD = standard deviation, VC = vinyl chloride.

somatosensory evoked potential (TSEP) following stimulation of the trigeminal nerve and reported statistically significantly delayed response on evoked potentials among exposed subjects compared to nonexposed individuals (Barret et al., 1982; Barret et al., 1984; Barret et al., 1987; Mhiri et al., 2004). Two studies which also measured trigeminal nerve function did not find any effect (El Ghawabi et al., 1973; Rasmussen et al., 1993d) but the methods were not provided in either study (El Ghawabi et al., 1973; Rasmussen et al., 1993d) or an appropriate control group was not included (Rasmussen et al., 1993d). These studies and results are described below and summarized in detail in Table 4-20.

Integrity of the trigeminal nerve is commonly measured using blink and masseter reflexes. Five studies (Barret et al., 1984; Feldman et al., 1988; Feldman et al., 1992; Kilburn, 2002b; Kilburn and Warshaw, 1993) significant increase in the latency to respond to the stimuli generating the reflex. The latency increases in the blink reflex ranged from 0.4 ms (Kilburn, 2002b) to up to 3.44 ms (Feldman et al., 1988). The population groups in these studies were exposed by inhalation occupationally (Barret et al., 1984) and through drinking water environmentally (Feldman et al., 1988; Kilburn, 2002b; Kilburn and Warshaw, 1993). Feldman et al. (1992) demonstrated persistence in the increased latency of the blink reflex response. In one subject, exposure to TCE (levels not reported by authors) occurred through a

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 degreasing accident (high and acute exposure), and increased latency response times persisted  
2 20 years after the accident. Another two subjects, evaluated at 9 months and 1 month following  
3 a high occupational exposure (exposure not reported by authors), also had higher blink reflex  
4 latencies with an average increase of 2.8 ms over the average response time in the control group  
5 used in the study. Although one study (Ruijten et al., 1991) did not find these increases in male  
6 printing workers exposed to TCE, this study did find a statistically significant average increase  
7 of 0.32 ms ( $p < 0.05$ ) in the latency response time in TCE-exposed workers on the masseter  
8 reflex test, another test commonly used to measure the integrity of the trigeminal nerve.

9 Three studies (Barret et al., 1982; Barret et al., 1987; Mhiri et al., 2004) adopting TSEPs  
10 to measure trigeminal nerve function found significant abnormalities in these evoked potentials.  
11 These studies were conducted on volunteers who were occupationally exposed to TCE through  
12 metal degreasing operations (Barret et al., 1982; Barret et al., 1987) or through cleaning tanks in  
13 the phosphate industry (Mhiri et al., 2004). Barret et al. (1982) reported that in 8 of the 11  
14 workers, an increased voltage ranging from a 25 to a 45 volt increase was needed to generate a  
15 normal TSEP and two of workers had an increased TSEP latency. Three out of 11 workers had  
16 increases in TSEP amplitudes. In a later study, Barret et al. (1987) also reported abnormal  
17 TSEPs (increased latency and/or increased amplitude) in 38% of the degreasers that were  
18 evaluated. The individuals with abnormal TSEPs were significantly older (45 vs. 40.1 years;  
19  $p < 0.05$ ) and were exposed to TCE longer (9.9 vs. 5.6 years;  $p < 0.01$ ). Mhiri et al. (2004) was  
20 the only study to evaluate individual components of the TSEP and noted significant increases in  
21 latencies for all TSEP potentials (N1, P1, N2, P2, N3;  $p < 0.01$ ) and significant decreases in  
22 TSEP amplitude (P1,  $p < 0.02$ ; N2,  $p < 0.05$ ). A significant positive correlation was  
23 demonstrated between exposure duration and increased TSEP latency ( $p < 0.02$ ).

24 Two studies reported no statistically significant effect of TCE exposure on trigeminal  
25 nerve function (El Ghawabi et al., 1973; Rasmussen et al., 1993a). El-Ghawabi et al. (1973)  
26 conducted a study on 30 money printing shop workers occupationally exposed to TCE.  
27 Trigeminal nerve involvement was not detected, but the authors did not include the experimental  
28 methods that were used to measure trigeminal nerve involvement and did not provide any data as  
29 to how this assessment was made. Rasmussen et al. (1993a) conducted an historical cohort study  
30 on 99 metal degreasers, 70 exposed to TCE and 29 to the fluorocarbon, CFC113. It was reported  
31 that 1 out of 21 people (5%) in the low exposure, 2 out of 37 (5%) in the medium exposure and 4  
32 out of 41 (10%) in the high exposure group experienced abnormalities in trigeminal nerve  
33 sensory function, with a linear trend test  $p$ -value of 0.42. The mean urinary trichloroacetic acid  
34 concentration was reported for the high exposure group only and was 7.7 mg/L (maximum  
35 concentration, 26.1 mg/L). The trigeminal nerve function findings of high exposure group



1 subjects was compared to that of low exposure group since this study did not include an  
2 unexposed or no TCE exposure group, and decreased the sensitivity of the study.

3

#### 4.3.1.1.2. Nerve Conduction Velocity—Human Studies

4 Two occupational studies assessed ulnar and median nerve function using tests of  
5 conduction latencies (Triebig et al., 1983; Triebig et al., 1982) (see Table 4-20). The ulnar nerve  
6 and median nerves are major nerves located in the arm and forearm. Triebig (1982) studied 24  
7 healthy workers (20 males, 4 females) exposed to TCE occupationally (5–70 ppm) at three  
8 different plants and did not find statistically significant differences in ulnar or median nerve  
9 conduction velocities between exposed and unexposed subjects. This study has measured  
10 exposure data, but exposures/responses are not reported by dose levels. The Triebig (1983)  
11 study is similar in design to the previous study (Triebig et al., 1982) but of a larger number of  
12 subjects. In this study, a dose-response relationship was observed between lengths of exposure  
13 to mixed solvents that included TCE (at unknown concentration). A statistically significant  
14 reduction in nerve conduction velocities was observed for the medium- and long-term exposure  
15 groups for the sensory ulnar nerve as was a statistically significant reduction in mean nerve  
16 conduction velocity observed between exposed and control subjects.

17

#### 4.3.1.1.3. Trigeminal Nerve Function: Laboratory Animal Studies

18 There is little evidence that TCE disrupts trigeminal nerve function in animal studies.  
19 Two studies demonstrated TCE produces morphological changes in the trigeminal nerve at a  
20 dose of 2,500 mg/kg-day for 10 weeks (Barret et al., 1992; Barret et al., 1991). However,  
21 dichloroacetylene, a degradation product formed during the volatilization of TCE was found to  
22 produce more severe morphological changes in the trigeminal nerve and at a lower dose of  
23 17 mg/kg-day (Barret et al., 1992; Barret et al., 1991). Only one study (Albee et al., 2006) has  
24 evaluated the effects of TCE on trigeminal nerve function and a subchronic inhalation exposure  
25 did not result in any significant functional changes. A summary of these studies is provided in  
26 Table 4-21.

27 Barret et al. (1992; 1991) conducted two studies evaluating the effects of both TCE and  
28 dichloroacetylene on trigeminal nerve fiber diameter and internodal length as well as several  
29 markers for fiber myelination. Female Sprague-Dawley rats ( $n = 7/\text{group}$ ) were dosed with  
30 2,500 mg/kg TCE or 17 mg/kg-day dichloroacetylene by gavage for 5 days/week for 10 weeks.

1 TCE-dosed animals only exhibited changes in the smaller Class A fibers where internode length  
2 increased marginally (<2%) and fiber diameter increased by 6%. Conversely,  
3

1  
2

**Table 4-21. Summary of animal trigeminal nerve studies**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL <sup>a</sup>	Effects
Barret et al. (1991)	Direct gastric administration	Rat, Sprague-Dawley, female, 7/group	0, 2.5 g/kg, acute administration. 17 mg/kg dichloroacetylene.	LOAEL: 2.5 g/kg	Morphometric analysis was used for analyzing the trigeminal nerve. Increase in external and internal fiber diameter as well as myelin thickness was observed in the trigeminal nerve after TCE treatment.
<b>Barret et al. (1992)</b>	<b>Direct gastric administration</b>	<b>Rat, Sprague-Dawley, female, 7/group</b>	<b>0, 2.5 g/kg; one dose/d, 5 d/wk, 10 wk. 17 mg/kg dichloroacetylene</b>	<b>LOAEL: 2.5 g/kg</b>	<b>Trigeminal nerve analyzed using morphometric analysis. Increased internode length and fiber diameter in class A fibers of the trigeminal nerve observed with TCE treatment. Changes in fatty acid composition also noted.</b>
Albee et al. (1997)	Inhalation	Rat, Fischer 344, male, 6	0 or 300-ppm dichloroacetylene, 2.25 h	LOAEL: 300 ppm dichloroacetylene	Dichloroacetylene (TCE byproduct) exposure impaired the TSEP up to 4 d postexposure.
Albee et al. (2006)	Inhalation	Rat, Fischer 344, male and female, 10/sex/group	0, 250, 800, or 2,500 ppm	N OAEL: 2,500 ppm	No effect on TSEPs was noted at any exposure level.

3  
4  
5  
6  
7

<sup>a</sup> NOAEL = no-observed-adverse-effect level, LOAEL = lowest-observed-adverse-effect-level.  
**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

8 dichloroacetylene-treated rats exhibited significant and more robust decreases in internode length  
9 and fiber diameter in both fiber classes A (decreased 8%) and B (decreased 4%).

10 Albee et al. (2006) evaluated the effects of a subchronic inhalation TCE exposure in  
11 Fischer 344 rats (10/sex/group). Rats were exposed to 0, 250, 800, and 2,500 ppm TCE for  
12 6 hours/day, 5 days/week for 13 weeks. TCE exposures were adequate to produce permanent  
13 auditory impairment even though TSEPs were unaffected. While TCE appears to be negative in  
14 disrupting the trigeminal nerve, the TCE breakdown product, dichloroacetylene, does impair  
15 trigeminal nerve function. Albee et al. (1997) showed that a single inhalation exposure of rats to

1 300-ppm dichloroacetylene, for 2.25 hours, disrupted trigeminal nerve evoked potentials for at  
2 least 4 days post exposure.

#### **4.3.1.1.4. Discussion and Conclusions: Trichloroethylene (TCE)-Induced Trigeminal Nerve Impairment**

3 Epidemiologic studies of exposure to TCE found impairment of trigeminal nerve  
4 function, assessed by the blink reflex test or the TSEP, in humans exposed occupationally by  
5 inhalation or environmentally by ingestion (see Table 4-20). Mean inhalational exposures  
6 inferred from biological monitoring or from a range of atmospheric monitoring in occupational  
7 studies was approximately 50 ppm to <150 ppm TCE exposure. Residence location is the  
8 exposure surrogate in geographical-base studies of contaminated water supplies with several  
9 solvents. Well water contaminant concentrations of TCE ranged from <0.2 ppb to 10,000 ppb  
10 and do not provide an estimate of TCE concentrations in drinking water to studied individuals.  
11 Two occupational studies, each including more than 100 subjects, reported statistically  
12 significant dose-response trends based on ambient TCE concentrations, duration of exposure,  
13 and/or urinary concentrations of the TCE metabolite TCA (Barret et al., 1984; Barret et al.,  
14 1987). Three geographical-based studies of environmental exposures to TCE via contaminated  
15 drinking water are further suggestive of trigeminal nerve function decrements; however, these  
16 studies are more limited than occupational studies due to questions of subject selection. Both  
17 exposed subjects who were litigants and control subjects who may not be representative of  
18 exposed (Kilburn, 2002a, b; Kilburn and Warshaw, 1993); referents in Kilburn and Warshaw  
19 (1993) were histology technicians and subjects in a previous study of formaldehyde and other  
20 solvent exposures and neurobehavioral effects (Kilburn et al., 1987; Kilburn and Warshaw,  
21 1992). Results were mixed in a number of smaller studies. Two of these studies reported  
22 changes in trigeminal nerve response (Barret et al., 1982; Mhiri et al., 2004), including evidence  
23 of a correlation with duration of exposure and increased latency in one study (Mhiri et al., 2004).  
24 Ruijten et al. (1991) reported no significant change in the blink reflex, but did report an increase  
25 in the latency of the masseter reflex, which also may reflect effects on the trigeminal nerve.  
26 Two other studies reported no observed effect on trigeminal nerve impairment, but the authors  
27 failed to provide assessment of trigeminal nerve function (El Ghawabi et al., 1973; Rasmussen et  
28 al., 1993a) or there was not a control (nonexposed) group included in the study (Rasmussen et  
29 al., 1993a). Therefore, because of limitations in statistical power, the possibility of exposure  
30 misclassification, and possible differences in measurement methods, these studies are not judged  
31 to provide substantial evidence against a causal relationship between TCE exposure and

1 trigeminal nerve impairment. Overall, the weight of evidence supports a relationship between  
2 TCE exposure and trigeminal nerve dysfunction in humans.

3 Impairment of trigeminal nerve function is observed in studies of laboratory animal  
4 studies. Although one subchronic animal study demonstrated no significant impairment of  
5 trigeminal nerve function following TCE exposure up to 2,500 ppm (no observed-adverse-effect  
6 level [NOAEL]; Albee et al., 2006), morphological analysis of the nerve revealed changes in its  
7 structure (Barret et al., 1992; Barret et al., 1991). However, the dose at which an effect was  
8 observed by Barret et al. (1992; 1991) was high (2,500 mg/kg-day—lowest-observed-adverse-  
9 effect level [LOAEL]) compared to any reasonable occupational or environmental setting,  
10 although no lower doses were used. The acute or subchronic duration of these studies, as  
11 compared to the much longer exposure duration in many of the human studies, may also  
12 contribute to the apparent disparity between the epidemiologic and (limited) laboratory animal  
13 data.

14 The subchronic study of Barret et al. (1992) and the acute exposure study of Albee et al.  
15 (Albee et al., 1997) also demonstrated that dichloroacetylene, a (ex vivo) TCE degradation  
16 product, also induces trigeminal nerve impairment, at much lower doses than TCE. It is possible  
17 that under some conditions, coexposure to dichloroacetylene from TCE degradation may  
18 contribute to the changes observed to be associated with TCE exposure in human studies, and  
19 this issue is discussed further below in Section 4.3.10.

20 Overall evidence from numerous epidemiologic studies supports a conclusion that TCE  
21 exposure induces trigeminal nerve impairment in humans. Laboratory animal studies provide  
22 limited additional support, and do not provide strong contradictory evidence. Persistence of  
23 these effects after cessation of exposure cannot be determined since exposure was ongoing in the  
24 available human and laboratory animal studies.

## 4.3.2. Auditory Effects

### 4.3.2.1.1. Auditory Function: Human Studies

26 The TCE Subregistry from the National Exposure Registry developed by the ATSDR was  
27 the subject of three studies (ATSDR, 2003b; Burg and Gist, 1999; Burg et al., 1995). A  
28 fourth study (Rasmussen et al., 1993a) of degreasing workers exposed to either TCE or CFC113  
29 also indirectly evaluated auditory function. These studies are discussed below and presented in  
30 detail in Table 4-22.

1           Burg et al. (1999; 1995) reviewed the effects of TCE on 4,281 individuals (TCE  
2 Subregistry) residentially exposed to this solvent for more than 30 consecutive days. Face-to-  
3 face interviews were conducted with the TCE subregistry population and self-reported hearing  
4 loss was evaluated based on personal assessment through the interview (no clinical evaluation  
5 was conducted). TCE registrants that were 9 years old or younger had a statistically significant  
6 increase in hearing impairment as reported by the subjects. The relative risk (RR) in this age  
7 group for hearing impairments was 2.13 (95% confidence interval [CI]: 1.12–4.06) which  
8 decreased to 1.12 (95% CI: 0.52–2.24) for the 10–17 age group and 0.32 (95% CI: 0.10–1.02)  
9

1  
2

**Table 4-22. Summary of human auditory function studies**

Reference	Subjects	Exposure	Effect
ATSDR (2003b)	116 children, under 10 yr of age, residing near six Superfund sites. Further study of children in Burg et al. (1999; 1995). Control: 182 children.	TCE and other solvents in ground water supplies. Exposures were modeled using tap water TCE concentrations and GIS for spatial interpolation, and LaGrange for temporal interpolation to estimate exposures from gestation to 1990 across the area of subject residences. Control = 0 ppb; low exposure group = 0 < 23 ppb-yr; and high exposure group = >23 ppb-yr.	Auditory screening revealed increased incidence of abnormal middle ear function in exposed groups as indicated from acoustic reflex test. Adjusted odds ratios for right ear ipsilateral acoustic reflects control, OR: 1.0, low exposure group, OR: 5.1, $p < 0.05$ ; high exposure group, OR: 7.2, $p < 0.05$ . ORs adjusted for age, sex, medical history and other chemical contaminants. No significant decrements reported in the pure tone and tympanometry screening.
Burg et al. (1995)	From an NHIS TCE subregistry of 4,281 (4,041 living and 240 deceased) residents.	Environmentally exposed to TCE and other solvents via well water in Indiana, Illinois, and Michigan.	Increase in self-reported hearing impairments for children $\leq 9$ yr.
Burg et al. (1999)	3,915 white registrants. Mean age 34 yr (SD = 19.9 yr).	Cumulative TCE exposure subgroups: <50 ppb, $n = 2,867$ ; 50–500 ppb, $n = 870$ ; 500–5,000 ppb, $n = 190$ ; >5,000 ppb, $n = 35$ .  Exposure duration subgroups: <2 yr, 2–5 yr, 5–10 yr., >10 yr.	A statistically significant association (adjusted for age and sex) between duration of exposure and self-reported hearing impairment was found.

<p>Rasmussen et al. (1993c)</p>	<p>96 Danish metal degreasers. Age range: 19–68 yr; . No unexposed controls; low exposed group is referent.</p>	<p>Average exposure duration: 7.1 yr.); range of full-time degreasing: 1 mo to 36 yr. Exposure to TCE or and CFC113.</p> <p>(1) Low exposure: <math>n = 19</math>, average full-time exposure 0.5 yr.</p> <p>(2) Medium exposure: <math>n = 36</math>, average full-time exposure 2.1 yr.</p> <p>(3) High exposure: <math>n = 41</math>, average full-time exposure 11 yr. Mean U-TCA in high exposure group = 7.7 mg/L (max = 26.1 mg/L).</p>	<p>Auditory impairments noted through several neurological tests. Significant relationship of exposure was found with Acoustic-motor function (<math>p &lt; 0.001</math>), Paced Auditory Serial Addition Test (<math>p &lt; 0.001</math>), Rey Auditory Verbal Learning Test (<math>p &lt; 0.001</math>).</p>
---------------------------------	---	--	--

1  
2 GIS = geographic information system, NHIS = National Health Interview Survey, SD = standard deviation,  
3 U-TCA = urinary trichloroacetic acid.  
4



1 for all older age groups. A statistically significant association (when adjusted for age and sex)  
2 was found between duration of exposure, in these studies this was length of residency, and  
3 reported hearing impairment. The odds ratio (OR) was 2.32 (95% CI: 1.18–4.56) for subjects  
4 exposed to TCE >2 years and ≤5 years, 1.17 (95% CI: 0.55–2.49) for exposure >5 years and  
5 ≤10 years, 2.46 (95% CI: 1.30–5.02) for exposure durations greater than 10 years.

6 ATSDR (2003b) conducted a follow-up study to the TCE subregistry findings (Burg and  
7 Gist, 1999; Burg et al., 1995) and focused on the subregistry children located in Elkhart, IN,  
8 Rockford, IL and Battle Creek, MI using clinical tests for oral motor, speech, and hearing  
9 function. Exposures were modeled using tap water TCE concentrations and geographic  
10 information system (GIS) for spatial interpolation, and LaGrange for temporal interpolation to  
11 estimate exposures from gestation to 1990 across the area of subject residences. Modeled data  
12 were used to estimate lifetime exposures (ppb-years) to TCE in residential wells. The median  
13 TCE exposure for the children was estimated from drinking water as 23 ppb/year of exposure  
14 (ranging from 0–702 ppb/year). Approximately 20% (ranged from 17–21% depending on  
15 ipsilateral or contralateral test reflex) of the children in the TCE subregistry and 5–7% in the  
16 control group exhibited an abnormal acoustic reflex (involuntary muscle contraction that  
17 measures movement of the stapedius muscle in the middle ear following a noise stimulus) which  
18 was statistically significant ( $p = 0.003$ ). Abnormalities in this reflex could be an early indicator  
19 of more serious hearing impairments. No significant decrements were reported in the pure tone  
20 and tympanometry screening.

21 Rasmussen et al. (1993c) used a psychometric test to measure potential auditory effects  
22 of TCE exposure in an occupational study. Results from 96 workers exposed to TCE and other  
23 solvents were presented in this study. Details of the exposure groups and exposure levels are  
24 provided in Table 4-22. The acoustic motor function test was used for evaluation of auditory  
25 function. Significant decrements ( $p < 0.05$ ) in acoustic motor function performance scores  
26 (average decrement of 2.5 points on a 10-point scale) was reported for TCE exposure.

#### 27 28 **4.3.2.1.2. Auditory Function: Laboratory Animal Studies**

29 The ability of TCE to permanently disrupt auditory function and produce abnormalities in  
30 inner ear histopathology has been demonstrated in several studies using a variety of test methods.  
31 Two different laboratories have identified NOAELs following inhalation exposure for auditory  
32 function of 1,600 ppm for 12 hours/day for 13 weeks in Long Evans rats ( $n = 6–10$ ) (Rebert et  
33 al., 1991) and 1,500 ppm for 18 hours/day, 5 days/week for 3 weeks in Wistar-derived rats  
( $n = 12$ ) (Jaspers et al., 1993). The LOAELs identified in these and similar studies are

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 2,500–4,000 ppm TCE for periods of exposure ranging from 4 hours/day for 5 days to  
2 12 hours/day for 13 weeks, (e.g., Albee et al., 2006; Boyes et al., 2000; Crofton and Zhao, 1997;  
3 Crofton et al., 1994; Fechter et al., 1998; Muijser et al., 2000; Rebert et al., 1993; Rebert et al.,  
4 1995). Rebert et al. (1993) estimated acute blood TCE levels associated with permanent hearing  
5 impairment at 125 µg/mL by methods that probably underestimated blood TCE values (rats were  
6 anaesthetized using 60% carbon dioxide [CO<sub>2</sub>]). A summary of these studies is presented in  
7 Table 4-23.

8 Reflex modification was used in several studies to evaluate the auditory function in  
9 TCE-exposed animals (Boyes et al., 2000; Crofton and Zhao, 1997; Crofton et al., 1994; Crofton  
10 and Zhao, 1993; Fechter et al., 1998; Jaspers et al., 1993; Muijser et al., 2000; Yamamura et al.,  
11 1983). These studies collectively demonstrate significant decreases in auditory function at  
12 midfrequency tones (8–20 kHz tones) for TCE exposures greater than 1,500 ppm after acute,  
13 short-term, and chronic durations. Only one study (Yamamura et al., 1983) did not demonstrate  
14 impairment in auditory function from TCE exposures as high as 17,000 ppm for 4 hours/day over  
15 5 days. This was the only study to evaluate auditory function in guinea pigs, whereas the other  
16 studies used various strains of rats. Despite the negative finding in Yamamura et al. (1983),  
17 auditory testing was not performed in an audiometric sound attenuating chamber and extraneous  
18 noise could have influenced the outcome. It is also important to note that the guinea pig has  
19 been reported to be far less sensitive than the rat to the effects of ototoxic aromatic hydrocarbons  
20 such as toluene.

21 Crofton and Zhao (1997) also presented a benchmark dose for which the calculated dose  
22 of TCE would yield a 15 dB loss in auditory threshold. This benchmark response was selected  
23 because a 15 dB threshold shift represents a significant loss in threshold sensitivity for humans.  
24 The benchmark concentrations for a 15 dB threshold shift are 5,223 ppm for 1 day, 2,108 ppm  
25 for 5 days, 1,418 ppm for 20 days and 1,707 ppm for 65 days of exposure. While more sensitive  
26 test methods might be used and other definitions of a benchmark effect chosen with a strong  
27 rationale, these data provide useful guidance for exposure concentrations that do yield hearing  
28 loss in rats.

29 Brainstem auditory-evoked potentials (BAERs) were also measured in several studies  
30 (Albee et al., 2006; Rebert et al., 1993; Rebert et al., 1991; Rebert et al., 1995) following at  
31 exposures ranging from 3–13 weeks. Rebert et al. (1991) measured BAERs in male Long Evans  
32 rats ( $n = 10$ ) and F344 rats ( $n = 4-5$ ) following stimulation with 4, 8, and 16 kHz sounds. The  
33 Long-Evans rats were exposed to 0, 1,600, or 3,200 ppm TCE, 12 hours/day for 12 weeks and  
34 the F344 rats were exposed to 0, 2,000, or 3,200 ppm TCE, 12 hours/day for 3 weeks. BAER

1 amplitudes were significantly decreased at all frequencies for F344 rats exposed to 2,000 and  
2 3,000 ppm TCE and for Long Evans rats exposed to 3,200 ppm TCE. These data identify a  
3

1  
2

**Table 4-23. Summary of animal auditory function studies**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL <sup>a</sup>	Effects
Rebert et al. (1991)	Inhalation	Rat, Long Evans, male, 10/group	Long Evans: 0, 1,600, and 3,200 ppm; 12 h/d, 12 wk	Long Evans: NOAEL: 1,600 ppm; LOAEL: 3,200 ppm	BAERs were measured. Significant decreases in BAER amplitude and an increase in latency of appearance of the initial peak (P1).
		Rat, F344, male, 4–5/group	F344: 0, 2,000, 3,200 ppm; 12 h/d, 3 wk	F344: LOAEL: 2,000 ppm	
Rebert et al. (1993)	Inhalation	Rat, Long Evans, male, 9/group	0, 2,500, 3,000, 3,500 ppm; 8 h/d, 5 d	NOAEL: 2,500 ppm; LOAEL: 3,000 ppm.	BAERs were measured 1–2 wk postexposure to assess auditory function. Significant decreases in BAERs were noted with TCE exposure.
Rebert et al. (1995)	Inhalation	Rat, Long Evans, male, 9/group	0, 2,800 ppm; 8 h/d, 5 d	LOAEL: 2,800 ppm	BAER measured 2–14 ds postexposure at a 16 kHz tone. Hearing loss ranged from 55–85 dB.
Crofton et al. (1994)	Inhalation	Rat, Long Evans, male, 7–8/group	0, 3,500 ppm TCE; 8 h/d, 5 d	LOAEL: 3,500 ppm	BAER measured and auditory thresholds determined 5–8 wk postexposure. Selective impairment of auditory function for mid-frequency tones (8 and 16 kHz).
Crofton and Zhou (1997); Boyes et al. (2000)	Inhalation	Rat, Long Evans, male, 9–12/group	0, 4,000, 6,000, 8,000 ppm; 6 h	NOAEL: 6,000 ppm; LOAEL: 8,000 ppm	<b>Auditory thresholds as measured by BAERs for the 16 kHz tone increased with TCE exposure. Measured 3–5 wk post exposure.</b>
		Rat, Long Evans, male, 8–10/group	0, 1,600, 2,400, and 3,200 ppm; 6 h/d, 5 d	NOAEL: 2,400 ppm; LOAEL: 3,200 ppm	
		Rat, Long Evans, male, 8–10/group	0, 800, 1,600, 2,400, and 3,200 ppm; 6 h/d, 5 d/wk, 4 wk	NOAEL: 2,400 ppm; LOAEL: 3,200 ppm	
		Rat, Long Evans, male, 8–10/group	0, 800, 1,600, 2,400, and 3,200 ppm; 6 h/d, 5 d/wk, 13 wk	NOAEL: 1,600 ppm; LOAEL: 2,400 ppm	

3

**Table 4-23. Summary of animal auditory function studies (continued)**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Fechter et al. (1998)	Inhalation	Rat, Long Evans, male, 12/group	0, 4,000 ppm; 6 h/d, 5 d	LOAEL: 4,000 ppm	Cochlear function measured 5–7 wk after exposure. Loss of spiral ganglion cells noted. Three wk postexposure, auditory function was significantly decreased as measured by compound action potentials and reflex modification.
Jaspers et al. (1993)	Inhalation	Rat, Wistar derived WAG-Rii/MBL, male, 12/group	0, 1,500, and 3,000 ppm; 18 h/d, 5 d/wk, 3 wk	NOAEL: 1,500 ppm	Auditory function assessed repeatedly 1–5 wk postexposure for 5, 20, and 35 kHz tones; no effect at 5 or 35 kHz; decreased auditory sensitivity at 20 kHz, 3,000 ppm.
Muijser et al. (2000)	Inhalation	Rat, Wistar derived WAG-Rii/MBL, male, 8	0, 3,000 ppm; 18 h/d, 5 d/wk, 3 wk	LOAEL: 3,000 ppm	Auditory sensitivity decreased with TCE exposure at 4, 8, 16, and 20 kHz tones. White noise potentiated the decrease in auditory sensitivity.
Albee et al. (2006)	Inhalation	Rat, Fischer 344, male and female, 10/sex/group	0, 250, 800, 2,500 ppm; 6 h/d, 5 d/wk, 13 wk	N OAEL: 800 ppm; LO AEL: 2,500 ppm	<b>Mild frequency specific hearing deficits; focal loss of cochlear hair cells.</b>
Yamamura et al. (1983)	Inhalation	Guinea Pig, albino Hartley, male, 7–10/group	0, 6,000, 12,000, 17,000 ppm; 4 h/d, 5 d	N OAEL: 17,000 ppm	No change in auditory sensitivity at any exposure level as measured by cochlear action potentials and microphonics. Study was conducted in guinea pig and species is less sensitive to auditory toxicity than rats. Studies were also not conducted in a sound-isolation chamber and effects may be impacted by background noise.

1  
2  
3  
4  
5  
6

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

BAER = brainstem auditory-evoked potential.

1 LOAEL at 2,000 ppm for the F344 rats and a NOAEL at 1,600 ppm for the Long Evans  
2 rats. In subsequent studies Rebert et al. (1993; 1995) again demonstrated TCE significantly  
3 decreases BAER amplitudes and also significantly increases the latency of appearance. Similar  
4 results were obtained by Albee et al. (2006) for male and female F344 rats exposed to TCE for  
5 13 weeks. The NOAEL for this study was 800 ppm based on ototoxicity at 2,500 ppm.

6 Notable physiological changes were also reported in a few auditory studies. Histological  
7 data from cochleas in Long-Evans rats exposed to 4,000 ppm TCE indicated that there was a loss  
8 in spiral ganglion cells (Fechter et al., 1998). Similarly, there was an observed loss in hair cells  
9 in the upper basal turn of the cochlea in F344 rats exposed to 2,500-ppm TCE (Albee et al.,  
10 2006).

#### 11 **4.3.2.1.3. Summary and Conclusion of Auditory Effects**

12 Human and animal studies indicated that TCE produces decrements in auditory function.  
13 In the human epidemiological studies (ATSDR, 2003b; Burg and Gist, 1999; Burg et al., 1995;  
14 Rasmussen et al., 1993a) it is suggested that auditory impairments result from both an inhalation  
15 and oral TCE exposure. A LOAEL of approximately 23 ppb-years TCE (extrapolated from  
16  $\leq 23$  ppb-years group in the ATSDR (2003b)) from oral intake is noted for auditory effects in  
17 children. The only occupational study where auditory effects were seen reported mean urinary  
18 trichloroacetic acid concentration, a nonspecific metabolite of TCE, of 7.7 mg/L for the high  
19 cumulative exposure group only (Rasmussen et al., 1993a). A NOAEL or a LOAEL for auditory  
20 changes resulting from inhalational exposure to TCE cannot be interpolated from average urinary  
21 trichloroacetic acid (U-TCA) concentration of subjects in the high exposure group because of a  
22 lack of detailed information on long-term exposure levels and duration (Rasmussen et al.,  
23 1993a). Two studies (Burg and Gist, 1999; Burg et al., 1995) evaluated self-reported hearing  
24 effects in people included in the TCE subregistry comprised of people residing near Superfund  
25 sites in Indiana, Illinois, and Michigan. In Burg et al. (1995), interviews were conducted with  
26 the TCE exposed population and it was found that children aged 9 years or younger had  
27 statistically significant hearing impairments in comparison to nonexposed children. This  
28 significant increase in hearing impairment was not observed in any other age group that was  
29 included in this epidemiological analysis. This lack of effect in other age groups may suggest  
30 association with another exposure other than drinking water ; however, it may also suggest that  
31 children may be more susceptible than adults. In a follow-up analysis, Burg et al. (1999)  
32 adjusted the statistical analysis of the original data (Burg et al., 1995) for age and sex. When  
33 these adjustments were made, a statistically significant association was reported self-reported for

1 auditory impairment and duration of residence. These epidemiological studies provided only  
2 limited information given their use of an indirect exposure metric of residence location, no  
3 auditory testing of this studied population and self-reporting of effects. ATSDR (2003b) further  
4 tested the findings in the Burg studies (Burg and Gist, 1999; Burg et al., 1995) by contacting the  
5 children that were classified as having hearing impairments in the earlier study and conducting  
6 several follow-up auditory tests. Significant abnormalities were reported for the children in the  
7 acoustic reflex test which suggested effects to the lower brainstem auditory pathway with the  
8 large effect measure, the odds ratio, was reported for the high cumulative exposure group.  
9 Strength of analyses was its adjustment for potential confounding effects of age, sex, medical  
10 history and other chemical contaminants in drinking water supplies. The ATSDR findings were  
11 important in that the results supported Burg et al. (1999; 1995). Rasmussen et al. (1993c) also  
12 evaluated auditory function in metal workers with inhalation exposure to either TCE or CFC113.  
13 Results from tasks including an auditory element suggested that these workers may have some  
14 auditory impairment. However, the tasks did not directly measure auditory function.

15 Animals strongly indicated that TCE produces deficits in hearing and provides biological  
16 context to the epidemiological study observations. Although there is a strong association  
17 between TCE and ototoxicity in the animal studies, most of the effects began to occur at higher  
18 inhalation exposures. NOAELs for ototoxicity ranged from 800–1,600 ppm for exposure  
19 durations of at least 12 weeks (Albee et al., 2006; Boyes et al., 2000; Crofton and Zhao, 1997;  
20 Rebert et al., 1991). Inhalation exposure to TCE was the route of administration in all the animal  
21 studies. These studies either used reflex modification audiometry (Crofton and Zhao, 1997;  
22 Crofton et al., 1994; Jaspers et al., 1993; Muijser et al., 2000) procedures or measured brainstem  
23 auditory evoked potentials (Rebert et al., 1993; Rebert et al., 1991; Rebert et al., 1995) to  
24 evaluate hearing in rats. Collectively, the animal database demonstrates that TCE produces  
25 ototoxicity at midfrequency tones (4–24 kHz) and no observed changes in auditory function  
26 were observed at either the low (<4 kHz) or high (>24 kHz) frequency tones. Additionally,  
27 deficits in auditory effects were found to persist for at least 7 weeks after the cessation of TCE  
28 exposure (Boyes et al., 2000; Crofton and Zhao, 1997; Fechter et al., 1998; Jaspers et al., 1993;  
29 Rebert et al., 1991). Decreased amplitude and latency were noted in the BAERs (Rebert et al.,  
30 1993; Rebert et al., 1991; Rebert et al., 1995) suggesting that TCE exposure affects central  
31 auditory processes. Decrements in auditory function following reflex modification audiometry  
32 (Crofton and Zhao, 1997; Crofton et al., 1994; Jaspers et al., 1993; Muijser et al., 2000)  
33 combined with changes observed in cochlear histopathology (Albee et al., 2006; Fechter et al.,  
34 1998) suggest that ototoxicity is occurring at the level of the cochlea and/or brainstem.

1 Changes in auditory function are noteworthy considering that TCE exposure is also  
2 associated with immunotoxicity and inflammatory-based diseases (discussed in Section 4.6).  
3 Autoimmune sensorineural hearing loss is a rare condition, sometimes seen with systemic  
4 autoimmune diseases (Bovo et al., 2006; Ruckenstein, 2004). The potential role of  
5 immunotoxicity in the observed auditory impairment seen with TCE is an area that requires  
6 additional research.  
7

### 4.3.3. Vestibular Function

#### 4.3.3.1.1. Vestibular Function: Human Studies

8 The earliest reports of neurological effects resulting from TCE exposures focused on  
9 subjective vestibular system symptoms, such as headaches, dizziness, and nausea. These  
10 symptoms are subjective and self-reported. However, there is little doubt that these effects can  
11 be caused by exposures to TCE, as they have been reported extensively in the literature, resulting  
12 from occupational exposures (Grandjean et al., 1955; Liu et al., 1988; Rasmussen and Sabroe,  
13 1986; Smith, 1970), environmental exposures (Hirsch et al., 1996), and in chamber studies  
14 (Smith, 1970; Stewart et al., 1970).

15 Kylin et al. (1967) exposed 12 volunteers to 1,000 ppm (5,500 mg/m<sup>3</sup>) TCE for 2 hours  
16 in a 1.5 × 2 × 2 meters chamber. Volunteers served as their own controls since 7 of the 12 were  
17 pretested prior to exposure and the remaining 5 were post-tested days after exposure. Subjects  
18 were tested for optokinetic nystagmus, which was recorded by electronystogmography, that is,  
19 “the potential difference produced by eye movements between electrodes placed in lateral angles  
20 between the eyes.” Venous blood was also taken from the volunteers to measure blood TCE  
21 levels during the vestibular task. The authors concluded that there was an overall reduction in  
22 the limit (“fusion limit”) to reach optokinetic nystagmus when individuals were exposed to TCE.  
23 Reduction of the “fusion limit” persisted for up to 2 hours after the TCE exposure was stopped  
24 and the blood TCE concentration was 0.2 mg/100 mL.  
25

#### 4.3.3.1.2. Vestibular Function: Laboratory Animal Data

26 The effect of TCE on vestibular function was evaluated by either (1) promoting  
27 nystagmus (vestibular system dysfunction) and comparing the level of effort required to achieve  
28 nystagmus in the presence and absence of TCE or (2) using an elevated beam apparatus and



1 measuring the balance. Overall, it was found that TCE disrupts vestibular function as presented  
2 below and summarized in Table 4-24.

3 Niklasson et al. (1993) showed acute impairment of vestibular function in male- and  
4 female-pigmented rats during acute inhalation exposure to TCE (2,700–7,200 ppm) and to  
5 trichloroethane (500–2,000 ppm). Both of these agents were able to promote nystagmus during  
6 optokinetic stimulation in a dose related manner. While there were no tests performed to assess  
7 persistence of these effects, Tham et al. (1984; 1979) did find complete recovery of vestibular  
8 function in rabbits ( $n = 19$ ) and female Sprague-Dawley rats ( $n = 11$ ) within minutes of  
9 terminating a direct arterial infusion with TCE solution.

1  
2

**Table 4-24. Summary of vestibular system studies**

Reference	Exposure route	Species/strain/ sex/number	Dose level/ exposure duration	NOAEL; LOAEL	Effects
<i>Vestibular system studies—humans</i>					
Kylin et al. (1967)	Inhalation	Humans, male and female, 12	1,000 ppm; 2 h	LOAEL: 1,000 ppm	Reduction in potential to reach nystagmus following TCE exposure.
<i>Vestibular system studies—animals</i>					
Tham et al. (1979)	Intravenous	Rabbit, strain unknown, sex unspecified, 19	1–5 mg/kg/min	---	Positional nystagmus developed once blood levels reached 30 ppm.
Tham et al. (1984)	Intravenous	Rat, Sprague-Dawley, female, 11	80 µg/kg/min	---	Excitatory effects on the vestibule-oculomotor reflex. Threshold effect at blood (TCE) of 120 ppm or 0.9 mM/L.
Niklasson et al. (1993)	Inhalation	Rat, strain unknown, male and female, 28	0, 2,700, 4,200, 6,000, 7,200 ppm; 1 h	LOAEL: 2,700 ppm	Increased ability to produce nystagmus.
Umezu et al. (1997)	Intraperitoneal	Mouse, ICR, male, 116	0, 250, 500, or 1,000 mg/kg, single dose and evaluated 30 min postadministration	NOAEL: 250 mg/kg LOAEL: 500 mg/kg	Decreased equilibrium and coordination as measured by the Bridge test (staying time on an elevated balance beam).

3  
4  
5  
6  
7  
8  
9  
10

The finding that trichloroethylene can yield transient abnormalities in vestibular function is not unique. Similar impairments have also been shown for toluene, styrene, along with trichloroethane (Niklasson et al., 1993) and by Tham et al. (1984) for a broad range of aromatic hydrocarbons. The concentration of TCE in blood at which effects were observed for TCE (0.9 mM/L) was quite close to that observed for most of these other vestibulo-active solvents.

**4.3.3.1.3. Summary and Conclusions for the Vestibular Function Studies**

11 Studies of TCE exposure in both humans and animals reported abnormalities in vestibular  
12 function. Headaches, dizziness, nausea, motor incoordination, among other subjective symptoms  
13 are reported in occupational epidemiological studies of TCE exposure (Grandjean et al., 1955;  
14 Hirsch et al., 1996; Liu et al., 1988; Rasmussen and Sabroe, 1986; Smith, 1970; Stewart et al.,  
15 1970). One human exposure study (Kylin et al., 1967) found that vestibular function was  
16 affected following an acute exposure to 1,000-ppm TCE (LOAEL). Individuals had a decreased

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 threshold to reach nystagmus than when exposed to TCE than to air. Animal studies also  
2 evaluated the threshold to reach nystagmus and reported that TCE decreased the threshold to  
3 produce nystagmus in rats (LOAEL: 2,700 ppm; Niklasson et al., 1993; Tham et al., 1984) and  
4 rabbits (Tham et al., 1984).

#### 4.3.4. Visual Effects

##### 4.3.4.1.1. Visual Effects: Human Studies

6 Visual impairment in humans has been demonstrated following exposures through  
7 groundwater (Kilburn, 2002b; Reif et al., 2003) from occupational exposure through inhalation  
8 (Rasmussen et al., 1993c; Troster and Ruff, 1990) and from a controlled inhalation exposure  
9 study (Vernon and Ferguson, 1969). Visual functions such as color discrimination and  
10 visuospatial learning tasks are impaired in TCE-exposed individuals. Additionally, an acute  
11 exposure can impair visual depth perception. Details of the studies are provided below and  
12 summarized in Table 4-25.

13 Geographical-based studies utilized color discrimination and contrast sensitivity tests to  
14 determine the effect of TCE exposure on vision. In these studies it was reported that TCE  
15 exposure significantly increased color discrimination errors (Kilburn, 2002b) or decreases in  
16 contrast sensitivity tests approached statistical significance after adjustments for several possible  
17 confounders ( $p = 0.06$  or  $0.07$ ; Reif et al., 2003). Exposure in Kilburn (2002b) is poorly  
18 characterized, and for both studies, TCE is one of several contaminants in drinking water  
19 supplies; neither study provides an estimate of an individual's exposure to TCE.

20 Rasmussen et al. (1993c) evaluated visual function in 96 metal workers, working in  
21 degreasing at various factories and with exposure to TCE or CFC113. Visual function was tested  
22 through the visual gestalts test (visual perception) and a visual recall test. In the visual gestalts  
23 test, the number of total errors significantly increased from the low group (3.4 errors) to the high  
24 exposure group (6.5 errors;  $p = 0.01$ ). No significant changes were observed in the visual recall  
25 task. Troster and Ruff (1990) presented case studies conducted on two occupationally exposed  
26 workers to TCE. Both patients presented with a visual-spatial task and neither could complete  
27 the task within the number of trials allowed suggesting visual function deficits as a measure of  
28 impaired visuospatial learning.

29 In a chamber exposure study (Vernon and Ferguson, 1969), eight male volunteers (ages  
30 21–30) were exposed to 0, 100, 300, and 1,000-ppm TCE for 2 hours. Each individual was  
31 exposed to all TCE concentrations and a span of at least 3 days was given between exposures.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 When the individuals were exposed to 1,000-ppm TCE (5,500 mg/m<sup>3</sup>), significant abnormalities  
2 were noted in depth perception as measured by the Howard-Dolman test ( $p < 0.01$ ). There were  
3 no effects on the flicker fusion frequency test (threshold frequency at which the individual sees a

1  
2

**Table 4-25. Summary of human visual function studies**

Reference	Subjects	Exposure	Effect
Kilburn et al. (2002b)	236 residents near a microchip plant in Phoenix, AZ. Controls: 67 local referents from Phoenix, AZ and 161 regional referents from Wickenburg, AZ.	TCE, TCA, 1,1-DCE, 1,2-DCE, PCE, and VC detected in well water up to 260,000 ppm; TCE concentrations in well water were 0.2–10,000 ppb. Exposure duration ranged from 2–37 yr.  Exposure duration ranged from 2–37 yr.	Color discrimination errors were increased among residents compared to regional referents ( $p < 0.01$ ). No adjustment for possible confounding factors.
Reif et al. (2003)	143 residents of the Rocky Mountain Arsenal community of Denver. Referent group at lowest concentration (<5 ppb).	Exposure modeling of TCE concentrations in groundwater and in distribution system to estimate mean TCE concentration by census block of residence. High exposure group >15 ppb. Medium exposure group $\geq 5$ ppb and $\leq 15$ ppb. Low exposure referent group <5 ppb.	Contrast sensitivity test performances (C and D) was marginally statistically significant ( $p = 0.06$ and $0.07$ , respectively). No significant effects reported for the Benton visual retention test. Significant decrements ( $p = 0.02$ ) were reported in the Benton visual retention test when stratified with alcohol consumption.

Rasmussen et al. (1993c)	96 Danish metal degreasers. Age range: 19–68; no unexposed controls; low exposure group was referent.	Average exposure duration: 7.1 yr); range of full-time degreasing: 1 mo to 36 yr. Exposure to TCE or CFC113.  1) Low exposure: $n = 19$ , average full-time expo 0.5 yr.  2) Medium exposure: $n = 36$ , average full-time exposure 2.1 yr.  3) high exposure: $n = 41$ , average full-time exposure 11 yr. TCA in high exposure group = 7.7 mg/L (max = 26.1 mg/L).	Statistically significant relationship of exposure was found with the Visual Gestalts learning and retention test (cognitive test) indicating deficits in visual performance.
Troster and Ruff (1990)	Two occupationally TCE-exposed workers. Controls: two groups of $n = 30$ matched controls; (all age and education matched).	Exposure concentration unknown. Exposure duration, 3–8 mos.	Both workers experienced impaired visuospatial learning.
Vernon and Ferguson (1969)	8 male volunteers age range 21–30; self controls.	0, 100, 300, and 1,000 ppm of TCE for 2 h.	Statistically significant effects on visual depth perception as measured by the Howard-Dolman test. NOAEL: 300 ppm; LOAEL: 1,000 ppm; No significant changes in any of the other visual test measurements.

1  
2  
3  
4  
5  
6

DCE = dichloroethylene.

flicker as a single beam of light) or on the form perception illusion test (volunteers presented with an illusion diagram).

**4.3.4.1.2. Visual Effects: Laboratory Animal Data**

7  
8

Changes in visual function have been demonstrated in animal studies during acute (Boyes et al., 2003, 2005) and subchronic exposure (Blain et al., 1994; Rebert et al., 1991). In

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 these studies, the effect of TCE on visual evoked responses to patterns (Boyes et al., 2003, 2005;  
2 Rebert et al., 1991) or a flash stimulus (Blain et al., 1994; Rebert et al., 1991) were evaluated.  
3 Overall, the studies demonstrated that exposure to TCE results in significant changes in the  
4 visual evoked response, which is reversible once TCE exposure is stopped. Details of the studies  
5 are provided below and are summarized in Table 4-26.

6 Boyes et al. (2003; 2005) exposed adult, male Long-Evans rats were to TCE in a  
7 head-only exposure chamber while pattern onset/offset visual evoked potentials (VEPs) were  
8 recorded. Exposure conditions were designed to provide concentration × time products of  
9 0 ppm/hours (0 ppm for 4 hours) or 4,000 ppm/hours (see Table 4-26 for more details). VEP  
10 amplitudes were depressed by TCE exposure during the course of TCE exposure. The degree of  
11 VEP depression showed a high correlation with the estimated brain TCE concentration for all  
12 levels of atmospheric TCE exposure.

13 In a subchronic exposure study, Rebert et al. (1991) exposed male Long Evans rats to  
14 1,600- or 3,200-ppm TCE, for 12 weeks, 12 hours/day. No significant changes in flash evoked  
15 potential measurements were reported following this exposure paradigm. Decreases in pattern  
16 reversal visual evoked potentials (NIP1 amplitude) reached statistical significance following 6,  
17 9, and 12 weeks of exposure. The drop in response amplitude ranged from approximately 20%  
18 after 8 weeks to nearly 50% at Week 14 but recovered completely within 1 week postexposure.

19 This transient effect of TCE on the peripheral visual system has also been reported by  
20 Blain (1994) in which New Zealand albino rabbits were exposed by inhalation to 350- and  
21 700-ppm TCE 4 hours/day, 4 days/week for 12 weeks. Electroretinograms (ERG) and  
22 oscillatory potentials (OPs) were recorded weekly under mesopic conditions. Recordings from  
23 the 350- and 700-ppm exposed groups showed a significant increase in the amplitude of the a-  
24 and b-waves (ERG). The amplitude of the OPs was significantly decreased at 350 ppm (57%)  
25 and increased at 700 ppm (117%). These electroretinal changes returned to pre-exposure  
26 conditions within 6 weeks after the inhalation stopped.

1  
2

**Table 4-26. Summary of animal visual system studies**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Rebert et al. (1991)	Inhalation	Rat, Long Evans, male, 10/group	0, 1,600, and 3,200 ppm; 12 h/d, 12 wk	NOAEL: 1,600 ppm	Significant amplitude decreases in pattern reversal evoked potentials (N1P1 amplitude) at 6, 9, and 12 wk.
Boyes et al. (2003)	Inhalation	Rat, Long Evans, male, 9–10/group	0 ppm, 4 h; 1,000 ppm, 4 h; 2,000 ppm, 2 h; 3,000 ppm, 1.3 h; 4,000 ppm, 1 h	LOAEL: 1,000 ppm, 4 h	Visual function significantly affected as measured by decreased amplitude (F2) in Fourier-transformed visual evoked potentials. Peak brain TCE concentration correlated with dose-response.
Boyes et al. (2005)	Inhalation	Rat, Long Evans, male, 8–10/group	0 ppm, 4 h; 500 ppm, 4 h; 1,000 ppm, 4 h; 2,000 ppm, 2 h; 3,000 ppm, 1.3 h; 4,000 ppm, 1 h; 5,000 ppm, 0.8 h	LOAEL: 500 ppm, 4 h	Visual function significantly affected as measured by decreased amplitude (F2) in Fourier-transformed visual evoked potentials. Peak brain TCE concentration correlated with dose-response.
<b>Blain et al. (1994)</b>	<b>Inhalation</b>	<b>Rabbit, New Zealand albino, male, 6–8/group</b>	<b>0, 350, 700 ppm; 4 h/d, 4 d/wk, 12 wk</b>	<b>LOAEL: 350 ppm</b>	<b>Significant effects noted in visual function as measured by ERG and OPs immediately after exposure. No differences in ERG or OP measurements were noted at 6 wk post-TCE exposure.</b>

3  
4  
5  
6  
7  
8

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

ERG = electroretinogram, OP = oscillatory potential.

**4.3.4.1.3. Summary and Conclusion of Visual Effects**

9 Changes in visual function are reported in human studies. Although central visual function  
10 was not evaluated in the human studies (such as electroretinograms, evoked potential  
11 measurements), clinical tests indicated deficits in color discrimination (Kilburn, 2002b), visual  
12 depth perception (Vernon and Ferguson, 1969) and contrast sensitivity (Reif et al., 2003). These  
13 changes in visual function were observed following both an acute exposure (Vernon and Ferguson,

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 1969) and residence in areas with groundwater contamination with TCE and other chemicals  
2 (Kilburn, 2002b; Reif et al., 2003). The exposure assessment approach of Reif et al., who adopted  
3 exposure modeling and information on water distribution patterns, is considered superior to that of  
4 Kilburn (2002b) who used residence location as a surrogate for exposure. In the one acute,  
5 inhalation study (Vernon and Ferguson, 1969), a NOAEL of 300 ppm and a LOAEL of 1,000 ppm  
6 for 2 hours was reported for visual effects. A NOAEL is not available from the drinking water  
7 studies since well water TCE concentration is a poor surrogate for an individual's TCE ingestion  
8 (2002b) and limited statistical analysis comparing high exposure group to low exposure group  
9 (Reif et al., 2003).

10 Animal studies have also demonstrated changes in visual function. All of the studies  
11 evaluated central visual function by measuring changes in evoked potential response following a  
12 visual stimulus that was presented to the animal. Two acute exposure inhalation studies (Boyes et  
13 al., 2003; Boyes et al., 2005) exposed Long Evans rats to TCE based on a concentration  $\times$  time  
14 schedule (Haber's law) and reported decreases in visual evoked potential amplitude. All of the  
15 exposures from these two studies resulted in decreased visual function with a LOAEL of 500 ppm  
16 for 4 hours. Another important finding that was noted is the selection of the appropriate  
17 dose-metric for visual function changes following an acute exposure. Boyes et al. (2003; 2005)  
18 found that among other potential dose-metrics, brain TCE concentration was best correlated with  
19 changes in visual function as measured by evoked potentials under acute exposure conditions.  
20 Two subchronic exposure studies (Blain et al., 1994; Rebert et al., 1991) demonstrated visual  
21 function changes as measured by pattern reversal evoked potentials (Rebert et al., 1991) or  
22 electroretinograms/oscillatory potentials (Blain et al., 1994). Unlike the other three visual function  
23 studies conducted with rats, Blain et al., demonstrated these changes in rabbits. Significant  
24 changes in ERGs and oscillatory potentials were noted following a 12-week exposure at 350 ppm  
25 (LOAEL) in rabbits (Blain et al., 1994) and in rats exposed to 3,200-ppm TCE for 12 weeks there  
26 were significant decreases in pattern reversal evoked potentials but no effect was noted in the  
27 1,600-ppm exposure group (Rebert et al., 1991). Both subchronic studies examined visual  
28 function following an exposure-free period of either 2 weeks (Rebert et al., 1991) or 6 weeks  
29 (Blain et al., 1994) and found that visual function returned to pre-exposure levels and the changes  
30 are reversible.

31

### 4.3.5. Cognitive Function

#### 4.3.5.1.1. Cognitive Effects: Human Studies

1           Effects of TCE on learning and memory have been evaluated in populations  
2 environmentally exposed to TCE through well water, in workers occupationally exposed through  
3 inhalation and under controlled exposure scenarios. Details of the studies are provided in  
4 Table 4-27 and discussed briefly below. In the geographical-based studies (Kilburn, 2002b;  
5 Kilburn and Warshaw, 1993) cognitive function was impaired in both studies and was evaluated  
6 by testing verbal recall and digit span memory among other measures. In Arizona residents  
7 involved in a lawsuit (Kilburn and Warshaw, 1993), significant impairments in all  
8 three cognitive measures were reported; verbal recall ( $p = 0.001$ ), visual recall ( $p = 0.03$ ) and  
9 digit span test ( $p = 0.07$ ), although a question exists whether the referent group was comparable  
10 to exposed subjects and the study's lack of consideration of possible confounding exposures in  
11 statistical analyses. Significant decreases in verbal recall ability was also reported in another  
12 environmental exposure study where 236 residents near a microchip plant with TCE  
13 concentration in well water ranging from 0.2–10,000 ppb (Kilburn, 2002b).

14           Cognitive impairments are assessed in the occupational exposure and case studies  
15 (Rasmussen et al., 1993c; Rasmussen et al., 1993d; Troster and Ruff, 1990). In metal degreasers  
16 occupationally exposed to TCE and CFC113, significant cognitive performance decreases were  
17 noted in verbal recall testing ( $p = 0.03$ ) and verbal learning ( $p = 0.04$ ; Rasmussen et al., 1993a).  
18 No significant effects were found in the visual recall or digit span test for these workers. Troster  
19 and Ruff (1990) reported decrements (no statistical analysis performed) in cognitive performance  
20 as measured in verbal and visual recall tests that were conducted immediately after presentation  
21 (learning phase) and one hour after original presentation (retention/memory phase) for two case  
22 studies.

23           Several controlled (chamber) exposure studies were conducted to cognitive ability during  
24 TCE exposure and most did not find any significant decrements in the neurobehavioral  
25 measurement. Only Salvini et al. (1971) found significant decrements in cognitive function.  
26 Six males were exposed to 110 ppm (550 mg/m<sup>3</sup>) TCE for 4-hour intervals, twice per day.  
27 Statistically significant results were observed for perception tests learning ( $p < 0.001$ ), mental  
28 fatigue ( $p < 0.01$ ), subjects ( $p < 0.05$ ); and choice reaction time (CRT) learning ( $p < 0.01$ ),  
29 mental fatigue ( $p < 0.01$ ), subjects ( $p < 0.05$ ). Triebig et al. (1977a, b) exposed seven total  
30 subjects (male and female) to 100 ppm TCE for 6 hours/day, 5 days/week and did not report any  
31 decreases in cognition but details on the experimental procedures were not provided.  
32 Additionally, Gamberale et al. (1976) found that subjects exposed to TCE as high as 194 ppm for

1 70 minutes did not exhibit any impairments on a short term memory test in comparison to an air  
2 exposure.

3

#### 4.3.5.1.2. Cognitive Effects: Laboratory Animal Studies

4 Many reports have demonstrated significant differences in performance of learning tasks  
5 such as the speed to complete the task. However, there is little evidence that learning and  
6 memory function are themselves impaired by exposure. There are also limited data that suggest  
7 alterations in the hippocampus of laboratory animals exposed to TCE. Given the important role  
8 that this structure plays in memory formation, such data may be relevant to the question of  
9 whether TCE impairs memory. The studies are briefly discussed below and details are provided  
10 in Table 4-28.

11 Two studies (Kulig, 1987; Umezu et al., 1997) reported decreased performance in  
12 operant-conditioning cognitive tasks for rodents. Kishi et al. (1993) acutely exposed Wistar rats  
13 to TCE at concentrations of 250, 500, 1,000, 2,000, and 4,000 ppm for 4 hours. Rats exposed to  
14 250 ppm TCE and higher showed a significant decrease both in the total number of lever presses  
15 and in avoidance responses compared with controls. The rats did not recover their pre-exposure  
16 performance until about 2 hours after exposure. Likewise, Umezu et al. (1997) reported a  
17 depressed rate of operant responding in male ICR strain mice ( $n = 6$ , exposed to all TCE doses,  
18 see Table 4-28) in a conditioned avoidance task that reached significance with i.p. injections of  
19 1,000 mg/kg. Increased responding during the signaled avoidance period at lower doses (250  
20 and 500 mg/kg) suggests an impairment in ability to inhibit responding or failure to attend to the  
21 signal.

22 Although cognitive impairments are noted, two additional studies indicate no change in  
23 cognition with continuous TCE exposure or improvements in cognitive tasks. No decrements in  
24 cognitive function as measured by the radial arm maze were observed in Mongolian gerbils  
25 exposed continuously by inhalation to 320 ppm TCE for 9 months (Kjellstrand et al., 1980).  
26 Improved performance was noted in a Morris swim test for weanling rats orally dosed with  
27 5.5 mg/day for 4 weeks followed by 2 weeks of no exposure and an additional 2 weeks of  
28 8.5 mg/day (Isaacson et al., 1990). This improved performance occurred despite a loss in  
29 hippocampal myelination.

30

#### **4.3.5.1.3. Summary and Conclusions of Cognitive Function Studies**

1 Human environmental and occupational exposure studies suggest impairments in  
2 cognitive function. Kilburn and Warshaw (1993) and Kilburn (2002b) reported memory deficits  
3 individuals although a question exists whether the referent group was comparable to exposed  
4 subjects and these studies lack of consideration of possible confounding exposures in statistical  
5 analyses. Significant impairments were found in visual and verbal recall and with the digit span  
6 test. Similarly, in occupational exposure studies (Rasmussen et al., 1993c; Rasmussen et al.,  
7 1993d; Troster and Ruff, 1990), short term memory tests indicated that immediate memory and  
8 learning were impaired in the absence of an effect on digit span performance. In controlled  
9 exposure and/or chamber studies, two studies did not report any cognitive impairment

1  
2

**Table 4-27. Summary of human cognition effect studies**

Reference	Subjects	Exposure	Effect
Kilburn and Warshaw (1993)	170 residents living in Southwest Tucson with TCE, other solvents, and chromium in groundwater. Control: 68 residential referents matched to subjects from two previous studies of waste oil and oil refinery exposures.	>500 ppb of TCE in well water before 1981 and 25–100 ppb afterwards.  Exposure duration ranged from 1–25 yr.	Decreased performance in the digit span memory test and story recall ability.
Kilburn 2002a	236 residents near a microchip plant. Controls: 67 local referents from Phoenix, AZ and 161 regional referents from Wickenburg, AZ.	<0.2–10,000 ppb of TCE, <0.2–260,000 ppb TCA, <0.2–6,900 ppb 1,1-DCE, <0.2–1,600 1,2-DCE, <0.2–23,000 ppb PCE, <0.02–330 ppb VC in well water.  Exposure duration ranged from 2–37 yr. Exposure duration ranged from 2–37 yr.	Cognitive effects decreased as measured by lower scores on Culture Fair 2A, vocabulary, grooved pegboard (dominant hand), trail making test, and verbal recall (i.e., memory).

Rasmussen (1993c; 1993d)	96 Danish metal degreasers. Age range: 19–68; No external controls.	<p>Average exposure duration: 7.1 yr.); range of full-time degreasing: 1 mo to 36 yr.</p> <p>1) Low exposure: <math>n = 19</math>, average full-time expo 0.5 yr.</p> <p>2) Medium exposure: <math>n = 36</math>, average full-time exposure 2.1 yr.</p> <p>3) High exposure: <math>n = 41</math>, average full-time exposure 11 yr. TCA in high exposure group = 7.7 mg/L (max = 26.1 mg/L).</p>	Cognitive impairment (psycho-organic syndrome) prevalent in exposed individuals. The incidence of this syndrome was 10.5% in the low exposure, 39.5% for medium exposure, and 63.4% for high exposure. Age is a confounder. Dose-response with 9 of 15 tests; Controlling for confounds, significant relationship of exposure was found with Acoustic-motor function ( $p < 0.001$ ), Paced Auditory Serial Addition Test ( $p < 0.001$ ), Rey Auditory Verbal-Learning Test ( $p < 0.001$ ), vocabulary ( $p < 0.001$ ) and visual gestalts ( $p < 0.001$ ); significant age effects. Age is a confounder.
Troster and Ruff (1990)	Two occupationally TCE-exposed workers. Controls: two groups of $n = 30$ matched controls; (all age and education matched.	Exposure concentration unknown; Exposure duration, 3–8 mo.	Both TCE cases exhibited significant deficits in verbal recall and visuospatial learning.

1

**Table 4-27. Summary of human cognition effect studies (continued)**

Reference	Subjects	Exposure	Effect
Triebig (1976)	Controlled exposure study four females, three males. Controls: four females, three males.	0, 100 ppm (550 mg/m <sup>3</sup> ), 6 h/d, 5 d.	There was no correlation seen between exposed and unexposed subjects for any measured psychological test results. No methods description was provided.
Triebig (1977b)	Seven men and one woman occupationally exposed with an age range from 23–38 yr. No control group.	50 ppm (260 mg/m <sup>3</sup> ). Exposure duration not reported.	The psychological tests showed no statistically significant difference in the results before or after the exposure-free time period. No methods description was provided.
Triebig (1977a)	Controlled exposure study on three male and four female students. Control: three male and four female students.	0, 100 ppm (550 mg/m <sup>3</sup> ), 6 h/d, 5 d	No significantly different changes were obtained. No methods description was provided.
Salvini et al. (1971)	Controlled exposure study six students, male. Self used as control.	TCE concentration was 110 ppm for 4-h intervals, twice per d. 0 ppm control exposure for all as self controls	Statistically significant results were observed for perception tests learning ( $p < 0.001$ ) and CRT learning ( $p < 0.01$ ).
Gamberale et al. (1976)	15 healthy men aged 20–31 yr old. Controls: Within Subjects (15 self-controls).	0 mg/m <sup>3</sup> , 540 mg/m <sup>3</sup> (97 ppm), 1,080 mg/m <sup>3</sup> (194 ppm), 70 min	Repetition of the testing led to a pronounced improvement in performance as a result of the training effect; No interaction effects between exposure to TCE and training.
Stewart et al. (1970)	130 (108 males, 22 females). Controls: 63 unexposed men.	TCA metabolite levels in urine were measured: 60.8% had levels up to 20 mg/L, and 82.1% had levels up to 60 mg/L	No significant effect on cognitive tests noted, but more effort required to perform the test in exposed group.
Chalupa (1960)	Case study—6 subjects. Average age 38.	No exposure data were reported	80% of those with pathological EEG displayed memory loss; 30% of those with normal EEGs displayed memory loss.

1  
2  
3

DCE = dichloroethylene, EEG = electroencephalogram.

1  
2  
3  
4  
5  
6  
7  
8

**Table 4-28. Summary of animal cognition effect studies**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Kjellstrand et al. (1980)	Inhalation	Gerbil, Mongolian, males and females, 12/sex/dose	0, 320 ppm; 9 mo, continuous (24 h/d) except 1–2 h/wk for cage cleaning	NOAEL: 320 ppm	No significant effect on spatial memory (radial arm maze).
<b>Isaacson et al. (1990)</b>	<b>Oral, drinking water</b>	<b>Rat, Sprague-Dawley, male weanlings, 12/dose</b>	(1) 0 mg/kg-day, 8 wk. (2) 5.5 mg-day (47 mg/kg-day <sup>a</sup> ), 4 wk + 0 mg/kg-day, 4 wk. (3) 5.5 mg/dd, 4 wk (47 mg/kg-day <sup>b</sup> ) + 0 mg/kg-day, 2 wk + 8.5 mg/dd (24 mg/kg-day <sup>b</sup> ), 2 wk	<b>NOAEL: 5.5 mg/d, 4 wk—spatial learning</b> <b>LOAEL: 5.5 mg/d—hippocampal demyelination</b>	<b>Decreased latency to find platform in the Morris water maze (Group #3); Hippocampal demyelination observed in all TCE-treated groups.</b>
Kishi et al. (1993)	Inhalation	Rats, Wistar, male, number not specified	0, 250,500, 1,000, 2,000, and 4,000 ppm, 4 h	LOAEL: 250 ppm	Decreased lever presses and avoidance responses in a shock avoidance task.
Umezu et al. (1997)	Intra-peritoneal	Mouse, ICR, male, six exposed to all treatments (repeated exposure)	0, 125, 250, 500, and 1,000 mg/kg, single dose and evaluated 30 min postadministration	NOAEL: 500 mg/kg LOAEL: 1,000 mg/kg	Decreased response rate in an operant response—condition avoidance task.
Oshiro et al. (2004)	Inhalation	Rat, Long Evans, male, 24	0, 1,600, and 2,400 ppm; 6 h/d, 5 d/wk, 4 wk	NOAEL: 2,400 ppm	No change in reaction time in signal detection task and when challenged with amphetamine, no change in response from control.

9  
10 <sup>a</sup> mg/kg-day conversion estimated from average male Sprague-Dawley rat body weight from ages 21–49 d (118 g)  
11 for the 5.5 mg dosing period and ages 63–78 d (354 g) for the 8.5 mg dosing period.  
12

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

*This document is a draft for review purposes only and does not constitute Agency policy.*



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

(Gamberale et al., 1976; Stewart et al., 1970) and one study (Salvini et al., 1971) reported significant impairments in learning memory and complex choice reaction tasks. All of the controlled exposure studies were acute and/or short-term exposure studies and the sensitivity of test procedures is unknown due to the lack of methodologic information provided in the reports. Despite identified study deficiencies, these studies collectively suggest cognitive function impairment.

The animal studies measured cognitive function through spatial memory and operant responding tasks. In the two studies where spatial memory was evaluated, there was either no effect at 320 ppm TCE (Kjellstrand et al., 1980) or improved cognitive performance in weanling rats at a dose of 5.5 mg/day for four weeks (Isaacson et al., 1990). Improved cognitive performance was observed in weanling rats (Isaacson et al., 1990) and could be due to continuing neurodevelopment as well as compensation from other possible areas in the brain since there was a significant loss in hippocampal myelination. Significant decreases in operant responding (avoidance/punished responding) during TCE exposure were reported in two studies (Kishi et al., 1993; Umezu et al., 1997). When TCE exposure was discontinued operant responding return to control levels and it is unclear if the significant effects are due to decreased motor function or decreased cognitive ability.

**4.3.6. Psychomotor Effects**

25 There is considerable evidence in the literature for both animals and humans on  
26 psychomotor testing although human and laboratory animal studies utilize very different  
27 measures of motor behavior. Generally, the human literature employs a wide variety of  
28 psychomotor tasks and assesses error rates and reaction time in the performance of the task. The  
29 laboratory animal data, by contrast, tend to include unlearned naturalistic behaviors such as  
30 locomotor activity, gait changes, and foot splay to assess neuromuscular ability.  
31

#### 4.3.6.1.1. Psychomotor Effects: Human Studies

1           The effects of TCE exposure on psychomotor response have been studied primarily as a  
2 change in reaction time (RT) with studies on motor dyscoordination resulting from TCE  
3 exposure providing subjective reporting.

#### 4.3.6.1.2. Reaction time

4           Several studies have evaluated the effects of TCE on reaction time using simple and  
5 choice reaction time tasks (simple reaction time [SRT] and CRT tasks). The studies are  
6 presented below and summarized in more detail in Table 4-29.

7           Increases in reaction time were observed in environmental exposure studies by Kilburn  
8 (2002b), Kilburn and Warshaw (1993), and Kilburn and Thornton (1996) as well as in an  
9 occupational exposure study by Gun et al. (1978). All populations except that of Gun et al.  
10 (1978) were exposed through groundwater contaminated as the result of environmental spills and  
11 the exposure duration was for at least 1 year and exposure levels ranged from 0.2–10,000 ppb for  
12 the three studies. Kilburn and Warshaw (1993) reported that SRT significantly increased from  
13  $281 \pm 55$  msec to  $348 \pm 96$  msec in individuals ( $p < 0.0001$ ). CRT of the exposed subjects was  
14 93 msec longer ( $p < 0.0001$ ) than referents. Kilburn and Thornton (1996) evaluated SRT and  
15 CRT function and also found similar increases in reaction time. The average SRT and CRT for  
16 the combined control groups were 276 msec and 532 msec, respectively. These reaction times  
17 increased in the TCE exposure group where the average SRT was 334 msec and CRT was  
18 619 msec. Similarly, Kilburn (2002b) compared reaction times between 236 TCE-exposed  
19 persons and the 161 unexposed regional controls. SRTs significantly increased from  
20  $283 \pm 63$  msec in controls to  $334 \pm 118$  msec in TCE exposed individuals ( $p < 0.0001$ ).  
21 Similarly, CRTs also increased from  $510 \pm 87$  msec to  $619 \pm 153$  msec with exposure to TCE  
22 ( $p < 0.0001$ ).

23           No effect on SRT was reported in a geographical-based study by Reif et al. (2003). SRTs  
24 were 301msec for the lowest exposure group and 316 msec for the highest exposure group  
25 ( $p = 0.42$ ). When the SRT data were analyzed individuals that consumed at least on alcoholic  
26 drink per month ( $n = 80$ ), a significant increase (18%,  $p < 0.04$ ) in SRT times were observed  
27 between the lowest exposure and the highest exposure groups. In TCE exposed individuals who  
28 did not consume alcohol ( $n = 55$ ), SRTs decreased from 321 msec in the lowest exposed group to  
29 296 msec in the highest exposed group, but this effect was not statistically significantly different.  
30 A controlled exposure (chamber study) of 15 healthy men aged 20–31 years old, were exposed to  
31 0, 540, and 1,080  $\text{mg}/\text{m}^3$  TCE for 70 minutes or served as his own control, reported no  
32 statistically significant differences with the SRT or CRT tasks. However, in the RT-Addition

*This document is a draft for review purposes only and does not constitute Agency policy.*

- 1 test the level of performance varied between the different exposure conditions ( $F(2.24) = 4.35$ ;
- 2  $p < 0.05$ ) and between successive measurement occasions ( $F(2.24) = 19.25$ ;  $p < 0.001$ ).
- 3

1  
2

**Table 4-29. Summary of human choice reaction time studies**

Reference	Subjects	Exposure	Effect
Kilburn (2002b)	236 residents near a microchip plant in Phoenix, AZ. Controls: 161 regional referents from Wickenburg, AZ. 67 referents from Phoenix, AZ not residing near a plant.	0.2–10,000 ppb of TCE, chronic exposure.	Simple and choice reaction times were increased in the exposed group ( $p < 0.05$ ).
Kilburn and Warshaw (1993)	160 residents living in Southwest Tucson with TCE and other solvents in groundwater. Control: 68 residential referents matched to subjects from two previous studies of waste oil and oil refinery exposures.	>500 ppb of TCE in well-water before 1981 and 25–100 ppb afterwards. Exposure duration ranged from 1–25 yr.	Mean simple reaction time was 67 milliseconds (msec) longer than the referent group ( $p < 0.0001$ ). CRT of the exposed subjects was between 93–100 msec longer in three different trials ( $p < 0.0001$ ) compared to referents.
Reif et al. (2003)	143 residents of the Rocky Mountain Arsenal community of Denver. Referent group at lowest concentration (<5 ppb).	High exposure group >15 ppb. Medium exposure group $\geq 5$ ppb and $\leq 15$ ppb. Low exposure referent group <5 ppb.	Significant increase in reaction time as measured by the simple reaction time test ( $p < 0.04$ ) in only among subjects who reported alcohol use (defined as having at least one drink per mo).
Kilburn and Thornton (1996)	Group A: Registered voters from Arizona and Louisiana with no exposure to TCE: $n = 264$ , aged 18–83. Group B volunteers from California $n = 29$ (17 males and 12 females). Group C: exposed to TCE and other chemicals for 5 yr or more $n = 217$ .	No exposure or groundwater analyses reported.	Significant increase in simple and choice reaction time in exposed group compared to the unexposed populations.
Gamberale et al. (1976)	15 healthy men aged 20–31 yr old. Controls: Within subjects (15 self-controls).	0 mg/m <sup>3</sup> , 540 mg/m <sup>3</sup> (97 ppm), 1,080 mg/m <sup>3</sup> (194 ppm), 70 min.	No change in CRT or SRT. Increase in time required to perform the RT-Addition Test (task for adding numbers) ( $p < 0.05$ ).

3

**Table 4-29. Summary of human choice reaction time studies (continued)**

Reference	Subjects	Exposure	Effect
Gun et al. (1978)	Four female workers from one plant exposed to TCE and four female workers from another plant exposed to TCE + nonhalogenated hydrocarbon solvent. Control: ( $n = 8$ ) four unexposed female workers from each plant.	3–419 ppm, duration not specified.	TCE-only exposure increased reaction time in comparison to controls. In TCE + solvent group, ambient TCE was lower and mean reaction time shortened in Session 2, then rose subsequently to be greater than at the start.

1  
2

**4.3.6.1.3. Muscular dyscoordination**

3 Three studies examined motor dyscoordination effects from TCE exposure using  
4 subjective and self-reported individual assessment. Rasmussen et al. (1993a) presented findings  
5 on muscular dyscoordination for 96 metal degreasers exposed to either TCE or CFC113. A  
6 statistically significant increasing trend of dyscoordination with TCE exposure was observed  
7 ( $p = 0.01$ ) in multivariate regression analyses which adjusted for the effects of age, neurological  
8 disease, arteriosclerotic disease, and alcohol abuse. Furthermore, a greater number of abnormal  
9 coordination tests were observed in the higher exposure group compared to the low exposure  
10 group ( $p = 0.003$ ).

11 Gash et al. (2008) reported fine motor hand movement times in subjects who had filed  
12 workman compensation claims were significantly slower ( $p < 0.0001$ ) than age-matched  
13 nonexposed controls. Exposures were based on self-reported information, and no information on  
14 the control group is presented. Troster and Ruff (1990) reported a case study conducted on  
15 two occupationally exposed workers to TCE. Mild deficits in motor speed were reported for  
16 both cases. In the first case, manual dexterity was impaired in a male exposed to TCE (unknown  
17 concentration) for eight months. In the second case study where a female was exposed to TCE  
18 (low concentration; exact level not specified) for 3 months, there was weakness in the quadriceps  
19 muscle as evaluated in a neurological exam and a decreased sensation to touch on one hand.  
20 Both Gash et al. (2008) and Troster and Ruff (1990) provide very limited information given their  
21 deficiencies related to lack of exposure data, self-reported information, and limited reporting of  
22 referents and statistical analysis.

23

#### 4.3.6.1.4. Psychomotor Effects: Laboratory Animal Data

1 Several animal studies have demonstrated that TCE exposure produces changes in  
2 psychomotor function. At high doses ( $\geq 2,000$  mg/kg) TCE causes mice to lose their righting  
3 reflex when the compound is injected intraperitoneally (Shih et al., 2001; Umezu et al., 1997).  
4 At lower exposures (inhalation and oral), TCE produces alterations in neurobehavioral measures  
5 including locomotor activity, gait, operant responding, and reactivity. The studies are described  
6 in Sections 4.3.6.2.1–4.3.6.2.3 and summarized in Tables 4-30 and 4-31.  
7

#### 4.3.6.1.5. Loss of righting reflex

8 Umezu et al. (1997) studied disruption of the righting reflex following acute injection  
9 (i.p.) of 2,000, 4,000, and 5,000 mg/kg TCE in male ICR mice. TCE disrupted the righting  
10 reflex at doses of 2,000 mg/kg and higher. At 2,000 mg/kg, loss of righting reflex (LORR) was  
11 observed in only 2/10 animals injected. At 4,000 mg/kg, 9/10 animals experienced LORR and  
12 100% of the animals experienced LORR at 5,000 mg/kg.

13 Shih et al. (2001) reported impaired righting reflexes at exposure doses of 5,000 mg/kg  
14 (i.p.) in male MF1 mice. Mice pretreated with dimethyl sulfoxide or disulfuram (CYP2E1  
15 inhibitor) delayed LORR in a dose related manner. By contrast, the alcohol dehydrogenase  
16 inhibitor, 4-methylpyridine did not delay LORR that resulted from 5,000 mg/kg TCE. These data  
17 suggest that the anesthetic properties of TCE involve its oxidation via CYP2E1 to an active  
18 metabolite.  
19

#### 4.3.6.1.6. Activity, sensory-motor and neuromuscular function

20 Changes in sensory-motor and neuromuscular activity was reported in three studies  
21 (Kishi et al., 1993; Moser et al., 1995; Moser et al., 2003). Kishi et al. (1993) exposed male  
22 Wistar rats to 250, 500, 1,000, 2,000, and 4,000 ppm TCE for 4 hours. Rats exposed to 250-ppm  
23 TCE showed a significant decrease both in the total number of lever presses and in avoidance  
24 responses at 140 minutes of exposure compared with controls. Moser et al. (1995) evaluated the  
25 effects of acute and short-term (14 day) administration of TCE in adult female Fischer 344 rats  
26 ( $n = 8-10/\text{dose}$ ) on activity level, neuromuscular function and sensorimotor function as part of a  
27 larger functional observational battery (FOB) testing. The NOAEL levels identified by the  
28 authors are 500 mg/kg (10% of the limit dose) for the acute treatment and 150 mg/kg (3% of the  
29 limit dose) for the 14-day study. In the acute study, TCE produced the most significant effects in

1 motor activity (activity domain), gait (neuromuscular domain), and click response (sensorimotor  
2 domain). In the 14-day study, only the activity domain (rearing) and neuromuscular domain  
3 (forelimb grip strength) were significantly different ( $p < 0.05$ ) from control animals. In a  
4

1  
2  
3

**Table 4-30. Summary of animal psychomotor function and reaction time studies**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Savolainen et al. (1977)	Inhalation	Rat, Sprague-Dawley, male, 10	0, 200 ppm; 6 h/d, 4 d	LOAEL: 200 ppm	Increased frequency of preening, rearing, and ambulation. Increased preening time.
Kishi et al. (1993)	Inhalation	Rats, Wistar, male, number not specified	0, 250, 500, 1,000, 2,000, and 4,000 ppm, 4 hours	LOAEL: 250 ppm	Decreased lever presses and increased responding when lever press coupled with a 10-s electric shock (decreased avoidance response).
Kulig et al. (1987)	Inhalation	Rat, Wistar, male, 8/dose	0, 500, 1,000, and 1,500 ppm; 16 h/d, 5 d/wk, 18 wk	NOAEL: 1,500 ppm	No change in spontaneous activity, grip strength, or hindlimb movement.
Moser et al. (1995)	Oral	Rat, Fischer 344, female, 8/dose	0, 150, 500, 1,500, and 5,000 mg/kg, one dose	NOAEL: 500 mg/kg LOAEL: 1,500 mg/kg	<b>Decreased motor activity; Neuro-muscular and sensorimotor impairment.</b>
			0, 50, 150, 500, and 1,500 mg/kg-day, 14 d	NOAEL: 150 mg/kg-day LOAEL: 500 mg/kg-day	<b>Increased rearing activity and decreased forelimb grip strength.</b>
Bushnell (1997)	Inhalation	Rat, Long Evans, male, 12	0, 400, 800, 1,200, 1,600, 2,000, or 2,400 ppm, 1 h/test d, 4 consecutive test days, 2 wk	NOAEL: 800 ppm LOAEL: 1,200 ppm	Decreased sensitivity and increased response time in the signal detection task.
Shih et al. (2001)	Intra-peritoneal	Mouse, MF1, male, 6	0, 5,000 mg/kg, acute	LOAEL: 5,000 mg/kg	Impairment of righting reflex.
Umezu et al. (1997)	Intra-peritoneal	Mouse, ICR, male, 10/group	0, 2,000, 4,000, 5,000 mg/kg—loss of righting reflex measure	LOAEL: 2,000 mg/kg—loss of righting reflex	Loss of righting reflex.
		Mouse, ICR, male, 6–10/group	0, 62.5, 125, 250, 500, and 1,000 mg/kg, single dose and evaluated 30 min postadministration	NOAEL: 500 mg/kg LOAEL: 1,000 mg/kg—operant behavior NOAEL: 125 mg/kg LOAEL: 250 mg/kg—punished responding	Decreased responses (lever presses) in an operant response task for food reward. Increased responding when lever press coupled with a 20-V electric shock (punished responding).

4



**Table 4-30. Summary of animal psychomotor function and reaction time studies (continued)**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Bushnell and Oshiro (2000)	Inhalation	Rat, Long Evans, male, 32	0, 2,000, 2,400 ppm; 70 min/d, 9 d	LOAEL: 2,000 ppm	Decreased performance on the signal detection task. Increased response time and decreased response rate.
<b>Nunes et al. (2001)</b>	<b>Oral</b>	<b>Rat, Sprague-Dawley, male, 10/group</b>	<b>0, 2,000 mg/kg-day, 7 d</b>	<b>LOAEL: 2,000 mg/kg-day</b>	<b>Increased foot splay. No change in any other FOB parameter (e.g., piloerection, activity, reactivity to handling).</b>
Moser et al. (2003)	Oral	Rat, Fischer 344, female, 10/group	0, 40, 200, 800, and 1,200 mg/kg-day, 10 d	---	Decreased motor activity; Decreased sensitivity to tail pinch; Increased abnormality in gait; Decreased grip strength; Adverse changes in several FOB parameters.
Albee et al. (2006)	Inhalation	Rat, Fischer 344, male and female, 10/sex/group	0, 250, 800, 2,500 ppm; 6 h/d, 5 d/wk, 13 wk	NOAEL: 2,500 ppm	No change in any FOB measured parameter.

1  
2  
3  
4  
5

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

FOB = functional observational battery.

1  
2

**Table 4-31. Summary of animal locomotor activity studies**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Wolff and Siegmund (1978)	Intra-peritoneal	Mouse, AB, male, 18	0, 182 mg/kg, tested 30 min after injection	LOAEL: 182 mg/kg	Decreased spontaneous motor activity.
<b>Kulig et al. (1987)</b>	<b>Inhalation</b>	<b>Rat, Wistar, male, 8/dose</b>	<b>0, 500, 1,000, and 1,500 ppm; 16 h/d, 5 d/wk, 18 wk</b>	<b>NOAEL: 500 ppm LOAEL: 1,000 ppm</b>	<b>No change in spontaneous activity, grip strength or hindlimb movement. Increased latency time in the 2-choice visual discrimination task (cognitive disruption and/or motor activity related effect).</b>
Moser et al. (1995)	Oral	Rat, Fischer 344, female, 8/dose	0, 150, 500, 1,500, and 5,000 mg/kg, one dose	NOAEL: 500 mg/kg LOAEL: 1,500 mg/kg	Decreased motor activity; Neuro-muscular and sensorimotor impairment.
			0, 50, 150, 500, and 1,500 mg/kg-day, 14 d	NOAEL: 150 mg/kg-day LOAEL: 500 mg/kg-day	Increased rearing activity.
<b>Waseem et al. (2001)</b>	Oral	Rat, Wistar, male, 8/group	0, 350, 700, and 1,400 ppm in drinking water for 90 d	NOAEL: 1,400 ppm	No significant effect on spontaneous locomotor activity.
	<b>Inhalation</b>	<b>Rat, Wistar, male, 8/group</b>	<b>0, 376 ppm for up to 180 d; 4 h/d, 5 d/wk</b>	<b>LOAEL: 376 ppm</b>	<b>Changes in locomotor activity and vary by timepoint when measured over the 180-d period.</b>
Moser et al. (2003)	Oral	Rat, Fischer 344, female, 10/group	0, 40, 200, 800, and 1,200 mg/kg-day, 10 d	—	Decreased motor activity; Decreased sensitivity; Increased abnormality in gait; Adverse changes in several FOB parameters.

3  
4  
5  
6  
7  
8  
9  
10  
11  
12

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

FOB = functional observational battery.

separate 10-day study (Moser et al., 2003), TCE administration significantly ( $p < 0.05$ ) reduced motor activity, tail pinch responsiveness, reactivity to handling, hind limb grip strength and body weight. Significant increases ( $p < 0.05$ ) in piloerection, gait scores, lethality, body weight loss, and lacrimation was also reported in comparison to controls.

1           There are also two negative studies which used adequate numbers of subjects in their  
2 experimental design but used lower doses than did Moser et al. (2003). Albee et al. (2006)  
3 exposed male and female Fischer 344 rats ( $n = 10/\text{sex}$ ) to TCE by inhalation at exposure doses of  
4 250, 800, and 2,500 ppm, for 6 hours/day, 5 days/week, for 13 weeks. The FOB was performed  
5 monthly although it is not certain how much time elapsed from the end of exposure until the  
6 FOB test was conducted. No treatment related differences in grip strength or landing foot splay  
7 were demonstrated in this study. Kulig et al. (1987) also failed to show significant effects of  
8 TCE inhalation exposure on markers of motor behavior. Wistar rats ( $n = 8$ ) exposed to 500,  
9 1,000, and 1,500 ppm, for 16 hours/day, 5 days/week, for 18 weeks failed to show changes in  
10 spontaneous activity, grip strength, or coordinated hind limb movement. Measurements were  
11 made every three weeks during the exposure period and occurred between 45 and 180 minutes  
12 following the previous TCE inhalation exposure.  
13

#### **4.3.6.1.7. Locomotor activity**

14           The data, with regard to locomotor activity, are inconsistent. Several studies showed that  
15 TCE exposure can decrease locomotor activity including Wolff and Siegmund (1978) where AB  
16 mice ( $n = 18$ ) were treated acutely with a dose of 182 mg/kg, i.p. at one of four time points  
17 during a 24-hour day. Moser et al. (1995; 2003) reported reduced locomotor activity in female  
18 Fischer 344 rats ( $n = 8-10$ ) gavaged with TCE over an acute (LOAEL = 5,000 mg/kg TCE) or  
19 subacute period (LOAEL = 500 but no effect at 5,000 mg/kg). In the Moser et al. (2003), it  
20 appears that 200-mg/kg TCE yielded a significant reduction in locomotor activity and that the  
21 degree of impairment at this dose represented a maximal effect on this measure. That is, higher  
22 doses of TCE appear to have produced equivalent or slightly less of an effect on this behavior.  
23 While this study identifies a LOAEL of 200-mg/kg TCE by gavage over a 10-day period, this is  
24 a much more lower dose effect than that reported in Moser et al. (1995). Both studies (Moser et  
25 al., 1995; Moser et al., 2003) demonstrate a depression in motor activity that occurs acutely  
26 following TCE administration. Kulig et al. (1987) demonstrated that rats had increased response  
27 latency to a two choice visual discrimination following 1,000- and 1,500-ppm TCE exposures for  
28 18 weeks. However, no significant changes in grip strength, hindlimb movement, or any other  
29 motor activity measurements were noted.

30           There are also a few studies (Fredriksson et al., 1993; Waseem et al., 2001) generally  
31 conducted using lower exposure doses that failed to demonstrate impairment of motor activity or  
32 ability following TCE exposure. Waseem et al. (2001) failed to demonstrate changes in  
33 locomotor activity in male Wistar rats ( $n = 8$ ) dosed with TCE (350, 700, and 1,400 ppm) in

1 drinking water for 90 days. Wistar rats ( $n = 8$ ) exposed to 500, 1,000, and 1,500 ppm for  
2 16 hours/day, 5 days/week, for 18 weeks failed to show changes in spontaneous activity. No  
3 changes in locomotor activity were observed for 17-day-old male NMRI mice that were dosed  
4 postnatally with 50 or 290 mg/kg-day from Day 10–16 (Fredriksson et al., 1993). However,  
5 rearing activity was significantly decreased in the NMRI mice at Day 60.  
6

#### 4.3.6.1.8. Summary and Conclusions for Psychomotor Effects

7 In human studies, psychomotor effects such as reaction time and muscular  
8 dyscoordination have been examined following TCE exposure. In the reaction time studies,  
9 statistically significant increases in CRT and SRT were reported in the Kilburn studies (Kilburn,  
10 2002b; Kilburn and Thornton, 1996; Kilburn and Warshaw, 1993). All of these studies were  
11 geographically based and it was suggested that the results were used for litigation and the  
12 differences between exposed and referent groups on other factors influencing reaction speed time  
13 may introduce a bias to the findings. Additionally, in these studies exposure to TCE and other  
14 chemicals occurred through drinking water for at least 1 year and TCE concentrations in well  
15 water ranged from 0.2 ppb to 10,000 ppb. Reif et al. (2003) whose exposure assessment  
16 approach included exposure modeling of water distribution system to estimate TCE  
17 concentrations in tap water at census track of residence found that residents with drinking water  
18 containing TCE (up to >15 ppb—the highest level not specified) and other chemicals did not  
19 significantly increase CRTs or SRTs. Inhalation studies also demonstrated increased reaction  
20 times. An acute exposure chamber study (Gamberale et al., 1976) tested for CRT, SRT, and  
21 RT-addition following a 70-minute exposure to TCE. A concentration-dependent significant  
22 decrease in performance was observed with the RT-addition test and not for CRT or SRT tasks.  
23 An occupational exposure study on eight female workers exposed to TCE (Gun et al., 1978) also  
24 reported increased reaction time in the females exposed to TCE-only. Muscular dyscoordination  
25 for humans following TCE exposure has been reported in a few studies as a subjective  
26 observation. The studies indicated that exposure resulted in decreased motor speed and dexterity  
27 (Rasmussen et al., 1993a; Troster and Ruff, 1990) and self-reported faster asymptomatic fine  
28 motor hand movements (Gash et al., 2008).

29 Animal studies evaluated psychomotor function by examining locomotor activity, operant  
30 responding, changes in gait, loss of righting reflex, and general motor behavior (see Tables 4-30  
31 and 4-31 for references). Overall, the studies demonstrated that TCE causes loss of righting  
32 reflex at injection doses of 2,000 mg/kg or higher (Shih et al., 2001; Umezu et al., 1997).  
33 Regarding general psychomotor testing, significant decreases in lever presses and avoidance

1 were observed at inhalation exposures as low as 250 ppm for 4 hours (LOAEL; Kishi et al.,  
2 1993). Following subchronic inhalation exposures, no significant changes in psychomotor  
3 activity were noted at up to 2,500 ppm for 13 weeks (Albee et al., 2006) or at 1,500 ppm for  
4 18 weeks (Kulig, 1987). In the oral administration studies (Moser et al., 1995; Moser et al.,  
5 2003), psychomotor effects were evaluated using an FOB. More psychomotor domains were  
6 significantly affected by TCE treatment in the acute study in comparison to the 14-day study, but  
7 a lower NOAEL (150 mg/kg-day) was reported for the 14-day study in comparison to the acute  
8 study (500 mg/kg; Moser et al., 1995). Upon closer examination of the data, a biphasic effect in  
9 one measure of the FOB (rearing) was resulting in the lower NOAEL for the 14-day study and  
10 doses that were higher and lower than the NOAEL did not produce a statistically significant  
11 increase in the number of rears. Therefore, it can be surmised that acute exposure to TCE results  
12 in significant changes in psychomotor function. However, there may be some tolerance to these  
13 psychomotor changes in increased exposure duration to TCE as evidenced by the results noted in  
14 the short-term and subchronic exposure studies.

15

#### **4.3.7. Mood Effects and Sleep Disorders**

##### **4.3.7.1.1. Effects on Mood: Human Studies**

16 Reports of mood disturbance (depression, anxiety) resulting from TCE exposure are  
17 numerous in the human literature. These symptoms are subjective and difficult to quantify.  
18 Studies by Gash et al. (2008), Kilburn and Warshaw (1993), Kilburn (2002a, b), McCunney et al.  
19 (1988), Mitchell et al. (1969), Rasmussen and Sabroe (1986), and Troster and Ruff (1990)  
20 reported mood disturbances in humans. Reif et al. (2003) and Triebig (1976) reported no effect  
21 on mood following TCE exposures.

22

##### **4.3.7.1.2. Effects on Mood: Laboratory Animal Findings**

23 It is difficult to obtain comparable data of emotionality in laboratory studies. However,  
24 Moser et al. (2003) and Albee et al. (2006) both report increases in handling reactivity among  
25 rats exposed to TCE. In the Moser study, female Fischer 344 rats received TCE by oral gavage  
26 for periods of 10 days at doses of 0, 40, 200, 800, and 1,200 while Albee et al. (2006) exposed  
27 Fischer 344 rats to TCE by inhalation at exposure doses of 250, 800, and 2,500 ppm for

1 6 hours/day, 5 days/week, for 13 weeks. These studies are summarized and described in  
 2 Table 4-32.  
 3

**4.3.7.1.3. Sleep Disturbances**

4 Arito et al. (1994) exposed male Wistar rats to 50-, 100-, and 300-ppm TCE for  
 5 8 hour/day, 5 days/week, for 6 weeks and measured electroencephalographic (EEG) responses  
 6 (see Table 4-32). EEG responses were used as a measure to determine the number of awake  
 7  
 8

**Table 4-32. Summary of animal mood effect and sleep disorder studies**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
<i>Mood Effects</i>					
Albee et al. (2006)	Inhalation	Rat, Fischer 344, male and female, 10/sex/group	0, 250, 800, 2,500 ppm; 6 h/d, 5 d/wk, 13 wk	NOAEL: 800 ppm	Increased handling reactivity.
Moser et al. (2003)	Oral	Rat, Fischer 344, female, 10/group	0, 40, 200, 800, and 1,200 mg/kg-day, 10 d	---	Decreased handling reactivity score.
<i>Sleep Disorder</i>					
Arito et al. (1994)	Inhalation	Rat, Wistar, male, 5/group	0, 50, 100, and 300 ppm; 8 h/d, 5 d/wk, for 6 wk	LOAEL: 50 ppm	Significant changes in sleep cycle as measured through EEG changes; Significant decreases in wakefulness.

9  
 10 **Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).  
 11

12  
 13 (wakefulness hours) and sleep hours. Exposure to all the TCE levels significantly decreased  
 14 amount of time spent in wakefulness (W) during the exposure period. Some carry over was  
 15 observed in the 22 hours post exposure period with significant decreases in wakefulness seen at  
 16 100-ppm TCE. Significant changes in W-sleep elicited by the long-term exposure appeared at  
 17 lower exposure levels. These data seem to identify a low dose effect of TCE and established a  
 18 LOAEL of 50 ppm for sleep changes.  
 19

### 4.3.8. Developmental Neurotoxicity

#### 4.3.8.1.1. Human Studies

1 In humans, CNS birth defects were observed in a few studies (ATSDR, 2001; Bove,  
2 1996; Bove et al., 1995; Lagakos et al., 1986). Postnatally, observed adverse effects in humans  
3 include delayed newborn reflexes following exposure to TCE during childbirth (Beppu, 1968),  
4 impaired learning or memory (Bernad et al., 1987, abstract; White et al., 1997); aggressive  
5 behavior (Bernad et al., 1987, abstract); hearing impairment (Burg and Gist, 1999); speech  
6 impairment (Burg and Gist, 1999; White et al., 1997); encephalopathy (White et al., 1997);  
7 impaired executive and motor function (White et al., 1997); attention deficit (Bernad et al., 1987,  
8 abstract; White et al., 1997); and autism spectrum disorder (Windham et al., 2006). The human  
9 developmental neurotoxicity studies are discussed in more detail in Section 4.8.2.1.2, and  
10 summarized in Table 4-33.

11 **Table 4-33. Summary of human developmental neurotoxicity associated with**  
12 **TCE exposures**  
13

Finding	Species	Citations
CNS defects, neural tube defects	Human	ATSDR (2001)
		Bove (1996); Bove et al. (1995)
		Lagakos et al. (1986)
Delayed newborn reflexes	Human	Beppu (1968)
Impaired learning or memory	Human	Bernad et al. (1987, abstract)
		White et al. (1997)
Aggressive behavior	Human	Bernad et al., (1987, abstract)
Hearing impairment	Human	Burg and Gist (1999)
Speech impairment	Human	Burg and Gist (1999)
		White et al. (1997)
Encephalopathy	Human	White et al. (1997)
Impaired executive function	Human	White et al. (1997)
Impaired motor function	Human	White et al. (1997)
Attention deficit	Human	White et al. (1997)
	Human	Bernad et al. (1987, abstract)
Autism spectrum disorder (ASD)	Human	Windham et al. (2006)

14  
15

#### 4.3.8.1.2. Animal Studies

1           There are a few studies demonstrating developmental neurotoxicity following  
2 trichloroethylene exposure (range of exposures) to experimental animals. These studies  
3 collectively suggest that developmental neurotoxicity result from TCE exposure, however, some  
4 types of effects such as learning and memory measures have not been evaluated. Most of the  
5 studies demonstrate either spontaneous motor activity changes (Taylor et al., 1985) or  
6 neurochemical changes such as decreased glucose uptake and changes in the specific gravity of  
7 the cortex and cerebellum (Isaacson and Taylor, 1989; Noland-Gerbec et al., 1986; Westergren et  
8 al., 1984). In addition, in most of these studies there is no assessment of the exposure to TCE or  
9 metabolites in the pups/offspring. Details of the studies are presented below and summarized in  
10 Table 4-34.

11           Taylor et al. (1985) administered TCE to female Sprague-Dawley rats in their drinking  
12 water from 14 days before breeding throughout gestation and until pups were weaned at 21 days.  
13 Measured TCE concentrations in the dams ranged from 312–646 mg/L, 625–1,102 mg/L, and  
14



1  
2  
3

**Table 4-34. Summary of mammalian in vivo developmental neurotoxicity studies—oral exposures**

Reference	Species/strain/ sex/number	Dose level/ exposure duration	NOAEL; LOAEL <sup>a</sup>	Effects
Fredriksson et al. (1993)	Mouse, NMRI, male pups, 12 pups from 3–4 different litters/group	0, 50, or 290 mg/kg-day PND 10–16	LOAEL: 50 mg/kg-day	Rearing activity sig. ↓ at both dose levels on PND 60.
George et al. (1986)	Rat, F334, male and female, 20 pairs/treatment group, 40 controls/sex	0, 0.15, 0.30, or 0.60% microencapsulated TCE in diet. Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females throughout pregnancy (i.e., 18 wk total).	LOAEL: 0.15%	Open field testing in pups: a sig. dose-related trend toward ↑ time required for male and female pups to cross the first grid in the test device.
Isaacson and Taylor (1989)	Rat, Sprague-Dawley, females, six dams/group	0, 312, or 625 mg/L (0, 4.0, or 8.1 mg/d) <sup>b</sup> . Dams (and pups) exposed from 14 d prior to mating until end of lactation.	LOAEL: 312 mg/L	Sig. ↓ myelinated fibers in the stratum lacunosum-moleculare of pups. Reduction in myelin in the CA1 region of the hippocampus.
Noland-Gerbec et al. (1986)	Rat, Sprague-Dawley, females, 9–11 dams/group	0, 312 mg/L. (Avg. total intake of dams: 825 mg TCE over 61 d.). Dams (and pups) exposed from 14 d prior to mating until end of lactation.	LOAEL: 312 mg/L	Sig. ↓ uptake of <sup>3</sup> H-2-DG in whole brains and cerebella (no effect in hippocampus) of exposed pups at 7, 11, and 16 d, but returned to control levels by 21 d.
Taylor et al. (1985)	Rat, Sprague-Dawley, females, no. dams/group not reported	0, 312, 625, and 1,250 mg/L in drinking water. Dams (and pups) exposed from 14 d prior to mating until end of lactation.	LOAEL: 312 mg/L	Exploratory behavior sig. ↑ in 60- and 90-d old male rats at all treatment levels. Locomotor activity (measured through the wheel-running tasks) was higher in rats from dams exposed to 1,250 mg/L TCE.
Blossom et al. (2008)	Mouse, MRL +/+, dams and both sexes offspring, eight litters/group; three–eight pups/group	Drinking water, from GD 0 to PND 42; 0 or 0.1 mg/mL; maternal dose = 25.7 mg/kg-day; offspring PND 24–42 dose = 31.0 mg/kg-day.	LOAEL: 31 mg/kg-day for offspring	Righting reflex, bar holding, and negative geotaxis were not impaired. Significant association between impaired nest quality and TCE exposure. Lower GSH levels and GSH:GSSG ratios with TCE exposure.

4  
5  
6  
7  
8

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup> LOEL (lowest-observed-effect level) are based upon reported study findings.

<sup>b</sup> Dose conversions provided by study author(s).

*This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2 GD = gestation day, GSSG = oxidized GSH, PND = postnatal day.  
3

1 1,250–1,991 mg/L in the low, mid, and high-dose groups as measured from the drinking water.  
2 Pups were evaluated for exploratory activity at 28, 60, or 90 days. No significant differences  
3 were noted between control and treated pups at 28 days. At 60 days, all TCE-treated animals  
4 had significantly increased exploratory activity in comparison to age-matched controls, but only  
5 the high group had increased activity at 90 days. A significant increase in spontaneous motor  
6 activity (as measured by a wheel-running task) was noted in only the high dose TCE  
7 (1,250–1,991 mg/L) group during the onset of the darkness period. This study demonstrated that  
8 both spontaneous and open field activities are significantly affected by developmental TCE  
9 exposure.

10 Spontaneous behavioral changes were also investigated in another study by Fredriksson  
11 et al. (1993). Male and female NMRI pups (mice) were orally administered 50 or  
12 290 mg/kg-day for 7 days starting at PND 10. Spontaneous motor activity was investigated in  
13 male mice at ages 17 and 60 days. TCE-treated animals tested at Day 17 did not demonstrate  
14 changes in any spontaneous activity measurements in comparison to control animals. Both doses  
15 of TCE (50 and 290 mg/kg-day) significantly decreased rearing in 60 day-old male mice.

16 Westergren et al. (1984) examined the brain specific gravity of litters from mice exposed  
17 to TCE. NMRI mice (male and female) were exposed to 150-ppm TCE (806.1 mg/m<sup>3</sup>) for  
18 30 days prior to mating. Exposure in males continued until the end of mating and females were  
19 exposed until the litters were born. Brains were removed from the offspring at either PNDs 1,  
20 10, 20–22, or 29–31. At PNDs 1 and 10, significant decreases were noted in the specific gravity  
21 of the cortex. Significant decreases in the specific gravity of the cerebellum were observed at  
22 PND 10 (decrease from  $1.0429 \pm 0.00046$ – $1.0405 \pm 0.00030$ ) and 20–22 (decrease from  $1.0496$   
23  $\pm 0.00014$ – $1.0487 \pm 0.00060$ ). Cerebellum measurements were not reported for PND 29–31  
24 animals. Neurobehavioral assessments were not conducted in this study. Additionally,  
25 decreased brain specific gravity is suggestive of either decreased brain weight or increased brain  
26 volume (probably from edema) or a combination of the two factors and is highly suggestive of an  
27 adverse neurological effect. The effects of TCE on the cortical specific gravity were not  
28 persistent since cortices from PND 29–31 animals did not exhibit any significant changes. It is  
29 unclear if the effects on the cerebellum were persistent since results were not reported for the  
30 PND 29–31 animals. However, the magnitude of the change in the specific gravity of the  
31 cerebellum is decreased from PND 10 to PND 20–22 suggesting that the effect may be reversible  
32 given a longer recovery period from TCE.

33 The effect of TCE on glucose uptake in the brain was evaluated in rat pups exposed to  
34 TCE during gestation and through weaning. The primary source of energy utilized in the CNS is  
35 glucose. Changes in glucose uptake in the brain are a good indicator for neuronal activity

1 modification. Noland-Grebec et al. (1986) administered 312 mg/L TCE through drinking water  
2 to female Sprague-Dawley rats from 2 weeks before breeding and up until pups reached 21 days  
3 of age. To measure glucose uptake, 2-deoxyglucose was administered intraperitoneally to male  
4 pups at either postnatal day (PND) 7, 11, 16, or 21. Significant decreases in glucose uptake were  
5 noted in whole brain and cerebellum at all postnatal days tested. Significant decreases in glucose  
6 uptake were also observed in the hippocampus except for animals tested at PND 21. The  
7 observed decrease in glucose uptake suggests decreased neuronal activity.

8 Female Sprague-Dawley rats (70 days old) were administered TCE in drinking water at a  
9 level of either 4.0 or 8.1 mg/day for 14 days prior to mating and continuing up through lactation  
10 (Isaacson and Taylor, 1989). Only the male pups were evaluated in the studies. At PND 21,  
11 brains were removed from the pups, sectioned, and stained to evaluate the changes in myelin.  
12 There was a significant decrease (40% decrease) in myelinated fibers in the CA1 region of the  
13 hippocampus of the male pups. This effect appeared to be limited to the CA1 region of the  
14 hippocampus since other areas such as the optic tract, fornix, and cerebral peduncles did not have  
15 decreases in myelinated fibers.

16 Neurological changes were found in pups exposed to TCE in a study conducted by the  
17 National Toxicology Program (NTP) in Fischer 344 rats (George et al., 1986). TCE was  
18 administered to rats at dietary levels of 0, 0.15, 0.30, or 0.60%. No intake calculations were  
19 presented for the rat study and therefore, a dose rate is unavailable for this study. Open field  
20 testing revealed a significant ( $p < 0.05$ ) dose-related trend toward an increase in the time required  
21 for male and female F1 weanling pups (PND 21) to cross the first grid in the testing device,  
22 suggesting an effect on the ability to react to a novel environment.

23 Blossom et al. (2008) treated male and female MRL  $+/+$  mice with 0 or 0.1 mg/mL TCE  
24 in the drinking water. Treatment was initiated at the time of mating, and continued in the  
25 females (8/group) throughout gestation and lactation. Behavioral testing consisted of righting  
26 reflex on PNDs 6, 8, and 10; bar-holding ability on PNDs 15 and 17; and negative geotaxis on  
27 PNDs 15 and 17. Nest building was assessed and scored on PND 35, the ability of the mice to  
28 detect and distinguish social odors was examined with an olfactory habituation/dishabituation  
29 method at PND 29, and a resident intruder test was performed at PND 40 to evaluate social  
30 behaviors. Righting reflex, bar holding, and negative geotaxis were not impaired by treatment.  
31 There was a significant association between impaired nest quality and TCE exposure in tests of  
32 nest-building behavior; however, TCE exposure did not have an effect on the ability of the mice  
33 to detect social and nonsocial odors using habituation and dishabituation methods. Resident  
34 intruder testing identified significantly more aggressive activities (i.e., wrestling and biting) in  
35 TCE-exposed juvenile male mice as compared to controls, and the cerebellar tissue from the

1 male TCE-treated mice had significantly lower GSH levels and GSH:oxidized GSH  
2 (GSH:GSSG) ratios, indicating increased oxidative stress and impaired thiol status, which have  
3 been previously reported to be associated with aggressive behaviors (Franco et al., 2006).  
4 Histopathological examination of the brain did not identify alterations indicative of neuronal  
5 damage or inflammation.  
6

#### **4.3.8.1.3. Summary and Conclusions for the Developmental Neurotoxicity Studies**

7 Gestational exposure to TCE in humans has resulted in several developmental  
8 abnormalities. These changes include neuroanatomical changes such as neural tube defects  
9 (ATSDR, 2001; Bove et al., 1995, 1996; Lagakos et al., 1986) and encephalopathy (White et al.,  
10 1997). Clinical neurological changes such as impaired cognition (Bernad et al., 1987; White et  
11 al., 1997), aggressive behavior (Bernad et al., 1987), and speech and hearing impairment (Burg  
12 and Gist, 1999; White et al., 1997) are also observed when TCE exposure occurs in utero.

13 In animal studies, anatomical and clinical developmental neurotoxicity is also observed.  
14 Following inhalation exposures of 150 ppm to mice during mating and gestation, the specific  
15 gravity of offspring brains was significantly decreased at postnatal time points through the age of  
16 weaning; this effect did not persist to 1 month of age (Westergren et al., 1984). In studies  
17 reported by Taylor et al. (1985), Isaacson and Taylor (1989), and Noland-Gerbec et al. (1986),  
18 312 mg/L exposures in drinking water that were initiated 2 weeks prior to mating and continued  
19 to the end of lactation resulted in (a) significant increase in exploratory behavior at PNDs 60 and  
20 90, (b) reductions in myelination in the CA1 hippocampal region of offspring at weaning, and  
21 (c) significantly decreased uptake of 2-deoxyglucose in the rat brain at PND 21. Gestational  
22 exposures to mice (Fredriksson et al., 1993) resulted in significantly decreased rearing activity  
23 on PND 60, and dietary exposures during the course of a continuous breeding study in rats  
24 (George et al., 1986) found a significant trend toward increased time to cross the first grid in  
25 open field testing. In a study by Blossom et al. (2008), male mice exposed gestationally to TCE  
26 exhibited lower GSH levels and lower GSH:GSSG ratios which is also observed in mice that  
27 have more aggressive behaviors (Franco et al., 2006).  
28

### **4.3.9. Mechanistic Studies of Trichloroethylene (TCE) Neurotoxicity**

#### **4.3.9.1.1. Dopamine Neuron Disruption**

1           There are very recent laboratory animal findings resulting from short-term TCE  
2 exposures that demonstrate vulnerability of dopamine neurons in the brain to this chlorinated  
3 hydrocarbon. The key limitation of these laboratory animal studies is that only one dosing  
4 regimen was included in each study. Moreover, there has been no systematic body of data to  
5 show that other chlorinated hydrocarbons such as tetrachloroethylene or aromatic solvents  
6 similarly target this cell type. Confidence in the limited data regarding dopamine neuron death  
7 and in vivo TCE exposure would be greatly enhanced by identifying a dose-response  
8 relationship. If indeed TCE can target dopamine neurons it would be anticipated that human  
9 exposure to this agent would result in elevated rates of parkinsonism. There are no systematic  
10 studies of this potential relationship in humans although one limited report attempted to address  
11 this possibility. Difficulties in subject recruitment into that study limit the weight that can be  
12 given to the results.

13           Endogenously formed chlorinated tetrahydro-beta-carbolines (TaClo) have been  
14 suggested to contribute to the development of Parkinson-like symptoms Bringmann et al. 1992  
15 (Bringmann et al., 1995; Kochen et al., 2003; Riederer et al., 2002). TaClo can be formed  
16 endogenously from metabolites of TCE such as trichloroacetaldehyde. TaClo has been  
17 characterized as a potent neurotoxicant to the dopaminergic system. Some research groups have  
18 hypothesized that Parkinson-like symptoms resulting from TCE exposure may occur through the  
19 formation of TaClo, but not enough evidence is available to determine if this mechanism occurs.  
20

#### **4.3.9.1.2. Dopamine neuron disruption: human studies**

21           There are no human studies that present evidence of this effect. Nagaya et al. (1990)  
22 examined serum dopamine  $\beta$ -hydroxylase activity without differences observed in mean  
23 activities between control and exposed subjects. In the study, 84 male workers exposed to TCE  
24 were compared to 83 male age-matched controls. The workers had constantly used TCE in their  
25 jobs and their length of employment ranged from 0.1–34 years.  
26

#### **4.3.9.1.3. Dopamine neuron disruption: animal studies**

27           There are limited data from mice and rats that suggest the potential for TCE to disrupt  
28 dopamine neurons in the basal ganglia (see Table 4-35). Gash et al. (2008) showed that TCE

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 gavage in Fischer 344 rats ( $n = 9$ ) at an exposure level of 1,000 mg/kg-day, 5 days/week, for  
 2 6 weeks yielded degeneration of dopamine neurons in the substantia nigra and alterations in  
 3 dopamine turnover as reflected in a shift in dopamine metabolite to parent compound ratios.  
 4 Guehl et al. (1999) reported similar findings in OF1 mice ( $n = 10$ ) that were injected i.p. with  
 5 400 mg/kg-day TCE 5 days/week for 4 weeks. Each of these studies evaluated only a single dose  
 6 level of TCE so that establishing a dose-response relationship is not possible. Consequently,  
 7 these data are of limited utility in risk assessment because they do not establish the potency of  
 8 TCE to damage dopamine neurons. They are important, however, in identifying a potential  
 9 permanent impairment that might occur following TCE exposure at relatively high exposure  
 10 doses. They also identify a potential mechanism by which TCE could produce CNS injury.

11 **Table 4-35. Summary of animal dopamine neuronal studies**

12

Reference	Exposure route	Species/strain/ sex/number	Dose level/ exposure duration	NOAEL; LOAEL	Effects
Guehl et al. (1999)	Intraperitoneal Administration	Mouse, OF1, male, 10	0 and 400 mg/kg; 5 d/wk, 4 wk	LOAEL: 400 mg/kg	Significant dopaminergic neuronal death in substantia nigra.
<b>Gash et al. (2008)</b>	<b>Oral gavage</b>	<b>Rat, Fischer 344, male, 9/group</b>	<b>0 and 1,000 mg/kg; 5 d/wk, 6 wk</b>	<b>LOAEL: 1,000 mg/kg</b>	<b>Degeneration of dopamine-containing neurons in substantia nigra. Change in dopamine metabolism.</b>

13 **Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).  
 14  
 15  
 16

#### 4.3.9.1.4. Summary and conclusions of dopamine neuron studies

17 Only two animal studies have reported changes in dopamine neuron effects from TCE  
 18 exposure (Gash et al., 2008; Guehl et al., 1999). Both studies demonstrated toxicity to  
 19 dopaminergic neurons in the substantia nigra in rats or mice. LOAELs of 400 mg/kg (mice;  
 20 Guehl et al., 1999) and 1,000 mg/kg (rats; Gash et al., 2008) were reported for this effect.  
 21 Dopaminergic neuronal degeneration following TCE exposure has not been studied in humans.  
 22 However, there were no changes in serum dopamine  $\beta$ -hydroxylase activity in TCE-exposed and  
 23 control individuals (Nagaya et al., 1990). Loss of dopaminergic neurons in the substantia nigra  
 24 also occurs in patients with Parkinson's disease and the substantia nigra is an important region in  
 25 helping to control movements. As a result, loss of dopaminergic neurons in the substantia nigra

1 may be one of the potential mechanisms involved in the clinical psychomotor effects that are  
 2 observed following TCE exposure.

3

#### 4.3.9.1.5. Neurochemical and Molecular Changes

4 There are limited data obtained only from laboratory animals that TCE exposure may  
 5 have consequences on GABAergic (gamma-amino butyric acid [GABA]) and glutamatergic  
 6 neurons (Briving et al., 1986; Shih et al., 2001; see Table 4-36). However, the data obtained are  
 7 limited with respect to brain region examined, persistence of effect, and whether there might be  
 8 functional consequences to these changes. The data of Briving et al. (1986) demonstrating  
 9 changes in cerebellar high affinity uptake for GABA and glutamate following chronic low level  
 10 (50 and 150 ppm) TCE exposure do not appear to be reflected in the only other brain region  
 11 evaluated (hippocampus). However, glutamate levels were increased in the hippocampus. The  
 12 data of Shih et al. (2001) are indirect in that it shows an altered response to GABAergic

13 **Table 4-36. Summary of neurophysiological, neurochemical, and**  
 14 **neuropathological effects with TCE exposure**

15

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Neurophysiological studies					
Shih et al. (2001)	Intra-peritoneal	Mouse, MF1, male, 6/group	0, 250 500, 1,000, or 2,000 mg/kg, 15 min; followed by tail infusion of PTZ (5 mg/mL), picrotoxin (0.8 mg/mL), bicuculline (0.06 mg/mL), strychnine (0.05 mg/mL), 4-AP (2 mg/mL), or NMDA (8 mg/mL).	---	Increased threshold for seizure appearance with TCE pretreatment for all convulsants. Effects strongest on the GABA <sub>A</sub> antagonists, PTZ, picrotoxin, and bicuculline suggesting GABA <sub>A</sub> receptor involvement. NMDA and glycine Rc involvement also suggested.
Ohta et al. (2001)	Intra-peritoneal	Mouse, ddY, male, 5/group	0, 300, or 1,000 mg/kg, sacrificed 24 h after injection.	LOAEL: 300 mg/kg	Decreased response (LTP response) to tetanic stimulation in the hippocampus.
Neurochemical studies					



Briving et al. (1986)	Inhalation	Gerbils, Mongolian, male and female, 6/group	0, 50, or 150 ppm, continuous, 24 h/d, 12 mos.	NOAEL: 50 ppm; LOAEL: 150 ppm for glutamate levels in hippocampus. NOAEL: 150 ppm for glutamate and GABA uptake in hippocampus. LOAEL: 50 ppm for glutamate and GABA uptake in cerebellar vermis.	Increased glutamate levels in the hippocampus. Increased glutamate and GABA uptake in the cerebellar vermis.
Subramoniam et al. (1989)	Oral	Rat, Wistar, female	0 or 1,000 mg/kg, 2 or 20 h. 0 or 1,000 mg/kg-day, 5 d/wk, 1 yr.	---	PI and PIP2 decreased by 24 and 17% at 2 h. PI and PIP2 increased by 22 and 38% at 20 h. PI, PIP, and PIP2 reduced by 52, 23, and 45% in 1 yr study.

1

**Table 4-36. Summary of neurophysiological, neurochemical, and neuropathological effects with TCE exposure (continued)**

Reference	Exposure route	Species/strain/sex/number	Dose level/exposure duration	NOAEL; LOAEL	Effects
Haglid et al. (1981)	Inhalation	Gerbil, Mongolian, male and female, 6–7/group	0, 60, or 320 ppm, 24 h/d, 7 d/wk, 3 mo.	LOAEL: 60 ppm, brain protein changes. NOAEL: 60 ppm; LOAEL: 320 ppm, brain DNA changes.	(1) Decreases in total brain soluble protein whereas increase in S100 protein. (2) Elevated DNA in cerebellar vermis and sensory motor cortex.
Neuropathological studies					
Kjellstrand et al. (1987)	Inhalation	Mouse, NMRI, male	0, 150, or 300 ppm, 24 h/d, 4 or 24 d.	LOAEL: 150 ppm, 4 and 24 d	Sciatic nerve regeneration was inhibited in both mice and rats.
		Rat, Sprague-Dawley, female	0, 300 ppm, 24 h/d, 4 or 24 d.	NOAEL: 300 ppm, 4 d. LOAEL: 300 ppm, 24 d.	
Isaacson and Taylor (1989)	Oral	Rat, Sprague-Dawley, females, six dams/group	0, 312, or 625 mg/L. (0, 4.0, or 8.1 mg/d). Dams (and pups) exposed from 14 d prior to mating until end of lactation.	LOAEL: 312 mg/L	Significant ↓ myelinated fibers in the stratum lacunosum-moleculare of pups. Reduction in myelin in the hippocampus.

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

NMDA = N-nitrosodimethylamine, PTZ = pentylenetetrazole.

antagonist drugs in mice treated by acute injection with 250, 500, 1,000, and 2,000 mg/kg TCE. However, these data do show some dose dependency with significant findings observed with TCE exposure as low as 250 mg/kg.

The development and physiology of the hippocampus has also been evaluated in two different studies (Isaacson and Taylor, 1989; Ohta et al., 2001). Isaacson and Taylor (1989) found a 40% decrease in myelinated fibers from hippocampi dissected from neonatal Sprague-Dawley rats ( $n = 2-3$ ) that were exposed to TCE (4 and 8.1 mg/day) in utero and during the preweaning period. Ohta et al. (2001) injected male ddY mice with 300 mg/kg TCE and found a significant reduction in response to titanic stimuli in excised hippocampal slices. Both of these studies demonstrated that there is some interaction with TCE and the hippocampal area in the brain.

1           Impairment of sciatic nerve regeneration was demonstrated in mice and rats exposed to  
2 TCE (Kjellstrand et al., 1987). Under heavy anesthesia, the sciatic nerve of the animals was  
3 artificially crushed to create a lesion. Prior to the lesion, some animals were pre-exposed to TCE  
4 for 20 days and then for an additional 4 days after the lesion. Another set of animals were only  
5 exposed to TCE for 4 days following the sciatic nerve lesion. For mice, regeneration of the  
6 sciatic nerve in comparison to air-exposed animals was 20 and 33% shorter in groups exposed to  
7 150- and 300-ppm TCE for 4 days, respectively. This effect did not significantly increase in  
8 mice pre-exposed to TCE for 20 days, and the regeneration was 30% shorter in the 150-ppm  
9 group and 22% shorter in the 300-ppm group. Comparatively, a 10% reduction in sciatic nerve  
10 regeneration length was observed in rats exposed to TCE for 20 days prior to the lesion plus the  
11 4 days after the sciatic nerve lesion.

12           There are also a few in vitro studies (summarized in Table 4-37) that have demonstrated  
13 that TCE exposure alters the function of inhibitory ion channels such as GABA<sub>A</sub> and glycine  
14 receptors (Beckstead et al., 2000; Krasowski and Harrison, 2000), and serotonin receptors  
15 (Lopreato et al., 2003). Krasowski and Harrison (2000) and Beckstead et al. (2000) were able to  
16 demonstrate that human GABA<sub>A</sub> and glycine receptors could be potentiated by TCE when a  
17 receptor agonist was coapplied. Krasowski and Harrison (2000) conducted an additional  
18 experiment in order to determine if TCE was interacting with the receptor or perturbing the  
19 cellular membrane (bilipid layer). Specific amino acids on the GABA<sub>A</sub> and glycine receptors  
20 were mutated and in the presence of a receptor agonist (GABA for GABA<sub>A</sub> and glycine for  
21 glycine receptors) and in these mutated receptors TCE-mediated potentiation was significantly  
22 decreased or abolished suggesting that there was an interaction between TCE and these  
23 receptors. Lopreato et al. (2003) conducted a similar study with the 5HT<sub>3A</sub> serotonin receptor  
24 and found that when TCE was coapplied with serotonin, there was a potentiation in receptor  
25 response. Additionally, TCE has been demonstrated to alter the function of voltage sensitive  
26 calcium channels (VSCCs) by inhibiting the calcium mediated-current at a holding potential of  
27 -70 mV and shifting the activation of the channels to a more hyperpolarizing potential (Shafer et  
28 al., 2005).

29

#### **4.3.10. Potential Mechanisms for Trichloroethylene (TCE)-Mediated Neurotoxicity**

30           The mechanisms of TCE neurotoxicity have not been established despite a significant  
31 level of research on the outcomes of TCE exposure. Results from several mechanistic studies  
32 can be used to help elucidate the mechanism(s) involved in TCE-mediated neurological effects.

33

1 **Table 4-37. Summary of in vitro ion channel effects with TCE exposure**  
2

Reference	Cellular system	Neuronal channel/receptor	Concentrations	Effects
In vitro studies				
Shafer et al. (2005)	PC12 cells	VSCC	0, 500, 1,000, 1,500, or 2,000 $\mu$ M	Shift of VSCC activation to a more hyperpolarizing potential. Inhibition of VSCCs at a holding potential of $-70$ mV.
Beckstead et al. (2000)	<i>Xenopus</i> oocytes	Human recombinant: glycine receptor $\alpha$ 1, GABA <sub>A</sub> receptors, $\alpha$ 1 $\beta$ 1, $\alpha$ 1 $\beta$ 2 $\gamma$ 2L	0 or 390 $\mu$ M	50% potentiation of the GABA <sub>A</sub> receptors; 100% potentiation of the glycine receptor.
Lopreato et al. (2003)	<i>Xenopus</i> oocytes	Human recombinant serotonin 3A receptor	0 or 390 $\mu$ M	Potentiation of serotonin receptor function.
Krasowski and Harrison (2000)	Human embryonic kidney 293 cells	Human recombinant Glycine receptor $\alpha$ 1, GABA <sub>A</sub> receptors $\alpha$ 2 $\beta$ 1	Not provided	Potentiation of glycine receptor function with an EC <sub>50</sub> of $0.65 \pm 0.05$ mM. Potentiation of GABA <sub>A</sub> receptor function with an EC <sub>50</sub> of $0.85 \pm 0.2$ mM.

3  
4 EC<sub>50</sub> = concentration of the chemical at which 50% of the maximal effect is produced.  
5  
6

7 The disruption of the trigeminal nerve appears to be a highly idiosyncratic outcome of  
8 TCE exposure. There are limited data to suggest that it might entail a demyelination  
9 phenomenon, but similar demyelination does not appear to occur in other nerve tracts. In this  
10 regard, then, TCE is unlike a variety of hydrocarbons that have more global demyelinating  
11 action. There are some data from central nervous system that focus on shifts in lipid profiles as  
12 well as data showing loss of myelinated fibers in the hippocampus. However, the changes in  
13 lipid profiles are both quite small and, also, inconsistent. And the limited data from  
14 hippocampus are not sufficient to conclude that TCE has significant demyelinating effects in this  
15 key brain region. Indeed, the bulk of the evidence from studies of learning and memory function  
16 (which would be tied to hippocampal function) suggests no clear impairments due to TCE.

17 Some researchers (Albee et al., 1997; Albee et al., 2006; Barret et al., 1992; Barret et al.,  
18 1991; Laurenno, 1993) Laurenno, 1988 have indicated that changes in trigeminal nerve function  
19 may be due to dichloroacetylene which is formed under nonbiological conditions of high  
20 alkalinity or temperature during volatilization of TCE. In experimental settings, trigeminal nerve  
21 function (Albee et al., 1997) and trigeminal nerve morphology (Barret et al., 1992; Barret et al.,  
22 1991) was found to be more altered following a low exposure to dichloroacetylene in comparison

1 to the higher TCE exposure. Barret et al. (1992; 1991) also demonstrated that TCE  
2 administration results in morphological changes in the trigeminal nerve. Thus, dichloroacetylene  
3 may contribute to trigeminal nerve impairment may be plausible following an inhalation  
4 exposure under conditions favoring its formation. Examples of such conditions include passing  
5 through a carbon dioxide scrubber containing alkaline materials, application to remove a wax  
6 coating from a concrete-lined stone floor, or mixture with alkaline solutions or caustic (Saunders,  
7 1967) Greim et al., 1984 Bingham et al., 2001. However, dichloroacetylene exposures have not  
8 been identified or measured in human epidemiologic studies with TCE exposure, and thus, do  
9 not appear to be common to occupational or residential settings (Lash and Green, 1993).  
10 Moreover, changes in trigeminal nerve function have also been consistently reported in humans  
11 exposed to TCE following an oral exposure (Kilburn, 2002a); across many human studies of  
12 occupational and drinking water exposures under conditions with highly varying potentials for  
13 dichloroacetylene formation individuals (Barret et al., 1982; Barret et al., 1984; Barret et al.,  
14 1987; Feldman et al., 1988). As a result, the mechanism(s) for trigeminal nerve function  
15 impairment following TCE exposure is unknown(Kilburn, 2002b; Kilburn and Warshaw, 1993;  
16 Mhiri et al., 2004; Ruijten et al., 1991). The varying dichloroacetylene exposure potential across  
17 these studies suggests TCE exposure, which is common to all of them, as the most likely  
18 etiologic agent for the observed effects.

19 The clearest consequences of TCE are permanent impairment of hearing in animal  
20 models and disruption of trigeminal nerve function in humans with animal models showing  
21 comparable changes following administration of a TCE metabolite. With regard to hearing loss,  
22 the effect of TCE has much in common with the effects of several aromatic hydrocarbons  
23 including ethylbenzene, toluene, and *p*-xylene. Many studies have attempted to determine how  
24 these solvents damage the cochlea. Of the hypotheses that have been advanced, there is little  
25 evidence to suggest oxidative stress, changes in membrane fluidity, or impairment of central  
26 efferent nerves whose endings innervate receptor cells in the cochlea. Rather, for reasons that  
27 are still uncertain these solvents seem to preferentially target supporting cells in the cochlea  
28 whose death then alters key structural elements of the cochlea resulting ultimately in hair cell  
29 displacement and death. Recently, potential modes of action resulting in ototoxicity have been  
30 speculated to be due to blockade of neuronal nicotinic receptors present on the auditory cells  
31 (Campo et al., 2007) and potentially changes in calcium transmission (Campo et al., 2008) from  
32 toluene exposure. Although these findings were reported following an acute toluene exposure, it  
33 is speculated that this mechanism may be a viable mechanism for TCE-mediated ototoxicity.

34 A few studies have tried to relate TCE exposure with selective impairments of dopamine  
35 neurons. Two studies (Gash et al., 2008; Guehl et al., 1999) demonstrated dopaminergic

1 neuronal death and/or degeneration following an acute TCE administration. However, the only  
2 human TCE exposure study examining dopamine neuronal activity found no changes in serum  
3 dopamine  $\beta$ -hydroxylase activity in comparison to nonexposed individuals (Nagaya et al., 1990).  
4 It is thought that TaClo, which can be formed from TCE metabolites such as  
5 trichloroacetaldehyde, may be the potent neurotoxicant that selectively targets the dopaminergic  
6 system. More studies are needed to confirm the dopamine neuronal function disruption and if  
7 this disruption is mediated through TaClo.

8         There is good evidence that TCE and certain metabolites such as chloral hydrate have  
9 CNS depressant properties and may account for some of the behavioral effects (such as  
10 vestibular effects, psychomotor activity changes, central visual changes, sleep and mood  
11 changes) that have been observed with TCE. Specifically, in vitro studies have demonstrated  
12 that TCE exposure results in changes in neuronal receptor function for the GABA<sub>A</sub>, glycine, and  
13 serotonin receptors (Beckstead et al., 2000; Krasowski and Harrison, 2000; Lopreato et al.,  
14 2003). All of these inhibitory receptors that are present in the CNS are potentiated when  
15 receptor-specific agonist and TCE are applied. These results are similar to other anesthetics and  
16 suggest that some of the behavioral functions are mediated by modifications in ion channel  
17 function. However, it is quite uncertain whether there are persistent consequences to such high  
18 dose TCE exposure. Additionally, with respect to the GABAergic system, acute administration  
19 of TCE increased the seizure threshold appearance and this effect was the strongest with  
20 convulsants that were GABA receptor antagonists (Shih et al., 2001). Therefore, this result  
21 suggests that TCE interacts with the GABA receptor and that was also verified in vitro  
22 (Beckstead et al., 2000; Krasowski and Harrison, 2000).

23         Also, TCE exposure has been linked to decreased sensitivity to titanic stimulation in the  
24 hippocampus (Ohta et al., 2001) as well as significant reduction in myelin in the hippocampus in  
25 a developmental exposure (Isaacson et al., 1990). These effects are notable since the  
26 hippocampus is highly involved in memory and learning functions. Changes in the hippocampal  
27 physiology may correlate with the cognitive changes that were reported following TCE  
28 exposure.

#### 4.3.11. Overall Summary and Conclusions—Weight of Evidence

30         Both human and animal studies have associated TCE exposure with effects on several  
31 neurological domains. The strongest neurological evidence of hazard in humans is for changes  
32 in trigeminal nerve function or morphology and impairment of vestibular function. Fewer and  
33 more limited evidence exists in humans on delayed motor function, and changes in auditory,

1 visual, and cognitive function or performance. Acute and subchronic animal studies show  
2 morphological changes in the trigeminal nerve, disruption of the peripheral auditory system  
3 leading to permanent function impairments and histopathology, changes in visual evoked  
4 responses to patterns or flash stimulus, and neurochemical and molecular changes. Additional  
5 acute studies reported structural or functional changes in hippocampus, such as decreased  
6 myelination or decreased excitability of hippocampal CA1 neurons, although the relationship of  
7 these effects to overall cognitive function is not established. Some evidence exists for  
8 motor-related changes in rats/mice exposed acutely/subchronically to TCE, but these effects have  
9 not been reported consistently across all studies.

10 Epidemiologic evidence supports a relationship between TCE exposure and trigeminal  
11 nerve function changes, with multiple studies in different populations reporting abnormalities in  
12 trigeminal nerve function in association with TCE exposure (Barret et al., 1982; Barret et al.,  
13 1984; Barret et al., 1987; Feldman et al., 1988; Feldman et al., 1992; Kilburn, 2002b; Kilburn  
14 and Warshaw, 1993; Mhiri et al., 2004) Ruitjen et al., 1991. Of these, two well conducted  
15 occupational cohort studies, each including more than 100 TCE-exposed workers without  
16 apparent confounding from multiple solvent exposures, additionally reported statistically  
17 significant dose-response trends based on ambient TCE concentrations, duration of exposure,  
18 and/or urinary concentrations of the TCE metabolite TCA (Barret et al., 1984; Barret et al.,  
19 1987). Limited additional support is provided by a positive relationship between prevalence of  
20 abnormal trigeminal nerve or sensory function and cumulative exposure to TCE (most subjects)  
21 or CFC-113 (<25% of subjects) (Rasmussen et al., 1993a). Test for linear trend in this study was  
22 not statistically significant and may reflect exposure misclassification since some subjects  
23 included in this study did not have TCE exposure. The lack of association between TCE  
24 exposure and overall nerve function in three small studies (trigeminal: El Ghawabi et al., 1973;  
25 ulnar and medial: Triebig et al., 1983; Triebig et al., 1982) does not provide substantial evidence  
26 against a causal relationship between TCE exposure and trigeminal nerve impairment because of  
27 limitations in statistical power, the possibility of exposure misclassification, and differences in  
28 measurement methods. Laboratory animal studies have also shown TCE-induced changes in the  
29 trigeminal nerve. Although one study reported no significant changes in trigeminal  
30 somatosensory evoked potential in rats exposed to TCE for 13 weeks (Albee et al., 2006), there  
31 is evidence of morphological changes in the trigeminal nerve following short-term exposures in  
32 rats (Barret et al., 1992; Barret et al., 1991).

33 Human chamber, occupational, geographic based/drinking water, and laboratory animal  
34 studies clearly established TCE exposure causes transient impairment of vestibular function.  
35 Subjective symptoms such as headaches, dizziness, and nausea resulting from occupational

1 (Grandjean et al., 1955; Liu et al., 1988; Rasmussen and Sabroe, 1986; Smith, 1970)  
2 environmental (Hirsch et al., 1996), or chamber exposures (Smith, 1970; Stewart et al., 1970)  
3 have been reported extensively. A few laboratory animal studies have investigated vestibular  
4 function, either by promoting nystagmus or by evaluating balance (Niklasson et al., 1993; Tham  
5 et al., 1984; Tham et al., 1979; Umezu et al., 1997).

6 In addition, mood disturbances have been reported in a number of studies, although these  
7 effects also tend to be subjective and difficult to quantify (Gash et al., 2008; Kilburn, 2002a, b;  
8 Kilburn and Warshaw, 1993; McCunney, 1988; Mitchell and Parsons-Smith, 1969; Rasmussen  
9 and Sabroe, 1986; Troster and Ruff, 1990), and a few studies have reported no effects from TCE  
10 on mood (Reif et al., 2003; Triebig et al., 1976; Triebig et al., 1977a). Few comparable mood  
11 studies are available in laboratory animals, although both Moser et al. (2003) and Albee et al.  
12 (2006) report increases in handling reactivity among rats exposed to TCE. Finally, significantly  
13 increased number of sleep hours was reported by Arito et al. (1994) in rats exposed via  
14 inhalation to 50–300-ppm TCE for 8 hours/day for 6 weeks.

15 Four epidemiologic studies of chronic exposure to TCE observed disruption of auditory  
16 function. One large occupational cohort study showed a statistically significant difference in  
17 auditory function with cumulative exposure to TCE or CFC-113 as compared to control groups  
18 after adjustment for possible confounders, as well as a positive relationship between auditory  
19 function and increasing cumulative exposure (Rasmussen et al., 1993c). Of the three studies  
20 based on populations from ATSDR’s TCE Subregistry from the National Exposure Registry,  
21 more limited than Rasmussen et al. (1993c) due to inferior exposure assessment, Burg et al.  
22 (1995) and Burg and Gist (1999) reported a higher prevalence of self-reported hearing  
23 impairments. The third study reported that auditory screening revealed abnormal middle ear  
24 function in children less than 10-years-of-age, although a dose-response relationship could not be  
25 established and other tests did not reveal differences in auditory function (ATSDR, 2003b).  
26 Further evidence for these effects is provided by numerous laboratory animal studies  
27 demonstrating that high dose subacute and subchronic TCE exposure in rats disrupts the auditory  
28 system leading to permanent functional impairments and histopathology.

29 Studies in humans exposed under a variety of conditions, both acutely and chronically,  
30 report impaired visual functions such as color discrimination, visuospatial learning tasks, and  
31 visual depth perception in subjects with TCE exposure. Abnormalities in visual depth perception  
32 were observed with a high acute exposure to TCE under controlled conditions (Vernon and  
33 Ferguson, 1969). Studies of lower TCE exposure concentrations also observed visuofunction  
34 effects. One occupational study (Rasmussen et al., 1993c) reported a statistically significant  
35 positive relationship between cumulative exposure to TCE or CFC-113 and visual gestalts



1 learning and retention among Danish degreasers. Two studies of populations living in a  
2 community with drinking water containing TCE and other solvents furthermore suggested  
3 changes in visual function (Kilburn, 2002b; Reif et al., 2003). These studies used more direct  
4 measures of visual function as compared to Rasmussen et al. (1993c), but their exposure  
5 assessment is more limited because TCE exposure is not assigned to individual subjects  
6 (Kilburn, 2002b), or because there are questions regarding control selection (Kilburn, 2002b) and  
7 exposure to several solvents (Kilburn, 2002b; Reif et al., 2003).

8 Additional evidence of effects of TCE exposure on visual function is provided by a  
9 number of laboratory animal studies demonstrating that acute or subchronic TCE exposure  
10 causes changes in visual evoked responses to patterns or flash stimulus (Blain et al., 1994; Boyes  
11 et al., 2003; Boyes et al., 2005). Animal studies have also reported that the degree of some  
12 effects is correlated with simultaneous brain TCE concentrations (Boyes et al., 2003; Boyes et  
13 al., 2005) and that, after a recovery period, visual effects return to control levels (Blain et al.,  
14 1994; Rebert et al., 1991). Overall, the human and laboratory animal data together suggest that  
15 TCE exposure can cause impairment of visual function, and some animal studies suggest that  
16 some of these effects may be reversible with termination of exposure.

17 Studies of human subjects exposed to TCE either acutely in chamber studies or  
18 chronically in occupational settings have observed deficits in cognition. Five chamber studies  
19 reported statistically significant deficits in cognitive performance measures or outcome measures  
20 suggestive of cognitive effects (Gamberale et al., 1976; Stewart et al., 1970; Triebig et al., 1976;  
21 Triebig et al., 1977b). Danish degreasers with high cumulative exposure to TCE or CFC-113  
22 had a high risk (OR: 13.7, 95% CI: 2.0–92.0) for psychoorganic syndrome characterized by  
23 cognitive impairment, personality changes, and reduced motivation, vigilance, and initiative  
24 compared to workers with low cumulative exposure. Studies of populations living in a  
25 community with contaminated groundwater also reported cognitive impairments (Kilburn,  
26 2002b; Kilburn and Warshaw, 1993), although these studies carry less weight in the analysis  
27 because TCE exposure is not assigned to individual subjects and their methodological design is  
28 weaker.

29 Laboratory studies provide some additional evidence for the potential for TCE to affect  
30 cognition, though the predominant effect reported has been changes in the time needed to  
31 complete a task, rather than impairment of actual learning and memory function (Kishi et al.,  
32 1993; Kulig, 1987; Umezu et al., 1997). In addition, in laboratory animals, it can be difficult to  
33 distinguish cognitive changes from motor-related changes. However, several studies have  
34 reported structural or functional changes in the hippocampus, such as decreased myelination  
35 (Isaacson and Taylor, 1989; Isaacson et al., 1990) or decreased excitability of hippocampal CA1

1 neurons (Ohta et al., 2001), although the relationship of these effects to overall cognitive  
2 function is not established.

3 Two studies of TCE exposure, one chamber study of acute exposure duration and  
4 one occupational study of chronic duration, reported changes in psychomotor responses. The  
5 chamber study of Gamberale et al. (1976) reported a dose-related decrease in performance in a  
6 choice reaction time test in healthy volunteers exposed to 100- and 200-ppm TCE for 70 minutes  
7 as compared to the same subjects without exposure. Rasmussen et al. (1993a) reported a  
8 statistically significant association with cumulative exposure to TCE or CFC-113 and  
9 dyscoordination trend among Danish degreasers. Observations in a third study (Gun et al., 1978)  
10 are difficult to judge given the author's lack of statistical treatment of data. In addition, Gash  
11 et al. (2008) reported that 14 out of 30 TCE-exposed workers exhibited significantly slower fine  
12 motor hand movements as measured through a movement analysis panel test. Studies of  
13 population living in communities with TCE and other solvents detected in groundwater supplies  
14 reported significant delays in simple and choice reaction times in individuals exposed to TCE in  
15 contaminated groundwater as compared to referent groups (Kilburn, 2002b; Kilburn and  
16 Thornton, 1996; Kilburn and Warshaw, 1993). Observations in these studies are more uncertain  
17 given questions of the representativeness of the referent population, lack of exposure assessment  
18 to individual study subjects, and inability to control for possible confounders including alcohol  
19 consumption and motivation. Finally, in a presentation of two case reports, decrements in motor  
20 skills as measured by the grooved pegboard and finger tapping tests were observed (Troster and  
21 Ruff, 1990).

22 Laboratory animal studies of acute or subchronic exposure to TCE observed psychomotor  
23 effects, such as loss of righting reflex (Shih et al., 2001; Umezu et al., 1997) and decrements in  
24 activity, sensory-motor function, and neuromuscular function (Kishi et al., 1993; Moser et al.,  
25 1995; Moser et al., 2003). However, two studies also noted an absence of significant changes in  
26 some measures of psychomotor function (Albee et al., 2006; Kulig, 1987). In addition, less  
27 consistent results have been reported with respect to locomotor activity in rodents. Some studies  
28 have reported increased locomotor activity after an acute i.p. dosage (Wolff and Siegmund,  
29 1978) or decreased activity after acute or short term oral gavage dosing (Moser et al., 1995,  
30 2003). No change in activity was observed following exposure through drinking water (Waseem  
31 et al., 2001), inhalation (Kulig, 1987) or orally during the neurodevelopment period (Fredriksson  
32 et al., 1993).

33 Several neurochemical and molecular changes have been reported in laboratory  
34 investigations of TCE toxicity. Kjellstrand et al. (1987) reported inhibition of sciatic nerve  
35 regeneration in mice and rats exposed continuously to 150-ppm TCE via inhalation for 24 days.

1 Two studies have reported changes in GABAergic and glutamatergic neurons in terms of GABA  
2 or glutamate uptake (Briving et al., 1986) or response to GABAergic antagonistic drugs (Shih et  
3 al., 2001) as a result of TCE exposure, with the Briving et al. (1986) conducted at 50 ppm for  
4 12 months. Although the functional consequences of these changes is unclear, Tham et al.  
5 (1984; 1979) described central vestibular system impairments as a result of TCE exposure that  
6 may be related to altered GABAergic function. In addition, several in vitro studies have  
7 demonstrated that TCE exposure alters the function of inhibitory ion channels such as receptors  
8 for GABA<sub>A</sub> glycine, and serotonin (Beckstead et al., 2000; Krasowski and Harrison, 2000;  
9 Lopreato et al., 2003) or of voltage-sensitive calcium channels (Shafer et al., 2005).  
10

#### 4.4. KIDNEY TOXICITY AND CANCER

##### 4.4.1. Human Studies of Kidney

###### 4.4.1.1.1. Nonspecific Markers of Nephrotoxicity

11 Investigations of nephrotoxicity in human populations show that workers highly exposed  
12 to TCE exhibit evidence of damage to the proximal tubule (NRC, 2006). The magnitude of  
13 exposure needed to produce kidney damage is not clear. Several kidney early biological effect  
14 markers, or biomarkers, are examined in these studies as are less sensitive clinical kidney  
15 outcomes such as glomerular filtration rate and end stage disease. Observation of elevated  
16 excretion of urinary proteins in the four studies of TCE exposure (Bolt et al., 2004; Brüning et  
17 al., 1999a; Brüning et al., 1999b; Green et al., 2004) indicates the occurrence of a toxic insult  
18 among TCE-exposed subjects compared to unexposed controls. Two studies are of subjects with  
19 previously diagnosed kidney cancer (Bolt et al., 2004; Brüning et al., 1999a), with limited  
20 interpretation if effects are associated with exposure or to the disease process. Subjects in  
21 Brüning et al. (1999b) and Green et al. (2004) are disease free. Urinary proteins are considered  
22 nonspecific markers of nephrotoxicity and include  $\alpha$ 1-Microglobulin, albumin, and *N*-acetyl- $\beta$ -  
23 D-glucosaminidase (NAG; Lybarger et al., 1999; Price et al., 1996; Price et al., 1999). Four  
24 studies measure  $\alpha$ 1-microglobulin with elevated excretion observed in the German studies (Bolt  
25 et al., 2004; Brüning et al., 1999a; Brüning et al., 1999b) but not Green et al. (2004). However,  
26 Green et al. (2004) found statistically significant group mean differences in NAG, another  
27 nonspecific marker of tubular toxicity, in disease free subjects. Observations in Green et al.  
28 (2004) provide evidence of tubular damage among workers exposed to trichloroethylene at  
29 32 ppm (mean) (range, 0.5–252 ppm). Elevated excretion of NAG as a nonspecific marker of

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 tubular damage has also been observed with acute TCE poisoning (Carrieri et al., 2007). These  
2 and other studies relevant to evaluating TCE nephrotoxicity are discussed in more detail below.

3 Biological monitoring of persons who previously experienced “high” exposures to  
4 trichloroethylene (100–500 ppm) in the workplace show altered kidney function evidenced by  
5 urinary excretion of proteins suggestive of renal tubule damage. Similar results were observed in  
6 the only study available of subjects with TCE exposure at current occupational limits (NRC,  
7 2006). Table 4-38 provides details and results from these studies. Brüning et al. (1999a) report a  
8 statistically significantly higher prevalence of elevated proteinuria suggestive of severe tubular  
9 damage ( $n = 24$ , 58.5%,  $p < 0.01$ ) and an elevated excretion of  $\alpha 1$ -microglobulin, another urinary  
10 biomarker of renal tubular function, was observed in 41 renal cell carcinoma cases with prior  
11 trichloroethylene exposure and with pending workman’s compensation claims compared with the  
12 nonexposed renal cell cancer patients ( $n = 14$ , 28%) and to hospitalized surgical patients  $n = 2$ ,  
13 2%). Statistical analyses did not adjust for differences in median systolic and diastolic blood  
14 pressure that appeared higher in exposed renal cell carcinoma cases compared to nonexposed  
15 controls. Similarly, severe tubular proteinuria is seen in 14 of 39 workers (35%) exposed to  
16 trichloroethylene in the electrical department, fitters shop and through general degreasing  
17 operations of felts and sieves in a cardboard manufacturing factory compared to no subjects of  
18 46 nonexposed males office and administrative workers from the same factory ( $p < 0.01$ )  
19 (Brüning et al., 1999b). Furthermore, slight tubular proteinuria is seen in 20% of exposed  
20 workers and in 2% of nonexposed workers (Brüning et al., 1999b). Exposed subjects also had  
21 statistically significantly elevated levels of  $\alpha 1$ -microglobulin compared to unexposed controls.  
22 Furthermore, subjects with tubular damage as indicated by urinary protein patterns had higher  
23 GST-alpha concentrations than nonexposed subjects ( $p < 0.001$ ). Both sex and use of spot or  
24 24-hour urine samples are shown to influence  $\alpha 1$ -microglobulin (Andersson et al., 2008);  
25 however, these factors are not considered to greatly influence observations given only males  
26 were subjects and  $\alpha 1$ -microglobulin levels in spot urine sample are adjusted for creatinine  
27 concentration.

28 Bolt et al. (2004) measured  $\alpha 1$ -microglobulin excretion in living subjects from the renal  
29 cell carcinoma case-control study by Brüning et al. (2003). Some subjects in this study were  
30 highly exposed. Of the 134 with renal cell cancer, 19 reported past exposures that led to narcotic  
31 effects and 18 of the 401 controls, experienced similar effects (OR: 3.71, 95% CI: 1.80–7.54)  
32 (Brüning et al., 2003). Bolt et al. (2004) found that  $\alpha 1$ -microglobulin excretion increased in  
33 exposed renal cancer patients compared with nonexposed patients controls. A lower proportion  
34 of exposed cancer patients had normal  $\alpha 1$ -microglobulin excretion, less than 5 mg/L, the  
35 detection level for the assay and the level considered by these investigators as associated with no

1 clinical or subclinical tubule damage, and a higher proportion of high values, defined as  
 2  $\geq 45$  mg/L, compared to cases who did not report TCE occupational exposure and to nonexposed  
 3 controls ( $p < 0.05$ ). Exposed cases, additionally, had statistically significantly higher median  
 4 concentration of  $\alpha 1$ -microglobulin compared to unexposed cases in creatinine-unadjusted spot  
 5 urine specimens ( $p < 0.05$ ). Reduced clearance of creatinine attributable to renal cancer does not

6 **Table 4-38. Summary of human kidney toxicity studies**  
 7

Subjects	Effect	Exposure	Reference
206 subjects— 104 male workers exposed to TCE; 102 male controls (source not identified)	Increased $\beta 2$ -microglobulin and total protein in spot urine specimen. $\beta 2$ -microglobulin: Exposed, $129.0 \pm 113.3$ mg/g creatinine (Cr) Controls, $113.6 \pm 110.6$ mg/g Cr. Total protein: Exposed, $83.4 \pm 113.2$ mg/g creatinine (Cr) Controls, $54.0 \pm 18.6$ mg/g Cr.	TCE exposure was through degreasing activities in metal parts factory or semiconductor industry. U-total trichloro compounds: Exposed, 83.4 mg/g Cr (range, 2–66.2 mg/g Cr). Controls, N.D. 8.4 $\pm$ 7.9 yr mean employment duration.	Nagaya et al. (1989b)
29 metal workers	NAG in morning urine specimen, $0.17 \pm 0.11$ U/mmol Cr.	Breathing zone monitoring, 3 ppm (median) and 5 ppm (mean).	Seldén et al. (1993)
191 subjects— 41 renal cell carcinoma cases pending cases involving compensation with TCE exposure; 50 unexposed renal cell carcinoma cases from same area as TCE-exposed cases; 100 nondiseased control and hospitalized surgical patients	Increased urinary proteins patterns, $\alpha 1$ -microglobulin, and total protein in spot urine specimen. Slight/severe tubular damage: TCE RCC cases, 93%. Nonexposed RCC cases, 46%. Surgical controls, 11%. $p < 0.01$ . $\alpha 1$ -microglobulin (mg/g creatinine): Exposed RCC cases, $24.6 \pm [SD] 13.9$ Unexposed RCC cases, $11.3 \pm [SD] 9.8$ . Surgical controls, $5.5 \pm [SD] 6.8$ .	All exposed RCC cases exposed to ‘high’ and ‘very high’ TCE intensity. 18 yr mean exposure duration.	Brüning et al. (1999a)

<p>85 male workers employed in cardboard manufacturing factory (39 TCE exposed, 46) nonexposed office and administrative controls)</p>	<p>Increased urinary protein patterns and excretion of proteins in spot urine specimen.  Slight/severe tubular damage:  TCE exposed, 67%  Nonexposed, RCC cases, 9%  <math>p &lt; 0.001</math>.  <math>\alpha 1</math>-microglobulin (mg/g creatinine):  Exposed, <math>16.2 \pm [SD] 10.3</math>  Unexposed, <math>7.8 \pm [SD] 6.9</math>  <math>p &lt; 0.001</math>.  GST-alpha (<math>\mu\text{g/g creatinine}</math>):  Exposed <math>6.0 \pm [SD] 3.3</math>  Unexposed, <math>2.0 \pm [SD] 0.57</math>  <math>p &lt; 0.001</math>.  No group differences in total protein or GST-pi.</p>	<p>“High” TCE exposure to workers in the fitters shop and electrical department.  “Very high” TCE exposure to workers through general degreasing operations in carton machinery section.</p>	<p>Brüning et al. (1999b)</p>
--	---	--	-------------------------------

1

**Table 4-38. Summary of human kidney toxicity studies (continued)**

Subjects	Effect	Exposure	Reference
<p>99 renal cell carcinoma cases and 298 hospital controls (from Brüning et al. (2003) and alive at the time of interview)</p>	<p>Increased excretion of <math>\alpha</math>1-microglobulin in spot urine specimen.                      Proportion of subjects with <math>\alpha</math>1-microglobulin &lt;5.0 mg/L:                      Exposed cases, 15%                      Unexposed cases, 51%                      Exposed controls, 55%                      Unexposed controls, 55%  <math>p &lt; 0.05</math>, prevalence of exposed cases compared to prevalences of either exposed controls or unexposed controls.                      Mean <math>\alpha</math>1-microglobulin:                      Exposed cases, 18.1 mg/L                      Unexposed cases, &lt;5.0 mg/L  <math>p &lt; 0.05</math>.</p>	<p>All exposed RCC cases exposed to ‘high’ and ‘very high’ TCE intensity .</p>	<p>Bolt et al. (2004)</p>
<p>124 subjects (70 workers currently exposed to TCE and 54 hospital and administrative staff controls)</p>	<p>Analysis of urinary proteins in spot urine sample obtained 4 d after exposure.                      Increased excretion of albumin, NAG, and formate in spot urine specimen.                      Albumin (mg/g creatinine):<sup>a</sup>                      Exposed, <math>9.71 \pm [SD] 11.6</math>                      Unexposed, <math>5.50 \pm [SD] 4.27</math>  <math>p &lt; 0.05</math>.                      Total NAG (U/g creatinine):                      Exposed, <math>5.27 \pm [SD] 3.78</math>                      Unexposed, <math>2.41 \pm [SD] 1.91</math>  <math>p &lt; 0.01</math>.                      Format (mg/g creatinine):                      Exposed, <math>9.45 \pm [SD] 4.78</math>                      Unexposed, <math>5.55 \pm [SD] 3.00</math>  <math>p &lt; 0.01</math>.                      No group mean differences in GST-alpha, retinol binding protein, <math>\alpha</math>1-microglobulin, <math>\beta</math>2-microglobulin, total protein, and methylmalonic acid.</p>	<p>Mean U-TCA of exposed workers was <math>64 \pm [SD] 102</math> (Range, 1–505).                      Mean U-TCOH of exposed workers was <math>122 \pm [SD] 119</math> (Range, 1–639).                      Mean TCE concentration to exposed subjects was estimated as 32 ppm (range, 0.5–252 ppm) and was estimated by applying the German occupational exposure limit (maximale arbeitsplatz konzentration, MAK) standard to U-TCA and assuming that the linear relationship holds for exposures above 100 ppm.                      86% of subjects with exposure to &lt;50 ppm TCE.</p>	<p>Green et al. (2004)</p>

**Table 4-38. Summary of human kidney toxicity studies (continued)**

Subjects	Effect	Exposure	Reference
101 cases or deaths from ESDR among male and female subjects in Hill Air Force Base aircraft maintenance worker cohort of Blair et al. (1998)	TCE exposure: Cox Proportional Hazard Analysis: Ever exposed to TCE, <sup>b</sup> 1.86 (1.02, 3.39). Logistic regression: <sup>b</sup> No chemical exposure (referent group): 1.0 <5 unit-yr, 1.73 (0.86, 3.48) 5–25 unit-yr, 1.65 (0.82, 3.35) >25 unit-yr, 1.65 (0.82, 3.35) Monotonic trend test, <i>p</i> > 0.05. Indirect low-intermittent TCE exposure, 2.47 (1.17, 5.19) Indirect peak/infrequent TCE exposure 3.55 (1.25, 10.74) Direct TCE exposure, “not statistically significant” but hazard ratio and confidence intervals were not presented in paper.	Cumulative TCE exposure (intensity × duration) identified using three categories, <5 unit-yr, 5–25 unit-yr, >25 unit-yr per job exposure matrix of Stewart et al. (1991).	Radican et al. (2006)
269 cases of IgA nephropathy or membranous nephropathy glomerulonephritis followed 5 yr (mean) for progression to ESRD	TCE exposure: Cox Proportional Hazard Analysis: Ever exposed to TCE, <sup>b</sup> 2.5 (0.9, 6.5) High exposure level to TCE, <sup>b</sup> 2.7 (0.7, 10.1).	Exposure to TCE assigned using job title and job-exposure matrix; two dose surrogates, ever exposed and high exposure level.	Jacob et al. (2007)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15

<sup>a</sup> For a urine sample, 10–17 mg of albumin per g of creatinine is considered to be suspected albuminuria in males (15–25 in females) (de Jong and Brenner, 2004).

<sup>b</sup> Hazard ratio and 95% CI.

ESDR = end-stage renal disease, N.D. = not detectable, SD = standard deviation.

explain the lower percentage of normal values among exposed cases given findings of similar prevalence of normal excretion among unexposed renal cell cases and controls.

In their study of 70 current employees (58 males, 12 females) of an electronic factory with trichloroethylene exposure and 54 (50 males, 4 females) age-matched subjects drawn from hospital or administrative staff, Green et al. (2004) found that urinary excretion of albumin, total NAG and formate were increased in the exposed group compared with the unexposed group.<sup>4</sup> No differences between exposed and unexposed subjects were observed in other urinary proteins,

<sup>4</sup> Elevation of NAG in urine is a sign of proteinuria, and proteinuria is both a sign and a cause of kidney malfunction (Zandi-Nejad et al., 2004). For a urine sample, 10–17 mg of albumin per g of creatinine is considered to be suspected albuminuria in males (15–25 in females) (de Jong and Brenner, 2004).



1 including  $\alpha$ 1-microglobulin,  $\beta$ 2-microglobulin, and GST-alpha. Green et al. (2004) stated that  
2 NAG is not an indicator of nephropathy, or damage, but rather is an indicator of functional  
3 change in the kidney. Green et al. (2004) further concluded that increased urinary albumin or  
4 NAG was not related to trichloroethylene exposure; analyses to examine the exposure-response  
5 relationship found neither NAG or albumin concentration correlated to U-TCA or employment  
6 duration (years). The National Research Council (NRC, 2006) did not consider U-TCA as  
7 sufficiently reliable to use as a quantitative measure of TCE exposure, concluding that the data  
8 reported by Green et al. (2004) were inadequate to establish exposure-response information  
9 because the relationship between U-TCA and ambient TCE intensity is highly variable and  
10 nonlinear, and conclusions about the absence of association between TCE and nephrotoxicity can  
11 not be made based on U-TCA. Moreover, use of employment duration does not consider  
12 exposure intensity differences between subjects with the same employment duration, and bias  
13 introduced through misclassification of exposure may explain the Green et al. (2004) findings.

14 Seldén et al. (1993) in their study of 29 metal workers (no controls) reported a correlation  
15 between NAG and U-TCA ( $r = 0.48$ ,  $p < 0.01$ ) but not with other exposure metrics of recent or  
16 long-term exposure. Personal monitoring of worker breath indicated median and mean  
17 time-weighted-average TCE exposures of 3 and 5 ppm, respectively. Individual NAG  
18 concentrations were within normal reference values. Rasmussen et al. (1993b), also, reported a  
19 positive relationship ( $p = 0.05$ ) between increasing urinary NAG concentration (adjusted for  
20 creatinine clearance) and increasing duration in their study of 95 metal degreasers (no controls)  
21 exposed to either TCE (70 subjects) or CFC113(25 subjects). Multivariate regression analyses  
22 which adjusted for age were suggestive of an association between NAG and exposure duration  
23 ( $p = 0.011$ ). Mean urinary NAG concentration was higher among subjects with annual exposure  
24 of >30 hours/week, defined as peak exposure, compared to subjects with annual exposure of less  
25 than <30 hours/week ( $72.4 \pm 44.1$   $\mu\text{g/g}$  creatinine compared to  $45.9 \pm 30.0$   $\mu\text{g/g}$  creatinine,  
26  $p < 0.01$ ).

27 Nagaya et al. (1989b) did not observe statistically significant group differences in urinary  
28  $\beta$ 2-microglobulin and total protein in spot urine specimens of male degreasers and their controls,  
29 nor were these proteins correlated with urinary total trichloro-compounds (U-TTC). The paper  
30 lacks details on subject selection, whether urine collection was at start of work week or after  
31 sufficient exposure, and presentation of  $p$ -values and correlation coefficients. The presentation  
32 of urinary protein concentrations stratified by broad age groups is less statistically powerful than  
33 examination of this confounder using logistic regression. Furthermore, although valid for  
34 pharmacokinetic studies, examination of renal function using U-TTC as a surrogate for TCE  
35 exposure is uncertain, as discussed above for Green et al. (2004).

#### 4.4.1.1.2. End-Stage Renal Disease

1 End-stage renal disease is associated with hydrocarbon or organic solvent exposures in  
2 two studies examining this endpoint (Jacob et al., 2007; Radican et al., 2006). Table 4-38  
3 provides details and results from Radican et al. (2006) and Jacob et al. (2007). Radican et al.  
4 (2006) assessed end-stage renal disease in a cohort of aircraft maintenance workers at Hill Air  
5 Force Base (Blair et al., 1998) with strong exposure assessment to trichloroethylene (NRC, 2006)  
6 and reported a twofold risk with overall TCE exposure and end stage renal disease (1.86, 95%  
7 CI: 1.02, 3.39). A second study, the GN-PROGRESS retrospective cohort study, observed a  
8 twofold elevated risk for progression of glomerulonephritis to ESRD from TCE (overall  
9 exposure: 2.5, 95% CI: 0.9–6.5; high level TCE exposure: 2.7, 95% CI: 0.7, 10.1) (Jacob et al.,  
10 2007). Statistical power was more limited in Jacob et al. (2007) because of its smaller number of  
11 exposed cases, 21 with overall exposure, compared to 56 exposed cases in Radican et al. (2006).  
12 Other occupational studies do not examine end-stage renal disease specifically, instead reporting  
13 relative risks associated with deaths due to nephritis and nephrosis (ATSDR, 2004; Boice et al.,  
14 1999; Boice et al., 2006b), all genitourinary system deaths (Costa et al., 1989; Garabrant et al.,  
15 1988; Ritz, 1999a), or providing no information on renal disease mortality in the published paper  
16 (Blair et al., 1998; Chang et al., 2003; Morgan et al., 1998).  
17

#### 4.4.2. Human Studies of Kidney Cancer

18 Cancer of the kidney and renal pelvis is the 6<sup>th</sup> leading cause of cancer in the  
19 United States with an estimated 54,390 (33,130 men and 21,260 women) newly diagnosed cases  
20 and 13,010 deaths (Jemal et al., 2008; NCI, 2008). Age-adjusted incidence rates based on cases  
21 diagnosed in 2001–2005 from 17 Surveillance, Epidemiology, and End Results (SEER)  
22 geographic areas are 18.3 per 100,000 for men and 9.2 per 100,000 for women. Age-adjusted  
23 mortality rates are much lower; 6.0 per 100,000 for men and 2.7 for women.

24 Cohort, case-control, and geographical studies have examined trichloroethylene and  
25 kidney cancer, defined either as cancer of kidney and renal pelvis in cohort and geographic based  
26 studies or as renal cell carcinoma, the most common type of kidney cancer, in case-control  
27 studies. Appendix C identifies these studies' design and exposure assessment characteristics.  
28 Observations in these studies are presented below in Table 4-39. Rate ratios for incidence  
29 studies in Table 4-39 are, generally, larger than for mortality studies.

30 Additionally, a large body of evidence exists on kidney cancer risk and either job or  
31 industry titles where trichloroethylene usage has been documented. TCE has been used as a

1 degreasing solvent in a number of jobs, task, and industries, some of which include metal,  
2 electronic, paper and printing, leather manufacturing and aerospace/aircraft manufacturing or  
3

1  
2

**Table 4-39. Summary of human studies on TCE exposure and kidney cancer**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort and PMR studies—incidence</b>				
Aerospace workers (Rocketdyne)				Zhao et al. (2005)
	Any exposure to TCE	Not reported		
	Low cum TCE score	1.00 <sup>a</sup>	6	
	Med cum TCE score	1.87 (0.56, 6.20)	6	
	High TCE score	4.90 (1.23, 19.6)	4	
	<i>p</i> for trend	<i>p</i> = 0.023		
TCE, 20 yr exposure lag <sup>b</sup>				
	Low cum TCE score	1.00 <sup>a</sup>	6	
	Med cum TCE score	1.19 (0.22, 6.40)	7	
	High TCE score	7.40 (0.47, 116)	3	
	<i>p</i> for trend	<i>p</i> = 0.120		
All employees at electronics factory (Taiwan)				Chang et al. (2005)
	Males	1.06 (0.45, 2.08) <sup>c</sup>	8	
	Females	1.09 (0.56, 1.91) <sup>c</sup>	12	
	Females	1.10 (0.62, 1.82) <sup>c</sup>	15	Sung et al. (2008)
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure, all subjects	1.2 (0.98, 1.46)	103	
	Any exposure, males	1.2 (0.97, 1.48)	93	
	Any exposure, females	1.2 (0.55, 2.11)	10	
Exposure lag time				
	20 yr	1.3 (0.86, 1.88)	28	
Employment duration				
	<1 yr	0.8 (0.5, 1.4)	16	
	1–4.9 yr	1.2 (0.8, 1.7)	28	
	≥5 yr	1.6 (1.1, 2.3)	32	
Subcohort w/higher exposure				
	Any TCE exposure	1.4 (1.0, 1.8)	53	
Employment duration				
	1–4.9 yr	1.1 (0.7, 1.7) <sup>d</sup>	23	
	≥5 yr	1.7 (1.1, 2.4) <sup>d</sup>	30	

3

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-39. Summary of human studies on TCE exposure and kidney cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Biologically monitored Danish workers		1.1 (0.3, 2.8)	4	Hansen et al. (2001)
	Any TCE exposure, males	0.9 (0.2, 2.6)	3	
	Any TCE exposure, females	2.4 (0.03, 14)	1	
	Cumulative exp (Ikeda)	Not reported		
	<17 ppm-yr			
	≥17 ppm-yr			
	Mean concentration (Ikeda)	Not reported		
	<4 ppm			
	4+ ppm			
	Employment duration	Not reported		
	<6.25 yr			
	≥6.25			
Aircraft maintenance workers from Hill Air Force Base				Blair et al. (1998)
	TCE subcohort	Not reported		
	Males, cumulative exp			
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	1.4 (0.4, 4.7)	9	
	5–25 ppm-yr	1.3 (0.3, 4.7)	5	
	>25 ppm-yr	0.4 (0.1, 2.3)	2	
	Females, cumulative exp			
	0	1.0 <sup>a</sup>		
	<5 ppm-yr		0	
	5–25 ppm-yr		0	
	>25 ppm-yr	3.6 (0.5, 25.6)	2	
Biologically-monitored Finnish workers				Anttila et al. (1995)
	All subjects	0.87 (0.32, 1.89)	6	
	Mean air-TCE (Ikeda extrapolation)			
	<6 ppm	Not reported		
	6+ ppm	Not reported		
Cardboard manufacturing workers in Arnsberg, Germany				Henschler et al. (1995)
	Exposed workers	7.97 (2.59, 8.59) <sup>c</sup>	5	
Biologically-monitored Swedish workers				Axelson et al. (1994)
	Any TCE exposure, males	1.16 (0.42, 2.52)	6	
	Any TCE exposure, females	Not reported		

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-39. Summary of human studies on TCE exposure and kidney cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Cardboard manufacturing workers, Atlanta area, GA				Sinks et al. (1992)
	All subjects	3.7 (1.4, 8.1)	6	
	All departments	$\infty$ (3.0, $\infty$ ) <sup>f</sup>	5	
	Finishing department	16.6 (1.7, 453.1) <sup>f</sup>	3	
<b>Cohort and PMR studies—mortality</b>				
Computer manufacturing workers (IBM), NY				Clapp and Hoffman (2008)
	Males	1.64 (0.45, 4.21) <sup>g</sup>	4	
	Females		0	
Aerospace workers (Rocketdyne)				Boice et al. (2006b) Zhao et al. (2005)
	Any TCE (utility/eng flush)	2.22 (0.89, 4.57)	7	
	Any exposure to TCE	Not reported		
	Low cum TCE score	1.00 <sup>a</sup>	7	
	Med cum TCE score	1.43 (0.49, 4.16)	7	
	High TCE score	2.13 (0.50, 8.32)	3	
	<i>p</i> for trend	<i>p</i> = 0.31		
	TCE, 20 yr exposure lag <sup>b</sup>			
	Low cum TCE score	1.00 <sup>a</sup>	10	
	Med cum TCE score	1.69 (0.29, 9.70)	6	
	High TCE score	1.82 (0.09, 38.6)	1	
	<i>p</i> for trend	<i>p</i> = 0.635		
View-Master employees				
	Males	2.76 (0.34, 9.96) <sup>g</sup>	2	
	Females	6.21 (2.68, 12.23) <sup>g</sup>	8	
United States Uranium-processing workers (Fernald)				Ritz (1999a) (as reported in NRC, 2006)
	Any TCE exposure	Not reported		
	Light TCE exposure, 2-10 yr duration <sup>d</sup>	1.94 (0.59, 6.44)	5	
	Light TCE exposure, >10 yr duration <sup>d</sup>	0.76 (0.14, 400.0)	2	
	Mod TCE exposure, >2 yr duration <sup>d</sup>		0	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-39. Summary of human studies on TCE exposure and kidney cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference	
Aerospace workers (Lockheed)				Boice et al. (1999)	
Routine Exp	0.99 (0.40, 2.04)	7			
Routine-Intermittent <sup>a</sup>	Not presented	11			
Duration of exposure					
0 yr	1.0	22			
<1 yr	0.97 (0.37, 2.50)	6			
1–4 yr	0.19 (0.02, 1.42)	1			
≥5 yr	0.69 (0.22, 2.12)	4			
<i>p</i> for trend					
Aerospace workers (Hughes)				Morgan et al. (1998)	
TCE subcohort	1.32 (0.57, 2.60)	8			
Low intensity (<50 ppm) <sup>c</sup>	0.47 (0.01, 2.62)	1			
High intensity (>50 ppm) <sup>c</sup>	1.78 (0.72, 3.66)	7			
TCE subcohort (Cox analysis)					
Never exposed	1.00 <sup>a</sup>	24			
Ever exposed	1.14 (0.51, 2.58) <sup>h</sup>	8			
Peak					
No/Low	1.00 <sup>a</sup>	24			
Med/Hi	1.89 (0.85, 4.23) <sup>h</sup>	8			
Cumulative					
Referent	1.00 <sup>a</sup>	24			
Low	0.31 (0.04, 2.36) <sup>h</sup>	1			
High	1.59 (0.68, 3.71) <sup>h</sup>	7			
Aircraft maintenance workers (Hill AFB, Utah)				Blair et al. (1998)	
TCE subcohort	1.6 (0.5, 5.1) <sup>a</sup>	15			
Males, cumulative exp					
0	1.0 <sup>a</sup>				
<5 ppm-yr	2.0 (0.5, 7.6)	8			
5–25 ppm-yr	0.4 (0.1, 4.0)	1			
>25 ppm-yr	1.2 (0.3, 4.8)	4			
Females, cumulative exp					
0	1.0 <sup>a</sup>				
<5 ppm-yr		0			
5–25 ppm-yr	9.8 (0.6, 157)	1			
>25 ppm-yr	3.5 (0.2, 56.4)	1			
TCE subcohort					Radican et al. (2008)
Males, cumulative exp	1.18 (0.47, 2.94) <sup>i</sup>	18			
0	1.24 (0.41, 3.71) <sup>i</sup>	16			
	1.0 <sup>i</sup>				

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-39. Summary of human studies on TCE exposure and kidney cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Aircraft maintenance workers (Hill AFB, Utah) (continued)				Blair et al. (1998)
	<5 ppm-yr	1.87 (0.59, 5.97) <sup>i</sup>	10	
	5–25 ppm-yr	0.31 (0.03, 2.75) <sup>i</sup>	1	
	>25 ppm-yr	1.16 (0.31, 4.32) <sup>i</sup>	5	
	Females, cumulative exp	0.93 (0.15, 5.76) <sup>i</sup>	2	
	0	1.0 <sup>a</sup>		
	<5 ppm-yr		0	
	5–25 ppm-yr	2.86 (0.27, 29.85) <sup>i</sup>	1	
	>25 ppm-yr	0.97 (0.10, 9.50) <sup>i</sup>	1	
Cardboard manufacturing workers in Arnsberg, Germany				Henschler et al. (1995)
	TCE exposed workers	3.28 (0.40, 11.84)	2	
	Unexposed workers	- (0.00, 5.00)	0	
Deaths reported to among GE pension fund (Pittsfield, MA)			12	Greenland et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA				Sinks et al. (1992)
		1.4 (0.0, 7.7)	1	
U.S. Coast Guard employees				Blair et al. (1989)
	Marine inspectors	1.06 (0.22, 3.10)	3	
	Noninspectors	1.03 (0.21, 3.01)	3	
Aircraft manufacturing plant employees (Italy)				Costa et al. (1989)
	All subjects	Not reported		
Aircraft manufacturing plant employees (San Diego, CA)				Garabrant et al. (1988)
	All subjects	0.93 (0.48, 1.64)	12	
<b>Case-control studies</b>				
Population of four countries in central and eastern Europe				Moore et al. (2010)
	Any TCE exposure	1.63 (1.04, 2.54)	48	
	Any TCE exposure (High confidence exposure)	2.05 (1.13, 3.73)	29	
	Cumulative TCE exposure			
	<1.58 ppm-yr	1.19 (0.61, 2.35)	17	
	≥1.58 ppm-yr	2.02 (1.14, 3.59) <sup>j</sup>	31	
	<i>p</i> for trend	<i>p</i> = 0.02		
	Average intensity			
	<0.076 ppm	1.38 (0.81, 2.35)	31	
	≥0.076 ppm	2.34 (1.05, 5.21)	17	
	<i>p</i> for trend	<i>p</i> = 0.02		

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-39. Summary of human studies on TCE exposure and kidney cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Population of Arve Valley, France				Charbotel et al. (2006; 2009; 2007)
Any TCE exposure	1.64 (0.95, 2.84)	37		
Any TCE exposure (High confidence exposure)	1.88 (0.89, 3.98)	16		
Cumulative TCE exposure				
Referent/nonexposed	1.00 <sup>a</sup>	49		
Low, 62.4 ppm-yr <sup>k</sup>	1.62 (0.75, 3.47)	12		
Medium, 253.2 ppm-yr <sup>k</sup>	1.15 (0.47, 2.77)	9		
High, 925 ppm-yr <sup>k</sup>	2.16 (1.02, 4.60) <sup>l</sup>	16		
Test for trend	$p = 0.04$			
Cumulative TCE exposure + peak				
Referent/nonexposed	1.00 <sup>a</sup>	49		
Low/medium, no peaks	1.35 (0.69, 2.63)	18		
Low/medium + peaks	1.61 (0.36, 7.30)	3		
High, no peaks	1.76 (0.65, 4.73)	8		
High + peaks	2.73 (1.06, 7.07) <sup>l</sup>	8		
Cumulative TCE exposure, 10-yr lag				
Referent/nonexposed	1.00 <sup>a</sup>	49		
Low/medium, no peaks	1.44 (0.69, 2.80)	19		
Low/medium + peaks	1.38 (0.32, 6.02)	3		
High, no peaks	1.50 (0.53, 4.21)	7		
High + peaks	3.15 (1.19, 8.38)	8		
Time-weighted-average TCE exposure <sup>m</sup>				
Referent/nonexposed	1.00 <sup>a</sup>	46		
Any TCE without cutting fluid	1.62 (0.76, 3.44)	15		
Any cutting fluid without TCE	2.39 (0.52, 11.03)	3		
<50 ppm TCE + cutting fluid	1.14 (0.49, 2.66)	12		
50 + ppm TCE + cutting fluid	2.70 (1.02, 7.17)	10		
Population of Arnsberg Region, Germany				Brüning et al. (2003)
Longest job held—TCE/PERC (CAREX)	1.80 (1.01, 3.20)	117		
Self-assessed exposure to TCE	2.47 (1.36, 4.49)	25		
Duration of self-assessed TCE exposure				
0	1.00 <sup>a</sup>	109		
<10 yr	3.78 (1.54, 9.28)	11		
10–20 yr	1.80 (0.67, 4.79)	7		
>20 yr	2.69 (0.84, 8.66)	8		

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-39. Summary of human studies on TCE exposure and kidney cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Population in five German Regions				Pesch et al., (2000a)
Any TCE Exposure	Not reported			
Males	Not reported			
Females	Not reported			
TCE exposure (Job Task Exposure Matrix)				
Males				
Medium	1.3 (1.0, 1.8)	68		
High	1.1 (0.8, 1.5)	59		
Substantial	1.3 (0.8, 2.1)	22		
Females				
Medium	1.3 (0.7, 2.6)	11		
High	0.8 (0.4, 1.9)	7		
Substantial	1.8 (0.6, 5.0)	5		
Population of Minnesota				Dosemeci et al. (1999)
Ever exposed to TCE, NCI JEM				
Males	1.04 (0.6, 1.7)	33		
Females	1.96 (1.0, 4.0)	22		
Males + Females	1.30 (0.9, 1.9)	55		
Population of Arnsberg Region, Germany				Vamvakas et al. (1998)
Self-assessed exposure to TCE	10.80 (3.36, 34.75)	19		
Population of Montreal, Canada				Siemiatycki et al. (1991)
Any TCE exposure	0.8 (0.4, 2.0) <sup>n</sup>	4		
Substantial TCE exposure	0.8 (0.2, 2.6) <sup>n</sup>	2		
<b>Geographic based studies</b>				
Residents in two study areas in Endicott, NY		1.90 (1.06, 3.13)	15	ATSDR (2006a) (2008)
Residents of 13 census tracts in Redlands, CA		0.80 (0.54, 1.12) <sup>o</sup>	54	Morgan and Cassady (2002)
Finnish residents				Vartiainen et al. (1993)
Residents of Hausjarvi		Not reported		
Residents of Huttula		Not reported		

1  
2 <sup>a</sup> Internal referents, workers not exposed to TCE.  
3 <sup>b</sup> Relative risks for TCE exposure after adjustment for 1<sup>st</sup> employment, socioeconomic status, age at event, and all  
4 other carcinogens, including hydrazine.  
5 <sup>c</sup> Chang et al. (2005)—urinary organs combined.  
6 <sup>d</sup> SIR for renal cell carcinoma.  
7 <sup>e</sup> Henschler et al. (1995) Expected number of incident cases calculated using incidence rates from the Danish Cancer  
8 Registry.  
9 <sup>f</sup> Odds ratio from nested case-control analysis.  
10 <sup>g</sup> Proportional mortality ratio.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2 **Table 4-39. Summary of human studies on TCE exposure and kidney cancer**  
3 **(continued)**  
4

5 <sup>h</sup> Risk ratio from Cox Proportional Hazard Analysis, stratified by age, sex and decade (EHS, 1997).

6 <sup>l</sup> In Radican et al. (2008), kidney cancer defined as renal cell carcinoma (ICDA 8 code 189.0) and estimated relative  
7 risks from Cox proportional hazard models were adjusted for age and sex.

8 <sup>j</sup> The odds ratio, adjusted for age, sex and center, for subjects with high-confidence exposure assessment with  
9 cumulative exposure,  $\geq 1.58$  ppm-yr, was 2.23 (95% CI: 1.07, 4.64) and *p*-value for trend = 0.02.

10 <sup>k</sup> Mean cumulative exposure score in Charbotel et al. (2006) (personal communication from Barbara Charbotel,  
11 University of Lyon, to Cheryl Scott, U.S. EPA, 11 April 2008).

12 <sup>l</sup> In Charbotel et al. (2006) analyses adjusted for age, sex, smoking and body mass index. The odds ratio, adjusted  
13 for age, sex, smoking, body mass index and exposure to cutting fluids and other petroleum oils, for high  
14 cumulative TCE exposure was 1.96 (95% CI: 0.71, 5.37) and for high cumulative + peak TCE exposure was 2.63  
15 (95% CI: 0.79, 8.83). The odds ratio for, considering only job periods with high confidence TCE exposure  
16 assessment, adjusted for age, sex, smoking and body mass index, for high cumulative dose plus peaks was 3.80  
17 (95% CI: 1.27, 11.40).

18 <sup>m</sup> The exposure surrogate is calculated for one occupational period only and is not the average exposure  
19 concentration over the entire employment period.

20 <sup>n</sup> 90% CI.

21 <sup>o</sup> 99% CI.

22  
23 GE = General Electric, IBM = International Business Machines Corporation, JEM = job-exposure matrix, NCI =  
24 National Cancer Institute, PERC = perchloroethylene, PMR = proportionate mortality ratio.  
25  
26

27 maintenance industries and job title of degreaser, metal workers, electrical worker, and machinist  
28 (IARC, 1995b; Purdue et al., 2011). NRC (2006) identifies characteristics for kidney cancer  
29 case-control studies that assess job title or occupation in their Table 3-8. Relative risks and  
30 95% CIs reported in these studies are found in Table 4-40 below.  
31

**4.4.2.1.1. Studies of Job Titles and Occupations with Historical Trichloroethylene (TCE)  
Usage**

32 Elevated risks are observed in many of the cohort or case-control studies between kidney  
33 cancer and industries or job titles with historical use of trichloroethylene (Brüning et al., 2003;  
34 Charbotel et al., 2006; Mandel et al., 1995; Mattioli et al., 2002; McCredie and Stewart, 1993;  
35 Parent et al., 2000a; Partanen et al., 1991; Pesch et al., 2000a; Schlehofer et al., 1995; Wilson et  
36 al., 2008; Zhang et al., 2004). Overall, these studies, although indicating association with metal  
37 work exposures and kidney cancer, are insensitive for identifying a TCE hazard. The use of job  
38 title or industry as a surrogate for exposure to a chemical is subject to substantial  
39 misclassification that will attenuate rate ratios due to exposure variation and differences among  
40 individuals with the same job title. Several small case-control studies (Auperin et al., 1994;  
41 Harrington et al., 1989; Jensen et al., 1988; Parent et al., 2000a; Sharpe et al., 1989; Vamvakas et

1 al., 1998) have insufficient statistical power to detect modest associations due to their small size  
 2 and potential exposure misclassification (NRC, 2006). For these reasons, statistical

3 **Table 4-40. Summary of case-control studies on kidney cancer and**  
 4 **occupation or job title**  
 5

Case ascertainment area/exposure group		Relative risk (95% CI)	No. exposed cases	Reference
Swedish Cancer Registry Cases				Wilson et al. (2008)
Machine/electronics industry		1.30 (1.08, 1.55) <sup>a</sup> [M]	120	
		1.75 (1.04, 2.76) <sup>a</sup> [F]	18	
Shop and construction metal work		1.19 (1.00, 1.40) <sup>a</sup> [M]	143	
Machine assembly		1.62 (0.94, 2.59) <sup>a</sup> [M]		
Metal plating work		2.70 (0.73, 6.92) <sup>a</sup> [M]	4	
Shop and construction metal work		1.66 (0.71, 3.26) <sup>a</sup> [F]	8	
Arve Valley, France				Charbotel et al. (2006)
Metal industry		1.02 (0.59, 1.76)	28	
Metal workers, job title		1.00 (0.56, 1.77)	25	
Metal industry, screw-cutting workshops		1.39 (0.75, 2.58)	22	
Machinery, electrical and transportation equipment manufacture		1.19 (0.61, 2.33)	15	
Iowa Cancer Registry Cases				Zhang et al. (2004)
Assemblers		2.5 (0.8, 7.6)	5	
>10 yr employment		4.2 (1.2, 15.3)	4	
Arnsberg Region, Germany				Brüning et al. (2003)
Iron/steel		1.15 (0.29, 4.54)	3	
Occupations with contact to metals		1.53 (0.97, 2.43)	46	
Longest job held		1.14 (0.66, 1.96)	24	
Metal greasing/degreasing		5.57 (2.33, 13.32)	15	
Degreasing agents				
Low exposure		2.11 (0.86, 5.18)	9	
High exposure		1.01 (0.40, 2.54)	7	
Bologna, Italy				Mattioli et al. (2002)
Metal workers		2.21 (0.99, 5.37)	37	
Printers		1.55 (0.17, 13.46)	7	
Solvents		0.79 (0.31, 1.98) [M]	17	
		1.47 (0.12, 17.46) [F]	3	

6  
7

**Table 4-40. Summary of case-control studies on kidney cancer and occupation or job title (continued)**

Case ascertainment area/exposure group		Relative risk (95% CI)	No. exposed cases	Reference
Montreal, Canada				Parent et al. (2000a)
Metal fabricating and machining industry	1.0 (0.6, 1.8)	14		
Metal processors	1.2 (0.4, 3.4)	4		
Printing and publishing industry	1.1 (0.4, 3.0)	4		
Printers	3.0 (1.2, 7.5)	6		
Aircraft mechanics	2.8 (1.0, 8.4)	4		
5 Regions in Germany				Pesch et al. (2000a)
Electrical and electronic equipment assembler	3.2 (1.0, 10.3) [M]	5		
	2.7 (1.3, 5.8) [F]	11		
Printers	3.5 (1.1, 11.2)[M]	5		
	2.1 (0.4, 11.7) [F]	2		
Metal cleaning/degreasing, job task	1.3 (0.7, 2.3) [M]	15		
	1.5 (0.3, 7.7) [F]	2		
New Zealand Cancer Registry				Delahunt et al. (1995)
Toolmakers and blacksmiths	1.48 (0.72, 3.03)	No info		
Printers	0.67 (0.25, 1.83)			
Minnesota Cancer Surveillance System				Mandel et al. (1995)
Iron or steel	1.6 (1.2, 2.2)	8		
Rhein-Neckar-Odenwald Area, Germany				Schlehofer et al. (1995)
Metal				
Industry	1.63 (1.07, 2.48)	71		
Occupation	1.38 (0.89, 2.12)			
Electronic				
Industry	0.51 (0.26, 1.01)	14		
Occupation	0.57 (0.25, 1.33)	9		
Chlorinated solvents	2.52 (1.23, 5.16)	27		
Metal and metal compounds	1.47 (0.94, 2.30)	62		
Danish Cancer Registry				Mellempgaard et al. (1994)
Iron and steel	1.4 (0.8, 2.4) [M]	31		
	1.0 (0.1, 3.2) [F]	1		
Solvents	1.5 (0.9, 2.4) [M]	50		
	6.4 (1.8, 23) [F]	16		

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-40. Summary of case-control studies on kidney cancer and occupation or job title (continued)**

Case ascertainment area/exposure group		Relative risk (95% CI)	No. exposed cases	Reference
France				Aupérin et al. (1994)
	Machine fitters, assemblers, and precision instrument makers	0.7 (0.3, 1.9)	16	
New South Wales, Australia				McCredie and Stewart (1993)
	Iron and steel	1.18 (0.75, 1.85) <sup>b</sup>	52	
		2.39 (1.26, 4.52) <sup>c</sup>	19	
	Printing or graphics	1.18 (0.87, 2.08) <sup>b</sup>	29	
		0.82 (0.32, 2.11) <sup>d</sup>	6	
	Machinist or tool maker	1.15 (0.72, 1.86) <sup>b</sup>	48	
		1.83 (0.92, 3.61) <sup>c</sup>	16	
	Solvents	1.54 (1.11, 2.14) <sup>b</sup>	109	
		1.40 (0.82, 2.40) <sup>c</sup>	24	
Finnish Cancer Registry				Partanen et al. (1991)
	Iron and metalware work	1.87 (0.94, 3.76)	22	
		2.33 (0.83, 6.51)	10	
	Paper and pulp; printing/publishing	2.20 (1.02, 4.72) [M]	18	
		5.95 (1.21, 29.2) [F]	7	
	Nonchlorinated solvents	3.46 (0.91, 13.2) [M]	9	
West Midlands UK Cancer Registry				Harrington et al. (1989)
	Organic solvents	Ever exposed	1.30 (0.31, 8.50)	
		Intermediate exposure	1.54 (0.69, 4.10)	
			3	
Montreal, Canada				Sharpe et al. (1989)
	Organic solvents	Degreasing solvents	1.68 (0.83, 2.22)	
			33	
Oklahoma				Asal et al. (1988a; 1988b)
	Metal degreasing	Machining	1.7 (0.7, 3.8) [M]	
			19	
		Painter, paint manufacture	1.7 (0.7, 4.3) [M]	
Missouri Cancer Registry				Brownson (1988)
	Machinists	1.3 (0.7, 2.6) [M]	22	
Danish Cancer Registry				Jensen et al. (1988)
	Machinists	2.2 (0.5, 10.3)	3	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-40. Summary of case-control studies on kidney cancer and occupation or job title (continued)**

Case ascertainment area/exposure group		Relative risk (95% CI)	No. exposed cases	Reference
	Iron and metal, blacksmith	1.4 (0.7, 2.9) <sup>d</sup>	17	
	Painter, paint manufacture	1.8 (0.7, 4.6)	10	

<sup>a</sup> Renal pelvis, Wilson et al. (2008).  
<sup>b</sup> Renal cell carcinoma, McCredie and Stewart (1993).  
<sup>c</sup> Renal pelvis, McCredie and Stewart (1993).  
<sup>d</sup> Renal pelvis and ureter, Jensen et al. (1988).

UK = United Kingdom.

variation in the risk estimate is large and observation of statistically significantly elevated risks associated with metal work in many of these studies is noteworthy. Some studies also examined broad chemical grouping such as degreasing solvents or chlorinated solvents. Observations in studies that assessed degreasing agents or chlorinated solvents reported statistically significant elevated kidney cancer risk (Asal et al., 1988a; Asal et al., 1988b; Brüning et al., 2003; Harrington et al., 1989; McCredie and Stewart, 1993; Mellemgaard et al., 1994; Pesch et al., 2000a; Schlehofer et al., 1995). Observations of association with degreasing agents together with job title or occupations where TCE has been used historically provide a signal and suggest an etiologic agent common to degreasing activities.

**4.4.2.1.2. Cohort and Case-Controls Studies of Trichloroethylene (TCE) Exposure**

Cohort and case-controls studies that include job-exposure matrices for assigning TCE exposure potential to individual study subjects show associations with kidney cancer, specifically renal cell carcinoma, and trichloroethylene exposure. Support for this conclusion derives from findings of increased risks in cohort studies (Henschler et al., 1995; Raaschou-Nielsen et al., 2003; Zhao et al., 2005) and in case-control studies from the Arnsberg region of Germany (Brüning et al., 2003; Pesch et al., 2000a; Vamvakas et al., 1998), the Arve Valley region in France (Charbotel et al., 2006; Charbotel et al., 2009), the United States (Dosemeci et al., 1999; Sinks et al., 1992) and the four central and eastern Europe countries of Czech Republic, Poland, Romania, and Russia (Moore et al., 2010).

A consideration of a study’s statistical power and exposure assessment approach is necessary to interpret observations in Table 4-39. Most cohort studies are underpowered to

1 detect a doubling of kidney cancer risks including the essentially null studies by Greenland et al.  
2 (1994), Axelson et al. (1994 [incidence]), Anttila et al. (1995 [incidence]), Blair et al. (1998  
3 [incidence and mortality]), Morgan et al. (1998), Boice et al. (1999) and Hansen et al. (2001).  
4 Only the exposure duration-response analysis of Raaschou-Nielsen et al. (2003) had over  
5 80% statistical power to detect a doubling of kidney cancer risk (NRC, 2006), and they observed  
6 a statistically significant association between kidney cancer and  $\geq 5$ -year employment duration.  
7 Rate ratios estimated in the mortality cohort studies of kidney cancer (e.g., (Axelson et al., 1994;  
8 Blair et al., 1998; Boice et al., 2006b; Garabrant et al., 1988; Greenland et al., 1994; Morgan et  
9 al., 1998; Ritz, 1999a, b; Sinks et al., 1992) are likely underestimated to some extent because  
10 their reliance on death certificates and increased potential of nondifferential misclassification of  
11 outcome in these studies, although the magnitude is difficult to predict (NRC, 2006). Cohort or  
12 proportionate mortality ratio (PMR) studies with more uncertain exposure assessment  
13 approaches, e.g., studies of all subjects working at a factory (ATSDR, 2004; Chang et al., 2003;  
14 Chang et al., 2005; Clapp and Hoffman, 2008; Costa et al., 1989; Garabrant et al., 1988; Sung et  
15 al., 2007), do not show association but are quite limited given their lack of attribution of higher  
16 or lower exposure potentials; risks are likely diluted due to their inclusion of no or low exposed  
17 subjects.

18 Two studies were carried out in geographic areas with a high frequency and a high degree  
19 of TCE exposure and were designed with a priori hypotheses to test for the effects of TCE  
20 exposure on renal cell cancer risk (Brüning et al., 2003; Charbotel et al., 2006; Charbotel et al.,  
21 2009) and a third study carried out in four central and eastern European countries with high renal  
22 cell carcinoma rates unexplained by established risk factors Ferlay et al., 2008 (Moore et al.,  
23 2010). For these reasons, their observations have important bearing to the epidemiologic  
24 evidence evaluation. These studies found a twofold elevated risk with any TCE exposure after  
25 adjustment for several possible confounding factors including smoking (2.47, 95% CI: 1.36,  
26 4.49) for self-assessed exposure to TCE (Brüning et al., 2003); any confidence job with high  
27 cumulative TCE exposure, 925 ppm-years (2.16, 95% CI: 1.02, 4.60) with a positive and  
28 statistically significant trend test,  $p = 0.04$ , high confidence jobs with high cumulative TCE  
29 exposure (3.34, 95% CI: 1.27, 8.74) (Charbotel et al., 2006); high confidence assessment of high  
30 TCE cumulative exposure  $\geq 1.58$  ppm-years (2.23, 95% CI: 1.07, 4.64) with a positive and  
31 statistically significant trend test,  $p = 0.02$  (Moore et al., 2010). Furthermore, renal cell  
32 carcinoma risk in Charbotel et al. (2005) increased to over threefold (95% CI: 1.19, 8.38) in  
33 statistical analyses which considered a 10-year exposure lag period. An exposure lag period is  
34 often adopted in analysis of cancer epidemiology to reduce exposure measurement biases  
35 (Salvan et al., 1995). Most exposed cases in this study were exposed to TCE below any current



1 occupational standard (26 of 37 cases [70%]) had held a job with a highest time-weighted  
2 average (TWA [ $<50$  ppm]) (Charbotel et al., 2009). A subsequent analysis of Charbotel et al.  
3 (2009) using an exposure surrogate defined as the highest TWA for any job held, an inferior  
4 surrogate given TCE exposures in other jobs were not considered, reported an almost threefold  
5 elevated risk (2.80, 95% CI: 1.12, 7.03) adjusted for age, sex, body mass index (BMI), and  
6 smoking with exposure to TCE in any job to  $\geq 50$ -ppm TWA (Charbotel et al., 2009).  
7 Considering all jobs, Moore et al. (2010) reported a risk of 2.34 (95% CI: 1.05, 5.21) for average  
8 TCE intensity ( $>0.76$  ppm), an exposure metric similar to a TWA exposure category. Zhao et al.  
9 (2005) compared 2,689 TCE-exposed workers at a California aerospace company to nonexposed  
10 workers from the same company as the internal referent population, and found a monotonic  
11 increase in incidence of kidney cancer by increasing cumulative TCE exposure. In addition, a  
12 fivefold increased incidence was associated with high cumulative TCE exposure. This  
13 relationship for high cumulative TCE exposure, lagged 20 years, was accentuated with  
14 adjustment for other occupational exposures (RR = 7.40, 95% CI: 0.47, 116), although the  
15 confidence intervals were increased. An increased confidence interval with adjustments is not  
16 unusual in occupational studies, as exposure is usually highly correlated with them, so that  
17 adjustments often inflate standard error without removing any bias (NRC, 2006). Observed risks  
18 were lower for kidney cancer mortality and because of reliance on cause of death on death  
19 certificates are likely underestimated because of nondifferential misclassification of outcome  
20 (Percy et al., 1981). Boice et al. (2006b), another study of 1,111 workers with potential TCE  
21 exposure at this company and which overlaps with Zhao et al. (2005), found a twofold increase  
22 in kidney cancer mortality (standardized mortality ratio [SMR] = 2.22, 95% CI: 0.89, 4.57). This  
23 study examined mortality in a cohort whose definition date differs slightly from Zhao et al.  
24 (2005), working between 1948–1999 with vital status as of 1999 (Boice et al., 2006b) compared  
25 to working between 1950–1993 with follow-up for mortality as of 2001 (Zhao et al., 2005), and  
26 used a qualitative approach for TCE exposure assessment. Boice et al. (2006b) is a study of  
27 fewer subjects identified with potential TCE exposure, of fewer kidney cancer deaths [7 deaths;  
28 10 incident cases, 10 deaths in Zhao et al. (2005)], of subjects with more recent exposures, and  
29 with a inferior exposure assessment approach compared to Zhao et al. (2005); a finding of a  
30 twofold mortality increase (95% CI: 0.89, 4.57) is noteworthy given the insensitivities.

31 Zhao et al. (2005), Charbotel et al. (2006) and Moore et al. (2010), furthermore, are three  
32 of the few studies to conduct a detailed assessment of exposure that allowed for the development  
33 of a job-exposure matrix that provided rank-ordered levels of exposure to TCE and other  
34 chemicals. NRC (2006) discussed the inclusion of rank-ordered exposure levels is a strength  
35 increasing precision and accuracy of exposure information compared to more inferior exposure

1 assessment approaches in some other studies such as duration of exposure or a grouping of all  
2 exposed subjects.

3 The finding in Raaschou-Nielsen et al. (2003) of an elevated renal cell carcinoma risk  
4 with longer employment duration is noteworthy given this study's use of a relatively insensitive  
5 exposure assessment approach. One strength of this study is the presentation of incidence ratios  
6 for a subcohort of higher exposed subjects, those with at least 1-year duration of employment  
7 and first employment before 1980, as a sensitivity analysis for assessing the effect of possible  
8 exposure misclassification bias. Renal cell carcinoma risk was higher in this subcohort  
9 compared to the larger cohort and indicated some potential for misclassification bias in the  
10 grouped analysis. For both the cohort and subcohort analyses, risk appeared to increase with  
11 increasing employment duration, although formal statistical tests for trend are not presented in  
12 the published paper.

#### 13 14 **4.4.2.1.3. Discussion of controversies on studies in the Arnsberg region of Germany**

15 Two previous studies of workers in this region, a case-control study of Vamvakas et al.  
16 (1998) and Henschler et al. (1995), a study prompted by a kidney cancer case cluster, observed  
17 strong associations between kidney cancer and TCE exposure. A fuller discussion of the studies  
18 from the Arnsberg region and their contribution to the overall weight of evidence on cancer  
19 hazard is warranted in this evaluation given the considerable controversy (Bloemen and  
20 Tomenson, 1995; Cherrie et al., 2001; Green and Lash, 1999; Mandel, 2001; McLaughlin and  
21 Blot, 1997; Swaen, 1995) surrounding Henschler et al. (1995) and Vamvakas et al. (1998).

22 Criticisms of Henschler et al. (1995) and Vamvakas et al. (1998) relate, in part, to  
23 possible selection biases that would lead to inflating observed associations and limited inferences  
24 of risk to the target population. Specifically, these include (1) the inclusion of kidney cancer  
25 cases first identified from a cluster and the omission of subjects lost to follow-up from Henschler  
26 et al. (1995); (2) use of a Danish population as referent, which may introduce bias due to  
27 differences in coding cause of death and background cancer rate differences (Henschler et al.,  
28 1995); (3) follow-up of some subjects outside the stated follow-up period (Henschler et al.,  
29 1995); (4) differences between hospitals in the identification of cases and controls in Vamvakas  
30 et al. (1998); (5) lack of temporality between case and control interviews (Vamvakas et al.,  
31 1998); (6) lack of blinded interviews (Vamvakas et al., 1998); (7) age differences in Vamvakas  
32 et al. (1998) cases and controls that may lead to a different TCE exposure potential; (8) inherent  
33 deficiencies in Vamvakas et al. (1998) as reflected by its inability to identify other known kidney  
cancer risk factors; and, (9) exposure uncertainty, particularly unclear intensity of TCE exposure.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Overall, NRC (2006) noted that some of the points above may have contributed to an  
2 underestimation of the true exposure distribution of the target population (points 5, 6, and 7),  
3 other points would underestimate risk (points 3), and that these effects could not have explained  
4 the entire excess risk observed in these studies (points 1, 2, and 4). The NRC (2006) furthermore  
5 disagreed with the exposure uncertainty criticism (point 9), and concluded TCE exposures,  
6 although of unknown intensity, were substantial and, clearly showed graded differences on  
7 several scales in Vamvakas et al. (1998) consistent with this study's semiquantitative exposure  
8 assessment.

9 Brüning et al. (2003) was carried out in a broader region in southern Germany, which  
10 included the Arnsberg region and a different set of cases and control identified from a later time  
11 period than Vamvakas et al. (1998). The TCE exposure range in this study was similar to that in  
12 Vamvakas et al. (1998), although at a lower exposure prevalence because of the larger and more  
13 heterogeneous ascertainment area for cases and controls. For "ever exposed" to TCE,  
14 Brüning et al. (2003) observed a risk ratio of 2.47 (95% CI: 1.36, 4.49) and a fourfold increase in  
15 risk (95% CI: 1.80, 7.54) among subjects with any occurrence of narcotic symptom and a sixfold  
16 increase in risk (95% CI: 1.46, 23.99) for subjects who had daily occurrences of narcotic  
17 symptoms; risks which are lower than observed in Vamvakas et al. (1998). The lower rate ratio  
18 in Brüning et al. (2003) might indicate bias in the Vamvakas et al., study or statistical variation  
19 between studies related to the broader base population included in Brüning et al. (2003).

20 Observational studies such as epidemiologic studies are subject to biases and  
21 confounding which can be minimized but never completely eliminated through a study's design  
22 and statistical analysis methods. While Brüning et al. (2003) overcomes many of the  
23 deficiencies of Henschler et al. (1995) and Vamvakas et al. (1998), nonetheless, possible biases  
24 and measurement errors could be introduced through their use of prevalent cases and residual  
25 noncases, use of controls from surgical and geriatric clinics, nonblinding of interviewers, a  
26 2-year difference between cases and controls in median age, use of proxy or next-of-kin  
27 interviews, and self-reported occupational history.

28 The impact of any one of the above points could either inflate or depress observed  
29 associations. Biases related to a longer period for case compared to control ascertainment could  
30 go in either direction. Next-of-kin interviewers for deceased cases, all controls being alive at the  
31 time of interview, would be expected to underestimate risk if exposures were not fully reported  
32 and thus, misclassified. On the other hand, the control subjects who were enrolled when the  
33 interviews were conducted might not represent the true exposure distribution of the target  
34 population through time and would lead to overestimate of risk. Selection of controls from  
35 clinics is not expected to greatly influence observed associations since these clinics specialized

1 in the type of care they provided (NRC, 2006). Brüning et al. (2003) is not the only kidney  
2 case-control study where interviewers were not blinded; in fact, only the study of Charbotel et al.  
3 (2006) included blinding of interviewers. Blinding of interviewers is preferred to reduce  
4 possible bias. Brüning et al.'s use of frequency matching using 5-year age groupings is common  
5 in epidemiologic studies and any biases introduced by age difference between cases and controls  
6 is expected to be minimal because the median age difference was 3 years.

7 Despite these issues, the three studies of the Arnsberg region, with very high apparent  
8 exposure and different base populations showed a significant elevation of risk and all have  
9 bearing on kidney cancer hazard evaluations. The emphasis provided by each study for  
10 identifying a kidney cancer hazard depends on its strengths and weaknesses. Brüning et al.  
11 (2003) overcomes many of the deficiencies in Henschler et al. (1995) and Vamvakas et al.  
12 (1998). The finding of a statistically significantly approximately threefold elevated odds ratio  
13 with occupational TCE exposure in Brüning et al. (2003) strengthens the signal previously  
14 reported by Henschler et al. (1995) and Vamvakas et al. (1998). A previous study of cardboard  
15 workers in the United States (Sinks et al., 1992), a study like Henschler et al. (1995) which was  
16 prompted by a reported cancer cluster, had observed association with kidney cancer incidence,  
17 particularly with work in the finishing department where TCE use was documented. Henschler  
18 et al. (1995), Vamvakas et al. (1998) and Sinks et al. (1992) are less likely to provide a precise  
19 estimate of the magnitude of the association given greater uncertainty in these studies compared  
20 to Brüning et al. (2003). For this reason, Brüning et al. (2003) is preferred for meta-analysis  
21 treatment since it is considered to better reflect risk in the target population than the two other  
22 studies. Another study (Charbotel et al., 2006) of similar exposure conditions of a different base  
23 population and of different case and control ascertainment methods as the Arnsberg region  
24 studies has become available since the Arnsberg studies. This study shows a statistically  
25 significant elevation of risk and high cumulative TCE exposure in addition to a positive trend  
26 with rank-order exposure levels. Charbotel et al. (2006) adds evidence to observations from  
27 earlier studies on high TCE exposures in Southern Germany and suggests that peak exposure  
28 may add to risk associated with cumulative TCE exposure.

#### 4.4.2.1.4. Examination of Possible Confounding Factors

30 Examination of potential confounding factors is an important consideration in the  
31 evaluation of observations in the epidemiologic studies on TCE and kidney cancer. A known  
32 risk factor for kidney cancer is cigarette smoking. Obesity, diabetes, hypertension and  
33 antihypertensive medications, and analgesics are linked to kidney cancer, but causality has not

1 been established (McLaughlin et al., 2006; Moore et al., 2005). On the other hand, fruit and  
2 vegetable consumption is considered protective of kidney cancer risk (McLaughlin et al., 2006).  
3 Studies by Asal et al. (1988a; 1988b), Partanen et al. (1991), McCredie and Stewart (1993),  
4 Aupérin et al. (1994), Chow et al. (1994), Mellemegaard et al. (1994), Mandel et al. (1995),  
5 Vamvakas et al. (1998), Dosemeci et al. (1999), Pesch et al. (2000a), Brüning et al. (2003), and  
6 Charbotel et al. (2006) controlled for smoking and all studies except Pesch et al. (2000a)  
7 controlled for BMI. Moore et al. (2010) examined but did not find smoking or BMI as potential  
8 confounders because statistical examination of cigarette smoking and BMI altered risk  
9 estimates for the association between TCE exposure and kidney cancer by less than 10%.  
10 Vamvakas et al. (1998) and Dosemeci et al. (1999) controlled for hypertension and or diuretic  
11 intake in the statistical analysis. Because it is unlikely that exposure to trichloroethylene is  
12 associated with smoking, body mass index, hypertension, or diuretic intake, these possible  
13 confounders do not significantly affect the estimates of risk (NRC, 2006).

14 Direct examination of possible confounders is less common in cohort studies than in  
15 case-control studies where information is obtained from study subjects or their proxies. Use of  
16 internal controls, such as for Zhao et al. (2005), in general minimizes effects of potential  
17 confounding due to smoking or socioeconomic status since exposed and referent subjects are  
18 drawn from the same target population. Information on possible confounding due to BMI  
19 (obesity) and to diabetes is lacking in cohort studies; however, any uncertainties are likely small  
20 given the generally healthy nature of an employed population and its favorable access to medical  
21 care.

22 The effect of smoking as a possible confounder may be assessed indirectly through  
23 (1) examination of risk ratios for other smoking-related sites, (2) examination of the expected  
24 contribution by smoking to cancer risks and (3) examination of lung cancer in nine TCE cohort  
25 studies in which there is a high likelihood of TCE exposure in individual study subjects (and  
26 which met, to a sufficient degree, the standards of epidemiologic design and analysis in a  
27 systematic review using meta-analysis methods. Some information on smoking-related lung and  
28 kidney cancer risks may be obtained from IARC (2004b) for indirectly evaluating the expected  
29 magnitude by smoking on kidney cancer risks in TCE cohort studies. Five cohort studies of  
30 cigarette smoking reported risk estimates for both lung and kidney cancers with an observed ratio  
31 of lung:kidney cancer risks of 3.5–10.6 for active smokers, who will have higher  
32 smoking-related risks than former smokers (see Table 4-41). The nNine -cohort studies (Anttila  
33 et al., 1995; Axelson et al., 1994; Boice et al., 1999; Greenland et al., 1994; Hansen et al., 2001;  
34 Morgan et al., 1998; Raaschou-Nielsen et al., 2003; Radican et al., 2008; Zhao et al., 2005)  
35 present lung cancer risks and reported risks for overall TCE exposure range from 0.69 (95% CI:

1 0.31, 1.30) by Axelson et al. (1994) to 1.4 (95% CI: 1.32, 1.52) by Raaschou-Nielsen et al.  
2 (2003) (see Table 4-70). Smoking was more prevalent in the Raaschou-Nielsen et al. (2003)  
3 cohort than the background population as suggested by the elevated risks for lung and other  
4 smoking-related sites.

1 **Table 4-41. Summary of lung and kidney cancer risks in active smokers**  
 2 **(from IARC, 2004b)**  
 3

Cohort	Relative risk		Ratio lung; kidney	Reference
	Lung	Kidney		
MRFIT (USA) 1975–1985, men	6.7	1.9	3.5	Kuller et al. (1991)
British Doctor's Study (UK) 1957–1991, men	14.9 <sup>a</sup>	1.4 <sup>a</sup>	10.6	Doll et al. (1994)
U.S. Veterans Study (USA) 1954–1980, men	11.6	1.5	7.7	McLaughlin et al. (1995)
Swedish Census Study (Sweden) 1963–1989, women	4.7	1.1	4.3	Nordlund et al. (1997, 1999)
Cancer Prevention Study II (USA) 1982–1986, women	12.4	1.4	9.1	Garfinkel and Stellman (1988); Heath et al. (1997)

4  
 5 <sup>a</sup> Relative mortality rate compared to nonsmokers.

6  
 7 UK = United Kingdom.  
 8  
 9

10 If smoking fully contributes to the observed 40 % excess lung cancer risk in this study  
 11 and based on observations in the five smoking cohorts, the expected contribution by smoking to  
 12 renal cell carcinoma risk is estimated as 1–6 % and far smaller than the 20 and 40% excess in  
 13 renal cell carcinoma risk in the cohort and subcohort. The use of internal referents who are  
 14 unexposed subjects drawn from the occupational settings as TCE exposed subjects in  
 15 three studies will reduce any confounding related to smoking as referents (Morgan et al., 1998;  
 16 Radican et al., 2008; Zhao et al., 2005). In the other cohort studies lacking direct adjustment for  
 17 smoking and internal referents, difference in cigarette smoking between exposed and referent  
 18 subjects is not sufficient to fully explain observed excess kidney cancer risks associated with  
 19 TCE, particularly, high TCE exposure. Lung cancer risk estimates are lower than or equal to  
 20 kidney cancer risk estimates and inconsistent with observations in the five smoking cohorts  
 21 (Axelson et al., 1994; Boice et al., 1999; Hansen et al., 2001).

22 Meta-analysis methods were adopted, additionally, as a tool for examining risk estimates  
 23 from the nine cohort studies in which there is a high likelihood of TCE exposure in individual  
 24 study subjects (e.g., based on job-exposure matrices or biomarker monitoring) and which met, to  
 25 a sufficient degree, the standards of epidemiologic design and analysis in a systematic review  
 26 reporting lung cancer to assess the presence of potential systematic error related to confounding  
 27 from smoking. Significant heterogeneity was observed across the nine studies of overall

1 exposure ( $I^2 = 90\%$ ) and for six of the nine studies with highest exposure groups ( $I^2 = 80\%$ ).  
2 Although the appropriateness of conducting any meta-analysis without attempting to explain the  
3 heterogeneity is arguable, the summary estimate from the primary random effects meta-analysis  
4 of the nine studies was 0.96 (95% CI: 0.76, 1.21) for overall TCE exposure, and 0.96 (95% CI:  
5 0.72, 1.27) for the highest group exposure reported by six studies. These observations suggest  
6 potential confounding by smoking of kidney cancer summary risk estimates can be reasonably  
7 excluded in cohort studies of TCE exposure.

8 Mineral oils such as cutting fluids or hydrazine common to some job titles with potential  
9 TCE exposures (such as machinists, metal workers, and test stand mechanics) were included as  
10 covariates in statistical analyses of Zhao et al. (2005), Boice et al. (2006b) and Charbotel et al.  
11 (2006; 2009) or evaluated as a single exposure for cases and controls in Moore et al. (Karami et  
12 al., 2011; 2010). A TCE effect on kidney cancer incidence was still evident although effect  
13 estimates were often imprecise due to lowered statistical power (Charbotel et al., 2006;  
14 Charbotel et al., 2009; Zhao et al., 2005). Observed associations were similar in analyses  
15 including chemical coexposures in both Zhao et al. (2005) and Charbotel et al. (2006; 2009)  
16 compared to chemical coexposure unadjusted risks. The association or OR between high TCE  
17 score and kidney cancer incidence in Zhao et al. (2005) was 7.71 (95% CI: 0.65, 91.4) after  
18 adjustment for other carcinogens including hydrazine and cutting oils, compared to analyses  
19 unadjusted for chemical coexposures (4.90, 95% CI: 1.23, 19.6).

20 In Charbotel et al. (2006), exposure to TCE was strongly associated with exposure to  
21 cutting fluids and petroleum oils (22 of the 37 TCE-exposed cases were exposed to both).  
22 Statistical modeling of all factors significant at 10% threshold showed the OR for cutting fluids  
23 to be almost equal to one, whereas the OR for the highest level of TCE exposure was close to  
24 two (Charbotel et al., 2006). Moreover, when exposure to cutting oils was divided into  
25 three levels, a decrease in OR with level of exposure was found. In conditional logistic  
26 regression adjusted for cutting oil exposure, the relative risk (OR) for renal cell carcinoma and  
27 TCE was similar to relative risks unadjusted for cutting fluid exposures (high cumulative TCE  
28 exposure: 1.96 [95% CI: 0.71–5.37] compared to 2.16 [95% CI: 1.02–4.60]; high cumulative and  
29 peak: 2.63 [95% CI: 0.79–8.83] compared to 2.73 [95% CI: 1.06–7.07] (Charbotel et al., 2006).  
30 Charbotel et al. (2009) further examined TCE exposure defined as the highest TWA in any job  
31 held, inferior to cumulative exposure given its lack of consideration of TCE exposure potential in  
32 other jobs, either as exposure to TCE alone, cutting fluids alone, or to both after adjusting for  
33 smoking, body mass index, age, sex, and exposure to other oils (TCE alone: 1.62 [95% CI: 0.75,  
34 3.44]); cutting fluids alone: 2.39 (95% CI: 0.52, 11.03); TCE >50-ppm TWA + cutting fluids:  
35 2.70 (95% CI: 1.02, 7.17). There were few cases exposed to cutting fluids alone ( $n = 3$ ) or to



1 TCE alone ( $n = 15$ ), all of whom had TCE exposure (in the highest exposed job held) of  
2 <35-ppm TWA, and the subgroup analyses were of limited statistical power. A finding of higher  
3 risk for both cutting oil and TCE exposure  $\geq 50$  ppm compared to cutting oil alone supports a  
4 TCE effect for kidney cancer. Adjustment for cutting oil exposures, furthermore, did not greatly  
5 affect the magnitude of TCE effect measures in the many analyses presented by Charbotel et al.  
6 (2006; 2009) suggesting cutting fluid exposure as not greatly confounding TCE effect measures.  
7 Two other kidney case-control studies of TCE exposure examined the effect of cutting oil as a  
8 single occupational exposure on kidney cancer risk (Brüning et al., 2003; Karami et al., 2011).  
9 Although Brüning et al. (2003) reported an odds ratio of 2.11 (95% CI: 0.66, 6.70) for  
10 self-reported cutting oil exposure and kidney cancer, cutting oil exposure did not appear highly  
11 correlated with TCE exposure as only 5 cases reported exposure to cutting oils compared to 25  
12 cases reporting TCE exposure. Karami et al. (2011), who examined mineral oil or cutting fluid  
13 exposure among cases and controls in Moore et al. (2010), reported an odds ratio of 0.8 (95% CI:  
14 0.6, 1.1) and 1.1 (95% CI: 0.8, 1.4), for cutting oil mists or other mineral oil mists respectively,  
15 and provides evidence that the reported association with TCE exposure in Moore et al. (2010) is  
16 not likely confounded by cutting or mineral oil exposures. Moreover, cutting oils and mineral  
17 oils have not been associated with kidney cancer in other cohort or case-control studies (Mirer,  
18 2010; NIOSH, 1998), and provides additional support for potential confounding by cutting oils  
19 as of minimal concern.

20 Boice et al. (2006b) was unable to directly examine hydrazine exposure on TCE effect  
21 measures because of a lack of model convergence in statistical analyses. Three of  
22 7 TCE-exposed kidney cancer cases were identified with hydrazine exposure of 1.5 years or less  
23 and the absence of exposure to the other four cases suggested confounding related to hydrazine  
24 was unlikely to greatly modify observed association between TCE and kidney cancer.

25

#### 4.4.2.1.5. Susceptible Populations—Kidney Cancer and Trichloroethylene (TCE) Exposure

26 Two studies of kidney cancer cases from the Arnsberg region in Germany and the study  
27 of kidney cancer cases from three Central and Eastern European countries have examined the  
28 influence of polymorphisms of the glutathione-S-transferase metabolic pathway on renal cell  
29 carcinoma risk and TCE exposure (Brüning et al., 1997a; Moore et al., 2010; Wiesenhütter et al.,  
30 2007). In their study of 45 TCE-exposed male and female renal cell carcinoma cases pending  
31 legal compensation and 48 unmatched male TCE-exposed controls, Brüning et al. (1997a)  
32 observed a higher prevalence of exposed cases homozygous and heterozygous for GSTM1  
33 positive, 60%, than the prevalence for this genotype among exposed controls, 35%. The

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 frequency of GSTM1 positive was lower among this control series than the frequency found in  
2 other European population studies, 50% (Brüning et al., 1997a). The prevalence of the GSTT1  
3 positive genotype was 93% among exposed cases and 77% among exposed controls. The  
4 prevalence of GSTT1 positive genotype in the European population is 75% (Brüning et al.,  
5 1997a).

6 Wiesenhütter et al. (2007) compares the frequency of genetic polymorphism among  
7 subjects from the renal cancer case-control study of Brüning et al. (2003) and to the frequencies  
8 of genetic polymorphisms in the areas of Dortmund and Lutherstadt Wittenberg, Germany.  
9 Wiesenhütter et al. (2007) identified the genetic frequencies of GSTM1 and GSTT1 phenotypes  
10 for 98 of the original 134 cases (73%) and 324 of the 401 controls (81%). The prevalence of  
11 GSTM1 positive genotype was 48% among all renal cell carcinoma cases, 40% among  
12 TCE-exposed cases, and 52% among all controls. The prevalence of GSTT1 positive genotypes  
13 was 81% among all cases and 81% among all controls. The prevalence of GSTT1 positive  
14 genotypes reported in this paper for all TCE-exposed cases was 20%. Wiesenhütter et al. (2007)  
15 noted background frequencies in the German population in the expanded control group were  
16 50% for GSTM1 positive and 81% for GSTT1 positive genotypes. The observations are limited  
17 as the paper is sparsely reported and numbers of exposed ( $n = 4$ ) and unexposed ( $n = 15$ ) GSTT1  
18 positive cases does not sum to the 79 cases with the GSTT1 positive genotype identified in the  
19 table's first row.

20 Moore et al. (2010) presents associations between TCE exposure and renal cell  
21 carcinoma risk stratified by GSTT genotype and for single nucleotide polymorphisms (SNPs) of  
22 the renal cysteine conjugate  $\beta$ -lyase gene. Genotyping was available for 925 of the 1,097 cases  
23 and 1,192 of the 1,476 controls. The percentage of cases and controls genotyped did not  
24 significantly differ among TCE-exposed and unexposed subjects nor was the active GSTT1  
25 genotype association with kidney cancer risk (0.94, 95% CI: 0.75, 1.19). However, adopting  
26 statistical analysis examining TCE exposure and kidney cancer that stratified on GSTT1  
27 polymorphism as null (deleted allele) or active ( $\geq 1$  intact allele), Moore et al. (2010) reported  
28 significant associations for GSTT1 active genotype and no association suggested for subjects  
29 with GSTT1 null genotype. The risk estimate for the association for TCE exposure and kidney  
30 cancer among subjects with an active GSTT1 genotype, was 1.88 (95% CI: 1.06, 3.33), with  
31 higher risk estimates for long exposure duration, cumulative exposure and average exposure  
32 intensity [ $\geq 13.5$  years, 2.13 (95% CI: 1.04, 4.39);  $\geq 1.58$  ppm-years, 2.59 (95% CI: 1.25, 5.35);  
33  $\geq 0.076$  ppm, 2.77 (95% CI: 1.01, 7.58)] and a positive trend with increasing exposure duration,  
34 cumulative exposure or average intensity categories ( $p \leq 0.03$ ) (Moore et al., 2010). The  
35 associations between TCE exposure and kidney cancer was stronger for subjects with a

1 functionally active GSTT1 than those for all subjects (both genotypes combined) (see  
2 Table 4-39). Moore et al. (2010) tested but did not find statistical interaction between GSTT1  
3 genotype and TCE exposure ( $p \geq 0.17$ ). Moore et al. (2010) also examined the effect of  
4 polymorphisms of the cysteine conjugate  $\beta$ -lyase gene on TCE risk and reported interaction  
5 between TCE exposure and four minor alleles (SNPs rs2293968, rs2280841, rs2259043, and  
6 rs941960) ( $p < 0.05$ ). Associations with TCE exposure and kidney cancer were threefold higher  
7 compared to unexposed subjects with these SNPs.

8 Observations in Brüning et al. (1997a) and Wiesenhütter et al. (2007) must be interpreted  
9 cautiously. Few details are provided in these studies on selection criteria and not all subjects  
10 from the Brüning et al. (2003) case-control study are included. For GSTM1 positive, the higher  
11 prevalence among exposed cases in Brüning et al. (1997a) compared Wiesenhütter et al. (2007)  
12 and the lower prevalence among controls compared to background frequency in the European  
13 population may reflect possible selection biases. On the other hand, the broader base population  
14 included in Brüning et al. (2003) may explain the observed lower frequency of GSTM1 positive  
15 cases in Wiesenhütter et al. (2007). Moreover, Wiesenhütter et al. (2007) does not report  
16 genotype frequencies for controls by exposure status and this information is essential to an  
17 examination of whether renal cell carcinoma risk and TCE exposure may be modified by  
18 polymorphism status. The statistical analyses in both studies was a simple comparison of  
19 exposure prevalence between cases and controls and did not include analyses that stratified on  
20 exposure status. An examination of exposure prevalence is limited as Moore et al. (2010), too,  
21 reported TCE exposure prevalence as similar between exposed cases and controls. Associations  
22 between TCE exposure and kidney cancer for GSTT1 active genotype, however, were reported  
23 in stratified analyses. The more rigorous study design and statistical methods in Moore et al.  
24 (2010) affords more weight to their reported observations than for Brüning et al. (1997a) and  
25 Wiesenhütter et al. (2007). Moore et al. (2010) provides evidence of greater susceptibility to  
26 TCE exposure and kidney cancer among subjects with a functionally active GSTT  
27 polymorphism, particularly among those with certain alleles in single nucleotide polymorphisms  
28 of the cysteine conjugation  $\beta$ -lyase gene region.

29 Of the three larger (in terms of number of cases) studies that did provide results  
30 separately by sex, Dosemeci et al. (1999) suggest that there may be a sex difference for TCE  
31 exposure and renal cell carcinoma (OR: 1.04, [95% CI: 0.6, 1.7]) in males and 1.96 (95% CI:  
32 1.0, 4.0 in females), while Raaschou-Nielsen et al. (2003) report the same standardized incidence  
33 ratio (SIR = 1.2) for both sexes and crude ORs calculated from data from the Pesch et al. (2000a)  
34 study (provided in a personal communication from Beate Pesch, Forschungsinstitut für  
35 Arbeitsmedizin, to Cheryl Scott, EPA, 21 February 2008) are 1.28 for males and 1.23 for

1 females. Whether the Dosemeci et al. (1999) observations are due to susceptibility differences or  
2 to exposure differences between males and females cannot be evaluated. Blair et al. (1998) and  
3 Hansen et al. (2001) also present some results by sex, but these two studies have too few cases to  
4 be informative about a sex difference for kidney cancer.  
5


#### 4.4.2.1.6. Meta-Analysis for Kidney Cancer

6 Meta-analysis (detailed methodology in Appendix C) was adopted as a tool for  
7 examining the body of epidemiologic evidence on kidney cancer and TCE exposure and to  
8 identify possible sources of heterogeneity. The meta-analyses of the overall effect of TCE  
9 exposure on kidney cancer suggest a small, statistically significant increase in risk that was  
10 stronger in a meta-analysis of the highest exposure group. There was no observable  
11 heterogeneity for any of the meta-analyses of the 15 studies and no indication of publication bias.  
12 Thus, these findings of increased risks of kidney cancer associated with TCE exposure are  
13 robust.

14 The meta-analysis of kidney cancer examines 15 cohort and case-control studies  
15 identified through a systematic review and evaluation of the epidemiologic literature on TCE  
16 exposure (Anttila et al., 1995; Axelson et al., 1994; Blair et al., 1998; Boice et al., 1999; Brüning  
17 et al., 2003; Charbotel et al., 2006; Dosemeci et al., 1999; Greenland et al., 1994; Hansen et al.,  
18 2001; Moore et al., 2010; Morgan et al., 1998; Pesch et al., 2000a; Raaschou-Nielsen et al.,  
19 2003; Siemiatycki, 1991; Zhao et al., 2005). Details of the systematic review and meta-analysis  
20 of the TCE studies are fully discussed in Appendix B and C.

21 The summary relative risk (RR<sub>m</sub>) estimate from the primary random effects  
22 meta-analysis of the 15 studies was 1.27(95% CI: 1.13, 1.43). The analysis was dominated by  
23 two (contributing almost 70% of the weight) or three (almost 80% of the weight) large studies  
24 (Dosemeci et al., 1999; Pesch et al., 2000a; Raaschou-Nielsen et al., 2003). Figure 4-1 arrays  
25 individual studies by their weight. No single study was overly influential; removal of individual  
26 studies resulted in RR<sub>m</sub> estimates that were all statistically significant ( $p < 0.005$ ) and that  
27 ranged from 1.24 (with the removal of Brüning et al. (2003)) to 1.30 (with the removal of  
28 Raaschou-Nielsen et al. (2003)). Similarly, the overall RR<sub>m</sub> estimate was not highly sensitive to  
29 alternate RR estimate selections nor was publication bias apparent. There was no apparent  
30 heterogeneity across the 15 studies, i.e., the random effects model and the fixed effect model  
31 gave the same results ( $p_{hetero} = 0.67$ ;  $I = 0\%$ ). Nonetheless, subgroup analyses were done  
32 examining the cohort and case-control studies separately with the random effects model; the  
33 resulting RR<sub>m</sub> estimates were 1.16 (95% CI: 0.96, 1.40) for the cohort studies and 1.48 (1.15,

- 1 1.91) for the case-control studies. There was no heterogeneity in the cohort subgroup ( $p = 0.998$ ;
- 2  $I^2 = 0\%$ ). There was heterogeneity in the case-control subgroup, but it was not statistically




**Figure 4-1. Meta-analysis of kidney cancer and overall TCE exposure (the summary estimate is in the bottom row, represented by the diamond).** Random effects model; fixed effect model same. Symbol sizes reflect relative weights of the studies.

1 significant ( $p = 0.14$ ) and the  $I^2$  value of 41% suggests that the extent of the heterogeneity in this  
2 subgroup was low-to-moderate.

3 Ten studies reported risks for higher exposure groups (Blair et al., 1998; Boice et al.,  
4 1999; Brüning et al., 2003; Charbotel et al., 2006; Dosemeci et al., 1999; Moore et al., 2010;  
5 Morgan et al., 1998; Parent et al., 2000a; Pesch et al., 2000a; Raaschou-Nielsen et al., 2003;  
6 Siemiatycki, 1991; Zhao et al., 2005). Different exposure metrics were used in the various  
7 studies, and the purpose of combining results across the different highest exposure groups was  
8 not to estimate an RRM associated with some level of exposure. Instead, the focus on the highest  
9 exposure category was meant to result in an estimate less affected by exposure misclassification.  
10 In other words, it is more likely to represent a greater differential TCE exposure compared to  
11 people in the referent group than the exposure differential for the overall (typically any vs. none)  
12 exposure comparison. Thus, if TCE exposure increases the risk of kidney cancer, the effects  
13 should be more apparent in the highest exposure groups.

14 The RRM estimate from the random effects meta-analysis of the studies with results  
15 presented for higher exposure groups was 1.64 (95% CI: 1.31, 2.04), higher than the RRM from  
16 the overall kidney cancer meta-analysis. As with the overall analyses, the meta-analyses of the  
17 highest-exposure groups were dominated by Pesch et al. (2000a) and Raaschou-Nielsen et al.  
18 (2003), which provided about 60% of the weight. Axelson et al. (1994), Anttila et al. (1995) and  
19 Hansen et al. (2001) do not report risk ratios for kidney cancer by higher exposure and a  
20 sensitivity analysis was carried out to address reporting bias. The RRM estimate from the  
21 primary random effects meta-analysis with null RR estimates (i.e., RR = 1.0) included for  
22 Axelson et al. (1994), Anttila et al. (1995) and Hansen et al. (2001) to address reporting bias  
23 associated with ever exposed was 1.58 (95% CI: 1.28, 1.96). Figure 4-2 arrays individual studies  
24 by their weight. The inclusion of these three additional studies contributed less than 7% of the  
25 total weight. No single study was overly influential; removal of individual studies resulted in  
26 RRM estimates that were all statistically significant ( $p < 0.005$ ) and that ranged from 1.52 (with  
27 the removal of Raaschou-Nielsen et al. (2003)) to 1.64 (with the removal of Pesch et al. (2000a)).  
28 Similarly, the RRM estimate was not highly sensitive to alternate RR estimate selections (all with  
29  $p < 0.0005$ ) and other than a negligible amount of heterogeneity observed in the sensitivity  
30 analysis with the Pesch job-exposure matrix (JEM) alternate ( $I^2 = 0.64\%$ ), there was no  
31 observable heterogeneity across the studies for any of the meta-analyses conducted with the  
32 highest-exposure groups, including those in which RR values for Anttila, Axelson, and Hansen  
33 were assumed ( $I^2 = 0\%$ ). For Pesch, the job-task exposure matrix (JTEM) approach is preferred  
34 because it seemed to be a more comprehensive and discriminating approach, taking actual job



**Figure 4-2. Meta-analysis of kidney cancer and TCE exposure—highest exposure groups.** With assumed null RR estimates for Antilla, Axelson, and Hansen (see Appendix C text). Random effects model; fixed effect model same. The summary estimate is in the bottom row, represented by the diamond. Symbol sizes reflect relative weights of the studies.



1 tasks into account, rather than just larger job categories. No subgroup analyses (e.g., cohort vs.  
2 case-control studies) were done with the highest exposure group results.

3 NRC (2006) deliberations on trichloroethylene commented on two prominent evaluations  
4 of the then-current TCE epidemiologic literature using meta-analysis techniques, Wartenberg  
5 et al. (2000) and Kelsh et al. (2005), submitted by Exponent-Health Sciences to NRC during  
6 their deliberations and who updated their analysis by including subsequently published studies of  
7 Boice et al. (2006b) and Charbotel et al. (2006), but not Radican et al. (2008), and presented  
8 summary relative risk estimates for cohort and case-control studies, separately, and combined  
9 (Kelsh et al., 2010). Wartenberg et al. (2000) reported an RR<sub>m</sub> of 1.7 (95% CI: 1.1, 2.7) for  
10 kidney cancer incidence in the TCE subcohorts (Anttila et al., 1995; Axelson et al., 1994; Blair et  
11 al., 1998; Henschler et al., 1995). For kidney cancer mortality in TCE subcohorts (Blair et al.,  
12 1998; Boice et al., 1999; Henschler et al., 1995; Morgan et al., 1998; Ritz, 1999a), Wartenberg  
13 et al. (2000) reported an RR<sub>m</sub> of 1.2 (95% CI: 0.8, 1.7). Kelsh et al. (2010) examined a slightly  
14 different grouping of cohort studies as did Wartenberg et al. (2000), presenting a summary  
15 relative risk estimate for kidney cancer incidence and mortality combined. The RR<sub>m</sub> for kidney  
16 cancer in Group I cohort studies (Anttila et al., 1995; Axelson et al., 1994; Blair et al., 1998;  
17 Boice et al., 1999; Hansen et al., 2001; Morgan et al., 1998; Raaschou-Nielsen et al., 2003) was  
18 1.34 (95% CI: 1.07–1.67) with no evidence of heterogeneity and, in Group II cohort studies,  
19 studies lacking documented TCE exposure (Blair et al., 1989; Chang et al., 2003; Costa et al.,  
20 1989; Garabrant et al., 1988; Henschler et al., 1995; Seldén and Ahlborg, 1991; Sinks et al.,  
21 1992), was 1.58 (95% CI: 0.75, 3.32) with evidence of heterogeneity. Removing both Henschler  
22 et al. (1995) and Sinks et al. (1992), considered by Kelsh et al. (2010) as outliers, eliminated  
23 observed heterogeneity and the summary risk estimate was 0.88 (95% CI: 0.8, 1.33). Kelsh et al.  
24 (2010), also, presented separately a summary relative risk for renal cancer case-control studies  
25 and TCE. For case-control studies Charbotel et al, 2006 (Brüning et al., 2003; Dosemeci et al.,  
26 1999; Greenland et al., 1994; Pesch et al., 2000a; Siemiatycki, 1991; Vamvakas et al., 1998), the  
27 RR<sub>m</sub> for renal cell carcinoma was 1.57 (95% CI: 1.06, 2.30) with evidence of heterogeneity, and  
28 RR<sub>m</sub> of 1.33 (95% CI: 1.02, 1.73) and no evidence of heterogeneity in a sensitivity analysis  
29 removing Vamvakas et al. (1998), a study Kelsh et al. (2010) considered as an outlier. Last,  
30 Kelsh et al. (2010) presented three summary relative risk estimates for renal cell cancer Groups I  
31 and II cohort and case-control studies combined: 1.30 (95% CI: 1.04, 1.61) with evidence of  
32 heterogeneity and included 23 studies with kidney cancer risk estimates for all subjects, those  
33 with documented TCE exposure and those unexposed to TCE, and Ritz (1999a) in Group I  
34 studies; 1.42 (95% CI 1.13, 1.77) with evidence of heterogeneity and included 23 studies, with  
35 TCE subcohort kidney cancer risk estimates replacing the total cohort estimate for Group I

1 studies; and, 1.24 (95% CI; 1.06, 1.45) with no evidence of heterogeneity and included  
2 20 studies, counting TCE subcohort kidney cancer risk estimates in Group I studies and  
3 removing the three studies Kelsh et al. (2010) considered as outliers.

4 The present analysis was conducted according to NRC (2006) suggestions for  
5 transparency, systematic review criteria, and examination of both cohort and case-control  
6 studies. EPA's meta-analysis has several advantages to previous ones of TCE exposure and  
7 cancer. The selection criteria adopted in this meta-analysis were intended to identify informative  
8 studies for the evaluation of TCE exposure and cancer, studies with reduced systematic errors.  
9 Neither Henschler et al. (1995) nor Vamvakas et al. (1998), two studies with incomplete cohort  
10 identification or potential selection bias of study controls, met our inclusion criteria and their  
11 inclusion in other meta-analysis may have contributed to the observed heterogeneity in kidney  
12 cancer RRm (Kelsh et al., 2010). Studies with background or low TCE exposure potential also  
13 did not meet another selection criterion as our analysis focused on TCE exposure potential  
14 inferred to each subject by reference to industrial hygiene records, individual biomarkers,  
15 job-exposure matrices, water distribution models, or questionnaire responses that likely had  
16 fewer biases associated with exposure misclassification, although this bias would not have been  
17 completely minimized. Inclusion of studies of lower exposure potential in meta-analyses can  
18 have important implications for identifying a cancer hazard (Steinmaus et al., 2008; Zhang et al.,  
19 2008; Vlaanderen et al., 2011). The present analysis includes the recently published studies of  
20 Charbotel et al. (2006), Moore et al. (2010), and updated mortality of the Blair et al. (1998)  
21 cohort by Radican et al. (2008). As discussed above, the summary estimate from the primary  
22 random effects meta-analysis of the 15 studies was 1.27 (95% CI: 1.13, 1.43). Additionally,  
23 EPA examined kidney cancer risk for higher exposure group. The RRm estimate from the  
24 random effects meta-analysis of the studies with results presented for higher exposure groups  
25 was 1.64 (95% CI: 1.31, 2.04), higher than the RRm from the overall kidney cancer  
26 meta-analysis, and 1.58 (95% CI: 1.28, 1.96) in the meta-analysis with null RR estimates (i.e.,  
27 RR = 1.0) to address possible reporting bias for three studies.

#### 4.4.3. Human Studies of Somatic Mutation of von Hippel-Lindau (VHL) Gene

29 Studies have been conducted to identify mutations in the *VHL* gene in renal cell  
30 carcinoma patients, with and without TCE exposures (Brauch et al., 1999; Charbotel et al., 2007;  
31 Furge et al., 2007; Kenck et al., 1996; Schraml et al., 1999; Toma et al., 2008; Wells et al.,  
32 2009). Inactivation of the *VHL* gene through mutations, LOH and imprinting has been observed  
33 in about 70% of sporadic renal clear cell carcinomas, the most common renal cell carcinoma

1 subtype (Kenck et al., 1996). Other genes or pathways, including c-myc activation and VEGF,  
2 have also been examined as to their role in various renal cell carcinoma subtypes (Furge et al.,  
3 2007; Toma et al., 2008). Furge et al. (2007) reported that there are molecularly distinct forms  
4 of RCC and possibly molecular differences between clear-cell renal cell carcinoma subtypes.  
5 This study was performed using tissues obtained from paraffin blocks. These results are  
6 supported by a more recent study which examined the genetic abnormalities of clear cell renal  
7 cell carcinoma using frozen tissues from 22 cc-RCC patients and paired normal tissues (Toma et  
8 al., 2008). This study found that 20 (91%) of the 22 cases had LOH on chromosome 3p  
9 (harboring the *VHL* gene). Alterations in copy number were also found on chromosome 9 (32%  
10 of cases), chromosome arm 14q (36% of cases), chromosome arm 5q (45% of cases) and  
11 chromosome 7 (32% of cases), suggesting roles for multiple genetic changes in RCC, and is also  
12 supported by genomes-wide single-nucleotide polymorphism analysis (Toma et al., 2008).

13 Several papers link mutation of the *VHL* gene in renal cell carcinoma patients to TCE  
14 exposure. These reports are based on comparisons of *VHL* mutation frequencies in TCE exposed  
15 cases from renal cell carcinoma case-control studies or from comparison to background mutation  
16 rates among renal cell carcinoma case series (see Table 4-42). Brüning et al. (1997b) first  
17 reported a high somatic mutation frequency (100%) in a series of 23 renal cell carcinomas cases  
18 with medium to high intensity TCE exposure as determined by an abnormal single stand  
19 conformation polymorphism (SSCP) pattern, with most variations found in exon two. Only four  
20 samples were sequenced at the time of publication and showed mutations in exon one, two and  
21 three (see Table 4-42). Some of the cases in this study were from the case-control study of  
22 Vamvakas et al. (1998) (see Section 4.4.3 and Appendix C).

23 Brauch et al. (1999; 2004) analyzed renal cancer cell tissues for mutations of the *VHL*  
24 gene and reported increased occurrence of mutations in patients exposed to high concentrations  
25 of TCE. In the first study (Brauch et al., 1999), an employer's liability or worker's  
26 compensation registry was used to identify 44 renal cell carcinoma cases, 18 of whom were also  
27 included in Brüning et al. (1997b). Brauch et al. (1999) found multiple mutations in 42% of the  
28 exposed patients who experienced any mutation and 57% showed loss of heterozygosity. A hot  
29 spot mutation of cytosine to thymine at nucleotide 454 (C454T) was found in 39% of samples  
30 that had a *VHL* mutation and was not found in renal cell cancers from nonexposed patients or in  
31 lymphocyte DNA from either exposed or nonexposed cases or controls. As discussed above,  
32 little information was given on how subjects were selected and whether there was blinding of  
33 from the renal cell carcinoma case-control study of Vamvakas et al. (1998). Brauch et al. (2004)  
34 compared age at diagnosis and histopathologic parameters of tumors as well as somatic mutation

- 1 characteristics in the *VHL* tumor suppressor gene between the TCE-exposed and non-TCE
- 2 exposed renal cell carcinoma patient groups (TCE-exposed from their previous 1999 publication)

**Table 4-42. Summary of human studies on somatic mutations of the VHL gene<sup>a</sup>**

TCE exposure status	Brüning et al. (1997b)	Brauch et al. (1999)		Schraml et al. (1999)		Brauch et al. (2004)		Charbotel et al. (2007)	
	Exposed	Exposed	Unexposed	Exposed	Unexposed	Exposed	Unexposed	Exposed	Unexposed
Number of subjects/ Number with mutations (%)	23/23 (100%)	44/33 (75%)	73/42 (58%)	9/3 (33%)	113/38 (34%)	17/14 (82%)	21/2 (10%)	25/2 (9%)	23/2 (8%)
Renal cell carcinoma subtype	Unknown	Unknown		Clear cell 9 (75%) Papillary 2 (18%) Oncocytomas 1 (8%)	Unknown	Clear cell 37 (%) Oncocytic adenoma 1 (%) Bilateral metachronous 1 (%)	Clear cell 51 (75%) Papillary 10 (10–15%) Chromophobe 4 (5%) Oncocytomas 4 (5%)		
Tissue type analyzed	Paraffin	Paraffin, fresh (lymphocyte)		Paraffin		Paraffin		Paraffin, frozen tissues, Bouin's fixative	
Assay	SSCP, <sup>b</sup> sequencing <sup>b</sup>	SSCP, sequencing, restriction enzyme digestion		CGH, sequencing		Sequencing		Sequencing	
Number of mutations	23	50	42	4	50	24	2	2	2
Type of mutation									
Missense	1	27	NA	1	Unknown	17	2	1	1
Nonmissense <sup>c</sup>	3	23	NA	3	Unknown	7	0	1	1

<sup>a</sup> Adapted from NRC (2006) with addition of Schraml et al. (1999) and Charbotel et al. (2007).

<sup>b</sup> By SSCP. Four (4) sequences confirmed by comparative genomic hybridization.

<sup>c</sup> Includes insertions, frameshifts, and deletions.

CGH = comparative genomic hybridization.

1 to the non TCE-exposed cases newly sequenced in this study). Renal cell carcinoma did not  
2 differ with respect to histopathologic characteristics in either patient group. Comparing results  
3 from TCE-exposed and nonexposed patients revealed clear differences with respect to  
4 (1) frequency of somatic *VHL* mutations, (2) incidence of C454T transition, and (3) incidence of  
5 multiple mutations. The C454T hot spot mutation at codon 81 was exclusively detected in  
6 tumors from TCE-exposed patients, as were multiple mutations. Also, the incidence of *VHL*  
7 mutations in the TCE-exposed group was at least twofold higher than in the nonexposed group.  
8 Overall, these findings support the view that the effect of TCE is not limited to clonal expansion  
9 of cells mutated spontaneously or by some other agent.

10        Brauch et al. (2004) were not able to analyze all RCCs from the Vamvakas study  
11 (Vamvakas et al., 1998), in part because samples were no longer available. Using the data  
12 described by Brauch et al. (2004) (*VHL* mutation found in 15 exposed and 2 nonexposed  
13 individuals, and *VHL* mutation not found in 2 exposed and 19 unexposed individuals), the  
14 calculated OR is 71.3. The lower bound of the OR including the excluded RCCs is derived from  
15 the assumption that all 20 cases that were excluded were exposed but did not have mutations in  
16 *VHL* (*VHL* mutations were found in 15 exposed and 2 unexposed individuals and *VHL* was not  
17 found in 22 exposed and 18 unexposed individuals), leading to an OR of 6.5 that remains  
18 statistically significant.

19        Charbotel et al. (2007) examines somatic mutations in the three *VHL* coding exons in  
20 RCC cases from their case-control study (Charbotel et al., 2006). Of the 87 RCCs in the  
21 case-control study, tissue specimens were available for 69 cases (79%) of which 48 were  
22 cc-RCC. *VHL* sequencing was carried out for only the cc-RCC cases, 66% of the 73 cc-RCC  
23 cases in Charbotel et al. (2006). Of the 48 cc-RCC cases available for *VHL* sequencing,  
24 15 subjects were identified with TCE exposure (31%), an exposure prevalence lower than 43%  
25 observed in the case-control study. Partial to full sequencing of the *VHL* gene was carried out  
26 using polymerase chain reaction (PCR) amplification and *VHL* mutation pattern recognition  
27 software of Bérout et al. (1998). Full sequencing of the *VHL* gene was possible for only  
28 26 RCC cases (36% of all RCC cases). Single point mutations were identified in four cases  
29 (8% prevalence): two unexposed cases, a G > C mutation in exon 2 splice site and a G > A in  
30 exon 1; one case identified with low/medium exposure, T > C mutation in exon 2, and, one case  
31 identified with high TCE exposure, T > C in exon 3. It should be noted that the two cases with  
32 T > C mutations were smokers unlike the cases with G > A or G > C mutations. The prevalence  
33 of somatic *VHL* mutation in this study is quite low compared to that observed in other RCC case  
34 series from this region; around 50% (Bailly et al., 1995; Gallou et al., 2001). To address possible  
35 bias from misclassification of TCE exposure, Charbotel et al. (2006) examined renal cancer risk

1 for jobs associated with a high level of confidence for TCE exposure. As would be expected if  
2 bias was a result of misclassification, they observed a stronger association between higher  
3 confidence TCE exposure and RCC, suggesting that some degree of misclassification bias is  
4 associated with their broader exposure assessment approach. Charbotel et al. (2007) do not  
5 present findings on *VHL* mutations for those subjects with higher level of confidence TCE  
6 exposure assignment.

7 Schraml et al. (1999) did not observe statistically significant differences in DNA  
8 sequence or mutation type in a series of 12 renal cell carcinomas from subjects exposed to  
9 solvents including varying TCE intensity and a parallel series of 113 clear cell carcinomas from  
10 non-TCE exposed patients. Only nine of the RCC were cc-RCC and were sequenced for  
11 mutations. *VHL* mutations were observed in clear cell tumors only; four mutations in three  
12 TCE-exposed subjects compared to 50 mutations in tumors of 38 nonexposed cases. Details as  
13 to exposure conditions are limited to a statement that subjects had been exposed to high doses of  
14 solvents, potential for mixed solvent exposures, and that exposure included a range of TCE  
15 concentrations. Limitations of this study include having a wider range of TCE exposure  
16 intensities as compared to the studies described above (Brauch et al., 1999; Brüning et al.,  
17 1997b), which focused on patients exposed to higher levels of TCE, and the limited number of  
18 TCE-exposed subjects analyzed, being the smallest of all available studies on RCC, TCE and  
19 *VHL* mutation. For these reasons, Schraml et al. (1999) is quite limited for examining the  
20 question of *VHL* mutations and TCE exposure.

21 Szymańska et al. (2010) examines somatic mutations in three *VHL* coding exons in 359  
22 RCC cases, 334 of whom with clear-cell carcinomas, from the case-control study of Moore et al.  
23 (2010) as part of a pilot examination of mutation in three other genes, TP53, EGFR, and KRAS.  
24 The prevalence of *VHL* mutations was high in the RCC series, 72% of the tumors carried at least  
25 one function mutation, Although occupational exposures were not examined and data was not  
26 presented, Szymańska et al. (2010) reported that *VHL* mutations were not associated with TCE  
27 exposure.

28 A number of additional methodological issues need to be considered in interpreting these  
29 studies. Isolation of DNA for mutation detection has been performed using various tissue  
30 preparations, including frozen tissues, formalin fixed tissues and tissue sections fixed in Bouin's  
31 solution. Ideally, studies would be performed using fresh or freshly frozen tissue samples to  
32 limit technical issues with the DNA extraction. When derived from other sources, the quality  
33 and quantity of the DNA isolated can vary, as the formic acid contained in the formalin solution,  
34 fixation time and period of storage of the tissue blocks often affect the quality of DNA. Picric  
35 acid contained in Bouin's solution is also known to degrade nucleic acids resulting in either low

1 yield or poor quality of DNA. In addition, during collection of tumor tissues, contamination of  
2 neighboring normal tissue can easily occur if proper care is not exercised. This could lead to the  
3 ‘dilution effect’ of the results—i.e., because of the presence of some normal tissue, frequency of  
4 mutations detected in the tumor tissue can be lower than expected. These technical difficulties  
5 are discussed in these papers, and should be considered when interpreting the results.  
6 Additionally, selection bias is possible given tissue specimens were not available for all RCC  
7 cases in Vamvakas et al. (1998) or in Charbotel et al. (2006). Some uncertainty associated with  
8 misclassification bias is possible given the lack of TCE exposure information to individual  
9 subjects in Schraml et al. (1999) and in Charbotel et al. (2007) from their use of broader  
10 exposure assessment approach compared to that associated with the higher confident exposure  
11 assignment approach. A recent study by Nickerson et al. (2008) addresses many of these  
12 concerns by utilizing more sensitive methods to look at both the genetic and epigenetic issues  
13 related to *VHL* inactivation. This study was performed on DNA from frozen tissue samples and  
14 used a more sensitive technique for analysis for mutations (endonuclease scanning) as well as  
15 analyzing for methylation changes that may lead to inactivation of the *VHL* gene. This method  
16 of analysis was validated on tissue samples with known mutations. Of the 205 cc-RCC samples  
17 analyzed, 169 showed mutations in the *VHL* gene (82.4%). Of those 36 without mutation, 11  
18 were hypermethylated in the promoter region, which will also lead to inactivation of the *VHL*  
19 gene. Therefore, this study showed inactivating alterations in the *VHL* gene (either by mutation  
20 or hypermethylation) in 91% tumor samples analyzed.

21 The limited animal studies examining the role of *VHL* mutation following exposure to  
22 chemicals including TCE are described below in Section 4.4.6.1.1. Conclusions as to the role of  
23 *VHL* mutation in TCE-induced kidney cancer, taking into account both human and experimental  
24 data, are presented below in Section 4.4.7.

25

#### 4.4.4. Kidney Noncancer Toxicity in Laboratory Animals

26 Acute, subchronic, and chronic exposures to TCE cause toxicity to the renal tubules in  
27 rats and mice of both sexes, via both inhalation (see Table 4-43) and oral (see Table 4-44)  
28 exposures. Nephrotoxicity from acute exposures to TCE has only been reported at relatively  
29 high doses, although histopathological changes have not been investigated in these experiments.  
30 Information about specific location of lesions is presented where available. TCE exposure for  
31 13-weeks (corn oil gavage) led to increased nephrotoxicity but no significant increases in  
32 preneoplastic or neoplastic lesions as compared to controls (Mally et al., 2006). Chronic  
33 nephropathy was also observed in both sexes of Osborne-Mendel rats following exposure to TCE

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 (549 and 1,097 mg/kg-day, 78 week). Chakrabarti and Tuchweber (1988) found that TCE  
2 administered to male F344 rats by intraperitoneal injection (723–2,890 mg/kg) or by inhalation  
3 (1,000–2,000 ppm for 6 hours) produced elevated urinary NAG,  $\gamma$ -glutamyl transpeptidase  
4 (GGT), glucose excretion, blood urea nitrogen (BUN), and high molecular weight protein  
5 excretion, characteristic signs of proximal tubular, and possibly glomerular injury, as soon as  
6 24 hours postexposure. In the intraperitoneal injection experiments, inflammation was observed,  
7 although some inflammation is expected due to the route of exposure, and nephrotoxicity effects  
8 were only statistically significantly elevated at the highest dose (2,890 mg/kg). In the inhalation  
9 experiments, the majority of the effects were statistically significant at both 1,000 and  
10 2,000 ppm. Similarly, at these exposures, renal cortical slice uptake of *p*-aminohippurate was  
11 inhibited, indicating reduced proximal tubular function. Cojocel et al. (1989) found similar  
12 effects in mice administered TCE by intraperitoneal injection (120–1,000 mg/kg) at 6 hours  
13 postexposure, such as the dose-dependent increase in plasma BUN concentrations and decrease  
14 in *p*-aminohippurate accumulation in renal cortical slices. In addition, malondialdehyde (MDA)  
15 and ethane production were increased, indicating lipid peroxidation.

16       Kidney weight increases have been observed following inhalation exposure to TCE in  
17 both mice (Kjellstrand et al., 1983b) and rats (Woolhiser et al., 2006) and following lifetime  
18 drinking water exposure in a genetically-prone murine model (Peden-Adams et al., 2008).  
19 Kjellstrand et al. (1983b) demonstrated an increase in kidney weights in both male (20%  
20 compared to control) and female (10% compared to control) mice following intermittent and  
21 continuous TCE whole-body inhalation exposure (up to 120 days). This increase was significant  
22 in males as low as 75 ppm exposure and in females starting at 150-ppm exposure. The latter  
23 inhalation study, an unpublished report by Woolhiser et al. (2006), was designed to examine  
24 immunotoxicity of TCE but also contains information regarding kidney weight increases in  
25 female Sprague Dawley (S-D) rats exposed to 0-, 100-, 300-, and 1,000-ppm TCE for  
26 6 hours/day, 5 days/week, for 4 weeks. Relative kidney weights were significantly elevated

**Table 4-43. Inhalation studies of kidney noncancer toxicity in laboratory animals**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Kidney effect(s) discussed in Section 4.4.4
Chakrabarti and Tuchweber (1988)	Fischer 344 rats (M)	Inhalation	0–20,00 ppm, 6 h	6/group	Increased signs of proximal tubular damage.
Green et al. (1998)	Fischer 344 rats (M)	Inhalation	0, 250, 500 ppm, 6 h/d for 1, 7, 15, 21, 28 d	3–5/group	Increased formic acid excretion; plasma and urinary markers of nephrotoxicity unchanged.
<b>Kjellstrand et al. (1983b)</b>	<b>NMRI mice (M and F)</b>	<b>Inhalation</b>	<b>0–3,600 ppm, variable time periods of 1–24 h/d, for 30 or 120 d</b>	<b>10–20/group</b>	<b>Increased kidney weight.</b>
Maltoni et al. (1986)	Sprague-Dawley rats, (M and F) B6C3F1 mice (M and F)	Inhalation	0, 100, 300, 600 ppm, 7 h/d, 5 d/wk, 104 wk exposure, observed for lifespan	116–141/group	Meganeucleocytosis in male rats (Details in Table 4.49).
Mensing et al. (2002)	Long-Evans rats (M)	Inhalation	0–500 ppm, 6 h/5 d/wk, 6 mo	5/group	Increased signs of nephrotoxicity.
<b>Woolhiser et al. (2006)</b>	<b>Sprague-Dawley rats (F)</b>	<b>Inhalation</b>	<b>0, 100, 300, and 1,000 ppm, 6 h/d, 5 d/wk, 4 wk</b>	<b>16/group</b>	<b>Increased kidney weight.</b>

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

**Table 4-44. Oral and i.p. studies of kidney noncancer toxicity in laboratory animals**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Kidney effect(s) discussed in Section 4.4.4
Chakrabarti and Tuchweber (1988)	Fischer 344 rats (M)	Intraperitoneal injection	0–2,890 mg/kg-day	6/group	Increased signs of proximal tubular damage.
Cojocel et al. (1989)	NMRI mice (M)	Intraperitoneal injection (sesame oil)	0–1,000 mg/kg	4/group	Increased signs of nephrotoxicity.
Green et al. (1997a)	Fischer 344 rats (M) B6C3F1 mice (M)	Gavage (corn oil)	0, 500, 2,000 mg/kg-day, 1 or 10 d	5 or 10/group	Increases in biochemical markers of kidney damage.
Green et al. (2003)	Fischer 344 rats (M)	Drinking water	0–54.3 mg/kg-day, 52 wk	60/group	Increased kidney weights and tubular degeneration.
Mally et al. (2006)	Eker rat (M)	Gavage (corn oil)	0–1,000 mg/kg BW, 5 d/wk, 13 wk	10/group	Increased nephrotoxicity.
Maltoni et al. (1986)	Sprague-Dawley rats (M and F)	Gavage (olive oil)	0, 50, 250 mg/kg-day 4–5 d/wk, 52 wk	30/group	Megakaryocytosis in male rats (Details in Table 4.47).
<b>NCI (1976)</b>	<b>Osborne-Mendel rats (M and F) B6C3F1 mice (M and F)</b>	<b>Gavage (corn oil)</b>	<b>0–2,339 mg/kg-day, variable doses, 5 d/wk, 78 wk</b>	<b>50/group</b>	<b>Toxic nephrosis in all exposed animals (Details in Table 4.46).</b>
<b>NTP (1988)</b>	<b>ACI, August, Marshall, and Osborne-Mendel rats (M and F)</b>	<b>Gavage (corn oil)</b>	<b>0, 500, 1,000 mg/kg-day, 5 d/wk, 103 wk</b>	<b>50/group</b>	<b>Cytomegaly and toxic nephropathy observed in all exposed rats (Details in Table 4-48).</b>
NTP (1990)	Fischer 344 rats (M and F) B6C3F1 mice (M and F)	Gavage (corn oil)	Rats: 0–2,000 mg/kg-day, Mice: 0–6,000 mg/kg-day, 5d/wk, 13 wk	10/group	Cytomegaly and karyomegaly of renal tubular epithelium in mice and rats (Details in Table 4.45).
Peden-Adams et al. (2008)	MRL mice (M and F)	Drinking water	0, 1,400, 14,000 ppb, lifetime	6/group	Increased kidney weight in male mice.

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

1 (17.4% relative to controls) at 1,000-ppm TCE exposure. However, the small number of animals  
2 and the variation in initial animal weight limit the ability of this study to determine statistically  
3 significant increases. The Peden-Adams et al. (2008) study was designed to assess the effects of  
4 TCE exposure in a genetically-prone murine lupus model. Although the study did not  
5 demonstrate an increase in the development of autoimmune disease markers (for details see  
6 Section 4.6.2), changes in body weight and organ weights in males were observed. Following  
7 lifetime exposure to TCE (14,000 ppb) in drinking water, males exhibited a decreasing trend in  
8 body mass of 12% from controls (female body weights not altered). Spleen, thymic and kidney  
9 mass in females were not altered following exposure to TCE, while an 18% increase in kidney  
10 mass was observed in the high dose treatment group (14,000 ppb) in males.

11 Similarly, overt signs of subchronic nephrotoxicity, such as changes in blood or urinary  
12 biomarkers, are also primarily a high dose phenomenon, although histopathological changes are  
13 evident at lower exposures. Green et al. (1997a) reported administration of 2,000 mg/kg-day  
14 TCE by corn oil gavage for 42 days in F344 rats caused increases of around twofold of control  
15 results in urinary markers of nephrotoxicity such as urine volume and protein (both 1.8×), NAG  
16 (1.6×), glucose (2.2×) and alkaline phosphatase (ALP; 2.0×), similar to the results of the acute  
17 study of Chakrabarti and Tuchweber (1988), above. No morphological changes were observed  
18 in kidneys from any animals (Green et al., 1997a). At lower dose levels, Green et al. (1998)  
19 reported that plasma and urinary markers of nephrotoxicity were unchanged. In particular, after  
20 1–28 day exposures to 250 or 500 ppm TCE for 6 hours/day, there were no statistically  
21 significant differences in plasma levels of BUN or in urinary levels of creatinine, protein, ALP,  
22 NAG, or GGT. However, increased urinary excretion of formic acid, accompanied by changes  
23 in urinary pH and increased ammonia, was found at these exposures. Interestingly, at the same  
24 exposure level of 500 ppm (6 hours/day, 5 days/week, for 6 months), Mensing et al. (2002)  
25 reported elevated excretion of low molecular weight proteins and NAG, biomarkers of  
26 nephrotoxicity, but after the longer exposure duration of 6 months.

27 Numerous studies have reported histological changes from TCE exposure for subchronic  
28 and chronic durations (Maltoni et al., 1988, 1986; Mensing et al., 2002; NTP, 1990, 1988). As  
29 summarized in Table 4-45, in 13-week studies in F344 rats and B6C3F1 mice, NTP (1990)  
30 reported relatively mild cytomegaly and karyomegaly of the renal tubular epithelial cells at the  
31 doses 1,000–6,000 mg/kg-day (at the other doses, tissues were not examined). The NTP report  
32 noted that “these renal effects were so minimal that they were diagnosed only during a  
33 reevaluation of the tissues...prompted by the production of definite renal toxicity in the 2-year  
34 study.” In the 6 month, 500-ppm inhalation exposure experiments of Mensing et al. (2002),  
35 some histological changes were noted in the glomeruli and tubuli of exposed rats, but they

**Table 4-45. Summary of renal toxicity and tumor findings in gavage studies of trichloroethylene by NTP (1990)**

Sex	Dose (mg/kg) <sup>a</sup>	Cytomegaly and karyomegaly incidence (severity <sup>b</sup> )	Adenoma (overall; terminal)	Adenocarcinoma (overall; terminal)
1/d, 5 d/wk, 13-wk study, F344/N rats				
Male	0, 125, 250, 500, 100	Tissues not evaluated	None reported	
	2,000	8/9 (Minimal/mild)		
Female	0, 62.5, 125, 250, 500	Tissues not evaluated		
	1,000	5/10 (Equivocal/minimal)		
1/d, 5 d/wk, 13-wk study, B6C3F1 mice				
Male	0, 375, 750, 1,500	Tissues not evaluated	None reported	
	3,000	7/10 <sup>c</sup> (Mild/moderate)		
	6,000	- <sup>d</sup>		
Female	0, 375, 750, 1,500	Tissues not evaluated		
	3,000	9/10 (Mild/moderate)		
	6,000	1/10 (Mild/moderate)		
1/d, 5 d/wk, 103-wk study, F344/N rats				
Male	0	0% (0)	0/48; 0/33	0/48; 0/33
	500	98% (2.8)	2/49; 0/20	0/49; 0/20
	1,000	98% (3.1)	0/49; 0/16	3/49; 3/16 <sup>e</sup>
Female	0	0% (0)	0/50; 0/37	0/50; 0/37
	500	100% (1.9)	0/49; 0/33	0/49; 0/33
	1,000	100% (2.7)	0/48; 0/26	1/48; 1/26
1/d, 5 d/wk, 103-wk study, B6C3F1 mice				
Male	0	0% (0)	1/49; 1/33	0/49; 0/33
	1,000	90% (1.5)	0/50; 0/16	1/50; 0/16
Female	0	0% (0)	0/48; 0/32	0/48; 0/32
	1,000	98% (1.8)	0/49; 0/23	0/49; 0/23

Study carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup> Corn oil vehicle.

<sup>b</sup> Numerical scores reflect the average grade of the lesion in each group (1, slight; 2, moderate; 3, well marked; and 4, severe).

<sup>c</sup> Observed in four mice that died after 7–13 wk and in three that survived the study.

<sup>d</sup> All mice died during the first wk.

<sup>e</sup>  $p = 0.028$ .

1 provided no detailed descriptions beyond the statement that “perivascular, interstitial infections  
2 and glomerulonephritis could well be detected in kidneys of exposed rats.”

3 After 1–2 years of chronic TCE exposure by gavage (NCI, 1976; NTP, 1990, 1988) or  
4 inhalation (Maltoni et al., 1988) (see Tables 4-45 to 4-49), both the incidence and severity of  
5 these effects increases, with mice and rats exhibiting lesions in the tubular epithelial cells of the  
6 inner renal cortex that are characterized by cytomegaly, karyomegaly, and toxic nephrosis. As  
7 with the studies at shorter duration, these chronic studies reported cytomegaly and karyomegaly  
8 of tubular cells. NTP (1990) specified the area of damage as the pars recta, located in the  
9 corticomedullary region. It is important to note that these effects are distinct from the chronic  
10 nephropathy and inflammation observed in control mice and rats (Lash et al., 2000b; Maltoni et  
11 al., 1988; NCI, 1976).

12 These effects of TCE on the kidney appear to be progressive. Maltoni et al. (1988) noted  
13 that the incidence and degree of renal toxicity increased with increased exposure time and  
14 increased time from the start of treatment. As mentioned above, signs of toxicity were present in  
15 the 13 week study (NTP, 1988), and NTP (1990) noted cytomegaly at 26 weeks. NTP (1990)  
16 noted that as “exposure time increased, affected tubular cells continued to enlarge and additional  
17 tubules and tubular cells were affected,” with toxicity extending to the cortical area as kidneys  
18 became more extensively damaged. NTP (1988, 1990) noted additional lesions that increased in  
19 frequency and severity with longer exposure, such as dilation of tubules and loss of tubular cells  
20 lining the basement membrane (“stripped appearance” (NTP, 1988) or flattening of these cells  
21 (NTP, 1990)). NTP (1990) also commented on the intratubular material and noted that the  
22 tubules were empty or “contained wisps of eosinophilic material.”

23 With gavage exposure, these lesions were present in both mice and rats of both sexes, but  
24 were on average more severe in rats than in mice, and in male rats than in female rats (NTP,  
25 1990). Thus, it appears that male rats are most sensitive to these effects, followed by female rats  
26 and then mice. This is consistent with the experiments of Maltoni et al. (1988), which only  
27 reported these effects in male rats. The limited response in female rats or mice of either sex in  
28 these experiments may be related to dose or strain. The lowest chronic gavage doses in the  
29 National Cancer Institute (NCI, 1976) and NTP (1988, 1990) F344 rat experiments was  
30 500 mg/kg-day, and in all these cases at least 80% (and frequently 100%) of the animals showed  
31 cytomegaly or related toxicity. By comparison, the highest gavage dose in the Maltoni et al.  
32 (1988) experiments (250 mg/kg-day) showed lower incidences of renal cytomegaly and  
33 karyomegaly in male Sprague-Dawley rats (47 and 67%, overall and corrected incidences) and  
34 none in female rats. The B6C3F1 mouse strain was used in the NCI (1976), NTP (1990), and  
35

1  
2  
3

**Table 4-46. Summary of renal toxicity and tumor findings in gavage studies of trichloroethylene by NCI (1976)**

<b>Sex</b>	<b>Dose (mg/kg)<sup>a</sup></b>	<b>Toxic nephrosis (overall; terminal)</b>	<b>Adenoma or adenocarcinoma (overall; terminal)<sup>b</sup></b>
1/d, 5 d/wk, 2-yr study, Osborn-Mendel rats			
Males	0	0/20; 0/2	0/20; 0/2
	549	46/50; 7/7	1/50; <sup>c</sup> 0/7
	1,097	46/50; 3/3	0/50; 0/3
Females	0	0/20; 0/8	0/20; 0/8
	549	39/48; 12/12	0/48; 0/12
	1,097	48/50; 13/13	0/50; 0/13
1/d, 5 d/wk, 2-yr study, B6C3F1 mice			
Males	0	0/20; 0/8	0/20; 0/8
	1,169	48/50; 35/35	0/50; 0/35
	2,339	45/50; 20/20	1/50; <sup>d</sup> 1/20
Females	0	0/20; 0/17	0/20; 0/17
	869	46/50; 40/40	0/50; 0/40
	1,739	46/47; <sup>e</sup> 39/39	0/47; 0/39

4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14

Study carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup> Treatment period was 48 wk for rats, 66 wk for mice. Doses were changed several times during the study based on monitoring of body weight changes and survival. Dose listed here is the time-weighted average dose over the days on which animals received a dose.

<sup>b</sup> A few malignant mixed tumors and hamartomas of the kidney were observed in control and low dose male rats, but are not counted here.

<sup>c</sup> Tubular adenocarcinoma.

<sup>d</sup> Tubular adenoma.

<sup>e</sup> One mouse was reported with “nephrosis,” but not “nephrosis toxic,” and so was not counted here.

1  
2  
3

**Table 4-47. Summary of renal toxicity findings in gavage studies of trichloroethylene by Maltoni et al. (1988)**

<b>Sex</b>	<b>Dose (mg/kg)<sup>a</sup></b>	<b>Megalonucleocytosis<sup>b</sup> (overall; corrected<sup>c</sup>)</b>
1/d, 4–5 d/wk, 52-wk exposure, observed for lifespan, Sprague-Dawley rats		
Males	0	0/20; 0/22
	50	0/30; 0/24
	250	14/30; 14/21
Females	0	0/30; 0/30
	50	0/30; 0/29
	250	0/30; 0/26

4  
5  
6  
7  
8  
9  
10  
11

<sup>a</sup> Olive oil vehicle.

<sup>b</sup> Renal tubuli megalonucleocytosis is the same as cytomegaly and karyomegaly of renal tubuli cells (Maltoni et al., 1988).

<sup>c</sup> Denominator for “corrected” incidences is the number of animals alive at the time of the first kidney lesion in this experiment (39 wk).



1  
2  
3

**Table 4-48. Summary of renal toxicity and tumor incidence in gavage studies of trichloroethylene by NTP (1988)**

Sex	Dose (mg/kg) <sup>a</sup>	Cytomegaly	Toxic nephropathy	Adenoma (overall; terminal)	Adenocarcinoma (overall; terminal)
1/d, 5 d/wk, 2-yr study, ACI rats					
Male	0	0/50	0/50	0/50; 0/38	0/50; 0/38
	500	40/49	18/49	0/49; 0/19	1/49; 0/19
	1,000	48/49	18/49	0/49; 0/11	0/49; 0/11
Female	0	0/48	0/48	0/48; 0/34	0/48; 0/34
	500	43/47	21/47	2/47; 1/20	1/47; 1/20
	1,000	42/43	19/43	0/43; 0/19	1/43; 0/19
1/d, 5 d/wk, 2-yr study, August rats					
Male	0	0/50	0/50	0/50; 0/21	0/50; 0/21
	500	46/50	10/50	1/50; 0/13	1/50; 1/13
	1,000	46/49	31/49	1/49; 1/16	0/49; 0/16
Female	0	0/49	0/49	1/49; 1/23	0/49; 0/23
	500	46/48	8/48	2/48; 1/26	2/48; 2/26
	1,000	50/50	29/50	0/50; 0/25	0/50; 0/25
1/d, 5 d/wk, 2-yr study, Marshall rats					
Male	0	0/49	0/49	0/49; 0/26	0/49; 0/26
	500	48/50	18/50	1/50; 0/12	0/50; 0/12
	1,000	47/47	23/47	0/47; 0/6	1/47; 0/6
Female	0	0/50	0/50	1/50; 0/30	0/50; 0/30
	500	46/48	30/48	1/48; 1/12	1/48; 0/12
	1,000	43/44	30/44	0/44; 0/10	1/44; 1/10
1/d, 5 d/wk, 2-yr study, Osborne-Mendel rats					
Male	0	0/50	0/50	0/50; 0/22	0/50; 0/22
	500	48/50	39/50	6/50; 5/17	0/50; 0/17
	1,000	49/50	35/50	1/50; 1/15	1/50; 0/15
Female	0	0/50	0/50	0/50; 0/20	0/50; 0/20
	500	48/50	30/50	0/50; 0/11	0/50; 0/11
	1,000	49/49	39/49	1/49; 0/7	0/49; 0/7

4  
5  
6  
7

Study carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup> Corn oil vehicle.

**Table 4-49. Summary of renal toxicity and tumor findings in inhalation studies of trichloroethylene by Maltoni et al. (1988)<sup>a</sup>**

Sex	Concentration (ppm)	Meganeucleocytosis <sup>b</sup> (overall; corrected)	Adenoma (overall; corrected)	Adenocarcinoma (overall; corrected)
7 h/d, 5 d/wk, 2-yr exposure, observed for lifespan, Sprague-Dawley rats <sup>c</sup>				
Male	0	0/135; 0/122	0/135; 0/122	0/135; 0/122
	100	0/130; 0/121	1/130; 1/121	0/130; 0/121
	300	22/130; 22/116	0/130; 0/116	0/130; 0/116
	600	101/130; 101/124	1/130; 1/124	4/130; 4/124
Female	0	0/145; 0/141	0/145; 0/141	0/145; 0/141
	100	0/130; 0/128	1/130; 1/128	0/130; 0/128
	300	0/130; 0/127	0/130; 0/127	0/130; 0/127
	600	0/130; 0/127	0/130; 0/127	1/130; 1/127
7 h/d, 5 d/wk, 78-wk exposure, observed for lifespan, B6C3F1 mice <sup>d</sup>				
Male	0	0/90	0/90	0/90
	100	0/90	0/90	1/90
	300	0/90	0/90	0/90
	600	0/90	0/90	0/90
Female	0	0/90	0/90	1/90
	100	0/90	0/90	0/90
	300	0/90	0/90	0/90
	600	0/90	0/90	0/90

Study carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup> Three inhalation experiments in this study found no renal megalonucleocytosis, adenomas, or adenocarcinomas: BT302 (8-wk exposure to 0, 100, 600 ppm in Sprague-Dawley rats); BT303 (8-wk exposure to 0, 100, or 600 ppm in Swiss mice); and BT305 (78-wk exposure to 0, 100, 300, or 600 ppm in Swiss mice).

<sup>b</sup> Renal tubuli meganeucleocytosis is the same as cytomegaly and karyomegaly of renal tubuli cells (Maltoni et al., 1988).

<sup>c</sup> Combined incidences from experiments BT304 and BT304bis. Corrected incidences reflect number of rats alive at 47 wk, when the first renal tubular megalonucleocytosis in these experiments appeared.

<sup>d</sup> Female incidences are from experiment BT306, while male incidences are from experiment BT306bis, which was added to the study because of high, early mortality due to aggressiveness and fighting in males in experiment BT306. Corrected incidences not show, because only the renal adenocarcinomas appeared at 107 wk in the male and 136 in the female, when the most of the mice were already deceased.

1 Maltoni et al. (1988) studies (see Tables 4-45–4-49). While the two gavage studies (NCI, 1976;  
2 NTP, 1990) were consistent, reporting at least 90% incidence of cytomegaly and karyomegaly at  
3 all studied doses, whether dose accounts for the lack of kidney effects in Maltoni et al. (1988)  
4 requires comparing inhalation and gavage dosing. Such comparisons depend substantially on the  
5 internal dose-metric, so conclusions as to whether dose can explain differences across studies  
6 cannot be addressed without dose-response analysis using PBPK modeling. Some minor  
7 differences were found in the multistrain NTP study (1988), but the high rate of response makes  
8 distinguishing among them difficult. Soffritti (personal communication with JC Caldwell,  
9 February 14, 2006) did note that the colony from which the rats in Maltoni et al. (1988; 1986)  
10 experiments were derived had historically low incidences of chronic progressive nephropathy  
11 and renal cancer.  
12

#### **4.4.5. Kidney Cancer in Laboratory Animals**

##### **4.4.5.1.1. Inhalation Studies of Trichloroethylene (TCE)**

13 A limited number of inhalation studies examined the carcinogenicity of TCE, with no  
14 statistically-significantly increases in kidney tumor incidence reported in mice or hamsters  
15 (Fukuda et al., 1983; Henschler et al., 1980; Maltoni et al., 1988; 1986). The cancer bioassay by  
16 Maltoni et al. (1988; 1986) reported no statistically significant increase in kidney tumors in mice  
17 or hamsters, but renal adenocarcinomas were found in male (4/130) and female (1/130) rats at  
18 the high dose (600 ppm) after 2 years exposure and observation at natural death. In males, these  
19 tumors seemed to have originated in the tubular cells, and were reported to have never been  
20 observed in over 50,000 Sprague-Dawley rats (untreated, vehicle-treated, or treated with  
21 different chemicals) examined in previous experiments in the same laboratory (Maltoni et al.,  
22 1986). The renal adenocarcinoma in the female rat was cortical and reported to be similar to that  
23 seen infrequently in historical controls. This study also demonstrated the appearance of  
24 increased cytomegaly or megalonucleocytosis in the tubular cells, a lesion that was  
25 significantly and dose-dependently increased in male rats only (see Table 4-49). Maltoni et al.  
26 (1986) noted that some considerations supported either the hypothesis that these were precursor  
27 lesions of renal adenocarcinomas cancer or the hypothesis that these are not precursors but rather  
28 the morphological expression of TCE-induced regressive changes. The inhalation studies by  
29 Fukuda et al. (1983) in Sprague-Dawley rats and female ICR mice, reported one clear cell  
30 carcinoma in rats exposed to the highest concentration (450 ppm) but saw no increase in kidney  
31 tumors in mice. This result was not statistically significant (see Table 4-50) and no details are

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 given about the specific location of the tumors. One negative study (Henschler et al., 1980)  
2 tested NMRI mice, Wistar rats, and Syrian hamsters of both sexes (60 animals per strain), and  
3

**Table 4-50. Summary of renal tumor findings in inhalation studies of trichloroethylene by Henschler et al. (1980)<sup>a</sup> and Fukuda et al. (1983)<sup>b</sup>**

Sex	Concentration (ppm)	Adenomas	Adenocarcinomas
6 h/d, 5 d/wk, 18-mo exposure, 30-mo observation, Han:NMRI mice (Henschler et al., 1980)			
Males	0	4/30	1/30
	100	1/29	0/30
	500	1/29	0/30
Females	0	0/29	0/29
	100	0/30	0/30
	500	0/28	0/28
6 h/d, 5 d/wk, 18-mo exposure, 36-mo observation, Han:WIST rats (Henschler et al., 1980)			
Males	0	2/29	0/29
	100	1/30	0/30
	500	2/30	1/30
Females	0	0/28	0/28
	100	0/30	0/30
	500	1/30	0/30
7 h/d, 5 d/wk, 2-yr study, Crj:CD (S-D) rats (Fukuda et al., 1983)			
Females	0	0/50	0/50
	50	0/50	0/50
	150	0/47	0/47
	450	0/51	1/50

<sup>a</sup> Henschler et al. (1980) observed no renal tumors in control or exposed Syrian hamsters.

<sup>b</sup> Fukuda et al. (1983) observed no renal tumors in control or exposed Crj:CD-1 (ICR) mice.

observed no significant increase in renal tubule tumors any of the species tested. Benign adenomas were observed in male mice and rats, a single adenocarcinoma was reported in male rats at the highest dose, and no renal adenocarcinomas reported in females of either species (see Table 4-50). Renal cell carcinomas appear to be very rare in Wistar rats, with historical control rates reported to be about 0.4% in males and 0.2% in females (Potracki and Walsh, 1998), so these data are very limited in power to detect small increases in their incidence.

#### 4.4.5.1.2. Gavage and Drinking Water Studies of Trichloroethylene (TCE)

1           Several chronic gavage studies exposing multiple strains of rats and mice to  
2 0–3,000 mg/kg TCE for at least 52 weeks have been conducted (see Tables 4-45 to 4-48, 4-51)  
3 (Henschler et al., 1984; Maltoni et al., 1986; NCI, 1976; NTP, 1988, 1990; Van Duuren et al.,  
4 1979). Van Duuren et al. (1979) examined TCE and 14 other halogenated compounds for  
5 carcinogenicity in both sexes of Swiss mice. While no excess tumors were observed, the dose  
6 rate (0.5 mg once per week, or an average dose rate of approximately 2.4 mg/kg-day for a 30 g  
7 mouse) is about 400-fold lower than that in the other gavage studies. Inadequate design and  
8 reporting of this study limit the ability to use the results as an indicator of TCE carcinogenicity.  
9 In the NCI (1976) study, the results for Osborne-Mendel rats were considered by the authors to  
10 be inconclusive due to significant early mortality. Two male rats demonstrated kidney lesions  
11 (dilated renal pelvis and dark red renal medulla), but in rats of both sexes, no increase was seen  
12 in primary tumor induction over that observed in controls. While both sexes of B6C3F1 mice  
13 showed a compound-related increase in nephropathy, no increase in tumors over controls was  
14 observed. The NCI study (1976) used technical grade TCE which contained two known  
15 carcinogenic compounds as stabilizers (epichlorohydrin and 1,2-epoxybutane). However, a  
16 subsequent study by Henschler et al. (1984) in mice reported no significant differences in  
17 systemic tumorigenesis between pure, industrial, and stabilized TCE, suggesting that  
18 concentrations of these stabilizers are too low to be the cause of tumors. A later gavage study by  
19 NTP (1988), using TCE stabilized with diisopropylamine, observed an increased incidence of  
20 renal tumors in all four strains of rats (ACI, August, Marshall, and Osborne-Mendel). All  
21 animals exposed for up to 2 years (rats and mice) had non-neoplastic kidney lesions (tubular cell  
22 cytomegaly), even if they did not later develop kidney cancer (see Table 4-48). This study was  
23 also considered inadequate by the authors because of chemically induced toxicity, reduced  
24 survival, and incomplete documentation of experimental data. The final NTP study (1990) in  
25 male and female F344 rats and B6C3F1 mice used epichlorohydrin-free TCE. Only in the  
26 highest-dose group (1,000 mg/kg) of male F344 rats was renal carcinoma statistically significant  
27 increased. The results for detecting a carcinogenic response in rats were considered by the  
28 authors to be equivocal because both groups receiving TCE showed significantly reduced  
29 survival compared to vehicle controls and because of a high rate (e.g., 20% of the animals in the  
30 high-dose group) of death by gavage error. However, historical control incidences at NTP of  
31

1  
2  
3

**Table 4-51. Summary of renal tumor findings in gavage studies of trichloroethylene by Henschler et al. (1984)<sup>a</sup> and Van Duuren et al. (1979)<sup>b</sup>**

<b>Sex (TCE dose)</b>	<b>Control or TCE exposed (stabilizers if present)</b>	<b>Adenomas</b>	<b>Adenocarcinomas</b>
5 d/wk, 18-mo exposure, 24-mo observation, Swiss mice (Henschler et al., 1984)			
Males (2.4g/kg BW)	Control (none)	1/50	1/50
	TCE (triethanolamine)	1/50	1/50
	TCE (industrial)	0/50	0/50
	TCE (epichlorohydrin (0.8%))	0/50	0/50
	TCE (1,2-epoxybutane (0.8%))	2/50	2/50
	TCE (both epichlorohydrin (0.25%) and 1,2-epoxybutane (0.25%))	0/50	0/50
Females (1.8 g/kg BW)	Control (none)	0/50	1/50
	TCE (triethanolamine)	4/50	0/50
	TCE (industrial)	0/50	0/50
	TCE (epichlorohydrin (0.8%))	0/50	0/50
	TCE (1,2-epoxybutane (0.8%))	0/50	0/50
	TCE (both epichlorohydrin (0.25%) and 1,2-epoxybutane (0.25%))	0/50	0/50
1 d/wk, 89-wk exposure, Swiss rats (Van Duuren et al., 1979)			
Males (0.5mg)	Control	0/30	0/30
	TCE (unknown)	0/30	0/30
Females (0.5mg)	Control	0/30	0/30
	TCE(unknown)	0/30	0/30

4  
5  
6  
7  
8  
9

<sup>a</sup> Henschler et al. (1984). Due to poor condition of the animals resulting from the nonspecific toxicity of high doses of TCE and/or the additives, gavage was stopped for all groups during wk 35–40, 65, and 69–78, and all doses were reduced by a factor of 2 from the 40<sup>th</sup> wk on.

<sup>b</sup> Van Duuren et al. (1979) observed no renal tumors in control or exposed Swiss mice.

1 kidney tumors in F344 rats is very low,<sup>5</sup> lending biological significance to their occurrence in  
2 this study, despite the study's limitations. Cytomegaly and karyomegaly were also increased,  
3 particularly in male rats. The toxic nephropathy (specific location in kidney not stated) observed  
4 in both rats and mice and contributed to the poor survival rate (see Table 4-45). As discussed  
5 previously, this toxic nephropathy was clearly distinguishable from the spontaneous chronic  
6 progression nephropathy commonly observed in aged rats.

7

#### 4.4.5.1.3. Conclusions: Kidney Cancer in Laboratory Animals

8 Chronic TCE carcinogenicity bioassays have shown evidence of neoplastic lesions in the  
9 kidney in rats (mainly in males, with less evidence in females), treated via inhalation and gavage.  
10 As discussed above, individual studies have a number of limitations and have shown limited  
11 increases in kidney tumors. However, given the rarity of these tumors as assessed by historical  
12 controls and the repeatability of this result, these are considered biologically significant.

13

#### 4.4.6. Role of Metabolism in Trichloroethylene (TCE) Kidney Toxicity

14 It is generally thought that one or more TCE metabolites rather than the parent compound  
15 are the active moieties for TCE nephrotoxicity. As reviewed in Section 3.3, oxidation by CYPs,  
16 of which CYP2E1 is thought to be the most active isoform, results in the production of chloral  
17 hydrate, trichloroacetic acid, dichloroacetic acid and trichloroethanol. The glutathione  
18 conjugation pathway produces metabolites such as DCVG, DCVC, dichlorovinylthiol, and  
19 NAcDCVC, although, as discussed in Section 3.3.3.2, the quantitative estimates of the amount  
20 systemically produced following TCE exposure remains uncertain. Because several of the steps  
21 for generating these reactive metabolites occur in the kidney, the GSH conjugation pathway has  
22 been thought to be responsible for producing the active moiety or moieties of TCE  
23 nephrotoxicity. A comparison of TCE's nephrotoxic effects with the effects of TCE metabolites,  
24 both in vivo and in vitro, thus, provides a basis for assessing the relative roles of different  
25 metabolites. While most of the available data have been on metabolites from GSH conjugation,  
26 such as DCVC, limited information is also available on the major oxidative metabolites TCOH  
27 and TCA.

---

<sup>5</sup> NTP (1990) reported a historical control incidence of 0.4% in males. The NTP web site reports historical control rates of renal carcinomas for rats dosed via corn oil gavage on the NIH-07 diet (used before 1995, when the TCE studies were conducted) to be 0.5% (2/400) for males and 0% (0/400) for females ([http://ntp-server.niehs.nih.gov/ntp/research/database\\_searches/historical\\_controls/path/r\\_gavco.txt](http://ntp-server.niehs.nih.gov/ntp/research/database_searches/historical_controls/path/r_gavco.txt)). In addition, the two occurrences in males came from the same study, with all other studies reporting 0/50 carcinomas.

*This document is a draft for review purposes only and does not constitute Agency policy.*



1

#### 4.4.6.1.1. In Vivo Studies of the Kidney Toxicity of Trichloroethylene (TCE) Metabolites

2           Studies of kidney toxicity of TCE metabolites discussed in this section are shown in  
3 Table 4-52.

4

#### 4.4.6.1.2. Role of GSH conjugation metabolites of trichloroethylene (TCE)

5           In numerous studies, DCVC has been shown to be acutely nephrotoxic in rats and mice.  
6 Mice receiving a single dose of 1 mg/kg DCVC (the lowest dose tested in this species) exhibited  
7 karyolytic proximal tubular cells in the outer stripe of the outer medulla, with some sloughing of  
8 cells into the lumen and moderate desquamation of the tubular epithelium (Eyre et al., 1995b).  
9 Higher doses in mice were associated with more severe histological changes similar to those  
10 induced by TCE, such as desquamation and necrosis of the tubular epithelium (Darnerud et al.,  
11 1989; TERRACINI and PARKER, 1965; Vaidya et al., 2003a, b). In rats, no histological  
12 changes in the kidney were reported after single doses of 1, 5, and 10 mg/kg DCVC (Eyre et al.,  
13 1995a; Green et al., 1997a), but cellular debris in the tubular lumen was reported at 25 mg/kg  
14 (Eyre et al., 1995b) and slight degeneration and necrosis were seen at 50 mg/kg (Green et al.,  
15 1997a). Green et al. (1997a) reported no histological changes were noted in rats after 10 doses  
16 of 0.1–5.0 mg/kg DCVC (although increases in urinary protein and GGT were found), but some  
17 karyomegaly was noted in mice after 10 daily doses of 1 mg/kg. Therefore, mice appear more  
18 sensitive than rats to the nephrotoxic effects of acute exposure to DCVC, although the number of  
19 animals used at each dose in these studies was limited (10 or less). Although the data are not  
20 sufficient to assess the relative sensitivity of other species, it is clear that multiple species,  
21 including rabbits, guinea pigs, cats, and dogs, are responsive to DCVC’s acute nephrotoxic  
22 effects (Jaffe et al., 1984; Krejci et al., 1991; TERRACINI and PARKER, 1965; Wolfgang et al.,  
23 1989b).

24           Very few studies are available at longer durations. Terracini and Parker (1965) gave  
25 DCVC in drinking water to rats at a concentration of 0.01% for 12 weeks (approximately  
26 10 mg/kg-day), and reported consistent pathological and histological changes in the kidney. The  
27 progression of these effects was as follows: (1) during the first few days, completely necrotic  
28 tubules, with isolated pyknotic cells being shed into the lumen; (2) after 1 week, dilated tubules  
29 in the inner part of the cortex, lined with flat epithelial cells that showed thick basal membranes,  
30 some with big hyperchromatic nuclei; (3) in the following weeks, increased prominence of  
31 tubular cells exhibiting karyomegaly, seen in almost all animals, less pronounced tubular

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 dilation, and cytomegaly in the same cells showing karyomegaly. In addition, increased mitotic  
2 activity was reported the first few days, but was not evident for the rest of the experiment.  
3 Terracini and Parker (1965) also reported the results of a small experiment (13 male and

**Table 4-52. Laboratory animal studies of kidney noncancer toxicity of TCE metabolites**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Kidney effect(s) discussed in Section 4.4.4
Dow and Green (2000)	Fischer 344 rats (M)	Drinking water	0, 0.5, 1 g/L Trichloroethanol, 12 wk	3/group	Increased formic acid in urine.
Jaffe et al. (1984)	Swiss-Webster mice (M)	Drinking water	0–22 mg/kg-day DCVC, 37 wk	5/group	Cytomegaly and tubular degeneration.
Mather et al. (1990)	Sprague-Dawley rats (M)	Drinking water	0–355 mg/kg-day TCA, 90 d	10/group	Increased kidney weight.
Terracini and Parker (1965)	Wistar rats (Gender not specified) Grey mice (Gender not specified)	Drinking water	0, 0.01% DCVC, 12 wk	35/group	Necrosis of tubular epithelium in mice and rats.

1 5 female rats) given the same concentration of DCVC in drinking water for 46 weeks, and  
2 observed for 87 weeks. They noted renal tubular cells exhibiting karyomegaly and cytomegaly  
3 consistently throughout the experiment. Moreover, a further group of eight female rats given  
4 DCVC in drinking water at a concentration of 0.001% (approximately 1 mg/kg-day) also  
5 exhibited similar, though less severe, changes in the renal tubules. In mice, Jaffe et al. (1984)  
6 gave DCVC in drinking water at concentrations of 0.001, 0.005, and 0.01% (estimated daily dose  
7 of 1–2, 7–13, and 17–22 mg/kg-day), and reported similar effects in all dose groups, including  
8 cytomegaly, nuclear hyperchromatism, and multiple nucleoli, particularly in the pars recta  
9 section of the kidney. Thus, effects were noted in both mice and rats under chronic exposures at  
10 doses as low as 1–2 mg/kg-day (the lowest dose tested). Therefore, while limited, the available  
11 data do not suggest differences between mice and rats to the nephrotoxic effects of DCVC under  
12 chronic exposure conditions, in contrast to the greater sensitivity of mice to acute and subchronic  
13 DCVC-induced nephrotoxicity.

14 Importantly, as summarized in Table 4-53, the histological changes and their location in  
15 these subchronic and chronic experiments with DCVC are quite similar to those reported in  
16 chronic studies of TCE, described above, particularly the prominence of karyomegaly and  
17 cytomegaly in the pars recta section of the kidney. Moreover, the morphological changes in the  
18 tubular cells, such as flattening and dilation, are quite similar. Similar pathology is not observed  
19 with the oxidative metabolites alone (see Section 4.4.6.1.2).

20 Additionally, it is important to consider whether sufficient DCVC may be formed from  
21 TCE exposure to account for TCE nephrotoxicity. While direct pharmacokinetic measurements,  
22 such as the excretion of NAcDCVC, have been used to argue that insufficient DCVC would be  
23 formed to be the active moiety for nephrotoxicity (Green et al., 1997a), as discussed in Section 3,  
24 urinary NAcDCVC is a poor marker of the flux through the GSH conjugation pathway because  
25 of the many other possible fates of metabolites in that pathway. In another approach, Eyre et al.  
26 (1995b) using acid-labile adducts as a common internal dosimeter between TCE and DCVC, and  
27 reported that a single TCE dose of 400 mg/kg in rats (similar to the lowest daily doses in the NCI  
28 and NTP rat bioassays) and 1,000 mg/kg (similar to the lowest daily doses in the NCI and NTP  
29 mouse bioassays) corresponded to a single equivalent DCVC dose of 6 and 1 mg/kg-day in rats  
30 and mice, respectively. These equivalent doses of DCVC are greater or equal to those in which  
31 nephrotoxicity has been reported in these species under chronic conditions. Therefore, assuming  
32 that this dose correspondence is accurate under chronic conditions, sufficient DCVC would be  
33 formed from TCE exposure to explain the observed histological changes in the renal tubules.  
34 Nevertheless, direct estimates of how much DCVC is formed after TCE exposure are lacking.

**Table 4-53. Summary of histological changes in renal proximal tubular cells induced by chronic exposure to TCE, DCVC, and TCOH**

Effects	TCE	DCVC	TCOH
Karyomegaly	Enlarged, hyperchromatic nuclei, irregular to oblong in shape. Vesicular nuclei containing prominent nucleoli.	Enlarged, hyperchromatic nuclei with and multiple nucleoli. Nuclear pyknosis and karyorrhexis.	None reported.
Cytomegaly	Epithelial cells were large, elongated and flattened.	Epithelial cells were large, elongated and flattened cells.	No report of enlarged cells.
Cell necrosis/hyperplasia	Stratified epithelium that partially or completely filled the tubular lumens. Cells in mitosis were variable in number or absent. Cells had abundant eosinophilic or basophilic cytoplasm.	Thinning of tubular epithelium, frank tubular necrosis, re-epitheliation. Tubular atrophy, interstitial fibrosis and destruction of renal parenchyma. More basophilic and finely vacuolated.	No flattening or loss of epithelium reported. Increased tubular cell basophilia, followed by increased cellular eosinophilia, tubular cell vacuolation.
Morphology / content of tubules	Some tubules enlarged/dilated to the extent that they were difficult to identify. Portions of basement membrane had a stripped appearance. Tubules were empty or contained “wisps of eosinophilic material.”	Tubular dilation, denuded tubules. Thick basal membrane. Focal areas of dysplasia, intraluminal casts.	No tubular dilation reported. Intratubular cast formation.

Sources: NCI (1976); NTP (1988, 1990); Maltoni et al. (1988); Terracini and Parker (1965); Jaffe et al. (1985); Green et al. (2003).

1           The Eker rat model (*Tsc-2*<sup>±</sup>) is at increased risk for the development of spontaneous renal  
2 cell carcinoma and as such has been used to understand the mechanisms of renal carcinogenesis  
3 (Stemmer et al., 2007; Wolf et al., 2000). One study has demonstrated similar pathway  
4 activation in Eker rats as that seen in humans with *VHL* mutations leading to renal cell  
5 carcinoma, suggesting *Tsc-2* inactivation is analogous to inactivation of *VHL* in human renal cell  
6 carcinoma (Liu et al., 2003). Although the Eker rat model is a useful tool for analyzing  
7 progression of renal carcinogenesis, it has some limitations in analysis of specific genetic  
8 changes, particularly given the potential for different genetic changes depending on type of  
9 exposure and tumor. The results of short-term assays to genotoxic carcinogens in the Eker rat  
10 model (Morton et al., 2002; Stemmer et al., 2007) reported limited preneoplastic and neoplastic  
11 lesions which may be related to the increased background rate of renal carcinomas in this animal  
12 model.

13           Recently, Mally et al. (2006) exposed male rats carrying the Eker mutation to TCE  
14 (0–1,000 mg/kg BW) by corn oil gavage and demonstrated no increase in renal preneoplastic  
15 lesions or tumors. Primary Eker rat kidney cells exposed to DCVC in this study did induce an  
16 increase in transformants in vitro but no DCVC-induced *vhl* or *Tsc-2* mutations were observed.  
17 In vivo exposure to TCE (5 days/week for 13 weeks), decreased body weight gain and increased  
18 urinary excretion at the two highest TCE concentrations analyzed (500 and 1,000 mg/kg BW)  
19 but did not change standard nephrotoxicity markers (GGT, creatinine and urinary protein).  
20 Renal tubular epithelial cellular proliferation as measured by BrdU incorporation was  
21 demonstrated at the three highest concentrations of TCE (250, 500 and 1,000 mg/kg-day). A  
22 minority of these cells also showed karyomegaly at the two higher TCE concentrations.  
23 Although renal cortical tumors were demonstrated in all TCE exposed groups, these were not  
24 significantly different from controls (13 weeks). These studies were complemented with in vitro  
25 studies of DCVC (10–50 µM) in rat kidney epithelial (RKE) cells examining proliferation at 8,  
26 24, and 72 hours and cellular transformation at 6–7 weeks. Treatment of RKE cells from  
27 susceptible rats with DCVC gave rise to morphologically transformed colonies consistently  
28 higher than background (Mally et al., 2006). Analyzing ten of the renal tumors from the TCE  
29 exposed rats and nine of the DCVC transformants from these studies for alterations to the *VHL*  
30 gene that might lead to inactivation found no alterations to *VHL* gene expression or mutations.

31           One paper has linked the *VHL* gene to chemical-induced carcinogenesis. Shiao et al.  
32 (1998) demonstrated *VHL* gene somatic mutations in *N*-nitrosodimethylamine-induced rat kidney  
33 cancers that were of the clear cell type. The clear cell phenotype is rare in rat kidney cancers,  
34 but it was only the clear cell cancers that showed *VHL* somatic mutation (three of eight tumors  
35 analyzed). This provided an additional link between *VHL* inactivation and clear cell kidney

1 cancer. However, this study examined archived formalin fixed paraffin embedded tissues from  
2 previous experiments. As described previously (see Section 4.4.2), DNA extraction from this  
3 type of preparation creates some technical issues. Similarly, archived formalin-fixed paraffin  
4 embedded tissues from rats exposed to potassium bromide were analyzed in a later study by  
5 Shiao et al. (2002). This later study examined the *VHL* gene mutations following exposure to  
6 potassium bromide, a rat renal carcinogen known to induce clear cell renal tumors. Clear cell  
7 renal tumors are the most common form of human renal epithelial neoplasms, but are extremely  
8 rare in animals. Although F344 rats exposed to potassium bromide in this study did develop  
9 renal clear cell carcinomas, only two of nine carried the same C to T mutation at the core region  
10 of the Sp1 transcription-factor binding motif in the *VHL* promoter region, and one of  
11 four untreated animals had a C to T mutation outside the conserved core region. Mutation in the  
12 *VHL* coding region was only detected in one tumor, so although the tumors developed following  
13 exposure to potassium bromide were morphologically similar to those found in humans; no  
14 similarities were found in the genetic changes.

15 Elfarra et al. (1984) found that both DCVG and DCVC administered to male F344 rats by  
16 intraperitoneal injections in isotonic saline resulted in elevations in BUN and urinary glucose  
17 excretion. Furthermore, inhibition of renal GGT activity with acivicin protected rats from  
18 DCVG-induced nephrotoxicity. In addition, both the  $\beta$ -lyase inhibitor AOAA and the renal  
19 organic anion transport inhibitor probenecid provided protection from DCVC, demonstrating a  
20 requirement for metabolism of DCVG to the cysteine conjugate by the action of renal GGT and  
21 dipeptidase, uptake into the renal cell by the organic anion transporter, and subsequent activation  
22 by the  $\beta$ -lyase. This conclusion was supported further by showing that the methyl analog of  
23 DCVC, which cannot undergo a  $\beta$ -elimination reaction due to the presence of the methyl group,  
24 was not nephrotoxic.

25 Korrapati et al. (2005) builds upon a series of investigations of hetero- (by mercuric  
26 chloride [ $\text{HgCl}_2$ ]) and homo-(by DCVC, 15 mg/kg) protection against a lethal dose of DCVC  
27 (75 mg/kg). Priming, or preconditioning, with pre-exposure to either  $\text{HgCl}_2$  or DCVC of male  
28 Swiss-Webster mice was said to augment and sustain cell division and tissue repair, hence  
29 protecting against the subsequent lethal DCVC dose (Vaidya et al., 2003a, b; 2003c). Korrapati  
30 et al. (2005) showed that a lethal dose of DCVC downregulates phosphorylation of endogenous  
31 retinoblastoma protein (pRb), which is considered critical in renal proximal tubular and  
32 mesangial cells for the passage of cells from G1 to S-phase, thereby leading to a block of renal  
33 tubule repair. Priming, in contrast, upregulated P-pRB which was sustained even after the  
34 administration of a lethal dose of DCVC, thereby stimulating S-phase DNA synthesis, which was  
35 concluded to result in tissue repair and recovery from acute renal failure and death. These

1 studies are more informative about the mechanism of autoprotection than on the mechanism of  
2 initial injury caused by DCVC. In addition, the priming injury (not innocuous, as it caused  
3 25–50% necrosis and elevated blood urea nitrogen) may have influenced the toxicokinetics of  
4 the second DCVC injection.  
5

#### 4.4.6.1.3. Role of oxidative metabolites of Trichloroethylene (TCE)

6 Some investigators (Dow and Green, 2000; Green et al., 2003; Green et al., 1998) have  
7 proposed that TCE nephrotoxicity is related to formic acid formation. They demonstrated that  
8 exposure to either trichloroethanol or trichloroacetic acid causes increased formation and urinary  
9 excretion of formic acid (Green et al., 1998). The formic acid does not come from  
10 trichloroethylene. Rather, trichloroethylene (or a metabolite) has been proposed to cause a  
11 functional depletion of vitamin B<sub>12</sub>, which is required for the methionine salvage pathway of  
12 folate metabolism. Vitamin B<sub>12</sub> depletion results in folate depletion. Folate is a cofactor in  
13 one-carbon metabolism and depletion of folate allows formic acid to accumulate, and then to be  
14 excreted in the urine (Dow and Green, 2000).

15 TCE (1 and 5 g/L), TCA (0.25, 0.5 and 1 g/L) and TCOH (0.5 and 1.0 g/L) exposure in  
16 male Fischer rats substantially increased excretion of formic acid in urine, an effect suggested as  
17 a possible explanation for TCE-induced renal toxicity in rats (Green et al., 1998). Green et al.  
18 (2003) reported tubular toxicity as a result of chronic (1 year) exposure to TCOH (0, 0.5, and  
19 1.0 g/L). Although TCOH causes tubular degeneration in a similar region of the kidney as TCE,  
20 there are several dissimilarities between the characteristics of nephrotoxicity between the  
21 two compounds, as summarized in Table 4-53. In particular, Green et al. (1998) did not observe  
22 TCOH causing karyomegaly and cytomegaly. These effects were seen as early as 13 weeks after  
23 the commencement of TCE exposure (NTP, 1990), with 300 ppm inhalation exposures to TCE  
24 (Maltoni et al., 1988), as well as at very low chronic exposures to DCVC (Jaffe et al., 1984;  
25 TERRACINI and PARKER, 1965). In addition, Green et al. (2003) reported neither flattening  
26 nor loss of the tubular epithelium nor hyperplasia, but suggested that the increased early  
27 basophilia was due to newly divided cells, and therefore, represented tubular regeneration in  
28 response to damage. Furthermore, they noted that such changes were seen with the spontaneous  
29 damage that occurs in aging rats. However, several of the chronic studies of TCE noted that the  
30 TCE-induced damage observed was distinct from the spontaneous nephropathy observed in rats.  
31 A recent in vitro study of rat hepatocytes and primary human renal proximal tubule cells from  
32 two donors measured formic acid production following exposure to CH (0.3–3 mM, 3–10 days)  
33 (Lock et al., 2007). This study observed increased formic acid production at day 10 in both



1 human renal proximal tubule cell strains, but a similar level of formic acid was measured when  
2 CH was added to media alone. The results of this study are limited by the use of only two  
3 primary human cell strains, but suggest exposure to CH does not lead to significant increases in  
4 formic acid production in vivo.

5 Interestingly, it appears that the amount of formic acid excreted reaches a plateau at a  
6 relatively low dose. Green et al. (2003) added folic acid to the drinking water of the group of  
7 rats receiving the lower dose of TCOH (18.3 mg/kg-day) in order to modulate the excretion of  
8 formic acid in that dose group, and retain the dose-response in formic acid excretion relative to  
9 the higher-dose group (54.3 mg/kg-day). These doses of TCOH are much lower than what  
10 would be expected to be formed in vivo at chronic gavage doses. For instance, after a single  
11 500-mg/kg dose of TCE (the lower daily dose in the NTP rat chronic bioassays), Green and  
12 Prout (1985) reported excretion of about 41% of the TCE gavage dose in urine as TCOH or  
13 trichloroethanol-glucuronide conjugate (TCOG) in 24 hours. Thus, using the measure of  
14 additional excretion after 24 hours and the TCOH converted to TCA as a lower bound as to the  
15 amount of TCOH formed by a single 500 mg/kg dose of TCE, the amount of TCOH would be  
16 about 205 mg/kg, almost fourfold greater than the high dose in the Green et al. (2003) study. By  
17 contrast, these TCOH doses are somewhat smaller than those expected from the inhalation  
18 exposures of TCE. For instance, after 6 hour exposure to 100 and 500 ppm TCE (similar to the  
19 daily inhalation exposures in Maltoni et al. (1988)), male rats excreted 1.5 and 4.4 mg of TCOH  
20 over 48 hours, corresponding to 5 and 15 mg/kg for a rat weighing 0.3 kg (Kaneko et al., 1994a).  
21 The higher equivalent TCOH dose is similar to the lower TCOH dose used in Green et al.  
22 (2003), so it is notable that while Maltoni et al. (1988) reported a substantial incidence of  
23 cytomegaly and karyomegaly after TCE exposure (300 and 600 ppm), none was reported in  
24 Green et al. (2003).

25 TCOH alone does not appear sufficient to explain the range of renal effects observed  
26 after TCE exposure, particularly cytomegaly, karyomegaly, and flattening and dilation of the  
27 tubular epithelium. However, given the studies described above, it is reasonable to conclude that  
28 TCOH may contribute to the nephrotoxicity of TCE, possibly due to excess formic acid  
29 production, because (1) there are some similarities between the effects observed with TCE and  
30 TCOH and (2) the dose at which effects with TCOH are observed overlap with the approximate  
31 equivalent TCOH dose from TCE exposure in the chronic studies.

32 Dow and Green (2000) noted that TCA also induced formic acid accumulation in rats,  
33 and suggested that TCA may therefore, contribute to TCE-induced nephrotoxicity. However,  
34 TCA has not been reported to cause any similar histologic changes in the kidney. Mather et al.  
35 (1990) reported an increase of kidney-weight to body-weight ratio in rats after 90 days of

1 exposure to trichloroacetic acid in drinking water at 5,000 ppm (5 g/L) but reported no  
2 histopathologic changes in the kidney. DeAngelo et al. (1997) reported no effects of  
3 trichloroacetic acid on kidney weight or histopathology in rats in a 2-year cancer bioassay.  
4 Dow and Green (2000) administered TCA at quite high doses (1 and 5 g/L in drinking water),  
5 greater than the subsequent experiments of Green et al. (2003) with TCOH (0.5 and 1 g/L in  
6 drinking water), and reported similar amounts of formic acid produced (about 20 mg/day for  
7 each compound). However, cytotoxicity or karyomegaly did not appear to be analyzed.  
8 Furthermore, much more TCOH is formed from TCE exposure than TCA. Therefore, if TCA  
9 contributes substantially to the nephrotoxicity of TCE, its contribution would be substantially  
10 less than that of TCOH. Lock et al. (2007) also measured formic acid production in human renal  
11 proximal tubule cells exposed to 0.3–3 mM CH for 10 days CH. This study measured  
12 metabolism of CH to TCOH and TCA as well as formic acid production and subsequent  
13 cytotoxicity. Increased formic acid was not observed in this study, and limited cytotoxicity was  
14 observed. However, this study was performed in human renal proximal tubular cells from only  
15 two donors, and there is potential for large interindividual variability in response, particularly  
16 with CYP enzymes.

17 In order to determine the ability of various chlorinated hydrocarbons to induce  
18 peroxisomal enzymes, Goldsworthy and Popp (1987) exposed male Fischer 344 rats and male  
19 B6C3F1 mice to TCE (1,000 mg/kg BW) and TCA (500 mg/kg BW) by corn oil gavage for  
20 10 consecutive days. Peroxisomal activation was measured by palmitoyl coenzyme A (CoA)  
21 oxidase activity levels. TCE led to increased peroxisomal activation in the kidneys of both rats  
22 (300% of control) and mice (625% of control), while TCA led to an increase only in mice (280%  
23 of control). A study by Zanelli et al. (1996) exposed Sprague-Dawley rats to TCA for 4 days and  
24 measured both renal and hepatic peroxisomal and cytochrome P450 enzyme activities.  
25 TCA-treated rats had increased activity in CYP 4A subfamily enzymes and peroxisomal  
26 palmitoyl-CoA oxidase. Both of these acute studies focused on enzyme activities and did not  
27 further analyze resulting histopathology.

28

#### **4.4.6.1.4. In Vitro Studies of Kidney Toxicity of Trichloroethylene (TCE) and Metabolites**

29 Generally, it is believed that TCE metabolites are responsible for the bulk of kidney  
30 toxicity observed following exposure. In particular, studies have demonstrated a role for DCVG  
31 and DCVC in kidney toxicity, though, as discussed in Section 3.3.3.2, the precise metabolic yield  
32 of these metabolites following TCE exposure remains uncertain. The work by Lash and  
33 colleagues (Cummings and Lash, 2000; Cummings et al., 2000b; Cummings et al., 2000c; Lash

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 et al., 2000b) examined the effect of trichloroethylene and its metabolites in vitro.  
2 Trichloroethylene and DCVC are toxic to primary cultures of rat proximal and distal tubular cells  
3 (Cummings et al., 2000a) while the TCE metabolites DCVG and DCVC have been demonstrated  
4 to be cytotoxic to rat and rabbit kidney cells in vitro (Groves et al., 1993; Hassall et al., 1983;  
5 Wolfgang et al., 1989a) Lash et al., 2000b, 2001b. Glutathione-related enzyme activities were  
6 well maintained in the cells, whereas CYP activities were not. The enzyme activity response to  
7 DCVC was greater than the response to trichloroethylene; however, the proximal and distal  
8 tubule cells had similar responses even though the proximal tubule is the target in vivo. The  
9 authors attributed this to the fact that the proximal tubule is exposed before the distal tubule in  
10 vivo and to possible differences in uptake transporters. They did not address the extent to which  
11 transporters were maintained in the cultured cells.

12 In further studies, Lash et al. (2001b) assessed the toxicity of trichloroethylene and its  
13 metabolites DCVC and DCVG using in vitro techniques (Lash et al., 2001b) as compared to in  
14 vivo studies. Experiments using isolated cells were performed only with tissues from  
15 Fischer 344 rats, and lactate dehydrogenase release was used as the measure of cellular toxicity.  
16 The effects were greater in males. DCVC and trichloroethylene had similar effects, but DCVG  
17 exhibited increased efficacy compared with trichloroethylene and DCVC.

18 In vitro mitochondrial toxicity was assessed in renal cells from both Fischer 344 rats and  
19 B6C3F1 mice following exposure to both DCVC and DCVG (Lash et al., 2001b). Renal  
20 mitochondria from male rats and mice responded similarly; a greater effect was seen in cells  
21 from the female mice. These studies show DCVC to be slightly more toxic than  
22 trichloroethylene and DCVG, but species differences are not consistent with the effects observed  
23 in long-term bioassays. This suggests that in vitro data be used with caution in risk assessment,  
24 being mindful that in vitro experiments do not account for in vivo pharmacokinetic and  
25 metabolic processes.

26 In LLC-PK1 cells, DCVC causes loss of mitochondrial membrane potential,  
27 mitochondrial swelling, release of cytochrome c, caspase activation, and apoptosis (Chen et al.,  
28 2001). Thus, DCVC is toxic to mitochondria, resulting in either apoptosis or necrosis.  
29 DCVC-induced apoptosis also has been reported in primary cultures of human proximal tubule  
30 cells (Lash et al., 2001a).

31 DCVC was further studied in human renal proximal tubule cells for alterations in gene  
32 expression patterns related to proposed modes of action in nephrotoxicity (Lock et al., 2006). In  
33 cells exposed to subtoxic levels of DCVC to better mimic workplace exposures, the expression  
34 of genes involved with apoptosis (caspase 8, FADD-like regulator) was increased at the higher  
35 dose (1  $\mu$ M) but not at the lower dose (0.1  $\mu$ M) of DCVC exposure. Genes related to oxidative

1 stress response (SOD, NF-κB, p53, c-Jun) were altered at both subtoxic doses, with genes  
2 generally upregulated at 0.1 μM DCVC being downregulated at 1 μM DCVC. The results of this  
3 study support the need for further study, and highlight the involvement of multiple pathways and  
4 variability of response based on different concentrations.

5 Lash et al. (2007) examined the effect of modulation of renal metabolism on toxicity of  
6 TCE in isolated rat cells and microsomes from kidney and liver. Following exposure to  
7 modulating chemicals, lactate dehydrogenase (LDH) was measured as a marker of cytotoxicity,  
8 and the presence of specific metabolites was documented (DCVG, TCA, TCOH, and CH).  
9 Inhibition of the CYP stimulated an increase of GSH conjugation of TCE and increased  
10 cytotoxicity in kidney cells. This modulation of CYP had a greater effect on TCE-induced  
11 cytotoxicity in liver cells than in kidney cells. Increases in GSH concentrations in the kidney  
12 cells led to increased cytotoxicity following exposure to TCE. Depletion of GSH in hepatocytes  
13 exposed to TCE, however, led to an increase in hepatic cytotoxicity. The results of this study  
14 highlight the role of different bioactivation pathways needed in both the kidney and the liver,  
15 with the kidney effects being more affected by the GSH conjugation pathways metabolic  
16 products.

17 In addition to the higher susceptibility of male rats to TCE-induced  
18 nephrocarcinogenicity and nephrotoxicity, isolated renal cortical cells from male F344 rats are  
19 more susceptible to acute cytotoxicity from TCE than cells from female rats. TCE caused a  
20 modest increase in LDH release from male rat kidney cells but had no significant effect on LDH  
21 release from female rat kidney cells. These results on male susceptibility to TCE agree with the  
22 in vivo data.

#### 23 **4.4.6.1.5. Conclusions as to the Active Agents of Trichloroethylene (TCE)-Induced Nephrotoxicity**

24 In summary, the TCE metabolites DCVC, TCOH, and TCA have all been proposed as  
25 possible contributors to the nephrotoxicity of TCE. Both in vivo and in vitro data strongly  
26 support the conclusion that DCVC and related GSH conjugation metabolites are the active agents  
27 of TCE-induced nephrotoxicity. Of these, DCVC induces effects in renal tissues, both in vivo  
28 and in vitro, that are most similar to those of TCE, and formed in sufficient amounts after TCE  
29 exposure to account for those effects. A role for formic acid due to TCOH or TCA formation  
30 from TCE cannot be ruled out, as it is known that substantial TCOH and TCA are formed from  
31 TCE exposure, that formic acid is produced from all three compounds, and that TCOH exposure  
32 leads to toxicity in the renal tubules. However, the characteristics of TCOH-induced

1 nephrotoxicity do not account for the range of effects observed after TCE exposure while those  
2 of DCVC-induced nephrotoxicity do. Also, TCOH does not induce the same pathology as TCE  
3 or DCVC. TCA has also been demonstrated to induce peroxisomal proliferation in the kidney  
4 (Goldsworthy and Popp, 1987), but this has not been associated with kidney cancer. Therefore,  
5 although TCOH and possibly TCA may contribute to TCE-induced nephrotoxicity, their  
6 contribution is likely to be small compared to that of DCVC. However, as discussed in  
7 Section 3.3.3.2, the precise metabolic yield of these DCVC following TCE exposure remains  
8 uncertain.

#### 4.4.7. Mode(s) of Action for Kidney Carcinogenicity

10 This section will discuss the evidentiary support for several hypothesized modes of action  
11 for kidney carcinogenicity, including mutagenicity, cytotoxicity and regenerative proliferation,  
12 peroxisome proliferation,  $\alpha$ 2 $\mu$ -related nephropathy and formic acid-related nephropathy,  
13 following the framework outlined in the *Cancer Guidelines* (U.S. EPA, 2005c, d).<sup>6</sup> The data and  
14 conclusions for the MOAs with the greatest experimental support are summarized in Table 4-54.

##### 4.4.7.1.1. Hypothesized Mode of Action: Mutagenicity

16 One hypothesis is that a mutagenic mode of action is operative in TCE-induced renal  
17 carcinogenesis. According to this hypothesis, the key events leading to TCE-induced kidney  
18 tumor formation constitute the following: TCE GSH conjugation metabolites (e.g., DCVG,  
19 DCVC, NAcDCVC, and/or other reactive metabolites derived from subsequent beta-lyase, flavin  
20 monooxygenases [FMO], or CYP metabolism) derived from the GSH-conjugation pathway, after  
21 being either produced in situ in or delivered systemically to the kidney, cause direct alterations to  
22 DNA (e.g., mutation, DNA damage, and/or micronuclei induction). Mutagenicity is a  
23 well-established cause of carcinogenicity.

---

<sup>6</sup> As recently reviewed ([Guyton et al., 2008](#)) the approach to evaluating mode of action information described in EPA's *Cancer Guidelines* ([2005c, d](#)) considers the issue of human relevance of a hypothesized mode of action in the context of hazard evaluation. This excludes, for example, consideration of toxicokinetic differences across species; specifically, the *Cancer Guidelines* state, "the toxicokinetic processes that lead to formation or distribution of the active agent to the target tissue are considered in estimating dose but are not part of the mode of action." In addition, information suggesting quantitative differences in the occurrence of a key event between test species and humans are noted for consideration in the dose-response assessment, but is not considered in human relevance determination. In keeping with these principles, a formal analysis of the dose-response of key events in the hypothesized modes of action is not presented unless it would aid in the overall weight of evidence analysis for carcinogenicity, as presented in Section 4.11.



**Table 4-54. Summary of major mode of action conclusions for TCE kidney carcinogenesis**

Hypothesized MOA <i>postulated key events</i>	Experimental support	Human relevance	Weight-of-evidence conclusion
<b>Mutagenicity</b>			
<i>GSH conjugation-derived metabolites produced in situ or delivered systemically to kidney</i>	Multiple in vitro and in vivo studies demonstrate GSH conjugation of TCE, and availability to the kidney (see Section 3.3.3). Uncertainties are quantitative (precise amount of flux), not qualitative.	Yes: demonstrated in humans in vivo and in human cells in vitro.	Highly likely that both human and rodent kidneys are exposed to the GSH-conjugation derived metabolites.
<i>GSH conjugation-derived metabolites cause direct alterations to DNA, resulting in mutations</i>	GSH conjugation derived metabolites (DCVG, DCVC, NAcDCVC) demonstrated to be genotoxic in most in vitro assays in which they have been tested, including Ames test (see Section 4.2.1.4.1). Kidney-specific genotoxicity in rats and rabbits after in vivo administration of TCE or DCVC. Not seen in mice, but may be due to species differences in metabolism and in sensitivity to renal carcinogenesis.	Yes: no basis for discounting in vitro genotoxicity results. Inconsistent results with respect to VHL mutation and TCE-induced kidney tumors.	Predominance of positive genotoxicity data consistent with GSH-conjugation derived metabolites causing mutations in the kidney.
<i>Mutations cause cancer</i>	Well established.	Yes: well established.	Mutagenicity is a well-established cause of cancer.
<b>Cytotoxicity and regenerative proliferation</b>			
<i>GSH conjugation-derived metabolites produced in situ or delivered systemically to kidney</i>	Multiple in vitro and in vivo studies demonstrate GSH conjugation of TCE, and availability to the kidney (see Section 3.3.3). Uncertainties are quantitative (precise amount of flux), not qualitative.	Yes: demonstrated in humans in vivo and in human cells in vitro.	Highly likely that both human and rodent kidneys are exposed to the GSH-conjugation derived metabolites.

**Table 4-54. Summary of major mode of action conclusions for TCE kidney carcinogenesis (continued)**

Hypothesized MOA <i>postulated key events</i>	Experimental support	Human relevance	Weight-of-evidence conclusion
<b>Mutagenicity</b>			
<i>GSH conjugation-derived metabolites cause cell death (cytotoxicity) in the kidney.</i>	Multiple human and laboratory animal studies demonstrating TCE to be nephrotoxic, including chronic studies (see Sections 4.4.1 and 4.4.4). Multiple laboratory animal studies and in vitro studies in rat and human kidney cells demonstrating DCVC to be nephrotoxic (see Sections 4.4.6.1.1 and 4.4.6.2). Some evidence that TCOH is nephrotoxic, but histological changes caused by TCE more similar to those caused by DCVC (see Section 4.4.6.1 and 4.4.6.3).	Yes: demonstrated human nephrotoxicity of TCE in vivo and DCVC in vitro.	TCE is nephrotoxic in humans, and DCVC is likely the predominant moiety responsible.
<i>Cell proliferation increases in the kidney to repair damage.</i>	No data specific to TCE or GSH conjugation-derived metabolites.	Yes	
<i>Increased cell turnover increases the rate of mutations.</i>	No data specific to TCE or GSH conjugation-derived metabolites.	Yes	
<i>Increased proliferation cause clonal expansion of initiated (pre-malignant) cells.</i>	No increase in preneoplastic or neoplastic lesions in Eker rats exposed to TCE for 13 wk, but no data on longer durations or from other rat strains sensitive to TCE renal carcinogenesis.	Yes	
<i>Increased number of mutations and/or initiated cells causes cancer.</i>	Well established.	Yes: well established.	Increases in mutations and in initiated cells are well-established causes of cancer.



#### 4.4.7.1.2. Experimental support for the hypothesized mode of action

1 Evidence for the hypothesized mode of action for TCE includes (1) the formation of  
2 GSH-conjugation pathway metabolites in the kidney demonstrated in TCE toxicokinetics studies;  
3 and (2) the genotoxicity of these GSH-conjugation pathway metabolites demonstrated in most  
4 existing in vitro and in vivo assays of gene mutations (i.e., Ames test) and in assays of  
5 unscheduled DNA synthesis, DNA strand breaks, and micronuclei using both “standard” systems  
6 and renal cells/tissues.<sup>7</sup> Additional relevant data come from analyses of *VHL* mutations in  
7 human kidney tumors and studies using the Eker rat model. These lines of evidence are  
8 elaborated below.

9 Toxicokinetic data are consistent with these genotoxic metabolites either being delivered  
10 to or produced in the kidney. As discussed in Section 3, following in vivo exposure to TCE, the  
11 metabolites DCVG, DCVC, and NAcDCVC have all been detected in the blood, kidney, or urine  
12 of rats, and DCVG in blood and NAcDCVC in urine have been detected in humans (Bernauer et  
13 al., 1996; Birner et al., 1993; Lash et al., 1999a; 2006). In addition, in vitro data have shown  
14 DCVG formation from TCE in cellular and subcellular fractions from the liver, from which it  
15 would be delivered to the kidney via systemic circulation, and from the kidney (see  
16 Tables 3-23–3-24, and references therein). Furthermore, in vitro data in both humans and  
17 rodents support the conclusion that DCVC is primarily formed from DCVG in the kidney itself,  
18 with subsequent in situ transformation to NAcDCVC by *N*-Acetyl transferase or to reactive  
19 metabolites by beta-lyase, FMO, or CYPs (see Sections 3.3.3.2.2–3.3.3.2.5). Therefore, it is  
20 highly likely that both human and rodent kidneys are exposed to these TCE metabolites.

21 As discussed in Section 4.2.1.4.2, DCVG, DCVC, and NAcDCVC have been  
22 demonstrated to be genotoxic in most available in vitro assays.<sup>8</sup> In particular, DCVC was

---

<sup>7</sup> The EPA *Cancer Guidelines* (2005c, d) note reliance on “evaluation of in vivo or in vitro short-term testing results for genetic endpoints” and evidence that “the carcinogen or a metabolite is DNA-reactive and/or has the ability to bind to DNA” as part of this weight of evidence supporting a mutagenic mode of action. While evidence from hypothesis-testing experiments that mutation is an early step in the carcinogenic process is considered if available, it is not required for determination of a mutagenic mode of action; rather, reliance on short-term genotoxicity tests is emphasized. Thus, such tests are the focus of this analysis, which also includes an analysis of other available data from humans and animals. In keeping with these principles, a formal analysis of the temporal concordance of key events in the hypothesized modes of action is not presented unless it would aid in the overall weight of evidence analysis for carcinogenicity, as presented in Section 4.11.

<sup>8</sup> Evaluation of genotoxicity data entails a weight of evidence approach that includes consideration of the various types of genetic damage that can occur. In acknowledging that genotoxicity tests are by design complementary evaluations of different mechanisms of genotoxicity, a recent IPCS publication (Eastmond et al., 2009) notes that “multiple negative results may not be sufficient to remove concern for mutagenicity raised by a clear positive result in a single mutagenicity assay.” These considerations inform the present approach. In addition, consistent with EPA’s *Cancer Guidelines* (2005c, d), the approach does not address relative potency (e.g., among TCE metabolites, or of such metabolites with other known genotoxic carcinogens) *per se*, nor does it consider quantitative issues

1 mutagenic in the Ames test in three of the tested strains of *S. typhimurium* (TA100, TA2638,  
2 TA98) (Dekant et al., 1986c; Vamvakas et al., 1988b) and caused dose-dependent increases in  
3 unscheduled DNA synthesis in the two available assays: porcine kidney tubular epithelial cell  
4 line (Vamvakas et al., 1996) and Syrian hamster embryo fibroblasts (Vamvakas et al., 1988a).  
5 DCVC has also been shown to induce DNA strand breaks in both available studies (Jaffe et al.,  
6 1985; Robbiano et al., 2004), and induce micronucleus formation in primary kidney cells from  
7 rats and humans (Robbiano et al., 2004) but not in Syrian hamster embryo fibroblasts (Vamvakas  
8 et al., 1988a). Only one study each is available for DCVG and *N*-AcDCVC, but notably both  
9 were positive in the Ames test (Vamvakas et al., 1987; 1988b). Although the number of test  
10 systems was limited, these results are consistent.

11 These in vitro results are further supported by studies reporting kidney-specific  
12 genotoxicity after in vivo administration of TCE or DCVC. In particular, Robbiano et al. (1998)  
13 reported increased numbers of micronucleated cells in the rat kidney following oral TCE  
14 exposure. Oral exposure to DCVC in both rabbits (Jaffe et al., 1985) and rats (Clay, 2008)  
15 increased DNA strand breaks in the kidney. However, in one inhalation exposure study in rats,  
16 TCE did not increase DNA breakage in the rat kidney, possibly due to study limitations (limited  
17 exposure time [6 hours/day for only 5 d] and small number of animals exposed [ $n = 5$ ]; Clay,  
18 (2008). One study of TCE exposure in the Eker rat, a rat model heterozygous for the tumor  
19 suppressor gene *Tsc-2*, reported no significant increase in kidney tumors as compared to controls  
20 (Mally et al., 2006). Inactivation of *Tsc-2* in this rat model is associated with spontaneous renal  
21 cell carcinoma with activation of pathways similar to that of *VHL* inactivation in humans (Liu et  
22 al., 2003). TCE exposure for 13-weeks (corn oil gavage) led to increased nephrotoxicity but no  
23 significant increases in preneoplastic or neoplastic lesions as compared to controls (Mally et al.,  
24 2006). This lack of increased incidence of neoplastic or preneoplastic lesions reported by Mally  
25 et al. (2006) in the tumor-prone Eker rat is similar to lack of significant short-term response  
26 exhibited by other genotoxic carcinogens in the Eker rat (Morton et al., 2002; Stemmer et al.,  
27 2007) and may be related to the increased background rate of renal carcinomas in this animal  
28 model. Mally et al. (2006) also exposed primary kidney epithelial cells from the Eker rat to  
29 DCVC in vitro and demonstrated increased transformation similar to that of other renal  
30 carcinogens (Horesovsky et al., 1994).

31 As discussed in Section 4.2.1.4.1, although Douglas et al. (1999) did not detect increased  
32 mutations in the kidney of *lacZ* transgenic mice exposed to TCE for 12 days, these results are not

---

related to the probable production of these metabolites in vivo. Instead, the analysis of genetic toxicity data presented in Section 4.2 and summarized here focuses on the identification of a genotoxic hazard of these metabolites; a quantitative analysis of TCE metabolism to reactive intermediates, via PBPK modeling, is presented in Section 3.5.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 highly informative as to the role of mutagenicity in TCE-induced kidney tumors, given the  
2 uncertainties in the production in genotoxic GSH conjugation metabolites in mice and the low  
3 carcinogenic potency of TCE for kidney tumors in rodents relative to what is detectable in  
4 experimental bioassays. Limited, mostly in vitro, toxicokinetic data do not suggest mice have  
5 less GSH conjugation or subsequent renal metabolism/bioactivation (see Section 3.3.3.2.7), but  
6 quantitatively, the uncertainties in the flux through these pathways remain significant (see  
7 Section 3.5). In additional, similar to other genotoxic renal carcinogens analyzed by NTP, there  
8 is limited evidence of mouse kidney tumors following TCE exposure. However, given the  
9 already low incidences of kidney tumors observed in rats, a relatively small difference in potency  
10 in mice would be undetectable in available chronic bioassays. Notably, of seven chemicals  
11 categorized as direct-acting genotoxic carcinogens that induced rat renal tumors in NTP studies,  
12 only two also led to renal tumors in the mouse (tris[2,3-dibromopropyl]phosphate and  
13 ochratoxin A) (Kanisawa and Suzuki, 1978; Reznik et al., 1979), so the lack of detectable  
14 response in mouse bioassays does not preclude a genotoxic MOA.

15 *VHL* inactivation (via mechanisms such as deletion, silencing or mutation) observed in  
16 human renal clear cell carcinomas, is the basis of a hereditary syndrome of kidney cancer  
17 predisposition, and is hypothesized to be an early and causative event in this disease (e.g., 2008).  
18 Therefore, specific actions of TCE metabolites that produce or select for mutations of the *VHL*  
19 suppressor gene could lead to kidney tumorigenesis. Several studies have compared *VHL*  
20 mutation frequencies in cases with TCE exposures with those from control or background  
21 populations. Brüning et al. (1997b) and Brauch et al. (1999; 2004) reported differences between  
22 TCE-exposed and nonexposed renal cell carcinoma patients in the frequency of somatic *VHL*  
23 mutations, the incidence of a hot spot mutation of cytosine to thymine at nucleotide 454, and the  
24 incidence of multiple mutations. These data suggest that kidney tumor genotype data in the form  
25 of a specific mutation pattern may potentially serve to discriminate TCE-induced tumors from  
26 other types of kidney tumors in humans. If validated, this would also suggest that TCE-induced  
27 kidney tumors are dissimilar from those occurring in unexposed individuals. Thus, while not  
28 confirming a mutation MOA, these data suggest that TCE-induced tumors may be distinct from  
29 those induced spontaneously in humans. However, it has not been examined whether a possible  
30 linkage exists between *VHL* loss or silencing and mutagenic TCE metabolites.

31 By contrast, Schraml et al. (1999) and Charbotel et al. (2007) reported that TCE-exposed  
32 renal cell carcinoma patients did not have significantly higher incidences of *VHL* mutations  
33 compared to nonexposed patients. However, details as to the exposure conditions were lacking  
34 in Schraml et al. (1999). In addition, the sample preparation methodology employed by  
35 Charbotel et al. (2007) and others (Brauch et al., 1999; Brüning et al., 1997b) often results in

1 poor quality and/or low quantity DNA, leading to study limitations (less than 100% of samples  
2 were able to be analyzed). Therefore, further investigations are necessary to either confirm or  
3 contradict the validity of the genetic biomarkers for TCE-related renal tumors reported by  
4 Brüning et al. (1997b) and Brauch et al. (1999; 2004).

5 In addition, while exposure to mutagens is certainly associated with cancer induction (as  
6 discussed with respect to the liver in Appendix E, Sections E.3.1 and E.3.2), examination of  
7 end-stage tumor phenotype or genotype has limitations concerning determination of early key  
8 events. The mutations that are observed with the progression of neoplasia are associated with  
9 increased genetic instability and an increase in mutation rate. Further, inactivation of the *VHL*  
10 gene also occurs through other mechanisms in addition to point mutations, such as loss of  
11 heterozygosity or hypermethylation (Kenck et al., 1996; Nickerson et al., 2008) not addressed in  
12 these studies. Recent studies examining the role of other genes or pathways suggest roles for  
13 multiple genes in renal cell carcinoma development (Furge et al., 2007; Toma et al., 2008).  
14 Therefore, the inconsistent results with respect to *VHL* mutation status do not constitute negative  
15 evidence for a mutational MOA and the positive studies are suggestive of a TCE-induced kidney  
16 tumor genotype.

17 In sum, the predominance of positive genotoxicity data in the database of available  
18 studies of TCE metabolites derived from GSH conjugation (in particular the evidence of  
19 kidney-specific genotoxicity following in vivo exposure to TCE or DCVC), coupled with the  
20 toxicokinetic data consistent with the in situ formation of these GSH-conjugation metabolites of  
21 TCE in the kidney, is consistent with the hypothesis that a mutagenic MOA is operative in  
22 TCE-induced kidney tumors. Mutagenicity is a well-established cause of carcinogenicity.  
23 Available data on the *VHL* gene in humans add biological plausibility to these conclusions.  
24 Quantitatively, however, as discussed in Section 3.3.3.2, the precise metabolic yield of the GSH  
25 conjugation metabolites following TCE exposure remains uncertain.

#### 4.4.7.1.3. Hypothesized Mode of Action: Cytotoxicity and Regenerative Proliferation

27 Another hypothesis is that TCE acts by a cytotoxicity mode of action in TCE-induced  
28 renal carcinogenesis. According to this hypothesis, the key events leading to TCE-induced  
29 kidney tumor formation comprise the following: the TCE GSH-conjugation metabolite DCVC,  
30 after being either produced in situ in or delivered systemically to the kidney, causes cytotoxicity,  
31 leading to compensatory cellular proliferation and subsequently increased mutations and clonal  
32 expansion of initiated cells.

#### 4.4.7.1.4. Experimental support for the hypothesized mode of action

1 Evidence for the hypothesized MOA consist primarily of (1) the demonstration of  
2 nephrotoxicity following TCE exposure at current occupational limits in human studies and  
3 chronic TCE exposure in animal studies; (2) the relatively high potential of the TCE metabolite  
4 DCVC to cause nephrotoxicity; and (3) toxicokinetic data demonstrating that DCVC is formed in  
5 the kidney following TCE exposure. Data on nephrotoxicity of TCE and DCVC are discussed in  
6 more detail below, while the toxicokinetic data were summarized previously in the discussion of  
7 mutagenicity. Thus, the data are consistent with the hypothesized MOA, and therefore do not  
8 rule out a contribution from cytotoxicity and regenerative proliferation to TCE-induced kidney  
9 carcinogenesis. However, there are a lack of experimental data supporting a causal link between  
10 TCE nephrotoxicity combined with sustained cellular proliferation and TCE-induced  
11 nephrocarcinogenicity.

12 There is substantial evidence that TCE is nephrotoxic in humans and laboratory animals  
13 and that its metabolite DCVC is nephrotoxic in laboratory animals. Epidemiological studies  
14 have consistently demonstrated increased excretion of nephrotoxicity markers (NAG, protein,  
15 albumin) at occupational (Green et al., 2004) and higher (Bolt et al., 2004; Brüning et al., 1999a;  
16 1999b) levels of TCE exposure. However, direct evidence of tubular toxicity, particularly in  
17 renal cell carcinoma cases, is not available. These studies are supported by the results of  
18 multiple laboratory animal studies. Chronic bioassays have reported very high (nearly 100%)  
19 incidences of nephrotoxicity of the proximal tubule in rats (NTP, 1988, 1990) and mice (NCI,  
20 1976; NTP, 1990) at the highest doses tested. In vivo studies examining the effect of TCE  
21 exposure on nephrotoxicity showed increased proximal tubule damage following intraperitoneal  
22 injection and inhalation of TCE in rats (Chakrabarti and Tuchweber, 1988) and intraperitoneal  
23 injection in mice (Cojocel et al., 1989). Studies examining DCVC exposure in rats (Elfarra et  
24 al., 1986; TERRACINI and PARKER, 1965) and mice (Darnerud et al., 1989; Jaffe et al., 1984)  
25 have also shown increases in kidney toxicity. The greater potency for kidney cytotoxicity for  
26 DCVC compared to TCE was shown by in vitro studies (Lash et al., 1986; 1995; Stevens et al.,  
27 1986). These studies also further confirmed the higher susceptibility of male rats or mice to  
28 DCVC-induced cytotoxicity. Cytokaryomegaly (an effect specific to TCE and not part of the  
29 chronic progressive nephropathy or the pathology that occurs in aging rat kidneys) was observed  
30 in the majority of rodent studies and may or may not progress to carcinogenesis. Finally, as  
31 discussed extensively in Section 4.4.6.1, a detailed comparison of the histological changes in the  
32 kidney caused by TCE and its metabolites supports the conclusion that DCVC is the predominant  
33 moiety responsible for TCE-induced nephrotoxicity.

1           Because it is known that not all cytotoxins are carcinogens (i.e., cytotoxicity is not a  
2 specific predictor of carcinogenicity), additional experimental support is required to causally link  
3 nephrotoxicity to nephrocarcinogenicity. For chemicals that bind to  $\alpha_2\mu$ -globulin, a MOA  
4 involving cell necrosis followed by subsequent regenerative proliferation has been hypothesized  
5 to cause kidney tumors in the absence of genotoxicity (Short, 1993). However, for other  
6 chemicals, toxicity and increased cell proliferation have been observed in the kidney without  
7 inducing tumors even after chronic exposure (Tennant et al., 1991). Similarly, in the liver,  
8 partial hepatectomy leading to regenerative hyperplasia does not by itself lead to increased  
9 hepatocarcinogenicity, and requires administration of a mutagen to exhibit enhanced  
10 carcinogenic effects. By analogy, a biologically plausible MOA may involve a combination of  
11 mutagenicity and cytotoxicity, with mutagenicity increasing the rate of mutation and  
12 regenerative proliferation induced by cytotoxicity enhancing the selection, survival or clonal  
13 expansion of mutated cells.

14           For TCE and kidney cancer, clearly, cytotoxicity occurs at doses below those causing  
15 carcinogenicity, as the incidence of nephrotoxicity in chronic bioassays is an order of magnitude  
16 higher than that of renal tumors. Thus, these data are consistent with cytotoxicity being a  
17 precursor to carcinogenicity (i.e., if the opposite were the case—carcinogenicity without  
18 cytotoxicity—then the hypothesis would be falsified). While chronic nephrotoxicity was reported  
19 in the same bioassays showing increased kidney tumor incidences, the use of such data to inform  
20 MOA is indirect and associative, and do not offer a test of the hypothesis (Short, 1993).  
21 Nephrotoxicity is observed in both mice and rats, in some cases with nearly 100% incidence in  
22 all dose groups, but kidney tumors are only observed at low incidences in rats at the highest  
23 tested doses (NCI, 1976; NTP, 1990). Therefore, nephrotoxicity alone appears to be insufficient,  
24 or at least not rate-limiting, for rodent renal carcinogenesis, since maximal levels of toxicity are  
25 reached before the onset of tumors. Furthermore, there are multiple mechanisms by which TCE  
26 has been hypothesized to induce cytotoxicity, including oxidative stress, disturbances in calcium  
27 ion homeostasis, mitochondrial dysfunction, and protein alkylation (Lash et al., 2000b). Some of  
28 these effects may therefore, have ancillary consequences related to tumor induction which are  
29 independent of cytotoxicity *per se*. Therefore, data currently cannot distinguish as to whether  
30 cytotoxicity is causally related to tumorigenesis or merely associated by virtue of being a marker  
31 for a different, key causal event.

32           Under the hypothesized MOA, cytotoxicity leads to the induction of repair processes and  
33 compensatory proliferation that could lead to an increased production or clonal expansion of  
34 cells previously initiated by mutations occurred spontaneously, from coexposures, or from TCE  
35 or its metabolites. Data on compensatory cellular proliferation and the subsequent hypothesized

1 key events in the kidney are few, with no data from rat strains used in chronic bioassays. In rats  
2 carrying the Eker mutation, Mally et al. (2006) reported increased DNA synthesis as measured  
3 by BrdU incorporation in animals exposed to the high dose of TCE (1,000 mg/kg-day) for  
4 13 weeks, but there was no evidence of clonal expansion or tumorigenesis in the form of  
5 increased preneoplastic or neoplastic lesions as compared to controls. Therefore, in both rodent  
6 and human studies of TCE, data demonstrating a causal link between compensatory proliferation  
7 and the induction of kidney tumors are lacking.

8 In sum, the predominance of positive nephrotoxicity data in the database of available  
9 studies of TCE metabolites derived from GSH conjugation (in particular the evidence of  
10 kidney-specific cytotoxicity following in vivo exposure to TCE or DCVC), coupled with the  
11 toxicokinetic data consistent with the in situ formation of these GSH-conjugation metabolites of  
12 TCE in the kidney, is consistent with the hypothesis that a MOA involving cytotoxicity and  
13 regenerative proliferation contributes TCE-induced kidney tumors, either independently or in  
14 combination with a mutagenic MOA. However, nephrotoxicity is not in itself predictive of  
15 tumorigenesis, and experimental data supporting for a causal link between TCE nephrotoxicity  
16 combined with sustained cellular proliferation and TCE-induced nephrocarcinogenicity are  
17 lacking. A more biologically plausible MOA may involve a combination of mutagenicity and  
18 cytotoxicity, with mutagenicity increasing the rate of mutation and regenerative proliferation  
19 enhancing the selection, survival or clonal expansion of mutated cells. However, this hypothesis  
20 has yet to be tested experimentally.

#### 21 **4.4.7.1.5. Additional Hypothesized Modes of Action with Limited Evidence or Inadequate Experimental Support**

22 Along with metabolites derived from GSH conjugation of TCE, oxidative metabolites are  
23 also present and could induce toxicity in the kidney. After TCE exposure, the oxidative  
24 metabolite and peroxisome proliferator TCA is present in the kidney and excreted in the urine as  
25 a biomarker of exposure. Hypotheses have also been generated regarding the roles of  
26  $\alpha_2\mu$ -globulin or formic acid in nephrotoxicity induced by TCE oxidative metabolites TCA or  
27 TCOH. However, the available data are limited or inadequate for supporting these hypothesized  
28 MOAs.

#### 4.4.7.1.6. Peroxisome proliferation

1           Although not as well studied as the effects of glutathione metabolites in the kidney, there  
2 is evidence that oxidative metabolites affect the kidney after TCE exposure. Both TCA and  
3 DCA are peroxisome proliferator activated receptor alpha (PPAR $\alpha$ ) agonists although most  
4 activity has been associated with TCA production after TCE exposure. Exposure to TCE has  
5 been found to induce peroxisome proliferation not only in the liver but also the kidney.  
6 Peroxisome proliferation in the kidney has been evaluated by only one study of TCE  
7 (Goldsworthy and Popp, 1987), using increases in cyanide-insensitive palmitoyl-CoA oxidation  
8 (PCO) activity as a marker. Increases in renal PCO activity were observed in rats (3.0-fold) and  
9 mice (3.6-fold) treated with TCE at 1,000 mg/kg-day for 10 days, with smaller increases in both  
10 species from TCA treatment at 500 mg/kg-day for 10 days. However, no significant increases in  
11 kidney/body weight ratios were observed in either species. There was no relationship between  
12 induction of renal peroxisome proliferation and renal tumors (i.e., a similar extent of peroxisome  
13 proliferation-associated enzyme activity occurred in species with and without TCE-induced renal  
14 tumors). However, the increased peroxisomal enzyme activities due to TCE exposure are  
15 indicative of oxidative metabolites being present and affecting the kidney. Such metabolites  
16 have been associated with other tumor types, especially liver, and whether coexposures to  
17 oxidative metabolites and glutathione metabolites contribute to kidney tumorigenicity has not  
18 been examined.  
19

#### 4.4.7.1.7. $\alpha$ 2 $\mu$ -Globulin-related nephropathy

20           Induction of  $\alpha$ 2 $\mu$ -globulin nephropathy by TCE has been investigated by Goldsworthy  
21 et al. (1988), who reported that TCE did not induce increases in this urinary protein, nor did it  
22 stimulate cellular proliferation in rats. In addition, whereas kidney tumors associated with  
23  $\alpha$ 2 $\mu$ -globulin nephropathy are specific to the male rat, as discussed above, nephrotoxicity is  
24 observed in both rats and mice and kidney tumor incidence is elevated (though not always  
25 statistically significant) in both male and female rats. TCOH was recently reported to cause  
26 hyaline droplet accumulation and an increase in  $\alpha$ 2 $\mu$ -globulin, but these levels were insufficient  
27 to account for the observed nephropathy as compared to other exposures (Green et al., 2003).  
28 Therefore, it is unlikely that  $\alpha$ 2 $\mu$ -globulin nephropathy contributes significantly to TCE-induced  
29 renal carcinogenesis.  
30



#### **4.4.7.1.8. Formic acid-related nephrotoxicity**

1 Another MOA hypothesis proposes that TCE nephrotoxicity is mediated by increased  
2 formation and urinary excretion of formic acid mediated by the oxidative metabolites TCA or  
3 TCOH (Dow and Green, 2000; Green et al., 2003; 1998). The subsequent hypothesized key  
4 events are the same as those for DCVC-induced cytotoxicity, discussed above (see  
5 Section 4.4.7.2). As discussed extensively in Section 4.4.6.1.2, these oxidative metabolites do  
6 not appear sufficient to explain the range of renal effects observed after TCE exposure,  
7 particularly cytomegaly, karyomegaly, and flattening and dilation of the tubular epithelium.  
8 Although TCOH and possibly TCA may contribute to the nephrotoxicity of TCE, perhaps due to  
9 excess formic acid production, these metabolites do not show the same range of cytotoxic effects  
10 observed following TCE exposure (see Table 4-53). Therefore, without specific evidence  
11 linking the specific nephrotoxic effects caused by TCOH or TCA to carcinogenesis, and in light  
12 of the substantial evidence that DCVC itself can adequately account for the nephrotoxic effects  
13 of TCE, the weight of evidence supports a conclusion that cytotoxicity mediated by increased  
14 formic acid production induced by oxidative metabolites TCOH and possibly TCA is not  
15 responsible for the majority of the TCE-induced cytotoxicity in the kidneys, and therefore, would  
16 not be the major contributor to the other hypothesized key events in this MOA, such as  
17 subsequent regenerative proliferation.  
18

#### **4.4.7.1.9. Conclusions About the Hypothesized Modes of Action**

##### **4.4.7.1.10. Is the hypothesized mode of action sufficiently supported in the test animals**

###### **4.4.7.1.10.1. Mutagenicity**

19 The predominance of positive genotoxicity data in the database of available studies of  
20 TCE metabolites derived from GSH conjugation (in particular the evidence of kidney-specific  
21 genotoxicity following in vivo exposure to TCE or DCVC), coupled with the toxicokinetic data  
22 consistent with the in situ formation of these GSH-conjugation metabolites of TCE in the kidney,  
23 supports the conclusion that a mutagenic MOA is operative in TCE-induced kidney tumors.  
24

#### 4.4.7.1.10.2. Cytotoxicity

1 As reviewed above, in vivo and in vitro studies have shown a consistent nephrotoxic  
2 response to TCE and its metabolites in proximal tubule cells from male rats. Therefore, it has  
3 been proposed that cytotoxicity seen in this region of the kidney is a precursor to carcinogenicity.  
4 Available data are consistent with the hypothesis that a MOA involving cytotoxicity and  
5 regenerative proliferation contributes to TCE-induced kidney tumors, either independently or in  
6 combination with a mutagenic MOA. However, it has not been determined whether tubular  
7 toxicity is a necessary precursor of carcinogenesis, and there is a lack of experimental support for  
8 causal links, such as compensatory cellular proliferation or clonal expansion of initiated cells,  
9 between nephrotoxicity and kidney tumors induced by TCE. Nephrotoxicity alone appears to be  
10 insufficient, or at least not rate-limiting, for rodent renal carcinogenesis, since maximal levels of  
11 toxicity are reached before the onset of tumors. A more biologically plausible MOA may  
12 involve a combination of mutagenicity and cytotoxicity, with mutagenicity increasing the rate of  
13 mutation and regenerative proliferation enhancing the survival or clonal expansion of mutated  
14 cells. However, this hypothesis has yet to be tested experimentally.  
15

#### 4.4.7.1.10.3. Additional hypotheses

16 The kidney is also exposed to oxidative metabolites that have been shown to be  
17 carcinogenic in other target organs. TCA is excreted in kidney after its metabolism from TCE  
18 and also can cause peroxisome proliferation in the kidney, but there are inadequate data to define  
19 a MOA for kidney tumor induction based on peroxisome proliferation. TCE induced little or no  
20  $\alpha_2\mu$ -globulin and hyaline droplet accumulation to account for the observed nephropathy, so  
21 available data do not support this hypothesized MOA. The production of formic acid following  
22 exposure to TCE and its oxidative metabolites TCOH and TCA may also contribute to  
23 nephrotoxicity; however, the available data indicate that TCOH and TCA are minor contributors  
24 to TCE-induced nephrotoxicity, and therefore, do not support this hypothesized MOA. Because  
25 these additional MOA hypotheses are either inadequately defined or are not supported by the  
26 available data, they are not considered further in the conclusions below.  
27

#### **4.4.7.1.11. 2. Is the hypothesized mode of action relevant to humans**

##### **4.4.7.1.11.1. Mutagenicity**

1           The evidence discussed above demonstrates that TCE GSH-conjugation metabolites are  
2 mutagens in microbial as well as test animal species. Therefore, the presumption that they would  
3 be mutagenic in humans. Available data on the *VHL* gene in humans add biological plausibility  
4 to this hypothesis. The few available data from human studies concerning the mutagenicity of  
5 TCE and its metabolites suggest consistency with this MOA, but are not sufficiently conclusive  
6 to provide direct supporting evidence for a mutagenic MOA. Therefore, this MOA is considered  
7 relevant to humans.

##### **4.4.7.1.11.2. Cytotoxicity**

9           Although data are inadequate to determine that the MOA is operative, none of the  
10 available data suggest that this MOA is biologically precluded in humans. Furthermore, both  
11 animal and human studies suggest that TCE causes nephrotoxicity at exposures that also induce  
12 renal cancer, constituting positive evidence of the human relevance of this hypothesized MOA.

#### **4.4.7.1.12. 3. Which populations or lifestages can be particularly susceptible to the hypothesized mode of action**

##### **4.4.7.1.12.1. Mutagenicity**

14           The mutagenic MOA is considered relevant to all populations and lifestages. According  
15 to EPA's *Cancer Guidelines* (U.S. EPA, 2005c) and *Supplemental Guidance* (U.S. EPA, 2005d),  
16 there may be increased susceptibility to early-life exposures for carcinogens with a mutagenic  
17 mode of action. Therefore, because the weight of evidence supports a mutagenic mode of action  
18 for TCE carcinogenicity and in the absence of chemical-specific data to evaluate differences in  
19 susceptibility, early-life susceptibility should be assumed and the age-dependent adjustment  
20 factors (ADAFs) should be applied, in accordance with the *Supplemental Guidance*.

21           In addition, because the MOA begins with GSH-conjugation metabolites being delivered  
22 systemically or produced in situ in the kidney, toxicokinetic differences—i.e., increased  
23 production or bioactivation of these metabolites—may render some individuals more susceptible  
24 to this MOA. However, as discussed in Section 3.3.3.2, quantitative estimates of the amount of

1 GSH conjugation following TCE exposure remain uncertain. Toxicokinetic-based susceptibility  
2 is discussed further in Section 4.10.

3 In rat chronic bioassays, TCE-treated males have higher incidence of kidney tumors than  
4 similarly treated females. However, the basis for this sex-difference is unknown, and whether it  
5 is indicative of a sex difference in human susceptibility to TCE-induced kidney tumors is  
6 likewise unknown. The epidemiologic studies generally do not show sex differences in kidney  
7 cancer risk. Lacking exposure-response information, it is not known if the sex-difference in one  
8 renal cell carcinoma case-control study (Dosemeci et al., 1999) may reflect exposure differences  
9 or susceptibility differences.

#### 10 4.4.7.1.12.2. **Cytotoxicity**

11 Populations which may be more susceptible based on the toxicokinetics of the production  
12 of GSH conjugation metabolites and the sex differences observed in rat chronic bioassays are the  
13 same as for a mutagenic MOA. No data are available as to whether other factors may lead to  
14 different populations or lifestages being more susceptible to a cytotoxic MOA for TCE-induced  
15 kidney tumors. For instance, it is not known how the hypothesized key events in this MOA  
16 interact with known risk factors for human renal cell carcinoma.

17 The weight of evidence sufficiently supports a mutagenic MOA for TCE in the kidney,  
18 based on supporting data that GSH-metabolites are genotoxic and produced in sufficient  
19 quantities in the kidney to lead to tumorigenesis. Cytotoxicity and regenerative proliferation  
20 were considered as an alternate MOA, however, there are inadequate data to support a causal  
21 association between cytotoxicity and kidney tumors. Further, hypothesized MOAs relating to  
22 peroxisomal proliferation,  $\alpha_2\mu$ -globulin nephropathy and formic acid-related nephrotoxicity  
23 were considered and rejected due to limited evidence and/or inadequate experimental support.

#### 24 **4.4.8. Summary: Trichloroethylene (TCE) Kidney Toxicity, Carcinogenicity, and Mode-of-Action**

25 Human studies have shown increased levels of proximal tubule damage in workers  
26 exposed to high levels of TCE (NRC, 2006). These studies analyzed workers exposed to TCE  
27 alone or in mixtures and reported increases in various urinary biomarkers of kidney toxicity or  
28 end-stage renal disease ( $\beta_2$ -microglobulin, total protein, NAG,  $\alpha_1$ -microglobulin) (Bolt et al.,  
29 2004; Brüning et al., 1999a; 1999b; Green et al., 2004; Jacob et al., 2007; Nagaya et al., 1989b;  
30 Radican et al., 2006; Selden et al., 1993). Laboratory animal studies examining TCE exposure

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 provide additional support, as multiple studies by both gavage and inhalation exposure show that  
2 TCE causes renal toxicity in the form of cytomegaly and karyomegaly of the renal tubules in  
3 male and female rats and mice. By gavage, incidences of these effects under chronic bioassay  
4 conditions approach 100%, with male rats appearing to be more sensitive than either female rats  
5 or mice of either sex based on the severity of effects. Under chronic inhalation exposures, only  
6 male rats exhibited these effects. Further studies with TCE metabolites have demonstrated a  
7 potential role for DCVC, TCOH, and TCA in TCE-induced nephrotoxicity. Of these, DCVC  
8 induces the renal effects that are most like TCE, and it is formed in sufficient amounts following  
9 TCE exposure to account for these effects.

10 Kidney cancer risk from TCE exposure has been studied related to TCE exposure in  
11 cohort, case-control and geographical studies. These studies have examined TCE in mixed  
12 exposures as well as alone. Elevated risks are observed in many of the cohort and case-control  
13 studies examining kidney cancer incidence in industries or job titles with historical use of TCE  
14 (see Table 4-39 and 4-40), particularly among subjects ever exposed to TCE (Brüning et al.,  
15 2003; Dosemeci et al., 1999; Moore et al., 2010; Raaschou-Nielsen et al., 2003) or subjects with  
16 TCE surrogate for high exposure (Brüning et al., 2003; Charbotel et al., 2006; Moore et al.,  
17 2010; Raaschou-Nielsen et al., 2003; Zhao et al., 2005). Greater susceptibility to TCE exposure  
18 and kidney cancer is observed among subjects with a functionally active GSTT polymorphism,  
19 particularly among those with certain alleles in single nucleotide polymorphisms of the cysteine  
20 conjugation  $\beta$ -lyase gene region (Moore et al., 2010). Although there are some controversies  
21 related to deficiencies of the epidemiological studies (Henschler et al., 1995; Vamvakas et al.,  
22 1998), many of these are overcome in later studies (Brüning et al., 2003; Charbotel et al., 2006;  
23 Moore et al., 2010). A meta-analysis of the overall effect of TCE exposure on kidney cancer,  
24 additionally, suggests a small, statistically significant increase in risk (RRm = 1.27 95% CI: 1.13,  
25 1.43) with a summary relative risk estimate in the higher exposure group of 1.64, (95% CI: 1.31,  
26 2.04), robust in sensitivity to alternatives and lacking observed statistical heterogeneity among  
27 17 studies meeting explicitly-defined inclusion criteria.

28 In vivo laboratory animal studies to date suggest a small increase in renal tubule tumors  
29 in male rats and, to a lesser extent, in female rats, with no increases seen in mice or hamsters.  
30 These results are based on limited studies of both oral and inhalation routes, some of which were  
31 deemed insufficient to determine carcinogenicity based on various experimental issues.  
32 However, because of the rarity of kidney tumors in rodents, the repeatability of this finding  
33 across strains and studies supports their biological significance despite the limitations of  
34 individual studies and relatively small increases in reported tumor incidence.

1           Some but not all human studies have suggested a role for *VHL* mutations in TCE-induced  
2 kidney cancer (Brauch et al., 1999; 2004; Brüning et al., 1997b; Charbotel et al., 2007; Schraml  
3 et al., 1999). Certain aspects of these studies may explain some of these discrepant results. The  
4 majority of these studies have examined paraffinized tissue that may lead to technical difficulties  
5 in analysis, as paraffin extractions yield small quantities of often low-quality DNA. The  
6 chemicals used in the extraction process itself may also interfere with enzymes required for  
7 further analysis (PCR, sequencing). Although these studies do not clearly show mutations in all  
8 TCE-exposed individuals, or in fact in all kidney tumors examined, this does not take into  
9 account other possible means of *VHL* inactivation, including silencing or loss, and other potential  
10 targets of TCE mutagenesis were not systematically examined. A recent study by Nickerson  
11 et al. (2008) analyzed both somatic mutation and promoter hypermethylation of the *VHL* gene in  
12 cc-RCC frozen tissue samples using more sensitive methods. The results of this study support  
13 the hypothesis that *VHL* alterations are an early event in clear cell RCC carcinogenesis, but these  
14 alterations may not be gene mutations. No experimental animal studies have been performed  
15 examining *vhl* inactivation following exposure to TCE, although one in vitro study examined *vhl*  
16 mutation status following exposure to the TCE-metabolite DCVC (Mally et al., 2006). This  
17 study found no mutations following DCVC exposure, although this does not rule out a role for  
18 DCVC in *vhl* inactivation by some other method or *vhl* alterations caused by other TCE  
19 metabolites.

20           Although not encompassing all of the actions of TCE and its metabolites that may be  
21 involved in the formation and progression of neoplasia, available evidence supports the  
22 conclusion that a mutagenic MOA mediated by the TCE GSH-conjugation metabolites  
23 (predominantly DCVC) is operative in TCE-induced kidney cancer. This conclusion is based on  
24 substantial evidence that these metabolites are genotoxic and are delivered to or produced in the  
25 kidney, including evidence of kidney-specific genotoxicity following in vivo exposure to TCE or  
26 DCVC. Cytotoxicity caused by DCVC leading to compensatory cellular proliferation is also a  
27 potential MOA in renal carcinogenesis. A combination of mutagenicity and cytotoxicity, with  
28 mutagenicity increasing the rate of mutation and regenerative proliferation enhancing the  
29 survival or clonal expansion of mutated cells, while biologically plausible, has yet to be tested  
30 experimentally. The additional MOA hypotheses of peroxisome proliferation, accumulation of  
31  $\alpha_2\mu$ -globulin, and cytotoxicity mediated by TCE-induced excess formic acid production are not  
32 supported by the available data.

33

## 4.5. LIVER TOXICITY AND CANCER

### 4.5.1. Liver Noncancer Toxicity in Humans

1           The complex of chronic liver disease is a spectrum of effects and comprises nonalcoholic  
2 fatty liver disease (nonalcoholic steatohepatitis) and cirrhosis, more rare anomalies ones such as  
3 autoimmune hepatitis, primary biliary cirrhosis, and primary sclerosing cholangitis, and  
4 hepatocellular and cholangiocarcinoma (intrahepatic bile duct cancer) (Juran and Lazaridis,  
5 2006). Chronic liver disease and cirrhosis, excluding neoplasia, is the 12<sup>th</sup> leading cause of death  
6 in the United States in 2005 with 27,530 deaths (Kung et al., 2008) with a morality rate of 9.0  
7 per 100,000 (Jemal et al., 2008).

8           Eight studies reported on liver outcomes and TCE exposure and are identified in  
9 Table 4-55. Three studies are suggestive of effects on liver function tests in metal degreasers  
10 occupationally exposed to trichloroethylene (Nagaya et al., 1993; Rasmussen et al., 1993b; Xu et  
11 al., 2009). Nagaya et al. (1993) in their study of 148 degreasers in metal parts factories,  
12 semiconductor factors, or other factories, observed total mean serum cholesterol concentration,  
13 mean serum high density lipoprotein-cholesterol (HDL-C) concentrations to increase with  
14 increasing TCE exposure, as defined by U-TTC), although a statistically significant linear trend  
15 was not found. Nagaya et al. (1993) estimated subjects in the low exposure group had TCE  
16 exposure to 1 ppm-, 6-ppm TCE in the moderate exposure group, and 210-ppm TCE in the high  
17 exposure group. No association was noted between serum liver function tests and U-TTC, a  
18 finding not surprising given individuals with a history of hepatobiliary disease were excluded  
19 from this study. Nagaya et al. (1993) follows 13 workers with higher U-TTC concentrations  
20 over a 2-year period; serum HDL-C and two hepatic function enzymes, GGT and aspartate  
21 aminotrasferase (AST) concentrations were highest during periods of high level exposure, as  
22 indicated from U-TTC concentrations. Similarly, in a study of 95 degreasers, 70 exposed to  
23 trichloroethylene exposure and 25 to CFC113 (Rasmussen et al., 1993b), mean serum GGT  
24 concentration for subjects with the highest TCE exposure duration was above normal reference  
25 values and were about threefold higher compared to the lowest exposure group. Rasmussen  
26 et al. (1993b) estimated mean urinary TCE concentration in the highest exposure group as  
27 7.7 mg/L with past exposures estimated as equivalent to 40–60 mg/L. Multivariate regression  
28 analysis showed a small statistically nonsignificant association due to age and a larger effect due  
29 to alcohol abuse that reduced and changed direction of a TCE exposure affect. The inclusion of  
30 CFC113 exposed subjects introduces a downward bias since liver toxicity is not associated with  
31 CFC113 exposure (U.S. EPA, 2008a) and would underestimate any possible TCE effect. Xu  
32 et al. (2009) reported symptoms and liver function tests of 21 metal degreasers with severe

1 hypersensitivity dermatitis (see last paragraph in this section for discussion of other liver effects  
 2 in hypersensitivity dermatitis cases). TCE concentration of agent used to clean metal parts  
 3 ranged from 10.2–63.5% with workplace ambient monitoring time-weighted-average TCE  
 4 concentrations of 18–683 mg/m<sup>3</sup> (3–127 ppm). Exposure was further documented by urinary  
 5 TCA levels in 14 of 21 cases above the recommended occupation level of 50 mg/L. The  
 6 prevalence of elevated liver enzymes among these subjects was 90% (19 cases) for alanine  
 7 aminotrasferase, 86% (18 cases) for asparatate aminotrasferase, and 76% (16 cases) for total  
 8 bilirubin (Xu et al., 2009). Two studies provide evidence of plasma or serum bile acids changes  
 9 among TCE-exposed degreasers. Neghab et al. (1997) in a small prevalence study of 10 healthy  
 10 workers (5 unexposed controls and 5 exposed) observed statistically significantly elevated total  
 11 serum bile acids, particularly deoxycholic acid and the subtotal of free bile acids, among TCE  
 12 subjects at postexposure compared to their pre-exposure concentrations and serum bile acid  
 13 levels correlated well with TCE exposure ( $r = 0.94$ ). Total serum bile acid concentration did not  
 14 change in control subjects between pre- and postexposure, nor did enzyme markers of liver  
 15 function in either unexposed or exposed subjects differ between pre and postexposure period.  
 16 However, the statistical power of this study is quite limited and the prevalence design does not  
 17 include subjects who may have left employment because of possible liver problems. The paper  
 18 provides minimal details of subject selection and workplace exposure conditions, except that  
 19 pre-exposure testing was carried out on the 1<sup>st</sup> work day of the week (pre-exposure), repeated  
 20 sampling after 2 days (postexposure), and a postexposure 8-hour time-weighted-average TCE  
 21 concentration of 9 ppm for exposed subjects; no exposure information is provided for control  
 22 subjects. Driscoll et al. (1992) in a study of 22 subjects (6 unexposed and 16 exposed) employed  
 23 at a factory manufacturing small appliances reported statistically significant group differences in  
 24 logistic regression analyses controlling for age and alcohol consumption in mean fasting plasma  
 25 bile acid concentrations. Other indicators of liver function such as plasma enzyme levels were  
 26 statistically significant different between exposed and unexposed subjects. Laboratory samples  
 27 were obtained at the start of subject’s work shift. Exposure data are not available on the  
 28 22 subjects and assignment of exposed and unexposed was based on work duties. Limited

29 **Table 4-55. Summary of human liver toxicity studies**  
 30

Subjects	Effect	Exposure	Reference
----------	--------	----------	-----------



148 male metal degreasers in metal parts, semiconductor and other factories	Serum liver function enzyme (HDL-C, AST, and GGT) concentrations did not correlate with TCE exposure assessed in a prevalence study but did correlate with TCE concentration over a 2-yr follow-up period	U-TTC levels obtained from spot urine sample obtained during working h used to assign exposure category included the following: High: $209 \pm 99$ mg/g Cr Medium: $35 \pm 27$ mg/g Cr Low: $5 \pm 2$ mg/g Cr Note: this study does not include an unexposed referent group	Nagaya et al. (1993)
95 workers (70 TCE exposed, 25 CFC113 exposed) selected from a cohort of 240 workers at 72 factors engaged in metal degreasing with chlorinated solvents	Increased serum GGT concentration with increasing cumulative exposure	4 groups (cumulative number of yr exposed over a working life): I: 0.6 (0–0.99) II: 1.9 (1–2.8) III: 4.4 (2.9–6.7) IV: 14.4 (6.8–35.6)	Rasmussen et al. (1993b)
21 metal degreasers with severe hypersensitivity dermatitis	High prevalence of serum liver function enzymes above normal levels: ALT, 19 of 21 cases; AST, 18 of 21 cases, and T-Bili, 16 of 21 cases	TWA mean ambient TCE concentration occupational setting of cases, $18 \text{ mg/m}^3$ – $683 \text{ mg/m}^3$ 14 of 21 cases with U-TCE above recommended occupational level of 50 mg/L	Xu et al. (2009)
Five healthy workers engaged in decreasing activities in steel industry and five healthy workers from clerical section of same company	Total serum bile acid concentration increased between pre- and postexposure (2-d period)	8-h TWA mean personal air: $8.9 \pm 3.2$ ppm postexposure	Neghab et al. (1997)
22 workers at a factory manufacturing small appliances	Increased in several bile acids	Regular exposure to <5 ppm TCE; peak exposure for two workers to >250 00m	Driscoll et al. (1992)
4,489 males and female residents from 15 Superfund site and identified from ATSDR Trichloroethylene Exposure Subregistry	Liver problems diagnosed with past yr	Residency in community with Superfund site identified with TCE and other chemicals	Davis et al., 2005
Case reports from eight countries of individuals with idiosyncratic generalized skin disorders	Hepatitis in 46–94% of cases; other liver effects includes hepatomegaly and elevated liver function enzymes; and in rare cases, acute liver failure	If reported, TCE, from <50 mg/m <sup>3</sup> to more than 4,000 mg/m <sup>3</sup> . Symptoms developed within 2–5 wk of initial exposure, with some intervals up to 3 mo	Kamijima et al. (2007)

1

**Table 4-55. Summary of human liver toxicity studies (continued)**

Subjects	Effect	Exposure	Reference
Deaths in California between 1979–1981 due to cirrhosis	SMR of 211 (95% CI: 136, 287) for white male sheet metal workers and SMR = 174 (95% CI: 150–197) for metal workers	Occupational title on death certificate	Leigh and Jiang (1993)

1  
2 ALT = alanine aminotrasferase.  
3  
4

5 personal monitoring from other nonparticipating workers at this facility indicated TCE exposure  
6 as low, less than 5 ppm, with occasional peaks over 250 ppm although details are lacking  
7 whether these data represent exposures of study subjects.

8         Davis et al. 2005 in their analysis of subjects from the TCE subregistry of ATSDR’s  
9 National Exposure Registry examined the prevalence of subjects reporting liver problems  
10 (defined as seeking treatment for the problem from a physician within the past year) using rates  
11 for the equivalent health condition from the National Health Interview Survey (a nationwide  
12 multipurpose health survey conducted by the National Center for Health Statistics, Centers for  
13 Disease Control and Prevention). The TCE subregistry is a cohort of exposed persons from  
14 15 sites in 5 states. The shortest time interval from inclusion in the exposure registry and last  
15 follow-up was 5 years for one site and 10 years for seven sites. Excess in past-year liver  
16 disorders relative to the general population persisted for much of the lifetime of follow-up.  
17 SMRs for liver problems were 3<sup>rd</sup> follow-up, SMR = 2.23 (99% CI: 1.13, 3.92); 4<sup>th</sup> follow-up,  
18 SMR = 3.25 (99% CI: 1.82, 5.32); and, 5<sup>th</sup> follow-up, SMR = 2.82 (99% CI: 1.46, 4.89).  
19 Examination by TCE exposure, duration or cumulative exposure to multiple organic solvents did  
20 not show exposure-response patterns. Overall, these observations are suggestive of liver  
21 disorders as associated with potential TCE exposure, but whether TCE caused these conditions is  
22 not possible to determine given the study’s limitations. These limitations include a potential for  
23 misclassification bias, the direction of which could dampen observations in a negative direction,  
24 and lack of adjustment in statistical analyses for alcohol consumption, which could bias  
25 observations in a positive direction.

26         Evaluation in epidemiologic studies of risk factors for cirrhosis other than alcohol  
27 consumption and Hepatitis A, B, and C is quite limited. NRC (2006) cited a case report of  
28 cirrhosis developing in an individual exposed occupationally to TCE for 5 years from a  
29 hot-process degreaser and to 1,1,1-trichloroethane for 3 months thereafter (Thiele et al., 1982).  
30 One cohort study on cirrhosis deaths in California between 1979 and 1981 and occupational risk

1 factors as assessed using job title observed elevated risks with occupational titles of sheet metal  
2 workers and metalworkers and cirrhosis among white males who comprised the majority of  
3 deaths (Leigh and Jiang, 1993). This analysis lacks information on alcohol patterns by  
4 occupational title in addition to specific chemical exposures. Few deaths attributable to cirrhosis  
5 are reported for nonwhite male and for both white and nonwhite female metalworkers with  
6 analyses examining these individuals limited by low statistical power. Some but not all  
7 trichloroethylene mortality studies report risk ratios for cirrhosis (see Table 4-56). A statistically  
8 significant deficit in cirrhosis mortality is observed in three studies (Boice et al., 1999; Boice et  
9 al., 2006b; Morgan et al., 1998) and with risk ratios including a risk of 1.0 in the remaining  
10 studies (ATSDR, 2004; Blair et al., 1989; 1998; Garabrant et al., 1988; Ritz, 1999a). These  
11 results do not rule out an effect of TCE on liver cirrhosis since disease misclassification may  
12 partly explain observations. Available studies are based on death certificates where a high  
13 degree of underreporting, up to 50%, is known to occur (Blake et al., 1988).

14 A number of case reports exist of liver toxicity including hepatitis accompanying  
15 immune-related generalized skin diseases described as a variation of erythema multiforme,  
16 Stevens-Johnson syndrome, toxic epiderma necrolysis patients, and hypersensitivity syndrome  
17 (Section 4.6.1.2 describes these disorders and evidence on TCE) (Kamijima et al., 2007).  
18 Kamijima et al. (2007) reported hepatitis was seen in 92–94% of cases presenting with an  
19 immune-related generalized skin diseases of variation of erythema multiforme, Stevens-Johnson  
20 syndrome, and toxic epiderma necrolysis patients, but the estimates within the hypersensitivity  
21 syndrome group were more variable (46–94%). Many cases developed with a short time after  
22 initial exposure and presented with jaundice, hepatomegaly or hepatosplenomegaly, in addition,  
23 to hepatitis. Hepatitis development was of a nonviral etiology, as antibody titers for Hepatitis A,  
24 B, and C viruses were not detectable, and not associated with alcohol consumption (Huang et al.,  
25 2002; Kamijima et al., 2007). Liver failure was moreover a leading cause of death among these  
26 subjects. Kamijima et al. (2007) note the similarities between specific skin manifestations and  
27 accompanying hepatic toxicity and case presentations of TCE-related generalized skin diseases  
28 and conditions that have been linked to specific medications (e.g., carbamezepine, allupurinol,  
29 antibacterial sulfonamides), possibly in conjunction with reactivation of specific latent viruses.  
30 However, neither cytomegalovirus or Epstein-Barr viruses are implicated in the few reports  
31 which did include examination of viral antibodies.

32

#### 4.5.2. Liver Cancer in Humans

1 Primary hepatocellular carcinoma and cholangiocarcinoma (intrahepatic and extrahepatic  
2 bile ducts) are the most common primary hepatic neoplasms (Blechacz and Gores, 2008; El-  
3 Serag, 2007). Primary hepatocellular carcinoma is the 5<sup>th</sup> most common of cancer deaths in  
4 males and 9<sup>th</sup> in females (Jemal et al., 2008). Age-adjusted incidence rates of hepatocellular  
5 carcinoma (HCC) and intrahepatic cholangiocarcinoma (ICC) are increasing, with a twofold  
6 increase in HCC over the past 20 years. This increase is higher than expected from an expanded  
7 definition of liver cancer to include primary or secondary neoplasms since International  
8 Classification of Disease (ICD)-9, incorrect classification of hilar cholangiocarcinomas in ICD-O  
9 as ICC, or to improved detection methods (El-Serag, 2007). It is estimated that  
10 21,370 Americans will be diagnosed in 2008 with liver and intrahepatic bile cancer; age-adjusted  
11 incidence rates for liver and intrahepatic bile duct cancer for all races are 9.9 per 100,000 for  
12 males and 3.5 per 100,000 for females Ries et al., 2008. Survival for liver and biliary tract  
13 cancers remains poor and age-adjusted mortality rates are just slightly lower than incidence rates.  
14 While hepatitis B and C viruses and heavy alcohol consumption are believed major risk factors  
15 for HCC and intrahepatic cholangiocarcinoma, these risk factors cannot fully account for roughly  
16 10 and 20% of HCC cases Kulkarni et al., 2004. Cirrhosis is considered a premalignant  
17 condition for HCC, however, cirrhosis is not a sufficient cause for HCC since 10–25% of HCC  
18 cases lack evidence of cirrhosis at time of detection (Chiesa et al., 2000; Fattovich et al., 2004;  
19 Kumar et al., 2007). Nonalcoholic steatohepatitis reflecting obesity and metabolic syndrome is  
20 recently suggested as contributing to liver cancer risk (El-Serag, 2007).

21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

1  
2  
3  
4

*This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2  
3

**Table 4-56. Selected results from epidemiologic studies of TCE exposure and cirrhosis**

Study population	Exposure group	Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort and PMR-mortality</b>				
Aerospace workers (Rocketdyne)				
	Any TCE (utility/eng flush)	0.39 (0.16, 0.80)	7	Boice et al. (2006b)
	Low cumulative TCE score	Not reported		Zhao et al. (2005)
	Medium cumulative TCE score			
	High TCE score			
	<i>p</i> for trend			
View-master workers				
	Males	0.76 (0.16, 2.22)	3	ATSDR (2003a)
	Females	1.51 (0.72, 2.78)	10	
Electronic workers (Taiwan)				
	Primary liver, males	Not reported		Chang et al. (2003; 2005)
	Primary liver, females	Not reported		
Uranium-processing workers				
	Any TCE exposure	0.91 (0.63, 1.28)	33	Ritz (1999a)
	Light TCE exposure, >2 yr duration	Not reported		
	Mod TCE exposure, >2 yr duration	Not reported		
Aerospace workers (Lockheed)				
	TCE routine exposure	0.61 (0.39, 0.91)	23	Boice et al. (1999)
	TCE routine-intermittent	Not reported	13	
Aerospace workers (Hughes)				
	TCE subcohort	0.55 (0.30, 0.93)	14	Morgan et al. (1998, 2000b)
	Low intensity (<50 ppm)	0.95 (0.43, 1.80)	9	
	High intensity (>50 ppm)	0.32 (0.10, 0.74)	5	
Aircraft maintenance workers (Hill AFB, Utah)				
	TCE subcohort	1.1 (0.6, 1.9) <sup>a</sup>	44	Blair et al. (1998)
	Males, cumulative exposure			
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	0.6 (0.2, 1.3)	10	
	5–25 ppm-yr	0.8 (0.3, 1.9)	9	
	>25 ppm-yr	1.2 (0.6, 2.4)	17	

4

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-56. Selected results from epidemiologic studies of TCE exposure and cirrhosis (continued)**

Study population	Exposure group	Relative risk (95% CI)	No. obs. events	Reference	
Aircraft maintenance workers (continued)	Females, cumulative exposure				
	0	1.0 <sup>a</sup>		Blair et al. (1998) (continued)	
	<5 ppm-yr	2.4 (1.4, 13.7)	6		
	5–25 ppm-yr	1.8 (0.2, 15.0)	1		
	>25 ppm-yr	0.6 (0.1, 4.8)	1		
	TCE subcohort				Radican et al. (2008)
	Males, cumulative exposure		1.04 (0.56, 1.93) <sup>a, b</sup>	37	
	0	0.87 (0.43, 1.73)	31		
	0	1.0 <sup>a, b</sup>			
	<5 ppm-yr	0.56 (0.23, 1.40)	8		
	5–25 ppm-yr	1.07 (0.45, 2.53)	10		
	>25 ppm-yr	1.06 (0.48, 2.38)	13		
	Females, cumulative exposure		1.79 (0.54, 5.93)	6	
	0	1.00 <sup>a</sup>			
<5 ppm-yr	3.30 (0.88, 12.41)	4			
5–25 ppm-yr	2.20 (0.26, 18.89)	1			
>25 ppm-yr	0.59 (0.97, 5.10)	1			
Deaths reported to GE pension fund (Pittsfield, MA)		Not reported		Greenland et al. (1994)	
U.S. Coast Guard employees				Blair et al. (1989)	
Marine inspectors		1.36 (0.79, 2.17)	17		
Noninspectors		0.53 (0.23, 1.05)	8		
Aircraft manufacturing plant employees (Italy)				Costa et al. (1989)	
All subjects		Not reported			
Aircraft manufacturing plant employees (San Diego, CA)				Garabrant et al. (1988)	
All subjects		0.86 (0.67, 1.11)	63		

<sup>a</sup> Referent group are subjects from the same plant or company, or internal referents.

<sup>b</sup> Numbers of cirrhosis deaths in Radican et al. (2008) are fewer than Blair et al. (1989) because Radican et al. (2008) excluded cirrhosis deaths due to alcohol.

GE = General Electric.

1  
2  
3  
4  
5  
6  
7  
8  
9

1  
2 All cohort studies, except Zhao et al. (2005), present risk ratios (SIRs or SMRs) for liver  
3 and biliary tract cancer. More rarely reported in cohort studies are risk ratios for primary liver  
4 cancer (HCC) or for gallbladder and extrahepatic bile duct cancer. Four community studies also  
5 presented risk ratios for liver and biliary tract cancer including a case-control study of primary  
6 liver cancer of residents of Taiwanese community with solvent-contaminated drinking water  
7 wells (ATSDR, 2006a; Lee et al., 2003; Morgan and Cassady, 2002; Vartiainen et al., 1993).  
8 Several population case-control studies examine liver cancer and organic solvents or  
9 occupational job titles with possible TCE usage (Austin et al., 1987; Døssing et al., 1997;  
10 Hardell et al., 1984; Heinemann et al., 2000; Hernberg et al., 1988; 1984; Ji and Hemminki,  
11 2005; Kvam et al., 2005; Lindbohm et al., 2009; Porru et al., 2001; Stemhagen et al., 1983;  
12 Weiderpass et al., 2003); however, the lack of detailed exposure assessment to TCE, specifically  
13 in the population case-control studies as well as in geographic-based studies, or, too few exposed  
14 cases and controls in those studies that do present some information limits their usefulness for  
15 evaluating hepatobiliary or gall bladder cancer and TCE exposure. Table 4-57 presents  
16 observations from cohort, case-control, and community studies on liver and biliary tract cancer,  
17 primary liver, and gallbladder and extrahepatic bile duct cancer and trichloroethylene.

18 Excess liver cancer incidence is observed in most studies in which there is a high  
19 likelihood of TCE exposure in individual study subjects (e.g., based on job-exposure matrices or  
20 biomarker monitoring) and which met, to a sufficient degree, the standards of epidemiologic  
21 design and analysis were identified (Anttila et al., 1995; Axelson et al., 1994; Hansen et al.,  
22 2001; Raaschou-Nielsen et al., 2003) as is mortality (ATSDR, 2004; Blair et al., 1998; Boice et  
23 al., 2006b; Morgan et al., 1998; 2008; Ritz, 1999a). Risks for primary liver cancer and for  
24 gallbladder and biliary tract cancers in females were statistically significantly elevated only in  
25 Raaschou-Nielsen et al. (2003), the study with the largest number of observed cases without  
26 suggestion of exposure duration-response patterns. Cohort studies with more uncertain exposure  
27 assessment approaches, e.g., studies of all subjects working at a factory (Blair et al., 1989; Chang  
28 et al., 2003; Chang et al., 2005; Costa et al., 1989; Garabrant et al., 1988), do not show  
29 association but are quite limited given their lacking attribution of who may have higher or lower  
30 exposure potentials. Ritz (1999a), the exception, found evidence of an exposure-response  
31 relationship; mortality from hepatobiliary cancer was found to increase with



**Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer**

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts		Reference
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	
<b>Cohort and PMR studies—incidence</b>								
Aerospace workers (Rocketdyne)								
	Low cumulative TCE score	Not reported						Zhao et al. (2005)
	Medium cumulative TCE score	Not reported						
	High TCE score	Not reported						
	<i>p</i> for trend							
Danish blue-collar workers with TCE exposure								
	Males + females	1.3 (1.0, 1.6) <sup>a</sup>	82					Raaschou-Nielsen et al. (2003)
	Males + females	1.4 (1.0, 1.8) <sup>b</sup>	57					
	Males, any exposure	1.1 (0.8, 1.5) <sup>b</sup>	41	1.1 (0.7, 1.6)	27	1.1 (0.6, 1.9)	14	
	<1 yr employment duration	1.2 (0.7, 2.1) <sup>b</sup>	13	1.3 (0.6, 2.5)	9	1.1 (0.3, 2.9)	4	
	1–4.9 yr employment duration	0.9 (0.5, 1.6) <sup>b</sup>	13	1.0 (0.5, 1.9)	9	0.8 (0.2, 2.1)	4	
	≥5 yr employment duration	1.1 (0.6, 1.7) <sup>b</sup>	15	1.1 (0.5, 2.1)	9	1.4 (0.5, 3.1)	6	
	Females, any exposure	2.8 (1.6, 4.6) <sup>b</sup>	16	2.8 (1.1, 5.8)	7	2.8 (1.3, 5.3)	9	
	<1 yr employment duration	2.5 (0.7, 6.5) <sup>b</sup>	4	2.8 (0.3, 10.0)	2	2.3 (0.3, 8.4)	2	
	1–4.9 yr employment duration	4.5 (2.2, 8.3) <sup>b</sup>	10	4.1 (1.1, 10.5)	4	4.8 (1.7, 10.4)	6	
	≥5 yr employment duration	1.1 (0.1, 3.8) <sup>b</sup>	2	1.3 (0.0, 7.1)	1	0.9 (0.0, 5.2)	1	

**Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)**

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts		Reference
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	
Biologically-monitored Danish workers								
	Males + females	2.1 (0.7, 5.0) <sup>b</sup>	5	1.7 (0.2, 6.0)	2	2.5 (0.5, 7.3)	3	Hansen et al. (2001)
	Males	2.6 (0.8, 6.0) <sup>b</sup>	5	1.8 (0.2, 6.6)	2	3.3 (0.7, 9.7)	3	
	Females		0 (0.4 exp)		0 (0.1 exp)		0 (0.3 exp)	
	Cumulative exposure (Ikeda)	Not reported						
	<17 ppm-yr							
	≥17 ppm-yr							
	Mean concentration (Ikeda)	Not reported						
	<4 ppm							
	4+ ppm							
	Employment duration	Not reported						
	<6.25 yr							
	≥6.25							

Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts			
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Reference	
Aircraft maintenance workers from Hill Air Force Base									
	TCE subcohort	Not reported	9	Not reported				Blair et al. (1998)	
	Males, cumulative exposure								
	0	1.0 <sup>c</sup>		1.03					
	<5 ppm-yr	0.6 (0.1, 3.1)	3	1.2 (0.1, 2.1)	2				
	5–25 ppm-yr	0.6 (0.1, 3.8)	2	1.0 (0.1, 16.7)	1				
	>25 ppm-yr	1.1 (0.2, 4.8)	4	2.6 (0.3, 25.0)	3				
	Females, cumulative exposure		0		0				
Biologically-monitored Finnish workers									
	All subjects	1.89 (0.86, 3.59) <sup>b</sup>	9	2.27 (0.74, 5.29)	5	1.56 (0.43, 4.00)	4	Anttila et al. (1995)	
	Mean air-TCE (Ikeda extrapolation from U-TCA)								
	<6 ppm	Not reported		1.64 (0.20, 5.92)	2				
	6+ ppm			2.74 (0.33, 9.88)	2				
Biologically-monitored Swedish workers									
	Males	1.41 (0.38, 3.60) <sup>b</sup>	4					Axelson et al. (1994)	
	Females	Not reported							
<b>Cohort and PMR-mortality</b>									
	Computer manufacturing workers (IBM), NY	Not reported	1					Clapp and Hoffman (2008)	

**Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)**

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts		
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Reference
Aerospace workers (Rocketdyne)								
	Any TCE (utility/eng flush)	1.28 (0.35, 3.27)	4					Boice et al. (2006b)
	Low cumulative TCE score	Not reported						Zhao et al. (2005)
	Med cumulative TCE score							
	High TCE score							
	<i>p</i> for trend							
View-Master workers								
	Males	2.45 (0.50, 7.12) <sup>d</sup>	3	1.01 (0.03, 5.63) <sup>d</sup>	1	8.41 (1.01, 30.4) <sup>d</sup>	2	ATSDR (2003a)
	Females		0 (2.61 exp)		0 (1.66 exp)		0 (0.95 exp)	
Electronic workers (Taiwan)								
	Primary liver, males	Not reported			0 (0.69 exp)			Chang et al. (2003; 2005)
	Primary liver, females	Not reported			0 (0.57 exp)			

Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts			
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Reference	
Uranium-processing workers									
	Any TCE exposure	Not reported						Ritz (1999a)	
	Light TCE exposure, >2 yr duration	0.93 (0.19, 4.53) <sup>c</sup>	3						
	Mod TCE exposure, >2 yr duration	4.97 (0.48, 51.1) <sup>e</sup>	1						
	Light TCE exposure, >5 yr duration	2.86 (0.48, 17.3) <sup>f</sup>	3						
	Mod TCE exposure, >5 yr duration	12.1 (1.03, 144) <sup>f</sup>	1						
Aerospace workers (Lockheed)									
	TCE routine exposure	0.54 (0.15, 1.38)	4					Boice et al. (1999)	
	TCE routine-intermittent								
	0 yr	1.00 <sup>c</sup>	22						
	Any exposure	Not reported	13						
	<1 yr	0.53 (0.18, 1.60)	4						
	1–4 yr	0.52 (0.15, 1.79)	3						
	≥5 yr	0.94 (0.36, 2.46)	6						
	<i>p</i> for trend	>0.20							

**Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)**

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts			
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Reference	
Aerospace workers (Hughes)									
	TCE subcohort	0.98 (0.36, 2.13)	6					Morgan et al. (1998, 2000b)	
	Low intensity (<50 ppm) <sup>c</sup>	1.32 (0.27, 3.85)	3						
	High intensity (>50 ppm) <sup>c</sup>	0.78 (0.16, 2.28)	3						
	TCE subcohort (Cox analysis)								
	Never exposed	1.00 <sup>c</sup>	14						
	Ever exposed	1.48 (0.56, 3.91) <sup>g,h</sup>	6						
	Cumulative								
	Low	2.12 (0.59, 7.66) <sup>h</sup>	3						
	High	1.19 (0.34, 4.16) <sup>h</sup>	3						
	Peak								
	No/low	1.00 <sup>c</sup>	17						
	Medium/high	0.98 (0.29, 3.35) <sup>h</sup>	3						

Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts		Reference
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	
Aircraft maintenance workers (Hill AFB, Utah)								Blair et al. (1998)
TCE subcohort		1.3 (0.5, 3.4) <sup>c</sup>	15	1.7 (0.2, 16.2) <sup>3</sup>	4			
Males, cumulative exposure								
0		1.0 <sup>c</sup>						
<5 ppm-yr		1.1 (0.3, 4.1)	6					
5–25 ppm-yr		0.9 (0.2, 4.3)	3					
>25 ppm-yr		0.7 (0.2, 3.2)	3					
Females, cumulative exposure								
0		1.0 <sup>c</sup>						
<5 ppm-yr		1.6 (0.2, 18.2)	1					
5–25 ppm-yr			0					
>25 ppm-yr		2.3 (0.3, 16.7)	2					
TCE subcohort		1.12 (0.57, 2.19) <sup>c,i</sup>	31	1.25 (0.31, 4.97) <sup>c,i</sup>	8			Radican et al. (2008)
Males, cumulative exposure								
0		1.0 <sup>c</sup>		1.03				
<5 ppm-yr		1.17 (0.45, 3.09)	10	3.28 (0.37, 29.45)	4			
5–25 ppm-yr		1.16 (0.39, 3.46)	6		0			
>25 ppm-yr		1.72 (0.68, 4.38)	12	4.05 (0.45, 36.41)	4			

Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts		
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Reference
Aircraft maintenance workers (continued)	Females, cumulative exposure	0.74 (0.18, 2.97) <sup>c</sup>	3		0			Radican et al. (2008) (continued)
	0	1.03						
	<5 ppm-yr	0.69 (0.08, 5.74)	1					
	5–25 ppm-yr		0					
	>25 ppm-yr	0.98 (0.20, 4.90)	2					
Deaths reported to GE pension fund (Pittsfield, MA)		0.54 (0.11, 2.63) <sup>j</sup>	9					Greenland et al. (1994)
U.S. Coast Guard employees								
	Marine inspectors	1.12 (0.23, 3.26)	3					Blair et al. (1989)
	Noninspectors	Not reported	0 (2 exp)					
Aircraft manufacturing plant employees (Italy)								
	All subjects	0.70 (0.23, 1.64)	5					Costa et al. (1989)
Aircraft manufacturing plant employees (San Diego, CA)								
	All subjects	0.94 (0.40, 1.86)	8					Garabrant et al. (1988)
<b>Case-control studies</b>								
Residents of community with contaminated drinking water (Taiwan)								
	Village of residency, males							Lee et al. (2003)
	Upstream	1.00						
	Downstream	2.57 (1.21, 5.46)	26					



**Table 4-57. Selected results from epidemiologic studies of TCE exposure and liver cancer (continued)**

Study population	Exposure group	Liver and intrahepatic bile ducts		Primary liver		Gallbladder and extrahepatic bile ducts		
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Reference
<b>Geographic studies</b>								
Residents in two study areas in Endicott, NY		0.71 (0.09, 2.56)	<6					ATSDR (2006b)
Residents in 13 census tracts in Redlands, CA		1.29 (0.74, 2.05) <sup>k</sup>	28					Morgan and Cassidy (2002)
<b>Finnish residents</b>								
	Residents of Hausjarvi	0.76 (0.3, 1.4)	7					Vartiainen et al. (1993)
	Residents of Huttula	0.6 (0.2, 1.3)	6					

<sup>a</sup> ICD-7, 155 and 156; Primary liver (155.0), gallbladder, and biliary passages (155.1), and liver secondary and unspecified (156).

<sup>b</sup> ICD-7, 155; Primary liver, gallbladder, and biliary passages.

<sup>c</sup> Internal referents, workers without TCE exposure.

<sup>d</sup> Proportional mortality ratio (PMR).

<sup>e</sup> Logistic regression analysis with a 0-yr lag for TCE exposure.

<sup>f</sup> Logistic regression analysis with a 15-yr lag for TCE exposure.

<sup>g</sup> Risk ratio from Cox Proportional Hazard Analysis, stratified by age, sex, and decade Environmental Health Strategies (1997).

<sup>h</sup> Morgan et al. (1998) do not identify if SIR is for liver and biliary passage or primary liver cancer; identified as primary liver in NRC (2006).

<sup>l</sup> Radican et al. (2008) provide results for TCE exposure for follow-up through 1990, comparing the Poisson model rate ratios as reported by Blair et al. (1998) with Cox model hazard ratios. Relative risk from Cox model adjusted for age and gender for liver and intrahepatic bile duct cancer was 1.2 (95% CI: 0.5, 3.4) and for primary liver cancer was 1.3 (95% CI: 0.1, 12.0).

<sup>j</sup> Odds ratio.

<sup>k</sup> 99% CIs.


exp = exposures, GE = General Electric, IBM = International Business Machines Corporation.

1 degree and duration of exposure and time since first exposure with a statistically significant but  
2 imprecise (wide confidence intervals) liver cancer risk for those with the highest exposure and  
3 longest time since first exposure. This observation is consistent with association with TCE, but  
4 with uncertainty given one TCE exposed case in the highest exposure group and correlation  
5 between TCE, cutting fluids, and radiation exposures.

6 Observations in these studies provide some evidence of susceptibility of liver, gallbladder  
7 and biliary tract; observations consistent with pharmacokinetic processing of TCE and the  
8 extensive intra- and extrahepatic recirculation of metabolites. Magnitude of risk of gallbladder  
9 and biliary tract cancer is slightly higher than that for primary liver cancer in Raaschou-Nielsen  
10 et al. (2003) the study with the most cases. Observations in Blair et al. (1998), Hansen et al.  
11 (2001), and Radican et al. (2008), three smaller studies, suggest slightly larger risk ratios for  
12 primary liver cancer compared to gallbladder and biliary tract cancer. Overall, these studies are  
13 not highly informative for cross-organ comparison of relative magnitude of susceptibility.

14 The largest geographic studies (Lee et al., 2003; Morgan and Cassady, 2002) are also  
15 suggestive of association with the risk ratio (mortality odds ratio) in Lee et al. (2003) as  
16 statistically significantly elevated. The geographic studies do not include a characterization of  
17 TCE exposure to individual subjects other than residency in a community with groundwater  
18 contamination by TCE with potential for exposure misclassification bias dampening  
19 observations; these studies lack characterization of TCE concentrations in drinking water and  
20 exposure characteristics such as individual consumption patterns. For this reason, observations  
21 in Morgan and Cassidy (2002) and Lee et al. (2003) are noteworthy, particularly if positive bias  
22 leading to false positive finding is considered minimal, and the lack of association with liver  
23 cancer in the two other community studies (ATSDR, 2006b; Vartiainen et al., 1993) does not  
24 detract from Morgan and Cassidy (2002) or Lee et al. (2003). Lee et al. (2003), however, do not  
25 address possible confounding related to hepatitis viral infection status, a risk factor for liver  
26 cancer, or potential misclassification due to the inclusion of secondary liver cancer among the  
27 case series, factors which may amplify observed association.

28 Meta-analysis is adopted as a tool for examining the body of epidemiologic evidence on  
29 liver cancer and TCE exposure, to identify possible sources of heterogeneity and as an additional  
30 means to identify cancer hazard. The meta-analyses of the overall effect of TCE exposure on  
31 liver (and gall bladder/biliary passages) cancer suggest a small, statistically significant increase  
32 in risk. The summary estimate from the primary random effects meta-analysis of the 9 (all  
33 cohort) studies is 1.29 (95% CI: 1.07, 1.56) (see Figure 4-3). The study of Raaschou-Nielsen  
34 et al. (2003) contributes about 57% of the weight; its removal from the analysis decreases  
35 somewhat the RR<sub>m</sub> estimate and is no longer statistically significant (RR<sub>m</sub> = 1.22; 95% CI:




**Figure 4-3. Relative risk estimates of liver and biliary tract cancer and overall TCE exposure.** Random effects model; fixed effect model same. The summary estimate is in the bottom row, represented by the diamond. Symbol sizes reflect relative weights of the studies.

0.93, 1.61). The summary estimate was not overly influenced by any other single study, nor was it overly sensitive to individual RR estimate selections. There is no evidence of publication bias in this data set, and no observable heterogeneity ( $I^2 = 0\%$ ) across the study results.

Examination of sites individually (i.e., primary liver and intrahepatic bile ducts separate from the combined liver and gallbladder/biliary passage grouping) resulted in the RRm estimate for liver cancer alone (for the three studies for which the data are available; for the other studies, results for the combined grouping were used) slightly lower than the one based entirely on results from the combined cancer categories and was just short of statistical significance (1.25; 95% CI: 0.99, 1.57). This result is driven by the fact that the risk ratio estimate from the large Raaschou-Nielsen et al. (2003) study decreased from 1.35 for liver and gall bladder/biliary passage cancers combined to 1.28 for liver cancer alone.

The RRm estimate from the random effects meta-analysis of liver cancer in the highest exposure groups in the six studies which provide risk estimates associated with highest exposure.



1

**Figure 4-4. Meta-analysis of liver cancer and TCE exposure—highest exposure groups, with assumed null RR estimates for Hansen and Zhao (see Appendix C text).** Random effects model; fixed effect model same. The summary estimate is in the bottom row, represented by the diamond. Symbol sizes reflect relative weights of the studies.

2

3

4 primary liver cancer is 1.32 (95% CI: 0.93, 1.86), slightly lower than the RRM estimate for liver  
 5 and gallbladder/biliary cancer and any TCE exposure of 1.33 (95% CI: 1.09, 1.64), and not  
 6 statistically significant (see Figure 4-4). Again, the RRM estimate of the highest-exposure  
 7 groups is dominated by one study (Raaschou-Nielsen et al., 2003). Two studies lack reporting of  
 8 liver cancer risk associated with highest exposure, so consideration of reporting bias (considered  
 9 the primary analysis) lead to a result of 1.28 (95% CI: 0.93, 1.77), similar to that estimated in the  
 10 more restricted set of studies presenting risk ratios association with highest exposure groups in  
 11 published papers.

12

13

14

15

Different exposure metrics are used in the various studies, and the purpose of combining results across the different highest exposure groups is not to estimate an RRM associated with some level of exposure, but rather to examine impacts of combining RR estimates that should be less affected by exposure misclassification. In other words, the highest exposure category is

1 more likely to represent a greater differential TCE exposure compared to people in the referent  
2 group than the exposure differential for the overall (typically any vs. none) exposure comparison.  
3 Thus, if TCE exposure increases the risk of liver and gallbladder/biliary cancer, the effects  
4 should be more apparent in the highest exposure groups. The findings of a lower RR<sub>m</sub>  
5 associated with highest exposure group reflects observations in Radican et al. (2008) and  
6 Raaschou-Nielsen et al. (2003), the study contributing greatest weight to the meta-analysis, that  
7 RR estimates for the highest-exposure groups, although greater than 1.0, are less than the RR  
8 estimates with any TCE exposure.

9 Thus, while the finding of an elevated and statistically significant RR<sub>m</sub> for liver and  
10 gallbladder/biliary cancer and any TCE exposure provides evidence of association, the statistical  
11 significance of the summary estimates is dependent on one study, which provides the majority of  
12 the weight in the meta-analyses. Furthermore, combining results from the highest-exposure  
13 groups yields lower RR<sub>m</sub> estimates than for an overall effect. These results do not rule out an  
14 effect of TCE on liver cancer, because the liver cancer results are relatively underpowered with  
15 respect to numbers of studies and number of cases; overall, the meta-analysis provides only  
16 minimal support for association between TCE exposure and liver and gallbladder/biliary cancer.

17 NRC (2006) deliberations on trichloroethylene commented on two prominent evaluations  
18 of the then-current TCE epidemiologic literature using meta-analysis techniques, Wartenberg  
19 et al. (2000) and Kelsh et al. (2005), submitted by Exponent-Health Sciences to NRC during  
20 their deliberations and published afterwards in the open literature as Alexander et al. (2007a)  
21 adding the then-published study of Boice et al. (2006b). NRC (2006) found weaknesses in the  
22 techniques used in Wartenberg et al. (2000) and the Exponent analyses. EPA staff conducted  
23 their analysis according to NRC (2006) suggestions for transparency, systematic review criteria,  
24 and examination of both cohort and case-control studies. The EPA analysis of liver cancer  
25 considered a similar set of studies as Alexander et al. (2007a) although treatment of these studies  
26 differs between analyses. Alexander et al. (2007a) in their Table 2, for example, present  
27 summary relative risk estimates, grouping of studies with differing exposure potentials, for  
28 example, including liver and biliary cancer risk estimates for all subjects, those exposed and  
29 unexposed to TCE, in Boice et al. (1999), Blair et al. (1998), Morgan et al. (1998) and Boice  
30 et al. (2006b), with biomarker studies (Anttila et al., 1995; Axelson et al., 1994; Hansen et al.,  
31 2001). The inclusion of risk estimates for subjects who have little to no TCE exposure over  
32 background levels has the potential to introduce misclassification bias and dampen observed risk  
33 ratios. Potential bias from exposure misclassification may be substantial in Alexander et al. (2007a)  
34 since the percentage of TCE exposed subjects to all cohort subjects in the four studies was 3, 23,  
35 51 and 68% in Boice et al. (1999), Morgan et al. (1998), Blair et al. (1998) and Boice et al.

1 (2006b), respectively, and is a likely alternative explanation for observed inconsistency across  
2 occupational groups reported by the authors. Another difference between the EPA and previous  
3 meta-analyses is their treatment of Ritz (1999a), included in Wartenberg et al. (2000), Kelsh  
4 et al. (2005), and Alexander et al. (2007a), but not in this analysis. For a grouping of studies  
5 with subcohorts most similar to those in EPA's analysis, summary liver and gall bladder/biliary  
6 tract cancer risk estimates for overall TCE exposure for TCE subcohorts is of a similar  
7 magnitude as that observed in EPA's updated and expanded analysis, Wartenberg et al. (2000),  
8 1.1 (95% CI: 0.3, 4.8) for incidence and 1.1 (95% CI: 0.7, 1.7) for mortality, Kelsh et al. (2005),  
9 1.32 (95% CI: 1.05, 1.66) and Alexander et al. (2007a), 1.30 (95% CI: 1.09–1.55).

#### 4.5.3. Experimental Studies of Trichloroethylene (TCE) in Rodents—Introduction

12 The previous sections have described available human data for TCE-induced noncancer  
13 effects (e.g., disturbances in bile production) and whether an increased risk of liver cancer in  
14 humans has been established from analysis of the epidemiological literature. A primary concern  
15 for effects on the liver comes from a large database in rodents indicating that, not only TCE, but  
16 a number of its metabolites are capable of inducing hepatocellular adenomas and carcinomas in  
17 rodent species. Thus, many of rodent bioassays have focused on the study of liver cancer for  
18 TCE and its metabolites and possible early effects specifically that may be related to tumor  
19 induction.

20 This section describes the hazard data for TCE effects in the rodent liver and inferences  
21 from studies of its metabolites. For more detailed descriptions of the issues providing context for  
22 these data in terms the state of the science of liver physiology (see Section E.1), cancer (see  
23 Section E.3), liver cancer (see Section E.3), and the MOA of liver cancer and other TCE-induced  
24 effects (see Section E.3.4), please see Appendix E. A more comprehensive review of individual  
25 studies of TCE-induced liver effects in laboratory animals is also provided in Section E.2 that  
26 includes detailed analyses of the strengths and the limitations of these studies. Issues have been  
27 raised regarding the relevance of mouse liver tumor data to human liver cancer risk that are  
28 addressed in Sections E.3.2 and E.3.3. Given that activation of the PPAR $\alpha$  receptor has received  
29 great attention as a potential MOA for TCE induced liver tumors, the current status of that  
30 hypothesis is reviewed in Section E.3.4.1. Finally, comparative studies of TCE metabolites and  
31 the similarities and differences of such study results are described in summary sections of  
32 Appendix E (i.e., Section E.2.4) as well as discussions of proposed MOAs for TCE-induced liver  
33 cancer (i.e., Sections E.2.4 and E.3.4.2).

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 A number of acute and subchronic studies have been undertaken to describe the early  
2 changes in the rodent liver after TCE administration with the majority using the oral gavage  
3 route of administration. Several key issues affect the interpretation of these data. The few  
4 drinking water studies available for TCE have recorded significant loss of TCE through  
5 volatilization in drinking water solutions and thus, this route of administration is generally not  
6 used. Some short-term studies of TCE have included detailed examinations while others have  
7 reported primarily liver weight changes as a marker of TCE response. The matching and  
8 recording of age, but especially initial and final body weight, for control and treatment groups is  
9 of particular importance for studies using liver weight gain as a measure of TCE response as  
10 differences in these parameters affect TCE-induced liver weight gain. Most data are for TCE  
11 exposures of at least 10 days to 42 days. For many of the subchronic inhalation studies  
12 (Kjellstrand et al., 1981, 1983a, b), issues associated with whole-body exposures make  
13 determination of dose levels more difficult. The focus of the long-term studies of TCE is  
14 primarily detection and characterization of liver tumor formation.

15 For gavage experiments, death due to gavage errors and specifically from use of this  
16 route of administration, especially at higher TCE exposure concentrations, has been a recurring  
17 problem, especially in rats. Unlike inhalation exposures, the effects of vehicle can also be an  
18 issue for background liver effects in gavage studies. Concerns regarding effects of oil vehicles,  
19 especially corn oil, have been raised (Charbonneau et al., 1991; Kim et al., 1990a). Several oral  
20 studies in particular document that use of corn oil as the vehicle for TCE gavage dosing induces  
21 a different pattern of toxicity, especially in male rodents (see Merrick et al., 1989, Section  
22 E.2.2.1). Several studies also report the effects of corn oil on hepatocellular DNA synthesis and  
23 indices of lipid peroxidation (Channel et al., 1998; Rusyn et al., 1999). For example, Rusyn  
24 et al. (1999) report that a single dose of dietary corn oil increases hepatocyte DNA synthesis 24  
25 hours after treatment by ~3.5-fold of control, activates of NF- $\kappa$ B to a similar extent ~2 hours  
26 after treatment almost exclusively in Kupffer cells, and induces an ~3–fourfold increase of  
27 control NF- $\kappa$ B in hepatocytes after 8 hours and an increase in tumor necrosis factor (TNF)- $\alpha$   
28 mRNA between 8 and 24 hours after a single dose in female rats.

29 In regard to studies that have used the i.p. route of administration, as noted by  
30 Kawamoto et al. (1988b), injection of TCE may result in paralytic ileus and peritonitis and that  
31 subcutaneous treatment paradigm will result in TCE not immediately being metabolized but  
32 retained in the fatty tissue. Wang and Stacey (1990) state that “intraperitoneal injection is not  
33 particularly relevant to humans” and suggest that intestinal interactions require consideration in  
34 responses such as increase serum bile acid.

1 While studies of TCE metabolites have been almost exclusively conducted via drinking  
2 water, and thus, have avoided vehicle effects and gavage error, they have issues of palatability at  
3 high doses and decreased drinking water consumption as a result that not only raises issues of the  
4 resulting internal dose of the agent but also of effects of drinking water reduction.

5 Although there are data for both mice and rats for TCE exposure and studies of its  
6 metabolites, the majority of the available information has been conducted in mice. This is  
7 especially the case for long-term studies of DCA and TCA in rats. There is currently one study  
8 each available for TCA and DCA in rats and both were conducted with such few numbers of  
9 animals that the ability to detect and discern whether there was a treatment-related effect are very  
10 limited (DeAngelo et al., 1996, 1997; Richmond et al., 1995).

11 With regard to the sensitivity of studies used to detect a response, there are issues  
12 regarding not only the number of animals used but also the strain and weight of the animals. For  
13 some studies of TCE strains were used that have less background rate of liver tumor  
14 development and carcinogenic response. As for the B6C3F1 mouse, the strain most used in the  
15 bioassays of TCE metabolites, the susceptibility of the B6C3F1 to hepatocarcinogenicity has  
16 made the strain a sensitive biomarker for a variety of hepatocarcinogens. Moreover, Leakey  
17 et al. (2003b) demonstrated that increased body weight at 45 weeks of life is an accurate  
18 predictor of large background tumor rates. Unfortunately a 2-year study of chloral hydrate  
19 (George et al., 2000) and the only available 2-year study of TCA (DeAngelo et al., 2008), which  
20 used the same control animals, were both conducted in B6C3F1 mice that grew very large  
21 (~50 g) and prone to liver cancer (64% background incidence of hepatocellular adenomas and  
22 carcinomas) and premature mortality. Thus, these bioassays are of limited value for  
23 determination of the dose-response for carcinogenicity.

24 Finally, as discussed below, the administration of TCE to laboratory animals as well as  
25 environmental exposure of TCE in humans are effectively coexposure studies. TCE is  
26 metabolized to a number of hepatoactive as well as hepatocarcinogenic agents. A greater  
27 variability of response is expected than from exposure to a single agent making it particularly  
28 important to look at the TCE database in a holistic fashion rather than the results of a single  
29 study, especially for quantitative inferences. This approach is particularly useful given that the  
30 number of animals in treatment groups in a variety of TCE and TCE metabolite studies have  
31 been variable and small for control and treatment groups. Thus, their statistical power was not  
32 only limited for detection of statistically significant changes but also in many cases to be able to  
33 determine whether there is not a treatment related effect (i.e., Type II error for power  
34 calculation). Section E.2.4.2 provides detailed analyses of the database for liver weight  
35 induction by TCE and its metabolites in mice and the results of those analyses are described



1 below. Specifically, the relationship of liver weight induction, but also other endpoints such as  
2 peroxisomal enzyme activation and increases in DNA synthesis to liver tumor responses are also  
3 addressed as well.  
4

#### 4.5.4. Trichloroethylene (TCE)-Induced Liver Noncancer Effects

5 A number of effects have been studied as indicators of TCE effects on the liver but also  
6 as proposed events whose sequellae could be associated with resultant liver tumors after chronic  
7 TCE exposure in rodents. Similar effects have been studied in rodents exposed to TCE  
8 metabolites which may be useful for not only determining whether such effects are associated  
9 with liver tumors induced by these metabolites but also if they are similar to what has been  
10 observed for TCE. Summaries of the laboratory animal studies of TCE noncancer effects in the  
11 liver are provided in Table 4-58 (oral studies) and Table 4-59 (inhalation studies), along with the  
12 types of effects discussed in the subsections below for each study.  
13

##### 4.5.4.1.1. Liver Weight

14 Increases in liver weight in mice, rats, and gerbils have been reported as a result of acute  
15 and short-term, and subchronic TCE treatment by inhalation and oral routes of exposure (Berman  
16 et al., 1995; Buben and O'Flaherty, 1985; Dees and Travis, 1993; Elcombe et al., 1985; Goel et  
17 al., 1992; Goldsworthy and Popp, 1987; Kjellstrand et al., 1983a; Kjellstrand et al., 1983b;  
18 Kjellstrand et al., 1981b; Laughter et al., 2004; Melnick et al., 1987; Merrick et al., 1989;  
19 Nakajima et al., 2000; Nunes et al., 2001; Tao et al., 2000a; Tucker et al., 1982). The extent of  
20 TCE-induced liver weight gain is dependent on species, strain, gender, nutrition status, duration  
21 of exposure, route of administration, vehicle used in oral studies, and the concentration of TCE  
22 administered. Of great importance to the determination of the magnitude of response is whether  
23 the dose of TCE administered also affects whole-body weight, and thus, liver weight and the  
24 percentage liver/body weight ratio. Therefore, studies which employed high enough doses to  
25 induce whole-body weight loss generally showed a corresponding decrease in percentage  
26 liver/body weight at such doses and "flattening" of the dose-response curve, while studies which  
27 did not show systemic toxicity reported liver/body weight ratios generally proportional to dose.  
28 Chronic studies, carried out for longer durations, that examine liver weight are few and often  
29 confounded by the presence of preneoplastic foci or tumors that also affect liver weight after an  
30 extended period of TCE exposure. The number of studies that examine liver weight changes in  
31 the rat are much fewer than for mouse. Overall, the database for mice provides data for

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 examination of the differences in TCE-induced effects from differing exposure levels, durations  
2 of exposure, vehicle, strain, and gender. One study provided a limited examination of  
3 TCE-induced liver weight changes in gerbils.

**Table 4-58. Oral studies of TCE-induced liver effects in mice and rats**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Section(s) where noncancer liver effects are discussed
Berman et al. (1995)	F344 rats (F)	Corn oil gavage	0, 150, 500, 1,500 or 5,000 mg/kg for once d 0, 50, 150, 500 or 1,500 mg/kg-day for 14 d]	8/group]	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology E.2.1.11. Berman et al. (1995)
<b>Buben and O'Flaherty (1985)</b>	<b>Swiss-Cox mice (M)</b>	<b>Corn oil gavage</b>	<b>0, 100, 200, 400, 800, 1,600, 2,400, or 3.200 mg/kg-day, 5 d/wk for 6 wk</b>	<b>12–15/group</b>	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology E.2.2.7. Buben and O'Flaherty (Buben and O'Flaherty, 1985)
Channel et al. (1998)	B6C3F1/CrlBR mice (M)	Corn oil gavage	0 (water), 0 (corn oil), 400, 800, or 1,200 mg/kg-day, 5 d/wk for up to 8 wk	77/group	4.5.4.2 Cytotoxicity and histopathology 4.5.4.3 Measures of DNA Synthesis, Cellular Proliferation, and Apoptosis 4.5.4.4 Peroxisome proliferation and related effects 4.5.4.5 Oxidative stress E.2.2.8. Channel et al. (1998)
Dees and Travis (1993)	B6C3F1 mice (M and F)	Corn oil gavage	0, 100, 250, 500, or 1,000 mg/kg-day for 10 d	5/group	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology 4.5.4.3 Measures of DNA Synthesis E.2.1.9. Dees and Travis (1993)
Elcombe et al. (1985)	B6C3F1 and Alderley Park (Swiss) mice (M) Osborne-Mendel and Alderley Park (Wistar) rats (M)	Corn oil gavage	0, 500, 1,000, or 1,500 mg/kg-day for 10 d	6–10/group	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology 4.5.4.3 Measures of DNA Synthesis, Cellular Proliferation, and Apoptosis 4.5.4.4 Peroxisome proliferation and related effects E.2.1.8. Elcombe et al. (1985)
Goel et al. (1992)	Swiss albino mice (M)	Groundnut oil gavage	0, 500, 1,000, or 2,000 mg/kg-day, 5 d/wk for 28 d.	6/group	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology 4.5.4.4 Peroxisome proliferation and related effects E.2.2.2. Goel et al. (1992)

**Table 4-58. Oral studies of TCE-induced liver effects in mice and rats (continued)**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Section(s) where noncancer liver effects are discussed
Goldsworthy and Popp (1987)	F344 rats (M) B6C3F1 mice (M)	Corn oil or methyl cellulose gavage	1,000 mg/kg-day for 10 d	5–7/group	4.5.4.1 Liver weight 4.5.4.4 Peroxisome proliferation and related effects E.2.1.7. Goldsworthy and Popp (1987)
Laughter et al. (2004)	Sv/129 and PPAR $\alpha$ -null mice (M)	Methyl-cellulose gavage	0–1,500 mg/kg-day for 3 d; and 5 d/wk for 3 wk	4–5/group	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology 4.5.4.3 Measures of DNA Synthesis, Cellular Proliferation, and Apoptosis 4.5.4.4 Peroxisome proliferation and related effects E.2.1.13. Laughter et al. (2004)
Melnick et al. (1987)	F344 rats (M)	Micro-encapsulated in feed Corn oil gavage	0, 0.055, 1.10, 2.21, or 4.41% in feed for 14 d, equivalent to 0, 600, 1,300, 2,200, or 4,800 mg/kg-day	10/group	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology 4.5.4.4 Peroxisome proliferation and related effects E.2.1.12. Melnick et al. (1987)
Merrick et al. (1989)	B6C3F1 mice (M and F)	Corn oil and 20% emulphor in water gavage	Males: 0, 600, 1,200, or 2,400 mg/kg-day Females: 0, 450, 900, or 1,800 mg/kg-day	12/group	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology E.2.2.1. Merrick et al. (1989)
Mirsalis et al. (1989)	B6C3F1 mice (M and F) F344 rats (M)	Corn oil gavage	0, 50, 200, or 1,000 mg/kg (single dose)	3/group	4.5.4.3 Measures of DNA Synthesis, Cellular Proliferation, and Apoptosis E.2.4.1. Summary of Results for Short-term Effects of Trichloroethylene (TCE)

**Table 4-58. Oral studies of TCE-induced liver effects in mice and rats (continued)**

<b>Reference</b>	<b>Animals (sex)</b>	<b>Exposure route</b>	<b>Dose/exposure concentration</b>	<b>Exposed</b>	<b>Section(s) where noncancer liver effects are discussed</b>
Nakajima et al. (2000)	Sv/129 and PPAR $\alpha$ -null mice (M and F)	Corn oil gavage	0 or 750 mg/kg-day for 14 d	6/sex/group	4.5.4.1 Liver weight 4.5.4.4 Peroxisome proliferation and related effects E.2.1.10. Nakajima et al. (2000)
NTP (1990)	B6C3F1 mice (M and F) F334/N rats (M and F)	Corn oil gavage	Mice: 0, 375–6,000 mg/kg-day, 5 d/wk, 13 wk Rats: 0, 62.5–1,000 mg/kg-day, 5 d/wk, 13 wk	10/group	4.5.4.2 Cytotoxicity and histopathology E.2.2.12.1 13-wk studies
NTP (1990)	B6C3F1 mice (M and F) F334/N rats (M and F)	Corn oil gavage	Mice: 0, or 1,000 mg/kg-day, 5 d/wk, 103 wk Rats: 0, 500, or 1,000 mg/kg-day, 5 d/wk, 103 wk	50/group	4.5.4.2 Cytotoxicity and histopathology E.2.2.12.2 2-yr studies
Nunes et al. (2001)	Sprague-Dawley rats (M)	Corn oil gavage	2,000 mg/kg-day on d 10–16 (with and without lead carbonate pretreatment for 9 d)	10/group	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology E.2.1.4. Nunes et al. (2001)
Tao et al. (2000a)	B6C3F1 mice (F)	Corn oil gavage	1,000 mg/kg-day for 5 d	4–6/group	4.5.4.1 Liver weight E.2.1.5. Tao et al., (2000a)
Tucker et al. (1982)	CD-1 mice (M and F)	Drinking water with 1% emulphor	0 (untreated), 0 (vehicle), 0.1, 1.0, 2.5, or 5 mg/ml for 4 or 6 mo. M: 0, 0, 18.4, 216.7, 393.0, or 660.2 mg/kg-day F: 0, 0, 17.9, 193.0, 437.1, or 793.3 mg/kg-day	140/group untreated and TCE-treated 260/group vehicle-treated	4.5.4.1 Liver weight E.2.1.6. Tucker et al. (1982)

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

**Table 4-59. Inhalation and i.p. studies of TCE-induced liver effects in mice and rats**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Section(s) where noncancer liver effects are discussed
Hamdan and Stacey (1993)	Sprague-Dawley rats (M)	i.p. in corn oil	0 or 131 mg/kg	6/group	4.5.4.6 Bile production E.2.6. Serum Bile Acid Assays
Kaneko et al. (2000)	MRL-lpr/lpr mice (M)	Inhalation	0, 500, 1,000, or 2,000 ppm, 4 h/d, 6 d/wk, for 8 wk	5/group	4.5.4.2 Cytotoxicity and histopathology
Kjellstrand et al. (1981b)	NMRI mice. Sprague-Dawley rats Mongolian gerbils	Inhalation	150 ppm continuous for 2–30 d	4–12/group	4.5.4.1 Liver weight E.2.2.3. Kjellstrand et al., (1981b)
Kjellstrand et al. (1983b)	wild, C57Bl, DBA, B6CBA, A/sn, NZB, and NMRI mice (M and F)	Inhalation	150 ppm continuous for 30 d	6/group	4.5.4.1 Liver weight E.2.2.5. Kjellstrand et al., (1983b)
<b>Kjellstrand et al. (1983a)</b>	<b>NMRI mice (M and F)</b>	<b>Inhalation</b>	<b>0–3,600 ppm, variable time periods of 1–24 h/d, for 30 or 120 d.</b>	<b>10–20/group</b>	<b>4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology E.2.2.6. Kjellstrand et al., (1983a)</b>
Kumar et al. (2001a)	Wistar rats (M)	Inhalation	376 ppm, 4 h/d, 5 d/wk, 8–24 wk	6/group	4.5.4.2 Cytotoxicity and histopathology E.2.2.10. Kumar et al., 2001
Okino et al. (1991)	Wistar rats (M)	Inhalation	0, 500 (8 h), 2,000 (2 h or 8 h), or 8,000 ppm (2 h) (single exposure)	5/group	4.5.4.2 Cytotoxicity and histopathology E.2.1.3. Okino et al. (1991)
Ramdhan et al. (2008)	SV/129 mice (M) CYP2E1-null mice (M)	Inhalation	0, 1,000, or 2,000 ppm, 8 h/d, 7 d	6/group	4.5.4.2 Cytotoxicity and histopathology 4.5.6.2.1. Hepatomegally- qualitative and quantitative comparisons 4.5.6.2.2. Cytotoxicity E.2.1.14. Ramdhan et al. (2008)

**Table 4-59. Inhalation and i.p. studies of TCE-induced liver effects in mice and rats (continued)**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Section(s) where noncancer liver effects are discussed
Ramdhan et al. (2010)	Sv/129, PPAR $\alpha$ -null, and hPPAR $\alpha$ mice (M)	Inhalation	0, 1,000, or 2,000 ppm. 8 h/d, 7 d	6/group]	4.5.4.1 Liver weight 4.5.4.2 Cytotoxicity and histopathology 4.5.6.2.1. Hepatomegally- qualitative and quantitative comparisons 4.5.6.2.2. Cytotoxicity 4.5.7.2. Peroxisome Proliferator Activated Receptor Alpha (PPAR $\alpha$ ) Receptor Activation E.2.1.15. Ramdhan et al. (2010)
Toraason et al. (1999)	Fischer rats (M)	i.p. in Alkamuls/ water	0, 100, 500, or 1,000 mg/kg	6/group	4.5.4.5 Oxidative stress E.2.4.3. Summary of Trichloroethylene (TCE) Subchronic and Chronic Studies E.3.4.2.3. Oxidative Stress
Wang and Stacey (1990)	Sprague-Dawley rats (M)	i.p. in corn oil Inhalation	i.p.: 0, 1.3–1,314 mg/kg-day for 3 d Inhalation: 0, 200, or 1,000 ppm, 6 h/d for 28 d	4–6/group	4.5.4.6 Bile production E.2.2. Subchronic and Chronic Studies of Trichloroethylene (TCE)
Watanabe and Fukui (2000)	ddY mice (M)	i.p. in corn oil	0, 158 mg/kg (single dose)	4/group	4.5.4.4 Peroxisome proliferation and related effects
<b>Woolhiser et al. (2006)</b>	<b>Sprague-Dawley rats (F)</b>	<b>Inhalation</b>	<b>0, 100, 300, or 1,000 ppm, 6 h/d, 5 d/wk, for 4 wk</b>	<b>16/group</b>	4.5.4.1 Liver weight E.2.2.4. Woolhiser et al. (2006)

**Bolded study(ies)** carried forward for consideration in dose-response assessment (see Section 5).

1 TCE-induced increases in liver weight have been reported to occur quickly.  
2 Kjellstrand et al. (1981b) reported liver weight increases after 2 days inhalation exposure in  
3 NMRI mice, Laughter et al. (2004) reported increased liver weight in SV129 mice in their 3-days  
4 study (see below), and Tao et al. (2000a) reported a increased in percentage liver/body weight  
5 ratio in female B6C3F1 mice for after 5 days. Elcombe et al. (1985) and Dees and Travis (1993)  
6 reported gavage results in mice and rats after 10 days exposure to TCE which showed  
7 TCE-induced increases in liver weight. Tucker et al. (1982) reported that 14 days of exposure to  
8 24 mg/kg and 240 mg/kg TCE via gavage to induce a dose-related increase in liver weight in  
9 male CD-1 mice but did not show the data.

10 For mice, the inhalation studies of Kjellstrand et al provided the most information on the  
11 affect of duration of exposure, dose of exposure, strain tested, gender, initial weight, and  
12 variability in response between experiments on TCE-induced liver weight increases. These  
13 experiments also provided results that were independent of vehicle effect. Although the  
14 determination of the exact magnitude of response is limited by experimental design,  
15 Kjellstrand et al. (1981b) reported that in NMRI mice, continuous TCE inhalation exposure  
16 induced increased percentage liver/body weight by 2 days and that by 30 days (the last recorded  
17 data point) the highest percentage liver/body weight ratio was reported (~1.75-fold over controls)  
18 in both male and female mice. Kjellstrand et al. (1983b) exposed seven different strains of mice  
19 (wild, C57BL, DBA, B6CBA, A/sn, NZB, NMRI) to 150-ppm TCE for 30 days and  
20 demonstrated that strain, gender, and toxicity, as reflected by changes in whole-body weight,  
21 affected the percentage liver/body weight ratios induced by 30 days of continuous TCE  
22 exposure. In general for the seven strains of mice examined, female mice had the less variable  
23 increases in TCE-induced liver weight gain across duplicate experiments than male mice. For  
24 instance, in strains that did not exhibit changes in body weight (reflecting systemic toxicity) in  
25 either gender (wild-type and DBA), 150-ppm TCE exposure for 30 days induced 1.74- to  
26 1.87-fold of control percentage liver/body weight ratios in female mice and 1.45- to 2.00-fold of  
27 control percentage liver/body weight ratios in male mice. The strain with the largest  
28 TCE-induced increase in percentage liver/body weight increase was the NZB strain (~2.08-fold  
29 of control for females and 2.34- to 3.57-fold of control for males). Kjellstrand et al. (1983a)  
30 provided dose-response information for the NMRI strain of mice (A Swiss-derived strain) that  
31 indicated dose-related increases in percentage liver/body weight ratios between 37- and 300-ppm  
32 TCE exposure for 30 days. The 150-ppm dose was reported to induce a 1.66- and 1.69-fold  
33 increases in percentage liver/body weight ratios in male and female mice, respectively.  
34 Interestingly, they also reported similar liver weight increases among groups with the same



1 cumulative exposure, but with different daily exposure durations (1 hour/day at 3,600 ppm to  
2 24 hours/day at 150 ppm for 30 days).

3 Not only have most gavage experiments have been carried out in male mice, which  
4 Kjellstrand et al. (1983b) had demonstrated to have more variability in response than females,  
5 but also vehicle effects were noted to occur in experiments that examined them. Merrick et al.  
6 (1989) reported that corn oil induced a similar increase in percentage liver/body weight ratios in  
7 female mice fed TCE in emulphor and corn oil for 4 weeks, male mice TCE administered in the  
8 corn oil vehicle induced a greater increase in liver weight than emulphor but less mortality at a  
9 high does.

10 Buben and O'Flaherty (1985) treated male, outbred Swiss-Cox mice for 6 weeks at doses  
11 ranging from 100–3,200 mg/kg-day, and reported increased liver/body-weight ratios at all tested  
12 doses (1.12- to 1.75-fold of controls). Given the large strain differences observed by Kjellstrand  
13 et al. (1983b), the use of predominantly male mice, and the effects of vehicle in gavage studies,  
14 interstudy variability in dose-response relationships is not surprising.

15 Dependence of PPAR $\alpha$  activation for TCE-liver weight gain has been investigated in  
16 PPAR $\alpha$  null mice by Nakajima et al. (2000), Laughter et al. (2004), and Ramdhan et al. (2010),  
17 the latter of which also investigated PPAR $\alpha$  null mice with human PPAR $\alpha$  inserted. Nakajima  
18 et al. (2000) reported that after 2 weeks of 750 mg/kg TCE exposure to carefully matched SV129  
19 wild-type or PPAR $\alpha$ -null male and female mice ( $n = 6$  group), there was a reported 1.50-fold  
20 increase in wild-type and 1.26-fold of control percentage liver/body weight ratio in PPAR $\alpha$ -null  
21 male mice. For female mice, there was ~1.25-fold of control percentage liver/body weight ratios  
22 for both wild-type and PPAR $\alpha$ -null mice. Thus, TCE-induced liver weight gain was not  
23 dependent on a functional PPAR $\alpha$  receptor in female mice and some portion of it may have been  
24 in male mice. Both wild-type male and female mice were reported to have similar increases in  
25 the number of peroxisome in the pericentral area of the liver and TCE exposure and, although  
26 increased twofold, were still only ~4% of cytoplasmic volume. Female wild-type mice were  
27 reported to have less TCE-induced elevation of very long chain acyl-CoA synthetase, D-type  
28 peroxisomal bifunctional protein, mitochondrial trifunctional protein  $\alpha$  subunits  $\alpha$  and  $\beta$ , and  
29 cytochrome P450 4A1 than males mice, even though peroxisomal volume was similarly elevated  
30 in male and female mice. The induction of PPAR $\alpha$  protein by TCE treatment was also reported  
31 to be slightly less in female than male wild-type mice (2.17- vs. 1.44-fold of control induction,  
32 respectively). Thus, differences between genders in this study were for increased liver weight  
33 were not associated with differences in peroxisomal volume in the hepatocytes but there was a  
34 gender-related difference in induction of enzymes and proteins associated with PPAR $\alpha$ .

1 The study of Laughter et al. (2004) used SV129 wild-type and PPAR $\alpha$ -null male mice  
2 treated with three daily doses of TCE in 0.1% methyl cellulose for either 3 days (1,500 mg/kg  
3 TCE) or 3 weeks (0, 10, 50, 125, 500, 1,000, or 1,500 mg/kg TCE 5 days a week). However, the  
4 paradigm is not strictly comparable to other gavage paradigms due to the different dose vehicle  
5 and the documented impacts of vehicles such as corn oil on TCE-induced effects. In addition, no  
6 initial or final body weights of the mice were reported and thus, the influence of differences in  
7 initial body weight on percentage liver/body weight determinations could not be ascertained.  
8 While control wild-type and PPAR $\alpha$ -null mice were reported to have similar percentage  
9 liver/body weight ratios (i.e., ~4.5%) at the end of the 3-day study, at the end of the 3-week  
10 experiment the percentage liver/body weight ratios were reported to be larger in the control  
11 PPAR $\alpha$ -null male mice (5.1%). TCE treatment for 3 days was reported for percentage liver/body  
12 weight ratio to be 1.4-fold of control in the wild-type mice and 1.07-fold of control in the null  
13 mice. After 3 weeks of TCE exposure at varying concentrations, wild-type mice were reported  
14 to have percentage liver/body weight ratios that were within ~2% of control values with the  
15 exception of the 1,000 mg/kg and 1,500 mg/kg treatment groups (~1.18- and 1.30-fold of  
16 control, respectively). For the PPAR $\alpha$ -null mice the variability in percentage liver/body weight  
17 ratios were reported to be greater than that of the wild-type mice in most of the TCE groups and  
18 the baseline levels of percentage liver/body weight ratio for control mice 1.16-fold of that of  
19 wild-type mice. TCE exposure was apparently more toxic in the PPAR $\alpha$ -null mice. Decreased  
20 survival at the 1,500 mg/kg TCE exposure level resulted in the prevention of recording of  
21 percentage liver/body weight ratios for this group. At 1,000 mg/kg TCE exposure level, there  
22 was a reported 1.10-fold of control percentage liver/body weight ratio in the PPAR $\alpha$ -null mice.  
23 None of the increases in percentage liver/body weight in the null mice were reported to be  
24 statistically significant by Laughter et al. (2004). However, the power of the study was limited  
25 due to low numbers of animals and increased variability in the null mice groups. The percentage  
26 liver/body weight ratio after TCE treatment reported in this study was actually greater in the  
27 PPAR $\alpha$ -null mice than the wild-type male mice at the 1,000 mg/kg TCE exposure level  
28 ( $5.6 \pm 0.4\%$  vs.  $5.2 \pm 0.5\%$ , for PPAR $\alpha$ -null and wild-type mice, respectively) resulting in a  
29 1.18-fold of wild-type and 1.10-fold of PPAR $\alpha$ -null mice. Although the results reported in  
30 Laughter et al. (2004) for DCA and TCA were not conducted in experiments that used the same  
31 paradigm, the TCE-induced increase in percentage liver/body weight more closely resembled the  
32 dose-response pattern for DCA than for DCA wild-type SV129 and PPAR $\alpha$ -null mice.  
33 Ramdhan et al. (2010) examined TCE-induced hepatic steatosis and toxicity using male  
34 wild type, PPAR $\alpha$ -null, and human PPAR $\alpha$  inserted (“humanized”) mice exposed to high  
35 inhalation concentrations of TCE for 7 days. Significant differences were observed among

1 control mice for each genotype with reduced body weight in untreated humanized mice.  
2 Liver/body weight ratios were 11% higher in untreated PPAR $\alpha$ -null mice than wild type mice.  
3 Higher levels of liver triglycerides and hepatic steatosis were reported in the untreated  
4 humanized mice and PPAR $\alpha$  null mice than wild type mice. Background expression of a number  
5 of genes and protein expression levels were significantly different between the untreated strains.  
6 In particular, human PPAR $\alpha$  protein levels were >10-fold greater in the humanized mice than  
7 mouse PPAR $\alpha$  in untreated wild type mice. Insertion of human PPAR $\alpha$  in the null mice did not  
8 return the mice to a normal state. Both PPAR $\alpha$  null and humanized mice were more susceptible  
9 to TCE toxicity. Hepatomegaly was induced in all strains to a similar extent after TCE  
10 exposure. However, urinary TCA concentrations were reported to be significantly lower and  
11 trichloroethanol levels significantly higher in TCE-treated PPAR $\alpha$ -null mice in comparison to  
12 treated wild type mice. This difference was not related to changes in expression of metabolic  
13 enzymes.

14 No study examined strain differences among rats, and cross-study comparisons are  
15 confounded by heterogeneity in the age of animals, dosing regimen, and other design  
16 characteristics that may affect the degree of response. For rats, TCE-induced percentage  
17 liver/body weight ratios were reported to range from 1.16- to 1.46-fold of control values  
18 depending on the study paradigm. The studies which employed the largest range of exposure  
19 concentrations (Berman et al., 1995; Melnick et al., 1987) examined four doses in the rat. In  
20 general, there was a dose-related increase in percentage liver/body weight in the rat, especially at  
21 doses that did not cause concurrent decreased survival or significant body weight loss. For  
22 gerbils, Kjellstrand et al. (1981b) reported a similar value of ~1.25-fold of control percentage  
23 liver/body weight as for S-D rats exposed to 150 ppm TCE continuously for 30 days. Woolhiser  
24 et al. (2006) also reported inhalation TCE exposure to increase the percentage liver/body weight  
25 ratios in female Sprague-Dawley rats although this strain appeared to be less responsive than  
26 others tested for induction of hepatomegaly from TCA exposure and to also be less prone to  
27 spontaneous liver cancer.

28 The size of the liver is under tight control and after cessation of a mitogenic stimulus or  
29 one inducing hepatomegaly, the liver will return to its preprogrammed size (see Appendix E).  
30 The increase in liver weight from TCE-exposure also appears to be reversible. Kjellstrand et al.  
31 (1981b) reported a reduction in liver weight gain increases after cessation of TCE exposure for 5  
32 or 30 days in male and female mice. However, experimental design limitations precluded  
33 discernment of the magnitude of decrease. Kjellstrand et al. (1983a) reported that mice exposed  
34 to 150 ppm TCE for 30 days and then examined 120 days after the cessation of exposure, had  
35 liver weights were 1.09-fold of control for TCE-exposed female mice and the same as controls

1 for TCE-exposed male mice. However, the livers were not the same as untreated liver in terms  
2 of histopathology. The authors reported that “after exposure to 150 ppm for 30 days, followed  
3 by 120 days of rehabilitation, the morphological picture was similar to that of the air-exposure  
4 controls except for changes in cellular and nuclear sizes.” Qualitatively, the reduction in liver  
5 weight after treatment cessation is consistent with the report of Elcombe et al. (1985) in Alderly  
6 Park mice. The authors report that the reversibility of liver effects after the administration of  
7 TCE to Alderly Park mice for 10 consecutive days. Effects upon liver weight, DNA  
8 concentration, and tritiated thymidine incorporation 24 and 48 hours after the last dose of TCE  
9 were reported to still be apparent. However, 6 days following the last dose of TCE, all of these  
10 parameters were reported to return to control values with the authors not showing the data to  
11 support this assertion. Thus, cessation of TCE exposure would have resulted in a 75% reduction  
12 in liver weight by 4 days in mice exposed to the highest TCE concentration. Quantitative  
13 comparisons are not possible because Elcombe et al. (1985) did not report data for these results  
14 (e.g., how many animals, what treatment doses, and differences in baseline body weights) and  
15 such a large decrease in such a short period of time needs to be verified.  
16

#### 4.5.4.1.2. Cytotoxicity and Histopathology

17 Acute exposure to TCE appears to induce low cytotoxicity below subchronically lethal  
18 doses. Relatively high doses of TCE appear necessary to induce cytotoxicity after a single  
19 exposure with two available studies reported in rats. Okino et al. (1991) reported small increases  
20 in the incidence of hepatocellular necrosis in male Wistar rats exposed to 2,000 ppm (8 hours)  
21 and 8,000 ppm (2 hours), but not at lower exposures. In addition, “swollen” hepatocytes were  
22 noted at the higher exposure when rats were pretreated with ethanol or Phenobarbital. Serum  
23 transaminases increased only marginally at the 8,000-ppm exposure, with greater increases with  
24 pretreatments. Berman et al. (1995) reported hepatocellular necrosis, but not changes in serum  
25 markers of necrosis, after single gavage doses of 1,500 and 5,000 mg/kg TCE in female F344  
26 rats. However, they did not report any indications of necrosis after 14 days of treatment at  
27 50–1,500 mg/kg-day nor the extent of necrosis.

28 At acute and subchronic exposure periods to multiple doses, the induction of cytotoxicity,  
29 though usually mild, appears to differ depending on rodent species, strain, dosing vehicle and  
30 duration of exposure, and the extent of reporting to vary between studies. For instance,  
31 Elcombe et al. (1985) and Dees and Travis (1993), which used the B6C3F1 mouse strain and  
32 corn oil vehicle, reported only slight or mild necrosis after 10 days of treatment with TCE at  
33 doses up to 1,500 mg/kg-day. Elcombe et al. (1985) also reported cell hypertrophy in the

1 centrilobular region. Dees and Travis (1993) reported some loss of vacuolization in hepatocytes  
2 of mice treated at 1,000 mg/kg-day. Laughter et al. (2004) reported that “wild-type” SV129  
3 mice exposed to 1,500 mg/kg TCE exposure for 3 weeks exhibited mild granuloma formation  
4 with calcification or mild hepatocyte degeneration but gave no other details or quantitative  
5 information as to the extent of the lesions or what parts of the liver lobule were affected. The  
6 authors noted that “wild-type mice administered 1,000 and 1,500 mg/kg exhibited centrilobular  
7 hypertrophy” and that “the mice in the other groups did not exhibit any gross pathological  
8 changes” after TCE exposure. Channel et al. (1998) reported no necrosis in B6C3F1 mice  
9 treated by 400–1,200 mg/kg-day TCE by corn oil gavage for 2 days to 8 weeks.

10 However, as stated above, Merrick et al. (1989) reported that corn oil resulted in more  
11 hepatocellular necrosis, as described by small focal areas of 3–5 hepatocytes, in male B6C3F1  
12 mice than use of emulphor as a vehicle for 4-week TCE gavage exposures. Necrotic hepatocytes  
13 were described as surrounded by macrophages and polymorphonuclear cells. The authors  
14 reported that visible necrosis was observed in 30–40% of male mice administered TCE in corn  
15 oil but not that there did not appear to be a dose-response. For female mice, the extent of  
16 necrosis was reported to be 0 for all control and TCE treatment groups using either vehicle.  
17 Serum enzyme activities for alanine aminotransferase (ALT), AST, and LDH (markers of liver  
18 toxicity) showed that there was no difference between vehicle groups at comparable TCE  
19 exposure levels for male or female mice. Except for LDH levels in male mice exposed to TCE  
20 in corn oil there was not a correlation with the extent of necrosis and the patterns of increases in  
21 ALT and AST enzyme levels.

22 Ramdhan et al. (2008) assessed TCE-induced hepatotoxicity by measuring plasma ALT  
23 and AST activities and histopathology in Sv/129 mice treated by inhalation exposure, which are  
24 not confounded by vehicle effects. Despite high variability and only six animals per dose group,  
25 all three measures showed statistically significant increases at the high dose of 2,000 ppm  
26 (8 hours/day for 7 days), although a nonstatistically significant elevation is evident at the low  
27 dose of 1,000 ppm. Even at the highest dose, cytotoxicity was not severe, with ALT and AST  
28 measures increased twofold or less and an average histological score less than two (range 0–4).

29 Using the same paradigm, Ramdhan et al. (2010) also reported increased in AST and  
30 ALT liver injury biomarkers to be significantly increased in all exposed mice (Sv/129 wild type,  
31 PPAR $\alpha$ -null and humanized PPAR $\alpha$  mice) relative to controls (41–74% and 36–79% higher,  
32 respectively) with mean levels within each group higher, though not statistically significantly  
33 different, with exposure to 2,000 versus 1,000 ppm TCE. . Steatosis scores were reported to be  
34 significantly higher in the 2,000 versus 1,000 ppm TCE exposures to PPAR $\alpha$ -null mice. The  
35 authors reported steatosis scored to be significantly correlated with liver triglyceride levels of all

1 mice examined in the study ( $r = 0.75$ ). Macrovesicular steatosis was reported to occur more  
2 frequently in hPPAR $\alpha$  than PPAR $\alpha$ -null mice. Necrosis scores were reported to be significantly  
3 higher in TCE exposed mice relative to controls in all three genotype mice and to be significantly  
4 higher with 2,000 versus 1,000 ppm TCE exposure in wild type mice and hPPAR $\alpha$  mice.  
5 Inflammation scores were reported to be significantly higher with exposed group than control  
6 with 2,000 ppm TCE exposure than controls for each genotype group with a difference between  
7 the 2,000 ppm and 1,000 ppm exposure groups in wild type mice.

8 Kjellstrand et al. (1983a) exposed male and female NRM1 mice to 150 ppm for  
9 30–120 days. Kjellstrand et al. (1983a) reported more detailed light microscopic findings from  
10 their study and stated that

11  
12  
13 After 150 ppm exposure for 30 days, the normal trabecular arrangement of the liver cells  
14 remained. However, the liver cells were generally larger and often displayed a fine vacuolization  
15 of the cytoplasm. The nucleoli varied slightly to moderately in size and shape and had a finer,  
16 granular chromatin with a varying basophilic staining intensity. The Kupffer cells of the sinusoid  
17 were increased in cellular and nuclear size. The intralobular connective tissue was infiltrated by  
18 inflammatory cells. There was not sign of bile stasis. Exposure to TCE in higher or lower  
19 concentrations during the 30 days produced a similar morphologic picture. After intermittent  
20 exposure for 30 days to a time-weighted-average concentration of 150 ppm or continuous  
21 exposure for 120 days, the trabecular cellular arrangement was less well preserved. The cells had  
22 increased in size and the variations in size and shape of the cells were much greater. The nuclei  
23 also displayed a greater variation in basophilic staining intensity, and often had one or two  
24 enlarged nucleoli. Mitosis was also more frequent in the groups exposed for longer intervals. The  
25 vacuolization of the cytoplasm was also much more pronounced. Inflammatory cell infiltration in  
26 the interlobular connective tissue was more prominent. After exposure to 150 ppm for 30 days,  
27 followed by 120 days of rehabilitation, the morphological picture was similar to that of the  
28 air-exposure controls except for changes in cellular and nuclear sizes.

29  
30  
31 Although not reporting comparisons between male and female mice in the results section  
32 of the paper for TCE-induced histopathological changes, the authors stated in the discussion  
33 section that “However, liver mass increase and the changes in liver cell morphology were similar  
34 in TCE-exposed male and female mice.” Kjellstrand et al. (1983a) did not present any  
35 quantitative data on the lesions they describe, especially in terms of dose-response. Most of the  
36 qualitative description presented was for the 150-ppm exposure level and the authors suggest that  
37 lower concentrations of TCE give a similar pathology as those at the 150-ppm level, but do not  
38 present data to support that conclusion. Although stating that Kupffer cells were reported to be  
39 increased in cellular and nuclear size, no differential staining was applied light microscopy  
40 sections to distinguish Kupffer from endothelial cells lining the hepatic sinusoid in this study.  
41 Without differential staining such a determination is difficult at the light microscopic level.

1           Indeed, Goel et al. (1992) describe proliferation of “sinusoidal endothelial cells” after  
2 1,000 and 2,000 mg/kg-day TCE exposure for 28 days in male Swiss mice. They reported that  
3 histologically, “the liver exhibits swelling, vacuolization, widespread degeneration/necrosis of  
4 hepatocytes as well as marked proliferation of endothelial cells of hepatic sinusoids at 1,000 and  
5 2,000 mg/kg TCE doses.” Only one figure is given, at the light microscopic level, in which it is  
6 impossible to distinguish endothelial cells from Kupffer cells and no quantitative measures or  
7 proliferation were examined or reported to support the conclusion that endothelial cells are  
8 proliferating in response to TCE treatment. Similarly, no quantitative analysis regarding the  
9 extent or location of hepatocellular necrosis was given. The presence or absence of  
10 inflammatory cells were not noted by the authors as well. In terms of white blood cell count, the  
11 authors note that it is slightly increased at 500 mg/kg-day but decreased at 1,000 and  
12 2,000 mg/kg-day TCE, perhaps indicating macrophage recruitment from blood to liver and  
13 kidney, which was also noted to have pathology at these concentrations of TCE.

14           The inflammatory cell infiltrates described in the Kjellstrand et al. (1983a) study are  
15 consistent with invasion of macrophages and well as polymorphonuclear cells into the liver,  
16 which could activate resident Kupffer cells. Although not specifically describing the changes as  
17 consistent with increased polyploidization of hepatocytes, the changes in cell size and especially  
18 the continued change in cell size and nuclear staining characteristics after 120 days of cessation  
19 of exposure are consistent with changes in polyploidization induced by TCE. Of note is that in  
20 the histological description provided by the authors, although vacuolization is reported and  
21 consistent with hepatotoxicity or lipid accumulation, which is lost during routine histological  
22 slide preparation, there is no mention of focal necrosis or apoptosis resulting from these  
23 exposures to TCE.

24           Buben and O’Flaherty (1985) reported liver degeneration “as swollen hepatocytes” and to  
25 be common with treatment of TCE to Male Swiss-Cox mice after 6 weeks. They reported that  
26 “Cells had indistinct borders; their cytoplasm was clumped and a vesicular pattern was apparent.  
27 The swelling was not simply due to edema, as wet weight/dry weight ratios did not increase.”  
28 Karyorrhexis (the disintegration of the nucleus) was reported to be present in nearly all  
29 specimens and suggestive of impending cell death. No Karyorrhexis, necrosis, or polyploidy  
30 was reported in controls, but a low score Karyorrhexis was given for 400 mg/kg TCE and a  
31 slightly higher one given for 1,600 mg/kg TCE. Central lobular necrosis reported to be present  
32 only at the 1,600 mg/kg TCE exposure level and assigned a low score. Polyploidy was described  
33 as characteristic in the central lobular region but with low score for both 400 mg/kg and  
34 1,600 mg/kg TCE exposures. The authors reported that “hepatic cells had two or more nuclei or  
35 had enlarged nuclei containing increased amounts of chromatin, suggesting that a regenerative

1 process was ongoing” and that there were no fine lipid droplets in TCE exposed animals. The  
2 finding of “no polyploidy” in control mouse liver in the study of Buben and O’Flaherty (1985) is  
3 unexpected given that binucleate and polyploid hepatocytes are a common finding in the mature  
4 mouse liver. It is possible that the authors were referring to unusually high instances of  
5 “polyploidy” in comparison to what would be expected for the mature mouse. The score given  
6 by the authors for polyploidy did not indicate a difference between the two TCE exposure  
7 treatments and that it was of the lowest level of severity or occurrence. No score was given for  
8 centrilobular hypertrophy although the DNA content and liver weight changes suggested a  
9 dose-response. The “Karyrrhexis” described in this study could have been a sign of cell death  
10 associated with increased liver cell number or dying of maturing hepatocytes associated with the  
11 increased ploidy, and suggests that TCE treatment was inducing polyploidization. Consistent  
12 with enzyme analyses, centrilobular necrosis was only seen at the highest dose and with the  
13 lowest qualitative score, indicating that even at the highest dose there was little toxicity.

14 At high doses, Kaneko et al. (2000) reported sporadic necrosis in male Mrl-lpr/lpr mice,  
15 which are “genetically liable to autoimmune disease,” exposed to 500–2,000 ppm, 4hours/day,  
16 6 days/week, for 8 weeks ( $n = 5$ ). Dose-dependent mild inflammation and associated changes  
17 were reported to be found in the liver. The effects on hepatocytes were reported to be minimal  
18 by the authors with 500-ppm TCE inducing sporadic necrosis in the hepatic lobule. Slight  
19 mobilization and activation of sinusoid lining cells were also noted. These pathological features  
20 were reported to increase with dose.

21 NTP (1990), which used the B6C3F1 mouse strain, reported centrilobular necrosis in  
22 6/10 male and 1/10 female B6C3F1 mice treated at a dose of 6,000 mg/kg-day for up to  
23 13 weeks (all the male mice and 8 of the 10 female mice died in the first week of treatment). At  
24 3,000 mg/kg-day exposure level, although centrilobular necrosis was not observed, 2/10 males  
25 had multifocal areas of calcification in their livers, which the authors suggest is indicative of  
26 earlier hepatocellular necrosis. However, only 3/10 male mice at this dose survived to the end of  
27 the 13-week study.

28 For the NTP (1990) 2-year study, B6C3F1 mice were reported to have no  
29 treatment-related increase in necrosis in the liver. A slight increase in the incidence of focal  
30 necrosis was noted TCE-exposed male mice (8 vs. 2%) with a slight reduction in fatty  
31 metamorphosis in treated male mice (zero treated vs. two control animals) and in female mice a  
32 slight increase in focal inflammation (29 vs. 19% of animals) and no other changes. Therefore,  
33 this study did not show concurrent evidence of liver toxicity with TCE-induced neoplasia after  
34 2 years of TCE exposure in mice.



1 For the more limited database in rats, there appears to be variability in reported TCE  
2 induced cytotoxicity and pathology. Nunes et al. (2001) reported no gross pathological changes  
3 in rats gavaged with corn oil or with corn oil plus 2,000 mg/kg TCE for 7 days. Goldsworthy  
4 and Popp (1987) gave no descriptions of liver histology given in this report for TCE-exposed  
5 animals or corn-oil controls. Kjellstrand et al. (1981b) gave also did not give histological  
6 descriptions for livers of rats in their inhalation study.

7 Elcombe et al. (1985) provided a description of the histopathology at the light  
8 microscopy level in Osborne-Mendel rats, and Alderly Park rats exposed to TCE via gavage for  
9 10 days. However, they did not provide a quantitative analysis or specific information regarding  
10 the variability of response between animals within group and there was no indication by the  
11 authors regarding how many rats were examined by light microscopy. Hematoxylin and eosin  
12 sections from Osborne-Mendel rats were reported to show that

13  
14  
15 Livers from control rats contained large quantities of glycogen and isolated inflammatory foci, but  
16 were otherwise normal. The majority of rats receiving 1,500 mg/kg body weight TCE showed  
17 slight changes in centrilobular hepatocytes. The hepatocytes were more eosinophilic and  
18 contained little glycogen. At lower doses these effects were less marked and were restricted to  
19 fewer animals. No evidence of treatment-related hepatotoxicity (as exemplified by single cell or  
20 focal necrosis) was seen in any rat receiving TCE. H&E [hematoxylin and eosin] sections from  
21 Alderly Park Rats showed no signs of treatment-related hepatotoxicity after administration of  
22 TCE. However, some signs of dose-related increase in centrilobular eosinophilia were noted.  
23  
24

25 Thus, both mice and rats were reported to exhibit pericentral hypertrophy and  
26 eosinophilia as noted from the histopathological examination in Elcombe et al. (1985).

27 Berman et al. (1995) reported that for female rats exposed to TCE for 14 days  
28 hepatocellular necrosis was noted to occur in the 1,500 and 5,000 mg/kg groups in 6/7 and  
29 6/8 female rats, respectively but not to occur in lower doses. The extent of necrosis was not  
30 noted by the authors for the two groups exhibiting a response after 1 day of exposure. Serum  
31 enzyme levels, indicative of liver necrosis, were not presented and because only positive results  
32 were presented in the paper, presumed to be negative. Therefore, the extent of necrosis was not  
33 of a magnitude to affect serum enzyme markers of cellular leakage.

34 Melnick et al. (1987) reported that the only treatment-related lesion observed  
35 microscopically in rats from either dosed-feed or gavage groups was individual cell necrosis of  
36 the liver with the frequency and severity of this lesion similar at each dosage levels of TCE  
37 microencapsulated in the feed or administered in corn oil. The severity for necrosis was only  
38 mild at the 2.2 and 4.8 g/kg feed groups and for the six animals in the 2.8 g/kg group corn oil  
39 group. The individual cell necrosis was reported to be randomly distributed throughout the liver

1 lobule with the change to not be accompanied by an inflammatory response. The authors also  
2 reported that there was no histologic evidence of cellular hypertrophy or edema in hepatic  
3 parenchymal cells. Thus, although there appeared to be TCE-treatment related increases in focal  
4 necrosis after 14 days of exposure, the extent was mild even at the highest doses and involved  
5 few hepatocytes.

6 For the 13-week NTP study (1990), only control and high dose F344/N rats were  
7 examined histologically. Pathological results were reported to reveal that 6/10 males and  
8 6/10 female rats had pulmonary vasculitis at the highest concentration of TCE. This change was  
9 also reported to have occurred in 1/10 control male and female rats. Most of those animals were  
10 also reported to have had mild interstitial pneumonitis. The authors report that viral titers were  
11 positive during this study for Sendai virus.

12 Kumar et al. (2001) reported that male Wistar rats exposed to 376 ppm, 4 hours/day,  
13 5 days/week for 8–24 weeks showed evidence of hepatic toxicity. The authors stated that, “after  
14 8 weeks of exposure enlarged hepatocytes, with uniform presence of fat vacuoles were found in  
15 all of the hepatocytes affecting the periportal, midzonal, and centrilobular areas, and fat  
16 vacuoles pushing the pyknotic nuclei to one side of hepatocytes. Moreover, congestion was not  
17 significant. After exposure of 12 and 24 weeks, the fatty changes became more progressive with  
18 marked necrosis, uniformly distributed in the entire organ.” No other description of pathology  
19 was provided in this report. In regard to the description of fatty change, the authors only did  
20 conventional H&E staining of sections with no precautions to preserve or stain lipids in their  
21 sections. However, as noted below, the NCI study also reports long-term TCE exposure in rats  
22 to result in hepatocellular fatty metamorphosis. The authors provided a table with histological  
23 scoring of simply + or – for minimal, mild or moderate effects and do not define the criteria for  
24 that scoring. There is also no quantitative information given as to the extent, nature, or location  
25 of hepatocellular necrosis. The authors report “no change was observed in glutamic oxoacetate  
26 transaminase and glutamic pyruvate transaminase levels of liver in all the three groups. The  
27 GSH level was significantly decreased while “total sulphhydryl” level was significantly increased  
28 during 8, 12, and 24 weeks of TCE exposure. The acid and alkaline phosphatases were  
29 significantly increased during 8, 12, and 24 weeks of TCE exposure.” The authors present a  
30 series of figures that are poor in quality to demonstrate histopathological TCE-induced changes.  
31 No mortality was observed from TCE exposure in any group despite the presence of liver  
32 necrosis.

33 Thus, in this limited database that spans durations of exposure from days to 24 weeks and  
34 uses differing routes of administration, generally high doses for long durations of exposure are  
35 required to induce hepatotoxicity from TCE exposure in the rat. The focus of 2-year bioassays in

1 rats has been the detection of a cancer response with little or no reporting of noncancer pathology  
2 in most studies. Henschler et al. (1984) and Fukuda et al. (1983) do not report noncancer  
3 histopathology, but do both report rare biliary cell derived tumors in rats in relatively insensitive  
4 assays. For male rats, noncancer pathology in the NCI (1976) study was reported to include  
5 increased fatty metamorphosis after TCE exposure and angiectasis or abnormally enlarged blood  
6 vessels. Angiectasis can be manifested by hyperproliferation of endothelial cells and dilatation  
7 of sinusoidal spaces. For the NTP (1990) study there was little reporting of non-neoplastic  
8 pathology or toxicity and no report of liver weight at termination of the study. In the NTP  
9 (1988) study, the 2 year study of TCE exposure reported no evidence of TCE-induced liver  
10 toxicity described as non-neoplastic changes in ACI, August, Marshal, and Osborne-Mendel rats.  
11 Interestingly, for the control animals of these four strains there was, in general, a low background  
12 level of focal necrosis in the liver of both genders. Obviously, the negative results in this  
13 bioassay for cancer are confounded by the killing of a large portion of the animals accidentally by  
14 experimental error but TCE-induced overt liver toxicity was not reported.

15 In sum, the cytotoxic effects in the liver of TCE treatment appear include little or no  
16 necrosis in the rodent liver, but rather, a number of histological changes such as mild focal  
17 hepatocyte degeneration at high doses, cellular “swelling” or hypertrophy, and enlarged nuclei.  
18 Histological changes consistent with increased polyploidization and specific descriptions of  
19 TCE-induced polyploidization have been noted in several experiments. Several studies note  
20 proliferation of nonparenchymal cells after TCE exposure as well. These results are more  
21 consistently reported in mice, but also have been reported in some studies at high doses in rats,  
22 for which fewer studies are available. In addition, the increase in cellular and nuclear sizes  
23 appeared to persist after cessation of TCE treatment. In neither rats nor mice is there evidence  
24 that TCE treatment results in marked necrosis leading to regenerative hyperplasia.

25

#### **4.5.4.1.3. Measures of DNA Synthesis, Cellular Proliferation, and Apoptosis**

26 The increased liver weight observed in rodents after TCE exposure may result from either  
27 increased numbers of cells in the liver, increased size of cells in the liver, or a combination of  
28 both. Studies of TCE in rodents have studied whole liver DNA content of TCE-treated animals  
29 to determine whether the concentration of DNA per gram of liver decreases as an indication of  
30 hepatocellular hypertrophy (Buben and O'Flaherty, 1985; Dees and Travis, 1993; Elcombe et al.,  
31 1985). While the slight decreases observed in some studies are consistent with hypertrophy, the  
32 large variability in controls and lack of dose-response limits the conclusions that can be drawn  
33 from these data. In addition, multiple factors beyond hypertrophy affect DNA concentration in

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 whole-liver homogenates, including changes in ploidy and the number of hepatocytes and  
2 nonparenchymal cells.

3 The incorporation of tritiated thymidine or BrdU has also been analyzed in whole liver  
4 DNA and in individual hepatocytes as a measure of DNA synthesis. Such DNA synthesis can  
5 occur from either increased numbers of hepatocytes in the liver or by increased polyploidization.  
6 Section E.1.1 describes polyploidization in human and rodent liver and its impacts on liver  
7 function, while Sections E.3.1.2 and E.3.3.1 discuss issues of target cell identification for liver  
8 cancer and changes in ploidy as a key event in liver cancer using animals models, respectively.  
9 Along with changes in cell size (hypertrophy), cell number (cellular proliferation), and the DNA  
10 content per cell (cell ploidy), the rate of apoptosis has also been noted or specifically examined  
11 in some studies of TCE and its metabolites. All of these phenomena have been identified in  
12 proposed hypotheses as key events possibly related to carcinogenicity. In particular, changes in  
13 cell proliferation and apoptosis have been postulated to be part of the MOA for PPAR $\alpha$ -agonists  
14 by Klaunig et al. (2003) (see Section E.3.4).

15 In regard to early changes in DNA synthesis, the data for TCE are very limited  
16 Mirsalis et al. (1989) reported measurements of in vivo-in vitro hepatocyte DNA repair and  
17 S-phase DNA synthesis in primary hepatocytes from male Fischer 344 rats and male and female  
18 B6C3F1 mice administered single doses of TCE by gavage in corn oil. They reported negative  
19 results 2–12 hours after treatment from 50–1,000 mg/kg TCE in rats and mice (male and female)  
20 for unscheduled DNA synthesis and repair using three animals per group. After 24 and 48 hours  
21 of 200 or 1,000 mg/kg TCE in male mice ( $n = 3$ ) and after 48 hours of 200 ( $n = 3$ ) or  
22 1,000 ( $n = 4$ ) mg/kg TCE in female mice, similar values of 0.30–0.69% of hepatocytes were  
23 reported as undergoing DNA synthesis in primary culture. Only the 1,000 mg/kg TCE dose in  
24 male mice at 48 hours was reported to give a result considered to be positive (~2.2% of  
25 hepatocytes) but no statistical analyses were performed on these measurements. These results  
26 are limited by both the number of animals examined and the relevance of the paradigm.

27 As noted above, TCE treatment in rodents has been reported to result in hepatocellular  
28 hypertrophy and increased centrilobular eosinophilia. Elcombe et al. (1985) reported a small  
29 decrease in DNA content with TCE treatment (consistent with hepatocellular hypertrophy) that  
30 was not dose-related, increased tritiated thymidine incorporation in whole mouse liver DNA that  
31 was that was treatment but not dose-related (i.e., a two-, two-, and fivefold of control in mice  
32 treated with 500, 1,000, and 1,500 mg/kg TCE), and slightly increased numbers of mitotic  
33 figures that were treatment but not dose-related and not correlated with DNA synthesis as  
34 measured by thymidine incorporation. Elcombe et al., reported no difference in response  
35 between 500 and 1,000 mg/kg TCE treatments for tritiated thymidine incorporation. Dees and

1 Travis (1993) also reported that incorporation of tritiated thymidine in DNA from mouse liver  
2 was elevated after TCE treatment with the mean peak level of tritiated thymidine incorporation  
3 occurred at 250 mg/kg TCE treatment level and remaining constant for the 500 and 1,000 mg/kg  
4 treated groups. Dees and Travis (1993) specifically report that mitotic figures, although very  
5 rare, were more frequently observed after TCE treatment, found most often in the intermediate  
6 zone, and found in cells resembling mature hepatocytes. They reported that there was little  
7 tritiated thymidine incorporation in areas near the bile duct epithelia or close to the portal triad in  
8 liver sections from both male and female mice. Channel et al. (1998) reported proliferating cell  
9 nuclear antigen (PCNA) positive cells, a measure of cells that have undergone DNA synthesis,  
10 was elevated only on Day 10 (out of the 21 studied) and only in the 1,200 mg/kg-day TCE  
11 exposed group with a mean of ~60 positive nuclei per 1,000 nuclei for six mice (~6%). Given  
12 that there was little difference in PCNA positive cells at the other TCE doses or time points  
13 studied, the small number of affected cells in the liver could not account for the increase in liver  
14 size reported in other experimental paradigms at these doses. The PCNA positive cells as well as  
15 “mitotic figures” were reported to be present in centrilobular, midzonal, and periportal regions  
16 with no observed predilection for a particular lobular distribution. No data were shown  
17 regarding any quantitative estimates of mitotic figures and whether they correlated with PCNA  
18 results. Thus, whether the DNA synthesis phases of the cell cycle indicated by PCNA staining  
19 were indentifying polyploidization or increased cell number cannot be determined.

20 For both rats and mice, the data from Elcombe et al. (1985) showed that tritiated  
21 thymidine incorporation in total liver DNA observed after TCE exposure did not correlate with  
22 mitotic index activity in hepatocytes. Both Elcombe et al. (1985) and Dees and Travis (1993)  
23 reported a small mitotic indexes and evidence of periportal hepatocellular hypertrophy from TCE  
24 exposure. Neither mitotic index or tritiated thymidine incorporation data support a correlation  
25 with TCE-induced liver weight increase in the mouse, but rather the increase to be most likely  
26 due to hepatocellular hypertrophy. If higher levels of hepatocyte replication had occurred  
27 earlier, such levels were not sustained by 10 days of TCE exposure. These data suggest that  
28 increased tritiated thymidine levels were targeted to mature hepatocytes and in areas of the liver  
29 where greater levels of polyploidization occur (see Section E.1.1). Both Elcombe et al. (1985)  
30 and Dees and Travis (1993) show that tritiated thymidine incorporation in the liver was ~twofold  
31 greater than controls between 250–1,000 mg/kg TCE, a result consistent with a doubling of  
32 DNA. Thus, given the normally quiescent state of the liver, the magnitude of this increase over  
33 control levels, even if a result of proliferation rather than polyplodization, would be confined to  
34 a very small population of cells in the liver after 10 days of TCE exposure.

1           Laughter et al. (2004) reported that there was an increase in DNA synthesis after aqueous  
2 gavage exposure to 500 and 1,000 mg/kg TCE given as three boluses a day for 3 weeks with  
3 BrdU given for the last week of treatment. An examination of DNA synthesis in individual  
4 hepatocytes was reported to show that 1 and 4.5% of hepatocytes had undergone DNA synthesis  
5 in the last week of treatment for the 500 and 1,000 mg/kg doses, respectively. Again, this level  
6 of DNA synthesis is reported for a small percentage of the total hepatocytes in the liver and not  
7 reported to be a result of regenerative hyperplasia.

8           Finally, Dees and Travis (1993) and Channel et al. (1998) reported evaluating changes in  
9 apoptosis with TCE treatment. Dees and Travis (1993) enumerated identified by either  
10 hematoxylin and eosin or feulgen staining in male and female mice after 10 days of TCE  
11 treatment by. Only zero or one apoptosis was observed per 100 high power (400×) fields in  
12 controls and all dose groups except for those given 1,000 mg/kg-day, in which eight or nine  
13 apoptoses per 100 fields were reported. None of the apoptoses were in the intermediate zones  
14 where mitotic figures were observed, and all were located near the central veins. This is the  
15 same region where one would expect endogenous apoptoses as hepatocytes “stream” from the  
16 portal triad toward the central vein (Schwartz-Arad et al., 1989). In addition, this is the same  
17 region where Buben and O’Flaherty (1985) noted necrosis and polyploidy. By contrast Channel  
18 et al. (1998) reported no significant differences in apoptosis at any treatment dose (400–1,200  
19 mg/kg-day) examined after any time from 2 days to 4 weeks.

#### 4.5.4.1.4. Peroxisomal Proliferation and Related Effects

21           Numerous studies have reported that TCE administered to mice and rats by gavage leads  
22 to proliferation of peroxisomes in hepatocytes. Some studies have measured changes in the  
23 volume and number of peroxisomes as measures of peroxisome proliferation while others have  
24 measured peroxisomal enzyme activity such catalase and cyanide-insensitive PCO. Like liver  
25 weight, the determination of a baseline level of peroxisomal volume, number, or enzyme activity  
26 can be variable and have great effect on the ability to determine the magnitude of a  
27 treatment-related effect.

28           Elcombe et al. (1985) reported increases in the percentage of the cytoplasm occupied by  
29 peroxisomes in B6C3F1 and Alderley Park mice treated for 10 days at 500–1,500 mg/kg-day.  
30 Although the increase over controls appeared larger in the B6C3F1 strain, this is largely due to  
31 the twofold smaller control levels in that strain, as the absolute percentage of peroxisomal  
32 volume was similar between strains after treatment. All these results showed high variability, as  
33 evidenced from the reported standard deviations. Channel et al. (1998) found a similar absolute

1 percentage of peroxisomal volume after 10 days treatment in the B6C3F1 mouse at  
2 1,200 mg/kg-day TCE but with the percentage in vehicle controls similar to the Alderley-Park  
3 mice in the Elcombe et al. (1985) study. Interestingly, Channel et al. (1998) found that the  
4 increase in peroxisomes peaked at 10 days, with lower values after 6 and 14 days of treatment.  
5 Furthermore, the vehicle control levels also varied almost twofold depending on the number of  
6 days of treatment. Nakajima et al. (2000), who treated male wild-type SV129 mice at  
7 750 mg/kg-day for 14 days, found even higher baseline values for the percentage of peroxisomal  
8 volume, but with an absolute level after treatment similar to that reported by Channel et al.  
9 (1998) in B6C3F1 mice treated at 1,200 mg/kg-day TCE for 14 days. Nakajima et al. (2000)  
10 also noted that the treatment-related increases were smaller for female wild-type mice, and that  
11 there were no increases in peroxisomal volume in male or female PPAR $\alpha$ -null mice, although  
12 vehicle control levels were slightly elevated (not statistically significant). Only Elcombe et al.  
13 (1985) examined peroxisomal volume in rats, and reported smaller treatment-related increases in  
14 two strains (OM and AP), but higher baseline levels. In particular, at 1,000 mg/kg-day, after  
15 10 days treatment, the percentage peroxisomal volume was similar in OM and AP rats, with  
16 similar control levels as well. While the differences from treatment were not statistically  
17 significant, only five animals were used in each group, and variability, as can be seen by the  
18 standard deviations, was high, particularly in the treated animals.

19 The activities of a number of different hepatic enzymes have also been as markers for  
20 peroxisome proliferation and/or activation of PPAR $\alpha$ . The most common of these are catalase  
21 and cyanide-insensitive PCO. In various strains of mice (B6C3F1, Swiss albino, SV129  
22 wild-type) treated at doses of 500–2,000 mg/kg-day for 10–28 days, increases in catalase activity  
23 have tended to be more modest (1.3- to 1.6-fold of control) as compared to increases in PCO  
24 (1.4- to 7.9-fold of control) (Elcombe et al., 1985; Goel et al., 1992; Goldsworthy and Popp,  
25 1987; Laughter et al., 2004; Nakajima et al., 2000; Watanabe and Fukui, 2000). In rats, Elcombe  
26 et al. (1985) reported no increases in catalase or PCO activity in Alderley-Park rats treated at  
27 1,000 mg/kg-day TCE for 10 days. In F344 rats, Goldsworthy and Popp (1987) and Melnick  
28 et al. (1987) reported increases of up to twofold in catalase and 4.1-fold in PCO relative to  
29 controls treated at 600–4,800 mg/kg-day for 10–14 days. The changes in catalase were similar  
30 to those in mice at similar treatment levels, with 1.1- to 1.5-fold of control enzyme activities at  
31 doses of 1,000–1,300 mg/kg-day (Elcombe et al., 1985; Melnick et al., 1987). However, the  
32 changes in PCO were smaller, with 1.1- to 1.8-fold of control activity at these doses, as  
33 compared to 6.3- to 7.9-fold of control in mice (Goldsworthy and Popp, 1987; Melnick et al.,  
34 1987).

1 In SV129 mice, Nakajima et al. (2000) and Laughter et al. (2004) investigated the  
2 dependence of these changes on PPAR $\alpha$  by using a null mouse. Nakajima et al. reported that  
3 neither male nor female wild-type or PPAR $\alpha$  null mice had significant increases in catalase after  
4 14 days of treatment at 750 mg/kg-day. However, given the small number of animals (4 per  
5 group) and the relatively small changes in catalase observed in other (wild-type) strains of mice,  
6 this study had limited power to detect such changes. Several other markers of peroxisome  
7 proliferation, including acyl-CoA oxidase and CYP4A1 (PCO was not investigated), were  
8 induced by TCE in male wild-type mice, but not in male null mice or female mice of either type.  
9 Unfortunately, none of these markers have been investigated using TCE in female mice of any  
10 other strain, so it is unclear whether the lack of response is characteristic of female mice in  
11 general, or just in this strain. Interestingly, as noted above, liver/body weight ratio increases  
12 were observed in both sexes of the null mice in this study. Laughter et al. (2004) only quantified  
13 activity of the peroxisome proliferation marker PCO in their study, and found in null mice a  
14 slight decrease (0.8-fold of control) at 500 mg/kg-day TCE and an increase (1.5-fold of control)  
15 at 1,500 mg/kg-day TCE after 3 weeks of treatment, with neither statistically significant  
16 (4–5 mice per group). However, baseline levels of PCO were almost twofold higher in the null  
17 mice, and the treated wild-type and null mice differed in PCO activity by only about 1.5-fold.

18 In sum, oral administration of TCE for up to 28 days causes proliferation of peroxisomes  
19 in hepatocytes along with associated increases in peroxisomal enzyme activities in both mice and  
20 rats. Male mice tend to be more sensitive in that at comparable doses, rats and female mice tend  
21 to exhibit smaller responses. For example, for peroxisomal volume and PCO, the fold-increase  
22 in rats appears to be lower by three- to sixfold than that in mice, but, for catalase, the changes  
23 were similar between mice in F344 rats. No inhalation or longer-term studies were located, and  
24 only one study examined these changes at more than one time-point. Therefore, little is known  
25 about the route-dependence, time course, and persistence of these changes. Finally, two studies  
26 in PPAR $\alpha$ -null mice (Laughter et al., 2004; Nakajima et al., 2000) found diminished responses in  
27 terms of increased peroxisomal volume and peroxisomal enzyme activities as compared to  
28 wild-type mice, although there was some confounding due to baseline differences between null  
29 and wild-type control mice in several measures.

30

#### 4.5.4.1.5. Oxidative Stress

31 Several studies have attempted to study the possible effects of “oxidative stress” and  
32 DNA damage resulting from TCE exposures. The effects of induction of metabolism by TCE, as  
33 well as through coexposure to ethanol, have been hypothesized to in itself increase levels of



1 “oxidative stress” as a common effect for both exposures (see Sections E.3.4.2.3 and E.4.2.4).  
2 Oxidative stress has been hypothesized to be a key event or MOA for peroxisome proliferators as  
3 well, but has been found to neither be correlated with cell proliferation nor carcinogenic potency  
4 of peroxisome proliferators (see Section E.3.4.1.1). As a MOA, it is not defined or specific as  
5 the term “oxidative stress” is implicated as part of the pathophysiologic events in a multitude of  
6 disease processes and is part of the normal physiologic function of the cell and cell signaling.

7 In regard to measures of oxidative stress, Rusyn et al. (2006) noted that although an  
8 overwhelming number of studies draw a conclusion between chemical exposure, DNA damage,  
9 and cancer based on detection of 8-hydroxy-2’ deoxyguanosine (8-OHdG), a highly mutagenic  
10 lesion, in DNA isolated from organs of in vivo treated animals, a concern exists as to whether  
11 increases in 8-OHdG represent damage to genomic DNA, a confounding contamination with  
12 mitochondrial DNA, or an experimental artifact. As noted in Sections E.2.1.1 and E.2.2.11,  
13 studies of TCE which employ the i.p. route of administration can be affected by inflammatory  
14 reactions resulting from that routes of administration and subsequent toxicity that can involve  
15 oxygen radical formation from inflammatory cells. Finally, as described in Section E.2.2.8, the  
16 study by Channel et al. (1998) demonstrated that corn oil as vehicle had significant effects on  
17 measures of “oxidative stress” such as thiobarbiturate acid-reactive substances (TBARS).

18 The TBARS results presented by Channel et al. (1998) indicate suppression of TBARS  
19 with increasing time of exposure to corn oil alone with data presented in such a way for 8-OHdG  
20 and total free radical changes that the pattern of corn oil administration was obscured. It was not  
21 apparent from that study that TCE exposure induced oxidative damage in the liver.

22 Toraason et al. (1999) measured 8-OHdG and a “free radical-catalyzed isomer of  
23 arachidonic acid and marker of oxidative damage to cell membranes, 8-Epi-prostaglandin F2 $\alpha$   
24 (8-epiPGF)”, excretion in the urine and TBARS (as an assessment of malondialdehyde and  
25 marker of lipid peroxidation) in the liver and kidney of male Fischer rats exposed to single i.p.  
26 injections in of TCE in Alkamuls vehicle. Using this paradigm, 500-mg/kg TCE was reported to  
27 induce Stage II anesthesia and a 1,000 mg/kg TCE to induce Level III or IV (absence of reflex  
28 response) anesthesia and burgundy colored urine with 2/6 rats at 24 hours comatose and  
29 hypothermic. The animals were sacrificed before they could die and the authors suggested that  
30 they would not have survived another 24 hours. Thus, using this paradigm there was significant  
31 toxicity and additional issues related to route of exposure. Urine volume declined significantly  
32 during the first 12 hours of treatment and while water consumption was not measured, it was  
33 suggested by the authors to be decreased due to the moribundity of the rats. Given that this study  
34 examined urinary markers of “oxidative stress” the effects on urine volume and water  
35 consumption, as well as the profound toxicity induced by this exposure paradigm, limit the

1 interpretation of the study. The issues of bias in selection of the data for this analysis, as well as  
2 the issues stated above for this paradigm limit interpretation of these data while the authors  
3 suggest that evidence of oxidative damage was equivocal.  
4

#### 4.5.4.1.6. Bile Production

5 Effects of TCE exposure in humans and in experimental animals is presented in  
6 Section E.2.6. Serum bile acids (SBA) have been suggested as a sensitive indicator of  
7 hepatotoxicity to a variety of halogenated solvents with an advantage of increased sensitivity and  
8 specificity over conventional liver enzyme tests that primarily reflect the acute perturbation of  
9 hepatocyte membrane integrity and “cell leakage” rather than liver functional capacity (i.e.,  
10 uptake, metabolism, storage, and excretion functions of the liver) (Neghab et al., 1997) Bai et al.,  
11 1992b. While some studies have reported negative results, a number of studies have reported  
12 elevated SBA in organic solvent-exposed workers in the absence of any alterations in normal  
13 liver function tests. These variations in results have been suggested to arise from failure of some  
14 methods to detect some of the more significantly elevated SBA and the short-lived and reversible  
15 nature of the effect (Neghab et al., 1997). Neghab et al. (1997) have reported that occupational  
16 exposure to 1,1,2-trichloro-1,2,2-trifluoroethane and trichloroethylene has resulted in elevated  
17 SBA and that several studies have reported elevated SBA in experimental animals to chlorinated  
18 solvents such as carbon tetrachloride, chloroform, hexachlorobutadiene, tetrachloroethylene,  
19 1,1,1-trichloroethane, and trichloroethylene at levels that do not induce hepatotoxicity (Bai et al.,  
20 1992a; Bai et al., 1992b; Hamdan and Stacey, 1993; Wang and Stacey, 1990). Toluene, a  
21 nonhalogenated solvent, has also been reported to increase SBA in the absence of changes in  
22 other hepatobiliary functions (Neghab and Stacey, 1997). Thus, disturbance in SBA appears to  
23 be a generalized effect of exposure to chlorinated solvents and nonchlorinated solvents and not  
24 specific to TCE exposure.

25 Wang and Stacey (1990) administered TCE in corn oil via i.p. injection to male  
26 Sprague-Dawley rats with liver enzymes and SBA examined 4 hours after the last TCE  
27 treatment. The limitations of i.p injection experiments have already been discussed. While  
28 reporting no overt liver toxicity there was, generally, a reported dose-related increase in cholic  
29 acid, chenodeoxycholic acid, deoxycholic acid, taurocholic acid, tauroursodeoxycholic acid with  
30 cholic acid and taurocholic acid increased at the lowest dose. The authors report that  
31 “examination of liver sections under light microscopy yielded no consistent effects that could be  
32 ascribed to trichloroethylene.” In the same study a rats were also exposed to TCE via inhalation  
33 and using this paradigm, cholic acid and taurocholic acid were also significantly elevated but the

1 large variability in responses between rats and the low number of rats tested in this paradigm  
2 limit its ability to determine quantitative differences between groups. Nevertheless, without the  
3 complications associated with i.p. exposure, inhalation exposure of TCE at relatively low  
4 exposure levels that were not associated with other measures of toxicity *were* associated with  
5 increased SBA level.

6 Hamdan and Stacey (1993) administered TCE in corn oil (1 mmol/kg) in male Sprague-  
7 Dawley rats and followed the time-course of SBA elevation, TCE concentration, and  
8 trichloroethanol in the blood up to 16 hours. Liver and blood concentration of TCE were  
9 reported to peak at 4 hours while those of trichloroethanol peaked at 8 hours after dosing. TCE  
10 levels were not detectable by 16 hours in either blood or liver while those of trichloroethanol  
11 were still elevated. Elevations of SBA were reported to parallel those of TCE with cholic acid  
12 and taurochloate acid reported to show the highest levels of bile acids. The authors state that  
13 liver injury parameters were checked and found unaffected by TCE exposure but did not show  
14 the data. Thus, it was TCE concentration and not that of its metabolite that was most closely  
15 related to changes in SBA and after a single exposure and the effect appeared to be reversible. In  
16 an in vitro study by Bai and Stacey (1993), TCE was studied in isolated rat hepatocytes with  
17 TCE reported to cause a dose-related suppression of initial rates of cholic acid and taurocholic  
18 acid but with no significant effects on enzyme leakage and intracellular calcium contents, further  
19 supporting a role for the parent compound in this effect.

#### 20 **4.5.4.1.7. Summary: Trichloroethylene (TCE)-Induced Noncancer Effects in Laboratory Animals**

21 In laboratory animals, TCE leads to a number of structural changes in the liver, including  
22 increased liver weight, small transient increases in DNA synthesis, cytomegaly in the form of  
23 “swollen” or enlarged hepatocytes, increased nuclear size probably reflecting polyploidization,  
24 and proliferation of peroxisomes. Liver weight increases proportional to TCE dose are  
25 consistently reported across numerous studies, and appear to be accompanied by periportal  
26 hepatocellular hypertrophy. There is also evidence of increased DNA synthesis in a small  
27 portion of hepatocytes at around 10 days in vivo exposure. The lack of correlation of  
28 hepatocellular mitotic figures with whole liver DNA synthesis or DNA synthesis observed in  
29 individual hepatocytes supports the conclusion that cellular proliferation is not the predominant  
30 cause of increased DNA synthesis. The lack of correlation of whole liver DNA synthesis and  
31 those reported for individual hepatocytes suggests that nonparenchymal cells also contribute to  
32 such synthesis. Indeed, nonparenchymal cell activation or proliferation has been noted in several

1 studies. Moreover, the histological descriptions of TCE exposed liver are consistent with and in  
2 some cases specifically note increased polyploidy after TCE exposure. Interestingly, changes in  
3 TCE-induced hepatocellular ploidy, as indicated by histological changes in nuclei, have been  
4 noted to remain after the cessation of exposure. In regard to apoptosis, TCE has been reported to  
5 either not change apoptosis or to cause a slight increase at high doses. Some studies have also  
6 noted effects from dosing vehicle alone (such as corn oil in particular) not only on liver  
7 pathology, but also on DNA synthesis.

8 Available data also suggest that TCE does not induce substantial cytotoxicity, necrosis, or  
9 regenerative hyperplasia, as only isolated, focal necroses and mild to moderate changes in serum  
10 and liver enzyme toxicity markers having been reported. Data on peroxisome proliferation,  
11 along with increases in a number of associated biochemical markers, show effects in both mice  
12 and rats. These effects are consistently observed across rodent species and strains, although the  
13 degree of response at a given mg/kg-day dose appears to be highly variability across strains, with  
14 mice on average appearing to be more sensitive.

15 In addition, like humans, laboratory animals exposed to TCE have been observed to have  
16 increased serum bile acids, though the toxicologic importance of these effects is unclear.

17

#### 4.5.5. Trichloroethylene (TCE)-Induced Liver Cancer in Laboratory Animals

18 For 2-year or lifetime studies of TCE exposure a consistent hepatocarcinogenic response  
19 has been observed using mice of differing strains and genders and from differing routes of  
20 exposure. However, some rat studies have been confounded by mortality from gavage error or  
21 the toxicity of the dose of TCE administered. In some studies, a relative insensitive strain of rat  
22 has been used. However, in general it appears that the mouse is more sensitive than the rat to  
23 TCE-induced liver cancer. Three studies give results the authors consider to be negative for  
24 TCE-induced liver cancer in mice, but have either design and/or reporting limitations, or are in  
25 strains and paradigms with apparent low ability for liver cancer induction or detection. Findings  
26 from these studies are shown in Tables 4-60 through 4-65, and discussed below.

27

##### 4.5.5.1.1. Negative or Inconclusive Studies of Mice and Rats

28 Fukuda et al. (1983) reported a 104-week inhalation bioassay in female Crj:CD-1 (ICR)  
29 mice and female Crj:CD (S-D) rats exposed to 0-, 50-, 150-, and 450-ppm TCE ( $n = 50$ ). There  
30 were no reported incidences of mice or rats with liver tumors for controls indicative of relatively  
31 insensitive strains and gender used in the study for liver effects. While TCE was reported to

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 induce a number of other tumors in mice and rats in this study, the incidence of liver tumors was  
2 less than 2% after TCE exposure. Of note is the report of cystic cholangioma reported in one  
3 group of rats.

4 Henschler et al. (1980) exposed NMRI mice and WIST random bred rats to 0-, 100-, and  
5 500-ppm TCE for 18 months ( $n = 30$ ). Control male mice were reported to have  
6 one hepatocellular carcinoma and one hepatocellular adenoma with the incidence rate unknown.  
7 In the 100-ppm TCE exposed group, two hepatocellular adenomas and one mesenchymal liver  
8 tumor were reported. No liver tumors were reported at any dose of TCE in female mice or  
9 controls. For male rats, only one hepatocellular adenomas at 100 ppm was reported. For female  
10 rats no liver tumors were reported in controls, but one adenoma and one cholangiocarcinoma was  
11 reported at 100-ppm TCE and at 500-ppm TCE, two cholangioadenomas, a relatively rare biliary  
12 tumor, was reported. The difference in survival in mice, did not affect the power to detect a  
13 response, as was the case for rats. However, the low number of animals studied, abbreviated  
14

1  
2  
3

**Table 4-60. Summary of liver tumor findings in gavage studies of trichloroethylene by NTP (1990)<sup>a</sup>**

<b>Sex</b>	<b>Dose (mg/kg)<sup>b</sup></b>	<b>Adenoma (overall; terminal<sup>c</sup>)</b>	<b>Adenocarcinoma (overall; terminal<sup>c</sup>)</b>
1/d, 5 d/wk, 103-wk study, F344/N rats			
Male	0	NA <sup>d</sup>	0/49
	500	NA	0/49
	1,000	NA	1/49
Female	0	NA	0/50
	500	NA	1/48
	1,000	NA	1/48
1/d, 5 d/wk, 103-wk study, B6C3F1 mice			
Male	0	7/48; 6/33	8/48; 6/33
	1,000	14/50; 6/16	31/50; 14/16 <sup>f</sup>
Female	0	4/48; 4/32	2/48; 2/32
	1,000	16/49; 11/23 <sup>e</sup>	13/49; 8/23 <sup>g</sup>

4  
5  
6  
7  
8  
9  
10  
11  
12

<sup>a</sup> Liver tumors not examined in 13-wk study, so data shown only for 103-wk study.

<sup>b</sup> Corn oil vehicle.

<sup>c</sup> Terminal values not available for rats.

<sup>d</sup> Data not available.

<sup>e</sup>  $p < 0.003$ .

<sup>f</sup>  $p < 0.001$ .

<sup>g</sup>  $p \leq 0.002$ .

1  
2  
3

**Table 4-61. Summary of liver tumor findings in gavage studies of trichloroethylene by NCI (1976)**

<b>Sex</b>	<b>Dose (mg/kg)<sup>a</sup></b>	<b>Hepatocarcinoma</b>
1/d, 5 d/wk, 2-yr study, Osborn-Mendel rats		
Males	0	0/20
	549	0/50
	1,097	0/50
Females	0	0/20
	549	1/48
	1,097	0/50
1/d, 5 d/wk, 2-yr study, B6C3F1 mice		
Males	0	1/20
	1,169	26/50 <sup>b</sup>
	2,339	31/48 <sup>b</sup>
Females	0	0/20
	869	4/50
	1,739	11/47 <sup>b</sup>

4  
5  
6  
7  
8  
9

<sup>a</sup>Treatment period was 48 wk for rats, 66 wk for mice. Doses were changed several times during the study based on monitoring of body weight changes and survival. Dose listed here is the time-weighted average dose over the days on which animals received a dose.

<sup>b</sup> $p < 0.01$ .

1  
2  
3

**Table 4-62. Summary of liver tumor incidence in gavage studies of trichloroethylene by NTP (1988)**

<b>Sex</b>	<b>Dose (mg/kg)<sup>a</sup></b>	<b>Adenoma</b>	<b>Adenocarcinoma</b>
1/d, 5 d/wk, 2-yr study, ACI rats			
Male	0	0/50	1/50
	500	0/49	1/49
	1,000	0/49	1/49
Female	0	0/49	2/49
	500	0/46	0/46
	1,000	0/39	0/39
1/d, 5 d/wk, 2-yr study, August rats			
Male	0	0/50	0/50
	500	0/50	1/50
	1,000	0/48	1/48
Female	0	0/48	2/48
	500	0/48	0/48
	1,000	0/50	0/50
1/d, 5 d/wk, 2-yr study, Marshall rats			
Male	0	1/49	1/49
	500	0/50	0/50
	1,000	0/47	1/47
Female	0	0/49	0/49
	500	0/48	0/48
	1,000	0/46	0/46
1/d, 5 d/wk, 2-yr study, Osborne-Mendel rats			
Male	0	1/50	1/50
	500	1/50	0/50
	1,000	1/49	2/49
Female	0	0/50	0/50
	500	0/48	2/48
	1,000	0/49	2/49

4



1   <sup>a</sup> Corn oil vehicle.  
2

1  
2  
3

**Table 4-63. Summary of liver tumor findings in inhalation studies of trichloroethylene by Maltoni et al. (1988)<sup>a</sup>**

Sex	Concentration (ppm)	Hepatoma
7 h/d, 5 d/wk, 8-wk exposure, observed for lifespan, Swiss mice		
Male	0	1/100
	100	3/60
	600	4/72
Female	0	1/100
	100	1/60
	600	0/72
7 h/d, 5 d/wk, 78-wk exposure, observed for lifespan, Swiss mice		
Male	0	4/90
	100	2/90
	300	8/90
	600	13/90
Female	0	0/90
	100	0/90
	300	0/90
	600	1/90
7 h/d, 5 d/wk, 78-wk exposure, observed for lifespan, B6C3F1 mice <sup>b</sup>		
Male	0	1/90
	100	1/90
	300	3/90
	600	6/90
Female	0	3/90
	100	4/90
	300	4/90
	600	9/90

4  
5  
6  
7  
8  
9  
10  
11

<sup>a</sup>Three inhalation experiments in this study found no hepatomas: BT302 (8-wk exposure to 0, 100, 600 ppm in Sprague-Dawley rats); BT303 (8-wk exposure to 0, 100, or 600 ppm in Swiss mice); and BT304 (78-wk exposure to 0, 100, 300, or 600 ppm in Sprague-Dawley rats).

<sup>b</sup>Female incidences are from experiment BT306, while male incidences are from experiment BT306bis, which was added to the study because of high, early mortality due to aggressiveness and fighting in males in experiment BT306.

1  
2  
3

**Table 4-64. Summary of liver tumor findings in inhalation studies of trichloroethylene by Henschler et al. (1980)<sup>a</sup> and Fukuda et al. (1983)**

Sex	Concentration (ppm)	Adenomas	Adenocarcinomas
6 h/d, 5 d/wk, 18-mo exposure, 30-mo observation, Han:NMRI mice (Henschler et al., 1980)			
Males	0	1/30 <sup>b</sup>	1/30
	100	2/29 <sup>b</sup>	0/30
	500	0/29	0/30
Females	0	0/29	0/29
	100	0/30	0/30
	500	0/28	0/28
6 h/d, 5 d/wk, 18-mo exposure, 36-mo observation, Han:WIST rats (Henschler et al., 1980)			
Males	0	1/29	0/29
	100	1/30	0/30
	500	0/30	0/30
Females	0	0/28	0/28
	100	1/30	1/30
	500	2/30	0/30
7 h/d, 5 d/wk, 2-yr study, Crj:CD (S-D) rats (Fukuda et al., 1983)			
Females	0	0/50	0/50
	50	1/50	0/50
	150	0/47	0/47
	450	0/51	1/50
7 h/d, 5 d/wk, 2-yr study, Crj:CD (ICR) mice (Fukuda et al., 1983)			
Females	0	0/49	0/49
	50	0/50	0/50
	150	0/50	0/50
	450	1/46	0/46

4  
5  
6  
7

<sup>a</sup> Henschler et al. (1980) observed no liver tumors in control or exposed Syrian hamsters.

<sup>b</sup> One additional hepatic tumor of undetermined class not included.

**Table 4-65. Summary of liver tumor findings in gavage studies of trichloroethylene by Henschler et al. (1984)<sup>a</sup>**

<b>Sex (TCE conc.)</b>	<b>TCE (Stabilizers if present)</b>	<b>Benign<sup>b</sup></b>	<b>Malignant<sup>c</sup></b>
5 d/wk, 18-mo exposure, 24-mo observation, Swiss mice (Henschler et al., 1984)			
Males (2.4g/kg BW)	Control (none)	5/50	0/50
	TCE (triethanolamine)	7/50	0/50
	TCE (industrial)	9/50	0/50
	TCE (epichlorohydrin (0.8%))	3/50	1/50
	TCE (1,2-epoxybutane [0.8%])	4/50	0/50
	TCE (both epichlorohydrin [0.25%] and 1,2-epoxybutane [0.25%])	5/50	0/50
Females (1.8 g/kg BW)	Control (none)	1/50	0/50
	TCE (triethanolamine)	7/50	0/50
	TCE (industrial)	9/50	0/50
	TCE (epichlorohydrin (0.8%))	3/50	0/50
	TCE (1,2-epoxybutane (0.8%))	2/50	0/50
	TCE (both epichlorohydrin (0.25%) and 1,2-epoxybutane (0.25%))	4/50	1/50

<sup>a</sup> Henschler et al. (1984) Due to poor condition of the animals resulting from the nonspecific toxicity of high doses of TCE and/or the additives, gavage was stopped for all groups during wk 35–40, 65 and 69–78, and all doses were reduced by a factor of 2 from the 40<sup>th</sup> wk on.

<sup>b</sup> Includes hepatocellular adenomas, hemangioendothelioma, cholangiocellular adenoma.

<sup>c</sup> Includes hepatocellular carcinoma, malignant hemangiosarcoma, cholangiocellular carcinoma.

Conc. = concentration.

exposure duration, low survival in rats, and absent background response (suggesting low intrinsic sensitivity to this endpoint) suggest a study of limited ability to detect a TCE carcinogenic liver response. Of note is that despite their limitations, both Fukuda et al. (1983) and Henschler et al. (1980) report rare biliary cell derived tumors in TCE-exposed rats.

Van Duuren et al. (1979), exposed mice to 0.5 mg/mouse to TCE via gavage once a week in 0.1 mL trioctanion (*n* = 30). Inadequate design and reporting of this study limit that ability to use the results as an indicator of TCE carcinogenicity.

1 The NCI (1976) study of TCE was initiated in 1972 and involved the exposure of  
2 Osborn-Mendel rats to varying concentrations of TCE. A low incidence of liver tumors was  
3 reported for controls and carbon tetrachloride positive controls in rats from this study. The  
4 authors concluded that due to mortality, “the test is inconclusive in rats.” They note the  
5 insensitivity of the rat strain used to the positive control of carbon tetrachloride exposure.

6 The NTP (1990) study of TCE exposure in male and female F344/N rats, and B6C3F1  
7 mice (500 and 1,000 mg/kg for rats) is limited in the ability to demonstrate a dose-response for  
8 hepatocarcinogenicity. For rats, the NTP (1990) study reported no treatment-related  
9 non-neoplastic liver lesions in males and a decrease in basophilic cytological change reported  
10 from TCE-exposure in female rats. The results for detecting a carcinogenic response in rats were  
11 considered to be equivocal because both groups receiving TCE showed significantly reduced  
12 survival compared to vehicle controls and because of a high rate (e.g., 20% of the animals in the  
13 high-dose group) of death by gavage error.

14 The NTP (1988) study of TCE exposure in four strains of rats to  
15 “diisopropylamine-stabilized TCE” was also considered inadequate for either comparing or  
16 assessing TCE-induced liver carcinogenesis in these strains of rats because of chemically  
17 induced toxicity, reduced survival, and incomplete documentation of experimental data. TCE  
18 gavage exposures of 0, 500, or 1,000 mg/kg-day (5 days/week, for 103 weeks) male and female  
19 rats was also marked by a large number of accidental deaths (e.g., for high-dose male Marshal  
20 rats 25 animals were accidentally killed).

21 Maltoni et al. (1986) reported the results of several studies of TCE via inhalation and  
22 gavage in mice and rats. A large number of animals were used in the treatment groups but the  
23 focus of the study was detection of a neoplastic response with only a generalized description of  
24 tumor pathology phenotype given and limited reporting of non-neoplastic changes in the liver.  
25 Accidental death by gavage error was reported not to occur in this study. In regards to effects of  
26 TCE exposure on rat survival, “a nonsignificant excess in mortality correlated to TCE treatment  
27 was observed only in female rats (treated by ingestion with the compound)”.

28 For rats, Maltoni et al. (1986) reported four liver angiosarcomas (one in a control male  
29 rat, one both in a TCE-exposed male and female at 600 ppm TCE for 8 weeks, and one in a  
30 female rat exposed to 600-ppm TCE for 104 weeks), but the specific results for incidences of  
31 hepatocellular “hepatomas” in treated and control rats were not given. Although the Maltoni  
32 et al. (1986) concluded that the small number was not treatment related, the findings were  
33 brought forward because of the extreme rarity of this tumor in control Sprague-Dawley rats,  
34 untreated or treated with vehicle materials. In rats treated for 104 weeks, there was no report of a  
35 TCE treatment-related increase in liver cancer in rats. This study only presented data for positive

1 findings so it did not give the background or treatment-related findings in rats for liver tumors in  
2 this study. Thus, the extent of background tumors and sensitivity for this endpoint cannot be  
3 determined. Of note is that the Sprague-Dawley strain used in this study was also noted in the  
4 Fukuda et al. (1983) study to be relatively insensitive for spontaneous liver cancer and to also be  
5 negative for TCE-induced hepatocellular liver cancer induction in rats. However, like Fukuda  
6 et al. (1983) and Henschler et al. (1980), that reported rare biliary tumors in insensitive strains of  
7 rat for hepatocellular tumors, Maltoni et al. (1986) reported a relatively rare tumor type,  
8 angiosarcoma, after TCE exposure in a relatively insensitive strain for “hepatomas.” As noted  
9 above, many of the rat studies were limited by premature mortality due to gavage error or  
10 premature mortality (Henschler et al., 1980; NCI, 1976; NTP, 1988, 1990), which was reported  
11 not occur in Maltoni et al. (1986).  
12

#### **4.5.5.1.2. Positive Trichloroethylene (TCE) Studies of Mice**

13 In the NCI (1976) study of TCE exposure in B6C3F1 mice, TCE was reported to increase  
14 incidence of hepatocellular carcinomas in both doses and both genders of mice (~1,170 and  
15 2,340 mg/kg for males and 870 and 1,740 mg/kg for female mice). Hepatocellular carcinoma  
16 diagnosis was based on histologic appearance and metastasis to the lung. The tumors were  
17 described in detail and to be heterogeneous “as described in the literature” and similar in  
18 appearance to tumors generated by carbon tetrachloride. The description of liver tumors in this  
19 study and tendency to metastasize to the lung are similar to descriptions provided by  
20 Maltoni et al. (1986) for TCE-induced liver tumors in mice via inhalation exposure.

21 The NTP (1990) study of TCE exposure in male and female B6C3F1 mice (1,000 mg/kg  
22 for mice) reported decreased latency of liver tumors, with animals first showing carcinomas at  
23 57 weeks for TCE-exposed animals and 75 weeks for control male mice. The administration of  
24 TCE was also associated with increased incidence of hepatocellular carcinoma (tumors with  
25 markedly abnormal cytology and architecture) in male and female mice. Hepatocellular  
26 adenomas were described as circumscribed areas of distinctive hepatic parenchymal cells with a  
27 perimeter of normal appearing parenchyma in which there were areas that appeared to be  
28 undergoing compression from expansion of the tumor. Mitotic figures were sparse or absent but  
29 the tumors lacked typical lobular organization. Hepatocellular carcinomas had markedly  
30 abnormal cytology and architecture with abnormalities in cytology cited as including increased  
31 cell size, decreased cell size, cytoplasmic eosinophilia, cytoplasmic basophilia, cytoplasmic  
32 vacuolization, cytoplasmic hyaline bodies, and variations in nuclear appearance. Furthermore, in  
33 many instances several or all of the abnormalities were present in different areas of the tumor

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 and variations in architecture with some of the hepatocellular carcinomas having areas of  
2 trabecular organization. Mitosis was variable in amount and location. Therefore, the phenotype  
3 of tumors reported from TCE exposure was heterogeneous in appearance between and within  
4 tumors. However, because it consisted of a single-dose group in addition to controls, this study  
5 is limited of limited utility for analyzing the dose-response for hepatocarcinogenicity. There was  
6 also little reporting of non-neoplastic pathology or toxicity and no report of liver weight at  
7 termination of the study.

8 Maltoni et al. (1986) reported the results of several studies of TCE in mice. A large  
9 number of animals were used in the treatment groups but the focus of the study was detection of  
10 a neoplastic response with only a generalized description of tumor pathology phenotype given  
11 and limited reporting of non-neoplastic changes in the liver. There was no accidental death by  
12 gavage error reported to occur in mice but, a “nonsignificant” excess in mortality correlated to  
13 TCE treatment was observed in male B6C3F1 mice. TCE-induced effects on body weight were  
14 reported to be absent in mice except for one experiment (BT 306 bis) in which a slight nondose  
15 correlated decrease was found in exposed animals. “Hepatoma” was the term used to describe  
16 all malignant tumors of hepatic cells, of different subhistotypes, and of various degrees of  
17 malignancy and were reported to be unique or multiple, and have different sizes (usually  
18 detected grossly at necropsy) from TCE exposure. In regard to phenotype tumors were described  
19 as usual type observed in Swiss and B6C3F1 mice, as well as in other mouse strains, either  
20 untreated or treated with hepatocarcinogens and to frequently have medullary (solid), trabecular,  
21 and pleomorphic (usually anaplastic) patterns. Swiss mice from this laboratory were reported to  
22 have a low incidence of hepatomas without treatment (1%). The relatively larger number of  
23 animals used in this bioassay ( $n = 90-100$ ), in comparison to NTP standard assays, allows for a  
24 greater power to detect a response.

25 TCE exposure for 8 weeks via inhalation at 100 or 600 ppm may have been associated  
26 with a small increase in liver tumors in male mice in comparison to concurrent controls during  
27 the life span of the animals. In Swiss mice exposed to TCE via inhalation for 78 weeks, there a  
28 reported increase in hepatomas associated with TCE treatment that was dose-related in male but  
29 not female Swiss mice. In B6C3F1 mice exposed via inhalation to TCE for 78 weeks, increases  
30 in hepatomas were reported in both males and females. However, the experiment in males was  
31 repeated with B6C3F1 mice from a different source, since in the first experiment more than half  
32 of the mice died prematurely due to excessive fighting. Although the mice in the two  
33 experiments in males were of the same strain, the background level of liver cancer was  
34 significantly different between mice from the different sources (1/90 vs. 19/90), though the early  
35 mortality may have led to some censoring. The finding of differences in response in animals of

1 the same strain but from differing sources has also been reported in other studies for other  
2 endpoints. However, for both groups of male B6C3F1 mice the background rate of liver tumors  
3 over the lifetime of the mice was no greater than about 20%.

4 There were other reports of TCE carcinogenicity in mice from chronic exposures that  
5 were focused primarily on detection of liver tumors with limited reporting of tumor phenotype or  
6 non-neoplastic pathology. Herren-Freund et al. (1987) reported that male B6C3F1 mice given  
7 40 mg/L TCE in drinking water had increased tumor response after 61 weeks of exposure.  
8 However, concentrations of TCE fell by about ½ at this dose of TCE during the twice a week  
9 change in drinking water solution so the actual dose of TCE the animals received was less than  
10 40 mg/L. The percentage liver/body weight was reported to be similar for control and  
11 TCE-exposed mice at the end of treatment. However, despite difficulties in establishing  
12 accurately the dose received, an increase in adenomas per animal and an increase in the number  
13 of animals with hepatocellular carcinomas were reported to be associated with TCE exposure  
14 after 61 weeks of exposure and without apparent hepatomegaly. Anna et al. (1994) reported  
15 tumor incidences for male B6C3F1 mice receiving 800 mg/kg-day TCE via gavage (5 days/week  
16 for 76 weeks). All TCE-treated mice were reported to be alive after 76 weeks of treatment.  
17 Although the control group contained a mixture of exposure durations (76–134 weeks) and  
18 concurrent controls had a very small number of animals, TCE-treatment appeared to increase the  
19 number of animals with adenomas, the mean number of adenomas and carcinomas, but with no  
20 concurrent TCE-induced cytotoxicity.

#### 21 **4.5.5.1.3. Summary: Trichloroethylene (TCE)-Induced Cancer in Laboratory Animals**

22 Chronic TCE bioassays have consistently reported increased liver tumor incidences in  
23 both sexes of B6C3F1 mice treated by inhalation and gavage exposure in a number of bioassays.  
24 The only inhalation study of TCE in Swiss mice also showed an effect in males. Data in the rat,  
25 while not reporting statistically significantly increased risks, are not entirely adequate due to low  
26 numbers of animals, inadequate reporting, use of insensitive bioassays, increased systemic  
27 toxicity, and/or increased mortality. Notably, several studies in rats noted a few very rare types  
28 of liver or biliary tumors (cystic cholangioma, cholangiocarcinoma, or angiosarcomas) in treated  
29 animals.

#### **4.5.6. Role of Metabolism in Liver Toxicity and Cancer**

30 It is generally thought that TCE oxidation by CYPs is necessary for induction of  
31 hepatotoxicity and hepatocarcinogenicity (Bull, 2000). Direct evidence for this hypothesis is

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 limited, e.g., the potentiation of hepatotoxicity by pretreatment with CYP inducers such as  
2 ethanol and phenobarbital (Nakajima et al., 1988; Okino et al., 1991). Rather the presumption  
3 that CYP-mediated oxidation is necessary for TCE hepatotoxicity and hepatocarcinogenicity is  
4 largely based on similar effects (e.g., increases in liver weight, peroxisome proliferation, and  
5 hepatocarcinogenicity) having been observed with TCE's oxidative metabolites. The discussion  
6 below focuses the similarities and differences between the major effects in the liver of TCE and  
7 of the oxidative metabolites CH, TCA, and DCA. In addition, CH is largely converted to TCOH,  
8 TCA, and possibly DCA. DCA has been used in human clinical practice for a variety of severe  
9 illnesses and no data on liver effects in humans have been reported (U.S. EPA, 2003b).  
10 However, as noted in EPA (2003b), data on DCA in humans are scarce and complicated by the  
11 fact that available studies have predominantly focused on individuals who have a pre-existing  
12 (usually severe) disease.  
13

#### **4.5.6.1.1. Pharmacokinetics of Chloral Hydrate (CH), Trichloroacetic Acid (TCA), and Dichloroacetic Acid (DCA) From Trichloroethylene (TCE) Exposure**

14 As discussed in Section 3, in vivo data confirm that CH and TCA, are oxidative  
15 metabolites of TCE, with available data on TCA incorporated into the PBPK modeling. In  
16 addition, there are indirect data suggesting the formation of DCA. However, direct in vivo  
17 evidence of the formation of DCA is confounded by its rapid clearance at low concentrations,  
18 and analytical artifacts in its detection in vivo that have yet to be entirely resolved. PBPK  
19 modeling (see Section 3.5) predicts that the proportions of TCE metabolized to CH and TCA  
20 varies considerably in mice (ranging from 15–97 and 4–38%, respectively) and rats (ranging  
21 7–75 and 0.5–22%, respectively). Therefore, a range of smaller concentrations of TCA or CH  
22 may be relevant for comparisons with TCE-induced liver effects. For example, for  
23 1,000 mg/kg-day oral doses of TCE, the relevant comparisons would be approximately  
24 0.25–1.5 g/L in drinking water for TCA and CH. For DCA a corresponding range is harder to  
25 determine and has been suggested to be an upper limit of about 12% following oral exposures  
26 (Barton et al., 1999). This is consistent with the range estimated from PBPK modeling  
27 attributing all of the “untracked” oxidation (i.e., not producing TCOH or TCA) to DCA (95% CI:  
28 0.2–16%, see Figure 3-22).

29 Two studies have used analytic methods for DCA that are considered more reliable and  
30 less confounded by artifactual formation. Kim et al. (2009), which was published too late to be  
31 incorporated into the PBPK model, used an empirical pharmacokinetic model to analyze data on  
32 male B6C3F1 mice exposed to a single dose of 2,100 mg/kg TCE by gavage. Peak levels of

1 TCA and DCA were found to be 64 µg/ml and 18 ng/ml, respectively, a difference of more than  
2 3,000-fold. The kinetic rate constant they estimated for TCE → DCA were more than  
3 five orders of magnitude smaller than the kinetic rate constant estimated for TCE → TCA.  
4 These data all suggest that DCA is a minor metabolite of TCE as compared to TCA at high doses  
5 of around 2,000 mg/kg. Delinsky et al. (2005) reported that in male Sprague-Dawley rats, after a  
6 single 2,000 mg/kg dose by oral gavage, peak levels of DCA were 39.5 ng/ml. Delinsky et al.  
7 (2005) did not report TCA levels for comparison. The only data available in rats in this range of  
8 oral gavage doses (coincidentally also in male Sprague-Dawley rats) reported peak levels of  
9 TCA of 24 and 60 mg/ml at oral gavage doses of 600 and 3,000 mg/kg, respectively (Larson and  
10 Bull, 1992b). This suggests a difference between DCA and TCA levels in rats exposed to TCE  
11 of about 1,000-fold, albeit with more uncertainty as compared to Kim et al. (2009), in which both  
12 were measured simultaneously in the same animals. However, liver toxicity in both rats and  
13 mice is evident at much lower doses, so additional data are needed to inform whether the relative  
14 amount of TCA and DCA changes at lower exposures.  
15

#### **4.5.6.1.2. Comparisons Between Trichloroethylene (TCE) and Trichloroacetic Acid (TCA), Dichloroacetic Acid (DCA), and Chloral Hydrate (CH) Noncancer Effects**

##### **4.5.6.1.3. Hepatomegaly—qualitative and quantitative comparisons**

16 As discussed above, TCE causes hepatomegaly in rats, mice, and gerbils under both acute  
17 and chronic dosing. Data from a few available studies suggest that oxidative metabolism is  
18 important for mediating these effects. Buben and O’Flaherty (1985) collected limited  
19 pharmacokinetic data in a sample of the same animals for which liver weight changes were being  
20 assessed. While liver weight increases had similarly strong correlations with applied dose and  
21 urinary metabolites for doses up to 1,600 mg/kg-day ( $R^2$  of 0.97 for both), above that dose, the  
22 linear relationship was maintained with urinary metabolites but not with applied dose. Ramdhan  
23 et al. (2008) conducted parallel experiments at TCE 1,000 and 2,000 ppm (8 hours/day, 7 days)  
24 in wild-type and CYP2E1-null mice, which did not exhibit increased liver/body weight ratios  
25 with TCE treatment and excreted twofold lower amounts of oxidative metabolites TCA and  
26 TCOH in urine as compared to wild-type mice. However, among control mice, those with the  
27 null genotype had 1.32-fold higher absolute liver weights and 1.18-fold higher liver/body weight  
28 ratios than wild-type mice, reducing the sensitivity of the experiment, particularly with only  
29 six mice per dose group.

1 Ramdhan et al. (2008) reported that stated that urinary TCA levels in wild type mice were  
2 incorrectly reported by Ramdhan et al. (2008) but were corrected in this study. The authors  
3 reported no differences in urinary volume by genotype or exposure but did not show the data.  
4 TCA and trichloroethanol were detected in all exposed mice with no significant differences  
5 between the 1,000 and 2,000 ppm TCE levels. TCA concentrations were reported to be  
6 significantly lower and trichloroethanol levels significantly higher in PPAR $\alpha$ -null mice relative  
7 to wild type mice with no differences in genotype between the sum of total TCA and  
8 trichloroethanol concentrations between genotypes. The authors reported that they measured  
9 hepatic protein expression of CYP2E1 and ALDH2 enzymes and did not observe a significant  
10 difference among controls (data not shown) and that TCE exposure did not alter hepatic CYP2E1  
11 expression but did decrease ALDH2 expression to a comparable extent in all mouse lines (data  
12 not shown). Thus, changes in urinary TCA levels in the differing strains were not related to  
13 changes in expression of these metabolic enzymes.

14 As stated above, hepatomegally was increased by TCE exposure in all three strains. TCE  
15 at both 1,000 and 2,000 ppm significantly increased liver weight in the three mouse lines to a  
16 similar extent (i.e., 38 and 49% in wild type mice, 20 and 37% in PPAR-null mice, and 28 and  
17 32% in hPPAR $\alpha$  mice). The increases were not statistically significant between doses within  
18 each strain. Liver/body weight ratios were also significantly increased with TCE exposure at  
19 1,000 and 2,000 ppm relative to controls (i.e., 38 and 43% in wild type mice, 24 and 36% in  
20 PPAR $\alpha$ -null mice, and 27 and 39% in hPPAR $\alpha$  mice, respectively). The difference between  
21 2,000 and 1,000 ppm TCE exposure was statistically significant in PPAR $\alpha$ -null mice. As to the  
22 nature of the hepatomegally induced under these conditions, hepatic triglyceride levels were  
23 reported to be significantly correlated with liver/body weight ratios of all mice used in the study  
24 ( $r = 0.54$ ).

25 With respect to oxidative metabolites themselves, data from CH studies are not  
26 informative—either because data were not shown (Sanders et al., 1982a) or, because at the time  
27 points measured, liver weight increases are substantially confounded by foci and carcinogenic  
28 lesions (Leahey et al., 2003a). TCA and DCA have both been found to cause hepatomegaly in  
29 mice and rats, with mice being more sensitive to this effect. DCA also increases liver/body  
30 weight ratios in dogs, but TCE and TCA have not been tested in this species (Cicmanec et al.,  
31 1991).

32 As noted above, TCE-induced changes in liver weight appear to be proportional to the  
33 exposure concentration across route of administration, gender and rodent species. As an  
34 indication of the potential contribution of TCE metabolites to this effect, a quantitative


1 comparison of the shape of the dose-response curves for liver weight induction for TCE and its  
2 metabolites is informative. The analysis below was reported in Evans et al. (2009).

3 A number of short-term (<4 weeks) studies of TCA and DCA in drinking water have  
4 attempted to measure changes in liver weight induction, with the majority of these studies being  
5 performed in male B6C3F1 mice. Studies conducted from 14–30 days show a consistent  
6 increase in percentage liver/body weight induction by TCA or DCA. However, as stated in  
7 many of the discussions of individual studies (see Appendix E), there is a limited ability to detect  
8 a statistically significant change in liver weight change in experiments that use a relatively small  
9 number of animals or do not match control and treatment groups for age and weight. The  
10 experiments of Buben and O’Flaherty used 12–14 mice per group giving it a greater ability to  
11 detect a TCE-induced dose-response. However, many experiments have been conducted with  
12 4–6 mice per dose group. For example, the data from DeAngelo et al. (2008) for TCA-induced  
13 percentage liver/body weight ratio increases in male B6C3F1 mice were only derived from  
14 five animals per treatment group after 4 weeks of exposure. The 0.05 and 0.5 g/L exposure  
15 concentrations were reported to give a 1.09- and 1.16-fold of control percentage liver/body  
16 weight ratios which were consistent with the increases noted in the cross-study database above.  
17 However, a power calculation shows that the Type II error (which should be >50% and thus,  
18 greater than the chances of “flipping a coin”) was only a 6 and 7% and therefore, the designed  
19 experiment could accept a false null hypothesis. In addition, some experiments took greater care  
20 to age and weight match the control and treatment groups before the start of treatment.

21 Therefore, given these limitations and the fact that many studies used a limited range of  
22 doses, an examination of the combined data from multiple studies (Carter et al., 1995; DeAngelo  
23 et al., 1989; 2008; Kato-Weinstein et al., 2001; Parrish et al., 1996; Sanchez and Bull, 1990) can  
24 best inform/discern differences in DCA and TCA dose-response relationships for liver weight  
25 induction (described in more detail in Section E.2.4.2). The dose-response curves for similar  
26 concentrations of DCA and TCA are presented in Figure 4-5 for durations of exposure from  
27 14–28 days in the male B6C3F1 mouse, which was the most common sex and strain used. As  
28 noted in Appendix E, there appears to be a linear correlation between dose in drinking water and  
29 liver weight induction up to 2 g/L of DCA. However, the shape of the dose-response curve for  
30 TCA appears to be quite different. Lower concentrations of TCA induce larger increase that  
31 does DCA, but the TCE response reaches an apparent plateau while that of DCA continues to  
32 increase the response. TCA studies did not show significant duration-dependent difference in  
33 liver weight induction in this duration range. Short duration studies (10–42 days) were selected  
34 because (1) in chronic studies, liver weight increases are confounded by tumor burden,

1 (2) multiple studies are available, and (3) TCA studies do not show significant  
2 duration-dependent differences in this duration range.

3 Of interest is the issue of how the dose-response curves for TCA and DCA compare to  
4 that of TCE in a similar model and dose range. Since TCA and DCA have strikingly different  
5 dose-response curves, which one if either best fits that of TCE and thus, can give insight as to  
6 which is causative agent for TCE's effects in the liver? The carcinogenicity of chronic TCE  
7 exposure has been predominantly studied in two mouse strains, Swiss and B6C3F1, both of  
8 which reportedly developed liver tumors. Rather than administered in drinking water, oral TCE  
9 studies have been conducted via oral gavage and generally in corn oil for 5 days of exposure per




10  
11 **Figure 4-5. Comparison of average fold-changes in relative liver weight to**  
12 **control and exposure concentrations of 2 g/L or less in drinking water for**  
13 **TCA and DCA in male B6C3F1 mice for 14–30 days (Carter et al., 1995;**  
14 **DeAngelo et al., 1989; 2008; Kato-Weinstein et al., 2001; Parrish et al., 1996;**  
15 **Sanchez and Bull, 1990).**


16  
17  
18 week. Factors adding to the increased difficulty in establishing the dose-response relationship  
19 for TCE across studies and for comparisons to the DCA and TCA database include vehicle  
20 effects, the difference between daily and weekly exposures, the dependence of TCE effects in the  
21 liver on its metabolism to a variety of agents capable inducing effects in the liver, differences in  
22 response between strains, and the inherent increased variability in use of the male mouse model.  
23 In particular, these factors would add variability to any effort at a combined analysis, and make a

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 consistent dose-response pattern more difficult to discern. Nonetheless, despite such differences  
2 in exposure route, vehicle, etc., a consistent pattern of dose-response emerges from combining  
3 the available TCE data. The effects of oral exposure to TCE from 10–42 days on liver weight  
4 induction is shown below in Figure 4-6 using the data of Elcombe et al. (1985), Dees and Travis  
5 (1993), Goel et al. (1992), Merrick et al. (1989), Goldsworthy and Popp (1987), and Buben and  
6 O’Flaherty (1985). Oral TCE administration in male B6C3F1 and Swiss mice appeared to  
7 induce a dose-related increase in percentage liver/body weight that was generally proportional to  
8 the increase in magnitude of dose, though as expected, with more variability than observed for a  
9 similar exercise for DCA or TCA in drinking water. Some of the variability is due to the



10



11


*This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2 **Figure 4-6. Comparisons of fold-changes in average relative liver weight and**  
3 **gavage dose of (top panel) male B6C3F1 mice for 10–28 days of exposure**  
4 **(Dees and Travis, 1993; Elcombe et al., 1985; Goldsworthy and Popp, 1987;**  
5 **Merrick et al., 1989) and (bottom panel) in male B6C3F1 and Swiss mice.**  
6  
7

8 inclusion of the 10 day studies, since as discussed in Section E.2.4.2, there was a greater increase  
9 in TCE-induced liver weight at 28–42 days of exposure Swiss mice than the 10-day data in  
10 B6C3F1 mice, and Kjellstrand et al. (1981b) noted that TCE-induced liver weight increases are  
11 still increasing at 10 days inhalation exposure. A strain difference is not evident between the  
12 Swiss and B6C3F1 males, as both the combined TCE data and that for only B6C3F1 mice show  
13 similar correlation with the magnitude of dose and magnitude of percentage liver/body weight  
14 increase. The correlation coefficients for the linear regressions presented for the B6C3F1 data  
15 are  $R^2 = 0.861$  and for the combined data sets is  $R^2 = 0.712$ . Comparisons of the slopes of the  
16 dose-response curves suggest a greater consistency between TCE and DCA than between TCE  
17 and TCA. There did not appear to be evidence of a plateau with higher TCE doses, and the  
18 degree of fold-increase rises to higher levels with TCE than with TCA in the same strain of  
19 mouse.

20 A more direct comparison would be on the basis of dose rather than drinking water  
21 concentration. The estimations of internal dose of DCA or TCA from drinking water studies,  
22 while varying considerably (DeAngelo et al., 1989; 2008), nonetheless suggest that the doses of  
23 TCE used in the gavage experiments were much higher than those of DCA or TCA. However,  
24 only a fraction of ingested TCE is metabolized to DCA or TCA, as, in addition to oxidative  
25 metabolism, TCE is also cleared by GSH conjugation and by exhalation. While DCA dosimetry  
26 is highly uncertain (see Sections 3.3 and 3.5), the mouse PBPK model, described in Section 3.5  
27 was calibrated using extensive in vivo data on TCA blood, plasma, liver, and urinary excretion  
28 data from inhalation and gavage TCE exposures, and makes robust predictions of the rate of  
29 TCA production. If TCA were predominantly responsible for TCE-induced liver weight  
30 increases, then replacing administered TCE dose (e.g., mg TCE/kg/day) by the rate of TCA  
31 produced from TCE (mg TCA/kg/day) should lead to dose-response curves for increased liver  
32 weight consistent with those from directly administered TCA. Figure 4-7 shows this comparison  
33 using the PBPK model-based estimates of TCA production for four TCE studies from  
34 28–42 days in the male NMRI, Swiss, and B6C3F1 mice (Kjellstrand et al., 1983a (Buben and  
35 O'Flaherty, 1985; Goel et al., 1992; Merrick et al., 1989) and four oral TCA studies in B6C3F1  
36 male mice at 2 g/L or lower drinking water exposure (DeAngelo et al., 1989; 2008; Kato-  
37 Weinstein et al., 2001; Parrish et al., 1996) from 14–28 days of exposure. The selection of the

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 28–42 day data for TCE was intended to address the decreased opportunity for full expression of  
 2 response at 10 days. PBPK modeling predictions of daily internal doses of TCA in terms of  
 3 mg/kg-day via produced via TCE metabolism would be are indeed lower than the TCE  
 4 concentrations in terms of mg/kg-day given orally by gavage. The predicted internal dose of  
 5 TCA from TCE exposure studies are of a comparable range to those predicted from TCA  
 6 drinking water studies at exposure concentrations in which palpability has not been an issue for  
 7 estimation of internal dose. Thus, although the TCE data are for higher exposure concentrations,  
 8 they are predicted to produce comparable levels of TCA internal dose estimated from direct TCA  
 9 administration in drinking water.

10



1 **Figure 4-7. Comparison of fold-changes in relative liver weight for data sets**  
2 **in male B6C3F1, Swiss, and NRM1 mice between TCE studies Kjellstrand**  
3 **et al., 1983a (Buben and O'Flaherty, 1985; Goel et al., 1992; Merrick et al.,**  
4 **1989) [duration 28–42 days] and studies of direct oral TCA administration**  
5 **to B6C3F1 mice (DeAngelo et al., 1989; DeAngelo et al., 2008; Kato-**  
6 **Weinstein et al., 2001; Parrish et al., 1996)[duration 14–28 days]).** Abscissa  
7 for TCE studies consists of the median estimates of the internal dose of TCA  
8 predicted from metabolism of TCE using the PBPK model described in  
9 Section 3.5 of the TCE risk assessment. Lines show linear regression with  
10 intercept fixed at unity. All data were reported fold-change in mean liver  
11 weight/body weight ratios, except for Kjellstrand et al. (1983b), with were the  
12 fold-change in the ratio of mean liver weight to mean body weight. In addition, in  
13 Kjellstrand et al. (1983b), some systemic toxicity as evidence by decreased total  
14 body weight was reported in the highest-dose group.  
15  
16

17 Figure 4-7 clearly shows that for a given amount of TCA produced from TCE, but going  
18 through intermediate metabolic pathways, the liver weight increases are substantially greater  
19 than, and highly inconsistent with, that expected based on direct TCA administration. In  
20 particular, the response from direct TCA administration appears to "saturate" with increasing  
21 TCA dose at a level of about 1.4-fold, while the response from TCE administration continues to  
22 increase with dose to 1.75-fold at the highest dose administered orally in Buben and O'Flaherty  
23 (1985) and over twofold in the inhalation study of Kjellstrand et al. (1983b). Because TCA liver  
24 concentrations are proportional to the dose TCA, and do not depend on whether it is  
25 administered in drinking water or internally produced in the liver, the results of the comparison  
26 using the TCA liver dose-metric are identical.

27 Furthermore, while as noted previously, oral studies appear to report a linear relationship  
28 between TCE exposure concentration and liver weight induction, the inclusion of inhalation  
29 studies on the basis of internal dose led to a highly consistent dose-response curve for among  
30 TCE study. Therefore, it is unlikely that differing routes of exposure can explain the  
31 inconsistencies in dose-response.


32 The bioavailability of TCA, which in the above analysis is assumed to be 100%, is  
33 another factor that may impact the dose-response. Sweeney et al. (2009), in an analysis of the  
34 potential role of TCA in the liver carcinogenesis of tetrachloroethylene, identified a number of  
35 previously unpublished TCA kinetic data in mice exposed to TCA via drinking water for  
36 3–14 days. They concluded that fractional absorption of TCA via drinking water exposures is  
37 much less than 100%—about 29% at low exposures and decreasing with increasing dose.  
38 However, the conclusions of the Sweeney et al. (2009) were based on the Hack et al. (2006) TCE  
39 PBPK model, which had a number of deficiencies, as noted in Section 3.5 and Appendix A.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Therefore, as discussed in Appendix A, Chiu (Chiu, In Press)(in press) reanalyzed those data  
2 using the updated TCE PBPK model of Evans et al. (2009) and Chiu et al. (2009) and concluded  
3 that while there was evidence of reduced absorption (80–90% at low exposures, and decreasing  
4 with increasing dose), it was not as low as that estimated by Sweeney et al. (2009). As discussed  
5 in Appendix A, it may be more accurate to characterize the fractional absorption as an empirical  
6 parameter reflecting unaccounted-for biological processes as well as experimental variation.

7 Chiu (Chiu, In Press) also reanalyzed the data on TCE- and TCA-induced hepatomegaly  
8 using the central estimates of the fractional absorption of TCA inferred from the analysis  
9 described above. Figure 4-8 shows the results, comparing a fixed fractional absorption of 95%  
10 with the fitted fractional absorption from Chiu (Chiu, In Press), here plotted using area-under-  
11 the-curve (AUC) of TCA in the liver as the dose-metric. For reference, the dose-response for  
12 administered TCA with an assumption of fixed, nearly complete absorption (analogous to the  
13 results from Evans et al., 2009, Figure 4-7) is also included. While the reduced fractional  
14 absorption inferred from drinking water data reported by Sweeney et al. (2009) accounts for part  
15 of the difference in dose-responses between TCE- and TCA-induced hepatomegaly reported by  
16 Evans et al. (2009), it does not appear to be able to account for the entire difference. In  
17 particular, the fraction of hepatomegaly contributed by TCA is about 0.20 assuming nearly  
18 complete absorption, as compared to about 0.33 assuming the best-fitting fractional absorption  
19 inferred from the PBPK

20



1 **Figure 4-8. Comparison of hepatomegaly as a function of AUC of TCA in**  
 2 **liver, using values for the TCA drinking water fractional absorption (Fabs).**  
 3 Fold-changes in relative liver weight for data sets in male B6C3F1, Swiss, and  
 4 NRMI mice between TCE studies Kjellstrand et al., 1983a (Buben and  
 5 O'Flaherty, 1985; Goel et al., 1992; Merrick et al., 1989) [duration 28–42 days]  
 6 and studies of direct oral TCA administration to B6C3F1 mice (DeAngelo et al.,  
 7 1989; DeAngelo et al., 2008; Kato-Weinstein et al., 2001; Parrish et al., 1996)  
 8 Green, 2003b [duration 14–28 days]. Linear regressions were compared using  
 9 ANOVA to assess whether the TCE studies were consistent with the TCA studies,  
 10 using TCA as the dose-metric. For each analysis of drinking water fraction  
 11 absorption, ANOVA  $p$ -values were  $<10^{-4}$  when comparing the assumption that all  
 12 the data had a common slope with the assumption that TCE and TCA data had  
 13 different slopes.

14  
 15  
 16 model-based analysis. The inability of TCA to account for TCE-induced hepatomegaly is  
 17 confirmed statistically by ANOVA, with  $p$ -values of  $<10^{-4}$ . Therefore, assuming a reduced TCA  
 18 bioavailability does not change the conclusion that the available data are inconsistent with the  
 19 toxicological hypothesis that TCA can fully account for TCE-induced hepatomegaly.

1 Additional analyses do, however, support a role for oxidative metabolism in  
2 TCE-induced liver weight increases, and that the parent compound TCE is not the likely active  
3 moiety (suggested previously by Buben and O'Flaherty (1985)). In particular, the same studies  
4 are shown in Figure 4-9 using PBPK-model based predictions of the AUC of TCE in blood and  
5 total oxidative metabolism, which produces chloral, TCOH, DCA, and other metabolites in  
6 addition to TCA. The dose-response relationship between TCE blood levels and liver weight  
7 increase, while still having a significant trend, shows substantial scatter and a low  $R^2$  of 0.43.  
8 On the other hand, using total oxidative metabolism as the dose-metric leads to substantially  
9 more consistency dose-response across studies, and a much tighter linear trend with an  $R^2$  of  
10 0.90 (see Figure 4-9). A similar consistency is observed using liver-only oxidative metabolism  
11 as the dose-metric, with  $R^2$  of 0.86 (not shown). Thus, while the slope is similar between liver  
12 weight increase and TCE concentration in the blood and liver weight increase and rate of total  
13 oxidative metabolism, the data are a much better fit for total oxidative metabolism.  
14

15  
16 **Figure 4-9. Fold-changes in relative liver weight for data sets in male**  
17 **B6C3F1, Swiss, and NRMI mice reported by TCE studies of duration**  
18 **28–42 days (Kjellstrand et al., 1983a(Buben and O'Flaherty, 1985; Goel et**  
19 **al., 1992; Merrick et al., 1989) using internal dose-metrics predicted by the**  
20 **PBPK model described in Section 3.5: (A) dose-metric is the median estimate**  
21 **of the daily AUC of TCE in blood, (B) dose-metric is the median estimate of**  
22 **the total daily rate of TCE oxidation.** Lines show linear regression. Use of  
23 liver oxidative metabolism as a dose-metric gives results qualitatively similar to  
24 (B), with  $R^2 = 0.86$ .  
25

1           Although the qualitative similarity to the linear dose-response relationship between DCA  
2 and liver weight increases is suggestive of DCA being the predominant metabolite responsible  
3 for TCE liver weight increases, due to the highly uncertain dosimetry of DCA derived from  
4 TCE, this hypothesis cannot be tested on the basis of internal dose. Similarly, another TCE  
5 metabolite, chloral hydrate, has also been reported to induce liver tumors in mice, however, there  
6 are no adequate comparative data to assess the nature of liver weight increases induced by this  
7 TCE metabolite (see Section E.2.5 and Section 4.5.1.2.4 below). Whether its formation in the  
8 liver after TCE exposure correlates with TCE-induced liver weight changes cannot be  
9 determined.

#### 4.5.6.1.4. Cytotoxicity

11           As discussed above, TCE has sometimes been reported to cause minimal/mild focal  
12 hepatocellular necrosis or other signs of hepatic injury, albeit of low frequency and mostly at  
13 doses  $\geq 1,000$  mg/kg-day (Dees and Travis, 1993; Elcombe et al., 1985) or at exposures  
14  $\geq 1,000$  ppm in air (Ramdhan et al., 2010; Ramdhan et al., 2008) from 7–10 days of exposure.  
15 Data from available studies are supportive of a role for oxidative metabolism in TCE-induced  
16 cytotoxicity in the liver, though they are not informative as to the actual active moiety(ies).  
17 Buben and O’Flaherty (1985) noted a strong correlation (R-squared of between  
18 glucose-6-phosphatase inhibition and total urinary oxidative metabolites). Ramdhan et al. (2008)  
19 conducted parallel experiments at TCE 1,000 and 2,000 ppm (8 hours/day, 7 days) in wild-type  
20 and CYP2E1-null mice, the latter of which did not exhibit hepatotoxicity (assessed by serum  
21 ALT, AST, and histopathology) and excreted twofold lower amounts of oxidative metabolites  
22 TCA and TCOH in urine as compared to wild-type mice. In addition, urinary TCA and TCOH  
23 excretion was correlated with serum ALT and AST measures, though the R-squared values  
24 (square of the reported correlation coefficients) were relatively low (0.54 and 0.67 for TCOH and  
25 TCA, respectively). Ramdhan et al. (2010) reported that TCA and TCOH were detected in the  
26 urine of wild type and PPAR $\alpha$ -null and humanized mice after TCE exposure with no significant  
27 differences between the 1,000 and 2,000 ppm TCE treatments. TCA concentrations were  
28 significantly lower and TCOH concentrations higher in exposed PPAR $\alpha$ -null mice relative to  
29 wild type mice. They stated that urinary TCA levels in wild type mice were incorrectly reported  
30 by Ramdhan et al. (2008) but have been corrected in this study. AST and ALT levels were  
31 significantly increased in all exposed mice relative to control with mean levels between 1,000  
32 and 2,000 ppm TCE exposures higher but not significantly different (41–74% and 36–79%  
33 higher, respectively). Although increased, such increases were small. Necrosis scores were

1 reported to be significantly higher in TCE exposed mice relative to controls in all three genotype  
2 mice and to be significantly higher with 2,000 versus 1,000 ppm TCE exposure in wild type mice  
3 and hPPAR $\alpha$  mice. Inflammation scores were reported to be significantly higher with exposed  
4 group than control with 2,000 ppm TCE exposure than controls for each genotype group with a  
5 difference between the 2,000 ppm and 1,000 ppm exposure groups in wild type mice. However,  
6 necrosis and inflammation score means at the highest TCE exposure levels in any mouse strain  
7 were minimal (only occasional necrotic cells in any lobule) for necrosis and mild for  
8 inflammation (<2 foci/field).

9 With respect to CH (166 mg/kg-day) and DCA (~90 mg/kg-day), Daniel et al. (1992)  
10 reported that after drinking water treatment, hepatocellular necrosis and chronic active  
11 inflammation were reported to be mildly increased in both prevalence and severity in all treated  
12 groups after 104 weeks of exposure. The histological findings, from interim sacrifices ( $n = 5$ ),  
13 were considered by the authors to be unremarkable and were not reported. TCA has not been  
14 reported to induce necrosis in the liver under the conditions tested. Relatively high doses of  
15 DCA ( $\geq 1$  g/L in drinking water) appear to result in mild focal necrosis with attendant reparative  
16 proliferation at lesion sites, but no such effects were reported at lower doses ( $\leq 0.5$  g/L in  
17 drinking water) more relevant for comparison with TCE (DeAngelo et al., 1999; Sanchez and  
18 Bull, 1990; Stauber et al., 1998). Enlarged nuclei and changes consistent with increased ploidy,  
19 are further discussed below in the context of DNA synthesis.

20

#### 4.5.6.1.5. DNA synthesis and polyploidization

21 The effects on DNA synthesis and polyploidization observed with TCE treatment have  
22 similarly been observed with TCA and DCA. With respect to CH, George et al. (2000) reported  
23 that CH exposure did not alter DNA synthesis in rats and mice at any of the time periods  
24 monitored (all well past 2 weeks), with the exception of 0.58 g/L chloral hydrate at 26 weeks  
25 slightly increasing hepatocyte labeling (~two–threefold of controls) in rats and mice but the  
26 percentage labeling still representing 3% or less of hepatocytes.

27 In terms of whole liver or hepatocyte label incorporation, the most comparable exposure  
28 duration between TCE, TCA, and DCA studies is the 10- and 14-day period. Several studies  
29 have reported that in this time period, peak label incorporation into individual hepatocytes and  
30 whole liver for TCA and DCA have already passed (Carter et al., 1995; Pereira, 1996; Sanchez  
31 and Bull, 1990; Styles et al., 1991). A direct time-course comparison is difficult, since data at  
32 earlier times for TCE are more limited.

1           There are conflicting reports of DNA synthesis induction in individual hepatocytes for up  
2 to 14 days of DCA or TCA exposure. In particular, Sanchez and Bull (1990) reported tritiated  
3 thymidine incorporation in individual hepatocytes up to 2 g/L exposure to DCA or TCA induced  
4 little increase in DNA synthesis except in instances and in close proximity to areas of  
5 proliferation/necrosis for DCA treatment after 14 days of exposure in male mice. The largest  
6 percentage of hepatocytes undergoing DNA synthesis for any treatment group was less than  
7 1% of hepatocytes. However, they reported treatment- and exposure duration-changes in hepatic  
8 DNA incorporation of tritiated thymidine for DCA and TCA. For TCA treatment, the largest  
9 increases over control levels for hepatic DNA incorporation (at the highest dose) was a threefold  
10 increase after 5 days of treatment and a twofold increase over controls after 14 days of treatment.  
11 For DCA whole-liver tritiated thymidine incorporation was only slightly elevated at necrogenic  
12 concentrations and decreased at the 0.3 g/L non-necrogenic level after 14 days of treatment. In  
13 contrast to Sanchez and Bull (1990), Stauber and Bull (1997) reported increased tritiated  
14 thymidine incorporation for individual hepatocytes after 14 days of treatment with 2 g/L DCA or  
15 TCA in male mice. They used a more extended period of tritiated thymidine exposure of  
16 3–5 days and so these results represent aggregate DNA synthesis occurring over a more extended  
17 period of time. A “1-day labeling index” was reported as less than 1% for the highest level of  
18 increased incorporation. However, after 14 days, the labeling index was reported to be increased  
19 by ~3.5-fold for TCA and ~5.5-fold for DCA over control values. After 28 days, the labeling  
20 index was reported to be decreased ~2.3-fold by DCA and increased ~2.5-fold after treatment  
21 with TCA. Pereira (1996) reported that for female B6C3F1 mice, 5-day incorporation of BrDU,  
22 as a measure of DNA synthesis, was increased at 0.86 g/L and 2.58 g/L DCA treatment for  
23 5 days (~twofold at the highest dose) but that by Day 12 and 33 levels had fallen to those of  
24 controls. For TCA exposures, 0.33 g/L, 1.10 g/L and 3.27 g/L TCA all gave a similar ~threefold  
25 increase in BrdU incorporation by 5 days, but that by 12 and 33 days were not changed from  
26 controls. Nonetheless, what is consistent is that these data report that, similar to TCE-exposed  
27 mice at 10 days of exposure, cells undergoing DNA synthesis in DCA- or TCA-exposed mice for  
28 up to 14 days of exposure to be confined to a very small population of cells in the liver. Thus,  
29 these data are consistent with hypertrophy being primarily responsible for liver weight gains as  
30 opposed to increases in cell number in mice.

31           Interestingly, a lack of correlation between whole liver label incorporation and that in  
32 individual hepatocytes has been reported by several studies of DCA (Carter et al., 1995; Sanchez  
33 and Bull, 1990). For example, Carter et al. (1995) reported no increase in labeling of  
34 hepatocytes in comparison to controls for any DCA treatment group from 5–30 days of DCA  
35 exposure. Rather than increase hepatocyte labeling, DCA induced no change from Days 5

1 though 15 but significantly decreased levels between days 20 and 30 for 0.5 g/L that were similar  
2 to those observed for the 5 g/L exposures. However, for whole liver DNA tritiated thymidine  
3 incorporation, Carter et al. (1995) reported 0.5g/L DCA treatments to show trends of initial  
4 inhibition of DNA tritiated thymidine incorporation followed by enhancement of labeling that  
5 was not statistically significant from 5–30 days of exposure. Examination of individual  
6 hepatocytes does not include the contribution of nonparenchymal cell DNA synthesis that would  
7 be detected in whole liver DNA. As noted above, proliferation of the nonparenchymal cell  
8 compartment of the liver has been noted in several studies of TCE in rodents, and thus, this is  
9 one possible reason for the reported discrepancy.

10 Another possible reason for this inconsistency with DCA treatment is polyploidization, as  
11 was suggested above for TCE. Although this was not examined for DCA or TCA exposure by  
12 Sanchez and Bull (1990), Carter et al. (1995) reported that hepatocytes from both 0.5 and 5 g/L  
13 DCA treatment groups had enlarged, presumably polyploidy nuclei, with some hepatocyte nuclei  
14 labeled in the mid-zonal area. There were statistically significant changes in cellularity, nuclear  
15 size, and multinucleated cells during 30 days exposure to DCA. The percentage of  
16 mononucleated cells hepatocytes was reported to be similar between control and DCA treatment  
17 groups at 5- and 10-day exposure. However, at 15 days and beyond DCA treatments were  
18 reported to induce increases in mononucleated hepatocytes with later time periods to also  
19 showing DCA-induced increases nuclear area, consistent with increased polyploidization without  
20 mitosis. The consistent reporting of an increasing number of mononucleated cells between 15  
21 and 30 days could be associated with clearance of mature hepatocytes as suggested by the report  
22 of DCA-induced loss of cell nuclei. The reported decrease in the numbers of binucleate cells in  
23 favor of mononucleate cells is not typical of any stage of normal liver growth (Brodsky and  
24 Uryvaeva, 1977). The pattern of consistent increase in percentage liver/body weight induced by  
25 0.5 g/L DCA treatment from days 5 though 30 was not consistent with the increased numbers of  
26 mononucleate cells and increase nuclear area reported from Day 20 onward. Specifically, the  
27 large differences in liver weight induction between the 0.5 g/L treatment group and the 5 g/L  
28 treatment groups at all times studied also did not correlate with changes in nuclear size and  
29 percentage of mononucleate cells. Thus, increased liver weight was not a function of cellular  
30 proliferation, but probably included both aspects of hypertrophy associated with polyploidization  
31 and increased glycogen deposition (see below) induced by DCA. Carter et al. (1995) suggested  
32 that although there is evidence of DCA-induced cytotoxicity (e.g., loss of cell membranes and  
33 apparent apoptosis), the 0.5 g/L exposure concentration has been shown to increase  
34 hepatocellular lesions after 100 weeks of treatment without concurrent peroxisome proliferation  
35 or cytotoxicity (DeAngelo et al., 1999).



1 In sum, the observation of TCE-treatment related changes in DNA content, label  
2 incorporation, and mitotic figures are generally consistent with patterns observed for both TCA  
3 and DCA. In all cases, hepatocellular proliferation is confined to a very small fraction of  
4 hepatocytes, and hepatomegaly observed with all three treatments probably largely reflects  
5 cytomegaly rather than cell proliferation. Moreover, label incorporation likely largely reflects  
6 polyploidization rather than hepatocellular proliferation, with a possible contribution from  
7 nonparenchymal cell proliferation. As with TCE, histological changes in nuclear sizes and  
8 number also suggest a significant degree of treatment-related polyploidization, particularly for  
9 DCA.

#### 10 **4.5.6.1.6. Apoptosis**

11 As for apoptosis, Both Elcombe et al. (1985) and Dees and Travis (1993) reported no  
12 changes in apoptosis other than increased apoptosis only at a treatment level of 1,000-mg/kg  
13 TCE. Dees and Travis (1993) reported that increased apoptoses from TCE exposure “did not  
14 appear to be in proportion to the applied TCE dose given to male or female mice.” Channel et al.  
15 (1998) reported that there was no significant difference in apoptosis between TCE treatment and  
16 control groups with data not shown. However, the extent of apoptosis in any of the treatment  
17 groups, or which groups and timepoints were studied for this effect cannot be determined. While  
18 these data are quite limited, it is notable that peroxisome proliferators have been suggested  
19 inhibit, rather than increase, apoptosis as part of their carcinogenic MOA (Klaunig et al., 2003).

20 However, for TCE metabolites, DCA has been most studied, though it is clear that age  
21 and species affect background rates of apoptosis. Snyder et al. (1995), in their study of DCA,  
22 report that control mice were reported to exhibit apoptotic frequencies ranging from  
23 ~0.04–0.085%, that over the 30-day period of their study the frequency rate of apoptosis  
24 declined, and suggest that this pattern is consistent with reports of the livers of young animals  
25 undergoing rapid changes in cell death and proliferation. They reported rat liver to have a  
26 greater the estimated frequency of spontaneous apoptosis (~0.1%) and therefore, greater than that  
27 of the mouse. Carter et al. (1995) reported that after 25 days of 0.5 g/L DCA treatment apoptotic  
28 bodies were reported as well as fewer nuclei in the pericentral zone and larger nuclei in central  
29 and midzonal areas. This would indicate an increase in the apoptosis associated with potential  
30 increases in polyploidization and cell maturation. However, Snyder et al. (1995) report that mice  
31 treated with 0.5 g/L DCA over a 30-day period had a similar trend as control mice of decreasing  
32 apoptosis with age. The percentage of apoptotic hepatocytes decreased in DCA-treated mice at  
33 the earliest time point studied and remained statistically significantly decreased from controls

1 from 5–30 days of exposure. Although the rate of apoptosis was very low in controls, treatment  
2 with 0.5g/L DCA reduced it further (~30–40% reduction) during the 30-day study period. The  
3 results of this study not only provide a baseline of apoptosis in the mouse liver, which is very  
4 low, but also to show the importance of taking into account the effects of age on such  
5 determinations. The significance of the DCA-induced reduction in apoptosis reported in this  
6 study, from a level that is already inherently low in the mouse, for the MOA for induction of  
7 DCA-induce liver cancer is difficult to discern.

8

#### 4.5.6.1.7. Glycogen accumulation

9 As discussed in Sections E.3.2 and E.3.4.2.1, glycogen accumulation has been described  
10 to be present in foci in both humans and animals as a result from exposure to a wide variety of  
11 carcinogenic agents and predisposing conditions in animals and humans. The data from  
12 Elcombe et al. (1985) included reports of TCE-induced pericentral hypertrophy and eosinophilia  
13 for both rats and mice but with “fewer animals affected at lower doses.” In terms of glycogen  
14 deposition, Elcombe report “somewhat” less glycogen pericentrally in the livers of rats treated  
15 with TCE at 1,500 mg/kg than controls with less marked changes at lower doses restricted to  
16 fewer animals. They do not comment on changes in glycogen in mice. Dees and Travis (1993)  
17 reported TCE-induced changes to “include an increase in eosinophilic cytoplasmic staining of  
18 hepatocytes located near central veins, accompanied by loss of cytoplasmic vacuolization.”  
19 Since glycogen is removed using conventional tissue processing and staining techniques, an  
20 increase in glycogen deposition would be expected to increase vacuolization and thus, the report  
21 from Dees and Travis is consistent with less not more glycogen deposition. Neither study  
22 produced a quantitative analysis of glycogen deposition changes from TCE exposure. Although  
23 not explicitly discussing liver glycogen content or examining it quantitatively in mice, these  
24 studies suggest that TCE-induced liver weight increases did not appear to be due to glycogen  
25 deposition after 10 days of exposure and any decreases in glycogen were not necessarily  
26 correlated with the magnitude of liver weight gain either.

27 For TCE and TCA 500 mg/kg treatments in mice for 10 days, changes in glycogen were  
28 not reported in the general descriptions of histopathological changes (Dees and Travis, 1993;  
29 Elcombe et al., 1985; Styles et al., 1991) or were specifically described by the authors as being  
30 similar to controls (Nelson et al., 1989). However, for DCA, glycogen deposition was  
31 specifically noted to be increased with treatment, although no quantitative analyses was  
32 presented that could give information as to the nature of the dose-response (Nelson et al., 1989).

1 In regard to cell size, although increased glycogen deposition with DCA exposure was  
2 noted by Sanchez and Bull (1990) to occur to a similar extent in B6C3F1 and Swiss Webster  
3 male mice despite differences in DCA-induced liver weight gain. Lack of quantitative analyses  
4 of that accumulation in this study precludes comparison with DCA-induced liver weight gain.  
5 Carter et al. (1995) reported that in control mice there was a large variation in apparent glycogen  
6 content and also did not perform a quantitative analysis of glycogen deposition. The variability  
7 of this parameter in untreated animals and the extraction of glycogen during normal tissue  
8 processing for light microscopy make quantitative analyses for dose-response difficult unless  
9 specific methodologies are employed to quantitatively assess liver glycogen levels as was done  
10 by Kato-Weinstein et al. (2001) and Pereira et al. (2004b).

11 Bull et al. (1990) reported that glycogen deposition was uniformly increased from 2 g/L  
12 DCA exposure with photographs of TCA exposure showing slightly less glycogen staining than  
13 controls. However, the abstract and statements in the paper suggest that there was increased  
14 PAS positive material from TCA treatment that has caused confusion in the literature in this  
15 regard. Kato-Weinstein et al. (2001) reported that in male B6C3F1 mice exposed to DCA and  
16 TCA, the DCA treatment increased glycogen and TCA decreased glycogen content of the liver  
17 by using both chemical measurement of glycogen in liver homogenates and by using  
18 ethanol-fixed sections stained with PAS, a procedure designed to minimize glycogen loss.

19 Kato-Weinstein et al. (2001) reported that glycogen rich and poor cells were scattered  
20 without zonal distribution in male B6C3F1 mice exposed to 2 g/L DCA for 8 weeks. For TCA  
21 treatments, they reported centrilobular decreases in glycogen and ~25% decreases in whole liver  
22 by 3 g/L TCA. Kato-Weinstein et al. (2001) reported whole liver glycogen to be increased  
23 ~1.50-fold of control (90 vs. 60 mg glycogen/g liver) by 2 g/L DCA after 8 weeks exposure male  
24 B6C3F1 mice with a maximal level of glycogen accumulation occurring after 4 weeks of DCA  
25 exposure. Pereira et al. (2004b) reported that after 8 weeks of exposure to 3.2 g/L DCA liver  
26 glycogen content was 2.20-fold of control levels (155.7 vs. 52.4 mg glycogen/g liver) in female  
27 B6C3F1 mice. Thus, the baseline level of glycogen content reported by (~60 mg/g) and the  
28 increase in glycogen after DCA exposure was consistent between Kato-Weinstein et al. (2001)  
29 and Pereira et al. (2004b). However, the increase in liver weight reported by Kato-Weinstein  
30 et al. (2001) of 1.60-fold of control percentage liver/body weight cannot be accounted for by the  
31 1.50-fold of control glycogen content. Glycogen content only accounts for 5% of liver mass so  
32 that 50% increase in glycogen cannot account for the 60% increase liver mass induced by 2 g/L  
33 DCA exposure for 8 weeks reported by Kato-Weinstein (2001). Thus, DCA-induced increases  
34 in liver weight are occurring from other processes as well. Carter et al. (2003) and DeAngelo  
35 et al. (1999) reported increased glycogen after DCA treatment at much lower doses after longer

1 periods of exposure (100 weeks). Carter reported increased glycogen at 0.5 g/L DCA and  
2 DeAngelo et al. (1999) at 0.03 g/L DCA in mice. However, there is no quantitation of that  
3 increase.  
4

#### 4.5.6.1.8. Peroxisome proliferation and related effects

5 TCA and DCA have both been reported to induce peroxisome proliferation or increase in  
6 related enzyme markers in rodent hepatocytes (DeAngelo et al., 1989; 1997; Mather et al., 1990;  
7 Parrish et al., 1996). Between TCA and DCA, both induce peroxisome proliferation in various  
8 strains of mice, but it clear that TCA and DCA are weak PPAR $\alpha$  agonists and that DCA is  
9 weaker than TCA in this regard (Nelson et al., 1989) using a similar paradigm.

10 George et al. (2000) reported that CH exposure did not hepatic PCO activity in rats and  
11 mice at any of the time periods monitored. It is notable that the only time at which DNA  
12 synthesis index was (slightly) increased, at 26 weeks, there remained a lack of induction of PCO.  
13 A number of measures that may be related to peroxisome proliferation were investigated in  
14 Leakey et al. (2003a). Of the enzymes associated with PPAR $\alpha$  agonism (total CYP, CYP2B  
15 isoform, CYP4A, or lauric acid  $\beta$ -hydroxylase activity), only CYP4A and lauric acid  
16  $\beta$ -hydroxylase activity were significantly increased at 15 months of exposure in the  
17 dietary-restricted group administered the highest dose (100 mg/kg CH) with no other groups  
18 reported showing a statistically significant increased response ( $n = 12$ /group). There is an issue  
19 of interpretation of peroxisomal enzyme activities and other enzymes associated with PPAR $\alpha$   
20 receptor activation to be a relevant event in liver cancer induction at a time period in which  
21 tumors or foci are already present. Although not statistically significant, the 100 mg/kg CH  
22 exposure group of *ad libitum*-fed mice also had an increase in CH-induced increases of CYP4A  
23 and lauric acid  $\beta$ -hydroxylase activity. Seng et al. (2003) described CH toxicokinetics and  
24 peroxisome proliferation-associated enzymes in mice at doses up to 1,000 mg/kg-day for  
25 2 weeks with dietary control or caloric restriction. Lauric acid  $\beta$ -hydroxylase and PCO activities  
26 were reported to be induced only at doses >100 mg/kg in all groups, with dietary-restricted mice  
27 showing the greatest induction. Differences in serum levels of TCA, the major metabolite  
28 remaining 24 hours after dosing, were reported not to correlate with hepatic lauric acid  
29  $\beta$ -hydroxylase activities across groups.

30 Direct quantitative inferences regarding the magnitude of response in these studies in  
31 comparison to TCE, however, are limited by possible variability and confounding. In particular,  
32 many studies used cyanide-insensitive PCO as a surrogate for peroxisome proliferation, but the  
33 utility of this marker may be limited for a number of reasons. First, several studies have shown

1 that this activity is not well correlated with the volume or number of peroxisomes that are  
2 increased as a result of exposure to TCE or its metabolites (Elcombe et al., 1985; Nakajima et al.,  
3 2000; Nelson et al., 1989). In addition, this activity appears to be highly variable both as a  
4 baseline measure and in response to chemical exposures. Laughter et al. (2004) presented data  
5 showing WY-14,643 induced increases in PCO activity that varied up to sixfold between  
6 different experiments in wild-type mice. They also showed that, in some instances, PCO activity  
7 in untreated PPAR $\alpha$ -null mice was up to sixfold greater than that in wild-type mice. Parrish  
8 et al. (1996) noted that control values between experiments varied as much as a factor of twofold  
9 for PCO activity and thus, their data were presented as percentage of concurrent controls.  
10 Furthermore, Melnick et al. (1987) reported that corn oil administration alone can elevate PCO  
11 (as well as catalase) activity, and corn oil has also been reported to potentiate the induction of  
12 PCO activity of TCA in male mice (DeAngelo et al., 1989). Thus, quantitative inferences  
13 regarding the magnitude of response in these studies are limited by a number of factors. For  
14 example, in the studies reported in DeAngelo et al. (2008) a small number of animals was  
15 studied for PCO activity at interim sacrifices ( $n = 5$ ). PCO activity varied 2.7-fold as baseline  
16 controls. Although there was a 10-fold difference in TCA exposure concentration, the increase  
17 in PCO activity at 4 weeks was 1.3-, 2.4-, and 5.3-fold of control. More information on the  
18 relationship of PCO enzyme activity and its relationship to carcinogenicity is discussed in  
19 Section E.3.4 and below.

20

#### 4.5.6.1.9. Oxidative stress

21 Very limited data are available as to oxidative stress and related markers induced by the  
22 oxidative metabolites of TCE. As discussed in Appendix E, above, there are limited data that do  
23 not indicate significant oxidative stress and associated DNA damage associated with acute and  
24 subacute TCE treatment. In regard to DCA and TCA, Larson and Bull (1992b) exposed male  
25 B6C3F1 mice or Fischer 344 rats to single doses TCA or DCA in distilled water by oral gavage  
26 ( $n = 4$ ). In the first experiment, TBARS was measured from liver homogenates and assumed to  
27 be malondialdehyde. The authors stated that a preliminary experiment had shown that maximal  
28 TBARS was increased 6 hours after a dose of DCA and 9 hours after a dose of TCA in mice and  
29 that by 24 hours TBARS concentrations had declined to control values. Time-course  
30 information in rats was not presented. A dose of 100 mg/kg DCA (rats or mice) or TCA (mice)  
31 did not elevate TBARS concentrations over that of control liver with this concentration of TCA  
32 not examined in rats. For TCA, there was a slight dose-related increase in TBARS over control  
33 values starting at 300 mg/kg in mice with the increase in TBARS increasing at a rate that was

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 lower than the magnitude of increase in dose. Of note, is the report that the induction of TBARS  
2 in mice is transient and has subsided within 24 hours of a single dose of DCA or TCA, that the  
3 response in mice appeared to be slightly greater with DCA than TCA at similar doses, and that  
4 for DCA, there was similar TBARS induction between rats and mice at similar dose levels.

5 Austin et al. (1996) appears to a follow-up publication of the preliminary experiment  
6 cited in Larson and Bull (1992b). Male B6C3F1 mice were treated with single doses of DCA or  
7 TCA via gavage with liver examined for 8-OHdG. The authors stated that in order to conserve  
8 animals, controls were not employed at each time point. There was a statistically significant  
9 increase over controls in 8-OHdG for the 4- and 6-hour time points for DCA (~1.4- and 1.5-fold  
10 of control, respectively) but not at 8 hours in mice. For TCA, there was a statistically significant  
11 increase in 8-OHdG at 8 and 10 hours for TCA (~1.4- and 1.3-fold of control, respectively).

12 Consistent results as to low, transient increases in markers of “oxidative stress” were also  
13 reported by Parrish et al. (1996), who in addition to examining oxidative stress alone, attempted  
14 to examine its possible relationship to PCO and liver weight in male B6C3F1 mice exposed to  
15 TCA or DCA for 3 or 10 weeks ( $n = 6$ ). The dose-related increase in PCO activity at 21 days for  
16 TCA was reported to not be increased similarly for DCA. Only the 2.0 g/L dose of DCA was  
17 reported to induce a statistically significant increase at 21-days of exposure of PCO activity over  
18 control (~1.8-fold of control). After 71 days of treatment, TCA induced dose-related increases in  
19 PCO activities that were approximately twice the magnitude as that reported at 21 days.  
20 Treatments with DCA at the 0.1 and 0.5 g/L exposure levels produced statistically significant  
21 increase in PCO activity of ~1.5- and 2.5-fold of control, respectively. The administration of  
22 1.25 g/L clofibric acid in drinking water, used as a positive control, gave ~six–sevenfold of  
23 control PCO activity at 21 and 71 days exposure. Parrish et al. (1996) reported that laurate  
24 hydroxylase activity was reported to be elevated significantly only by TCA at 21 days and to  
25 approximately the same extent (~1.4- to 1.6-fold of control) increased at all doses tested and at  
26 71 days both the 0.5 and 2.0 g/L TCA exposures to a statistically significant increase in laurate  
27 hydroxylase activity (i.e., 1.6- and 2.5-fold of control, respectively). No change was reported  
28 after DCA exposure. Laurate hydroxylase activity within the control values varying 1.7-fold  
29 between 21 and 71 days experiments. Levels of 8-OHdG in isolated liver nuclei were reported to  
30 not be altered from 0.1, 0.5, or 2.0 g/L TCA or DCA after 21 days of exposure and this negative  
31 result was reported to remain even when treatments were extended to 71 days of treatment. The  
32 authors noted that the level of 8-OHdG increased in control mice with age (i.e., ~twofold  
33 increase between 71-day and 21-day control mice). Thus, the increases in PCO activity noted for  
34 DCA and TCA were not associated with 8-OHdG levels (which were unchanged) and also not  
35 with changes laurate hydrolase activity observed after either DCA or TCA exposure. Of note, is

1 that the authors report taking steps to minimize artifactual responses for their 8-OHdG  
2 determinations. The authors concluded that their data suggest that peroxisome proliferative  
3 properties of TCA were not linked to oxidative stress or carcinogenic response.  
4

#### **4.5.6.1.10. Comparisons of Trichloroethylene (TCE)-Induced Carcinogenic Responses With Trichloroacetic Acid (TCA), Dichloroacetic Acid (DCA), and Chloral Hydrate (CH) Studies**

#### **4.5.6.1.11. Studies in rats**

5 As discussed above, data on TCE carcinogenicity in rats, while not reporting statistically  
6 significantly increased risks, are not entirely adequate due to low numbers of animals, increased  
7 systemic toxicity, and/or increased treatment-related or accidental mortality. Notably, several  
8 studies in rats noted a few very rare types of liver or biliary tumors (cystic cholangioma,  
9 cholangiocarcinoma, or angiosarcomas) in treated animals. For TCA, DCA and CH, there are  
10 even fewer studies in rats, so there is a very limited ability to assess the consistency or lack  
11 thereof in rat carcinogenicity among these compounds.

12 For TCA, the only available study in rats (DeAngelo et al., 1997) has been frequently  
13 cited in the literature to indicate a lack of response in this species for TCA-induced liver tumors.  
14 However, this study does report an apparent dose-related increase in multiplicity of adenomas  
15 and an increase in carcinomas over control at the highest dose. The use by DeAngelo et al.  
16 (1997) of a relatively low number of animals per treatment group ( $n = 20-24$ ) limits this study's  
17 ability to determine a statistically significant increase in tumor response. Its ability to determine  
18 an absence of treatment-related effect is similarly limited. In particular, a power calculation of  
19 the study shows that for most endpoints (incidence and multiplicity of all tumors at all exposure  
20 DCA concentrations), the Type II error, which should be  $>50\%$ , was less than 8%. The only  
21 exception was for the incidence of adenomas and adenomas and carcinomas for the 0.5 g/L  
22 treatment group (58%), at which, notably, there was a reported increase in reported adenomas or  
23 adenomas and carcinomas combined over control (15 vs. 4%). Therefore, the likelihood of a  
24 false null hypothesis was not negligible. Thus, while suggesting a lower response than for mice  
25 for liver tumor induction, this study is inconclusive for determining of whether TCA induces a  
26 carcinogenic response in the liver of rats.

27 For DCA, there are two reported long-term studies in rats (DeAngelo et al., 1996;  
28 Richmond et al., 1995) that appear to have reported the majority of their results from the same  
29 data set and which consequently were subject to similar design limitations and DCA-induced

1 neurotoxicity in this species. DeAngelo et al. (1996) reported increased hepatocellular adenomas  
2 and carcinomas in male F344 rats exposed to DCA for 2 years. However, the data from  
3 exposure concentrations at a 5 g/L dose had to be discarded and the 2.5 g/L DCA dose had to be  
4 continuously lowered during the study due to neurotoxicity. There was a DCA-induced  
5 increased in adenomas and carcinomas combined reported for the 0.5 g/L DCA (24.1 vs.  
6 4.4% adenomas and carcinomas combined in treated vs. controls) and an increase at a variable  
7 dose started at 2.5 g/L DCA and continuously lowered (28.6 vs. 3.0% adenomas and carcinomas  
8 combined in treated vs. controls). Only combined incidences of adenomas and carcinomas for  
9 the 0.5 g/L DCA exposure group was reported to be statistically significant by the authors  
10 although the incidence of adenomas was 17.2 versus 4% in treated versus control rats.  
11 Hepatocellular tumor multiplicity was reported to be increased in the 0.5 g/L DCA group  
12 (0.31 adenomas and carcinomas/animal in treated vs. 0.04 in control rats) but was reported by the  
13 authors to not be statistically significant. At the starting dose of 2.5 g/L that was continuously  
14 lowered due to neurotoxicity, the increased multiplicity of hepatocellular carcinomas was  
15 reported by the authors to be to be statistically significant (0.25 carcinomas/animals vs. 0.03 in  
16 control) as well as the multiplicity of combined adenomas and carcinomas (0.36 adenomas and  
17 carcinomas/animals vs. 0.03 in control rats). Issues that affect the ability to determine the nature  
18 of the dose-response for this study include (1) the use of a small number of animals ( $n = 23$ ,  
19  $n = 21$ , and  $n = 23$  at final sacrifice for the 2.0 g/L NaCl control, 0.05 g/L and 0.5 g/L treatment  
20 groups) that limit the power of the study to both determine statistically significant responses and  
21 to determine that there are not treatment-related effects (i.e., power) (2) apparent addition of  
22 animals for tumor analysis not present at final sacrifice (i.e., 0.05 and 0.5 g/L treatment groups),  
23 and (3) most of all, the lack of a consistent dose for the 2.5 g/L DCA exposed animals.

24 Similar issues are present for the study of Richmond et al. (1995) which was conducted  
25 by the same authors as DeAngelo et al. (1996) and appeared to be the same data set. There was a  
26 small difference in reports of the results between the two studies for the same data for the 0.5 g/L  
27 DCA group in which Richmond et al. (1995) reported a 21% incidence of adenomas and  
28 DeAngelo et al. (1996) reported a 17.2% incidence. The authors did not report any of the results  
29 of DCA-induced increases of adenomas and carcinomas to be statistically significant. The same  
30 issues discussed above for DeAngelo et al. (1996) apply to this study. Similar to the DeAngelo  
31 et al. (1997) study of TCA in rats, the use in these DCA studies (DeAngelo et al., 1996;  
32 Richmond et al., 1995) of relatively small numbers of rats limits the detection of  
33 treatment-related effects and the ability to determine whether there was no treatment related  
34 effects (Type II error), especially at the low concentrations of DCA exposure.



1 For CH, George et al. (2000) exposed male F344/N rats to CH in drinking water for  
2 2 years. Groups of animals were sacrificed at 13, 26, 52, and 78 weeks following the initiation  
3 of dosing, with terminal sacrifices at Week 104. Only a few animals received a complete  
4 pathological examination. The number of animals surviving >78 weeks and the number  
5 examined for hepatocellular proliferative appeared to differ (42–44 animals examined but  
6 32–35 surviving till the end of the experiment). Only the lowest treatment group had increased  
7 liver tumors which were marginally significantly increased.

8 Leuschner and Beuscher (1998) examined the carcinogenic effects of CH in male and  
9 female Sprague-Dawley rats (69–79 g, 25–29 days old at initiation of the experiment)  
10 administered 0, 15, 45, and 135 mg/kg CH in unbuffered drinking water 7 days/week  
11 ( $n = 50/\text{group}$ ) for 124 weeks in males and 128 weeks in females. Two control groups were  
12 noted in the methods section without explanation as to why they were conducted as two groups.  
13 The authors report no substance-related influence on organ weights and no macroscopic evidence  
14 of tumors or lesions in male or female rats treated with CH for 124 or 128 weeks. However, no  
15 data are presented on the incidence of tumors in either treatment or control groups. The authors  
16 did report a statistically significant increase in the incidence of hepatocellular hypertrophy in  
17 male rats at the 135 mg/kg dose (14/50 animals vs. 4/50 and 7/50 in Controls I and II). For  
18 female rats, the incidence of hepatocellular hypertrophy was reported to be 10/50 rats (Control I)  
19 and 16/50 (Control II) rats with 18/50, 13/50 and 12/50 female rats having hepatocellular  
20 hypertrophy after 15, 45, and 135 mg/kg CH, respectively. The lack of reporting in regard to  
21 final body weights, histology, and especially background and treatment group data for tumor  
22 incidences, limit the interpretation of this study. Whether this paradigm was sensitive for  
23 induction of liver cancer cannot be determined.

24 Therefore, given the limitations in the available studies, a comparison of rat liver  
25 carcinogenicity induced by TCE, TCA, DCA, and CH reveals no strong inconsistencies, but nor  
26 does it provide much insight into the relative importance of different TCE metabolites in liver  
27 tumor induction.  
28

#### 4.5.6.1.12. Studies in mice

29 Similar to TCE, the bioassay data in mice for DCA, TCA, and CH are much more  
30 extensive and have shown that all three compounds induce liver tumors in mice. Several 2-year  
31 bioassays have been reported for CH (Daniel et al., 1992; George et al., 2000; Leakey et al.,  
32 2003a). For many of the DCA and TCA studies, the focus was not carcinogenic dose-response  
33 but rather investigation of the nature of the tumors and potential MOAs in relation to TCE. As a

1 result, studies often employed relatively high concentrations of DCA or TCA and/or were  
2 conducted for a year or less. As shown previously in Section 4.5.4.2.1, the dose-response curves  
3 for increased liver weight for TCE administration in male mice are more similar to those for  
4 DCA administration and TCE oxidative metabolism than for direct TCA administration  
5 (inadequate data were available for CH). An analogous comparison for DCA-, TCA-, and  
6 CH-induced tumors would be informative, ideally using data from 2-year studies.  
7

#### 4.5.6.1.12.1. Trichloroethylene (TCE) carcinogenicity dose-response data

8 Unfortunately, the database for TCE, while consistently showing an induction of liver  
9 tumors in mice, is very limited for making inferences regarding the shape of the dose-response  
10 curve. For many of these experiments multiplicity was not given only liver tumor incidence.  
11 NTP (1990), Bull et al. (2002), Anna et al. (1994) conducted gavage experiments in which they  
12 only tested one dose of ~1,000 mg/kg-day TCE. NCI (1976) tested two doses that were adjusted  
13 during exposure to an average of 1,169 and 2,339 mg/kg-day in male mice with only twofold  
14 dose spacing in only two doses tested. Maltoni et al. (1986) conducted inhalation experiments in  
15 two sets of B6C3F1 mice and one set of Swiss mice at three exposure concentrations that were  
16 threefold apart in magnitude between the low and mid-dose and twofold apart in magnitude  
17 between the mid- and high-dose. However, for one experiment in male B6C3F1 mice (BT306),  
18 the mice fought and suffered premature mortality and for two the experiments in B6C3F1 mice,  
19 although using the same strain, the mice were obtained from differing sources with very different  
20 background liver tumor levels. For the Maltoni et al. (1988) study a general descriptor of  
21 “hepatoma” was used for liver neoplasia rather than describing hepatocellular adenomas and  
22 carcinomas so that comparison of that data with those from other experiments is difficult. More  
23 importantly, while the number of adenomas and carcinomas may be the same between treatments  
24 or durations of exposure, the number of adenomas may decrease as the number of carcinomas  
25 increase during the course of tumor progression. Such information is lost by using only a  
26 hepatoma descriptor.

27 Given the limited database, it would be useful if different studies could be combined to  
28 yield a more comprehensive dose-response curve, as was done for liver weight, above. However,  
29 this is probably not appropriate for several reasons. First, only NTP (1990) was performed with  
30 dosing duration and time of sacrifice both being the “standard” 104 weeks. NCI (1976), Maltoni  
31 et al. (1986), Anna et al. (1994), and Bull et al. (2002) all had shorter dosing periods and either  
32 longer (Maltoni et al., 1986) or shorter (the other three studies) observation times. Therefore,  
33 because of potential dose-rate effects and differences in the degree of expression of TCE-induced

1 tumors, it is difficult to even come up with a comparable administered dose-metric across studies.  
2 Moreover, the background tumor incidences are substantially different across experiments, even  
3 controlling for mouse strain and sex. For example, across gavage studies in male B6C3F1 mice,  
4 the incidence of hepatocellular carcinomas ranged from 1.2–16.7% (Anna et al., 1994; NCI,  
5 1976; NTP, 1990) and the incidence of adenomas ranged from 1.2–14.6% (Anna et al., 1994;  
6 NTP, 1990) in control B6C3F1 mice. After ~1,000 mg/kg-day TCE treatment, the incidence of  
7 carcinomas ranged from 19.4–62% (Anna et al., 1994; Bull et al., 2002; NCI, 1976; NTP, 1990),  
8 with three of the studies (Anna et al., 1994; NCI, 1976; NTP, 1990) reporting a range of  
9 incidences between 42.8–62.0%). The incidence of adenomas ranged from 28–66.7% (Anna et  
10 al., 1994; Bull et al., 2002; NTP, 1990). In the Maltoni et al. (1986) inhalation study as well,  
11 male B6C3F1 mice from two different sources had very different control incidences of hepatomas  
12 (~2% vs. about ~20%).

13 Therefore, only data from the same experiment in which more than a single exposed dose  
14 group was used provide reliable data on the dose-response relationship for TCE  
15 hepatocarcinogenicity, and incidences from these experiments are shown in Figures 4-10 and  
16 4-11. Except for one of the two Maltoni et al. (1986) inhalation experiments in male B6C3F1  
17 mice, all of these data sets show relatively proportional increases with dose, albeit with  
18 somewhat different slopes as may be expected across strains and sexes. Direct comparison is  
19 difficult, since the “hepatomas” reported by Maltoni et al. (1986) are much more heterogeneous,  
20 including neoplastic nodules, adenomas, and carcinomas, than the carcinomas reported by NCI  
21 (1976). Nonetheless, although the data limitations preclude a conclusive statement, these data  
22 are generally consistent with the linear relationship observed with TCE-induced liver weight  
23 changes.

24

#### 4.5.6.1.12.2. **Dichloroacetic acid (DCA) carcinogenicity dose-response data**

25 With respect to DCA, Pereira (1996) reported that for 82 week exposure to DCA in  
26 female B6C3F1 mice, DCA exposure concentrations of 0, 2, 6.67, and 20 mmol/L (0, 0.26, 0.86,  
27 and 2.6 g/L) led to close proportionally increasing adenoma prevalences of 2.2, 6, 25, and 84.2%,  
28 though adenoma multiplicity increased more than linearly between the highest two doses.  
29 Unfortunately, too few carcinomas were observed at these doses and duration to meaningfully  
30 inform the shape of the dose-response relationship. More useful is DeAngelo et al. (1999),  
31 which reported on a study of DCA hepatocarcinogenicity in male B6C3F1 mice over a lifetime  
32 exposure. DeAngelo et al. (1999) used 0.05 g/L, 0.5 g/L, 1.0 g/L, 2.0 g/L and 3.5 g/L exposure  
33 concentrations of DCA in their 100-week drinking water study. The number of animals at final

1 sacrifice was generally low in the DCA treatment groups and variable. The multiplicity or  
2 number of hepatocellular carcinomas/animals was reported to be significantly increased over  
3 controls in a dose-related manner at all DCA treatments including 0.05 g/L DCA, and a  
4 no-observed-effect level (NOEL) reported not to be observed by the authors. Between the 0.5  
5 g/L and 3.5 g/L exposure concentrations of DCA the magnitude of increase in multiplicity was  
6 similar to the increases in magnitude in dose. The incidence of hepatocellular carcinomas were  
7 reported to be increased at all doses as well but not reported to be statistically significant at the  
8 0.05 g/L exposure concentration. However, given that the number of mice examined for this  
9 response ( $n = 33$ ), the power of the experiment at this dose was only 16.9% to be able to

10  
11  
12  
13  
14  
15  
16  
17  
18  
19

**Figure 4-10. Dose-response relationship, expressed as (A) percentage incidence and (B) fold-increase over controls, for TCE hepatocarcinogenicity in NCI (1976).** For comparison, incidences of carcinomas for NTP (1990), Anna et al. (1994), and Bull et al. (2002) are included, but without connecting lines since they are not appropriate for assessing the shape of the dose-response relationship.

1  
2  
3  
4  
5  
6  
7

**Figure 4-11. Dose-response relationship, expressed as (A) incidence and (B) fold-increase over controls, for TCE hepatocarcinogenicity in Maltoni et al. (1986).** Note that the BT306 experiment reported excessive mortality due to fighting, and so the paradigm was repeated in experiment BT306bis using mice from a different source.

1 determine that there was not a treatment related effect. Indeed, Figure 4-12 replots the data from  
2 DeAngelo et al. (1999) with an abscissa drawn to scale (unlike the figure in the original paper,  
3 which was not to scale), suggests even a slightly greater than linear effect at the lowest dose  
4 (0.05 g/L, or 8 mg/kg-day) as compared to the next lowest dose (0.5 g/L, or 84 mg/kg-day),  
5 though of course the power of such a determination is limited. The authors did not report the  
6 incidence or multiplicity of adenomas for the 0.05 g/L exposure group in the study or the  
7 incidence or multiplicity of adenomas and carcinomas in combination. For the animals surviving  
8 from 79–100 weeks of exposure, the incidence and multiplicity of adenomas peaked at 1 g/L  
9 while hepatocellular carcinomas continued to increase at the higher doses. This would be  
10 expected where some portion of the adenomas would either regress or progress to carcinomas at  
11 the higher doses.  
12

13  
14 **Figure 4-12. Dose-response data for hepatocellular carcinomas (HCC)**  
15 **(A) incidence and (B) multiplicity, induced by DCA from DeAngelo et al.**  
16 **(1999).** Drinking water concentrations were 0, 0.05, 0.5, 1, 2, and 3.5 g/L, from  
17 which daily average doses were calculated using observed water consumption in the  
18 study.  
19

20  
21 Associations of DCA carcinogenicity with various noncancer, possibly precursor, effects  
22 was also investigated. Importantly, the doses that induced tumors in DeAngelo et al. (1999)  
23 were reported to not induce widespread cytotoxicity. An attempt was also made to relate  
24 differing exposure levels to subchronic changes and peroxisomal enzyme induction.  
25 Interestingly, DeAngelo et al. (1999) reported that peroxisome proliferation was significantly  
26 increased at 3.5 g/L DCA only at 26 weeks, not correlated with tumor response, and to not be

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 increased at either 0.05 g/L or 0.5 g/L treatments. The authors concluded that DCA-induced  
2 carcinogenesis was not dependent on peroxisome proliferation or chemically sustained  
3 proliferation, as measured by DNA synthesis. Slight hepatomegaly was present by 26 weeks in  
4 the 0.5 g/L group and decreased with time. By contrast, increases in both percentage liver/body  
5 weight and the multiplicity of hepatocellular carcinomas increased proportionally with DCA  
6 exposure concentration after 79–100 weeks of exposure. DeAngelo et al. (1999) presented a  
7 figure comparing the number of hepatocellular carcinomas/animal at 100 weeks compared with  
8 the percentage liver/body weight at 26 weeks that showed a linear correlation ( $r^2 = 0.9977$ ) while  
9 peroxisome proliferation and DNA synthesis did not correlate with tumor induction profiles.  
10 The proportional increase in liver weight with DCA exposure was also reported for shorter  
11 durations of exposure as noted previously. Therefore, for DCA, both tumor incidence and liver  
12 weight appear to increase proportionally with dose.  
13

#### 4.5.6.1.12.3. **Trichloroacetic acid (TCA) carcinogenicity dose-response data**

14 With respect to TCA, Pereira (1996) reported that for 82 week exposure to TCA in  
15 female B6C3F1 mice, TCA exposure concentrations of 0, 2, 6.67, and 20 mmol/L (0, 0.33, 1.1,  
16 and 3.3 g/L) led to increasing incidences and multiplicity of adenomas and of carcinomas (see  
17 Figure 4-13). DeAngelo et al. (2008) reported the results of three experiments exposing male  
18 B6C3F1 mice to neutralized TCA in drinking water (incidences also in Figure 4-13). Rather  
19 than using five exposure levels that were generally twofold apart, as was done in DeAngelo et al.  
20 (1999) for DCA, DeAngelo et al. (2008) studied only three doses of TCA that were an order of  
21 magnitude apart which limits the elucidation of the shape of the dose-response curve. In  
22 addition, the 104-week data, DeAngelo et al. (2008) contained two studies, each conducted in a  
23 separate laboratories—the two lower doses were studied in one study and the highest dose in  
24 another. The first 104-week study was conducted using 2 g/L NaCl, or 0.05, 0.5, or 5 g/L TCA  
25 in drinking water for 60 weeks (Study #1) while the other two were conducted for a period of  
26 104 weeks (Study #2 with 2.5 g/L neutralized acetic acid or 4.5 g/L TCA exposure groups and  
27 Study #3 with deionized water, 0.05 g/L TCA and 0.5 g/L TCA exposure groups). In addition, a  
28 relatively small number of animals were used for the determination of a tumor response ( $n \sim 30$   
29 at final necropsy).

1  
2  
3 **Figure 4-13. Reported incidences of hepatocellular carcinomas (HCC) and**  
4 **hepatocellular adenomas plus carcinomas (HCA + HCC) in various studies**  
5 **in B6C3F1 mice (DeAngelo et al., 2008; Pereira, 1996).** Combined  
6 HCA + HCC were not reported in Pereira (1996).  
7  
8

9 In Study #1, the incidence data for adenomas observed at 60 weeks at 0.05 g/L, 0.5 g/L  
10 and 5.0 g/L TCA were 2.1-, 3.0- and 5.4-fold of control values, with similar fold increases in  
11 multiplicity. As shown by Pereira (1996), 60 weeks does not allow for full tumor expression, so  
12 whether the dose-response relationship is the same at 104 weeks is not certain. For instance,  
13 Pereira (1996) examined the tumor induction in female B6C3F1 mice and demonstrated that foci,  
14 adenoma, and carcinoma development in mice are dependent on duration of exposure (period of  
15 observation in controls). In control female mice a 360- versus 576-day observation period  
16 showed that at 360 days no foci or carcinomas and only 2.5% of animals had adenomas whereas  
17 by 576 days of observation, 11% had foci, 2% adenomas, and 2% had carcinomas. For DCA and  
18 TCA treatments, foci, adenomas, and carcinoma incidence and multiplicity did not reach full  
19 expression until 82 weeks at the three doses employed. Although the numbers of animals were  
20 relatively low and variable at the two highest doses (18–28 mice) there were 50–53 mice studied  
21 at the lowest dose level and 90 animals studied in the control group.

22 Therefore, the 104-week DeAngelo et al. (2008) data from Studies #2 and #3 would  
23 generally be preferred for elucidating the TCA dose-response relationship. However, Study #2  
24 was only conducted at one dose, and although Study #3 used lower doses, it exhibited  
25 extraordinarily high control incidences of liver tumors. In particular, while the incidence of  
26 adenomas and carcinomas was 12% in Study #2, it was reported to be 64% in Study #3. The

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 mice in Study #3 were of very large size (weighing ~50 g at 45 weeks) as compared to Study #1,  
2 Study #2, or most other bioassays in general, and the large background rate of tumors reported is  
3 consistent with the body-weight-dependence observed by Leakey et al. (2003b).


4 To put into context the 64% incidence data for carcinomas and adenomas reported in  
5 DeAngelo et al. (2008) for the control group of Study #3, other studies cited in this review for  
6 male B6C3F1 mice show a much lower incidence in liver tumors with (1) NCI (1976) study of  
7 TCE reporting a colony control level of 6.5% for vehicle and 7.1% incidence of hepatocellular  
8 carcinomas for untreated male B6C3F1 mice ( $n = 70-77$ ) at 78 weeks, (2) Herren-Freund et al.  
9 (1987) reporting a 9% incidence of adenomas in control male B6C3F1 mice with a multiplicity  
10 of  $0.09 \pm 0.06$  and no carcinomas ( $n = 22$ ) at 61 weeks, (3) NTP (1990) reporting an incidence of  
11 14.6% adenomas and 16.6% carcinomas in male B6C3F1 mice after 103 weeks ( $n = 48$ ), and  
12 (4) Maltoni et al. (1986) reporting that B6C3F1 male mice from the “NCI source” had a  
13 1.1% incidence of “hepatoma” (carcinomas and adenomas) and those from “Charles River Co.”  
14 had a 18.9% incidence of “hepatoma” during the entire lifetime of the mice ( $n = 90$  per group).  
15 The importance of examining an adequate number of control or treated animals before  
16 confidence can be placed in those results is illustrated by Anna et al. (1994) in which at  
17 76 weeks 3/10 control male B6C3F1 mice that were untreated and 2/10 control animals given  
18 corn oil were reported to have adenomas but from 76–134 weeks, 4/32 mice were reported to  
19 have adenomas (multiplicity of  $0.13 \pm 0.06$ ) and 4/32 mice were reported to have carcinomas  
20 (multiplicity of  $0.12 \pm 0.06$ ). Thus, the reported combined incidence of carcinomas and  
21 adenomas of 64% reported by DeAngelo et al. (2008) for the control mice of Study # 3, not only  
22 is inconsistent and much higher than those reported in Studies #1 and #2, but also much higher  
23 than reported in a number of other studies of TCE.

24 Therefore, this large background rate and the increased mortality for these mice limit  
25 their use for determining the nature of the dose-response for TCA liver carcinogenicity. At the  
26 two lowest doses of 0.05 g/L and 0.5 g/L TCA from Study #3, the differences in the incidences  
27 and multiplicities for all tumors were twofold at 104 weeks. However, there was no difference in  
28 any of the tumor results (i.e., adenoma, carcinoma, and combinations of adenoma and carcinoma  
29 incidence and multiplicity) between the 4.5 g/L dose group in Study #2 and the 0.5 g/L dose  
30 group in Study #3 at 104 weeks. By contrast, at 60 weeks of exposure, but within the same study  
31 (Study #1), there was a twofold increase in multiplicity for adenomas, and for adenomas and  
32 carcinomas combined between the 0.5 and 5.0 g/L TCA exposure groups. These results are  
33 consistent with the two highest exposure levels reaching a plateau of response after a long  
34 enough duration of exposure for full expression of the tumors (i.e., ~90% of animals having liver  
35 tumors at the 0.5 g/L and 5 g/L exposures). However, whether such a plateau would have been

1 observed in mice with a more “normal” body weight, and hence a lower background tumor  
2 burden cannot be determined.

3 Because of the limitations of different studies, it is difficult to discern whether the liver  
4 tumor dose-response curves of TCA and DCA are different in a way analogous to that for liver  
5 weight (see Figure 4-14). Certainly, it is clear that at the same concentration in drinking water or  
6 estimated applied dose, DCA is more potent than TCA, as DCA induces nearly 100% incidence  
7 of carcinomas at a lower dose than TCA. Therefore, like with liver weight gains, DCA has a  
8 steeper dose-response function than TCA. However, the evidence for a “plateau” in tumor  
9 response at high doses with TCA, as was observed for liver weight, is equivocal, as it is  
10 confounded by the highly varying background tumor rates and the limitations of the available  
11 study paradigms.

12



13

14 **Figure 4-14. Reported incidence of hepatocellular carcinomas induced by**  
15 **DCA and TCA in 104-week studies (DeAngelo et al., 2008; DeAngelo et al.,**  
16 **1999).** Only carcinomas were reported in DeAngelo et al. (1999), so combined  
17 adenomas and carcinomas could not be compared.

18

19

20 DeAngelo et al. (2008) attempt to identify a NOEL for tumorigenicity using tumor  
21 multiplicity data and estimated TCA dose. However, it is not an appropriate descriptor for these  
22 data, especially given that “statistical significance” of the tumor response is the determinant used  
23 by the authors to support the conclusions regarding a dose in which there is no TCA-induced  
24 effect. Due to issues related to the appropriateness of use of the concurrent control in Study #3,

1 only the 60-week experiment (i.e., Study # 1) is useful for the determination of tumor  
2 dose-response. Not only is there not allowance for full expression of a tumor response at the  
3 60-week time point but a power calculation of the 60-week study shows that the Type II error,  
4 which should be >50% and thus, greater than the chances of “flipping a coin,” was 41 and 71%  
5 for incidence and 7 and 15% for multiplicity of adenomas for the 0.05 and 0.5 g/L TCA exposure  
6 groups. For the combination of adenomas and carcinomas, the power calculation was 8 and 92%  
7 for incidence and 6 and 56% for multiplicity at 0.05 and 0.5 g/L TCA exposure. Therefore, the  
8 designed experiment could accept a false null hypothesis, especially in terms of tumor  
9 multiplicity, at the lower exposure doses and erroneously conclude that there is no response due  
10 to TCA treatment.

11 In terms of correlations with other noncancer, possibly precursor effects, DeAngelo et al.  
12 (2008) also reported that PCO activity, which varied 2.7-fold as baseline controls, was 1.3-, 2.4-,  
13 and 5.3-fold of control for the 0.05, 0.5, and 5 g/L TCA exposure groups in Study #1 at 4 weeks  
14 was for adenomas incidence 2.1-, 3.0-, and 5.4-fold of control and not similar at the lowest dose  
15 level at 60 weeks. However, it is not clear whether the similarity between PCO and  
16 carcinogenicity at 60 weeks would persist for tumor incidence at 104 weeks. DeAngelo et al.  
17 (2008) report a regression analyses that compare “percent of hepatocellular neoplasia,” indicated  
18 by tumor multiplicity, with TCA dose, represented by estimations of the TCA dose in  
19 mg/kg-day, and with PCO activity for the 60-week and 104-week data. Whether adenomas and  
20 carcinomas combined or individual tumor type were used in these analysis was not reported by  
21 the authors. However, it would be preferable to compare “precursor” levels of PCO at earlier  
22 time points, rather than at a time when there was already a significant tumor response. In  
23 addition, linear regression analyses of these data are difficult to interpret because of the wide  
24 dose spacing of these experiments. In such a situation, for a linear regression, control and 5 g/L  
25 exposure levels will basically determine the shape of the dose-response curve since the 0.05 g/L  
26 and 0.5 g/L exposure levels are so close to the control (zero) value. Thus, dose-response appears  
27 to be linear between control and the 5.0 g/L value with the two lowest doses not affectively  
28 changing the slope of the line (i.e., “leveraging” the regression). Moreover, at the 5 g/L dose  
29 level, there is potential for effects due to palatability, as reported in one study in which drinking  
30 water consumption declined at this concentration (DeAngelo et al., 2008). Thus, the value of  
31 these analyses is limited by (1) use of data from Study # 3 in a tumor prone mouse that is not  
32 comparable to those used in Studies #1 and #2, (2) the appropriateness of using PCO values from  
33 later time points and the variability in PCO control values, (3) the uncertainty of the effects of  
34 palatability on the 5 g/L TCA results which were reported in one study to reduce drinking water  
35 consumption, and (4) the dose-spacing of the experiment.

#### 4.5.6.1.12.4. Chloral hydrate (CH) carcinogenic dose-response

1 Although a much more limited database in rodents than for TCA or DCA, there is  
2 evidence that chloral hydrate is also a rodent liver hepatocarcinogen (see also Section E.2.5 and  
3 Caldwell and Keshava [2006]).


4 Daniel et al. (1992) exposed adult male B6C3F1 28-day-old mice to 1 g/L CH in drinking  
5 water for 30 and 60 weeks ( $n = 5$  for interim sacrifice) and for 104 weeks ( $n = 40$ ). The  
6 concentration of CH was 1 g/L and estimated to provide a 166-mg/kg-day dose. It is not clear  
7 from the report what control group better matched the CH group, as the mean initial body  
8 weights of the groups as well as the number of animals varied considerably in each group (i.e.,  
9 ~40% difference in mean body weights at the beginning of the study). Liver tumors were  
10 increased by CH treatment. The percentage incidence of liver carcinomas and adenomas in the  
11 surviving animals was 15% in control and 71% in CH-treated mice and the incidence of  
12 hepatocellular carcinoma reported to be 46% in the CH-treated group. The number of  
13 tumors/animals was also significantly increased with CH treatment. However, because this was  
14 a single dose study, a comparison with the dose-response relationship with TCE, TCA, or DCA  
15 is not feasible.

16 George et al. (2000) exposed male B6C3F1 mice to CH in drinking water for 2 years.  
17 Groups of animals were sacrificed at 26, 52, and 78 weeks following the initiation of dosing,  
18 with terminal sacrifices at Week 104. Only a few animals received a complete pathological  
19 examination. Preneoplastic foci and adenomas were reported to be increased in the livers of all  
20 CH treatment groups at 104 weeks. The percentage incidence of hepatocellular adenomas was  
21 reported to be 21.4, 43.5, 51.3, and 50% in control, 13.5, 65.0 and 146.6 mg/kg-day CH  
22 treatment groups, respectively. The percentage incidence of hepatocellular carcinomas was  
23 reported to be 54.8, 54.3, 59.0 and 84.4% in these same groups. The resulting percentage  
24 incidence of hepatocellular adenomas and carcinomas was reported to be 64.3, 78.3, 79.5 and  
25 90.6%. Of concern is the reporting of a 64% incidence of hepatocellular carcinomas and  
26 adenomas in the control group of mice for this experiment, which is the same as that for another  
27 study published by this same laboratory (DeAngelo et al., 2008). DeAngelo et al. (2008) did not  
28 identify them as being contemporaneous studies or sharing controls, but a comparison of the  
29 control data published by DeAngelo et al. (2008) for TCA and that published by George et al.  
30 (2000) for the CH studies shows them to be the same data set. Therefore, as discussed above,  
31 this data set was derived from B6C3F1 mice that were large (~50 g) and resultantly tumor prone,  
32 making determinations of the dose-response of CH from this experiment difficult. Therefore, for  
33 the purposes of comparison of dose-response relationships, this study has the same limitations as  
34 the DeAngelo et al. (2008) study, discussed above.

1           Leakey et al. (2003a) studied the effects of CH exposure (0, 25, 50, and 100 mg/kg-day,  
2 5 days/week, 104–105 weeks via gavage) in male B6C3F1 mice with dietary control used to  
3 manipulate body growth ( $n = 48$  for 2-year study and  $n = 12$  for the 15-month interim study).  
4 Dietary control was reported to decrease background liver tumor rates (decreased by 15–20%)  
5 and was reported to be associated with decreased variation in liver-to-body weight ratios, thereby  
6 potentially increasing assay sensitivity. In dietary-controlled groups and groups fed *ad libitum*,  
7 liver adenomas and carcinomas (combined) were reported to be increased with CH treatment.  
8 With dietary restriction there was a more discernable CH tumor-response with overall tumor  
9 incidence reduced, and time-to-tumor increased by dietary control in comparison to *ad libitum*  
10 fed mice. Incidences of hepatocellular adenoma and carcinoma overall rates were reported to be  
11 33, 52, 49, and 46% for control, 25, 50, and 100 mg/kg *ad libitum*-fed mice, respectively. For  
12 dietary controlled mice the incidence rates were reported to be 22.9, 22.9, 29.2, and 37.5% for  
13 controls, 25, 50, and 100 mg/kg CH, respectively. Body weights were matched and carefully  
14 controlled in this study. These data are shown in Figure 4-15, relative to control incidences. It is  
15 evident from these data that dietary control significantly changes the apparent shape of the  
16 dose-response curve, presumably by reducing variability between animals. While the *ad libitum*  
17 dose groups had an apparent “saturation” of response, this was not evident with the dietary  
18 controlled group. Of note all the other bioassays for TCE, TCA, DCA, and CH were in *ad*  
19 *libitum* fed mice. Therefore, it is difficult to compare the dose-response curves for CH-treated  
20 mice on dietary restriction to those fed *ad libitum*. However, the rationale for dietary restriction  
21 in the B6C3F1 mouse is to prevent the types of weight gain and corresponding high background  
22 tumor levels observed in DeAngelo et al. (2008) and George et al. (2000). As stated previously,  
23 most other studies of TCA, DCA, and TCE had background levels that, while varied, were lower  
24 than the *ad libitum* fed mice studied in Leakey et al. (2003a).

25           Of note is that incidences of adenomas and carcinomas combined do not show  
26 differences in tumor progression as carcinomas may increase and adenomas may regress. Liver  
27 weight increases at 15-months did not correlate with 2-year tumor incidences in the *ad libitum*  
28 group, but a consistent dose-response shape between these two measures is evident in the dietary  
29 controlled group. However, of note is the reporting of liver weight at 15 months is for a time  
30 period in which foci and liver tumors have been reported to have already occurred in other  
31 studies, so hepatomegaly in the absence of these changes is hard to detect.

32           In terms of other noncancer effects that may be associated with tumor induction, it is  
33 notable that while dietary restriction reduced the overall level of CH-mediated tumor induction,  
34 it led to greater CH-mediated induction of peroxisome proliferation-associated enzymes.  
35 Moreover, between control groups, dietary restricted mice appeared to have higher levels of



1

1 **Figure 4-15. Effects of dietary control on the dose-response curves for**  
2 **changes in liver tumor incidences induced by CH in diet (Leakey et al.,**  
3 **2003a).**  
4  
5

6 lauric acid  $\omega$ -hydroxylase activity than *ad libitum*-fed mice. Seng et al. (2003) report that lauric  
7 acid  $\beta$ -hydroxylase and PCO were induced only at exposure levels >100 mg/kg CH, again with  
8 dietary restricted groups showing the greatest induction. Such data argue against the role of  
9 peroxisome proliferation in CH-liver tumor induction in mice.

10 Leakey et al. (2003a) gave no descriptions of liver pathology were given other than  
11 incidence of mice with fatty liver changes. Hepatic malondialdehyde concentration in *ad libitum*  
12 fed and dietary controlled mice did not change with CH exposure at 15 months but the dietary  
13 controlled groups were all approximately half that of the *ad libitum*-fed mice. Thus, while  
14 overall increased tumors observed in the *ad libitum* diet correlated with increased  
15 malondialdehyde concentration, there was no association between CH dose and malondialdehyde  
16 induction for either diet.

17 Overall, from the CH studies in mice, there is an apparent increase in liver adenomas and  
18 carcinomas induced by CH treatment by either drinking water or gavage with all available  
19 studies performed in male B6C3F1 mice. However, the background levels of hepatocellular  
20 adenomas and carcinomas in these mice in George et al. (2000) and body-weight data from this  
21 study are high, consistent with the association between large body weight and background tumor  
22 susceptibility shown with dietary control (Leakey et al., 2003a). With dietary control, Leakey  
23 et al. (2003a) report a dose-response relationship between exposure and tumor incidence that is  
24 proportional to dose.  
25

4.5.6.1.12.5. **Degree of concordance among trichloroethylene (TCE), trichloroacetic acid (TCA), dichloroacetic acid (DCA), and chloral hydrate (CH) dose-response relationships**

26 A quantitative comparison of the carcinogenicity dose-response relationships among  
27 TCE, TCA, DCA, and CH—analogue to the quantitative comparison between TCE and TCA  
28 hepatomegaly—was considered. This first step in such a comparison would an examination of  
29 the dose-response data for TCE alone to see if they are consistent with a single dose-response  
30 relationship. As shown in Figure 4-10 and 4-11, there is substantial variability among the  
31 available liver tumor dose-response data that was not observed for hepatomegaly. The strain of  
32 mice used in the bioassays not only had a difference in TCE liver tumor response but also a  
33 difference in background liver tumor incidence. Differences in exposure paradigms in the

1 bioassays also leads to difference in tumor incidence and reporting. In addition, unlike the case  
2 with TCE hepatomegaly data in mice, the TCE dose-response data for liver tumors in mice  
3 exposed via inhalation and gavage are not consistent with a common dose-response curve even  
4 on an internal dose basis (e.g., Rhomberg, 2000; Section 5.2). This heterogeneity is also evident  
5 for the TCA dose-response data, as shown in Figure 4-13, which may in part be due to the  
6 differences in study duration. Furthermore, among all the available cancer bioassay data for  
7 TCE, TCA, DCA, and CH, the control incidences for background liver tumors vary from about  
8 1% to over 50%, and difference of more than 50-fold that adds substantial uncertainty to any  
9 joint analysis. Therefore, differences within and across the databases of these compounds, such  
10 as the comparability of study durations, control tumor incidences, and carcinogenic potency,  
11 preclude either a quantitative analysis or a definitive conclusion. This question may be better  
12 addressed experimentally where similar animals are exposed to different compounds in the same  
13 experimental setting.  
14

#### **4.5.6.1.13. Inferences from liver tumor phenotype and genotype**

15 A number of studies have investigation tumor phenotypes, such as c-Jun staining,  
16 tincture, and dysplacidity, or genotypes, such as H-ras mutations, to inform both the identification  
17 of the active agents of TCE liver tumor induction as well as what MOA(s) may be involved.  
18

##### **4.5.6.1.13.1. Tumor phenotype—staining and appearance**

19 The descriptions of tumors in mice reported by the NCI, NTP, and Maltoni et al studies  
20 are also consistent with phenotypic heterogeneity as well as spontaneous tumor morphology (see  
21 Section E.3.4.1.5). As noted in Section E.3.1, hepatocellular carcinomas observed in humans are  
22 also heterogeneous. For mice, Maltoni et al. (1986) described malignant tumors of hepatic cells  
23 to be of different subhistotypes, and of various degrees of malignancy and were reported to be  
24 unique or multiple, and have different sizes (usually detected grossly at necropsy) from TCE  
25 exposure. In regard to phenotype, tumors were described as usual type observed in Swiss and  
26 B6C3F1 mice, as well as in other mouse strains, either untreated or treated with  
27 hepatocarcinogens and to frequently have medullary (solid), trabecular, and pleomorphic  
28 (usually anaplastic) patterns. For the NCI (1976) study, the mouse liver tumors were described  
29 in detail and to be heterogeneous “as described in the literature” and similar in appearance to  
30 tumors generated by carbon tetrachloride. The description of liver tumors in this study and  
31 tendency to metastasize to the lung are similar to descriptions provided by Maltoni et al. (1986)

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 for TCE-induced liver tumors in mice via inhalation exposure. The NTP (1990) study reported  
2 TCE exposure to be associated with increased incidence of hepatocellular carcinoma (tumors  
3 with markedly abnormal cytology and architecture) in male and female mice. Hepatocellular  
4 adenomas were described as circumscribed areas of distinctive hepatic parenchymal cells with a  
5 perimeter of normal appearing parenchyma in which there were areas that appeared to be  
6 undergoing compression from expansion of the tumor. Mitotic figures were sparse or absent but  
7 the tumors lacked typical lobular organization. Hepatocellular carcinomas were reported to have  
8 markedly abnormal cytology and architecture with abnormalities in cytology cited as including  
9 increased cell size, decreased cell size, cytoplasmic eosinophilia, cytoplasmic basophilia,  
10 cytoplasmic vacuolization, cytoplasmic hyaline bodies and variations in nuclear appearance.  
11 Furthermore, in many instance several or all of the abnormalities were reported to be present in  
12 different areas of the tumor and variations in architecture with some of the hepatocellular  
13 carcinomas having areas of trabecular organization. Mitosis was variable in amount and  
14 location. Therefore, the phenotype of tumors reported from TCE exposure was heterogeneous in  
15 appearance between and within tumors from all three of these studies.

16 Caldwell and Keshava (2006) report “that Bannasch (2001) and Bannasch et al. (2001)  
17 describe the early phenotypes of preneoplastic foci induced by many oncogenic agents  
18 (DNA-reactive chemicals, radiation, viruses, transgenic oncogenes and local hyperinsulinism) as  
19 insulinomimetic. These foci and tumors have been described by tincture (after hematoxylin and  
20 eosin staining of structural contents) as primarily eosinophilic (pink, reflecting eosin staining,  
21 e.g., staining of intracellular and extracellular protein), basophilic (blue, reflecting hematoxylin  
22 staining, e.g., staining of ribosomes and arginine rich basic nucleorprotein such as histones), and  
23 to be heterogeneous. Primary eosin staining is associated with a less malignant state of the  
24 hepatocyte with increased ribosomal content, decreased glycogen content, and increased  
25 basophilia of the cytoplasm by hematoxylin staining to be indicative of a more malignant state or  
26 tumor progression (Bannasch, 2001; Carter et al., 2003). Several studies do identify foci and  
27 tumors as primarily eosinophilic or basophilic, but do not give specific criteria for how a foci or  
28 tumor (which can be and usually is made up of a mixture of phenotypically heterogenous cells)  
29 are assigned to be one category or another. Caldwell and Keshava (2006) noted that the tumors  
30 observed after TCE exposure are consistent with the description for the main tumor lines of  
31 development described by Bannasch et al. (2001) (see Section 3.4.1.5). Thus, the response of  
32 liver to DCA (glycogenesis with emergence of glycogen poor tumors) is similar to the  
33 progression of preneoplastic foci to tumors induced from a variety of agents and conditions  
34 associated with increased cancer risk.” Furthermore Caldwell and Keshava (2006) note that Bull  
35 et al. (2002) report expression of insulin receptor to be elevated in tumors of control mice or

1 mice treated with TCE, TCA and DCA but not in nontumor areas suggesting that this effect is  
2 not specific to DCA.

3 There is a body of literature that has focused on the effects of TCE and its metabolites  
4 after rats or mice have been exposed to “mutagenic” agents to “initiate” hepatocarcinogenesis  
5 and this is discussed in Section E.4.2. TCE and its metabolites were reported to affect tumor  
6 incidence, multiplicity, and phenotype when given to mice as a coexposure with a variety of  
7 “initiating” agents and with other carcinogens. Pereira and Phelps (1996) reported that methyl  
8 nitrosourea (MNU) alone induced basophilic foci and adenomas. MNU and low concentrations  
9 of DCA or TCA in female mice were reported to induce heterogeneous for foci and tumor with a  
10 higher concentration of DCA inducing more eosinophilic and a higher concentration of TCA  
11 inducing more tumors that were basophilic. Pereira et al. (2001) reported that not only dose, but  
12 gender also affected phenotype in mice that had already been exposed to MNU and were then  
13 exposed to DCA. As for other phenotypic markers, Lantendresse and Pereira (1997) reported  
14 that exposure to MNU and TCA or DCA induced tumors that had some commonalities, were  
15 heterogeneous, but for female mice were overall different between DCA and TCA as  
16 coexposures with MNU.

17 With regard to the phenotype of TCA and DCA-induced tumors, Stauber and Bull (1997)  
18 reported the for male B6C3F1 mice, DCA-induced “lesions” contained a number of smaller  
19 lesions that were heterogeneous and more eosinophilic with larger “lesions” tending to less  
20 numerous and more basophilic. For TCA results using this paradigm, the “lesions” were  
21 reported to be less numerous, more basophilic, and larger than those induced by DCA. Carter  
22 et al. (2003) used tissues from the DeAngelo et al. (1999) and examined the heterogeneity of the  
23 DCA-induced lesions and the type and phenotype of preneoplastic and neoplastic lesions pooled  
24 across all time points. Carter et al. (2003) examined the phenotype of liver tumors induced by  
25 DCA in male B6C3F1 mice and the shape of the dose-response curve for insight into its MOA.  
26 They reported a dose-response of histopathologic changes (all classes of premalignant lesions  
27 and carcinomas) occurring in the livers of mice from 0.05–3.5 g/L DCA for 26–100 weeks and  
28 suggest foci and adenomas demonstrated neoplastic progression with time at lower doses than  
29 observed DCA genotoxicity. Preneoplastic lesions were identified as eosinophilic, basophilic  
30 and/or clear cell (grouped with clear cell and mixed cell) and dysplastic. Altered foci were  
31 50% eosinophilic with about 30% basophilic. As foci became larger and evolved into  
32 carcinomas they became increasingly basophilic. The pattern held true throughout the exposure  
33 range. There was also a dose and length of exposure related increase in atypical nuclei in  
34 “noninvolved” liver. Glycogen deposition was also reported to be dose-dependent with  
35 periportal accumulation at the 0.5 g/L exposure level. Carter et al. (2003) suggested that size and

1 evolution into a more malignant state are associated with increasing basophilia, a conclusion  
2 consistent with those of Bannasch (1996) and that there a greater periportal location of lesions  
3 suggestive as the location from which they arose. Consistent with the results of DeAngelo et al.  
4 (1999), Carter et al. (2003) reported that DCA (0.05–3.5 g/L) increased the number of lesions  
5 per animal relative to animals receiving distilled water, shortened the time to development of all  
6 classes of hepatic lesions, and that the phenotype of the lesions were similar to those  
7 spontaneously arising in controls. Along with basophilic and eosinophilic lesions or foci,  
8 Carter et al. (2003) concluded that DCA-induced tumors also arose from isolated, highly  
9 dysplastic hepatocytes in male B6C3F1 mice chronically exposed to DCA suggesting another  
10 direct neoplastic conversion pathway other than through eosinophilic or basophilic foci.

11 Rather than male B6C3F1 mice, Pereira (1996) studied the dose-response relationship for  
12 the carcinogenic activity of DCA and TCA and characterized their lesions (foci, adenomas and  
13 carcinomas) by tincture in females (the generally less sensitive gender). Like the studies of TCE  
14 by Maltoni et al. (1986), female mice were also reported to have increased liver tumors after  
15 TCA and DCA exposures. Pereira (1996) pool lesions were pooled for phenotype analysis so the  
16 affect of duration of exposure could not be determined nor adenomas separated from carcinomas  
17 for “tumors.” However, as the concentration of DCA was decreased the number of foci was  
18 reported by Pereira (1996) to be decreased but the phenotype of the foci to go from primarily  
19 eosinophilic foci (i.e., ~95% eosinophilic at 2.58 g/L DCA) to basophilic foci (~57%  
20 eosinophilic at 0.26 g/L). For TCA the number of foci was reported to ~40 basophilic and  
21 ~60 eosinophilic regardless of dose. Spontaneously occurring foci were more basophilic by a  
22 ratio of 7/3. Pereira (1996) described the foci of altered hepatocytes and tumors induced by  
23 DCA in female B6C3F1 mice to be eosinophilic at higher exposure levels but at lower or  
24 intermittent exposures to be half eosinophilic and half basophilic. Regardless of exposure level,  
25 half of the TCA-induced foci were reported to be half eosinophilic and half basophilic with  
26 tumors 75% basophilic. In control female mice, the limited numbers of lesions were mostly  
27 basophilic, with most of the rest being eosinophilic with the exception of a few mixed tumors.  
28 The limitations of descriptions tincture and especially for inferences regarding peroxisome  
29 proliferator from the description of “basophilia” is discussed in Section E.3.4.1.5.

30 Thus, the results appear to differ between male and female B6C3F1 mice in regard to  
31 tincture for DCA and TCA at differing doses. What is apparent is that the tincture of the lesions  
32 is dependent on the stage of tumor progression, agent (DCA or TCA), gender, and dose. Also  
33 what is apparent from these studies is the both DCA and TCA are heterogeneous in their tinctoral  
34 characteristics.

1 Overall, tumors induced by TCA, DCA, CH, and TCE are all heterogeneous in their  
2 physical and tinctural characteristics in a manner this not markedly distinguishable from  
3 spontaneous lesions or those induced by a wide variety of chemical carcinogens. For instance,  
4 Daniel et al. (1992), which studies DCA and CH carcinogenicity (discussed above) noted that  
5 morphologically, there did not appear to be any discernable differences in the visual appearance  
6 of the DCA- and CH-induced tumors. Therefore, these data do not provide strong insights into  
7 elucidating the active agent(s) for TCE hepatocarcinogenicity or their MOA(s).

#### 8 4.5.6.1.13.2. **C-Jun staining**

9 Stauber and Bull (1997) reported that in male B6C3F1 mice, the oncoproteins c-Jun and  
10 c-Fos were expressed in liver tumors induced by DCA but not those induced by TCA. Although  
11 Bull et al. (2004) have suggested that the negative expression of c-Jun in TCA-induced tumors  
12 may be consistent with a characteristic phenotype shown in general by peroxisome proliferators  
13 as a class, as pointed out by Caldwell and Keshava (2006), there is no supporting evidence of  
14 this. Nonetheless, the observation that TCA and DCA have different levels of oncogene  
15 expression led to a number of follow-up studies by this group. No data on oncoprotein  
16 immunostaining are available for CH.

17 Stauber et al. (1998) studied induction of “transformed” hepatocytes by DCA and TCE  
18 treatment in vitro, including an examination of c-Jun staining. Stauber et al. (1998) isolated  
19 primary hepatocytes from 5–8 week old male B6C3F1 mice ( $n = 3$ ) and subsequently cultured  
20 them in the presence of DCA or TCA. In a separate experiment 0.5 g/L DCA was given to mice  
21 as pretreatment for 2 weeks prior to isolation. The authors assumed that the  
22 anchorage-independent growth of these hepatocytes was an indication of an “initiated cell.”  
23 After 10 days in culture with DCA or TCA (0, 0.2, 0.5 and 2.0 mM), concentrations of 0.5 mM  
24 or more DCA and TCA both induced an increase in the number of colonies that was statistically  
25 significant, with DCA showing dose-dependence as well as slightly greater overall increases than  
26 TCA. In a time course experiment the number of colonies from DCA treatment in vitro peaked  
27 by 10 days and did not change through Days 15–25 at the highest dose and, at lower  
28 concentrations of DCA, increased time in culture induced similar peak levels of colony  
29 formation by Days 20–25 as that reached by 10 days at the higher dose. Therefore, the number  
30 of colonies formed was independent of dose if the cells were treated long enough in vitro.  
31 However, not only did treatment with DCA or TCA induce anchorage independent growth but  
32 untreated hepatocytes also formed larger numbers of colonies with time, although at a lower rate  
33 than those treated with DCA. The level reached by untreated cells in tissue culture at 20 days

1 was similar to the level induced by 10 days of exposure to 0.5 mM DCA. The time course of  
2 TCA exposure was not tested to see if it had a similar effect with time as did DCA. The colonies  
3 observed at 10 days were tested for c-Jun expression with the authors noting that “colonies  
4 promoted by DCA were primarily c-Jun positive in contrast to TCA promoted colonies that were  
5 predominantly c-Jun negative.” Of the colonies that arose spontaneously from tissue culture  
6 conditions, 10/13 (76.9%) were reported to be c-Jun +, those treated with DCA 28/34 (82.3%)  
7 were c-Jun +, and those treated with TCA 5/22 (22.7%) were c-Jun +. Thus, these data show  
8 heterogeneity in cell in colonies but with more that were c-Jun + colonies occurring by tissue  
9 culture conditions alone than in the presence of DCA, rather than in the presence of TCA.

10 Bull et al. (2002) administered TCE, TCA, DCA, and combinations of TCA and DCA to  
11 male B6C3F1 mice by daily gavage (TCE) or drinking water (TCA, DCA, and TCA + DCA) for  
12 52–79 weeks, in order to compare a number of tumor characteristics, including c-Jun expression,  
13 across these different exposures. Bull et al. (2002) reported lesion reactivity to c-Jun antibody to  
14 be dependent on the proportion of the DCA and TCA administered after 52 weeks of exposure.  
15 Given alone, DCA was reported to produce lesions in mouse liver for which approximately half  
16 displayed a diffuse immunoreactivity to a c-Jun antibody, half did not, and none exhibited a  
17 mixture of the two. After TCA exposure alone, no lesions were reported to be stained with this  
18 antibody. When given in various combinations, DCA and TCA coexposure induced a few  
19 lesions that were only c-Jun+, many that were only c-Jun–, and a number with a mixed  
20 phenotype whose frequency increased with the dose of DCA. For TCE exposure of 79 weeks,  
21 TCE-induced lesions were reported to also have a mixture of phenotypes (42% c-Jun+, 34%  
22 c-Jun–, and 24% mixed) and to be most consistent with those resulting from DCA and TCA  
23 coexposure but not either metabolite alone.

24 A number of the limitations of the experiment are discussed in Caldwell et al. (2008).  
25 Specifically, for the DCA and TCA exposed animals, the experiment was limited by low  
26 statistical power, a relatively short duration of exposure, and uncertainty in reports of lesion  
27 prevalence and multiplicity due to inappropriate lesions grouping (i.e., grouping of hyperplastic  
28 nodules, adenomas, and carcinomas together as “tumors”), and incomplete histopathology  
29 determinations (i.e., random selection of gross lesions for histopathology examination). For  
30 determinations of immunoreactivity to c-Jun, Bull et al. (2002) combined hyperplastic nodules,  
31 adenomas, and carcinomas in most of their treatment groups, so differences in c-Jun expression  
32 across differing types of lesions were not discernable.

33 Nonetheless, these data collectively strongly suggest that TCA is not the sole agent of  
34 TCE-induced mouse liver tumors. In particular, TCE-induced tumors that were, in order of  
35 frequency, c-Jun+, c-Jun–, and of mixed phenotype, while c-Jun+ tumors have never been

1 observed with TCA treatment. Nor do these data support DCA as the sole contributor, since  
2 mixed phenotypes were not observed with DCA treatment.

#### 3 4.5.6.1.13.3. **Tumor genotype: H-ras mutation frequency and spectrum**

4 An approach to determine the potential MOAs of DCA and TCA through examination of  
5 the types of tumors each “induced” or “selected” was to examine H-ras activation (Anna et al.,  
6 1994; Bull et al., 2002; Ferreira-Gonzalez et al., 1995; Nelson et al., 1989). No data of this type  
7 were available for CH. This approach has also been used to try to establish an H-ras activation  
8 pattern for “genotoxic” and “nongenotoxic” liver carcinogens compounds and to make  
9 inferences concerning peroxisome proliferator-induced liver tumors. However, as noted by  
10 Stanley et al. (1994), the genetic background of the mice used and the dose of carcinogen may  
11 affect the number of activated H-ras containing tumors which develop. In addition, the stage of  
12 progression of “lesions” (i.e., foci vs. adenomas vs. carcinomas) also has been linked the  
13 observance of H-ras mutations. Fox et al. (1990) note that tumors induced by phenobarbital  
14 (0.05% drinking water [H<sub>2</sub>O], 1 year), chloroform (200 mg/kg corn oil gavage, two times weekly  
15 for 1 year) or ciprofibrate (0.0125% diet, 2 years) had a much lower frequency of H-ras gene  
16 activation than those that arose spontaneously (2-year bioassays of control animals) or induced  
17 with the “genotoxic” carcinogen benzidine-2 hydrochloric acid (HCl) (120 ppm, drinking H<sub>2</sub>O,  
18 1 year) in mice. In that study, the term “tumor” was not specifically defined but a correlation  
19 between the incidence of H-ras gene activation and development of either a hepatocellular  
20 adenoma or hepatocellular carcinoma was reported to be made with no statistically significant  
21 difference between the frequency of H-ras gene activation in the hepatocellular adenomas and  
22 carcinomas. Histopathological examination of the spontaneous tumors, tumors induced with  
23 benzidine-2 HCl, Phenobarbital, and chloroform was not reported to reveal any significant  
24 changes in morphology or staining characteristics. Spontaneous tumors were reported to have  
25 64% point mutation in codon 61 (*n* = 50 tumors examined) with a similar response for Benzidine  
26 of 59% (*n* = 22 tumors examined), whereas for Phenobarbital the mutation rate was  
27 7% (*n* = 15 tumors examined), chloroform 21% (*n* = 24 tumors examined) and ciprofibrate  
28 21% (*n* = 39 tumors examined). The ciprofibrate-induced tumors were reported to be more  
29 eosinophilic as were the surrounding normal hepatocytes.

30 Hegi et al. (1993) tested ciprofibrate-induced tumors in the NIH3T3 cotransfection-nude  
31 mouse tumorigenicity assay, which the authors state is capable of detecting a variety of activated  
32 protooncogenes. The tumors examined (ciprofibrate-induced or spontaneously arising) were  
33 taken from the Fox et al. study (1990), screened previously, and found to be negative for H-ras

1 activation. With the limited number of samples examined, Hegi et al concluded that ras  
2 protooncogene activation or activation of other protooncogenes using the nude mouse assay were  
3 not frequent events in ciprofibrate-induced tumors and that spontaneous tumors were not  
4 promoted with it. Using the more sensitive methods, the H-ras activation rate was reported to be  
5 raised from 21–31% for ciprofibrate-induced tumors and from 64–66% for spontaneous tumors.  
6 Stanley et al. (1994) studied the effect of methyleclofenapate (MCP) (25 mg/kg for up to 2 years),  
7 a peroxisome proliferator, in B6C3F1 (relatively sensitive) and C57BL/10J (relatively resistant)  
8 mice for H-ras codon 61 point mutations in MCP-induced liver tumors (hepatocellular adenomas  
9 and carcinomas). In the B6C3F1 mice the number of tumors with codon 61 mutations was 11/46  
10 and for C57BL/10J mice 4/31. Unlike the findings of Fox et al. (1990), Stanley et al. (1994)  
11 reported an increase in the frequency of mutation in carcinomas, which was reported to be twice  
12 that of adenomas in both strains of mice, indicating that stage of progression was related to the  
13 number of mutations in those tumors, although most tumors induced by MCP did not have this  
14 mutation.

15 Anna et al. (1994) reported that the H-ras codon 61 mutation frequency was not  
16 statistically different in liver tumors from DCA and TCE-treated mice from a highly variable  
17 number of tumors examined. From their concurrent controls, they reported that H-ras codon  
18 61 mutations in 17% ( $n = 6$ ) of adenomas and 100% ( $n = 5$ ) of carcinomas. For historical  
19 controls (published and unpublished), they reported mutations in 73% ( $n = 33$ ) of adenomas and  
20 mutations in 70% ( $n = 30$ ) of carcinomas. For tumors from TCE-treated animals, they reported  
21 mutations in 35% ( $n = 40$ ) of adenomas and 69% ( $n = 36$ ) of carcinomas, while for DCA-treated  
22 animals, they reported mutations in 54% ( $n = 24$ ) of adenomas and in 68% ( $n = 40$ ) of  
23 carcinomas. Anna et al. (1994) reported more mutations in TCE-induced carcinomas than  
24 adenomas. In regard to mutation spectra in H-ras oncogenes in control or spontaneous tumors,  
25 the patterns were slightly different but those from TCE treatment were mostly similar to that of  
26 DCA-induced tumors (0.5% in drinking water).

27 The study of Ferreira -Gonzalez (1995) in male B6C3F1 mice has the advantage of  
28 comparison of tumor phenotype at the same stage of progression (hepatocellular carcinoma), for  
29 allowance of the full expression of a tumor response (i.e., 104 weeks), and an adequate number  
30 of spontaneous control lesions for comparison with DCA or TCA treatments. However, tumor  
31 phenotype at an end stage of tumor progression may not be indicative of earlier stages of the  
32 disease process. In spontaneous liver carcinomas, 58% were reported to show mutations in H-61  
33 as compared with 50% of tumor from 3.5 g/L DCA-treated mice and 45% of tumors from  
34 4.5 g/L TCA-treated mice. A number of peroxisome proliferators have been reported to have a  
35 much smaller mutation frequency than spontaneous tumors (e.g., 13–24% H-ras codon 61

1 mutations after methylclofenopate depending on mouse strain, Stanely et al. [1994]: 21–31% for  
2 ciprofibrate-induced tumors and from 64–66% for spontaneous tumors, Fox et al. [1990] and  
3 Hegi et al [1993]). Thus, there was a heterogeneous response for this phenotypic marker for the  
4 spontaneous, DCA-, and TCA- treatment induced hepatocellular carcinomas had similar patterns  
5 H-ras mutations that differed from the reduced H-ras mutation frequencies reported for a number  
6 of peroxisome proliferators.

7 In his review, Bull (2000) suggested “the report by Anna et al. (1994) indicated that  
8 TCE-induced tumors possessed a different mutation spectra in codon 61 of the H-ras oncogene  
9 than those observed in spontaneous tumors of control mice.” Bull (2000) stated that “results of  
10 this type have been interpreted as suggesting that a chemical is acting by a mutagenic  
11 mechanism” but went on to suggest that it is not possible to a priori rule out a role for selection  
12 in this process and that differences in mutation frequency and spectra in this gene provide some  
13 insight into the relative contribution of different metabolites to TCE-induced liver tumors. Bull  
14 (2000) noted that data from Anna et al. (1994), Ferreira-Gonzalez et al. (1995), and Maronpot  
15 et al. (1995b) indicated that mutation frequency in DCA-induced tumors did not differ  
16 significantly from that observed in spontaneous tumors. Bull (2000) also noted that the mutation  
17 spectra found in DCA-induced tumors has a striking similarity to that observed in TCE-induced  
18 tumors, and DCA-induced tumors were significantly different than that of TCA-induced liver  
19 tumors.

20 Bull et al. (2002) reported that mutation frequency spectra for the H-ras codon 61 in  
21 mouse liver “tumors” induced by TCE ( $n = 37$  tumors examined) were reported to be  
22 significantly different than that for TCA ( $n = 41$  tumors examined), with DCA-treated mice  
23 tumors giving an intermediate result ( $n = 64$  tumors examined). In this experiment,  
24 TCA-induced “tumors” were reported to have more mutations in codon 61 (44%) than those  
25 from TCE (21%) and DCA (33%). This frequency of mutation in the H-ras codon 61 for TCA is  
26 the opposite pattern as that observed for a number of peroxisome proliferators in which the  
27 number of mutations at H-ras codon 61 in tumors has been reported to be much lower than  
28 spontaneously arising tumors (see above). Bull et al. (2002) noted that the mutation frequency  
29 for all TCE, TCA or DCA tumors was lower in this experiment than for spontaneous tumors  
30 reported in other studies (they had too few spontaneous tumors to analyze in this study), but that  
31 this study utilized lower doses and was of shorter duration than that of Ferreira-Gonzalez (1995).  
32 Furthermore, the disparities from previous studies may also be impacted by lesion grouping,  
33 mentioned above, in which lower stages of progression are grouped with more advanced stages.

34 Overall, in terms of H-ras mutation, TCE-induced tumors appears to be more like  
35 DCA-induced tumors (which are consistent with spontaneous tumors), or those resulting from a



1 coexposure to both DCA and TCA (Bull et al., 2002), than from those induced by TCA. As  
2 noted above, Bull et al. (2002) reported the mutation frequency spectra for the H-ras codon 61 in  
3 mouse liver tumors induced by TCE to be significantly different than that for TCA, with  
4 DCA-treated mice tumors giving an intermediate result and for TCA-induced tumors to have a  
5 H-ras profile that is the opposite than those of a number of other peroxisome proliferators. More  
6 importantly, however, these data, along with the measures discussed above, show that mouse  
7 liver tumors induced by TCE are heterogeneous in phenotype and genotype in a manner similar  
8 to that observed in spontaneous tumors.  
9

#### 4.5.6.1.14. “Stop” experiments

10 Several stop experiments, in which treatment is terminated early in some dose groups,  
11 have attempted to ascertain the whether progression differences exist between TCA and DCA.  
12 After 37 weeks of treatment and then a cessation of exposure for 15 weeks, Bull et al. (1990)  
13 reported that after combined 52 week period, liver weight and percentage liver/body weight were  
14 reported to still be statistically significantly elevated after DCA or TCA treatment. The authors  
15 partially attribute the remaining increases in liver weight to the continued presence of  
16 hyperplastic nodules in the liver. In terms of liver tumor induction, the authors stated that  
17 “statistical analysis of tumor incidence employed a general linear model ANOVA with contrasts  
18 for linearity and deviations from linearity to determine if results from groups in which treatments  
19 were discontinued after 37 weeks were lower than would have been predicted by the total dose  
20 consumed.” The multiplicity of tumors (incidence was not used) observed in male mice exposed  
21 to DCA or TCA at 37 weeks and then sacrificed at 52 weeks were compared with those exposed  
22 for a full 52 weeks. The response in animals that received the shorter duration of DCA exposure  
23 was very close to that which would be predicted from the total dose consumed by these animals.  
24 By contrast, the response to TCA exposure for the shorter duration was reported by the authors  
25 to deviate significantly ( $p = 0.022$ ) from the linear model predicted by the total dose consumed.  
26 However, in the prediction of “dose-response,” foci, adenomas, and carcinomas were combined  
27 into one measure. Therefore, foci, a certain percentage of which have been commonly shown to  
28 spontaneously regress with time, were included in the calculation of total “lesions.” Moreover,  
29 only a sample of lesions were selected for histological examination, and as is evident in the  
30 sample, some lesions appeared “normal” upon microscopic examination (see below). Therefore,  
31 while suggesting that cessation of exposure diminished the number of “lesions,” methodological  
32 limitations temper any conclusions regarding the identity and progression of lesion with  
33 continuous versus noncontinuous DCA and TCA treatment.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1           Additionally, Bull et al. (1990) noted that after stopping treatment, DCA lesions appeared  
2 to arrest their progression in contrast to TCA lesions, which appeared to progress. In particular,  
3 among those in the stop treatment group (at 2 g/L) with 0/19 lesions examined histologically  
4 were carcinomas, while in the continuous treatment groups, a significant fraction of lesions  
5 examined were carcinomas at the higher exposure (6/23 at 2 g/L). By contrast, at terminal  
6 sacrifice, TCA lesions a larger fraction of the lesions examined were carcinomas in the stop  
7 treatment group (3/5 at 2 g/L) than in the continuous treatment group (2/7 and 4/16 at 1 g/L and  
8 2 g/L, respectively).

9           However, as mentioned above, these inferences are based on examination of only a  
10 subset of lesions. Specifically, for TCA treatment the number of animals examined for  
11 determination of which “lesions” were foci, adenomas, and carcinomas was 11 out of the  
12 19 mice with “lesions” at 52 weeks while all four mice with lesions after 37 weeks of exposure  
13 and 15 weeks of cessation were examined. For DCA treatment the number of animals examined  
14 was only 10 out of 23 mice with “lesions” at 52 weeks while all 7 mice with lesions after 37  
15 weeks of exposure and 15 weeks of cessation were examined. Most importantly, when lesions  
16 were examined microscopically, some did not all turn out to be preneoplastic or neoplastic—for  
17 example, two lesions appeared “to be histologically normal” and one necrotic.

18           While limited, the conclusions of Bull et al. (1990) are consistent with later experiments  
19 performed by Pereira and Phelps (1996). They noted that in MNU-treated mice that were then  
20 treated with DCA, the yield of altered hepatocytes decreases as the tumor yields increase  
21 between 31 and 51 weeks of exposure suggesting progression of foci to adenomas, but that  
22 adenomas did not appear to progress to carcinomas. For TCA, Pereira and Phelps (1996)  
23 reported that “MNU-initiated” adenomas promoted with TCA continued to progress. However,  
24 the use of MNU initiation complicates direct comparisons with treatment with TCA or DCA  
25 alone.

26           No similar data comparing stop and continued treatment of TCE are available to assess  
27 the consistency or lack-thereof with TCA or DCA. Moreover, the informative of such a  
28 comparison would be limited by designs of the available TCA and DCA studies, which have  
29 used higher concentrations in conjunction with the much lower durations of exposure. While  
30 higher doses allow for responses to be more easily detected, it introduces uncertainty as to the  
31 effects of the higher doses alone. In addition, because the overall duration of the experiments is  
32 also generally much less than 104 weeks, it is not possible to discern whether the differences in  
33 results between those animals in which treatment was suspended in comparison to those in which  
34 had not had been conducted would persist with longer durations.

#### 4.5.6.1.15. Conclusions Regarding the Role of Trichloroacetic Acid (TCA), Dichloroacetic Acid (DCA), and Chloral Hydrate (CH) in Trichloroethylene (TCE)-Induced Effects in the Liver

1           In summary, it is likely that oxidative metabolism is necessary for TCE-induced effects in  
2 the liver. However, the specific metabolite or metabolites responsible for both noncancer and  
3 cancer effects is less clear. TCE, TCA, and DCA exposures have all been associated with  
4 induction of peroxisomal enzymes but are all weak PPAR $\alpha$  agonists. The available data strongly  
5 support TCA *not* being the sole or predominant active moiety for TCE-induced liver effects.  
6 With respect to hepatomegaly, TCE and TCA dose-response relationships are quantitatively  
7 inconsistent, for TCE leads to greater increases in liver/body weight ratios than expected from  
8 predicted rates of TCA production. In fact, above a certain dose of TCE, liver/body weight  
9 ratios are greater than that observed under any conditions studied so far for TCA. Histological  
10 changes and effects on DNA synthesis are generally consistent with contributions from either  
11 TCA or DCA, with a degree of polyploidization, rather than cell proliferation, likely to be  
12 significant for TCE, TCA, and DCA. With respect to liver tumor induction, TCE leads to a  
13 heterogeneous population of tumors, not unlike those that occur spontaneously or that are  
14 observed following TCA-, DCA-, or CH-treatment. Moreover, some liver phenotype  
15 experiments, particularly those utilizing immunostaining for c-Jun, support a role for both DCA  
16 and TCA in TCE-induced tumors, with strong evidence that TCA cannot solely account for the  
17 characteristics of TCE-induced tumors. In addition, H-ras mutation frequency and spectrum of  
18 TCE-induced tumors more closely resembles that of spontaneous tumors or of those induced by  
19 DCA, and were less similar in comparison to that of TCA-induced tumors. The heterogeneity of  
20 TCE-induced tumors is similar to that observed to be induced by a broad category of  
21 carcinogens, and to that observed in human liver cancer. Overall, then, it is likely that multiple  
22 TCE metabolites, and therefore, multiple pathways, contribute to TCE-induced liver tumors.

#### 4.5.7. Mode of Action (MOA) for Trichloroethylene (TCE) Liver Carcinogenicity

23           This section will discuss the evidentiary support for several hypothesized modes of action  
24 for liver carcinogenicity (including mutagenicity and peroxisome proliferation, as well as several  
25 additional proposed hypotheses and key events with limited evidence or inadequate experimental  
26 support), following the framework outlined in the *Cancer Guidelines* (U.S. EPA, 2005c, d).<sup>9</sup>

---

<sup>9</sup> As recently reviewed (Guyton et al., 2008) the approach to evaluating mode of action information described in EPA's *Cancer Guidelines* (U.S. EPA, 2005c, d) considers the issue of human relevance of a hypothesized mode of action in the context of hazard evaluation. This excludes, for example, consideration of toxicokinetic differences across species; specifically, the Cancer Guidelines state, "the toxicokinetic processes that lead to formation or distribution of the active agent to the target tissue are considered in estimating dose but are not part of the mode of action." In addition, information suggesting quantitative differences in the occurrence of a key event between test

*This document is a draft for review purposes only and does not constitute Agency policy.*

1

#### 4.5.7.1.1. Mutagenicity

2           The hypothesis is that TCE acts by a mutagenic mode of action in TCE-induced  
3 hepatocarcinogenesis. According to this hypothesis, the key events leading to TCE-induced liver  
4 tumor formation constitute the following: TCE oxidative metabolite CH, after being produced in  
5 the liver, cause direct alterations to DNA (e.g., mutation, DNA damage, and/or micronuclei  
6 induction). Mutagenicity is a well established cause of carcinogenicity.  
7

#### 4.5.7.1.2. Experimental support for the hypothesized mode of action

8           The genotoxicity, as described by the ability of TCE, CH, TCA, and DCA to induce  
9 mutations, was discussed previously in Section 4.2. The strongest data for mutagenic potential  
10 are for CH, thought to be a relatively short-lived intermediate in the metabolism of TCE that is  
11 rapidly converted to TCA and TCOH in the liver (see Section 3.3). CH causes a variety of  
12 genotoxic effects in available in vitro and in vivo assays, with particularly strong data as to its  
13 ability to induce aneuploidy. It has been argued that CH mutagenicity is unlikely to be the cause  
14 of TCE carcinogenicity because the concentrations required to elicit these responses are  
15 generally quite high, several orders of magnitude higher than achieved in vivo (Moore and  
16 Harrington-Brock, 2000). For example, peak concentrations of CH in the liver of around  
17 2–3 mg/kg have been reported after TCE administration at doses that are hepatocarcinogenic in  
18 chronic bioassays (Abbas and Fisher, 1997; Greenberg et al., 1999). Assuming a liver density of  
19 about 1 kg/L, these concentrations are orders of magnitude less than the minimum concentrations  
20 reported to elicit genotoxic responses in the Ames test and various in vitro measures of  
21 micronucleus, aneuploidy, and chromosome aberrations, which are in the 100–1,000 mg/L range.  
22 However, it is not clear how much of a correspondence is to be expected from concentrations in  
23 genotoxicity assays in vitro and concentrations in vivo, as reported in vivo CH concentrations are  
24 in whole-liver homogenate while in vitro concentrations are in culture media. In addition, a few  
25 in vitro studies have reported positive results at concentrations as low as 1 or 10 mg/L, including  
26 Furnus et al. (1990) for aneuploidy in Chinese hamster CHED cells (10 mg/L), Eichenlaub-Ritter  
27 et al. (1996) for bivalent chromosomes in meiosis I in MF1 mouse oocytes (10 mg/L), and  
28 Gibson et al. (1995) for cell transformation in Syrian hamster embryo cells after 7 day treatment.  
29 Moreover, some in vivo genotoxicity assays of CH reported positive results at doses similar to

---

species and humans are noted for consideration in the dose-response assessment, but is not considered in human

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 those eliciting a carcinogenic response in chronic bioassays. For example, Nelson and Bull  
2 (1988) reported increased DNA single strand breaks at 100 CH mg/kg (oral) in male B6C3F1  
3 mice, although the result was not replicated by Chang et al. (1992). In another example, four of  
4 six in vivo mouse genotoxicity studies reported that CH induced micronuclei in mouse  
5 bone-marrow erythrocytes, with the lowest effective doses in positive studies ranging from  
6 83–500 mg/kg (positive: Russo and Levis (1992), Russo et al. (1992), Marrazzini et al. (1994),  
7 Beland et al. (1999); negative: Leuschner and Leuschner (1991), Leopardi et al. (1993)).  
8 However, the use of i.p. administration in these and many other in vivo genotoxicity assays  
9 complicates the comparison with carcinogenicity data. Also, it is difficult with the available data  
10 to assess the contributions from the genotoxic effects of CH along with those from the genotoxic  
11 and nongenotoxic effects of other oxidative metabolites (discussed below in Sections 4.5.5.2 and  
12 4.5.5.3).

13 Furthermore, altered DNA methylation, another heritable mechanism by which gene  
14 expression may be altered, is discussed below in the in Section 4.5.1.3.2.6. As discussed  
15 previously, the differential patterns of H-ras mutations observed in liver tumors induced by TCE,  
16 TCA, and DCA may be more indicative of tumor selection and tumor progression resulting from  
17 exposure to these agents rather than a particular mechanism of tumor induction. The state of the  
18 science of cancer and the role of epigenetic changes, in addition to genetic changes, in the  
19 initiation and progression of cancer and specifically liver cancer, are discussed in Section E.3.1.

20 Therefore, while data are insufficient to conclude that a mutagenic MOA mediated by CH  
21 is operant, a mutagenic MOA, mediated either by CH or by some other oxidative metabolite of  
22 TCE, cannot be ruled out.  
23

#### 4.5.7.1.3. Peroxisome Proliferator Activated Receptor Alpha (PPAR $\alpha$ ) Receptor Activation

24 The hypothesis is that TCE acts by a PPAR $\alpha$  agonism MOA in TCE-induced  
25 hepatocarcinogenesis. According to this hypothesis, the key events leading to TCE-induced liver  
26 tumor formation constitute the following: the TCE oxidative metabolite TCA, after being  
27 produced in the liver, activates the PPAR $\alpha$  receptor, which then causes alterations in cell  
28 proliferation and apoptosis and clonal expansion of initiated cells. This MOA is assumed to  
29 apply only to the liver.  
30

---

relevance determination.

#### 4.5.7.1.4. Experimental support for the hypothesized mode of action

1 Proliferation of peroxisomes and increased activity of a number of related marker  
2 enzymes has been observed in rodents treated with TCE, TCA, and DCA. The  
3 peroxisome-related effects of TCE are most likely mediated primarily through TCA based on  
4 TCE metabolism producing more TCA than DCA and the lower doses of TCA required to elicit  
5 a response relative to DCA. However, Bull (2004a) and Bull et al. (2004) have recently  
6 suggested that peroxisome proliferation occurs at higher exposure levels than those that induce  
7 liver tumors for TCE and its metabolites. They report that a direct comparison in the no-effect  
8 level or low-effect level for induction of liver tumors in the mouse and several other endpoints  
9 shows that, for TCA, liver tumors occur at lower concentrations than peroxisome proliferation in  
10 vivo but that PPAR $\alpha$  activation occurs at a lower dose than either tumor formation or peroxisome  
11 proliferation. A similar comparison for DCA shows that liver tumor formation occurs at a much  
12 lower exposure level than peroxisome proliferation or PPAR $\alpha$  activation. In vitro transactivation  
13 studies have shown that human and murine versions of PPAR $\alpha$  are activated by TCA and DCA,  
14 while TCE itself is relatively inactive in the in vitro system, at least with mouse PPAR $\alpha$   
15 (Maloney and Waxman, 1999; Zhou and Waxman, 1998). In addition, Laughter et al. (2004)  
16 reported that the responses of acyl CoA oxidase (ACO), PCO, and CYP4A induction by TCE,  
17 TCA, and DCA were substantially diminished in PPAR $\alpha$ -null mice. Therefore, evidence  
18 suggests that TCE, through its metabolites TCA and DCA, activate PPAR $\alpha$ , and that at doses  
19 relevant to TCE-induced hepatocarcinogenesis, the role of TCA in PPAR $\alpha$  agonism is likely to  
20 predominate.

21 It has been suggested that PPAR $\alpha$  receptor activation is both the MOA for TCA liver  
22 tumor induction as well as the MOA for TCE liver tumor induction, as a result of the metabolism  
23 of TCE to TCA (Corton, 2008; NRC, 2006). Section E.3.4 addressed the status of the PPAR $\alpha$   
24 MOA hypothesis for liver tumor induction and provides a more detailed discussion. However, as  
25 discussed previously and in Section E.2.1.10, TCE-induced increases in liver weight have been  
26 reported in male and female mice that do not have a functional PPAR $\alpha$  receptor (Nakajima et al.,  
27 2000). The dose-response for TCE-induced liver weight increases differs from that of TCA (see  
28 Section E.2.4.2). The phenotype of the tumors induced by TCE have been described to differ  
29 from those by TCA and to be more like those occurring spontaneously in mice, those induced by  
30 DCA, or those resulting from a combination of exposures to both DCA and TCA (see  
31 Section E.2.4.4). As to whether TCA induces tumors through activation of the PPAR $\alpha$  receptor,  
32 the tumor phenotype of TCA-induced mouse liver tumors has been reported to have a different  
33 pattern of H-ras mutation frequency from other peroxisome proliferators (see Section E.2.4.4,  
34 Bull et al., 2002; Fox et al., 1990; Hegi et al., 1993; Stanley et al., 1994). While TCE, DCA, and

1 TCA are weak peroxisome proliferators, liver weight induction from exposure to these agents  
2 has not correlated with increases in peroxisomal enzyme activity (e.g., PCO activity) or changes  
3 in peroxisomal number or volume. By contrast, as discussed above, liver weight induction from  
4 subchronic exposures appears to be a more accurate predictor of carcinogenic response for DCA,  
5 TCA and TCE in mice (see also Section E.2.4.4). The database for cancer induction in rats is  
6 much more limited than that of mice for determination of a carcinogenic response to these  
7 chemicals in the liver and the nature of such a response.

8 While many compounds known to cause rodent liver tumors with long-term treatment  
9 also activate the nuclear receptor PPAR $\alpha$ , the mechanisms by which PPAR $\alpha$  activation  
10 contributes to tumorigenesis are not completely known (Klaunig et al., 2003; NRC, 2006; Yang  
11 et al., 2007). As reviewed by Keshava and Caldwell (2006), PPAR $\alpha$  activation leads to a highly  
12 pleiotropic response and may play a role in toxicity in multiple organs as well as in multiple  
13 chronic conditions besides cancer (obesity, atherosclerosis, diabetes, inflammation). Klaunig  
14 et al. (2003) and NRC (2006) proposed that the key causal events for PPAR $\alpha$  agonist-induced  
15 liver carcinogenesis, after PPAR $\alpha$  activation, are perturbation of cell proliferation and/or  
16 apoptosis, mediated by gene expression changes, and selective clonal expansion. It has also been  
17 proposed that sufficient evidence for this MOA consists of evidence of PPAR $\alpha$  agonism (i.e., in  
18 a receptor assay) in combination with either light- or electron-microscopic evidence for  
19 peroxisome proliferation or both increased liver weight and one more of the in vivo markers of  
20 peroxisome proliferation (Klaunig et al., 2003). However, it should be noted that peroxisome  
21 proliferation and in vivo markers such as PCO are not considered causal events (Klaunig et al.,  
22 2003; NRC, 2006), and that their correlation with carcinogenic potency is poor (Marsman et al.,  
23 1988). Therefore, for the purposes of this discussion, peroxisome proliferation and its markers  
24 are considered indicators of PPAR $\alpha$  activation, as it is well established that these highly specific  
25 effects are mediated through PPAR $\alpha$  (Klaunig et al., 2003; Peters et al., 1997).

26 As recently reviewed by Guyton et al. (2009), recent data suggest that PPAR $\alpha$  activation  
27 along with these hypothesized causal events may not be sufficient for carcinogenesis. In  
28 particular, Yang et al. (2007) reported comparisons between mice treated with Wy-14643 and  
29 transgenic mice in which PPAR $\alpha$  was constitutively activated in hepatocytes without the  
30 presence of ligand. Yang et al. (2007) reported that, in contrast to Wy-14643-treatment, the  
31 transgene did not induce liver tumors at 11 months, despite inducing PPAR $\alpha$ -mediated effects of  
32 a similar type and magnitude seen in response to tumorigenic doses of Wy-14643 in wild-type  
33 mice (decreased serum fatty acids, induction of PPAR $\alpha$  target genes, altered expression of  
34 cell-cycle control genes, and a sustained increase in cellular proliferation). Nonetheless, it is  
35 important to discuss the extent to which PPAR $\alpha$  activation mediates the effects proposed by

1 Klaunig et al. (2003) and NRC (2006), even if the hypothesized sequence of key events may not  
2 be sufficient for carcinogenesis. Investigation continues into additional events that may also  
3 contribute, such as nonparenchymal cell activation and micro-RNA-based regulation of  
4 protooncogenes (Shah et al., 2007; Yang et al., 2007). Specifically addressed below are gene  
5 expression changes, proliferation, clonal expansion, and mutation frequency or spectrum.

6 With respect to gene expression changes due to TCE, Laughter et al. (2004) evaluated  
7 transcript profiles induced by TCE in wild-type and PPAR $\alpha$ -null mice. As noted in  
8 Sections E.3.4.1.3 and E.3.1.2, there are limitations to the interpretation of such studies, some of  
9 which are discussed below. Also noted in Appendix E are discussions of how studies of  
10 peroxisome proliferators, indicate of the need for phenotypic anchoring, especially since gene  
11 expression is highly variable between studies and within studies using the same experimental  
12 paradigm. Section E.3.4 in also provides detailed discussions of the status of the PPAR $\alpha$   
13 hypothesis. Of note, all null mice at the highest TCE dose (1,500 mg/kg-day) were moribund  
14 prior to the end of the planned 3-week experiment (Laughter et al., 2004), and it was proposed  
15 that this may reflect a greater sensitivity in PPAR $\alpha$ -null mice to hepatotoxins due to defects in  
16 tissue repair abilities. Laughter et al. (2004) also noted that four genes known to be regulated by  
17 other peroxisome proliferators also had altered expression with TCE treatment in wild-type, but  
18 not null mice. Ramdhan et al. (2010) report that not only do PPAR $\alpha$ -null mice, but also  
19 humanized mice (PPAR $\alpha$ -null mice with inserted human PPAR $\alpha$ ) have underlying dysregulation  
20 of lipid metabolism and gene expression. However, in a comparative analysis, Bartosiewicz  
21 et al. (2001) concluded that TCE induced a different pattern of transcription than two other  
22 peroxisome proliferators, di(2-ethylhexyl) phthalate (DEHP) and clofibrate. In addition,  
23 Keshava and Caldwell (2006) compared gene expression data from Wy-14643, dibutyl phthalate  
24 (DBP), gemfibrozil (GEM), and DEHP, and noted a lack of consistent results across PPAR $\alpha$   
25 agonists. Thus, available data are insufficient to conclude that TCE gene expression changes are  
26 similar to other PPAR agonists, or even that there are consistent changes (beyond the in vivo  
27 markers of peroxisome proliferation, such as ACO, PCO, CYP4A, etc.) among different  
28 agonists. It should also be noted that Laughter et al. (2004) did not compare baseline (i.e.,  
29 control levels of) gene expression between null and wild-type control mice, hindering  
30 interpretation of these results (Keshava and Caldwell, 2006). The possible relationship between  
31 PPAR $\alpha$  activation and hypomethylation are discussed below in Section 4.5.7.1.9.

32 In terms of proliferation, mitosis itself has not been examined in PPAR $\alpha$ -null mice, but  
33 BrdU incorporation, a measure of DNA synthesis that may reflect cell division, polyploidization,  
34 or DNA repair, was observed to be diminished in null mice as compared to wild-type mice at 500  
35 and 1,000 mg/kg-day TCE (Laughter et al., 2004). However, BrdU incorporation in null mice



1 was still about threefold higher than controls, although it was not statistically significantly  
2 different due to the small number of animals, high variability, and the two- to threefold higher  
3 baseline levels of BrdU incorporation in control null mice as compared to control wild-type  
4 mice. Therefore, while PPAR $\alpha$  appears to contribute to the short-term increase in DNA  
5 synthesis observed with TCE treatment, these results cannot rule out other contributing  
6 mechanisms. However, since it is likely that both cellular proliferation and increased ploidy  
7 contribute to the observed TCE-induced increases in DNA synthesis, it is not clear to whether the  
8 observed decrease in BrdU incorporation is due to reduced proliferation, reduced  
9 polyploidization, or both.

10 With respect to clonal expansion, it has been suggested that tumor characteristics such as  
11 tincture (i.e., the staining characteristics light microscopy sections of tumor using H&E stains)  
12 and oncogene mutation status can be used to associate chemical carcinogens with a particular  
13 MOA such as PPAR $\alpha$  agonism (Klaunig et al., 2003; NRC, 2006). This approach is problematic  
14 primarily because of the lack of specificity of these measures. For example, with respect to  
15 tincture, it has been suggested that TCA-induced foci and tumors resemble those of other  
16 peroxisome proliferators in basophilia and lack of expression of GGT and GST-pi. However, as  
17 discussed in Caldwell and Keshava (2006), the term “basophilic” in describing foci and tumors  
18 can be misleading, because, for example, multiple lineages of foci and tumors exhibit basophilia,  
19 including those not associated with peroxisome proliferators (Bannasch, 1996; Bannasch et al.,  
20 2001; Carter et al., 2003). Moreover, a number of studies indicate that foci and tumors induced  
21 by other “classic” peroxisome proliferators may have different phenotypic characteristics from  
22 that attributed to the class through studies of WY-14643, including DEHP (Voss et al., 2005) and  
23 clofibric acid (Michel et al., 2007). Furthermore, even the combination of GGT and GST-pi  
24 negative, basophilic foci are nonspecific to peroxisome proliferators, as they have been observed  
25 in rats treated with AFB1 and AFB1 plus PB, none of which are peroxisome proliferators (Grasl-  
26 Kraupp et al., 1993)(Kraupp-Grasl et al., 1990). Finally, while Bull et al. (2004) suggested that  
27 negative expression of *c-jun* in TCA-induced tumors may be consistent with a characteristic  
28 phenotype of peroxisome proliferators, no data could be located to support this statement.  
29 Therefore, of phenotypic information does not appear to be reliable for associating a chemical  
30 with a PPAR $\alpha$  agonism MOA.

31 Mutation frequency or spectrum in oncogenes has also been suggested to be an indicator  
32 of a PPAR $\alpha$  agonism MOA being active (NRC, 2006), with the idea being that specific  
33 genotypes are being promoted by PPAR $\alpha$  agonists. Although not a highly specific marker, *H-ras*  
34 codon 61 mutation frequency and spectra data do not support a similarity between mutations in  
35 TCE-induced, TCA-, or DCA- tumors and those due to other peroxisome proliferators. For

1 example, while ciprofibrate and methylclofenopate had lower mutation frequencies than  
2 historical controls (Hegi et al., 1993; Stanley et al., 1994), TCA-induced tumors had mutation  
3 frequencies similar to or higher than historical controls (Bull et al., 2002; Ferreira-Gonzalez et  
4 al., 1995). Anna et al. (1994) and Ferreira-Gonzalez et al. (1995) also reported TCE and  
5 DCA-induced tumors to have mutation frequencies similar to historical controls, although Bull  
6 et al. (2002) reported lower frequencies for these chemicals. However, the data reported by Bull  
7 et al. (2002) consist of mixed lesions at different stages of progression, and such differing stages,  
8 in addition to differences in genetic background and dose, can influence the frequency of *H-ras*  
9 mutations (Stanley et al., 1994). In addition, a greater frequency of mutations was reported in  
10 carcinomas than adenomas, and Bull et al. (2002) stated that this suggested that *H-ras* mutations  
11 were a late event. Moreover, Fox et al. (1990) noted that tumors induced by phenobarbital,  
12 chloroform, and ciprofibrate all had a much lower frequency of *H-ras* gene activation than those  
13 that arose spontaneously, so this marker does not have good specificity. Mutation spectrum is  
14 similarly of low utility for supporting a PPAR $\alpha$  agonism MOA. First, because many peroxisome  
15 proliferators been reported to have low frequency of mutations, the comparison of mutation  
16 spectrum would be limited to a small fraction tumors. In addition to the low power due to small  
17 numbers, the mutation spectrum is relatively nonspecific, as Fox et al. (1990) reported that of the  
18 tumors with mutations, the spectra of the peroxisome proliferator ciprofibrate, historical controls,  
19 and the genotoxic carcinogen benzidine-2 HCl were similar.

20 In summary, TCE clearly activates PPAR $\alpha$ , and some of the effects contributing to  
21 tumorigenesis that Klaunig et al. (2003) and NRC (2006) propose to be the result of PPAR $\alpha$   
22 agonism are observed with TCE, TCA, or DCA treatment. While this consistency is supportive a  
23 role for PPAR $\alpha$ , all of the proposed key causal effects with the exception of PPAR $\alpha$  agonism  
24 itself are nonspecific, and may be caused by multiple mechanisms. There is more direct  
25 evidence that several of these effects, including alterations in gene expression and changes in  
26 DNA synthesis, are mediated by multiple mechanisms in the case of TCE, and a causal linkage  
27 to PPAR $\alpha$  specifically is lacking. Therefore, because, as discussed further in the MOA  
28 discussion below, there are multiple lines of evidence supporting the role of multiple pathways  
29 of TCE-induced tumorigenesis, the hypothesis that PPAR $\alpha$  agonism and the key causal events  
30 proposed by Klaunig et al. (2003) and NRC (2006) constitute the sole or predominant MOA for  
31 TCE-induced carcinogenesis is considered unlikely.

32 Furthermore, as reviewed by Guyton et al. (2009), recent data strongly suggest that  
33 PPAR $\alpha$  and key events hypothesized by Klaunig et al. (2003) are not sufficient for  
34 carcinogenesis induced by the purported prototypical agonist Wy-14643. Therefore, the

1 proposed PPAR $\alpha$  MOA is likely “incomplete” in the sense that the sequence of key events<sup>10</sup>  
2 necessary for cancer induction has not been identified. A recent 2-year bioassay of the  
3 peroxisome proliferator DEHP showed that it can induce a liver tumor response in mice lacking  
4 PPAR $\alpha$  similar to that in wild-type mice (Ito et al., 2007). Klaunig et al. (2003) previously  
5 concluded that PPAR $\alpha$  agonism was the sole MOA for DEHP-induced liver tumorigenesis based  
6 on the lack of tumors in PPAR $\alpha$ -null mice after 11 months treatment with Wy-14643 (Peters et  
7 al., 1997). They also assumed that due to the lack of markers of PPAR $\alpha$  agonism in PPAR $\alpha$ -null  
8 mice after short-term treatment with DEHP (Ward et al., 1998), a long-term study of DEHP in  
9 PPAR $\alpha$ -null mice would yield the same results as for Wy-14643. However, due the finding by  
10 Ito et al. (2007) that PPAR $\alpha$ -null mice exposed to DEHP do develop liver tumors, they  
11 concluded that DEHP can induce liver tumors by multiple mechanisms (Ito et al., 2007;  
12 Takashima et al., 2008). Hence, since there is no 2-year bioassay in PPAR $\alpha$ -null mice exposed  
13 to TCE or its metabolites, it is not justifiable to use a similar argument based on Peters et al.  
14 (1997) and short-term experiments to suggest that the PPAR $\alpha$  MOA is operative. Therefore, the  
15 conclusion is supported that the hypothesized PPAR $\alpha$  MOA is inadequately specified because  
16 the data do not adequately show the proposed key events individually being required for  
17 hepatocarcinogenesis, nor do they show the sequence of key events collectively to be sufficient  
18 for hepatocarcinogenesis.  
19

#### 4.5.7.1.5. Quantitative relationships between key events and tumor induction

20 The issues of whether there is a quantitative relationship between hypothesized key  
21 events and tumor induction were recently examined in Guyton et al. (2009) and are discussed  
22 below. Furthermore, IARC has recently concluded that additional mechanistic information has  
23 become available, including studies with DEHP in PPAR $\alpha$ -null mice, studies with several  
24 transgenic mouse strains, carrying human *PPARalpha* or with hepatocyte-specific constitutively  
25 activated PPAR $\alpha$  and a study in humans exposed to DEHP from the environment that has  
26 changed its conclusions regarding the relevance of rodent tumor data to human risk (Grosse et  
27 al., 2011). Data from these new studies suggest that many molecular signals and pathways in  
28 several cell types in the liver, rather than a single molecular event, contribute to cancer

---

10 As defined by the EPA *Cancer Guidelines* (2005a, b) a “key event” is “an empirically observable precursor step that is itself a necessary element of the mode of action or is a biologically based marker for such an element,” and the term “mode of action” (MOA) is defined as “a sequence of key events and processes, starting with interaction of an agent with a cell, proceeding through operational and anatomical changes, and resulting in cancer formation.” Therefore, a single key event alone is necessary, but not necessarily sufficient for carcinogenesis; however, the *sequence* of key events constituting a MOA needs to be sufficient for carcinogenesis.

1 development in rodents with IARC concluding that the human relevance of the molecular events  
2 leading to DEHP-induced cancer in several target tissues (e.g., liver and testis) in rats or mice  
3 could not be ruled out, resulting in the evaluation of DEHP as a Group-2B agent, rather than  
4 Group 3.

5 This following discussion is from Guyton et al. (2009):

6  
7  
8 **Are key or associative events in the PPAR- $\alpha$  activation MOA quantitatively predictive of**  
9 **hepatocarcinogenicity?**

10 Another question to consider is whether potency for PPAR- $\alpha$  activation or its attendant  
11 sequelae is quantitatively associated with carcinogenic activity or potency. If so, differences in  
12 sensitivity for carcinogenesis (such as may occur across species) could be predicted using  
13 quantitative information about the key events alone. If robust correlations were established, then  
14 they could potentially be used either to quantitatively account for pharmacodynamic differences  
15 that impact carcinogenic potency or as precursor events in nonlinear dose response assessment.  
16 However, there are limitations in the dose-response data available for analyses of quantitative  
17 relationships between potencies for precursor events in the proposed PPAR- $\alpha$  activation MOA and  
18 for liver tumor induction. Most tumor data, including for the best characterized PPAR- $\alpha$  agonists,  
19 are for exposure concentrations inducing well above 50% tumor incidence with less-than-lifetime  
20 administration. Precursor events have typically been studied at a single dose, often eliciting a near  
21 maximal response, thus precluding benchmark-based comparisons across studies. This is  
22 especially true for Wy-14,643, which has been administered most often at only one exposure  
23 concentration (1,000 ppm) that elicits a 100% tumor incidence after 1 year or less (Peters et al.,  
24 1997) and that also appears to be necrogenic (Woods et al., 2007a). On the other hand,  
25 hypothesized precursor events such as hepatomegaly, peroxisome proliferation, and increased  
26 DNA synthesis appear to have reached their maximal responses at 50 ppm Wy-14,643, with some  
27 statistically significant responses as low as 5 ppm (Marsman et al., 1992; Wada et al., 1992).  
28 Potencies across compounds have rarely been compared in a single study using the same  
29 experimental paradigm. These deficits in the database notwithstanding, provided below is an  
30 assessment of the quantitative predictive power of the potency for four proposed data elements for  
31 establishing the hypothesized MOA for hepatocarcinogenesis: PPAR- $\alpha$  activation in mice; and  
32 hepatomegaly, DNA synthesis, and increased peroxisome proliferation in rats.

33  
34 **PPAR-  $\alpha$  activation in mice**

35 Table 2 [reproduced as Table 4-66] presents data for four peroxisome proliferators in order of  
36 decreasing potency for inducing mouse liver tumors. These compounds were selected because of  
37 their importance to environmental human health risk assessments and because data to derive  
38 receptor activation potency indicators were available from a single study (Maloney and Waxman,  
39 1999). The transactivation potencies of MEHP, Wy-14,643, dichloroacetic acid (DCA), and TCA  
40 for the mouse PPAR- $\alpha$  were monitored using a luciferase reporter gene containing multiple PPAR  
41 response elements derived from the rat hydratase/dehydrogenase promoter in transiently transfected  
42 COS-1 monkey kidney cells. The derived potency indicators were compared to the TD<sub>50</sub> (i.e., the  
43 daily dose inducing tumors in half of the mice that would otherwise have remained tumor-free)  
44 from the Carcinogenic Potency Database (CPDB) of Gold et al. (2005). Note that for Wy-14,643,  
45 the dose listed yielded a maximal response and thus represents an upper limit to the TD<sub>50</sub>  
46 (indicated by “<”). Two estimates of PPAR- $\alpha$  transactivation potency are given, the first based on  
47 50% of the maximal response (i.e., EC<sub>50</sub>) and the second based on the effective concentration  
48 required for a 2-fold increase in activity (i.e., EC<sub>2-fold</sub>) (Maloney and Waxman, 1999). Orally  
49 administered DEHP undergoes presystemic hydrolysis catalyzed by lipase to MEHP in the gut,  
50 with mice exhibiting higher lipase activities in the small intestine compared to rats and marmosets  
51 (Kessler et al., 2004; Pollack et al., 1985) Ito et al. 2005). Therefore, because the mouse liver is  
52 likely exposed predominantly to MEHP rather than DEHP and unmetabolized -----be

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 explained by pharmacokinetics, i.e., hepatic conversion of DEHP to its mono-ester MEHP, since  
 2 studies in rats demonstrate that orally administered DEHP undergoes presystemic hydrolysis to  
 3 MEHP in the gut (Kessler et al., 2004; Pollack et al., 1985). Possible explanations for these  
 4 results include one or more of the following: (1) the transactivation assay is not an accurate  
 5 quantitative indicator of in vivo receptor activation, (2) the rate and nature of effects downstream  
 6 of PPAR- $\alpha$  activation depends on the ligand or, (3) there are rate-limiting events independent of  
 7 PPAR- $\alpha$  activation that contribute to mouse hepatocarcinogenesis by the agonists examined.  
 8

9 **Hepatomegaly, DNA synthesis, and peroxisome proliferation in rats**

10 Table 1 [reproduced as 4-67] compares potency indicators for various precursor effects at  
 11 the TD<sub>50</sub> for four PPAR- $\alpha$  agonists and rat hepatocarcinogens. Our analysis of whether there are  
 12 consistent levels of in vivo precursor effect induction across peroxisome proliferators at the TD<sub>50</sub>  
 13 does not include all of the data from a similar, prior analysis by Ashby et al. (1994) for several  
 14 reasons. First, unlike the CPDB, Ashby et al. (1994) did not adjust carcinogenicity data for  
 15 less-than-lifetime dosing, which is relevant for most compounds. Second, for those mouse  
 16 carcinogens reported in the CPDB, only acute data are available regarding DNA synthesis effects  
 17 from Ashby et al.. Therefore, our analysis was restricted to rat precursor and potency data for the  
 18 four compounds Wy-14,643, nafenopin, clofibrate, and DEHP and included both 1-week and  
 19 13-week data to separately address transient and sustained changes in DNA synthesis. Even for  
 20 this small set of compounds, several limitations in the rat database were apparent. Because no  
 21 single study provided comparative data for the precursor endpoints of interest, four separate  
 22 reports were used. In the Wada et al. (1992) and Tanaka et al. (1992) studies of Wy-14,643 and  
 23 clofibrate, respectively, administered doses were within 10% of the TD<sub>50</sub>. However, nafenopin  
 24 data were only available at a single dose of 500 ppm (Lake et al., 1993), which was linearly  
 25 interpolated to the TD<sub>50</sub>. The highest administered dose of DEHP was 12,500 ppm (David et al.,  
 26 1999), a dose notably below the TD<sub>50</sub>, and thus a lower limit based on the assumption of  
 27 monotonicity with dose is shown. A further data limitation is that in the CPDB, only the TD<sub>50</sub> for  
 28 one of the four compounds, DEHP, incorporates data from studies administering more than one  
 29 dose for two years.

30 The results shown in Table 1 [reproduced as Table 4-67] indicate that potency for the  
 31 occurrence of short-term in vivo markers of PPAR- $\alpha$  activation varies widely in magnitude and  
 32 lacks any apparent correlation with carcinogenic potency. Such differences have been noted  
 33 previously. Similar to the results presented in Table 1 [reproduced as Table 4-67], Marsman et al.  
 34 (1988) noted that although DEHP (12,000 ppm) and Wy-14,643 (1,000 ppm) induced a similar  
 35 extent of hepatomegaly and peroxisome proliferation (measured either morphologically or  
 36 biochemically) after 1 year, the frequency of hepatocellular lesions was over 100-fold higher in  
 37 Wy-14,643- relative to DEHP-exposed rats. In addition, a higher labeling index was reported for  
 38 12,500 ppm DEHP than the maximal level attained after 50–1,000 ppm Wy-14,643 (David et al.,  
 39 1999; Tanaka et al., 1992; Wada et al., 1992). We did not examine such differences in maximal  
 40 responses in our analysis. We also do not present differences in response with dose and time seen  
 41 among PPAR- $\alpha$  agonists, which are prominent enough to prevent displaying dose-response data  
 42 on a common scale. For instance, labeling index is increased in a dose-dependent manner at 1  
 43 week by clofibrate (1,500, 4,500 and 9,000 ppm) but is decreased compared with controls at 13  
 44 weeks at the two higher doses (Tanaka et al., 1992). Together, these findings underscore the  
 45 significant chemical-specific quantitative differences in these markers that limit their utility for  
 46 predicting carcinogenic dose-response relationships.  
 47

48 **Table 4-66. Potency indicators for mouse hepatocarcinogenicity and in vitro**  
 49 **transactivation of mouse PPAR $\alpha$  for four PPAR $\alpha$  agonists**  
 50

	<b>Carcinogenic potency indicators</b>	<b>Transactivation potency indicators (<math>\mu</math>M)</b>
--	--	---

*This document is a draft for review purposes only and does not constitute Agency policy.*

	(mg/kg-day)		
Chemical	TD <sub>50</sub>	EC <sub>50</sub>	EC <sub>twofold</sub>
Hepatocarcinogens			
Wy-14,643	<10.8	0.63	~0.4
DCA	119	~300	~300
TCA	584	~300	~300
DEHP/MEHP	700	~0.7	~0.7

Source: reproduced from Table 2 of Guyton et al. (2009).

Note: TD<sub>50</sub> = the daily dose inducing tumors in half of the mice that would otherwise have remained tumor-free, estimated from the Carcinogenic Potency Database (Gold et al. 2005). EC<sub>50</sub> = the effective concentration yielding 50% of the maximal response; EC<sub>twofold</sub> = the effective concentration required for a twofold increase in activity. Transactivation potencies were estimated from Maloney and Waxman (1999). The “<” symbol denotes an upper limit due to maximal response. A “~” symbol indicates that the transactivation potency was approximated from figures in Maloney and Waxman (1999).

MEHP = monoethylhexyl phthalate.

**Table 4-67. Potency indicators for rat hepatocarcinogenicity and common short-term markers of PPAR $\alpha$  activation for four PPAR $\alpha$  agonists**

Chemical	Tumor TD <sub>50</sub> (ppm in diet)	Fold-increase over control at tumor TD <sub>50</sub>					
		1 wk			13 wk		
		RLW	LI	PCO	RLW	LI	PCO
Wy-14,643	109	1.8	12	13	2.6	6.8	39
Nafenopin	275	1.4	3.6	7.6	1.5	1.12	6.7
Clofibrate	4.225	1.4	4.4	4.2	1.4	0.95	3.7
DEHP	17.900	≥1.4	≥19	≥3.6	≥1.9	≥1.25	≥4.9

Source: reproduced from Table 1 of Guyton et al. (2009).

Note: For ease of comparison with precursor effect studies, administered doses for the tumor TD<sub>50</sub>s in the Carcinogenic Potency Database were back-converted to equivalent ppm in diet using the formula of Gold et al. (2005), *i.e.*, TD<sub>50</sub> (mg/kg-day) = TD<sub>50</sub> (ppm in diet) × 0.04 (for male rats). Administered doses for precursor data on Wy-14,643 (Wada et al., 1992) and clofibrate (Tanaka et al., 1992) were within 10% of the TD<sub>50</sub>. Because nafenopin precursor data were only available at 0 and 500 ppm (Lake et al., 1993), these doses were linearly interpolated to the TD<sub>50</sub>. Because the highest administered dose of DEHP in precursor effect studies was 12,500 ppm (David et al., 1999), a lower limit is shown, based on the assumption of monotonicity with dose. RLW = relative liver weight, LI = labeling index, PCO = cyanide insensitive palmitoyl CoA oxidation.

#### **4.5.7.1.6. Additional Proposed Hypotheses and Key Events with Limited Evidence or Inadequate Experimental Support**

1           Several effects that been hypothesized to be associated with liver cancer induction are  
2 discussed in more detail below, including increased liver weight, DNA hypomethylation, and  
3 pathways involved in glycogen accumulation such as insulin signaling proteins. As discussed  
4 above, TCE and its metabolites reportedly increase nuclear size and ploidy in hepatocytes, and  
5 these effects likely account for much of the increases in labeling index and DNA synthesis  
6 caused by TCE. Importantly, these changes appear to persist with cessation of treatment, with  
7 liver weights, but not nuclear sizes, returning to control levels (Kjellstrand et al., 1983a). In  
8 addition, glycogen deposition, DNA synthesis, increases in mitosis, or peroxisomal enzyme  
9 activity do not appear correlated with TCE-induced liver weight changes.

10

#### **4.5.7.1.7. Increased liver weight**

11           Increased liver weight or liver/body weight ratios (hepatomegaly) is associated with  
12 increased risk of liver tumors in rodents, but it is relatively nonspecific (Allen et al., 2004). The  
13 evidence presented above for TCE and its metabolites suggest a similarity in dose-response  
14 between liver weight increases at short-term durations of exposure and liver tumor induction  
15 observed from chronic exposure. Liver weight increases may results from several concurrent  
16 processes that have been associated with increase cancer risk (e.g., hyperplasia, increased ploidy,  
17 and glycogen accumulation) and when observed after chronic exposure may result from the  
18 increased presence of foci and tumors themselves. Therefore, there are inadequate data to  
19 adequately define a MOA hypothesis for hepatocarcinogenesis based on liver weight increases.

20

#### **4.5.7.1.8. “Negative selection”**

21           As discussed above, TCE, TCA, and DCA all cause transient increases in DNA synthesis.  
22 This DNA synthesis has been assumed to result from proliferation of hepatocytes. However, the  
23 dose-related TCA- and DCA-induced increases in liver weight not correlate with patterns of  
24 DNA synthesis; moreover, there have been reports that DNA synthesis in individual hepatocytes  
25 does not correlate with whole liver DNA synthesis measures (Carter et al., 1995; Sanchez and  
26 Bull, 1990). With continued treatment, decreases in DNA synthesis have been reported for DCA  
27 (Carter et al., 1995). More importantly, several studies show that transient DNA synthesis is  
28 confined to a very small population of cells in the liver in mice exposed to TCE for 10 days or to

1 DCA or TCA for up to 14 days of exposure. Therefore, generalized mitogenic stimulation is not  
2 likely to play a role in TCE-induced liver carcinogenesis.

3 Bull has proposed that the TCE metabolites TCA and DCA may contribute to liver tumor  
4 induction through so-called “negative selection” by way of several possible processes (Bull,  
5 2000). First, it is hypothesized that the mitogenic stimulation by continued TCA and DCA  
6 exposure is down-regulated in normal hepatocytes, conferring a growth advantage to initiated  
7 cells that either do not exhibit the down-regulation of response or are resistant to the  
8 down-regulating signals. This is implausible as both the normal rates of cell division in the liver  
9 and the TCE-stimulated increases are very low. Polyploidization has been reported to decrease  
10 the normal rates of cell division even further. That the transient and relatively low level of DNA  
11 synthesis reported for TCE, DCA, and TCA is reflective of proliferation rather than  
12 polyploidization is not supported by data on mitosis. A mechanism for such “down-regulation”  
13 has not been identified experimentally.

14 A second proposed contributor to “negative-selection” is direct enhancement by TCA and  
15 DCA in the growth of certain populations of initiated cells. While differences in phenotype of  
16 end stage tumors have been reported between DCA and TCA, the role of selection and  
17 emergence of potentially different foci has not been elucidated. Neither have pathway  
18 perturbations been identified that are common to liver cancer in human and rodent for TCE,  
19 DCA, and TCA. The selective growth of clones of hepatocytes that may progress fully to cancer  
20 is a general feature of cancer and not specific to at TCE, TCA, or DCA MOA.

21 A third proposed mechanism by which TCE may enhance liver carcinogenesis within this  
22 “negative selection” paradigm is through changing apoptosis. However, as stated above, TCE  
23 has been reported to either not change apoptosis or to cause a slight increase at high doses.  
24 Rather than increases in apoptosis, peroxisome proliferators have been suggested to inhibit  
25 apoptosis as part of their carcinogenic MOA. However, the age and species studied appear to  
26 greatly affect background rates of apoptosis (Snyder et al., 1995) with the rat having a greater  
27 rate of apoptosis than the mouse. DCA has been reported to induce decreases in apoptosis in the  
28 mouse (Carter et al., 1995; Snyder et al., 1995). However, the significance of the DCA-induced  
29 reduction in apoptosis, from a level that is already inherently low in the mouse, for the MOA for  
30 induction of DCA-induce liver cancer is difficult to discern.

31 Therefore, for a MOA for hepatocarcinogenesis based on “negative selection,” there are  
32 inadequate data to adequately define the MOA hypothesis, or the available data do not support  
33 such a MOA being operative.

34



#### 4.5.7.1.9. Polyploidization

1            Polyploidization may be an important key event in tumor induction. For example, in  
2 addition to TCE, partial hepatectomy, nafenopin, methylclofenopate, DEHP, diethylnitrosamine,  
3 *N*-nitrosomorpholine, and various other exposures that contribute to liver tumor induction also  
4 shift the hepatocyte ploidy distribution to be increasingly diploid or polyploid (Hasmall and  
5 Roberts, 2000; Melchiorri et al., 1993; Miller et al., 1996; Styles et al., 1988; Vickers and Lucier,  
6 1996). As discussed by Gupta (2000), “[w]orking models indicate that extensive polyploidy  
7 could lead to organ failure, as well as to oncogenesis with activation of precancerous cell  
8 clones.” However, the mechanism(s) by which increased polyploidy enhances carcinogenesis is  
9 not currently understood. Due to increased DNA content, polyploid cells will generally have  
10 increased gene expression. However, polyploid cells are considered more highly differentiated  
11 and generally divide more slowly and are more likely to undergo apoptosis, perhaps thereby  
12 indirectly conferring a growth advantage to initiated cells (see Section E.1). Of note is that  
13 changes in ploidy have been observed in transgenic mouse models that are also prone to develop  
14 liver cancer (see Section E.3.3.1). It is likely that polyploidization occurs with TCE exposure  
15 and it is biologically plausible that polyploidization can contribute to liver carcinogenesis,  
16 although the mechanism(s) is (are) not known. However, whether polyploidization is necessary  
17 for TCE-induced carcinogenesis is not known, as no experiment in which polyploidization  
18 specifically is blocked or diminished has been performed and the extent of polyploidization has  
19 not been quantified. Therefore, there are inadequate data to adequately define a MOA  
20 hypothesis for hepatocarcinogenesis based on polyploidization.  
21

#### 4.5.7.1.10. Glycogen storage

22            As discussed above, several studies have reported that DCA causes accumulation of  
23 glycogen in mouse hepatocytes. Such glycogen accumulation has been suggested to be  
24 pathogenic, as it is resistant to mobilization by fasting (Kato-Weinstein et al., 1998). In humans,  
25 glycogenesis due to glycogen storage disease or poorly controlled diabetes has been associated  
26 with increased risk of liver cancer (Adami et al., 1996; La Vecchia et al., 1994; Rake et al., 2002;  
27 Wideroff et al., 1997). Glycogen accumulation has also been reported to occur in rats exposed to  
28 DCA.

29            For TCE exposure in mice or rats, glycogen content of hepatocytes has been reported to  
30 be somewhat less than or the same as controls, or not remarked upon in the studies. TCA  
31 exposure has been reported to decrease glycogen content in rodent hepatocytes while DCA has  
32 been reported to increase it (Kato-Weinstein et al., 2001). There is also evidence that

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 DCA-induced increases in glycogen accumulation are not proportional to liver weight increases  
2 and only account for a relatively small portion of increases in liver mass. DCA-induced  
3 increases in liver weight are not a function of cellular proliferation but probably include  
4 hypertrophy associated with polyploidization, increased glycogen deposition and other factors.

5 While not accounting for increases in liver weight, excess glycogen can still be not only  
6 be pathogenic but a predisposing condition for hepatocarcinogenesis. Some hypotheses  
7 regarding the possible relationship between glycogenesis and carcinogenesis have been posed  
8 that lend them biological plausibility. Evert et al. (2003), using an animal model of hepatocyte  
9 exposure to a local hyperinsulinemia from transplanted islets of Langerhans with remaining  
10 tissue is hypoinsulinemic, reported that insulin induces alterations resembling preneoplastic foci  
11 of altered hepatocytes that develop into hepatocellular tumors in later stages of carcinogenesis.  
12 Lingohr et al. (2001) suggest that normal hepatocytes down-regulate insulin-signaling proteins in  
13 response to the accumulation of liver glycogen caused by DCA and that the initiated cell  
14 population, which does not accumulate glycogen and is promoted by DCA treatment, responds  
15 differently from normal hepatocytes to the insulin-like effects of DCA. Bull et al. (2002)  
16 reported increased insulin receptor protein expression in tumor tissues regardless of whether they  
17 were induced by TCE, TCA, or DCA. Given the greater activity of DCA relative to TCA on  
18 carbohydrate metabolism, it is unclear whether changes in these pathways are causes or simply  
19 reflect the effects of tumor progression. Therefore, it is biologically plausible that changes in  
20 glycogen status may occur from the opposing actions of TCE metabolites, but changes in  
21 glycogen content due to TCE exposure has not been quantitatively studied. The possible  
22 contribution of these effects to TCE-induced hepatocarcinogenesis is unclear. Therefore, there  
23 are inadequate data to adequately define a MOA hypothesis for TCE-induced  
24 hepatocarcinogenesis based on changes in glycogen storage or even data to support increased  
25 glycogen storage to result from TCE exposure.

#### 26 **4.5.7.1.11. Inactivation of GST-zeta**

27 DCA has been shown to inhibit its own metabolism in that pretreatment in rodents prior  
28 to a subsequent challenge dose leads to a longer biological half-life (Schultz et al., 2002). This  
29 self-inhibition is hypothesized to occur through inactivation of GST-zeta (Schultz et al., 2002).  
30 In addition, TCE has been shown to cause the same prolongation of DCA half-life in rodents,  
31 suggesting that TCE inhibits GST-zeta, probably through the formation of DCA (Schultz et al.,  
32 2002). DCA-induced inhibition of GST-zeta has also been reported in humans, with GST-zeta  
33 polymorphisms reported to influence the degree of inactivation (Blackburn et al., 2001;

1 Blackburn et al., 2000; Tzeng et al., 2000). Board et al. (2001) report one variant to have  
2 significantly higher activity with DCA as a substrate than other GST-zeta isoforms, which could  
3 affect DCA susceptibility.

4 GST-zeta, which is identical to maleylacetoacetate isomerase, is part of the tyrosine  
5 catabolism pathway which is disrupted in Type 1 hereditary tyrosinemia, a disease associated  
6 with the development of hepatocellular carcinoma at a young age (Tanguay et al., 1996). In  
7 particular, GST-zeta metabolizes maleylacetoacetate (MAA) to fumarylacetoacetate (FAA) and  
8 maleylacetone (MA) to fumarylacetone (Cornett et al., 1999; Tanguay et al., 1996). It has been  
9 suggested that the increased cancer risk with this disease, as well as through DCA exposure,  
10 results from accumulation of MAA and MA, both alkylating agents, or FAA, which displays  
11 apoptogenic, mutagenic, aneugenic, and mitogenic activities (Bergeron et al., 2003; Cornett et  
12 al., 1999; Jorquera and Tanguay, 2001; Kim et al., 2000; Tanguay et al., 1996). However, the  
13 possible effects of DCA through this pathway will depend on whether MAA, MA, or FAA is the  
14 greater risk factor, since inhibition of GST-zeta will lead to greater concentrations of MAA and  
15 MA and lower concentrations of FAA. Therefore, if MAA is the more active agent, DCA may  
16 increase carcinogenic risk, while if FAA is the more active, DCA may decrease carcinogenic  
17 risk. Tzeng et al. (2000) propose the later based on the greater genotoxicity of FAA, and in fact  
18 suggest that DCA may “merit consideration for trial in the clinical management of hereditary  
19 tyrosinemia type 1.”

20 Therefore, TCE-induced inactivation GST-zeta, probably through formation of DCA,  
21 may play a role in TCE-induced hepatocarcinogenesis. However, this mode of action is not  
22 sufficiently delineated at this point for further evaluation, as even the question of whether its  
23 actions through this pathway may increase or decrease cancer risk has yet to be experimentally  
24 tested.

#### 25 **4.5.7.1.12. Oxidative stress**

26 Several studies have attempted to study the possible effects of “oxidative stress” and  
27 DNA damage resulting from TCE exposures. The effects of induction of metabolism by TCE, as  
28 well as through coexposure to ethanol, have been hypothesized to in itself increase levels of  
29 “oxidative stress” as a common effect for both exposures (see Section E.4.2.4). In terms of  
30 contributing to a carcinogenic MOA, the term “oxidative stress” is a somewhat nonspecific term,  
31 as it is implicated as part of the pathophysiologic events in a multitude of disease processes and  
32 is part of the normal physiologic function of the cell and cell signaling. Commonly, it appears to  
33 refer to the formation of reactive oxygen species leading to cellular or DNA damage. As

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 discussed above, however, measures of oxidative stress induced by TCE, TCA, and DCA appear  
2 to be either not apparent, or at the very most transient and nonpersistent with continued treatment  
3 (Channel et al., 1998; Larson and Bull, 1992b; Parrish et al., 1996; Toraason et al., 1999).

4 Therefore, while the available data are limited, there is insufficient evidence to support a role for  
5 such effects in TCE-induced liver carcinogenesis.

6 Oxidative stress has been hypothesized to be part of the MOA for peroxisome  
7 proliferators, but has been found to neither be correlated with cell proliferation nor carcinogenic  
8 potency of peroxisome proliferators (see Section E.3.4.1.1). For instance, Parrish et al. (1996)  
9 reported that increases in PCO activity noted for DCA and TCA were not associated with  
10 8-OHdG levels (which were unchanged) and also not with changes laurate hydrolase activity  
11 observed after either DCA or TCA exposure. The authors concluded that their data do not  
12 support an increase in steady state oxidative damage to be associated with TCA initiation of  
13 cancer and that extension of treatment to time periods sufficient to insure peroxisome  
14 proliferation failed to elevate 8-OHdG in hepatic DNA. The authors thus, suggested that  
15 peroxisome proliferative properties of TCA were not linked to oxidative stress or carcinogenic  
16 response.

#### 4.5.7.1.13. Changes in gene expression (e.g., hypomethylation)

18 Studies of gene expression as well as considerations for interpretation of studies of using  
19 the emerging technologies of DNA, siRNA, and miRNA microarrays for MOA analyses are  
20 included in Sections E.3.1.2 and E.3.4.2.2. Caldwell and Keshava (2006) and Keshava and  
21 Caldwell (2006) report on both genetic expression studies and studies of changes in methylation  
22 status induced by TCE and its metabolites as well as differences and difficulties in the patterns of  
23 gene expression between differing PPAR $\alpha$  agonists. In particular are concerns for the  
24 interpretation of studies which employ pooling of data as well as interpretation of “snapshots in  
25 time of multiple gene changes.” For instance, in the Laughter et al. (2004) study, it is not clear  
26 whether transcription arrays were performed on pooled data as well as the issue of phenotypic  
27 anchoring as data on percentage liver/body weight indicates significant variability within TCE  
28 treatment groups, especially in PPAR $\alpha$ -null mice. For studies of gene expression using  
29 microarrays Bartosiewicz et al. (2001) used a screening analysis of 148 genes for  
30 xenobiotic-metabolizing enzymes, DNA repair enzymes, heat shock proteins, cytokines, and  
31 housekeeping gene expression patterns in the liver in response TCE. The TCE-induced gene  
32 induction was reported to be highly selective; only Hsp 25 and 86 and Cyp2a were up-regulated  
33 at the highest dose tested. Collier et al. (2003) reported differentially expressed mRNA

1 transcripts in embryonic hearts from Sprague-Dawley rats exposed to TCE with sequences  
2 down-regulated with TCE exposure appearing to be those associated with cellular housekeeping,  
3 cell adhesion, and developmental processes. TCE was reported to induce up-regulated  
4 expression of numerous stress-response and homeostatic genes.

5 For the Laughter et al. (2004) study, transcription profiles using macroarrays containing  
6 approximately 1,200 genes were reported in response to TCE exposure with 43 genes reported to  
7 be significantly altered in the TCE-treated wild-type mice and 67 genes significantly altered in  
8 the TCE-treated PPAR $\alpha$  knockout mice. However, the interpretation of this information is  
9 difficult because in general, PPAR $\alpha$  knockout mice have been reported to be more sensitive to a  
10 number of hepatotoxins partly because of defects in the ability to effectively repair tissue damage  
11 in the liver (Mehendale, 2000; Shankar et al., 2003) and because a comparison of gene  
12 expression profiles between controls (wild-type and PPAR $\alpha$  knockout) were not reported. As  
13 reported by Voss et al. (2006), dose-, time course-, species-, and strain-related differences should  
14 be considered in interpreting gene array data. The comparison of differing PPAR $\alpha$  agonists  
15 presented in Keshava and Caldwell (2006) illustrate the pleiotropic and varying liver responses  
16 of the PPAR $\alpha$  receptor to various agonists, but did not imply that these responses were  
17 responsible for carcinogenesis.

18 As discussed above in Section E.3.3.5, Aberrant DNA methylation is a common hallmark  
19 of all types of cancers, with hypermethylation of the promoter region of specific tumor  
20 suppressor genes and DNA repair genes leading to their silencing (an effect similar to their  
21 mutation) and genome-wide hypomethylation (Ballestar and Esteller, 2002; Berger and  
22 Daxenbichler, 2002; Herman et al., 1998; Pereira et al., 2004a; Rhee et al., 2002). Whether  
23 DNA methylation is a consequence or cause of cancer is a long-standing issue (Ballestar and  
24 Esteller, 2002). Fraga et al. (2005; 2004) reported global loss of monoacetylation and  
25 trimethylation of histone H4 as a common hallmark of human tumor cells; they suggested,  
26 however, that genomewide loss of 5-methylcytosine (associated with the acquisition of a  
27 transformed phenotype) exists not as a static predefined value throughout the process of  
28 carcinogenesis but rather as a dynamic parameter (i.e., decreases are seen early and become more  
29 marked in later stages).

30 DNA methylation is a naturally occurring epigenetic mechanism for modulating gene  
31 expression, and disruption of this mechanism is known to be relevant to human carcinogenesis.  
32 As reviewed by Calvisi et al. (2007),

33  
34  
35 [a]berrant DNA methylation occurs commonly in human cancers in the forms of genome-wide  
36 hypomethylation and regional hypermethylation. Global DNA hypomethylation (also known as  
37 demethylation) is associated with activation of protooncogenes, such as c-Jun, c-Myc, and

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 c-HA-Ras, and generation of genomic instability. Hypermethylation on CpG islands located in the  
2 promoter regions of tumor suppressor genes results in transcriptional silencing and genomic  
3 instability.  
4  
5

6 While clearly associated with cancer, it has not been conclusively established whether these  
7 epigenetic changes play a causative role or are merely a consequence of transformation  
8 (Tryndyak et al., 2006). However, as Calvisi et al. (2007) note, “Current evidence suggests that  
9 hypomethylation might promote malignant transformation via multiple mechanisms, including  
10 chromosome instability, activation of protooncogenes, reactivation of transposable elements, and  
11 loss of imprinting.”

12 Although little is known about how it occurs, a hypothesis has also been proposed that  
13 that the toxicity of TCE and its metabolites may arise from its effects on DNA methylation  
14 status. In regard to methylation studies, many are coexposure studies as they have been  
15 conducted in initiated animals with some studies being very limited in their reporting and  
16 conduct. Caldwell and Keshava (2006) review the body of work regarding TCE, DCA, and  
17 TCA. Methionine status has been noted to affect the emergence of liver tumors (Counts et al.,  
18 1996). Tao et al. (2000a) and Pereira et al. (2004b) have studied the effects of excess methionine  
19 in the diet to see if it has the opposite effects as a deficiency (i.e., and reduction in a carcinogenic  
20 response rather than enhancement). However, Tao et al. (2000a) report that the administration of  
21 excess methionine in the diet is not without effect and can result in percentage liver/body weight  
22 ratios. Pereira et al. (2004b) report that methionine treatment alone at the 8 g/kg level was  
23 reported to increase liver weight, decrease lauryl-CoA activity and to increase DNA methylation.

24 Pereira et al. (2004b) reported that very high level of methionine supplementation to an  
25 AIN-760A diet, affected the number of foci and adenomas after 44 weeks of coexposure to  
26 3.2 g/L DCA. However, while the highest concentration of methionine (8.0 g/kg) was reported  
27 to decrease both the number of DCA-induce foci and adenomas, the lower level of methionine  
28 coexposure (4.0 g/kg) increased the incidence of foci. Coexposure of methionine (4.0 or  
29 8.0 g/kg) with 3.2 g/L DCA was reported to decrease by ~25% DCA-induced glycogen  
30 accumulation, increase mortality, but not to have much of an effect on peroxisome enzyme  
31 activity (which was not elevated by more than 33% over control for DCA exposure alone). The  
32 authors suggested that their data indicate that methionine treatment slowed the progression of  
33 foci to tumors. Given that increasing hypomethylation is associated with tumor progression,  
34 decreased hypomethylation from large doses of methionine are consistent with a slowing of  
35 progression. Whether, these results would be similar for lower concentrations of DCA and lower  
36 concentrations of methionine that were administered to mice for longer durations of exposure,  
37 cannot be ascertained from these data. It is possible that in a longer-term study, the number of

1 tumors would be similar. Finally, a decrease in tumor progression by methionine  
2 supplementation is not shown to be a specific event for the MOA for DCA-induced liver  
3 carcinogenicity.

4 Tao et al. (2000a) reported that 7 days of gavage dosing of TCE (1,000 mg/kg in corn  
5 oil), TCA (500 mg/kg, neutralized aqueous solution), and DCA (500 mg/kg, neutralized aqueous  
6 solution) in 8-week old female B6C3F1 mice resulted in not only increased liver weight but also  
7 increased hypomethylation of the promoter regions of *c-jun* and *c-myc* genes in whole liver  
8 DNA. However, data were shown for 1–2 mice per treatment. Treatment with methionine was  
9 reported to abrogate this response only at a 300 mg/kg i.p dose with 0–100 mg/kg doses of  
10 methionine having no effect. Ge et al. (2001a) reported DCA- and TCA-induced DNA  
11 hypomethylation and cell proliferation in the liver of female mice at 500 mg/kg and decreased  
12 methylation of the *c-myc* promoter region in liver, kidney and urinary bladder. However,  
13 increased cell proliferation preceded hypomethylation. Ge et al. (2002) also reported  
14 hypomethylation of the *c-myc* gene in the liver after exposure to the peroxisome proliferators  
15 2,4-dichlorophenoxyacetic acid (1,680 ppm), DBP (20,000 ppm), gemfibrozil (8,000 ppm), and  
16 Wy-14,643 (50–500 ppm, with no effect at 5 or 10 ppm) after 6 days in the diet. Caldwell and  
17 Keshava (2006) concluded that hypomethylation did not appear to be a chemical-specific effect  
18 at these concentrations. As noted Section E.3.3.5, chemical exposure to a number of differing  
19 carcinogens have been reported to lead to progressive loss of DNA methylation..

20 After initiation by *N*-methyl-*N*-nitrosourea (25 mg/kg) and exposure to 20 mmol/L DCA  
21 or TCA (46 weeks), Tao et al. (2004a) report similar hypomethylation of total mouse liver DNA  
22 by DCA and TCA with tumor DNA showing greater hypomethylation. A similar effect was  
23 noted for the differentially methylated region-2 of the insulin-like growth factor-II (IGF-II) gene.  
24 The authors suggest that hypomethylation of total liver DNA and the IGF-II gene found in  
25 nontumorous liver tissue would appear to be the result of a more prolonged activity and not cell  
26 proliferation, while hypomethylation of tumors could be an intrinsic property of the tumors. As  
27 pointed out by Caldwell and Keshava (2006) over expression of IGF-II gene in liver tumors and  
28 preneoplastic foci has been shown in both animal models of hepatocarcinogenesis and humans,  
29 and may enhance tumor growth, acting via the over-expressed IGF-I receptor (Scharf et al.,  
30 2001; Werner and Le Roith, 2000).

31 Diminished hypomethylation was observed in Wy-14643-treated PPAR $\alpha$ -null mice as  
32 compared to wild-type mice, suggestive of involvement of PPAR $\alpha$  in mediating hypomethylation  
33 (Pogribny et al., 2007), but it is unclear how relevant these results are to TCE and its metabolites.  
34 First, the doses of Wy-14643 administered are associated with substantial liver necrosis and  
35 mortality with long-term treatment (Woods et al., 2007a), adding confounding factors the

1 interpretation of their results. Hypomethylation by Wy-14643 progressively increased with time  
2 up to 5 months (Pogribny et al., 2007), consistent with the sustained DNA synthesis caused by  
3 Wy-14643 and a role for proliferation in causing hypomethylation. Regardless, as discussed  
4 above, it is unlikely that PPAR $\alpha$  is the mediator of the observed transient increase in DNA  
5 synthesis by DCA, so even if it is important for hypomethylation by TCA, there may be more  
6 than one pathway for this effect.

7 To summarize, aberrant DNA methylation status, including hypomethylation, is clearly  
8 associated with both human and rodent carcinogenesis. Hypomethylation itself appears to be  
9 sufficient for carcinogenesis, as diets deficient in choline and methionine that induce  
10 hypomethylation have been shown to cause liver tumors in both rats and mice (Ghoshal and  
11 Farber, 1984; Henning and Swendseid, 1996; Mikol et al., 1983; Wainfan and Poirier, 1992).  
12 However, it is not known to what extent hypomethylation is necessary for TCE-induced  
13 carcinogenesis. However, as noted by Bull (2004a) and Bull et al. (2004), the doses of TCA and  
14 DCA that have been tested for induction of hypomethylation are quite high compared to doses at  
15 which tumor induction occurs—at least 500 mg/kg-day. Whether these effects are still manifest  
16 at lower doses relevant to TCE carcinogenicity, particularly with respect to DCA, has not been  
17 investigated. Finally, the role of PPAR $\alpha$  in modulating hypomethylation, possibly through  
18 increased DNA synthesis as suggested by experiments with Wy-14643, are unknown for TCE  
19 and its metabolites.

#### 4.5.7.1.14. Cytotoxicity

21 Cytotoxicity and subsequent induction of reparative hyperplasia have been proposed as  
22 key events for a number of chlorinated solvents, such as chloroform and carbon tetrachloride..  
23 However, as discussed above and discussed by Bull (2004a) and Bull et al. (2004), TCE  
24 treatment at doses relevant to liver carcinogenicity results in relatively low cytotoxicity. While a  
25 number of histological changes with TCE exposure are observed, in most cases necrosis is  
26 minimal or mild, associated with vehicle effects, and with relatively low prevalence. This is  
27 consistent with the low prevalence of necrosis observed with TCA and DCA treatment at doses  
28 relevant to TCE exposure. Therefore, it is unlikely that cytotoxicity and reparative hyperplasia  
29 play a significant role in TCE carcinogenicity



#### **4.5.7.1.15. Mode of Action (MOA) Conclusions**

1           The conclusions regarding the MOA for TCE-induced liver carcinogenesis described in  
2 the preceding sections are summarized in Table 4-68. Overall, although a role for many of the  
3 proposed key events discussed above cannot be ruled out, there are inadequate data to support  
4 the conclusion that any of the particular MOA hypotheses reviewed above are operant. The  
5 available data do suggest that the MOA of liver tumors induced by TCE is complex, as it is  
6 likely that key events from several pathways may operate. Nonetheless, because a collection of  
7 key events sufficient to induce liver tumors has not been identified, the answer to the first key

**Table 4-68. Summary of mode of action conclusions for TCE-induced liver carcinogenesis**

Hypothesized MOA and key events	Evidence that TCE or TCE metabolites induces key events	Necessity of MOAs key events for carcinogenesis	Sufficiency of MOA for carcinogenesis	Summary and section reference
Mutagenicity induced by one or more metabolites advances acquisition of multiple critical traits contributing to carcinogenesis	In rodents, TCE binds to and/or induces damage in DNA and chromosome structure. TCE has a limited ability to induce mutation in bacterial systems. Oxidative metabolites, particularly CH, can cause a variety of genotoxic effects (including aneuploidy) in available in vitro and in vivo assays.	No TCE-specific studies	No TCE-specific studies; mutagenicity is assumed to cause cancer, as a sufficient cause	Data are inadequate to support a conclusion that a mutagenic MOA mediated by CH is operant; however, a mutagenic MOA, mediated either by CH or other oxidative metabolites of TCE, cannot be ruled out. Section 4.5.7.1 ; see also 4.2.1.5
PPAR $\alpha$ activation: <ul style="list-style-type: none"> <li>• TCE oxidative metabolites (e.g., TCA), after being produced in the liver, activate PPAR<math>\alpha</math></li> <li>• Alterations in cell proliferation and apoptosis</li> <li>• Clonal expansion of initiated cells</li> </ul>	TCE, TCA and DCA activate PPAR $\alpha$ , induce peroxisome proliferation and hepatocyte proliferation in mice and rats (e.g., DeAngelo et al., 2008; Dees and Travis, 1994; Elcombe et al., 1985; Goel et al., 1992; Goldsworthy and Popp, 1987; Laughter et al., 2004; Nakajima et al., 2000; Pereira, 1996; Sanchez and Bull, 1990; Stauber and Bull, 1997; Watanabe and Fukui, 2000).	No studies of TCE or its metabolites (e.g., cancer bioassays in PPAR $\alpha$ -null mice). TCE induces increases in liver weight in male and female mice lacking a functional PPAR $\alpha$ receptor (Nakajima et al., 2000; Ramdhan et al., 2010) and in humanized null mice (Ramdhan et al., 2010). Liver tumor response from WY dramatically diminished in PPAR $\alpha$ -null mice (Peters et al., 1997); however, liver tumor response from DEHP unchanged in PPAR $\alpha$ -null mice (Ito et al., 2007). Thus, inferences regarding TCE are not possible.	No TCE-specific studies; PPAR $\alpha$ activation in a transgenic mouse model caused all the key events in the MOA, but not carcinogenesis, suggesting that the MOA is not sufficient for carcinogenesis (Yang et al., 2007). Consistent with hypothesis that TCE liver carcinogenesis involves multiple mechanisms.	It is unlikely that PPAR $\alpha$ agonism and its sequelae constitute the sole or predominant MOA for TCE-induced carcinogenesis. Section 4.5.7.2

Table 4-68. Summary of mode of action conclusions for TCE-induced liver carcinogenesis (continued)

Hypothesized MOA and key events	Evidence that TCE or TCE metabolites induces key events	Necessity of MOAs key events for carcinogenesis	Sufficiency of MOA for carcinogenesis	Summary and section reference
Liver weight increases	TCE increases liver weight in mice, rats and gerbils following acute, short-term and subchronic exposures (Berman et al., 1995; Buben and O'Flaherty, 1985; Dees and Travis, 1993, 1994; Elcombe et al., 1985; Goel et al., 1992; Goldsworthy and Popp, 1987; Kjellstrand et al., 1983a; Kjellstrand et al., 1983b; Kjellstrand et al., 1981b; Laughter et al., 2004; Melnick et al., 1987; Merrick et al., 1989; Nakajima et al., 2000; Nunes et al., 2001; Ramdhan et al., 2010; Tao et al., 2000a; Tucker et al., 1982). Similarity in dose-response between liver weight increases at short-term durations of exposure and liver tumor induction observed from chronic exposure to TCE and metabolites.	No TCE-specific studies	No TCE-specific studies	Data are inadequate to define a MOA hypothesis for hepatocarcinogenesis based on liver weight increases. Section 4.5.7.3.1
“Negative selection” confers a growth advantage to initiated cells	Transient DNA synthesis is confined to a very small population of cells in mouse liver (e.g., Dees and Travis, 1993; Elcombe et al., 1985; Laughter et al., 2004).	No TCE-specific studies	No TCE-specific studies	Data are inadequate to define a MOA hypothesis for hepatocarcinogenesis based on “negative selection”. Section 4.5.7.3.3
Polyploidization	Polyploidization likely occurs with TCE exposure, although the evidence is limited (Buben and O'Flaherty, 1985).	No TCE-specific studies	No TCE-specific studies	Although it is biologically plausible that polyploidization can contribute to liver carcinogenesis, inadequate data are available to support this hypothesized MOA for TCE. Section 4.5.7.3.3

**Table 4-68. Summary of mode of action conclusions for TCE-induced liver carcinogenesis (continued)**

<b>Hypothesized MOA and key events</b>	<b>Evidence that TCE or TCE metabolites induces key events</b>	<b>Necessity of MOAs key events for carcinogenesis</b>	<b>Sufficiency of MOA for carcinogenesis</b>	<b>Summary and section reference</b>
Glycogen storage	DCA increases glycogen deposition (Nelson et al., 1989); For TCE and TCA, effects on glycogen were either not reported (Dees and Travis, 1993; Elcombe et al., 1985; Styles et al., 1991) or were described as similar to controls (Nelson et al., 1989).	No TCE-specific studies	No TCE-specific studies	Data are inadequate to define a MOA hypothesis for TCE-induced hepatocarcinogenesis based on changes in glycogen storage. Section 4.5.7.3.4
Inactivation of GST-zeta	TCE prolongs DCA half-life in rodents, suggesting that TCE may inhibit GST-zeta, likely through the formation of DCA (Schultz et al., 2002).	No TCE-specific studies	No TCE-specific studies	Data are inadequate to define a MOA hypothesis for TCE-induced hepatocarcinogenesis based on inactivation of GST-zeta. Section 4.5.7.3.5
Oxidative stress	Measures of oxidative stress induced by TCE, TCA, and DCA either do not occur, or are transient and do not persistent with continued treatment (Channel et al., 1998; Larson and Bull, 1992b; Parrish et al., 1996).	No TCE-specific studies	No TCE-specific studies	Available data are limited to support a role for oxidative stress in TCE-induced liver carcinogenesis. Section 4.5.7.3.6
Epigenetic changes, particularly DNA methylation, induced by one or more metabolites (TCA, DCA, and other reactive species) advance acquisition of multiple critical traits contributing to carcinogenesis	TCE, TCA and DCA decrease global DNA methylation and promoter hypomethylation (e.g., of c-myc) in mouse liver (Ge et al., 2001b; Tao et al., 1998; Tao et al., 2004a)	No TCE-specific studies	No TCE-specific studies	Although it is biologically plausible that epigenetic changes contribute to liver carcinogenesis, inadequate data are available to support this hypothesized MOA for TCE. Section 4.5.7.3.7

**Table 4-68. Summary of mode of action conclusions for TCE-induced liver carcinogenesis (continued)**

<b>Hypothesized MOA and key events</b>	<b>Evidence that TCE or TCE metabolites induces key events</b>	<b>Necessity of MOAs key events for carcinogenesis</b>	<b>Sufficiency of MOA for carcinogenesis</b>	<b>Summary and section reference</b>
Cytotoxicity and reparative hyperplasia: <ul style="list-style-type: none"> <li>• One or more reactive intermediates induces hepatotoxicity</li> <li>• Reparative hyperplasia ensues</li> </ul>	TCE treatment at doses relevant to liver carcinogenicity results in relatively low cytotoxicity (Bull, 2004a; Bull et al., 2004).	No TCE-specific studies	No TCE-specific studies	It is unlikely that cytotoxicity and reparative hyperplasia play a significant role in TCE carcinogenicity. Section 4.5.7.3.8

1 question “*1. Is the hypothesized mode of action sufficiently supported in the test animals?*” is  
2 “no” at this time. Consequently, the other key questions of “*2. Is the hypothesized mode of*  
3 *action relevant to humans?*” and “*3. Which populations or lifestages can be particularly*  
4 *susceptible to the hypothesized mode of action?*” will not be discussed in a MOA-specific  
5 manner. Rather, they are discussed below in more general terms, first qualitatively and then  
6 quantitatively, using available relevant data.  
7

#### 4.5.7.1.16. Qualitative human relevance and susceptibility

8 No data exist that suggests that TCE-induced liver tumorigenesis is caused by processes  
9 that irrelevant in humans. In addition, as discussed above, several of the other effects such as  
10 polyploidization, changes in glycogen storage, and inhibition of GST-zeta—are either clearly  
11 related to human carcinogenesis or areas of active research as to their potential roles. For  
12 example, the effects of DCA on glycogen storage parallel the observation that individuals with  
13 conditions that lead to glycogenesis appear to be at an increased risk of liver cancer (Adami et  
14 al., 1996; La Vecchia et al., 1994; Rake et al., 2002; Wideroff et al., 1997). In addition, there  
15 may be some relationship between the effects of DCA and the mechanism of increased liver  
16 tumor risk in childhood in those with Type 1 hereditary tyrosinemia, though the hypotheses  
17 needs to be tested experimentally. Similarly, with respect to PPAR $\alpha$  activation and downstream  
18 events hypothesized to be causally related to liver carcinogenesis, it is generally acknowledged  
19 that “a point in the rat/mouse key events cascade where the pathway is biologically precluded in  
20 humans cannot be identified, in principle” (Klaunig et al., 2003; NRC, 2006).

21 In terms of human relevance and susceptibility, it is also useful to briefly review what is  
22 known about human HCC. A number of risk factors have been identified for human  
23 hepatocellular carcinoma, including ethanol consumption, hepatitis B and C virus infection,  
24 aflatoxin B1 exposure, and, more recently, diabetes and perhaps obesity (El-Serag and Rudolph,  
25 2007). However, it is also estimated that a substantial minority of HCC patients, perhaps  
26 15–50%, have no established risk factors (El-Serag and Rudolph, 2007). In addition, cirrhosis is  
27 present in a large proportion of HCC patients, but the prevalence of HCC without underlying  
28 cirrhosis, while not precisely known, is still significant, with estimates based on relatively small  
29 samples ranging from 7–54% (Fattovich et al., 2004).

30 However, despite the identification of numerous factors that appear to play a role in the  
31 human risk of HCC, the mechanisms are still largely unclear (Yeh et al., 2007). Interestingly,  
32 the observation by Leakey et al. (2003a; 2003b) that body weight significantly and strongly  
33 impacts background liver tumor rates in B6C3F1 mice parallels the observed epidemiologic

1 associations between liver cancer and obesity (review in 2007). This concordance suggests that  
2 similar pathways may be involved in spontaneous liver tumor induction between mice and  
3 humans. The extent to which TCE exposure may interact with known risk factors for HCC  
4 cannot be determined at this point, but several hypotheses can be posed based on existing data.  
5 If TCE affects some of the same pathways involved in human HCC, as suggested in the  
6 discussion of several TCE-induced effects above, then TCE exposure may lead a risk that is  
7 additive to background.

8 As discussed above, there are several parallels between the possible key events in  
9 TCE-induced liver tumors in mice and what is known about mechanisms of human HCC, though  
10 none have been experimentally tested. Altered ploidy distribution and DNA hypomethylation  
11 are commonly observed in human HCC (Calvisi et al., 2007; Lin et al., 2003; Zeppa et al., 1998).  
12 Interestingly, El-Serag and Rudolph (2007) have been suggested that the risk of HCC increases  
13 with cirrhosis in part because the liver parenchymal cells have decreased proliferative capacity,  
14 resulting in an altered milieu that promotes tumor cell proliferation. This description suggests a  
15 similarity in mode of action, though via different mechanisms, with the “negative selection”  
16 hypothesis proposed by Bull (2000) for TCE and its metabolites although for TCE changes in  
17 apoptosis and cell proliferation have not been noted or examined to such an extent to provide  
18 evidence of a similar environment. Increased ploidy decreases proliferative capacity, so that  
19 may be another mechanism through which the effects of TCE mimic the conditions thought to  
20 facilitate the induction of human HCC.

21 In sum, from the perspective of hazard characterization, the available data support the  
22 conclusion that the mode of action for TCE-induced mouse liver tumors is relevant to humans.  
23 No data suggest that any of the key events are biologically precluded in humans, and a number of  
24 qualitative parallels exist between hypotheses for the mode of action in mice and what is known  
25 about the etiology and induction of human HCC. A number of risk factors have been identified  
26 that appear to modulate the risk of human HCC, and these may also modulate the susceptibility  
27 to the effects from TCE exposure. As noted in Section E.4, TCE exposure in the human  
28 population is accompanied not only by external exposures to its metabolites, but brominated  
29 analogues of those metabolites that are also rodent carcinogens, a number of chlorinate solvents  
30 that are hepatocarcinogenic and alcohol consumption. The types of tumors and the heterogeneity  
31 of tumors induced by TCE in rodents parallel those observed in humans (see Section E.3.1.8).  
32 The pathways identified for induction of cancer in humans for cancer are similar to those for the  
33 induction of liver cancer (see Section E.3.2.1). However, while risk factors have been identified  
34 for human liver cancer that have similarities to TCE-induced effects and those of its metabolites,

1 both the mechanism for human liver cancer induction and that for TCE-induced liver  
2 carcinogenesis in rodents are not known.

#### 4.5.7.1.17. Quantitative species differences

3 As a precursor to the discussion of quantitative differences between humans and rodents  
4 and among humans, it should be noted that an adequate explanation for the difference in  
5 response for TCE-liver cancer induction between rats and mice has yet to be established or for  
6 that difference to be adequately described given the limitations in the rat database. For TCA,  
7 there is only one available long-term study in rats that, while suggestive that TCA is less potent  
8 in rats than mice, is insufficient to determine if there was a TCA-induced effect or what its  
9 magnitude may be. While some have proposed that the lower rate of TCA formation in rats  
10 relative to mice would explain the species difference, PBPK modeling suggests that the  
11 differences (three–fivefold) may be inadequate to fully explain the differences in carcinogenic  
12 potency. Moreover, inferences from comparing the effects of TCE and TCA on liver weight,  
13 using PBPK model-based estimates of TCA internal dose-metrics as a result of TCE or TCA  
14 administration, indicate that TCA is not likely to play a predominant role in hepatomegaly.  
15 Combined with the qualitative correlation between rodent hepatomegaly and  
16 hepatocarcinogenesis observed across many chemicals, this suggests that TCA similarly is not a  
17 predominant factor in TCE-induced hepatocarcinogenesis. Indeed, there are multiple lines of  
18 evidence that TCA is insufficient to account for TCE-induced tumors, including data on tumor  
19 phenotype (e.g., c-Jun immunostaining) and genotype (e.g., H-ras mutation frequency and  
20 spectrum). For DCA, only a single experiment in rats is available (reported in two publications),  
21 and although it suggests lower hepatocarcinogenic potency in rats relative to mice, its relatively  
22 low power limits the inferences that can be made as to species differences.

23 As TCA induces peroxisome proliferation in the mouse and the rat, some have suggested  
24 that difference in peroxisomal enzyme induction is responsible for the difference in susceptibility  
25 to TCA liver carcinogenesis. The study of DeAngelo et al. (1989) has been cited in the literature  
26 as providing evidence of differences between rats and mice for peroxisomal response to TCA.  
27 However, data from the most resistant strain of rat (Sprague-Dawley) have been cited in  
28 comparisons of peroxisomal enzyme effects but the Osborne-Mendel and F344 rat were not  
29 refractory and showed increased PCO activity so it is not correct to state that the rat is refractory  
30 to TCA-induction of peroxisome activity (see Section E.2.3.1.5). In addition, as discussed  
31 above, inferences based on PCO activity are limited by its high variability, even in control  
32 animals, as well as its not necessarily being predictive of the peroxisome number or cytoplasmic  
33 volume.



1 The same assumption of lower species sensitivity by measuring peroxisome proliferation  
2 has been applied to humans, as peroxisome proliferation caused by therapeutic PPAR $\alpha$  agonists  
3 such as fibrates in humans is generally lower (<twofold induction) than that observed in rodents  
4 (20- to 50-fold induction). However, as mentioned above, it is known that peroxisome  
5 proliferation is not a good predictor of potency (Marsman et al., 1988).

6 Limited data exist on the relative sensitivity of the occurrence of key events for liver  
7 tumor induction between mice and humans and among humans. Pharmacokinetic differences are  
8 addressed with PBPK modeling to the extent that data allow, so the discussion here will  
9 concentrate on pharmacodynamic differences. Most striking is the difference in “background”  
10 rates of liver tumors. Data from NTP indicates that control B6C3F1 mice in 2-year bioassays  
11 have a background incidence of hepatocellular carcinomas of 26% in males and 10% in females,  
12 with higher incidences for combined hepatocellular adenomas and carcinomas (Maronpot, 2007).  
13 However, as discussed above, Leakey et al. (2003a; 2003b) report that the background incidence  
14 rates are very dependent on the weight of the mice. By contrast, the estimated lifetime risk of  
15 liver and biliary tract cancer in the United States (about 75% of which are hepatocellular  
16 carcinomas) is 0.97% for men and 0.43% for women (Ries et al., 2008). However, regions of the  
17 world where additional risk factors (hepatitis infection, aflatoxin exposure) have high  
18 prevalence have liver cancer incidences up to more than sixfold greater than the United States  
19 (Ferlay et al., 2004). Therefore, one possible quantitative difference that can be flagged for use  
20 in dose-response assessment is the background rate of liver tumors between species.  
21 Biologically-based dose-response modeling by Chen (2000) suggested that the data were  
22 consistent with a purely promotional model in which potency would be proportional to  
23 background tumor incidence. However, it is notable that male Swiss mice, which have lower  
24 background liver tumor rates than the B6C3F1 strain, were also positive in one long-term  
25 bioassay (Maltoni et al., 1986).

26 Similarly, in terms of intraspecies susceptibility, to the extent that TCE may  
27 independently promote pre-existing initiated cells, it can be hypothesized that those with greater  
28 risk for developing HCC due to one more of the known risk factors would have a proportional  
29 increase in the any contributions from TCE exposure. In addition, in both humans and mice,  
30 males appear to be at increased risk of liver cancer, possibly due to sexually dimorphism in  
31 inflammatory responses (Lawrence et al., 2007; Naugler et al., 2007; Rakoff-Nahoum and  
32 Medzhitov, 2007), suggesting that men may also be more susceptible to TCE-induced liver  
33 tumorigenesis than women. It has been observed that human HCC is highly heterogeneous  
34 histologically, but within patients and between patients, studies are only beginning to distinguish

1 the different pathways that may be responsible for this heterogeneity (Chen et al., 2002;  
2 Feitelson et al., 2002; Yeh et al., 2007).

3 Appropriate quantitative data are generally lacking on interspecies differences in the  
4 occurrence of most other proposed key events, although many have argued that there are  
5 significant quantitative differences between rodents and humans related to PPAR $\alpha$  activation  
6 (Klaunig et al., 2003; NRC, 2006). For instance, it has been suggested that lower levels of  
7 PPAR $\alpha$  receptor in human hepatocytes relative to rodent hepatocytes contributes to lower human  
8 sensitivity (Klaunig et al., 2003; Palmer et al., 1998; Tugwood et al., 1996). However, out of a  
9 small sample of human livers ( $n = 6$ ) show similar protein levels to mice (Walgren et al., 2000b).  
10 Another proposed species difference has been ligand affinity, but while transactivation assays  
11 showed greater affinity of Wy-14643 and perfluorooctanoic acid for rodent relative to human  
12 PPAR $\alpha$ , they showed TCA and DCA had a similar affinities between species (Maloney and  
13 Waxman, 1999). Furthermore, it is not clear that receptor-ligand kinetics (capacity and affinity)  
14 are rate-limiting for eliciting hepatocarcinogenic effects, as it is known that maximal receptor  
15 occupation is not necessary for a maximal receptor mediated response (Stephenson, 1956) (see  
16 also review by Danhof et al., 2007).

17 There is also limited in vivo and in vitro data suggesting that increases in cell  
18 proliferation mediated by PPAR $\alpha$  agonists are diminished in humans and other primates relative  
19 to rodents (Hoivik et al., 2004; Klaunig et al., 2003; NRC, 2006). However, Walgren et al.  
20 (2000a) reported that TCA and DCA were not mitogenic in either human or rodent hepatocytes  
21 in vitro. Furthermore, TCE, TCA, and DCA all induce only transient increases in cell  
22 proliferation, so the relevance to TCE of interspecies differences from PPAR $\alpha$  agonists that to  
23 produce sustained proliferation, such as Wy-14643, is not clear. In addition, comparisons  
24 between primate and rodent models should take into account the differences in the ability to  
25 respond to any mitogenic stimulation (see Section E.3.2). Primate and human liver respond  
26 differently (and much more slowly) to a stimulus such as partial hepatectomy.

27 Recent studies in “humanized” mice (PPAR $\alpha$ -null mice in which a human PPAR $\alpha$  gene  
28 was subsequently inserted and expressed in the liver) reported that treatment with a PPAR $\alpha$   
29 agonist lead to greatly lower incidence of liver tumors as compared to wild-type mice (Morimura  
30 et al., 2006). However, these experiments were performed with WY-14643 at a dose causing  
31 systemic toxicity (reduced growth and survival), had a duration of less than 1 year, and involved  
32 a limited number of animals. In addition, because liver tumors in mice at less than 1 year are  
33 extremely rare, the finding a one adenoma in WY-14643-treated humanized mice suggests  
34 carcinogenic potential that could be further realized with continued treatment (Keshava and  
35 Caldwell, 2006). In addition, Yang et al. (2007) recently noted that let-7C, a microRNA

1 involved in cell growth and thought to be a regulatory target of PPAR $\alpha$  (Shah, 2008), was  
2 inhibited by Wy-14643 in wild-type mice, but not in “humanized mice” in which had human  
3 PPAR $\alpha$  was expressed throughout the body on a PPAR $\alpha$ -null background. However, these  
4 humanized mice had about a 20-fold higher baseline expression of let-7C, as reported in control  
5 mice, potentially masking any treatment effects. More generally, it is not known to what extent  
6 PPAR $\alpha$ -related events are rate-limiting in TCE-induced liver tumorigenesis, for which multiple  
7 pathways appear to be operative. So even if quantitative differences mediated by PPAR $\alpha$  were  
8 well estimated, they would not be directly usable for dose-response assessment in the absence of  
9 way to integrate the contributions from the different pathways.

10 In sum, the only quantitative data and inter- and intraspecies susceptibility suitable for  
11 consideration in dose-response assessment are differences background liver tumor risk. These  
12 may modulate the effects of TCE if relative risk, rather than additional risk, is the appropriate  
13 common inter- and intraspecies metric. However, the extent to which relative risk would provide  
14 a more accurate estimate of human risk is unknown.

#### 4.6. IMMUNOTOXICITY AND CANCERS OF THE IMMUNE SYSTEM

15 Chemical exposures may result in a variety of adverse immune-related effects, including  
16 immunosuppression (decreased host resistance), autoimmunity, and allergy-hypersensitivity, and  
17 may result in specific diseases such as infections, systemic or organ-specific autoimmune  
18 diseases, or asthma. Cell-mediated immune response, such as activation of macrophages, natural  
19 killer cells, and cytokine production, can also influence a broader range of diseases, such as  
20 cancer. Measures of immune function (e.g., T-cell counts, immunoglobulin [Ig] E levels,  
21 specific autoantibodies, cytokine levels) may provide evidence of an altered immune response  
22 that precedes the development of clinically expressed diseases. The first section of this section  
23 discusses effects relating to immunotoxicity, including risk of autoimmune diseases, allergy and  
24 hypersensitivity, measures of altered immune response, and lymphoid cancers. Studies  
25 pertaining to effects in humans are presented first, followed by a section discussing relevant  
26 studies in animals. The second section of this section discusses evidence pertaining to  
27 trichloroethylene in relation to lymphoid tissue cancers, including childhood leukemia.

## 4.6.1. Human Studies

### 4.6.1.1.1. Noncancer Immune-Related Effects

#### 4.6.1.1.2. Immunosuppression, asthma, and allergies

1           In 1982, Lagakos et al. conducted a telephone survey of residents of Woburn,  
2 Massachusetts, collecting information on residential history and history of 14 types of medically  
3 diagnosed conditions (Lagakos et al., 1986). The survey included 4,978 children born since  
4 1960 who lived in Woburn before age 19. Completed surveys were obtained from  
5 approximately 57% of the town residences with listed phone numbers. Two of the wells  
6 providing the town's water supply from 1964–1979 had been found to be contaminated with a  
7 number of solvents, including tetrachloroethylene (21 ppb) and trichloroethylene (267 ppb) (as  
8 cited in Lagakos et al., 1986). Lagakos et al. used information from a study by the  
9 Massachusetts Department of Environmental Quality and Engineering to estimate the  
10 contribution of water from the two contaminated wells to the residence of each participant, based  
11 on zones within the town receiving different mixtures of water from various wells, for the period  
12 in which the contaminated wells were operating. This exposure information was used to  
13 estimate a cumulative exposure based on each child's length of residence in Woburn. A higher  
14 cumulative exposure measure was associated with conditions indicative of immunosuppression  
15 (e.g., bacterial or viral infections) or hypersensitivity (e.g., asthma). In contrast, a recent study  
16 using the National Health and Nutrition Examination Survey data collected from 1999–2000 in a  
17 representative sample of the U.S. population ( $n = 550$ ) did not find an association between TCE  
18 exposure and self-report of a history of physician-diagnosed asthma (OR: 0.94, 95% CI: 0.77,  
19 1.14) (Arif and Shah, 2007). TCE exposure, as well as exposure to 9 other volatile organic  
20 compounds, was determined through a passive monitor covering a period of 48–72 hours. No  
21 clear trend was seen with self-reported wheeze episodes (OR: 1.29, 95% CI: 0.98, 1.68 for one to  
22 two episodes; OR: 0.21, 95% CI: 0.04, 10.05 for three or more episodes in the past 12 months).

23           Allergy and hypersensitivity, as assessed with measures of immune system parameters or  
24 immune function tests (e.g., atopy) in humans, have not been extensively studied with respect to  
25 the effects of trichloroethylene (see Table 4-69). Lehmann et al. reported data pertaining to  
26 immunoglobulin E (IgE) levels and response to specific antigens in relation to indoor levels of  
27 volatile organic compounds among children (age 36 months) selected from a birth cohort study  
28 in Leipzig, Germany (Lehmann et al., 2001). Enrollment into the birth cohort occurred between  
29 1995 and 1996. The children in this allergy study represent a higher-risk group for development  
30 of allergic disease, with eligibility criteria that were based on low birth weight (between 1,500

1 and 2,500 g), or cord blood IgE greater than 0.9 kU/L with double positive family history of  
2 atopy. These eligibility criteria were met by 429 children; 200 of these children participated in  
3 the allergy study described below, but complete data (IgE and volatile organic compound  
4 measurements) were available for only 121 of the study participants. Lehmann et al. measured  
5 26 volatile organic compounds via passive indoor sampling in the child's bedroom for a period  
6 of 4 weeks around the age of 36 months. The median exposure of trichloroethylene was 0.42  
7  $\mu\text{g}/\text{m}^3$  (0.17  $\mu\text{g}/\text{m}^3$  and 0.87  $\mu\text{g}/\text{m}^3$  for the 25<sup>th</sup> and 75<sup>th</sup> percentiles, respectively). Blood samples  
8 were taken at the 36-month-study examination and were used to measure the total IgE and  
9 specific IgE antibodies directed to egg white, milk, indoor allergens (house dust mites, cats, and  
10 molds), and outdoor allergens (timothy-perennial grass and birch trees). There was no  
11 association between trichloroethylene exposure and any of the allergens tested in this study,  
12 although some of the other volatile organic compounds (e.g., toluene, 4-ethyltoluene) were  
13 associated with elevated total IgE levels and with sensitization to milk or eggs.  
14

#### **4.6.1.1.3. Generalized hypersensitivity skin diseases, with or without hepatitis**

15 Occupational exposure to trichloroethylene has been associated with a severe,  
16 generalized skin disorder that is distinct from contact dermatitis in the clinical presentation of the  
17 skin disease (which often involves mucosal lesions), and in the accompanying systemic effects  
18 that can include lymphadenopathy, hepatitis, and other organ involvement. Kamijima et al.  
19 recently reviewed case reports describing 260 patients with trichloroethylene-related generalized

**Table 4-69. Studies of immune parameters (IgE antibodies and cytokines) and trichloroethylene in humans**

<b>Parameter, source of data</b>	<b>Results</b>	<b>Reference, location, diagnosis period, sample size, age</b>
<b>IgE antibodies</b> blood sample, indoor air sampling of 28 volatile organic chemicals in child's bedroom	Trichloroethylene exposure not associated with sensitization to indoor or outdoor allergens	Lehmann et al. (2001) Germany. 1997–1999. <i>n</i> = 121 36-mo old children
<b>Cytokine secreting CD3+ T-cell populations</b> cord blood, indoor air sampling of 28 volatile organic chemicals in child's bedroom 4 wk after birth	In CD3+ cord blood cells, some evidence of association between increasing trichloroethylene levels and decreased IL-4 >75 <sup>th</sup> percentile OR: 0.6 (95% CI: 0.2, 2.1), <25 <sup>th</sup> percentile OR 4.4 (95% CI: 1.1, 17.8) increased IFN- $\gamma$ >75 <sup>th</sup> percentile OR: 3.6 (95% CI: 0.9, 14.9) <25 <sup>th</sup> percentile OR: 0.7 (95% CI: 0.2, 2.2) Similar trends not seen with tumor necrosis factor- $\alpha$ or IL-2	Lehmann et al. (2002) Germany. 1995–1996. <i>n</i> = 85 newborns
<b>Cytokine secreting CD3+ and CD8+ T-cell populations</b> blood sample, indoor air sampling of 28 volatile organic chemicals in child's bedroom	Trichloroethylene exposure not associated with percentages of IL-4 CD3+ or IFN- $\gamma$ CD8+ T-cells	Lehmann et al. (2001) Germany. 1995–1999. <i>n</i> = 200 36-mo old children
<b>Cytokine concentration—serum</b> urine sample (trichloroacetic acid concentration), blood sample, questionnaire (smoking history, age, residence), workplace TCE measures (personal samples, 4 exposed and 4 nonexposed workers)	Nonexposed workers similar to office controls for all cytokine measures. Compared to nonexposed workers, the trichloroethylene exposed workers had decreased IL-4 (mean 3.9 vs. 8.1 pg/mL) increased IL-2 (mean 798 vs. 706 pg/mL) increased IFN- $\gamma$ (mean 37.1 vs. 22.9 pg/mL)	Iavicoli et al. (2005) Italy. <i>n</i> = 35 printers using TCE, 30 nonexposed workers (in same factory, did not use or were not near TCE), 40 office worker controls. All men. Mean age ~33 yr

IFN = interferon, IL = interleukin.

1 skin disorders (Kamijima et al., 2007). Six of the patients were from the United States or  
2 Europe, with the remainder occurring in China, Singapore, Philippines, and other Asian  
3 countries. One study in Guangdong province, in southeastern China, included more than 100 of  
4 these cases in a single year (Huang et al., 2002). Kamijima et al. categorized the case  
5 descriptions as indicative of hypersensitivity syndrome ( $n = 124$ ) or a variation of erythema  
6 multiforme, Stevens-Johnson syndrome, and toxic epiderma necrolysis ( $n = 115$ ), with 21 other  
7 cases unclassified in either category. The fatality rate, approximately 10%, was similar in the  
8 two groups, but the prevalence of fever and lymphadenopathy was higher in the hypersensitivity  
9 syndrome patients. Hepatitis was seen in 92–94% of the multiforme, Stevens-Johnson  
10 syndrome, and toxic epiderma necrolysis patients, but the estimates within the hypersensitivity  
11 syndrome group were more variable (46–94%) (Kamijima et al., 2007).

12 Some of the case reports reviewed by Kamijima et al. provided information on the total  
13 number of exposed workers, working conditions, and measures of exposure levels. From the  
14 available data, generalized skin disease within a worksite occurred in 0.25–13% of workers in  
15 the same location, doing the same type of work (Kamijima et al., 2007). The measured  
16 concentration of trichloroethylene ranged from  $<50 \text{ mg/m}^3$  to more than  $4,000 \text{ mg/m}^3$ , and  
17 exposure scenarios included inhalation only and inhalation with dermal exposures. Disease  
18 manifestation generally occurred within 2–5 weeks of initial exposure, with some intervals up to  
19 3 months. Most of the reports were published since 1995, and the geographical distribution of  
20 cases reflects the newly industrializing areas within Asia.

21 Kamijima and colleagues recently conducted an analysis of urinary measures of  
22 trichloroethylene metabolites (trichloroacetic acid and trichloroethanol) in 25 workers  
23 hospitalized for hypersensitivity skin disease in 2002 (Kamijima et al., 2008). Samples taken  
24 within 15 days of the last exposure to trichloroethylene exposure were available for 19 of the  
25 25 patients, with a mean time of 8.4 days. Samples from the other patients were not used in the  
26 analysis because the half life of urinary trichloroacetic acid is 50–100 hours. In addition,  
27 3–6 healthy workers doing the same type of work in the factories of the affected worker, and  
28 2 control workers in other factories not exposed to trichloroethylene were recruited in  
29 2002–2003 for a study of breathing zone concentration of volatile organochlorines and urinary  
30 measures of trichloroethylene metabolites. Worksite measures of trichloroethylene concentration  
31 were also obtained. Adjusting for time between exposure and sample collection, mean urinary  
32 concentration at the time of last exposure among the 19 patients was 206 mg/mL for  
33 trichloroacetic acid. Estimates for trichloroethanol were not presented because of the shorter  
34 half-life for this compound. Urinary trichloroacetic acid levels in the healthy exposed workers  
35 varied among the 4 factories, with means ( $\pm$ standard deviation [SD]) of 41.6 ( $\pm$ 18.0),

1 131 ( $\pm 90.2$ ), 180 ( $\pm 92$ ), and 395 ( $\pm 684$ ). The lower values were found in a factory in which the  
2 degreasing machine had been partitioned from the workers after the illnesses had occurred.  
3 Trichloroethylene concentrations (personal time-weighted averages) at the factories of the  
4 affected workers ranged from 164–2,330 mg/m<sup>3</sup> (30–431 ppm). At the two factories with no  
5 affected workers in the past 3 years, the mean personal time-weighted average trichloroethylene  
6 concentrations were 44.9 mg/m<sup>3</sup> (14 ppm) and 1,803 mg/m<sup>3</sup> (334 ppm). There was no  
7 commonality of additives or impurities detected among the affected factories that could explain  
8 the occurrence of the hypersensitivity disorder.

9 To examine genetic influences on disease risk, Dai et al. conducted a case-control study  
10 of 111 patients with trichloroethylene-related severe generalized dermatitis and  
11 152 trichloroethylene-exposed workers who did not develop this disease (Dai et al., 2004).  
12 Patients were recruited from May 1999 to November 2003 in Guangdong Province, and were  
13 employed in approximately 80 electronic and metal-plating manufacturing plants. Initial  
14 symptoms occurred within 3 months of exposure. The comparison group was drawn from the  
15 same plants as the cases, and had worked for more than 3 months without development of skin or  
16 other symptoms. Mean age in both groups was approximately 23 years. A blood sample was  
17 obtained from study participants for genotyping of TNF- $\alpha$ , TNF- $\beta$ , and interleukin (IL)-4  
18 genotypes. The genes were selected based on the role of TNF and of interleukin-4 in  
19 hypersensitivity and inflammatory responses. The specific analyses included two  
20 polymorphisms in the promoter region of TNF- $\alpha$  (G  $\rightarrow$  A substitution at position -308)  
21 designated as TNFAII, with wild-type designated TNFAI; and a G  $\rightarrow$  A substitution at position  
22 -238), a polymorphism at the first intron on TNF- $\beta$ , and a polymorphism in the promoter region  
23 of IL-4 (C  $\rightarrow$  T substitution at -590). There was no difference in the frequency of the  
24 TNF- $\alpha$ <sup>-238</sup>, TNF- $\beta$ , or IL-4 polymorphisms between cases and controls, but the wild-type  
25 TNF- $\alpha$ <sup>-308</sup> genotype was somewhat more common among cases (TNF A I/I genotype 94% in  
26 cases and 86% in controls).

27 Kamijima et al. note the similarities, particular with respect to specific skin  
28 manifestations, of the case presentations of trichloroethylene-related generalized skin diseases to  
29 conditions that have been linked to specific medications (e.g., carbamazepine, allopurinol,  
30 antibacterial sulfonamides), possibly in conjunction with reactivation of specific latent herpes  
31 viruses (Kamijima et al., 2007). A previous review by these investigators discusses insights with  
32 respect to drug metabolism that may be useful in developing hypotheses regarding susceptibility  
33 to trichloroethylene-related generalized skin disorders (Nakajima et al., 2003). Based on  
34 consideration of metabolic pathways and intermediaries, variability in CYP2E1,  
35 UDP-glucuronyltransferase, glutathione-S transferase, and N-acetyl transferase (NAT) activities



1 could be hypothesized to affect the toxicity of trichloroethylene. NAT2 is most highly expressed  
2 in liver, and the “slow” acetylation phenotype (which arises from a specific mutation) has been  
3 associated with adverse effects of medications, including drug-induced lupus (Lemke and  
4 McQueen, 1995) and hypersensitivity reactions (Spielberg, 1996). There are limited data  
5 pertaining to genetic or other sources of variability in these enzymes on risk of trichloroethylene-  
6 related generalized skin diseases, however. In a study in Guangdong province, CYP1A1,  
7 GSTM1, GSTP1, GSTT1, and NAT2 genotypes in 43 cases of trichloroethylene-related  
8 generalized skin disease were compared to 43 healthy trichloroethylene-exposed workers (Huang  
9 et al., 2002). The authors reported that the NAT2 slow acetylation genotype was associated with  
10 disease, but the data pertaining to this finding were not presented.

11

#### 4.6.1.1.4. Cytokine profiles and lymphocyte subsets

12 Cytokines are produced by many of the immune regulatory cells (e.g., macrophages,  
13 dendritic cells), and have many different effects on the immune system. The T-helper Type 1  
14 (Th1) cytokines, are characterized as “pro-inflammatory” cytokines, and include TNF- $\alpha$  and  
15 interferon (IFN)- $\gamma$ . Although this is a necessary and important part of the innate immune  
16 response to foreign antigens, an aberrant pro-inflammatory response may result in a chronic  
17 inflammatory condition and contribute to development of scarring or fibrotic tissue, as well as to  
18 autoimmune diseases. Th2 cytokines are important regulators of humoral (antibody-related)  
19 immunity. IL-4 stimulates production of IgE and thus influences IgE-mediated effects such as  
20 allergy, atopy, and asthma. Th2 cytokines can also act as “brakes” on the inflammatory  
21 response, so the balance between different types of cytokine production is also important with  
22 respect to risk of conditions resulting from chronic inflammation. Several studies have examined  
23 cytokine profiles in relation to occupational or environmental TCE exposure (see Table 4-69).

24 The 2001 Lehmann et al. study of 36-month old children (described above) also included  
25 a blood sample taken at the 3-year study visit, which was used to determine the percentages of  
26 specific cytokine producing T-cells in relation to the indoor volatile organic compounds  
27 exposures measured at birth. There was no association between trichloroethylene exposure and  
28 either IL-4 CD3+ or IFN- $\gamma$  CD8+ T-cells (Lehmann et al., 2001).

29 Another study by Lehmann et al. examined the relationship between indoor exposures to  
30 volatile organic compounds and T-cell subpopulations measured in cord blood of newborns  
31 (Lehmann et al., 2002). The study authors randomly selected 85 newborns (43 boys and  
32 42 girls) from a larger cohort study of 997 healthy, full-term babies, recruited between 1997 and  
33 1999 in Germany. Exclusion criteria included a history in the mother of an autoimmune disease

1 or infectious disease during the pregnancy. Twenty-eight volatile organic compounds were  
2 measured via passive indoor sampling in the child's bedroom for a period of 4 weeks after the  
3 birth (a period which is likely to reflect the exposures during the prenatal period close to the time  
4 of delivery). The levels were generally similar or slightly higher than the levels seen in the  
5 previous study using samples from the bedrooms of the 36-month-old children. The highest  
6 levels of exposure were seen for limonene (median 24.3  $\mu\text{g}/\text{m}^3$ ),  $\alpha$ -pinene (median 19.3  $\mu\text{g}/\text{m}^3$ )  
7 and toluene (median 18.3  $\mu\text{g}/\text{m}^3$ ), and the median exposure of trichloroethylene was 0.6  $\mu\text{g}/\text{m}^3$   
8 (0.2  $\mu\text{g}/\text{m}^3$  and 1.0  $\mu\text{g}/\text{m}^3$  for the 25<sup>th</sup> and 75<sup>th</sup> percentiles, respectively). Flow cytometry was  
9 used to measure the presence of CD3 T-cells obtained from the cord blood labeled with  
10 antibodies against IFN- $\gamma$ , tumor necrosis factor- $\alpha$ , IL-2, and IL-4. There was some evidence of a  
11 decreased level of IL-2 with higher trichloroethylene exposure in the univariate analysis, with  
12 median percentage of IL-2 cells of 46.1 and 33.0% in the groups that were below the 75<sup>th</sup>  
13 percentile and above the 75<sup>th</sup> percentile of trichloroethylene exposure, respectively. In analyses  
14 adjusting for family history of atopy, gender and smoking history of the mother during  
15 pregnancy, there was little evidence of an association with either IL-2 or IFN- $\gamma$ , but there was a  
16 trend of increasing trichloroethylene levels associated with decreased IL-4 and increased IFN- $\gamma$ .

17 Iavicoli et al. examined cytokine levels in 35 trichloroethylene-exposed workers (Group  
18 A) from a printing area of a factory in Italy. Their work involved use of trichloroethylene in  
19 degreasing (Iavicoli et al., 2005). Two comparison groups were included. Group B consisted of  
20 30 other factory workers who were not involved in degreasing activities and did not work near  
21 this location, and Group C consisted of 40 office workers at the factory. All study participants  
22 were male and had worked at their present position for at least 3 years, and all were considered  
23 healthy. Personal breathing zone air samples from four workers in Group A and four workers in  
24 Group B were obtained in three consecutive shifts (24 total samples) to determine air  
25 concentration of trichloroethylene. A urine sample was obtained from each Group A and Group  
26 B worker (end of shift at end of work week) for determination of trichloroacetic acid  
27 concentrations (corrected for creatinine), and blood samples were collected for assessment of  
28 IL-2, IL-4, and IFN- $\gamma$  concentrations in serum using enzyme-linked immunosorbent assays.  
29 Among exposed workers, the mean trichloroethylene concentration was approximately 35  $\text{mg}/\text{m}^3$   
30 ( $30.75 \pm \text{SD } 9.9$ ,  $37.75 \pm 23.0$ , and  $36.5 \pm 8.2 \text{ mg}/\text{m}^3$  in the morning, evening, and night shifts,  
31 respectively). The urinary trichloroacetic acid concentrations were much higher in exposed  
32 workers compared with nonexposed workers (mean  $\pm$  SD, Group A  $13.3 \pm 5.9 \text{ mg}/\text{g}$  creatinine;  
33 Group B  $0.02 \pm 0.02 \text{ mg}/\text{g}$  creatinine). There was no difference in cytokine levels between the  
34 two control groups, but the exposed workers differed significantly (all  $p$ -values  $< 0.01$  using  
35 Dunnett's test for multiple comparisons) from each of the two comparison groups. The observed

1 differences were a decrease in IL-4 levels (mean 3.9, 8.1, and 8.1 pg/mL for Groups A, B, and C,  
2 respectively), and an increase in IL-2 levels (mean 798, 706, and 730 pg/mL for Groups A, B,  
3 and C, respectively) and in IFN- $\gamma$  levels (mean 37.1, 22.9, and 22.8 pg/mL for Groups A, B, and  
4 C, respectively).

5 The available data from these studies (Iavicoli et al., 2005; Lehmann et al., 2001;  
6 Lehmann et al., 2002) provide some evidence of an association between increased  
7 trichloroethylene exposure and modulation of immune response involving an increase in pro-  
8 inflammatory cytokines (IL-2, IFN- $\gamma$ ) and a decrease in Th2 (allergy-related) cytokines (e.g., IL-  
9 4). These observations add support to the influence of trichloroethylene in immune-related  
10 conditions affected by chronic inflammation.

11 Lan et al. (2010) examined lymphocyte subsets among 80 TCE-exposed workers and  
12 96 controls in Guangdong, China. Six factories using TCE for cleaning metals, optical lenses, or  
13 circuit boards were included in this study. These factories did not use other solvents (benzene,  
14 styrene, ethylene oxide, formaldehyde, or epichlorohydrin), based on an exposure screening  
15 using Dräger tubes and 3M Badges. Eighty workers from these factories and 96 unexposed  
16 controls (frequency matched by sex and 5-year age groups to controls) from clothes  
17 manufacturers, a food production factory, and a hospital, were included in the study. The study  
18 was conducted in 2006. Study participants provided a blood sample, buccal cells, postshift and  
19 overnight urine samples, and completed a questionnaire with demographic, alcohol and smoking  
20 history, and occupational history data. A blood sample was used for a complete blood count and  
21 differential lymphocyte subset analysis. At the time of the blood draw, a clinical examination,  
22 including measurement of height and weight, and symptoms of recent respiratory infection  
23 (which could affect the differential blood cell counts) was conducted. TCE monitoring was  
24 conducted using full-shift personal air exposure measurements. The median level of exposure,  
25 based on the mean of two measurements taken for each participant in the month before the blood  
26 draw, among the 80 TCE-exposed workers was 12 ppm. The analysis used this level to  
27 categorize workers into high ( $\geq 12$  ppm; mean 38 ppm) and low ( $< 12$  ppm; mean 5 ppm)  
28 exposures. Among the controls, the mean TCE exposure was  $< 0.03$  ppm. The total number of  
29 lymphocytes, T cells, CD4+ T cells, CD8+ T cells, B cells and natural killer (NK) cells was  
30 significantly lower among TCE-exposed workers compared with controls, with the largest  
31 decrease seen in the higher exposure group. For example, the age- and sex-adjusted lymphocyte  
32 count was 2,154, 2,012, and 1,671 cells/ $\mu$ L blood in the controls,  $< 12$  and  $\geq 12$  ppm groups,  
33 respectively (trend  $p = < 0.0001$ ). Plasma concentrations of soluble CD27 and CD30, two  
34 costimulators involved in the regulation of T cells, were also decreased in both exposure groups  
35 compared with controls. Similar patterns were seen when limited to the 77 workers with

1 exposure levels <100 ppm, and when limited to the 60 workers with exposure levels <25 ppm.  
2 Granulocytes, monocytes and platelet counts did not differ by exposure. The authors note that  
3 the immunosuppression and decreased lymphocyte activation seen in this study provide support  
4 the biological plausibility of a role of TCE exposure in NHL.  
5

#### 4.6.1.1.5. Autoimmune disease

##### 4.6.1.1.5.1. Disease clusters and geographic-based studies

6 Reported clusters of diseases have stimulated interest in environmental influences on  
7 systemic autoimmune diseases. These descriptions include investigations into reported clusters  
8 of systemic lupus erythematosus (Balluz et al., 2001; Dahlgren et al., 2007) and Wegener  
9 granulomatosis (Albert et al., 2005). Wegener granulomatosis, an autoimmune disease involving  
10 small vessel vasculitis, usually with lung or kidney involvement, is a very rare condition, with an  
11 incidence rate of 3–14 per million per year (Mahr et al., 2006). Trichloroethylene was one of  
12 several ground water contaminants identified in a recent study investigating a cluster of seven  
13 cases of Wegener granulomatosis around Dublin, Pennsylvania. Because of the multiple  
14 contaminants, it is difficult to attribute the apparent disease cluster to any one exposure.

15 In addition to the study of asthma and infectious disease history among residents of  
16 Woburn, Massachusetts (Lagakos et al., 1986) (see Section 4.6.1.1.1), Byers et al. provide data  
17 pertaining to immune function from 23 family members of leukemia patients in Woburn,  
18 Massachusetts (Byers et al., 1988). Serum samples were collected in May and June of 1984 and  
19 in November of 1985 (several years after 1979, when the contaminated wells had been closed).  
20 Total lymphocyte counts and lymphocyte subpopulations (CD3, CD4, and CD8) and the  
21 CD4/CD8 ratio were determined in these samples, and in samples from a combined control  
22 group of 30 laboratory workers and 40 residents of Boston selected through a randomized  
23 probability area sampling process. The study authors also assessed the presence of antinuclear  
24 antibodies (ANA) or other autoantibodies (antismooth muscle, antiovarian, antithyroglobulin,  
25 and antimicrosomal antibodies) in the family member samples and compared the results with  
26 laboratory reference values. The age distribution of the control group, and stratified analyses by  
27 age, are not provided. The lymphocyte subpopulations (CD3, CD4, and CD8) were higher and  
28 the CD4/CD8 ratio was lower in the Woburn family members compared to the controls in both  
29 of the samples taken in 1984. In the 1985 samples, however, the lymphocyte subpopulation  
30 levels had decreased and the CD4/CD8 ratio had increased; the values were no longer  
31 statistically different from the controls. None of the family member serum samples had

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 antithyroglobulin or antimicrobial antibodies, but 10 family-member serum samples (43%) had  
2 ANA (compared to <5% expected based on the reference value). Because the initial blood  
3 sample was taken in 1984, it is not possible to determine the patterns at a time nearer to the time  
4 of the exposure. The coexposures that occurred also make it difficult to infer the exact role of  
5 trichloroethylene in any alterations of the immunologic parameters.

6 Kilburn and Warshaw reported data from a study of contamination by metal-cleaning  
7 solvents (primarily trichloroethylene) and heavy metals (e.g., chromium) of the aquifer of the  
8 Santa Cruz River in Tucson, Arizona (Kilburn and Warshaw, 1992b). Exposure concentrations  
9 above 5 ppb (6–500 ppb) had been documented in some of the wells in this area. A study of  
10 neurological effects was undertaken between 1986 and 1989 (Kilburn and Warshaw, 1993b), and  
11 two of the groups within this larger study were also included in a study of symptoms relating to  
12 systemic lupus erythematosus. Residents of Tucson ( $n = 362$ ) were compared to residents of  
13 southwest Arizona ( $n = 158$ ) recruited through a Catholic parish. The Tucson residents were  
14 selected from the neighborhoods with documented water contamination (>5 ppb  
15 trichloroethylene for at least 1 year between 1957 and 1981). Details of the recruitment strategy  
16 are not clearly described, but the process included recruitment of patients with lupus or other  
17 rheumatic diseases (Kilburn and Warshaw, 1992b, 1993b). The prevalence of some self-reported  
18 symptoms (malar rash, arthritis/arthritis, Raynaud syndrome, skin lesions, and seizure or  
19 convulsion) was significantly higher in Tucson, but there was little difference between the groups  
20 in the prevalence of oral ulcers, anemia, low white blood count or low platelet count, pleurisy,  
21 alopecia, or proteinuria. The total number of symptoms reported was higher in Tucson than in  
22 the other southwest Arizona residents (14.3 vs. 6.4% reported four or more symptoms,  
23 respectively). Low-titer (1:80) ANA were seen in 10.6 and 4.7% of the Tucson and other  
24 Arizona residents, respectively ( $p = 0.013$ ). However, since part of the Tucson study group was  
25 specifically recruited based on the presence of rheumatic diseases, it is difficult to interpret these  
26 results.

#### 4.6.1.1.5.2. Case-control studies

28 Interest in the role of organic solvents, including trichloroethylene, in autoimmune  
29 diseases was spurred by the observation of a scleroderma-like disease characterized by skin  
30 thickening, Raynaud's phenomenon, and acroosteolysis and pulmonary involvement in workers  
31 exposed to vinyl chloride (Gama and Meira, 1978). A case report in 1987 described the  
32 occurrence of a severe and rapidly progressive case of systemic sclerosis in a 47-year-old woman

1 who had cleaned X-ray tubes in a tank of trichloroethylene for approximately 2.5 hours (Lockey  
2 et al., 1987).

3 One of the major impediments to autoimmune disease research is the lack of disease  
4 registries, which make it difficult to identify incident cases of specific diseases. There are no  
5 cohort studies of the incidence of autoimmune diseases in workers exposed to trichloroethylene.  
6 Most of the epidemiologic studies of solvents and autoimmune disease rely on general measures  
7 of occupational exposures to solvents, organic solvents, or chlorinated solvents exposures. A  
8 two- to threefold increased risk of systemic sclerosis (scleroderma) (Aryal et al., 2001; Garabrant  
9 et al., 2003; Maitre et al., 2004), **rheumatoid arthritis** (Lundberg et al., 1994; [Sverdrup et al.,  
10 2005](#)), undifferentiated connective tissue disease (Lacey et al., 1999), and antineutrophil-  
11 cytoplasmic antibody (ANCA)-related vasculitis (Beaudreuil et al., 2005; Lane et al., 2003) has  
12 generally been seen in these studies, but there was little evidence of an association between  
13 solvent exposure and systemic lupus erythematosus in two recent case-control studies (Cooper et  
14 al., 2004; Finckh et al., 2006).

15 Two case-control studies of scleroderma (Bovenzi et al., 2004; Maitre et al., 2004) and  
16 two of rheumatoid arthritis (Olsson et al., 2004; Olsson et al., 2000) provide data concerning  
17 solvent exposure that occurred among metal workers or in jobs that involved cleaning metal (i.e.,  
18 types of jobs which were likely to use trichloroethylene as a solvent). There was a twofold  
19 increased risk among male workers in the two studies of rheumatoid arthritis from Sweden  
20 (Olsson et al., 2004; Olsson et al., 2000). The results from the smaller studies of scleroderma  
21 were more variable, with no exposed cases seen in one study with 93 cases and 206 controls  
22 (Maitre et al., 2004), and an odds ratio of 5.2 (95% CI: 0.7, 37) seen in a study with 56 cases and  
23 171 controls (Bovenzi et al., 2004).

24 Five other case-control studies provide data specifically about trichloroethylene exposure,  
25 based on industrial hygienist review of job history data (see Table 4-70). Three of these studies  
26 are of scleroderma (Diot et al., 2002; Garabrant et al., 2003; Nietert et al., 1998), one is of  
27 undifferentiated connective tissue disease (Lacey et al., 1999), and one is of small vessel  
28 vasculitides involving ANCAs (Beaudreuil et al., 2005).

29 These studies included some kind of expert review of job histories, but only two studies  
30 included a quantification of exposure (e.g., a cumulative exposure metric, or a “high” exposure  
31 group) (Diot et al., 2002; Nietert et al., 1998). Most of the studies present data stratified by sex,  
32 and as expected, the prevalence of exposure (either based on type of job or on industrial  
33 hygienist assessment) is considerably lower in women compared with men. In men the studies  
34 generally reported odds ratios between 2.0 and 8.0, and in women, the odds ratios were between  
35 1.0 and 2.0. The incidence rate of scleroderma in the general population is approximately

- 1 5–10 times higher in women compared with men, which may make it easier to detect large
- 2 relative risks in men.

**Table 4-70. Case-control studies of autoimmune diseases with measures of trichloroethylene exposure**

Disease, source of data	Results: exposure prevalence, OR, 95% CI	Reference, location, sample size, age
Scleroderma		
Structured interview (specific jobs and materials; jobs held 1 or more years). Exposure classified by self-report and by expert review (job exposure matrix).	<p>Men</p> <p>Maximum intensity 30% cases, 10% controls OR: 3.3 (95% CI: 1.0, 10.3)</p> <p>Cumulative intensity 32% cases, 21% controls OR: 2.0 (95% CI: 0.7, 5.3)</p> <p>Maximum probability 16% cases, 3% controls OR: 5.1 (95% CI: not calculated)</p> <p>Women</p> <p>Maximum intensity 6% cases, 7% controls OR: 0.9 (95% CI: 0.3, 2.3)</p> <p>Cumulative intensity 10% cases, 9% controls OR: 1.2 (95% CI: 0.5, 2.6)</p> <p>Maximum probability 4% cases, 5% controls OR: 0.7 (95% CI: 0.2, 2.2)</p>	Nietert et al. (1998) South Carolina. Prevalent cases, 178 cases (141 women, 37 men), 200 hospital-based controls. Mean age at onset 45.2 yr
Structured interview (specific jobs and materials; jobs held 6 or more months). Exposure classified by expert review.	<p>Men and women</p> <p>Any exposure: cases 16%, controls 8% OR: 2.4 (95% CI: 1.0, 5.4)</p> <p>High exposure:<sup>a</sup> cases 9%, controls 1% OR: 7.6 (95% CI: 1.5, 37.4)</p> <p>Men</p> <p>Any exposure: cases 64%, controls 27% OR: 4.7 (95% CI: 0.99, 22.0)</p> <p>Women</p> <p>Any exposure: cases 9%, controls 4% OR: 2.1 (95% CI: 0.65, 6.8)</p>	Diot et al. (2002) France. Prevalent cases, 80 cases (69 women, 11 men), 160 hospital controls. Mean age at diagnosis 48 yr
Structured interview (specific jobs and materials; jobs held 3 or more months). Exposure classified by self-report and by expert review.	<p>Women</p> <p>Self report: cases 1.3%, controls 0.7% OR: 2.0 (95% CI: 0.8, 4.8)</p> <p>Expert review: cases 0.7%, controls 0.4% OR: 1.9 (95% CI: 0.6, 6.6)</p>	Garabrant et al. (2003) Michigan and Ohio. Prevalent cases, 660 cases (all women), 2,227 population controls. <sup>b</sup> Ages 18 and older
Undifferentiated connective tissue disease		



Structured interview (specific jobs and materials; jobs held 3 or more months). Exposure classified by self-report and by expert review.	Women Self report: cases 0.5%, controls 0.7% OR: 0.88 (95% CI: 0.11, 6.95) Expert review: cases 0.5%, controls 0.4% OR: 1.67 (95% CI: 0.19, 14.9)	Lacey et al. (1999), Michigan and Ohio. Prevalent cases, 205 cases (all women), 2,095 population controls. Ages 18 and older
--	---	--

**Table 4-70. Case-control studies of autoimmune diseases with measures of trichloroethylene exposure (continued)**

Disease, source of data	Results: exposure prevalence, OR, 95% CI	Reference, location, sample size, age
ANCA-related diseases <sup>c</sup>		
Structured interview (specific jobs and materials; jobs held 6 or more months). Exposure classified by expert review.	Men and women (data not presented separately by sex) cases 18.3%, controls 17.5% OR: 1.1 (95% CI: 0.5, 2.4)	Beaudreuil et al. (2005) France. Incident cases, 60 cases (~50% women), 120 hospital controls. Mean age 61 yr

<sup>a</sup>Cumulative exposure defined as product of probability × intensity × frequency × duration scores, summed across all jobs; scores of >1 classified as “high.”

<sup>b</sup>Total *n*; *n* with TCE data: self-report 606 cases, 2,138 control; expert review 606 cases, 2,137 controls.

<sup>c</sup>Diseases included Wegener glomerulonephritis (*n* = 20), microscopic polyangiitis (*n* = 8), pauci-immune glomerulonephritis (*n* = 10), uveitis (*n* = 6), Churg-Strauss syndrome (*n* = 4), stroke (*n* = 4) and other diseases (no more than 2 each).

1 The EPA conducted a meta-analysis of the three scleroderma studies with specific  
2 measures of trichloroethylene (Diot et al., 2002; Garabrant et al., 2003; Nietert et al., 1998),  
3 examining separate estimates for males and for females. The resulting combined estimate for  
4 “any” exposure, using a random effects model to include the possibility of nonrandom error  
5 between studies (DerSimonian and Laird, 1986), was OR: 2.5 (95% CI: 1.1, 5.4) for men and  
6 OR: 1.2 (95% CI: 0.58, 2.6) in women. (Because the “any” exposure variable was not included  
7 in the published report, Dr. Paul Nietert provided the EPA with a new analysis with these results,  
8 e-mail communication from Paul Nietert to Glinda Cooper, November 28, 2007.)

9 Specific genes may influence the risk of developing autoimmune diseases, and genes  
10 involving immune response (e.g., cytokines, major histocompatibility complex, B- and T-cell  
11 activation) have been the focus of research pertaining to the etiology of specific diseases. The  
12 metabolism of specific chemical exposures may also be involved (Cooper et al., 1999).  
13 Povey et al. (2001) examined polymorphisms of two cytochrome CYP genes, CYP2E1 and  
14 CYP2C19, in relation to solvent exposure and risk of developing scleroderma. These specific  
15 genes were examined because of their hypothesized role in metabolism of many solvents,  
16 including trichloroethylene. Seven scleroderma patients who reported a history of solvent  
17 exposure were compared to 71 scleroderma patients with no history of solvent exposure and to  
18 106 population-based controls. The CYP2E1\*3 allele and the CYP2E1\*4 allele were more  
19 common in the 7 solvent-exposed patients (each seen in 2 of the 7 patients; 29%) than in either  
20 of the comparison groups (approximately 5% for CYP2E1\*3 and 14% for CYP2E1\*4). The  
21 authors present these results as observations that require a larger study for corroboration and  
22 further elucidation of specific interactions.

#### 4.6.1.1.6. Cancers of the Immune System, Including Childhood Leukemia

#### 4.6.1.1.7. Description of studies

24 Human studies have reported cancers of the immune system resulting from TCE  
25 exposure. Lymphoid tissue neoplasms arise in the immune system and result from events that  
26 occur within immature lymphoid cells in the bone marrow or peripheral blood (leukemias), or  
27 more mature cells in the peripheral organs (NHL). As such, the distinction between lymphoid  
28 leukemia and NHL is largely distributional with overlapping entities, such that a particular  
29 lymphoid neoplasm may manifest both lymphomatous and leukemic features during the course  
30 of the disease (Weisenburger, 1992). The broad category of lymphomas can be divided into  
31 specific types of cancers, including non-Hodgkin lymphoma, Hodgkin lymphoma, multiple

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 myeloma, and various types of leukemia (e.g., acute and chronic forms of lymphoblastic and  
2 myeloid leukemia). The classification criteria for these cancers has changed over the past 30  
3 years, reflecting improved understanding of the underlying stem cell origins of these specific  
4 subtypes. Lymphomas are grouped according to the World Health Organization (WHO)  
5 classification as B-cell neoplasms, T-cell/NK-cell neoplasms, and Hodgkin lymphoma, formerly  
6 known as Hodgkin disease (Harris et al., 2000). For example, hairy cell leukemia, chronic  
7 lymphocytic leukemia, non-Hodgkin lymphoma, and multiple myeloma may arise from mature  
8 B cells and are types of NHLs according to the WHO's lymphoma classification system (Morton  
9 et al., 2007, 2006). Most of the studies of TCE exposure evaluate NHL defined as  
10 lymphosarcoma, reticulum-cell sarcoma, and other lymphoid tissue neoplasms with recently  
11 published studies reporting on total B-cell or specific B-cell neoplasms.

12 Numerous studies are found in the published literature on NHL and either broad exposure  
13 categories or occupational title. The NHL studies generally report positive associations with  
14 organic solvents or job title as aircraft mechanic, metal cleaner or machine tool operator, and  
15 printers, although associations are not observed consistently across all studies, specific solvents  
16 are not identified, and different lymphoma classifications are adopted ('t Mannetje et al., 2008;  
17 Alexander et al., 2007b; Blair et al., 1993; Boffetta and de Vocht, 2007; Chiu and Weisenburger,  
18 2003; Cocco et al., 2010; Dryver et al., 2004; Figgs et al., 1995; Karunanayake et al., 2008;  
19 Lynge et al., 1997; Orsi et al., 2010; Richardson et al., 2008; Schenk et al., 2009; Seidler et al.,  
20 2007; Tatham et al., 1997; Vineis et al., 2007; Wang et al., 2009). A major use of TCE is the  
21 degreasing, as vapor or cold state solvent, of metal and other products with potential exposure in  
22 jobs in the metal industry, printing industry and aircraft maintenance or manufacturing industry  
23 (Bakke et al., 2007). The recent NHL case-control study of Purdue et al. (2009) examined  
24 degreasing tasks, specifically, and reported an increasing positive trend between NHL risk in  
25 males and three degreasing exposure surrogates: average frequency (hours/year) ( $p = 0.02$ ),  
26 maximal frequency (hours/year), ( $p = 0.06$ ), or cumulative number of hours ( $p = 0.04$ ).

27 As described in Appendix B, the EPA conducted a thorough and systematic search of  
28 published epidemiological studies of cancer risk and trichloroethylene exposure using the  
29 PubMed, TOXNET<sup>®</sup>, and EMBASE<sup>®</sup> bibliographic database. The EPA also requested  
30 unpublished data pertaining to trichloroethylene from studies that may have collected these data  
31 but did not include it in their published reports. ATSDR and state health department peer-  
32 reviewed studies were also reviewed. Information from each of these studies relating to  
33 specified design and analysis criteria was abstracted. These criteria included aspects of study  
34 design, representativeness of study subjects, participation rate/loss to follow-up, latency  
35 considerations, potential for biases related to exposure misclassification, disease

1 misclassification, and surrogate information, consideration of possible confounding, and  
2 approach to statistical analysis. All studies are considered for hazard identification but those  
3 studies more fully meeting the objective criteria provided the greater weight for identifying a  
4 cancer hazard.

5 The body of evidence on NHL and trichloroethylene is comprised of occupational cohort  
6 studies, population-based case-control studies and geographic studies. Four case-control studies  
7 and four geographic studies also examine childhood leukemia and trichloroethylene. Most  
8 studies report observed risk estimates and associated confidence intervals for NHL and overall  
9 TCE exposure. The studies included a broad but sometimes slightly different group of  
10 lymphosarcoma, reticulum-cell sarcoma, and other lymphoid tissue neoplasms, with the  
11 exception of the Nordstrom et al. (1998) case-control study, which examined hairy cell leukemia,  
12 now considered a NHL, the Zhao et al. (2005) cohort study, which reported only results for *all*  
13 lymphohematopoietic cancers, including nonlymphoid types and excluding chronic lymphocytic  
14 leukemia, and the Greenland et al. (1994) nested case-control study which reported results for  
15 NHL and Hodgkin lymphoma combined. Persson and Fredrikson (1999) do not identify the  
16 classification system for defining NHL, and Hardell et al. (1994) define NHL using the  
17 Rappaport classification system. Miligi et al. (2006) used an NCI classification system and  
18 considered chronic lymphocytic leukemias and NHL, classified as lymphosarcoma,  
19 reticulosarcoma, and other lymphoid tissue neoplasms, together, while Cocco et al. (2010), used  
20 the World Health Organization classification system, which reclassifies lymphocytic leukemias  
21 and NHLs as lymphomas of B-cell or T-cell origin. EPA staff, additionally, was able to obtain  
22 results generally consistent with the traditional NHL definition from Dr. Cocco, although  
23 lymphomas not otherwise specified were excluded (personal communication, Pierluigi Cocco to  
24 Cheryl Siegel Scott). The cohort studies (except for Zhao et al., 2005) and the nested case-  
25 control study of Greenland et al. (1994) have some consistency in coding NHL, with NHL  
26 defined as lymphosarcoma and reticulum-cell sarcoma (ICD code 200) and other lymphoid tissue  
27 neoplasms (ICD code 202) using the ICD Revisions 7, 8, or 9. Revisions 7 and 8 are essentially  
28 the same with respect to NHL; under Revision 9, the definition of NHL was broadened to  
29 include some neoplasms previously classified as Hodgkin lymphomas (Banks, 1992). Wang et  
30 al. (2009) refer to their cases as “NHL” cases and according to the ICD-O classification system  
31 that they used, their cases are more specifically NHL subtypes such as diffuse, lymphosarcoma,  
32 or follicular lymphoma (9590–9642, 9690–9701) or mast cell tumors (9740–9750) which is  
33 consistent with the traditional definition of NHL (i.e., ICD-7, -8, -9 codes 200 + 202) (Morton et  
34 al., 2003). NHL cases in Purdue et al. (2011) were also classified according to ICD-O (2<sup>nd</sup>  
35 Edition converted to ICO-O 3<sup>rd</sup> Edition codes), included diffuse, follicular T-cell and all other

1 NHL subtypes, which is generally consistent with the traditional definition of NHL, although this  
2 grouping does not include the malignant lymphomas of unspecified type coded as M-9590-9599.  
3 Fewer studies presented in published papers this information for cell-specific lymphomas,  
4 leukemia, leukemia cell type, or multiple myeloma (Anttila et al., 1995; Axelson et al., 1994;  
5 Boice et al., 1999; Boice et al., 2006b; Cocco et al., 2010; Costantini et al., 2008; Gold et al., In  
6 Press; Hansen et al., 2001; Morgan et al., 1998; Raaschou-Nielsen et al., 2001; Radican et al.,  
7 2008).

8 The seven cohort studies with data on the incidence of lymphopoietic and hematopoietic  
9 cancer in relation to trichloroethylene exposure range in size (803 (Hansen et al., 2001) to 86,868  
10 (Chang et al., 2005)), and were conducted in Denmark, Sweden, Finland, Taiwan and the United  
11 States (see Table 4-71; for additional study descriptions, see Appendix B). Some subjects in the  
12 Hansen et al. study are also included in a study reported by Raaschou-Nielsen et al. (2003);  
13 however, any contribution from the former to the latter are minimal given the large differences in  
14 cohort sizes of these studies (Hansen et al., 2001; Raaschou-Nielsen et al., 2003). The exposure  
15 assessment techniques used in all studies except Chang et al. (2005) and Sung et al. (2007)  
16 included a detailed job exposure matrix (Blair et al., 1998; Zhao et al., 2005), biomonitoring data  
17 (Anttila et al., 1995; Axelson et al., 1994; Hansen et al., 2001), or reference to industrial hygiene  
18 records on TCE exposure patterns and factors that affected exposure, indicating a high  
19 probability of TCE exposure potential (Raaschou-Nielsen et al., 2003) with high probability of  
20 TCE exposure to individual subjects. Subjects in Chang et al. (2005) and Sung et al. (2007), two  
21 studies with overlapping subjects employed at an electronics plant in Taiwan, have potential  
22 exposure to several solvents including TCE; all subjects are presumed as “exposed” because of  
23 employment in the plant although individual subjects would be expected to have differing  
24 exposure potentials. The lack of attribution of exposure intensity to individual subjects yields a  
25 greater likelihood for exposure misclassification compared to the six other studies with exposure  
26 assessment approaches supported by information on job titles, tasks, and industrial hygiene  
27 monitoring data. Incidence ascertainment in two cohorts began 21 (Blair et al., 1998) and  
28 38 years (Zhao et al., 2005) after the inception of the cohort. Specifically, Zhao et al. (2005)  
29 noted that their results may not accurately reflect the effects of carcinogenic exposure that  
30 resulted in nonfatal cancers before 1988. Because of the issues concerning case ascertainment  
31 raised by this incomplete coverage, observations must be interpreted in light of possible bias  
32 reflecting incomplete ascertainment of incident cases.

33 Eighteen cohort or PMR studies describing mortality risks from lymphopoietic and  
34 hematopoietic cancer are summarized in Table 4-72 (for additional study descriptions, see  
35 Appendix B). Two studies examined cancer incidence, Radican et al. (2008), who updated

1 mortality in Blair et al. (1998) cohort, and Zhao et al. (2005), and are identified above. In 10 of  
2 the 18 studies presenting mortality risks (ATSDR, 2004a; Blair et al., 1989; Chang et al., 2003b;  
3 Clapp and Hoffman, 2008; Costa et al., 1989; Garabrant et al., 1988; Henschler et al., 1995;  
4 Sinks et al., 1992; Sung et al., 2007; Wilcosky et al., 1984)

**Table 4-71. Incidence cohort studies of TCE exposure and lymphopoietic and hematopoietic cancer risk**

Population exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	
Aerospace workers (Rocketdyne), CA									Zhao et al. (2005)
Any TCE exposure	Not reported		Not reported						<i>n</i> = 5,049 (2,689 with high cumulative TCE exposure), began work before 1980, worked at least 2 yr, alive with no cancer diagnosis in 1988, follow-up from 1988–2000, job exposure matrix (intensity), internal referents (workers with no TCE exposure). Leukemia and multiple myeloma observations included in non-Hodgkin lymphoma category.
Low cumulative TCE score			1.0 (referent)	28					
Medium cumulative TCE score			0.88 (0.47, 1.65)	16					
High cumulative TCE score			0.20 (0.03, 1.46)	1					
( <i>p</i> for trend)			(0.097)						
Electronic workers, Taiwan									Chang et al. (2005); Sung et al. (2007)
All employees	0.67 (0.42, 1.01)	22							<i>n</i> = 88,868 ( <i>n</i> = 70,735 female), follow-up 1979–1997, does not identify TCE exposure to individual subjects (Chang et al., 2005).
Males	0.73 (0.27, 1.60)	6	Not reported		Not reported		Not reported		
Females	0.65 (0.37, 1.05)	16	Not reported		Not reported		Not reported		
Females					0.78 (0.49, 1.17)	23	Not reported		<i>n</i> = 63,982 females, follow-up 1979–2001, does not identify TCE exposure to individual subjects (Sung et al., 2007).

**Table 4-71. Incidence cohort studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	
Blue-collar workers, Denmark									Raaschou-Nielsen et al. (2003)
Any exposure	1.1 (1.0, 1.6)	229	1.2 (1.0, 1.5)	96	1.2 (0.9, 1.4)	82	1.03 (0.70, 1.47)	31	<i>n</i> = 40,049 (14,360 with presumed higher level exposure to TCE), worked for at least 3 mo, follow-up from 1968–1997, documented TCE use <sup>c</sup> . EPA based the lymphopoietic cancer category on summation of ICD codes 200–204.
Subcohort w/higher exposure <sup>d</sup>	Not reported		1.5 (1.2, 2.0)	65	Not reported		Not reported		
Employment duration									
1–4.9 yr			1.5 (1.1, 2.1)	35					
≥5 yr			1.6 (1.1, 2.2)	30					
Biologically-monitored workers, Denmark									Hansen et al. (2001)
Any TCE exposure	2.0 (1.1, 3.3)	15	3.1 (1.3, 6.1)	8	2.0 (0.7, 4.4)	6	0.71 (0.02, 3.98)	1	<i>n</i> = 803, U-TCA or air TCE samples, follow-up 1968–1996 (subset of Raaschou-Nielsen et al. (2003) cohort). EPA based the lymphopoietic cancer category on summation of ICD codes 200–204.
Cumulative exposure (Ikeda), males	Not reported				Not reported		Not reported		
<17 ppm-yr			3.9 (0.8, 11)	3					
≥17 ppm-yr			3.1 (0.6, 9.1)	3					
Mean concentration (Ikeda), males	Not reported				Not reported		Not reported		
<4 ppm			3.9 (1.1, 10)	4					
4+ ppm			3.2 (1.1, 10)	4					
Employment duration, males	Not reported				Not reported		Not reported		
<6.25 yr			2.5 (0.3, 9.2)	2					
≥6.25 yr			4.2 (1.1, 11)	4					



**Table 4-71. Incidence cohort studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	
Aircraft maintenance workers, Hill Air Force Base, UT									Blair et al. (1998)
TCE Subcohort	Not reported		Not reported		Not reported		Not reported		<i>n</i> = 10,461 men and 3,605 women (total <i>n</i> = 14,066, <i>n</i> = 7,204 with TCE exposure), employed at least 1 yr from 1952–1956, follow-up 1973–1990, job exposure matrix (intensity), internal referent (workers with no chemical exposures).
Males, cumulative exposure									
0	1.0 (referent)		1.0 (referent)		1.0 (referent)		1.0 (referent)	9	
<5 ppm-yr	0.8 (0.4, 1.7)	12	0.9 (0.3, 2.6)	8	0.4 (0.1, 2.0)	2	0.8 (0.1, 12.7)	1	
5–25 ppm-yr	0.7 (0.3, 1.8)	7	0.7 (0.2, 2.6)	4		0	3.8 (0.4, 37.4)	3	
>25 ppm-yr	1.4 (0.6, 2.9)	17	1.0 (0.4, 2.9)	7	0.9 (0.2, 3.7)	4	5.1 (0.6, 43.7)	5	
Females, cumulative exposure									
0	1.0 (referent)		1.0 (referent)		1.0 (referent)		1.0 (referent)		
<5 ppm-yr	1.2 (0.3, 4.4)	3	0.6 (0.1, 5.0)	1		0	Not reported	2	
5–25 ppm-yr	1.9 (0.4, 8.8)	2		0	2.4 (0.3, 21.8)	1	Not reported	1	
>25 ppm-yr	0.9 (9.2, 3.3)	3	0.9 (0.2, 4.5)	2		0	Not reported	1	
Biologically-monitored workers, Finland									Anttila et al. (1995)
Any TCE exposure	1.51 (0.92, 2.33)	20	1.81 (0.78, 3.56)	8	1.08 (0.35, 2.53)	5	1.62 (0.44, 4.16)	4	<i>n</i> = 3,089 men and women, U-TCA samples, follow-up 1967–1992.
Mean air-TCE (Ikeda extrapolation)									
<6 ppm	1.36 (0.65, 2.49)	10	2.01 (0.65, 4.69)	5	0.39 (0.01, 2.19)	1	1.48 (0.18, 5.35)	2	
6+ ppm	2.08 (0.95, 3.95)	9	1.40 (0.17, 5.04)	2	2.65 (0.72, 6.78)	4	2.41 (0.29, 6.78)	2	

**Table 4-71. Incidence cohort studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	Relative risk (95% CI) <sup>a</sup>	<i>n</i> <sup>a</sup>	
Biologically-monitored workers, Sweden									Axelsson et al. (1994)
Males	1.17 (0.47, 2.40)	7	1.56 (0.51, 3.64)	5	Not reported		0.57 (0.01, 3.17)	1	<i>n</i> = 1,421 men and 249 women (total 1,670), U-TCA samples, follow-up 1958–1987. EPA based the lymphopoietic cancer category includes ICD-7 200–203.
0–17 ppm (Ikeda extrapolation)	Not reported		1.44 (0.30, 4.20)	3	Not reported				
18–35 ppm (Ikeda extrapolation)			(0, 8.58)	0					
≥36 ppm (Ikeda extrapolation)			6.25 (0.16, 34.8)	1					
Females	Not reported		Not reported		Not reported				

<sup>a</sup>*n* = number of observed cases.

<sup>b</sup>Standardized incidence ratios using an external population referent group unless otherwise noted.

<sup>c</sup>Exposure assessment based on industrial hygiene data on TCE exposure patterns and factors that affect such exposure (Raaschou-Nielsen et al., 2002), with high probability of TCE exposure potential to individual subjects. Companies included iron and metal (48%), electronics (11%), painting (11%), printing (8%), chemical (5%), dry cleaning (5%), and other industries.

<sup>d</sup>Defined as at least 1 yr duration and first employed before 1980.

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
Computer manufacturing workers (IBM), NY									Clapp and Hoffman (2008)
Males	2.24 (1.01, 4.19)	9	Not reported		Not reported		Not reported	3	<i>n</i> = 115 cancer deaths from 1969–2001, proportional cancer mortality ratio, does not identify TCE exposure to individual subjects. EPA based the lymphopoietic cancer category on “all lymphatic cancers.”
Females	Not reported	0	Not reported		Not reported		Not reported	0	
Aerospace workers (Rocketdyne), CA									Boice et al. (2006b)
Any TCE (utility/eng flush)	0.74 (0.34, 1.40)	9	0.21 (0.01, 1.18)	1	1.08 (0.35, 2.53)	5	0.50 (0.01, 2.77)	1	<i>n</i> = 41,351 (1,111 Santa Susana workers with any TCE exposure), employed on or after 1948–1999, worked ≥6 mo, follow-up to 1999, job exposure matrix without quantitative estimate of TCE intensity.

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
Aerospace workers (Rocketdyne), CA (continued)									Zhao et al. (2005)
Any TCE exposure	Not reported		Not reported	60	Not reported		Not reported		<i>n</i> = 6,044 ( <i>n</i> = 2,689 with high cumulative level exposure to TCE), began work and worked at least 2 yr in 1950 or later - 1993, follow-up to 2001, job exposure matrix (intensity), internal referents (workers with no TCE exposure). Leukemia and multiple myeloma observations included in non-Hodgkin lymphoma category.
Low cumulative TCE score			1.0 (referent)	27					
Medium cumulative TCE score			1.49 (0.86, 2.57)	27					
High TCE score			1.30 (0.52, 3.23)	6					
( <i>p</i> for trend)			(0.370)						
View-Master employees, OR									ATSDR (2004a)
Males	0.58 (0.11, 1.69)	3	0.69 (0.08, 2.49)	2	0.50 (0.01, 2.79)	1			<i>n</i> = 616 deaths from 1989–2001, proportional mortality ratio, does not identify TCE exposure to individual subjects. EPA based the non-Hodgkin lymphoma cancer category on “other lymphopoietic tissue” which included NHL and multiple myeloma.
Females	0.64 (0.28, 1.26)	8	0.52 (0.14, 1.33)	4	0.67 (0.14, 1.96)	3			

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
Electronic workers, Taiwan									Chang et al. (2003b)
All employees									<i>n</i> = 88,868 ( <i>n</i> = 70,735 female), began work 1978–1997, follow-up 1985–1997, does not identify TCE exposure to individual subjects.
Males	Not reported		1.27 (0.41, 2.97)	5	0.44 (0.05, 1.59)	2	Not reported		
Females	Not reported		1.14 (0.55, 2.10)	10	0.54 (0.23, 1.07)	8	Not reported		
Aerospace workers (Lockheed), CA									Boice et al. (1999)
Routine TCE									<i>n</i> = 77,965 ( <i>n</i> = 2,267 with routine TCE exposure and <i>n</i> = 3,016 with intermittent-routine TCE exposure), began work ≥1960, worked at least 1 yr, follow-up from 1960–1996, job exposure matrix without quantitative estimate of TCE intensity.
Any TCE exposure	1.5 (0.81, 1.60)	36	1.19 (0.65, 1.99)	14	1.05 (0.54, 1.84)	12	0.91 (0.34, 1.99)	6	
Routine-intermittent									
Any TCE exposure	Not reported		Not reported		Not reported				
Exposure duration	Not reported				Not reported				
0 yr			1.0 (referent)	32			1.0 (referent)	24	
<1 yr			0.74 (0.32, 1.72)	7			0.45 (0.13, 1.54)	3	
1–4 yr			1.33 (0.64, 2.78)	10			1.48 (0.64, 3.41)	8	
≥5 yr			1.62 (0.82, 3.22)	14			0.51 (0.15, 1.76)	3	
<i>p</i> for trend			0.20				>0.20		

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
Uranium-processing workers (Fernald), OH									Ritz (1999a)
Any TCE exposure	Not reported		Not reported		Not reported		Not reported		<i>n</i> = 3,814 ( <i>n</i> = 2,971 with TCE), began work 1951–1972, worked ≥3 mo, follow-up to 1989, internal referents (workers with no TCE exposure).
No TCE exposure	1.0 (referent)		Not reported		Not reported		Not reported		
Light TCE exposure, >2 yr	1.45 (0.68, 3.06) <sup>c</sup>	18	Not reported		Not reported		Not reported		
Moderate TCE exposure, >2 yr	1.17 (0.15, 9.00) <sup>c</sup>	1	Not reported		Not reported		Not reported		
Aerospace workers (Hughes), CA									Morgan et al. (1998)
TCE subcohort	0.99 (0.64, 1.47)	25	0.96 (0.20, 2.81) <sup>d</sup>	3	1.05 (0.50, 1.93)	10	1.08 (0.35, 2.53) <sup>c</sup>	5	<i>n</i> = 20,508 (4,733 with TCE exposure), worked ≥6 mo 1950–1985, follow-up to 1993, external and internal (all non-TCE exposed workers) workers referent, job exposure matrix (intensity).
TCE subcohort			1.01 (0.46, 1.92) <sup>c</sup>	9					
Low intensity (<50 ppm)	1.07 (0.51, 1.96)	10	1.79 (0.22, 6.46) <sup>d</sup>	2	0.85 (0.17, 2.47)	3			
High intensity (>50 ppm)	0.95 (0.53, 1.57)	15	0.50 (0.01, 2.79) <sup>d</sup>	1	1.17 (0.47, 2.41)	7			
TCE subcohort (Cox Analysis)									
Never exposed	1.0 (referent)	82	1.0 (referent)	8	1.0 (referent)	32			
Ever exposed	1.05 (0.67, 1.65) <sup>f</sup>	25	1.36 (0.35, 5.22) <sup>d, f</sup>	3	0.99 (0.48, 2.03) <sup>f</sup>	10			
Peak									
No/Low	1.0 (referent)	90	1.0 (referent)	9	1.0 (referent)	35			
Medium/High	1.08 (0.64, 1.82)	17	1.31 (0.28, 6.08) <sup>d</sup>	2	1.10 (0.49, 2.49)	7			

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>	
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>		
Aerospace workers (Hughes), CA (continued)										
Cumulative										
Referent	1.0 (referent)	82	1.0 (referent)	8	1.0 (referent)	32				
Low	1.09 (0.56, 2.14)	10	2.25 (0.46, 11.1) <sup>d</sup>	2	0.69 (0.21, 2.32)	3				
High	1.03 (0.59, 1.79)	15	0.81 (0.10, 6.49) <sup>d</sup>	1	1.14 (0.5, 2.60)	7				
Aircraft maintenance workers, Hill Air Force Base, UT										
TCE subcohort	1.1 (0.7, 1.8) <sup>g</sup>	66	2.0 (0.9, 4.6) <sup>g</sup>	28	0.6 (0.3, 1.2) <sup>g</sup>	16	1.3 (0.5, 3.4)	14	<i>n</i> = 14,066 ( <i>n</i> = 7,204 ever exposed to TCE), employed at least 1 yr from 1952–1956, follow-up to 1990 (Blair et al., 1998) or to 2000 (Radican et al., 2008), job exposure matrix, internal referent (workers with no chemical exposures).	
Males, cumulative exposure										
0	1.0 (referent)		1.0 (referent)		1.0 (referent)		1.0 (referent)			
<5 ppm-yr	1.1 (0.6, 2.1)	21	1.8 (0.6, 5.4)	10	1.0 (0.3, 3.2)	7	1.0 (0.2, 4.2)	4		
5–25 ppm-yr	1.0 (0.4, 2.1)	11	1.9 (0.6, 6.3)	6		0	0.8 (0.1, 4.4)	2		
>25 ppm-yr	1.3 (0.7, 2.5)	21	1.1 (0.3, 3.8)	5	1.2 (0.4, 3.6)	7	1.2 (0.3, 4.7)	4		
Females, cumulative exposure										
0	1.0 (referent)				1.0 (referent)		1.0 (referent)			
<5 ppm-yr	1.5 (0.6, 4.0)	6	3.8 (0.8, 18.9)	3	0.4 (0.1, 3.2)	1	3.2 (0.5, 19.8)	2		
5–25 ppm-yr	0.7 (0.1, 4.9)	1		0		0	4.3 (0.4, 23.4)	1		
>25 ppm-yr	1.1 (0.4, 3.0)	6	3.6 (0.8, 16.2)	4	0.3 (0.1, 2.4)	1	1.3 (0.1, 13.2)	1		
TCE subcohort	1.06 (0.75, 1.51) <sup>h</sup>	106	1.36 (0.77, 2.39) <sup>h</sup>	46	0.64 (0.35, 1.18) <sup>h</sup>	27	1.35 (0.62, 2.93)	25		

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
Aircraft maintenance workers, Hill Air Force Base, UT (continued)									
Males, cumulative exposure	1.12 (0.72, 1.73)	88	1.56 (0.79, 4.21)	37	0.77 (0.37, 1.62)	24	1.08 (0.43, 2.71)	19	
0	1.0 (referent)		1.0 (referent)		1.0 (referent)		1.0 (referent)		
<5 ppm-yr	1.04 (0.63, 1.74)	34	1.83 (0.79, 4.21)	18	0.86 (0.36, 2.02)	11	0.69 (0.21, 2.27)	5	
5–25 ppm-yr	1.06 (0.49, 1.88)	21	1.17 (0.42, 3.24)	7	0.51 (0.16, 1.63)	4	1.58 (0.53, 4.71)	7	
>25 ppm-yr	1.25 (0.75, 2.09)	33	1.50 (0.61, 3.69)	12	0.87 (0.35, 2.14)	9	1.19 (0.40, 3.54)	7	
Females, cumulative exposure	1.00 (0.55, 1.83)	18	1.18 (0.49, 2.85)	9	0.36 (0.10, 1.32)	3	2.37 (0.67, 8.44)	6	
0	1.0 (referent)		1.0 (referent)		1.0 (referent)		1.0 (referent)		
<5 ppm-yr	1.10 (0.48, 2.54)	7	1.48 (0.47, 4.66)	4	0.35 (0.05, 2.72)	1	2.20 (0.40, 12.02)	2	
5–25 ppm-yr	0.38 (0.05, 2.79)	1		0		0	2.79 (0.31, 25.05)	1	
>25 ppm-yr	1.11 (0.53, 2.31)	10	1.30 (0.45, 3.77)	5	0.48 (0.10, 2.19)	2	2.38 (0.53, 10.67)	3	
Cardboard manufacturing workers, Arnsburg, Germany									Henschler et al. (1995)
TCE-exposed subjects	1.10 (0.03, 6.12)	1							<i>n</i> = 169 TCE exposed and <i>n</i> = 190 unexposed men, employed ≥1 yr from 1956–1975, follow-up to 1992, local population referent, qualitative exposure assessment.
Unexposed subjects from same factory	1.11 (0.03, 6.19)	1							



**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
General Electric plant, Pittsfield, MA			0.76 (0.24, 2.42) <sup>i,j</sup>	15	1.1 (0.46, 2.66) <sup>i</sup>	22			Greenland et al. (1994)
									Nested case-control study, <i>n</i> = 512 cancer (cases) and 1,202 noncancer (controls) male deaths reported to pension fund between 1969–1984 among workers employed <1984 and with job history record, job exposure matrix-ever held job with TCE exposure. Hodgkin lymphoma in NHL grouping.
Cardboard manufacturing workers, Atlanta, GA									Sinks et al. 1992
	0.3 (0.0, 1.6)	1	Not reported		Not reported		Not reported		<i>n</i> = 2,050, employed on or before 1957–1988, follow- up to 1988, Material Data Safety Sheets used to identify chemicals used in work areas.

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
U.S. Coast Guard employees									Blair et al. (1989)
Marine inspectors	1.57 (0.91, 2.51)	17	1.75 (0.48, 4.49)	4	1.55 (0.62, 3.19)	7	Not reported		<i>n</i> = 3,781 males (1,767 marine inspectors), employed 1942–1970, follow-up to 1980. TCE and nine other chemicals identified as potential exposures; no exposure assessment to individual subjects.
Noninspectors	0.60 (0.24, 1.26)	7	0.41 (0.01, 2.30)	1	0.66 (0.14, 1.94)	3	Not reported		
Aircraft manufacturing employees, Italy									Costa et al. (1989)
All male subjects	0.80 (0.41, 1.40)	12	Not reported		Not reported		Not reported		<i>n</i> = 7,676 males, employed on or before 1954–1981, followed to 1981, job titles of white- and blue-collar workers, technical staff, and administrative clerks, does not identify TCE exposure to individual subjects.
Aircraft manufacturing, San Diego, CA									Garabrant et al. (1988)
All employees	0.82 (0.56, 1.15)	32	0.82 (0.44, 1.41) <sup>d</sup>	13	0.82 (0.47, 1.32)	10	Not reported		<i>n</i> = 14,067, employed at least 4 yr with company and ≥1 d at San Diego plant from 1958–1982, followed to 1982, does not identify TCE exposure to individual subjects.
			0.65 (0.21, 1.52) <sup>k</sup>	5					

**Table 4-72. Mortality cohort and PMR studies of TCE exposure and lymphopoietic and hematopoietic cancer risk (continued)**

Population, exposure group	Lymphopoietic cancer		non-Hodgkin lymphoma		Leukemia		Multiple myeloma		Reference(s) and study description <sup>b</sup>
	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	Relative risk (95% CI)	<i>n</i> <sup>a</sup>	
Solvent-exposed rubber workers									Wilcosky et al. (1984)
	2.4 <sup>i</sup>	3	0.81	3					Nested case-control study, <i>n</i> = 9 lymphosarcoma and 10 leukemia (cases) and 20% random sample of all other deaths (controls) between 1964–1973 in cohort of <i>n</i> = 6,678, exposure assessment by company record for use in work area.

<sup>a</sup>*n* = number of observed cases.

<sup>b</sup>Unless otherwise noted, all studies reported standardized mortality ratios using an external population referent group.

<sup>c</sup>Logistic regression analysis with 15 lag for TCE exposure (Ritz, 1999a).

<sup>d</sup>In Morgan et al. (1998) and Garabrant et al. (1988), this category was based on lymphosarcoma and reticulosarcoma.

<sup>e</sup>As presented in Mandel et al. (2006) for NHL, this category defined as ICD-7, ICDA-8, and ICD-9 codes of 200 and 202. As presented in Alexander et al. (2006) for multiple myeloma.

<sup>f</sup>Risk ratio from Cox Proportional Hazard Analysis, stratified by age and sex, from Environmental Health Strategies (1997) Final Report to Hughes Corporation (Communication from Paul A. Cammer, President, Trichloroethylene Issues Group to Cheryl Siegel Scott, U.S. EPA, December 22, 1997).

<sup>g</sup>Estimated relative risks from Blair et al. (1998) from Poisson regression models adjusted for date of hire, calendar year of death and sex.

<sup>h</sup>Estimated relative risks from Radican et al. (2008) from Cox proportional hazard models adjusted for age and sex.

<sup>i</sup>Odds ratio from nested case-control analysis.

<sup>j</sup>Lymphomas, lymphosarcomas, reticulosarcomas, and Hodgkin lymphoma (ICDA-8 200-202) in Greenland et al. (1994).

<sup>k</sup>Other lymphatic and hematopoietic tissue neoplasms (Garabrant et al., 1988).

IBM = International Business Machines Corporation.

1 a relatively limited exposure assessment methodology was used, study participants may not  
2 represent the underlying population, or there was a low exposure prevalence of TCE exposure.  
3 For reasons identified in the systematic review, these studies are given less weight in the overall  
4 evaluation of the literature than the eight other cohort studies that better met the ideals of  
5 evaluation criteria (Blair et al., 1998 and extended follow-up by; Boice et al., 1999; Boice et al.,  
6 2006b; Greenland et al., 1994; Morgan et al., 1998; Radican et al., 2008; Ritz, 1999a; Zhao et al.,  
7 2005).

8 Case-control studies of NHL from United States (Connecticut), Germany, Italy, Sweden,  
9 and Canada were identified, and are summarized in Table 4-73 (for additional study descriptions,  
10 see Appendix B). These studies identified cases from hospital records (Cocco et al., 2010;  
11 Costantini et al., 2008; Hardell et al., 1994; Mester et al., 2006; Miligi et al., 2006; Persson and  
12 Fredrikson, 1999; Seidler et al., 2007; Siemiatycki, 1991); the Surveillance, Epidemiology, and  
13 End Results Cancer Registry—Connecticut residents (Wang et al., 2009), Iowa, Los Angeles  
14 County, and Seattle and Detroit metropolitan area residents (Purdue et al., 2011), or Seattle and  
15 Detroit metropolitan area residents (Gold et al., In Press); or the Swedish Cancer Registry  
16 (Nordström et al., 1998), and hospital or population controls. These studies assign potential  
17 occupational TCE exposure to cases and controls using self-reported information obtained from a  
18 mailed questionnaire (Hardell et al., 1994; Nordström et al., 1998; Persson and Fredrikson, 1999)  
19 or from direct interview with study subjects, with industrial hygienist ratings of exposure  
20 potential and a job exposure matrix (Cocco et al., 2010; Costantini et al., 2008; Miligi et al.,  
21 2006; Purdue et al., 2011; Seidler et al., 2007; Siemiatycki, 1991; Wang et al., 2009).  
22 Additionally, large multiple center lymphoma case-control studies examine specific types of  
23 NHL (Cocco et al., 2010; Miligi et al., 2006; Purdue et al., 2011; Wang et al., 2009), leukemia  
24 (Costantini et al., 2008), or multiple myeloma (Cocco et al., 2010; Costantini et al., 2008; Purdue  
25 et al., 2011).

26 Four geographic based studies on NHL in adults are summarized in Table 4-74 (for  
27 additional study descriptions, see Appendix B) and subjects in three studies are identified based  
28 upon their residence in a community where TCE was detected in water serving the community  
29 (ATSDR, 2006b; Cohn et al., 1994b; Vartiainen et al., 1993). Both Cohn et al. (1994b) and  
30 ATSDR (2006b) also present estimates for childhood leukemia and these observations are  
31 discussed below with other studies reporting on childhood leukemia. A subject is assumed to  
32 have a probability of exposure due to residence likely receiving water containing TCE. Most  
33 studies do not include statistical models of water distribution networks, which may influence  
34 TCE concentrations delivered to a home, nor a subject's ingestion rate to estimate TCE exposure

- 1 to individual study subjects. ATSDR (2006b) adopts exposure modeling of soil vapor
- 2 contamination to define

**Table 4-73. Case-control studies of TCE exposure and lymphopietic cancer, leukemia or multiple myeloma**

Population	Cancer type and exposure group	Odds ratio (95% CI)	n exposed cases	Reference(s)
Men and women aged 20–74 in Iowa, Los Angeles County (CA), Seattle and Detroit metropolitan areas	<b>Non-Hodgkin lymphoma</b>			Gold et al. (In Press); Purdue et al. (2011)
	Any TCE exposure			
	Possible	1.1 (0.9, 1.3)	545	
	Probable	1.4 (0.8, 2.4)	45	
	Average weekly exposure <sup>a</sup>			
	0 ppm-h per week	1.0	341	
	1–60 ppm-h per week	1.6 (0.7, 3.8)	15	
	61–150 ppm-h per week	0.5 (0.2, 1.4)	7	
	>150 ppm-h per week	2.5 (1.1, 6.1)	23	
	( <i>p</i> for linear trend)	0.02		
	Cumulative exposure <sup>a</sup>			
	0	1.0	341	
	1–46,800 ppm-h	1.4 (0.6, 3.3)	14	
	46,801–112,320 ppm-h	0.6 (0.2, 1.7)	7	
	>112,320 ppm-h	2.3 (1.0, 5.0)	24	
	( <i>p</i> for linear trend)	0.08		
	<b>Non-Hodgkin lymphoma types</b>			
	Probable TCE exposure			
	Diffuse	0.9 (0.5, 2.0)	155	
	Follicular	2.1 (1.0, 4.2)	13	
Chronic lymphocytic leukemia	2.7 (1.2, 5.8)	11		

**Table 4-73. Case-control studies of TCE exposure and lymphopoietic cancer, leukemia or multiple myeloma (continued)**

Population	Cancer type and exposure group	Odds ratio (95% CI)	n exposed cases	Reference(s)
Men and women aged 20–74 (continued)	<b>Multiple myeloma</b>			Gold et al. (In Press); Purdue et al. (2011) (continued)
	Any TCE exposure	1.4 (0.9, 2.1)	66	
	High confidence exposure <sup>b</sup>	1.7 (1.0, 2.7)	43	
	Cumulative exposure <sup>b</sup>			
	0	1.0	139	
	1–471 ppm-h	1.1 (0.4, 2.9)	6	
	472–3,000 ppm-h	1.6 (0.7, 3.5)	11	
	3,001–7,644 ppm-h	1.5 (0.6, 3.9)	7	
	7,645–570,000 ppm-h	2.3 (1.1, 5.0)	17	
	(p for linear trend)	0.03		
Men and women aged ≥17 yr in Czech Republic, Finland, France, Germany, Ireland, Italy, and Spain (Epilymph study)	<b>All Centers:</b>			Cocco et al. (2010)
	<b>B-cell NHL<sup>b</sup></b>			
	Any TCE exposure	0.8 (0.6, 1.1)	71	
	Cumulative Exposure			
	Low	0.9 (0.6, 1.6)	26	
	Medium	0.5 (0.3, 0.9)	16	
	High	1.0 (0.6, 1.6)	29	
	(p for linear trend)	0.16		

**Table 4-73. Case-control studies of TCE exposure and lymphopietic cancer, leukemia or multiple myeloma (continued)**

Population	Cancer type and exposure group	Odds ratio (95% CI)	n exposed cases	Reference(s)
Men and women aged $\geq 17$ yr in Czech Republic, Finland, France, Germany, Ireland, Italy, and Spain (EpiLymph study) (continued)	<b>Non-Hodgkin lymphoma types<sup>c</sup></b>			Cocco et al. (2010) (continued)
	Diffuse large B-cell	0.7 (0.4, 1.1)	17	
	Follicular	1.2 (0.6, 2.3)	11	
	Chronic lymphocytic leukemia	0.9 (0.5, 1.5)	18	
	Multiple myeloma	0.6 (0.3, 1.2)	9	
	T-cell lymphoma	0.9 (0.4, 2.2)	6	
	<b>German Centers:</b>			Seidler et al. (2007); Mester et al. (2006)
	<b>Non-Hodgkin lymphoma</b>			
	Any TCE exposure	Not reported		
	Cumulative TCE			
	0 ppm-yr	1.0	610	
	>0– $\leq 4$ ppm-yr	0.7 (0.4, 1.1)	40	
	4.4– $< 35$ ppm-yr	0.7 (0.5, 1.2)	32	
	High exposure, $> 35$ ppm-yr	2.1 (1.0, 4.8)	21	
( <i>p</i> for linear trend)	0.14			
>35 ppm-yr, 10 yr lag	2.2 (1.0, 4.9)			
Women aged 21–84 in Connecticut, U.S.	<b>Non-Hodgkin lymphoma</b>			Wang et al. (2009)
	Any TCE exposure	1.2 (0.9, 1.8)	77	
	Low intensity TCE exposure	1.1 (0.8, 1.6)	64	
	Medium-high intensity TCE exposure	2.2 (0.9, 5.4)	13	
	( <i>p</i> for linear trend)	0.06		
	Low probability TCE exposure	1.1 (0.7, 1.8)	43	
	Medium-high probability TCE exposure	1.4 (0.9, 2.4)	34	
( <i>p</i> for linear trend)	0.37			



**Table 4-73. Case-control studies of TCE exposure and lymphopoietic cancer, leukemia or multiple myeloma (continued)**

Population	Cancer type and exposure group	Odds ratio (95% CI)	n exposed cases	Reference(s)
Women aged 21–84 in Connecticut, U.S. (continued)	Low intensity TCE exposure/low probability	0.9 (0.6, 1.5)	30	Wang et al. (2009) (continued)
	Low intensity/medium-high probability	1.4 (0.9, 2.4)	34	
	Medium-high intensity/low probability	2.2 (0.9, 5.4)	13	
	Medium-high intensity/medium-high probability		0	
Population in 8 Italian regions	<b>Non-Hodgkin lymphoma</b>			Miligi et al. (2006); Costantini et al. (2008)
	Any TCE exposure	Not reported		
	TCE exposure intensity			
	Very low/low	0.8 (0.5, 1.3)	35	
	Medium/high	1.2 (0.7, 2.0)	35	
	(p for linear trend)	0.8		
	Duration exposure, medium/high TCE intensity			
	≤15 yr	1.1 (0.6, 2.1)	22	
	>15 yr	1.0 (0.5, 2.6)	12	
	(p for linear trend)	0.72		
	<b>Other non-Hodgkin lymphoma</b>			
	TCE exposure intensity, medium/high			
	Small lymphocytic NHL	0.9 (0.4, 2.1)	7	
	Follicular NHL	Not presented	3	
	Diffuse NHL	1.9 (0.9, 3.7)	13	
	Other NHL	1.2 (0.6, 2.4)	11	
Multiple myeloma	0.9 (0.3, 2.4)	27		

**Table 4-73. Case-control studies of TCE exposure and lymphopoietic cancer, leukemia or multiple myeloma (continued)**

Population	Cancer type and exposure group	Odds ratio (95% CI)	n exposed cases	Reference(s)	
Population in 8 Italian regions (continued)	<b>Leukemia</b>			Miligi et al. (2006); Costantini et al. (2008) (continued)	
	Any TCE exposure	Not reported			
	TCE exposure intensity				
	Very low/low	1.0 (0.5, 1.8)	17		
	Medium/high	0.7 (0.4, 1.5)	11		
	<b>Chronic lymphocytic leukemia</b>				
	Any TCE exposure	Not reported			
	TCE exposure intensity				
	Very low/low	1.2 (0.5, 2.7)	8		
Medium/high	0.9 (0.3, 2.6)	4			
Population of Örebro and Linköping, Sweden	<b>B-cell non-Hodgkin lymphoma</b>			Persson and Fredrikson (1999)	
	Any TCE exposure	1.2 (0.5, 2.4)	16		
Population of Sweden	<b>Hairy cell lymphoma</b>			Nordstrom et al. (1998)	
	Any TCE exposure	1.5 (0.7, 3.3)	9		
Population of Umea, Sweden	<b>Non-Hodgkin lymphoma</b>			Hardell et al. (1994)	
	Any exposure to TCE	7.2 (1.3, 42)	4		
Population of Montreal, Canada	<b>Non-Hodgkin lymphoma</b>			Siemiatycki et al. (1991)	
	Any TCE exposure	1.1 (0.6, 2.3) <sup>d</sup>	6		
	Substantial TCE exposure	0.8 (0.2, 2.5) <sup>d</sup>	2		

<sup>a</sup>For Purdue et al. (2011), OR for subjects interviewed using computer-assisted personal interview with job modules and includes subjects assessed as unexposed or with probably exposure, defined as holding one or more jobs with an assigned probability of TCE exposure of  $\geq 50\%$ .

<sup>b</sup>For Gold et al. (In Press) subjects with jobs assessed with low confidence considered as unexposed.

<sup>c</sup>For Cocco et al. (2010), OR for subjects with high confidence assessment of TCE exposure.

<sup>d</sup>90% CI.

**Table 4-74. Geographic-based studies of TCE and non-Hodgkin lymphoma or leukemia in adults**

Population	Exposure group	non-Hodgkin lymphoma		Leukemia		Reference <sup>a</sup>
		Relative risk (95% CI)	<i>n</i> exposed cases	Relative risk (95% CI)	<i>n</i> exposed cases	
Two study areas in Endicott, NY		0.54 (0.22, 1.12)	7	0.79 (0.34, 1.55)	8	ATSDR (2006b)
Residents of 13 census tracts in Redlands, CA		1.09 (0.84, 1.38)	111	1.02 (0.74, 1.35)	77	Morgan and Cassady (2002)
Population in New Jersey	Males, maximum estimated TCE concentration (ppb) in municipal drinking water					Cohn et al. (1994b)
	<0.1	1.00	493	1.00	438	
	0.1–0.5	1.28 (1.10, 1.48)	272	0.85 (0.71, 1.02)	162	
	≥5.0	1.20 (0.94, 1.52)	78	1.10 (0.84, 1.90)	63	
	Females, maximum estimated TCE concentration (ppb) in municipal drinking water					
	<0.1	1.00	504	1.00; 315		
	0.1–0.5	1.02 (0.87, 1.2)	26	1.13 (0.93, 1.37)	156	
	>5.0	1.36 (1.08, 1.70)	87	1.43 (1.43, 1.90)	56	
Population in Finland	Residents of Hausjarvi	0.6 (0.3, 1.1)	14	1.2 (0.8, 1.7)	33	Vartiainen et al. (1993)
	Residents of Huttula	1.4 (1.0, 2.0)	13	0.7 (0.4, 1.1)	19	

<sup>a</sup>No geographic-based study reported a relative risk estimate for multiple myeloma except Vartiainen et al. (1993) who observed standardized incidence ratios of 0.7 (95% CI: 0.3, 1.3) and 0.6 (95% CI: 0.2, 1.3) for residents of Hausjarvi and Huttula, respectively.

1 study area boundaries and to identify census tracts with a higher probability of exposure to  
2 volatile organic solvents without identifying exposure concentrations to TCE and other solvents.  
3 In these studies, one level of exposure to all subjects in a geographic area is assigned, although  
4 there is some inherent measurement error and misclassification bias because not all subjects are  
5 exposed uniformly.

6 NHL risk is statistically significantly elevated in three studies in which there is a high  
7 likelihood of TCE exposure in individual study subjects (e.g., based on job-exposure matrices or  
8 biomarker monitoring) and which met, to a sufficient degree, the standards of epidemiologic  
9 design and analysis in a systematic review (3.1, 95% CI: 1.3, 6.1 (Hansen et al., 2001); 1.5, 95%  
10 CI: 1.2, 2.0, subcohort with higher exposure (Raaschou-Nielsen et al., 2003), 2.3, 95% CI: 1.0,  
11 5.0, >112,320-ppm hours cumulative TCE exposure, 2.5, 95% CI: 1.1, 6.1, >150-ppm hours  
12 average weekly TCE exposure (Purdue et al., 2011)). Two of these incidence studies report  
13 statistically significant associations for NHL for subjects with longer employment duration as a  
14 surrogate of TCE exposure ( $\geq 6.25$  year, 4.2, 95% CI: 1.1, 11 (Hansen et al., 2001);  $\geq 5$  year, 1.6,  
15 95% CI: 1.1, 2.2, (Raaschou-Nielsen et al., 2003)) and Purdue et al. (2011) report a positive  
16 trend with NHL and cumulative TCE exposure ( $p = 0.08$ ) or average weekly TCE exposure ( $p =$   
17  $0.02$ ). Hansen et al. (2001) also examined two other exposure surrogates, cumulative exposure  
18 and exposure intensity, with estimated risk larger in low exposure groups than for high exposure  
19 groups. A fourth study from Sweden reports a large and imprecise risk with TCE (7.2, 95% CI:  
20 1.3, 42 (Hardell et al., 1994)) based on four exposed cases. Cohort mortality studies and other  
21 case-control studies, except Cocco et al. (2010), observed a 10–50% increased risk between  
22 NHL and any TCE exposure (1.2, 95% CI: 0.65, 1.99 (Boice et al., 1999); 1.36, 95% CI: 0.28,  
23 6.08 (Morgan et al., 1998); 1.5, 95% CI: 0.7, 3.3 (Nordström et al., 1998); 1.2, 95% CI: 0.5, 2.4  
24 (Persson and Fredrikson, 1999); 1.36, 95% CI: 0.77, 2.39 (Radican et al., 2008); 1.1,  
25 95% CI: 0.6, 2.3 (Siemiatycki, 1991); 1.2, 95% CI: 0.9, 1.8 (Wang et al., 2009)).

26 Odds ratios are higher for diffuse or follicular NHL, primarily B-cell lymphomas, than  
27 for all non-Hodgkin lymphomas in both studies which examine forms of lymphoma, although  
28 based on few exposed cases and inconsistently reported (Cocco et al., 2010; Miligi et al., 2006;  
29 Purdue et al., 2011) (see Table 4-74). Observations in the two other studies of B-cell lymphomas  
30 (Persson and Fredrikson, 1999; Wang et al., 2009) appear consistent with Miligi et al. (2006) and  
31 Purdue et al. (2011). Together, these observations suggest that the associations between  
32 trichloroethylene and specific NHL types are stronger than the associations seen with other  
33 forms of NHL, and that disease misclassification may be introduced in studies examining  
34 trichloroethylene and NHL as a broader category. Mortality observations in other occupational  
35 cohorts (Costa et al., 1989; Garabrant et al., 1988; Greenland et al., 1994; Wilcosky et al., 1984)

1 (ATSDR, 2004a; Boice et al., 2006b; Chang et al., 2003b; Henschler et al., 1995; Ritz, 1999a;  
2 Sung et al., 2007) included a risk estimate of 1.0 in 95% CIs; these studies neither add to nor  
3 detract from the overall weight of evidence given their lower likelihood for TCE exposure due to  
4 inferior exposure assessment approaches, lower prevalence of exposure, lower statistical power,  
5 and fewer exposed deaths.

6 Seven studies presented estimated risks for leukemia and overall TCE exposure  
7 (Anttila et al., 1995; Blair et al., 1998 and its update by Radican et al., 2008; Morgan et al., 1998;  
8 Boice et al., 1999, 2006; Hansen et al., 2001; Raaschou-Nielsen et al., 2003); only three studies  
9 also presented estimated risks for a high exposure category (Anttila et al., 1995; Blair et al.,  
10 1998; Morgan et al., 1998). Three case-control studies presented estimated risk for leukemia  
11 categories and overall TCE exposure or low or high TCE exposure category (Cocco et al., 2010;  
12 Costantini et al., 2008; Purdue et al., 2011). Risk estimates in these cohort studies ranged from  
13 0.64 (95% CI: 0.35, 1.18) (Radican et al., 2008) to 2.0 (95% CI: 0.7, 4.44) (Hansen et al., 2001).  
14 The largest study, with 82 observed incident leukemia cases, reported a relative risk estimate of  
15 1.2 (95% CI: 0.9, 1.4) (Raaschou-Nielsen et al., 2003). Case-control studies which examined all  
16 leukemias (Costantini et al., 2008) or chronic lymphocytic leukemia (Cocco et al., 2010;  
17 Costantini et al., 2008; Purdue et al., 2011), and TCE exposure are quite limited in statistical  
18 power. Risk estimates in the four case-control studies ranged from 0.7 (95% CI: 0.4, 1.5) for all  
19 leukemias and medium to high exposure intensity (Costantini et al. (2008) to 2.7 (95% CI: 1.2,  
20 5.8) for chronic lymphocytic leukemia and probable TCE exposure (Purdue et al., 2011).

21 Eight cohort studies presented estimated risks for multiple myeloma and overall TCE  
22 exposure (Anttila et al., 1995; Axelson et al., 1994) (Blair et al., 1998; and its update by Radican  
23 et al., 2008) (Boice et al., 1999; Boice et al., 2006b; Hansen et al., 2001; Morgan et al., 1998;  
24 Raaschou-Nielsen et al., 2003); only three studies also presented estimated risks for a high  
25 exposure category (Anttila et al., 1995; Boice et al., 1999; Radican et al., 2008). Three case-  
26 control studies presented estimated risk for multiple myeloma and overall TCE exposure or low  
27 or high TCE exposure category (Cocco et al., 2010; Costantini et al., 2008; Gold et al., In Press).  
28 Risk estimates in these cohort studies ranged from 0.57 (95% CI: 0.01, 3.17) (Axelson et al.,  
29 1994) to 1.62 (95% CI: 0.44, 4.16) (Anttila et al., 1995). The largest cohort study, with 31  
30 observed incident multiple myeloma cases, reported a relative risk estimate of 1.03 (95% CI:  
31 0.70, 1.47) (Raaschou-Nielsen et al., 2003). The largest case-control study of 43 exposed  
32 multiple myeloma cases with high confidence TCE exposure reported an odds ratio of 1.7 (95%  
33 CI: 1.0, 2.7) and a positive trend with increasing cumulative TCE exposure ( $p = 0.03$ ) (Gold et  
34 al., In Press).

1 The number of studies of childhood lymphoma including acute lymphatic leukemia and  
2 trichloroethylene is much smaller than the number of studies of trichloroethylene and adult  
3 lymphomas, and consists of four case-control studies (Costas et al., 2002; Lowengart et al., 1987;  
4 McKinney et al., 1991; Shu et al., 1999) and four geographic based studies (ADHS, 1990, 1995;  
5 Aickin et al., 1992; ATSDR, 2006a, 2008; Cohn et al., 1994b), (see Table 4-75). An additional  
6 publication, focusing on ras mutations, based on one of the case-control studies is also available  
7 (Shu et al., 2004). All four case-control studies evaluate maternal exposure, and three studies  
8 also examine paternal occupational exposure (Lowengart et al., 1987; McKinney et al., 1991;  
9 Shu et al., 2004; Shu et al., 1999). There are relatively few cases with maternal exposure (range  
10 0–16) in these case-control studies, and only Shu et al. have a large number ( $n = 136$ ) of cases  
11 with paternal exposure (Shu et al., 2004; Shu et al., 1999). The small numbers of exposed case  
12 parents limit examination of possible susceptibility time windows. Overall, evidence for  
13 association between parental trichloroethylene exposure and childhood leukemia is not robust or  
14 conclusive.

15 The results from the studies of Costas et al. (2002) and Shu et al. (2004; 1999) suggest a  
16 fetal susceptibility to maternal exposure during pregnancy, with relative risks observed for this  
17 time period equal or higher than the relative risks observed for periods before conception or after  
18 birth (see Table 4-75). The studies by Lowengart et al. (1987) and McKinney et al. (1991) do  
19 not provide informative data pertaining to this issue due to the small number ( $n = <3$ ) of exposed  
20 case mothers. A recent update of a cohort study of electronics workers at a plant in Taiwan  
21 (Chang et al., 2003b; 2005) reported a fourfold increased risk (3.83; 95% CI: 1.17, 12.55 (Sung  
22 et al., 2008)) for childhood leukemia risk among the offspring of female workers employed  
23 during the three months before to three months after conception. Exposures at this factory  
24 included trichloroethylene, perchloroethylene, and other organic solvents (Sung et al., 2008).  
25 The lack of TCE assignment to individual subjects in this study decrease its weight in the overall  
26 analysis.

27 The evidence for an association between childhood leukemia and paternal exposure to  
28 solvents is quite strong (Colt and Blair, 1998); however, for studies of TCE exposure, the small  
29 numbers of exposed case fathers in two studies (Lowengart et al., 1987; McKinney et al., 1991)  
30 and, for all three studies, likelihood of misclassification resulting from a high percentage of  
31 paternal occupation information obtained from proxy interviews, limits observation  
32 interpretations. Both Lowengart et al. (1987) and McKinney et al. (1991) provide some evidence  
33 for a two- to fourfold increase of childhood leukemia risk and paternal occupational exposure  
34 although the population study of Shu et al. (2004; 1999), with 13% of case father's occupation  
35 reported by proxy respondents, does not appear to support the earlier and smaller studies.



1  
2  
3

**Table 4-75. Selected results from epidemiologic studies of TCE exposure and childhood leukemia**

		Relative risk (95% CI)	n observed events	Reference(s)
<b>Cohort studies (solvents)</b>				
Childhood leukemia among offspring of electronic workers				Sung et al. (2008)
	Nonexposed	1.0 <sup>a</sup>	9	
	Exposed pregnancy to organic solvents	3.83 (1.17, 12.55)	6	
<b>Case-control studies</b>				
Children's Cancer Group Study (children ≤15 yr age)				
<b>Acute lymphocytic leukemia</b>				
Maternal occupational exposure to TCE				Shu et al. (1999)
	Anytime	1.8 (0.8, 4.1)	15	
	Preconception	1.8 (0.8, 5.2)	9	
	During pregnancy	1.8 (0.5, 6.4)	6	
	Postnatal	1.4 (0.5, 4.1)	9	
Paternal occupational exposure to TCE				
	Anytime	1.1 (0.8, 1.5)	136	
	Preconception	1.1 (0.8, 1.5)	100	
	During pregnancy	0.9 (0.6, 1.4)	56	
	Postnatal	1.0 (0.7, 1.3)	77	
<b>K-ras + acute lymphocytic leukemia</b>				Shu et al., (2004)
Maternal occupational exposure to TCE				
	Anytime	1.8 (0.6, 4.8)	5	
	Preconception	2.0 (0.7, 6.3)	4	
	During pregnancy	3.1 (1.0, 9.7)	4	
	Postnatal		0	
Paternal occupational exposure to TCE				
	Anytime	0.6 (0.3, 1.4)	9	
	Preconception	0.6 (0.3, 1.5)	8	
	During pregnancy	0.3 (0.1, 1.2)	2	
	Postnatal	0.4 (0.1, 1.4)	3	

4



**Table 4-75. Selected results from epidemiologic studies of TCE exposure and childhood leukemia (continued)**

	Relative risk (95% CI)	n observed events	Reference(s)
Residents of ages ≤19 in Woburn, MA			Costas et al. (2002)
Maternal exposure 2 yr before conception to diagnosis			
Never	1.00	3	
Least	5.00 (0.75, 33.5)	9	
Most	3.56 (0.51, 24.8)	7	
(p for linear trend)	≥0.05		
Maternal exposure 2 yr before conception			
Never	1.00	11	
Least	2.48 (0.42, 15.2)	4	
Most	2.82 (0.30, 26.4)	4	
(p for linear trend)	≥0.05		
Birth to diagnosis			
Never	1.00	7	
Least	1.82 (0.31, 10.8)	7	
Most	0.90 (0.18, 4.56)	5	
(p for linear trend)	≥0.05		
Maternal exposure during pregnancy			
Never	1.00	9	
Least	3.53 (0.22, 58.1)	3	
Most	14.3 (0.92, 224)	7	
(p for linear trend)	<0.05		
Population ≤14 yr of age in 3 areas north England, United Kingdom			McKinney et al. (1991)
<b>Acute lymphocytic leukemia and NHL</b>			
Maternal occupation exposure to TCE			
Preconception	1.16 (0.13, 7.91)	2	
Paternal occupational exposure to TCE			
Preconception	2.27 (0.84, 6.16)	9	
Periconception and gestation	4.49 (1, 15, 21)	7	
Postnatal	2.66 (0.82, 9.19)	7	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-75. Selected results from epidemiologic studies of TCE exposure and childhood leukemia (continued)**

	Relative risk (95% CI)	n observed events	Reference(s)
Los Angeles Cancer Surveillance Program			Lowengart et al. (1987)
<b>Acute lymphocytic and nonlymphocytic leukemia, ≤10 yr of age</b>			
Maternal occupational exposure to TCE		0	
Paternal occupational exposure to TCE			
One year before pregnancy	2.0 ( <i>p</i> = 0.16)	6/3 <sup>b</sup>	
During pregnancy	2.0 ( <i>p</i> = 0.16)	6/3 <sup>b</sup>	
After delivery	2.7 (0.64, 15.6)	8/3 <sup>b</sup>	
<b>Geographic based studies</b>			
Two study areas in Endicott, NY			ATSDR (2006a)
<b>Leukemia, ≤19 yr of age</b>	Not reported	<6	
Population in New Jersey			
<b>Acute lymphocytic leukemia</b>			
Maximum estimated TCE concentration (ppb) in municipal drinking water			Cohn et al. (1994b)
Males			
<0.1	1.00	45	
0.1–0.5	0.91 (0.53, 1.57)	16	
≥5.0	0.54 (0.17, 17.7)	3	
Females			
<0.1	1.00	25	
0.1–0.5	1.85 (1.03, 3.70)	22	
≥5.0	2.36 (1.03, 5.45)	7	
Resident of Tucson Airport Area, AZ			ADHS (1990, 1995)
<b>Leukemia, ≤19 yr of age</b>			
1970–1986	1.48 (0.74, 2.65)	11	
1987–1991	0.80 (0.31, 2.05)	3	
Resident of West Central Phoenix, AZ			Aickin et al. (1992)
<b>Leukemia, ≤19 yr of age</b>	1.95 (1.43, 2.63)	38	

1  
2  
3  
4  
5

<sup>a</sup>Internal referents, live born children among female workers not exposed to organic solvents.

<sup>b</sup>Discordant pairs.


1           The geographic based studies for adult lymphopoietic (see Table 4-74) or childhood  
2 leukemias (see Table 4-75) do not greatly contribute to the overall weight of evidence. While  
3 some studies observed statistically significantly elevated risks for NHL or childhood cancer,  
4 these studies generally fulfilled only the minimal of evaluation criteria with questions raised  
5 about subject selection (Morgan and Cassady, 2002), their use of less sophisticated exposure  
6 assessment approaches and associated assumption of an average exposure to all subjects (all  
7 studies), and few cases with high level parental exposure (all studies).

#### 4.6.1.1.8. Meta-analysis of NHL risk

9           Meta-analysis is adopted as a tool for examining the body of epidemiologic evidence on  
10 NHL and TCE exposure and to identify possible sources of heterogeneity. The meta-analysis of  
11 NHL examines 17 cohort and case-control studies identified through a systematic review and  
12 evaluation of the epidemiologic literature on TCE exposure (Anttila et al., 1995; Axelson et al.,  
13 1994; Boice et al., 1999; Cocco et al., 2010; Greenland et al., 1994; Hansen et al., 2001; Hardell  
14 et al., 1994; Miligi et al., 2006; Morgan et al., 1998; Nordström et al., 1998; Persson and  
15 Fredrikson, 1999; Purdue et al., 2011; Raaschou-Nielsen et al., 2003; Radican et al., 2008;  
16 Siemiatycki, 1991; Wang et al., 2009; Zhao et al., 2005) and two studies as alternatives (Blair et  
17 al., 1998; Boice et al., 2006b). These 19 studies of NHL and TCE had high likelihood of  
18 exposure, were judged to have met, to a sufficient degree, the criteria of epidemiologic design  
19 and analysis, and reported estimated risks for overall TCE exposure; 13 of these studies, also,  
20 presented estimated NHL risk with high level TCE exposure (Anttila et al., 1995; Axelson et al.,  
21 1994; Boice et al., 1999; Cocco et al., 2010; Hansen et al., 2001; Miligi et al., 2006; Morgan et  
22 al., 1998; Purdue et al., 2011; Raaschou-Nielsen et al., 2003; Radican et al., 2008; Siemiatycki,  
23 1991; Wang et al., 2009; Zhao et al., 2005). Full details of the systematic review, criteria to  
24 identify studies for including in the meta-analysis, and meta-analysis methodology and findings  
25 are discussed in Appendices B and C.

26           The meta-analyses of the overall effect of TCE exposure on NHL suggest a small, robust,  
27 and statistically significant increase in NHL risk. The summary estimate from the primary  
28 random effect meta-analysis (RR<sub>m</sub>) was 1.23 (95% CI: 1.07, 1.42) (see Figure 4-16). This result  
29 and its statistical significance were not influenced by individual studies. Removal of individual  
30 studies resulted in RR<sub>m</sub> estimates between 1.18 (with the removal of Hansen et al., 2001) to 1.27  
31 (with the removal of Miligi et al. (2006) or Cocco et al. (2010)), and lower 95% CIs excluded 1.0  
32 (all *p*-values were *p* < 0.02). The result is similarly not sensitive to individual risk ratio estimate


1 selections. Use of six alternative selections, individually, resulted in RRm estimate that ranged  
2 from 1.20 (95% CI: 1.03, 1.39) (with estimated overall RR for incidence in Zhao et al., 2005) to



**Figure 4-16. Meta-analysis of NHL and overall TCE exposure.** The summary estimate is in the bottom row. Symbol sizes reflect relative weights of the studies. The horizontal midpoint of the bottom diamond represents the RRM estimate.

1 1.28 (95% CI: 1.09, 1.49) (with Raaschou-Nielsen et al. (2003) subgroup). Nor was the RRM  
2 estimate highly sensitive to restriction of the meta-analysis to only those studies for which RR  
3 estimates for the traditional definition of NHL were available. An alternate analysis which  
4 omitted Miligi (which included CLLs), Nordstrom (which was a study of hairy cell leukemias),  
5 Persson and Frederikson (for which the classification system not specified), and Greenland  
6 (which included Hodgkin lymphomas) and which included Boice (2006b) instead of Zhao  
7 (which included all lymphohematopoietic cancers) yielded an RRM estimate of 1.27 (95% CI:  
8 1.05, 1.55). Meta-analysis of the highest exposure groups, either duration, intensity, or their  
9 product, cumulative exposure, results in an RRM of 1.43 (95% CI: 1.13, 1.82), which is greater  
10 than the RRM from the overall exposure analysis, and provides additional support for an  
11 association between NHL and TCE (see Figure 4-17). No single study was overly influential;  
12 removal of individual studies resulted in RRM estimates that were all statistically significant (all  
13 with  $p \leq 0.025$ ) and that ranged from 1.38 (with the removal of Purdue et al. (2011)) to 1.57  
14 (with the removal of Cocco et al. (2010)). In addition, the RRM estimate was not highly  
15 sensitive to alternate RR estimate selections. Use of the 9 alternate selections, individually,  
16 resulted in RRM estimates that were all statistically significant (all with  $p < 0.025$ ) and all in the  
17 narrow range from 1.40 (95% CI: 09, 1.80) (with Blair et al. (1998) incidence relative risk  
18 instead of Radican et al. (2008) mortality hazard ratio) to 1.49 (95% CI: 1.14, 1.93) (with Hansen  
19 et al. [2001] duration). The highest exposure category groups have a reduced likelihood for  
20 exposure misclassification because they are believed to represent a greater differential TCE  
21 exposure compared to people identified with overall TCE exposure. Observation of greater risk  
22 associated with higher exposure category compared to overall (typically any versus none)  
23 exposure comparison additionally suggests an exposure-response gradient between NHL and  
24 TCE, although estimation of a level of exposure associated with the summary or meta-relative  
25 risk is not possible.

26 Low-to-moderate heterogeneity in RRM is observed across the results of the 17 studies in  
27 the meta-analysis of the overall effect of TCE and the 13 studies with highest exposure groups,  
28 but it was not statistically significant ( $p = 0.16$  and  $p = 0.30$ , respectively). The  $I^2$ -values were  
29 26% for overall exposure and 14% for highest exposure groups, suggesting low-to-moderate and  
30 low heterogeneity, respectively. To investigate the heterogeneity, subgroup analyses were done  
31 examining the cohort and case-control studies separately. Difference between cohort and case-  
32 control studies could explain much of the observed heterogeneity. In the subgroup analysis of  
33 overall exposure and of highest exposure groups, increased risk of NHL was strengthened in  
34 analysis limited to cohort studies and reduced in the case-control study analysis. Examination of  
35 heterogeneity in cohort and case-control studies of overall exposure separately was not



**Figure 4-17. Meta-analysis of NHL and TCE exposure—highest exposure groups.** The summary estimate is in the bottom row. Symbol sizes reflect relative weights of the studies. The horizontal midpoint of the bottom diamond represents the RRm estimate.

1 statistically significant in either case ( $I^2$ -values for the cohort studies were 12%, suggesting low  
2 heterogeneity and 27% for the case-control studies, suggesting low-to-moderate heterogeneity)  
3 although some may be present given that statistical tests of heterogeneity are generally  
4 insensitive in cases of minor heterogeneity. Subgroup analyses examining the cohort and case-  
5 control studies highest exposure groups, separately, showed no residual heterogeneity in the  
6 cohort subgroup ( $I^2 = 0\%$ ) and moderate heterogeneity in the case-control subgroup ( $I^2$ -value  
7 was 53%) that was not statistically significant ( $p = 0.08$ ). Although no further attempt was made  
8 to quantitatively investigate potential sources of heterogeneity, the removal of the Cocco et al.  
9 (2010) study, an influential study, eliminates all of the heterogeneity, suggesting that the RR  
10 estimate for the highest exposure group from that study is a relative outlier.

11 In general, sources of heterogeneity are uncertain and may reflect several features known  
12 to influence epidemiologic studies. Study design itself is unlikely to be an underlying cause of  
13 heterogeneity and, to the extent that it may explain some of the differences across studies, is  
14 more probably a surrogate for some other difference(s) across studies that may be associated  
15 with study design. Furthermore, other potential sources of heterogeneity may be masked by the  
16 broad study design subgroupings. The true source(s) of heterogeneity across these studies is an  
17 uncertainty.

18 One reason may be differences in exposure assessment and in overall TCE exposure  
19 concentration between cohort and case-control studies. Several cohort and case-control studies  
20 included TCE assignment from information on job and task exposures, e.g., a JEM (Boice et al.,  
21 1999; Boice et al., 2006b; Cocco et al., 2010; Miligi et al., 2006; Morgan et al., 1998; Purdue et  
22 al., 2011; Radican et al., 2008; Siemiatycki, 1991; Wang et al., 2009; Zhao et al., 2005), or from  
23 an exposure biomarker in either breath or urine (Anttila et al., 1995; Axelson et al., 1994; Hansen  
24 et al., 2001). Three case-control studies (Hardell et al., 1994; Nordström et al., 1998; Persson  
25 and Fredrikson, 1999) relied on self-reported TCE exposure. No information is available to  
26 judge the degree of possible misclassification bias associated with a particular exposure  
27 assessment approach; it is quite possible that in some cohort studies, in which past exposure is  
28 inferred from various data sources, exposure misclassification may be as great as in population-  
29 based or hospital-based case-control studies. In addition, a low overall TCE exposure prevalence  
30 is anticipated in population case-control studies which would typically assess a large number of  
31 workplaces and operations, where exposures are less well defined, and where case and control  
32 subjects identified as exposed to TCE probably have minimal contact (NRC, 2006). Observed  
33 higher risk ratios with higher exposure categories in NHL case-control studies support exposure  
34 differences as a source of heterogeneity.



1 Diagnostic inaccuracies are likely another source of heterogeneity in the meta-analysis  
2 through study differences in NHL groupings and in lymphoma classification schemes, although  
3 restricting the meta-analysis to only those studies for which RR estimates based on the traditional  
4 NHL definition were available did not eliminate all heterogeneity. All studies include a broad  
5 but slightly different group of lymphosarcoma, reticulum-cell sarcoma, and other lymphoid  
6 tissue neoplasms (Codes 200 and 202), except Nordstrom et al. (1998), Zhao et al. (2005), and  
7 Greenland et al. (1994). Cohort studies have some consistency in coding NHL, with NHL  
8 defined as lymphosarcoma and reticulum-cell sarcoma (200) and other lymphoid tissue  
9 neoplasms (202) using the ICD, Revision 7, 200 and 202—four studies (Anttila et al., 1995;  
10 Axelson et al., 1994; Hansen et al., 2001; Raaschou-Nielsen et al., 2003), ICD-Adapted,  
11 Revision 8 (Blair et al., 1998), and ICD-7, -8, -9, and -10, per the version in use at the time of  
12 death Morgen et al., 1998 (as presented in Boice et al., 1999; Mandel et al., 2006; Radican et al.,  
13 2008), as does the case-control study of Siemiatycki (1991) whose coding scheme for NHL is  
14 consistent with ICD 9, 200 and 202. Case-control studies, on the other hand, have adopted other  
15 classification systems for defining NHL including the NCI Working Formulation (Miligi et al.,  
16 2006), Rappaport (Hardell et al., 1994), or else do not identify the classification system for  
17 defining NHL (Persson and Fredrikson, 1999). Cocco et al. (2010) used the WHO/Revised  
18 European-American Lymphoma (REAL) classification system, which reclassifies lymphocytic  
19 leukemias and NHLs as lymphomas of B-cell or T-cell origin and considers CLLs and multiple  
20 myelomas as (non-Hodgkin) lymphomas; however, we were able to obtain results generally  
21 consistent with the traditional NHL definition from Dr. Cocco, although lymphomas not  
22 otherwise specified were excluded. Wang et al. (2009) defined NHL using ICD-O-2 codes  
23 (M-9590-9595, 9670-9688, 9690-9698, 9700-9723), which is consistent with the traditional  
24 definition of NHL (i.e., ICD-7, -8, -9 codes 200 + 202). Purdue et al. (2011) used ICD-O-3  
25 codes 967-972, which is generally consistent with the traditional definition of NHL, although  
26 this grouping does not include the malignant lymphomas of unspecified type coded as  
27 M-9590-9599.

28 There is some evidence of potential publication bias in this data set; however, it is  
29 uncertain that this is actually publication bias rather than an association between standard error  
30 and effect size resulting for some other reason, e.g., a difference in study populations or  
31 protocols in the smaller studies. Furthermore, if there is publication bias in this data set, it does  
32 not appear to account completely for the finding of an increased NHL risk.

33 NRC (2006) deliberations on trichloroethylene commented on two prominent evaluations  
34 of the then-current epidemiologic literature using meta-analysis techniques. These studies were  
35 by Wartenberg et al. (2000), and by Kelsh et al. (2005), submitted by Exponent-Health Sciences

1 to NRC during their deliberations and subsequently published in a paper on NHL (Mandel et al.,  
2 2006) and a paper on multiple myeloma and leukemia (Alexander et al., 2006). The NRC found  
3 weaknesses in the techniques used in each of these studies, and suggested that EPA conduct a  
4 new meta-analysis of the epidemiologic data on trichloroethylene using objective and transparent  
5 criteria so as to improve on the past analyses. EPA staff conducted their analysis according to  
6 NRC (2006) suggestions for transparency, systematic review criteria, and examination of both  
7 cohort and case-control studies. The EPA analysis of NHL analysis considered a larger number  
8 of studies than in the previous analyses (Mandel et al., 2006; Wartenberg et al., 2000), includes  
9 recently published studies (Boice et al., 2006b; Cocco et al., 2010; Miligi et al., 2006; Purdue et  
10 al., 2011; Radican et al., 2008; Wang et al., 2009; Zhao et al., 2005), and combines both cohort  
11 and case-control studies.  
12

#### 4.6.2. Animal Studies

13 The immunosuppressive and immunomodulating potential of TCE has not been fully  
14 evaluated in animal models across various exposure routes, over various relevant durations of  
15 exposure, across representative life stages, and/or across a wide variety of endpoints.  
16 Nevertheless, the studies that have been conducted indicate a potential for TCE-induced  
17 immunotoxicity, both following exposures in adult animals and during immune system  
18 development (i.e., in utero and preweaning exposures).  
19

##### 4.6.2.1.1. Immunosuppression

20 A number of animal studies have indicated that moderate to high concentrations of TCE  
21 over long periods have the potential to result in immunosuppression in animal models, dependant  
22 on species and gender. These studies are described in detail below and summarized in  
23 Table 4-76.  
24

##### 4.6.2.1.2. Inhalation exposures

25 Mature cross-bred dogs (5/group) were exposed to 0-, 200-, 500-, 700-, 1,000-, 1,500-, or  
26 2,000-ppm TCE for 1-hour or to 700 ppm TCE for 4 hours, by tracheal intubation under  
27 intravenous sodium pentobarbital anesthesia. An additional group of dogs was exposed by  
28 venous injection of 50 mg/kg TCE administered at a rate of 1 mL/minute (Hobara et al., 1984).  
29 Blood was sampled pre- and postexposure for erythrocyte and leukocyte counts. Marked,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 transient decreases in leukocyte counts were observed at all exposure levels 30 minutes after  
2 initiation of exposure. At the end of the exposure period, all types of leukocytes were decreased  
3 (by 85%); neutrophils were decreased 33%, and lymphocytes were increased 40%. There were  
4 no treatment-related changes in erythrocyte counts, hematocrit values, or thrombocyte counts.

**Table 4-76. Summary of TCE immunosuppression studies**

Exposure route/vehicle, duration, dose	NOAEL; LOAEL <sup>a</sup>	Results	Reference, species/strain sex/number
<b>Inhalation exposure studies</b>			
Single 1-h exposure to all dose groups; plus single 4-h exposure at 700 ppm <sup>b</sup> 0, 200, 500, 700, 1,000, 1,500, or 2,000 ppm	LOAEL: 200 ppm	Marked transient ↓ leukocyte counts at all exposure levels 30 min after initiating exposure. At end of exposure, 85% ↓ leukocyte counts (33% ↓ neutrophils, 40% ↓ lymphocytes).	Hobara et al. (1984) Dog, cross-bred, both sexes, 5/group
Single 3-h exposure. Also, 3 h/d on 5 d at lowest dose 0, 2.6, 5.2, 10.6, 25.6, or 48 ppm	NOAEL: 2.6 ppm LOAEL: 5.2 ppm	Challenged with <i>Streptococcus zooepidemicus</i> to assess susceptibility to infection and <i>Klebsiella pneumoniae</i> to assess bacterial clearance. For single exposure: dose-related sig. ↑ mortality at ≥5.2 ppm over 14 d. Sig. ↓ in bactericidal activity at 10.6 ppm.	Aranyi et al. (1986) Mouse, CD-1 females, 4–5 wk old, approx. 30 mice/group, 5–10 replications; for pulmonary bactericidal activity assay, 17–24 mice/group
Single 3-h exposure. 0, 5, 10, 25, 50, 100, 200 ppm	NOAEL: 25 ppm LOAEL: 50 ppm	Challenged with <i>Streptococcus zooepidemicus</i> to assess susceptibility to infection and bacterial clearance. For single exposure: dose-related sig. ↑ mortality at ≥50 ppm over 20 d. Dose dependent responses also observed in the clearance of bacteria from the lung at ≥50 ppm, the number of mice with delayed bacterial clearance at various postinfection time points at ≥50 ppm, and the phagocytic function of alveolar macrophages at 200 ppm.	Selgrade and Gilmour (2010) Mouse, CD-1 females, 5–6 wk old, at least 38 mice/group
Single 3-h exposure, 50–200 ppm <sup>c</sup>		Challenged with <i>Streptococcus zooepidemicus</i> . Dose-related ↑ mortality, bacterial antiphagocytic capsule formation, and bacterial survival. Dose-related impairment of alveolar macrophages; increased neutrophils in bronchoalveolar fluid at 3 d postinfection.	Park et al. (1993) (abstract) Mouse, CD-1, (sex and #/group not specified)
<b>4-wk, 6 h/d, 5 d/wk 0, 100, 300, or 1,000 ppm</b>	<b>NOAEL: 300 ppm LOAEL: 1,000 ppm</b>	<b>At 1,000 ppm, 64% ↓ plaque-forming cell assay response.</b>	<b>Woolhiser et al. (2006) Rat, Sprague-Dawley, female, 16/group</b>

**Table 4-76. Summary of TCE immunosuppression studies (continued)**

Exposure route/vehicle, duration, dose	NOAEL; LOAEL <sup>a</sup>	Results	Reference, species/strain sex/number
<b>Oral exposure studies</b>			
Gavage in 10% emulphor, 14 d, daily, 0, 24, or 240 mg/kg-day	LOAEL: 24 mg/kg-day	Sig. ↓ cell-mediated immune response to SRBC at both dose levels.	Sanders et al. (1982b) Mouse, CD-1, male, 9–12/group
<b>Drinking water with 1% emulphor, 4–6 mo 0, 0.1, 1.0, 2.5, or 5.0 mg/mL</b>	<b>LOAEL: 0.1 mg/kg-day</b>	<b>In females, humoral immunity ↓ at 2.5 and 5 mg/mL TCE, whereas cell-mediated immunity ↓ and bone marrow stem cell colonization ↓ at all four concentrations. The males were relatively unaffected after both 4 and 6 mo.</b>	<b>Sanders et al. (1982b) Mouse, CD-1, male and female, 7–25/group</b>
Gavage, 14 d, 0, 14.4, or 144 mg/kg-day chloral hydrate	NOAEL: 144 mg/kg-day	No treatment-related effects.	Kauffmann et al. (1982) Mouse, CD-1, male, 12/group
Drinking water, 90 d, 0, 0.07, or 0.7 mg/mL chloral hydrate. (M: 0, 16, or 160 mg/kg-day; F: 0, 18, or 173 mg/kg-day)	NOAEL: 0.07 mg/mL LOAEL: 0.7 mg/mL	Sig. ↓ cell-mediated immune response (plasma hemagglutination titers and spleen antibody-producing cells of mice sensitized to SRBC) in females at 0.7 mg/mL.	Kauffmann et al. (1982) Mouse, CD-1, male and female, 15–20/group
Drinking water, From mating to PND 21 or PND 56, (emulphor concentration not provided) 0 (emulphor), 1, or 10 ppm	LOAEL: 1 ppm	At 10 ppm, ↓ body weight and length at PND 21. IgM antibody response to SRBC challenge suppressed in both ♂ and ♀ pups at 10 ppm, and ♂ pups at 1 ppm, ↓ in splenic CD4+CD8-T-cells. At 56 PND, striking ↑ in natural killer cell activity seen at both doses.	Adams et al. (2003) (abstract) Mouse, B6C3F1, both sexes, numbers of pups not stated
Drinking water, from GD 0–3 or 8 wk of age, 0, 1,400, or 14,000 ppb	LOAEL: 1,400 ppb	Suppressed PFC responses in both sexes and ages at 14,000 ppb, in males at both ages at 1,400 ppb, and in females at 8 wk at 1,400 ppb. Numbers of spleen B220+ cells ↓ at 3 wk at 14,000 ppb. Pronounced ↑ thymus T-cell populations at 8 wk.	Peden-Adams et al. (2006) Mouse, B6C3F1, dams and both sexes offspring, 5 litters/group; 5–7 pups/group at 3 wk; 4–5 pups/sex/group at 8 wk

**Table 4-76. Summary of TCE immunosuppression studies (continued)**

Exposure route/vehicle, duration, dose	NOAEL; LOAEL <sup>a</sup>	Results	Reference, species/strain sex/number
Drinking water, from GD 0 to 7–8 wk of age; 0, 0.5, or 2.5 mg/mL	LOAEL: 0.5 mg/mL	At 0.5 mg/mL: Sig ↓ postweaning weight; sig. ↑ IFN $\gamma$ produced by splenic CD4+ cells at 5–6 wk; sig ↓ splenic CD8+ and B220+ lymphocytes; sig. ↑ IgG2a and histone; sig. altered CD4–/CD8– and CD4+/CD8+ thymocyte profile At 2.5 mg/mL: Sig ↓ postweaning weight; sig. ↑ IFN $\gamma$ produced by splenic CD4+ and CD8+ cells at 4–5 and 5–6 wk; sig ↓ splenic CD4+, CD8+, and B220+ lymphocytes; sig. altered CD4+/CD8+ thymocyte profile.	Blossom and Doss (2007) Mouse, MRL +/+, dams and both sexes offspring, 3 litters/group; 8–12 pups/group
Drinking water, from GD 0 to PND 42; 0 or 0.1 mg/mL; maternal dose = 25.7 mg/kg-day; offspring PND 24–42 dose = 31.0 mg/kg-day	LOAEL: 0.1 mg/mL	At 0.1 mg/mL: at PND 20, sig. ↑ thymocyte cellularity and distribution, associated with sig. ↑ in thymocyte subset distribution; sig. ↑ reactive oxygen species generation in total thymocytes; sig. ↑ in splenic CD4+ T-cell production of IFN- $\gamma$ and IL-2 in females and TNF- $\alpha$ in males at PND 42.	Blossom et al. (2008) Mouse, MRL +/+, dams and both sexes offspring, 8 litters/group; 3–8 pups/group
Drinking water, from GD 0 to 12 mo of age; 0 (1% emulphor), 1,400, or 14,000 ppb	LOAEL: 1,400 ppb	At 1,400 ppb: splenic CD4–/CD8– cells sig. ↑ in females; thymic CD4+/CD8+ cells sig. ↓ in males; 18% ↑ in male kidney weight. At 14,000 ppb: thymic T-cell subpopulations (CD8+, CD4/CD8–, CD4+) sig. ↓ in males.	Peden-Adams et al. (2008) Mouse, MRL +/+, dams and both sexes offspring, unknown # litters/group, 6–10 offspring/sex/group
<b>Intraperitoneal injection exposure studies</b>			
3 d, single daily injection, 0, 0.05, 0.5, or 5 mmol/kg/day	NOAEL: 0.05 mmol/kg/day LOAEL: 0.5 mmol/kg/day	↓ natural killer cell activity at 0.5 and 5 mmol/kg/day. ↓ splenocyte counts at 5 mmol/kg/day.	Wright et al. (1991) Rat, Sprague-Dawley
3 d, single daily injection, 0 or 10 mmol/kg/day	LOAEL: 10 mmol/kg/day	↓ natural killer cell activity and ↓ spleen weights at 10 mmol/kg/day.	Wright et al. (1991) Mouse, B6C3F1

**Bolded studies** carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup>NOAEL and LOAEL are based upon reported study findings.

<sup>b</sup>Inhalation, tracheal intubation under anesthesia.

<sup>c</sup>Exact dose levels not specified.

↓, ↑ = decreased, increased; GD = gestation day; PFC = plaque-forming cell; sig. = statistically significant; SRBC = sheep red blood cells.

1 In a study that examined the effects of a series of inhaled organic chemical air  
2 contaminants on murine lung host defenses, Aranyi et al. exposed female CD-1 mice to single  
3 3-hour exposures of TCE at time-weighted concentrations of 0, 2.6, 5.2, 10.6, 25.6, or 48 ppm  
4 (Aranyi et al., 1986). Additionally, at the dose at which no adverse treatment-related effect  
5 occurred with a single exposure (i.e., 2.6 ppm), a multiple exposure test (5 days, 3 hours/day)  
6 was conducted. Susceptibility to *Streptococcus zooepidemicus* aerosol infection and pulmonary  
7 bactericidal activity to inhaled *Klebsiella pneumoniae* were evaluated. There was a significant  
8 ( $p < 0.0001$ ) treatment by concentration interaction for mortality, with the magnitude of the  
9 effect increasing with concentration. A significant ( $p < 0.0001$ ) treatment by concentration  
10 interaction was also found for bactericidal activity. Single 3-hour exposures at 10.6, 25.6, and  
11 48 ppm resulted in significant increases in mortality, although increases observed after single  
12 exposures at 5.2 or 2.6 ppm or five exposures at 2.6 ppm were not significant. Pulmonary  
13 bactericidal activity was significantly decreased after a single exposure at 10.6 ppm, but single  
14 exposures to 2.6 or 5.2 ppm resulted in significant increases.

15 Suppression of pulmonary host defenses and enhanced susceptibility to respiratory  
16 bacterial infection was studied in female CD-1 mice by Selgrade and Gilmour (2010). The mice  
17 (5–6 weeks of age; at least 38 per exposure group) were exposed via inhalation for 3 hours to  
18 concentrations of 0, 5, 10, 25, 50, 100, or 200 ppm TCE. The mice were then challenged by  
19 aerosol doses of *Streptococcus zooepidemicus*. Bacterial clearance (based upon organisms  
20 present in lung lavage fluid) and a phagocytic index (percentage of phagocytic cells in lung  
21 lavage fluid and the number of bacteria ingested per phagocytic cell) were assessed. Mortality  
22 due to infection was significantly increased with TCE exposure concentration at exposures of  
23 50 ppm and higher (NOAEL = 25 ppm). Dose-dependent responses were also observed for the  
24 clearance of bacteria from the lung at  $\geq 25$  ppm, the number of mice with delayed bacterial  
25 clearance at various postinfection time points at  $\geq 25$  ppm, and the phagocytic function of  
26 alveolar macrophages at 200 ppm. The higher NOAEL for mortality observed in this study  
27 compared to Aranyi et al. (1986), i.e., 25 ppm versus 5 ppm, was attributed to the use of  
28 unencapsulated bacteria in this study; the study authors suggested that this may be more  
29 representative of the human condition.

30 In a host-resistance assay, CD-1 mice (sex and number/group not specified) exposed to  
31 TCE by inhalation for 3 hours at 50–200 ppm were found to be more susceptible to increased  
32 infection following challenge with *Streptococcus zooepidemicus* administered via aerosol (Park  
33 et al., 1993). Dose-related increases in mortality, bacterial antiphagocytic capsule formation, and  
34 bacterial survival were observed. Alveolar macrophage phagocytosis was impaired in a dose-

1 responsive manner, and an increase in neutrophils in bronchoalveolar lavage fluid was observed  
2 in exposed mice 3 days post infection.

3 A guideline (OPPTS 870.3800) 4-week inhalation immunotoxicity study was conducted  
4 in female Sprague-Dawley rats (Woolhiser et al., 2006). The animals (16/group) were exposed  
5 to TCE at nominal levels of 0, 100, 300, or 1,000 ppm for 6 hours/day, 5 days/week. Effects on  
6 the immune system were assessed using an antigen response assay, relevant organs weights,  
7 histopathology of immune organs, and hematology parameters. Four days prior to study  
8 termination, the rats were immunized with sheep red blood cells (SRBC), and within 24 hours  
9 following the last exposure to TCE, a plaque forming cell assay was conducted to determine  
10 effects on splenic anti-SRBC IgM response. Minor, transient effects on body weight and food  
11 consumption were noted in treated rats for the first 2 weeks of exposure. Mean relative liver and  
12 kidney weights were significantly ( $p = 0.05$ ) increased at 1,000 ppm as compared to control,  
13 while lung, spleen, and thymus weights were similar to control. No treatment-related effects  
14 were observed for hematology, white blood cell differential counts, or histopathological  
15 evaluations (including spleen, thymus, and lung-associated lymph nodes). At 1,000 ppm, rats  
16 demonstrated a 64% decrease in plaque forming cell assay response. Lactate dehydrogenase,  
17 total protein levels, and cellular differentiation counts evaluated from bronchoalveolar lavage  
18 (BAL) samples were similar between control and treated groups. A phagocytic assay using BAL  
19 cells showed no alteration in phagocytosis, although these data were not considered fully reliable  
20 since (1) the number of retrieved macrophage cells was lower than expected and pooling of  
21 samples was conducted and (2) samples appear to have been collected at 24 hours after the last  
22 exposure (rather than within approximately 2 hours of the last exposure), thereby allowing for  
23 possible macrophage recovery. The NOAEL for this study was considered by the study authors  
24 to be 300 ppm, and the LOAEL was 1,000 ppm; however, the effect level may have actually  
25 been lower. It is noted that the outcome of this study does not agree with the studies by Aranyi  
26 et al. (1986) and Park et al. (1993), both of which identified impairment of macrophage  
27 phagocytic activity in BAL following inhalation TCE exposures.

#### 28 **4.6.2.1.3. Oral exposures**

29 In a study by Sanders et al., TCE was administered to male and female CD-1 mice for 4  
30 or 6 months in drinking water at concentrations of 0, 0.1, 1, 2.5, or 5 mg/mL (Sanders et al.,  
31 1982b). In females, humoral immunity was suppressed at 2.5 and 5 mg/mL, while cell-mediated  
32 immunity and bone marrow stem cell activity were inhibited at all dose levels. Male mice were  
33 relatively unaffected either at 4 or 6 months, even though a preliminary study in male CD-1 mice

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 (exposed to TCE for 14 days by gavage at 0, 24, or 240 mg/kg-day) had demonstrated a decrease  
2 in cell-mediated immune response to SRBC in male mice at both treatment levels.

3 A significant decrease in humoral immunity (as measured by plasma hemagglutination  
4 titers and the number of spleen antibody producing cells of mice sensitized to sheep  
5 erythrocytes) was observed by Kaufmann et al. (1982) in female CD-1 mice (15–20/group)  
6 following a 90-day drinking water exposure to 0, 0.07, or 0.7 mg/mL (equivalent to 0, 18, or  
7 173 mg/kg) chloral hydrate, a metabolite of TCE. Similar responses were not observed in male  
8 CD-1 mice exposed for 90 days in drinking water (at doses of 0, 16, or 160 mg/kg-day), or when  
9 administered chloral hydrate by gavage to 12/group for 14 days at 14.4 or 144 mg/kg-day.

10 The potential for developmental immunotoxicity was assessed in B6C3F1 mice  
11 administered TCE in drinking water at dose levels of 0, 1,400 or 14,000 ppb from gestation day  
12 (GD) 0 to either 3 or 8 weeks of age (Adams et al., 2003 [preliminary data]; Peden-Adams et al.,  
13 2006). At 3 and 8 weeks of age, offspring lymphocyte proliferation, NK cell activity, SRBC-  
14 specific IgM production (plaque-forming cell [PFC] response), splenic B220+ cells, and thymus  
15 and spleen T-cell immunophenotypes were assessed. Delayed-typed hypersensitivity and  
16 autoantibodies to double-stranded DNA (dsDNA) were evaluated in offspring at 8 weeks of age.  
17 Observed positive responses consisted of suppressed PFC responses in males at both ages and  
18 both TCE treatment levels, and in females at both ages at 14,000 ppb and at 8 weeks of age at  
19 1,400 ppb. Spleen numbers of B220+ cells were decreased in 3-week old pups at 14,000 ppb.  
20 Pronounced increases in all thymus T-cell subpopulations (CD4+, CD8+, CD4+/CD8+, and  
21 CD4-/CD8-) were observed at 8 weeks of age. Delayed hypersensitivity response was  
22 increased in 8-week old females at both treatment levels and in males at 14,000 ppb only. No  
23 treatment-related increase in serum anti-dsDNA antibody levels was found in the offspring at 8  
24 weeks of age.

25 In a study designed to examine potential susceptibility of the young (Blossom and Doss,  
26 2007), TCE was administered to groups of pregnant MRL +/+ mice in drinking water at  
27 occupationally-relevant levels of 0, 0.5, or 2.5 mg/mL. A total of 3 litters per treatment group  
28 were maintained following delivery (i.e., a total of 11 pups at 0 mg/mL TCE, 8 pups at  
29 0.5 mg/mL TCE, and 12 pups at 2.5 mg/mL TCE), and TCE was continuously administered to  
30 the offspring until young adulthood (i.e., 7–8 weeks of age). Although there were no effects on  
31 reproduction, offspring postweaning body weights were significantly decreased in both treated  
32 groups. Additionally, TCE exposure was found to modulate the immune system following  
33 developmental and early life exposures. Decreased spleen cellularity and reduced numbers of  
34 CD4+, CD8+, and B220+ lymphocyte subpopulations were observed in the postweaning  
35 offspring. Thymocyte development was altered by TCE exposures, as evidenced by significant

1 alterations in the proportions of double-negative subpopulations and inhibition of in vitro  
2 apoptosis in immature thymocytes. TCE was also shown to induce a dose-dependent increase in  
3 CD4+ and CD8+ T-lymphocyte IFN $\gamma$  in peripheral blood by 4–5 weeks of age, although these  
4 effects were no longer observed at 7–8 weeks of age. Serum antihistone autoantibodies and total  
5 IgG<sub>2a</sub> were significantly increased in treated offspring; however, no histopathological signs of  
6 autoimmunity were observed in the liver and kidneys at sacrifice.

7 This increase in T-cell hyperactivity was further explored in a study by Blossom et al.  
8 (2008). In this study, MRL +/+ mice were treated with 0 or 0.1 mg/mL TCE in the drinking  
9 water. Based on drinking water consumption data, average maternal doses of TCE were  
10 25.7 mg/kg-day, and average offspring (PND 24–42) doses of TCE were 31.0 mg/kg-day.  
11 Treatment was initiated at the time of mating, and continued in the females (8/group) throughout  
12 gestation and lactation. Pups were weaned at PND 24, and the offspring were continued on  
13 drinking water treatment in a group-housed environment until study termination (PND 42).  
14 Subsets of offspring were sacrificed at PND 10 and 20, at which time developmental and  
15 functional endpoints in the thymus were evaluated (i.e., total cellularity, CD4+/CD8+ ratios,  
16 CD24 differentiation markers, and double-negative subpopulation counts). Indicators of  
17 oxidative stress were measured in the thymus at PND 10 and 20, and in the brain at PND 42.  
18 Mitogen-induced intracellular cytokine production by splenic CD4+ and CD8+ T-cells was  
19 evaluated in juvenile mice and brain tissue was examined at PND 42 for evidence of  
20 inflammation. Behavioral testing was also conducted; these methods and results are described in  
21 Section 4.3. TCE treatment did not affect reproductive capacity, parturition, or ability of dams to  
22 maintain litters. The mean body weight of offspring was not different between the control and  
23 treated groups. Evaluation of the thymus identified a significant treatment-related increase in  
24 cellularity, accompanied by alterations in thymocyte subset distribution, at PND 20 (sexes  
25 combined). TCE treatment also appeared to promote T-cell differentiation and maturation at  
26 PND 42, and ex vivo evaluation of cultured thymocytes indicated increased reactive oxygen  
27 species (ROS) generation. Evaluation of peripheral blood indicated that splenic CD4+ T-cells  
28 from TCE-exposed PND 42 mice produced significantly greater levels of IFN- $\gamma$  and IL-2 in  
29 males and TNF- $\alpha$  in both sexes. There was no effect on cytokine production on PND 10 or 20.  
30 The dose of TCE that resulted in adverse offspring outcomes in this study (i.e., 0.1 mg/mL,  
31 equivalent to 25.7–31.0 mg/kg-day) is comparable to that which has been previously  
32 demonstrated to result in immune system alterations and autoimmunity in adult MRL +/+ mice  
33 (i.e., 0.1 mg/mL, equivalent to 21 mg/kg-day; (Griffin et al., 2000b).

34 Another study that examined the effects of developmental exposure to TCE on the  
35 MRL+/+ mouse was conducted by Peden-Adams et al. (2008). In this study, MRL/MpJ (i.e.,

1 MRL +/-) mice (unspecified number of dams/group) were exposed to TCE (solubilized with 1%  
2 emulphor) in drinking water at levels of 0, 1,400, or 14,000 ppb from GD 0 and continuing until  
3 the offspring were 12 months of age. TCE concentrations in the drinking water were reported to  
4 be analytically confirmed. Endpoints evaluated in offspring at 12 months of age included final  
5 body weight; spleen, thymus, and kidney weights; spleen and thymus lymphocyte  
6 immunophenotyping (CD4 or CD8); splenic B-cell counts; mitogen-induced splenic lymphocyte  
7 proliferation; serum levels of autoantibodies to dsDNA and glomerular antigen (GA),  
8 periodically measured from 4–12 months of age; and urinary protein measures. Reported sample  
9 sizes for the offspring measurements varied from 6–10 per sex per group; the number of source  
10 litters represented within each sample was not specified. The only organ weight alteration was  
11 an 18% increase in kidney weight in the 1,400 ppb males. Splenic CD4–/CD8– cells were  
12 altered in female mice (but not males) at 1,400 ppm only. Splenic T-cell populations, numbers  
13 of B220+ cells, and lymphocyte proliferation were not affected by treatment. Populations of  
14 thymic T-cell subpopulations (CD8+, CD4–/CD8–, and CD4+) were significantly decreased in  
15 male but not female mice following exposure to 14,000-ppb TCE, and CD4+/CD8+ cells were  
16 significantly reduced in males by treatment with both TCE concentrations. Autoantibody levels  
17 (anti-dsDNA and anti-GA) were not increased in the offspring over the course of the study,  
18 indicating that TCE did not contribute to the development of autoimmune disease markers  
19 following developmental exposures that continued into adult life.

20 Overall, the studies by Peden-Adams et al. (2006; 2008), Blossom and Doss (2007), and  
21 Blossom et al. (2008), which examined various immunotoxicity endpoints following exposures  
22 that spanned the critical periods of immune system development in the rodent, were generally  
23 not designed to assess issues such as posttreatment recovery, latent outcomes, or differences in  
24 severity of response that might be attributed to the early life exposures.

25

#### **4.6.2.1.4. Intraperitoneal administration**

26 Wright et al. reported that following 3 days of single intraperitoneal injections of TCE in  
27 Sprague-Dawley rats at 0, 0.05, 0.5, or 5 mmol/kg/day and B6C3F1 mice at 0 or 10  
28 mmol/kg/day, NK cell activity was depressed in the rats at the mid- and high-dose levels, and in  
29 the mice at the high dose level (Wright et al., 1991). Also at the highest dose levels tested,  
30 decreased splenocyte counts and relative spleen weight were observed in the rats and mice,  
31 respectively. In vitro assays demonstrated treatment-related decreases in splenocyte viability,  
32 inhibition of lipopolysaccharide-stimulated lymphocyte mitogenesis, and inhibited NK cell

1 activity suggesting the possibility that compromised immune function may play a role in  
2 carcinogenic responses of experimental animals treated with TCE.

#### 4.6.2.1.5. Hypersensitivity

3 Evidence of a treatment-related increase in delayed hypersensitivity response has been  
4 observed in guinea pigs following dermal exposures with TCE and in mice following exposures  
5 that occurred both during development and postnatally (see Table 4-77).

6 In a modified guinea pig maximization test, Tang et al. (2002) evaluated the contact  
7 allergenicity potential of TCE and three metabolites (trichloroacetic acid, trichloroethanol, and  
8 chloral hydrate) in 4 animals (FMMU strain, sex not specified) per group (Tang et al., 2002).  
9 Edema and erythema indicative of skin sensitization (and confirmed by histopathology) were  
10 observed. Sensitization rates were reported to be 71.4% for TCE and 58.3% for trichloroacetic  
11 acid, as compared to a reference positive control response rate (i.e., 100% for 2,4-  
12 dinitrochlorobenzene). In this study, the mean response scores for TCE, trichloroacetic acid, and  
13 2,4-dinitrochlorobenzene were 2.3, 1.1, and 6.0, respectively. TCE was judged to be a strong  
14 allergen and TCA was a moderate allergen, according to the criteria of Magnusson and Kligman  
15 (Magnusson and Kligman, 1969). Trichloroethanol and chloral hydrate were not found to elicit a  
16 dermal hypersensitivity response.

17 Immune-mediated hepatitis associated with dermal hypersensitivity reactions in the  
18 guinea pig following TCE exposures was characterized by Tang et al. (2008). In this study,  
19 FMMU strain female guinea pigs (5–6/group) were treated with intradermal injection of 0, 167,  
20 500, 1,500, or 4,500 mg/kg TCE or with a dermal patch containing 0 or 900 mg/kg TCE and  
21 sacrificed at 48 hours posttreatment. At the intradermal dose of 1,500 mg/kg, a significant  
22 increase ( $p < 0.05$ ) in serum AST level was observed. At 4,500 mg/kg, significantly ( $p < 0.01$ )  
23 increased ALT and AST levels were reported, and total protein and globulin decreased  
24 significantly ( $p < 0.05$ ). Histopathological examination of the liver revealed fatty degeneration,  
25 hepatic sinusoid dilation, and inflammatory cell infiltration. No changes were observed at the  
26 intradermal doses of 500 mg/kg or below, or the dermal patch dose of 900 mg/kg. A Guinea Pig  
27 Maximization Test was also conducted according to the procedures of Magnusson and Kligman  
28 on 10 FMMU females/group, in which the total TCE dosage from induction through challenge  
29 phases was below 340 mg/kg. TCE treatment resulted in dermal erythema and edema, and the  
30 sensitization rate was 66% (i.e., classified as a strong sensitizer). Significant increases ( $p < 0.05$ )  
31 in ALT, AST, lactate dehydrogenase, and relative liver weight, and significant decreases  
32 ( $p < 0.05$ ) in albumin, IgA, and GGT were observed. Additionally, hepatic lesions (diffuse  
33 ballooning changes without lymphocyte infiltration and necrotic hepatocytes) were noted. It was

1 concluded that TCE exposure to guinea pigs resulted in delayed type hypersensitivity reactions  
2 with hepatic injury that was similar to occupational medicamentosa-like dermatitis disorders  
3 observed in human occupational studies.

**Table 4-77. Summary of TCE hypersensitivity studies**

<b>Exposure route/vehicle, duration, dose</b>	<b>NOAEL; LOAEL<sup>a</sup></b>	<b>Results</b>	<b>Reference, species/strain sex/number</b>
Induction by single intradermal injection, then challenge by dermal application at 21 d 0 or 0.1 mL induction; 0 or 0.2 mL challenge TCE, TCA, TCOH, and chloral hydrate		Edema and erythema (confirmed by histopathology) indicative of skin sensitization for TCE (strong sensitizer) and TCA (moderate sensitizer)	Tang et al. (2002) Guinea pig, FMMU strain, sex not specified, 4/group
Intradermal injection, 0, 167, 500, 1,500, or 4,500 mg/kg  Dermal patch, 0 or 900 mg/kg  Hypersensitivity: total dose from induction through challenge <340 mg/kg	Intradermal NOAEL: 500 mg/kg Intradermal LOAEL: 1,500 mg/kg  Dermal patch NOAEL: 900 mg/kg	Intradermal injection: At 1,500 mg/kg: Sig. ↑ AST; at 4,500 mg/kg, sig. ↑ ALT and AST, sig. ↓ total protein and globulin; fatty degeneration of liver  Dermal patch: no effects of treatment  Hypersensitivity: sensitization rate of 66% (strong sensitizer), with edema and erythema; sig. ↑ ALT, AST, and lactate dehydrogenase; sig. ↑ relative liver weight; sig. ↓ albumin, IgA, and GGT; hepatic lesions (ballooning changes)	Tang et al., (2008) Guinea pig, FMMU strain, female, 5–6/group for intradermal/dermal patch study, 10/group for hypersensitivity study, female
<b>Drinking water, from GD 0 to 8 wk of age 0, 1,400, or 14,000 ppb</b>	<b>LOAEL: 1,400 ppb</b>	<b>Sig. ↑ swelling of foot pad in females at 1,400 and in both sexes at 14,000 ppb</b>	<b>Peden-Adams et al. (2006) Mouse, B6C3F1, both sexes, 5 litters/group; 4–5 pups/sex/group at 8 wks<sup>b</sup></b>

**Bolded study** carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup>NOAEL and LOAEL are based upon reported study findings.

<sup>b</sup>Subset of immunosuppression study.

↓, ↑ = decreased, increased, sig. = statistically significant.

5/12/11

*This document is a draft for review purposes only and does not constitute Agency policy.*

466

DRAFT—DO NOT CITE OR QUOTE

1 Also, as indicated in Section 4.6.2.1.2 above, in a developmental immunotoxicity-type  
2 study in B6C3F1 mice, administration of TCE in drinking water at dose levels of 0, 1,400, or  
3 14,000 ppb from GD 0 through to 8 weeks of age resulted in an increased delayed  
4 hypersensitivity response in 8-week old female offspring at both treatment levels and in males at  
5 the high dose of 14,000 ppb (Peden-Adams et al., 2006).

6 In an in vitro study that evaluated a number of chlorinated organic solvents, nonpurified  
7 rat peritoneal mast cells (NPMC) and rat basophilic leukemia (RBL-2H3) cells were sensitized  
8 with anti-DNP (dinitrophenol) monoclonal IgE antibody and then stimulated with  
9 DNP-conjugated bovine serum albumin plus TCE (Seo et al., 2008). TCE enhanced antigen-  
10 induced histamine release from NPMC and RBL-2H3 cells in a dose-related manner, and  
11 increased IL-4 and TNF- $\alpha$  production from the RBL-2H3 cells. In an in vivo study, i.p.-injected  
12 TCE was found to markedly enhance passive cutaneous anaphylaxis reaction in antigen-  
13 challenged rats. These results suggest that TCE increases histamine release and inflammatory  
14 mediator production from antigen-stimulated mast cells via the modulation of immune  
15 responses; TCE exposure may lead to the enhancement of allergic disease through this response.  
16

#### 4.6.2.1.6. Autoimmunity

17 A number of studies have been conducted to examine the effects of TCE exposure in  
18 mouse strains (i.e., MRL +/+, MRL -lpr, or NZB  $\times$  NZW) which are all known to be genetically  
19 susceptible to autoimmune disease. The studies have demonstrated the potential for TCE to  
20 induce autoimmune disease (as demonstrated in Table 4-78 which summarizes those studies  
21 which assessed serology, ex vivo assays of cultured splenocytes, and/or clinical or  
22 histopathology). These and other studies conducted in susceptible mouse strains have proven to  
23 be useful tools in exploring various aspects of the mode of action for this response.

24 Khan et al. used the MRL +/+ mouse model to evaluate the potential for TCE and one of  
25 its metabolites, dichloroacetyl chloride (DCAC) to elicit an autoimmune response (Khan et al.,  
26 1995). Female mice (4–5/group) were dosed by intraperitoneal injection with 10 mmol/kg TCE  
27 or 0.2 mmol/kg DCAC every 4<sup>th</sup> day for 6 weeks and then sacrificed. Spleen weights and IgG  
28 were increased. ANA and anti-ssDNA (single-stranded DNA) antibodies were detected in the  
29 serum of TCE- and DCAC-treated mice; anticardiolipin antibodies were detected in the serum of  
30 DCAC-treated mice. A greater magnitude of response observed with DCAC treatment suggested  
31 that the metabolite may be important to the mechanism of TCE-induced autoimmunity.

32 Other studies in female MRL +/+ mice (8/group) examined exposure via drinking water.  
33 In one of these studies, mice were treated with 2.5 or 5.0 mg/mL (455 or 734 mg/kg-day) TCE in



- 1 drinking water for up to 22 weeks (Gilbert et al., 1999; Griffin et al., 2000a). Serial sacrifices
- 2 were conducted at Weeks 4, 8, and 22. Significant increases in ANA and total serum

**Table 4-78. Summary of autoimmune-related studies of TCE and metabolites in mice and rats (by sex, strain, and route of exposure)<sup>a</sup>**

Number/group, vehicle, dose, duration	NOAEL; LOAEL <sup>b</sup>	Results			Reference
		Serology	Ex vivo assays of cultured splenocytes	Clinical and histopathology	
<b>Autoimmune-prone: female MRL +/- mice, drinking water</b>					
8 per group, 0, 2.5, or 5 mg/mL TCE (average 0, 455, or 734 mg/kg-day), 4, 8, or 22 wk	LOAEL: 2.5 mg/mL	Increased ANA at 4 and 8 wk, no difference between groups at 22 wk	Increased activated CD4+ T-cells and IFN- $\gamma$ secretion across doses at 4 wk, these effects were reversed at 22 wk; decreased IL-4 secretion (4 and 22 wk)	No evidence of liver or renal damage, based on serum alanine aminotransferase, sorbitol dehydrogenase, and blood urea nitrogen.	Griffin et al., (2000a)
8 per group, 0, 0.1, 0.5, or 2.5 mg/mL TCE (0, 21, 100, or 400 mg/kg-day), 4 or 32 wk	LOAEL: 0.1 mg/mL	Increased ANA in all treated groups at 4 wk, but not at 32 wk	Increased activated CD4+ T-cells (32 wk), IFN- $\gamma$ secretion (4 and 32 wk), no effect on IL-4 secretion	Extensive hepatic mononuclear cellular infiltrate in 0.5 and 2.5 mg/mL groups, and hepatocyte reactive changes in all treated groups at 32 wk.	Griffin et al., (2000b)
6–8 per group, 0, 0.1, or 0.9 mg/mL trichloroacetaldehyde hydrate (0, 24, or 220 mg/kg-day) or trichloroacetic acid (0, 27, or 205 mg/kg-day), 4 wk	LOAEL: 0.1 mg/mL	Increased ANA and antihistone antibodies at 0.9 mg/mL trichloroacetaldehyde hydrate <sup>c</sup>	Increased activated CD4+ T-cells at 0.1 and 0.9 g/mL doses of both metabolites. At 0.9 mg/mL, increased IFN- $\gamma$ secretion, no effect on IL-4 secretion	No evidence of liver or kidney damage, based on serum alanine aminotransferase, liver and kidney histology..	Blossom et al. (2004)
8 per group, 0, 0.1, 0.3, or 0.9 mg/mL trichloroacetaldehyde hydrate (0, 13, 46, or 143 mg/kg-day), 40 wk	LOAEL: 0.9 mg/mL	Slightly suppressed anti-ssDNA, anti-dsDNA, and antihistone antibody expression; differences not statistically significant	Increased activated CD4+ T-cells and increased INF- $\gamma$ secretion, no effect on IL-4 secretion	Diffuse alopecia, skin inflammation and ulceration, mononuclear cell infiltration, mast cell hyperplasia, dermal fibrosis. Statistically significant increase at 0.9 mg/mL dose group, but also increased at lower doses. No liver or kidney histopathology effects seen.	Blossom et al. (2007)

**Table 4-78. Summary of autoimmune-related studies of TCE and metabolites (by sex, strain, and route of exposure) (continued)**

Number/group, vehicle, dose, duration	NOAEL; LOAEL <sup>b</sup>	Results			Reference
		Serology	Ex vivo assays of cultured splenocytes	Clinical and histopathology	
5 per group, 0 or 0.5 mg/mL TCE (mean 60 µg/g-d), 48 wk	LOAEL: 0.5 mg/mL	Increased ANA after 24 wk but not statistically significant	Increased INF-γ secretion after 36 wk but not statistically significant	Hepatic necrosis; hepatocyte proliferation; leukocyte infiltrate in the liver, lungs, and kidneys; no difference in serum aminotransferase liver enzymes	Cai et al. (2008)
<b>Autoimmune-prone: male and female offspring MRL +/+ mice, drinking water</b>					
3 litters/group, 8–12 offspring/group; 0, 0.5, or 2.5 mg/mL, GD 0 to 7–8 wk of age	LOAEL: 0.5 mg/mL	Increased antihistone antibodies and total IgG <sub>2a</sub> in treated groups	Dose-dependent increase in INF-γ secretion at 4–5 wk of age but not 7–8 wk of age	No histopathological effects in liver or kidneys	Blossom and Doss (2007)
8 litters/group, 8–12 offspring/group; 0 or 0.1 mg/mL; maternal dose = 25.7 mg/kg-day; offspring PND 24–42 dose = 31.0 mg/kg-day; GD 0 to PND 42	LOAEL: 0.1 mg/mL	Not evaluated	Increased INF-γ and IL-2 in females, increased TNF-α in both sexes	Not evaluated	Blossom et al. (2008)
Unknown # litters/group, 6–10 offspring/sex/group; 0 (1% emulphor), 1,400, or 14,000 ppb; GD 0 to 12 mo of age	NOAEL: 1,400 ppb	No increase in autoantibody levels	Not evaluated	Not evaluated	Peden-Adams et al. (2008)

**Table 4-78. Summary of autoimmune-related studies of TCE and metabolites (by sex, strain, and route of exposure) (continued)**

Number/group, vehicle, dose, duration	NOAEL; LOAEL <sup>b</sup>	Results			Reference
		Serology	Ex vivo assays of cultured splenocytes	Clinical and histopathology	
<b>Autoimmune-prone: female MRL +/+ mice, intraperitoneal injection</b>					
4–5 per group, 0 (corn oil), 10 mmol/kg TCE, or 0.2 mmol/kg dichloroacetyl chloride, every 4 <sup>th</sup> day for 6 wk	LOAEL: 10 mmol/kg TCE, 0.2 mmol/kg dichloroacetyl chloride	In both groups, increased ANA and anti-ssDNA antibodies. In dichloroacetyl chloride group, anticardiolipin antibodies. No difference in antihistone, -Sm, or -DNA antibodies	Not evaluated	Not evaluated	Khan et al. (1995)
6 per group, 0 (corn oil), 0.2 mmol/kg dichloroacetyl chloride, or 0.2 mmol/kg dichloroacetic anhydride, 2 times per wk for 6 wk	LOAEL: 0.2 mmol/kg TCE, 0.2 mmol/kg dichloroacetic anhydride	In both treated groups, increased ANA	In both treated groups, increased IL-1 $\alpha$ , IL-1 $\beta$ , IL-3, IL-6, IFN- $\gamma$ , G-CSF and KC secretion; decreased IL-5. In dichloroacetyl chloride group, increased IL-17 and INF- $\alpha^d$	In both treated groups, increased lymphocytes in spleen, thickening of alveolar septa with lymphocytic interstitial infiltration	Cai et al. (2006)
<b>Autoimmune-prone: female NZB <math>\times</math> NZW mice, drinking water</b>					
6 per group, 0, 1,400, or 14,000 ppb TCE <sup>e,f</sup> , 27 wk exposure	LOAEL: 1,400 ppb	Increased anti-dsDNA antibodies at 19 wk and at 32–32 wk in the 1,400 ppb group	Not evaluated	At 14,000 ppb, proteinuria increased beginning at 20 wk; renal pathology scores increased, no evidence of liver disease	Gilkeson et al. (2004)
10 per group, 0, 1,400, or 14,000 ppb TCE <sup>f</sup> , 27 wk exposure	LOAEL: 1,400 ppb	Increased anti-dsDNA antibodies at 19 wk and at 32–32 wk in the 1,400 ppb group	No effect on splenocyte NK activity	No effect on renal pathology score; liver disease not examined	Keil et al. (2009)

**Table 4-78. Summary of autoimmune-related studies of TCE and metabolites (by sex, strain, and route of exposure) (continued)**

Number/group, vehicle, dose, duration	NOAEL; LOAEL <sup>b</sup>	Results			Reference
		Serology	Ex vivo assays of cultured splenocytes	Clinical and histopathology	
<b>Autoimmune-prone: male MRL—<i>lpr/lpr</i> mice, inhalation</b>					
5 per group, 0, 500, 1,000, or 2,000 ppm TCE, 4 h/d, 6 d/wk, 8 wk	LOAEL: 500 ppm			At ≥500 ppm, dose-related liver inflammation, splenomegaly and hyperplasia of lymphatic follicles; at 1,000 ppm, immunoblastic cell formation in lymphatic follicles, no changes in thymus	Kaneko et al. (2000)
<b>Autoimmune-inducible: female brown Norway Rat, gavage</b>					
6–8 per group, 0, 100, 200, 400 mg/kg, 5 d/wk, 6 wk followed by 1 mg/kg HgCl <sub>2</sub> challenge	NOAEL 500 mg/kg	Not reported <sup>g</sup>	Not evaluated	Not evaluated	White et al. (2000)
<b>Nonautoimmune-prone: female B6C3F1 mice, drinking water</b>					
6 per group, 0, 1,400, or 14,000 ppb TCE, <sup>e,f</sup> 30 wk exposure	LOAEL: 1,400 ppb	Anti-dsDNA increased in 1,400 ppb group beginning at age 32 wk and in the 14,000 ppb group beginning at age 26 wk	No effect on splenocyte NK activity	No renal disease observed	Gilkeson et al. (2004)

**Table 4-78. Summary of autoimmune-related studies of TCE and metabolites (by sex, strain, and route of exposure) (continued)**

Number/group, vehicle, dose, duration	NOAEL; LOAEL <sup>b</sup>	Results			Reference
		Serology	Ex vivo assays of cultured splenocytes	Clinical and histopathology	
<b>10 per group, 0, 1,400, or 14,000 ppb TCE,<sup>f</sup> 30 wk exposure</b>	<b>LOAEL: 1,400 ppb</b>	<b>Anti-dsDNA increased beginning at 26 wk in the 14,000 ppb group and at 32 wk of age in the 1,400 ppb group; increases in anti-ssDNA antibodies seen in both groups at 32 wk. Anti-GA were not affected</b>	<b>No effect on splenocyte NK activity</b>	<b>Increased renal pathology scores in 1,400 ppb group; Significant decrease in thymus weight in both groups</b>	<b>Keil et al. (2009)</b>

**Bolded studies** carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup>Selected endpoints, based on those reported across the majority of studies. Lupus-prone mouse strains develop lupus-like condition spontaneously, with virtually complete penetrance. The autoimmune-inducible (Brown Norway) rat has been used as a model of mercuric chloride induced glomerulonephritis and experimental autoimmune myasthenia gravis.

<sup>b</sup>NOAEL and LOAEL are based upon reported study findings.

<sup>c</sup>No difference reported in anti-dsDNA, -ssDNA, -ribonucleosome, -SSA, -SSB, -Sm, -Jo-1, or -Scl-70 antibodies.

<sup>d</sup>No difference reported in secretion of other cytokines measured: IL-2, IL-4, IL-10, IL-12, TNF- $\alpha$ , granulocyte monocyte colony stimulating factor, macrophage inflammatory protein-1 $\alpha$ , and RANTES (CCL-5).

<sup>e</sup>Dose levels cited in the report (Gilkeson et al., 2004) were incorrect; corrections provided by personal communication from Margie Peden-Adams (Medical University of South Carolina) to Glinda Cooper (U.S. EPA) on 13 August 2008; dose levels in this table are correctly report.

<sup>f</sup>Dose in mg/kg-day not given.

<sup>g</sup>Anti-dsDNA tests were described in the methods section; no effect of TCE on serum IgE levels was seen, and it is not clear if the additional serological tests were conducted in the TCE portion of this study or if they were conducted but not reported because no effect was seen.

G-CSF = granulocyte colony stimulating factor, KC = keratinocyte-derived chemokine.

1 immunoglobulin were found at 4 weeks of TCE treatment (indicating an autoimmune response),  
2 but not at 32 weeks. Increased expression of the activation marker C44 on splenic CD4+ cells  
3 was observed at 32 weeks. In addition, at 4 and 32 weeks, splenic T-cells from treated mice  
4 secreted more IFN- $\gamma$  than control T-cells (significant at 0.5 and 2.5 mg/mL), consistent with a  
5 Th1 immune or inflammatory response. By 22 weeks of TCE treatment, a specific immune  
6 serum antibody response directed against dichloroacetylated proteins was activated in hepatic  
7 tissues, indicating the presence of protein adducts. There was a slight but statistically significant  
8 increase in serum alanine aminotransferase levels at 32 weeks at 0.5 mg/mL. Histopathological  
9 evaluation at 32 weeks revealed extensive hepatic lymphocytic cell infiltration at 0.5 and  
10 2.5 mg/mL; all treated groups contained significantly more hepatocyte reactive changes (i.e.,  
11 presence of multinucleated hepatocytes, variations in hepatocyte morphology, and hepatocytes in  
12 mitosis) than controls.

13 In a subsequent study which assessed occupationally relevant concentrations, TCE was  
14 administered to female MRL +/+ mice (8/group) in drinking water at treatment levels of 0.1, 0.5,  
15 or 2.5 mg/mL (21, 100, or 400 mg/kg-day) for 4 and 32 weeks (Griffin et al., 2000b). At 4  
16 weeks, significant increases in serum antinuclear antibody levels were observed at 0.1 and  
17 0.5 mg/kg-day; at 32 weeks, the effects were observed at all three treatment levels. A dose-  
18 related increase in the percentage of activated CD4+ T-cells in spleens and lymph nodes of  
19 treated mice was observed at 32 weeks, and the CD4+ T-cells were found to secrete Th1-type  
20 cytokines at 4 and 32 weeks.

21 A similar response was observed by Cai et al. following chronic (48 weeks) exposure of  
22 TCE to female MRL +/+ mice (5/group) in drinking water at 0 or 0.5 mg/mL (approximately  
23 60  $\mu$ g/g/day) (Cai et al., 2008). After 11 weeks of treatment, a statistically significant decrease  
24 in body weight gain was observed. After 24 weeks of exposure, serum ANA were consistently  
25 elevated in treated mice as compared to control, although statistical significance was not  
26 achieved. Apparent treatment-related effects on serum cytokines included decreased IL-6 after  
27 36 and 48 weeks, decreased TNF- $\alpha$  after 48 weeks, and increased granulocyte colony stimulating  
28 factor (G-CSF) after 36 weeks of treatment. After 36 weeks of treatment, ex vivo cultured  
29 splenocytes secreted higher levels of IFN- $\gamma$  than control splenocytes. Although there were no  
30 observed effects on serum aminotransferase liver enzymes at termination, statistically significant  
31 incidences of hepatocytic necrosis and leukocyte infiltration (including CD3+ T lymphocytes)  
32 into liver lobules were observed in treated mice after 48 weeks of exposure. Hepatocyte  
33 proliferation was also increased. TCE treatment for 48 weeks also induced necrosis and  
34 extensive infiltration of leukocytes in the pancreas, infiltration of leukocytes into the perivascular  
35 and peribronchial regions of the lungs, and thickening of the alveolar septa in the lungs. At 36

1 and 48 weeks of exposure, massive perivascular infiltration of leukocytes (including CD3+  
2 T lymphocytes) was observed in the kidneys, and immunoglobulin deposits were found in the  
3 glomeruli.

4 To examine the role of metabolic activation in the autoimmune response, Griffin et al.  
5 (2000c) treated MRL +/+ mice with 2.5 mg/mL (300 mg/kg-day) TCE in drinking water for  
6 4 weeks (Griffin et al., 2000c). Immune responses were examined in the presence or absence of  
7 subcutaneous doses of 200 mg/kg-day diallyl sulfide, a specific inhibitor of CYP2E1 which is  
8 known to be a primary CYP that is active in TCE metabolism. With diallyl sulfide cotreatment  
9 that resulted in a decreased level of CYP2E1 apoprotein in liver microsomes, the enhanced  
10 mitogen-induced proliferative capacity of T-cells was inhibited and the reduction in IL-4 levels  
11 secreted by CD4+ T-cells was reversed for TCE-treated MRL +/+ mice. This study suggests that  
12 metabolism of TCE by CYP2E1 is responsible, at least in part, for the treatment-related CD4+ T-  
13 cell alterations.

14 The TCE metabolite, trichloroacetaldehyde (TCAA) or trichloroacetaldehyde hydrate  
15 (TCAH), was also evaluated in MRL +/+ mice (Blossom et al., 2007; Blossom and Gilbert,  
16 2006; Gilbert et al., 2004) in order to determine if outcomes similar to the immunoregulatory  
17 effects of TCE would be observed, and to attempt to further characterize the role of metabolism  
18 in the mode of action for TCE. At concentrations ranging from 0.04–1 mM, TCAA stimulated  
19 proliferation of murine Th1 cells treated with anti-CD3 antibody or antigen in vitro. At similar  
20 concentrations, TCAA induced phenotypic alterations consistent with upregulation of CD28 and  
21 downregulation of CD62L in cloned memory Th1 cells and DC4+ T-cells from untreated MRL  
22 +/+ mice. Phosphorylation of activating transcription factor 2 (ATF-2) and c-Jun (two  
23 components of the activator protein-1 transcription factor) was also observed with  
24 TCAA-induced Th1 cell activation. Higher concentrations of TCAA formed a Schiff base on  
25 T-cells, which suppressed the ability of TCAA to phosphorylate ATF-2. These findings  
26 suggested that TCAA may promote T-cell activation by stimulating the mitogen-activated  
27 protein kinase pathway in association with Schiff base formation on T-cell surface proteins  
28 (Gilbert et al., 2004).

29 In order to determine whether metabolites of TCE could mediate the immunoregulatory  
30 effects previously observed with TCE treatment (i.e., the generation of lupus and autoimmune  
31 hepatitis, associated with activation of IFN- $\gamma$ -producing CD4+ T-cells), Blossom et al. (2004)  
32 administered TCE metabolites, TCAH and trichloroacetic acid (TCA), to MRL +/+ mice  
33 (6–8/group) in drinking water for 4 weeks. Drinking water concentrations were 0, 0.1, or  
34 0.9 mg/mL; average daily doses were calculated as 0, 24, or 220 mg/kg-day for TCAH and 0, 27,  
35 or 205 mg/kg-day for TCA. These treatment levels were considered to be physiologically



1 relevant and to reflect occupational exposure. A phenotypic analysis of splenic and lymph node  
2 cells, cytokine profile analysis, evaluation of apoptosis in CD4<sup>+</sup> T-cells, and examination of  
3 serum markers of autoimmunity (anti-ssDNA, antihistone, or ANA) were conducted. Exposure  
4 to TCAH or TCA at both treatment levels was found to promote CD4<sup>+</sup> T-cell activation, as  
5 shown by significant ( $p < 0.05$ ) increases in the percentage of CD62L<sup>lo</sup> CD4<sup>+</sup> T-cells in the  
6 spleens and lymph nodes of the MRL +/+ mice. Increased levels of IFN- $\gamma$  were secreted by  
7 CD4<sup>+</sup> T-cells from mice treated by TCAH and TCA. No significant changes in body weight  
8 were observed; spleen weights were similar between control and treated mice with the exception  
9 of a significant decrease in spleen weight from mice treated with 0.9 mg/mL TCA. Liver and  
10 kidney histology were not affected, and serum alanine aminotransferase levels were similar for  
11 control and treated mice. A generalized trend towards an increase in serum autoantibodies  
12 (anti-ssDNA) was observed in TCAH-treated mice, and slight but significant increases in  
13 antihistone and antinuclear antibody production were observed in mice treated with  
14 0.9 mg/mL-day TCAH.

15 The autoimmune response of female MRL +/+ mice to DCAC, a metabolite of TCE, and  
16 to dichloroacetic anhydride (DCAA) a similar acylating agent, was evaluated by Cai et al.  
17 (2006). Six mice/group were injected intraperitoneally, twice weekly for 6 weeks, with  
18 0.2 mmol/kg DCAC or DCAA in corn oil. Body weight gain was significantly decreased after 5  
19 or 6 weeks treatment with DCAC and DCAA. DCAC treatment resulted in significant increases  
20 in total serum IgG (77% increase over control) and IgG1 (172% increase over control), as well as  
21 the induction of DCAC-specific IgG and IgG1. Serum IgM levels were significantly decreased  
22 by 25 and 18% in DCAC and DCAA-treated mice, respectively. IgE levels were increased  
23 100% over controls in DCAC-treated mice. Of eight Th1/Th2 cytokines measured, only IL-5  
24 was decreased in DCAC- and DCAA-treated mice. Serum ANA were detected in both DCAC-  
25 and DCAA-treated mice. Treatment-related increases in cytokine and chemokine secretion in  
26 cultured splenocytes were observed for DCAC and DCAA (IL-1, G-CSF, keratinocyte-derived  
27 chemokine, IL-3, and IL-6). DCAC-treated splenocytes also secreted more IL-17 and IFN- $\alpha$   
28 than controls. Histopathological changes were observed in the spleens of DCAC and DCAA-  
29 treated mice (lymphocyte population increases in the red pulp). With both DCAC and DCAA  
30 treatment, the alveolar septa were thickened in the lungs, moderate levels of lymphocytic  
31 interstitial infiltrates were present in tissues, and alveolar capillaries were clogged with  
32 erythrocytes. These findings were attributed both to the predisposition of the MRL +/+ mice  
33 towards autoimmune disease, and to the treatment-related induction of autoimmune responses.

34 Fas-dependant activation-induced cell death leading to autoimmune disease has been  
35 shown to be related to impaired Fas or FasL ligand expression in humans and mice, and defects

1 in the Fas-signaling pathways have been described in autoimmune disease models. The study by  
2 Blossom and Gilbert examined the effects of TCAH on Fas-dependent autoimmune cell death  
3 (Blossom and Gilbert, 2006). In this study, TCAH (1) inhibited apoptosis of antigen-activated  
4 cells, (2) did not protect CD4+ T-cells from Fas-independent apoptosis, (3) did not inhibit  
5 autoimmune cell death induced by direct engagement of the Fas receptor, (4) inhibited the  
6 expression of FasL but not Fas on the surface of activated CD4+ T-cell, (5) increased release of  
7 FasL from CD4+ cells in a metalloprotein-dependent manner, and (6) increased metalloprotein  
8 MMP-7 expression.

9         Gilbert et al. (2006) studied the effect of treatment on apoptosis in CD4+ T-lymphocytes  
10 isolated from MRL +/+ female mice that had been exposed to TCE (0, 0.1, 0.5, or 2.5 mg/mL) in  
11 the drinking water for 4 or 32 weeks or to TCAH (0.1, 0.3, or 0.9 mg/mL) in drinking water for 4  
12 or 40 weeks. After only 4 weeks, decreased activation-induced apoptosis was associated with  
13 decreased FasL expression in the CD4+ T-cells, suggesting that TCE- and TCAH-induced  
14 autoimmune disease was promoted through suppression of the process that would otherwise  
15 delete activated self-reactive T-lymphocytes. By 32 weeks of treatment, TCE had induced  
16 autoimmune hepatitis, which was associated with the promotion of oxidative stress, the  
17 formation of liver protein adducts, and the stimulated production of antibodies to those adducts.  
18 TCAH-treated mice did not exhibit autoimmune hepatitis by 40 weeks, but developed a dose-  
19 dependent alopecia and skin inflammation (Blossom et al., 2007). TCAH appeared to modulate  
20 the CD4+ T-cell subset by promoting the expression of an activated/effector phenotype with an  
21 increased capacity to secrete the proinflammatory cytokine IFN- $\gamma$ . A 4-week exposure to TCAH  
22 attenuated activation-induced cell death and the expression of the death receptor Fas in CD4+  
23 cells; these effects were not seen after a 40-week exposure period. Differences in response were  
24 tentatively attributed to higher levels of metalloproteinases (specifically MMP-7) at 4-weeks of  
25 treatment, suggesting a possible mechanism for the promotion of skin pathology by TCAH.

26         The role of protein adduct formation in autoimmune response has been pursued by  
27 various researchers. Halmes et al. administered a single i.p. dose of TCE in corn oil to male  
28 Sprague-Dawley rats (2/group) at 0 or 1,000 mg/kg (Halmes et al., 1997). Using antiserum that  
29 recognizes TCE covalently bound to protein, a single 50 kDa microsomal adduct was detected by  
30 Western blot in livers of treated rats. Using affinity chromatography, a 50 kDa dichloroacetyl  
31 protein was also isolated from rat plasma. The protein was reactive immunochemically with  
32 anti-CYP2E1 antibodies. The data suggest that the protein adduct may be CYP2E1 that has been  
33 released from TCE-damaged hepatocytes.

34         Cai et al. examined the role of protein haptenization in the induction of immune  
35 responses (Cai et al., 2007). In this study, MRL +/+ mice were immunized with albumin adducts

1 of various TCE reactive intermediates of oxidative metabolism. Serum immunoglobulins and  
2 cytokine levels were measured to evaluate immune responses against the haptenized albumin.  
3 Antigen-specific IgG responses (subtypes: IgG1, IgG2a, and IgG2b) were found. Serum levels  
4 of G-CSF were increased in immunized mice, suggesting macrophage activation. Following  
5 immunization with formyl-albumin, lymphocyte infiltration in the hepatic lobule and portal area  
6 was increased. This study suggests that proteins that are haptenized by metabolites of TCE may  
7 act as antigens to induce humoral immune responses and T-cell-mediated hepatitis.

8 A possible role for oxidative stress in inflammatory autoimmune disease was proposed by  
9 Khan et al. (2001). A study was performed in which female MRL +/+ mice were treated with  
10 10 mmol/kg TCE or 0.2 mmol/kg DCAC via intraperitoneal injection every 4<sup>th</sup> day for 2, 4, 6, or  
11 8 weeks. Antimalondialdehyde serum antibodies, a marker of lipid peroxidation and oxidative  
12 stress, were measured and were found to increase by 4 weeks of treatment, marginally for TCE  
13 and significantly for DCAC. It was reported that antimalondialdehyde antibodies has also been  
14 found to be present in the serum of systemic lupus erythematosus-prone MRL-lpr/lpr mice.

15 In another study that addressed the association of oxidative and nitrosative stress, and the  
16 role of lipid peroxidation and protein nitration, in TCE-mediated autoimmune response,  
17 Wang et al. treated female MRL +/+ mice with 0.5 mg/mL TCE in drinking water for 48 weeks  
18 (Wang et al., 2007b). The formation of antibodies in the serum to lipid peroxidation-derived  
19 aldehyde protein adducts was evaluated. With TCE treatment, the serum levels of  
20 antimalondialdehyde and anti-4-hydroxynonenal protein adduct antibodies, inducible nitric oxide  
21 synthase, and nitrotyrosine were increased. These were associated with increases in antinuclear-,  
22 anti-ssDNA- and anti-dsDNA antibodies. The involvement of lipid peroxidation-derived  
23 aldehyde protein adducts in TCE autoimmunity was further explored, using female MRL +/+  
24 mice that were administered by i.p. injections of TCE at 10 mmol/kg, either every 4<sup>th</sup> day for 6 or  
25 12 weeks (Wang et al., 2007a) or once per week for 4 weeks (Wang et al., 2008). Significant  
26 increases in malondialdehyde and 4-hydroxynonenal protein adducts, as well as significant  
27 induction of specific antibodies directed against these antigens were observed in both studies.  
28 Wang et al. also demonstrated a significant proliferation of CD4<sup>+</sup> T-cells in TCE-treated mice,  
29 and splenic lymphocytes from TCE-treated mice released more IL-2 and IFN- $\gamma$  when stimulated  
30 with MDA- or 4-hydroxynonenal-adducted mouse serum albumin (Wang et al., 2008). Overall,  
31 the result of these studies suggest a role for lipid peroxidation aldehydes in the induction and/or  
32 exacerbation of autoimmune response in the MRL +/+ animal model, and the involvement of  
33 Th1 cell activation.

34 In studies conducted in other rodent strains, less consistent outcomes have been observed.  
35 Inhalation exposure of an autoimmune-prone strain of male mice (MRL-lpr/lpr) to 0-, 500-,

1 1,000-, or 2,000-ppm TCE for 4 hours/day, 6 days/week, for 8 weeks resulted in depressed serum  
2 IgG levels and increased numbers of lymphoblastoid cells (Kaneko et al., 2000). Also at  
3 2,000 ppm, changes in T-cell helper to suppressor cell ratios were observed. At  
4 histopathological evaluation, dose-dependent inflammation and associated changes were noted in  
5 the liver at  $\geq 500$  ppm, hyperplasia of the lymphatic follicles of the spleen and splenomegaly  
6 were observed at  $\geq 500$  ppm, and the spleen exhibited the development of an immunoblastic-cell-  
7 like structure at 1,000 ppm.

8 A 26-week drinking water study of TCE in NZB  $\times$  NZW (NZBWF1) autoimmune-prone  
9 mice demonstrated an increase in anti-dsDNA antibodies at 19 weeks and at 32 and 34 weeks in  
10 the 1,400 ppb group, and increased kidney disease at 14,000 ppb (i.e., increased proteinuria at  
11 20 weeks; increased renal pathology scores at termination, based upon glomerular proliferation,  
12 inflammation, and necrosis) (Gilkeson et al., 2004).<sup>11</sup> Also in that study, a small increase in  
13 anti-dsDNA antibody production, without kidney disease, was observed in B6C3F1 mice, with  
14 statistically significant ( $p < 0.05$ ) or borderline ( $p = 0.07$ ) effects seen in the 1,400-ppb group at  
15 observations between 32 and 39 weeks of age, and in the 14,000 ppb group at observations  
16 between 26 and 39 weeks of age.

17 Keil et al. (2009) also assessed the effects of TCE exposure on NZBWF1 mice,  
18 comparing the responses to those of TCE-exposed B6C3F1 mice, which are not autoimmune  
19 prone (Keil et al., 2009). In this study, groups of NZBWF1 and B6C3F1 female mice (10/dose  
20 level) were administered 0, 1,400, or 14,000 ppb TCE in the drinking water. Treatment was  
21 initiated at 9 weeks of age and continued until 36 weeks of age for the NZBWF1 and until  
22 39 weeks of age for the B6C3F1 mice. Body weight; spleen, thymus, liver, and kidney weight;  
23 spleen and thymus cellularity; and renal pathology were assessed. Splenic lymphocyte  
24 proliferation, autoantibody production (anti-dsDNA, anti-ssDNA, and antiglomerular), total  
25 serum IgG, NK cell activity, and mitogen-induced lymphocyte proliferation were conducted.  
26 Administration of TCE did not result in alterations in NK cell activity or T- or B-cell  
27 proliferation in either strain of mice. In the NZBWF1 mice, there was little evidence of an  
28 increase or of an acceleration in ssDNA antibody production with TCE exposure, but as was seen  
29 in the earlier study by these investigators (Gilkeson et al., 2004), dsDNA antibodies were  
30 increased at 19 weeks and at 32–34 weeks in the 1,400 ppb group. However, antiglomerular  
31 antibody levels were increased in NZBWF1 mice early in the study, returning to control levels  
32 by 23 weeks of age. In the B6C3F1 mice the number of activated T-cells (CD4<sup>++</sup>/CD44<sup>+</sup>) was

---

<sup>11</sup>The study was reported in symposium proceedings. Dose levels cited in the proceedings were incorrect; however, corrections were provided by personal communication from Margie Peden-Adams (Medical University of South Carolina) to Glinda Cooper (U.S. EPA) on 13 August 2008, and dose levels are correctly reported here.

1 increased (significantly at 14,000 ppm;  $p \leq 0.05$ ) and thymus weights were significantly  
2 decreased ( $p \leq 0.05$ ) in a dose-responsive manner. Renal pathology (as indicated by renal score  
3 based on assessment of glomerular inflammation, proliferation, crescent formation and necrosis)  
4 was significantly increased ( $p \leq 0.05$ ) at 1,400 ppm. Also in the B6C3F1 mice, autoantibodies to  
5 dsDNA were increased relative to controls beginning at 26 weeks in the 14,000-ppb group and at  
6 32 weeks of age in the 1,400 ppb group; increases in anti-ssDNA antibodies were seen in both  
7 groups at 32 weeks. Antiglomerular antibodies were not affected in B6C3F1 mice. In summary,  
8 the authors concluded that this study showed that 27–30 weeks of TCE drinking water  
9 administration to NZBWF1 (autoimmune-prone) mice did not contribute to the progression of  
10 autoimmune disease, while similar administration to B6C3F1 (nonautoimmune-prone) mice  
11 increased the expression of a number of markers that are associated with autoimmune disease.  
12 This study is important in that it demonstrates that autoimmune responses to TCE exposure in  
13 animal models are not solely dependent upon a genetic predisposition to autoimmune disease.

14 White et al. conducted a study in female Brown Norway rats, which have been shown to  
15 be susceptible to development of chemically-induced IgE mediated glomerulonephritis that is  
16 similar to the nephritic damage seen in systemic lupus erythematosus (White et al., 2000). TCE  
17 administered by gavage 5 days/week at 100, 200, or 400 mg/kg did not increase in IgE levels  
18 after 6 weeks exposure, or after an additional challenge with 1 mg/kg HgCl<sub>2</sub>.

19 Several studies have examined the potential for autoimmune response following oral  
20 exposures during pre- and postnatal immune system development, as described in  
21 Section 4.6.2.1.2 above. Peden-Adams et al. conducted two such studies. In the first study,  
22 B6C3F1 mice were treated with either 1,400 or 14,000 ppb TCE in drinking water from GD 0 to  
23 postnatal Week 8 (Peden-Adams et al., 2006). No treatment-related increases in serum anti-  
24 dsDNA antibody levels were observed in the 8-week old offspring, although it is noted that the  
25 mouse strain used in the experiment is not an autoimmune-prone animal model. A more recent  
26 study (Peden-Adams et al., 2008) exposed pregnant MRL +/+ mice to TCE in drinking water at  
27 levels of 0, 1,400, or 14,000 ppb from GD 0 and continued the exposures until the offspring were  
28 12 months of age. Consistent with the findings of the 2006 publication, autoantibody levels  
29 (anti-dsDNA and antiglomerular) were not increased in the offspring over the course of the  
30 study. Contrasting with these negative studies, the lupus-prone MRL +/+ mouse model was  
31 utilized in two additional drinking water studies with developmental exposures in which there  
32 was some indication of a positive association between developmental exposures to TCE and the  
33 initiation of autoimmune disease. Blossom and Doss (2007) administered TCE to pregnant MRL  
34 +/+ mice in drinking water at levels of 0, 0.5, or 2.5 mg/mL and continued administration to the  
35 offspring until approximately 7–8 weeks of age. TCE exposure induced a dose-dependent

1 increase in T-lymphocyte IFN- $\gamma$  in peripheral blood at 4–5 weeks of age, but this effect was not  
2 observed in splenic T-lymphocytes at 7–8 weeks of age. Serum antihistone autoantibodies and  
3 total IgG<sub>2a</sub> were significantly increased in the TCE-treated offspring; however, histopathological  
4 evaluation of the liver and kidneys did not reveal any treatment-related signs of autoimmunity.  
5 In a study by Blossom et al. (2008), pregnant MRL +/+ mice were administered TCE in the  
6 drinking water at levels of 0 or 0.1 mg/mL from GD 0 through lactation, and continuing  
7 postweaning in the offspring until GD 42. Significant treatment-related increases in pro-  
8 inflammatory cytokines (IFN- $\gamma$  and IL-2 in males and TNF- $\alpha$  in both sexes) produced by splenic  
9 CD4+ T-cells were observed in PND 42 offspring.

10 In summary, TCE treatment induces and exacerbates autoimmune disease in genetically  
11 susceptible strains of mice, and has also been shown to induce signs of autoimmune disease in a  
12 nongenetically predisposed strain. Although the mechanism for this response is not fully  
13 understood, a number of studies have been conducted to examine this issue. The primary  
14 conclusion to date is that metabolism of the TCE to its chloral or dichloroacetic acid metabolites  
15 is at least partially responsible for activating T-cells or altering T-cell regulation and survival  
16 associated with polyclonal disease in susceptible mice strains.

#### 4.6.2.1.7. Cancers of the Immune System

17 Cancers of the immune system that have been observed in animal studies and are  
18 associated with TCE exposure are summarized in Tables 4-79 and 4-80. The specific tumor  
19 types observed are malignant lymphomas, lymphosarcomas, and reticulum cell sarcomas in mice  
20 and leukemias in rats.

21 In the NCI (1976) study, the results for Osborne-Mendel rats were considered  
22 inconclusive due to significant early mortality, but exposure to B6C3F1 mice were also  
23 analyzed. Limited increases in lymphomas over controls were observed in both sexes of mice  
24 exposed (see Table 4-79). The NCI study (1976) used technical grade TCE which contained  
25 two known carcinogenic compounds as stabilizers (epichlorohydrin and 1,2-epoxybutane). A  
26 later study (Henschler et al., 1984) in which mice were given TCE that was pure, industrial, and  
27 stabilized with one or both of these stabilizers did not find significant increases in lymphomas  
28 over historical controls. A gavage study by NTP (1988), which used TCE stabilized with  
29 diisopropylamine, did not see an increase in lymphomas in all four strains of rats (ACI, August,  
30 Marshall, and Osborne-Mendel). The final NTP study (1990) in male and female F344 rats and  
31 B6C3F1 mice, using epichlorohydrin-free TCE, again experienced early mortality in male rats.  
32 This study did not observe significant increase in lymphomas over that of controls. Henschler et

1 al. (1980) tested NMRI mice, WIST rats and Syrian hamsters of both sexes, and observed a  
 2 variety of tumors in both sexes (Henschler et al., 1980), consistent with the spontaneous tumor

3 **Table 4-79. Malignant lymphomas incidence in mice exposed to TCE in**  
 4 **gavage and inhalation exposure studies**  
 5

Cancer type, species, and sex	Prevalence in exposure groups: <i>n</i> affected/ <i>n</i> total (% affected)						Reference
	Vehicle control		Low dose		High dose		
<b>Gavage exposure</b>							
Malignant lymphomas	Vehicle control		1,000 mg/kg-day				NTP (1990)
B6C3F1 mice, male	11/50 (22%)		13/50 (26%)				
B6C3F1 mice, female	7/48 (15%)		13/49 (27%)				
Lymphosarcomas and reticulum cell sarcomas	Vehicle control		Low dose		High dose		NCI (1976) <sup>b</sup>
B6C3F1 mice, male	1/20 (5%)		4/50 (8%)		2/48 (4%)		
B6C3F1 mice, female	1/20 (5%)		5/50 (10%)		5/47 (11%)		
Malignant lymphomas	Control	TCE-pure	TCE-indust	TCE-EPC	TCE-BO	TCE-EPC-BO	Henschler et al. (1984) <sup>c</sup>
Swiss (ICR/HA) mice, male	19/50 (38%)	16/50 (32%)	17/49 (35%)	11/49 (22%)	11/49 (22%)	12/49 (24%)	
Swiss (ICR/HA) mice, female	28/50 (56%)	21/50 (42%)	19/50 (38%)	20/50 (40%)	23/48 (48%)	18/50 (36%)	
<b>Inhalation exposure</b>							
Malignant lymphomas	Control		96		480		Henschler et al. (1980) <sup>d</sup>
Han:NMRI mice, male	7/30 (23%)		7/29 (24%)		6/30 (20%)		
Han:NMRI mice, female <sup>e</sup>	9/29 (31%)		17/30 (57%)		18/28 (64%)		

6  
 7 <sup>a</sup>After 103 wk gavage exposure, beginning at 8 wk of age.

8 <sup>b</sup>After 90 wk gavage exposure, beginning at 5 wk of age. Low dose is 1,200 mg/kg-day for male mice,  
 9 900 mg/kg-day for female mice (5 d/wk). High dose is 2,400 mg/kg-day for male mice, 1,800 mg/kg-day for  
 10 female mice (5 d/wk).

11 <sup>c</sup>After 72 wk gavage exposure (corn oil), beginning at 5 wk of age. Male mice received 2,400 mg/kg-day, female  
 12 mice received 1,800 mg/kg-day. Stabilizers were added in the percentage w/w: TCE-EPC, 0.8%, TCE-BO, 0.8%,  
 13 TCE-EPC-BO, 0.25 and 0.25%.

14 <sup>d</sup>After 78 wk inhalation exposure. Administered daily concentration: low dose is 96 (mg/m<sup>3</sup>) and high dose is 480  
 15 (mg/m<sup>3</sup>), equivalent to 100 and 500 ppm (100 ppm = 540 mg/m<sup>3</sup>), adjusted for 6 h/d, 5 d/wk exposure.

16 <sup>e</sup>Statistically significant by Cochran-Armitage trend test (*p* < 0.05).

17  
 18 Sources: NTP (1990) Tables 8, 9; NCI (1976) Table XXXa; Henschler et al. (1980) Table 3a.  
 19

1 **Table 4-80. Leukemia incidence in rats exposed to TCE in gavage and**  
 2 **inhalation exposure studies**  
 3

Species and sex	Prevalence in exposure groups: <i>n</i> affected/ <i>n</i> total (% affected)				Reference
	Control	50 mg/kg	250 mg/kg	600 ppm	
<b>Gavage exposure</b>					Maltoni et al. (1986) <sup>a</sup>
Sprague-Dawley rats, male	0/30 (0%)	2/30 (6.7%)	3/30 (10.0%)		
Sprague-Dawley rats, female	1/30 (3.3%)	0/30 (0%)	0/30 (0%)		
	Control	500 mg/kg	1,000 mg/kg		
August rats, female	0/50 (0%)	1/50 (2%)	5/50 (10%)		NTP (1988) <sup>b</sup>
<b>Inhalation exposure</b>	Control	100 ppm	300 ppm	600 ppm	Maltoni et al. (1988) <sup>c</sup>
Sprague-Dawley rats, male	9/135 (6.7)	13/130 (10.0)	14/130 (10.8)	15/130 (11.5)	
Sprague-Dawley rats, female	7/145 (4.8)	9/130 (6.9)	2/130 (1.5)	11/130 (8.5)	

4  
 5 <sup>a</sup>After 52 wk gavage exposure, beginning at 13 wk of age, olive oil vehicle. Percentage affected and starting *n* given  
 6 in reported; EPA calculated *n* affected.

7 <sup>b</sup>After 104 wk gavage exposure, beginning at 6.5–8 wk of age, corn oil vehicle.

8 <sup>c</sup>After 104 wk inhalation exposure, BT304 and BT304bis. Percentage affected and starting *n* given in reported; EPA  
 9 calculated *n* affected.

10  
 11  
 12 incidence in this strain (Deerberg and Müller-Peddinghaus, 1970; Deerberg et al., 1974).  
 13 Henschler et al. did not show an increase in lymphomas in rats or hamsters of either sex  
 14 (Henschler et al., 1980). Background levels of lymphomas in this mouse strain are high, making  
 15 it difficult to determine if the increased lymphomas in female mice is a treatment effect. In a  
 16 follow-up study, Henschler et al. (1984) examined the role of stabilizers of TCE in the  
 17 lymphomas demonstrated in female mice in the 1980 paper. Each exposure group had  
 18 ~50 SPF-bred ICR/HA-Swiss mice and exposure was for 18 months. Background incidence of  
 19 tumors was high in all groups. Focusing just on malignant lymphomas (see Table 4-79), the high  
 20 background incidence in unexposed animals again makes it difficult to determine if there is TCE  
 21 and/or stabilizer-related incidence of lymphomas. There are no data at any other timepoint than  
 22 18 months. A high mortality rate in all animals as well as the increased incidence of  
 23 ‘background’ lymphomas in that report was also a problem and may have been related to the  
 24 shorter time frame.

25 Maltoni et al. reported a nonsignificant increase in leukemias in male rats exposed via  
 26 inhalation (Maltoni et al., 1988, 1986). Maltoni et al. (1986) demonstrates a borderline higher



1 frequency of leukemias in male Sprague-Dawley rats following exposure by ingestion for  
2 52 weeks, believed by the authors to be related to an increase in lymphoblastic lymphosarcomas  
3 (see Table 4-80). The gavage study by NTP (1988), which used TCE stabilized with  
4 diisopropylamine, observed leukemia in female August rats with a positive trend, but was not  
5 significantly greater than the vehicle controls.

6 In summary, overall there is limited available data in animals on the role of TCE in  
7 lymphomas and leukemias. There are few studies that analyze for lymphomas and/or leukemias.  
8 Lymphomas were described in four studies (Henschler et al., 1984; Henschler et al., 1980; NCI,  
9 1976; NTP, 1990) but study limitations (high background rate) in most studies make it difficult  
10 to determine if these are TCE-induced. Three studies found positive trends in leukemia in  
11 specific strains and/or gender (Maltoni et al., 1988; Maltoni et al., 1986; NTP, 1988). Due to  
12 study limitations, these trends cannot be determined to be TCE-induced.

### 4.6.3. Summary

#### 4.6.3.1.1. Noncancer Effects

14 The human and animal studies of TCE and immune-related effects provide strong  
15 evidence for a role of TCE in autoimmune disease and in a specific type of generalized  
16 hypersensitivity syndrome. The data pertaining to immunosuppressive effects is weaker. It  
17 should also be noted that immune-related and inflammatory effects, particularly cell-mediated  
18 immunity involving cytokine production and activation of macrophages and natural killer cells,  
19 may influence a variety of other conditions of considerable public health importance, including  
20 cancer (tumor surveillance) and atherosclerosis. Thus the relevance of immune-related effects of  
21 TCE should are not only limited to diseases affecting organs and tissues within the immune  
22 system. The relation between systemic autoimmune diseases, such as scleroderma, and  
23 occupational exposure to TCE has been reported in several recent studies. A meta-analysis of  
24 scleroderma studies (Diot et al., 2002; Garabrant et al., 2003; Nietert et al., 1998) conducted by  
25 the EPA resulted in a statistically significant combined odds ratio for any exposure in men  
26 (OR: 2.5, 95% CI: 1.1, 5.4), with a lower relative risk seen in women in women (OR: 1.2,  
27 95% CI: 0.58, 2.6). The incidence of systemic sclerosis among men is very low (approximately  
28 1 per 100,000 per year), and is approximately 10 times lower than the rate seen in women  
29 (Cooper and Stroehla, 2003). Thus, the human data at this time do not allow us to determine if  
30 the difference in effect estimates between men and women reflects the relatively low background  
31 risk of scleroderma in men, gender-related differences in exposure prevalence or in the reliability

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 of exposure assessment (Messing et al., 2003), a gender-related difference in susceptibility to the  
2 effects of TCE, or chance. Changes in levels of inflammatory cytokines were reported in an  
3 occupational study of degreasers exposed to TCE (Iavicoli et al., 2005) and a study of infants  
4 exposed to TCE via indoor air (Lehmann et al., 2001; Lehmann et al., 2002). Experimental  
5 studies support the biological plausibility of these effects. Numerous studies have demonstrated  
6 accelerated autoimmune responses in autoimmune-prone mice (Blossom et al., 2007; Blossom et  
7 al., 2004; Cai et al., 2008; Griffin et al., 2000a; Griffin et al., 2000b). With shorter exposure  
8 periods, effects include changes in cytokine levels similar to those reported in human studies.  
9 More severe effects, including autoimmune hepatitis, inflammatory skin lesions, and alopecia,  
10 were manifest at longer exposure periods, and interestingly, these effects differ somewhat from  
11 the “normal” expression in these mice. Immunotoxic effects, including increases in anti-dsDNA  
12 antibodies in adult animals and decreased plaque forming cell response with prenatal and  
13 neonatal exposure, have been also reported in B6C3F1 mice, which do not have a known  
14 particular susceptibility to autoimmune disease (Gilkeson et al., 2004; Peden-Adams et al.,  
15 2006). Recent mechanistic studies have focused on the roles of various measures of oxidative  
16 stress in the induction of these effects by TCE (Wang et al., 2007b; Wang et al., 2008).

17       There have been a large number of case reports of a severe hypersensitivity skin disorder,  
18 distinct from contact dermatitis and often accompanied by hepatitis, associated with occupational  
19 exposure to TCE, with prevalences as high as 13% of workers in the same location (Kamijima et  
20 al., 2007; Kamijima et al., 2008). Evidence of a treatment-related increase in delayed  
21 hypersensitivity response accompanied by hepatic damage has been observed in guinea pigs  
22 following intradermal injection (Tang et al., 2008), 2002), and hypersensitivity response was also  
23 seen in mice exposed via drinking water pre- and postnatally (GD 0 through to 8 weeks of age)  
24 (Peden-Adams et al., 2006).

25       Human data pertaining to TCE-related immunosuppression resulting in an increased risk  
26 of infectious diseases is limited to the report of an association between reported history of  
27 bacteria of viral infections in Woburn, MA (Lagakos et al., 1986). Evidence of localized  
28 immunosuppression, as measured by pulmonary response to bacterial challenge (i.e., risk of  
29 Streptococcal pneumonia-related mortality and clearance of *Klebsiella* bacteria) was seen in an  
30 acute exposure study in CD-1 mice (Aranyi et al., 1986). A 4-week inhalation exposure in  
31 Sprague-Dawley rats reported a decrease in plaque forming cell response at exposures of  
32 1,000 ppm (Woolhiser et al., 2006).

33

#### 4.6.3.1.2. Cancer

1 Associations observed in epidemiologic studies of lymphoma and TCE exposure suggest  
2 a causal relation between trichloroethylene exposure and NHL. Issues of study heterogeneity,  
3 potential publication bias, and weaker exposure-response results contribute uncertainty to the  
4 evaluation of the available data.

5 In a review of the NHL studies, studies in which there is a high likelihood of TCE  
6 exposure in individual study subjects (e.g., based on job-exposure matrices, biomarker  
7 monitoring, or industrial hygiene data on TCE exposure patterns and factors that affect such  
8 exposure) and which met, to a sufficient degree, the standards of epidemiologic design and  
9 analysis were identified. These studies generally reported excess relative risk estimates for NHL  
10 between 0.8 and 3.1 for overall TCE exposure. Statistically significant elevated relative risk  
11 estimates with NHL and overall TCE exposure were observed in two cohort (Hansen et al., 2001;  
12 Raaschou-Nielsen et al., 2003) and one case-control (Hardell et al., 1994) study. Both cohort  
13 studies reported statistically significant associations with NHL for subjects with longer  
14 employment duration as a surrogate of TCE exposure as does a second case-control study with  
15 high-quality exposure-assessment methodology reported statistically significant associations  
16 with highest cumulative TCE exposure or highest average-weekly TCE exposure (Purdue et al.,  
17 2011). Hardell et al. (1994) reported a strong but imprecise association, in part reflecting  
18 possible bias from subject-reported exposure history and few exposed cases. Other identified  
19 studies reported a 10–50% elevated relative risk estimate with overall TCE exposure that were  
20 not statistically significant, except for two population case-control studies of NHL, one of which  
21 did not report relative risk estimates with overall TCE exposure but did for medium-high  
22 intensity or cumulative TCE exposure (Anttila et al., 1995; Axelson et al., 1994; Boice et al.,  
23 1999; Cocco et al., 2010; Greenland et al., 1994; Miligi et al., 2006; Morgan et al., 1998;  
24 Nordström et al., 1998; Persson and Fredrikson, 1999; Purdue et al., 2011; Radican et al., 2008;  
25 Siemiatycki, 1991; Wang et al., 2009; Zhao et al., 2005). Fifteen additional studies were given  
26 less weight because of their lesser likelihood of TCE exposure and other design limitations that  
27 would decrease study power and sensitivity (ATSDR, 2004a, 2006a; Blair et al., 1989; Chang et  
28 al., 2003b; Chang et al., 2005; Clapp and Hoffman, 2008; Cohn et al., 1994b; Costa et al., 1989;  
29 Garabrant et al., 1988; Henschler et al., 1995; Morgan and Cassady, 2002; Ritz, 1999a;  
30 Vartiainen et al., 1993; Wilcosky et al., 1984) Sinks et al., 1992. The observed lack of  
31 association with NHL in these studies likely reflects study design and exposure assessment  
32 limitations and is not considered inconsistent with the overall evidence on TCE and NHL.

33 Consistency of the association between TCE exposure and NHL is further supported by  
34 the results of meta-analyses of 17 studies reporting risk estimates for overall TCE exposure that

1 met the meta-analysis inclusion criteria. These meta-analyses found a statistically significant  
2 increased summary relative risk estimate for NHL of 1.23 (95% CI: 1.07, 1.42) for overall TCE  
3 exposure. The analysis of NHL was robust to the removal of individual studies and the use of  
4 alternate relative risk estimates from individual studies, and in only one cases was the resulting  
5 summary relative risk no longer statistically significant (lower 95% confidence bounds of 1.00).  
6 Some evidence heterogeneity was observed, particularly between cohort and case-control  
7 studies, but it was not statistically significant ( $p = 0.10$ ); and, in addition, there was some  
8 evidence of potential publication bias. Analyzing the cohort and case-control studies separately  
9 resolved most of the heterogeneity, but the result for the summary case-control studies was only  
10 a 7% increased relative risk estimate and was not statistically significant. The sources of  
11 heterogeneity are uncertain but may be the result of some bias associated with exposure  
12 assessment and/or disease classification, or from differences between cohort and case-control  
13 studies in average TCE exposure.

14 Exposure-response relationships are examined in the TCE epidemiologic studies only to a  
15 limited extent. Many studies examined only overall “exposed” versus “unexposed” groups and  
16 did not provide exposure information by level of exposure. Others do not have adequate  
17 exposure assessments to confidently distinguish between levels of exposure. The NHL case-  
18 control study of Purdue et al. (2011) reported a statistically significant trend with TCE exposure  
19 ( $p = 0.02$  for average-weekly TCE exposure), and NHL risk in Boice et al. (1999) appeared to  
20 increase with increasing exposure duration ( $p = 0.20$  for routine-intermittent exposed subjects).  
21 The borderline statistically significant trend with TCE intensity in the case-control study of  
22 Wang et al. (2009 [ $p = 0.06$ ]) and with cumulative TCE exposure in the case-control study of  
23 Purdue et al. (2011 [ $p = 0.08$ ]) is consistent with that observed with average weekly TCE  
24 exposure in Purdue et al. (2011). Further support was provided by meta-analyses using only the  
25 highest exposure groups, which yielded a higher summary relative risk estimate (1.57 [95% CI:  
26 1.27, 1.94]) than for overall TCE exposure (1.27 [95% CI: 1.04, 1.44]).

27 Few risk factors are recognized for NHL, with the exception of viruses,  
28 immunosuppression or smoking, which are associated with specific NHL subtypes (Besson et al.,  
29 2006). Associations between NHL and TCE exposure are based on groupings of several  
30 subtypes. Two of the seven NHL case-control studies adjusted for age, sex and smoking in  
31 statistical analyses (Miligi et al., 2006; Wang et al., 2009), two others adjusted for age and sex  
32 (Cocco et al., 2010; Purdue et al., 2011), and the other three case-control studies presented only  
33 unadjusted estimates of the odds ratio (Hardell et al., 1994; Nordström et al., 1998; Persson and  
34 Fredrikson, 1999).

1           Animal studies describing rates of lymphomas and/or leukemias in relation to TCE  
2 exposure (Henschler et al., 1984; Henschler et al., 1980; Maltoni et al., 1988; Maltoni et al.,  
3 1986; NCI, 1976; NTP, 1988, 1990) are available. Henschler et al. (1980) reported statistically  
4 significant increases in lymphomas in female Han:NMRI mice treated via inhalation. While  
5 Henschler et al. (1980) suggested these lymphomas were of viral origin specific to this strain,  
6 subsequent studies reported increased lymphomas in female B6C3F1 mice treated via corn oil  
7 gavage (NTP, 1990) and leukemias in male Sprague-Dawley and female August rats (Maltoni et  
8 al., 1986; NTP, 1988). However, these tumors had relatively modest increases in incidence with  
9 treatment, and were not reported to be increased in other studies.  
10

## **4.7. RESPIRATORY TRACT TOXICITY AND CANCER**

### **4.7.1. Epidemiologic Evidence**

#### **4.7.1.1.1. Chronic Effects: Inhalation**

11           Two reports of a study of 1,091 gun-manufacturing workers are found on noncancer  
12 pulmonary toxicity (Cakmak et al., 2004; Saygun et al., 2007). A subset of these workers  
13 ( $n = 411$ ) had potential exposure to multiple organic solvents including toluene, acetone, butanol,  
14 xylene, benzene and TCE used to clean gun parts; however, both papers lacked information on  
15 exposure concentration. Mean exposure duration in Cakmak et al. (2004) was 17 years  
16 (SD = 7.9) for nonsmokers and 16 years (SD = 7.1) for smokers. Cakmak et al. (2004) indicated  
17 effects of smoking and exposure to solvents, with smoking having the most important effect on  
18 asthma-related symptoms (smoking, OR = 2.8, 95% CI: 2.0, 3.8; solvent exposure, OR = 1.4,  
19 95% CI: 1.1, 1.9). Similarly, smoking, but not solvent exposure, was shown as a statistically  
20 significantly predictor of lung function decrements. Saygun et al. (2007) reported on a 5-year  
21 follow-up of 393 of the original 1,091 subjects, 214 of who were exposed to solvents. Of the  
22 393 original subjects, the prevalence of definitive asthma symptoms, a more rigorous definition  
23 than used by Cakmak et al. (2004), was 3.3% among exposed and 1.1% among nonexposed  
24 subjects,  $p > 0.05$ . Saygun et al. (2007) presents observations on lung function tests for 697  
25 current workers, a group which includes the 393 original study subjects. Smoking, but not  
26 solvent exposure, was a predictor of mean annual forced expiratory volume (FEV<sub>1</sub>) decrease.  
27

#### 4.7.1.1.2. Cancer

1 Cancers of the respiratory tract including the lung, bronchus, and trachea are examined in  
2 25 cohort, community studies and case-control studies of TCE. Twelve of the 25 studies  
3 approached standards of epidemiologic design and analysis identified in the review of the  
4 epidemiologic body of literature on TCE and cancer (Anttila et al., 1995; see Appendix B;  
5 Axelson et al., 1994; Blair et al., 1998; Boice et al., 1999; Boice et al., 2006b; Greenland et al.,  
6 1994; Hansen et al., 2001; Morgan et al., 1998; Raaschou-Nielsen et al., 2003; Radican et al.,  
7 2008; Siemiatycki, 1991; Zhao et al., 2005). Cancers at other sites besides lung, bronchus, and  
8 trachea in the respiratory system are more limitedly reported in these studies. Some information  
9 is available on laryngeal cancer; however, only 9 of the 16 occupational cohort studies providing  
10 information on lung cancer also reported findings for this site. Case-control studies of lung or  
11 laryngeal cancers and occupational title or organic solvent exposure were found in the literature.  
12 Two case-control studies of lung cancer, one population-based and the other nested within a  
13 cohort, were of TCE exposure specifically. Lung and laryngeal cancer risk ratios reported in  
14 cohort, community and case-control studies are found in Table 4-81.

15 Lung cancer relative risks were reported in 11 of 12 cohort studies of aircraft  
16 manufacturing, aircraft maintenance, aerospace, and metal workers, with potential exposure to  
17 TCE as a degreasing agent, and in occupational cohort studies employing biological markers of  
18 TCE exposures. All 11 studies had a high likelihood of TCE exposure in individual study  
19 subjects and were judged to have met, to a sufficient degree, the standards of epidemiologic  
20 design and analysis (Anttila et al., 1995; Axelson et al., 1994; Blair et al., 1998; Boice et al.,  
21 1999; Boice et al., 2006b; Greenland et al., 1994; Hansen et al., 2001; Morgan et al., 1998;  
22 Raaschou-Nielsen et al., 2003; Radican et al., 2008; Zhao et al., 2005). Lung cancer risks were  
23 not reported for Fernald uranium processing workers with potential TCE exposure (Ritz, 1999a),  
24 a study of less weight than the other 11 studies.. The incidence study of Raaschou-Nielsen et al.  
25 (2003) was the largest cohort, with 40,049 subjects identified as potentially exposed to TCE in  
26 several industries (primarily, in the iron/metal and electronic industries), including 14,360 of  
27 whom had presumably higher level exposures to TCE. The study included 632 lung cancer cases  
28 and reported a 40% elevated incidence in TCE exposed males and females combined (95% CI:  
29 1.32, 1.55), with no exposure duration gradient. The 95% CIs in other studies of lung cancer  
30 incidence included a risk ratio of 1.0 (Anttila et al., 1995; Axelson et al., 1994; Blair et al., 1998;  
31 Hansen et al., 2001; Zhao et al., 2005). Lung cancer mortality risks in studies of TCE exposure  
32 to aircraft manufacturing, aircraft maintenance, and aerospace workers included a relative risk of  
33 1.0 in their 95% CIs (Blair et al., 1998; Boice et al., 2006b; Morgan et al., 1998; Radican et al.,  
34 2008; Zhao et al., 2005). Boice et al. (1999) observed a 24% decrement (95% CI: 0.60, 0.95) for

1 subjects with routine TCE exposure. Exposure-response analyses using internal controls  
2 (unexposed subjects at the same company) showed a statistically significant decreasing trend  
3 between lung cancer risk and routine or intermittent TCE exposure duration. The routine or  
4 intermittent category is broader and includes more subjects with potential TCE exposure. Five  
5 other studies with internal controls do not provide evidence of either an increasing or decreasing  
6 pattern between TCE and lung cancer incidence or mortality (Blair et al., 1998; Boice et al.,  
7 2006b; Morgan et al., 1998; Radican et al., 2008; Zhao et al., 2005).

8

1  
2

**Table 4-81. Selected results from epidemiologic studies of TCE exposure and lung cancer**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort studies—incidence</b>				
Aerospace workers (Rocketdyne)				Zhao et al. (2005)
	Any exposure to TCE	Not reported		
	Low cumulative TCE score	1.00 <sup>a</sup>	43	
	Medium cumulative TCE score	1.36 (0.86, 2.14)	35	
	High TCE score	1.11 (0.60, 2.06)	14	
	<i>p</i> for trend	0.60		
All employees at electronics factory (Taiwan)				Chang et al. (2005)
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure, all subjects	1.4 (1.32, 1.55)	632	
	Any exposure, males	1.4 (1.28, 1.51)	559	
	Any exposure, females	1.9 (1.48, 2.35)	73	
	Employment duration			
	<1 yr	1.7 (1.46, 1.93)	209	
	1–4.9 yr	1.3 (1.16, 1.52)	218	
	≥5 yr	1.4 (1.23, 1.63)	205	
Biologically-monitored Danish workers				Hansen et al. (2001)
	Any TCE exposure, males	0.8 (0.5, 1.3)	16	
	Any TCE exposure, females	0.7 (0.01, 3.8)	1	
	Cumulative exposure (Ikeda)			
	<17 ppm-yr			
	≥17 ppm-yr			
	Mean concentration (Ikeda)			
	<4 ppm			
	4+ ppm			
	Employment duration			
	<6.25 yr			
	≥6.25 yr			
Aircraft maintenance workers (Hill Air Force Base, UT)				Blair et al. (1998)
	TCE subcohort	Not reported		
	Males, cumulative exposure			
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	1.0 (0.6, 2.0)	24	
	5–25 ppm-yr	0.8 (0.4, 1.6)	11	
	>25 ppm-yr	0.8 (0.4, 1.7)	15	

3

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-81. Selected results from epidemiologic studies of TCE exposure and lung cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Females, cumulative exposure				Blair et al. (1998) (continued)
0		1.0 <sup>a</sup>		
<5 ppm-yr			1	
5–25 ppm-yr			1	
>25 ppm-yr			1	
Biologically-monitored Finnish workers				Anttila et al. (1995)
All subjects		0.92 (0.59, 1.35)	25	
Mean air-TCE (Ikeda extrapolation)				
<6 ppm		1.02 (0.58, 1.66)	16	
6+ ppm		0.83 (0.33, 1.71)	7	
Biologically-monitored Swedish workers				Axelson et al. (1994)
Any TCE exposure, males		0.69 (0.31, 1.30)	9	
Any TCE exposure, females		Not reported		
<b>Cohort and PMR-mortality</b>				
Computer manufacturing workers (IBM), NY				Clapp and Hoffman (2008)
Males		1.03 (0.71, 1.42)	35	
Females		0.95 (0.20, 2.77)	3	
Aerospace workers (Rocketdyne)				
Any TCE (utility or engine flush workers)		1.24 (0.92, 1.63)	51	Boice et al. (2006b)
Engine flush—duration of exposure				
Referent		1.0 <sup>a</sup>	472	
0 yr (utility workers with TCE exposure)		0.5 (0.22, 1.00)	7	
<4 yr		0.8 (0.50, 1.26)	27	
≥4 yr		0.8 (0.46, 1.41)	24	
Any exposure to TCE		Not reported		
Low cumulative TCE score		1.00 <sup>a</sup>	99	Zhao et al. (2005)
Medium cumulative TCE score		1.05 (0.76, 1.44)	62	
High TCE score		1.02 (0.68, 1.53)	33	
<i>p</i> for trend		0.91		
View-Master employees				ATSDR (2004a)
Males		0.81 (0.42, 1.42) <sup>b</sup>	12	
Females		0.99 (0.71, 1.35) <sup>b</sup>	41	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-81. Selected results from epidemiologic studies of TCE exposure and lung cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
United States uranium-processing workers (Fernald)				Ritz (1999a)
	Any TCE exposure	Not reported		
	Light TCE exposure, >2 yr duration <sup>c</sup>	Not reported		
	Moderate TCE exposure, >2 yr duration <sup>c</sup>	Not reported		
Aerospace workers (Lockheed)				Boice et al. (1999)
	Routine exposure	0.76 (0.60, 0.95)	78	
	Routine-intermittent exposure <sup>a</sup>	Not reported	173	
Duration of exposure				
	0 yr	1.0	288	
	<1 yr	0.85 (0.65, 1.13)	66	
	1–4 yr	0.98 (0.74, 1.30)	63	
	≥5 yr	0.64 (0.46, 0.89)	44	
	Trend test	$p < 0.05$		
Aerospace workers (Hughes)				Morgan et al. (1998)
	TCE subcohort	1.10 (0.89, 1.34)	97	
	Low intensity (<50 ppm)	1.49 (1.09, 1.99)	45	
	High intensity (>50 ppm)	0.90 (0.67, 1.20)	52	
TCE subcohort (Cox Analysis) <sup>b</sup>				
	Never exposed	1.00 <sup>a</sup>	291	
	Ever exposed	1.14 (0.90, 1.44)	97	
Peak				
	No/Low	1.00 <sup>a</sup>	324	
	Medium/High	1.07 (0.82, 1.40)	64	
Cumulative				
	Referent	1.00 <sup>a</sup>	291	
	Low	1.47 (1.07, 2.03)	45	
	High	0.96 (0.72, 1.29)	52	
Aircraft maintenance workers (Hill Air Force Base, UT)				Blair et al. (1998)
	TCE subcohort			
	Any TCE exposure	0.9 (0.6, 1.3) <sup>a</sup>	109	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-81. Selected results from epidemiologic studies of TCE exposure and lung cancer (continued)**

Exposure group	Relative risk (95% CI)	No. obs. events	Reference
Males, cumulative exposure			Blair et al. (1998) (continued)
0	1.0 <sup>a</sup>	51	
<5 ppm-yr	1.0 (0.7, 1.6)	43	
5–25 ppm-yr	0.9 (0.5, 1.6)	23	
>25 ppm-yr	1.1 (0.7, 1.8)	38	
Females, Cumulative exp			
0	1.0 <sup>a</sup>	2	
<5 ppm-yr	0.6 (0.1, 2.4)	2	
5–25 ppm-yr	0.6 (0.1, 4.7)	11	
>25 ppm-yr	0.4 (0.1, 1.8)	2	
TCE subcohort			Radican et al. (2008)
Any TCE exposure	0.83 (0.63, 1.08)	166	
Males, cumulative exposure	0.91 (0.67, 1.24)	155	
0	1.0 <sup>a</sup>	66	
<5 ppm-yr	0.96 (0.67, 1.37)		
5–25 ppm-yr	0.71 (0.46, 1.11)	31	
>25 ppm-yr	1.00 (0.69, 1.45)	58	
Females, cumulative exposure	0.53 (0.27, 1.07)	11	
0	1.0 <sup>a</sup>		
<5 ppm-yr	0.69 (0.27, 1.77)	5	
5–25 ppm-yr	0.65 (0.16, 2.73)	2	
>25 ppm-yr	0.39 (0.14, 1.11)	4	
Cardboard manufacturing workers in Arnsburg, Germany			
TCE-exposed workers	1.38 (0.55, 2.86)	7	
Unexposed workers	1.06 (0.34, 2.47)	5	
Deaths reported to GE pension fund (Pittsfield, MA)	1.01 (0.69, 1.47) <sup>d</sup>	139	Greenland et al. (1994)
U.S. Coast Guard employees			Blair et al. (1998)
Marine inspectors	0.52 (0.31, 0.82)	18	
Noninspectors	0.81 (0.55, 1.16)	30	
Aircraft manufacturing employees (Italy)			Costa et al. (1989)
All employees	0.99 (0.73, 1.32)	99	
Aircraft manufacturing plant employees (San Diego, CA)			Garabrant et al. (1988)
All subjects	0.80 (0.68, 0.95)	138	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-81. Selected results from epidemiologic studies of TCE exposure and lung cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Lamp manufacturing workers (GE)		0.58 (0.27, 1.27)	6	Shannon et al. (1988)
Rubber industry workers (Ohio)		0.64 ( $p > 0.05$ ) <sup>c</sup>	11	Wilcosky et al. (1984)
<b>Case-control studies</b>				
Population of Montreal, Canada				Siemiatycki et al. (1991)
	Any TCE exposure	0.9 (0.6, 1.5) <sup>c</sup>	21	
	Substantial TCE exposure	0.6 (0.3, 1.2) <sup>c</sup>	9	
<b>Geographic based studies</b>				
Two study areas in Endicott, NY		1.28 (0.99, 1.62)	68	ATSDR (2006a)
Residents of 13 census tracts				Morgan and Cassidy (2002)
	In Redlands, CA	0.71 (0.61, 0.81) <sup>f</sup>	356	
Iowa residents with TCE in water supply				Isacson et al. (1985)
	Males			
	<0.15 µg/L	343.1 <sup>g</sup>	1,181	
	≥0.15 µg/L	345.7 <sup>g</sup>	299	
	Females			
	<0.15 µg/L	58.7 <sup>g</sup>	289	
	≥0.15 µg/L	47.8 <sup>g</sup>	59	

<sup>a</sup>Internal referents, workers not exposed to TCE.

<sup>b</sup>Risk ratio from Cox Proportional Hazard Analysis, stratified by age, sex, and decade (EHS, 1997).

<sup>c</sup>Odds ratio from nested case-control study.

<sup>d</sup>Odds ratio from nested case-control analysis.

<sup>e</sup>90% CI.

<sup>f</sup>99% CI.

<sup>g</sup>Average annual age-adjusted incidence (per 100,000).

GE = General Electric, IBM = International Business Machines Corporation, No. obs. events = number of observed events.

The population studied by Garabrant et al. (1988), ATSDR (2004a) and Chang et al. (2005) are all employees (white- and blue-collar) at a manufacturing facility or plant with potential TCE exposures. Garabrant et al. (1988) observed a 20% deficit in lung cancer mortality (95% CI: 0.68, 0.95) in their study of all employees working for 4 or more years at an aircraft manufacturing company. Blair et al. (1998), a study of Coast Guard marine inspectors with potential for TCE exposure but lacking assessment to individual subjects, observed a 48% deficit in lung cancer mortality (95% CI: 0.31, 0.82). Confidence intervals (95% CI) in Costa et

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 al. (1989), Chang et al. (2005) and ATSDR (2004a) included a risk of 1.0. TCE exposure was  
2 not known for individual subjects in these studies. A wide potential for TCE exposure is likely  
3 ranging from subjects with little to no TCE exposure potential to those with some TCE exposure  
4 potential. Exposure misclassification bias, typically considered as a negative bias, is likely  
5 greater in these studies compared to studies adopting more sophisticated exposure assessment  
6 approaches, which are able to assign quantitative exposure metrics to individual study subjects.  
7 All three studies were of lower likelihood for TCE exposure, in addition to limited statistical  
8 power and other design limitations, and these aspects, in addition to potential exposure  
9 misclassification bias were alternative explanations of observed findings.

10 One population case-control study examined the relationship between lung cancer and  
11 TCE exposure (Siemiatycki, 1991) with risk ratios of 0.9 (95% CI: 0.6, 1.5) for any TCE  
12 exposure and 0.6 (95% CI: 0.3, 1.2) for substantial TCE exposure after adjustment for cigarette  
13 smoking. TCE exposure prevalence in cases in this study was 2.5% for any exposure. Only 1%  
14 had “substantial” (author’s term) exposure, limiting the sensitivity of this study. Relative risks  
15 above 2.0 could only be detected with sufficient (80%) statistical power. The finding of no  
16 association of lung cancer with TCE exposure, therefore, is not surprising. One nested case-  
17 control study of rubber workers observed a smoking unadjusted risk of 0.64 (95% CI: not  
18 presented in paper) in those who had >1 year cumulative exposure to TCE (Wilcosky et al.,  
19 1984).

20 Three geographic based studies reported lung cancer incidence or mortality risks for  
21 drinking water contamination with TCE (ATSDR, 2006a; Isacson et al., 1985; Morgan and  
22 Cassady, 2002). Morgan and Cassidy (2002) observed a relative risk of 0.71 (99% CI: 0.61,  
23 0.81) for lung cancer among residents of Redlands County, CA, whose drinking water was  
24 contaminated with TCE and perchlorate. However, ATSDR (2006a) reported a 28% increase  
25 (95% CI: 0.99, 1.62) in lung cancer incidence among residents living in a area in Endicott, NY,  
26 whose drinking water was contaminated with TCE and other solvents. No information on  
27 smoking patterns is available for individual lung cancer cases as identified by the New York  
28 State Department of Health (NYS DOH) for other cancer cases in this study (ATSDR, 2008).  
29 Isacson et al. (1985) presented lung cancer age-adjusted incidence rates for Iowa residents by  
30 TCE level in drinking water supplies and did not observe an exposure-response gradient.  
31 Exposure information is inadequate in all three of these studies, with monitoring data, if  
32 available, based on few samples and for current periods only, and no information on water  
33 distribution, consumption patterns, or temporal changes. Thus, TCE exposure potential to  
34 individual subjects was not known with any precision, introducing misclassification bias, and  
35 greatly limiting their ability to inform evaluation of TCE and lung cancer.

1 Laryngeal cancer risks are presented in a limited number of cohort studies involving TCE  
2 exposure. No case-control or geographic based studies of TCE exposure were found in the  
3 published literature. All but one of the cohort studies providing information on laryngeal cancer  
4 observed less than five incident cases or deaths. Accordingly, these studies are limited for  
5 examining the relationship between TCE exposure and laryngeal cancer. Risk ratios for  
6 laryngeal cancer are found in Table 4-82.

7 In summary, studies in humans examining lung and laryngeal cancer and TCE exposure  
8 are inconclusive and do not support either a positive or a negative association between TCE  
9 exposure and lung cancer or laryngeal cancer. Raaschou-Nielsen et al. (2003), with the largest  
10 numbers of lung cancer cases of all studies, was the only one to observe a statistically  
11 significantly elevated lung cancer risk with TCE exposure. Raaschou-Nielsen et al. (2003) also  
12 noted several factors that may have confounded or biased their results in either a positive or  
13 negative direction. This study and other cohort studies, as with almost any occupational study,  
14 were not able to control confounding by exposure to chemicals other than TCE (although no  
15 such chemical was apparent in the reports). Information available for factors related to  
16 socioeconomic status (e.g., diet, smoking, alcohol consumption) was also not available. Such  
17 information may positively confound smoking-related cancers such as lung cancer, particularly  
18 in those studies, which adopted national rates to derive expected numbers of site-specific cancer,  
19 if greater smoking rates were over-represented in blue-collar workers or residents of lower socio-  
20 economic status. The finding of a larger risk among subjects with shortest exposure also argues  
21 against a causal interpretation for the observed association for all subjects (NRC, 2006).

22 Four studies reported a statistically significant deficit in lung cancer incidence (Blair et  
23 al., 1998; Boice et al., 1999; Garabrant et al., 1988; Morgan and Cassady, 2002). Absence of  
24 smoking information in these studies would introduce a negative bias if the studied population  
25 smoked less than the referent population and may partially explain the lung cancer decrements  
26 observed in these studies. Morgan and Cassidy (2002) noted the relatively high education high  
27 income levels, and high access to health care of subjects in this study compared to the averages  
28 for the county as a whole, likely leading to a lower smoking rate compared to their referent  
29 population. Garabrant et al. (1988) similarly attributed their observations to negative selection  
30 bias introduced when comparison is made to national mortality rates, also known as a “healthy  
31 worker effect.” The statistically significant decreasing trend in Boice et al. (1999) with exposure  
32 duration to intermittent or routine exposure may reflect a protective effect between TCE and lung  
33 cancer. The use of internal controls in this analysis reduces bias associated with use of an  
34 external population who may have different smoking patterns than an employed population.  
35 However, the exposure assessment approach in this study is limited due to inclusion of subjects

1  
2

**Table 4-82. Selected results from epidemiologic studies of TCE exposure and laryngeal cancer**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort studies—incidence</b>				
Aerospace workers with TCE exposure		Not reported		Zhao et al. (2005)
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure, males	1.2 (0.87, 1.52)	53	
	Any exposure, females	1.7 (0.33, 4.82)	3	
	Employment duration	Not reported		
	<1 yr			
	1–4.9 yr			
	≥5 yr			
Biologically-monitored Danish workers				Hansen et al. (2001)
	Any TCE exposure, males	1.1 (0.1, 3.9)	2	
	Any TCE exposure, females		0 (0.1 exp)	
	Cumulative exposure (Ikeda)	Not reported		
	<17 ppm-yr			
	≥17 ppm-yr			
	Mean concentration (Ikeda)	Not reported		
	<4 ppm			
	4+ ppm			
	Employment duration	Not reported		
	<6.25 yr			
	≥6.25 yr			
Aircraft maintenance workers (Hill Air Force Base, Utah)				Blair et al. (1998)
	TCE subcohort			
	Any exposure	Not reported		
	Males, cumulative exposure	Not reported		
	0			
	<5 ppm-yr			
	5–25 ppm-yr			
	>25 ppm-yr			
	Females, cumulative exposure	Not reported		
	0			
	<5 ppm-yr			
	5–25 ppm-yr			
	>25 ppm-yr			

3

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-82. Selected results from epidemiologic studies of TCE exposure and laryngeal cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Biologically-monitored Finnish workers		Not reported		Anttila et al. (1995)
	Mean air-TCE (Ikeda extrapolation from U-TCA)	Not reported		
	<6 ppm			
	6+ ppm			
Biologically-monitored Swedish workers				Axelson et al. (1994)
	Any TCE exposure, males	1.39 (0.17, 5.00)	2	
	Any TCE exposure, females	Not reported		
<b>Cohort and PMR-mortality</b>				
Computer manufacturing workers (IBM), NY		Not reported		Clapp and Hoffman (2008)
Aerospace workers (Rocketdyne)				
	Any TCE (utility or engine flush workers)	1.45 (0.18, 5.25)	2	Boice et al. (2006b)
	Engine flush—duration of exposure	Not reported		
	Referent			
	0 yr (utility workers with TCE exposure)			
	<4 yr			
	≥4 yr			
	Any exposure to TCE	Not reported		
View-Master employees		Not reported		ATSDR (2004a)
	Males			
	Females			
All employees at electronic factory (Taiwan)				Chang et al. (2003b)
	Males		0 (0.90 exp)	
	Females	0	0 (0.23 exp)	
United States uranium-processing workers (Fernald)				Ritz (1999a)
	Any TCE exposure	Not reported		
	Light TCE exposure, >2 yr duration <sup>4</sup>	Not reported		
	Moderate TCE exposure, >2 yr duration	Not reported		
Aerospace workers (Lockheed)				Boice et al. (1999)
	Routine exposure	1.10 (0.30, 2.82)	4	
	Routine-intermittent exposure	Not reported		

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-82. Selected results from epidemiologic studies of TCE exposure and laryngeal cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Aerospace workers (Hughes)				Morgan et al. (1998)
TCE subcohort	Not reported			
Low intensity (<50 ppm)				
High intensity (>50 ppm)				
Peak	Not reported			
No/low				
Medium/high				
Cumulative	Not reported			
Referent				
Low				
High				
Aircraft maintenance workers (Hill Air Force Base, UT)				Blair et al. (1998)
TCE subcohort	Not reported			
Males, cumulative exposure	Not reported			
0				
<5 ppm-yr				
5–25 ppm-yr				
>25 ppm-yr				
Females, cumulative exposure	Not reported			
0				
<5 ppm-yr				
5–25 ppm-yr				
>25 ppm-yr				
Cardboard manufacturing workers in Arnsburg, Germany	Not reported			Henschler et al. (1995)
Deaths reported to GE pension fund (Pittsfield, MA)	Not examined			Greenland et al. (1994)
U. S. Coast Guard employees				Blair et al. (1998)
Marine inspectors	0.57 (0.01, 3.17)	1		
Noninspectors	0.58 (0.01, 3.20)	1		
Aircraft manufacturing employees (Italy)				Costa et al. (1989)
All employees	0.27 (0.03, 0.98)	2		
Aircraft manufacturing plant employees (San Diego, CA)				Garabrant et al. (1988)
All subjects		0 (7.41 exp)		

1 GE = General Electric, IBM = International Business Machines Corporation, No. obs. events = number of observed  
2 events.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 identified with intermittent TCE exposure (i.e., workers who would be exposed only during  
2 particular shop runs or when assisting other workers during busy periods) (Boice et al., 1999).  
3 The Boice et al. (1999) analysis is based on twice as many lung cancer deaths (i.e., 173 lung  
4 cancer deaths) among subjects with routine or intermittent TCE exposure compared to only  
5 routinely exposed subjects (78 deaths). Subjects identified as intermittently exposed are  
6 considered as having a lower exposure potential than routinely exposed subject and their  
7 inclusion in exposure-response analyses may introduce exposure misclassification bias. Such  
8 bias is a possible explanation for the decreasing trend observation, particularly if workers with  
9 lower potential for TCE exposure have longer exposure (employment) durations.

10 Thus, a qualitative assessment suggests the epidemiological literature on respiratory  
11 cancer and TCE, although limited and of sufficient power to detect only large relative risks, does  
12 not provide strong evidence for any association between TCE exposure and lung cancer. These  
13 studies can only rule out risks of a magnitude of 2.0 or greater for lung cancer and relative risks  
14 greater than 3.0 or 4.0 for laryngeal cancer for exposures to studied populations.

## 4.7.2. Laboratory Animal Studies

### 4.7.2.1.1. Respiratory Tract Animal Toxicity

16 Limited studies are available to determine the effects of TCE exposure on the respiratory  
17 tract (summarized in Table 4-83). Many of these studies in mice have examined acute effects  
18 following intraperitoneal administration at relatively high TCE doses. However, effects on the  
19 bronchial epithelium have been noted in mice and rats with TCE administered via gavage, with  
20 doses 1,000 mg/kg-day and higher reported to cause rales and dyspnea (Narotsky et al., 1995)  
21 and pulmonary vasculitis (NTP, 1990) in rats. Mice appear to be more sensitive than rats to  
22 histopathological changes in the lung via inhalation; pulmonary effects are also seen in rats with  
23 gavage exposure. It is difficult to compare intraperitoneal to oral and inhalation routes of  
24 exposure given the risk of peritonitis and paralytic ileus. Any inflammatory response from this  
25 route of administration can also affect the pulmonary targets of TCE exposure such as the Clara  
26 cells.

27 This section reviews the existing literature on TCE, and the role of the various TCE  
28 metabolites in TCE-induced lung effects. The most prominent toxic effect reported is damage to  
29 Clara cells in mouse lung. The nonciliated, columnar Clara cells comprise the majority of the  
30 bronchiolar and terminal bronchiolar epithelium in mice, and alveolar Type I and Type II cells  
31 constitute the alveolar epithelium. These cells have been proposed as a progenitor of lung

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 adenocarcinomas in both humans and mice (Kim et al., 2005). Long-term studies have not  
2 focused on the detection of pulmonary adenoma carcinomas but have shown a consistently

**Table 4-83. Animal toxicity studies of trichloroethylene**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Results
Green et al. (1997b)	CD-1 mice (F)	Inhalation	450-ppm, 6 h/d, 5 d with 2 d break then 5 more days; sacrificed 18 h after 1, 5, 6, or 10 exposures	5/group	Increased vacuolation and proliferation of Clara cells caused by accumulation of chloral.
Forkert and Forkert (1994)	CD-1 mice (M)	Intraperitoneal injection	2,000 mg/kg in corn oil (0.01 mL/g BW); sacrificed 15, 30, 60 and 90 d after single exposure	10/group	Increased fibrotic lesions, with early signs visible at 15 d postexposure.
Villaschi et al. (1991)	BC3F1 mice (M)	Single inhalation	30 min 500, 1,000, 2,000, 3,500, and 7,000 ppm; sacrificed 2 h, 24 h, 2, 5, or 7 d post exposure	3/group	Increased vacuolation and proliferation of nonciliated bronchial cells. Injury was maximal at 24 h with some repair occurring between 24 h and 48 h.
Odum et al. (1992)	CD-1 mice (F)	Inhalation	6 h/d; separate repeated study in mice: 450 ppm for 6 h/d, 5 d/wk for 2 wk; sacrificed 24 h after exposure; repeat study sacrificed at 2, 5, 6, 8, 9, 12, or 13 d; mice: 20, 100, 200, 450, 1,000, or 2,000 ppm	4/group	Dose-dependent increase in Clara cell vacuolation in mice after a single exposure, resolved after 5 d repeated exposures but recurred following a 2-d break from exposure. Changes accompanied by decrease in CYP activity in mice. Exposure to chloral alone demonstrated similar response as TCE exposure in mice. No changes were seen in rats.
	Alpk APfSD rats (F)	Inhalation	6 h/d; repeat study sacrificed at 2, 5, 6, 8, 9, 12, or 13 d; rats: 500, or 1,000 ppm	4/group	
Kurasawa (1988) (translation)	Ethanol-treated (130) and nontreated (110) Wistar rats (M)	Inhalation	500, 1,000, 2,000, 4,000, and 8,000 ppm for 2 h; sacrificed 22 h after exposure	10/group	TCE exposure resulted in highly selective damage to Clara cells that occurred between 8 and 22 h after the highest exposure with repair by 4 wk post exposure.
Forkert et al. (2006)	CD-1 mice (M); wild-type (mixed 129/Sv and C57BL) and CYP2E1-null mice (M)	Intraperitoneal injection	500, 750, and 1,000 mg/kg in corn oil; for inhibition studies mice pretreated with 100 mg/kg diallyl sulfone; for immunoblotting, 250, 500, 750, and 1,000 mg/kg; for PNP hydroxylation, 50, 100, 250, 500, 750, and 1,000 mg/kg; sacrificed 4 h after exposure	4/group	TCE bioactivation by CYP2E1 and/or 2F2 correlated with bronchiolar cytotoxicity in mice.

**Table 4-83. Animal toxicity studies of trichloroethylene (continued)**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Results
Forkert et al. (1985)	CD-1 mice (M)	Intraperitoneal injection	2,000, 2,500 or 3,000 mg/kg in mineral oil; sacrificed 24 h postexposure for dose response; time course sacrificed 1, 2, 12, and 24 h postexposure	10/group	Clara cell injury was increased following exposure at all doses tested; time course demonstrated a rapid and marked reduction in pulmonary microsomal cytochrome P450 content and aryl hydrocarbon hydroxylase activity. Alveolar Type II cells were also affected.
Forkert and Birch (1989)	CD-1 mice (M)	Intraperitoneal injection	2,000 mg/kg in corn oil; sacrificed 1, 2, 4, 8, 12, and 24 h postexposure	10/group	Necrotic changes seen in Clara cells as soon as 1 h postexposure; increased vacuolation was seen by 4 h postexposure; covalent binding of TCE to lung macromolecules peaked at 4 h and reached a plateau at 12 and 24 h post exposure.
Stewart et al. (1979) Le Mesurier et al., (1980)	Wistar Rats (F)	Inhalation (whole-body chamber)	30 min, 48.5 g/m <sup>3</sup> (9,030 ppm); sacrificed at 5 and 15 d postexposure	5/group	Decreased recovery of pulmonary surfactant (dose-dependent).
Lewis et al. (1984)	Mice	Inhalation (Pyrex bell jars)	10,000 ppm, 1–4 h daily for 5 consecutive days; sacrificed 24 h after last exposure	~28/group	Increased vacuolation and reduced activity of pulmonary mixed function oxidases.
Scott et al. (1988)	CD-1 mice (M)	Intraperitoneal injection	single injection of 2,500–3,000 mg/kg, sacrificed 24 h postexposure	4/group	Clara cells were damaged and exfoliated from the epithelium of the lung.
NTP (1990)	F344 rats (M,F) B6C3F1 mice (M,F)	Gavage	Male rats: 0, 125, 250, 500, 1,000, and 2,000 mg/kg BW (corn oil); female rats: 0, 62.5, 125, 250, 500 or 1,000 mg/kg BW (corn oil); Mice: 0, 375, 750, 1,500, 3,000, and 6,000 mg/kg BW (corn oil); dosed 5d/w for 13 wk	10/group	Increased pulmonary vasculitis in the high-dose groups of male and female rats (6/10 group as compared to 1/10 in controls). No pulmonary effects described in mice at this time point.

**Table 4-83. Animal toxicity studies of trichloroethylene (continued)**

Reference	Animals (sex)	Exposure route	Dose/exposure concentration	Exposed	Results
Prendergast et al. (1967)	Sprague-Dawley or Long-Evans rats; Hartley Guinea pigs; New Zealand albino rabbits; beagle dogs; squirrel monkeys (sex not given for any species)	Inhalation	730 ppm for 8 h/d, 5 d/w, 6 wk or 35 ppm for 90 d constant	Rats (15); guinea pigs (15); rabbit (3); dog (2); monkey (3)	No histopathological changes observed, although rats were described to show a nasal discharge in the 6 wk study. No quantification was given.
Narotsky et al. (1995)	F344 rats (F)	Gavage	0, 1,125, 1,500 mg/kg-day	21, 16, or 17 per group	Rales and dyspnea were observed in the TCE high-dose group; two females with dyspnea subsequently died.

1 positive response in mice but not rats. However, chronic toxicity data on noncancer effects are  
2 very limited.  
3

#### 4.7.2.1.2. Acute and short-term effects: inhalation

4 Relatively high-dose single and multiple inhalation exposures to TCE result in dilation of  
5 endoplasmic reticulum and vacuolation of nonciliated (Clara) cells throughout the bronchial tree  
6 in mice. A single study in rats reported similar findings. In mice, single exposure experiments  
7 show vacuolation at all dose levels tested with the extent of damage increasing with dose.  
8 Villaschi et al. (1991) reported similar degrees of vacuolation in B6C3F1 mice (3/group) at  
9 24 hours after the start of exposure across all tested doses (500, 1,000, 2,000, 3,500, and  
10 7,000 ppm, 30 minutes), with the percentage of the nonciliated cells remaining vacuolated at  
11 48 hours increasing with dose. Clara cell vacuolation was reported to be resolved 7 days after  
12 single 30 minute exposure to TCE. Odum et al. (1992) reported that, when observed 24 hours  
13 after the start of 6 hours exposure, the majority of Clara cells in mice were unaffected at the  
14 lowest dose of 20 ppm exposures, while marked vacuolation was observed at 200 ppm (no  
15 quantitative measures of damage given and only 3 animals per group were examined).

16 In rats, Odum et al. (1992) reported no morphological changes in the female Alpk APfSD  
17 rat epithelium after 6 hours exposure (500 or 1,000 ppm) when observed 24 hours after the start  
18 of exposure ( $n = 3$ /group). However, Kurasawa reported pronounced dose-related morphological  
19 changes in Clara cells at the highest dose (8,000 ppm) for 2 hours in Wistar rats ( $n = 10$  per  
20 group). At 500 and 1,000 ppm, slight dilation of the apical surface was reported, but  
21 morphological measurements (the ratio of the lengths of the apical surface to that of the base line  
22 of apical cytoplasm) were not statistically-significantly different from controls. From  
23 2,000–8,000 ppm, a progressively increasing flattening of the apical surface was observed. In  
24 addition, at 2,000 ppm, slight dilation of the smooth endoplasmic reticulum was also observed,  
25 with marked dilation and possible necrosis at 8,000 ppm. Kurasawa (1988) also examined the  
26 time-course of Clara cell changes following a single 8,000-ppm exposure, reporting the greatest  
27 effects at 1 day to 1 week, repair at 2 weeks, and nearly normal morphology at 4 weeks. The  
28 only other respiratory effect that has been reported from one study in rats exposed via inhalation  
29 is a reduction in pulmonary surfactant yield following 30 minute exposures at 9,030 ppm for 5 or  
30 15 days (Stewart et al., 1979). Therefore, single inhalation experiments (Kurasawa, 1988; Odum  
31 et al., 1992; Villaschi et al., 1991) suggest that the Clara cell is the target for TCE exposure in  
32 both rats and mice and that mice are more susceptible to these effects. However, the database is

1 limited in its ability to discern quantitative differences in susceptibility or the nature of the dose-  
2 response after a single dose of TCE.

3 Other experiments examined the effects of several days of TCE inhalation exposure in  
4 mice and potential recovery. While single exposures require 1–4 weeks for complete recovery,  
5 after short-term repeated exposure, the bronchial epithelium in mice appears to either adapt to or  
6 become resistant to damage Odum et al. (1992) and Green et al. (1997b) observed Clara cells in  
7 mice to be morphologically normal at the end of exposures 6 hours/day for 4 or 5 days. As with  
8 single dose experiments, the extent of recovery in multidose exposures may be dose-dependent.  
9 Using a very high dose, Lewis et al. (1984) report vacuolation of bronchial epithelial cells after  
10 4 hours/day, but not 1 hours/day, (10,000 ppm) for 5 days in mice. In addition, Odum et al.  
11 (1992) reported that the damage to Clara cells that resolved after repeated exposures of 5 days, a  
12 sign of adaptation to TCE exposure, returned when exposure was resumed after 2 days.

13 In rats, only one inhalation study reported in two published articles (Le Mesurier et al.,  
14 1980; Stewart et al., 1979) using repeated exposures examined pulmonary histopathology.  
15 Interestingly, this study reported vacuolation in Type 1 alveolar cells, but not in Clara cells, after  
16 5 days of exposure to approximately 9,030 ppm for 30 minutes/day (only dose tested). In  
17 addition, abnormalities were observed in the endothelium (bulging of thin endothelial segments  
18 into the microcirculatory lumen) and minor morphological changes in Type 2 alveolar cells.  
19 Although exposures were carried out for 5 consecutive days, histopathology was recorded up to  
20 15 days post exposure, giving cell populations time to recover. Because earlier time points were  
21 not examined, it is not possible to discern whether the lack of reported Clara cell damage in rats  
22 following repeated exposure is due to recovery or lack of toxicity in this particular experiment.

23 Although recovery of individual damaged cells may occur, cell proliferation, presumed  
24 from labeling index data suggestive of increased DNA synthesis, contributes, at least in part, to  
25 the recovery of the bronchial epithelium in mice. Villaschi et al. (1991) observed a dose-  
26 dependent increase in labeling index as compared to controls in the mouse lung at 48 hours after  
27 a single TCE exposure (30 minutes; 500, 1,000, 2,000, 3,500, or 7,000 ppm), which decreased to  
28 baseline values at 7 days postexposure. Morphological analysis of cells was not performed,  
29 although the authors stated the dividing cells had the appearance of Clara cells. Interestingly,  
30 Green et al. (1997b) reported no increase in BrdU labeling 24 hours after a single exposure  
31 (6 hours 450 ppm), but did see increased BrdU labeling at the end of multiple exposures  
32 (1/day, 5 days) while Villaschi et al. (1991) reported increased [<sup>3</sup>H]Thymidine labeling 2, 5, and  
33 7 days after single 30 minute exposures to 500–7,000 ppm. Therefore, the data for single  
34 exposures at 450–500 ppm may be consistent if increased cell proliferation occurred only for a  
35 short period of time around 48 hours postexposure, and was thereby effectively washed-out by



1 the longer “averaging time” in the experiments by Green et al. (1997b). Also, these  
2 contradictory results may be due to differences in methodology. Green et al. (1997b) and  
3 Villaschi et al. (1991) reported very different control labeling indices (6 and 0%, respectively)  
4 while reporting similar absolute labeling indices at 450–500 ppm (6.5 and 5.2%, respectively).  
5 The different control values may be a result of substantially-different times over which the label  
6 was incorporated: the mice in Green et al. (1997b) were given BrdU via a surgically-implanted  
7 osmotic pump over 4 days prior to sacrifice, while the mice in Villaschi et al. (1991) were given  
8 a single intraperitoneal dose of [<sup>3</sup>H]Thymidine 1 hour prior to sacrifice. Stewart et al. (1979)  
9 observed no stimulation of thymidine incorporation after daily exposure to TCE (9,000 ppm) for  
10 up to 15 days. This study did, however, report a nonstatistically significant reduction in orotate  
11 incorporation, an indicator of RNA synthesis, after 15 days, although the data was not shown.

12 At the biochemical level, changes in pulmonary metabolism, particularly with respect to  
13 CYP activity, have been reported following TCE exposure via inhalation or intraperitoneal  
14 administration in mice. Odum et al. (1992) reported reduced enzyme activity in Clara cell  
15 sonicates of ethoxycoumarin *O*-deethylase, aldrin epoxidation, and nicotinamide adenine  
16 dinucleotide phosphate-oxidase (NADPH) cytochrome c reductase after 6 hour exposures to  
17 20–2,000 ppm TCE, although the reduction at 20 ppm was not statistically significant. No  
18 reduction of GST activity as determined by chlorodinitrobenzene as a substrate was detected.  
19 With repeated exposure at 450 ppm, the results were substrate-dependent, with ethoxycoumarin  
20 *O*-deethylase activity remaining reduced, while aldrin epoxidation and NADPH cytochrome c  
21 reductase activity showing some eventual recovery by 2 weeks. The results reported by Odum et  
22 al. (1992) for NADPH cytochrome c reductase were consistent with those of Lewis et al. (1984),  
23 who reported similarly reduced NADPH cytochrome c reductase activity following a much  
24 larger dose of 10,000 ppm for 1 and 4 hours/day for 5 days in mice (strain not specified). TCE  
25 exposure has also been associated with a decrease in pulmonary surfactant. Repeated exposure  
26 of female Wistar rats to TCE (9,000 ppm, 30 minutes/day) for 5 or 15 days resulted in a  
27 significant decrease in pulmonary surfactant as compared to unexposed controls (Le Mesurier et  
28 al., 1980).

29

#### 4.7.2.1.2.1. Acute and short-term effects: intraperitoneal injection and gavage exposure

30 As stated above the intraperitoneal route of administration is not a relevant paradigm for  
31 human exposure. A number of studies have used this route of exposure to study the effects of  
32 acute TCE exposure in mice. In general, similar lung targets are seen following inhalation or  
33 intraperitoneal treatment in mice (Forkert et al., 2006; Forkert and Birch, 1989; 1985; Scott et al.,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 1988). Inhalation studies generally reported the Clara cell as the target in mice. No lung  
2 histopathology from intraperitoneal injection studies in rats is available. Forkert et al. (1985) and  
3 Forkert and Birch (1989) reported vacuolation of Clara cells as soon as 1 hour following  
4 intraperitoneal administration of a single dose of 2,000 mg/kg in mice. At 2,500 mg/kg, both  
5 Forkert et al. (1985) and Scott et al. (1988) reported exfoliation of Clara cells and parenchymal  
6 changes, with morphological distortion in alveolar Type II cells and inconsistently observed  
7 minor swelling in Type I cells at 24 hours postexposure. Furthermore, at 3,000 mg/kg,  
8 Scott et al. (1988) also reported a significant (85%) decrease in intracellularly stored surfactant  
9 phospholipids at 24 hours postexposure. These data indicate that both Clara cells and alveolar  
10 Type I and II cells are targets of TCE toxicity at these doses and using this route of  
11 administration. Recently, Forkert et al. (2006) reported Clara cell toxicity that showed increased  
12 severity with increased dose (pyknotic nuclei, exfoliation) at 500–1,000 mg/kg intraperitoneal  
13 doses as soon as 4 hours postexposure in mice. Even at 500 mg/kg, a few Clara cells were  
14 reported with pyknotic nuclei that were in the process of exfoliation. Damage to alveolar Type II  
15 cells was not observed in this dose range. The study by Scott et al. (1988) examined surfactant  
16 phospholipids and phospholipase A2 activity in male CD-1 mice exposed by intraperitoneal  
17 injection of TCE (2,500 or 3,000 mg/kg, 24 hours). The lower concentration led to damage to  
18 and exfoliation of Clara cells from the epithelial lining into the airway lumen, while only the  
19 higher concentration led to changes in surfactant phospholipids. This study demonstrated an  
20 increase in total phospholipid content in the lamellar body fractions in the mouse lung.

21 The study by Narotsky et al. (1995) exposed F344 timed-pregnant rats to TCE (0, 1,125,  
22 and 1,500 mg/kg BW) by gavage and examined both systemic toxicity and developmental effects  
23 at 14 days postexposure. Rales and dyspnea in the dams were observed in the high-dose group,  
24 with two of the animals with dyspnea subsequently dying. The developmental effects observed  
25 in this study are discussed in more detail in Section 4.8.

26

#### 4.7.2.1.2.2. Subchronic and chronic effects

27 There are a few reports of the subchronic and chronic noncancer effects of TCE on the  
28 respiratory system from intraperitoneal exposure in mice and from gavage exposure in rats.  
29 Forkert and Forkert (1994) reported pulmonary fibrosis in mice 90 days after intraperitoneal  
30 administration of a single 2,000 mg/kg dose of TCE. The effects were in the lung parenchyma,  
31 not the bronchioles where Clara cell damage has been observed after acute exposure. It is  
32 possible that fibrotic responses in the alveolar region occur irrespective of where acute injury  
33 occurs. Effects upon Clara cells can also impact other areas of the lung via cytokine regulation

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 (Elizur et al., 2008). Alternatively, the alveolar and/or capillary components of the lung may  
2 have been affected by TCE in a manner that was not morphologically apparent in short-term  
3 experiments. In addition effects from a single or a few short-term exposures may take longer to  
4 manifest. The latter hypothesis is supported by the alveolar damage reported by Odum et al.  
5 (1992) after chloral administration by inhalation, and by the adducts reported in alveolar Type II  
6 cells by Forkert et al. (2006) after 500–1,000 mg/kg TCE intraperitoneal administration.

7 As noted previously, rats have responded to short-term inhalation exposures of TCE with  
8 Clara cell and alveolar Type I and II effects. After repeated inhalation exposures over 6 weeks  
9 (8 hours/day, 5 days/week, 730 ppm) and continuous exposures over 90 days (35 ppm),  
10 Prendergast et al. (1967) noted no histopathologic changes in rats, guinea pigs, rabbits, dogs, or  
11 monkeys after TCE exposure, but did describe qualitatively observing some nasal discharge in  
12 the rats exposed for 6 weeks. The study details in Prendergast et al. (1967) are somewhat  
13 limited. Exposed animals are described as “typically” 15 Long-Evans or Sprague-Dawley rats,  
14 15 Hartley guinea pigs, 3 squirrel monkeys, 3 New Zealand albino rabbits, and 2 beagle dogs.  
15 Controls were grouped between studies. In a 13-week NTP study in F344/N rats ( $n = 10/\text{group}$ )  
16 exposed to TCE (0–2,000 mg/kg-day 5 days/week) by gavage, pulmonary vasculitis was  
17 observed in 6/10 animals of each sex of the highest dose group (2,000 mg/kg-day), in contrast  
18 to 1/10 in controls of each sex (NTP, 1990).

#### 4.7.2.1.3. Respiratory Tract Cancer

20 Limited studies have been performed examining lung cancer following TCE exposure  
21 (summarized in Table 4-84). TCE inhalation exposure was reported to cause statistically  
22 significant increase in pulmonary tumors (i.e., pulmonary adenocarcinomas) in some studies in  
23 mice, but not in studies in rats and hamsters. Oral administration of TCE frequently resulted in  
24 elevated lung tumor incidences in mice, but not in any tested species was there a statistically  
25 significant increase. This section will describe the data regarding TCE induction of pulmonary  
26 tumors in rodent models. The next sections will consider the role of metabolism and potential  
27 MOAs for inhalation carcinogenicity, primarily in mice.

#### 4.7.2.1.4. Inhalation

29 There are three published inhalation studies examining the carcinogenicity of TCE at  
30 exposures from 0–600 ppm, two of which reported statistically significantly increased lung  
31 tumor incidence in mice at the higher concentrations (Fukuda et al., 1983; Henschler et al., 1980;

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 1988; Maltoni et al., 1986). Rats and hamsters did not show an increase in lung tumors  
2 following exposure.  
3

**Table 4-84. Animal carcinogenicity studies of trichloroethylene**

Reference	Animals (sex)	Exposure route	Dose/exp conc (stabilizers, if any)	Pulmonary tumor incidences	
				Benign + malignant	Malignant only
Fukuda et al. (1983)	ICR mice (F) S-D rats (F)	Inhalation, 7 h/d, 5 d/wk, 104 wk, hold until 107 wk	0, 50, 150, or 450 ppm (epichlorohydrin)	Mice: 6/49, 5/50, 13/50, 11/46 Rats: 0/50, 0/50, 1/47, 1/51	Mice: 1/49, 3/50, 8/50 <sup>a</sup> , 7/46 <sup>a</sup> Rats: none
Maltoni et al. (1988; 1986)	S-D rats (M, F) Swiss mice (M, F) B6C3F1 mice (M, F)	Inhalation, 7 h/d, 5 d/wk, 104 wk, hold until death	0, 100, 300, or 600 ppm	Rats: 0/280, 0/260, 0/260, 0/260 Swiss Mice: M: 10/90, 11/90, 23/90 <sup>a</sup> , 27/90 <sup>b</sup> ; F: 15/90, 15/90, 13/90, 20/90 B6C3F1 Mice: M: 2/90, 2/90, 3/90, 1/90; F: 4/90, 6/90, 7/90, 15/90 <sup>a</sup>	Rats: 0/280, 0/260, 0/260, 0/260 Swiss Mice: M: 0/90, 0/90, 0/90, 1/90; F: 2/90, 0/90, 0/90, 2/90 B6C3F1 Mice M: 0/90, 0/90, 0/90, 0/90; F: 0/90, 1/90, 0/90, 0/90
Henschler et al. (1980)	Wistar rats (M, F) Syrian hamsters (M, F) NMRI mice	Inhalation, 6 h/d, 5 d/wk, 78 wk, hold until 130 wk (mice and hamsters) or 156 wk (rats)	0, 100, or 500 ppm (triethanolamine)	Rats: M: 1/29, 1/30, 1/30; F: 0/28; 1/30; 0/30 Hamsters: 0/60, 0/59, 0/60 Mice: M: 1/30, 3/29, 1/30; F: 3/29, 0/30, 1/28	Rats: M: 1/29, 1/30, 1/30; F: 0/28; 1/30; 0/30 Hamsters: 0/60, 0/59, 0/60 Mice: M: 5/30, 3/29, 1/30; F: 1/29, 3/30, 0/28
Henschler et al. (1984)	Swiss mice (M, F)	Gavage, 5/wk, 72 wk hold 104 wk	2.4 g/kg BW (M), 1.8 g/kg BW (F) all treatments; (control, triethanolamine, industrial, epichlorohydrin, 1,2-epoxybutane, both)	M: 18/50, 17/50, 14/50, 21/50, 15/50, 18/50; F: 12/50, 20/50, 21/50, 17/50, 18/50, 18/50	M: 8/50, 6/50, 7/50, 5/50, 7/50, 7/50; F: 5/50, 11/50, 8/50, 3/50, 7/50, 7/50
Van Duuren et al. (1979)	Swiss mice (M, F)	Gavage, 1/wk, 89 wk	0 or 0.5 mg (unknown)	0/30 for all groups	0/30 for all groups
NCI (1976)	Osborne-Mendel rats (M, F) B6C3F1 mice (M, F)	Gavage, 5/wk, 78 wk, hold until 110 wk (rats) or 90 wk (mice)	Rats: TWA: 0, 549, or 1,097 mg/kg Mice: TWA: M: 0, 1,169, or 2,339 mg/kg; F: 0, 869, or 1,739 mg/kg (epoxybutane, epichlorohydrin)	Rats: M: 1/20, 0/50, 0/50; F: 0/20, 1/47, 0/50 Mice: M: 0/20, 5/50, 2/48; F: 1/20, 4/50, 7/47	Rats: M: 0/20, 0/50, 0/50; F: 0/20, 1/47, 0/50 Mice: M: 0/20, 0/50, 1/48; F: 0/20, 2/50, 2/47

**Table 4-84. Animal carcinogenicity studies of trichloroethylene (continued)**

Reference	Animals (sex)	Exposure route	Dose/exp conc (stabilizers, if any)	Pulmonary tumor incidences	
				Benign + malignant	Malignant only
NTP (1988)	ACI, August, Marshall, Osborne-Mendel rats	Gavage, 1/d, 5 d/wk, 103 wk	0, 500, or 1,000 mg/kg (diisopropylamine)	ACI M: 1/50, 4/47, 0/46; F: 0/49, 2/47, 2/42 August M: 1/50, 1/50, 0/49; F: 1/50, 1/50, 0/50 Marshall M: 3/49, 2/50, 2/47; F: 3/49, 3/49, 1/46 Osborne-Mendel M: 2/50, 1/50, 1/50; F: 0/50, 3/50, 2/50	ACI M: 1/50, 2/47, 0/46; F: 0/49, 1/47, 2/42 August M: 0/50, 1/50, 0/49; F: 1/50, 0/50, 0/50 Marshall M: 3/49, 2/50, 2/47; F: 3/49, 3/49, 1/46 Osborne-Mendel M: 1/50, 1/50, 0/50; F: 0/50, 3/50, 1/50
NTP (1990)	F344 rats (M, F) B6C3F1 mice (M, F)	Gavage, 1/d, 5 d/wk, 103 wk	Mice: 0 or 1,000 mg/kg Rats: 0, 500, 1,000 mg/kg	Mice: M: 7/49, 6/50; F: 1/48, 4/49 Rats: M: 4/50, 2/50, 3/49; F: 1/50, 1/49, 4/50	Mice: M: 3/49, 1/50; F: 1/48, 0/49 Rats: M: 3/50, 2/50, 3/49; F: 0/50, 0/49, 2/50
Maltoni et al. (1986)	S-D rats (M, F)	Gavage, 1/d, 4–5 d/wk, 56 wk; hold until death	0, 50 or 250 mg/kg	M: 0/30, 0/30, 0/30; F: 0/30, 0/30, 0/30	M: 0/30, 0/30, 0/30; F: 0/30, 0/30, 0/30

<sup>a</sup> Statistically-significantly different from controls by Fisher's exact test ( $p < 0.05$ ).

<sup>b</sup> Statistically-significantly different from controls by Fisher's exact test ( $p < 0.01$ ).

1 The inhalation studies by Fukuda et al. (1983), which involved female ICR mice and  
2 Sprague-Dawley rats, observed a threefold increase in lung tumors per mouse in those exposed  
3 to the two higher concentrations (150–450 ppm) but reported no increase in lung tumors in the  
4 rats. Maltoni et al. (1988; 1986) reported statistically-significantly increased pulmonary tumors  
5 in male Swiss and female B6C3F1 mice at the highest dose of 600 ppm, but no significant  
6 increases in any of the other species/strains/sexes tested. Henschler et al. (1980) tested NMRI  
7 mice, Wistar rats and Syrian hamsters of both sexes, and reported no observed increase in  
8 pulmonary tumors any of the species tested (see Section 4.4 and Appendix E for details of the  
9 conduct of these studies).

#### 4.7.2.1.5. Gavage

11 None of the six chronic gavage studies (Henschler et al., 1984; Maltoni et al., 1986; NCI,  
12 1976; NTP, 1988, 1990; Van Duuren et al., 1979), which exposed multiple strains of rats and  
13 mice to 0–3,000 mg/kg TCE for at least 56 weeks, reported a statistically-significant excess in  
14 lung tumors, although nonstatistically-significant increases were frequently observed in mice.

15 The study by Van Duuren et al. (1979) examined TCE along with 14 other halogenated  
16 compounds for carcinogenicity in both sexes of Swiss mice. While no excess tumors were  
17 observed, the dose rate of 0.5 mg once per week is equivalent to an average dose rate of  
18 approximately 2.4 mg/kg-day for a mouse weighing 30 g, which is about 400-fold smaller than  
19 that in the other gavage studies. In the NCI (1976) study, the results for Osborne-Mendel rats  
20 were considered inconclusive due to significant early mortality, but female B6C3F1 mice  
21 (though not males) exhibited a nonstatistically-significant elevation in pulmonary tumor  
22 incidence. The NCI study (1976) used technical grade TCE which contained two known  
23 carcinogenic compounds as stabilizers (epichlorohydrin and 1,2-epoxybutane), but a later study  
24 by Henschler et al. (1984) in which mice were given TCE that was either pure, industrial, and  
25 stabilized with one or both of these stabilizers found similar pulmonary tumors regardless of the  
26 presence of stabilizers. In this study, female mice ( $n = 50$ ) had elevated, but again not  
27 statistically-significant, increases in pulmonary tumors. A later gavage study by NTP (1988),  
28 which used TCE stabilized with diisopropylamine, observed no pulmonary tumors, but chemical  
29 toxicity and early mortality rendered this study inadequate for determining carcinogenicity. The  
30 final NTP study (1990) in male and female F344 rats and B6C3F1 mice, using epichlorohydrin-  
31 free TCE, again showed early mortality in male rats. Similar to the other gavage studies, a  
32 nonstatistically significant elevation in (malignant) pulmonary tumors was observed in mice, in  
33 this case in both sexes. These animal studies show that while there is a limited increase in lung

1 tumors following gavage exposure to TCE in mice, the only statistically significant increase in  
2 lung tumors occurs following inhalation exposure in mice.

### 4.7.3. Role of Metabolism in Pulmonary Toxicity

3 TCE oxidative metabolism has been demonstrated to play a main role in TCE pulmonary  
4 toxicity in mice. However, data are not available on the role of specific oxidative metabolites in  
5 the lung. The Clara cell is thought to be the cell type responsible for much of the CYP  
6 metabolism in the lung. Therefore, damage to this cell type would be expected to also affect  
7 metabolism. More direct measures of CYP and isozyme-specific depression following TCE  
8 exposure have been reported following intraperitoneal administration in mice. Forkert et al.  
9 (1985) reported significant reduction in microsomal aryl hydrocarbon hydroxylase activity as  
10 well as CYP content between 1 and 24 hours after exposure (2,000–3,000 mg/kg i.p. TCE).  
11 Maximal depression occurred between 2 and 12 hours, with aryl hydrocarbon hydroxylase  
12 activity (a function of CYP) less than 50% of controls and CYP content less than 20% of  
13 controls. While there was a trend towards recovery from 12–24 hours, depression was still  
14 significant at 24 hours. Forkert et al. (2005) reported decreases in immunoreactive CYP2E1,  
15 CYP2F2, and CYP2B1 in the 4 hours after TCE treatment with 750 mg/kg intraperitoneal  
16 injection in mice. The amount and time of maximal reduction was isozyme dependent  
17 (CYP2E1: 30% of controls at 2 hours; CYP2F2: abolished at 30 minutes; CYP2B1: 43% of  
18 controls at 4 hours). Catalytic markers for CYP2E1, CYP2F2, and CYP2B enzymes showed  
19 rapid onset (15 minutes or less after TCE administration) of decreased activity, and continued  
20 depression through 4 hours. Decrease in CYP2E1 and CYP2F2 activity (measured by PNP  
21 hydroxylase activity) was greater than that of CYP2B (measured by pentoxyresorufin  
22 *O*-dealkylase activity). Forkert et al. (2006) reported similar results in which 4 hours after  
23 treatment, immunodetectable CYP2E1 protein was virtually abolished at doses 250–1,000 mg/kg  
24 and immunodetectable CYP2F2 protein, while still detectable, was reduced. PNP hydroxylase  
25 activity was also reduced 4 hours after treatment to 37% of controls at the lowest dose tested of  
26 50 mg/kg, with further decreases to around 8% of control levels at doses of 500 mg/kg and  
27 higher. These results correlate with previously described increases in Clara cell cytotoxicity, as  
28 well as dichloroacetyl lysine (DAL) protein adduct formation. DAL adducts were observed in  
29 the bronchiolar epithelium of CD-1 mice and most prominent in the cellular apices of Clara cells  
30 (Forkert et al., 2006). This study also examined the effect of TCE in vitro exposure on the  
31 formation of chloral hydrate in lung microsomes from male CD-1 mice and CYP2E1 knock-out  
32 mice. The rates of CH formation were the same for lysosomes from both CD-1 and CYP2E1



1 knockout mice from 0.25 mM to 0.75 mM, but the CH formation peaked earlier for in the wild-  
2 type lysosomes (0.75 mM) as compared to CYP2E1-null lysosomes (1 mM).

3 The strongest evidence for the necessary role of TCE oxidation is that pretreatment of  
4 mice with diallyl sulfone (DASO<sub>2</sub>), an inhibitor of CYP2E1 and CYP2F2, protected against  
5 TCE-induced pulmonary toxicity. In particular, following an intraperitoneal TCE dose of  
6 750 mg/kg, Clara cells and the bronchiolar epithelium in mice pretreated with the  
7 CYP2E1/CYP2F2 inhibitor appeared normal. In naive mice given the same dose, the epithelium  
8 was attenuated due to exfoliation and there was clear morphological distortion of Clara cells  
9 (Forkert et al., 2005). In addition, the greater susceptibility of mouse lungs relative to rat lungs  
10 is consistent with their larger capacity to oxidize TCE, as measured in vitro in lung microsomal  
11 preparations (Green et al., 1997b). Analysis by immunolocalization also found considerably  
12 higher levels of CYP2E1 in the mouse lung, heavily localized in Clara cells, as compared to rat  
13 lungs, with no detectable CYP2E1 in human lung samples (Green et al., 1997b). In addition,  
14 both Green et al. (1997b) and Forkert et al. (2006) report substantially lower metabolism of TCE  
15 in human lung microsomal preparations than either rats or mice. It is clear that CYP2E1 is not  
16 the only CYP enzyme involved in pulmonary metabolism, as lung microsomes from CYP2E1-  
17 null mice showed greater or similar rates of CH formation compared to those from wild-type  
18 mice. Recent studies have suggested a role for CYP2F2 in TCE oxidative metabolism, although  
19 more work is needed to make definitive conclusions. In addition, there may be substantial  
20 variability in human lung oxidative metabolism, as Forkert et al. (2006) reported that in  
21 microsomal samples from eight individuals, five exhibited no detectable TCE oxidation (<0.05  
22 pmol/mg protein/20 minutes), while others exhibited levels well above the limit of detection  
23 (0.4–0.6 pmol/mg protein/minute).

24 In terms of direct pulmonary effects of TCE metabolites, Odum et al. (1992) reported that  
25 mice exposed to 100 ppm via inhalation of chloral for 6 hours resulted in bronchiolar lesions  
26 similar to those seen with TCE, although with a severity equivalent to 1,000 ppm TCE  
27 exposures. In addition, some alveolar necrosis, alveolar oedema, and desquamation of the  
28 epithelium were evident. In the same study, TCOH (100 and 500 ppm) also produced Clara cell  
29 damage, but with lower incidence than TCE, and without alveolar lesions, while TCA treatment  
30 produced no observable pulmonary effects. Therefore, it has been proposed that chloral is the  
31 active metabolite responsible for TCE pulmonary toxicity, and the localization of damage to  
32 Clara cells (rather than to other cell types, as seen with direct exposure to chloral) is due to the  
33 localization of oxidative metabolism in that cell type (Green, 2000; Green et al., 1997b; Odum et  
34 al., 1992). However, the recent identification by Forkert et al. (2006) of DAL adducts, also  
35 localized with Clara cell, suggests that TCE oxidation to dichloroacetyl chloride, which is not  
36 believed to be derived from chloral, may also contribute to adverse health effects.

1 Due to the histological similarities between TCE- and chloral-induced pulmonary  
2 toxicity, consistent with chloral being the active moiety, it has been proposed that the limited or  
3 absent capacity for reduction of chloral (rapidly converted to CH in the presence of water) to  
4 TCOH and glucuronidation of TCOH to TCOG in mouse lungs leads to “accumulation” of  
5 chloral in Clara cells. However, the lack of TCOH glucuronidation capacity of Clara cells  
6 reported by Odum et al. (1992), while possibly an important determinant of TCOH  
7 concentrations, should have no bearing on CH concentrations, which depend on the production  
8 and clearance of CH only. While isolated mouse Clara cells form smaller amounts of TCOH  
9 relative to CH (Odum et al., 1992), the cell-type distribution of the enzymes metabolizing CH is  
10 not clear. Indeed, cytosolic fractions of mouse, rat and human whole lungs show significant  
11 activity for CH conversion to TCOH (Green et al., 1997b). In particular, in mouse lung  
12 subcellular fractions, 1 micromole of TCE in a 1.3 mL reactival was converted to CH at a rate of  
13 1 nmol/minute/mg microsomal protein, while 10 nmol CH in a 1.3 mL reactival was converted  
14 to TCOH at a rate of 0.24 nmol/minute/mg cytosolic protein (Green et al., 1997b). How this  
15 fourfold difference in activity would translate in vivo is uncertain given the 100-fold difference  
16 in substrate concentrations, lack of information as to the concentration-dependence of activity,  
17 and uncertain differences between cytosolic and microsomal protein content in the lung. It is  
18 unclear whether local pulmonary metabolism of chloral is the primary clearance process in vivo,  
19 as in the presence of water, chloral rapidly converts to chloral hydrate, which is soluble in water  
20 and hence can rapidly diffuse to surrounding tissue and to the blood, which also has the capacity  
21 to metabolize chloral hydrate (Lipscomb et al., 1996). Nonetheless, experiments with isolated  
22 perfused lungs of rats and guinea pigs found rapid appearance of TCOH in blood following TCE  
23 inhalation exposure, with no detectable chloral hydrate or TCOG (Dalbey and Bingham, 1978).  
24 Therefore, it appears likely that chloral in the lung either is rapidly metabolized to TCOH, which  
25 then diffuses to blood, or diffuses to blood as CH and is rapidly metabolized to TCOH by  
26 erythrocytes (Lipscomb et al., 1996).

27 This hypothesis is further supported by in vivo data. No in vivo data in rats on CH after  
28 TCE administration were located, and Fisher et al. (1998) reported CH in blood of human  
29 volunteers exposed to TCE via inhalation were below detection limits. In mice, however, after  
30 both inhalation and oral gavage exposure to TCE, CH has been reported in whole lung tissue at  
31 concentrations similar to or somewhat greater than that in blood (Abbas and Fisher, 1997;  
32 Greenberg et al., 1999). A peak concentration (1.3 µg/g) of pulmonary CH was reported after  
33 inhalation exposure to 600 ppm—at or above exposures where Clara cell toxicity was reported in  
34 acute studies (Green et al., 1997b; Odum et al., 1992). However, this was fivefold *less* than the  
35 reported pulmonary CH concentration (6.65 µg/g) after gavage exposures of 1,200 mg/kg.  
36 Specifically, a 600-ppm exposure or 450-ppm exposure reported in the Maltoni et al. and

1 Fukuda et al. studies results in a greater incidence in lung tumors than the  
2 1,000–1,200 mg/kg-day exposures in the NTP (1990) and NCI (1976) bioassays. However, the  
3 peak CH levels measured in whole lung tissues after inhalation exposure to TCE at 600 ppm  
4 were reported to be about fivefold *lower* than that at 1,200 mg/kg by gavage, therefore, showing  
5 the *opposite* pattern (Abbas and Fisher, 1997; Greenberg et al., 1999). No studies of Clara cell  
6 toxicity after gavage exposures were located, but several studies in mice administered TCE via  
7 intraperitoneal injection did show Clara cell toxicity at around a dose of 750 mg/kg (Forkert et  
8 al., 2006) or above (e.g., Forkert and Birch, 1989; Forkert and Forkert, 1994). However, as  
9 noted previously, i.p. exposures are subject to an inflammatory response, confounding direct  
10 comparisons of dose via other routes of administration.

11 Although, whole lung CH concentrations may not precisely reflect the concentrations  
12 within specific cell types, as discussed above, the water solubility of CH suggests rapid  
13 equilibrium between cell types and between tissues and blood. Both Abbas and Fisher (1997)  
14 and Greenberg et al. (1999) were able to fit CH blood and lung levels using a PBPK model that  
15 did not include pulmonary metabolism, suggesting that lung CH levels may be derived largely by  
16 systemic delivery, i.e., from CH formed in the liver. However, a more detailed PBPK model-  
17 based analysis of this hypothesis has not been performed, as CH is not included in the PBPK  
18 model developed by Hack et al. (2006) that was updated in Section 3.5.

19 Two studies have reported formation of reactive metabolites in pulmonary tissues as  
20 assessed by macromolecular binding after TCE intraperitoneal administration. Forkert and Birch  
21 (1989) reported temporal correlations between the severity of Clara cell necrosis with increased  
22 levels of covalent binding macromolecules in the lung of TCE or metabolites with a single  
23 2,000 mg/kg dose of [<sup>14</sup>C]TCE. The amount of bound TCE or metabolites per gram of lung  
24 tissue, DNA, or protein peaked at 4 hours and decreased progressively at 8, 12, and 24 hours.  
25 The fraction of radioactivity in lung tissue macromolecules that was covalently bound reached a  
26 plateau of about 20% from 4–24 hours, suggesting that clearance of total and covalently bound  
27 TCE or metabolites was similar. The amount of covalent binding in the liver was three- to  
28 10-fold higher than in the lung, although hepatic cytotoxicity was not apparent. This tissue  
29 difference could either be due to greater localization of metabolism in the lung, so that  
30 concentrations reactive metabolites in individual Clara cells are greater than both the lung as a  
31 whole and hepatocytes, or because of greater sensitivity of Clara cells as compared to  
32 hepatocytes to reactive metabolites. More recently, Forkert et al. (2006) examined DAL adducts  
33 resulting from metabolism of TCE to dichloroacetyl chloride as an *in vivo* marker of production  
34 of reactive metabolites. Following intraperitoneal administration of 500–1,000 mg/kg TCE in  
35 CD-1 mice, they found localization of DAL adducts believed to be from oxidative metabolism  
36 within Clara cell apices, with dose-dependent increase in labeling with a polyclonal anti-DAL

1 antibody that correlated with increased Clara cell damage. Dose-dependent DAL adducts were  
2 also found in alveolar Type II cells, although no morphologic changes in those cells were  
3 observed. Both Clara cell damage (as discussed above) and DAL labeling were abolished in  
4 mice pretreated with DASO<sub>2</sub>, an inhibitor of CYP2E1 and CYP2F2. However, Clara cell  
5 damage in treated CYP2E1-null mice was more severe than in CD-1 mice. Although DAL  
6 labeling was less pronounced in CYP2E1-null mice as compared to CD-1 mice, this was due in  
7 part to the greater histopathologic damage leading to attenuation of the epithelium and loss of  
8 Clara cells in the null mice. In addition, protein immunoblotting with anti-DAL, anti-CYP2E1  
9 and anti-CYP2F2 antibodies suggested that a reactive TCE metabolite including dichloroacetyl  
10 chloride was formed that is capable of binding to CYP2E1 and CYP2F2 and changing their  
11 protein structures. Follow-up studies are needed in the lung and other target tissues to determine  
12 the potential role of the DAL adducts in TCE-induced toxicity.

13 Finally, although Green (2000) and others have attributed species differences in  
14 pulmonary toxicity to differences in the capacity for oxidative metabolism in the lung, it should  
15 be noted that the concentration of the active metabolite is determined by both its production and  
16 clearance (Clewel et al., 2000). Therefore, while the maximal pulmonary capacity to produce  
17 oxidative metabolites is clearly greater in the mouse than in rats or humans, there is little  
18 quantitative information as to species differences in clearance, whether by local chemical  
19 transformation/metabolism or by diffusion to blood and subsequent systemic clearance. In  
20 addition, existing in vitro data on pulmonary metabolism are at millimolar TCE concentrations  
21 where metabolism is likely to be approaching saturation, so the relative species differences at  
22 lower doses has not been characterized. Studies with recombinant CYP enzymes examined  
23 species differences in the catalytic efficiencies of CYP2E1, CYP2F, and CYP2B1, but the  
24 relative contributions of each isoform to pulmonary oxidation of TCE in vivo remains unknown  
25 (Forkert et al., 2005). Furthermore, systemic delivery of oxidative metabolites to the lung may  
26 contribute, as evidenced by respiratory toxicity reported with i.p. administration. Therefore,  
27 while the differences between mice and rats in metabolic capacity are correlated with their  
28 pulmonary sensitivity, it is not clear that differences in capacity alone are accurate quantitative  
29 predictors of toxic potency. Thus, while it is likely that the human lung is exposed to lower  
30 concentrations of oxidative metabolites, quantitative estimates for differential sensitivity made  
31 with currently available data and dosimetry models are highly uncertain.

32 In summary, it appears likely that pulmonary toxicity is dependent on in situ oxidative  
33 metabolism, however, the active agent has not been confidently identified. The similarities in  
34 histopathologic changes in Clara cells between TCE and chloral inhalation exposure, combined  
35 with the wider range of cell types affected by direct chloral administration relative to TCE, led  
36 some to hypothesize that chloral is the toxic moiety in both cases, but with that generated in situ

1 from TCE in Clara cells “accumulating” in those cells (Green, 2000). However, chemical and  
2 toxicokinetic data suggest that such “accumulation” is unlikely for several reasons. These  
3 include the rapid conversion of chloral to chloral hydrate in the presence of water, the water  
4 solubility of CH leading to rapid diffusion to other cell types and blood, the likely rapid  
5 metabolism of chloral hydrate to TCOH either in pulmonary tissue or in blood erythrocytes, and  
6 in vivo data showing lack of correlation across routes of exposure between whole-lung CH  
7 concentrations and pulmonary carcinogenicity and toxicity. However, additional possibilities for  
8 the active moiety exist, such as dichloroacetyl chloride, which is derived through a TCE  
9 oxidation pathway independent of chloral and which appears to result in adducts with lysine  
10 localized in Clara cells.  
11

#### 4.7.4. Mode of Action for Pulmonary Carcinogenicity

12 A number of effects have been hypothesized to be key events in the pulmonary  
13 carcinogenicity of TCE, including cytotoxicity leading to increased cell proliferation, formation  
14 of DAL protein adducts, and mutagenicity. As stated previously, the target cell for pulmonary  
15 adenocarcinoma formation has not been established. Much of the hazard and MOA information  
16 has focused on Clara cell effects from TCE which is a target in both susceptible and  
17 nonsusceptible rodent species for lung tumors. However, the role of Clara cell susceptibility to  
18 TCE-induced lung toxicity or to other potential targets such as lung stem cells that are activated  
19 to repopulate both Clara and Type II alveolar cells after injury, has not been determined for  
20 pulmonary carcinogenicity. While all of the events described above may be plausibly involved  
21 in the MOA for TCE pulmonary carcinogenicity, none have been directly shown to be necessary  
22 for carcinogenesis.  
23

##### 4.7.4.1.1. Mutagenicity via Oxidative Metabolism

24 The hypothesis is that TCE acts by a mutagenic MOA in TCE- induced lung tumors.  
25 According to this hypothesis, the key events leading to TCE-induced lung tumor formation  
26 constitute the following: the oxidative metabolism of TCE producing chloral/chloral hydrate  
27 delivered to pulmonary tissues, causes direct alterations to DNA (e.g., mutation, DNA damage,  
28 and/or micronuclei induction). Mutagenicity is a well-established cause of carcinogenicity.  
29

#### 4.7.4.1.2. Experimental support for the hypothesized mode of action

1 Pulmonary toxicity has been proposed to be dependent on in situ oxidative metabolism,  
2 however, the active agent has not been confidently identified. The similarities in histopathologic  
3 changes in Clara cells between TCE and chloral inhalation exposure, combined with the wider  
4 range of cell types affected by direct chloral administration relative to TCE, led some to  
5 hypothesize that chloral is the toxic moiety. Chloral that is formed from the metabolism of TCE  
6 is quickly converted to CH upon hydration under physiological conditions. As discussed in  
7 Section 4.2.4, CH clearly induces aneuploidy in multiple test systems, including bacterial and  
8 fungal assays in vitro (Crebelli et al., 1991; Kafer, 1986; Kappas, 1989), mammalian cells in  
9 vitro (Sbrana et al., 1993; Vagnarelli et al., 1990), and mammalian germ-line cells in vivo (Miller  
10 and Adler, 1992; Russo et al., 1984). Conflicting results were observed in in vitro and in vivo  
11 mammalian studies of micronuclei formation (Beland, 1999a; Degrassi and Tanzarella, 1988;  
12 Giller et al., 1995; Nessler and Marzin, 1999; Russo and Levis, 1992a; 1992b) with positive  
13 results in germ-line cells (Nutley et al., 1996)(Allen et al., 1994). In addition, it is mutagenic in  
14 the Ames bacterial mutation assay for some strains (Beland, 1999a; Giller et al., 1995; Haworth  
15 et al., 1983; Ni et al., 1994). Structurally related chlorinated aldehydes 2-chloroacetaldehyde  
16 and 2,2-dichloroacetaldehyde are both alkylating agents, are both positive in a genotoxic assay  
17 (Bignami et al., 1980), and both interact covalently with cellular macromolecules (Guengerich et  
18 al., 1979).

19 As discussed in the section describing the experimental support for the mutagenic MOA  
20 for liver carcinogenesis (see Section 4.5.7.1), it has been argued that CH mutagenicity is unlikely  
21 to be the cause of TCE carcinogenicity because the concentrations required to elicit these  
22 responses are several orders of magnitude higher than achieved in vivo (Moore and Harrington-  
23 Brock, 2000). Similar to the case of the liver, it is not clear how much of a correspondence is to  
24 be expected from concentrations in genotoxicity assays in vitro and concentrations in vivo, as  
25 reported in vivo CH concentrations are in whole lung homogenate while in vitro concentrations  
26 are in culture media. None of the available in vivo genotoxicity assays used the inhalation route  
27 that elicited the greatest lung tumor response under chronic exposure conditions, so direct in vivo  
28 comparisons are not possible. Finally, as discussed in Section 4.5.7.1, the use of i.p.  
29 administration in many other in vivo genotoxicity assays complicates the comparison with  
30 carcinogenicity data.

31 As discussed above (see Section 4.7.3), chemical and toxicokinetic data are not  
32 supportive of CH being the active agent of TCE-induced pulmonary toxicity, and directly  
33 contradict the hypothesis of chloral “accumulation.” Nonetheless, CH has been measured in the  
34 mouse lung following inhalation and gavage exposures to TCE (Abbas and Fisher, 1997;  
35 Greenberg et al., 1999), possibly the result of both in situ production and systemic delivery.

1 Therefore, in principle, CH could cause direct alterations in DNA in pulmonary tissue.  
2 However, as discussed above, the relative amounts of CH measured in whole lung tissue from  
3 inhalation and oral exposures do not appear to correlate with sensitivity to TCE lung tumor  
4 induction across exposure routes. While these data cannot rule out a role for mutagenicity  
5 mediated by CH due to various uncertainties, such as whether whole lung CH concentrations  
6 accurately reflect cell-type specific concentrations and possible confounding due to strain  
7 differences between inhalation and oral chronic bioassays, they do not provide support for this  
8 MOA.

9 Additional possibilities for the active moiety exist, such as dichloroacetyl chloride, which  
10 is derived through a TCE oxidation pathway independent of chloral and which appears to result  
11 in adducts with lysine localized in Clara cells (Forkert et al., 2006). DCA, which has some  
12 genotoxic activity, is, also, presumed to be formed through this pathway (see Section 3.3).  
13 Currently, however, there are insufficient data to support a role for these oxidative metabolites in  
14 a mutagenic MOA.  
15

#### 4.7.4.1.3. Cytotoxicity Leading to Increased Cell Proliferation

16 The hypothesis is that TCE acts by a cytotoxicity MOA in TCE-induced pulmonary  
17 carcinogenesis. According to this hypothesis, the key events leading to TCE-induced lung tumor  
18 formation constitute the following: TCE oxidative metabolism in situ leads to currently unknown  
19 reactive metabolites that cause cytotoxicity, leading to compensatory cellular proliferation and  
20 subsequently increased mutations and clonal expansion of initiated cells.  
21

#### 4.7.4.1.4. Experimental support for the hypothesized mode of action

22 Evidence for the hypothesized MOA consists primarily of (1) the demonstration of acute  
23 cytotoxicity and transient cell proliferation following TCE exposure in laboratory mouse studies;  
24 (2) toxicokinetic data supporting oxidative metabolism being necessary for TCE pulmonary  
25 toxicity; (3) the association of lower pulmonary oxidative metabolism and lower potency for  
26 TCE-induced cytotoxicity with the lack of observed pulmonary carcinogenicity in laboratory  
27 rats. However, there is a lack of experimental support linking TCE acute pulmonary cytotoxicity  
28 to sustained cellular proliferation of chronic exposures or clonal expansion of initiated cells.

29 As discussed above, a number of acute studies have shown that TCE is particularly  
30 cytotoxic to Clara cells in mice, which has been suggested to be involved in the development of  
31 mouse lung tumors (Buckpitt et al., 1995; Forkert and Forkert, 1994; Kim et al., 2005). In

1 addition, studies examining cell labeling by either BrdU (Green et al., 1997b) or 3H-thymidine  
2 incorporation (Villaschi et al., 1991) suggest increased cellular proliferation in mouse Clara cells  
3 following acute inhalation exposures to TCE. Moreover, in short-term studies, Clara cells appear  
4 to become resistant to cytotoxicity with repeated exposure, but regain their susceptibility after  
5 2 days without exposure. This observation led to the hypothesis that the 5 day/week inhalation  
6 dosing regime (Fukuda et al., 1983; Henschler et al., 1980; Maltoni et al., 1988; 1986) in the  
7 chronic mouse studies leads to periodic cytotoxicity in the mouse lung at the beginning of each  
8 week followed by cellular regeneration, and that the increased rate of cell division leads to  
9 increased incidence of tumors by increasing the overall mutation rate and by increasing the  
10 division rate of already initiated cells (Green, 2000). However, longer-term studies to test this  
11 hypothesis have not been carried out.

12 As discussed above (see Section 4.7.3), there is substantial evidence that pulmonary  
13 oxidative metabolism is necessary for TCE-induced pulmonary toxicity, although the active  
14 moiety remains unknown. In addition, the lower capacity for pulmonary oxidative metabolism  
15 in rats as compared to mice is consistent with studies in rats not reporting pulmonary cytotoxicity  
16 until exposures higher than those in the bioassays, and the lack of reported pulmonary  
17 carcinogenicity in rats at similar doses to mice. However, rats also have a lower background rate  
18 of lung tumors (Green, 2000), and so would be less sensitive to carcinogenic effects in that tissue  
19 to the extent that relative risks is the important metric across species. In addition, this MOA  
20 hypothesis requires a number of additional key assumptions for which there are currently no  
21 direct evidence. First, the cycle of cytotoxicity, repair, resistance to toxicity, and loss of  
22 resistance after exposure interruption, has not been documented and under the proposed MOA  
23 should continue under chronic exposure conditions. This cycle has thus, far only been observed  
24 in short term (up to 13-day) studies. In addition, although Clara cells have been identified as the  
25 target of toxicity whether they or endogenous stem cells in the lung are the cells responsible for  
26 mouse lung tumors has not been established. There is currently no data as to the cell type of  
27 origin for TCE-induced lung tumors.

28 This hypothesized MOA has been proposed for other compounds that induce mouse lung  
29 tumors, such as coumarin, naphthalene, and styrene (e.g., Cruzan et al., 2009). Among these,  
30 only for styrene have there been studies of chronic duration linking cytotoxicity with  
31 hyperplasia, and no studies appear to provide experimental linkage to clonal expansion of  
32 initiated cells.

33



#### **4.7.4.1.5. Additional Hypothesized Modes of Action with Limited Evidence or Inadequate Experimental Support**

#### **4.7.4.1.6. Role of formation of DAL protein adducts**

1           As discussed above, Forkert et al. (2006) recently observed dose-dependent formation of  
2 DAL protein adducts in the Clara cells of mice exposed to TCE via intraperitoneal injection.  
3 While adducts were highly localized in Clara cells, they were also found in alveolar Type II  
4 cells, though these cells did not show signs of cytotoxicity in this particular experimental  
5 paradigm. In terms of the MOA for TCE-induced pulmonary carcinogenicity, these adducts may  
6 either be causally important in and of themselves, or they may be markers of a different causal  
7 effect. For instance, it is possible that these adducts are a cause for the observed Clara cell  
8 toxicity, and Forkert et al. (2006) suggested that the lack of toxicity in alveolar Type II cells may  
9 indicate that “there may be a threshold in adduct formation and hence bioactivation at which  
10 toxicity is manifested.” In this case, they are an additional precursor event in the same causal  
11 pathway proposed above. Alternatively, these adducts may be indicative of effects related to  
12 carcinogenesis but unrelated to cytotoxicity. In this case, the Clara cell need not be the cell type  
13 of origin for mouse lung tumors.

14           Because of their recent discovery, there is little additional data supporting, refuting, or  
15 clarifying the potential role for DAL protein adducts in the MOA for TCE-induced pulmonary  
16 carcinogenesis. For instance, the presence and localization of such adducts in rats has not been  
17 investigated, and could indicate the extent to which the level of adduct formation is correlated  
18 with existing data on species differences in metabolism, cytotoxicity, and carcinogenicity. In  
19 addition, the formation of these adducts has only been investigated in a single dose study using  
20 i.p. injection. As stated above, i.p. injection may involve the initiation of a systemic  
21 inflammatory response that can activate lung macrophages or affect Clara cells. Experiments  
22 with repeated exposures over chronic durations and by inhalation or oral of administration would  
23 be highly informative. Finally, the biological effects of these adducts, whether cytotoxicity or  
24 something else, have not been investigated.

25

#### 4.7.4.1.7. Conclusions About the Hypothesized Modes of Action

#### 4.7.4.1.8. (1) Is the hypothesized mode of action sufficiently supported in the test animals?

##### 4.7.4.1.8.1. Mutagenicity

1 Chloral hydrate is clearly genotoxic, as there are substantial data from multiple in vitro  
2 and in vivo assays supporting its ability induce aneuploidy, with more limited data as to other  
3 genotoxic effects, such as point mutations. Chloral hydrate is also clearly present in pulmonary  
4 tissues of mice following TCE exposures similar to those inducing lung tumors in chronic  
5 bioassays. However, chemical and toxicokinetic data are not supportive of CH being the  
6 predominant metabolite for TCE carcinogenicity. Such data include the water solubility of CH  
7 leading to rapid diffusion to other cell types and blood, its likely rapid metabolism to TCOH  
8 either in pulmonary tissue or in blood erythrocytes, and in vivo data showing lack of correlation  
9 across routes of exposure between whole lung CH concentrations and pulmonary  
10 carcinogenicity. Therefore, while a role for mutagenicity via CH in the MOA of TCE-induced  
11 lung tumors cannot be ruled about, available evidence is inadequate to support the conclusion  
12 that direct alterations in DNA caused by CH produced in or delivered to the lung after TCE  
13 exposure constitute a MOA for TCE-induced lung tumors.  
14

##### 4.7.4.1.8.2. Cytotoxicity

15 The MOA hypothesis for TCE-induced lung tumors involving cytotoxicity is supported  
16 by relatively consistent and specific evidence for cytotoxicity at tumorigenic doses in mice.  
17 However, the majority of cytotoxicity-related key events have been investigated in studies less  
18 than 13 days, and none has been shown to be causally related to TCE-induced lung tumors. In  
19 addition, the cell type (or types) of origin for the observed lung tumors in mice has not been  
20 determined, so the contribution to carcinogenicity of Clara cell toxicity and subsequent  
21 regenerative cell division is not known. Similarly, the relative contribution from recently  
22 discovered dichloroacetyl-lysine protein adducts to the tumor response has not been investigated  
23 and has currently only been studied in i.p. exposure paradigms of short duration. In summary,  
24 while there are no data directly challenging the hypothesized MOA described above, the existing  
25 support for their playing a causal role in TCE-induced lung tumors is largely associative, and  
26 based on acute or short term studies. Therefore, there are inadequate data to support a cytotoxic  
27 MOA based on the TCE-induced cytotoxicity in Clara cells in the lungs of test animals.  
28

#### 4.7.4.1.8.3. **Additional hypothesis**

1 Inadequate data are available to develop a MOA hypothesis based on recently discovered  
2 DAL adducts induced by TCE inhalation and i.p. exposures. It will therefore, not be considered  
3 further in the conclusions below.

4 Overall, therefore, the MOA for TCE-induced lung tumors is considered unknown at this  
5 time.

6

#### 4.7.4.1.9. (2) **Is the hypothesized mode of action relevant to humans?**

##### 4.7.4.1.9.1. **Mutagenicity**

7 The evidence discussed above demonstrates that CH is mutagenic in microbial as well as  
8 test animal species. There is, therefore, the presumption that they would be mutagenic in  
9 humans. Therefore, this MOA is considered relevant to humans.

10

##### 4.7.4.1.9.2. **Cytotoxicity**

11 No data from human studies are available on the cytotoxicity of TCE and its metabolites  
12 in the lung, and no causal link between cytotoxicity and pulmonary carcinogenicity has been  
13 demonstrated in animal or human studies. Nonetheless, in terms of human relevance, no data  
14 suggest that the proposed key events are not biologically plausible in humans, therefore,  
15 qualitatively, TCE-induced lung tumors are considered relevant to humans. This conclusion that  
16 this hypothesized MOA is qualitatively relevant has also been reached for other compounds for  
17 which the MOA has been postulated (Cruzan et al., 2009). Information about the relative  
18 pharmacodynamic sensitivity between rodents and humans is absent, but information on  
19 pharmacokinetic differences in lung oxidative metabolism does exist and will be considered in  
20 dose-response assessment when extrapolating between species (see Section 5.2.1.2).

21

**4.7.4.1.10. (3) Which populations or lifestages can be particularly susceptible to the hypothesized mode of action?**

**4.7.4.1.10.1. Mutagenicity**

1           The mutagenic MOA is considered relevant to all populations and lifestages. According  
2 to EPA’s *Guidelines for Carcinogen Risk Assessment* (U.S. EPA, 2005c) and *Supplemental*  
3 *Guidance for Assessing Susceptibility From Early-Life Exposure to Carcinogens* (U.S. EPA,  
4 2005d), there may be increased susceptibility to early-life exposures for carcinogens with a  
5 mutagenic mode of action. However, because the weight of evidence is inadequate to support a  
6 mutagenic MOA for TCE pulmonary carcinogenicity, and in the absence of chemical-specific  
7 data to evaluate differences in susceptibility, the ADAFs should not be applied, in accordance  
8 with the *Supplemental Guidance*.  
9

**4.7.4.1.10.2. Cytotoxicity**

10           No information based is available as to which populations or lifestages may be  
11 particularly susceptible to TCE-induced lung tumors. However, pharmacokinetic differences in  
12 lung oxidative metabolism among humans do exist, and because of the association between lung  
13 oxidative metabolism and toxicity, will be considered in dose-response assessment when  
14 extrapolating within species.  
15

**4.7.5. Summary and Conclusions**

16           The studies described here show pulmonary toxicity found mainly in Clara cells in mice  
17 (Forkert and Birch, 1989; Forkert et al., 1985; Green et al., 1997b; Odum et al., 1992; Villaschi  
18 et al., 1991) and rats (Kurasawa, 1988). The most convincing albeit limited data regarding this  
19 type of toxicity was demonstrated predominantly in mice exposed via inhalation, although some  
20 toxicity was shown in intraperitoneal injection studies. Increased vacuolation of Clara cells was  
21 often seen within the first 24-hours-of-exposure, depending on dose, but with cellular repair  
22 occurring within days or weeks of exposure. Continued exposure led to resistance to  
23 TCE-induced Clara cell toxicity, but damage recurred if exposure was stopped after 5 days and  
24 then resumed after 2 days without exposure. However, Clara cell toxicity has only been  
25 observed in acute and short-term studies, and it is unclear whether they persist with subchronic  
26 or chronic exposure, particularly in mice, which are the more sensitive species. With respect to  
27 pulmonary carcinogenicity, statistically-significantly increased incidence of lung tumors from

1 chronic inhalation exposures to TCE was observed female ICR mice (Fukuda et al., 1983), male  
2 Swiss mice, and female B6C3F1 mice (Maltoni et al., 1986), though not in other sex/strain  
3 combinations, nor in rats (Henschler et al., 1980; Maltoni et al., 1986). However, lung toxicity  
4 and Clara cell effects have also been observed in rats. Overall, the limited carcinogenesis studies  
5 described above are consistent with TCE causing mild increases in pulmonary tumor incidence  
6 in mice, but not in other species tested such as rats and hamsters.

7 The epidemiologic studies are quite limited for examining the role of TCE in cancers of  
8 the respiratory system, with no studies found on TCE exposure specifically examining toxicity of  
9 the respiratory tract. The two studies found on organic solvent exposure which included TCE  
10 suggested smoking as a primary factor for observed lung function decreases among exposed  
11 workers. Animal studies have demonstrated toxicity in the respiratory tract, particularly damage  
12 to the Clara cells (nonciliated bronchial epithelial cells), as well as decreases in pulmonary  
13 surfactant following both inhalation and intraperitoneal exposures, especially in mice. Dose-  
14 related increases in vacuolation of Clara cells have been observed in mice and rats as early as  
15 24 hours postexposure (2006; Forkert and Birch, 1989; Forkert et al., 1985; Kurasawa, 1988;  
16 Odum et al., 1992; Scott et al., 1988). Mice appear to be more sensitive to these changes, but  
17 both species show a return to normal cellular morphology at four weeks postexposure (Odum et  
18 al., 1992). Studies in mice have also shown an adaptation or resistance to this damage after only  
19 4–5 days of repeated exposures (Green et al., 1997b; Odum et al., 1992). The limited  
20 epidemiological literature on lung and laryngeal cancer in TCE-exposed groups is inconclusive  
21 due to study limitations (low power, null associations, confidence intervals on relative risks that  
22 include 1.0). These studies can only rule out risks of a magnitude of 2.0 or greater for lung  
23 cancer and relative risks greater than 3.0 or 4.0 for laryngeal cancer for exposures to studied  
24 populations and thus, may not detect a level of response consistent with other endpoints. Animal  
25 studies demonstrated a statistically significant increase in pulmonary tumors in mice following  
26 chronic inhalation exposure to TCE (Fukuda et al., 1983; 1988; Maltoni et al., 1986). These  
27 results were not seen in other species tested (rats, hamsters; (Fukuda et al., 1983; Henschler et  
28 al., 1980; 1988; Maltoni et al., 1986). By gavage, elevated, but not statistically significant,  
29 incidences of benign and/or malignant pulmonary tumors have been reported in B6C3F1 mice  
30 (Henschler et al., 1984; NCI, 1976; NTP, 1990). No increased pulmonary tumor incidences have  
31 been reported in rats exposed to TCE by gavage (NCI, 1976; NTP, 1988, 1990), although all the  
32 studies suffered from early mortality in at least one sex of rat.

33 Although no epidemiologic studies on the role of metabolism of TCE in adverse  
34 pulmonary health effects have been published, animal studies have demonstrated the importance  
35 of the oxidative metabolism of TCE by CYP2E1 and/or CYP2F2 in pulmonary toxicity.  
36 Exposure to DASO<sub>2</sub>, an inhibitor of both enzymes protects against pulmonary toxicity in mice

1 following exposure to TCE (Forkert et al., 2005). The increased susceptibility in mice correlates  
2 with the greater capacity to oxidize TCE based on increased levels of CYP2E1 in mouse lungs  
3 relative to lungs of rats and humans (Forkert et al., 2006; Green et al., 1997b), but it is not clear  
4 that these differences in capacity alone are accurate quantitative predictors of sensitivity to  
5 toxicity. In addition, available evidence argues against the previously proposed hypothesis (e.g.,  
6 Green, 2000) that “accumulation” of chloral in Clara cells is responsible for pulmonary toxicity,  
7 since chloral is first converted the water-soluble compounds chloral hydrate and TCOH that can  
8 rapidly diffuse to surrounding tissue and blood. Furthermore, the observation of DAL protein  
9 adducts, likely derived dichloroacetyl chloride and not from chloral, that were localized in Clara  
10 cells suggests an alternative to chloral as the active moiety. While chloral hydrate has shown  
11 substantial genotoxic activity, chemical and toxicokinetic data on CH as well as the lack of  
12 correlation across routes of exposure between in vivo measurements of CH in lung tissues and  
13 reported pulmonary carcinogenicity suggest that evidence is inadequate to conclude that a  
14 mutagenic MOA mediated by CH is operative for TCE-induced lung tumors. Another MOA for  
15 TCE-induced lung tumors has been plausibly hypothesized to involve cytotoxicity leading to  
16 increased cell proliferation, but the available evidence is largely associative and based on short-  
17 term studies, so a determination of whether this MOA is operative cannot be made. The recently  
18 discovered formation of DAL protein adducts in pulmonary tissues may also play a role in the  
19 MOA of TCE-induced lung tumors, but an adequately defined hypothesis has yet to be  
20 developed. Therefore, the MOA for TCE-induced lung tumors is currently considered unknown,  
21 and this endpoint is thus, considered relevant to humans. Moreover, none of the available data  
22 suggest that any of the currently hypothesized mechanisms would be biologically precluded in  
23 humans.

24

## **4.8. REPRODUCTIVE AND DEVELOPMENTAL TOXICITY**

### **4.8.1. Reproductive Toxicity**

25 An assessment of the human and experimental animal data, taking into consideration the  
26 overall weight of the evidence, demonstrates a concordance of adverse reproductive outcomes  
27 associated with TCE exposures. Effects on male reproductive system integrity and function are  
28 particularly notable and are discussed below. Cancers of the reproductive system in both males  
29 and females have also been identified and are discussed below.

30

#### **4.8.1.1.1. Human Reproductive Outcome Data**

1           A number of human studies have been conducted that examined the effects of TCE on  
2 male and female reproduction following occupational and community exposures. These are  
3 described below and summarized in Table 4-85. Epidemiological studies of female human  
4 reproduction examined infertility and menstrual cycle disturbances related to TCE exposure.  
5 Other studies of exposure to pregnant women are discussed in the section on human  
6 developmental studies (see Section 4.8.2.1). Epidemiological studies of male human  
7

**Table 4-85. Human reproductive effects**

Subjects	Exposure	Effect	Reference
<b>Female and male combined effects</b>			
<i>Reproductive behavior</i>			
75 men and 71 women living near Rocky Mountain Arsenal, Colorado	Low: <5.0 ppb Medium: ≥5.0–<10.0 ppb High: <10.0 ppb Highest: <15 ppb	Altered libido <sup>a</sup> Low: referent Med: OR <sub>adj</sub> : 0.67 (95% CI: 0.18–2.49) High: OR <sub>adj</sub> : 1.65 (95% CI: 0.54–5.01) Highest: OR <sub>adj</sub> : 2.46 (95% CI: 0.59–10.28)	ATSDR (2001)
<b>Female effects</b>			
<i>Infertility</i>			
197 women occupationally exposed to solvents in Finland 1973–1983	U-TCA (μmol/L) <sup>b</sup> Median: 48.1 Mean: 96.2 ± 19.2	Reduced incidence of fecundability in the high exposure group <sup>c</sup> as measured by time to pregnancy Low: IDR = 1.21 (95%CI: 0.73–2.00) High: IDR = 0.61 (95%CI: 0.28–1.33)	Sallmén et al. (1995)
71 women living near Rocky Mountain Arsenal, Colorado	Low: <5.0 ppb Med: ≥5.0 to <10.0 ppb High: <10.0 ppb	No effect on lifetime infertility <sup>a</sup> Low: referent Med: OR <sub>adj</sub> : 0.45 (95% CI: 0.02–8.92) High: OR <sub>adj</sub> : 0.88 (95% CI: 0.13–6.22)	ATSDR (2001)
<i>Menstrual cycle disturbance</i>			
71 women living near Rocky Mountain Arsenal, Colorado	Low: <5.0 ppb Med: ≥5.0 to <10.0 ppb High: <10.0 ppb	Increase in abnormal menstrual cycle (defined as <26 d or >30 d) Low: referent Med: OR <sub>adj</sub> : 4.17 (95% CI: 0.31–56.65) High: OR <sub>adj</sub> : 2.39 (95% CI: 0.41–13.97)	ATSDR (2001)
184 women working in a factory assembling small electrical parts in Poland	Mean indoor air TCE: 200 mg/m <sup>3</sup>	18% reporting increase in amenorrhea in exposed group ( <i>n</i> = 140), compared to 2% increase in unexposed group ( <i>n</i> = 44)	Zielinski (1973)
32 women working in dry cleaning or metal degreasing in Czechoslovakia <sup>d</sup>	0.28–3.4 mg/L TCE for 0.5–25 yr	31% reporting increase in menstrual disturbances <sup>a</sup>	Bardodej and Vyskocil (1956)
20-yr-old woman was occupationally exposed to TCE via inhalation	Urine total trichloro-compounds 3.2 ng/mL (21–25 d after exposure)	Amenorrhea, followed by irregular menstruation and lack of ovulation	Sagawa et al. (1973)
<b>Male effects</b>			
<i>Reproductive behavior</i>			
43 men working in dry cleaning or metal degreasing in Czechoslovakia	0.28–3.4 mg/L TCE for 0.5–25 yr	30% reporting decreased potency <sup>a</sup>	Bardodej and Vyskocil (1956)
30 male workers in a money printing shop in Egypt	38–172 ppm TCE	Decreased libido reported in 10 men (33%), compared to 3 men in the control group (10%)	El Ghawabi et al. (1973)



**Table 4-85. Human reproductive effects (continued)**

Subjects	Exposure	Effect	Reference
42 yr-old male aircraft mechanic in UK	TCE exposure reported but not measured; exposure for 25 yr	Gynaecomastia, impotence	Saihan et al. (1978)
<i>Altered sperm quality</i>			
15 men working as metal degreasers in Denmark	TCE exposure reported but not measured	Nonsignificant increase in percentage of two YFF in spermatozoa; no effect on sperm count or morphology	Rasmussen et al. (1988)
85 men of Chinese descent working in an electronics factory	Mean personal air TCE: 29.6 ppm; Mean U-TCA: 22.4 mg/g creatinine	Decreased normal sperm morphology and hyperzoospermia	Chia et al. (1996)
<i>Altered endocrine function</i>			
85 men of Chinese descent working in an electronics factory	Mean personal air TCE: 29.6 ppm; Mean U-TCA: 22.4 mg/g creatinine	Increased DHEAS and decreased FSH, SHBG and testosterone levels; dose-response observed	Chia et al. (1997)
85 men of Chinese descent working in an electronics factory	Mean personal air TCE: 29.6 ppm; Mean U-TCA: 22.4 mg/g creatinine	Decreased serum levels of testosterone and SHBG were significantly correlated with years of exposure to TCE; increased insulin levels for exposure <2 yr	Goh et al. (1998)
<i>Infertility</i>			
282 men occupationally exposed to solvents in Finland 1973–1983	U-TCA (µmol/L): High exposure: <sup>c</sup> Mean: 45 (SD 42) Median 31 Low exposure: <sup>c</sup> Mean: 41 (SD 88) Median: 15	No effect on fecundability <sup>c</sup> (as measured by time to pregnancy) Low: FDR: 0.99 (95% CI: 0.63–1.56) Intermediate/High: FDR: <sup>c</sup> 1.03 (95% CI: 0.60–1.76)	Sallmén et al. (1998)
8 male mechanics seeking treatment for infertility in Canada	Urine (µmol/): TCA: <0.30–4.22 TCOH: <0.60–0.89 Seminal fluid (pg/extract): TCE: 20.4–5,419.0 Chloral: 61.2–1,739.0 TCOH 2.7–25.5 TCA: <100–5,504 DCA: <100–13,342	Infertility could not be associated with TCE as controls were 5 men also in treatment for infertility	Forkert et al. (2003)
75 men living near Rocky Mountain Arsenal, Colorado	Low: <5.0 ppb Med: ≥5.0 to <10.0 ppb High: <10.0 ppb	No effect on lifetime infertility (not defined) Low: referent Med: n/a High: OR <sub>adj</sub> : 0.83 (95% CI: 0.11–6.37)	ATSDR (2001)

1 <sup>a</sup>Not defined by the authors.

2 <sup>b</sup>As reported in Lindbohm et al. (1990).

3 <sup>c</sup>Low/intermediate exposure indicated use of TCE <1 or 1–4 d/wk, and biological measures indicated high exposure.

4 High exposure indicated daily use of TCE, or if biological measures indicated high exposure.

5 <sup>d</sup>Number inferred from data provided in Tables 2 and 3 in Bardodej and Vyskocil (1956).

6 UK = United Kingdom.

1 reproduction examined reproductive behavior, altered sperm morphology, altered endocrine  
2 function, and infertility related to TCE exposure.

3

#### 4.8.1.1.2. Female and male combined human reproductive effects

##### 4 *Reproductive behavior*

5 A residential study of individuals living near the Rocky Mountain Arsenal in Colorado  
6 examined the reproductive outcomes in 75 men and 71 women exposed to TCE in drinking water  
7 (ATSDR, 2001). TCE exposure was classified as high (>10.0 ppb), medium ( $\geq 5.0$  to <10.0 ppb),  
8 and low (<5.0 ppb). Altered libido for men and women combined was observed in a dose-  
9 response fashion, although the results were nonsignificant. The results were not stratified by  
10 gender.

11

#### 4.8.1.1.3. Female human reproductive effects

##### 4.8.1.1.3.1. Infertility

12 Sallmén et al. (1995) examined maternal occupational exposure to organic solvents and  
13 time-to-pregnancy. Cases of spontaneous abortion and controls from a prior study of maternal  
14 occupational exposure to organic solvents in Finland during 1973–1983 and pregnancy outcome  
15 (Lindbohm et al., 1990) were used to study time-to-pregnancy of 197 couples. Exposure was  
16 assessed by questionnaire during the first trimester and confirmed with employment records.  
17 Biological measurements of TCA in urine in 64 women who held the same job during pregnancy  
18 and measurement (time of measurement not stated) had a median value of 48.1  $\mu\text{mol/L}$  (mean:  
19  $96.2 \pm 19.2 \mu\text{mol/L}$ ) (Lindbohm et al., 1990). Nineteen women had low exposure to TCE (used  
20 <1 or 1–4 times/week), and 9 had high exposure to TCE (daily use). In this follow-up study, an  
21 additional questionnaire on time-to-pregnancy was answered by the mothers (Sallmén et al.,  
22 1995). The incidence density ratio (IDR) was used in this study to estimate the ratio of average  
23 incidence rate of pregnancies for exposed women compared to nonexposed women; therefore, a  
24 lower IDR indicates infertility. For TCE, a reduced incidence of fecundability was observed in  
25 the high exposure group (IDR: 0.61, 95% CI: 0.28–1.33) but not in the low exposure group  
26 (IDR: 1.21, 95% CI: 0.73–2.00). A similar study of paternal occupational exposure (Sallmén et  
27 al., 1998) is discussed in Section 4.2.1.2.

28 The residential study in Colorado discussed above did not observe an effect on lifetime  
29 infertility in the medium ( $\text{OR}_{\text{adj}}$ : 0.45; 95% CI: 0.02–8.92) or high exposure groups

1 (OR<sub>adj</sub>: 0.88; 95% CI: 0.13–6.22) (ATSDR, 2001). Curiously, exposed women had more  
2 pregnancies and live births than controls.

3

#### 4.8.1.1.3.2. Menstrual cycle disturbance

4 The ATSDR (2001) study discussed above also examined effects on the menstrual cycle  
5 (ATSDR, 2001). Nonsignificant associations without a dose-response were seen for abnormal  
6 menstrual cycle in women (OR<sub>adj</sub>: 2.23, 95% CI: 0.45–11.18).

7 Other studies have examined the effect of TCE exposure on the menstrual cycle. One  
8 study examined women working in a factory assembling small electrical parts (Zielinski, 1973,  
9 translated). The mean concentration of TCE in indoor air was reported to be 200 mg/m<sup>3</sup>. Of the  
10 140 exposed women, eighteen percent suffered from amenorrhea, compared to only 2% of the 44  
11 nonexposed workers. The other study examined 75 men and women working in dry cleaning or  
12 metal degreasing (Bardodej and Vyskocil, 1956). Exposures ranged from 0.28–3.4 mg/L, and  
13 length of exposure ranged from 0.5–25 years. This study reported that many women  
14 experienced menstrual cycle disturbances, with a trend for increasing air concentrations and  
15 increasing duration of exposure.

16 An additional case study of a 20-year-old woman was occupationally exposed to TCE via  
17 inhalation. The exposure was estimated to be as high as 10 mg/mL or several thousand ppm,  
18 based on urine samples 21–25 days after exposure of 3.2 ng/mL of total trichloro-compounds.  
19 The primary effect was neurological, although she also experienced amenorrhea, followed by  
20 irregular menstruation and lack of ovulation as measured by basal body temperature curves  
21 (Sagawa et al., 1973).

22

#### 4.8.1.1.4. Male human reproductive effects

##### 4.8.1.1.4.1. Reproductive behavior

23 One study reported on the effect of TCE exposure on the male reproductive behavior in  
24 75 men working in dry cleaning or metal degreasing (Bardodej and Vyskocil, 1956). Exposures  
25 ranged from 0.28–3.4 mg/L, and length of exposure ranged from 0.5–25 years. This study found  
26 that men experienced decreased potency or sexual disturbances; the authors speculated that the  
27 effects on men could be due to the CNS effects of TCE exposure. This study also measured  
28 serial neutral 17-ketosteroid determinations but they were found to be not statistically significant  
29 (Bardodej and Vyskocil, 1956).

1 An occupational study of 30 men working in a money printing shop were exposed to  
2 TCE for <1 year to 5 years (El Ghawabi et al., 1973). Depending on the job description, the  
3 exposures ranged from 38–172-ppm TCE. Ten (33%) men suffered from decreased libido,  
4 compared to three (10%) of unexposed controls. However, these results were not stratified by  
5 exposure level or duration. The authors speculate that decreased libido was likely due to the  
6 common symptoms of fatigue and sleepiness.

7 A case study described a 42 year-old man exposed to TCE who worked as an aircraft  
8 mechanic for approximately 25 years (Saihan et al., 1978). He suffered from a number of health  
9 complaints including gynaecomastia and impotence, along with neurotoxicity and  
10 immunotoxicity. In addition, he drank alcohol daily which could have increased his response to  
11 TCE.  
12

#### 4.8.1.1.4.2. Altered sperm quality

13 Genotoxic effects on male reproductive function were examined in a study evaluating  
14 occupational TCE exposure in 15 male metal degreasers (Rasmussen et al., 1988). No  
15 measurement of TCE exposure was reported. Sperm count, morphology, and spermatozoa  
16 Y-chromosomal nondisjunction during spermatogenesis were examined, along with  
17 chromosomal aberrations in cultured lymphocytes. A nonsignificant increase in percentage of  
18 two fluorescent Y-bodies (YFF) in spermatozoa were seen in the exposed group ( $p > 0.10$ ), and  
19 no difference was seen in sperm count or morphology compared to controls.

20 An occupational study of men using TCE for electronics degreasing (Chia et al., 1996,  
21 1997; Goh et al., 1998) examined subjects ( $n = 85$ ) who were offered a free medical exam if they  
22 had no prior history related to endocrine function, no clinical abnormalities, and normal liver  
23 function tests; no controls were used. These participants provided urine, blood, and sperm  
24 samples. The mean urine TCA level was 22.4 mg/g creatinine (range: 0.8–136.4 mg/g  
25 creatinine). In addition, 12 participants provided personal 8-hour air samples, which resulted in  
26 a mean TCE exposure of 29.6 ppm (range: 9–131 ppm). Sperm samples were divided into two  
27 exposure groups; low for urine TCE less than 25 mg/g creatinine, and high for urine TCA greater  
28 than or equal to 25 mg/g creatinine. A decreased percentage of normal sperm morphology was  
29 observed in the sperm samples in the high exposure group ( $n = 48$ ) compared to the low  
30 exposure group ( $n = 37$ ). However, TCE exposure had no effect on semen volume, sperm  
31 density, or motility. There was also an increased prevalence of hyperzoospermia (sperm density  
32 of >120 million sperm per mL ejaculate) with increasing urine TCA levels (Chia et al., 1996).  
33

#### 4.8.1.1.4.3. Altered endocrine function

1 Two studies followed up on the study by Chia et al. (1996) to examine endocrine function  
2 (Chia et al., 1997; Goh et al., 1998). The first examined serum testosterone, follicle-stimulating  
3 hormone (FSH), dehydroepiandrosterone sulphate (DHEAS), and sex-hormone binding globulin  
4 (SHBG) (Chia et al., 1997). With increased years of exposure to TCE, an increase in DHEAS  
5 levels were seen, from 255 ng/mL for <3 years to 717.8 ng/mL  $\geq$ 7 years exposure. Also with  
6 increased years of exposure to TCE, decreased FSH, SHBG and testosterone levels were seen.  
7 The authors speculated these effects could be due to decreased liver function related to TCE  
8 exposure (Chia et al., 1997).

9 The second follow-up study of this cohort studied the hormonal effects of chronic low-  
10 dose TCE exposure in these men (Goh et al., 1998). Because urine TCE measures only indicate  
11 short-term exposure, long-term exposure was indicated by years of exposure. Hormone levels  
12 examined include androstenedione, cortisol, testosterone, aldosterone, SHBG, and insulin.  
13 Results show that a decrease in serum levels of testosterone and SHBG were significantly  
14 correlated with years of exposure to TCE, and an increase in insulin levels were seen in those  
15 exposed for less than 2 years. Androstenedione, cortisol, and aldosterone were in normal ranges  
16 and did not change with years of exposure to TCE.

#### 4.8.1.1.4.4. Infertility

18 Sallmén et al. (1998) examined paternal occupational exposure and time-to-pregnancy  
19 among their wives. Cases of spontaneous abortion and controls from a prior study of pregnancy  
20 outcome (Taskinen et al., 1989) were used to study time-to-pregnancy of 282 couples. Exposure  
21 was determined by biological measurements of the father who held the same job during  
22 pregnancy and measurement (time of measurement not stated) and questionnaires answered by  
23 both the mother and father. An additional questionnaire on time-to-pregnancy was answered by  
24 the mother for this study six years after the original study (Sallmén et al., 1998). The level of  
25 exposure was determined by questionnaire and classified as “low/intermediate” if the chemical  
26 was used <1 or 1–4 days/week and biological measures indicated high exposure (defined as  
27 above the reference value for the general population), and “high” if used daily or if biological  
28 measures indicated high exposure. For 13 men highly exposed, mean levels of urine TCA were  
29 45  $\mu$ mol/L (SD 42  $\mu$ mol/L; median 31  $\mu$ mol/L); for 22 men low/intermediately exposed, mean  
30 levels of urine TCA were 41  $\mu$ mol/L (SD 88  $\mu$ mol/L; median 15  $\mu$ mol/L). The terminology IDR  
31 was replaced by fecundability density ratio (FDR) in order to reflect that pregnancy is a desired  
32 outcome; therefore, a high FDR indicates infertility. No effect was seen on fertility in the low

1 exposure group (FDR: 0.99, 95% CI: 0.63–1.56) or in the intermediate/high exposure group  
2 (FDR: 1.03, 95% CI: 0.60–1.76). However, the exposure categories were grouped by  
3 low/intermediate versus high, whereas the outcome categories were grouped by low versus  
4 intermediate/high, making a dose-response association difficult.

5 A small occupational study reported on eight male mechanics exposed to TCE for at least  
6 two years who sought medical treatment for infertility (Forkert et al., 2003). The wives were  
7 determined to have normal fertility. Samples of urine from two of the eight male mechanics  
8 contained TCA and/or TCOH, demonstrating the rapid metabolism in the body. However,  
9 samples of seminal fluid taken from all eight individuals detected TCE and the metabolites  
10 chloral hydrate and TCOH, with two samples detecting DCA and one sample detecting TCA.  
11 Five unexposed controls also diagnosed with infertility did not have any TCE or metabolites in  
12 samples of seminal fluid. There was no control group that did not experience infertility.  
13 Increased levels of TCE and its metabolites in the seminal fluid of exposed workers compared to  
14 lower levels found in their urine samples was explained by cumulative exposure and  
15 mobilization of TCE from adipose tissue, particularly that surrounding the epididymis. In  
16 addition, CYP2E1 was detected in the epididymis, demonstrating that metabolism of TCE can  
17 occur in the male reproductive tract. However, this study could not directly link TCE to the  
18 infertility, as both the exposed and control populations were selected due to their infertility.

19 The ATSDR (2001) study discussed above on the reproductive effects from TCE in  
20 drinking water of individuals living near the Rocky Mountain Arsenal in Colorado did not  
21 observe infertility or other adverse reproductive effects for the high exposure group compared to  
22 the low exposure group ( $OR_{adj}$ : 0.83; 95% CI: 0.11–6.37). Curiously, exposed men had more  
23 pregnancies and live births than controls.

#### 24 **4.8.1.1.5. Summary of human reproductive toxicity**

25 Following exposure to TCE, adverse effects on the female reproductive system observed  
26 include reduced incidence of fecundability (as measured by time-to-pregnancy) and menstrual  
27 cycle disturbances. Adverse effects on the male reproductive system observed include altered  
28 sperm morphology, hyperzoospermia, altered endocrine function, decreased sexual drive and  
29 function, and altered fertility. These are summarized in Table 4-85.

#### **4.8.1.1.6. Animal Reproductive Toxicity Studies**

1           A number of animal studies have been conducted that examined the effects of TCE on  
2 reproductive organs and function following either inhalation or oral exposures. These are  
3 described below and summarized in Tables 4-86 and 4-87. Other animal studies of offspring  
4 exposed during fetal development are discussed in the section on animal developmental studies  
5 (see Section 4.8.2.2).  
6

#### **4.8.1.1.7. Inhalation exposures**

7           Studies in rodents exposed to TCE via inhalation are described below and summarized in  
8 Table 4-86. These studies focused on various aspects of male reproductive organ integrity,  
9 spermatogenesis, or sperm function in rats or mice. In the studies published after the year 2000,  
10 the effects of either 376 or 1,000-ppm TCE were studied following exposure durations ranging  
11 from 1–24 weeks, and adverse effects on male reproductive endpoints were observed.  
12

1  
2  
3

**Table 4-86. Summary of mammalian in vivo reproductive toxicity studies—  
inhalation exposures**

Reference	Species/strain/ sex/number	Exposure level/duration	NOAEL; LOAEL <sup>a</sup>	Effects
Forkert et al. (2002)	Mouse, CD-1, male, 6/group	0 or 1,000 ppm (5,374 mg/m <sup>3</sup> ) <sup>b</sup>  6 h/d, 5 d/wk, 19 d over 4 wk	LOAEL: 1,000 ppm	U-TCA and U-TCOH increased by 2 <sup>nd</sup> and 3 <sup>rd</sup> wk, respectively. Cytochrome P450 2E1 and <i>p</i> -nitrophenol hydroxylation in epididymal epithelium > testicular Leydig cells. Choral also generated from TCE in epididymis > testis. Sloughing of epididymal epithelial cells after 4 wk exposure.
Kan et al. (2007)	Mouse, CD-1, male, 4/group	0 or 1,000 ppm  6 h/d, 5 d/wk, 1–4 wk	LOAEL: 1,000 ppm	Light microscopy findings: degeneration and sloughing of epididymal epithelial cells as early as 1 wk into exposure; more severe by 4 wk. Ultrastructural findings: vesiculation in cytoplasm, disintegration of basolateral cell membranes, sloughing of epithelial cells. Sperm found in situ in cytoplasm of degenerated epididymal cells. Abnormalities of the head and tail in sperm located in the epididymal lumen.
Kumar et al. (2000b)	Rat, Wistar, male, 12–13/group	0 or 376 ppm  4 h/d, 5 d/wk, 2–10 wk exposure, 2–8 wk rest period	LOAEL: 376 ppm	Alterations in testes histopathology (smaller, necrotic spermatogenic tubules), ↑ sperm abnormalities, and sig. ↑ pre- and/or postimplantation loss in litters observed in the groups with 2 or 10 wk of exposure, or 5 wk of exposure with 2 wk rest.
Kumar et al. (2000a)	Rat, Wistar, males, 12–13/group	0 or 376 ppm  4 h/d, 5 d/wk, 12 and 24 wk	LOAEL: 376 ppm	Sig. ↓ in total epididymal sperm count and sperm motility, with sig. ↓ in serum testosterone, sig. ↑ in testes cholesterol, sig. ↓ of glucose 6-phosphate dehydrogenase and 17-β-hydroxy steroid dehydrogenase at 12 and 24 wk exposure.
Kumar et al. (2000a)	Rat, Wistar, male, 6/group	0 or 376 ppm  4 h/d, 5 d/wk, 12 and 24 wk	LOAEL: 376 ppm	BW gain sig. ↓. Testis weight, sperm count and motility sig. ↓, effect stronger with exposure time. After 12 wk, numbers of spermatogenic cells and spermatids ↓, some of the spermatogenic cells appeared necrotic. After 24 wk testes were atrophied, tubules were smaller, had Sertoli cells and were almost devoid of spermatocytes and spermatids. Leydig cells were hyperplastic. SDH, G6PDH sig. ↓, GGT and β-glucuronidase sig. ↑; effects stronger with exposure time.

4



**Table 4-86. Summary of mammalian in vivo reproductive toxicity studies—  
inhalation exposures (continued)**

Reference	Species/strain/ sex/number	Exposure level/duration	NOAEL; LOAEL <sup>a</sup>	Effects
<b>Land et al. (1981)</b>	<b>Mouse, C57BlxC3H (F1), male, 5 or 10/group</b>	<b>0, 0.02%, or 0.2%  4 h/d, 5 d, 23 d rest</b>	<b>NOAEL: 0.02% LOAEL: 0.2%</b>	<b>Sig. ↑ percentage morphologically abnormal epididymal sperm.</b>
<b>Xu et al. (2004)</b>	<b>Mouse, CD-1, male, 4–27/group</b>	<b>0 or 1,000 ppm (5.37 mg/L)<sup>b</sup>  6 h/d, 5 d/wk, 1–6 wk</b>	<b>LOAEL: 1,000 ppm</b>	<b>Sig. ↓ in vitro sperm-oocyte binding and in vivo fertilization</b>

**Bolded studies** carried forward for consideration in dose-response assessment (see Section 5).

1  
2  
3  
4  
5  
6

<sup>a</sup>NOAEL and LOAEL are based upon reported study findings.

<sup>b</sup>Dose conversion calculations by study author(s).

G6PDH = glucose 6-p dehydrogenase.

1  
2  
3

**Table 4-87. Summary of mammalian in vivo reproductive toxicity studies—oral exposures**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
<b>Studies assessing male reproductive outcomes</b>					
DuTeaux et al. (2003)	Rat, Sprague-Dawley, male, 3/group	0, 0.2, or 0.4% (0, 143, or 270 mg/kg-day)	Drinking water; 3% ethoxylated castor oil vehicle	LOEL: 0.2%	TCE metabolite-protein adducts formed by a cytochrome P450-mediated pathway were detected by fluorescence immunohistochemistry in the epithelia of corpus epididymis and in efferent ducts.
<b>DuTeaux et al. (2004a)</b>	<b>Rat, Sprague-Dawley, male, 3/group, or Simonson albino (UC-Davis), male, 3/group</b>	<b>0, 0.2, or 0.4% (0, 143, or 270 mg/kg-day)</b> <b>14 d</b>	<b>Drinking water, 3% ethoxylated castor oil vehicle</b>	<b>LOAEL: 0.2%</b>	<b>Dose-dependent ↓ in ability of sperm to fertilize oocytes collected from untreated ♀s. Oxidative damage to sperm membrane in head and mid-piece was indicated by dose-related ↑ in oxidized proteins and lipid peroxidation.</b>
Veeramachani et al. (2001)	Rabbit, Dutch belted, females and offspring; 7–9 offspring/group	9.5- or 28.5-ppm TCE <sup>b</sup> GD 20 thru lactation, then to offspring thru postnatal wk 15	Drinking water	LOAEL: 9.5 ppm	Decreased copulatory behavior; acrosomal dysgenesis, nuclear malformations; sig. ↓ LH and testosterone.
Zenick et al. (1984)	Rat, Long-Evans, male, 10/group	0, 10, 100, or 1,000 mg/kg-day 6 wk, 5 d/wk; 4 wk recovery	Gavage, corn oil vehicle	NOAEL: 100 mg/kg-day LOAEL: 1,000 mg/kg-day	At 1,000 mg/kg, BW ↓, liver/BW ratios ↑, and impaired copulatory behavior. Copulatory performance returned to normal by 5 <sup>th</sup> wk of exposure. At wk 6, TCE and metabolites concentrated to a significant extent in male reproductive organs.
<b>Studies assessing female reproductive outcomes</b>					
Berger and Horner (2003)	Rat, Simonson (S-D derived), female, (5–6) × 3/group	0 or 0.45% 2 wk	Drinking water, 3% Tween vehicle	LOAEL: 0.45%	In vitro fertilization and sperm penetration of oocytes sig. ↓ with sperm harvested from untreated males.

4

**Table 4-87. Summary of mammalian in vivo reproductive toxicity studies—oral exposures (continued)**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
Cosby and Dukelow (1992)	Mouse, B6D2F1, female, 7–12/group	0, 24, or 240 mg/kg-day GD 1–5, 6–10, or 11–15	Gavage, corn oil vehicle	NOAEL: 240 mg/kg-day	No treatment-related effects on in vitro fertilization in dams or offspring.
Manson et al. (1984)	Rat, Long-Evans, female, 23–25/group	0, 10, 100, or 1,000 mg/kg-day 6 wk: 2 wk pre mating, 1 wk mating period, GD 1–21	Gavage, corn oil vehicle	NOAEL: 100 mg/kg-day LOAEL: 1,000 mg/kg-day	Female fertility and mating success was not affected. At 1,000 mg/kg-day group, 5/23 females died, gestation BW gain was sig. ↓. After subchronic oral TCE exposure, TCE was detected in fat, adrenals, and ovaries; TCA levels in uterine tissue were high.  At 1,000 mg/kg-day, neonatal deaths (female pups) were ↑ on PNDs 1, 10, and 14. Dose-related ↑ seen in TCA in blood, liver and milk in stomach of ♀ pups, not ♂s.
Wu and Berger (2007)	Rat, Simonson (S-D derived), female, (no./group not reported)	0 or 0.45% (0.66 g/kg-d) <sup>c</sup> Preovulation days 1–5, 6–10, 11–14, or 1–14	Drinking water, 3% Tween vehicle	LOAEL: 0.45%	In vitro fertilization and sperm penetration of oocytes sig. ↓ with sperm harvested from untreated males.
Wu and Berger (2008)	Rat, Simonson (S-D derived), female, (no./group not reported)	0 or 0.45% (0.66 g/kg-d) <sup>c</sup> 1 or 5 d	Drinking water, 3% Tween vehicle	NOEL: 0.45%	Ovarian mRNA expression for ALCAM and Cud21 protein were not altered.

**Table 4-87. Summary of mammalian in vivo reproductive toxicity studies—oral exposures (continued)**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
<b>Studies assessing fertility and reproductive outcome in both sexes</b>					
George et al. (1985)	Mouse, CD-1, male and female, 20 pairs/treatment group; 40 controls/sex	0, 0.15, 0.30, or 0.60% <sup>d</sup> micro-encapsulated TCE  (TWA dose estimates: 0, 173, 362, or 737 mg/kg-day) <sup>c</sup>  Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females throughout gestation (i.e., 18 wk total)	Dietary	Parental systemic toxicity: NOAEL: 0.30% LOAEL: 0.60%	At 0.60%, in F0: sig. ↑ liver weights in both sexes; sig. ↓ testis and seminal vesicle weight; histopathology of liver and kidney in both sexes.  At 0.60%, in F1: sig. ↓ BW on PND 74, and in postpartum F1 dams; sig. ↑ liver, testis, and epididymis weights in males, sig. ↑ kidney weights in both sexes; sig. ↓ testis and seminal vesicle weight; histopathology of liver and kidney in both sexes.
				Parental reproductive function: LOAEL: 0.60% c	At 0.60%, in F0 and F1 males: sig. ↓ sperm motility.
				Offspring toxicity: NOAEL: 0.30% LOAEL: 0.60%	At 0.60%, in F1 pups: sig. ↓ live birth weights, sig. ↓ PND 4 pup BW; perinatal mortality ↑ (PND 0–21).
George et al. (1986)	Rat, F334, males and female, 20 pairs/treatment group, 40 controls/sex	0, 0.15, 0.30 or 0.60% <sup>d</sup> micro-encapsulated TCE  Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females throughout gestation (i.e., 18 wk total)	Dietary	Parental systemic toxicity: LOAEL: 0.15%	At 0.60%, in F0: sig. ↓ postpartum dam BW; sig. ↓ term. BW in both sexes; sig. ↑ liver, and kidney/adrenal weights in both sexes; sig. ↑ testis/epididymis weights; in F1: sig. ↓ testis weight.  At all doses in F1: sig. ↓ postpartum dam BW; sig. ↓ term. BW in both sexes, sig. ↑ liver weight in both sexes.  At 0.30 and 0.60%, in F1: sig. ↑ liver weight in females.

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-87. Summary of mammalian in vivo reproductive toxicity studies—oral exposures (continued)**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
George et al. (1986) (continued)				Parental reproductive function: LOAEL: 0.60% <sup>d</sup>	At 0.60%, sig ↓ mating in F0 males and females (in cross-over mating trials).
				Offspring toxicity: LOAEL: 0.15%	At 0.60%, sig. ↓ F1 BW on PND 4 and 14. At all doses, sig. ↓ F1 BW on PND 21 and 80.  At 0.3 and 0.60%, sig. ↓ live F1 pups/litter. Sig. trend towards ↓ live litters per pair At 0.15 and 0.60%, trend toward ↓ F1 survival from PND 21–80.

**Bolded studies** carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup>NOAEL, LOAEL, NOEL, and LOEL (lowest-observed-effect level) are based upon reported study findings.

<sup>b</sup>Concurrent exposure to several ground water contaminants; values given are for TCE levels in the mixture.

<sup>c</sup>Dose conversion calculations by study author(s).

<sup>d</sup>Fertility and reproduction assessment of last litter from continuous breeding phase and cross-over mating assessment (rats only) were conducted for 0 or 0.60% dose groups only.

LH = luteinizing hormone.

Kumar et al. (2000b) exposed male Wistar rats in whole-body inhalation chambers to 376-ppm TCE for 4 hours/day, 5 days/week over several duration scenarios. These were 2-weeks (to observe the effect on the epididymal sperm maturation phase), 10 weeks (to observe the effect on the entire spermatogenic cycle), 5 weeks with 2 weeks rest (to observe the effect on primary spermatocytes differentiation to sperm), 8 weeks with 5 weeks rest (to observe effects on an intermediate stage of spermatogenesis), and 10 weeks with 8 weeks rest (to observe the effect on spermatogonial differentiation to sperm). Control rats were exposed to ambient air. Weekly mating with untreated females was conducted. At the end of the treatment/rest periods, the animals were sacrificed; testes and cauda epididymes tissues were collected. Alterations in testes histopathology (smaller, necrotic spermatogenic tubules), increased sperm abnormalities, and significantly increased pre- and/or postimplantation loss in litters were observed in the

1 groups with 2 or 10 weeks of exposure, or 5 weeks of exposure with 2 weeks rest. It was  
2 hypothesized that postmeiotic cells of spermatogenesis and epididymal sperm were affected by  
3 TCE exposure, leading to reproductive impairment.

4 To test the hypothesis that TCE exposure adversely affects sperm function and  
5 fertilization, Xu et al. (2004) conducted a study in which male CD-1 mice were exposed by  
6 inhalation to atmospheres containing 1,000 ppm (5.37 mg/L) TCE for 1–6 weeks (6 hours/day,  
7 5 days/week). After each TCE exposure, body weights were recorded. Following termination,  
8 the right testis and epididymis of each treated male were weighed, and sperm was collected from  
9 the left epididymis and vas deferens for assessment of the number of total sperm and motile  
10 sperm. Sperm function was evaluated in the following experiments: (1) suspensions of  
11 capacitated vas deferens/cauda epididymal sperm were examined for spontaneous acrosome  
12 reaction, (2) in vitro binding of capacitated sperm to mature eggs from female CF-1 mice  
13 (expressed as the number of sperm bound per egg) was assessed, and (3) in vivo fertilization was  
14 evaluated via mating of male mice to superovulated female CF-1 mice immediately following  
15 inhalation exposure; cumulus masses containing mature eggs were collected from the oviducts of  
16 the females, and the percentage of eggs fertilized was examined. Inhalation exposure to TCE did  
17 not result in altered body weight, testis and epididymis weights, sperm count, or sperm  
18 morphology or motility. Percentages of acrosome-intact sperm populations were similar  
19 between treated and control animals. Nevertheless, for males treated with TCE for 2 or more  
20 weeks decreases were observed in the number of sperm bound to the oocytes in vitro (significant  
21 at 2 and 6 weeks,  $p < 0.001$ ). In a follow-up assessment, control sperm were incubated for  
22 30-minutes in buffered solutions of TCE or metabolites (chloral hydrate or trichloroethanol);  
23 while TCE-incubation had no effect on sperm-oocyte binding, decreased binding capacity was  
24 noted for the metabolite-incubated sperm. The ability for sperm from TCE-exposed males to  
25 bind to and fertilize oocytes in vivo was also found to be significantly impaired ( $p < 0.05$ ).

26 A study designed to investigate the role of testosterone, and of cholesterol and ascorbic  
27 acid (which are primary precursors of testosterone) in TCE-exposed rats with compromised  
28 reproductive function was conducted by Kumar et al. (2000a). Male Wistar rats (12–13/group)  
29 were exposed (whole body) to 376 ppm TCE by inhalation for 4 hours/day, 5 days/week, for  
30 either 12 or 24 weeks and then terminated. Separate ambient-air control groups were conducted  
31 for the 12- and 24-week exposure studies. Epididymal sperm count and motility were evaluated,  
32 and measures of 17- $\beta$ -hydroxy steroid dehydrogenase (17- $\beta$ -HSD), testicular total cholesterol  
33 and ascorbic acid, serum testosterone, and glucose 6-p dehydrogenase (G6PDH) in testicular  
34 homogenate were assayed. In rats exposed to TCE for either 12 or 24 weeks, total epididymal  
35 sperm count and motility, serum testosterone concentration, and specific activities of both

1 17-β-HSD and G6PDH were significantly decreased ( $p < 0.05$ ), while total cholesterol content  
2 was significantly ( $p < 0.05$ ) increased. Ascorbic acid levels were not affected.

3 In another study, Kumar et al. (2001b) utilized the same exposure paradigm to examine  
4 cauda epididymal sperm count and motility, testicular histopathology, and testicular marker  
5 enzymes: sorbitol dehydrogenase (SDH), G6PDH, glutamyl transferase (GT), and glucuronidase,  
6 in Wistar rats (6/group). After 24 weeks of exposure, testes weights and epididymal sperm count  
7 and motility were significantly decreased ( $p < 0.05$ ). After 12 weeks of TCE exposure,  
8 histopathological examination of the testes revealed a reduced number of spermatogenic cells in  
9 the seminiferous tubules, fewer spermatids as compared to controls, and the presence of necrotic  
10 spermatogenic cells. Testicular atrophy, smaller tubules, hyperplastic Leydig cells, and a lack of  
11 spermatocytes and spermatids in the tubules were observed after 24 weeks of TCE exposure.  
12 After both 12 and 24 weeks of exposure, SDH and G6PDH were significantly ( $p < 0.05$ ) reduced  
13 while GT and β-glucuronidase were significantly ( $p < 0.05$ ) increased.

14 In a study by Land et al. (1981), 8–10 week old male mice (C57BlxC3H)F1 (5 or  
15 10/group) were exposed (whole body) by inhalation to a number of anesthetic agents for  
16 5 consecutive days at 4 hours/day and sacrificed 28 days after the first day of exposure.  
17 Chamber concentration levels for the TCE groups were 0.02 and 0.2%. The control group  
18 received ambient air. Epididymal sperm were evaluated for morphological abnormalities. At  
19 0.2% TCE, the percentage of abnormal sperm in a sample of 1,000 was significantly ( $p < 0.01$ )  
20 increased as compared to control mice; no treatment-related effect on sperm morphology was  
21 observed at 0.02% TCE.

22 Forkert et al. (2002) exposed male CD-1 mice by inhalation to 1,000-ppm TCE  
23 (6 hours/day, 5 day/week) for 4 consecutive weeks and observed sloughing of portions of the  
24 epithelium upon histopathological evaluation of testicular and epididymal tissues.

25 Kan et al. (2007) also demonstrated that damage to the epididymal epithelium and sperm  
26 of CD-1 mice (4/group) resulted from exposure to 0 or 1,000-ppm TCE by inhalation for  
27 6 hours/day, 5 days/week, for 1–4 weeks. Segments of the epididymis (caput, corpus, and  
28 cauda) were examined by light and electron microscope. As early as 1 week after TCE exposure,  
29 degeneration and sloughing of epithelial cells from all three epididymal areas were observed by  
30 light microscopy; these findings became more pronounced by 4 weeks of exposure. Vesiculation  
31 in the cytoplasm, disintegration of basolateral cell membranes, and epithelial cell sloughing were  
32 observed with electron microscopy. Sperm were found in situ in the cytoplasm of degenerated  
33 epididymal cells. A large number of sperm in the lumen of the epididymis were abnormal,  
34 including head and tail abnormalities.

35

#### 4.8.1.1.8. Oral exposures

1 A variety of studies were conducted to assess various aspects of male and/or female  
2 reproductive capacity in laboratory animal species following oral exposures to TCE. These are  
3 described below and summarized in Table 4-87. They include studies that focused on male  
4 reproductive outcomes in rats or rabbits following gavage or drinking water exposures (DuTeaux  
5 et al., 2004a; DuTeaux et al., 2003; Veeramachaneni et al., 2001; Zenick et al., 1984), studies  
6 that focused on female reproductive outcomes in rats following gavage or drinking water  
7 exposures (Berger and Horner, 2003; Cosby and Dukelow, 1992; Manson et al., 1984; Wu and  
8 Berger, 2007, 2008), and studies assessed fertility and reproductive outcome in both sexes  
9 following dietary exposures to CD-1 mice or F344 rats (George et al., 1985, 1986).

##### 4.8.1.1.8.1. Studies assessing male reproductive outcomes

11 Zenick et al. (1984) conducted a study in which sexually experienced Long-Evans  
12 hooded male rats were administered 0, 10, 100, or 1,000 mg/kg-day TCE by gavage in corn oil  
13 for 6 weeks. A 4-week recovery phase was also incorporated into the study design. Endpoints  
14 assessed on Weeks 1 and 5 of treatment included copulatory behavior, ejaculatory plug weights,  
15 and ejaculated or epididymal sperm measures (count, motility, and morphology). Sperm  
16 measures and plug weights were not affected by treatment, nor were Week 6 plasma testosterone  
17 levels found to be altered. TCE effects on copulatory behavior (ejaculation latency, number of  
18 mounts, and number of intromissions) were observed at 1,000 mg/kg-day; these effects were  
19 recovered by 1–4 weeks posttreatment. Although the effects on male sexual behavior in this  
20 study were believed to be unrelated to narcotic effects of TCE, a later study by Nelson and  
21 Zenick (1986) showed that naltrexone (an opioid receptor antagonist, 2.0 mg/kg, i.p.,  
22 administered 15 minutes prior to testing) could block the effect. Thus, it was hypothesized that  
23 the adverse effects of TCE on male copulatory behavior in the rat at 1,000 ppm may in fact be  
24 mediated by the endogenous opioid system at the CNS level.

25 In a series of experiments by DuTeaux et al. (2004a; 2003), adult male rats were  
26 administered 0, 0.2, or 0.4% TCE (v/v) (equivalent to 0, 2.73 mg/L, or 5.46 mg/L) in a solution  
27 of 3% ethoxylated castor oil in drinking water for 14 days. These concentrations were within the  
28 range of measurements obtained in formerly contaminated drinking water wells, as reported by  
29 ATSDR (1997c). The average ingested doses of TCE (based upon animal body weight and  
30 average daily water consumption of 28 mL) were calculated to be 143 or 270 mg/kg-day for the  
31 low and high-dose groups, respectively (DuTeaux et al., (2003). Cauda epididymal and vas  
32 deferens sperm from treated males were incubated in culture medium with oviductal cumulus



1 masses from untreated females to assess in vitro fertilization capability. Treatment with TCE  
2 resulted in a dose-dependent decrease in the ability of sperm to fertilize oocytes. Terminal body  
3 weights and testis/epididymal weights were similar between control and treated groups.  
4 Evaluation of sperm concentration or motility parameters did not reveal any treatment-related  
5 alterations; acrosomal stability and mitochondrial membrane potential were not affected by  
6 treatment. Although no histopathological changes were observed in the testis or in the caput,  
7 corpus, or cauda epididymis, exposure to 0.2 and 0.4% TCE resulted in slight cellular alterations  
8 in the efferent ductule epithelium.

9         Veeramachaneni et al. (2001) evaluated the effects of drinking water containing  
10 chemicals typical of ground water near hazardous waste sites (including 9.5- or 28.5-ppm TCE)  
11 on male reproduction. In this study, pregnant Dutch-belted rabbits were administered treated  
12 drinking water from GD 20; treatment continued through the lactation period and to weaned  
13 offspring (7–9/group) through postnatal Week 15. Deionized water was administered from  
14 postnatal weeks 16–61, at which time the animals were terminated. At 57–61 weeks of age,  
15 ejaculatory capability, and seminal, testicular, epididymal, and endocrine characteristics were  
16 evaluated. In both treated groups, long-term effects consisted of decreased copulatory behavior  
17 (interest, erection, and/or ejaculation), significant increases in acrosomal dysgenesis and nuclear  
18 malformations ( $p < 0.03$ ), and significant decreases in serum concentration of luteinizing  
19 hormone ( $p < 0.05$ ) and testosterone secretion after human chorionic gonadotropin  
20 administration ( $p < 0.04$ ). There were no effects on total spermatozoa per ejaculate or on daily  
21 sperm production. The contribution of individual drinking water contaminants to adverse male  
22 reproductive outcome could not be discerned in this study. Additionally, it was not designed to  
23 distinguish between adverse effects that may have resulted from exposures in late gestation (i.e.,  
24 during critical period of male reproductive system development) versus postnatal life.

#### 4.8.1.1.8.2. Studies assessing female reproductive outcomes

26         In a study that evaluated postnatal growth following gestational exposures, female  
27 B6D2F1 mice (7–12/group) were administered TCE at doses of 0, 1% LD<sub>50</sub> (24 mg/kg-day), and  
28 10% LD<sub>50</sub> (240 mg/kg-day) by gavage in corn oil from GDs 1–5, 6–10, or 11–15 (day of mating  
29 was defined as GD 1) (Cosby and Dukelow, 1992). Litters were examined for pup count, sex,  
30 weight, and crown-rump measurement until GD 21. Some offspring were retained to 6 weeks of  
31 age, at which time they were killed and the gonads were removed, weighed, and preserved. No  
32 treatment-related effects were observed in the dams or offspring. In a second series of studies  
33 conducted by (Cosby and Dukelow) and reported in the same paper, TCE and its metabolites

1 DCA, TCA, and TCOH were added to culture media with capacitated sperm and cumulus masses  
2 from B6D2F1 mice to assess effects on in vitro fertilization. Dose-related decreases in  
3 fertilization were observed for DCA, TCA, and TCOH at 100 and 1,000 ppm, but not with TCE.  
4 Synergistic effects were not observed with TCA and TCOH.

5 A study was conducted by Manson et al. (1984) to determine if subchronic oral exposure  
6 to TCE affected female reproductive performance, and if TCE or its metabolites trichloroacetic  
7 acid or trichloroethanol accumulated in female reproductive organs or neonatal tissues. Female  
8 Long-Evans hooded rats (22–23/group) were administered 0 (corn oil vehicle), 10, 100, or  
9 1,000 mg/kg-day of TCE by gavage for 2 weeks prior to mating, throughout mating, and to GD  
10 21. Delivered pups were examined for gross anomalies, and body weight and survival were  
11 monitored for 31 days. Three maternal animals per group and 8–10 neonates per group (killed  
12 on GDs 3 and 31) were analyzed for TCE and metabolite levels in tissues. TCE exposure  
13 resulted in 5 deaths and decreased maternal body weight gain at 1,000 mg/kg-day, but did not  
14 affect estrous cycle length or female fertility at any dose level. There were no evident  
15 developmental anomalies observed at any treatment level; however, at 1,000 mg/kg-day there  
16 was a significant increase in the number of pups (mostly female) born dead, and the cumulative  
17 neonatal survival count through PND 18 was significantly decreased as compared to control.  
18 TCE levels were uniformly high in fat, adrenal glands, and ovaries across treatment groups, and  
19 TCA levels were high in uterine tissue. TCE levels in the blood, liver, and milk contents of the  
20 stomach increased in female PND-3 neonates across treatment groups. These findings suggest  
21 that increased metabolite levels did not influence fertility, mating success, or pregnancy  
22 outcome.

23 In another study that examined the potential effect of TCE on female reproductive  
24 function, Berger and Horner (2003) conducted 2-week exposures of Sprague-Dawley derived  
25 female Simonson rats to tetrachloroethylene, trichloroethylene, several ethers, and  
26 4-vinylcyclohexene diepoxide in separate groups. The TCE-treated group received 0.45% TCE  
27 in drinking water containing 3% Tween vehicle; control groups were administered either  
28 untreated water, or water containing the 3% Tween vehicle. There were 5–6 females/group, and  
29 three replicates were conducted for each group. At the end of exposure, ovulation was induced,  
30 the rats were killed, and the ovaries were removed. The zona pellucida was removed from  
31 dissected oocytes, which were then placed into culture medium and inseminated with sperm from  
32 untreated males. TCE treatment did not affect female body weight gain, the percentage of  
33 females ovulating, or the number of oocytes per ovulating female. Fertilizability of the oocytes  
34 from treated females was reduced significantly (46% for TCE-treated females versus 56% for  
35 vehicle controls). Oocytes from TCE-treated females had reduced ability to bind sperm plasma  
36 membrane proteins compared with vehicle controls.

1 In subsequent studies, Wu and Berger (2007, 2008) examined the effect of TCE on  
2 oocyte fertilizability and ovarian gene expression. TCE was administered to female Simonson  
3 rats (number of subjects not reported) in the drinking water at 0 or 0.45% (in 3% Tween vehicle);  
4 daily doses were estimated to be 0.66 g TCE/kg body weight/day. In the oocyte fertilizability  
5 study (Wu and Berger, 2007), the female rats were treated on Days 1–5, 6–10, 11–14, or 1–14 of  
6 the 2-week period preceding ovulation (on Day 15). Oocytes were extracted and fertilized in  
7 vitro with sperm from a single male donor rat. With any duration of TCE exposure, fertilization  
8 (as assessed by the presence of decondensed sperm heads) was significantly ( $p < 0.05$ ) decreased  
9 as compared to controls. After exposure on Days 6–10, 11–14, or 1–14, the oocytes from  
10 TCE-treated females had a significantly decreased ability to bind sperm ( $p < 0.05$ ) in comparison  
11 to oocytes from vehicle controls. Increased protein carbonyls (an indicator of oxidatively  
12 modified proteins) were detected in the granulosa cells of ovaries from females exposed to TCE  
13 for 2 weeks. The presence of oxidized protein was confirmed by Western blot analysis.  
14 Microsomal preparations demonstrated the localization of cytochrome P450 2E1 and glutathione  
15 S-transferase (TCE-metabolizing enzymes) in the ovary. Ovarian mRNA transcription for  
16 ALCAM and Cuzd1 protein was not found to be altered after 1 or 5 days of exposure (Wu and  
17 Berger, 2008), suggesting that the posttranslational modification of proteins within the ovary  
18 may partially explain the observed reductions in oocyte fertilization.  
19

#### 4.8.1.1.8.3. Studies assessing fertility and reproductive outcomes in both sexes

20 Assessments of reproduction and fertility with continuous breeding were conducted in  
21 NTP studies in CD-1 mice (George et al., 1985) and Fischer 344 rats (George et al., 1986). TCE  
22 was administered to the mice and rats at dietary levels of 0, 0.15, 0.30, or 0.60%, based upon the  
23 results of preliminary 14-day dose-range finding toxicity studies. Actual daily intake levels for  
24 the study in mice were calculated from the results of dietary formulation analyses and body  
25 weight/food consumption data at several time points during study conduct; the most conservative  
26 were from the second week of the continuous breeding study: 0, 52.5, 266.3, and  
27 615.0 mg/kg-day. No intake calculations were presented for the rat study. In these studies,  
28 which were designed as described by Chapin and Sloane (1997), the continuous breeding phase  
29 in F0 adults consisted of a 7-day pre-mating exposure, 98-day cohabitation period, and 28-day  
30 segregation period. In rats, a crossover mating trial (i.e., control males × control females; 0.60%  
31 TCE males × control females; control males × 0.60% TCE females) was conducted to further  
32 elucidate treatment-related adverse reproductive trends observed in the continuous breeding  
33 phase. The last litter of the continuous breeding phase was raised to sexual maturity for an

1 assessment of fertility and reproduction in control and high-dose groups; for the rats, this  
2 included an open field behavioral assessment of F1 pups. The study protocol included terminal  
3 studies in both generations, including sperm evaluation (count morphology, and motility) in 10  
4 selected males per dose level, macroscopic pathology, organ weights, and histopathology of  
5 selected organs.

6 In the continuous breeding phase of the CD-1 mouse study (George et al., 1985), no  
7 clinical signs of toxicity were observed in the parental (F0) animals, and there were no treatment-  
8 related effects on the proportion of breeding pairs able to produce a litter, the number of live  
9 pups per litter, the percentage born live, the proportion of pups born live, the sex of pups born  
10 live, absolute live pup weights, or adjusted female pup weights. At the high dose level of 0.60%,  
11 a number of adverse outcomes were observed. In the parental animals, absolute and body-  
12 weight-adjusted male and female liver weight values were significantly increased ( $p < 0.01$ ), and  
13 right testis and seminal vesicle weights were decreased ( $p < 0.05$ ), but kidney/adrenal weights  
14 were not affected. Sperm motility was significantly ( $p < 0.01$ ) decreased by 45% in treated  
15 males as compared to controls. Histopathology examination revealed lesions in the liver  
16 (hypertrophy of the centrilobular liver cells) and kidneys (tubular degeneration and karyomegaly  
17 of the corticomedullary renal tubular epithelium) of F0 males and females. In the pups at 0.60%,  
18 adjusted live birth weights for males and both sexes combined were significantly decreased  
19 ( $p < 0.01$ ) as compared to control. The last control and high-dose litters of the continuous  
20 breeding assessment were raised to the age of sexual maturity for a further assessment of  
21 reproductive performance. In these F1 pups, body weights (both sexes) were significantly  
22 decreased at PND 4, and male offspring body weights were significantly ( $p < 0.05$ ) less than  
23 controls at PND 74 ( $\pm 10$ ). It was reported that perinatal mortality (PND 0–21) was increased,  
24 with a 61.3% mortality rate for TCE-treated pups versus a 28.3% mortality rate for control pups.  
25 Reproductive performance was not affected by treatment, and postmortem evaluations of the F1  
26 adult mice revealed significant findings at 0.60% TCE that were consistent with those seen in the  
27 F0 adults and additionally demonstrated renal toxicity, i.e., elevated liver and kidney/adrenal  
28 weights and hepatic and renal histopathological lesions in both sexes, elevated testis and  
29 epididymis weights in males, and decreased sperm motility (18% less than control).

30 The F344 rat study continuous breeding phase demonstrated no evidence of treatment-  
31 related effects on the proportion of breeding pairs able to produce a litter, percentage of pups  
32 born alive, the sex of pups born alive, or absolute or adjusted pup weights (George et al., 1986).  
33 However, the number of live pups per litter was significantly ( $p < 0.05$ ) decreased at 0.30 and  
34 0.60% TCE, and a significant ( $p < 0.01$ ) trend toward a dose-related decrease in the number of  
35 live litters per pair was observed; individual data were reported to indicate a progressive decrease  
36 in the number of breeding pairs in each treatment group producing third, fourth, and fifth litters.

1 The crossover mating trial conducted in order to pursue this outcome demonstrated that the  
2 proportion of detected matings was significantly depressed ( $p < 0.05$ ) in the mating pairs with  
3 TCE-treated partners compared to the control pairs. In the F0 adults at 0.60% TCE, postpartum  
4 dam body weights were significantly decreased ( $p < 0.01$  or  $0.05$ ) in the continuous breeding  
5 phase and the crossover mating trials, and terminal body weights were significantly decreased  
6 ( $p < 0.01$ ) for both male and female rats. Postmortem findings for F0 adults in the high-dose  
7 group included significantly increased absolute and body-weight-adjusted liver and  
8 kidney/adrenal weights in males, increased adjusted liver and kidney/adrenal weights in females,  
9 and significantly increased adjusted left testis/epididymal weights. Sperm assessment did not  
10 identify any effects on motility, concentration or morphology, and histopathological examination  
11 was negative. The last control and high-dose litters of the continuous breeding assessment were  
12 raised to the age of sexual maturity for assessment of open field behavior and reproductive  
13 performance. In these F1 pups at 0.60% TCE, body weights of male and females were  
14 significantly ( $p < 0.05$  or  $0.01$ , respectively) decreased at PND 4 and 14. By PND 21, pup  
15 weights in both sexes were significantly reduced in all treated groups, and this continued until  
16 termination (approximately PND 80). A tendency toward decreased postweaning survival (i.e.,  
17 from PND 21 to PND  $81 \pm 10$ ) was reported for F1 pups at the 0.15 and 0.60% levels. Open  
18 field testing revealed a significant ( $p < 0.05$ ) dose-related trend toward an increase in the time  
19 required for male and female F1 weanling pups to cross the first grid in the testing device,  
20 suggesting an effect on the ability to react to a novel environment. Reproductive performance  
21 assessments conducted in this study phase were not affected by treatment. Postpartum F1 dam  
22 body weights were significantly decreased ( $p < 0.05$  or  $0.01$ ) in all of the TCE-treated groups as  
23 compared to controls, as were terminal body weights for both adult F1 males and females.  
24 Postmortem evaluations of the F1 adult rats revealed significantly ( $p < 0.01$ ) decreased left  
25 testis/epididymis weight at 0.60% TCE, and significantly ( $p < 0.05$  or  $0.01$ ) increased adjusted  
26 mean liver weight in all treated groups for males and at 0.30 and 0.60% for females. Sperm  
27 assessments for F1 males revealed a significant increase ( $p < 0.05$ ) in the percentage of abnormal  
28 sperm in the 0.30% TCE group, but no other adverse effects on sperm motility, concentration, or  
29 morphology were observed. As with the F0 adults, there were no adverse treatment-related  
30 findings revealed at histopathological assessment. The study authors concluded that the  
31 observed effects to TCE exposure in this study were primarily due to generalized toxicity and not  
32 to a specific effect on the reproductive system; however, based upon the overall toxicological  
33 profile for TCE, which demonstrates that the male reproductive system is a target for TCE  
34 exposures, this conclusion is not supported.  
35

#### 4.8.1.1.9. Discussion/Synthesis of Noncancer Reproductive Toxicity Findings

1 The human epidemiological findings and animal study evidence consistently indicate that  
2 TCE exposures can result in adverse reproductive outcomes. Although the epidemiological data  
3 may not always be robust or unequivocal, they demonstrate the potential for a wide range of  
4 exposure-related adverse outcomes on female and male reproduction. In animal studies, there is  
5 some evidence for female-specific reproductive toxicity; but there is strong and compelling  
6 evidence for adverse effects of TCE exposure on male reproductive system and function.  
7

#### 4.8.1.1.10. Female reproductive toxicity

8 Although few epidemiological studies have examined TCE exposure in relation to female  
9 reproductive function (see Table 4-88), the available studies provide evidence of decreased  
10 fertility, as measured by time to pregnancy (Sallmén et al., 1995), and effects on menstrual cycle  
11 patterns, including abnormal cycle length (ATSDR, 2001), amenorrhea (Sagawa et al., 1973;  
12 Zielinski, 1973), and menstrual “disturbance” (Bardodej and Vyskocil, 1956). In experimental  
13 animals, the effects on female reproduction include evidence of reduced in vitro oocyte  
14 fertilizability in rats (Berger and Horner, 2003; Wu and Berger, 2007). However, in other  
15 studies that assessed reproductive outcome in female rodents (Cosby and Dukelow, 1992;  
16 George et al., 1985, 1986; Manson et al., 1984), there was no evidence of adverse effects of TCE  
17 exposure on female reproductive function. Overall, although the data are suggestive, there are  
18 inadequate data to make conclusions as to whether adverse effects on human female  
19 reproduction are caused by TCE.  
20  
21

22 **Table 4-88. Summary of adverse female reproductive outcomes associated**  
23 **with TCE exposures**  
24

Finding	Species	Citation
Menstrual cycle disturbance	Human	ATSDR (2001) <sup>a</sup>
		Bardodej and Vyskocil (1956)
		Sagawa et al. (1973)
		Zielinski (1973)
Reduced fertility	Human <sup>a</sup>	Sallmén et al. (1995)
	Rat <sup>b</sup>	Berger and Horner (2003)
		Wu and Berger (2007)

1 <sup>a</sup>Not significant.  
2 <sup>b</sup>In vitro oocyte fertilizability.  
3  
4

#### **4.8.1.1.11. Male reproductive toxicity**

5           Notably, the results of a number of studies in both humans and experimental animals  
6 have suggested that exposure to TCE can result in targeted male reproductive toxicity (see  
7 Table 4-89). The adverse effects that have been observed in both male humans and male animal  
8

1  
2  
3

**Table 4-89. Summary of adverse male reproductive outcomes associated with TCE exposures**

Finding	Species	Citation
Testicular toxicity/pathology	Rat	George et al. (1986)
		Kumar et al. (2000b)
		Kumar et al. (2001b)
	Mouse	Kan et al. (2007)
Epididymal toxicity/pathology	Mouse	Forkert et al. (2002)
Decreased sperm quantity/quality	Human	Chia et al. (1996)
		Rasmussen et al. (1988) <sup>a</sup>
	Rat	Kumar et al. (2001b; 2000a; 2000b)
	Mouse	George et al. (1985)
		Land et al. (1981)
	Rabbit	Veeramachaneni et al. (2001)
Altered in vitro sperm-oocyte binding or in vivo fertilization	Rat	DuTeaux et al. (2004a)
	Mouse	Cosby and Dukelow (1992) <sup>b</sup>
		Xu et al. (2004) <sup>b</sup>
Altered sexual drive or function	Human	El Ghawabi et al. (1973)
		Saihan et al. (1978) <sup>c</sup>
		Bardodej and Vyskocil (1956)
	Rat	George et al. (1986)
		Zenick et al. (1984)
	Rabbit	Veeramachaneni et al. (2001)
Altered serum testosterone levels	Human	Chia et al. (1997) <sup>d</sup>
		Goh et al. (1998) <sup>e</sup>
	Rat	Kumar et al. (2000a)
	Rabbit	Veeramachaneni et al. (2001)
Reduced fertility	Rat	George et al. (1986)
Gynaecomastia	Human	Saihan et al. (1978) <sup>c</sup>

4  
5  
6  
7

<sup>a</sup>Nonsignificant increase in percentage of two YFF in spermatozoa; no effect on sperm count or morphology.

<sup>b</sup>Observed with metabolite(s) of TCE only.

<sup>c</sup>Case study of one individual.

<sup>d</sup>Also observed altered levels of DHEAS, FSH, and SHBG.



1 °Also observed altered levels of SHBG.  
2

1 models include altered sperm count, morphology, or motility (Chia et al., 1996; George et al.,  
2 1985; Kumar et al., 2001b; Kumar et al., 2000a; Kumar et al., 2000b; Land et al., 1981;  
3 Rasmussen et al., 1988; Veeramachaneni et al., 2001); decreased libido or copulatory behavior  
4 (Bardodej and Vyskocil, 1956; El Ghawabi et al., 1973; George et al., 1986; Saihan et al., 1978;  
5 Veeramachaneni et al., 2001; Zenick et al., 1984); alterations in serum hormone levels (Chia et  
6 al., 1997; Goh et al., 1998; Kumar et al., 2000a; Veeramachaneni et al., 2001); and reduced  
7 fertility (George et al., 1986). However, other studies in humans did not see evidence of altered  
8 sperm count or morphology (Rasmussen et al., 1988) or reduced fertility (Forkert et al., 2003;  
9 Sallmen et al., 1998), and some animal studies also did not identify altered sperm measures  
10 (Cosby and Dukelow, 1992; George et al., 1986; Xu et al., 2004; Zenick et al., 1984). Additional  
11 adverse effects observed in animals include histopathological lesions of the testes (George et al.,  
12 1986; Kumar et al., 2001b; Kumar et al., 2000b) or epididymides (Forkert et al., 2002; Kan et al.,  
13 2007) and altered in vitro sperm-oocyte binding and/or in vivo fertilization for TCE and/or its  
14 metabolites (DuTeaux et al., 2004a; Xu et al., 2004).

15 In spite of the preponderance of studies demonstrating effects on sperm parameters, there  
16 is an absence of overwhelming evidence in the database of adverse effects of TCE on overall  
17 fertility in the rodent studies. That is not surprising, however, given the redundancy and  
18 efficiency of rodent reproductive capabilities. Nevertheless, the continuous breeding  
19 reproductive toxicity study in rats (George et al., 1986) did demonstrate a trend towards  
20 reproductive compromise (i.e., a progressive decrease in the number of breeding pairs producing  
21 third, fourth, and fifth litters).

22 It is noted that in the studies by George et al. (George et al., 1985, 1986), adverse  
23 reproductive outcomes in male rats and mice were observed at the highest dose level tested  
24 (0.060% TCE in diet) which was also systemically toxic (i.e., demonstrating kidney toxicity and  
25 liver enzyme induction and toxicity, sometimes in conjunction with body weight deficits).  
26 Because of this, the study authors concluded that the observed reproductive toxicity was a  
27 secondary effect of generalized systemic toxicity; however, this conclusion is not supported by  
28 the overall toxicological profile of TCE which provides significant evidence indicating that TCE  
29 is a reproductive toxicant.

30

#### 4.8.1.1.11.1. **The role of metabolism in male reproductive toxicity**

31 There has been particular focus on evidence of exposure to male reproductive organs by  
32 TCE and/or its metabolites, as well as the role of TCE metabolites in the observed toxic effects.

1 In humans, a few studies demonstrating male reproductive toxicity have measured levels  
2 of TCE in the body. U-TCA was measured in men employed in an electronics factory, and  
3 adverse effects observed included abnormal sperm morphology and hyperzoospermia and altered  
4 serum hormone levels (Chia et al., 1996, 1997; Goh et al., 1998). U-TCA was also measured as  
5 a marker of exposure to TCE in men occupationally exposed to solvents, although this study did  
6 not report any adverse effects on fertility (Sallmen et al., 1998).

7 In the study in Long-Evans male rats by Zenick et al. (1984), blood and tissue levels of  
8 TCE, TCA, and TCOH were measured in three rats/group following 6 weeks of gavage treatment  
9 at 0, 10, 100, and 1,000 mg/kg-day. Additionally the levels of TCE and metabolites were  
10 measured in seminal plugs recovered following copulation at Week 5. Marked increases in TCE  
11 levels were observed only at 1,000 mg/kg-day, in blood, muscle, adrenals, and seminal plugs. It  
12 was reported that dose-related increases in TCA and TCOH concentrations were observed in the  
13 organs evaluated, notably including the reproductive organs (epididymis, vas deferens, testis,  
14 prostate, and seminal vesicle), thus, creating a potential for interference with reproductive  
15 function.

16 This potential was explored further in a study by Forkert et al. (2002), in which male  
17 CD-1 mice were exposed by inhalation to 1,000-ppm TCE (6 hours/day, 5 day/week) for  
18 4 consecutive weeks. Urine was obtained on Days 4, 9, 14, and 19 of exposure and analyzed for  
19 concentrations of TCE and TCOH. Microsomal preparations from the liver, testis and  
20 epididymis were used for immunoblotting, determining *p*-nitrophenol hydroxylase and CYP2E1  
21 activities, and evaluating the microsomal metabolism of TCE.

22 Subsequent studies conducted by the same laboratory (Forkert et al., 2003) evaluated the  
23 potential of the male reproductive tract to accumulate TCE and its metabolites including chloral,  
24 TCOH, TCA, and DCA. Human seminal fluid and urine samples from eight mechanics  
25 diagnosed with clinical infertility and exposed to TCE occupationally were analyzed. Urine  
26 samples from two of the eight subjects contained TCA and/or TCOH, suggesting that TCE  
27 exposure and/or metabolism was low during the time just prior to sample collection. TCE,  
28 chloral, and TCOH were detected in seminal fluid samples from all eight subjects, while TCA  
29 was found in one subject, and DCA was found in two subjects. Additionally, TCE and its  
30 metabolites were assessed in the epididymis and testis of CD-1 mice (4/group) exposed by  
31 inhalation (6 hours/day, 5 days/week) to 1,000 ppm TCE for 1, 2, and 4 weeks. TCE, chloral,  
32 and TCOH were found in the epididymis at all timepoints, although TCOH levels were increased  
33 significantly (tripled) at 4 weeks of exposure. This study showed that the metabolic disposition  
34 of TCE in humans is similar to that in mice, indicating that the murine model is appropriate for  
35 investigating the effects of TCE-induced toxicity in the male reproductive system. These studies  
36 provide support for the premise that TCE is metabolized in the human reproductive tract, mainly

1 in the epididymis, resulting in the production of metabolites that cause damage to the epididymal  
2 epithelium and affect the normal development of sperm.

3 Immunohistochemical experiments (Forkert et al., 2002) confirmed the presence of  
4 CYP2E1 in the epididymis and testis of mice; it was found to be localized in the testicular  
5 Leydig cells and the epididymal epithelium. Similar results were obtained with the  
6 immunohistochemical evaluation of human and primate tissue samples. CYP2E1 has been  
7 previously shown by Lipscomb et al. (1998a) to be the predominant CYP enzyme catalyzing the  
8 hepatic metabolism of TCE in both animals and rodents. These findings support the role of  
9 CYP2E1 in TCE metabolism in the male reproductive tract of humans, primates, and mice.  
10

#### 4.8.1.1.11.2. **Mode of action for male reproductive toxicity**

11 A number of studies have been conducted to attempt to characterize various aspects of  
12 the mode of action for observed male reproductive outcomes.

13 Studies by Kumar et al. (2001b; 2000a) suggest that perturbation of testosterone  
14 biosynthesis may have some role in testicular toxicity and altered sperm measures. Significant  
15 decreases in the activity of G6PDH and accumulation of cholesterol are suggestive of an  
16 alteration in testicular steroid biosynthesis. Increased testicular lipids, including cholesterol,  
17 have been noted for other testicular toxicants such as lead (Saxena et al., 1987),  
18 triethylenemelamine (Johnson et al., 1967), and quinalphos (Ray et al., 1987), in association with  
19 testicular degeneration and impaired spermatogenesis. Since testosterone has been shown to be  
20 essential for the progression of spermatogenesis (O'Donnell et al., 1994), alterations in  
21 testosterone production could be a key event in male reproductive dysfunction following TCE  
22 exposure. Additionally, the observed TCE-related reduction of 17- $\beta$ -HSD, which is involved in  
23 the conversion of androstenedione to testosterone, has also been associated with male  
24 reproductive insufficiency following exposure to phthalate esters (Srivastava and Srivastava,  
25 1991), quinalphos (Ray et al., 1987), and lead (Saxena et al., 1987). Reductions in SDH, which  
26 are primarily associated with the pachytene spermatocyte maturation of germinal epithelium,  
27 have been shown to be associated with depletion of germ cells (Chapin et al., 1982; Mills and  
28 Means, 1972), and the activity of G6PDH is greatest in premeiotic germ cells and Leydig cells of  
29 the interstitium (Blackshaw, 1970). The increased GT and glucuronidase observed following  
30 TCE exposures appear to be indicative of impaired Sertoli cell function (Hodgen and Sherins,  
31 1973; Sherins and Hodgen, 1976). Based upon the conclusions of these studies, Kumar et al.  
32 (2001b) hypothesized that the reduced activity of G6PDH and SDH in testes of TCE-exposed

1 male rats is indicative of the depletion of germ cells, spermatogenic arrest, and impaired function  
2 of the Sertoli cells and Leydig cells of the interstitium.

3 In the series of experiments by DuTeaux et al. (2004a; 2003), protein dichloroacetyl  
4 adducts were found in the corpus epididymis and in the efferent ducts of rats administered TCE;  
5 this effect was also demonstrated following in vitro exposure of reproductive tissues to TCE.  
6 Oxidized proteins were detected on the surface of spermatozoa from TCE-treated rats in a  
7 dose-response pattern; this was confirmed using a Western blotting technique. Soluble (but not  
8 mitochondrial) cysteine-conjugate  $\beta$ -lyase was detected in the epididymis and efferent ducts of  
9 treated rats. Following a single intraperitoneal injection of DCVC, no dichloroacetylated protein  
10 adducts were detected in the epididymis and efferent ducts. The presence of CYP2E1 was found  
11 in epididymis and efferent ducts, suggesting a role of cytochrome P450-dependent metabolism  
12 in adduct formation. An in vitro assay was used to demonstrate that epididymal and efferent  
13 duct microsomes are capable of metabolizing TCE; TCE metabolism in the efferent ducts was  
14 found to be inhibited by anti-CYP2E1 antibody. Lipid peroxidation in sperm, presumably  
15 initiated by free radicals, was increased in a significant ( $p < 0.005$ ) dose-dependent manner after  
16 TCE-exposure.

17 Overall, it has been suggested (DuTeaux et al., 2004a) that reproductive organ toxicities  
18 observed following TCE exposure are initiated by metabolic bioactivation, leading to subsequent  
19 protein adduct formation. It has been hypothesized that epoxide hydrolases in the rat epididymis  
20 may play a role in the biological activation of metabolites (DuTeaux et al., 2004b). Disruption  
21 of colony stimulating factor and of macrophage development may also play a role in sperm  
22 production (Cohen et al., 1999), and thus may be another route through which immune-related  
23 effects of TCE may operate. In addition, the potential for epigenetic changes, through which  
24 heritable changes in gene mutations occur without changes in DNA sequencing, should also be  
25 considered in the evaluation of transgenerational effects (Guerrero-Bosagna and Skinner, 2009).

#### 26 **4.8.1.1.12. Summary of noncancer reproductive toxicity**

27 The toxicological database for TCE includes a number of studies that demonstrate  
28 adverse effects on the integrity and function of the reproductive system in females and males.  
29 Both the epidemiological and animal toxicology databases provide suggestive, but limited,  
30 evidence of adverse outcomes to female reproductive outcomes. However, much more extensive  
31 evidence exists in support of an association between TCE exposures and male reproductive  
32 toxicity. The available epidemiological data and case reports that associate TCE with adverse  
33 effects on male reproductive function are limited in size and provide little quantitative dose data

1 (Lamb and Hentz, 2006). However, the animal data provide extensive evidence of TCE-related  
2 male reproductive toxicity. Strengths of the database include the presence of both functional and  
3 structural outcomes, similarities in adverse treatment-related effects observed in multiple species,  
4 and evidence that metabolism of TCE in male reproductive tract tissues is associated with  
5 adverse effects on sperm measures in both humans and animals (suggesting that the murine  
6 model is appropriate for extrapolation to human health risk assessment). Additionally some  
7 aspects of a putative MOA (e.g., perturbations in testosterone biosynthesis) appear to have some  
8 commonalities between humans and animals.  
9

#### 4.8.2. Cancers of the Reproductive System

10 The effects of TCE on cancers of the reproductive system have been examined for males  
11 and females in both epidemiological and experimental animal studies. The epidemiological  
12 literature includes data on prostate in males and cancers of the breast and cervix in females. The  
13 experimental animal literature includes data on prostate and testes in male rodents; and uterus,  
14 ovary, mammary gland, vulva, and genital tract in female rodents. The evidence for these  
15 cancers is generally not robust.  
16

##### 4.8.2.1.1. Human Data

17 The epidemiologic evidence on TCE and cancer of the prostate, breast, and cervix is from  
18 cohort and geographic based studies. Two additional case-control studies of prostate cancer in  
19 males are nested within cohorts (Greenland et al., 1994; Krishnadasan et al., 2007). The nested  
20 case-control studies are identified in Tables 4-90 through 4-92 with cohort studies given their  
21 source population for case and control identification. One population-based case-control study  
22 examined on TCE exposure and prostate (Siemiatycki, 1991); however, no population case-  
23 control studies on breast or cervical cancers and TCE exposure were found in the peer-reviewed  
24 literature.  
25

##### 4.8.2.1.2. Prostate cancer

26 Sixteen cohort or PMR studies, two nested case-control, one population case-control, and  
27 two geographic-based studies present relative risk estimates for prostate cancer (Anttila et al.,  
28 1995; ATSDR, 2004a, 2006a; Axelson et al., 1994; Blair et al., 1989; Blair et al., 1998; Boice et  
29 al., 1999; Boice et al., 2006a; Chang et al., 2003a; Chang et al., 2005; Garabrant et al., 1988;

1 Greenland et al., 1994; Hansen et al., 2001; Krishnadasan et al., 2007; Morgan and Cassady,  
 2 2002; Morgan et al., 1998; Raaschou-Nielsen et al., 2003; Radican et al., 2008; Ritz, 1999a;  
 3 Shannon et al., 1988; Siemiatycki, 1991; Wilcosky et al., 1984). Three small cohort studies  
 4 (Costa et al., 1989; Henschler et al., 1995; Sinks et al., 1992), one multiple-site population case-  
 5 control (Siemiatycki, 1991) and one geographic-based study (Vartiainen et al., 1993) do not  
 6 report estimates for prostate cancer in their published papers. Twelve of the 19 studies with  
 7 prostate cancer relative risk estimates had high likelihood of TCE exposure in individual study  
 8 subjects and were judged to have met, to a sufficient degree, the standards of epidemiologic  
 9 design and analysis (Siemiatycki, 1991);

10  
 11 **Table 4-90. Summary of human studies on TCE exposure and prostate**  
 12 **cancer**  
 13

Studies	Exposure group	Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort studies—incidence</b>				
Aerospace workers (Rocketdyne)				Krishnadasan et al. (2007)
	Low/moderate TCE score	1.3 (0.81, 2.1) <sup>a,b</sup>	90	
	High TCE score	2.1 (1.2, 3.9) <sup>a,b</sup>	45	
	<i>p</i> for trend	0.02		
	Low/moderate TCE score	1.3 (0.81, 2.1) <sup>a,c</sup>		
	High TCE score	2.4 (1.3, 4.4) <sup>a,c</sup>		
	<i>p</i> for trend	0.01		
All employees at electronics factory (Taiwan)		0.14 (0.00, 0.76) <sup>d</sup>	1	Chang et al. (2005)
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure	0.9 (0.79, 1.08)	163	
Biologically-monitored Danish workers				Hansen et al. (2001)
	Any TCE exposure, females	0.6 (0.2, 1.3)	6	
Aircraft maintenance workers (Hill Air Force Base, UT)				
	TCE subcohort	Not reported	158	Blair et al. (1998)
	Cumulative exposure			
	0	1.0 <sup>e</sup>		
	<5 ppm-yr	1.1 (0.7, 1.6)	64	
	5–25 ppm-yr	1.0 (0.6, 1.6)	38	
	>25 ppm-yr	1.2 (0.8, 1.8)	56	
	TCE subcohort	1.2 (0.92, 1.76)	116	Radican et al. (2008)
	Cumulative exposure			

	0	1.0 <sup>e</sup>		
	<5 ppm-yr	1.03 (0.65, 1.62)	41	
	5–25 ppm-yr	1.33 (0.82, 2.15)	42	
	>25 ppm-yr	1.31 (0.84, 2.06)	43	
Biologically-monitored Finnish workers		1.38 (0.73, 2.35)	13	Anttila et al. (1995)
	Mean air-TCE (Ikeda extrapolation)			
	<6 ppm	1.43 (0.62, 2.82)	8	
	6+ ppm	0.68 (0.08, 2.44)	2	

1



**Table 4-90. Summary of human studies on TCE exposure and prostate cancer (continued)**

Studies	Exposure group	Relative risk (95% CI)	No. obs. Events	Reference
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
	Exposed workers	Not reported		
Biologically-monitored Swedish workers		1.25 (0.84, 1.84)	26	Axelsson et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA		Not reported		Sinks et al. (1992)
<b>Cohort and PMR-mortality</b>				
Aerospace workers (Rocketdyne)				Boice et al. (2006a)
	Any TCE (utility/eng flush)	0.82 (0.36, 1.62)	8	
View-Master employees		1.69 (0.68, 3.48) <sup>f</sup>	8	ATSDR (2004a)
All employees at electronics factory (Taiwan)		Not reported	0	Chang et al. (2003a)
Fernald workers				Ritz (1999a)
	Any TCE exposure	Not reported		
	Light TCE exposure, >2 yr duration	0.91 (0.38, 2.18) <sup>e,g</sup>	10	
	Moderate TCE exposure, >2 yr duration	1.44 (0.19, 11.4) <sup>e,g</sup>	1	
Aerospace workers (Lockheed)				Boice et al. (1999)
	Routine exposure to TCE	1.31 (0.52, 2.69)	7	
	Routine-intermittent	Not reported		
Aerospace workers (Hughes)				Morgan et al. (1998, 2000b)
	TCE subcohort	1.18 (0.73, 1.80)	21	
	Low intensity (<50 ppm)	1.03 (0.51, 1.84)	7	
	High intensity (>50 ppm)	0.47 (0.15, 1.11)	14	
TCE subcohort (Cox Analysis)				
	Never exposed	1.00 <sup>e</sup>		
	Ever exposed	1.58 (0.96, 2.62) <sup>h</sup>		
Peak				
	No/Low	1.00 <sup>e</sup>		
	Medium/high	1.39 (0.80, 2.41) <sup>h</sup>		
Cumulative				
	Referent	1.00 <sup>e</sup>		
	Low	1.72 (0.78, 3.80) <sup>h</sup>		
	High	1.53 (0.85, 2.75) <sup>h</sup>		

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-90. Summary of human studies on TCE exposure and prostate cancer (continued)**

Studies	Exposure group	Relative risk (95% CI)	No. obs. Events	Reference
Aircraft maintenance workers (Hill Air Force Base, UT)				Blair et al. (1998)
	TCE subcohort	1.1 (0.6, 1.8)	54	
	Cumulative exposure			
	0	1.0 <sup>c</sup>		
	<5 ppm-yr	0.9 (0.5, 1.8)	19	
	>25 ppm-yr	1.3 (0.7, 2.4)	22	
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
	TCE exposed workers	Not reported		
Deaths reported to GE pension fund (Pittsfield, MA)		0.82 (0.46, 1.46) <sup>a</sup>	58	Greenland et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA		Not reported	0	Sinks et al. (1992)
U. S. Coast Guard employee				Blair et al. (1989)
	Marine inspectors	1.06 (0.51, 1.95)	10	
	Noninspectors	0.57 (0.15, 1.45)	7	
Aircraft manufacturing plant employees (Italy)				Costa et al. (1989)
Aircraft manufacturing plant employees (San Diego, CA)		0.93 (0.60, 1.37)	25	Garabrant et al. (1988)
Lamp manufacturing workers (GE)		1.56 (0.63, 3.22)	7	Shannon et al. (1988)
Rubber workers				Wilcosky et al. (1984)
	Any TCE exposure	0.62 (not reported)	3	
<b>Case-control studies</b>				
Population of Montreal, Canada				Siemiatycki (1991)
	Any TCE exposure	1.1 (0.6, 2.1) <sup>i</sup>	11	
	Substantial TCE exposure	1.8 (0.8, 4.0) <sup>i</sup>	7	
<b>Geographic based studies</b>				
Residents in two study areas in Endicott, NY		1.05 (0.75, 1.43)	40	ATSDR (2006b)
Residents of 13 census tracts in Redlands, CA		1.11 (0.98, 1.25) <sup>j</sup>	483	Morgan and Cassady (2002)
Finnish residents				Vartiainen et al. (1993)
	Residents of Hausjarvi	Not reported		
	Residents of Huttula	Not reported		

1  
2 <sup>a</sup>Odds ratio from nested case-control study.  
3 <sup>b</sup>Odds ratio, zero lag.

1                   **Table 4-90. Summary of human studies on TCE exposure and prostate**  
2                   **cancer (continued)**

3  
4                   <sup>c</sup>Odds ratio, 20 yr lag.

5                   <sup>d</sup>Chang et al. (2005) presents SIRs for a category site of all cancers of male genital organs.

6                   <sup>e</sup>Internal referents, workers without TCE exposure.

7                   <sup>f</sup>Proportional mortality ratio.

8                   <sup>g</sup>Analysis for >2 yrs exposure duration and a lagged TCE exposure period of 15 yrs.

9                   <sup>h</sup>Risk ratio from Cox Proportional Hazard Analysis, stratified by age and sex, from Environmental Health Strategies  
10                   (1997) Final Report to Hughes Corporation (Communication from Paul A. Cammer, President, Trichloroethylene  
11                   Issues Group to Cheryl Siegel Scott, U.S. EPA, December 22, 1997).

12                   <sup>i</sup>90% CI.

13                   <sup>j</sup>99% CI.

14  
15                   GE = General Electric, No. obs. events = number of observed events.

16

1  
2

**Table 4-91. Summary of human studies on TCE exposure and breast cancer**

Studies	Exposure group	Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort studies—incidence</b>				
Aerospace workers (Rocketdyne)				Zhao et al. (2005)
	Any TCE exposure	Not reported		
	Low cumulative TCE score			
	Medium cumulative TCE score			
	High TCE score			
	<i>p</i> for trend			
All employees at electronics factory (Taiwan)				
	Females	1.09 (0.96, 1.22) <sup>a</sup>	286	Sung et al. (2007)
	Females	1.19 (1.03, 1.36)	215	Chang et al. (2005)
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure, males	0.5 (0.06, 1.90)	2	
	Any exposure, females	1.1 (0.89, 1.24)	145	
Biologically-monitored Danish workers				Hansen et al. (2001)
	Any TCE exposure, males		0 (0.2 exp)	
	Any TCE exposure, females	0.9 (0.2, 2.3)	4	
Aircraft maintenance workers (Hill Air Force Base, UT)				Blair et al. (1998)
	TCE subcohort	Not reported	34	
	Females, cumulative exposure			
	0	1.0 <sup>b</sup>		
	<5 ppm-yr	0.3 (0.1, 1.4)	20	
	5–25 ppm-yr	0.4 (0.1, 2.9)	11	
	>25 ppm-yr	0.4 (0.4, 1.2)	3	
Biologically-monitored Finnish workers		Not reported		Anttila et al. (1995)
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
	Exposed workers	Not reported		
Biologically-monitored Swedish workers		Not reported		Axelsson et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA		Not reported		Sinks et al. (1992)

3

**Table 4-91. Summary of human studies on TCE exposure and breast cancer (continued)**

Studies	Exposure group	Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort and PMR-mortality</b>				
Aerospace workers (Rocketdyne)				
	Any TCE (utility/eng flush)	Not reported		Boice et al. (2006a)
	Any exposure to TCE	Not reported		Zhao et al. (2005)
	Low cumulative TCE score	Not reported		
	Medium cumulative TCE score	Not reported		
	High TCE score	Not reported		
	<i>p</i> for trend			
View-Master employees				ATSDR (2004a)
	Males		0 (0.05 exp)	
	Females	1.02 (0.67, 1.49) <sup>c</sup>	27	
Fernald workers				Ritz (1999a)
	Any TCE exposure	Not reported		
	Light TCE exposure, >2 yr duration	Not reported		
	Moderate TCE exposure, >2 yr duration	Not reported		
Aerospace workers (Lockheed)				Boice et al. (1999)
	Routine exposure to TCE	1.31 (0.52, 2.69) <sup>d</sup>	7	
	Routine-intermittent <sup>a</sup>	Not reported		
Aerospace workers (Hughes)				Morgan et al. (1998)
	TCE subcohort	0.75 (0.43, 1.22) <sup>d</sup>	16	
	Low intensity (<50 ppm)	1.03 (0.51, 1.84) <sup>d</sup>	11	
	High intensity (>50 ppm)	0.47 (0.15, 1.11) <sup>d</sup>	5	
TCE subcohort (Cox Analysis)				
	Never exposed	1.00 <sup>d</sup>	NR	
	Ever exposed	0.94 (0.51, 1.75) <sup>d,e</sup>	NR	
Peak				
	No/Low	1.00 <sup>d</sup>		
	Medium/high	1.14 (0.48, 2.70) <sup>d,e</sup>	NR	
Cumulative				
	Referent	1.00 <sup>b</sup>		
	Low	1.20 (0.60, 2.40) <sup>d,e</sup>	NR	
	High	0.65 (0.25, 1.69) <sup>d,e</sup>	NR	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-91. Summary of human studies on TCE exposure and breast cancer (continued)**

Studies	Exposure group	Relative risk (95% CI)	No. obs. events	Reference
Aircraft maintenance workers (Hill Air Force Base, UT)				
	TCE subcohort (females)	2.0 (0.9, 4.6)	20	Blair et al. (1998)
	Females, cumulative exposure			
	0	1.0 <sup>b</sup>		
	<5 ppm-yr	2.4 (1.1, 5.2)	10	
	5–25 ppm-yr	1.2 (0.3, 5.4)	21	
	>25 ppm-yr	1.4 (0.6, 3.2)	8	
	Low level intermittent exposure	3.1 (1.5, 6.2)	15	
	Low level continuous exposure	3.4 (1.4, 8.0)	8	
	Frequent peaks	1.4 (0.7, 3.2)	10	
	TCE subcohort (females)	1.23 (0.73, 2.06)	26	Radican et al. (2008)
	Females, cumulative exposure			
	0	1.0 <sup>b</sup>		
	<5 ppm-yr	1.57 (0.81, 3.04)	12	
	5–25 ppm-yr	1.01 (0.31, 3.30)	3	
	>25 ppm-yr	1.05 (0.53, 2.07)	11	
	Low level intermittent exposure	1.92 (1.08, 3.43)	18	
	Low level continuous exposure	1.71 (0.79, 3.71)	8	
	Frequent peaks	1.08 (0.57, 2.02)	14	
Cardboard manufacturing workers in Arnsburg, Germany				
	TCE exposed workers	Not examined		Henschler et al. (1995)
Deaths reported to GE pension fund (Pittsfield, MA)		Not reported		Greenland et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA		Not reported	0	Sinks et al. (1992)
U. S. Coast Guard employees				
	Marine inspectors	Not reported		Blair et al. (1989)
	Noninspectors	Not reported		
Aircraft manufacturing plant employees (Italy)		Not reported <sup>f</sup>		Costa et al. (1989)
Aircraft manufacturing plant employees (San Diego, CA)				
	All subjects, females	0.81 (0.52, 1.48) <sup>d</sup>	16	Garabrant et al. (1988)
Lamp manufacturing workers (GE)				
	Coil/wire drawing	2.04 (0.88, 4.02)	8	Shannon et al. (1988)
	Other areas	0.97 (0.57, 1.66)	13	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-91. Summary of human studies on TCE exposure and breast cancer (continued)**

Studies	Exposure group	Relative risk (95% CI)	No. obs. events	Reference
<b>Case-control Studies</b>				
Population of Montreal, Canada				Siemiatycki (1991)
	Any TCE exposure	Not reported		
	Substantial TCE exposure	Not reported		
<b>Geographic Based Studies</b>				
Residents in two study areas in Endicott, NY		0.88 (0.65, 1.18)	46	ATSDR (2006b)
Residents of 13 census tracts in Redlands, CA		1.09 (0.97, 1.21)	536	Morgan and Cassady (2002)
Finnish residents				Vartiainen et al. (1993)
	Residents of Hausjarvi	Not reported		
	Residents of Huttula	Not reported		

1  
2 <sup>a</sup>15 yr lag.  
3 <sup>b</sup>Internal referents, workers not exposed to TCE.  
4 <sup>c</sup>Proportional mortality ratio.  
5 <sup>d</sup>In Garabrant et al. (1988), Morgan et al. (1998) and Boice et al. (1999), breast cancer risk is for males and females  
6 combined (ICD-9, 174, 175).  
7 <sup>e</sup>Risk ratio from Cox Proportional Hazard Analysis, stratified by age and sex, from Environmental Health Strategies  
8 (1997) Final Report to Hughes Corporation (Communication from Paul A. Cammer, President, Trichloroethylene  
9 Issues Group to Cheryl Siegel Scott, U.S. EPA, December 22, 1997).  
10 <sup>f</sup>The cohort of Blair et al. (1989) and Costa et al. (1989) are composed of males only.  
11  
12 GE = General Electric, No. obs. events = number of observed events; NR = not reported.  
13

1  
2  
3

**Table 4-92. Summary of human studies on TCE exposure and cervical cancer**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort studies—incidence</b>				
Aerospace workers (Rocketdyne)				Zhao et al. (2005)
	Any exposure to TCE	Not reported		
	Low cumulative TCE score	Not reported		
	Medium cumulative TCE score			
	High TCE score			
	<i>p</i> for trend			
All employees at electronics factory (Taiwan)		0.96 (0.86, 1.22) <sup>a</sup>	337	Sung et al. (2007)
Danish blue-collar worker w/TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure	1.9 (1.42, 2.37)	62	
	Exposure lag time			
	20 yr	1.5 (0.7, 2.9)	9	
	Employment duration			
	<1 yr	2.5 (1.7, 3.5)	30	
	1–4.9 yr	1.6 (1.0, 2.4)	22	
	≥5 yr	1.3 (0.6, 2.4)	10	
Biologically-monitored Danish workers				Hansen et al. (2001)
	Any TCE exposure	3.8 (1.0, 9.8)	4	
	Cumulative exposure (Ikeda)			
	<17 ppm-yr	2.9 (0.04, 16)	1	
	≥17 ppm-yr	2.6 (0.03, 14)	1	
	Mean concentration (Ikeda)			
	<4 ppm	3.4 (0.4, 12)	2	
	4+ ppm	4.3 (0.5, 16)	2	
	Employment duration			
	<6.25 yr	3.8 (0.1, 21)	1	
	≥6.25 yr	2.1 (0.03, 12)	1	
Aircraft maintenance workers from Hill Air Force Base, UT				Blair et al. (1998)
	TCE subcohort	Not reported		
	Cumulative exposure	Not reported		

4



**Table 4-92. Summary of human studies on TCE exposure and cervical cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Biologically-monitored Finnish workers				Anttila et al. (1995)
All subjects		2.42 (1.05, 4.77)	8	
Mean air-TCE (Ikeda extrapolation)				
<6 ppm		1.86 (0.38, 5.45)	3	
6+ ppm		4.35 (1.41, 10.1)	5	
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
Exposed workers		Not reported		
Biologically-monitored Swedish workers				Axelson et al. (1994)
Any TCE exposure		Not reported		
Cardboard manufacturing workers, Atlanta area, GA				Sinks et al. (1992)
All subjects		Not reported		
<b>Cohort studies-mortality</b>				
Aerospace workers (Rocketdyne)				
Any TCE (utility/eng flush)		Not reported		Boice et al. (2006a)
Any exposure to TCE		Not reported		Zhao et al. (2005)
View-Master employees				ATSDR (2004a)
Females		1.77 (0.57, 4.12) <sup>b</sup>	5	
United States uranium-processing workers (Fernald, OH)				Ritz (1999a)
Any TCE exposure		Not reported		
Light TCE exposure, >2 yr duration		Not reported		
Moderate TCE exposure, >2 yr duration		Not reported		
Aerospace workers (Lockheed)				Boice et al. (1999)
Routine exposure		-- (0.00, 5.47)	0	
Routine-intermittent		Not reported		
Aerospace workers (Hughes)				Morgan et al. (1998)
TCE subcohort		(0.00, 1.07)	0 (3.5 exp)	
Low intensity (<50 ppm)			0 (1.91 exp)	
High intensity (>50 ppm)			0 (1.54 exp)	

**Table 4-92. Summary of human studies on TCE exposure and cervical cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Aircraft maintenance workers (Hill AFB, UT)				
	TCE subcohort	1.8 (0.5, 6.5) <sup>c</sup>	5	Blair et al. (1998)
	Cumulative exposure			
	0	1.0 <sup>c</sup>		
	<5 ppm-yr	0.9 (0.1, 8.3)	1	
	5–25 ppm-yr		0	
	>25 ppm-yr	3.0 (0.8, 11.7)	4	
	TCE subcohort	1.67 (0.54, 5.22)	6	Radican et al. (2008)
	Cumulative exposure			
	0	1.0 <sup>c</sup>		
	<5 ppm-yr	0.76 (0.09, 6.35)	1	
	5–25 ppm-yr		0	
	>25 ppm-yr	2.83 (0.86, 9.33)	5	
Cardboard manufacturing workers in Arnsburg, Germany				
	TCE exposed workers	Not reported		Henschler et al. (1995)
	Unexposed workers	Not reported		
Deaths reported to GE pension fund (Pittsfield, MA)		Not examined <sup>d</sup>		Greenland et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA		Not reported		Sinks et al. (1992)
U. S. Coast Guard employees		Not reported <sup>e</sup>		Blair et al. (1989)
Aircraft manufacturing plant employees (Italy)		Not reported <sup>e</sup>		Costa et al. (1989)
Aircraft manufacturing plant employees (San Diego, CA)				
	All subjects	0.61 (0.25, 1.26) <sup>f</sup>	7	Garabrant et al. (1988)
Lamp manufacturing workers (GE)				
	Coil/wire drawing	1.05 (0.03, 5.86)	1	Shannon et al. (1988)
	Other areas	1.16 (0.32, 2.97)	4	
<b>Case-control studies</b>				
<b>Geographic based studies</b>				
Residents in two study areas in Endicott, NY		1.06 (0.29, 2.71)	<6	ATSDR (2006b)
Residents in Texas				Coyle et al. (2005)
	Counties reporting any air TCE release	66.4 <sup>g</sup>		
	Countries not reporting any air TCE release	60.8 <sup>g</sup>		
Residents of 13 census tracts in Redlands, CA		0.65 (0.38, 1.02)	29	Morgan and Cassady (2002)

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-92. Summary of human studies on TCE exposure and cervical cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Finnish residents				Vartiainen et al. (1993)
	Residents of Hausjarvi	Not reported		
	Residents of Huttula	Not reported		

<sup>a</sup>Standardized incidence ratio for females in Sung et al. (2007) reflects a 15-yr lag period.

<sup>b</sup>Proportional mortality ratio.

<sup>c</sup>Internal referents, workers not exposed to TCE.

<sup>d</sup>Nested case-control analysis.

<sup>e</sup>Males only in cohort.

<sup>f</sup>SMR is for cancer of the genital organs (cervix, uterus, endometrium, etc.).

<sup>g</sup> Median annual age-adjusted breast cancer rate (1995–2000).

GE = General Electric, No. obs. events = number of observed events.

(Anttila et al., 1995; Axelson et al., 1994; Blair et al., 1998; Boice et al., 1999; Boice et al., 2006a; Greenland et al., 1994; Hansen et al., 2001; Krishnadasan et al., 2007; Morgan et al., 1998, 2000b; Raaschou-Nielsen et al., 2003; Radican et al., 2008). Krishnadasan et al. (2007) in their nested case-control study of prostate cancer observed a twofold odds ratio estimate with high cumulative TCE exposure score (2.4, 95% CI: 1.3, 4.4, 20 year lagged exposure) and an increasing positive relationship between prostate cancer incidence and TCE cumulative exposure score ( $p = 0.02$ ). TCE exposure was positively correlated with several other occupational exposures, and Krishnadasan et al. (2007) adjusted for possible confounding from all other chemical exposures as well as age at diagnosis, occupational physical activity, and socio-economic status in statistical analyses. Relative risk estimates in studies other than Krishnadasan et al. (2007) were above 1.0 for overall TCE exposure (1.8, 95% CI: 0.8, 4.0 (Siemiatycki, 1991); 1.1, 95% CI: 0.6, 1.8 (Blair et al., 1998) and 1.20, 95% CI: 0.92, 1.76, with an additional 10-year follow-up (Radican et al., 2008); 1.58, 95% CI: 0.96, 2.62 (EHS, 1997; Morgan et al., 1998, 2000b); 1.3, 95% CI: 0.52, 2.69 (Boice et al., 1999); 1.38, 95% CI: 0.73, 2.35 (Anttila et al., 1995)) and prostate cancer risks did not appear to increase with increasing exposure. Four studies observed relative risk estimates below 1.0 for overall TCE exposure (0.93, 95% CI: 0.60, 1.37 (Garabrant et al., 1988); 0.6, 95% CI: 0.2, 1.30 (Hansen et al., 2001); 0.9, 95% CI: 0.79, 1.08 (Raaschou-Nielsen et al., 2003); 0.82, 95% CI: 0.36, 1.62 (Boice et al., 2006a)), and are not considered inconsistent because alternative explanations are possible and included observations are based on few subjects, lowering statistical power, or to poorer exposure assessment approaches that may result in a higher likelihood of exposure misclassification.

1 Seven other cohort, PMR, and geographic based studies were given less weight in the  
2 analysis because of their lesser likelihood of TCE exposure and other study design limitations  
3 that would decrease statistical power and study sensitivity (ATSDR, 2004a, 2006a; Blair et al.,  
4 1989; Chang et al., 2005; Morgan and Cassady, 2002; Shannon et al., 1988; Wilcosky et al.,  
5 1984). Chang et al. (2005) observed a statistically significant deficit in prostate cancer risk,  
6 based on one case, and an insensitive exposure assessment (0.14, 95% CI: 0.00, 0.76). Relative  
7 risks in the other five studies ranged from 0.62 (CI not presented in paper) (Wilcosky et al.,  
8 1984) to 1.11 (95% CI: 0.98, 1.25) (Morgan and Cassady, 2002).

9 Risk factors for prostate cancer include age, family history of prostate cancer, and  
10 ethnicity as causal with inadequate evidence for a relationship with smoking or alcohol (Wigle et  
11 al., 2008). All studies except Krishnadasan et al. (2007) were not able to adjust for possible  
12 confounding from other chemical exposures in the work environment. None of the studies  
13 including Krishnadasan et al. (2007) accounted for other well-established nonoccupational risk  
14 factors for prostate cancer such as race, prostate cancer screening and family history. There is  
15 limited evidence that physical activity may provide a protective effect for prostate cancer (Wigle  
16 et al., 2008). Krishnadasan et al. (2008) examined the effect of physical activity in the  
17 Rocketdyne aerospace cohort (Krishnadasan et al., 2007; Zhao et al., 2005). Their finding of a  
18 protective effect with high physical activity (0.55, 95% CI: 0.32, 0.95,  $p$  trend = 0.04) after  
19 control for TCE exposure provides additional evidence (Krishnadasan et al., 2008) and suggests  
20 underlying risk may be obscured in studies lacking adjustment for physical activity.  
21

#### 4.8.2.1.3. Breast cancer

22 Fifteen studies of TCE exposure reported findings on breast cancer in males and females  
23 combined (Boice et al., 1999; Garabrant et al., 1988; Greenland et al., 1994), in males and  
24 females, separately (ATSDR, 2004a; Clapp and Hoffman, 2008; Hansen et al., 2001; Raaschou-  
25 Nielsen et al., 2003), or in females only (ATSDR, 2006a; Blair et al., 1998; Chang et al., 2005;  
26 Coyle et al., 2005; Morgan et al., 1998; Radican et al., 2008; Shannon et al., 1988; Sung et al.,  
27 2007). Six studies have high likelihood of TCE exposure in individual study subjects and met, to  
28 a sufficient degree, the standards of epidemiologic design and analysis (Blair et al., 1998; Boice  
29 et al., 1999; Hansen et al., 2001; Morgan et al., 1998; Raaschou-Nielsen et al., 2003; Radican et  
30 al., 2008). Four studies with risk estimates for other cancer sites do not report risk estimates for  
31 breast cancer (Anttila et al., 1995; Axelson et al., 1994; Boice et al., 2006a; Siemiatycki, 1991).  
32 No case-control studies were found on TCE exposure, although several studies examine  
33 occupational title or organic solvent as a class (Band et al., 2000; Ji et al., 2008; Rennix et al.,  
34 2005; Weiderpass et al., 1999). While association is seen with occupational title or industry and

1 breast cancer (employment in aircraft and aircraft part industry, 2.48, 95% CI: 1.14, 5.39 (Band  
2 et al., 2000); solvent user: 1.48, 95% CI: 1.03, 2.12 (Rennix et al., 2005)), TCE exposure is not  
3 uniquely identified. The two studies suggest association between organic solvents and female  
4 breast cancer needs further investigation of possible risk factors.

5 Relative risk estimates in the five studies in which there is a high likelihood of TCE  
6 exposure in individual study subjects and which met, to a sufficient degree, the standards of  
7 epidemiologic design and analysis in a systematic review ranged from 0.75 (0.43, 1.22) (females  
8 and males; (Morgan et al., 1998)) to 2.0 (0.9, 4.6) (mortality in females; (Blair et al., 1998)).  
9 Blair et al. (1998), additionally, observed stronger risk estimates for breast cancer mortality  
10 among females with low level intermittent (3.1, 95% CI: 1.5, 6.2) and low level continuous (3.4,  
11 95% CI: 1.4, 8.0) TCE exposures, but not with frequent peaks (1.4, 95% CI: 0.7, 3.2). A similar  
12 pattern of risks was also observed by Radican et al. (2008) who studied mortality in this cohort  
13 and adding 10 years of follow-up, although the magnitude of breast cancer risk in females was  
14 lower than that observed in Blair et al. (1998). Risk estimates did not appear to increase with  
15 increasing cumulative exposure in the two studies that included exposure-response analyses  
16 (Blair et al., 1998; Morgan et al., 1998). None of these five studies reported a statistically  
17 significant deficit in breast cancer and confidence intervals on relative risks estimates included  
18 1.0 (no risk). Few female subjects in these studies appear to have high TCE exposure. For  
19 example, Blair et al. (1998) identified 8 of the 28 breast cancer deaths and 3 of the 34 breast  
20 cancer cases with high cumulative exposure.

21 Relative risk estimates in six studies of lower likelihood TCE exposure and other design  
22 deficiencies ranged from 0.81 (95% CI: 0.52, 1.48) (Garabrant et al., 1988) to 1.19 (1.03, 1.36)  
23 (Chang et al., 2005). These studies lack a quantitative surrogate for TCE exposure to individual  
24 subjects and instead classify all subjects as “potentially exposed,” with resulting large dilution of  
25 actual risk and decreased sensitivity (ATSDR, 2006a; Chang et al., 2005; Garabrant et al., 1988;  
26 Morgan and Cassady, 2002; NRC, 2006; Shannon et al., 1988; Sung et al., 2007).

27 Four studies reported on male breast cancer separately (ATSDR, 2004a; Clapp and  
28 Hoffman, 2008; Hansen et al., 2001; Raaschou-Nielsen et al., 2003) and a total of three cases  
29 were observed. Breast cancer in men is a rare disease and is best studied using a case-control  
30 approach (Weiss et al., 2005). Reports exist of male breast cancer among former residents of  
31 Camp Lejuene (ATSDR, 2007, 2010). Further assessment of TCE exposure and male breast  
32 cancer is warranted.

33 Overall, the epidemiologic studies on TCE exposure and breast cancer are quite limited in  
34 statistical power; observations are based on few breast cancer cases or on inferior TCE exposure  
35 assessment in studies with large numbers of observed cases. Additionally, adjustment for  
36 nonoccupational breast cancer risk factors is less likely in cohort and geographic based studies

1 given their use of employment and public records. Breast cancer mortality observations in Blair  
2 et al. (1998) and further follow-up of this cohort by Radican et al. (2008) of an elevated risk with  
3 overall TCE exposure, particularly low level intermittent and continuous TCE exposure, provide  
4 evidence of an association with TCE. No other study with high likelihood of TCE exposure in  
5 individual study subjects reported a statistically significant association with breast cancer,  
6 although few observed cases leading to lower statistical power or examination of risk for males  
7 and females combined are alternative explanations for the null observations in these studies.  
8 Both Chang et al. (2005) and Sung et al. (2007), two overlapping studies of female electronics  
9 workers exposed to TCE, perchloroethylene, and mixed solvents, reported association with  
10 breast cancer incidence, with breast cancer risk in Chang et al. (2005) appearing to increase with  
11 employment duration. Both studies, in addition to association provided by studies of exposure to  
12 broader category of organic solvents (Band et al., 2000; Rennix et al., 2005), support Blair et al.  
13 (1998) and Radican et al. (2008), although the lack of exposure assessment is an uncertainty.  
14 The epidemiologic evidence is limited for examining TCE and breast cancer, and while these  
15 studies do not provide any strong evidence for association with TCE exposure they in turn do not  
16 provide evidence of an absence of association.

17

#### **4.8.2.1.4. Cervical cancer**

18 Eleven cohort or PMR studies and 2 geographic based studies present relative risk  
19 estimates (Anttila et al., 1995; ATSDR, 2004a, 2006a; Blair et al., 1998; Boice et al., 1999;  
20 Garabrant et al., 1988; Hansen et al., 2001; Morgan and Cassady, 2002; Morgan et al., 1998;  
21 Raaschou-Nielsen et al., 2003; Radican et al., 2008; Shannon et al., 1988; Sung et al., 2007).  
22 Seven of these studies had high likelihood of TCE exposure in individual study subjects and  
23 were judged to have met, to a sufficient degree, the standards of epidemiologic design and  
24 analysis (Anttila et al., 1995; Blair et al., 1998; Boice et al., 1999; Hansen et al., 2001; Morgan et  
25 al., 1998; Raaschou-Nielsen et al., 2003; Radican et al., 2008). Three small cohort studies (Costa  
26 et al., 1989; Henschler et al., 1995; Sinks et al., 1992) as well as three studies with high  
27 likelihood of TCE exposure in individual study subjects (Axelson et al., 1994; Boice et al.,  
28 2006a; Zhao et al., 2005) did not present relative risk estimates for cervical cancer. Additionally,  
29 one population case-control and one geographic study of several site-specific cancers do not  
30 present information on cervical cancer (Siemiatycki, 1991; Vartiainen et al., 1993).

31 Five studies with high likelihood of TCE exposure in individual study subjects and which  
32 met, to a sufficient degree, the standards of epidemiologic design and analysis in a systematic  
33 review observed elevated risk for cervical cancer and overall TCE exposure (2.42, 95% CI: 1.05,

1 4.77 (Anttila et al., 1995); 1.8, 95% CI: 0.5, 6.5 (Blair et al., 1998) that changed little with an  
2 additional 10 years follow-up, 1.67, 95% CI: 0.54, 5.22 (Radican et al., 2008); 3.8, 95% CI: 1.42,  
3 2.37 (Hansen et al., 2001); 1.9, 95% CI: 1.42, 2.37 (Raaschou-Nielsen et al., 2003). The  
4 observations of a three- to fourfold elevated cervical cancer risk with high mean TCE exposure  
5 compared to subjects in the low exposure category (6+ ppm: 4.35, 95% CI: 1.41, 10.1 (Anttila et  
6 al., 1995); 4+ ppm: 4.3, 95% CI: 0.5, 16 (Hansen et al., 2001)) or with high cumulative TCE  
7 exposure (0.25-ppm year: 3.0, 95% CI: 0.8, 11.7 (Blair et al., 1998), 2.83, 95% CI: 0.86, 9.33  
8 (Radican et al., 2008)) provides additional support for association with TCE. Cervical cancer  
9 risk was lowest for subjects in the high exposure duration category (Hansen et al., 2001;  
10 Raaschou-Nielsen et al., 2003); however, duration of employment is a poor exposure metric  
11 given subjects may have differing exposure intensity with similar exposure duration (NRC,  
12 2006). No deaths due to cervical cancer were observed in two other studies (Boice et al., 1999;  
13 Morgan et al., 1998), less than 4 deaths were expected, suggesting these cohorts contained few  
14 female subjects with TCE exposure.

15 Human papilloma virus and low socioeconomic status are known risk factors for cervical  
16 cancer (ACS, 2008). Subjects in Raaschou-Nielsen et al. (2003) are blue-collar workers and low  
17 socioeconomic status likely explains observed associations in this and the other studies. The use  
18 of internal controls in Blair et al. (1998) who are similar in socioeconomic status as TCE subjects  
19 is believed to partly account for possible confounder related to socio-economic status; however,  
20 direct information on individual subjects is lacking.

21 Six other cohort, PMR, and geographic based studies were given less weight in the  
22 analysis because of their lesser likelihood of TCE exposure and other study design limitations  
23 that would decrease statistical power and study sensitivity (ATSDR, 2004a, 2006a; Garabrant et  
24 al., 1988; Morgan and Cassady, 2002; Shannon et al., 1988; Sung et al., 2007). Cervical cancer  
25 risk estimates in these studies ranged between 0.65 (95% CI: 0.38, 1.02) (Morgan and Cassady,  
26 2002) to 1.77 (proportional mortality ratio; 95% CI: 0.57, 4.12 (ATSDR, 2004a)). No study  
27 reported a statistically significant deficit in cervical cancer risk.

28

#### **4.8.2.1.5. Animal Studies**

29 Histopathology findings have been noted in reproductive organs in various cancer  
30 bioassay studies conducted with TCE. A number of these findings (summarized in Table 4-93)  
31 do not demonstrate a treatment-related profile.

1           Cancers of the reproductive system that are associated with TCE exposure and observed  
2 in animal studies are comprised of testicular tumors (interstitial cell and Leydig cell). A  
3 summary of the incidences of testicular tumors observed in male rats is presented in Table 4-94.  
4



**Table 4-93. Histopathology findings in reproductive organs**

<b>Tumor incidence in mice after 18 mo inhalation exposure<sup>a</sup></b>								
	<b>Tissue</b>	<b>Finding</b>	<b>Control</b>	<b>100 ppm</b>	<b>500 ppm</b>			
Males	Number examined:		30	29	30			
	Prostate	Myoma	1	0	0			
	Testis	Carcinoma	0	0	1			
		Cyst	0	0	1			
Females	No. examined:		29	30	28			
	Uterus	Adenocarcinoma	1	0	0			
	Ovary	Adenocarcinoma	1	0	0			
		Adenoma	3	1	3			
		Carcinoma	0	2	2			
		Granulosa cell tumor	4	0	2			
<b>Tumor incidence in rats after 18 mo inhalation exposure<sup>a</sup></b>								
	<b>Tissue</b>	<b>Finding</b>	<b>Control</b>	<b>100 ppm</b>	<b>500 ppm</b>			
Males	Number examined:		29	30	30			
	Testis	Interstitial cell tumors	4	0	3			
Females	No. examined:		28	30	30			
	Mammary	Fibroadenoma	2	0	0			
		Adenocarcinoma	3	2	2			
	Uterus	Adenocarcinoma	3	1	4			
	Ovary	Carcinoma	4	0	1			
		Granulosa cell tumor	1	0	0			
	Genital tract	Squamous cell carcinoma	0	2	0			
<b>Tumor incidence in hamsters after 18 mo inhalation exposure<sup>a</sup></b>								
	<b>Tissue</b>	<b>Finding</b>	<b>Control</b>	<b>100 ppm</b>	<b>500 ppm</b>			
Females	Number examined:		30	29	30			
	Ovary	Cystadenoma	1	0	0			
<b>Tumor incidence in mice after 18 mo gavage administration<sup>b</sup></b>								
	<b>Tissue</b>	<b>Finding</b>	<b>Con- trol</b>	<b>TCE Pure</b>	<b>TCE Industrial</b>	<b>TCE + EPC</b>	<b>TCE + BO</b>	<b>TCE + EPC + BO</b>
Females	Number examined:		50	50	50	50	48	50
	Mammary	Carcinoma	1	2	0	0	0	0
	Ovary	Granulosa cell tumor	0	1	0	0	0	0
	Vulva	Squamous cell carcinoma	0	0	0	0	1	1

2 <sup>a</sup>Henschler et al. (1980).3 <sup>b</sup>Henschler et al. (1984); EPC = epichlorohydrin; BO = 1,2-epoxybutane.

1 **Table 4-94. Testicular tumors in male rats exposed to TCE, adjusted for**  
 2 **reduced survival<sup>a</sup>**  
 3

<b>Interstitial cell tumors after 103 wk gavage exposure, beginning at 6.5–8 wk of age (NTP, 1988, 1990)</b>				
<b>Administered dose (mg/kg-day)</b>	<b>Untreated control</b>	<b>Vehicle control</b>	<b>500</b>	<b>1,000</b>
Male ACI rats	38/45 (84%)	36/44 (82%)	23/26 (88%)	17/19 (89%)
Male August rats	36/46 (78%)	34/46 (74%)	30/34 (88%)	26/30 (87%)
Male Marshall rats <sup>b</sup>	16/46 (35%)	17/46 (37%)	21/33 (64%)	32/39 (82%)
Male Osborne-Mendel rats	1/30 (3%)	0/28 (0%)	0/25 (0%)	1/19 (5%)
Male F344/N rats	44/47 (94%)	47/48 (98%)	47/48 (98%)	32/44 (73%)
<b>Leydig cell tumors after 104 wk inhalation exposure, beginning at 12 wk of age (Maltoni et al., 1986)</b>				
<b>Administered daily concentration (mg/m<sup>3</sup>)<sup>c</sup></b>	<b>Control</b>	<b>112.5</b>	<b>337.5</b>	<b>675</b>
Male Sprague-Dawley rats <sup>b</sup>	6/114 (5%)	16/105 (15%)	30/107 (28%)	31/113 (27%)

4  
 5 <sup>a</sup>ACI rats alive at Week 70, August rats at Week 65, Marshall rats at Week 32, Osborne-Mendel rats at Week 97,  
 6 F344/N rats at Week 32, Sprague-Dawley rats at Week 81 (except BT304) or Week 62 (except BT304 bis).

7 <sup>b</sup>Equivalent to 100, 300, or 600 ppm (100 ppm = 540 mg/m<sup>3</sup>), adjusted for 7 h/d, 5 d/wk exposure.

8 <sup>c</sup>Statistically significant by Cochran-Armitage trend test ( $p < 0.05$ ).

9  
 10 Sources: NTP (1988) Tables A2, C2, E2, G2; NTP (1990) Table A3; Maltoni et al. (1986) IV/IV Table 21, IV/V  
 11 Table 21.  
 12  
 13

#### 4.8.2.1.6. Mode of Action for Testicular Tumors

14 The database for TCE does not include an extensive characterization of the mode of  
 15 action for Leydig cell tumorigenesis in the rat, although data exist that are suggestive of  
 16 hormonal disruption in male rats. A study by Kumar et al. (2000a) found significant decreases in  
 17 serum testosterone concentration and in 17- $\beta$ -HSD, G6PDH, and total cholesterol and ascorbic  
 18 acid levels in testicular homogenate from male rats that had been exposed via inhalation to  
 19 376 ppm TCE for 12 or 24 weeks. In a follow-up study, Kumar et al. (2001b) also identified  
 20 decreases in SDH in the testes of TCE-treated rats. These changes are markers of disruption to  
 21 testosterone biosynthesis. Evidence of testicular atrophy, observed in the 2001 study by  
 22 Kumar et al., as well as the multiple in vivo and in vitro studies that observed alterations in  
 23 spermatogenesis and/or sperm function, could also be consistent with alterations in testosterone

1 levels. Therefore, while the available data are suggestive of a MOA involving hormonal  
2 disruption for TCE-induced testicular tumors, the evidence is inadequate to specify and test a  
3 hypothesized sequence of key events.

4       Leydig cell tumors can be chemically induced by alterations of steroid hormone levels,  
5 through mechanisms such as agonism of estrogen, gonadotropin releasing hormone, or dopamine  
6 receptors; antagonism of androgen receptors; and inhibition of 5 $\alpha$ -reductase, testosterone  
7 biosynthesis, or aromatase (Cook et al., 1999). For those plausible mechanisms that involve  
8 disruption of the hypothalamic-pituitary-testis (HPT) axis, decreased testosterone or estradiol  
9 levels or recognition is involved, and increased luteinizing hormone (LH) levels are commonly  
10 observed. Although there is evidence to suggest that humans are quantitatively less sensitive  
11 than rats in their proliferative response to LH, evidence of treatment-related Leydig cell tumors  
12 in rats that are induced via HPT disruption is considered to represent a potential risk to humans  
13 (with the possible exception of GnRh or dopamine agonists), since the pathways for regulation of  
14 the HPT axis are similar in rats and humans (Clegg et al., 1997).

### 4.8.3. Developmental Toxicity

16       An evaluation of the human and experimental animal data for developmental toxicity,  
17 considering the overall weight and strength of the evidence, suggests a potential for adverse  
18 outcomes associated with pre- and/or postnatal TCE exposures.

#### 4.8.3.1.1. Human Developmental Data

20       Epidemiological developmental studies (summarized in Table 4-95) examined the  
21 relationship between TCE exposure and prenatal developmental outcomes including spontaneous  
22 abortion and perinatal death; decreased birth weight, small for gestational age, and postnatal  
23 growth; congenital malformations; and other adverse birth outcomes. Postnatal developmental  
24 outcomes examined include developmental neurotoxicity, developmental immunotoxicity, other  
25 developmental outcomes, and childhood cancer.

#### **4.8.3.1.2. Adverse fetal/birth outcomes**

##### **4.8.3.1.2.1. Spontaneous abortion and perinatal death**

1           Spontaneous abortion or miscarriage is defined as nonmedically induced premature  
2 delivery of a fetus prior to 20 weeks gestation. Perinatal death is defined as stillbirths and deaths  
3 before 7 days after birth. Available data comes from several studies of occupational exposures in  
4 Finland and Santa Clara, California, and by geographic-based studies in areas with known  
5 contamination of water supplies in Woburn, MA; Tucson Valley, AZ; Rocky Mountain Arsenal,  
6 CO; Endicott, NY; and New Jersey.

7

**Table 4-95. Developmental studies in humans**

Subjects	Exposure	Effect	Reference
<b>Adverse fetal/birth outcomes</b>			
Spontaneous abortion and perinatal death			
371 men occupationally exposed to solvents in Finland 1973–1983	Questionnaire Low/rare used <1 d/wk; Intermediate used 1–4 d/wk or intermediate/low TCA urine levels; High/frequent used daily or high TCA urine levels	No risk of spontaneous abortion after paternal exposure, based on 17 cases and 35 controls exposed to TCE OR: 1.0, 95% CI: 0.6–2.0	Taskinen et al. (1989)
535 women occupationally exposed to solvents in Finland 1973–1986	Questionnaire Rare used 1–2 d/wk; Frequent used $\geq 3$ d/wk	Increased risk of spontaneous abortion among frequently-exposed women, based on 7 cases and 9 controls exposed to TCE OR: 1.6, 95% CI: 0.5–4.8	Taskinen et al. (1994)
3,265 women occupationally exposed to organic solvents in Finland 1973–1983	Questionnaire U-TCA: median: 48.1 $\mu\text{mol/L}$ ; mean 96.2 $\pm$ 19.2 $\mu\text{mol/L}$	No increased risk of spontaneous abortion based on 3 cases and 13 controls exposed to TCE OR: 0.6, 95% CI: 0.2–2.3	Lindbohm et al. (1990)
361 women occupationally and residentially exposed to solvents in Santa Clara County, CA June 1986–February 1987 (735 controls)	Questionnaire	Increased risk of spontaneous abortion based on 6 cases and 4 controls exposed to TCE <sup>a</sup> OR: 3.1, 95% CI: 0.92–10.4	Windham et al. (1991)
4,396 pregnancies among residents of Woburn, MA 1960–1982	TCE: 267 $\mu\text{g/L}$ Tetrachloroethylene: 21 $\mu\text{g/L}$ Chloroform: 12 $\mu\text{g/L}$	Increased risk of perinatal death ( $n = 67$ ) after 1970 ( $p = 0.55$ ) but not before 1970 (OR: 10, $p = 0.003$ ) No increased risk of spontaneous abortion ( $n = 520$ ; $p = 0.66$ )	Lagakos et al. (1986)
707 parents of children with congenital heart disease in Tucson Valley, AZ 1969–1987	6–239 ppb TCE, along with DCA and chromium	No increased risk of fetal death (not quantified) based on 246 exposed and 461 unexposed cases	Goldberg et al. (1990)
75 men and 71 women living near Rocky Mountain Arsenal, CO 1981–1986	Low: <5.0 ppb Medium: $\geq 5.0$ to <10.0 ppb High: <10.0 ppb	Increased risk of miscarriage OR <sub>adj</sub> : 4.44, 95% CI: 0.76–26.12 Increased risk of no live birth OR <sub>adj</sub> : 2.46, 95% CI: 0.24–24.95	ATSDR (2001)

**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
1,440 pregnancies among residents of Endicott, NY 1978–2002	indoor air from soil vapor: 0.18–140 mg/m <sup>3</sup>	No increase in spontaneous fetal death SIR: 0.66, 95% CI: 0.22–1.55	ATSDR (ATSDR, 2006b, 2008)
81,532 pregnancies among residents of 75 New Jersey towns 1985–1988 (3 control groups)	55 ppb TCE, along with many other compounds	No increased risk of fetal death for >10 ppb OR: 1.12	Bove (1996); Bove et al. (1995)
Decreased birth weight, small for gestational age, and postnatal growth			
361 women occupationally and residentially exposed to solvents in Santa Clara County, CA June 1986–February 1987 (735 controls)	Questionnaire	Increased risk of IUGR based on one case exposed to both TCE and tetrachloroethylene OR: 12.5	Windham et al. (1991)
3,462 births in Woburn, MA 1960–1982	267 µg/L TCE in drinking water, along with tetrachloroethylene and chloroform	No increase in low birth weight ( $p = 0.77$ )	Lagakos et al. (1986)
1,099 singleton births <sup>b</sup> to residents of 3 census tracts near Tucson International Airport 1979–1981 (877 controls)	<5–107 µg/L	No increase in full-term low birth weight (OR: 0.81) No increase in low birth weight (OR: 0.9) Increase in very low birth weight OR: 3.3, 95% CI: 0.53–20.6	Rodenbeck et al. (2000)
1,440 births <sup>c</sup> to residents of Endicott, NY 1978–2002	Indoor air from soil vapor: 0.18–140 mg/m <sup>3</sup>	Small increase in low birth weight OR: 1.26, 95% CI: 1.00–1.59 Small increase in small for gestational age OR: 1.22, 95% CI: 1.02–1.45 Increase in full-term low birth weight OR: 1.41, 95% CI: 1.01–1.95	ATSDR (ATSDR, 2006b, 2008)

**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
6,289 pregnancies among women residing at Camp Lejeune, NC 1968–1985 (141 short-term and 31 long-term TCE-exposed, 5,681 unexposed controls) <sup>d</sup>	Tarawa Terrace: TCE: 8 ppb 1,2-DCE: 12 ppb PCE: 215 ppb Hadnot Point: TCE: 1,400 ppb 1,2-DCE: 407 ppb	Change in mean birth weight Long-term total: -139 g, 90% CI: -277, -1 Long-term males: -312 g, 90% CI: -540, -85 Short term total: +70g, 90% CI: -6, 146 Increase in SGA Long-term total: OR: 1.5, 90% CI: 0.5, 3.8 Long-term males: OR: 3.9, 90% CI: 1.1–11.9 Short term total: OR: 1.1, 90% CI: 0.2–1.1	ATSDR (1998b)
81,532 pregnancies <sup>e</sup> among residents of 75 New Jersey towns 1985–1988	55 ppb TCE, along with many other compounds	Decreased birth weight at >5 ppb by 17.9g No increase in prematurity at >10 ppb: OR: 1.02 Increase in low birth weight, term >10 ppb: OR: 1.23, 50% CI: 1.09–1.39 No risk for very low birth weight	Bove (1996); Bove et al. (1995)
Congenital malformations			
1,148 men and 969 women occupationally exposed to TCE in Finland 1963–1976	U-TCA: <10 to >500 mg/L	No congenital malformations reported	Tola et al. (1980)
371 men occupationally exposed to solvents in Finland 1973–1983	Low/rare used <1 d/wk; Intermediate used 1–4 d/wk or if biological measures indicated high exposure; High/frequent used daily or if biological measures indicated high exposure	No increase in congenital malformations based on 17 cases and 35 controls exposed to TCE OR: 0.6, 95% CI: 0.2–2.0	Taskinen et al. (1989)
100 babies with oral cleft defects born to women occupationally exposed in Europe 1989–1992	Questionnaire	Increase in cleft lip based on 2 of 4 TCE-exposed women OR <sub>adj</sub> : 3.21, 95% CI: 0.49–20.9 Increase in cleft palate based on 2 of 4 TCE-exposed women OR <sub>adj</sub> : 4.47, 95% CI: 1.02–40.9	Lorente et al. (2000)

**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
4,396 pregnancies among residents of Woburn, MA 1960–1982	TCE: 267 µg/L Tetrachloroethylene: 21 µg/L Chloroform: 12 µg/L	Increase in eye/ear birth anomalies: OR: 14.9, $p < 0.0001$ Increase in CNS/chromosomal/oral cleft anomalies: OR: 4.5, $p = 0.01$ Increase in kidney/urinary tract disorders: OR: 1.35, $p = 0.02$ Small increase in lung/respiratory tract disorders: OR: 1.16, $p = 0.05$ No increase in cardiovascular anomalies ( $n = 5$ ): $p = 0.91$	Lagakos et al. (1986)
707 children with congenital heart disease in Tucson Valley, AZ 1969–1987 (246 exposed, 461 unexposed)	Wells contaminated with TCE (range: 6–239 ppb), along with DCA and chromium	Increase in congenital heart disease <1981: OR: $\approx 3$ ( $p < 0.005$ ) >1981: OR: $\approx 1$ Increased prevalence after maternal exposure during first trimester ( $p < 0.001$ , 95% CI: 1.14–4.14)	Goldberg et al. (1990)
75 men, 71 women living near Rocky Mountain Arsenal, CO 1981–1986	Low: <5.0 ppb Medium: $\geq 5.0$ to <10.0 ppb High: <10.0 ppb	Increase in total birth defects ( $n = 9$ ) OR: 5.87, 95% CI: 0.59–58.81	ATSDR (2001)
Births to residents of Endicott, NY 1983–2000 <sup>f</sup>	Indoor air from soil vapor: 0.18–140 mg/m <sup>3</sup>	No increase in total birth defects: RR: 1.08, 95% CI: 0.82–1.42 Increase in total cardiac defects: RR: 1.94, 95% CI: 1.21–3.12 Increase in major cardiac defects: RR: 2.52, 95% CI: 1.2–5.29 Increase in conotruncal heart defects: RR: 4.83, 95% CI: 1.81–12.89	ATSDR (ATSDR, 2006b, 2008)



**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
81,532 pregnancies among residents of 75 New Jersey towns 1985–1988	55 ppb TCE, along with many other compounds	No increase in total birth defects: >10 ppb: OR: 1.12 Increase in total CNS defects at high dose >1–5 ppb: OR: 0.93, 90% CI: 0.47–1.77 >10 ppb: OR: 1.68, 90% CI: 0.76–3.52 Increase in neural tube defects >1–5 ppb: OR: 1.58, 90% CI: 0.69–3.40 >10 ppb: OR: 2.53, 90% CI: 0.91–6.37 Increase in oral clefts: >5 ppb: OR: 2.24, 95% CI: 1.16–4.20 Increase in major cardiac defects: >10 ppb: OR: 1.24, 50% CI: 0.75–1.94 Increase in ventricular septal defects >5ppb: OR: 1.30, 95% CI: 0.88–1.87	Bove (1996); Bove et al. (1995)
1,623 children <20 yr old dying from congenital anomalies in Maricopa County, AZ 1966–1986	8.9 and 29 ppb TCE in drinking water	Increase in deaths due to congenital anomalies in East Central Phoenix 1966–1969: RR: 1.4, 95% CI: 1.1–1.7 1970–1981: RR: 1.5, 95% CI: 1.3–1.7 1982–1986: RR: 2.0, 95% CI: 1.5–2.5	AZ DHS (Flood, 1988)
4,025 infants born with congenital heart defects in Milwaukee, WI 1997–1999	Maternal residence within 1.32 miles from at least one TCE emissions source	Increase in congenital heart defects for mothers ≥38 yr old Exposed: OR: 6.2, 95% CI: 2.6–14.5 Unexposed: OR: 1.9, 95% CI: 1.1–3.5 No increase in congenital heart defects for exposed mothers <38 yr old: OR: 0.9, 95% CI: 0.6–1.2	Yauck et al. (2004)
12 children exposed to TCE in well water in Michigan	5–10 yr to 8–14 ppm	1 born with multiple birth defects	Bernad et al. (1987), abstract
Other adverse birth outcomes			
34 live births for which inhalation of TCE for anesthesia was used in Japan 1962–1967	2–8 mL (mean 4.3 mL) for 2–98 min (mean: 34.7 min)	1 case of asphyxia; 3 “sleepy babies” with Apgar scores of 5–9. Delayed appearance of newborn reflexes	Beppu (1968)

**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
51 UK women whose fetus was considered to be at risk for hypoxia during labor administered TCE as an analgesic (50 controls)	Amount and route of exposure not reported	TCE caused fetal pH to fall more, base deficit increased more, and PO <sub>2</sub> fell more than the control group by fourfold or more compared to other analgesics used	Phillips and Macdonald (1971)
<b>Postnatal developmental outcomes</b>			
Developmental neurotoxicity			
54 individuals from 3 residential cohorts in the United States exposed to TCE in drinking water	Woburn, MA 63–400 ppb for <1–12 yr Alpha, OH 3.3–330 ppb for 5–17 yr Twin Cities, MN 261–2,440 ppb for 0.25–25 yr	Woburn, MA Verbal naming/language impairment in 6/13 children (46%) Alpha, OH Verbal naming/language impairment in 1/2 children (50%) Twin Cities, MN Verbal naming/language impairment in 4/4 children (100%) Memory impairment in 4/4 children (100%) Academic impairment in 4/4 children (100%) Moderate encephalopathy in 4/4 children (100%) Poor performance on reading/spelling test in 3/4 children (75%) Poor performance on information test in 3/4 children (75%)	White et al. (1997)
284 cases of ASD diagnosed <9 yr old and 657 controls born in the San Francisco Bay Area 1994	Births geocoded to census tracts, and linked to HAPs data	Increase in ASD upper 3 <sup>rd</sup> quartile: OR: 1.37, 95% CI: 0.96–1.95 upper 4 <sup>th</sup> quartile: OR: 1.47, 95% CI: 1.03–2.08	Windham et al. (2006)
948 children (<18 yr) in the trichloroethylene Subregistry	0.4 to >5,000 ppb TCE	Increase in speech impairment: 0–9 yr old: RR: 2.45, 99% CI: 1.31–4.58 10–17 yr old: RR: 1.14, 99% CI: 0.46–2.85 Increase in hearing impairment: 0–9 yr old: RR: 2.13, 99% CI: 1.12–4.07 10–17 yr old: RR: 1.12, 99% CI: 0.52–2.24	ATSDR (2003b); Burg et al. (1995); Burg and Gist (1999)
12 children exposed to TCE in well water in Michigan	5–10 yr to 8–14 ppm	9 of 12 children (75%) had poor learning ability, aggressive behavior, and low attention span	Bernad et al. (1987), abstract

**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
Developmental immunotoxicity			
200 children aged 36 mo old born prematurely <sup>g</sup> and at risk of atopy <sup>h</sup> in Leipzig, Germany 1995–1996	Median air level in child's bedroom: 0.42 µg/m <sup>3</sup>	No association with allergic sensitization to egg white and milk, or to cytokine producing peripheral T-cells	Lehmann et al. (2001)
85 healthy <sup>i</sup> full-term neonates born in Leipzig, Germany 1997–1999	Median air level in child's bedroom 3–4 wk after birth: 0.6 µg/m <sup>3</sup>	Significant reduction of Th1 IL-2 producing T-cells	Lehmann et al. (2002)
Other developmental outcomes			
55 children (6 mo to 10 yr old) were anesthetized for operations to repair developmental defects of the jaw and face in Poland 1964	≥10 mL TCE	Reports of bradycardia, accelerated heart rate, and respiratory acceleration observed; no arrhythmia was observed	Jasinka (1965), translation
Childhood cancer			
98 children (<10 yr old) diagnosed with brain tumors in Los Angeles County 1972–1977	Questionnaire of parental occupational exposures	Two cases were reported for TCE exposure, one with methyl ethyl ketone	Peters and Preston-Martin (1981)
22 children (<19 yr old) diagnosed with neuroblastoma in United States and Canada 1992–1994 (12 controls)	Questionnaire of parental occupational exposures	Increase in neuroblastoma after paternal exposure OR: 1.4, 95% CI: 0.7–2.9 Maternal exposure not reported	De Roos et al. (2001)
61 boys and 62 girls (<10 yr old) diagnosed with leukemia and 123 controls in Los Angeles County 1980–1984	Questionnaire of parents for occupational exposure	Increase in leukemia after paternal exposure Preconception (1 yr): OR: 2.0, <i>p</i> = 0.16 Prenatal: OR: 2.0, <i>p</i> = 0.16 Postnatal: OR: 2.7, <i>p</i> = 0.7 Maternal exposure not reported	Lowengart et al. (1987)

**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
1,842 children (<15 yr old) diagnosed with ALL in United States and Canada 1989–1993 (1986 controls)	Questionnaire of parents for occupational exposure	Increase in ALL after maternal exposure Preconception: OR: 1.8, 95% CI: 0.6–5.2 Pregnancy: OR: 1.8, 95% CI: 0.5–6.4 Postnatal: OR: 1.4, 95% CI: 0.5–4.1 Anytime: OR: 1.8, 95% CI: 0.8–4.1 No increase in ALL after paternal exposure Anytime: OR: 1.1, 95% CI: 0.8–1.5	Shu et al. (1999)
109 children (<15 yr old) born in UK 1974–1988 (218 controls)	Questionnaire of parents for occupational exposure	Increase in leukemia and NHL after paternal exposure Preconception: OR: 2.27, 95% CI: 0.84–6.16 Prenatal: OR: 4.40, 95% CI: 1.15–21.01 Postnatal: OR: 2.66, 95% CI: 0.82–9.19 No increase in leukemia and NHL after maternal exposure Preconception: OR: 1.16, 95% CI: 0.13–7.91	McKinney et al. (1991)
22 children (<15 yr old) diagnosed with childhood cancer in California 1988–1998	0.09–97 ppb TCE in drinking water	No increase in total cancer: SIR: 0.83, 99% CI: 0.44–1.40 No increase in CNS cancer: SIR: 1.05, 99% CI: 0.24–2.70 No increase in leukemia: SIR: 1.09, 99% CI: 0.38–2.31	Morgan and Cassady (2002)
1,190 children (<20 yr old) diagnosed with leukemia in 4 counties in New Jersey 1979–1987	0–67 ppb TCE in drinking water	Increase in ALL in girls with >5 ppb exposure <20 yr old: RR: 3.36, 95% CI: 1.29–8.28 <5 yr old: RR: 4.54, 95% CI: 1.47–10.6	Cohn et al. (1994b)
24 children (<15 yr old) diagnosed with leukemia in Woburn, MA 1969–1997	267 µg/L TCE in drinking water, along with tetrachloroethylene, arsenic, and chloroform	Increase in childhood leukemia Preconception: OR <sub>adj</sub> : 2.61, 95% CI: 0.47–14.97 Pregnancy: OR <sub>adj</sub> : 8.33, 95% CI: 0.73–94.67 Postnatal: OR <sub>adj</sub> : 1.18, 95% CI: 0.28–5.05 Ever: OR <sub>adj</sub> : 2.39, 95% CI: 0.54–10.59	Costas et al. (2002); Cutler et al. (1986); Lagakos et al. (1986); MA DPH (1997a) <sup>1</sup>
347 children (<20 yr old) diagnosed with cancer in Endicott, NY 1980–2001	indoor air from soil vapor: 0.18–140 mg/m <sup>3</sup>	No increase in cancer (<6 cases, similar to expected)	ATSDR (ATSDR, 2006b, 2008)

**Table 4-95. Developmental studies in humans (continued)**

Subjects	Exposure	Effect	Reference
189 children (<20 yr old) diagnosed with cancer in Maricopa County, AZ 1965–1990	8.9 and 29 ppb TCE in drinking water	Increase in leukemia: 1965–1986: SIR: 1.67, 95% CI: 1.20–2.27 1982–1986: SIR: 1.91, 95% CI: 1.11–3.12 No increase in total childhood cancers, lymphoma, brain/CNS, or other cancers	AZ DHS (Flood, 1988; Flood, 1997) (1990) <sup>k</sup>
16 children (<20 yr old) diagnosed with cancer in East Phoenix, AZ 1965–1986	TCE, TCA, and other contaminants in drinking water	No increase in leukemia: SIR: 0.85, 95% CI: 0.50–1.35	AZ DHS (Kioski et al., 1990b)
37 children (<20 yr old) diagnosed with cancer in Pima County, AZ 1970–1986	1.1–239 ppb TCE, along with 1,1-DCE, chloroform and chromium in drinking water	Increase in leukemia ( $n = 11$ ): SIR: 1.50, 95% CI: 0.76–2.70 No increase in testicular cancer ( $n = 6$ ): SIR: 0.78, 95% CI: 0.32–1.59 No increase in lymphoma ( $n = 2$ ): SIR: 0.63, 95% CI: 0.13–1.80 No increase in CNS/brain cancer ( $n = 3$ ): SIR: 0.84, 95% CI: 0.23–2.16 Increase in other cancer ( $n = 15$ ): SIR: 1.40, 95% CI: 0.79–2.30	AZ DHS (Kioski et al., 1990a)

<sup>a</sup>Of those exposed to TCE, four were also exposed to tetrachloroethylene and one was also exposed to paint strippers and thinners.

<sup>b</sup>Full term defined as between 35 and 46 wk gestation, low birth weight as <2,501 g, and very low birth weight as <1,501 g.

<sup>c</sup>Low birth weight defined as <2,500, moderately low birth weight (1,500–<2,500 g), term low birth weight ( $\geq 37$  wk gestation and <25,000 g).

<sup>d</sup>Unexposed residents resided at locations not classified for long-term or short-term TCE exposure. Long-term TCE exposed mothers resided at Hospital Point during 1968–1985 for at least 1 wk prior to birth. Short-term TCE exposed mothers resided at Berkeley Manor, Midway Park, Paradise Point, and Wakins Village at the time of birth and at least 1 wk during January 27 to February 7, 1985. In addition, the mother's last menstrual period occurred on or before January 31, 1985 and the birth occurred after February 2, 1985.

<sup>e</sup>Low birth weight defined as <2,500 g, very low birth weight as <1,500 g.

<sup>f</sup>1,440 births reported for years 1978–2002, but number not reported for years 1983–2000.

<sup>g</sup>Premature defined as 1,500–2,500 g at birth.

<sup>h</sup>Risk of atopy defined as cord blood IgE >0.9 kU/L; double positive family atopy history.

<sup>i</sup>Healthy birth defined as  $\geq 2,500$  g and  $\geq 37$  wk gestation.

<sup>j</sup>Only results from Costas et al. (2002) are reported in the table.

<sup>k</sup>Only results from AZ DHS (1990) are reported in the table.

ALL = acute lymphoblastic leukemia, ASD = autism spectrum disorder, DCE = dichloroethylene, HAP = hazardous air pollutant, IUGR = intrauterine growth restriction, PCE = perchloroethylene, UK = United Kingdom.

#### 1 4.8.3.1.2.1.1. *Occupational studies*

2 The risks of spontaneous abortion and congenital malformations among offspring of men  
3 occupationally exposed to TCE and other organic solvents were examined by Taskinen et al.  
4 (1989). This nested case-control study was conducted in Finland from 1973–1983. Exposure  
5 was determined by biological measurements of the father and questionnaires answered by both  
6 the mother and father. The level of exposure was classified as “low/rare” if the chemical was  
7 used <1 days/week, “intermediate” if used 1–4 days/week or if TCA urine measurements  
8 indicated intermediate/low exposure, and “high/frequent” if used daily or if TCA urine  
9 measurements indicated clear occupational exposure (defined as above the RfV for the general  
10 population). There was no risk of spontaneous abortion from paternal TCE exposure (OR: 1.0,  
11 95% CI: 0.6–2.0), although there was a significant increase for paternal organic solvent exposure  
12 (OR: 2.7, 95% CI: 1.3–5.6) and a nonsignificant increase for maternal organic solvent exposure  
13 (OR: 1.4, 95% CI: 0.6–3.0). (Also see section below for results from this study for congenital  
14 malformations).

15 Another case-control study in Finland examined pregnancy outcomes in 1973–1986  
16 among female laboratory technicians aged 20–34 years (Taskinen et al., 1994). Exposure was  
17 reported via questionnaire, and was classified as “rare” if the chemical was used 1–2 days/week,  
18 and “frequent” if used at least 3 days/week. Cases of spontaneous abortion ( $n = 206$ ) were  
19 compared with controls who had delivered a baby and did not report prior spontaneous abortions  
20 ( $n = 329$ ). A nonstatistically significant increased risk was seen between spontaneous abortion  
21 and TCE use at least 3-days-a-week (OR: 1.6, 95% CI: 0.5–4.8).

22 The association between maternal exposure to organic solvents and spontaneous abortion  
23 was examined in Finland for births 1973–1983 (Lindbohm et al., 1990). Exposure was assessed  
24 by questionnaire and confirmed with employment records, and the level of exposure was either  
25 high, low or none based on the frequency of use and known information about typical levels of  
26 exposure for job type. Biological measurements of trichloroacetic acid in urine were also taken  
27 on 64 women, with a median value of 48.1  $\mu\text{mol/L}$  (mean:  $96.2 \pm 19.2 \mu\text{mol/L}$ ). Three cases and  
28 13 controls were exposed to TCE, with no increased risk seen for spontaneous abortion (OR: 0.6,  
29 95% CI: 0.2–2.3,  $p = 0.45$ ).

30 A case-control study in Santa Clara County, California, examined the association  
31 between solvents and adverse pregnancy outcomes in women  $\geq 18$  years old (Windham et al.,  
32 1991). For pregnancies occurring between June 1986 and February 1987, 361 cases of  
33 spontaneous abortion were compared to 735 women who had a live birth during this time period.  
34 Telephone interviews included detailed questions on occupational solvent exposure, as well as  
35 additional questions on residential solvent use. For TCE exposure, six cases of spontaneous

1 abortion were compared to four controls of live births; of these ten TCE-exposed individuals,  
2 four reported exposure to tetrachloroethylene, and one reported exposure to paint strippers and  
3 thinners. An increased risk of spontaneous abortions was seen with TCE exposure (OR: 3.1,  
4 95% CI: 0.92–10.4), with a statistically significant increased risk for those exposed  
5  $\geq 0.5$  hours/week (OR: 7.7, 95% CI: 1.3–47.4). An increased risk for spontaneous abortion was  
6 also seen for those reporting a more “intense” exposure based primarily on odor, as well as skin  
7 contact or other symptoms (OR: 3.9,  $p = 0.04$ ). (Also see section below from this study on low  
8 birth weight.)  
9

#### 10 **4.8.3.1.2.1.2. Geographic-based studies**

11 A community in Woburn, MA with contaminated well water experienced an increased  
12 incidence of adverse birth outcomes and childhood leukemia (Lagakos et al., 1986). In 1979, the  
13 wells supplying drinking water were found to be contaminated with 267 ppb TCE, 21 ppb  
14 tetrachloroethylene, 11.8 ppb, and 12 ppb chloroform, and were subsequently closed. Pregnancy  
15 and childhood outcomes were examined from 4,396 pregnancies among residents (Lagakos et  
16 al., 1986). No association between water access and incidence of spontaneous abortion ( $n = 520$ )  
17 was observed ( $p = 0.66$ ). The town’s water distribution system was divided into five zones,  
18 which was reorganized in 1970. Prior to 1970, no association was observed between water  
19 access and incidence of perinatal deaths ( $n = 46$  still births and 21 deaths before 7 days)  
20 ( $p = 0.55$ ). However, after 1970, a statistically significant positive association between access to  
21 contaminated water and perinatal deaths was observed (OR: 10.0,  $p = 0.003$ ). The authors could  
22 not explain why this discrepancy was observed, but speculated that contaminants were either not  
23 present prior to 1970, or were increased after 1970. (Also see sections below on decreased birth  
24 weight, congenital malformations, and childhood cancer for additional results from this cohort.)

25 A community in Tucson Valley, AZ with contaminated well water had a number of  
26 reported cases of congenital heart disease. The wells were found to be contaminated with TCE  
27 (range = 6–239 ppb), along with dichloroethylene and chromium (Goldberg et al., 1990). This  
28 study identified 707 children born with congenital heart disease during the years 1969–1987. Of  
29 the study participants, 246 families had parental residential and occupational exposure during  
30 one month prior to conception and during the first trimester of pregnancy, and 461 families had  
31 no exposure before the end of the first trimester. In addition to this control group, two others  
32 were used: (1) those that had contact with the contaminated water area, and (2) those that had  
33 contact with the contaminated water area and matched with cases for education, ethnicity, and  
34 occupation. Among these cases of congenital heart disease, no significant difference was seen

1 for fetal death (not quantified) for exposed cases compared to unexposed cases. (Also see  
2 section below on congenital malformations for additional results from this cohort.)

3 A residential study of individuals living near the Rocky Mountain Arsenal in Colorado  
4 examined the outcomes in offspring of 75 men and 71 women exposed to TCE in drinking water  
5 (ATSDR, 2001). TCE exposure was stratified by high (>10.0 ppb), medium ( $\geq$ 5.0 ppm to  
6 <10.0 ppb), and low (<5.0 ppb). Among women with >5 ppb exposure experiencing miscarriage  
7 ( $n = 22/57$ ) compared to unexposed women experiencing miscarriage ( $n = 2/13$ ) an elevated  
8 nonsignificant association was observed (OR<sub>adj</sub>: 4.44, 95% CI: 0.76–26.12). For lifetime number  
9 of miscarriages reported by men and women, results were increased but without dose-response  
10 for women (medium: OR<sub>adj</sub>: 8.56, 95% CI: 0.69–105.99; high: OR<sub>adj</sub>: 4.16, 95% CI: 0.61–25.99),  
11 but less for men (medium: OR<sub>adj</sub>: 1.68, 95% CI: 0.26–10.77; high: OR<sub>adj</sub>: 0.65,  
12 95% CI: 0.12–3.48). Among women with >5 ppb exposure experiencing no live birth ( $n = 9/57$ )  
13 compared to unexposed women experiencing no live birth ( $n = 1/13$ ) an elevated nonsignificant  
14 association was observed (OR<sub>adj</sub>: 2.46, 95% CI: 0.24–24.95). (Also see below for results from  
15 this study on birth defects.)

16 NYS DOH and ATSDR conducted a study in Endicott, NY to examine childhood cancer  
17 and birth outcomes in an area contaminated by a number of volatile organic compounds (VOCs),  
18 including “thousands of gallons” of TCE (ATSDR, 2006a). Soil vapor levels tested ranged from  
19 0.18–140 mg/m<sup>3</sup> in indoor air. A follow-up study by ATSDR (2008) reported that during the  
20 years 1978–1993 only five spontaneous fetal deaths occurring  $\geq$ 20 weeks gestation were  
21 reported when 7.5 were expected (SIR: 0.66, 95% CI: 0.22–1.55). (See sections on low birth  
22 weight, congenital malformations, and childhood cancer for additional results from this cohort.)

23 Women were exposed to contaminated drinking water while pregnant and living in 75  
24 New Jersey towns during the years 1985–1988 (Bove, 1996; Bove et al., 1995). The water  
25 contained multiple trihalomethanes, including an average of 55 ppb TCE, along with  
26 tetrachloroethylene, 1,1,1-trichloroethane, carbon tetrachloride, 1,2-dichloroethane, and benzene.  
27 A number of birth outcomes were examined for 81,532 pregnancies, which resulted in  
28 80,938 live births and 594 fetal deaths. No association was seen for exposure to >10 ppb TCE  
29 and fetal death (OR<sub>adj</sub>: 1.12). (See below for results from this study on decreased birth weight  
30 and congenital malformations.)

#### 4.8.3.1.2.2. Decreased birth weight, small for gestational age, and postnatal growth

32 Available data pertaining to birth weight and other growth-related outcomes come from  
33 the case-control study in Santa Clara, CA (discussed above), and by geographic-based studies as



1 well as geographic areas with known contamination of water supplies areas in Woburn, MA;  
2 Tucson, AZ, Endicott, NY; Camp Lejeune, NC; and New Jersey.

#### 4 **4.8.3.1.2.2.1. Occupational studies**

5 The case-control study of the relationship between solvents and adverse pregnancy  
6 outcomes discussed above (Windham et al., 1991) also examined intrauterine growth restriction  
7 (IUGR). Telephone interviews included detailed questions on occupational solvent exposure, as  
8 well as additional questions on residential solvent use. An increased risk of IUGR was observed  
9 (OR: 12.5), although this was based only on one case that was exposed to both TCE and  
10 tetrachloroethylene (also see section above on spontaneous abortion).

#### 12 **4.8.3.1.2.2.2. Geographic-based studies**

13 The study of Woburn, MA with contaminated well water discussed above (Lagakos et al.,  
14 1986) examined birth weight. Of 3,462 live births surviving to 7 days, 220 were less than  
15 6 pounds at birth (6.4%). No association was observed between water access and low birth  
16 weight ( $p = 0.77$ ). (See section on spontaneous abortion for study details, and see sections on  
17 spontaneous abortion, congenital malformations, and childhood cancer for additional results  
18 from this cohort.)

19 An ecological analysis of well water contaminated with TCE in Tucson and birth-weight  
20 was conducted by Rodenbeck et al. (2000). The source of the exposure was a U.S. Air Force  
21 plant and the Tucson International Airport. The wells were taken out of service in 1981 after  
22 concentrations of TCE were measured in the range of  $<5 \mu\text{g/L}$  to  $107 \mu\text{g/L}$ . The study  
23 population consisted of 1,099 babies born within census tracts between 1979 and 1981, and the  
24 comparison population consisted of 877 babies from nearby unexposed census tracts. There was  
25 a nonsignificant increased risk for maternal exposure to TCE in drinking water and very-low-  
26 birth-weight ( $<1,501 \text{ g}$ ) (OR: 3.3, 95% CI: 0.53–20.6). No increases were observed in the low-  
27 birth-weight ( $<2,501 \text{ g}$ ) (OR: 0.9) or full-term ( $>35$ -week and  $<46$ -week gestation) low-birth-  
28 weight (OR: 0.81).

29 The study of VOC exposure in Endicott, NY reported data on low birth weight and small  
30 for gestational age (ATSDR, 2006a), see section on spontaneous abortion for study details). For  
31 births occurring during the years 1978–2002, low birth weight was slightly but statistically  
32 elevated (OR: 1.26, 95% CI: 1.00–1.59), as was small for gestational age (SGA; OR: 1.22,  
33 95% CI: 1.02–1.45), and full-term low birth weight (OR: 1.41, 95% CI: 1.01–1.95). (Also see

1 sections on spontaneous abortion, congenital malformations, and childhood cancer for additional  
2 results from this cohort.)

3 Well water at the U.S. Marine Corps Base in Camp Lejeune, NC was identified to be  
4 contaminated with TCE, tetrachloroethylene, and 1,2-dichloroethane in April, 1982 and the wells  
5 were closed in December, 1984. ATSDR examined pregnancy outcomes among women living  
6 on the base during the years 1968–1985 (ATSDR, 1998b). Compared to unexposed residents<sup>12</sup>  
7 ( $n = 5,681$ ), babies exposed to TCE long-term<sup>13</sup> ( $n = 31$ ) had a lower mean birth weight after  
8 adjustment for gestational age ( $-139$  g, 90% CL =  $-277, -1$ ), and babies exposed short-term<sup>14</sup>  
9 ( $n = 141$ ) had a slightly higher mean birth weight ( $+70$  g, 90% CL =  $-6, 146$ ). For the long-term  
10 group, no effect was seen for very low birth weight ( $<1,500$  g) or prematurity ( $>5$  ppb,  
11 OR: 1.05). No preterm births were reported in the long-term group and those ( $n = 8$ ) in the  
12 short-term group did not have an increased risk (OR: 0.7, 90% CI: 0.3–1.2). A higher  
13 prevalence of SGA<sup>15</sup> was seen in the long-term exposed group ( $n = 3$ ; OR 1.5, 90% CL: 0.5, 3.8)  
14 compared to the short-term exposed group (OR: 1.1, 90% CI: 0.2–1.1). When the long-term  
15 group was stratified by gender, male offspring were at more risk for both reduced birth weight  
16 ( $-312$  g, 90% CL =  $-632, -102$ ) and SGA (OR: 3.9, 90% CL: 1.1–11.8). This study is limited  
17 due the mixture of chemicals in the water, as well as its small sample size. ATSDR is currently  
18 reanalyzing the findings because of an error in the exposure assessment related to the start-up  
19 date of a water treatment plant (ATSDR, 2007, 2009; U.S. GAO, 2007)

20 Pregnancy outcomes among women were exposed to contaminated drinking water while  
21 pregnant and living in 75 New Jersey towns during the years 1985–1988 was examined by  
22 Bove et al. (Bove, 1996; Bove et al., 1995). The water contained multiple trihalomethanes,  
23 including an average of 55 ppb TCE, along with tetrachloroethylene, 1,1,1-trichloroethane,  
24 carbon tetrachloride, 1,2-dichloroethane, and benzene. A number of birth outcomes were  
25 examined for 81,532 pregnancies, which resulted in 80,938 live births and 594 fetal deaths. A  
26 slight decrease of 17.9 g in birth weight was seen for exposure  $>5$  ppb, with a slight increase in  
27 risk for exposure  $>10$  ppb (OR: 1.23), but no effect was seen for very low birth weight or  
28 SGA/prematurity ( $>5$  ppb, OR: 1.05). However, due to the multiple contaminants in the water, it  
29 is difficult to attribute the results solely to TCE exposure. (See below for results from this study  
30 on congenital malformations.)  
31

---

12Unexposed residents resided at locations not classified for long-term or short-term TCE exposure.

13Long-term TCE exposed mothers resided at Hospital Point during 1968-1985 for at least 1 week prior to birth.

14Short-term TCE exposed mothers resided at Berkeley Manor, Midway Park, Paradise Point, and Wakins Village at the time of birth and at least 1 week during January 27 to February 7, 1985. In addition, the mother's last menstrual period occurred on or before January 31, 1985 and the birth occurred after February 2, 1985.

15The criteria for SGA being singleton births less than the 10<sup>th</sup> percentile of published sex-specific growth curves.

#### 4.8.3.1.2.3. Congenital malformations

1 Three studies focusing on occupational solvent exposure and congenital malformations  
2 from Europe provide data pertaining to TCE. Analyses of risk of congenital malformations were  
3 also included in the studies in the four geographic areas described above (Woburn, MA; Tucson,  
4 AZ, Rocky Mountain Arsenal, CO; Endicott, NY; and New Jersey), as well as additional sites in  
5 Phoenix, AZ; and Milwaukee, WI. Specific categories of malformations examined include  
6 cardiac defects, as well as cleft lip or cleft palate.  
7

##### 8 4.8.3.1.2.3.1. Occupational studies

9 A study of 1,148 men and 969 women occupationally exposed to TCE in Finland from  
10 1963–1976 to examined congenital malformations of offspring (Tola et al., 1980). Urinary  
11 trichloroacetic acid measurements available for 2,004 employees ranged from <10 to >500 mg/L,  
12 although 91% of the samples were below 100 mg/L. No congenital malformations were seen in  
13 the offspring of women between the ages of 15–49 years, although 3 were expected based on the  
14 national incidence. Expected number of cases for the cohort could not be estimated because the  
15 number of pregnancies was unknown.

16 Men from Finland occupationally exposed to organic solvents including TCE did not  
17 observe a risk of congenital malformations from paternal organic solvent exposure based on  
18 17 cases and 35 controls exposed to TCE (OR: 0.6, 95% CI: 0.2–2.0) (Taskinen et al., 1989).  
19 (Also see section above on spontaneous abortion for study details and additional results from this  
20 cohort.)

21 An occupational study of 100 women who gave birth to babies born with oral cleft  
22 defects and 751 control women with normal births were examined for exposure to a number of  
23 agents including TCE during the first trimester of pregnancy (Lorente et al., 2000). All women  
24 were participants in a multicenter European case-referent study whose children were born  
25 between 1989 and 1992. Four women were exposed to TCE, resulting in two cases of cleft lip  
26 (OR<sup>a</sup><sub>adj</sub>: 3.21, 95% CI: 0.49–20.9), and two cases of cleft palate (OR<sub>adj</sub>: 4.47, 95% CI: 1.02–40.9).  
27 Using logistic regression, the increased risk of cleft palate remained high (OR: 6.7, 95%  
28 CI: 0.9–49.7), even when controlling for tobacco and alcohol consumption (OR: 7.8, 95%  
29 CI: 0.8–71.8). However, the number of cases was small, and exposure levels were not known.  
30

##### 31 4.8.3.1.2.3.2. Geographic-based studies

32 A community in Woburn, MA with contaminated well water experienced an increased  
33 incidence of adverse birth outcomes and childhood leukemia (Lagakos et al., 1986, see section

1 on spontaneous abortion for study details). Statistically significant positive association between  
2 access to contaminated water and eye/ear birth anomalies (OR: 14.9,  $p < 0.0001$ ),  
3 CNS/chromosomal/oral cleft anomalies (OR: 4.5,  $p = 0.01$ ), kidney/urinary tract disorders  
4 (OR: 1.35,  $p = 0.02$ ) and lung/respiratory tract disorders (OR: 1.16,  $p = 0.05$ ) were observed.  
5 There were also five cases of cardiovascular anomalies, but there was not a significant  
6 association with TCE ( $p = 0.91$ ). However, since organogenesis occurs during gestational weeks  
7 3–5 in humans, some of these effects could have been missed if fetal loss occurred. (Also see  
8 sections on spontaneous abortion, perinatal death, decreased birth weight, and childhood cancer  
9 for additional results from this cohort.)

10 A high prevalence of congenital heart disease was found within an area of Tucson Valley,  
11 AZ (Goldberg et al., 1990, see section on spontaneous abortion for study details and additional  
12 results). Of the total 707 case families included, 246 (35%) were exposed to wells providing  
13 drinking water found to be contaminated with TCE (range = 6–239 ppb), along with  
14 dichloroethylene and chromium. Before the wells were closed after the contamination was  
15 discovered in 1981, the OR of congenital heart disease was 3 times higher for those exposed to  
16 contaminated drinking water compared to those not exposed; after the wells were closed, there  
17 was no difference seen. This study observed 18 exposed cases of congenital heart disease when  
18 16.4 would be expected (RR: 1.1). Prevalence of congenital heart disease in offspring after  
19 maternal exposure during the first trimester (6.8 in 1,000 live births) was significantly increased  
20 compared to nonexposed families (2.64 in 1,000 live births) ( $p < 0.001$ , 95% CI: 1.14–4.14). No  
21 difference in prevalence was seen if paternal data was included, and there was no difference in  
22 prevalence by ethnicity. In addition, no significant difference was seen for cardiac lesions.

23 A residential study of individuals living near the Rocky Mountain Arsenal in Colorado  
24 examined the outcomes in offspring of 75 men and 71 women exposed to TCE in drinking water  
25 (ATSDR, 2001). The risk was elevated for the nine birth defects observed (OR: 5.87,  
26 95% CI: 0.59–58.81), including one nervous system defect, one heart defect, and one incidence  
27 of cerebral palsy. The remaining cases were classified as “other,” and the authors speculate  
28 these may be based on inaccurate reports. (See above for study details and results on  
29 spontaneous abortion.)

30 The study of VOC exposure in Endicott, NY examined a number of birth defects during  
31 the years 1983–2000 (ATSDR, 2006a), see section on spontaneous for study details). These  
32 include total reportable birth defects, structural birth defects, surveillance birth defects, total  
33 cardiac defects, major cardiac defects, cleft lip/cleft palate, neural tube defects, and choanal  
34 atresia (blocked nasal cavities). There were 56 expected cases of all birth defects and 61 were  
35 observed resulting in no elevation of risk (rate ratio, RR: 1.08, 95% CI: 0.82–1.42). There were  
36 no cases of cleft lip/cleft palate, neural tube defects, or choanal atresia. Both total cardiac

1 defects ( $n = 15$ ; RR: 1.94, 95% CI: 1.21–3.12) and major cardiac defects ( $n = 6$ ; RR: 2.52,  
2 95% CI: 1.2–5.29) were statistically increased. A follow-up study by ATSDR (2008) reported  
3 that conotruncal heart malformations were particularly elevated ( $n = 4$ ; RR: 4.83, 95% CI:  
4 1.81–12.89). The results remained significantly elevated ( $\alpha$ RR: 3.74; 95% CI: 1.21–11.62)  
5 when infants with Down syndrome were excluded from the analysis. (Also see sections on  
6 spontaneous abortion, decreased birth weight, and childhood cancer for additional results from  
7 this cohort.)

8 In the New Jersey study described previously, the prevalence of birth defects reported by  
9 surveillance systems was examined among the women exposed to TCE and other contaminants  
10 in water while pregnant between 1985–1988 (Bove, 1996; Bove et al., 1995). For exposure  
11  $>10$  ppb ( $n = 1,372$ ), an increased risk, with relatively wide confidence intervals, was seen for all  
12 birth defects (OR: 2.53, 95% CI: 0.77–7.34). An increased risk was also seen for CNS defects  
13 ( $>10$  ppb: OR: 1.68), specifically 56 cases of neural tube defects ( $<1$ –5 ppb: 1.58,  
14 95% CI: 0.61–3.85;  $>10$  ppb: OR: 2.53, 95% CI: 0.77–7.34). A slight increase was seen in  
15 major cardiac defects ( $>10$  ppb: OR: 1.24, 50% CI: 0.75–1.94), including ventricular septal defects  
16 ( $>5$  ppb: OR: 1.30, 95% CI: 0.88–1.87). An elevated risk was seen for 9 cases of oral clefts  
17 ( $<5$  ppb: OR: 2.24, 95% CI: 1.04–4.66), although no dose-response was seen ( $>10$  ppb,  
18 OR: 1.30). However, due to the multiple contaminants in the water, it is difficult to attribute the  
19 results solely to TCE exposure. (See above for results from this study on fetal death and  
20 decreased birth weight.)

21 Arizona Department of Health Services (AZ DHS) conducted studies of contaminated  
22 drinking water and congenital malformations ( $<20$  years old) in Maricopa County, which  
23 encompasses Phoenix and the surrounding area (Flood, 1988). TCE contamination was  
24 associated with elevated levels of deaths in children less than 20 years old due to total congenital  
25 anomalies in East Central Phoenix from 1966–1969 (RR: 1.4, 95% CI: 1.1–1.7), from  
26 1970–1981 (RR: 1.5, 95% CI: 1.3–1.7), and from 1982–1986 (RR: 2.0, 95% CI: 1.5–2.5), as  
27 well as in other areas of the county. (See below for results from this study on childhood  
28 leukemia.)

29 A study was conducted of children born 1997–1999 with congenital heart defects in  
30 Milwaukee, WI (Yauck et al., 2004). TCE emissions data were ascertained from state and EPA  
31 databases, and distance between maternal residence and the emission source was determined  
32 using a GIS. Exposure was defined as those within 1.32 miles from at least one site. Results  
33 showed that an increased risk of congenital heart defects was seen for the offspring of exposed  
34 mothers 38 years old or older (OR: 6.2, 95% CI: 2.6–14.5), although an increased risk was also  
35 seen for offspring of unexposed mothers 38 years old or older (OR: 1.9, 95% CI: 1.1–3.5), and  
36 no risk was seen for offspring of exposed mothers younger than 38 years (OR: 0.9, 95%

1 CI: 0.6–1.2). The authors speculate that studies that did not find a risk only examined younger  
2 mothers. The authors also note that statistically-significant increased risk was seen for mothers  
3 with preexisting diabetes, chronic hypertension, or alcohol use during pregnancy.

4 An abstract reported that twenty-eight people living in a Michigan town were exposed for  
5 5–10 years to 8–14 ppm TCE in well water (Bernad et al., 1987, abstract). One child was born  
6 with multiple birth defects, with no further details.

#### 7 8 4.8.3.1.2.4. **Other adverse birth outcomes**

9 TCE was previously used as a general anesthetic during pregnancy. One study measured  
10 the levels of TCE in maternal and newborn blood after use during 34 vaginal childbirths (Beppu,  
11 1968). TCE was administered through a vaporizer from two to 98 minutes (mean 34.7 minutes)  
12 at volumes from 2–8 mL (mean 4.3 mL). Mean blood TCE concentrations were  $2.80 \pm 1.14$   
13 mg/dL in maternal femoral arteries;  $2.36 \pm 1.17$  mg/dL in maternal cubital veins;  $1.83 \pm 1.08$   
14 mg/dL in umbilical vein; and  $1.91 \pm 0.95$  mg/dL in the umbilical arteries. A significant  
15 correlation was seen for maternal arterial blood and infants' venous blood, and the concentration  
16 of the fetal blood was lower than that of the mother. Of these newborns, one had asphyxia and  
17 three "sleepy babies" had Apgar scores of 5–9; however, these results could not be correlated to  
18 length of inhalation and there was no difference in the TCE levels in the mother or newborn  
19 blood compared to those without adverse effects. Discussion included delayed newborn reflexes  
20 (raising the head and buttocks, bending the spine, and sound reflex), blood pressure, jaundice,  
21 and body weight gain; however, the results were compared to newborns exposed to other  
22 compounds, not to an unexposed population. This study also examined the concentration of TCE  
23 in one mother at 22-weeks gestation exposed for four minutes, after which the fetus was  
24 "artificially delivered." Maternal blood concentration was 3.0 mg/dL, and 0.9 mg/dL of TCE  
25 was found in the fetal heart, but not in other organs.

26 Another study of TCE administered during childbirth to the mother as an analgesic  
27 examined perinatal measures, including fetal pH, fetal partial pressure carbon dioxide (PCO<sub>2</sub>)  
28 fetal base deficit, fetal partial pressure oxygen (PO<sub>2</sub>), Apgar scores, and neonatal capillary blood  
29 (Phillips and Macdonald, 1971). The study consisted of 152 women whose fetus was considered  
30 to be at risk for hypoxia during labor. Out of this group, 51 received TCE (amount and route of  
31 exposure not reported). TCE caused fetal pH to fall more, base deficit increased more, and PO<sub>2</sub>  
32 fell more than the control group by fourfold or more compared to other analgesics used.

### 4.8.3.1.3. Postnatal developmental outcomes

#### 4.8.3.1.3.1. Developmental neurotoxicity

1 The studies examining neurotoxic effects from TCE exposure are discussed in  
2 Section 4.3, and the human developmental neurotoxic effects are reiterated here.

##### 4.8.3.1.3.1.1. Occupational studies

5 An occupational study examined the neurodevelopment of the offspring of 32 women  
6 exposed to various organic solvents during pregnancy (Laslo-Baker et al., 2004) Till et al., 2001.  
7 Three of these women were exposed to TCE; however, no levels were measured and the results  
8 for examined outcomes are for total organic solvent exposure, and are not specific to TCE.

##### 4.8.3.1.3.1.2. Geographic-based studies

10 A study of three residential cohorts (Woburn, MA; Alpha, OH; and Twin Cities, MN)  
11 examined the neurological effects of TCE exposure in drinking water (White et al., 1997). For  
12 Woburn, MA, 28 individuals ranging from 9–55 years old were assessed, with exposure from a  
13 tanning factor and chemical plant at levels 63–400 ppb for <1 to 12 years; the time between  
14 exposure and neurological examination was about 5 years. In this cohort, six of thirteen children  
15 (46%) had impairments in the verbal naming/language domain. For Alpha, OH, 12 individuals  
16 ranging from 12–68 years old were assessed, with exposure from degreasing used at a  
17 manufacturing operation at levels 3.3–330 ppb for 5–17 years; the time between exposure and  
18 neurological examination was 5–17 years. In this cohort, one of two children (50%) had  
19 impairments in the verbal naming/language domain. For Twin Cities, MN, 14 individuals  
20 ranging from 8–62 years old were assessed, with exposure from an army ammunition plant at  
21 levels 261–2,440 ppb for 0.25–25 years; the time between exposure and neurological  
22 examination was 4–22 years. In this cohort, four of four children (100%) had impairments in the  
23 verbal naming/language, memory, and academic domains and were diagnosed with moderate  
24 encephalopathy; and three of four children (75%) performed poorly on the WRAT-R Reading  
25 and Spelling and WAIS-R Information tests.

27 A case-control study was conducted to examine the relationship between multiple  
28 environmental agents and autism spectrum disorder (ASD) (Windham et al., 2006). Cases  
29 ( $n = 284$ ) and controls ( $n = 657$ ) were born in 1994 in the San Francisco Bay Area. Cases were  
30 diagnosed before age nine. Exposure was determined by geocoding births to census tracts, and  
31 linking to hazardous air pollutants data. An elevated risk was seen for TCE in the upper 3<sup>rd</sup>

1 quartile (OR: 1.37, 95% CI: 0.96–1.95), and a statistically significant elevated risk was seen for  
2 the upper 4<sup>th</sup> quartile (OR: 1.47, 95% CI: 1.03–2.08).

3 The Trichloroethylene Subregistry (Burg and Gist, 1999; Burg et al., 1995), including  
4 948 children <18 years old from 13 sites located in 3 states, was examined for any association of  
5 ingestion of drinking water contaminated with TCE and various health effects (ATSDR, 2003a;  
6 Burg and Gist, 1999; Burg et al., 1995). Exposure groups included (1) maximum TCE exposure,  
7 (2) cumulative TCE exposure, (3) cumulative chemical exposure, and (4) duration of exposure.  
8 Exposed children 0–9 years old had statistically increased hearing impairment compared to  
9 controls (RR: 2.13, 99% CI: 1.12–4.07), with children <5 having a 5.2-fold increase over  
10 controls. Exposed children 0–9 years old also had statistically increased speech impairment  
11 (RR: 2.45, 99% CI: 1.31–4.58). In addition, anemia and other blood disorders were statistically  
12 higher for males 0–9 years old. The authors noted that exposure could have occurred prenatally  
13 or postnatally. There was further analysis on the 116 exposed children and 182 controls who  
14 were under 10 years old at the time that the baseline study was conducted by ATSDR. This  
15 analysis did not find a continued association with speech and hearing impairment in these  
16 children; however, the absence of acoustic reflexes (contraction of the middle ear muscles in  
17 response to sound) remained significant (ATSDR, 2003a). No differences were seen when  
18 stratified by prenatal and postnatal exposure.

19 Twenty-eight people living in a Michigan town were exposed for 5–10 years to  
20 8–14 ppm TCE in well water (Bernad et al., 1987, abstract). Ten adults and 12 children  
21 completed a questionnaire on neurotoxic endpoints. Nine of the 12 children had poor learning  
22 ability, aggressive behavior, and low attention span.

#### 23 24 4.8.3.1.3.2. **Developmental immunotoxicity**

25 The studies examining human immunotoxic effects from TCE exposure are discussed in  
26 Section 4.6.1. The studies reporting developmental effects are reiterated briefly here.

27 Two studies focused on immunological development in children after maternal exposure  
28 to VOCs (Lehmann et al., 2001; Lehmann et al., 2002). The first examined premature neonates  
29 (1,500–2,500 g) and neonates at risk of atopy (cord blood IgE >0.9 kU/L; double positive family  
30 atopy history) at 36 months of age (Lehmann et al., 2001). Median air level in child's bedroom  
31 measured 0.42 µg/m<sup>3</sup>. There was no association with allergic sensitization to egg white and  
32 milk, or to cytokine producing peripheral T-cells. The second examined healthy, full-term  
neonates (≥2,500 g; ≥37 weeks gestation) born in Leipzig, Germany (Lehmann et al., 2002).



1 Median air level in the child’s bedroom 3–4 weeks after birth measured 0.6  $\mu\text{g}/\text{m}^3$ . A significant  
2 reduction of Th1 IL-2 producing T-cells was observed.

3 Byers et al. (1988) observed altered immune response in family members of children  
4 diagnosed with leukemia in Woburn, MA (Lagakos et al., 1986, see below for results of this  
5 study). The family members included 13 siblings under 19 years old at the time of exposure;  
6 however, an analysis looking at only these children was not done. This study is discussed in  
7 further detail in Section 4.6.1.

8

#### 4.8.3.1.3.3. **Other developmental outcomes**

9 A study demonstrated the adverse effects of TCE used as an anesthetic in children during  
10 operations during 1964 in Poland to repair developmental defects of the jaw and face (Jasińska,  
11 1965, translation). Fifty-five children ranging from 6 months to 10 years old were anesthetized  
12 with at least 10 mL TCE placed into an evaporator. Bradycardia occurred in 2 children, an  
13 accelerated heart rate of 20–25 beats per minute occurred in 7 children, no arrhythmia was  
14 observed, and arterial blood pressure remained steady or dropped by 10 mmHg only.  
15 Respiratory acceleration was observed in 25 of the children, and was seen more in infants and  
16 younger children.

17

#### 4.8.3.1.3.4. **Childhood cancer**

18 Several studies of parental occupational exposure were conducted in North America and  
19 the United Kingdom to determine an association with childhood cancer. A number of  
20 geographic-based studies were conducted in California; New Jersey; Woburn, MA; Endicott,  
21 NY; Phoenix, AZ; and Tucson, AZ. Specific categories of childhood cancers examined include  
22 leukemia, non-Hodgkin lymphoma, and CNS tumors.

23

##### 24 **4.8.3.1.3.4.1. Occupational studies**

25 Brain tumors in 98 children less than 10 years old at diagnosis from 1972–1977 in Los  
26 Angeles County have been observed in the offspring of fathers (Peters et al., 1985; Peters et al.,  
27 1981). Exposure was determined by questionnaire. Two cases with TCE exposure were  
28 reported: one case of oligodendroglioma in an 8-year-old whose father was a machinist, and  
29 astrocytoma in a 7-year-old whose father was an inspector for production scheduling and parts  
30 also exposed to methyl ethyl ketone (Peters et al., 1981). Peters et al. (1985) also briefly

1 mentioned 5 cases and no controls of paternal exposure to TCE and brain tumors in the offspring  
2 (resulting in an inability to calculate an odds ratio), but without providing any additional data.

3 A case-control study was conducted to assess an association between parental  
4 occupational exposure and neuroblastoma diagnosed in offspring <19 years old in the United  
5 States and Canada from May 1992 to April 1994 (De Roos et al., 2001). Paternal self-reported  
6 exposure to TCE was reported in 22 cases and 12 controls, resulting in an elevated risk of  
7 neuroblastoma in the offspring (OR: 1.4, 95% CI: 0.7–2.9). Maternal exposure to TCE was not  
8 reported.

9 A case-control study of parental occupational exposure and childhood leukemia was  
10 conducted in Los Angeles County (Lowengart et al., 1987). Children (61 boys and 62 girls)  
11 diagnosed less than 10 years old (mean age 4 years) from 1980–1984 were included in the  
12 analysis. Paternal occupation exposure to TCE was elevated for 1 year preconception (OR: 2.0,  
13  $p = 0.16$ ), prenatal (OR: 2.0,  $p = 0.16$ ), and postnatal (OR: 2.7,  $p = 0.7$ ). Maternal exposure to  
14 TCE was not reported.

15 A case-control study children diagnosed with acute lymphoblastic leukemia (ALL)  
16 examined parental occupational exposure to hydrocarbons in the United States and Canada (Shu  
17 et al., 1999). Children were under the age of 15 years at diagnosis during the years 1989–1993.  
18 Cases were confirmed with a bone marrow sample. 1,842 case-control pairs were given  
19 questionnaires on maternal and paternal exposures, resulting in 15 cases and 9 controls  
20 maternally exposed and 136 cases and 104 controls paternally exposed to TCE. There was an  
21 increased but nonsignificant risk for maternal exposure to TCE during preconception (OR: 1.8,  
22 95% CI: 0.6–5.2), pregnancy (OR: 1.8, 95% CI: 0.5–6.4), postnatally (OR: 1.4,  
23 95% CI: 0.5–4.1), or any of these periods (OR: 1.8, 95% CI: 0.8–4.1). However, there was no  
24 increased risk for paternal exposure to TCE.

25 Occupational exposure in communities in the United Kingdom was examined to  
26 determine an association with leukemia and non-Hodgkin lymphoma diagnosed in the offspring  
27 (McKinney et al., 1991). Paternal occupational exposure was elevated for exposure occurring  
28 during preconception (OR: 2.27, 95% CI: 0.84–6.16), prenatal (OR: 4.40, 95% CI: 1.15–21.01),  
29 and postnatal (OR: 2.66, 95% CI: 0.82–9.19). Risk from maternal preconception exposure was  
30 not elevated (OR: 1.16, 95% CI: 0.13–7.91). However, the number of cases examined in this  
31 study was low, particularly for maternal exposure.

#### 33 **4.8.3.1.3.4.2. Geographic-based studies**

34 A California community exposed to TCE (0.09–97 ppb) in drinking water from  
35 contaminated wells was examined for cancer (Morgan and Cassady, 2002). A specific emphasis

1 was placed on the examination of 22 cases of childhood cancer diagnosed before 15 years old.  
2 However, the incidence did not exceed those expected for the community for total cancer  
3 (SIR: 0.83, 99% CI: 0.44–1.40), CNS cancer (SIR: 1.05, 99% CI: 0.24–2.70), and leukemia  
4 (SIR: 1.09, 99% CI: 0.38–2.31).

5 An examination of drinking water was conducted in four New Jersey counties to  
6 determine an association with leukemia and non-Hodgkin lymphoma (Cohn et al., 1994b). A  
7 number of contaminants were reported, including VOCs and trihalomethanes. TCE was found as  
8 high as 67 ppb, and exposure categories were assigned to be >0.1, 0.1–5, and >5 ppb. A  
9 significantly elevated dose-response risk for ALL was observed for girls diagnosed before  
10 20 years old (RR: 3.36, 95% CI: 1.29–8.28), which was increased among girls diagnosed before  
11 5 years old (RR:4.54, 95% CI: 1.47–10.6). A significantly elevated dose-response risk for girls  
12 was also observed for total leukemia (RR: 1.43, 95% CI: 1.07–1.98).

13 The Woburn, MA community with contaminated well water experienced an increase in  
14 the incidence of childhood leukemia (Costas et al., 2002; Cutler et al., 1986; Lagakos et al.,  
15 1986; MDPH, 1997a). An initial study examined twelve cases of childhood leukemia diagnosed  
16 in children less than 15 years old between 1969–1979, when 5.2 cases were expected, and a  
17 higher risk was observed in boys compared to girls; however, no factors were observed to  
18 account for this increase (Cutler et al., 1986). Another study observed statistically significant  
19 positive association between access to contaminated water and 20 cases of childhood cancer  
20 were observed for both cumulative exposure metric (OR: 1.39,  $p = 0.03$ ), and none versus some  
21 exposure metric (OR: 3.03,  $p = 0.02$ ) (Lagakos et al., 1986). Massachusetts Department of  
22 Public Health (MDPH, 1997a) conducted a case-control study of children less than 20 years old  
23 living in Woburn and diagnosed with leukemia between 1969 and 1989 ( $n = 21$ ) and observed  
24 that consumption of drinking water increased the risk of leukemia (OR: 3.03, 95%  
25 CI: 0.82–11.28), with the highest risk from exposure during fetal development (OR: 8.33,  
26 95% CI: 0.73–94.67). This study found that paternal occupational exposure to TCE was not  
27 related to leukemia in the offspring (MDPH, 1997a). In the most recent update, Costas et al.  
28 (2002) reported that between the years 1969 and 1997, 24 cases of childhood leukemia were  
29 observed when 11 were expected. Risk was calculated for cumulative exposure to contaminated  
30 drinking water two years prior to conception (OR<sub>adj</sub>: 2.61, 95% CI: 0.47–14.97), during  
31 pregnancy (OR<sub>adj</sub>: 8.33, 95% CI: 0.73–94.67), postnatal (OR<sub>adj</sub>: 1.18, 95% CI: 0.28–5.05), and  
32 any of these time periods (OR<sub>adj</sub>: 2.39, 95% CI: 0.54–10.59). A dose response was observed  
33 during pregnancy only. Cases were more likely to be male (76%), <9 years old at diagnosis  
34 (62%), breast-fed (OR: 10.17, 95% CI: 1.22–84.50), and exposed during pregnancy (adjusted  
35 OR: 8.33, 95% CI: 0.73–94.67). A dose-response was seen during the pregnancy exposure  
36 period, with the most exposed having an adjusted OR of 14.30 (95% CI: 0.92–224.52). Other

1 elevated risks observed included maternal alcohol intake during pregnancy (OR: 1.50,  
2 95% CI: 0.54–4.20), having a paternal grandfather diagnosed with cancer (OR: 2.01,  
3 95% CI: 0.73–5.58), father employed in a high risk industry (OR: 2.55, 95% CI: 0.78–8.30), and  
4 public water being the subject’s primary beverage (OR: 3.03, 95% CI: 0.82–11.28). (Also see  
5 sections on spontaneous abortion, perinatal death, decreased birth weight, and congenital  
6 malformations for additional results from this cohort.)

7 The study of VOC exposure in Endicott, NY discussed above observed fewer than six  
8 cases of cancer that were diagnosed between 1980 and 2001 in children less than 20 years old,  
9 and did not exceed expected cases or types (ATSDR, 2006a). (See section on spontaneous  
10 abortion for study details, and sections on spontaneous abortion, decreased birth weight, and  
11 congenital malformations for additional results from this cohort.)

12 The AZ DHS conducted a number of studies of contaminated drinking water and 189  
13 cases of childhood cancer (<20 years old) (ADHS, 1990; Flood, 1988; Flood, 1997; Kioski et al.,  
14 1990a; Kioski et al., 1990b). In Maricopa County, which encompasses Phoenix and the  
15 surrounding area, TCE contamination (8.9 and 29 ppb in two wells) was associated with elevated  
16 levels of childhood leukemia ( $n = 67$ ) in west central Phoenix during 1965–1986 (SIR: 1.67,  
17 95% CI: 1.20–2.27) and 1982–1986 (SIR: 1.91, 95% CI: 1.11–3.12), but did not observe a  
18 significant increase in total childhood cancers, lymphoma, brain/CNS, or other cancers during  
19 these time periods (ADHS, 1990). (See above for results from this study on congenital  
20 anomalies.) A follow-up study retrospectively asked parents about exposures and found that  
21 residence within 2 miles of wells contaminated with TCE was not a risk factor for childhood  
22 leukemia, but identified a number of other risk factors (Flood, 1997). A further study of East  
23 Phoenix, reported on TCE contamination found along with 1,1,1-trichloroethane and 25 other  
24 contaminants in well water (levels not reported) and found no increase in incidence of childhood  
25 leukemia (SIR: 0.85, 95% CI: 0.50–1.35) based on 16 cases (Kioski et al., 1990b). There were  
26 also 16 cases of other types of childhood cancer, but were too few to be analyzed separately. In  
27 Pima County, which encompasses Tucson and the surrounding area, TCE was found in drinking  
28 wells (1.1–239 ppb), along with 1,1-dichloroethylene (DCE), chloroform and chromium and  
29 found a nonstatistically elevated risk of leukemia was observed (SIR: 1.50, 95% CI: 0.76–2.70),  
30 but no risk was observed for testicular cancer, lymphoma, or CNS/brain cancer (Kioski et al.,  
31 1990a).

#### 4.8.3.1.4. Summary of human developmental toxicity

1 Epidemiological developmental studies examined the association between TCE exposure  
 2 and a number of prenatal and postnatal developmental outcomes. Prenatal developmental  
 3 outcomes examined include spontaneous abortion and perinatal death; decreased birth weight,  
 4 small for gestational age, and postnatal growth; congenital malformations; and other adverse  
 5 birth outcomes. Postnatal developmental outcomes examined include developmental  
 6 neurotoxicity, developmental immunotoxicity, other developmental outcomes, and childhood  
 7 cancer related to TCE exposure.

8 More information on developmental outcomes is expected. A follow-up study of the  
 9 Camp Lejeune cohort (ATSDR, 1998b) for birth defects and childhood cancers was initiated in  
 10 1999 (ATSDR, 2003c) and expected to be completed soon (ATSDR, 2009; U.S. GAO, 2007).  
 11 Out of a total of 106 potential cases of either birth defects or childhood cancer, 57 have been  
 12 confirmed and will constitute the cases. These will be compared 548 control offspring of  
 13 mothers who also lived at Camp Lejeune during their pregnancy from 1968–1985. As part of  
 14 this study, a drinking water model was developed to determine a more accurate level and  
 15 duration of exposure to these pregnant women (ATSDR, 2007). Additional health studies have  
 16 been suggested, including adverse neurological or behavioral effects or pregnancy loss.

#### 4.8.3.1.5. Animal Developmental Toxicology Studies

18 A number of animal studies have been conducted to assess the potential for  
 19 developmental toxicity of TCE. These include studies conducted in rodents by prenatal  
 20 inhalation or oral exposures (summarized in Tables 4-96 and 4-97), as well as assessments in  
 21 nonmammalian species (e.g., avian, amphibian, and invertebrate species) exposed to TCE during  
 22 development. Studies have been conducted that provide information on the potential for effects

23 **Table 4-96. Summary of mammalian in vivo developmental toxicity**  
 24 **studies—inhalation exposures**

Reference	Species/strain/ sex/number	Exposure level/ duration	NOAEL; LOAEL <sup>a</sup>	Effects
Carney et al. (2006)	Rat, Sprague-Dawley, females, 27 dams/group	0, 50, 150, or 600 ppm (600 ppm = 3.2 mg/L) <sup>b</sup>	Mat. NOAEL: 150 ppm Mat. LOAEL: 600 ppm	↓ BW gain (22% less than control) on GD 6–9 at 600 ppm.
		6 h/d; GD 6–20	Dev. NOAEL: 600 ppm	No evidence of developmental toxicity, including heart defects.
Dorfmueller et al. (1979)	Rat, Long-Evans, females, 30	0 or 1,800 ± 200 ppm	Mat. NOAEL: 1,800 ± 200 ppm	No maternal abnormalities.

	dams/group	(9,674 ± 1,075 mg/m <sup>3</sup> ) <sup>b</sup>  2 wk, 6 h/d, 5 d/wk; prior to mating and/or on GD 0–20	Dev. LOAEL: 1,800 ± 200 ppm	Sig. ↑ skeletal and soft tissue anomalies in fetuses from dams exposed during pregnancy only. No sig. treatment effects on behavior of offspring 10, 20, or 100 d postpartum. BW gains sig. ↓ in pups from dams with pregestational exposure.
Hardin et al. (1981)	Rat, Sprague- Dawley, female, nominal 30/group	0 or 500 ppm  6–7 h/d; GD 1–19	Mat. NOAEL: 500 ppm	No maternal toxicity
			Dev. NOAEL: 500 ppm	No embryonic or fetal toxicity.
	Rabbit, New Zealand white, female, nominal 20/group	0 or 500 ppm  6–7 h/d; GD 1–24	Mat. NOAEL: 500 ppm	No maternal toxicity.
			Dev. LOAEL: 500 ppm	Hydrocephaly observed in 2 fetuses of 2 litters, considered equivocal evidence of teratogenic potential.
Healy et al. (1982)	Rat, Wistar, females, 31–32 dams/group	0 or 100 ppm  4 h/d; GD 8–21	Mat. NOAEL: 100 ppm	No maternal abnormalities.
			<b>Dev. LOAEL: 100 ppm</b>	<b>Litters with total resorptions sig. ↑. Sig. ↓ fetal weight, and ↑ bipartite or absent skeletal ossification centers.</b>
Schwetz et al. (1975)	Rat, Sprague- Dawley, female, 20–35/group Mouse, Swiss- Webster, females, 30–40 dams/group	0 or 300 ppm  7 h/d; GD 6–15	Mat. LOAEL: 300 ppm	4–5% ↓ maternal BW
			Dev. NOAEL: 300 ppm	No embryonic or fetal toxicity; not teratogenic.
Westergren et al. (1984)	Mouse, NMRI, male and female, 6–12 offspring/group	0 or 150 ppm  24 h/d; 30 d (during 7 d of mating and until GD 22)	Dev. LOAEL: 150 ppm <sup>c</sup>	Specific gravity of brains sig. ↓ at PND 0, 10, and 20–22. Similar effects at PND 20–22 in occipital cortex and cerebellum. No effects at 1 mo of age.

**Bolded studies** carried forward for consideration in dose-response assessment (see Section 5).

- 1 <sup>a</sup>NOAEL and LOAEL are based upon reported study findings.
- 2 <sup>b</sup>Dose conversions provided by study author(s).
- 3 <sup>c</sup>Parental observations not reported.
- 4 Dev. = developmental; Mat. = maternal; sig. = statistically significant.
- 5

1 **Table 4-97. Ocular defects observed (Narotsky et al., 1995)**  
2

<b>Dose TCE (mg/kg-day)</b>	<b>Incidence (number affected pups/total number pups)<sup>a</sup></b>	<b>Percentage of pups with eye defects</b>
0	1/197	0.51
10.1	0/71	0.00
32	0/85	0.00
101	3/68	4.41
320	3/82	3.66
475	6/100	6.00
633	6/100	6.00
844	7/58	12.07
1,125	12/44	27.27

3  
4 <sup>a</sup>Reported in Barton and Das (1996).  
5  
6

7 on specific organ systems, including the developing nervous, immune, and pulmonary systems.  
8 Additionally, a number of research efforts have focused on further characterization of the mode  
9 of action for cardiac malformations that have been reported to be associated with TCE exposure.  
10

#### 4.8.3.1.6. Mammalian studies

11 Studies that have examined the effects of TCE on mammalian development following  
12 either inhalation or oral exposures are described below and summarized in Tables 4-96 and 4-98,  
13 respectively.  
14

##### 4.8.3.1.6.1. Inhalation exposures

15 Dorfmueller et al. (1979) conducted a study in which TCE was administered by  
16 inhalation exposure to groups of approximately 30 female Long-Evans hooded rats at a  
17 concentration of 1,800 ± 200 ppm before mating only, during gestation only, or throughout the  
18 pre-mating and gestation periods. Half of the dams were killed at the end of gestation and half  
19 were allowed to deliver. There were no effects on body weight change or relative liver weight in  
20 the dams. The number of corpora lutea, implantation sites, live fetuses, fetal body weight,

1 resorptions, and sex ratio were not affected by treatment. In the group exposed only during  
2 gestation, a significant increase in four specific sternebral, vertebral, and rib findings, and a  
3



1  
2  
3

**Table 4-98. Summary of mammalian in vivo developmental toxicity studies—oral exposures**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
Blossom and Doss (2007)	Mouse, MRL +/+, dams and both sexes offspring, 3 litters/group, 8–12 offspring/group	0, 0.5, or 2.5 mg/mL  Parental mice and/or offspring exposed from GD 0 to 7–8 mo of age	Drinking water	Dev. LOAEL = 0.5 mg/mL <sup>b</sup>	At 0.5 mg/mL: sig ↓ postweaning weight; sig. ↑ IFN $\gamma$ produced by splenic CD4+ cells at 5–6 wk; sig ↓ splenic CD8+ and B220+ lymphocytes; sig. ↑ IgG2a and histone; sig. altered CD4-/CD8- and CD4+/CD8+ thymocyte profile. At 2.5 mg/mL: Sig ↓ postweaning weight; sig. ↑ IFN $\gamma$ produced by splenic CD4+ and CD8+ cells at 4–5 and 5–6 wk; sig ↓ splenic CD4+, CD8+, and B220+ lymphocytes; sig. altered CD4+/CD8+ thymocyte profile.
Blossom et al. (2008)	Mouse, MRL +/+, dams and both sexes offspring, 8 litters/group, 3–8 offspring/group	0 or 0.1 mg/mL (maternal dose = 25.7 mg/kg-day; offspring PND 24–42 dose—31.0 mg/kg-day  Parental mice and/or offspring exposed from GD 0 to PND 42	Drinking water	Dev. LOAEL = 1,400 ppb <sup>b</sup>	At 0.1 mg/mL: at PND 20, sig. ↑ thymocyte cellularity and distribution, associated with sig. ↑ in thymocyte subset distribution; sig. ↑ reactive oxygen species generation in total thymocytes; sig. ↑ in splenic CD4+ T-cell production of IFN- $\gamma$ and IL-2 in females and TNF- $\alpha$ in males at PND 42. Significantly impaired nest-building behaviors at PND 35. Increased aggressive activities, and increased oxidative stress and impaired thiol status in the cerebellar tissue of male offspring at PND 40.

4

**Table 4-98. Summary of mammalian in vivo developmental toxicity studies—oral exposures (continued)**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
Collier et al. (2003)	Rat, Sprague-Dawley, female, number dams/group not reported	0, 0.11, or 1.1 mg/mL  (0, 830, or 8,300 µgM) <sup>c</sup>  GD 0–11	Drinking water	Dev. LOEL: 0.11 mg/mL	Embryos collected between GD 10.5 and 11. Gene expression at 1.1 mg/mL TCE: 8 housekeeping genes ↑, and one gene ↓; 3 stress response genes ↑, IL-10 ↓; 2 cyto-skeletal/cell adhesion/blood related genes ↑, 3 genes ↓; 2 heart-specific genes ↑. Effects at 0.11 mg/mL reduced considerably. Two possible markers for fetal TCE exposure identified as Serca-2 Ca <sup>+2</sup> ATPase and GPI-p137.
Cosby and Dukelow (1992)	Mouse, B6D2F1, female, 28–62 dams/group	0, 24, or 240 mg/kg-day  GD 1–5, 6–10, or 11–15	Gavage in corn oil	Mat. NOAEL: 240 mg/kg-day	No maternal toxicity.
				Dev. NOAEL: 240 mg/kg-day	No effects on embryonic or fetal development.
Dawson et al. (1993)	Rat, Sprague-Dawley, 116 females allocated to 11 groups	0, 1.5, or 1,100 ppm  2 mo before mating and/or during gestation	Drinking water	Mat. NOAEL: 1,100 ppm	No maternal toxicity.
				Dev. LOAEL: 1.5 ppm	Sig. ↑ in heart defects, primarily atrial septal defects, found at both dose levels in groups exposed prior to pregnancy and during pregnancy, as well as in group exposed to 1,100 ppm dose during pregnancy only. No sig. ↑ in congenital heart defects in groups exposed prior to pregnancy only.
Fisher et al. (2001); Warren et al. (2006)	Rat, Sprague-Dawley, female, 20–25 dams/group	0 or 500 mg/kg-day  GD 6–15	Gavage in soybean oil	Mat. NOAEL: 500 mg/kg-day	No maternal toxicity.
				Dev. NOAEL: 500 mg/kg-day	No developmental toxicity. The incidence of heart malformations for fetuses from TCE-treated dams (3–5%) did not differ from negative controls. No eye defects observed.

**Table 4-98. Summary of mammalian in vivo developmental toxicity studies—oral exposures (continued)**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
Fredriksson et al. (1993)	Mouse, NMRI, male pups, 12 pups from 3–4 different litters/group	0, 50, or 290 mg/kg-day  PND 10–16	Gavage in a 20% fat emulsion prepared from egg lecithin and peanut oil	Dev. LOAEL: 50 mg/kg-day	Rearing activity sig. ↓ at both dose levels on PND 60.
George et al. (1986)	Rat, F334, male and female, 20 pairs/treatment group, 40 controls/sex	0, 0.15, 0.30 or 0.60% micro-encapsulated TCE  Breeders exposed 1 wk pre-mating, then for 13 wk; pregnant ♀s throughout pregnancy (i.e., 18 wk total)	Dietary	LOAEL: 0.15%	Open field testing in pups: a sig. dose-related trend toward ↑ time required for male and female pups to cross the first grid in the test device.
Isaacson and Taylor (1989)	Rat, Sprague-Dawley, females, 6 dams/group	0, 312, or 625 mg/L. (0, 4.0, or 8.1 mg/d) <sup>c</sup>  Dams (and pups) exposed from 14 d prior to mating until end of lactation.	Drinking water	Dev. LOAEL: 312 mg/L <sup>b</sup>	Sig. ↓ myelinated fibers in the stratum lacunosum-moleculare of pups. Reduction in myelin in the hippocampus.
Johnson et al. (2003)	Rat, Sprague-Dawley, female, 9–13/group, 55 in control group	0, 2.5 ppb, 250 ppb, 1.5 ppm, or 1,100 ppm  (0, 0.00045, 0.048, 0.218, or 129 mg/kg-day) <sup>c</sup>  GD 0–22	Drinking water	Dev. NOAEL: 2.5 ppb Dev. LOAEL: 250 ppb <sup>b</sup>	Sig. ↑ in percentage of abnormal hearts and the percentage of litters with abnormal hearts at ≥250 ppb.

**Table 4-98. Summary of mammalian in vivo developmental toxicity studies—oral exposures (continued)**

Reference	Species/strain/ sex/number	Dose level/ exposure duration	Route/ vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
Narotsky et al. (1995)	Rat, Fischer 344, females, 8–12 dams/group	0, 10.1, 32, 101, 320, 475, 633, 844, or 1,125 mg/kg-day  GD 6–15	Gavage in corn oil	Mat. LOAEL: 475 mg/kg-day	Sig. dose-related ↓ dam BW gain at all dose levels on GD 6–8 and 6–20. Delayed parturition at ≥475 mg/kg-day; ataxia at ≥633 mg/kg-day; mortality at 1,125 mg/kg-day.
Narotsky et al. (1995) (continued)				Dev. NOAEL: 32 mg/kg-day Dev. LOAEL: 101 mg/kg-day	↑ full litter resorption and postnatal mortality at ≥425 mg/kg-day. Sig. prenatal loss at 1,125 mg/kg-day. Pup BW ↓ (not sig.) on PND 1 and 6. Sig. ↑ in pups with eye defects at 1,125 mg/kg-day. Dose-related (not sig.) ↑ in pups with eye defects at ≥101 mg/kg-day.
Narotsky and Kavlock (1995)	Rat, Fischer 344, females, 16–21 dams/group	0, 1,125, or 1,500 mg/kg-day  GD 6–19	Gavage in corn oil	Mat. LOAEL: 1,125 mg/kg-day	Ataxia, ↓ activity, piloerection; dose-related ↓ BW gain.
				Dev. LOAEL: 1,125 mg/kg-day	Sig. ↑ full litter resorptions, ↓ live pups/litter; sig. ↓ pup BW on PND 1; sig. ↑ incidences of microphthalmia and anophthalmia.
Noland-Gerbec et al. (1986)	Rat, Sprague-Dawley, females, 9–11 dams/group	0 or 312 mg/L (Average total intake of dams: 825 mg TCE over 61 d) <sup>c</sup>  Dams (and pups) exposed from 14 d prior to mating until end of lactation	Drinking water	Dev. LOEL: 312 mg/L <sup>b</sup>	Sig. ↓ uptake of <sup>3</sup> H-2-DG in whole brains and cerebella (no effect in hippocampus) of exposed pups at 7, 11, and 16 d, but returned to control levels by 21 d.

**Table 4-98. Summary of mammalian in vivo developmental toxicity studies—oral exposures (continued)**

Reference	Species/strain/sex/number	Dose level/exposure duration	Route/vehicle	NOAEL; LOAEL <sup>a</sup>	Effects
Peden-Adams et al. (2006)	Mouse, B6C3F1, dams and both sexes offspring, 5 dams/group; 5–7 pups/group at 3 wk; 4–5 pups/sex/group at 8 wk	0, 1,400, or 14,000 ppb  Parental mice and/or offspring exposed during mating, and from GD 0 thru 3 or 8 wk of age	Drinking water	Dev. LOAEL: 1,400 ppb <sup>b</sup>	At 1,400 ppb: Suppressed plaque-forming cell (PFC) responses in males at 3 and 8 wk of age and in females at 8 wk of age. Delayed hypersensitivity response increased at 8 wk of age in females. At 14,000 ppb: Suppressed PFC responses in males and females at 3 and 8 wk of age. Splenic cell population decreased in 3 wk old pups. Increased thymic T-cells at 8 wk of age. Delayed hypersensitivity response increased at 8 wk of age in males and females.
Peden-Adams et al. (2008)	Mouse, MRL +/+, dams and both sexes offspring, unknown number litters/group, 6–10 offspring/sex/group	0, 1,400, or 14,000 ppb (vehicle = 1% emulphore)  Parental mice and/or offspring exposed from GD 0 to 12 mo of age	Drinking water	Dev. LOAEL = 1,400 ppb <sup>b</sup>	At 1,400 ppb: splenic CD4 <sup>-</sup> /CD8 <sup>-</sup> cells sig. ↑ in females; thymic CD4 <sup>+</sup> /CD8 <sup>+</sup> cells sig. ↓ in males; 18% ↑ in male kidney weight. At 14,000 ppb: thymic T-cell subpopulations (CD8 <sup>+</sup> , CD4/CD8 <sup>-</sup> , CD4 <sup>+</sup> ) sig. ↓ in males.
Taylor et al. (1985)	Rat, Sprague-Dawley, females, no. dams/group not reported	0, 312, 625, or 1,250 mg/L  Dams (and pups) exposed from 14 d prior to mating until end of lactation	Drinking water	Dev. LOAEL: 312 mg/L <sup>b</sup>	Exploratory behavior sig. ↑ in 60- and 90-d old male rats at all treatment levels. Locomotor activity was higher in rats from dams exposed to 1,250 ppm TCE.

**Bolded studies** carried forward for consideration in dose-response assessment (see Section 5).

<sup>a</sup>NOAEL, LOAEL, and LOEL (lowest-observed-effect level) are based upon reported study findings.

<sup>b</sup>Dose conversions provided by study author(s).

<sup>c</sup>Maternal observations not reported.

Dev. = developmental; Mat. = maternal; sig. = statistically significant.

1  
2  
3  
4  
5  
6  
7  
8

1 significant increase in displaced right ovary were observed upon fetal skeletal and soft tissue  
2 evaluation. Mixed function oxidase enzymes (ethoxycoumarin and ethoxyresorbin) which are  
3 indicative of cytochrome P450 and P448 activities, respectively, were measured in the livers of  
4 dams and fetuses, but no treatment-related findings were identified. Postnatal growth was  
5 significantly ( $p < 0.05$ ) decreased in the group with gestation-only exposures. Postnatal  
6 behavioral studies, consisting of an automated assessment of ambulatory response in a novel  
7 environment on GDs 10, 20, and 100, did not identify any effect on general motor activity of  
8 offspring following in utero exposure to TCE.

9 In a study by Schwetz et al. (1975), pregnant Sprague-Dawley rats and Swiss Webster  
10 mice (30–40 dams/group) were exposed to TCE via inhalation at a concentration of 300 ppm for  
11 7 hours/day on GDs 6–15. The only adverse finding reported was a statistically significant  
12 4–5% decrease in maternal rat body weight. There were no treatment related effects on pre- and  
13 postimplantation loss, litter size, fetal body weight, crown-rump length, or external, soft tissue,  
14 or skeletal findings.

15 Hardin et al. (1981) summarized the results of inhalation developmental toxicology  
16 studies conducted in pregnant Sprague-Dawley rats and New Zealand white rabbits for a number  
17 of industrial chemicals, including TCE. Exposure concentrations of 0 or 500 ppm TCE were  
18 administered for 6–7 hours/day, on gestations days 1–19 (rats) or 1–24 (rabbits), and cesarean  
19 sections were conducted on GDs 21 or 30, respectively. There were no adverse findings in  
20 maternal animals. No statistically significant increase in the incidence of malformations was  
21 reported for either species; however, the presence of hydrocephaly in two fetuses of two TCE-  
22 treated rabbit litters was interpreted as a possible indicator of teratogenic potential.

23 Healy et al. (1982) did not identify any treatment-related fetal malformations following  
24 inhalation exposure of pregnant inbred Wistar rats to 0 or 100 ppm (535 mg/m<sup>3</sup>) on GD 8–21. In  
25 this study, significant differences between control and treated litters were observed as an  
26 increased incidence of total litter loss ( $p < 0.05$ ), decreased mean fetal weight ( $p < 0.05$ ), and  
27 increased incidence of minor ossification variations ( $p = 0.003$ ) (absent or bipartite centers of  
28 ossification).

29 Carney et al. (2006) investigated the effects of whole-body inhalation exposures to  
30 pregnant Sprague-Dawley rats at nominal (and actual) chamber concentrations of 0, 50, 150, or  
31 600 ppm TCE for 6 hours/day, 7 days/week, on GDs 6–20. This study was conducted under  
32 Good Laboratory Practice regulations according to current EPA and Organisation for Economic  
33 Co-operation and Development (OECD) regulatory testing guidelines (i.e., OPPTS 870.3700 and  
34 OECD GD 414). Maternal toxicity consisted of a statistically significant decrease (22%) in body  
35 weight gain during the first 3 days of exposure to 600-ppm TCE, establishing a no-observed-  
36 effect concentration (NOEC) of 150 ppm for dams. No significant difference between control

1 and TCE-treated groups was noted for pregnancy rates, number of corpora lutea, implantations,  
2 viable fetuses per litter, percentage pre- and postimplantation loss, resorption rates, fetal sex  
3 ratios, or gravid uterine weights. External, soft tissue, and skeletal evaluation of fetal specimens  
4 did not identify any treatment-related effects. No cardiac malformations were identified in  
5 treated fetuses. The fetal NOEC for this study was established at 600 ppm.

6 Westergren et al. (1984) examined brain specific gravity of NMRI mice pups following  
7 developmental exposures to TCE. Male and female mice were separately exposed 24 hours/day  
8 (except for limited periods of animal husbandry activities) to 0- or 150-ppm TCE for 30 days and  
9 mated during exposure for 7 days. Exposure of the females was continued throughout gestation,  
10 until the first litter was born. Offspring (6–12/group; litter origin not provided in report) were  
11 terminated by decapitation on PND 1, 10, 21–22, or 30. The specific gravity of the brain frontal  
12 cortex, cortex, occipital cortex, and cerebellum were measured. The cortex specific gravity was  
13 significantly decreased at PND 1 ( $p < 0.001$ ) and 10 ( $p < 0.01$ ) in pups from exposed mice.  
14 There were also significant differences ( $p < 0.05$ ) in the occipital cortex and cerebellum at  
15 PND 20–22. This was considered suggestive of delayed maturation. No significant differences  
16 between control and treated pups were observed at 1 month of age.

#### 4.8.3.1.6.2. Oral exposures

18 A screening study conducted by Narotsky and Kavlock (1995) assessed the  
19 developmental toxicity potential of a number of pesticides and solvents, including TCE. In this  
20 study, Fischer 344 rats were administered TCE by gavage at 0, 1,125, and 1,500 mg/kg-day on  
21 GDs 6–19, and litters were examined on GDs 1, 3, and 6. TCE-related increased incidences of  
22 full-litter resorptions, decreased litter sizes, and decreased mean pup birth weights were observed  
23 at both treatment levels. Additionally, TCE treatment was reported to be associated with  
24 increased incidences of eye abnormalities (microphthalmia or anophthalmia). Increased  
25 incidences of fetal loss and percentage of pups with eye abnormalities were confirmed by  
26 Narotsky et al. (1995) in a preliminary dose-setting study that treated Fischer 344 rats with TCE  
27 by gavage doses of 475, 633, 844, or 1,125 mg/kg-day on GDs 6–15, and then in a  $5 \times 5 \times 5$   
28 mixtures study that used TCE doses of 0, 10.1, 32, 101, and 320 mg/kg-day on GD 6–15. In  
29 both studies, dams were allowed to deliver, and pups were examined postnatally. The incidence  
30 of ocular defects observed across all TCE treatment levels tested is presented in Table 4-97.

31 Other developmental findings in this study included increased full litter resorption at 475,  
32 844, and 1,125 mg/kg-day; increased postnatal mortality at 425 mg/kg-day. Pup body weights  
33 were decreased (not significantly) on PND 1 and 6 at 1,125 mg/kg-day. In both the Narotsky  
34 and Kavlock (1995) and Narotsky et al. (1995) studies, significantly decreased maternal body

1 weight gain was observed at the same treatment levels at which full litter resorption was noted.  
2 Additionally, in Narotsky et al. (1995) maternal observations included delayed parturition at 475,  
3 844, and 1,125 mg/kg-day, ataxia at 633 mg/kg-day, and mortality at 1,125 mg/kg-day.

4 Cosby and Dukelow (1992) administered TCE in corn oil by gavage to female B6D2F1  
5 mice (28–62/group) on GDs 1–5, 6–10, or 11–15 (where mating = GD 1). Dose levels were 0,  
6 1/100 and 1/10 of the oral LD<sub>50</sub> (i.e., 0, 24.02, and 240.2 mg/kg body weight). Dams were  
7 allowed to deliver; litters were evaluated for pup count sex, weight, and crown-rump length until  
8 weaning (PND 21). Some litters were retained until 6 weeks of age at which time gonads (from  
9 a minimum of 2 litters/group) were removed, weighed, and examined. No treatment-related  
10 reproductive or developmental abnormalities were observed.

11 A single dose of TCE was administered by gavage to pregnant CD-1 mice (9–19/group)  
12 at doses of 0, 0.1, or 1.0 µg/kg in distilled water, or 0, 48.3, or 483 mg/kg in olive oil, 24 hours  
13 after premating human chorionic gonadotropin (hCG) injection (Coberly et al., 1992). At  
14 53 hours after the hCG-injection, the dams were terminated, and the embryos were flushed from  
15 excised oviducts. Chimera embryos were constructed, cultured, and examined. Calculated  
16 proliferation ratios did not identify any differences between control and treated blastomeres. A  
17 lack of treatment-related adverse outcome was also noted when the TCE was administered by i.p.  
18 injection to pregnant mice (16–39/group) at 24 and 48 hours post-hCG at doses of 0, 0.01, 0.02,  
19 or 10 µg/kg body weight.

20 In a study intended to confirm or refute the cardiac teratogenicity of TCE that had been  
21 previously observed in chick embryos, Dawson et al. (1990b) continuously infused the gravid  
22 uterine horns of Sprague-Dawley rats with solutions of 0-, 15-, or 1,500-ppm TCE (or 1.5 or  
23 150-ppm dichloroethylene) on GDs 7–22. At terminal cesarean section on GD 22, the uterine  
24 contents were examined, and fetal hearts were removed and prepared for further dissection and  
25 examination under a light microscope. Cardiac malformations were observed in 3% of control  
26 fetuses, 9% of the 15-ppm TCE fetuses ( $p = 0.18$ ), and 14% of the 1,500-ppm TCE fetuses. ( $p =$   
27  $0.03$ ). There was a >60% increase in the percentage of defects with a 100-fold increase in dose.  
28 No individual malformation or combination of abnormalities was found to be selectively induced  
29 by treatment.

30 To further examine these TCE-induced cardiac malformations in rats, Dawson et al.  
31 (1993) administered 0, 1.5 or 1,100-ppm TCE in drinking water to female Sprague-Dawley rats.  
32 Experimental treatment regimens were (1) a period of approximately 2 months prior to  
33 pregnancy plus the full duration of pregnancy, (2) the full duration of pregnancy only, or (3) an  
34 average of 3 months before pregnancy only. The average total daily doses of TCE consumed for  
35 each exposure group at both dose levels were  
36



1

	1.5 ppm	1,100 ppm
Group 1	23.5 µL	1,206 µL
Group 2	0.78 µL	261 µL
Group 3	3.97 µL	1,185 µL

2

3 The study also evaluated 0, 0.15, or 110-ppm dichloroethylene in drinking water, with treatment  
4 administered (1) two months prior to pregnancy plus the full duration of pregnancy, or (2) an  
5 average of 2 months before pregnancy only. At terminal cesarean section, uterine contents were  
6 examined, fetuses were evaluated for external defects, and the heart of each fetus was removed  
7 for gross histologic examination under a dissecting microscope, conducted without knowledge of  
8 treatment group. There were no differences between TCE-treated and control group relative to  
9 percentage of live births, implants, and resorptions. The percentage of cardiac defects in  
10 TCE-treated groups ranged from 8.2–13.0%, and was statistically significant as compared to the  
11 control incidence of 3%. The dose-response was relatively flat, even in spite of the extensive  
12 difference between the treatment levels. There was a broad representation of various types of  
13 cardiac abnormalities identified, notably including multiple transposition, great artery, septal,  
14 and valve defects (see Table 4-99). No particular combination of defects or syndrome  
15 predominated. Exposure before pregnancy did not appear to be a significant factor in the  
16 incidence of cardiac defects.

17 In an attempt to determine a threshold for cardiac anomalies following TCE exposures,  
18 Johnson et al. (2003) 2005 compiled and reanalyzed data from five studies conducted from  
19 1989–1995. In these studies, TCE was administered in drinking water to Sprague-Dawley rats  
20 throughout gestation (i.e., a total of 22 days) at levels of 2.5 ppb (0.0025 ppm), 250 ppb  
21 (0.25-ppm), 1.5, or 1,100 ppm. The dams were terminated on the last day of pregnancy and  
22 fetuses were evaluated for abnormalities of the heart and great vessels. The control data from the  
23 five studies were combined prior to statistical comparison to the individual treated groups, which  
24 were conducted separately. The study author reported that significant increases in the percentage  
25 of abnormal hearts and the percentage of litters with abnormal hearts were observed in a  
26 generally dose-responsive manner at 250 ppb and greater (see Table 4-100).

27 In a study by Fisher et al. (2001), pregnant Sprague-Dawley rats were administered daily  
28 gavage doses on GDs 6–15 of TCE (500 mg/kg-day), TCA (300 mg/kg-day), or DCA  
29 (300 mg/kg-day). Cesarean delivery of fetuses was conducted on GD 21. Water and soybean oil  
30 negative control groups, and a retinoic acid positive control group were also conducted  
31 simultaneously. Maternal body weight gain was not significantly different from control for any  
32 of the treated groups. No significant differences were observed for number of implantations,

1  
2  
3  
4

**Table 4-99. Types of congenital cardiac defects observed in TCE-exposed fetuses (Dawson et al., 1993, Table 3)**

Cardiac abnormalities	Control	TCE concentrations					
		Premating		Premating/gestation		Gestation only	
		1,100 ppm	1.5 ppm	1,100 ppm	1.5 ppm	1,100 ppm	1.5 ppm
d-transposition (right chest)	2						
l-transposition (left chest)					2		1
Great artery defects				1	2		1
Atrial septal defects	1	7	3	19	5	7	4
Mitral valve defects				5	8		
Tricuspid valve defects		1		1	2		
Ventricular septal defects							
Subaortic	1			4	1	1	2
Membranous				2			
Muscular	2	1	1	4		4	1
Endocardial cushion defect	1					1	
Pulmonary valve defects			3	2	1		1
Aortic valve defects			1	2	2	2	
Situs inversus				1			
Total abnormalities	7	9	8	41	23	15	10
Total abnormal hearts	7	9	8	40	23	11	9

5  
6

**Table 4-100. Types of heart malformations per 100 fetuses (Johnson et al., 2003, Table 2, p. 290)**

Type of defect/100 fetuses	Control	TCE dose group			
		1,100 ppm	1.5 ppm	250 ppb	2.5 ppb
Abnormal looping	0.33		1		
Coronary artery/sinus				1.82	
Aortic hypoplasia			0.55		
Pulmonary artery hypoplasia			0.55		
Atrial septal defect	1.16	6.67	2.21	0.91	
Mitral valve defect	0.17			0.91	
Tricuspid valve defect				0.91	
Ventricular septal defect					
Perimembranous (subaortic)	0.33	2.86	1.66		
Muscular	0.33	0.95	0.55		
Atriventricular septal defect	0.17	0.95			
Pulmonary valve defect					
Aortic valve defects		1.9		0.91	
Fetuses with abnormal hearts ( <i>n</i> )	13	11	9	5	0
Total fetuses ( <i>n</i> )	606	105	181	110	144
Litters with fetuses with abnormal hearts/litter ( <i>n</i> )	9/55	6/9	5/13	4/9	0/12
Litter with fetuses with abnormal hearts/number litters (%)	16.4	66.7	38.5	44.4	0.0

resorptions, or litter size. Mean fetal body weight was reduced by treatment with TCA and DCA. The incidence of heart malformations was not significantly increased in treated groups as compared to controls. The fetal rate of cardiac malformations ranged from 3–5% across the TCE, TCA, and DCA dose groups and from 6.5–2.9% for the soybean and water control dose groups, respectively. It was suggested that the apparent differences between the results of this study and the Dawson et al. (1993) study may be related to factors such as differences in purity of test substances or in the rat strains, or differences in experimental design (e.g., oral gavage versus drinking water, exposure only during the period of organogenesis versus during the entire gestation period, or the use of a staining procedure). The rats from this study were also examined for eye malformations to follow-up on the findings of Narotsky (1995). As reported in Warren et al. (2006), gross evaluation of the fetuses as well as computerized morphometry

1 conducted on preserved and sectioned heads revealed no ocular anomalies in the groups treated  
2 with TCE. This technique allowed for quantification of the lens area, globe area, medial  
3 canthus, distance, and interocular distance. DCA treatment was associated with statistically  
4 significant reductions in the lens area, globe area, and interocular distance. All four measures  
5 were reduced in the TCA-treated group, but not significantly. The sensitivity of the assay was  
6 demonstrated successfully with the use of a positive control group that was dosed on GD 6–15  
7 with a known ocular teratogen, retinoic acid (15 mg/kg-day).

8 Johnson et al. (1998a; 1998b) conducted a series of studies to determine whether specific  
9 metabolites of TCE or dichloroethylene were responsible for the cardiac malformations observed  
10 in rats following administration during the period of organogenesis. Several metabolites of the  
11 two chemicals were administered in drinking water to Sprague-Dawley rats from GD 1–22.  
12 These included carboxy methylcystine, dichloroacetaldehyde, dichlorovinyl cystine,  
13 monochloroacetic acid, trichloroacetic acid, trichloroacetaldehyde, and trichloroethanol.  
14 Dichloroacetic acid, a primary common metabolite of TCE and dichloroethylene, was not  
15 included in these studies. The level of each metabolite administered in the water was based upon  
16 the dosage equivalent expected if 1,100 ppm (the limit of solubility) TCE broke down  
17 completely into that metabolite. Cesarean sections were performed on GD 22, uterine contents  
18 were examined, and fetuses were processed and evaluated for heart defects according to the  
19 procedures used by Dawson et al. (1993). No treatment-related maternal toxicity was observed  
20 for any metabolite group. Adverse fetal outcomes were limited to significantly increased  
21 incidences of fetuses with abnormal hearts (see Table 4-101). Significant increases in fetuses  
22 with cardiac defects (on a per-fetus and per-litter basis) were observed for only one of the  
23 metabolites evaluated, i.e., trichloroacetic acid (2,730 ppm, equivalent to a dose of 291  
24 mg/kg-day). Notably, significant increases in fetuses with cardiac malformations were also  
25 observed with 1.5 or 1,100-ppm TCE (0.218 or 129 mg/kg-day), or with 0.15 or 110-ppm DCE  
26 (0.015 or 10.64 mg/kg-day), but in each case only with pre-pregnancy-plus-pregnancy treatment  
27 regimens. The cardiac abnormalities observed were diverse and did not segregate to any  
28 particular anomaly or grouping. Dose related increases in response were observed for the overall  
29 number of fetuses with any cardiac malformation for both TCE and DCE; however, no dose-  
30 related increase occurred for any specific cardiac anomaly (Johnson et al., 1998b).

31 The TCE metabolites TCA and DCA were also studied by Smith et al. (1992; 1989).  
32 Doses of 0, 330, 800, 1,200, or 1,800 mg/kg TCA were administered daily by oral gavage to  
33 Long-Evan hooded rats on GDs 6–15. Similarly, DCA was administered daily by gavage to  
34 Long-Evans rats on GD 6–15 in two separate studies, at 0, 900, 1,400, 1,900, or  
35 2,400 mg/kg-day and 0, 14, 140, or 400 mg/kg-day. Embryo lethality and statistically or

- 1 biologically significant incidences of orbital anomalies (combined soft tissue and skeletal
- 2 findings) were observed for TCA at  $\geq 800$  mg/kg-day, and for DCA at  $\geq 900$  mg/kg-day. Fetal

**Table 4-101. Congenital cardiac malformations (Johnson et al., 1998a, Table 2, p. 997)**

Heart abnormalities	Treatment group												
	Normal water	TCE p+p 1,100 ppm	TCE p+p 1.5 ppm	TCE p 1,100 ppm	DCE p+p 110 ppm	DCE p+p 0.15 ppm	TCAA p 2,730 ppm	MCAA p 1,570 ppm	TCEth p 1,249 ppm	TCAld p 1,232 ppm	DCAlld p 174 ppm	CMC p 473 ppm	DCVC p 50 ppm
Abnormal looping	2	-	2	-	-	-	-	-	-	-	-	-	-
Aortic hypoplasia	-	1	1	-	1	-	1	-	1	-	1	-	1
Pulmonary artery hypoplasia	-	-	1	-	-	-	2	1	-	-	2	-	-
Atrial septal defects	7	19	5	7	11	7	3	3	-	2	-	-	1
Mitral valve defects, hypoplasia or ectasia	1	5	8	-	4	3	1	-	1	2	-	-	1
Tricuspid valve defects, hypoplasia or ectasia	-	1	1	-	1	-	-	-	1	-	-	-	-
Ventricular septal defects													
Perimembranous <sup>a</sup>	2	6	2	1	4	1	4	-	-	3	-	1	-
Muscular	2	4	-	4	2	1	1	-	1	-	-	2	2
Atrioventricular septal defects	1	-	-	1	1	-	-	-	-	-	-	-	-
Pulmonary valve defects	-	2	1	-	1	-	1	3	1	1	-	-	-
Aortic valve defects	-	2	2	2	2	3	-	-	1	-	-	1	-
Situs inversus	-	1	-	-	-	-	-	-	-	-	-	-	-
Total													
Abnormal hearts	15	41	23	15	25	15	13	7	6	8	3	4	5
Fetuses with abnormal hearts	13	40 <sup>a</sup>	22 <sup>a</sup>	11 <sup>a</sup>	24 <sup>a</sup>	14 <sup>a</sup>	12 <sup>a</sup>	6	5	8	3	4	5
Fetuses	605	434	255	105	184	121	114	132	121	248	101	85	140

<sup>a</sup>Subaortic.<sup>b</sup>Per-fetus statistical significance (Fisher's exact test).

p = pregnancy ; p+p = pregnancy; and prepregnancy.

1 growth (body weight and crown-rump length) was affected at  $\geq 330$  mg/kg-day for TCE and at  
2  $\geq 400$  mg/kg-day for DCA. For TCA, the most common cardiac malformations observed were  
3 levocardia at  $\geq 330$  mg/kg-day and interventricular septal defect at  $\geq 800$  mg/kg-day. For DCA,  
4 levocardia was observed at  $\geq 900$  mg/kg-day, interventricular septal defect was observed at  
5  $\geq 1,400$  mg/kg-day, and a defect between the ascending aorta and right ventricle was observed in  
6 all treated groups (i.e.,  $\geq 14$  mg/kg-day, although the authors appeared to discount the single fetal  
7 finding at the lowest dose tested). Thus, NOAELs were not definitively established for either  
8 metabolite, although it appears that TCA was generally more potent than DCA in inducing  
9 cardiac abnormalities.

10 These findings were followed up by a series of studies on DCA reported by Epstein et al.  
11 (1992), which were designed to determine the most sensitive period of development and further  
12 characterize the heart defects. In these studies, Long-Evans hooded rats were dosed by oral  
13 gavage with a single dose of 2,400 mg/kg-day on selected days of gestation (6–8, 9–11, or  
14 12–15); with a single dose of 2,400 mg/kg on Days 10, 11, 12, or 13; or with a single dose of  
15 3,500 mg/kg on Days 9, 10, 11, 12, or 13. The heart defects observed in these studies were  
16 diagnosed as high interventricular septal defects rather than membranous type interventricular  
17 septal defects. The authors hypothesized that high intraventricular septal defects are a specific  
18 type of defect produced by a failure of proliferating interventricular septal tissue to fuse with the  
19 right tubercle of the atrioventricular cushion tissue. This study identified GDs 9 through 12 as a  
20 particularly sensitive period for eliciting high interventricular septal defects. It was postulated  
21 that DCA interferes with the closure of the tertiary interventricular foramen, allowing the aorta to  
22 retain its embryonic connection with the right ventricle. Further, it was suggested that the  
23 selectivity of DCA in inducing cardiac malformations may be due to the disruption of a discrete  
24 cell population.

25 TCE and its metabolites DCE and TCAA were administered in drinking water to  
26 pregnant Sprague-Dawley rats from GDs 0–11 (Collier et al., 2003). Treatment levels were 0,  
27 110, or 1,100 ppm (i.e., 0, 830 or 8,300  $\mu\text{gM}$ ) TCE; 0, 11, or 110 ppm (i.e., 0, 110, or 1,100  
28  $\mu\text{gM}$ ) DCE; 0, 2.75, or 27.3 mg/mL (i.e., 0, 10, or 100 mM) TCAA. Embryos (including hearts)  
29 were harvested between embryonic days 10.5–11, since this is the stage at which the  
30 developmental processes of myoblast differentiation, cardiac looping, atrioventricular valve  
31 formation, and trabeculation would typically be occurring. A PCR based subtraction scheme  
32 was used to identify genes that were differentially regulated with TCE or metabolite exposure.  
33 Numerous differentially regulated gene sequences were identified. Up-regulated transcripts  
34 included genes associated with stress response (Hsp 70) and homeostasis (several ribosomal  
35 proteins). Down-regulated transcripts included extracellular matrix components (GPI-p137 and  
36 vimentin) and  $\text{Ca}^{2+}$  responsive proteins (Serca-2  $\text{Ca}^{2+}$ -ATPase and  $\beta$ -catenin). Serca-2  $\text{Ca}^{2+}$  and

1 GPI-p137 were identified as two possible markers for fetal TCE exposure. Differential  
2 regulation of expression of these markers by TCE was confirmed by dot blot analysis and  
3 semiquantitative real time PCR with decreased expression seen at levels of TCE exposure  
4 between 100 and 250 ppb (0.76 and 1.9  $\mu$ M).  
5

#### 6 **4.8.3.1.6.2.1. *Developmental neurotoxicity and developmental immunotoxicity***

7 Several studies were conducted that included assessments of the effects of TCE oral  
8 exposure on the developing nervous system (Blossom et al., 2008; Dorfmueller et al., 1979;  
9 Fredriksson et al., 1993; George et al., 1986; Isaacson and Taylor, 1989; Noland-Gerbec et al.,  
10 1986) or immune system (Blossom and Doss, 2007; Blossom et al., 2008; Peden-Adams et al.,  
11 2006; Peden-Adams et al., 2008). These studies, summarized below, are addressed in additional  
12 detail in Section 4.3 (nervous system) and Section 4.6.2.1.2 (immune system).  
13

#### 14 **4.8.3.1.6.2.2. *Developmental neurotoxicity***

15 Fredriksson et al. (1993) conducted a study in male NMRI weanling mice (12/group,  
16 selected from 3–4 litters), which were exposed to trichloroethylene by oral gavage at doses of 0  
17 (vehicle), 50, or 290 mg/kg-day TCE in a fat emulsion vehicle, on PNDs 10–16. Locomotor  
18 behavior (horizontal movement, rearing and total activity) were assessed over three 20-minute  
19 time periods at GDs 17 and 60. There were no effects of treatment in locomotor activity at PND  
20 17. At PND 60, the mice treated with 50 and 290 mg/kg-day TCE showed a significant  
21 ( $p < 0.01$ ) decrease in rearing behavior at the 0–20 and 20–40 minute time points, but not at the  
22 40–60 minute time point. Mean rearing counts were decreased by over 50% in treated groups as  
23 compared to control. Horizontal activity and total activity were not affected by treatment.

24 Open field testing was conducted in control and high-dose F1 weanling Fischer 344 rat  
25 pups in an NTP reproduction and fertility study with continuous breeding (George et al., 1986).  
26 In this study, TCE was administered at dietary levels of 0, 0.15, 0.30, or 0.60%. The open field  
27 testing revealed a significant ( $p < 0.05$ ) dose-related trend toward an increase in the time required  
28 for male and female pups to cross the first grid in the testing device, suggesting an effect on the  
29 ability to react to a novel environment.

30 Taylor et al. (1985) administered TCE in drinking water (0, 312, 625, or 1,250 ppm) to  
31 female Sprague-Dawley rats for 14 days prior to breeding, and from GD 0 through offspring GD  
32 21. The number of litters/group was not reported, nor did the study state how many pups per  
33 litter were evaluated for behavioral parameters. Exploratory behavior was measured in the pups  
34 in an automated apparatus during a 15-minute sampling period on PND 28, 60, and 90.



1 Additionally, wheel-running, feeding, and drinking behavior was monitored 24 hours/day on  
2 PND 55-60. The number of exploratory events was significantly increased by approximately  
3 25–50% in 60- and 90-day old male TCE-treated rats at all dose levels, with the largest effect  
4 observed at the highest dose level tested, although there were no effects of treatment on the  
5 number of infrared beam-breaks. No difference between control and treated rats was noted for  
6 pups tested on PND 28. Wheel-running activity was increased approximately 40% in 60-day old  
7 males exposed to 1,25-ppm TCE as compared to controls. It is notable that adverse outcomes  
8 reported in the developmentally-exposed offspring on this study were observed long after  
9 treatment ceased.

10 Using a similar treatment protocol, the effects of TCE on development of myelinated  
11 axons in the hippocampus was evaluated by Isaacson and Taylor (1989) in Sprague-Dawley rats.  
12 Female rats (6/group) were exposed in the drinking water from 14 days prior to breeding and  
13 through the mating period; then the dams and their pups were exposed throughout the prenatal  
14 period and until PND 21, when they were sacrificed. The dams received 0, 312 or 625 ppm (0,  
15 4, or 8.1 mg/day TCE in the drinking water. Myelinated fibers were counted in the hippocampus  
16 of 2–3 pups per treatment group at PND 21, revealing a decrease of approximately 40% in  
17 myelinated fibers in the CA1 area of the hippocampus of pups from dams at both treatment  
18 levels, with no dose-response relationship. There was no effect of TCE treatment on myelination  
19 in several other brain regions including the internal capsule, optic tract or fornix.

20 A study by Noland-Gerbec et al. (1986) examined the effect of pre- and perinatal  
21 exposure to TCE on 2-deoxyglucose (2-DG) uptake in the cerebellum, hippocampus and whole  
22 brain of neonatal rats. Sprague-Dawley female rats (9–11/group) were exposed via drinking  
23 water to 0 or 312 mg TCE/liter distilled water from 14 days prior to mating until their pups were  
24 euthanized at GD 21. The total TCE dose received by the dams was 825 mg over the 61-day  
25 exposure period. Pairs of male neonates were euthanized on PND 7, 11, 16, and 21. There was  
26 no significant impairment in neonatal weight or brain weight attributable to treatment, nor were  
27 other overt effects observed. 2-DG uptake was significantly reduced from control values in  
28 neonatal whole brain (9–11%) and cerebellum (8–16%) from treated rats at all ages studied, and  
29 hippocampal 2-DG uptake was significantly reduced (7–21% from control) in treated rats at all  
30 ages except at PND 21.

31 In a study by Blossom et al. (2008), MRL +/+ mice were treated in the drinking water  
32 with 0 or 0.1 mg/mL TCE from maternal GD 0 through offspring PND 42. Based on drinking  
33 water consumption data, average maternal doses of TCE were 25.7 mg/kg-day, and average  
34 offspring (PND 24–42) doses of TCE were 31.0 mg/kg-day. In this study, a subset of offspring  
35 (3 randomly selected neonates from each litter) was evaluated for righting reflex on PNDs 6, 8,  
36 and 10; bar-holding ability on PNDs 15 and 17; and negative geotaxis on PNDs 15 and 17; none

1 of these were impaired by treatment. In an assessment of offspring nest building on PND 35,  
2 there was a significant association between impaired nest quality and TCE exposure; however,  
3 TCE exposure did not have an effect on the ability of the mice to detect social and nonsocial  
4 odors on PND 29 using olfactory habituation and dishabituation methods. Resident intruder  
5 testing conducted on PND 40 to evaluate social behaviors identified significantly more  
6 aggressive activities (i.e., wrestling and biting) in TCE-exposed juvenile male mice as compared  
7 to controls. Cerebellar tissue homogenates from the male TCE-treated mice had significantly  
8 lower GSH levels and GSH:GSSG ratios, indicating increased oxidative stress and impaired thiol  
9 status; these have been previously reported to be associated with aggressive behaviors (Franco et  
10 al., 2006). Qualitative histopathological examination of the brain did not identify alterations  
11 indicative of neuronal damage or inflammation. Although the study author attempted to link the  
12 treatment-related alterations in social behaviors to the potential for developmental exposures to  
13 TCE to result in autism in humans, this association is not supported by data and is considered  
14 speculative at this time.

15 As previously noted, postnatal behavioral studies conducted by Dorfmueller et al. (1979)  
16 did not identify any changes in general motor activity measurements of rat offspring on PND 10,  
17 20, and 100 following maternal gestational inhalation exposure to TCE at  $1,800 \pm 200$  ppm.  
18

#### 19 **4.8.3.1.6.2.3. *Developmental immunotoxicity***

20 Peden-Adams et al. (2006) assessed the potential for developmental immunotoxicity  
21 following TCE exposures. In this study, B6C3F1 mice (5/sex/group) were administered TCE via  
22 drinking water at dose levels of 0, 1,400 or 14,000 ppb from maternal GD 0 to either postnatal 3  
23 or 8, when offspring lymphocyte proliferation, NK cell activity, SRBC-specific IgM production  
24 (PFC response), splenic B220+ cells, and thymus and spleen T-cell immunophenotypes were  
25 assessed. (A total of 5–7 pups per group were evaluated at Week 3, and the remainder were  
26 evaluated at Week 8.) Observed positive responses consisted of suppressed PFC responses in  
27 males at both ages and both TCE treatment levels, and in females at both ages at 14,000 ppb and  
28 at 8 weeks of age at 1,400 ppb. Spleen numbers of B220+ cells were decreased in 3-week old  
29 pups at 14,000 ppb. Pronounced increases in all thymus T-cell subpopulations (CD4+, CD8+,  
30 CD4+/CD8+, and CD4-/CD8-) were observed at 8 weeks of age. Delayed hypersensitivity  
31 response, assessed in offspring at 8 weeks of age, was increased in females at both treatment  
32 levels and in males at 14,000 ppb only. No treatment-related increase in serum anti-dsDNA  
33 antibody levels was found in the offspring at 8 weeks of age.

34 In a study by Blossom and Doss (2007), TCE was administered to groups of pregnant  
35 MRL +/+ mice in drinking water at levels of 0, 0.5, or 2.5 mg/mL. TCE was continuously

1 administered to the offspring until young adulthood (i.e., 7–8 weeks of age). Offspring  
2 postweaning body weights were significantly decreased in both treated groups. Decreased  
3 spleen cellularity and reduced numbers of CD4+, CD8+, and B220+ lymphocyte subpopulations  
4 were observed in the postweaning offspring. Thymocyte development was altered by TCE  
5 exposures (significant alterations in the proportions of double-negative subpopulations and  
6 inhibition of in vitro apoptosis in immature thymocytes). A dose-dependent increase in CD4+  
7 and CD8+ T-lymphocyte IFN $\gamma$  was observed in peripheral blood by 4–5 weeks of age, although  
8 these effects were no longer observed at 7–8 weeks of age. Serum antihistone autoantibodies  
9 and total IgG<sub>2a</sub> were significantly increased in treated offspring; however, no histopathological  
10 signs of autoimmunity were observed in the liver and kidneys at sacrifice.

11 Blossom et al. (2008) administered TCE to MRL +/+ mice (8 dams/group) in the drinking  
12 water at levels of 0 or 0.1 mg/mL from GD 0 through offspring GD 42. Average maternal doses  
13 of TCE were 25.7 mg/kg-day, and average offspring (PND 24–42) doses of TCE were 31.0  
14 mg/kg-day. Subsets of offspring were sacrificed at PND 10 and 20, and thymus endpoints (i.e.,  
15 total cellularity, CD4+/CD8+ ratios, CD24 differentiation markers, and double-negative  
16 subpopulation counts) were evaluated. Evaluation of the thymus identified a significant  
17 treatment-related increase in cellularity, accompanied by alterations in thymocyte subset  
18 distribution, at PND 20 (sexes combined). TCE treatment also appeared to promote T-cell  
19 differentiation and maturation at PND 42. Indicators of oxidative stress were measured in the  
20 thymus at PND 10 and 20, and in the brain at PND 42, and ex vivo evaluation of cultured  
21 thymocytes indicated increased ROS generation. Mitogen-induced intracellular cytokine  
22 production by splenic CD4+ and CD8+ T-cells was evaluated in juvenile mice and brain tissue  
23 was examined at PND 42 for evidence of inflammation. Evaluation of peripheral blood  
24 indicated that splenic CD4+ T-cells from TCE-exposed PND 42 mice produced significantly  
25 greater levels of IFN- $\gamma$  and IL-2 in males and TNF- $\alpha$  in both sexes. There was no effect on  
26 cytokine production on PND 10 or 20.

27 Peden-Adams et al. (2008) administered TCE to MRL+/+ mice (unspecified number of  
28 dams/group) in drinking water at levels of 0, 1,400, or 14,000 ppb from GD 0 and continuing  
29 until the offspring were 12 months of age. At 12 months of age, final body weight; spleen,  
30 thymus, and kidney weights; spleen and thymus lymphocyte immunophenotyping (CD4 or  
31 CD8); splenic B-cell counts; mitogen-induced splenic lymphocyte proliferation; serum levels of  
32 autoantibodies to dsDNA and GA, periodically measured from 4–12 months of age; and urinary  
33 protein measures were recorded. Reported sample sizes for the offspring measurements varied  
34 from 6–10 per sex per group; the number of source litters represented within each sample was  
35 not specified. The only organ weight alteration was an 18% increase in kidney weight in the  
36 1,400 ppb males. Splenic CD4-/CD8- cells were altered in female mice (but not males) at 1,400

1 ppm only. Splenic T-cell populations, numbers of B220+ cells, and lymphocyte proliferation  
2 were not affected by treatment. Populations of thymic T-cell subpopulations (CD8+,  
3 CD4-/CD8-, and CD4+) were significantly decreased in male but not female mice following  
4 exposure to 14,000 ppb TCE, and CD4+/CD8+ cells were significantly reduced in males by  
5 treatment with both TCE concentrations. Autoantibody levels (anti-dsDNA and anti-GA) were  
6 not increased in the offspring over the course of the study.

7 Although all of the developmental immunotoxicity studies with TCE (Peden-Adams et  
8 al., 2006, 2008; Blossom and Doss, 2007; Blossom et al., 2008) exposed the offspring during  
9 critical periods of pre- and postnatal immune system development, they were not designed to  
10 assess issues such as posttreatment recovery, latent outcomes, or differences in severity of  
11 response that might be attributed to the early life exposures.  
12

#### 4.8.3.1.6.3. Intrapерitoneal exposures

13 The effect of TCE on pulmonary development was evaluated in a study by Das and Scott  
14 (1994). Pregnant Swiss-Webster mice (5/group) were administered a single intraperitoneal  
15 injection of TCE in peanut oil at doses of 0 or 3,000 mg/kg on GD 17 (where mating = Day 1).  
16 Lungs from GD 18 and 19 fetuses and from neonates on PND 1, 5, and 10 were evaluated for  
17 phospholipid content, DNA, and microscopic pathology. Fetal and neonatal (PND 1) mortality  
18 was significantly increased ( $p < 0.01$ ) in the treated group. Pup body weight and absolute lung  
19 weight were significantly decreased ( $p < 0.05$ ) on PND 1, and mean absolute and relative (to  
20 body weight) lung weights were significantly decreased on GDs 18 and 19. Total DNA content  
21 ( $\mu\text{g}/\text{mg}$  lung) was similar between control and treated mice, but lung phospholipid was  
22 significantly ( $p < 0.05$ ) reduced on GD 19 and significantly increased ( $p < 0.05$ ) on PND 10 in  
23 the TCE-treated group. Microscopic examination revealed delays in progressive lung  
24 morphological development in treated offspring, first observed at GD 19 and continuing at least  
25 through PND 5.  
26

#### 4.8.3.1.7. Studies in nonmammalian species

##### 4.8.3.1.7.1. Avian

27 Injection of White Leghorn chick embryos with 1, 5, 10, or 25  $\mu\text{mol}$  TCE per egg on  
28 Days 1 and 2 of embryogenesis demonstrated mortality, growth defects, and morphological  
29 anomalies at evaluation on Day 14 (Bross et al., 1983). These findings were consistent with a

1 previous study that had been conducted by Elovaara et al. (1979). Up to 67% mortality was  
2 observed in the treated groups, and most of the surviving embryos were malformed (as compared  
3 to a complete absence of malformed chicks in the untreated and mineral-oil-treated control  
4 groups). Reported anomalies included subcutaneous edema, evisceration (gastroschisis), light  
5 dermal pigmentation, beak malformations, club foot, and patchy feathering. Retarded growth  
6 was observed as significantly ( $p < 0.05$ ) reduced crown-rump, leg, wing, toe, and beak lengths as  
7 compared to untreated controls. This study did not identify any liver damage or cardiac  
8 anomalies.

9 In a study by Loeber et al. (1988), 5, 10, 15, 20, or 25  $\mu\text{mol}$  TCE was injected into the air  
10 space of White Longhorn eggs at embryonic stages 6, 12, 18, or 23. Embryo cardiac  
11 development was examined in surviving chicks in a double-blinded manner at stages 29, 34, or  
12 44. Cardiac malformations were found in 7.3% of TCE-treated hearts, compared to 2.3% of  
13 saline controls and 1.5% of mineral oil controls. The observed defects included septal defects,  
14 cor biloculare, conotruncal abnormalities, atrioventricular canal defects, and abnormal cardiac  
15 muscle.

16 Drake et al. (2006a) injected embryonated White Leghorn chicken eggs (Babcock or  
17 Bovan strains) with 0, 0.4, 8, or 400 ppb TCE per egg during the period of cardiac valvuloseptal  
18 morphogenesis (i.e., 2–3.3 days incubation). The injections were administered in four aliquots at  
19 Hamberger and Hamilton (HH) stages 13, 15, 17, and 20, which spanned the major events of  
20 cardiac cushion formation, from induction through mesenchyme transformation and migration.  
21 Embryos were harvested 22 hours after the last injection (i.e., HH 24 or HH 30) and evaluated  
22 for embryonic survival, apoptosis, cellularity and proliferation, or cardiac function. Survival was  
23 significantly reduced for embryos at 8 and 400 ppb TCE at HH 30. Cellular morphology of  
24 cushion mesenchyme, cardiomyocytes, and endocardioocytes was not affected by TCE treatment;  
25 however, the proliferative index was significantly increased in the atrioventricular canal (AVC)  
26 cushions at both treatment levels and in the outflow tract (OFT) cushions at 8 ppb. This resulted  
27 in significant cushion hypercellularity for both the OFT and AVC of TCE-treated embryos.  
28 Similar outcomes were observed in embryos when TCA or TCOH was administered, and the  
29 effects of TCA were more severe than for TCE. Doppler ultrasound assessment of cardiac  
30 hemodynamics revealed no effects of TCE exposure on cardiac cycle length or heart rate;  
31 however, there was a reduction in dorsal aortic blood flow, which was attributed to a 30.5%  
32 reduction in the active component of atrioventricular blood flow. Additionally the passive-to-  
33 active atrioventricular blood flow was significantly increased in treated embryos, and there was a  
34 trend toward lower stroke volume. The overall conclusion was that exposure to 8 ppb TCE  
35 during cushion morphogenesis reduced the cardiac output of the embryos in this study. The  
36 findings of cardiac malformations and/or mortality following in ovo exposure to chick embryos

1 with 8 ppb TCE during the period of valvuloseptal morphogenesis has also been confirmed by  
2 Rufer et al. (2008; 2010).

3 In a follow-up study, Drake et al. (2006b) injected embryonated White Leghorn chicken  
4 eggs with TCE or TCA during the critical window of avian heart development, beginning at HH  
5 stage 3+ when the primary heart field is specified in the primitive streak and ending  
6 approximately 50 hours later at HH stage 17, at the onset of chambering. Total dosages of 0, 0.2,  
7 2, 4, 20, or 200 nmol (equivalent to 0, 0.4, 4, 8, 40, or 400 ppb) were injected in four aliquots  
8 into each egg yolk during this window (i.e., at stages 3+, 6, 13, and 17: hours 16, 24, 46, and 68).  
9 Embryos were harvested at 72 hours, 3.5 days, 4 days or 4.25 days (HH stages 18, 21, 23, or 24,  
10 respectively) and evaluated for embryonic survival, cardiac function, or cellular parameters.  
11 Doppler ultrasound technology was utilized to assess cardiovascular effects at HH 18, 21, and  
12 23. In contrast with the results of Drake et al. (2006a), all of the functional parameters assessed  
13 (i.e., cardiac cycle length, heart rate, stroke volume, and dorsal aortic and atrioventricular blood  
14 flow) were similar between control and TCE- or TCA-treated embryos. The authors attributed  
15 this difference in response between studies to dependence upon developmental stage at the time  
16 of exposure. In this case, the chick embryo was relatively resistant to TCE when exposure  
17 occurred during early cardiogenic stages, but was extremely vulnerable when TCE exposure  
18 occurred during valvuloseptal morphogenesis. It was opined that this could explain why some  
19 researchers have observed no developmental cardiac effects after TCE exposure to mammalian  
20 models, while others have reported positive associations.  
21

#### 4.8.3.1.7.2. **Amphibian**

22 The developmental toxicity of TCE was evaluated in the *Frog Embryo Teratogenesis*  
23 *Assay: Xenopus* by Fort et al. (1991; 1993). Late *Xenopus laevis* blastulae were exposed to TCE,  
24 with and without exogenous metabolic activation systems, or to TCE metabolites (dichloroacetic  
25 acid, trichloroacetic acid, trichloroethanol, or oxalic acid), and developmental toxicity ensued.  
26 Findings included alterations in embryo growth, and increased types and severity of induced  
27 malformations. Findings included cardiac malformations that were reportedly similar to those  
28 that had been observed in avian studies. It was suggested that a mixed function oxidase-  
29 mediated reactive epoxide intermediate (i.e., TCE-oxide) may play a significant role in observed  
30 developmental toxicity in in vitro tests.

31 Likewise, McDaniel et al. (2004) observed dose-dependent increases in developmental  
32 abnormalities in embryos of four North American amphibian species (wood frogs, green frogs,  
33 American toads, and spotted salamanders) following 96-hour exposures to TCE. Median  
34 effective concentrations (EC<sub>50</sub>) for malformations was 40 mg/L for TCE in green frogs, while

1 American toads were less sensitive (with no EC<sub>50</sub> at the highest concentration tested—85 mg/L).  
2 Although significant mortality was not observed, the types of malformations noted would be  
3 expected to compromise survival in an environmental context.  
4

#### 4.8.3.1.7.3. Invertebrate

5 The response of the daphnid *Ceriodaphnia dubia* to six industrial chemicals, including  
6 TCE, was evaluated by Niederlehner et al. (1998). Exposures were conducted for 6–7 days,  
7 according to standard EPA testing guidelines. Lethality, impairment of reproduction, and  
8 behavioral changes, such as narcosis and abnormal movement, were observed with TCE  
9 exposures. The reproductive sublethal effect concentration value for TCE was found to be  
10 82 µM.  
11

#### 4.8.3.1.8. In vitro studies

12 Rat whole embryo cultures were used by Saillenfait et al. (1995) to evaluate the  
13 embryotoxicity of TCE, tetrachloroethylene, and four metabolites (trichloroacetic acid,  
14 dichloroacetic acid, chloral hydrate, and trichloroacetyl chloride). In this study, explanted  
15 embryos of Sprague-Dawley rats were cultured in the presence of the test chemicals for 46 hours  
16 and subsequently evaluated. Concentration-dependant decreases in growth and differentiation,  
17 and increases in the incidence of morphologically abnormal embryos were observed for TCE at  
18  $\geq 5$  mM.

19 Whole embryo cultures were also utilized by Hunter et al. (1996) in evaluating the  
20 embryotoxic potential of a number of disinfection by-products, including the TCE metabolites  
21 DCA and TCA. CD-1 mouse conceptuses (GD 9; 3–6 somites) were cultured for 24–26 hours in  
22 treated medium. DCA levels assessed were 0, 734, 1,468, 4,403, 5,871, 7,339, 11,010, or  
23 14,680 µM; TCA levels assessed were 0, 500, 1,000, 2,000, 3,000, 4,000, 5,000 µM. For DCA,  
24 neural tube defects were observed at levels of  $\geq 5,871$  µM, heart defects were observed at  
25  $\geq 7,339$  µM, and eye defects were observed at levels of  $\geq 11,010$  µM. For TCA, neural tube  
26 defects were observed at levels of  $\geq 2,000$  µM, heart and eye defects were observed at  
27  $\geq 3,000$  µM. The heart defects for TCA were reported to include incomplete looping, a reduction  
28 in the length of the heart beyond the bulboventricular fold, and a marked reduction in the caliber  
29 of the heart tube lumen. Overall benchmark concentrations (i.e., the lower limit of the 95%  
30 confidence interval required to produce a 5% increase in the number of embryos with neural tube  
31 defects) were 2,451.9 µM for DCA and 1,335.8 µM for TCA (Richard and Hunter, 1996).

1 Boyer et al. (2000) used an in vitro chick-AVC culture to test the hypothesis that TCE  
2 might cause cardiac valve and septal defects by specifically perturbing epithelial-mesenchymal  
3 cell transformation of endothelial cells in the AVC and outflow tract areas of the heart. AV  
4 explants from Stage 16 White Leghorn chick embryos were placed in hydrated collagen gels,  
5 with medium and TCE concentrations of 0, 50, 100, 150, 200, or 250 ppm. TCE was found to  
6 block the endothelial cell-cell separation process that is associated with endothelial activation as  
7 well as to inhibit mesenchymal cell formation across all TCE concentrations tested. TCE did  
8 not, however, have an effect on the cell migration rate of fully formed mesenchymal cells. TCE-  
9 treatment was also found to inhibit the expression of transformation factor Mox-1 and  
10 extracellular matrix protein fibrillin 2, two protein markers of epithelial-mesenchyme cell  
11 transformation.

#### 4.8.3.1.9. Discussion/Synthesis of Developmental Data

12 In summary, an overall review of the weight of evidence in humans and experimental  
13 animals is suggestive of the potential for developmental toxicity with TCE exposure. A number  
14 of developmental outcomes have been observed in the animal toxicity and the epidemiological  
15 data, as discussed below. These include adverse fetal/birth outcomes including death  
16 (spontaneous abortion, perinatal death, pre- or postimplantation loss, resorptions), decreased  
17 growth (low birth weight, small for gestational age, intrauterine growth restriction, decreased  
18 postnatal growth), and congenital malformations, in particular cardiac defects. Postnatal  
19 developmental outcomes include developmental neurotoxicity, developmental immunotoxicity,  
20 and childhood cancer.

21

#### 4.8.3.1.10. Adverse fetal and early neonatal outcomes

22 Studies that demonstrate adverse fetal or early neonatal outcomes are summarized in  
23 Table 4-102. In human studies of prenatal TCE exposure, increased risk of spontaneous abortion  
24 was observed in some studies (ATSDR, 2001; Taskinen et al., 1994; Windham et al., 1991), but  
25 not in others (ATSDR, 2001, 2008; Goldberg et al., 1990; Lagakos et al., 1986; Lindbohm et al.,  
26 1990; Taskinen et al., 1989). In addition, perinatal deaths were observed after 1970, but not  
27 before 1970 (Lagakos et al., 1986). In rodent studies that examined offspring viability and  
28 survival, there was an indication that TCE exposure may have resulted in increased pre-and/or  
29 postimplantation loss (Healy et al., 1982; Kumar et al., 2000b; Narotsky and Kavlock, 1995), and  
30 in reductions in live pups born as well as in postnatal and postweaning survival (George et al.,  
31 1985, 1986).



1           Decreased birth weight and small for gestational age was observed (ATSDR, 1998b,  
 2 2006b; Rodenbeck et al., 2000; Windham et al., 1991), however, no association was observed in  
 3 other studies (Bove, 1996; Bove et al., 1995; Lagakos et al., 1986). While comprising both  
 4 occupational and environmental exposures, these human studies are overall not highly  
 5 informative due to their small numbers of cases and limited exposure characterization or to the  
 6 fact that exposures to mixed solvents were involved. However, decreased fetal weight, live birth  
 7 weights and postnatal growth were also observed in rodents, (George et al., 1985, 1986; Healy et  
 8 al., 1982; Narotsky and Kavlock, 1995), adding to the weight of evidence for this endpoint. It is  
 9 noted that the rat studies reporting effects on fetal or neonatal viability and growth used Fischer  
 10 344 or Wistar rats, while several other studies, which used Sprague-Dawley rats, reported no  
 11 increased risk in these developmental measures (Carney et al., 2006; Hardin et al., 1981;  
 12 Schwetz et al., 1975).

13  
 14           **Table 4-102. Summary of adverse fetal and early neonatal outcomes**  
 15           **associated with TCE exposures**  
 16

Positive finding	Species	Citation
Spontaneous abortion, miscarriage, pre- and/or postimplantation loss	Human	ATSDR (2001) <sup>a</sup> Taskinen et al. (1994) <sup>a</sup> Windham et al. (1991)
	Rat	Kumar et al. (2000b) Healy et al. (1982) Narotsky and Kavlock (1995) Narotsky et al. (1995)
Perinatal death, reduction in live births	Human	Lagakos et al. (1986) <sup>b</sup>
	Mouse	George et al. (1985)
	Rat	George et al. (1986)
Postnatal and postweaning survival	Mouse	George et al. (1985)
	Rat	George et al. (1986)
Decreased birth weight, small for gestational age, postnatal growth	Human	ATSDR (1998b) ATSDR (2006a) Rodenbeck et al. (2000) <sup>c</sup> Windham et al. (1991)
	Mouse	George et al. (1985)
	Rat	George et al. (1986) Healy et al. (1982) Narotsky and Kavlock (1995) Narotsky et al. (1995)

- 1 <sup>a</sup>Not significant.
- 2 <sup>b</sup>Observed for exposures after 1970, but not before.
- 3 <sup>c</sup>Increased risk for very low birth weight but not low birth weight or full-term low birth weight.

4  
5  
6  
7  
8  
9

Overall, based on weakly suggestive epidemiologic data and fairly consistent laboratory animal data, it can be concluded that TCE exposure poses a potential hazard for prenatal losses and decreased growth or birth weight of offspring.

**4.8.3.1.11. Cardiac malformations**

10 A discrete number of epidemiological studies and studies in laboratory animal models  
11 have identified an association between TCE exposures and cardiac defects in developing  
12 embryos and/or fetuses. These are listed in Table 4-103. Additionally, a number of avian and  
13 rodent in vivo studies and in vitro assays have examined various aspects of the induction of  
14 cardiac malformations.

15  
16  
17  
18

**Table 4-103. Summary of studies that identified cardiac malformations associated with TCE exposures**

<b>Finding</b>	<b>Species</b>	<b>Citations</b>
Cardiac defects	Human	ATSDR (2006a, 2008) Yauck et al. (2004)
	Rat	Dawson et al. (1993; 1990b) Johnson et al. (2003) (2005) Johnson et al. (1998a; 1998b) <sup>a</sup> Smith et al. (1989), <sup>a</sup> (1992) <sup>a</sup> Epstein et al. (1992) <sup>a</sup>
	Chicken	Bross et al. (1983) Boyer et al. (2000) Loeber et al. (1988) Drake et al. (2006a; 2006b) Mishima et al. (2006) Rufer et al. (2008; 2010)
Altered heart rate	Human	Jasinka (1965, translation)

19  
20  
21  
22

<sup>a</sup>Metabolites of TCE.

1 In humans, an increased risk of cardiac defects has been observed after exposure to TCE  
2 in studies reported by ATSDR (ATSDR, 2006a, 2008) and Yauck et al. (2004), although others  
3 saw no significant effect (Bove, 1996; Bove et al., 1995; Goldberg et al., 1990; Lagakos et al.,  
4 1986), possibly due to a small number of cases. In addition, altered heart rate was seen in one  
5 study (Jasińska, 1965, translation). A cohort of water contamination in Santa Clara County, CA  
6 is often cited as a study of TCE exposure and cardiac defects; however, the chemical of exposure  
7 is in fact trichloroethane, not TCE (Deane et al., 1989; Swan et al., 1989).

8 In laboratory animal models, avian studies were the first to identify adverse effects of  
9 TCE exposure on cardiac development. As described in Section 4.8.2.2.1, cardiac malformations  
10 have been reported in chick embryos exposed to TCE (Boyer et al., 2000; Bross et al., 1983;  
11 Drake et al., 2006a; Drake et al., 2006b; Loeber et al., 1988; Mishima et al., 2006; Rufer et al.,  
12 2008). Additionally, a number of studies were conducted in rodents in which cardiac  
13 malformations were observed in fetuses following the oral administration of TCE to maternal  
14 animals during gestation (Dawson et al., 1993; Dawson et al., 1990b; Johnson et al., 2003) 2005;  
15 see Section 4.8.2.2.1.2). Cardiac defects were also observed in rats following oral gestational  
16 treatment with metabolites of TCE (Epstein et al., 1992; Johnson et al., 1998a; Johnson et al.,  
17 1998b; Smith et al., 1992; Smith et al., 1989).

18 However, cardiac malformations were not observed in a number of other studies in  
19 laboratory animals in which TCE was administered during the period of cardiac organogenesis  
20 and fetal visceral findings were assessed. These included inhalation studies in rats (Carney et al.,  
21 2006; Dorfmueller et al., 1979; Hardin et al., 1981; Healy et al., 1982; Schwetz et al., 1975) and  
22 rabbits (Hardin et al., 1981), and oral gavage studies in rats (Fisher et al., 2001; Narotsky and  
23 Kavlock, 1995; Narotsky et al., 1995) and mice (Cosby and Dukelow, 1992).

24 It is generally recognized that response variability among developmental bioassays  
25 conducted with the same chemical agent may be related to factors such as the study design (e.g.,  
26 the species and strain of laboratory animal model used, the day(s) or time of day of dose  
27 administration in relation to critical developmental windows, the route of exposure, the vehicle  
28 used, the day of study termination), or the study methodologies (e.g., how fetuses were  
29 processed, fixed, and examined; what standard procedures were used in the evaluation of  
30 morphological landmarks or anomalies, and whether there was consistency in the fetal  
31 evaluations that were conducted). In the case of studies that addressed cardiac malformations,  
32 there is additional concern as to whether detailed visceral observations were conducted, whether  
33 or not cardiac evaluation was conducted using standardized dissection procedures (e.g., with the  
34 use of a dissection microscope or including confirmation by histopathological evaluation, and  
35 whether the examinations were conducted by technicians who were trained and familiar with  
36 fetal cardiac anatomy). Furthermore, interpretation of the findings can be influenced by the

1 analytical approaches applied to the data as well as by biological considerations such as the  
2 historical incidence data for the species and strain of interest. These issues have been critically  
3 examined in the case of the TCE developmental toxicity studies (Hardin et al., 2005; Watson et  
4 al., 2006).

5 In the available animal developmental studies with TCE, differences were noted in the  
6 procedures used to evaluate fetal cardiac morphology following TCE gestational exposures  
7 across studies, and some of these differences may have resulted in inconsistent fetal outcomes  
8 and/or the inability to detect cardiac malformations. Most of the studies that did not identify  
9 cardiac anomalies used a traditional free-hand sectioning technique (as described in Wilson,  
10 1965) on fixed fetal specimens (Dorfmueller et al., 1979; Hardin et al., 1981; Healy et al., 1982;  
11 Schwetz et al., 1975). Detection of cardiac anomalies can be enhanced through the use of a fresh  
12 dissection technique as described by Staples (1974) and Stuckhardt and Poppe (1984); a  
13 significant increase in treatment-related cardiac heart defects was observed by Dawson et al.  
14 (1990b) when this technique was used. Further refinement of this fresh dissection technique was  
15 employed by Dawson and colleagues at the University of Arizona (UA), resulting in several  
16 additional studies that reported cardiac malformations (Dawson et al., 1993; Johnson et al., 2003)  
17 2005. However, two studies conducted in an attempt to verify the teratogenic outcomes of the  
18 UA laboratory studies used the same or similar enhanced fresh dissection techniques and were  
19 unable to detect cardiac anomalies (Carney et al., 2006; Fisher et al., 2001). Although the  
20 Carney et al. study was administered via inhalation (a route which has not previously been  
21 shown to produce positive outcomes), the Fisher et al. study was administered orally and  
22 included collaboration between industry and UA scientists. It was suggested that the apparent  
23 differences between the results of the Fisher et al. study and the Dawson et al. (1993) and  
24 Johnson et al. studies may be related to factors such as differences in purity of test substances or  
25 in the rat strains, or differences in experimental design (e.g., oral gavage versus drinking water,  
26 exposure only during the period of organogenesis versus during the entire gestation period, or the  
27 use of a staining procedure).

28 It is notable that all studies that identified cardiac anomalies following gestational  
29 exposure to TCE or its metabolites were (1) conducted in rats and (2) dosed by an oral route of  
30 exposure (gavage or drinking water). Cross-species and route-specific differences in fetal  
31 response may be due in part to toxicokinetic factors. Although a strong accumulation and  
32 retention of TCA was found in the amniotic fluid of pregnant mice following inhalation  
33 exposures to TCE (Ghantous et al., 1986), other toxicokinetic factors may be critical. The  
34 consideration of toxicokinetics in determining the relevance of murine developmental data for  
35 human risk assessment is briefly discussed by Watson et al. (2006). There are differences in the  
36 metabolism of TCE between rodent and humans in that TCE is metabolized more efficiently in

1 rats and mice than humans, and a greater proportion of TCE is metabolized to DCA in rodents  
2 versus to TCA in humans. Studies that examined the induction of cardiac malformations with  
3 gestational exposures of rodents to various metabolites of TCE identified TCA and DCA as  
4 putative cardiac teratogens. Johnson et al. (1998a; 1998b) and Smith et al. (1989) reported  
5 increased incidences of cardiac defects with gestational TCA exposures, while Smith et al.  
6 (1992) and Epstein et al. (1992) reported increased incidences following DCA exposures.

7 In all studies that observed increased cardiac defects, either TCE or its metabolites were  
8 administered during critical windows of in utero cardiac development, primarily during the entire  
9 duration of gestation, or during the period of major organogenesis (e.g., GD 6–15 in the rat).  
10 The study by Epstein et al. (1992) used dosing with DCA on discrete days of gestation and had  
11 identified GDs 9 through 12 as a particularly sensitive period for eliciting high interventricular  
12 septal defects associated with exposures to TCE or its metabolites.

13 In the oral studies that identified increased incidences of cardiac malformations following  
14 gestational exposure to TCE, there was a broad range of administered doses at which effects  
15 were observed. In drinking water studies, Dawson et al. (1993) observed cardiac anomalies at  
16 1.5 and 1,100 ppm (with no NOAEL) and Johnson et al. (2003) 2005 reported effects at 250 ppb  
17 (with a NOAEL of 2.5 ppb). One concern is the lack of a clear dose-response for the incidence  
18 of any specific cardiac anomaly or combination of anomalies was not identified, a disparity for  
19 which no reasonable explanation for this disparity has been put forth.

20 The analysis of the incidence data for cardiac defects observed in the Johnson et al.  
21 (2003) 2005 studies has been critiqued (Watson et al., 2006). Issues of concern that have been  
22 raised include the statistical analyses of findings on a per-fetus (rather than the more appropriate  
23 per-litter) basis (Benson, 2004), and the use of nonconcurrent control data in the analysis (Hardin  
24 et al., 2004). In response, the study author has further explained procedures used (Johnson et al.,  
25 2004) and has provided individual litter incidence data to the EPA for independent statistical  
26 analysis (P. Johnson, personal communication, 2008) (see Section 5.1.2.8, dose-response). In  
27 sum, while the studies by Dawson et al. (1993) and Johnson et al. (2003)(2005) have significant  
28 limitations, there is insufficient reason to dismiss their findings.

#### 4.8.3.1.11.1. **Mode of action for cardiac malformations**

30 A number of in vitro studies have been conducted to further characterize the potential for  
31 alterations in cardiac development that have been attributed to exposures with TCE and/or its  
32 metabolites. It was noted that many of the cardiac defects observed in humans and laboratory  
33 species (primarily rats and chickens) involved septal and valvular structures.

1 During early cardiac morphogenesis, outflow tract and AV endothelial cells differentiate  
 2 into mesenchymal cells. These mesenchymal cells have characteristics of smooth muscle-like  
 3 myofibroblasts and form endocardial cushion tissue, which is the primordia of septa and valves  
 4 in the adult heart. Events that take place in cardiac valve formation in mammals and birds are  
 5 summarized by NRC (2006) and reproduced in Table 4-104.

6 Methods have been developed to extract the chick stage 16 atrioventricular canal from the  
 7 embryo and culture it on a hydrated collagen gel for 24–48 hours, allowing evaluation of the  
 8 described stages of cardiac development and their response to chemical treatment. Factors that  
 9 have been shown to influence the induction of endocardial cushion tissue include molecular  
 10 components such as fibronectin, laminin, and galactosyltransferase (Loeber and Runyan, 1990;  
 11 Mjaatvedt et al., 1987), components of the extracellular matrix (Mjaatvedt et al., 1991), and  
 12 smooth muscle  $\alpha$ -actin and transforming growth factor  $\beta$ 3 (Nakajima et al., 1997; Ramsdell and  
 13 Markwald, 1997).

14 Boyer et al. (2000) utilized the in vitro chick AVC culture system to examine the  
 15 molecular mechanism of TCE effects on cardiac morphogenesis. AVC explants from stage 16  
 16 chick embryos (15/treatment level) were placed onto collagen gels and treated with 0, 50, 100,  
 17 150, 200, or 250-ppm TCE and incubated for a total of 54 hours. Epithelial-mesenchymal  
 18 transformation, endothelial cell density, cell migration, and immunohistochemistry were  
 19 evaluated. TCE treatment was found to inhibit endothelial cell activation and normal

20 **Table 4-104. Events in cardiac valve formation in mammals and birds<sup>a</sup>**  
 21

Stage and event	Structural description <sup>b</sup>
Early cardiac development	The heart is a hollow, linear, tube-like structure with two cell layers. The outer surface is a myocardial cell layer, and the inner luminal surface is an endothelial layer. Extracellular matrix is between the two cell layers.
Epithelial-mesenchymal cell transformation	A subpopulation of endothelial cells lining the atrioventricular canal detaches from adjacent cells and invades the underlying extracellular matrix. Three events occur <ul style="list-style-type: none"> <li>➤ Endothelial cell activation (avian stage 14)</li> <li>➤ Mesenchymal cell formation (avian stage 16)</li> <li>➤ Mesenchymal cell migration into the extracellular matrix (avian stages 17 and 18).</li> </ul>
Mesenchymal cell migration and proliferation	Endothelial-derived mesenchymal cells migrate toward the surrounding myocardium and proliferate to populate the AVC extracellular matrix.
Development of septa and valvular structures	Cardiac mesenchyme provides cellular constituents for <ul style="list-style-type: none"> <li>➤ Septum intermedium</li> <li>➤ Valvular leaflets of the mitral and tricuspid AV valves.</li> </ul> The septum intermedium subsequently contributes to <ul style="list-style-type: none"> <li>➤ Lower portion of the interatrial septum</li> <li>➤ Membranous portion of the interventricular septum.</li> </ul>

22 <sup>a</sup>As summarized in NRC (2006).

23 <sup>b</sup>Markwald et al. (1996; 1984), Boyer et al. (2000).  
 24

1  
2  
3 mesenchymal cell transformation, endothelial cell-cell separation, and protein marker expression  
4 (i.e., transcription factor Mox-1 and extracellular matrix protein fibrillin 2). Mesenchymal cell  
5 migration was not affected, nor was the expression of smooth muscle  $\alpha$ -actin. The study authors  
6 proposed that TCE may cause cardiac valvular and septal malformations by inhibiting  
7 endothelial separation and early events of mesenchymal cell formation. Hoffman et al. (2004)  
8 has proposed alternatively that TCE may be affecting the adhesive properties of the endocardial  
9 cells. No experimental data are currently available that address the levels of TCE in cardiac  
10 tissue in vivo, resulting in some questions (Dugard, 2000) regarding the relevance of these  
11 mechanistic findings to human health risk assessment.

12 In a study by Mishima et al. (2006), White Leghorn chick whole embryo cultures (stage  
13 13 and 14) were used to assess the susceptibility of endocardial epithelial-mesenchymal  
14 transformation in the early chick heart to TCE at analytically determined concentrations of 0, 10,  
15 20, 40, or 80 ppm. This methodology maintained the anatomical relationships of developing  
16 tissues and organs, while exposing precisely staged embryos to quantifiable levels of TCE and  
17 facilitating direct monitoring of developmental morphology. Following 24 hours of incubation  
18 the numbers of mesenchymal cells in the inferior and superior AV cushions were counted. TCE  
19 treatment significantly reduced the number of mesenchymal cells in both the superior and  
20 inferior AV cushions at 80 ppm.

21 Ou et al. (2003) examined the possible role of endothelial nitric oxide synthase (which  
22 generates nitric oxide that has an important role in normal endothelial cell proliferation and  
23 hence normal blood vessel growth and development) in TCE-mediated toxicity. Cultured  
24 proliferating bovine coronary endothelial cells were treated with TCE at 0–100  $\mu$ M and  
25 stimulated with a calcium ionophore to determined changes in endothelial cells and the  
26 generation of endothelial nitric oxide synthase, nitric oxide, and superoxide anion. TCE was  
27 shown to alter heat shock protein interactions with endothelial nitric oxide synthase and induce  
28 endothelial nitric oxide synthase to shift nitric oxide to superoxide-anion generation. These  
29 findings provide insight into how TCE impairs endothelial proliferation.

30 Several studies have also identified a TCE-related perturbation of several proteins  
31 involved in regulation of intracellular  $\text{Ca}^{2+}$ . After 12 days of maternal exposure to TCE in  
32 drinking water, *Serca2a* (sarcoendoplasmic reticulum  $\text{Ca}^{2+}$  ATPase) mRNA expression was  
33 reduced in rat embryo cardiac tissues (Collier et al., 2003). Selmin et al. (2008) conducted a  
34 microarray analysis of a P19 mouse stem cell line exposed to 1-ppm TCE in vitro, identifying  
35 altered expression of *Ryr2* (ryanodine receptor isoform 2), a  $\text{Ca}^{2+}$  release channel that is  
36 important in normal rhythmic heart activity (Gyorke and Terentyev, 2008). Alterations in  $\text{Ca}^{2+}$

1 cycling and resulting contractile dysfunction is a recognized pathogenic mechanism of cardiac  
2 arrhythmias and sudden cardiac death (Leandri et al., 1995; Lehnart et al., 2008; Yano et al.,  
3 2008). Caldwell et al. (2008b) used real-time PCR and digital imaging microscopy to  
4 characterize the effects of various doses of TCE on gene expression and  $Ca^{2+}$  response to  
5 vasopressin in rat cardiac myocytes (H9c2) *Serca2a* and *Ryr2* expression were reduced at 12  
6 and 48 hours following exposure to TCE. Additionally,  $Ca^{2+}$  response to vasopressin was altered  
7 following TCE treatment. Makwana et al. (2010) dosed chick embryos in ovo with 8 or 800 ppb  
8 TCE; RT-PCR analysis of RNA isolated during specific windows of cardiac development  
9 demonstrated effects on the expression of genes associated with reduced blood flow. Although it  
10 has been hypothesized that TCE might interfere with the folic acid/methylation pathway in liver  
11 and kidney and alter gene regulation by epigenetic mechanisms, Caldwell et al. (2010) found that  
12 the effects of TCE exposure on normal gene expression in rat embryonic hearts was not altered  
13 by the administration of exogenous folate. Overall, these data suggest that TCE may disrupt the  
14 ability to regulate cellular  $Ca^{2+}$  fluxes, altering blood flow and leading to morphogenic  
15 consequences in the developing heart. This remains an open area of research.

16 Thus, in summary, a number of studies have been conducted in an attempt to characterize  
17 the MOA for TCE-induced cardiac defects. A major research focus has been on disruptions in  
18 cardiac valve formation, using avian in ovo and in vitro studies. These studies demonstrated  
19 treatment-related alterations in endothelial cushion development that could plausibly be  
20 associated with defects involving septal and valvular morphogenesis in rodents and chickens.  
21 However, a broad array of cardiac malformations has been observed in animal models following  
22 TCE exposures (Dawson et al., 1993; Johnson et al., 2003) 2005, and other evidence of  
23 molecular disruption of  $Ca^{2+}$  during cardiac development has been examined (Caldwell et al.,  
24 2008b; Collier et al., 2003; Selmin et al., 2008) suggesting the possible existence of multiple  
25 MOAs. The observation of defective myocardial development in a mouse model deficient for  
26 gp130, a signal transducer receptor for IL-6 (Yoshida et al., 1996), suggests the potential  
27 involvement of immune-mediated effects.

28

#### 4.8.3.1.11.2. Association of peroxisome proliferator activated receptor alpha (PPAR $\alpha$ ) with developmental outcomes

29 The PPARs are ligand activated receptors that belong to the nuclear hormone receptor  
30 family. Three isotypes have been identified (PPAR $\alpha$ , PPAR $\delta$  [also known as PPAR $\beta$ ], and  
31 PPAR $\gamma$ ). These receptors, upon binding to an activator, stimulate the expression of target genes  
32 implicated in important metabolic pathways. In rodents, all three isotypes show specific time  
33 and tissue-dependent patterns of expression during fetal development and in adult animals. In



1 development, they have been especially implicated in several aspects of tissue differentiation,  
2 e.g., of the adipose tissue, brain, placenta and skin. Epidermal differentiation has been linked  
3 strongly with PPAR $\alpha$  and PPAR $\delta$  (Michalik et al., 2002). PPAR $\alpha$  starts late in development,  
4 with increasing levels in organs such as liver, kidney, intestine, and pancreas; it is also  
5 transiently expressed in fetal epidermis and CNS (Braissant and Wahli, 1998) and has been  
6 linked to phthalate-induced developmental and testicular toxicity (Corton and Lapinskas, 2005).  
7 Liver, kidney, and heart are the sites of highest PPAR $\alpha$  expression (Toth et al., 2007). PPAR $\delta$   
8 and PPAR $\gamma$  have been linked to placental development and function, with PPAR $\gamma$  found to be  
9 crucial for vascularization of the chorioallantoic placenta in rodents (Wendling et al., 1999), and  
10 placental anomalies mediated by PPAR $\gamma$  have been linked to rodent cardiac defects (Barak et al.,  
11 2008). While it might be hypothesized that there is some correlation between PPAR signaling,  
12 fetal deaths, and/or cardiac defects observed following TCE exposures in rodents, no definitive  
13 data have been generated that elucidate a possible PPAR-mediated MOA for these outcomes.  
14

#### 4.8.3.1.11.3. **Summary of the weight of evidence on cardiac malformations**

15 The evidence for an association between TCE exposures in the human population and the  
16 occurrence of congenital cardiac defects is not particularly strong. Many of the epidemiological  
17 study designs were not sufficiently robust to detect exposure-related birth defects with a high  
18 degree of confidence. However, two well-conducted studies by ATSDR (2006a, 2008) clearly  
19 demonstrated an elevation in cardiac defects. It could be surmised that the identified cardiac  
20 defects were detected because they were severe, and that additional cases with less severe  
21 cardiac anomalies may have gone undetected.

22 The animal data provide strong, but not unequivocal, evidence of the potential for  
23 TCE-induced cardiac malformations following oral exposures during gestation. Strengths of the  
24 evidence are the duplication of the adverse response in several studies from the same laboratory  
25 group, detection of treatment-related cardiac defects in both mammalian and avian species (i.e.,  
26 rat and chicken), general cross-study consistency in the positive association of increased cardiac  
27 malformations with test species (i.e., rat), route of administration (i.e., oral), and the  
28 methodologies used in cardiac morphological evaluation (i.e., fresh dissection of fetal hearts).  
29 Furthermore, when differences in response are observed across studies they can generally be  
30 attributed to obvious methodological differences, and a number of in ovo and in vitro studies  
31 demonstrate a consistent and biologically plausible MOA for one type of malformation observed.  
32 Weaknesses in the evidence include lack of a clear dose-related response in the incidence of

1 cardiac defects, and the broad variety of cardiac defects observed, such that they cannot all be  
2 grouped easily by type or etiology.

3 Taken together, the epidemiological and animal study evidence raise sufficient concern  
4 regarding the potential for developmental toxicity (increased incidence of cardiac defects) with  
5 in utero TCE exposures.

#### 4.8.3.1.12. Other structural developmental outcomes

6 A summary of other structural developmental outcomes that have been associated with  
7 TCE exposures is presented in Table 4-105.

8 In humans, a variety of birth defects other than cardiac have been observed. These  
9 include total birth defects (ATSDR, 2001; Bove, 1996; Bove et al., 1995; Flood, 1988) CNS  
10 birth defects (ATSDR, 2001; Bove, 1996; Bove et al., 1995; Lagakos et al., 1986), eye/ear birth  
11 anomalies (Lagakos et al., 1986); oral cleft defects (Bove, 1996; Bove et al., 1995; Lagakos et  
12 al., 1986; Lorente et al., 2000); kidney/urinary tract disorders (Lagakos et al., 1986);  
13 musculoskeletal birth anomalies (Lagakos et al., 1986); anemia/blood disorders (Burg and Gist,  
14 1999); and lung/respiratory tract disorders (Lagakos et al., 1986). While some of these results  
15 were statistically significant, they have not been reported elsewhere. Occupational cohort  
16 studies, while not reporting positive results, are generally limited by the small number of  
17 observed or expected cases of birth defects (Lorente et al., 2000; Taskinen et al., 1989; Tola et  
18 al., 1980).

19  
20  
21 **Table 4-105. Summary of other structural developmental outcomes**  
22 **associated with TCE exposures**  
23

Finding	Species	Citations
Eye/ear birth anomalies	Human	Lagakos et al. (1986)
	Rat	Narotsky (1995) Narotsky and Kavlock (1995)
Oral cleft defects	Human	Bove (1996) Bove et al. (1995) Lagakos et al. (1986) Lorente et al. (2000)
Kidney/urinary tract disorders	Human	Lagakos et al. (1986)
Musculoskeletal birth anomalies	Human	Lagakos et al. (1986)

Anemia/blood disorders	Human	Burg and Gist (1999)
Lung/respiratory tract disorders	Human	Lagakos et al. (1986)
	Mouse	Das and Scott (1994)
Skeletal	Rat	Healy et al. (1982)
Other <sup>a</sup>	Human	ATSDR (2001)

<sup>a</sup>As reported by the authors.

In experimental animals, a statistically significant increase in the incidence of fetal eye defects, primarily microphthalmia and anophthalmia, manifested as reduced or absent eye bulge, was observed in rats following gavage administration of 1,125 mg/kg-day TCE during the period of organogenesis (Narotsky and Kavlock, 1995; Narotsky et al., 1995). Dose-related nonsignificant increases in the incidence of Fischer 344 rat pups with eye defects were also observed at lower dose levels (101, 320, 475, 633, and 844 mg/kg-day) in the Narotsky et al. (1995) study (also reported in Barton and Das, 1996). However, no other developmental or reproductive toxicity studies identified abnormalities of eye development following TCE exposures. For example, in a study reported by Warren et al. (2006), extensive computerized morphometric ocular evaluation was conducted in Sprague-Dawley rat fetuses that had been examined for cardiac defects by Fisher et al. (2001); the dams had been administered TCE (500 mg/kg-day), DCA (300 mg/kg-day), or TCA (300 mg/kg-day) during GDs 6–15. No ocular defects were found with TCE exposures; however, significant reductions in the lens area, globe area, and interocular distance were observed with DCA exposures, and nonsignificant decreases in these measures as well as the medial canthus distance were noted with TCA exposures. Developmental toxicity studies conducted by Smith et al. (1992; 1989) also identified orbital defects (combined soft tissue and skeletal abnormalities) in Long Evans rat fetuses following GD 6–15 exposures with TCA and DCA (statistically or biologically significant at  $\geq 800$  and  $\geq 900$  mg/kg-day, respectively). Overall, the study evidence indicates that TCE and its oxidative metabolites can disrupt ocular development in rats. In addition to the evidence of alteration to the normal development of ocular structure, these findings may also be an indicator of disruptions to nervous system development. It has been suggested by Warren et al. (2006) and Williams and DeSesso (2008) that the effects of concern (defined as statistically significant outcomes) are observed only at high dose levels and are not relevant to risk assessment for environmental exposures. On the other hand, Barton and Das (1996) point out that benchmark dose modeling of the quantal eye defect incidence data provides a reasonable approach to the development of oral toxicity values for TCE human health risk assessment. It is also noted that

1 concerns may exist not only for risks related to low level environmental exposures, but also for  
2 risks resulting from acute or short-term occupational or accidental exposures, which may be  
3 associated with much higher inadvertent doses.

4 It was also notable that a study using a single intraperitoneal dose of 3,000 mg/kg TCE to  
5 mice during late gestation (GD 17) identified apparent delays in lung development and increased  
6 neonatal mortality (Das and Scott, 1994). No further evaluation of this outcome has been  
7 identified in the literature.

8 Healy et al. (1982) did not identify any treatment-related fetal malformations following  
9 inhalation exposure of pregnant inbred Wistar rats to 0 or 100 ppm (535 mg/m<sup>3</sup>) on GD 8–21. In  
10 this study, significant differences between control and treated litters were observed as an  
11 increased incidence of minor ossification variations ( $p = 0.003$ ) (absent or bipartite centers of  
12 ossification).

#### 4.8.3.1.13. Developmental neurotoxicity

14 Studies that address effects of TCE on the developing nervous system are discussed in  
15 detail in Section 4.3, addressed above in the sections on human developmental toxicity (see  
16 Section 4.8.3) and on mammalian studies (see Section 4.8.3.2.1) by route of exposure, and  
17 summarized in Table 4-106. The available data collectively suggest that the developing brain is  
18 susceptible to TCE exposures.

19 In humans, CNS birth defects were observed in a few studies (ATSDR, 2001; Bove,  
20 1996; Bove et al., 1995; Lagakos et al., 1986). Postnatally, observed adverse effects in humans  
21 include delayed newborn reflexes following use of TCE during childbirth (Beppu, 1968),  
22 impaired learning or memory (Bernad et al., 1987, abstract; White et al., 1997); aggressive

23 **Table 4-106. Summary of developmental neurotoxicity associated with TCE**  
24 **exposures**  
25

Positive findings	Species	Citations
CNS defects, neural tube defects	Human	ATSDR (2001)
		Bove (1996); Bove et al. (1995)
		Lagakos et al. (1986)
Eye defects	Rat	Narotsky (1995); Narotsky and Kavlock (1995)
Delayed newborn reflexes	Human	Beppu (1968)
Impaired learning or memory	Human	Bernad et al. (1987), abstract

		White et al. (1997)
Aggressive behavior	Human	Bernad et al. (1987), abstract
	Rat	Blossom et al. (2008)
Hearing impairment	Human	ATSDR (2003a); Burg et al. (1995); Burg and Gist (1999)
		Beppu (1968)
Speech impairment	Human	ATSDR (2003a); Burg et al. (1995); Burg and Gist (1999)
		White et al. (1997)
Encephalopathy	Human	White et al. (1997)
Impaired executive function	Human	White et al. (1997)
Impaired motor function	Human	White et al. (1997)
Attention deficit	Human	Bernad et al. (1987), abstract
ASD	Human	Windham et al. (2006)
Delayed or altered biomarkers of CNS development	Rat	Isaacson and Taylor (1989); Noland-Gerbec et al. (1986); Westergren et al. (1984)
Behavioral alterations	Mice	Blossom et al. (2008); Fredriksson et al. (1993)
	Rat	George et al. (1986); Taylor et al. (1985)

1  
2  
3 behavior (Bernad et al., 1987, abstract); hearing impairment (ATSDR, 2003a; Beppu, 1968; Burg  
4 and Gist, 1999; Burg et al., 1995); speech impairment (Burg and Gist, 1999; Burg et al., 1995;  
5 White et al., 1997); encephalopathy (White et al., 1997); impaired executive and motor function  
6 (White et al., 1997); attention deficit (Bernad et al., 1987, abstract; White et al., 1997), and  
7 autism spectrum disorder (Windham et al., 2006). While there are broad developmental  
8 neurotoxic effects that have been associated with TCE exposure, there are many limitations in  
9 the studies.

10 More compelling evidence for the adverse effect of TCE exposure on the developing  
11 nervous system is found in the animal study data, although a rigorous evaluation of potential  
12 outcomes has not been conducted. For example, there has not been an assessment of cognitive  
13 function (i.e., learning and memory) following developmental exposures to TCE, nor have most

1 of the available studies characterized the pre- or postnatal exposure of the offspring to TCE or its  
2 metabolites. Nevertheless, there is evidence of treatment-related alterations in brain  
3 development and in behavioral parameters (e.g., spontaneous motor activity and social  
4 behaviors) associated with exposures during neurological development. The animal study  
5 database includes the following information: Following inhalation exposures of 150 ppm to mice  
6 during mating and gestation, the specific gravity of offspring brains were significantly decreased  
7 at postnatal time points through the age of weaning; however, this effect did not persist to  
8 1 month of age (Westergren et al., 1984). In studies reported by Taylor et al. (1985), Isaacson  
9 and Taylor (1989), and Noland-Gerbec et al. (1986), 312 mg/L exposures in drinking water that  
10 were initiated 2 weeks prior to mating and continued to the end of lactation resulted,  
11 respectively, in (a) significant increases in exploratory behavior at GDs 60 and 90, (b) reductions  
12 in myelination in the brains of offspring at weaning, and (c) significantly decreased uptake of  
13 2-deoxyglucose in the neonatal rat brain (suggesting decreased neuronal activity). Ocular  
14 malformations in rats observed by Narotsky (1995) and Narotsky and Kavlock (1995) following  
15 maternal gavage doses of 1,125 mg/kg-day during gestation may also be indicative of alterations  
16 of nervous system development. Gestational exposures to mice (Fredriksson et al., 1993)  
17 resulted in significantly decreased rearing activity on GD 60, and dietary exposures during the  
18 course of a continuous breeding study in rats (George et al., 1986) found a significant trend  
19 toward increased time to cross the first grid in open field testing. In a study by Blossom et al.  
20 (2008), alterations in social behaviors (deficits in nest-building quality and increased aggression  
21 in males) were observed in pubertal-age MRL +/- mice that had been exposed to 0.1 mg/mL  
22 TCE via drinking water during prenatal and postnatal development (until PND 42). Dorfmueller  
23 et al. (1979) was the only study that assessed neurobehavioral endpoints following in utero  
24 exposure (maternal inhalation exposures of 1,800 ± 200 ppm during gestation) and found no  
25 adverse effects that could be attributed to TCE exposure. Specifically, an automated assessment  
26 of ambulatory response in a novel environment on GDs 10, 20 and 100, did not identify any  
27 effect on general motor activity of offspring.

28

#### **4.8.3.1.14. Developmental immunotoxicity**

29 Studies that address the developmental immunotoxic effects of TCE are discussed in  
30 detail in Section 4.6, addressed above in the sections on human developmental toxicity (see  
31 Section 4.8.3) and on mammalian studies (see Section 4.8.3.2.1) by route of exposure, and  
32 summarized in Table 4-107.

33

34

**Table 4-107. Summary of developmental immunotoxicity associated with TCE exposures**

<b>Finding</b>	<b>Species (strain)</b>	<b>Citations</b>
Significant reduction in Th1 IL-2 producing cells	Human	Lehmann et al. (2002)
Altered immune response	Human	Byers et al. (1988)
Suppression of PFC responses, increased T-cell subpopulations, decreased spleen cellularity, and increased hypersensitivity response	Mouse (B6C3F1)	Peden-Adams et al. (2006)
Altered splenic and thymic T-cell subpopulations	Mouse (MRL +/+)	Peden-Adams et al. (2008)
Altered thymic T-cell subpopulations; transient increased proinflammatory cytokine production by T-cells; increased autoantibody levels and IgG	Mouse (MRL +/+)	Blossom and Doss (2007)
Increased proinflammatory cytokine production by T-cells	Mouse (MRL +/+)	Blossom et al. (2008)

Two epidemiological studies that addressed potential immunological perturbations in children that were exposed to TCE were reported by Lehmann et al. (2001; 2002). In the 2001 study, no association was observed between TCE and allergic sensitization to egg white and milk, or to cytokine producing peripheral T-cells, in premature neonates and 36-month-old neonates that were at risk of atopy. In the 2002 study, there was a significant reduction in Th1 IL-2 producing cells. Another study observed altered immune response in family members of those diagnosed with childhood leukemia, including 13 siblings under age 19 at the time of exposure, but an analysis looking at only these children was not done (Byers et al., 1988).

Several studies were identified (Blossom and Doss, 2007; Blossom et al., 2008; Peden-Adams et al., 2006; Peden-Adams et al., 2008) which assessed the potential for developmental immunotoxicity in mice following oral (drinking water) TCE exposures during critical pre- and postnatal stages of immune system development. Peden-Adams et al. (2006) noted evidence of immune system perturbation (suppression of PFC responses, increased T-cell subpopulations, decreased spleen cellularity, and increased hypersensitivity response) in B6C3F1 offspring following in utero and 8 weeks of postnatal exposures to TCE. Evidence of autoimmune response was not observed in the offspring of this nonautoimmune-prone strain of mice. However, in a study by Peden-Adams et al. (2008) MRL +/+ mice, which are autoimmune-prone, were exposed from conception until 12 months of age. Consistent with the Peden-Adams

1 et al. (2006) study, no evidence of increased autoantibody levels was observed in the offspring.  
2 In two other studies focused on autoimmune responses following drinking water exposures of  
3 MRL +/+ mice to TCE during in utero development and continuing until the time of sexual  
4 maturation, Blossom and Doss (2007) and Blossom et al. (2008) reported some peripheral blood  
5 changes that were indicative of treatment-related autoimmune responses in offspring. Positive  
6 response levels were 0.5 and 2.5 mg/mL for Blossom and Doss (2007) and 0.1 mg/mL for  
7 Blossom et al. (2008). None of these studies were designed to extensively evaluate recovery,  
8 latent outcomes, or differences in severity of response that might be attributed to the early life  
9 exposures. Consistency in response in these animal studies was difficult to ascertain due to the  
10 variations in study design (e.g., animal strain used, duration of exposure, treatment levels  
11 evaluated, timing of assessments, and endpoints evaluated). Likewise, the endpoints assessed in  
12 the few epidemiological studies that evaluated immunological outcomes following  
13 developmental exposures to TCE were dissimilar from those evaluated in the animal models, and  
14 so provided no clear cross-species correlation. The most sensitive immune system response  
15 noted in the studies that exposed developing animals were the decreased PFC and increased  
16 hypersensitivity observed by Peden-Adams et al. (2006); treatment-related outcomes were noted  
17 in mice exposed in the drinking water at a concentration of 1,400 ppb. None of the other studies  
18 that treated mice during immune system development assessed these same endpoints; therefore,  
19 direct confirmation of these findings across studies was not possible. It is noted, however, that  
20 similar responses were not observed in studies in which adult animals were administered TCE  
21 (e.g., Woolhiser et al., 2006), suggesting increased susceptibility in the young. Differential  
22 lifestage-related responses have been observed with other diverse chemicals (e.g.,  
23 diethylstilbestrol; diazepam; lead; 2,3,7,8-tetrachlorobenzo-*p* dioxin; and tributyltin oxide) in  
24 which immune system perturbations were observed at lower doses and/or with greater  
25 persistence when tested in developing animals as compared to adults (Luebke et al., 2006).  
26 Thus, such an adverse response with TCE exposure is considered biologically plausible and an  
27 issue of concern for human health risk assessment.  
28

#### 4.8.3.1.15. Childhood cancers

29 A summary of childhood cancers that have been associated with TCE exposures  
30 discussed above is presented in Table 4-108. A summary of studies that observed childhood  
31 leukemia is also discussed in detail in Section 4.6.1.3 and Section 4.8.3.1.2.4 contains details of  
32 epidemiologic studies on childhood brain cancer.  
33



1  
2

**Table 4-108. Summary of childhood cancers associated with TCE exposures**

Finding	Species	Citations
Leukemia	Human	AZ DHS (1990; Flood, 1988)
		AZ DHS (Kioski et al., 1990a)
		Cohn et al. (1994b)
		Cutler et al. (1986); Costas et al. (2002); Lagakos et al. (1986); MA DPH (1997a)
		Lowengart et al. (1987)
		McKinney et al. (1991)
		Shu et al. (1999)
Neuroblastoma	Human	De Roos et al. (2001)
		Peters et al. (1985; 1981)

3  
4

5 A nonsignificant increased risk of leukemia diagnosed during childhood has been  
6 observed in a number of studies examining TCE exposure (1990; Flood, 1988; Kioski et al.,  
7 1990a), (Cohn et al., 1994b; Costas et al., 2002; Lagakos et al., 1986; Lowengart et al., 1987;  
8 McKinney et al., 1991; MDPH, 1997a; Shu et al., 1999). However, other studies did not  
9 observed an increased risk for childhood leukemia after TCE exposure (Flood, 1997; Kioski et  
10 al., 1990b; Morgan and Cassady, 2002), possibly due to the limited number of cases or the  
11 analysis based on multiple solvents.

12 CNS cancers during childhood have been reported on in a few studies. Neuroblastomas were not  
13 statistically elevated in one study observing parental exposure to multiple chemicals, including  
14 TCE (De Roos et al., 2001). Brain tumors were observed in another study, but the odds ratio  
15 could not be determined (Peters et al., 1985; Peters et al., 1981). CNS cancers were not elevated  
16 in other studies (Kioski et al., 1990a; Morgan and Cassady, 2002). Other studies did not see an  
17 excess risk of total childhood cancers (ATSDR, 2006a; Morgan and Cassady, 2002).

18 A follow-up study of the Camp Lejeune cohort that will examine childhood cancers  
19 (along with birth defects) was initiated in 1999 (ATSDR, 2003c), is expected to be completed  
20 soon (ATSDR, 2009; U.S. GAO, 2007), and may provide additional insight.

21 No studies of cancers in experimental animals in early lifestages have been identified.

## 4.9. OTHER SITE-SPECIFIC CANCERS

### 4.9.1. Esophageal Cancer

1           Increasing esophageal cancer incidence has been observed in males, but not females in  
2 the United States between 1975 and 2002, a result of increasing incidence of esophageal  
3 adenocarcinoma (Ward et al., 2006). Males also have higher age-adjusted incidence and  
4 mortality rates (incidence, 7.8 per 100,000; mortality, 7.8 per 100,000) than females (incidence,  
5 2.0 per 100,000; mortality, 1.7 per 100,000) (Ries et al., 2008). Survival for esophageal cancer  
6 remains poor and age-adjusted mortality rates are just slightly lower than incidence rates. Major  
7 risk factors associated with esophageal cancer are smoking and alcohol for squamous cell  
8 carcinoma, typically found in the upper third of the esophagus, and obesity, gastroesophageal  
9 reflux, and Barrett’s esophagus for adenocarcinoma that generally occurs in the lower esophagus  
10 (Ward et al., 2006).

11           Seventeen epidemiologic studies on TCE exposure reported relative risks for esophageal  
12 cancer (ATSDR, 2004a, 2006b; Blair et al., 1989; Blair et al., 1998; Boice et al., 1999; Boice et  
13 al., 2006b; Clapp and Hoffman, 2008; Costa et al., 1989; Garabrant et al., 1988; Greenland et al.,  
14 1994; Hansen et al., 2001; Raaschou-Nielsen et al., 2003; Radican et al., 2008; Ritz, 1999a;  
15 Siemiatycki, 1991; Sung et al., 2007; Zhao et al., 2005). Ten studies had high likelihood of TCE  
16 exposure in individual study subjects and were judged to have met, to a sufficient degree, the  
17 standards of epidemiologic design and analysis (Blair et al., 1998; Boice et al., 1999; Boice et al.,  
18 2006b; Greenland et al., 1994; Hansen et al., 2001; Raaschou-Nielsen et al., 2003; Radican et al.,  
19 2008; Ritz, 1999a; Siemiatycki, 1991; Zhao et al., 2005). Four studies with TCE exposure  
20 potential assigned to individual subjects (Anttila et al., 1995; Axelson et al., 1994; Blair et al.,  
21 1998 [Incidence]; Morgan et al., 1998) do not present relative risk estimates for esophageal  
22 cancer and TCE exposure nor do two other studies which carry less weight in the analysis  
23 because of design limitations (Henschler et al., 1995; Sinks et al., 1992). Only Raaschou-  
24 Nielsen et al. (2003) examines esophageal cancer histologic type, an important consideration  
25 given differences between suspected risk factors for adenocarcinoma and those for squamous cell  
26 carcinoma. Appendix B identifies these studies’ design and exposure assessment characteristics.

27           Several population case-control studies (Engel et al., 2002; Gustavsson et al., 1998;  
28 Parent et al., 2000b; Ramanakumar et al., 2008; Santibanez et al., 2008; Weiderpass et al., 2003;  
29 Yu et al., 1988) examine esophageal cancer and organic solvents or occupational job titles with  
30 past TCE use documented (Bakke et al., 2006). Relative risk estimates in case-control studies  
31 that examine metal occupations or job titles, or solvent exposures are found in Table 4-109. The

- 1 lack of exposure assessment to TCE, low prevalence of exposure to chlorinated hydrocarbon
- 2 solvents, or few exposed cases and controls in those studies lowers their sensitivity for informing
- 3 evaluations of TCE and esophageal cancer.

**Table 4-109. Selected observations from case-control studies of TCE exposure and esophageal cancer**

Study population	Exposure group	All esophageal cancers		Squamous cell cancer		Adenocarcinoma		Reference
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	
Population of regions in Eastern Spain								Santibañez et al. (2008)
	Metal molders, welders, etc.	0.94 (0.14, 6.16)	3	0.40 (0.05, 3.18)	2	3.55 (0.28, 44.70)	1	
	Metal-processing plant operators	1.14 (0.29, 4.44)	5	1.23 (0.23, 6.51)	4	0.86 (0.08, 8.63)	1	
	Chlorinated hydrocarbon solvents							
	Low exposure	1.05 (0.15, 7.17)	2		0	4.92 (0.69, 34.66)	2	
	High exposure	1.76 (0.40, 7.74)	6	2.18 (0.41, 11.57)	5	3.03 (0.28, 32.15)	1	
Population of Montreal, Canada								Ramanakumar et al. (2008); Parent et al. (2000b)
	Painter, Metal coatings							
	Any exposure	1.3 (0.4, 4.2)	6					
	Substantial exposure	4.2 (1.1, 17.0)	4					
	Solvents							
	Any exposure	1.1 (0.7, 1.7)	39	1.4 (0.8, 2.5)	30			
	Nonsubstantial exposure	1.0 (0.5, 1.9)	16	1.3 (0.6, 2.6)	12			
	Substantial exposure	1.1 (0.6, 1.9)	39	1.4 (0.8, 2.5)	30			
Population of Sweden								Jansson et al., (2005; 2006)
	Organic solvents							
	No exposure			1.0	145	1.0	128	
	Moderate exposure			0.7 (0.4, 1.5)	15	1.2 (0.6, 2.3)	14	
	High exposure			1.3 (0.7, 2.3)	21	1.4 (0.7, 2.5)	18	
	Test for trend			$p = 0.47$		$p = 0.59$		
	No exposure			1.0		1.0		
	Moderate exposure			0.5 (0.1, 3.9) <sup>a</sup>	1	0.4 (0.1, 1.5) <sup>a</sup>	2	
	High exposure			0.4 (0.1, 1.8) <sup>a</sup>	2	0.9 (0.5, 1.6) <sup>a</sup>	12	
	Test for trend			$p = 0.44$		$p = 0.36$		

**Table 4-109. Selected observations from case-control studies of TCE exposure and esophageal cancer (continued)**

Study population	Exposure group	All esophageal cancers		Squamous cell cancer		Adenocarcinoma		Reference
		Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	Relative risk (95% CI)	No. obs. events	
Population of Finland (Females)								Weiderpass et al., (2003)
	Chlorinated hydrocarbon solvents							
	Low level exposure	0.95 (0.54, 1.66)	Not reported					
	High level exposure	0.62 (0.34, 1.13)	Not reported					
Population of NJ, CT, WA State								Engel et al. (2002)
	Precision metal workers	Not reported		0.7 (0.3, 1.5)	12	1.4 (0.8, 2.3)	25	
	Metal product manufacturing	Not reported		0.8 (0.3, 1.8)	15	1.3 (0.8, 2.3)	26	

<sup>a</sup> Jansson et al. (2006) is a registry-based study of the Swedish Construction Worker Cohort. Relative risks are incidence rate ratios from Cox regression analysis using calendar time and adjustment for attained age, calendar period at entry into the cohort, tobacco smoking status at entry into the cohort and BMI at entry into the cohort.

No. obs. events = number of observable events.

1 Table 4-110 presents risk estimates for TCE exposure and esophageal cancer observed in  
2 cohort, PMR, case-control, and geographic based studies. Ten studies in which there is a high  
3 likelihood of TCE exposure in individual study subjects (e.g., based on job-exposure matrices or  
4 biomarker monitoring) reported risk estimates for esophageal cancer (Blair et al., 1998; Boice et  
5 al., 1999; Boice et al., 2006b; Greenland et al., 1994; Hansen et al., 2001; Raaschou-Nielsen et  
6 al., 2003; Radican et al., 2008; Ritz, 1999a; Siemiatycki, 1991; Zhao et al., 2005). Some  
7 evidence for association with esophageal cancer and overall TCE exposure comes from studies  
8 with high likelihood of TCE exposure (5.6, 95% CI: 0.7, 44.5 (Blair et al., 1998) and 1.88, 95%  
9 CI: 0.61, 5.79 [Radican et al. (2008), which was an update of Blair et al. (1998) with an  
10 additional 10 years of follow-up]; 4.2, 95% CI: 1.5, 9.2, (Hansen et al., 2001); 1.2, 95% CI: 0.84,  
11 1.57 (Raaschou-Nielsen et al., 2003)). Two studies support an association with adenocarcinoma  
12 histologic type of esophageal cancer and TCE exposure (five of the six observed esophageal  
13 cancers were adenocarcinomas [less than 1 expected; Hansen et al. (2001)]; 1.8, 95% CI: 1.2, 2.7  
14 (Raaschou-Nielsen et al., 2003)). Risk estimates in other studies are based on few deaths, low  
15 statistical power to detect a doubling of esophageal cancer risk, and confidence intervals which  
16 include a risk estimate of 1.0 (no increased risk).

17 Seven other studies (ATSDR, 2004a, 2006a; Blair et al., 1989; Clapp and Hoffman, 2008;  
18 Costa et al., 1989; Garabrant et al., 1988; Sung et al., 2007) with lower likelihood for TCE  
19 exposure, in addition to limited statistical power and other design limitations, observed relative  
20 risk estimates between 0.21 (95% CI: 0.001, 1.17) (Costa et al., 1989) to 1.14 (95% CI: 0.62,  
21 1.92) (Garabrant et al., 1988). For these reasons, esophageal cancer observations in these studies  
22 are not inconsistent with Blair et al. (1998) and its update Radican et al. (2008), Hansen et al.  
23 (2001), or Raaschou-Nielsen et al. (2003). No study reported a statistically significant deficit in  
24 the esophageal cancer risk estimate and overall of TCE exposure. Of those studies with  
25 exposure-response analyses, a pattern of increasing esophageal cancer relative risk with  
26 increasing exposure metric is not generally noted (Blair et al., 1998; Boice et al., 1999; Radican  
27 et al., 2008; Siemiatycki, 1991; Zhao et al., 2005) except for Hansen et al. (2001) and  
28 Raaschou-Nielsen et al. (2003). In these last two studies, esophageal cancer relative risk  
29 estimates associated with long employment duration were slightly higher (SIR: 6.6, 95% CI: 1.8,  
30 7.0, 3.7 (Hansen et al., 2001); SIR: 1.9, 95% CO: 0.8, 3.7 (Raaschou-Nielsen et al., 2003)) than  
31 those for short employment duration (SIR: 4.4, 95% CI: 0.5, 19 (Hansen et al., 2001); SIR: 1.7,  
32 95% CI: 0.6, 3.6 (Raaschou-Nielsen et al., 2003)). Hansen et al. (2001) also reports risk for two  
33 other TCE exposure surrogates, average intensity and cumulative exposure, and in both cases  
34 observed lower risk estimates with the higher exposure surrogate.

35

1  
2  
3

**Table 4-110. Summary of human studies on TCE exposure and esophageal cancer**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference	
<b>Cohort studies—incidence</b>					
Aerospace workers (Rocketdyne)				Zhao et al. (2005)	
Any exposure to TCE	Not reported				
Low cumulative TCE score	1.00 <sup>a</sup>		9		
Med cumulative TCE score	1.66 (0.62, 4.41) <sup>b</sup>		8		
High TCE score	0.82 (0.17, 3.95) <sup>b</sup>		2		
<i>p</i> for trend	<i>p</i> = 0.974				
All employees at electronics factory (Taiwan)				Sung et al. (2007)	
Males	Not reported				
Females	1.16 (0.014, 4.20) <sup>c</sup>		2		
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)	
Any exposure, all subjects		1.2 (0.84, 1.57)			44
Any exposure, males		1.1 (0.81, 1.53)			40
Any exposure, females		2.0 (0.54, 5.16)			4
Any exposure, males		1.8 (1.15, 2.73) <sup>d</sup>			23
Any exposure, females					0 (0.4 exp) <sup>d</sup>
Exposure lag time					
20 yr		1.7 (0.8, 3.0) <sup>d</sup>			10
Employment duration					
<1 yr		1.7 (0.6, 3.6) <sup>d</sup>			6
1–4.9 yr		1.9 (0.9, 3.6) <sup>d</sup>			9
≥5 yr		1.9 (0.8, 3.7) <sup>d</sup>			8
Subcohort with higher exposure					
Any TCE exposure		1.7 (0.9, 2.9) <sup>d</sup>			13
Employment duration					
1–4.9 yr		1.6 (0.6, 3.4) <sup>d</sup>		6	
≥5 yr		1.9 (0.8, 3.8) <sup>d</sup>		7	

4

**Table 4-110. Summary of human studies on TCE exposure and esophageal cancer (continued)**

Exposure group	Relative risk (95% CI)	No. obs. events	Reference
Biologically-monitored Danish workers	4.0 (1.5, 8.72)	6	Hansen et al. (2001)
Any TCE exposure, males	4.2 (1.5, 9.2)	6	
Adenocarcinoma histologic type	3.6 (1.2, 8.3) <sup>e</sup>	5	
Any TCE exposure, females		0 (0.1 exp)	
Cumulative exposure (Ikeda)			
<17 ppm-yr	6.5 (1.3, 19)	3	
≥17 ppm-yr	4.2 (1.5, 9.2)	3	
Mean concentration (Ikeda)			
<4 ppm	8.0 (2.6, 19)	5	
4+ ppm	1.3 (0.02, 7.0)	1	
Employment duration			
<6.25 yr	4.4 (0.5, 16)	2	
≥6.25 yr	6.6 (1.8, 17)	4	
Aircraft maintenance workers from Hill Air Force Base			Blair et al. (1998)
TCE subcohort	Not reported		
Males, cumulative exposure			
0	1.0 <sup>a</sup>		
<5 ppm-yr	Not reported		
5–25 ppm-yr	Not reported		
>25 ppm-yr	Not reported		
Females, cumulative exposure			
0	1.0 <sup>a</sup>		
<5 ppm-yr	Not reported		
5–25 ppm-yr	Not reported		
>25 ppm-yr	Not reported		
Biologically-monitored Finnish workers			Anttila et al. (1995)
All subjects	Not reported		
Mean air-TCE (Ikeda extrapolation)			
<6 ppm	Not reported		
6+ ppm	Not reported		

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-110. Summary of human studies on TCE exposure and esophageal cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
	Exposed workers	Not reported		
Biologically-monitored Swedish workers				Axelson et al. (1994)
	Any TCE exposure, males	Not reported		
	Any TCE exposure, females	Not reported		
Cardboard manufacturing workers, Atlanta area, GA				Sinks et al. (1992)
	All subjects	Not reported		
<b>Cohort and PMR studies-mortality</b>				
Computer manufacturing workers (IBM), NY				Clapp and Hoffman (2008)
	Males	1.12 (0.30, 2.86) <sup>f</sup>		
		5.24 (0.13, 29.2) <sup>f</sup>		
Aerospace workers (Rocketdyne)				
	Any TCE (utility/eng flush)	0.88 (0.18, 2.58)	3	Boice et al. (2006b)
	Any exposure to TCE	Not reported		Zhao et al. (2005)
	Low cumulative TCE score	1.00 <sup>a</sup>	18	
	Medium cumulative TCE score	1.40 (0.70, 2.82) <sup>b</sup>	15	
	High TCE score	1.27 (0.52, 3.13) <sup>b</sup>	7	
	<i>p</i> for trend	<i>p</i> = 0.535		
View-Master employees				ATSDR (2004a)
	Males	0.62 (0.02, 3.45) <sup>f</sup>	1	
	Females		0 (1.45 exp) <sup>f</sup>	
All employees at electronics factory (Taiwan)				Chang et al. (2003b)
	Males		0 (3.34 exp)	
	Females		0 (0.83 exp)	
United States uranium-processing workers (Fernald)				Ritz (1999a)
	Any TCE exposure	Not reported		
	Light TCE exposure, >2 yr duration	2.61 (0.99, 6.88) <sup>g</sup>	12	
	Moderate TCE exposure, >2 yr duration		0	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-110. Summary of human studies on TCE exposure and esophageal cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Aerospace workers (Lockheed)				Boice et al. (1999)
Routine exposure	0.83 (0.34, 1.72)	7		
Routine-intermittent <sup>a</sup>	Not presented	11		
Duration of exposure				
0 yr	1.0 <sup>a</sup>	28		
<1 yr	0.23 (0.05, 0.99)	2		
1–4 yr	0.57 (0.20, 1.67)	4		
≥5 yr	0.91 (0.38, 2.22)	7		
<i>p</i> for trend	<i>p</i> > 0.20			
Aerospace workers (Hughes)				Morgan et al. (1998)
TCE subcohort	Not reported			
Low intensity (<50 ppm)				
High intensity (>50 ppm)				
TCE subcohort (Cox Analysis)	Not reported			
Never exposed				
Ever exposed				
Peak	Not reported			
No/Low				
Medium/high				
Cumulative	Not reported			
Referent				
Low				
High				
Aircraft maintenance workers (Hill AFB, UT)				Blair et al. (1998)
TCE subcohort	5.6 (0.7, 44.5) <sup>a</sup>	10		
Males, cumulative exposure				
0	1.0 <sup>a</sup>			
<5 ppm-yr	Not reported <sup>h</sup>	3		
5–25 ppm-yr	Not reported <sup>h</sup>	2		
>25 ppm-yr	Not reported <sup>h</sup>	4		

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-110. Summary of human studies on TCE exposure and esophageal cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Females, cumulative exposure				Blair et al. (1998) (continued)
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	3.6 (0.2, 58)	1	
	5–25 ppm-yr		0	
	>25 ppm-yr		0	
TCE subcohort		1.88 (0.61, 5.79)	17	Radican et al. (2008)
Males, cumulative exposure		1.66 (0.48, 5.74)	15	
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	1.84 (0.48, 7.14)	7	
	5–25 ppm-yr	1.33 (0.27, 6.59)	3	
	>25 ppm-yr	1.67 (0.40, 7.00)	5	
Females, cumulative exposure		2.81 (0.25, 31.10)	2	
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	3.99 (0.25, 63.94)	1	
	5–25 ppm-yr	9.59 (0.60, 154.14)	1	
	>25 ppm-yr		0	
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
	TCE exposed workers	Not reported		
	Unexposed workers	Not reported		
Deaths reported to among GE pension fund (Pittsfield, MA)		0.95 (0.1, 3.17) <sup>i</sup>	13	Greenland et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA		Not reported		Sinks et al. (1992)
U. S. Coast Guard employees				Blair et al. (1989)
	Marine inspectors	0.72 (0.09, 2.62)	2	
	Noninspectors	0.74 (0.09, 2.68)	2	
Aircraft manufacturing plant employees (Italy)				Costa et al. (1989)
	All subjects	0.21 (0.01, 1.17)	1	
Rubber Workers		Not reported <sup>i</sup>		Wilcosky et al. (1984)
Aircraft manufacturing plant employees (San Diego, CA)				Garabrant et al. (1988)
	All subjects	1.14 (0.62, 1.92)	14	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-110. Summary of human studies on TCE exposure and esophageal cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Case-control studies</b>				
Population of Montreal, Canada				Siemiatycki et al. (1991); Parent et al. (2000b)
	Any TCE exposure	0.5 (0.1, 2.5) <sup>j</sup>	1	
	Substantial TCE exposure	0.8 (0.1, 4.6) <sup>j</sup>	1	
<b>Geographic based studies</b>				
Residents in two study areas in Endicott, NY		0.78 (0.29, 1.70)	6	ATSDR (2006a)
Residents of 13 census tracts in Redlands, CA		Not reported		Morgan and Cassady (2002)
Finnish residents				Vartiainen et al. (1993)
	Residents of Hausjarvi	Not reported		
	Residents of Huttula	Not reported		

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

<sup>a</sup>Internal referents, workers not exposed to TCE.

<sup>b</sup>Ritz (1999a) and Zhao et al. (2005) reported relative risks for the combined site of esophagus and stomach.

<sup>c</sup>Sung et al. (2007) and Chang et al. (2005)—SIR for females and reflects a 10-yr lag period.

<sup>d</sup>SIR for adenocarcinoma of the esophagus.

<sup>e</sup>The SIR for adenocarcinoma histologic type cannot be calculated because Hansen et al. (2001) do not present expected numbers for adenocarcinoma histologic type of esophageal cancer. An approximation of the SIR for adenocarcinoma histologic type is presented using the expected number of total number of expected esophageal cancers for males ( $n = 1.4$ ). The expected numbers of esophageal adenocarcinomas in males will be lower; Hansen et al. (2001) noted the proportion of adenocarcinomas among the comparable Danish male population during the later period of the study (1990–1996) as 38%. A rough approximation of the expected number of esophageal carcinomas would be 0.5 expected cases and an approximated SIR of 9.4 (3.1, 22).

<sup>f</sup>Proportional mortality ratio.

<sup>g</sup>Adjusted relative risks for >2 yr exposure duration and 15 yr lag from 1<sup>st</sup> exposure.

<sup>h</sup>No esophageal cancer deaths occurred in the referent population in Blair et al. (1998) and relative risk in could not be calculated for this reason.

<sup>i</sup>Odds ratio from nested case-control analysis.

<sup>j</sup>90% CI.

GE = General Electric, IBM = International Business Machines Corporation, No. obs. events = number of observable events.

Meta-analysis is not adopted as a tool for examining the body of epidemiologic evidence on esophageal cancer and TCE exposure given the absence of reported relative risk estimates in several of the studies in which there is a high likelihood of TCE exposure in individual study subjects and which met, to a sufficient degree, the standards of epidemiologic design and analysis in a systematic review (Anttila et al., 1995; Axelson et al., 1994; Morgan et al., 1998).

1 Overall, three cohort studies in which there is a high likelihood of TCE exposure in  
2 individual study subjects and which met, to a sufficient degree, the standards of epidemiologic  
3 design and analysis in a systematic review provide some evidence of association for esophageal  
4 cancer and TCE exposure. The finding in two of these studies of esophageal risk estimates  
5 among subjects with long employment duration were higher than those associated with low  
6 employment duration provides additional evidence (Hansen et al., 2001; Raaschou-Nielsen et al.,  
7 2003). The cohort studies are unable to directly examine possible confounding due to suspected  
8 risk factors for esophageal cancer such as smoking, obesity and alcohol. The use of an internal  
9 referent group, similar in socioeconomic status as exposed subjects, is believed to minimize but  
10 may not completely control for possible confounding related to smoking and health status (Blair  
11 et al. (1998); its follow-up Radican et al. (2008); Zhao et al. (2005); Boice et al. (2006b).  
12 Observation of a higher risk for adenocarcinoma histologic type than for a combined category of  
13 esophageal cancer in Raaschou-Nielsen et al. (2003) also suggests minimal confounding from  
14 smoking. Smoking is not identified as a possible risk factor for the adenocarcinoma histologic  
15 type of esophageal cancer but is believed a risk factor for squamous cell histologic type.  
16 Furthermore, the magnitude of lung cancer risk in Raaschou-Nielsen et al. (2003) suggests a high  
17 smoking rate is unlikely. The lack of association with overall TCE exposure and the absence of  
18 exposure-response patterns in the other studies of TCE exposure may reflect limitations in  
19 statistical power, the possibility of exposure misclassification, and differences in measurement  
20 methods. These studies do not provide evidence against an association between TCE exposure  
21 and esophageal cancer.  
22

#### 4.9.2. Bladder Cancer

23 Twenty-five epidemiologic studies present risk estimates for bladder cancer Pesch et al.,  
24 2000a (Anttila et al., 1995; ATSDR, 2004a, 2006a; Axelson et al., 1994; Blair et al., 1989; Blair  
25 et al., 1998; Boice et al., 1999; Boice et al., 2006b; Chang et al., 2003b; Chang et al., 2005;  
26 Costa et al., 1989; Garabrant et al., 1988; Greenland et al., 1994; Hansen et al., 2001; Mallin,  
27 1990; Morgan and Cassady, 2002; Morgan et al., 1998; Raaschou-Nielsen et al., 2003; Radican  
28 et al., 2008; Shannon et al., 1988; Siemiatycki, 1991; Sinks et al., 1992; Sung et al., 2007; Zhao  
29 et al., 2005). Table 4-111 presents risk estimates for TCE exposure and bladder cancer observed  
30 in cohort, case-control, and geographic based studies. Thirteen studies, all either cohort or case-  
31 control studies, which there is a high likelihood of TCE exposure in individual study subjects  
32 (e.g., based on job-exposure matrices or biomarker monitoring) or which met, to a sufficient  
33 degree, the standards of epidemiologic design and analysis in a systematic review, reported

1 relative risk estimates for bladder or urothelial cancer between 0.6 (Siemiatycki, 1991) and 1.7  
2 (Boice et al., 2006b) and overall TCE exposure. Relative risk estimates were generally based on  
3 small numbers of cases or deaths, except for one study (Raaschou-Nielsen et al., 2003), with the  
4 result of wide confidence intervals on the estimates. Of these studies, two reported statistically  
5 significant elevated bladder or urothelial cancer risks with the highest cumulative TCE exposure  
6 category (2.71, 95% CI: 1.10, 6.65 (Morgan et al., 1998); 1.8, 95% CI: 1.2, 2.7 [Pesch et al.,  
7 2000b]) and five presented risk estimates and categories of increasing cumulative TCE exposure  
8 Pesch et al., 2000b (Blair et al., 1998; Morgan et al., 1998; Radican et al., 2008; Zhao et al.,  
9 2005). Risk estimates in  
10

1  
2  
3

**Table 4-111. Summary of human studies on TCE exposure and bladder cancer**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort studies—incidence</b>				
Aerospace workers (Rocketdyne)				Zhao et al. (2005)
	Any exposure to TCE	Not reported		
	Low cumulative TCE score	1.00 <sup>a</sup>	20	
	Medium cumulative TCE score	1.54 (0.81, 2.92) <sup>b</sup>	19	
	High TCE score	1.98 (0.93, 4.22) <sup>b</sup>	11	
	<i>p</i> for trend	<i>p</i> = 0.069		
TCE, 20 yr exposure lag				
	Low cumulative TCE score	1.00 <sup>a</sup>	20	
	Medium cumulative TCE score	1.76 (0.61, 5.10) <sup>c</sup>	20	
	High TCE score	3.68 (0.87, 15.5) <sup>c</sup>	10	
	<i>p</i> for trend	<i>p</i> = 0.064		
All employees at electronics factory (Taiwan)				
	Males	Not reported		Sung et al. (2007)
	Females	0.34 (0.07, 1.00)	10	
	Males	1.06 (0.45, 2.08) <sup>d</sup>	8	Chang et al. (2005)
	Females	1.09 (0.56, 1.91) <sup>d</sup>	12	
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure, all subjects	1.1 (0.92, 1.21)	220	
	Any exposure, males	1.0 (0.89, 1.18)	203	
	Any exposure, females	1.6 (0.93, 2.57)	17	
Biologically-monitored Danish workers		1.0 (0.48, 1.86)	10	Hansen et al. (2001)
	Any TCE exposure, males	1.1 (0.50, 2.0)	10	
	Any TCE exposure, females	0.5 expected	0	
Aircraft maintenance workers from Hill Air Force Base				Blair et al. (1998)
	TCE subcohort	Not reported		
Males, cumulative exposure				
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	1.7 (0.6, 4.4)	13	
	5–25 ppm-yr	1.7 (0.6, 4.9)	9	
	>25 ppm-yr	1.4 (0.5, 4.1)	9	

4

**Table 4-111. Summary of human studies on TCE exposure and bladder cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Females, cumulative exposure				Blair et al. (1998) (continued)
0		1.0 <sup>a</sup>		
<5 ppm-yr		1.1 (0.1, 10.8)	1	
5–25 ppm-yr			0	
>25 ppm-yr		1.0 (0.1, 9.1)	1	
Biologically-monitored Finnish workers				Anttila et al. (1995)
All subjects		0.82 (0.27, 1.90)	5	
Biologically-monitored Swedish workers				Axelson et al. (1994)
Any TCE exposure, males		1.02 (0.44, 2.00)	8	
Any TCE exposure, females		Not reported		
<b>Cohort and PMR studies-mortality</b>				
Aerospace workers (Rocketdyne)				
Any TCE (utility/eng flush)		1.66 (0.54, 3.87)	5	Boice et al. (2006b)
Any exposure to TCE		Not reported		Zhao et al. (2005)
Low cumulative TCE score		1.00 <sup>a</sup>	8	
Med cumulative TCE score		1.27 (0.43, 3.73) <sup>b</sup>	6	
High TCE score		1.15 (0.29, 4.51) <sup>b</sup>	3	
<i>p</i> for trend		<i>p</i> = 0.809		
TCE, 20 yr exposure lag				
Low cumulative TCE score		1.00 <sup>a</sup>	8	
Medium cumulative TCE score		0.95 (0.15, 6.02) <sup>c</sup>	7	
High TCE score		1.85 (0.12, 27.7) <sup>c</sup>	2	
<i>p</i> for trend		<i>p</i> = 0.533		
View-Master employees				ATSDR (2004a)
Males		1.22 (0.15, 4.40)		
Females		0.78 (0.09, 2.82)		
United States uranium-processing workers (Fernald)				Ritz (1999a)
Any TCE exposure		Not reported		
Light TCE exposure, >2 yr duration		Not reported		
Moderate TCE exposure, >2 yr duration		Not reported		
Aerospace workers (Lockheed)				Boice et al. (1999)
Routine exposure		0.55 (0.18, 1.28)	5	
Routine-intermittent <sup>a</sup>		Not reported		

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-111. Summary of human studies on TCE exposure and bladder cancer (continued)**

Exposure group	Relative risk (95% CI)	No. obs. events	Reference
Aerospace workers (Hughes)			Morgan et al. (1998)
TCE subcohort	1.36 (0.59, 2.68)	8	
Low intensity (<50 ppm)	0.51 (0.01, 2.83)	1	
High intensity (>50 ppm)	1.79 (0.72, 3.69)	7	
TCE subcohort (Cox Analysis)			
Never exposed	1.0 <sup>a</sup>		
Ever exposed	2.05 (0.86, 4.85) <sup>c</sup>	8	
Peak			
No/low	1.0 <sup>a</sup>		
Medium/high	1.41 (0.52, 3.81)	5	
Cumulative			
Referent	1.0 <sup>a</sup>		
Low	0.69 (0.09, 5.36)	1	
High	2.71 (1.10, 6.65)	7	
Aircraft maintenance workers (Hill AFB, UT)			Blair et al. (1998)
TCE subcohort	1.2 (0.5, 2.9) <sup>a</sup>	17	
Males, cumulative exposure			
0	1.0 <sup>a</sup>		
<5 ppm-yr	1.8 (0.5, 6.2)	7	
5–25 ppm-yr	2.1 (0.6, 8.0)	5	
>25 ppm-yr	1.0 (0.2, 5.1)	3	
Females, cumulative exposure			
0	1.0 <sup>a</sup>		
<5 ppm-yr		0	
5–25 ppm-yr		0	
>25 ppm-yr	0.8 (0.1, 7.5)	1	
TCE subcohort			
Males, cumulative exposure			
0	1.0 <sup>a</sup>		
<5 ppm-yr	0.96 (0.37, 2.51)	9	
5–25 ppm-yr	1.77 (0.70, 4.52)	10	
>25 ppm-yr	0.67 (0.15, 2.95)	5	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-111. Summary of human studies on TCE exposure and bladder cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
	Females, cumulative exposure	0.22 (0.03, 1.83)	1	Radican et al. (2008) (continued)
	0	1.0 <sup>a</sup>		
	<5 ppm-yr		0	
	5–25 ppm-yr	2.86 (0.27, 29.85)	1	
	>25 ppm-yr		0	
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
	TCE exposed workers	Not reported		
	Unexposed workers	Not reported		
Deaths reported to GE pension fund (Pittsfield, MA)		0.85 (0.32, 2.23) <sup>f</sup>	20	Greenland et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA				Sinks et al. (1992)
		0.3 (0.0, 1.6)	1	
U. S. Coast Guard employees				Blair et al. (1989)
	Marine inspectors	0.50 (0.06, 1.79)	2	
	Noninspectors	0.90 (0.18, 2.62)	3	
Aircraft manufacturing plant employees (Italy)				Costa et al. (1989)
	All subjects	0.74 (0.30, 1.53)	7	
Aircraft manufacturing plant employees (San Diego, CA)				Garabrant et al. (1988)
	All subjects	1.26 (0.74, 2.03)	17	
Lamp manufacturing workers (GE)		0.93 (0.19, 2.72)	3	Shannon et al. (1988)
<b>Case-control studies</b>				
Population of 5 regions in Germany				Pesch et al., 2000a
	Any TCE exposure	Not reported		
	Males	Not reported		
	Females	Not reported		
	Males			
	Medium	0.8 (0.6, 1.2) <sup>g</sup>	47	
	High	1.3 (0.8, 1.7) <sup>g</sup>	74	
	Substantial	1.8 (1.2, 2.7) <sup>g</sup>	36	
Population of Montreal, Canada				Siemiatycki (1991); Siemiatycki et al. (1994)
	Any TCE exposure	0.6 (0.3, 1.2)	8	
	Substantial TCE exposure	0.7 (0.3, 1.6)	5	
<b>Geographic based studies</b>				
Residents in two study areas in Endicott, NY				ATSDR (2006a)
		0.71 (0.38, 1.21)	13	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-111. Summary of human studies on TCE exposure and bladder cancer (continued)**

Exposure group	Relative risk (95% CI)	No. obs. events	Reference
Residents of 13 census tracts in Redlands, CA			Morgan and Cassady (2002)
	0.98 (0.71, 1.29) <sup>h</sup>	82	
Finnish residents			Vartiainen et al. (1993)
Residents of Hausjarvi	Not reported		
Residents of Huttula	Not reported		
Residents of 9 county area in Northwestern Illinois			Mallin (1990)
All zip codes in study area			
Males	1.4 (1.1, 1.9)	47	
Females	1.8 (1.2, 2.7)	21	
Cluster community			
Males	1.7 (1.1, 2.6)	21	
Females	2.6 (1.2, 4.7)	10	
Adjacent community			
Males	1.2 (0.6, 2.0)	12	
Females	1.6 (0.5, 3.8)	5	
Remainder of zip code areas			
Males	1.4 (0.8, 2.2)	14	
Females	1.4 (0.5, 3.0)	6	

<sup>a</sup>Internal referents, workers not exposed to TCE.

<sup>b</sup>Relative risk estimates for TCE exposure after adjustment for 1<sup>st</sup> employment, socioeconomic status, and age at event.

<sup>c</sup>Relative risk estimates for TCE exposure after adjustment for 1<sup>st</sup> employment, socioeconomic status, age at event, and all other carcinogen exposures, including hydrazine.

<sup>d</sup>Chang et al. (2005) and Costa et al. (1989) report estimated risks for a combined site of all urinary organ cancers.

<sup>e</sup>Risk ratio from Cox Proportional Hazard Analysis, stratified by age, sex and decade (EHS, 1997).

<sup>f</sup>Odds ratio from nested case-control analysis.

<sup>g</sup>Odds ratio for urothelial cancer, a category of bladder, ureter, and renal pelvis cancers) and cumulative TCE exposure, as assigned using a JTEM approach (Pesch et al., 2000a).

<sup>h</sup>99% CI.

GE = General Electric, No. obs. events = number of observable events.

Morgan et al. (1998), Pesch et al. (2000a), and Zhao et al. (2005) appeared to increase with increasing cumulative TCE exposure with the *p*-value for trend of 0.07 in Zhao et al. (2005), the only study to present a formal statistical test for linear trend. Risk estimates did not appear to either increase or decrease with increasing cumulative TCE exposure in Blair et al. (1998) or its update Radican et al. (2008), which added another 10 years of follow-up. Twelve additional

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 studies were given less weight because of their lesser likelihood of TCE exposure and other  
2 design limitations that would decrease statistical power and study sensitivity (ATSDR, 2004a,  
3 2006a; Blair et al., 1989; Chang et al., 2003b; Chang et al., 2005; Costa et al., 1989; Garabrant et  
4 al., 1988; Mallin, 1990; Morgan and Cassady, 2002; Shannon et al., 1988; Sinks et al., 1992;  
5 Sung et al., 2007).

6 Meta-analysis is not adopted as a tool for examining the body of epidemiologic evidence  
7 on bladder cancer and TCE.

8 Overall, three cohort or case-control studies in which there is a high likelihood of TCE  
9 exposure in individual study subjects and which met, to a sufficient degree, the standards of  
10 epidemiologic design and analysis in a systematic review provide some evidence of association  
11 for bladder or urothelial cancer and high cumulative TCE exposure Pesch et al., 2000a (Morgan  
12 et al., 1998; Zhao et al., 2005). The case-control study of Pesch et al. (2000a) adjusted for age,  
13 study center, and cigarette smoking, with a finding of a statistically significant risk estimate  
14 between urothelial cancer and the highest TCE exposure category. Cancer cases in this study are  
15 of several sites, bladder, ureter, and renal pelvis, and grouping different site-specific cancers with  
16 possible etiologic heterogeneity may introduce misclassification bias. The cohort studies are  
17 unable to directly examine possible confounding due to suspected risk factors for esophageal  
18 cancer such as smoking, obesity, and alcohol. The use of an internal referent group, similar in  
19 socioeconomic status as exposed subjects, by Morgan et al. (1998) and Zhao et al. (2005) is  
20 believed to minimize but may not completely control for possible confounding related to  
21 smoking and health status. The lack of association with overall TCE exposure in other studies  
22 and the absence of exposure-response patterns with TCE exposure in Blair et al. (1998) and  
23 Radican et al. (2008) may reflect limitations in statistical power, the possibility of exposure  
24 misclassification, and differences in measurement methods. These studies do not provide  
25 evidence against an association between TCE exposure and bladder cancer.  
26

#### **4.9.3. Central Nervous System and Brain Cancers**

27 Brain cancer is examined in most cohort studies and in one case-control study (Anttila et  
28 al., 1995; Blair et al., 1989; Blair et al., 1998; Boice et al., 1999; Boice et al., 2006b; Chang et  
29 al., 2003b; Chang et al., 2005; Clapp and Hoffman, 2008; Costa et al., 1989; Garabrant et al.,  
30 1988; Greenland et al., 1994; Hansen et al., 2001; Heineman et al., 1994; Henschler et al., 1995;  
31 Morgan et al., 1998; Raaschou-Nielsen et al., 2003; Radican et al., 2008; Ritz, 1999a; Sung et  
32 al., 2007; Zhao et al., 2005). Overall, these epidemiologic studies do not provide strong  
33 evidence for or against association between TCE and brain cancer in adults (see Table 4-112).

1 Relative risk estimates in well designed and conducted cohort studies, Axelson et al. (1994),  
2 Anttila et al. (1995), Blair et al. (1998), its follow-up reported in Radican et al. (2008), Morgan  
3 et al. (1998), Boice et al. (1999), Zhao et al. (2005), and Boice et al. (2006b), are near a risk of  
4 1.0 and imprecise, confidence intervals all include a risk estimate of 1.0. All studies except  
5 Raaschou-Nielsen et al. (2003),  
6

**Table 4-112. Summary of human studies on TCE exposure and brain cancer**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort studies—incidence</b>				
Aerospace workers (Rocketdyne)				Zhao et al. (2005)
	Any exposure to TCE	Not reported		
	Low cumulative TCE score	1.00 <sup>a</sup>	7	
	Medium cumulative TCE score	0.46 (0.09, 2.25) <sup>b</sup>	2	
	High TCE score	0.47 (0.06, 3.95) <sup>b</sup>	1	
	<i>p</i> for trend	<i>p</i> = 0.382		
All employees at electronics factory (Taiwan)				
	Males	Not reported		Sung et al. (2007)
	Females	1.07 (0.59, 1.80) <sup>c</sup>		
	Males	0.40 (0.05, 1.46)	2	Chang et al. (2005)
	Females	0.97 (0.54, 1.61)	15	
Danish blue-collar worker with TCE exposure				Raaschou-Nielsen et al. (2003)
	Any exposure, all subjects	1.0 (0.84, 1.24)	104	
	Any exposure, males	1.0 (0.76, 1.18)	85	
	Any exposure, females	1.1 (0.67, 1.74)	19	
Biologically-monitored Danish workers		0.3 (0.01, 1.86)	1	Hansen et al. (2001)
	Any TCE exposure, males	0.4 (0.01, 2.1)	1	
	Any TCE exposure, females	0.5 expected	0	
Aircraft maintenance workers from Hill Air Force Base				Blair et al. (1998)
	TCE subcohort	Not reported		
Males, cumulative exposure				
	0	1.0 <sup>a</sup>		
	<5 ppm-yr	2.0 (0.2, 19.7)	3	
	5–25 ppm-yr	3.9 (0.4, 34.9)	4	
	>25 ppm-yr	0.8 (0.1, 13.2)	1	
Females, cumulative exposure				
	0	1.0 <sup>a</sup>		
	<5 ppm-yr		0	
	5–25 ppm-yr		0	
	>25 ppm-yr		0	
Biologically-monitored Finnish workers				
	All subjects	1.09 (0.50, 2.07)	9	
Mean air-TCE (Ikeda extrapolation)				
	<6 ppm	1.52 (0.61, 3.13)	7	
	6+ ppm	0.76 (0.01, 2.74)	2	
Biologically-monitored Swedish workers				Axelson et al. (1994)
	Any TCE exposure, males	Not reported		
	Any TCE exposure, females	Not reported		

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-112. Summary of human studies on TCE exposure and brain cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Cohort and PMR studies-mortality</b>				
Computer manufacturing workers (IBM), NY				Clapp and Hoffman (2008)
	Males	1.90 (0.52, 4.85)	4	
	Females		0	
Aerospace workers (Rocketdyne)				
	Any TCE (utility/eng flush)	0.81 (0.17, 2.36)	3	Boice et al. (2006b)
	Any exposure to TCE	Not reported		Zhao et al. (2005)
	Low cumulative TCE score	1.00 <sup>a</sup>	12	
	Medium cumulative TCE score	0.42 (0.12, 1.50)	3	
	High TCE score	0.83 (0.23, 3.08)	3	
	<i>p</i> for trend	<i>p</i> = 0.613		
View-Master employees				ATSDR (2004a)
	Males	Not reported		
	Females	Not reported		
All employees at electronics factory (Taiwan)				Chang et al. (2003b)
	Males	0.96 (0.01, 5.36)	1	
	Females	0.96 (0.01, 5.33)	1	
United States uranium-processing workers (Fernald)				Ritz (1999a)
	Any TCE exposure	Not reported		
	Light TCE exposure, >2 yr duration, 0 lag	1.81 (0.49, 6.71) <sup>d</sup>	6	
	Moderate TCE exposure, >2 yr duration, 0 lag	3.26 (0.37, 28.9) <sup>d</sup>	1	
	Light TCE exposure, >5 yr duration, 15 yr lag	5.41 (0.87, 33.9) <sup>d</sup>	3	
	Moderate TCE exposure, >5 yr duration, 15 yr lag	14.4 (1.24, 167) <sup>d</sup>	1	
Aerospace workers (Lockheed)				Boice et al. (1999)
	Routine exposure	0.54 (0.15, 1.37)	4	
	Routine-intermittent <sup>a</sup>	Not presented		
Aerospace workers (Hughes)				Morgan et al. (1998)
	TCE subcohort	0.99 (0.64, 1.47)	4	
	Low intensity (<50 ppm) <sup>d</sup>	0.73 (0.09, 2.64)	2	
	High intensity (>50 ppm) <sup>d</sup>	0.44 (0.05, 1.58)	2	
Aircraft maintenance workers (Hill AFB, UT)				Blair et al. (1998)
	TCE subcohort	0.8 (0.2, 2.2) <sup>a</sup>	11	

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 4-112. Summary of human studies on TCE exposure and brain cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
Males, cumulative exposure				Blair et al. (1998) (continued)
0		1.0 <sup>a</sup>		
<5 ppm-yr		0.7 (0.7, 3.3)	3	
5–25 ppm-yr		2.0 (0.5, 8.4)	5	
>25 ppm-yr		0.9 (0.2, 4.4)	2	
Females, cumulative exposure				
0		1.0 <sup>a</sup>		
<5 ppm-yr			0	
5–25 ppm-yr			0	
>25 ppm-yr			0	
TCE subcohort		1.02 (0.39, 2.67)	17	Radican et al. (2008)
Males, cumulative exposure		1.26 (0.43, 3.75)	17	
0		1.0 <sup>a</sup>		
<5 ppm-yr		1.46 (0.44, 4.86)	8	
5–25 ppm-yr		1.74 (0.49, 6.16)	6	
>25 ppm-yr		0.66 (0.15, 2.95)	3	
Females, cumulative exposure			0	
0				
<5 ppm-yr				
5–25 ppm-yr				
>25 ppm-yr				
Cardboard manufacturing workers in Arnsburg, Germany				Henschler et al. (1995)
TCE exposed workers		3.70 (0.09, 20.64)	1	
Unexposed workers		9.38 (1.93, 27.27)	3	
Deaths reported to GE pension fund (Pittsfield, MA)		0.93 (0.32, 2.69) <sup>c</sup>	16	Greenland et al. (1994)
Cardboard manufacturing workers, Atlanta area, GA				Sinks et al. (1992)
		Not reported		
U. S. Coast Guard employees				Blair et al. (1989)
Marine inspectors		1.70 (0.55, 3.95)	5	
Noninspectors		1.36 (0.44, 3.17)	5	
Aircraft manufacturing plant employees (Italy)				Costa et al. (1989)
All subjects		0.79 (0.16, 2.31)	3	
Aircraft manufacturing plant employees (San Diego, CA)				Garabrant et al. (1988)
All subjects		0.78 (0.42, 1.34)	16	
<b>Case-control studies</b>				
Children's Cancer Group/Pediatric Oncology Group				De Roos et al. (2001)
Any TCE exposure		1.64 (0.95, 2.84)	37	

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 4-112. Summary of human studies on TCE exposure and brain cancer (continued)**

Exposure group		Relative risk (95% CI)	No. obs. events	Reference
<b>Neuroblastoma, ≤15 yr age</b>				
Paternal TCE exposure				De Roos et al. (2001) (continued)
	Self-reported exposure	1.4 (0.7, 2.9)	22	
	IH assignment of probable exposure	0.9 (0.3, 2.5)	9	
Population of So. LA, NJ, Philadelphia, PA				Heineman et al. (1994)
	Any TCE exposure	1.1 (0.8, 1.6)	128	
	Low exposure	1.1 (0.7, 1.7)	27	
	Medium exposure	1.1 (0.6, 1.8)	42	
	High exposure	1.1 (0.5, 2.8)	12	
	<i>p</i> for trend	0.45		
<b>Geographic based studies</b>				
Residents in two study areas in Endicott, NY				ATSDR (2006a)
	<b>Brain/CNS, ≤19 yr of age</b>	Not reported	<6	
Residents of 13 census tracts in Redlands, CA				Morgan and Cassady (2002)
	<b>Brain/CNS, &lt;15 yr of age</b>	1.05 (0.24, 2.70) <sup>f</sup>	6	
Resident of Tucson Airport Area, AZ				AZ DHS (1990, 1995)
	<b>Brain/CNS, ≤19 yr of age</b>			
	1970–1986	0.84 (0.23, 2.16)	3	
	1987–1991	0.78 (0.26, 2.39)	2	

<sup>a</sup>Internal referents, workers not exposed to TCE.

<sup>b</sup>Relative risks for TCE exposure after adjustment for 1<sup>st</sup> employment, socioeconomic status, and age at event.

<sup>c</sup>Standardized incidence ratio from analyses lagging exposure 10 yrs prior to end of follow-up or date of incident cancer.

<sup>d</sup>Relative risks for TCE exposure after adjustment for time since 1<sup>st</sup> hired, external and internal radiation dose, and same chemical at a different level.

<sup>e</sup>Odds ratio from nested case-control analysis.

<sup>f</sup>99% CI.

IBM = International Business Machines Corporation, No. obs. events = number of observable events.

observations are based on few events and lowered statistical power. Bias resulting from exposure misclassification is likely in these studies, although of a lower magnitude compared to other cohort studies identified in Table 4-112, and may partly explain observations. Exposure misclassification is also likely in the case-control study of occupational exposure of Heineman et al. (1994) who do not report association with TCE exposure.

Three geographic-based studies and one case-control study examined childhood brain cancer (ADHS, 1990, 1995; ATSDR, 2006a; De Roos et al., 2001; Morgan and Cassady, 2002).

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 The strongest study, De Roos et al. (2001), a population case-control study which examined  
2 paternal exposure, used expert judgment to evaluate the probably of TCE exposure from self-  
3 reported information in an attempt to reduce exposure misclassification bias. The odds ratio  
4 estimate in this study was 0.9 (95% CI: 0.3, 2.5). Like many population case-control studies, a  
5 low prevalence of TCE exposure was found, only 9 fathers were identified with probable TCE  
6 exposure by the industrial hygiene review, and greatly impacted statistical power. There is some  
7 concern for childhood brain cancer and organic solvent exposure based on Peters et al. (1981)  
8 whose case-control study of childhood brain cancer reported to the Los Angeles County Cancer  
9 Surveillance Program observed a high odds ratio estimate for paternal employment in the aircraft  
10 industry (OR:  $\infty$ ,  $p < 0.001$ ). This study does not present an odds ratio for TCE exposure only  
11 although it did identify two of the 14 case and control fathers with previous employment in the  
12 aircraft industry reported exposure to TCE.  
13

#### 4.10. SUSCEPTIBLE LIFESTAGES AND POPULATIONS

14 Variation in response among segments of the population may be due to age, genetics, and  
15 ethnicity, as well as to differences in lifestyle, nutrition, and disease status. These could be  
16 potential risk factors that play an important role in determining an individual's susceptibility and  
17 sensitivity to chemical exposures. Available studies on TCE toxicity in relation to some of these  
18 risk factors including lifestage, gender, genetics, race/ethnicity, preexisting health status, and  
19 lifestyle are discussed below. However, there is a general lack of data demonstrating the  
20 modulation of health effects from TCE exposure based on these factors. Additional data  
21 examining these factors would provide further understanding of the populations that may be  
22 more susceptible to the health effects from TCE exposure. Others have also reviewed factors  
23 related to human variability and their potential for susceptibility to TCE (ATSDR, 1997b, 1998a;  
24 Barton et al., 1996; Clewell et al., 2000; Davidson and Beliles, 1991; NRC, 2006; Pastino et al.,  
25 2000).  
26

##### 4.10.1. Lifestages

27 Individuals of different lifestages are physiologically, anatomically, and biochemically  
28 different. Early (infants and children) and later (the elderly) lifestages differ greatly from  
29 adulthood in body composition, organ function, and many other physiological parameters that  
30 can influence the toxicokinetics of chemicals and their metabolites in the body (ILSI, 1992). The  
31 limited data on TCE exposure among these segments of the population—particularly individuals

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 in early lifestages—suggest they may have greater susceptibility than does the general  
2 population. This section presents and evaluates the pertinent published literature available to  
3 assess how individuals of differing lifestages may respond differently to TCE.

#### 4.10.1.1.1. Early Lifestages

#### 4.10.1.1.2. Early lifestage-specific exposures

4 Section 2.4 describes the various exposure pathways of concern for TCE. For all  
5 postnatal lifestages, the primary exposure routes of concern include inhalation and contaminated  
6 drinking water. In addition, there are exposure pathways to TCE are unique to early lifestages.  
7 Fetal and infant exposure to TCE can occur through placental transfer and breast milk  
8 consumption if the mother has been exposed, and could potentially increase overall TCE  
9 exposure. Placental transfer of TCE has been demonstrated in humans (Beppu, 1968; Laham,  
10 1970), rats (Withey and Karpinski, 1985), mice (Ghantous et al., 1986), rabbits (Beppu, 1968),  
11 and sheep and goats (Helliwell and Hutton, 1950). Similarly, TCE has been found in breast milk  
12 in humans (Fisher et al., 1997; Pellizzari et al., 1982), goats (Hamada and Tanaka, 1995), and  
13 rats (Fisher et al., 1990). Pellizzari et al. (1982) conducted a survey of environmental  
14 contaminants in human milk, using samples from cities in the northeastern region of the United  
15 States and one in the southern region and detected TCE in 8 milk samples taken from 42  
16 lactating women. No details of when the samples were taken postpartum, milk lipid content, or  
17 TCE concentration in milk or blood were reported. Fisher et al. (1997) predicted that a nursing  
18 infant would consume 0.496 mg TCE during a 24-hour period. In lactating rats exposed to 600  
19 ppm (3,225 mg/m<sup>3</sup>) TCE for 4 hours resulted in concentrations of TCE in milk of 110 µg/mL  
20 immediately following the cessation of exposure (Fisher et al., 1990).

21 Direct childhood exposures to TCE from oral exposures may also occur. A  
22 contamination of infant formula resulted in levels of 13 ppb (Fan, 1988). Children consume high  
23 levels of dairy products, and TCE has been found in butter and cheese (Wu and Schaum, 2000).  
24 In addition, TCE has been found in food and beverages containing fats such as margarine  
25 (Wallace et al., 1984), grains and peanut butter (Wu and Schaum, 2000), all of which children  
26 consume in high amounts. A number of studies have examined the potential adverse effects of  
27 prenatal or postnatal exposure to drinking water contaminated with TCE (ATSDR, 1998a, 2001;  
28 Bernad et al., 1987, abstract; Bove, 1996; Bove et al., 1995; Burg and Gist, 1999; Goldberg et  
29 al., 1990; Lagakos et al., 1986; Rodenbeck et al., 2000; Sonnenfeld et al., 2001; White et al.,  
30 1997; see Section 4.10.2.1). TCE in residential water may also be a source of dermal or  
31 inhalation exposure during bathing and showering (Fan, 1988; Franco et al., 2007; Giardino and

1 Andelman, 1996; Lee et al., 2002; Weisel and Jo, 1996; Wu and Schaum, 2000); it has been  
2 estimated that showering and bathing scenarios in water containing 3-ppm TCE, a child of 22 kg  
3 receives a higher dose (about 1.5 times) on a mg/kg basis than a 70 kg adult (Fan, 1988).

4 Direct childhood inhalation exposure to TCE have been documented in both urban and  
5 rural settings. A study of VOCs measured personal, indoor and outdoor TCE in 284 homes, with  
6 72 children providing personal measures and time-activity diaries (Adgate et al., 2004a). The  
7 intensive-phase of the study found a mean personal level of  $0.8 \mu\text{g}/\text{m}^3$  and mean indoor and  
8 outdoor levels of  $0.6 \mu\text{g}/\text{m}^3$ , with urban homes have significantly higher indoor levels of TCE  
9 than nonurban homes ( $t = 2.3, p = 0.024$ ) (Adgate et al., 2004a). A similar study of personal,  
10 indoor and outdoor TCE was conducted in two inner-city elementary schools as well as in the  
11 homes of 113 children along with time-activity diaries, and found a median a median personal  
12 level of  $0.3 \mu\text{g}/\text{m}^3$ , a median school indoor level of  $0.2 \mu\text{g}/\text{m}^3$ , a median home indoor level of  
13  $0.3 \mu\text{g}/\text{m}^3$ , a median outdoor level of  $0.3 \mu\text{g}/\text{m}^3$  in the winter, with slightly lower levels in the  
14 spring (Adgate et al., 2004b). Studies from Leipzig, Germany measured the median air level of  
15 TCE in children's bedrooms to be  $0.42 \mu\text{g}/\text{m}^3$  (Lehmann et al., 2001) and  $0.6 \mu\text{g}/\text{m}^3$  (Lehmann et  
16 al., 2002). A study of VOCs in Hong Kong measured air levels in schools, including an 8-hour  
17 average of  $1.28 \mu\text{g}/\text{m}^3$ , which was associated with the lowest risk of cancer in the study (Guo et  
18 al., 2004). Another found air TCE levels to be highest in school/work settings, followed by  
19 outside, in home, in other, and in transit settings (Sexton et al., 2007). Measured indoor air  
20 levels ranged from  $0.18\text{--}140 \mu\text{g}/\text{m}^3$  for children exposed through vapor intrusion from soil vapor  
21 (ATSDR, 2006b). Contaminated soil may be a source of either dermal or ingestion exposure of  
22 TCE for children (Wu and Schaum, 2000).

23 Additional TCE exposure has also been documented to have occurred during medical  
24 procedures. TCE was used in the past as an anesthetic during childbirth (Beppu, 1968; Phillips  
25 and Macdonald, 1971) and surgery during childhood (Jasińska, 1965). These studies are  
26 discussed in more detail in Section 4.8.3.1.1. In addition, the TCE metabolite chloral hydrate has  
27 been used as an anesthetic for children for CAT scans (Steinberg, 1993).

28 Dose received per body weight for 3-ppm TCE via oral, dermal, dermal plus inhalation,  
29 and bathing scenarios was estimated for a 10-kg infant, a 22-kg child, and a 70-kg adult (Fan,  
30 1988; see Table 4-113). For the oral route (drinking water), an infant would receive a higher  
31 daily dose than a child, and the child more than the adult. For the dermal and dermal plus  
32 inhalation route, the child would receive more than the adult. For the bathing scenario, the infant  
33 and child would receive comparable amounts, more than the adult.

#### **4.10.1.1.3. Early lifestage-specific toxicokinetics**

1           Section 3 describes the toxicokinetics of TCE. However, toxicokinetics in developmental  
2 lifestages are distinct from toxicokinetics in adults (Benedetti et al., 2007; Ginsberg et al., 2004a;  
3 Ginsberg et al., 2004b; Ginsberg et al., 2002; Hattis et al., 2003) due to, for example, altered  
4 ventilation rates, percentage of adipose tissue, and metabolic enzyme expression. Early  
5 lifestage-specific information is described below for absorption, distribution, metabolism, and  
6 excretion, followed by available early lifestage-specific PBPK models.

7

**Table 4-113. Estimated lifestage-specific daily doses for TCE in water<sup>a</sup>**

	Body weight		
	Infant (10 kg)	Child (22 kg)	Adult (70 kg)
Drinking water	0.3 mg/kg	0.204 mg/kg	0.086 mg/kg
Showering—dermal	-	0.1 mg/kg	0.064 mg/kg
Showering—dermal and inhalation	-	0.129 mg/kg	0.083 mg/kg
Bathing—15 min	-	0.24 mg/kg	0.154 mg/kg
Bathing—5 min	0.08 mg/kg	0.08 mg/kg	0.051 mg/kg

<sup>a</sup> Adapted from Fan (1988).

#### 4.10.1.1.3.1. Absorption

As discussed in Section 3.1, exposure to TCE may occur via inhalation, ingestion, and dermal absorption. In addition, prenatal exposure may result in absorption via the transplacental route. Exposure via inhalation is proportional to the ventilation rate, duration of exposure, and concentration of expired air, and children have increased ventilation rates per kg body weight compared to adults, with an increased alveolar surface area per kg body weight for the first two years (U.S. EPA, 2008b). It is not clear to what extent dermal absorption may be different for children compared to adults; however, infants have a twofold increase in surface area compared to adults, although similar permeability (except for premature babies) compared to adults (U.S. EPA, 2008b).

#### 4.10.1.1.3.2. Distribution

Both human and animal studies provide clear evidence that TCE distributes widely to all tissues of the body (see Section 3.2). For lipophilic compounds such as TCE, percentage adipose tissue, which varies with age, will affect absorption and retention of the absorbed dose. Infants have a lower percentage of adipose tissue per body weight than adults, resulting in a higher concentration of the lipophilic compound in the fat of the child (NRC, 1993).

During pregnancy of humans and experimental animals, TCE is distributed to the placenta (Beppu, 1968; Ghantous et al., 1986; Helliwell and Hutton, 1950; Laham, 1970; Withey and Karpinski, 1985). In humans, TCE has been found in newborn blood after exposure to TCE during childbirth with ratios of concentrations in fetal:maternal blood ranging from

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 approximately 0.5 to approximately 2 (Laham, 1970). In childhood, blood levels concentrations  
2 of TCE were found to range from 0.01–0.02 ng/mL (Sexton et al., 2005). Pregnant rats exposed  
3 to TCE vapors on GD 17 resulted in concentrations of TCE in fetal blood approximately one-  
4 third the concentration in corresponding maternal blood, and was altered based upon the position  
5 along the uterine horn (Withey and Karpinski, 1985). TCE has also been found in the organs of  
6 prenatal rabbits including the brain, liver, kidneys and heart (Beppu, 1968). Rats prenatally  
7 exposed to TCE had increased levels measured in the brain at PND 10, compared to rats exposed  
8 as adults (Rodriguez et al., 2007). TCE can cross the blood:brain barrier during both prenatal  
9 and postnatal development, and may occur to a greater extent in younger children. It is also  
10 important to note that it has been observed in mice that TCE can cycle from the fetus into the  
11 amniotic fluid and back to the fetus (Ghantous et al., 1986).

12 Studies have examined the differential distribution by age to a mixture of six VOCs  
13 including TCE to children aged 3–10 years and adults aged 20–82 years old (Mahle et al., 2007)  
14 and in rats at PND 10, 2 months (adult), and 2 years (aged) (Mahle et al., 2007; Rodriguez et al.,  
15 2007). In humans, the blood:air partition coefficient for male or female children was  
16 significantly lower compared to adult males (Mahle et al., 2007). In rats, the difference in  
17 tissue:air partition coefficients increased with age (Mahle et al., 2007). Higher peak  
18 concentrations of TCE in the blood were observed in the PND 10 rat compared to the adult rat  
19 after inhalation exposure, likely due to the lower metabolic capacity of the young rats (Rodriguez  
20 et al., 2007).

21

#### 4.10.1.1.3.3. **Metabolism**

22 Section 3.3 describes the enzymes involved in the metabolism of TCE, including CYP  
23 and GST. Expression of these enzymes changes during various stages of fetal development  
24 (Dorne et al., 2005; Hakkola et al., 1996a; Hakkola et al., 1998a; Hakkola et al., 1996b; 1998b;  
25 Hines and McCarver, 2002; Shao et al., 2007; van Lieshout et al., 1998) and during postnatal  
26 development (Blake et al., 2005; Dorne et al., 2005; Tateishi et al., 1997), and may result in  
27 altered susceptibility.

28 Expression of CYP enzymes have been shown to play a role in decreasing the  
29 metabolism of TCE during pregnancy in rats, although metabolism increased in young rats  
30 (3-week-old) compared to adult rats (18-week-old) (Nakajima et al., 1992b). For TCE, CYP2E1  
31 is the main metabolic CYP enzyme, and expression of this enzyme has been observed in humans  
32 in prenatal brain tissue at low levels beginning at 8-weeks gestation and increasing throughout  
33 gestation (Brzezinski et al., 1999). Very low levels of CYP2E1 have been detected in some

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 samples fetal liver during the second trimester (37% of samples) and third trimester (80% of  
2 samples) (Carpenter et al., 1996; Johnsrud et al., 2003), although hepatic expression surges  
3 immediately after birth in most cases (Johnsrud et al., 2003; Vieira et al., 1996) and in most  
4 infants reaches adult values by 3 months of age (Johnsrud et al., 2003; Vieira et al., 1996).

5 Although there is some uncertainty as to which GST isoforms mediate TCE conjugation,  
6 it should be noted that their expression changes with fetal development (McCarver and Hines,  
7 2002; Rajmakers et al., 2001; van Lieshout et al., 1998).

8

#### 4.10.1.1.3.4. **Excretion**

9 The major processes of excretion of TCE and its metabolites are discussed in Section 3.4,  
10 yet little is known about whether there are age-related differences in excretion of TCE. The  
11 major pathway for elimination of TCE is via exhalation, and its metabolites via urine and feces,  
12 and it is known that renal processes are not mature until about 6 months of age (NRC, 1993).  
13 Only one case study was identified that measured TCE or its metabolites in exhaled breath and  
14 urine in a 17-year old who ingested a large quantity of TCE (Brüning et al., 1998). TCE has also  
15 been measured in the breast milk in lactating women (Fisher et al., 1997; Pellizzari et al., 1982),  
16 goats (Hamada and Tanaka, 1995), and rats (Fisher et al., 1990).

17

#### 4.10.1.1.3.5. **Physiologically based pharmacokinetic (PBPK) models**

18 Early lifestage-specific information regarding absorption, distribution, metabolism, and  
19 excretion needs to be considered for a child-specific and chemical-specific PBPK model. To  
20 adequately address the risk to infants and children, age-specific parameters for these values  
21 should be used in PBPK models that can approximate the internal dose an infant or child receives  
22 based on a specific exposure level (see Section 3.5).

23 Fisher et al. developed PBPK models to describe the toxicokinetics of TCE in the  
24 pregnant rat (Fisher et al., 1989), lactating rat and nursing pup (Fisher et al., 1990). The prenatal  
25 study demonstrates that approximately two-thirds of maternal exposure to both TCE and TCA  
26 reached the fetus after maternal inhalation, gavage, or drinking water exposure (Fisher et al.,  
27 1989). After birth, only 2% of maternal exposure to TCE reaches the pup; however, 15% and  
28 30% of maternal TCA reaches the pup after maternal inhalation and drinking water exposure,  
29 respectively (Fisher et al., 1990). One analysis of PBPK models examined the variability in  
30 response to VOCs including TCE between adults and children, and concluded that the

1 intraspecies uncertainty factor for pharmacokinetics is sufficient to capture variability between  
2 adults and children (Pelekis et al., 2001).

3

#### 4.10.1.1.4. Early lifestage-specific effects

4 Although limited data exist on TCE toxicity as it relates to early lifestages, there is  
5 enough information to discuss the qualitative differences. In addition to the evidence described  
6 below, Section 4.8 contains information on reproductive and developmental toxicity. In  
7 addition, Sections 4.3 on neurotoxicity and Section 4.6 on immunotoxicity characterize a wide  
8 array of postnatal developmental effects.

9

##### 4.10.1.1.4.1. Differential noncancer outcomes in early lifestages

10 Some adverse health outcomes, in particular birth defects, are observed only after early  
11 lifestage exposure to TCE. A summary of structural developmental outcomes that have been  
12 associated with TCE exposures is presented in Sections 4.8.2.3.

13 Cardiac birth defects have been observed after exposure to TCE in humans (ATSDR,  
14 2006a; Goldberg et al., 1990; Lagakos et al., 1986; Yauck et al., 2004), rodents (Johnson et al.,  
15 2005)(Dawson et al., 1990a, 1993; Johnson et al., 1998a; Johnson et al., 2003; Johnson et al.,  
16 1998b; Smith et al., 1992; Smith et al., 1989), and chicks (Boyer et al., 2000; Bross et al., 1983;  
17 Drake et al., 2006a; Drake et al., 2006b; Loeber et al., 1988; Mishima et al., 2006; Rufer et al.,  
18 2008). However, it is notable that cardiac malformations were not observed in a number of other  
19 studies in humans (Lagakos et al., 1986; Taskinen et al., 1989; Tola et al., 1980), rodents  
20 (Carney et al., 2006; Coberly et al., 1992; Cosby and Dukelow, 1992; Dorfmueller et al., 1979;  
21 Fisher et al., 2001; Hardin et al., 1981; Healy et al., 1982; Narotsky and Kavlock, 1995;  
22 Narotsky et al., 1995; Schwetz et al., 1975), and rabbits (Hardin et al., 1981). See Section  
23 4.8.2.3.2 for further discussion on cardiac malformations.

24 Structural CNS birth defects were observed in humans (ATSDR, 2001; Bove, 1996; Bove  
25 et al., 1995; Lagakos et al., 1986). In addition, a number of postnatal nonstructural adverse  
26 effects on the CNS system have been observed in humans and experimental animals following  
27 prenatal exposure to TCE. See Sections 4.3.10 and 4.8.2.3.3 for further discussion on  
28 developmental neurotoxicity.

29 A variety of other birth defects have been observed—including eye/ear birth anomalies in  
30 humans and rats (Lagakos et al., 1986; Narotsky and Kavlock, 1995; Narotsky et al., 1995);  
31 lung/respiratory tract disorders in humans and mice (Das and Scott, 1994; Lagakos et al., 1986);

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 and oral cleft defects (Bove, 1996; Bove et al., 1995; Lagakos et al., 1986), kidney/urinary tract  
2 disorders, musculoskeletal birth anomalies (Lagakos et al., 1986), and anemia/blood disorders  
3 (Burg and Gist, 1999) in humans. See Section 4.8.2.3.5 for further discussion on other structural  
4 developmental outcomes. A current follow-up study of the Camp Lejeune cohort will examine  
5 birth defects and may provide additional insight (ATSDR, 2003c, 2009; U.S. GAO, 2007).  
6

#### 4.10.1.1.4.2. **Susceptibility to noncancer outcomes in early lifestages**

7 There are a number of adverse health outcomes observed after exposure to TCE that are  
8 observed in both children and adults. Below is a discussion of differential exposure, incidence  
9 and/or severity in early lifestages compared to adulthood.

10 Occupational TCE poisonings via inhalation exposure resulted in an elevated percentage  
11 of cases in the adolescents aged 15–19 years old compared those  $\geq 20$  years old (McCarthy and  
12 Jones, 1983). In addition, there is concern for intentional exposure to TCE during adolescence,  
13 including a series of deaths involving inhaling typewriter correction fluid (King et al., 1985) a  
14 case of glue sniffing likely associated with cerebral infarction in a 12-year-old boy with a 2-year  
15 history of exposure (Parker et al., 1984), and a case of attempted suicide by ingestion of 70 mg  
16 TCE in a 17-year-old boy (Brüning et al., 1998).  
17

##### 18 **4.10.1.1.4.2.1. Neurotoxicity**

19 Adverse CNS effects observed after early lifestage exposure to TCE in humans include  
20 delayed newborn reflexes (Beppu, 1968); impaired learning or memory (Bernad et al., 1987,  
21 abstract; White et al., 1997); aggressive behavior (Bernad et al., 1987, abstract; Blossom et al.,  
22 2008); hearing impairment (Burg and Gist, 1999; Burg et al., 1995); speech impairment (Burg  
23 and Gist, 1999; Burg et al., 1995; White et al., 1997); encephalopathy (White et al., 1997)  
24 impaired executive and motor function (White et al., 1997); attention deficit (Bernad et al., 1987,  
25 abstract) (White et al., 1997), and autism spectrum disorder (Windham et al., 2006). One  
26 analysis observed a trend for increased adversity during development, with those exposed during  
27 childhood demonstrating more deficits than those exposed during adulthood (White et al., 1997).  
28 In experimental animals, observations include decreased specific gravity of newborn brains until  
29 weaning (Westergren et al., 1984), reductions in myelination in the brains at weaning,  
30 significantly decreased uptake of 2-deoxyglucose in the neonatal rat brain, significant increase in  
31 exploratory behavior (Isaacson and Taylor, 1989; Noland-Gerbec et al., 1986; Taylor et al.,

1 1985), decreased rearing activity (Fredriksson et al., 1993), and increased time to cross the first  
2 grid in open field testing (George et al., 1986).

3 Few studies addressed whether or not children are more susceptible to CNS effects  
4 compared to adults (Burg and Gist, 1999; Burg et al., 1995; White et al., 1997). An analysis of  
5 three residential exposures of TCE observed speech impairments in younger children and not at  
6 any other lifestage (White et al., 1997). A national TCE exposure registry also observed  
7 statistically significant speech impairment and hearing impairment in 0–9 year olds and no other  
8 age group (Burg and Gist, 1999; Burg et al., 1995). However, a follow-up study did not find a  
9 continued association with speech and hearing impairment in these children, although the  
10 absence of acoustic reflexes remained significant (ATSDR, 2003b). See Section 4.3 for further  
11 information on central nervous system toxicity, and Section 4.8.3.3.3 for further information on  
12 developmental neurotoxicity.  
13

#### 14 **4.10.1.1.4.2.2. *Liver toxicity***

15 No early lifestage-specific effects were observed after TCE exposure. See Section 4.4 for  
16 further information on liver toxicity.  
17

#### 18 **4.10.1.1.4.2.3. *Kidney toxicity***

19 Residents of Woburn, Massachusetts including 4,978 children were surveyed on  
20 residential and medical history to examine an association between observed adverse health  
21 outcomes and wells contaminated with TCE and other chemicals; among these children, an  
22 association was observed for higher cumulative exposure measure and history of kidney and  
23 urinary tract disorders (primarily kidney or urinary tract infections) and with lung and respiratory  
24 disorders (asthma, chronic bronchitis, or pneumonia) (Lagakos et al., 1986). Comparisons were  
25 not made for the adults living in this community. See Section 4.5 for further information on  
26 kidney toxicity.  
27

#### 28 **4.10.1.1.4.2.4. *Immunotoxicity***

29 Several studies in exposure to TCE in early lifestages of humans (Lehmann et al., 2001;  
30 Lehmann et al., 2002) and experimental animals (Adams et al., 2003; Blossom and Doss, 2007;  
31 Blossom et al., 2008; Peden-Adams et al., 2006; Peden-Adams et al., 2008) were identified that  
32 assessed the potential for developmental immunotoxicity. While some noted evidence of  
33 immune system perturbation (Adams et al., 2003; Blossom and Doss, 2007; Blossom et al.,

1 2008; Lehmann et al., 2002; Peden-Adams et al., 2006), others did not (Lehmann et al., 2001;  
2 Peden-Adams et al., 2008). However, none of these studies assessed whether exposure during  
3 early life resulted in evidence of increased susceptibility as compared to exposure during  
4 adulthood; this is an area for future research. See Section 4.6 for further information on  
5 immunotoxicity, and Section 4.8.2.3.4 for further discussion on developmental immunotoxicity.  
6

#### 7 **4.10.1.1.4.2.5. *Respiratory toxicity***

8 Residents of Woburn, Massachusetts including 4,978 children were surveyed on  
9 residential and medical history to examine an association between observed adverse health  
10 outcomes and wells contaminated with TCE and other chemicals; among these children, an  
11 association was observed for lung and respiratory disorders (asthma, chronic bronchitis, or  
12 pneumonia) (Lagakos et al., 1986). Comparisons were not made for the adults living in this  
13 community. See Section 4.7 for further information on respiratory tract toxicity.  
14

#### 4.10.1.1.4.3. **Susceptibility to cancer outcomes in early lifestages**

15 The epidemiologic and experimental animal evidence is limited regarding susceptibility  
16 to cancer from exposure to TCE during early lifestages. The human epidemiological evidence is  
17 summarized above for cancer diagnosed during childhood (see Sections 4.8.2.1 and 4.8.2.3.5),  
18 including a discussion of childhood cancers of the nervous system including neuroblastoma and  
19 the immune system including leukemia (see Section 4.6.1.3). A current follow-up study of the  
20 Camp Lejeune cohort will examine childhood cancers and may provide additional insight  
21 (ATSDR, 2003c, 2009; U.S. GAO, 2007). No studies of cancers in experimental animals in  
22 early lifestages have been observed.  
23

#### 24 **4.10.1.1.4.3.1. *Total childhood cancer***

25 Total childhood cancers have been examined in relationship to TCE exposure (ATSDR,  
26 2006a; Morgan and Cassady, 2002). Two studies examining total childhood cancer in relation to  
27 TCE in drinking water did not observe an association. A study in Endicott, NY contaminated by  
28 a number of VOCs, including “thousands of gallons” of TCE observed fewer than 6 cases of  
29 cancer diagnosed between 1980 and 2001 in children aged 0–19 years, and did not exceed  
30 expected cases or types (ATSDR, 2006a). A California community exposed to TCE in drinking  
31 water from contaminated wells was examined for cancer, with a specific emphasis on childhood

1 cancer (<15 years old); however, the incidence did not exceed those expected for the community  
2 (Morgan and Cassady, 2002). A third study of childhood cancer in relation to TCE in drinking  
3 water in Camp Lejeune, NC is currently underway (U.S. GAO, 2007).  
4

#### 5 **4.10.1.1.4.3.2. *Childhood leukemia***

6 Childhood leukemia has been examined in relationship to TCE exposure (Cohn et al.,  
7 1994b; Costas et al., 2002; Lagakos et al., 1986; Lowengart et al., 1987; McKinney et al., 1991;  
8 Shu et al., 1999). In a study examining drinking water exposure to TCE in 75 New Jersey towns,  
9 childhood leukemia, (including ALL) was significantly increased for girls ( $n = 6$ ) diagnosed  
10 before age 20 years, but this was not observed for boys (Cohn et al., 1994b). A community in  
11 Woburn, MA with contaminated well water including TCE experienced 20 cases of childhood  
12 leukemia, significantly more than expected (Lagakos et al., 1986); however, the incidence of  
13 leukemia among children was not compared to the incidence rate among adults living in this  
14 community. Further analysis by Costas et al. (2002) also observed a greater than twofold  
15 increase over expected cases of childhood leukemia. Cases were more likely to be male (76%),  
16 <9 years old at diagnosis (62%), breast-fed (OR: 10.17, 95% CI: 1.22–84.50), and exposed  
17 during pregnancy (adjusted OR: 8.33, 95% CI: 0.73–94.67). The highest risk was observed for  
18 exposure during pregnancy compared to preconception or postnatal exposure, and a dose-  
19 response was seen for exposure during pregnancy (Costas et al., 2002). In addition, family  
20 members of those diagnosed with childhood leukemia, including 13 siblings under age 19 at the  
21 time of exposure, had altered immune response, but an analysis looking at only these children  
22 was not done (Byers et al., 1988).

23 Case-control studies examined children diagnosed with ALL for parental occupational  
24 exposures and found a nonsignificant two- to fourfold increase of childhood leukemia risk for  
25 exposure to TCE during preconception, pregnancy, postnatally, or all developmental periods  
26 combined (Lowengart et al., 1987; McKinney et al., 1991; Shu et al., 1999). Some studies  
27 showed an elevated risk for maternal (Shu et al., 1999) or paternal exposure (Lowengart et al.,  
28 1987; McKinney et al., 1991), while others did not show an elevated risk for maternal  
29 (McKinney et al., 1991) or paternal exposure (Shu et al., 1999), possibly due to the small number  
30 of cases. No variability was observed in the developmental stages in Shu et al. (1999), although  
31 Lowengart et al. (1987) observed the highest risk to be paternal exposure to TCE after birth.  
32

1 **4.10.1.1.4.3.3. CNS tumors**

2 In a case-control study of parental occupational exposures, paternal self-reported  
3 exposure to TCE was not significantly associated with neuroblastoma in the offspring (OR = 1.4,  
4 95% CI: 0.7–2.9) (De Roos et al., 2001). Brain tumors have also been observed in the offspring  
5 of fathers exposed to TCE, but the odds ratio could not be determined (Peters et al., 1985; Peters  
6 et al., 1981).

7  
8 **4.10.1.1.4.3.4. Age-dependent adjustment factors (ADAFs)**

9 According to EPA’s *Supplemental Guidance for Assessing Susceptibility from Early-Life*  
10 *Exposure to Carcinogens* (U.S. EPA, 2005d) there may be increased susceptibility to early-life  
11 exposures for carcinogens with a mutagenic MOA. Therefore, because the weight of evidence  
12 supports a mutagenic MOA for TCE carcinogenicity in the kidney (see Section 4.4.7), the lack of  
13 data suggesting an absence of GSTT1 expression in neonates, and in the absence of chemical-  
14 specific data to evaluate differences in susceptibility, early-life susceptibility should be assumed  
15 and the ADAFs should be applied, in accordance with the *Supplemental Guidance*.

16  
**4.10.1.1.5. Later Lifestages**

17 Few studies examine the differential effects of TCE exposure for elderly adults  
18 (>65 years old). These limited studies suggest that older adults may experience increased  
19 adverse effects than younger adults. However, there is no further evidence for elderly  
20 individuals exposed to TCE beyond these studies.

21 Toxicokinetics in later lifestages can be distinct from toxicokinetics in younger adults  
22 (Benedetti et al., 2007; Ginsberg et al., 2005), although there is limited evidence showing a  
23 possible age-related difference in CYP expression (Dorne et al., 2005; George et al., 1995;  
24 Parkinson et al., 2004). GST expression has been observed to decrease with age in human  
25 lymphocytes, with the lowest expression in those aged 60–80 years old (van Lieshout and Peters,  
26 1998).

27 Studies have examined the age differences in TK after exposure to a mixture of six VOCs  
28 including TCE for humans (Mahle et al., 2007) and rats (Mahle et al., 2007; Rodriguez et al.,  
29 2007). In humans, the blood:air partition coefficient for adult males (20–82 years) was  
30 significantly ( $p \leq 0.05$ ) higher ( $11.7 \pm 1.9$ ) compared to male ( $11.2 \pm 1.8$ ) or female ( $11.0 \pm 1.6$ )  
31 children (3–10 years) (Mahle et al., 2007); when the data was stratified for adults above and  
32 below 55 years of age, there was no significant difference observed between adults (20–55

1 years) and aged (56–82) (data not reported). In rats, the difference in tissue:air partition  
2 coefficients also increased from PND10 to adult (2 months) to aged (2 years) rat (Mahle et al.,  
3 2007). TCE has also been measured in the brain of rats, with an increased level observed in  
4 older (2 year old) rats compared to adult (2 month old) rats (Rodriguez et al., 2007). It was also  
5 observed that aged rats reached steady state slower with higher concentrations compared to the  
6 adult rat; the authors suggest that the almost twofold greater percentage of body fat in the elderly  
7 is responsible for this response (Rodriguez et al., 2007).

8 One cohort of TCE exposed metal degreasers found an increase in psychoorganic  
9 syndrome and increased vibration threshold related to increasing age (Rasmussen et al., 1993b;  
10 Rasmussen et al., 1993c; Rasmussen et al., 1993d), although the age groups were  $\leq 29$  years,  
11 30–39 years, and 40+ years, but the age ranged only from 18–68 years and did not examine  $>65$   
12 years as a separate category.

#### 4.10.2. Other Susceptibility Factors

13 Aside from age, many other factors may affect susceptibility to TCE toxicity. A partial  
14 list of these factors includes gender, genetic polymorphisms, preexisting disease status,  
15 nutritional status, diet, and previous or concurrent exposures to other chemicals. The toxicity  
16 that results due to changes in multiple factors may be quite variable, depending on the exposed  
17 population and the type of exposure. Qualitatively, the presence of multiple susceptibility  
18 factors will increase the variability that is seen in a population response to TCE toxicity.

##### 4.10.2.1.1. Gender

19 Individuals of different genders are physiologically, anatomically, and biochemically  
20 different. Males and females can differ greatly in many physiological parameters such as body  
21 composition, organ function, and ventilation rate, which can influence the toxicokinetics of  
22 chemicals and their metabolites in the body (Gandhi et al., 2004; Gochfeld, 2007).

##### 4.10.2.1.2. Gender-specific toxicokinetics

24 Section 3 describes the toxicokinetics of TCE. Gender-specific information is described  
25 below for absorption, distribution, metabolism, and excretion, followed by available gender-  
26 specific PBPK models.



#### 4.10.2.1.2.1. **Absorption**

1 As discussed in Section 3.1, exposure to TCE may occur via inhalation, ingestion, and  
2 skin absorption. Exposure via inhalation is proportional to the ventilation rate, duration of  
3 exposure, and concentration of expired air, and women have increased ventilation rates during  
4 exercise compared to men (Gochfeld, 2007). Percentage of body fat varies with gender  
5 (Gochfeld, 2007), which for lipophilic compounds such as TCE will affect absorption and  
6 retention of the absorbed dose. After experimental exposure to TCE, women were found to  
7 absorb a lower dose due to lower alveolar intake rates compared to men (Sato, 1993; Sato et al.,  
8 1991b).

#### 4.10.2.1.2.2. **Distribution**

10 Both human and animal studies provide clear evidence that TCE distributes widely to all  
11 tissues of the body (see Section 3.2). The distribution of TCE to specific organs will depend on  
12 organ blood flow and the lipid and water content of the organ, which may vary between genders  
13 (Gochfeld, 2007). After experimental exposure to humans, higher distribution of TCE into fat  
14 tissue was observed in women leading to a greater blood concentration 16 hours after exposure  
15 compared to men (Sato, 1993; Sato et al., 1991b). In experimental animals, male rats generally  
16 have higher levels of TCE in tissues compared to female rats, likely due to gender differences in  
17 metabolism (Lash et al., 2006). In addition, TCE has been observed in the male reproductive  
18 organs (epididymis, vas deferens, testis, prostate, and seminal vesicle) (Zenick et al., 1984).

#### 4.10.2.1.2.3. **Metabolism**

20 Section 3.3 describes the metabolic processes involved in the metabolism of TCE,  
21 including CYP and GST enzymes. In addition, the role of metabolism in male reproductive  
22 toxicity is discussed in Section 4.8.1.3.2. In general, there is some indication that TCE  
23 metabolism is different between males and females, with females more rapidly metabolizing  
24 TCE after oral exposure to rats (Lash et al., 2006), intraperitoneal injections in rats (Verma and  
25 Rana, 2003), and in mouse, rat and human liver microsomes (Elfarra et al., 1998).

26 In general, CYP expression may differ between genders (Gandhi et al., 2004; Gochfeld,  
27 2007; Parkinson et al., 2004), although no gender-related difference in CYP2E1 activity is  
28 observed in the human liver microsomes (George et al., 1995; Parkinson et al., 2004). After  
29 exposure to TCE, CYP2E1 was detected in the epididymis and testes of mice (Forkert et al.,

1 2002), and CYP2E1 and GST-alpha has been detected in the ovaries of rats (Wu and Berger,  
2 2008), indicating that metabolism of TCE can occur in both the male and female reproductive  
3 tracts. One study of TCE exposure in mice observed induced CYP2E1 expression in the liver of  
4 males only (Nakajima et al., 2000). Male rats have been shown to have higher levels of TCE  
5 metabolites in the liver (Lash et al., 2006), and lower levels of TCE metabolites in the kidney  
6 (Lash et al., 2006) compared to female rats. However, another study did not observe any sex-  
7 related differences in the metabolism of TCE in rats (Nakajima et al., 1992b).

8 Unlike CYP-mediated oxidation, quantitative differences in the polymorphic distribution  
9 or activity levels of GST isoforms in humans are not presently known. However, the available  
10 data (Lash et al., 1999a; Lash et al., 1999b) do suggest that significant variation in GST-  
11 mediated conjugation of TCE exists in humans. One study observed that GSH conjugation is  
12 higher in male rats compared to female rats (Lash et al., 2000b); however, it has also been  
13 speculated that any gender difference may be due to a polymorphism in GSH conjugation of  
14 TCE rather than a true gender difference (Lash et al., 1999a). Also, induction of PPAR $\alpha$   
15 expression in male mice after TCE exposure was greater than that in females (Nakajima et al.,  
16 2000).

#### 17 4.10.2.1.2.4. Excretion

18 The major processes of excretion of TCE and its metabolites are discussed in Section 3.4.  
19 Two human voluntary inhalation exposure studies observed the levels of TCE and its metabolites  
20 in exhaled breath and urine (Kimmerle and Eben, 1973b; Nomiyama and Nomiyama, 1971).  
21 Increased levels of TCE in exhaled breath in males were observed in one human voluntary  
22 inhalation exposure study of 250–380 ppm for 160 minutes (Nomiyama and Nomiyama, 1971),  
23 but no difference was observed in another study of 40 ppm for 4 hours or 50 ppm for 4 hours for  
24 5 days (Kimmerle and Eben, 1973b).

25 After experimental exposure to TCE, women were generally found to excrete higher  
26 levels of TCE and TCA compared to men (Kimmerle and Eben, 1973b; Nomiyama and  
27 Nomiyama, 1971). However, other studies observed an increase in TCE in the urine of males  
28 (Inoue et al., 1989), an increase in TCA in the urine of males (Sato et al., 1991b), or no  
29 statistically significant ( $p > 0.10$ ) gender difference for TCA in the urine (Inoue et al., 1989).  
30 Others found that the urinary elimination half-life of TCE metabolites is longer in women  
31 compared to men (Ikeda, 1977; Ikeda and Imamura, 1973).

32 In addition to excretion pathways that occur in both genders, excretion occurs uniquely in  
33 men and women. In both humans and experimental animals, it has been observed that females

1 can excrete TCE and metabolites in breast milk (Fisher et al., 1997; Fisher et al., 1990; Hamada  
2 and Tanaka, 1995; Pellizzari et al., 1982), while males can excrete TCE and metabolites in  
3 seminal fluid (Forkert et al., 2003; Zenick et al., 1984).  
4

#### 4.10.2.1.2.5. **Physiologically based pharmacokinetic (PBPK) models**

5 Gender-specific differences in uptake and metabolism of TCE were incorporated into a  
6 PBPK model using human exposure data (Fisher et al., 1998). The chemical-specific parameters  
7 included cardiac output at rest, ventilation rates, tissue volumes, blood flow, and fat volume.  
8 This model found that gender differences for the toxicokinetics of TCE are minor.  
9

#### 4.10.2.1.3. **Gender -specific effects**

##### 4.10.2.1.3.1. **Gender susceptibility to noncancer outcomes**

###### 10 **4.10.2.1.3.1.1. *Liver toxicity***

11 No gender susceptibility to noncancerous outcomes in the liver was observed. A detailed  
12 discussion of the studies examining the effects of TCE on the liver can be found in Section 4.4.  
13

###### 14 **4.10.2.1.3.1.2. *Kidney toxicity***

15 A detailed discussion of the studies examining the noncancer effects of TCE on the  
16 kidney can be found in Section 4.5. A residential study found that females aged 55–64 years old  
17 had an elevated risk of kidney disease (RR = 4.57, 99% CI: 2.10–9.93) compared to males ,  
18 although an elevated risk of urinary tract disorders was reported for both males and females  
19 (Burg et al., 1995). Additionally, a higher rate of diabetes in females compared to males exposed  
20 to TCE was reported in two studies (Burg et al., 1995; Davis et al., 2005). In rodents, however,  
21 and kidney weights were increased more in male mice than in females (Kjellstrand et al., 1983a;  
22 Kjellstrand et al., 1983b), and male rats have exhibited increased renal toxicity to TCE compared  
23 to females (Lash et al., 2001b; Lash et al., 1998a).  
24

1 **4.10.2.1.3.1.3. Immunotoxicity**

2 A detailed discussion of the studies examining the immunotoxic effects of TCE can be  
3 found in Section 4.6. Most of the immunotoxicity studies present data stratified by sex. The  
4 prevalence of exposure to TCE is generally lower in women compared with men. In men, the  
5 studies generally reported odds ratios between 2.0 and 8.0, and in women, the odds ratios were  
6 between 1.0 and 2.0 (Cooper et al., 2009). Based on small numbers of cases, an occupational  
7 study of TCE exposure found an increased risk for systemic sclerosis for men (OR: 4.75,  
8 95% CI: 0.99–21.89) compared to women (OR: 2.10; 95% CI: 0.65–6.75) (Diot et al., 2002).  
9 Another study found similar results, with an elevated risk for men with a maximum intensity,  
10 cumulative intensity and maximum probability of exposure to TCE compared to women (Nietert  
11 et al., 1998). These two studies, along with one focused exclusively on the risk of scleroderma  
12 to women (Garabrant et al., 2003), were included in a meta-analysis conducted by the EPA  
13 resulting in a combined estimate for “any” exposure, was OR = 2.5 (95% CI: 1.1, 5.4) for men  
14 and OR = 1.2 (95% CI: 0.58, 2.6) in women.  
15

16 **4.10.2.1.3.1.4. Respiratory toxicity**

17 No gender susceptibility to noncancerous outcomes in the respiratory tract after TCE  
18 exposure was observed. A detailed discussion of the studies examining the respiratory effects of  
19 TCE can be found in Section 4.7.  
20

21 **4.10.2.1.3.1.5. Reproductive toxicity**

22 A detailed discussion of the studies examining the gender-specific noncancer  
23 reproductive effects of TCE can be found in Section 4.8.1.

24 Studies examining males after exposure to TCE observed altered sperm morphology and  
25 hyperzoospermia (Chia et al., 1996), altered endocrine function (Chia et al., 1997; Goh et al.,  
26 1998), decreased sexual drive and function (Bardodej and Vyskocil, 1956; El Ghawabi et al.,  
27 1973; Saihan et al., 1978), and altered fertility to TCE exposure. Infertility was not associated  
28 with TCE exposure in other studies (Forkert et al., 2003; Sallmen et al., 1998), and sperm  
29 abnormalities were not observed in another study (Rasmussen et al., 1988).

30 There is more limited evidence for reproductive toxicity in females. There are  
31 epidemiological indicators of a possible effect of TCE exposure on female fertility (Sallmen et  
32 al., 1998), increased rate of miscarriage (ATSDR, 2001), and menstrual cycle disturbance  
33 (ATSDR, 2001; Bardodej and Vyskocil, 1956; Zielinski, 1973). In experimental animals, the  
34 effects on female reproduction include evidence of reduced in vitro oocyte fertilizability in rats

1 (Berger and Horner, 2003; Wu and Berger, 2007, 2008). However, in other studies that assessed  
2 reproductive outcome in female rodents (Cosby and Dukelow, 1992; George et al., 1985, 1986;  
3 Manson et al., 1984), there was no evidence of adverse effects of TCE exposure on female  
4 reproductive function.  
5

#### 6 **4.10.2.1.3.1.6. *Developmental toxicity***

7 A detailed discussion of the studies examining the gender-specific noncancer  
8 developmental effects of TCE can be found in Section 4.8.3. Only one study of contaminated  
9 drinking water exposure in Camp Lejeune, NC observed a higher risk of SGA in males compared  
10 to females (ATSDR, 1998b; Sonnenfeld et al., 2001).  
11

#### 12 **4.10.2.1.3.2. *Gender susceptibility to cancer outcomes***

13 A detailed discussion of the studies examining the carcinogenic effects of TCE can be  
14 found on the liver in Section 4.4, on the kidney in Section 4.5, in the immune system in  
15 Section 4.6.4, in the respiratory system in Sections 4.7.1.2 and 4.7.3, and on the reproductive  
16 system in Section 4.8.2.

#### 17 **4.10.2.1.3.2.1. *Liver cancer***

18 An elevated risk of liver cancer was observed for females compared to males in both  
19 human (Raaschou-Nielsen et al., 2003) and rodent (Elfarra et al., 1998) studies. In addition,  
20 gallbladder cancer was significantly elevated for women compared to men (Raaschou-Nielsen et  
21 al., 2003). A detailed discussion of the studies examining the gender-specific liver cancer effects  
22 of TCE can be found in Section 4.4.  
23

#### 24 **4.10.2.1.3.2.2. *Kidney cancer***

25 One study of occupational exposure to TCE observed an increase in renal cell carcinoma  
26 for women compared to men (Dosemeci et al., 1999), but no gender difference was observed in  
27 other studies (Pesch et al., 2000b; Raaschou-Nielsen et al., 2003). Blair et al. (1998) and Hansen  
28 et al. (2001) also present some results by sex, but both of these studies have too few cases to be  
29 informative about a sex difference for kidney cancer. Exposure differences between males and  
30 females in Dosemeci et al. (1999) may explain their finding. These studies, however, provide  
31 little information to evaluate susceptibility between sexes because of their lack of quantitative

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 exposure assessment and lower statistical power. A detailed discussion of the studies examining  
2 the gender-specific kidney cancer effects of TCE can be found in Section 4.5.

#### 4 **4.10.2.1.3.2.3. *Cancers of the immune system***

5 Two drinking water studies suggest that there may be an increase of leukemia (Cohn et  
6 al., 1994b; Fagliano et al., 1990) and NHL (Cohn et al., 1994b) among females compared to  
7 males. An occupational study also observed an elevated risk of leukemia in females compared to  
8 males (Raaschou-Nielsen et al., 2003), although a study of contaminated drinking water in  
9 Woburn, MA observed an increased risk of childhood leukemia in males compared to females  
10 (Costas et al., 2002). A detailed discussion of the studies examining the gender-specific cancers  
11 of the immune system following TCE exposure can be found in Section 4.6.4.

#### 13 **4.10.2.1.3.2.4. *Respiratory cancers***

14 One study observed significantly elevated risk of lung cancer following occupational  
15 TCE exposure for both men and women, although the risk was found to be higher for women  
16 compared to men (Raaschou-Nielsen et al., 2003). This same study observed a nonsignificant  
17 elevated risk in both men and women for laryngeal cancer, again with an increased risk for  
18 women compared to men (Raaschou-Nielsen et al., 2003). Conversely, a study of Iowa residents  
19 with TCE-contaminated drinking water observed a sevenfold increased annual age-adjusted  
20 incidence for males compared to females (Isacson et al., 1985). However, other studies did not  
21 observe a gender-related difference (ATSDR, 2003b; Blair et al., 1998; Hansen et al., 2001). A  
22 detailed discussion of the studies examining the gender-specific respiratory cancers following  
23 TCE exposure can be found in Sections 4.7.1.2 and 4.7.3.

#### 25 **4.10.2.1.3.2.5. *Reproductive cancers***

26 Breast cancer in females and prostate cancer in males was reported after exposure to TCE  
27 in drinking water (Isacson et al., 1985). A statistically elevated risk for cervical cancer, but not  
28 breast, ovarian or uterine cancer, was observed in women in another study (Raaschou-Nielsen et  
29 al., 2003). This study also did not observe elevated prostate or testicular cancer (Raaschou-  
30 Nielsen et al., 2003). A detailed discussion of the studies examining the gender-specific  
31 reproductive cancers following TCE exposure can be found in Section 4.8.2.

1 **4.10.2.1.3.2.6. Other Cancers**

2 Bladder and rectal cancer was increased in men compared to women after exposure to  
3 TCE in drinking water, but no gender difference was observed for colon cancer (Isacson et al.,  
4 1985). After occupational TCE exposure, bladder, stomach, colon, and esophageal cancer was  
5 nonsignificantly elevated in women compared to men (Raaschou-Nielsen et al., 2003).

6  
**4.10.2.1.4. Genetic Variability**

7 Section 3.3 describes the metabolic processes involved in the metabolism of TCE.  
8 Human variation in response to TCE exposure may be associated with genetic variation. TCE is  
9 metabolized by both CYP and GST; therefore, it is likely that polymorphisms will alter the  
10 response to exposure (Garte et al., 2001; Nakajima and Aoyama, 2000), as well as exposure  
11 other chemicals that may alter the metabolism of TCE (Lash et al., 2007) (see Section 4.10.2.6).  
12 It is important to note that even with a given genetic polymorphism, metabolic expression is not  
13 static, and depends on lifestage (see Section 4.10.1.1.2), obesity (see Section 4.10.2.4.1), and  
14 alcohol intake (see Section 4.10.2.5.1).

**4.10.2.1.5. CYP genotypes**

15 In general, variability in CYP expression occurs within humans (Dorne et al., 2005), and  
16 variability in CYP expression has been observed in experimental animals exposed to TCE  
17 (Nakajima et al., 1993). In particular, increased CYP2E1 activity may lead to increased  
18 susceptibility to TCE (Lipscomb et al., 1997). The CYP2E1\*3 allele and the CYP2E1\*4 allele  
19 were more common among those who developed scleroderma who were exposed to solvents  
20 including TCE (Povey et al., 2001). A PBPK model of CYP2E1 expression after TCE exposure  
21 has been developed for rats and humans (Yoon et al., 2007).

22 In experimental animals, toxicokinetics of TCE differed among CYP2E1 knockout and  
23 wild-type mice (Kim and Ghanayem, 2006). This study found that exhalation was more  
24 prevalent among the knockout mice, whereas urinary excretion was more prevalent among the  
25 wild-type mice. In addition, the dose was found to be retained to a greater degree by the  
26 knockout mice compared to the wild-type mice.

**4.10.2.1.6. GST genotype**

27 There is a possibility that GST polymorphisms could play a role in variability in toxic  
28 response to TCE (Caldwell and Keshava, 2006), but this has not been sufficiently tested

1 (NRC, 2006). One study of renal cell cancer in workers exposed to TCE demonstrated a  
2 significant increased for those with GSTM1+ and GSTT1+ polymorphisms, compared to a  
3 negative risk for those with GSTM1– and GSTT1-polymorphisms (Brüning et al., 1997a).  
4 Another study of occupational TCE exposure found that renal cell carcinoma was significantly  
5 associated with the GSTT+ polymorphism but not with GSTT– (Moore et al., 2010). However,  
6 another study did not confirm this hypothesis, observing no clear relationship between GSTM1  
7 and GSTT1 polymorphisms and renal cell carcinoma among TCE-exposed individuals, although  
8 they did see a possible association with the homozygous wild-type allele GSTP1\*A  
9 (Wiesenhütter et al., 2007). Unrelated to TCE exposure, Sweeney et al. (2000) found GSTT1–  
10 to be associated with an increased risk of renal cell carcinoma, but no difference was seen for  
11 GSTM1 and GSTP1 alleles. The role of GST polymorphisms in the development of renal cell  
12 carcinoma is an area in need of future research.

13

#### 4.10.2.1.7. Other genotypes

14 Other genetic polymorphisms could play a role in variability in toxic response, in  
15 particular TCE-related skin disorders. Studies have found that many TCE-exposed patients  
16 diagnosed with skin conditions exhibited the slow-acetylator NAT2 genotype (Huang et al.,  
17 2002; Nakajima et al., 2003); whereas there was no difference in NAT2 status for those  
18 diagnosed with renal cell carcinoma (Wiesenhütter et al., 2007). Other studies have found that  
19 many TCE-exposed patients diagnosed with skin conditions expressed variant HLA alleles (Li et  
20 al., 2007; Yue et al., 2007), in particular HLA-B\*1301 which is more common in Asians  
21 compared to whites (Cao et al., 2001; Williams et al., 2001); or TNF  $\alpha$ -308 allele (Dai et al.,  
22 2004). Also, an in vitro study of human lung adenocarcinoma cells exposed to TCE varied in  
23 response based on their p53 status, with p53-wild-type cells resulting in severe cellular damage,  
24 but not the p53-null cells (Chen et al., 2002a).

25

#### 4.10.2.1.8. Race/Ethnicity

26 Different racial or ethnic groups may express metabolic enzymes in different ratios and  
27 proportions due to genetic variability (Garte et al., 2001). In particular, ethnic variability in CYP  
28 (Dorne et al., 2005; McCarver et al., 1998; Parkinson et al., 2004; Shimada et al., 1994; Stephens  
29 et al., 1994) and GST (Nelson et al., 1995) expression has been reported.

30 It has been observed that the metabolic rate for TCE may differ between the Japanese and  
31 Chinese (Inoue et al., 1989). Also, body size varies among ethnic groups, and increased body

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 size was related to increased absorption of TCE and urinary excretion of TCE metabolites (Sato  
2 et al., 1991b).

#### 3 4.10.2.1.9. Preexisting Health Status

4 It is known that kidney and liver diseases can affect the clearance of chemicals from the  
5 body, and therefore, poor health may lead to increased half-lives for TCE and its metabolites.  
6 There are some data indicating that obesity/metabolic syndrome, diabetes and hypertension may  
7 increase susceptibility to TCE exposure through altered toxicokinetics. In addition, some of  
8 these conditions lead to increased risk for adverse effects that have also been associated with  
9 TCE exposure, though the possible interaction between TCE and known risk factors for these  
10 effects is not understood.

#### 11 4.10.2.1.10. Obesity

12 TCE is lipophilic and stored in adipose tissue; therefore, obese individuals may  
13 experience altered toxicokinetics of TCE compared to thin individuals. The absorption of TCE  
14 is increased in obese individuals compared to thin individuals (Clewell et al., 2000), as observed  
15 by lower blood concentrations immediately after exposure in obese men compared to thin men  
16 (Sato, 1993; Sato et al., 1991b). Once absorbed, obese individuals have increased storage of  
17 TCE in the adipose tissue compared to thin men (Clewell et al., 2000) which prolongs internal  
18 exposures (Davidson and Beliles, 1991; Lash et al., 2000b). Obesity also likely alters TCE  
19 metabolism, since increased CYP2E1 expression has been observed in obese individuals  
20 compared to thin individuals (McCarver et al., 1998). Finally, delayed excretion has been  
21 observed in obese individuals compared to thin individuals in both exhaled air (Monster, 1979)  
22 and urine (Sato, 1993; Sato et al., 1991b). In sum, obese individuals have altered toxicokinetics  
23 of TCE compared to thin individuals due to increased storage of TCE, increased CYP2E1  
24 metabolism, and a slower rate of elimination.

25 In addition, individuals with high BMI are at increased risk of some of the same health  
26 effects associated with TCE exposure. For example, renal cell carcinoma, liver cancer, and  
27 prostate cancer may be positively associated with BMI or obesity (Asal et al., 1988a; Asal et al.,  
28 1988b; Benichou et al., 1998; El-Serag and Rudolph, 2007; Wigle et al., 2008). However,  
29 whether and how TCE interacts with known risk factors for such diseases is unknown, as  
30 existing epidemiologic studies have only examined these factors as possible confounders for  
31 effects associated with TCE, or vice versa (Charbotel et al., 2006; Krishnadasan et al., 2008).

*This document is a draft for review purposes only and does not constitute Agency policy.*

1

#### **4.10.2.1.11. Diabetes**

2           A higher rate of diabetes in females compared to males exposed to TCE was reported in  
3 two studies (Burg et al., 1995; Davis et al., 2005). Whether the TCE may have caused the  
4 diabetes or the diabetes may have increased susceptibility to TCE is not clear. However, it has  
5 been observed that CYP2E1 expression is increased in obese Type II diabetics (Wang et al.,  
6 2003), and in poorly controlled Type I diabetics (Song et al., 1990), which may consequently  
7 alter the metabolism of TCE.

8

#### **4.10.2.1.12. Hypertension**

9           One study found no difference in risk for renal cell carcinoma among those diagnosed  
10 with hypertension among those living in an area with high TCE exposure; however, a slightly  
11 elevated risk was seen for those being treated for hypertension (OR: 1.57, 95% CI: 0.90–2.72)  
12 (Charbotel et al., 2006). Unrelated to TCE exposure, hypertension has been associated with  
13 increased risk of renal cell carcinoma in women compared to men (Benichou et al., 1998).

14

#### **4.10.2.1.13. Lifestyle Factors and Nutrition Status**

#### **4.10.2.1.14. Alcohol intake**

15           A number of studies have examined the interaction between TCE and ethanol exposure in  
16 both humans (Bardodej and Vyskocil, 1956; Barret et al., 1984; McCarver et al., 1998; Muller et  
17 al., 1975; Sato, 1993; Sato et al., 1991a; Sato et al., 1981; Stewart et al., 1974) and experimental  
18 animals (Kaneko et al., 1994a; Larson and Bull, 1989; Nakajima et al., 1988; Nakajima et al.,  
19 1992a; Nakajima et al., 1990; Okino et al., 1991; Sato and Nakajima, 1985; Sato et al., 1980,  
20 1983; White and Carlson, 1981).

21           The coexposure causes metabolic inhibition of TCE in humans (Muller et al., 1975;  
22 Windemuller and Ettema, 1978), male rats (Kaneko et al., 1994a; Larson and Bull, 1989;  
23 Nakajima et al., 1988; Nakajima et al., 1990; Nakanishi et al., 1978; Okino et al., 1991; Sato and  
24 Nakajima, 1985; Sato et al., 1981), and rabbits (White and Carlson, 1981). Similarly, individuals  
25 exposed to TCE reported an increase in alcohol intolerance (Bardodej and Vyskocil, 1956;

1 Grandjean et al., 1955; Rasmussen and Sabroe, 1986). Disulfiram, used to treat alcoholism, has  
2 also been found to decrease the elimination of TCE and TCA (Bartonicek and Teisinger, 1962).

3 A “degreasers flush” has been described, reflecting a reddening of the face of those  
4 working with TCE after drinking alcohol, and measured an elevated level of TCE in exhaled  
5 breath compared to nondrinkers exposed to TCE (Stewart et al., 1974). This may be due to  
6 increased CYP2E1 expression in those that consume alcohol compared to nondrinkers, unrelated  
7 to TCE exposure (Caldwell et al., 2008a; Liangpunsakul et al., 2005; Lieber, 2004; McCarver et  
8 al., 1998; Parkinson et al., 2004; Perrot et al., 1989).

9 In experimental animals, male rats pretreated with ethanol experienced an induction of  
10 TCE metabolism (Nakajima et al., 1992a), although another study of male rats observed that  
11 pretreatment with ethanol did not decrease CYP activity (Okino et al., 1991). It is important to  
12 note that there a further increased response of TCE and ethanol has been reported when also  
13 combined with low fat diets or low carbohydrate diets in male rats (Sato et al., 1983).

14 Since the liver is a target organ for both TCE and alcohol, decreased metabolism of TCE  
15 could be related to cirrhosis of the liver as a result of alcohol abuse (McCarver et al., 1998), and  
16 an increase in clinical liver impairment along with degreasers flush has been observed (Barret  
17 et al., 1984).

18 The central nervous system may also be impacted by the coexposure. Individuals  
19 exposed to TCE and ethanol reported an increase in altered mood states (Reif et al., 2003),  
20 decreased mental capacity as described as small increases in functional load (Windemuller and  
21 Ettema, 1978), and those exposed to TCE and tetrachloroethylene who consumed alcohol had an  
22 elevated color confusion index (Valic et al., 1997).

#### 23 **4.10.2.1.15. Tobacco smoking**

24 Individuals who smoke tobacco may be at increased risk of the health effects from TCE  
25 exposure. One study examining those living in an area with high TCE exposure found an  
26 increasing trend of risk ( $p = 0.008$ ) for renal cell carcinoma among smokers, with the highest OR  
27 among those with  $\geq 40$  pack-years (OR = 3.27, 95% CI: 1.48–7.19) (Charbotel et al., 2006). It  
28 has been shown that renal cell carcinoma is independently associated with smoking in a dose-  
29 response manner (Yuan et al., 1998), particularly in men (Benichou et al., 1998). While  
30 Charbotel et al. (2006) adjusted for smoking effects in analyses examining TCE exposure and  
31 renal cell carcinoma, this study provides no information on potential effect modification of TCE  
32 exposure by smoking.

1 A number of factors correlated to smoking (e.g., socioeconomic status, diet, alcohol  
2 consumption) may positively confound results if greater smoking rates were over-represented, as  
3 observed in an occupational cohort exposed to TCE (Raaschou-Nielsen et al., 2003). Absence of  
4 smoking information, on the other hand, could introduce a negative bias. In a drinking water  
5 study with exposures to TCE and perchlorate, Morgan and Cassidy (2002) noted the relatively  
6 high education high income levels and high access to health care of subjects in this study  
7 compared to the averages for the county as a whole likely leads to a lower smoking rate.  
8

#### 4.10.2.1.16. Nutritional status

9 Malnutrition may also increase susceptibility to TCE. Bioavailability of TCE after oral  
10 and intravenous exposure increased with fasting from approximately 63% in nonfasted rats to  
11 greater than 90% in fasted rats, with blood levels in fasted rats were elevated two- to threefold,  
12 and increased half-life in the blood of fasted rats (D'Souza et al., 1985). Food deprivation (Sato  
13 and Nakajima, 1985) and carbohydrate restriction (Nakajima et al., 1982; Sato and Nakajima,  
14 1985) enhanced metabolism of TCE in male rats, but this was not observed for dietary changes  
15 in protein or fat levels (Nakajima et al., 1982).

16 Vitamin intake may also alter susceptibility to TCE. An in vitro study of cultured normal  
17 human epidermal keratinocyte demonstrated an increased lipid peroxidation in a dose-dependent  
18 manner after exposure to TCE, which were then attenuated by exposure to Vitamin E (Ding et  
19 al., 2006).  
20

#### 4.10.2.1.17. Physical activity

21 Increased inhalation during physical activity increases TCE concentrations in the alveoli  
22 when compared to inhalation in a resting state (Astrand, 1975). Studies have examined the time  
23 course of inhaled TCE and metabolites in blood and urine in individuals with different workloads  
24 (Astrand and Ovrum, 1976; Jakubowski and Wieczorek, 1988; Monster et al., 1976; Vesterberg  
25 and Astrand, 1976; Vesterberg et al., 1976). These studies demonstrate that an increase in  
26 pulmonary ventilation increases the amount of TCE taken up during exposure (Astrand and  
27 Ovrum, 1976; Jakubowski and Wieczorek, 1988; Monster et al., 1976; Sato, 1993).

28 The Rocketdyne aerospace cohort exposed to TCE (and other chemicals) found a  
29 protective effect with high physical activity, but only after controlling for TCE exposure and  
30 socioeconomic status (OR = 0.55, 95% CI: 0.32–0.95, *p* trend = 0.04) (Krishnadasan et al.,

1 2008). In general, physical activity may provide a protective effect for prostate cancer (Wigle et  
2 al., 2008) (see Section 4.8.3.1.1).

#### 4.10.2.1.18. Socioeconomic status

3 Socioeconomic status (SES) can be an indicator for a number of coexposures, such as  
4 increased tobacco smoking, poor diet, education, income, and health care access, which may play  
5 a role in the results observed in the health effects of TCE exposure (Morgan and Cassady, 2002).

6 Children's exposure to TCE was measured in a low SES community, as characterized by  
7 income, educational level, and receipt of free or reduced cost school meals (Sexton et al., 2005);  
8 however, this study did not compare data to a higher SES community, nor examine health  
9 effects.

10 An elevated risk of NHL and esophagus/adenocarcinoma after exposure to TCE was  
11 observed for blue-collar workers compared to white collar workers and workers with unknown  
12 SES (Raaschou-Nielsen et al., 2003). Authors speculate that these results could be confounded  
13 due to other related factors to SES such as smoking.

14

#### 4.10.2.1.19. Mixtures

15 TCE exposure often occurs concurrently with other chemical substances. In general, the  
16 effects of exposures to multiple chemicals is considered by EPA in the *Framework for*  
17 *Cumulative Risk Assessment* (U.S. EPA, 2003a). A summary of the interactive effects of TCE  
18 and other chemical coexposures is addressed in Caldwell et al. (2008a) and in Chapter 10 of the  
19 National Research Council's report *Assessing the Human Health Risks of Trichloroethylene: Key*  
20 *Scientific Issues* (NRC, 2006).

21 Section 2 discusses that other parent compounds produce similar metabolites to TCE (see  
22 Table 2-1) or have similar properties or industrial uses (see Tables 2-3 and 2-14). The metabolic  
23 pathway of TCE is discussed in Section 3.3; due its metabolism into multiple compounds,  
24 exposure to TCE itself can be considered as exposure to a mixture (NRC, 2006). Many of the  
25 studies discussed above in Section 4 demonstrate that exposure to TCE and other chemical  
26 substances often occur together in both occupational and nonoccupational settings.

27 Coexposures to other solvents may induce or saturate toxicokinetic pathways, altering the  
28 way in which TCE is metabolized and cleared from the body. The limited data summarized by  
29 the ATSDR in its interaction profile on TCE, 1,1,1-trichloroethane, 1,1-dichloroethane, and  
30 tetrachloroethylene suggest that additive joint action is plausible (ATSDR, 2004b; Pohl et al.,  
31 2003). Joint exposure to TCE and the fungicide fenarimol has been shown to alter TCE

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 metabolism and genetic expression in mice (Hrelia et al., 1994). Joint exposure to TCE, benzene  
2 and methyl mercury has been shown to induce genetic expression in the liver and the kidney of  
3 rats (Hendriksen et al., 2007). Metabolic competition was also observed for TCE and various  
4 agents in another study by Jakobson et al. (1986).

5 PBPK models have been developed demonstrating the interaction between  
6 1,1-dichloroethylene and TCE (Andersen et al., 1987b) and the interaction between TCE,  
7 tetrachloroethylene, and 1,1,1-trichloroethane in rats (Dobrev et al., 2001) and humans (Dobrev  
8 et al., 2002). Other PBPK models also showed metabolic inhibition at higher doses for TCE and  
9 toluene (Thrall and Poet, 2000), and for TCE and chloroform (Isaacs et al., 2004). Another  
10 PBPK model of TCE and multiple VOCs showed metabolic inhibition and induction when  
11 exposure occurs concurrently (Haddad et al., 2000).

#### 4.10.3. Uncertainty of Database and Research Needs for Susceptible Populations

13 There is some evidence that certain populations may be more susceptible to exposure to  
14 TCE. These populations include early and later lifestages, gender, genetic polymorphisms,  
15 race/ethnicity, preexisting health status, and lifestyle factors and nutrition status. In general, this  
16 database would be improved by future epidemiologic and toxicological studies of TCE exposure  
17 that provide data on effect modification, including the factors discussed here.

18 Although the toxicokinetic variability has been characterized by population PBPK  
19 modeling (see Section 3.5), the available data are limited due to the relative small numbers of  
20 individuals ( $n < 100$ ), their all being adults, and the fact that subjects were selected nonrandomly  
21 (healthy human volunteers).

22 Although there is more information on early life exposure to TCE than on other  
23 potentially susceptible populations, there remain a number of uncertainties and data gaps  
24 regarding children's susceptibility. Improved PBPK modeling for using childhood parameters  
25 early lifestages as recommended by the NRC (2006), and validation of these models, will aid in  
26 determining how variations in metabolic enzymes affect TCE metabolism. In particular, the  
27 NRC states that it is prudent to assume children need greater protection than adults—unless  
28 sufficient data are available to justify otherwise (NRC, 2006).

29 More studies specifically designed to evaluate effects in early and later lifestages are  
30 needed in order to more fully characterize potential life stage-related TCE toxicity. Because the  
31 neurological effects of TCE constitute the most sensitive endpoints of concern for noncancer  
32 effects, it is quite likely that the early lifestages may be more susceptible to these outcomes than  
33 are adults. Lifestage-specific neurotoxic effects, particularly in the developing fetus, need

1 further evaluation. It is important to consider the use of age-appropriate testing for assessment of  
2 these and other outcomes, both for cancer and noncancer outcomes. Data specific to the  
3 carcinogenic effects of TCE exposure during the critical periods of development of experimental  
4 animals and humans also are sparse.

5         There is a need to better characterize the implications of TCE exposures to susceptible  
6 populations. There is suggestive evidence that there may be greater susceptibility for exposures  
7 to the elderly. Gender and race/ethnic differences in susceptibility are likely due to variation in  
8 physiology and exposure, and genetic variation likely has an effect on the toxicokinetics of TCE.  
9 In particular, the relationship between genetic variation and generalized hypersensitivity skin  
10 diseases is relevant for future study (see Sections 4.6.1.1.2 and 4.10.2.2). Diminished health  
11 status (e.g., impaired kidney liver or kidney), alcohol consumption, tobacco smoking, and  
12 nutritional status will likely affect an individual’s ability to metabolize TCE. In addition, further  
13 evaluation of the effects due to coexposures to other compounds with similar or different MOAs  
14 need to be evaluated. Future research should better characterize possible susceptibility for  
15 certain lifestages or populations.

**4.11. HAZARD CHARACTERIZATION**

**4.11.1. Characterization of Noncancer Effects**

**4.11.1.1.1. Neurotoxicity**

16         Both human and animal studies have associated TCE exposure with effects on several  
17 neurological domains. The strongest neurological evidence of hazard in humans is for changes  
18 in trigeminal nerve function or morphology and impairment of vestibular function. Fewer and  
19 more limited evidence exists in humans on delayed motor function, and changes in auditory,  
20 visual, and cognitive function or performance. Acute and subchronic animal studies show  
21 morphological changes in the trigeminal nerve, disruption of the peripheral auditory system  
22 leading to permanent function impairments and histopathology, changes in visual evoked  
23 responses to patterns or flash stimulus, and neurochemical and molecular changes. Additional  
24 acute studies reported structural or functional changes in hippocampus, such as decreased  
25 myelination or decreased excitability of hippocampal CA1 neurons, although the relationship of  
26 these effects to overall cognitive function is not established. Some evidence exists for motor-  
27 related changes in rats/mice exposed acutely/subchronically to TCE, but these effects have not  
28 been reported consistently across all studies.

1 Epidemiologic evidence supports a relationship between TCE exposure and trigeminal  
2 nerve function changes, with multiple studies in different populations reporting abnormalities in  
3 trigeminal nerve function in association with TCE exposure Ruitjen et al., 2001;(Barret et al.,  
4 1982; Barret et al., 1984; Barret et al., 1987; Feldman et al., 1988; Feldman et al., 1992; Kilburn,  
5 2002b; Kilburn and Warshaw, 1993a; Mhiri et al., 2004). Of these, two well conducted  
6 occupational cohort studies, each including more than 100 TCE-exposed workers without  
7 apparent confounding from multiple solvent exposures, additionally reported statistically  
8 significant dose-response trends based on ambient TCE concentrations, duration of exposure,  
9 and/or urinary concentrations of the TCE metabolite TCA (Barret et al., 1984; 1987). Limited  
10 additional support is provided by a positive relationship between prevalence of abnormal  
11 trigeminal nerve or sensory function and cumulative exposure to TCE (most subjects) or CFC-  
12 113 (<25% of subjects) (Rasmussen et al., 1993d). Test for linear trend in this study was not  
13 statistically significant and may reflect exposure misclassification since some subjects included  
14 in this study did not have TCE exposure. The lack of association between TCE exposure and  
15 overall nerve function in three small studies (trigeminal: (El Ghawabi et al., 1973); ulnar and  
16 medial: (Triebig et al., 1983; 1982)) does not provide substantial evidence against a causal  
17 relationship between TCE exposure and trigeminal nerve impairment because of limitations in  
18 statistical power, the possibility of exposure misclassification, and differences in measurement  
19 methods. Laboratory animal studies have also shown TCE-induced changes in the trigeminal  
20 nerve. Although one study reported no significant changes in trigeminal somatosensory evoked  
21 potential in rats exposed to TCE for 13 weeks (Albee et al., 2006), there is evidence of  
22 morphological changes in the trigeminal nerve following short-term exposures in rats (Barret et  
23 al., 1992; 1991).

24 Human chamber, occupational, geographic based/drinking water, and laboratory animal  
25 studies clearly established TCE exposure causes transient impairment of vestibular function.  
26 Subjective symptoms such as headaches, dizziness, and nausea resulting from occupational  
27 (Grandjean et al., 1955; Liu et al., 1988; Rasmussen and Sabroe, 1986; Smith, 1970),  
28 environmental (Hirsch et al., 1996), or chamber exposures (Smith, 1970; Stewart et al., 1970)  
29 have been reported extensively. A few laboratory animal studies have investigated vestibular  
30 function, either by promoting nystagmus or by evaluating balance (Niklasson et al., 1993; Tham  
31 et al., 1984; 1979; Umezu et al., 1997).

32 In addition, mood disturbances have been reported in a number of studies, although these  
33 effects also tend to be subjective and difficult to quantify (Gash et al., 2008; Kilburn, 2002a,  
34 2002b; Kilburn and Warshaw, 1993a; McCunney, 1988; Mitchell and Parsons-Smith, 1969;  
35 Rasmussen and Sabroe, 1986; Troster and Ruff, 1990), and a few studies have reported no



1 effects from TCE on mood (Reif et al., 2003; Triebig et al., 1976; Triebig et al., 1977a). Few  
2 comparable mood studies are available in laboratory animals, although both Moser et al. (2003)  
3 and Albee et al. (2006) report increases in handling reactivity among rats exposed to TCE.  
4 Finally, significantly increased number of sleep hours was reported by Arito et al. (1994) in rats  
5 exposed via inhalation to 50–300-ppm TCE for 8 hours/day for 6 weeks.

6 Four epidemiologic studies of chronic exposure to TCE observed disruption of auditory  
7 function. One large occupational cohort study showed a statistically significant difference in  
8 auditory function with cumulative exposure to TCE or CFC-113 as compared to control groups  
9 after adjustment for possible confounders, as well as a positive relationship between auditory  
10 function and increasing cumulative exposure (Rasmussen et al., 1993b). Of the three studies  
11 based on populations from ATSDR’s TCE Subregistry from the National Exposure Registry,  
12 more limited than Rasmussen et al. (1993b) due to inferior exposure assessment, Burg et al.  
13 (1995) and Burg and Gist (1999) reported a higher prevalence of self-reported hearing  
14 impairments. The third study reported that auditory screening revealed abnormal middle ear  
15 function in children less than 10 years of age, although a dose-response relationship could not be  
16 established and other tests did not reveal differences in auditory function (ATSDR, 2003b).  
17 Further evidence for these effects is provided by numerous laboratory animal studies  
18 demonstrating that high dose subacute and subchronic TCE exposure in rats disrupts the auditory  
19 system leading to permanent functional impairments and histopathology.

20 Studies in humans exposed under a variety of conditions, both acutely and chronically,  
21 report impaired visual functions such as color discrimination, visuospatial learning tasks, and  
22 visual depth perception in subjects with TCE exposure. Abnormalities in visual depth perception  
23 were observed with a high acute exposure to TCE under controlled conditions (Vernon and  
24 Ferguson, 1969). Studies of lower TCE exposure concentrations also observed visuofunction  
25 effects. One occupational study (Rasmussen et al., 1993b) reported a statistically significant  
26 positive relationship between cumulative exposure to TCE or CFC-113 and visual gestalts  
27 learning and retention among Danish degreasers. Two studies of populations living in a  
28 community with drinking water containing TCE and other solvents furthermore suggested  
29 changes in visual function (Kilburn, 2002a)(Reif et al., 2003). These studies used more direct  
30 measures of visual function as compared to Rasmussen et al. (1993b), but their exposure  
31 assessment is more limited because TCE exposure is not assigned to individual subjects  
32 (Kilburn 2002a), or because there are questions regarding control selection (Kilburn 2002a) and  
33 exposure to several solvents (Kilburn, 2002a)(Reif et al., 2003).

34 Additional evidence of effects of TCE exposure on visual function is provided by a  
35 number of laboratory animal studies demonstrating that acute or subchronic TCE exposure

1 causes changes in visual evoked responses to patterns or flash stimulus (Boyes et al., 2005)(Blain  
2 et al., 1994; Boyes et al., 2003). Animal studies have also reported that the degree of some  
3 effects is correlated with simultaneous brain TCE concentrations (Boyes et al., 2003; Boyes et  
4 al., 2005) and that, after a recovery period, visual effects return to control levels (Blain et al.,  
5 1994; Rebert et al., 1991). Overall, the human and laboratory animal data together suggest that  
6 TCE exposure can cause impairment of visual function, and some animal studies suggest that  
7 some of these effects may be reversible with termination of exposure.

8 Studies of human subjects exposed to TCE either acutely in chamber studies or  
9 chronically in occupational settings have observed deficits in cognition. Five chamber studies  
10 reported statistically significant deficits in cognitive performance measures or outcome measures  
11 suggestive of cognitive effects (Gamberale et al., 1976; Stewart et al., 1970; Triebig et al., 1976;  
12 Triebig et al., 1977a). Danish degreasers with high cumulative exposure to TCE or CFC-113 had  
13 a high risk (OR = 13.7, 95% CI: 2.0–92.0) for psychoorganic syndrome characterized by  
14 cognitive impairment, personality changes, and reduced motivation, vigilance, and initiative  
15 compared to workers with low cumulative exposure. Studies of populations living in a  
16 community with contaminated groundwater also reported cognitive impairments (Kilburn,  
17 2002b; Kilburn and Warshaw, 1993a), although these studies carry less weight in the analysis  
18 because TCE exposure is not assigned to individual subjects and their methodological design is  
19 weaker.

20 Laboratory studies provide some additional evidence for the potential for TCE to affect  
21 cognition, although the predominant effect reported has been changes in the time needed to  
22 complete a task, rather than impairment of actual learning and memory function (Kishi et al.,  
23 1993; Kulig, 1987; Umezu et al., 1997). In addition, in laboratory animals, it can be difficult to  
24 distinguish cognitive changes from motor-related changes. However, several studies have  
25 reported structural or functional changes in the hippocampus, such as decreased myelination  
26 (Isaacson and Taylor, 1989; Isaacson et al., 1990) or decreased excitability of hippocampal CA1  
27 neurons (Ohta et al., 2001), although the relationship of these effects to overall cognitive  
28 function is not established.

29 Two studies of TCE exposure, one chamber study of acute exposure duration and one  
30 occupational study of chronic duration, reported changes in psychomotor responses. The  
31 chamber study of Gamberale et al. (1976) reported a dose-related decrease in performance in a  
32 choice reaction time test in healthy volunteers exposed to 100 and 200-ppm TCE for 70 minutes  
33 as compared to the same subjects without exposure. Rasmussen et al. (1993d) reported a  
34 statistically significant association with cumulative exposure to TCE or CFC-113 and  
35 dyscoordination trend among Danish degreasers. Observations in a third study (Gun et al., 1978)

1 are difficult to judge given the author's lack of statistical treatment of data. In addition, Gash et  
2 al. (2008) reported that 14 out of 30 TCE-exposed workers exhibited significantly slower fine  
3 motor hand movements as measured through a movement analysis panel test. Studies of  
4 populations living in communities with TCE and other solvents detected in groundwater supplies  
5 reported significant delays in simple and choice reaction times in individuals exposed to TCE in  
6 contaminated groundwater as compared to referent groups (Kilburn, 2002b; Kilburn and  
7 Warshaw, 1993a) (Kilburn and Thornton, 1996). Observations in these studies are more  
8 uncertain given questions of the representativeness of the referent population, lack of exposure  
9 assessment to individual study subjects, and inability to control for possible confounders  
10 including alcohol consumption and motivation. Finally, in a presentation of 2 case reports,  
11 decrements in motor skills as measured by the grooved pegboard and finger tapping tests were  
12 observed (Troster and Ruff, 1990).

13 Laboratory animal studies of acute or subchronic exposure to TCE observed psychomotor  
14 effects, such as loss of righting reflex (Shih et al., 2001; Umezu et al., 1997) and decrements in  
15 activity, sensory-motor function, and neuromuscular function (Kishi et al., 1993; Moser et al.,  
16 1995; 2003). However, two studies also noted an absence of significant changes in some  
17 measures of psychomotor function (Albee et al., 2006; Kulig, 1987). In addition, less consistent  
18 results have been reported with respect to locomotor activity in rodents. Some studies have  
19 reported increased locomotor activity after an acute i.p. dosage (Wolff and Siegmund, 1978) or  
20 decreased activity after acute or short term oral gavage dosing (Moser et al., 1995; 2003). No  
21 change in activity was observed following exposure through drinking water (Waseem et al.,  
22 2001), inhalation (Kulig, 1987) or orally during the neurodevelopment period (Fredriksson et al.,  
23 1993).

24 Several neurochemical and molecular changes have been reported in laboratory  
25 investigations of TCE toxicity. Kjellstrand et al. (1987) reported inhibition of sciatic nerve  
26 regeneration in mice and rats exposed continuously to 150-ppm TCE via inhalation for 24 days.  
27 Two studies have reported changes in GABAergic and glutamatergic neurons in terms of GABA  
28 or glutamate uptake (Briving et al., 1986) or response to GABAergic antagonistic drugs (Shih et  
29 al., 2001) as a result of TCE exposure, with the Briving et al. (1986) conducted at 50 ppm for  
30 12 months. Although the functional consequences of these changes is unclear, Tham et al.  
31 (1984; 1979) described central vestibular system impairments as a result of TCE exposure that  
32 may be related to altered GABAergic function. In addition, several in vitro studies have  
33 demonstrated that TCE exposure alters the function of inhibitory ion channels such as receptors  
34 for GABA<sub>A</sub> glycine, and serotonin (Beckstead et al., 2000; Krasowski and Harrison, 2000;  
35 Lopreato et al., 2003) or of voltage-sensitive calcium channels (Shafer et al., 2005).

#### 4.11.1.1.2. Kidney Toxicity

2           There are few human data pertaining to TCE-related noncancer kidney toxicity.  
3           Observation of elevated excretion of urinary proteins in the available studies (Bolt et al., 2004;  
4           Brüning et al., 1999a; Brüning et al., 1999b; Green et al., 2004; Rasmussen et al., 1993d)  
5           indicates the occurrence of a toxic insult among TCE-exposed subjects compared to unexposed  
6           controls. Two studies are of subjects with previously diagnosed kidney cancer (Bolt et al., 2004;  
7           Brüning et al., 1999a), while subjects in the other studies are disease free. Urinary proteins are  
8           considered nonspecific markers of nephrotoxicity and include  $\alpha$ 1-microglobulin, albumin, and  
9           NAG (Lybarger et al., 1999; Price et al., 1996; 1999). Four studies measure  $\alpha$ 1-microglobulin  
10          with elevated excretion observed in the German studies (Bolt et al., 2004; Brüning et al., 1999a;  
11          Brüning et al., 1999b) but not Green et al. (2004). However, Rasmussen et al. (1993d) reported a  
12          positive relationship between increasing urinary NAG, another nonspecific marker of tubular  
13          toxicity, and increasing exposure duration; and Green et al. (2004) found statistically significant  
14          group mean differences in NAG. Observations in Green et al. (2004) provide evidence of  
15          tubular damage among workers exposed to trichloroethylene at current occupational levels.  
16          Elevated excretion of NAG has also been observed with acute TCE poisoning (Carrieri et al.,  
17          2007). Some support for TCE nephrotoxicity in humans is provided by a study of end-stage  
18          renal disease in a cohort of workers at Hill Air Force Base (Radican et al., 2006), although  
19          subjects in this study were exposed to hydrocarbons, JP-4 gasoline, and solvents in addition to  
20          TCE, including 1,1,1-trichloroethane, and a second reporting a twofold elevated risk for  
21          progression of glomerulonephritis to ESRD with TCE exposure (Jacob et al., 2007).

22          Laboratory animal and in vitro data provide additional support for TCE nephrotoxicity.  
23          Multiple studies with both gavage and inhalation exposure show that TCE causes renal toxicity  
24          in the form of cytomegaly and karyomegaly of the renal tubules in male and female rats and  
25          mice (summarized in Section 4.4.4). Further studies with TCE metabolites have demonstrated a  
26          potential role for DCVC, TCOH, and TCA in TCE-induced nephrotoxicity. Of these, available  
27          data suggest that DCVC induced renal effects most like those of TCE and is formed in sufficient  
28          amounts following TCE exposure to account for these effects. TCE or DCVC have also been  
29          shown to be cytotoxic to primary cultures of rat and human renal tubular cells (Cummings et al.,  
30          2000a; Cummings and Lash, 2000; Cummings et al., 2000b).

31          Overall, multiple lines of evidence support the conclusion that TCE causes nephrotoxicity  
32          in the form of tubular toxicity, mediated predominantly through the TCE GSH conjugation  
33          product DCVC.

#### 4.11.1.1.3. Liver Toxicity

2 Few studies on liver toxicity and TCE exposure are found in humans. Of these, three  
3 studies reported significant changes in serum liver function tests, widely used in clinical settings  
4 in part to identify patients with liver disease, in metal degreasers whose TCE exposure was  
5 assessed using urinary trichloro-compounds as a biomarker (Nagaya et al., 1993; Rasmussen et  
6 al., 1993b; Xu et al., 2009). Two additional studies reported plasma or serum bile acid changes  
7 (Driscoll et al., 1992; Neghab et al., 1997). One study of subjects from the TCE subregistry of  
8 ATSDR's National Exposure Registry is suggestive of liver disorders but limitations preclude  
9 inferences whether TCE caused these conditions is not possible given the study's limitations  
10 (Davis et al., 2005). Furthermore, a number of case reports exist of liver toxicity including  
11 hepatitis accompanying immune-related generalized skin diseases described as a variation of  
12 erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis patients, and  
13 hypersensitivity syndrome (Kamijima et al., 2007) in addition to jaundice, hepatomegaly,  
14 hepatosplenomegaly, and liver failure TCE-exposed workers (Huang et al., 2002; Thiele et al.,  
15 1982). Cohort studies have examined cirrhosis mortality and either TCE exposure (ATSDR,  
16 2004a; Blair et al., 1989; 1998; Boice et al., 1999; Boice et al., 2006b; Garabrant et al., 1988;  
17 Morgan et al., 1998; Radican et al., 2008; Ritz, 1999a) or solvent exposure (Leigh and Jiang,  
18 1993), but are greatly limited by their use of death certificates where there is a high degree (up to  
19 50%) of underreporting (Blake et al., 1988), so these null findings do not rule out an effect of  
20 TCE on cirrhosis. Overall, while there some evidence exists of liver toxicity as assessed from  
21 liver function tests, the data are inadequate for making conclusions regarding causality.

22 In laboratory animals, TCE exposure is associated with a wide array of hepatotoxic  
23 endpoints. Like humans, laboratory animals exposed to TCE have been observed to have  
24 increased serum bile acids (Bai et al., 1992a; Neghab et al., 1997), although the toxicological  
25 importance of this effect is unclear. Most other effects in laboratory animals have not been  
26 studied in humans, but nonetheless provide evidence that TCE exposure leads to hepatotoxicity.  
27 These effects include increased liver weight, small transient increases in DNA synthesis,  
28 cytomegaly in the form of "swollen" or enlarged hepatocytes, increased nuclear size probably  
29 reflecting polyploidization, and proliferation of peroxisomes. Liver weight increases  
30 proportional to TCE dose are consistently reported across numerous studies and appear to be  
31 accompanied by periportal hepatocellular hypertrophy (Berman et al., 1995; Buben and  
32 O'Flaherty, 1985; Dees and Travis, 1993; Elcombe et al., 1985; Goel et al., 1992; Goldsworthy  
33 and Popp, 1987; Kjellstrand et al., 1983a; Kjellstrand et al., 1983b; Kjellstrand et al., 1981b;

1 Laughter et al., 2004; Melnick et al., 1987; Merrick et al., 1989; Nakajima et al., 2000; Nunes et  
2 al., 2001; Tao et al., 2000a; Tucker et al., 1982). There is also evidence of increased DNA  
3 synthesis in a small portion of hepatocytes at around 10 days in vivo exposure (Channel et al.,  
4 1998; Dees and Travis, 1993; Elcombe et al., 1985; Mirsalis et al., 1989). The lack of  
5 correlation of hepatocellular mitotic figures with whole liver DNA synthesis or DNA synthesis  
6 observed in individual hepatocytes (Dees and Travis, 1993; Elcombe et al., 1985) supports the  
7 conclusions that cellular proliferation is not the predominant cause of increased DNA synthesis  
8 and that nonparenchymal cells may also contribute to such synthesis. Indeed, nonparenchymal  
9 cell activation or proliferation has been noted in several studies (Goel et al., 1992; Kjellstrand et  
10 al., 1983a). Moreover, the histological descriptions of TCE-exposed livers are consistent with  
11 and, in some cases, specifically note increased polyploidy (Buben and O'Flaherty, 1985).  
12 Interestingly, changes in TCE-induced hepatocellular ploidy, as indicated by histological  
13 changes in nuclei, have been noted to remain after the cessation of exposure (Kjellstrand et al.,  
14 1983a). In regard to apoptosis, TCE has been reported either to have no effect or to cause a  
15 slight increase at high doses (Channel et al., 1998; Dees and Travis, 1993). Some studies have  
16 also noted effects from dosing vehicle alone (such as corn oil, in particular) not only on liver  
17 pathology, but also on DNA synthesis (Channel et al., 1998; Merrick et al., 1989). Available  
18 data also suggest that TCE does not induce substantial cytotoxicity, necrosis, or regenerative  
19 hyperplasia, as only isolated, focal necroses and mild to moderate changes in serum and liver  
20 enzyme toxicity markers having been reported (Channel et al., 1998; Dees and Travis, 1993;  
21 Elcombe et al., 1985). Data on peroxisome proliferation, along with increases in a number of  
22 associated biochemical markers, show effects in both mice and rats (Channel et al., 1998;  
23 Elcombe et al., 1985; Goldsworthy and Popp, 1987). These effects are consistently observed  
24 across rodent species and strains, although the degree of response at a given mg/kg-day dose  
25 appears to be highly variability across strains, with mice on average appearing to be more  
26 sensitive.

27 While it is likely that oxidative metabolism is necessary for TCE-induced effects in the  
28 liver, the specific metabolite or metabolites responsible is less clear. TCE, TCA, and DCA  
29 exposures have all been associated with induction of changes in liver weight, DNA synthesis,  
30 and peroxisomal enzymes. The available data strongly support TCA *not* being the sole or  
31 predominant active moiety for TCE-induced liver effects, particularly with respect to  
32 hepatomegaly. In particular, TCE and TCA dose-response relationships are quantitatively  
33 inconsistent, for TCE leads to greater increases in liver/body weight ratios that expected from  
34 predicted rates of TCA production (see analysis in Section 4.5.6.2.1). In fact, above a certain  
35 dose of TCE, liver/body weight ratios are greater than that observed under any conditions studied

1 so far for TCA. Histological changes and effects on DNA synthesis are generally consistent with  
2 contributions from either TCA or DCA, with a degree of polyploidization, rather than cell  
3 proliferation, likely to be significant for TCE, TCA, and DCA.

4 Overall, TCE, likely through its oxidative metabolites, clearly leads to liver toxicity in  
5 laboratory animals, with mice appearing to be more sensitive than other laboratory animal  
6 species, but there is only limited epidemiologic evidence of hepatotoxicity being associated with  
7 TCE exposure.  
8

#### 4.11.1.1.4. Immunotoxicity

9 Studies in humans provide evidence of associations between TCE exposure and a number  
10 of immunotoxicological endpoints. The relation between systemic autoimmune diseases, such as  
11 scleroderma, and occupational exposure to TCE has been reported in several recent studies. A  
12 meta-analysis of scleroderma studies (Diot et al., 2002; Garabrant et al., 2003; Nietert et al.,  
13 1998) conducted by the EPA resulted in a statistically significant combined odds ratio for any  
14 exposure in men (OR: 2.5, 95% CI: 1.1, 5.4), with a lower relative risk seen in women (OR: 1.2,  
15 95% CI: 0.58, 2.6). The incidence of systemic sclerosis among men is very low (approximately  
16 1 per 100,000 per year), and is approximately 10 times lower than the rate seen in women  
17 (Cooper and Stroehla, 2003). Thus, the human data at this time do not allow determination of  
18 whether the difference in effect estimates between men and women reflects the relatively low  
19 background risk of scleroderma in men, gender-related differences in exposure prevalence or in  
20 the reliability of exposure assessment (Messing et al., 2003), a gender-related difference in  
21 susceptibility to the effects of TCE, or chance. Changes in levels of inflammatory cytokines  
22 were reported in an occupational study of degreasers exposed to TCE (Iavicoli et al., 2005) and a  
23 study of infants exposed to TCE via indoor air (Lehmann et al., 2001; 2002).

24 Experimental studies provide additional support for these effects. Numerous studies have  
25 demonstrated accelerated autoimmune responses in autoimmune-prone mice (Blossom et al.,  
26 2007; Blossom et al., 2004; Cai et al., 2008; Griffin et al., 2000a; Griffin et al., 2000b). With  
27 shorter exposure periods, effects include changes in cytokine levels similar to those reported in  
28 human studies. More severe effects, including autoimmune hepatitis, inflammatory skin lesions,  
29 and alopecia, were manifest at longer exposure periods, and interestingly, these effects differ  
30 somewhat from the “normal” expression in these mice. Immunotoxic effects, including increases  
31 in anti-dsDNA antibodies in adult animals, decreased thymus weights, and decreased plaque  
32 forming cell response with prenatal and neonatal exposure, have been also reported in B6C3F1  
33 mice, which do not have a known particular susceptibility to autoimmune disease (Gilkeson et

1 al., 2004; Keil et al., 2009; Peden-Adams et al., 2006). Recent mechanistic studies have focused  
2 on the roles of various measures of oxidative stress in the induction of these effects by TCE  
3 (Wang et al., 2007b; Wang et al., 2008).

4 There have been a large number of case reports of a severe hypersensitivity skin disorder,  
5 distinct from contact dermatitis and often accompanied by hepatitis, associated with occupational  
6 exposure to TCE, with prevalences as high as 13% of workers in the same location (Kamijima et  
7 al., 2007; 2008). Evidence of a treatment-related increase in delayed hypersensitivity response  
8 accompanied by hepatic damage has been observed in guinea pigs following intradermal  
9 injection (Tang et al., 2002; Tang et al., 2008), and hypersensitivity response was also seen in  
10 mice exposed via drinking water pre- and postnatally (GD 0 through to 8 weeks of age) (Peden-  
11 Adams et al., 2006).

12 Human data pertaining to TCE-related immunosuppression resulting in an increased risk  
13 of infectious diseases is limited to the report of an association between reported history of  
14 bacteria of viral infections in Woburn, MA (Lagakos et al., 1986). Evidence of localized  
15 immunosuppression, as measured by pulmonary response to bacterial challenge (i.e., risk of  
16 Streptococcal pneumonia-related mortality and clearance of Klebsiella bacteria) was seen in an  
17 acute exposure study in CD-1 mice (Aranyi et al., 1986). A 4-week inhalation exposure in  
18 Sprague-Dawley rats reported a decrease in plaque forming cell response at exposures of  
19 1,000 ppm (Woolhiser et al., 2006).

20 Overall, the human and animal studies of TCE and immune-related effects provide strong  
21 evidence for a role of TCE in autoimmune disease and in a specific type of generalized  
22 hypersensitivity syndrome, while there are less data pertaining to immunosuppressive effects.  
23

#### 4.11.1.1.5. Respiratory Tract Toxicity

24 There are very limited human data on pulmonary toxicity and TCE exposure. Two recent  
25 reports of a study of gun manufacturing workers reported asthma-related symptoms and lung  
26 function decrements associated with solvent exposure (Cakmak et al., 2004; Saygun et al., 2007),  
27 but these studies are limited by multiple solvent exposures and the significant effect of smoking  
28 on pulmonary function. Laboratory studies in mice and rats have shown toxicity in the bronchial  
29 epithelium, primarily in Clara cells, following acute exposures to TCE by inhalation (see  
30 Section 4.7.2.1.1). A few studies of longer duration have reported more generalized toxicity,  
31 such as pulmonary fibrosis 90 days after a single 2,000 mg/kg i.p. dose in mice and pulmonary  
32 vasculitis after 13-week oral gavage exposures to 2,000 mg/kg-day in rats (Forkert and Forkert,  
33 1994; NTP, 1990). However, respiratory tract effects were not reported in other longer-term



1 studies. Acute pulmonary toxicity appears to be dependent on oxidative metabolism, although  
2 the particular active moiety is not known. While earlier studies implicated chloral produced in  
3 situ by CYP enzymes in respiratory tract tissue was responsible for toxicity (reviewed in Green,  
4 2000), the evidence is inconsistent, and several other possibilities are viable. First, substantial  
5 “accumulation” of chloral is unlikely, as it is likely either to be rapidly converted to TCOH in  
6 respiratory tract tissue or to diffuse rapidly into blood and be converted to TCOH in erythrocytes  
7 or the liver. Conversely, a role for systemically produced oxidative metabolites cannot be  
8 discounted, as CH and TCOH in blood have both been reported following inhalation dosing in  
9 mice. In addition, a recent study reported dichloroacetyl chloride protein adducts in the lungs of  
10 mice to which TCE was administered by i.p. injection, suggesting dichloroacetyl chloride, which  
11 is not believed to be derived from chloral, may also contribute to TCE respiratory toxicity.  
12 Although humans appear to have lower overall capacity for enzymatic oxidation in the lung  
13 relative to mice, CYP enzymes do reside in human respiratory tract tissue, suggesting that,  
14 qualitatively, the respiratory tract toxicity observed in rodents is biologically plausible in  
15 humans. However, quantitative estimates of differential sensitivity across species due to  
16 respiratory metabolism are highly uncertain due to limited data. Therefore, overall, data are  
17 suggestive of TCE causing respiratory tract toxicity, based primarily on short-term studies in  
18 mice and rats, and no data suggest that such hazards would be biologically precluded in humans.  
19

#### 4.11.1.1.6. Reproductive Toxicity

20 Reproductive toxicity related to TCE exposure has been evaluated in human and  
21 experimental animal studies for effects in males and females. Only a limited number of studies  
22 have examined whether TCE causes female reproductive toxicity. Epidemiologic studies have  
23 identified possible associations of TCE exposure with effects on female fertility (ATSDR, 2001;  
24 Sallmén et al., 1995) and with menstrual cycle disturbances (ATSDR, 2001; Bardodej and  
25 Vyskocil, 1956; Sagawa et al., 1973; Zielinski, 1973). Reduced in vitro oocyte fertilizability has  
26 been reported as a result of TCE exposure in rats (Berger and Horner, 2003; Wu and Berger,  
27 2007), but a number of other laboratory animal studies did not report adverse effects on female  
28 reproductive function (Cosby and Dukelow, 1992; George et al., 1985, 1986; Manson et al.,  
29 1984). Overall, there are inadequate data to conclude whether adverse effects on human female  
30 reproduction are caused by TCE.

31 By contrast, a number of human and laboratory animal studies suggest that TCE exposure  
32 has the potential for male reproductive toxicity. In particular, human studies have reported TCE  
33 exposure to be associated, in several cases statistically-significantly, with increased sperm

1 density and decreased sperm quality (Chia et al., 1996; Rasmussen et al., 1988), altered sexual  
2 drive or function (Bardodej and Vyskocil, 1956; El Ghawabi et al., 1973; Saihan et al., 1978), or  
3 altered serum endocrine levels (Chia et al., 1997; Goh et al., 1998). In addition, three studies  
4 that reported measures of fertility did not or could not report changes associated with TCE  
5 exposure (ATSDR, 2001; Forkert et al., 2003; Sallmen et al., 1998), although the statistical  
6 power of these studies is quite limited. Further evidence of similar effects is provided by several  
7 laboratory animal studies that reported effects on sperm (George et al., 1985; Kumar et al.,  
8 2001b; Kumar et al., 2000a; Kumar et al., 2000b; Land et al., 1981; Veeramachaneni et al.,  
9 2001), libido/copulatory behavior (George et al., 1986; Veeramachaneni et al., 2001; Zenick et  
10 al., 1984), and serum hormone levels (Kumar et al., 2000a; Veeramachaneni et al., 2001). As  
11 with the human database, some studies that assessed sperm measures did not report treatment-  
12 related alterations (Cosby and Dukelow, 1992; George et al., 1986; Xu et al., 2004; Zenick et al.,  
13 1984). Additional adverse effects on male reproduction have also been reported, including  
14 histopathological lesions in the testes or epididymides (Forkert et al., 2002; George et al., 1986;  
15 Kan et al., 2007; Kumar et al., 2001b; Kumar et al., 2000b) and altered in vitro sperm-oocyte  
16 binding or in vivo fertilization due to TCE or metabolites (DuTeaux et al., 2004b; Xu et al.,  
17 2004). While reduced fertility in rodents was only observed in one study (George et al., 1986),  
18 this is not surprising given the redundancy and efficiency of rodent reproductive capabilities.  
19 Furthermore, while George et al. (1986) proposed that the adverse male reproductive outcomes  
20 observed in rats were due to systemic toxicity, the database as a whole suggests that TCE does  
21 induce reproductive toxicity independent of systemic effects. Therefore, overall, the human and  
22 laboratory animal data together support the conclusion that TCE exposure poses a potential  
23 hazard to the male reproductive system.  
24

#### 4.11.1.1.7. Developmental Toxicity

25 The relationship between TCE exposure (direct or parental) and adverse developmental  
26 outcomes has been investigated in a number of epidemiologic and laboratory animal studies.  
27 Prenatal effects examined include death (spontaneous abortion, perinatal death, pre- or  
28 postimplantation loss, resorptions), decreased growth (low birth weight, small for gestational  
29 age, intrauterine growth restriction, decreased postnatal growth), and congenital malformations,  
30 in particular eye and cardiac defects. Postnatal developmental outcomes examined include  
31 growth and survival, developmental neurotoxicity, developmental immunotoxicity, and  
32 childhood cancers.

1 A few epidemiological studies have reported associations between parental exposure to  
2 TCE and spontaneous abortion or perinatal death (ATSDR, 2001; Taskinen et al., 1994;  
3 Windham et al., 1991), although other studies reported mixed or null findings (ATSDR, 2006b,  
4 2008; Bove, 1996; Bove et al., 1995; Goldberg et al., 1990; Lagakos et al., 1986; Lindbohm et  
5 al., 1990; Taskinen et al., 1989). Studies examining associations between TCE exposure and  
6 decreased birth weight or small for gestational age have reported small, often nonstatistically  
7 significant, increases in risk for these effects ATSDR, 2006b,(ATSDR, 2008; Windham et al.,  
8 1991). However, other studies observed mixed or no association (Bove, 1996; Bove et al., 1995;  
9 Lagakos et al., 1986; Rodenbeck et al., 2000). While comprising both occupational and  
10 environmental exposures, these studies are overall not highly informative due to their small  
11 numbers of cases and limited exposure characterization or to the fact that exposures to mixed  
12 solvents were involved. However, a number of laboratory animal studies show analogous effects  
13 of TCE exposure in rodents. In particular, pre- or postimplantation losses, increased resorptions,  
14 perinatal death, and decreased birth weight have been reported in multiple well-conducted  
15 studies in rats and mice (George et al., 1985, 1986; Healy et al., 1982; Kumar et al., 2000b;  
16 Narotsky and Kavlock, 1995; Narotsky et al., 1995). Interestingly, the rat studies reporting these  
17 effects used Fischer 344 or Wistar rats, while several other studies, all of which used  
18 Sprague-Dawley rats, reported no increased risk in these developmental measures (Carney et al.,  
19 2006; Hardin et al., 1981; Schwetz et al., 1975). Overall, based on weakly suggestive  
20 epidemiologic data and fairly consistent laboratory animal data, it can be concluded that TCE  
21 exposure poses a potential hazard for prenatal losses and decreased growth or birth weight of  
22 offspring.

23 Epidemiologic data provide some support for the possible relationship between maternal  
24 TCE exposure and birth defects in offspring, in particular cardiac defects. Other developmental  
25 outcomes observed in epidemiology and experimental animal studies include an increase in total  
26 birth defects (ATSDR, 2001; Flood, 1988), CNS defects (ATSDR, 2001; Bove, 1996; Bove et  
27 al., 1995; Lagakos et al., 1986), oral cleft defects (Bove, 1996; Bove et al., 1995; Lagakos et al.,  
28 1986; Lorente et al., 2000), eye/ear defects (Lagakos et al., 1986; Narotsky and Kavlock, 1995;  
29 Narotsky et al., 1995), kidney/urinary tract disorders (Lagakos et al., 1986), musculoskeletal  
30 birth anomalies (Lagakos et al., 1986), lung/respiratory tract disorders (Das and Scott, 1994;  
31 Lagakos et al., 1986), and skeletal defects (Healy et al., 1982). Occupational cohort studies,  
32 while not consistently reporting positive results, are generally limited by the small number of  
33 observed or expected cases of birth defects (Lorente et al., 2000; Taskinen et al., 1989; Tola et  
34 al., 1980).

1 While only one of the epidemiological studies specifically reported observations of eye  
2 anomalies (Lagakos et al., 1986), studies in rats have identified increases in the incidence of fetal  
3 eye defects following oral exposures during the period of organogenesis with TCE (Narotsky and  
4 Kavlock, 1995; Narotsky et al., 1995) or its oxidative metabolites DCA and TCA (Smith et al.,  
5 1989; Warren et al., 2006)(Smith et al., 1992). No other developmental or reproductive toxicity  
6 studies identified abnormalities of eye development following TCE exposures, which may have  
7 been related to the administered dose or other aspects of study design (e.g., level of detail applied  
8 to fetal ocular evaluation). Overall, the study evidence suggests a potential for the disruption of  
9 ocular development by exposure to TCE and its oxidative metabolites.

10 The epidemiological studies, while individually limited, as a whole show relatively  
11 consistent elevations, some of which were statistically significant, in the incidence of cardiac  
12 effects in TCE-exposed populations compared to reference groups (ATSDR, 2001, 2006a, 2008;  
13 Bove, 1996; Bove et al., 1995; Goldberg et al., 1990; Yauck et al., 2004). Interestingly,  
14 Goldberg et al. (1990) noted that the odds ratio for congenital heart disease in offspring declined  
15 from threefold to no difference as compared to controls after TCE-contaminated drinking water  
16 wells were closed, suggestive of a causal relationship. However, this study reported no  
17 significant differences in cardiac lesions between exposed and nonexposed groups (Goldberg et  
18 al., 1990). One additional community study reported that, among the 5 cases of cardiovascular  
19 anomalies, there was no significant association with TCE (Lagakos et al., 1986), but due to the  
20 small number of cases this does not support an absence of effect. In laboratory animal models,  
21 avian studies were the first to identify adverse effects of TCE exposure on cardiac development,  
22 and the initial findings have been confirmed multiple times (Boyer et al., 2000; Bross et al.,  
23 1983; Drake et al., 2006a; Drake et al., 2006b; Loeber et al., 1988; Mishima et al., 2006; Rufer et  
24 al., 2008). Additionally, administration of TCE and TCE metabolites TCA and DCA in maternal  
25 drinking water during gestation has been reported to induce cardiac malformations in rat fetuses  
26 (Johnson et al., 2005; Smith et al., 1989)(Dawson et al., 1993; Dawson et al., 1990b; Epstein et  
27 al., 1992; Johnson et al., 1998a; Johnson et al., 2003; Johnson et al., 1998b; Smith et al., 1992).  
28 However, it is notable that a number of other studies, several of which were well conducted, did  
29 not report induction of cardiac defects in rats or rabbits from TCE administered by inhalation  
30 (Carney et al., 2006; Dorfmueller et al., 1979; Hardin et al., 1981; Healy et al., 1982; Schwetz et  
31 al., 1975) or in rats and mice by gavage (Cosby and Dukelow, 1992; Fisher et al., 2001; Narotsky  
32 and Kavlock, 1995; Narotsky et al., 1995).

33 The potential importance of these effects warrants a more detailed discussion of possible  
34 explanations for the apparent inconsistencies in the laboratory animal studies. Many of the  
35 studies that did not identify cardiac anomalies used a traditional free-hand section technique on

1 fixed fetal specimens (Dorfmueller et al., 1979; Hardin et al., 1981; Healy et al., 1982; Schwetz  
2 et al., 1975). Detection of such anomalies can be enhanced through the use of a fresh dissection  
3 technique as described by Staples (1974) and Stuckhardt and Poppe (1984) and this was the  
4 technique used in the study by Dawson et al. (1990b) with further refinement of the technique  
5 used in the positive studies by Dawson et al. (1993) and Johnson et al. (2005)(2003). However,  
6 two studies that used the same or similar fresh dissection technique did not report cardiac  
7 anomalies (Carney et al., 2006; Fisher et al., 2001), although it has been suggested that  
8 differences in experimental design (e.g., inhalation versus gavage versus drinking water route of  
9 administration, exposure during organogenesis versus the entire gestational period, or varied  
10 dissection or evaluation procedures) may have been contributing factors to the differences in  
11 observed response. A number of other limitations in the studies by Dawson et al. (1993) and  
12 Johnson et al. (2005)(2003) have been suggested (Hardin et al., 2005; Watson et al., 2006). One  
13 concern is the lack of clear dose-response relationship for the incidence of any specific cardiac  
14 anomaly or combination of anomalies, a disparity for which no reasonable explanation has been  
15 put forth. In addition, analyses on a fetal- rather than litter-basis and the pooling of data  
16 collected over an extended period, including nonconcurrent controls, have been criticized. With  
17 respect to the first issue, the study authors provided individual litter incidence data to EPA for  
18 analysis (see Section 5, Dose-Response Assessment), and, in response to the second issue, the  
19 study authors provided further explanation as to their experimental procedures (Johnson et al.,  
20 2004). In sum, while the studies by Dawson et al. (1993) and Johnson et al. (2005)(2003) have  
21 significant limitations, there is insufficient reason to dismiss their findings.

22 Finally, mechanistic studies, particularly based on the avian studies mentioned above,  
23 provide additional support for TCE-induced fetal cardiac malformation, particularly with respect  
24 to defects involving septal and valvular morphogenesis. As summarized by NRC (2006), there is  
25 substantial concordance in the stages and events of cardiac valve formation between mammals  
26 and birds. While quantitative extrapolation of findings from avian studies to humans is not  
27 possible without appropriate kinetic data for these experimental systems, the treatment-related  
28 alterations in endothelial cushion development observed in avian in ovo and in vitro studies  
29 (Boyer et al., 2000; Mishima et al., 2006; Ou et al., 2003) provide a plausible mechanistic basis  
30 for defects in septal and valvular morphogenesis observed in rodents, and consequently support  
31 the plausibility of cardiac defects induced by TCE in humans.

32 Postnatal developmental outcomes examined after TCE prenatal and/or postnatal  
33 exposure in both humans and experimental animals include developmental neurotoxicity,  
34 developmental immunotoxicity, and childhood cancer. Effects on the developing nervous  
35 system included a broad array of structural and behavioral alterations in humans (ATSDR,

1 2003b; Beppu, 1968; Bernad et al., 1987; Burg and Gist, 1997; Burg et al., 1995; Laslo-Baker et  
2 al., 2004; Till et al., 2001a; White et al., 1997, abstract; Windham et al., 2006) and animals  
3 (Blossom et al., 2008; Fredriksson et al., 1993; George et al., 1986; Isaacson and Taylor, 1989;  
4 Narotsky and Kavlock, 1995; Noland-Gerbec et al., 1986; Taylor et al., 1985; Westergren et al.,  
5 1984). Adverse immunological findings in humans following developmental exposures to TCE  
6 were reported by Lehmann et al. (2002) and Byers et al. (1988). In mice, alterations in T-cell  
7 subpopulations, spleen and/or thymic cellularity, cytokine production, autoantibody levels (in an  
8 autoimmune-prone mouse strain), and/or hypersensitivity response were observed after  
9 exposures during development (Blossom and Doss, 2007; Blossom et al., 2008; Peden-Adams et  
10 al., 2006; Peden-Adams et al., 2008), Childhood cancers included leukemia and non-Hodgkin  
11 lymphoma (ADHS, 1990; Cohn et al., 1994b; Costas et al., 2002; Cutler et al., 1986; Flood,  
12 1988; Flood, 1997; Kioski et al., 1990a; Kioski et al., 1990b; Lagakos et al., 1986; Lowengart et  
13 al., 1987; McKinney et al., 1991; MDPH, 1997a; Morgan and Cassady, 2002; Shu et al., 1999),  
14 CNS tumors (ADHS, 1990; Flood, 1988; Flood, 1997; Kioski et al., 1990a)Peters and Preston-  
15 Martin, 1981)(De Roos et al., 2001; Morgan and Cassady, 2002; Peters et al., 1985; Peters et al.,  
16 1981), and total cancers ((ADHS, 1990; Flood, 1988; Flood, 1997; Porter, 1993){Morgan, 2002,  
17 707097;ATSDR, 2008, 730402}(ATSDR, 2006a). These outcomes are discussed in the other  
18 relevant sections for neurotoxicity, immunotoxicity, and carcinogenesis.

#### 4.11.2. Characterization of Carcinogenicity

19 Following EPA (2005c) *Guidelines for Carcinogen Risk Assessment*, TCE is  
20 characterized as “*carcinogenic to humans*” by all routes of exposure. This conclusion is based  
21 on convincing evidence of a causal association between TCE exposure in humans and kidney  
22 cancer. The kidney cancer association cannot be reasonably attributed to chance, bias, or  
23 confounding. The human evidence of carcinogenicity from epidemiologic studies of TCE  
24 exposure is compelling for NHL but less convincing than for kidney cancer, and more limited for  
25 liver and biliary tract cancer. In addition to the body of evidence pertaining to kidney cancer,  
26 NHL, and liver cancer, the available epidemiologic studies also provide more limited evidence of  
27 an association between TCE exposure and other types of cancer, including bladder, esophageal,  
28 prostate, cervical, breast, and childhood leukemia. Differences between these sets of data and  
29 the data for kidney cancer, NHL, and liver cancer are observations from fewer numbers of  
30 studies, a mixed pattern of observed risk estimates, and the general absence of exposure-response  
31 data from the studies using a quantitative TCE-specific exposure measure.

32 There are several lines of supporting evidence for TCE carcinogenicity in humans. First,  
33 TCE induces site-specific tumors in rodents given TCE by oral gavage and inhalation. Second,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 toxicokinetic data indicate that TCE absorption, distribution, metabolism, and excretion are  
2 qualitatively similar in humans and rodents. Finally, there is sufficient weight of evidence to  
3 conclude that a mutagenic MOA is operative for TCE-induced kidney tumors, and this MOA is  
4 clearly relevant to humans. MOAs have not been established for other TCE-induced tumors in  
5 rodents, and no mechanistic data indicate that any hypothesized key events are biologically  
6 precluded in humans.

7

#### **4.11.2.1.1. Summary Evaluation of Epidemiologic Evidence of Trichloroethylene (TCE) and Cancer**

8 The available epidemiologic studies provide convincing evidence of a causal association  
9 between TCE exposure and cancer. The strongest epidemiologic evidence consists of reported  
10 increased risks of kidney cancer, with more limited evidence for NHL and liver cancer, in  
11 several well-designed cohort and case-control studies (discussed below). The summary  
12 evaluation below of the evidence for causality is based on guidelines adapted from Hill (1965)  
13 by EPA (2005c), and focuses on evidence related to kidney cancer, NHL, and liver cancer.

14

#### **4.11.2.1.2. (a) Consistency of observed association**

15 Elevated risks for kidney cancer have been observed across many independent studies.  
16 Eighteen studies in which there is a high likelihood of TCE exposure in individual study subjects  
17 (e.g., based on job-exposure matrices or biomarker monitoring) and which were judged to have  
18 met, to a sufficient degree, the standards of epidemiologic design and analysis, were identified in  
19 a systematic review of the epidemiologic literature. Of the 15 of these studies reporting risks of  
20 kidney cancer (Anttila et al., 1995; Axelson et al., 1994; Boice et al., 1999; Brüning et al., 2003;  
21 Charbotel et al., 2006; Dosemeci et al., 1999; Greenland et al., 1994; Hansen et al., 2001; Moore  
22 et al., 2010; Morgan et al., 1998; Pesch et al., 2000b; Raaschou-Nielsen et al., 2003; Radican et  
23 al., 2008; Siemiatycki, 1991; Zhao et al., 2005), most estimated relative risks between 1.1 and  
24 1.9 for overall exposure to TCE. Six of these 15 studies reported statistically significant  
25 increased risks either for overall exposure to TCE (Brüning et al., 2003; Dosemeci et al., 1999;  
26 Moore et al., 2010; Raaschou-Nielsen et al., 2003) or for one of the highest TCE exposure group  
27 (Charbotel et al., 2006; Moore et al., 2010; Raaschou-Nielsen et al., 2003; Zhao et al., 2005).  
28 Thirteen other cohort, case-control, and geographic based studies were given less weight because  
29 of their lesser likelihood of TCE exposure and other study design limitations that would decrease  
30 statistical power and study sensitivity (see Sections 4.1 and 4.4.2).

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 The consistency of association between TCE exposure and kidney cancer is further  
2 supported by the results of the meta-analyses of the 15 cohort and case-control studies of  
3 sufficient quality and with high probability TCE exposure potential to individual subjects. These  
4 analyses observed a statistically significant increased summary relative risk estimate (RR<sub>m</sub>) for  
5 kidney cancer of 1.27 (95% CI: 1.13, 1.43) for overall TCE. The summary relative risk were  
6 robust and did not change appreciably with the removal of any individual study or with the use  
7 of alternate relative risk estimates from individual studies. In addition, there was no evidence for  
8 heterogeneity or publication bias.

9 The consistency of increased kidney cancer relative risk estimates across a large number  
10 of independent studies of different designs and populations from different countries and  
11 industries argues against chance, bias or confounding as the basis for observed associations.  
12 This consistency, thus, provides substantial support for a causal effect between kidney cancer  
13 and TCE exposure.

14 Some evidence of consistency is found between TCE exposure and NHL and liver  
15 cancer. In a weight-of-evidence review of the NHL studies, 17 studies in which there is a high  
16 likelihood of TCE exposure in individual study subjects (e.g., based on job-exposure matrices or  
17 biomarker monitoring) and which met, to a sufficient degree, the standards of epidemiologic  
18 design and analysis were identified. These studies generally reported excess relative risk  
19 estimates for NHL between 0.8 and 3.1 for overall TCE exposure. Statistically significant  
20 elevated relative risk estimates for overall exposure were observed in two cohort (Hansen et al.,  
21 2001; Raaschou-Nielsen et al., 2003) and one case-control (Hardell et al., 1994) studies. The  
22 other 14 identified studies reported elevated relative risk estimates with overall TCE exposure  
23 that were not statistically significant (Anttila et al., 1995; Axelson et al., 1994; Boice et al., 1999;  
24 Cocco et al., 2010; Greenland et al., 1994; Miligi et al., 2006; Morgan et al., 1998; Nordström et  
25 al., 1998; Persson and Fredrikson, 1999; Purdue et al., 2011; Radican et al., 2008; Siemiatycki,  
26 1991; Wang et al., 2009; Zhao et al., 2005). Fifteen additional studies were given less weight  
27 because of their lesser likelihood of TCE exposure and other design limitations that would  
28 decrease study power and sensitivity (see Sections 4.1 and 4.6.1.2). The observed lack of  
29 association with NHL in these studies likely reflects study design and exposure assessment  
30 limitations and is not considered inconsistent with the overall evidence on TCE and NHL.

31 Consistency of the association between TCE exposure and NHL is further supported by  
32 the results of meta-analyses. These meta-analyses found a statistically significant increased  
33 summary relative risk estimate for NHL of 1.23 (95% CI: 1.07, 1.42) for overall TCE exposure.  
34 This result and its statistical significance were not overly influenced by most individual studies.  
35 Some heterogeneity was observed across the 17 studies of overall exposure, though it was not



1 statistically significant ( $p = 0.16$ ). Analyzing the cohort and case-control studies separately  
2 resolved most of the heterogeneity, but the result for the summary case-control studies was only  
3 about a 7% increased relative risk estimate and was not statistically significant. The sources of  
4 heterogeneity are uncertain but may be the result of some bias associated with exposure  
5 assessment and/or disease classification, or from differences between cohort and case-control  
6 studies in average TCE exposure. In addition, there is some evidence of potential publication  
7 bias in this data set; however, it is uncertain that this is actually publication bias rather than an  
8 association between standard error and effect size resulting for some other reason, e.g., a  
9 difference in study populations or protocols in the smaller studies. Furthermore, if there is  
10 publication bias in this data set, it does not appear to account completely for the finding of an  
11 increased NHL risk.

12         There are fewer studies on liver cancer than for kidney cancer and NHL. Of nine studies,  
13 all of them cohort studies, in which there is a high likelihood of TCE exposure in individual  
14 study subjects (e.g., based on job-exposure matrices or biomarker monitoring) and which met, to  
15 a sufficient degree, the standards of epidemiologic design and analysis in a systematic review  
16 (Anttila et al., 1995; Axelson et al., 1994; Boice et al., 1999; Boice et al., 2006b; Greenland et  
17 al., 1994; Hansen et al., 2001; Morgan et al., 1998; Raaschou-Nielsen et al., 2003; Radican et al.,  
18 2008), most reported relative risk estimates for liver and gallbladder cancer between 0.5 and 2.0  
19 for overall exposure to TCE. Relative risk estimates were generally based on small numbers of  
20 cases or deaths, with the result of wide confidence intervals on the estimates, except for one  
21 study (Raaschou-Nielsen et al., 2003). This study has almost 6 times more cancer cases than the  
22 next largest study and observed a statistically significant elevated liver and gallbladder cancer  
23 risk with overall TCE exposure (RR = 1.35 [95% CI: 1.03, 1.77]). Ten additional studies were  
24 given less weight because of their lesser likelihood of TCE exposure and other design limitations  
25 that would decrease statistical power and study sensitivity (see Sections 4.1 and 4.5.2).

26         Consistency of the association between TCE exposure and liver cancer is further  
27 supported by the results of meta-analyses. These meta-analyses found a statistically significant  
28 increased summary relative risk estimate for liver and biliary tract cancer of 1.29 (95% CI: 1.07,  
29 1.56) with overall TCE exposure. Although there was no evidence of heterogeneity or  
30 publication bias and the summary estimate was fairly insensitive to the use of alternative relative  
31 risk estimates, the statistical significance of the summary estimate depends heavily on the one  
32 large study by Raaschou-Nielsen et al. (2003). However, there were fewer adequate studies  
33 available for meta-analysis of liver cancer (9 versus 17 for NHL and 15 for kidney), leading to  
34 lower statistical power, even with pooling. Moreover, liver cancer is comparatively rarer, with

1 age-adjusted incidences roughly half or less those for kidney cancer or NHL; thus, fewer liver  
2 cancer cases are generally observed in individual cohort studies.

3

#### 4.11.2.1.3. (b) Strength of the observed association

4 In general, the observed associations between TCE exposure and cancer are modest, with  
5 relative risks or odds ratios for overall TCE exposure generally less than 2.0, and higher relative  
6 risks or odds ratios for high exposure categories. Among the highest statistically significant  
7 relative risks were those reported for kidney cancer in the studies by Henschler et al. (1995)  
8 (7.97 [95% CI: 2.59, 8.59]) and Vamvakas et al. (1998) (10.80 [95% CI: 3.36, 34.75]). As  
9 discussed in Section 4.5.3, risk magnitude in both studies is highly uncertain due, in part, to  
10 possible selection biases, and neither was included in the meta-analyses. However, the findings  
11 of these studies were corroborated, though with lower reported relative risks, by later studies  
12 which overcame many of their deficiencies, such as Brüning et al. (2003) (2.47 [95% CI: 1.36,  
13 4.49]), Charbotel et al. (2006; 2009) (2.16 [95% CI: 1.02, 4.60] for the high cumulative exposure  
14 group), and Moore et al. (2010) (2.05 [95% CI: 1.13, 3.73] for high confidence assessment of  
15 TCE). In addition, the very high apparent exposure in the subjects of Henschler et al. (1995) and  
16 Vamvakas et al. (1998) may have contributed to their reported relative risks being higher than  
17 those in other studies. Exposures in most population case-control studies are of lower overall  
18 TCE intensity compared to exposures in Brüning et al. (2003) and Charbotel et al. (2006; 2009),  
19 and, as would be expected, observed relative risk estimates are lower (1.24 [95% CI: 1.03,  
20 1.49]), Pesch et al. (2000b); 1.30 [95% CI: 0.9, 1.9], Dosemeci et al. (1999)). A few high-quality  
21 cohort and case-control studies reported statistically significant relative risks of approximately  
22 2.0 with highest exposure, including Zhao et al. (2005) (4.9 [95% CI: 1.23, 19.6] for high TCE  
23 score), Raaschou-Nielsen et al. (2003) (1.7 [95% CI: 1.1, 2.4] for  $\geq 5$  year exposure duration,  
24 subcohort with higher exposure]), Charbotel et al. (2006) (2.16 [95% CI: 1.02, 4.60] for high  
25 cumulative exposure and 2.73 [95% CI: 1.06, 7.07] for high cumulative exposure plus peaks) and  
26 Moore et al. (2010) (2.23 [95% CI: 1.07, 4.64] for high cumulative exposure and 2.41 [95% CI:  
27 1.05, 5.56] for high average intensity TCE exposure).

28 Among the highest statistically significant relative risks reported for NHL were those of  
29 Hansen et al. (2001) (3.1 [95% CI: 1.3, 6.1]), Hardell et al. (1994) (7.2 [95% CI: 1.3, 42]), the  
30 latter a case-control study whose magnitude of risk is uncertain because of self-reported  
31 occupational TCE exposure. A similar magnitude of risk was reported in Purdue et al. (2011) for  
32 highest exposure (3.3 [95% CI: 1.1, 10.1], >234,000 ppm-hour, and 7.9 [95% CI: 1.8, 34.3],

1 >360 ppm-hour/week). Observed relative risk estimates for liver cancer and overall TCE  
2 exposure are generally more modest.

3 The strength of association between TCE exposure and cancer is modest with overall  
4 TCE exposure. Large relative risk estimates are considered strong evidence of causality;  
5 however, a modest risk does not preclude a causal association and may reflect a lower level of  
6 exposure, an agent of lower potency, or a common disease with a high background level (U.S.  
7 EPA, 2005c). Modest relative risk estimates have been observed with several well-established  
8 human carcinogens such as benzene and secondhand smoke. Chance cannot explain the  
9 observed association between TCE and cancer; statistically significant associations are found in a  
10 number of the studies that contribute greater weight to the overall evidence, given their design  
11 and statistical analysis approaches. In addition, other known or suspected risk factors cannot  
12 fully explain the observed elevations in kidney cancer relative risks. All kidney cancer case-  
13 control studies included adjustment for possible confounding effects of smoking, and some  
14 studies included body mass index, hypertension, and coexposure to other occupational agents  
15 such as cutting or petroleum oils. Cutting oils and petroleum oils, known as metalworking  
16 fluids, have not been associated with kidney cancer (Mirer, 2010; NIOSH, 1998), and potential  
17 confounding by this occupational co-exposure is unable to explain the observed association with  
18 TCE. Additionally, the associations between kidney cancer and TCE exposure remained in these  
19 studies after statistical adjustment for possible known and suspected confounders. Charbotel et  
20 al. (2005) observed a nonstatistically significantly kidney cancer risk with exposure to TCE  
21 adjusted for cutting or petroleum oil exposures (1.96 [95% CI: 0.71, 5.37] for the high-cumulative  
22 exposure group and 2.63 [95% CI: 0.79, 8.83] for high-exposure group with peaks).

23 All kidney cancer case-control studies adjusted for smoking except the Moore et al.  
24 (2010) study, which reported that smoking did not significantly change the overall association  
25 with TCE exposure. Although direct examination of smoking and other suspected kidney cancer  
26 risk factors is usually not possible in cohort studies, confounding is less likely in Zhao et al.  
27 (2005), given their use of an internal referent group and adjustment for socioeconomic status, an  
28 indirect surrogate for smoking, and other occupational exposures. In addition, the magnitude of  
29 the lung cancer risk in Raaschou-Nielsen et al. (2003) suggests a high smoking rate is unlikely  
30 and cannot explain their finding on kidney cancer. Last, a meta-analysis of the nine cohort  
31 studies that reported kidney cancer risks found a summary relative risk estimate for lung cancer  
32 of 0.96 (95% CI: 0.76, 1.21) for overall TCE exposure and 0.96 (95% CI: 0.72, 1.27) for the  
33 highest exposure group. These observations suggest that confounding by smoking is not an  
34 alternative explanation for the kidney cancer meta-analysis results.

1 Few risk factors are recognized for NHL, with the exception of viruses and suspected  
2 factors such as immunosuppression or smoking, which are associated with specific NHL  
3 subtypes. Associations between NHL and TCE exposure are based on groupings of several NHL  
4 subtypes. Three of the seven NHL case-control studies adjusted for age, sex and smoking in  
5 statistical analyses (Miligi et al., 2006; Wang et al., 2009) two others adjusted for age, sex and  
6 education (Cocco et al., 2010; Purdue et al., 2011), and the other three case-control studies  
7 adjusted for age only or age and sex (Hardell et al., 1994; Nordström et al., 1998; Persson and  
8 Fredrikson, 1999). Like for kidney cancer, direct examination of possible confounding in cohort  
9 studies is not possible. The use of internal controls in some of the cohort studies is intended to  
10 reduce possible confounding related to lifestyle differences, including smoking habits, between  
11 exposed and referent subjects.

12 Heavy alcohol use and viral hepatitis are established risk factors for liver cancer, with  
13 severe obesity and diabetes characterized as a metabolic syndrome associated with liver cancer.  
14 Only cohort studies for liver cancer are available, and they were not able to consider these  
15 possible risk factors.

#### 4.11.2.1.4. (c) Specificity of the observed association

16 Specificity is generally not as relevant as other aspects for judging causality. As stated in  
17 the EPA *Guidelines for Carcinogen Risk Assessment* (2005), based on our current understanding  
18 that many agents cause cancer at multiple sites, and cancers have multiple causes, the absence of  
19 specificity does not detract from evidence for a causal effect. Evidence for specificity could be  
20 provided by a biological marker in tumors that was specific to TCE exposure. There is some  
21 evidence suggesting particular VHL mutations in kidney tumors may be caused by TCE, but  
22 uncertainties in these data preclude a definitive conclusion.

23

#### 4.11.2.1.5. (d) Temporal relationship of the observed association

24 Each cohort study was evaluated for the adequacy of the follow-up period to account for  
25 the latency of cancer development. The studies with the greatest weight based on study design  
26 characteristics (e.g., those used in the meta-analysis) all had adequate follow-up to assess  
27 associations between TCE exposure and cancer. Therefore, the findings of those studies are  
28 consistent with a temporal relationship.

29

#### 4.11.2.1.6. (e) Biological gradient (exposure-response relationship)

1 Exposure-response relationships are examined in the TCE epidemiologic studies only to a  
2 limited extent. Many studies examined only overall “exposed” versus “unexposed” groups and  
3 did not provide exposure information by level of exposure. Others do not have adequate  
4 exposure assessments to confidently distinguish between levels of exposure. For example, many  
5 studies used duration of employment as an exposure surrogate; however, this is a poor exposure  
6 metric given subjects may have differing exposure intensity with similar exposure duration  
7 (NRC, 2006).

8 Three studies of kidney cancer reported a statistically significant trend of increasing risk  
9 with increasing TCE exposure, Zhao et al. (2005) ( $p = 0.023$  for trend with TCE score),  
10 Charbotel et al. (2005; 2007) ( $p = 0.04$  for trend with cumulative TCE exposure) and Moore et  
11 al. (2010) ( $p = 0.02$  for trend with cumulative TCE exposure). Charbotel et al. (2007) was  
12 specifically designed to examine TCE exposure and had a high-quality exposure assessment and  
13 the Moore et al. (2010) exposure assessment considered detailed information on jobs using  
14 solvents. Zhao et al. (2005) also had a relatively well-designed exposure assessment. A positive  
15 trend was also observed in one other study (Raaschou-Nielsen et al. (2003), with employment  
16 duration).

17 Biological gradient is further supported by meta-analyses for kidney cancer using only  
18 the highest exposure groups and accounting for possible reporting bias, which yielded a higher  
19 summary relative risk estimate (1.58 [95% CI: 1.28, 1.96]) than for overall TCE exposure (1.27  
20 [95% CI: 1.13, 1.43]). Although this analysis uses a subset of studies in the overall TCE  
21 exposure analysis, the finding of higher risk in the highest exposure groups, where such groups  
22 were available, is consistent with a trend of increased risk with increased exposure.

23 The NHL case-control study of Purdue et al. (2011) reported a statistically significant  
24 trend with TCE exposure ( $p = 0.02$  for trend with average-weekly TCE exposure), and NHL risk  
25 in Boice et al. (1999) appeared to increase with increasing exposure duration ( $p = 0.20$  for  
26 routine-intermittent exposed subjects). The borderline trend with TCE intensity in the case-  
27 control studies of Wang et al. (2009) ( $p = 0.06$ ) and Purdue et al. (2011) ( $p = 0.08$  for trend with  
28 cumulative TCE exposure) is consistent with their findings for average weekly TCE exposure.  
29 As with kidney cancer, further support was provided by meta-analyses using only the highest  
30 exposure groups, which yielded a higher summary relative risk estimate (1.43 [95% CI: 1.13,  
31 1.82]) than for overall TCE exposure (1.23 [95% CI: 1.07, 1.42]). For liver cancer, the meta-  
32 analyses using only the highest exposure groups yielded a lower, and nonstatistically significant,  
33 summary estimate (1.28 [95% CI: 0.93, 1.77]) than for overall TCE exposure (1.29 [95% CI:  
34 1.07, 1.56]). There were no case-control studies on liver cancer and TCE, and the cohort studies

1 generally had few liver cancer cases, making it more difficult to assess exposure-response  
2 relationships. The one large study (Raaschou-Nielsen et al., 2003) used only duration of  
3 employment, which is an inferior exposure metric.  
4

#### 4.11.2.1.7. (f) Biological plausibility

5 TCE metabolism is similar in humans, rats, and mice and results in reactive metabolites.  
6 TCE is metabolized in multiple organs and metabolites are systemically distributed. Several  
7 oxidative metabolites produced primarily in the liver, including CH, TCA and DCA, are rodent  
8 hepatocarcinogens. Two other metabolites, DCVC and DCVG, which can be produced and  
9 cleared by the kidney, have shown genotoxic activity, suggesting the potential for  
10 carcinogenicity. Kidney cancer, NHL, and liver cancer have all been observed in rodent  
11 bioassays (see below). The laboratory animal data for liver and kidney cancer are the most  
12 robust, corroborated in multiple studies, sexes, and strains, although each has only been reported  
13 in a single species and the incidences of kidney cancer are quite low. Lymphomas were only  
14 reported to be statistically significantly elevated in a single study in mice, but one additional  
15 mouse study reported elevated lymphoma incidence and one rat study reported elevated leukemia  
16 incidence. In addition, there is some evidence both in humans and laboratory animals for kidney,  
17 liver and immune system noncancer toxicity from TCE exposure. Several hypothesized modes  
18 of action have been presented for the rodent tumor findings, although there are insufficient data  
19 to support any one mode of action, and the available evidence does not preclude the relevance of  
20 the hypothesized modes of action to humans. Activation of macrophages, natural killer cells,  
21 and cytokine production (e.g., tumor necrosis factor), may also play an etiologic role in  
22 carcinogenesis, and so the immune-related effects of TCE should also be considered. In  
23 addition, the decreased in lymphocyte counts and subsets, including CD4+ T cells, and decreased  
24 lymphocyte activation seen in TCE-exposed workers (Lan et al., 2010) also support the  
25 biological plausibility of a role of TCE exposure in NHL.  
26

#### 4.11.2.1.8. (g) Coherence

27 Coherence is defined as consistency with the known biology. As discussed under  
28 biological plausibility, the observance of kidney and liver cancer, and NHL in humans is  
29 consistent with the biological processing and toxicity of TCE.  
30

#### 4.11.2.1.9. (h) Experimental evidence (from human populations)

1 Few experimental data from human populations are available on the relationship between  
2 TCE exposure and cancer. The only study of a “natural experiment” (i.e., observations of a  
3 temporal change in cancer incidence in relation to a specific event) notes that childhood  
4 leukemia cases appeared to be more evenly distributed throughout Woburn, MA, after closure of  
5 the two wells contaminated with trichloroethylene and other organic solvents (MDPH, 1997b).  
6

#### 4.11.2.1.10. (i) Analogy

7 Exposure to structurally related chlorinated solvents such as tetrachloroethylene and  
8 dichloromethane have also been associated with kidney, lymphoid, and liver tumors in human,  
9 although the evidence for TCE is considered stronger.  
10

#### 4.11.2.1.11. Conclusion

11 In conclusion, based on the weight-of-evidence analysis for kidney cancer and in  
12 accordance with EPA guidelines, TCE is characterized as “*carcinogenic to humans.*” This  
13 hazard descriptor is used when there is convincing epidemiologic evidence of a causal  
14 association between human exposure and cancer. Convincing evidence is found in the  
15 consistency of the kidney cancer findings. The consistency of increased kidney cancer relative  
16 risk estimates across a large number of independent studies of different designs and populations  
17 from different countries and industries provides compelling evidence given the difficulty, a  
18 priori, in detecting effects in epidemiologic studies when the relative risks are modest, the  
19 cancers are relatively rare, and therefore, individual studies have limited statistical power. This  
20 strong consistency argues against chance, bias, and confounding as explanations for the elevated  
21 kidney cancer risks. In addition, statistically significant exposure-response trends are observed  
22 in high-quality studies. These studies were designed to examine kidney cancer in populations  
23 with high TCE exposure intensity. These studies addressed important potential confounders and  
24 biases, further supporting the observed associations with kidney cancer as causal. In a meta-  
25 analysis of the 15 studies that met the inclusion criteria, a statistically significant summary  
26 relative risk estimate was observed for overall TCE exposure (RR<sub>m</sub>: 1.27 [95% CI: 1.13, 1.43]).  
27 The summary relative risk estimate was greater for the highest TCE exposure groups (RR<sub>m</sub>: 1.58  
28 [95% CI: 1.28, 1.96]; *n* = 13 studies). Meta-analyses investigating the influence of individual  
29 studies and the sensitivity of the results to alternate relative risk estimate selections found the

1 summary relative risk estimates to be highly robust. Furthermore, there was no indication of  
2 publication bias or significant heterogeneity. It would require a substantial amount of negative  
3 data from informative studies (i.e., studies having a high likelihood of TCE exposure in  
4 individual study subjects and which meet, to a sufficient degree, the standards of epidemiologic  
5 design and analysis in a systematic review) to contradict this observed association.

6 The evidence is less convincing for NHL and liver cancer. While the evidence is strong  
7 for NHL, issues of (nonstatistically significant) study heterogeneity, potential publication bias,  
8 and weaker exposure-response results contribute greater uncertainty. The evidence is more  
9 limited for liver cancer mainly because only cohort studies are available and most of these  
10 studies have small numbers of cases. In addition to the body of evidence described above  
11 pertaining to kidney cancer, NHL, and liver cancer, the available epidemiologic studies also  
12 provide suggestive evidence of an association between TCE exposure and other types of cancer,  
13 including bladder, esophageal, prostate, cervical, breast, and childhood leukemia, breast.  
14 Differences between these sets of data and the data for kidney cancer, NHL, and liver cancer are  
15 observations are from fewer numbers of studies, a mixed pattern of observed risk estimates and  
16 the general absence of exposure-response data from the studies using a quantitative TCE-specific  
17 cumulative exposure measure.  
18

#### **4.11.2.1.12. Summary of Evidence for Trichloroethylene (TCE) Carcinogenicity in Rodents**

19 Additional evidence of TCE carcinogenicity consists of increased incidences of tumors  
20 reported in multiple chronic bioassays in rats and mice. In total, this database identifies some of  
21 the same target tissues of TCE carcinogenicity also seen in epidemiological studies, including the  
22 kidney, liver, and lymphoid tissues.

23 Of particular note is the site-concordant finding of TCE-induced kidney cancer in rats. In  
24 particular, low, but biologically and sometimes statistically significant, increases in the incidence  
25 of kidney tumors were observed in multiple strains of rats treated with TCE by either inhalation  
26 or corn oil gavage (Maltoni et al., 1986; NTP, 1988, 1990). For instance, Maltoni et al. (1986)  
27 reported that although only 4/130 renal adenocarcinomas in rats in the highest dose group, these  
28 tumors had never been observed in over 50,000 Sprague-Dawley rats (untreated, vehicle-treated,  
29 or treated with different chemicals) examined in previous experiments in the same laboratory. In  
30 addition, the gavage study by NCI (1976) and two inhalation studies by Henschler et al. (1980),  
31 and Fukuda et al. (1983) each observed one renal adenoma or adenocarcinoma in some dose  
32 groups and none in controls. The largest (but still small) incidences were observed in treated  
33 male rats, only in the highest dose groups. However, given the small numbers, an effect in



1 females cannot be ruled out. Several studies in rats were limited by excessive toxicity,  
2 accidental deaths, or deficiencies in reporting (NCI, 1976; NTP, 1988, 1990). Individually,  
3 therefore, these studies provide only suggestive evidence of renal carcinogenicity. Overall,  
4 given the rarity of these types of tumors in the rat strains tested and the repeated similar results  
5 across experiments and strains, these studies taken together support the conclusion that TCE is a  
6 kidney carcinogen in rats, with males being more sensitive than females. No other tested  
7 laboratory species (i.e., mice and hamsters) have exhibited increased kidney tumors, although  
8 high incidences of kidney toxicity have been reported in mice (Maltoni et al., 1986; NCI, 1976;  
9 NTP, 1990). The GSH-conjugation-derived metabolites suspected of mediating TCE-induced  
10 kidney carcinogenesis have not been tested in a standard 2-year bioassay, so their role cannot be  
11 confirmed definitively. However, it is clear that GSH conjugation of TCE occurs in humans and  
12 that the human kidney contains the appropriate enzymes for bioactivation of GSH conjugates.  
13 Therefore, the production of the active metabolites thought to be responsible for kidney tumor  
14 induction in rats likely occurs in humans.

15 Statistically significant increases in TCE-induced liver tumors have been reported in  
16 multiple inhalation and gavage studies with male Swiss mice and B6C3F1 mice of both sexes  
17 (Anna et al., 1994; Bull et al., 2002; Herren-Freund et al., 1987; Maltoni et al., 1986; NCI, 1976;  
18 NTP, 1990). In female Swiss mice, on the other hand, Fukuda et al. (1983), in CD-1 (ICR,  
19 Swiss-derived) mice, and Maltoni et al. (1986) both reported small, nonsignificant increases at  
20 the highest dose by inhalation. Henschler et al. (1984; 1980) reported no increases in either sex  
21 of Han:NMRI (also Swiss-derived) mice exposed by inhalation and ICR/HA (Swiss) mice  
22 exposed by gavage. However, the inhalation study (Henschler et al., 1980) had only 30 mice per  
23 dose group and the gavage study (Henschler et al., 1984) had dosing interrupted due to toxicity.  
24 Studies in rats (Henschler et al., 1980; Maltoni et al., 1986; NCI, 1976; NTP, 1988, 1990) and  
25 hamsters (Henschler et al., 1980) did not report statistically significant increases in liver tumor  
26 induction with TCE treatment. However, several studies in rats were limited by excessive  
27 toxicity or accidental deaths (NCI, 1976; NTP, 1988, 1990), and the study in hamsters only had  
28 30 animals per dose group. These data are inadequate for concluding that TCE lacks  
29 hepatocarcinogenicity in rats and hamsters, but are indicative of a lower potency in these species.  
30 Moreover, it is notable that a few studies in rats reported low incidences (too few for statistical  
31 significance) of very rare biliary- or endothelial-derived tumors in the livers of some treated  
32 animals (Fukuda et al., 1983; Henschler et al., 1980; Maltoni et al., 1986). Further evidence for  
33 the hepatocarcinogenicity of TCE is derived from chronic bioassays of the TCE oxidative  
34 metabolites CH, TCA, and DCA in mice (e.g., Bull et al., 1990; DeAngelo et al., 1996;  
35 DeAngelo et al., 2008; DeAngelo et al., 1999; George et al., 2000; Leakey et al., 2003a) (Leakey

1 et al., 2003a), all of which reported hepatocarcinogenicity. Very limited testing of these TCE  
2 metabolites has been done in rats, with a single experiment reported in both Richmond et al.  
3 (1995) and DeAngelo et al. (1996) finding statistically significant DCA-induced  
4 hepatocarcinogenicity. With respect to TCA, DeAngelo et al. (1997), often cited as  
5 demonstrating lack of hepatocarcinogenicity in rats, actually reported elevated adenoma  
6 multiplicity and carcinoma incidence from TCA treatment. However, statistically, the role of  
7 chance could not be confidently excluded because of the low number of animals per dose group  
8 (20–24 per treatment group at final sacrifice). Overall, TCE and its oxidative metabolites are  
9 clearly carcinogenic in mice, with males more sensitive than females and the B6C3F1 strain  
10 appearing to be more sensitive than the Swiss strain. Such strain and sex differences are not  
11 unexpected, as they appear to parallel, qualitatively, differences in background tumor incidence.  
12 Data in other laboratory animal species are limited. Thus, except for DCA, which is  
13 carcinogenic in rats, inadequate evidence exists to evaluate the hepatocarcinogenicity of these  
14 compounds in rats or hamsters. However, to the extent that there is hepatocarcinogenic potential  
15 in rats, TCE is clearly less potent in the strains tested in this species than in B6C3F1 and Swiss  
16 mice.

17 Additionally, there is more limited evidence for TCE-induced lymphatic cancers in rats  
18 and mice, lung tumors in mice, and testicular tumors in rats. With respect to the lymphomas,  
19 Henschler et al. (1980) reported statistically significant increases in lymphomas in female  
20 Han:NMRI mice treated via inhalation. While Henschler et al. (1980) suggested these  
21 lymphomas were of viral origin specific to this strain, subsequent studies reported increased  
22 lymphomas in female B6C3F1 mice treated via corn oil gavage (NTP, 1990) and leukemias in  
23 male Sprague-Dawley and female August rats (Maltoni et al., 1986; NTP, 1988). However,  
24 these tumors had relatively modest increases in incidence with treatment, and were not reported  
25 to be increased in other studies. With respect to lung tumors, rodent bioassays have  
26 demonstrated a statistically significant increase in pulmonary tumors in mice following chronic  
27 inhalation exposure to TCE (Fukuda et al., 1983; Maltoni et al., 1988; Maltoni et al., 1986).  
28 Pulmonary tumors were not reported in other species tested (i.e., rats and hamsters; (Fukuda et  
29 al., 1983; Henschler et al., 1980; Maltoni et al., 1988; Maltoni et al., 1986)). Chronic oral  
30 exposure to TCE led to a nonstatistically significant increase in pulmonary tumors in mice but,  
31 again, not in rats or hamsters (Henschler et al., 1984; Maltoni et al., 1986; NCI, 1976; NTP,  
32 1988, 1990; Van Duuren et al., 1979). A lower response via oral exposure would be consistent  
33 with a role of respiratory metabolism in pulmonary carcinogenicity. Finally, increased testicular  
34 (interstitial cell and Leydig cell) tumors have been observed in rats exposed by inhalation and  
35 gavage (Maltoni et al., 1986; NTP, 1988, 1990). Statistically significant increases were reported

1 in Sprague-Dawley rats exposed via inhalation (Maltoni et al., 1986) and Marshall rats exposed  
2 via gavage (NTP, 1988). In three rat strains, ACI, August, and F344/N, a high (>75%) control  
3 rate of testicular tumors was observed, limiting the ability to detect a treatment effect (NTP,  
4 1988, 1990).

5 In summary, there is clear evidence for TCE carcinogenicity in rats and mice, with  
6 multiple studies showing TCE to cause tumors at multiple sites. The apparent lack of site  
7 concordance across laboratory animal species may be due to limitations in design or conduct in a  
8 number of rat bioassays and/or genuine interspecies differences in sensitivity. Nonetheless, these  
9 studies have shown carcinogenic effects across different strains, sexes, and routes of exposure,  
10 and site-concordance is not necessarily expected for carcinogens. Of greater import is the  
11 finding that there is site-concordance between the main cancers observed in TCE-exposed  
12 humans and those observed in rodent studies—in particular, cancers of the kidney, liver, and  
13 lymphoid tissues.

14

#### **4.11.2.1.13. Summary of Additional Evidence on Biological Plausibility**

15 Additional evidence from toxicokinetic, toxicity, and mechanistic studies supports the  
16 biological plausibility of TCE carcinogenicity in humans.

17

#### **4.11.2.1.14. Toxicokinetics**

18 As described in Section 3, there is no evidence of major qualitative differences across  
19 species in TCE absorption, distribution, metabolism, and excretion. In particular, available  
20 evidence is consistent with TCE being readily absorbed via oral, dermal, and inhalation  
21 exposures, and rapidly distributed to tissues via systemic circulation. Extensive in vivo and in  
22 vitro data show that mice, rats, and humans all metabolize TCE via two primary pathways:  
23 oxidation by CYPs and conjugation with glutathione via GSTs. Several metabolites and  
24 excretion products from both pathways, including TCA, DCA, TCOH, TCOG, NAcDCVC, and  
25 DCVG, have been detected in blood and urine from exposed humans as well as from at least  
26 one rodent species. In addition, the subsequent distribution, metabolism, and excretion of TCE  
27 metabolites are qualitatively similar among species. Therefore, humans possess the metabolic  
28 pathways that produce the TCE metabolites thought to be involved in the induction of rat kidney  
29 and mouse liver tumors, and internal target tissues of both humans and rodents experience a  
30 similar mix of TCE and metabolites.

1 As addressed in further detail elsewhere (see Sections 3 and 5), examples of quantitative  
2 interspecies differences in toxicokinetics include differences in partition coefficients, metabolic  
3 capacity and affinity in various tissues, and plasma binding of the metabolite TCA. These and  
4 other differences are addressed through PBPK modeling, which also incorporates physiological  
5 differences among species (see Section 3.5), and are accounted for in the PBPK model-based  
6 dose-response analyses (see Section 5). Importantly, these quantitative differences affect only  
7 interspecies extrapolations of carcinogenic potency, and do not affect inferences as to the  
8 carcinogenic hazard for TCE. In addition, available data on toxicokinetic differences do not  
9 appear sufficient to explain interspecies differences in target sites of TCE carcinogenicity  
10 (discussed further in Section 5: Dose-Response Assessment).

#### 11 **4.11.2.1.15. Toxicity and mode of action**

12 Many different MOAs have been proposed for TCE-induced carcinogenesis. With  
13 respect to genotoxicity, although it appears unlikely that TCE, as a pure compound, causes point  
14 mutations, there is evidence for TCE genotoxicity with respect to other genetic endpoints, such  
15 as micronucleus formation (see Section 4.2.1.4.4). In addition, as discussed further below,  
16 several TCE metabolites have tested positive in genotoxicity assays. The MOA conclusions for  
17 specific target organs in laboratory animals are summarized below. Only in the case of the  
18 kidney is it concluded that the data are sufficient to support a particular MOA being operative.  
19 However, the available evidence do not indicate that qualitative differences between humans and  
20 test animals would preclude any of the hypothesized key events in rodents from occurring in  
21 humans.

22 For the kidney, the predominance of positive genotoxicity data in the database of  
23 available studies of TCE metabolites derived from GSH conjugation (in particular DCVC, see  
24 Section 4.2.5), together with toxicokinetic data consistent with their systemic delivery to and in  
25 situ formation in the kidney, supports the conclusion that a mutagenic MOA is operative in  
26 TCE-induced kidney tumors (see Section 4.4.7.1). Relevant data include demonstration of  
27 genotoxicity in available in vitro assays of GSH conjugation metabolites and reported kidney-  
28 specific genotoxicity after in vivo administration of TCE or DCVC. Mutagenicity is a well-  
29 established cause of carcinogenicity. While supporting the biological plausibility of this  
30 hypothesized MOA, available data on the *VHL* gene in humans or transgenic animals do not  
31 conclusively elucidate the role of *VHL* mutation in TCE-induced renal carcinogenesis.  
32 Cytotoxicity and compensatory cell proliferation, also presumed to be mediated through  
33 metabolites formed after GSH-conjugation of TCE, have also been suggested to play a role in the

1 MOA for renal carcinogenesis, as high incidences of nephrotoxicity have been observed in  
2 animals at doses that also induce kidney tumors. Human studies have reported markers for  
3 nephrotoxicity at current occupational exposures, although data are lacking at lower exposures.  
4 Toxicity is observed in both mice and rats, in some cases with nearly 100% incidence in all dose  
5 groups, but kidney tumors are only observed at low incidences in rats at the highest tested doses.  
6 Therefore, nephrotoxicity alone appears to be insufficient, or at least not rate-limiting, for rodent  
7 renal carcinogenesis, since maximal levels of toxicity are reached before the onset of tumors. In  
8 addition, nephrotoxicity has not been shown to be necessary for kidney tumor induction by TCE  
9 in rodents. In particular, there is a lack of experimental support for causal links, such as  
10 compensatory cellular proliferation or clonal expansion of initiated cells, between nephrotoxicity  
11 and kidney tumors induced by TCE. Furthermore, it is not clear if nephrotoxicity is one of  
12 several key events in a MOA, if it is a marker for an “upstream” key event (such as oxidative  
13 stress) that may contribute independently to both nephrotoxicity and renal carcinogenesis, or if it  
14 is incidental to kidney tumor induction. Moreover, while toxicokinetic differences in the GSH  
15 conjugation pathway, along with their uncertainty, are addressed through PBPK modeling, no  
16 data suggest that any of the proposed key events for TCE-induced kidney tumors rats are  
17 precluded in humans. Therefore, TCE-induced rat kidney tumors provide additional support for  
18 the convincing human evidence of TCE-induced kidney cancer, with mechanistic data supportive  
19 of a mutagenic MOA.

20 The strongest data supporting the hypothesis of a mutagenic MOA in either the lung or  
21 the liver are those demonstrating the genotoxicity of CH (see Section 4.2.4), which is produced  
22 in these target organs as a result of oxidative metabolism of TCE. It has been suggested that CH  
23 mutagenicity is unlikely to be the cause of TCE hepatocarcinogenicity because the  
24 concentrations required to elicit these responses are several orders of magnitude higher than  
25 achieved in vivo (Moore and Harrington-Brock, 2000). However, it is not clear how much of a  
26 correspondence is to be expected from concentrations in genotoxicity assays in vitro and  
27 concentrations in vivo, as reported in vivo CH concentrations are in whole liver homogenate  
28 while in vitro concentrations are in culture media. The use of i.p. administration, which leads to  
29 an inflammatory response, in many other in vivo genotoxicity assays in the liver and lung  
30 complicates the comparison with carcinogenicity data. Also, it is difficult with the available data  
31 to assess the contributions from genotoxic effects of CH along with those from the genotoxic and  
32 nongenotoxic effects of other oxidative metabolites (e.g., DCA and TCA). Therefore, while data  
33 are insufficient to conclude that a mutagenic MOA mediated by CH is operant, a mutagenic  
34 MOA in the liver or lung, either mediated by CH or by some other oxidative metabolite of TCE,  
35 cannot be ruled out.

1 A second MOA hypothesis for TCE-induced liver tumors involves activation of the  
2 PPAR $\alpha$  receptor. Clearly, in vivo administration of TCE leads to activation of PPAR $\alpha$  in rodents  
3 and likely does so in humans as well (based on in vitro data for TCE and its oxidative  
4 metabolites). However, the evidence as a whole does not support the view that PPAR $\alpha$  is the  
5 sole operant MOA mediating TCE hepatocarcinogenesis. Although metabolites of TCE activate  
6 PPAR $\alpha$ , the data on the subsequent elements in the hypothesized MOA (e.g., gene regulation,  
7 cell proliferation, apoptosis, and selective clonal expansion), while limited, indicate significant  
8 differences between PPAR $\alpha$  agonists such as Wy-14643 and TCE or its metabolites. For  
9 example, compared with other agonists, TCE induces transient as opposed to persistent increases  
10 in DNA synthesis; increases (or is without effect on), as opposed to decreases, apoptosis; and  
11 induces a different H-ras mutation frequency or spectrum. These data support the view that  
12 mechanisms other than PPAR $\alpha$  activation may contribute to these effects; besides PPAR $\alpha$   
13 activation, the other hypothesized key events are nonspecific, and available data (e.g., using  
14 knockout mice) do not indicate that they are solely or predominantly dependent on PPAR $\alpha$ . A  
15 second consideration is whether certain TCE metabolites (e.g., TCA) that activate PPAR $\alpha$  are the  
16 sole contributors to its carcinogenicity. As summarized above (see Section 4.11.1.3), TCA is not  
17 the only metabolite contributing to the observed noncancer effects of TCE in the liver. Other  
18 data also suggest that multiple metabolites may also contribute to the hepatic carcinogenicity of  
19 TCE. Liver phenotype experiments, particularly those utilizing immunostaining for c-Jun,  
20 support a role for both DCA and TCA in TCE-induced tumors, with strong evidence that TCA  
21 cannot solely account for the characteristics of TCE-induced tumors (e.g., Bull et al., 2002). In  
22 addition, H-ras mutation frequency and spectrum of TCE-induced tumors more closely  
23 resembles that of spontaneous tumors or of those induced by DCA, and were less similar in  
24 comparison to that of TCA-induced tumors. The heterogeneity of TCE-induced tumors is similar  
25 to that observed to be induced by a diversity carcinogens including those that do not activate  
26 PPAR $\alpha$ , and to that observed in human liver cancer. Taken together, the available data indicate  
27 that, rather than being solely dependent on a single metabolite (TCA) and/or molecular target  
28 (PPAR $\alpha$ ) multiple TCE metabolites and multiple toxicity pathways contribute to TCE-induced  
29 liver tumors.

30 Other considerations as well as new data published since the NRC (2006) review are also  
31 pertinent to the liver tumor MOA conclusions. It is generally acknowledged that, qualitatively,  
32 there are no data to support the conclusion that effects mediated by the PPAR $\alpha$  receptor that  
33 contribute to hepatocarcinogenesis would be biologically precluded in humans (Klaunig et al.,  
34 2003; NRC, 2006). It has, on the other hand, been argued that due to quantitative toxicokinetic  
35 and toxicodynamic differences, the hepatocarcinogenic effects of chemicals activating this

1 receptor are “unlikely” to occur in humans (Klaunig et al., 2003; NRC, 2006); however, several  
2 lines of evidence strongly undermine the confidence in this assertion. With respect to  
3 toxicokinetics, as discussed above, quantitative differences in oxidative metabolism are  
4 accounted for in PBPK modeling of available in vivo data, and do not support interspecies  
5 differences of a magnitude that would preclude hepatocarcinogenic effects based on  
6 toxicokinetics alone. With respect to the MOA proposed by Klaunig et al. (2003), recent  
7 experiments have demonstrated that PPAR $\alpha$  activation and the sequence of key events in the  
8 hypothesized MOA are not sufficient to induce hepatocarcinogenesis (Yang et al., 2007).  
9 Moreover, the demonstration that the PPAR $\alpha$  agonist DEHP induces tumors in PPAR $\alpha$ -null mice  
10 supports the view that the events comprising the hypothesized MOA are not necessary for liver  
11 tumor induction in mice by this PPAR $\alpha$  agonist (Ito et al., 2007). Therefore, several lines of  
12 evidence, including experiments published since the NRC (2006) review, call into question the  
13 scientific validity of using the PPAR $\alpha$  MOA hypothesis as the basis for evaluating the relevance  
14 to human carcinogenesis of rodent liver tumors (Guyton et al., 2009).

15 In summary, available data support the conclusion that the MOA for TCE-induced liver  
16 tumors in laboratory animals is not known. However, a number of qualitative similarities exist  
17 between observations in TCE-exposed mice and what is known about the etiology and induction  
18 of human hepatocellular carcinomas. Polyploidization, changes in glycogen storage, inhibition  
19 of GST-zeta, and aberrant DNA methylation status, which have been observed in studies of mice  
20 exposed to TCE or its oxidative metabolites, are all either clearly related to human  
21 carcinogenesis or are areas of active research as to their potential roles (PPAR $\alpha$  activation is  
22 discussed below). The mechanisms by which TCE exposure may interact with known risk  
23 factors for human hepatocellular carcinomas are not known. However, available data do not  
24 suggest that TCE exposure to mice results in liver tumors that are substantially different in terms  
25 of their phenotypic characteristics either from human hepatocellular carcinomas or from rodent  
26 liver tumors induced by other chemicals.

27 Comparing various other, albeit relatively nonspecific, tumor characteristics between  
28 rodent species and humans provides additional support to the biologic plausibility of TCE  
29 carcinogenicity. For example, in the kidney and the liver, the higher incidences of background  
30 and TCE-induced tumors in male rats and mice, respectively, as compared to females parallels  
31 the observed higher human incidences in males for these cancers (Ries et al., 2008). For the  
32 liver, while there is a lower background incidence of liver tumors in humans than in rodents, in  
33 the United States there is an increasing occurrence of liver cancer associated with several factors,  
34 including viral hepatitis, higher survival rates for cirrhosis, and possibly diabetes (reviewed in  
35 El-Serag, 2007). In addition, Leakey et al. (2003b) reported that increased body weight in

1 B6C3F1 mice is strongly associated with increased background liver tumor incidences, although  
2 the mechanistic basis for this risk factor in mice has not been established. Nonetheless, it is  
3 interesting that recent epidemiologic studies have suggested obesity, in addition to associated  
4 disorders such as diabetes and metabolic syndrome, as a risk factor for human liver cancer (El-  
5 Serag, 2007; El-Serag and Rudolph, 2007). Furthermore, the phenotypic and morphologic  
6 heterogeneity of tumors seen in the human liver is qualitatively similar to descriptions of mouse  
7 liver tumors induced by TCE exposure, as well as those observed from exposure to a variety of  
8 other chemical carcinogens. These parallels suggest similar pathways (e.g., for cell signaling) of  
9 carcinogenesis may be active in mice and humans and support the qualitative relevance of mouse  
10 models of liver to human liver cancer.

11 For mouse lung tumors, MOA hypotheses have centered on TCE metabolites produced  
12 via oxidative metabolism in situ. As discussed above, the hypothesis that the mutagenicity of  
13 reactive intermediates or metabolites (e.g., CH) generated during CYP metabolism contributes to  
14 lung tumors cannot be ruled out, although available data are inadequate to conclusively support  
15 this MOA. An alternative MOA has been posited involving other effects of such oxidative  
16 metabolites, particularly CH, including cytotoxicity and regenerative cell proliferation.  
17 Experimental support for this alternative hypothesis remains limited, with no data on proposed  
18 key events in experiments of duration 2 weeks or longer. While the data are inadequate to  
19 support this MOA hypothesis, the data also do not suggest that any proposed key events would  
20 be biologically plausible in humans. Furthermore, the focus of the existing MOA hypothesis  
21 involving cytotoxicity has been CH, and, as summarized above (see Section 4.11.1.5), other  
22 metabolites may contribute to respiratory tract noncancer toxicity or carcinogenicity. In sum, the  
23 MOA for mouse lung tumors induced by TCE is not known.

24 A MOA subsequent to in situ oxidative metabolism, whether involving mutagenicity,  
25 cytotoxicity, or other key events, may also be relevant to other tissues where TCE would  
26 undergo CYP metabolism. For instance, CYP2E1, oxidative metabolites, and protein adducts  
27 have been reported in the testes of rats exposed to TCE, and, in some rat bioassays, TCE  
28 exposure increased the incidence of rat testicular tumors. However, inadequate data exist to  
29 adequately define a MOA hypothesis for this tumor site.

### 4.11.3. Characterization of Factors Impacting Susceptibility

30 As discussed in more detail in Section 4.10, there is some evidence that certain  
31 populations may be more susceptible to exposure to TCE. Factors affecting susceptibility  
32 examined include lifestage, gender, genetic polymorphisms, race/ethnicity, preexisting health  
33 status, and lifestyle factors and nutrition status.

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 Examination of early lifestages includes exposures such as transplacental transfer  
2 (Beppu, 1968; Ghantous et al., 1986; Helliwell and Hutton, 1950; Laham, 1970; Withey and  
3 Karpinski, 1985) and breast milk ingestion (Fisher et al., 1997; Fisher et al., 1990; Hamada and  
4 Tanaka, 1995; Pellizzari et al., 1982), early lifestage-specific toxicokinetics, PBPK models  
5 (Fisher et al., 1989; Fisher et al., 1990), and differential outcomes in early lifestages such as  
6 developmental cardiac defects. Although there is more information on susceptibility to TCE  
7 during early lifestages than on susceptibility during later lifestages or for other populations with  
8 potentially increased susceptibility, there remain a number of uncertainties and data gaps  
9 regarding children's susceptibility. Improved PBPK modeling for using childhood parameters  
10 for early lifestages as recommended by the NRC (2006), and validation of these models will aid  
11 in determining how variations in metabolic enzymes affect TCE metabolism. In particular, the  
12 NRC states that it is prudent to assume children need greater protection than adults, unless  
13 sufficient data are available to justify otherwise (NRC, 2006). Because the weight of evidence  
14 supports a mutagenic MOA for TCE carcinogenicity in the kidney (see Section 4.4.7), and there  
15 is an absence of chemical-specific data to evaluate differences in carcinogenic susceptibility,  
16 early-life susceptibility should be assumed and the ADAFs should be applied, in accordance with  
17 the Supplemental Guidance (discussed further in Section 5).

18 Fewer data are available on later lifestages, although there is suggestive evidence to  
19 indicate that older adults may experience increased adverse effects than younger adults (Mahle et  
20 al., 2007; Rodriguez et al., 2007). In general, more studies specifically designed to evaluate  
21 effects in early and later lifestages are needed in order to more fully characterize potential life  
22 stage-related TCE toxicity.

23 Examination of gender-specific susceptibility includes toxicokinetics, PBPK models  
24 (Fisher et al., 1998), and differential outcomes. Gender differences observed are likely due to  
25 variation in physiology and exposure.

26 Genetic variation likely has an effect on the toxicokinetics of TCE. In particular,  
27 differences in CYP2E1 activity may affect susceptibility of TCE due to effects on production of  
28 toxic metabolites (Kim and Ghanayem, 2006; Lipscomb et al., 1997; Povey et al., 2001; Yoon et  
29 al., 2007). GST polymorphisms could also play a role in variability in toxic response (Brüning et  
30 al., 1997a; Wiesenhütter et al., 2007), as well as other genotypes, but these have not been  
31 sufficiently tested. Differences in genetic polymorphisms related to the metabolism of TCE have  
32 also been observed among various race/ethnic groups (Inoue et al., 1989; Sato et al., 1991b).

33 Preexisting diminished health status may alter the response to TCE exposure. Individuals  
34 with increased body mass may have an altered toxicokinetic response (Clewell et al., 2000;  
35 Davidson and Beliles, 1991; Lash et al., 2000a; McCarver et al., 1998; Monster et al., 1979;

1 Sato, 1993; Sato et al., 1991b) resulting in changes the internal concentrations of TCE or in the  
2 production of toxic metabolites. Other conditions, including diabetes and hypertension, are risk  
3 factors for some of the same health effects that have been associated with TCE exposure, such as  
4 renal cell carcinoma. However, the interaction between TCE and known risk factors for human  
5 diseases is not known, and further evaluation of the effects due to these factors is needed.

6 Lifestyle and nutrition factors examined include alcohol consumption, tobacco smoking,  
7 nutritional status, physical activity, and socioeconomic status. In particular, alcohol intake has  
8 been associated with metabolic inhibition (altered CYP2E1 expression) of TCE in both humans  
9 and experimental animals (Bardodej and Vyskocil, 1956; Barret et al., 1984; Kaneko et al.,  
10 1994a; Larson and Bull, 1989; McCarver et al., 1998; Muller et al., 1975; Nakajima et al., 1988;  
11 Nakajima et al., 1992a; Nakajima et al., 1990; Okino et al., 1991; Sato, 1993; Sato et al., 1991a;  
12 Sato and Nakajima, 1985; Sato et al., 1980, 1981, 1983; Stewart et al., 1974; White and Carlson,  
13 1981b). In addition, such factors have been associated with increased baseline risks for health  
14 effects associated with TCE, such as kidney cancer (e.g., smoking) and liver cancer (e.g., alcohol  
15 consumption). However, the interaction between TCE and known risk factors for human  
16 diseases is not known, and further evaluation of the effects due to these factors is needed.

17 In sum, there is some evidence that certain populations may be more susceptible to  
18 exposure to TCE. Factors affecting susceptibility examined include lifestage, gender, genetic  
19 polymorphisms, race/ethnicity, preexisting health status, and lifestyle factors and nutrition status.  
20 However, except in the case of toxicokinetic variability characterized using the PBPK model  
21 described in Section 3.5, there are inadequate chemical-specific data to quantify the degree of  
22 differential susceptibility due to such factors.

## 5. DOSE-RESPONSE ASSESSMENT

### 5.1. DOSE-RESPONSE ANALYSES FOR NONCANCER ENDPOINTS

1           Because of the large number of noncancer health effects associated with trichloroethylene  
2 (TCE) exposure and the large number of studies reporting on these effects, a screening process,  
3 described below, was used to reduce the number of endpoints and studies to those that would  
4 best inform the selection of the critical effects for the inhalation reference concentration (RfC)  
5 and oral reference dose (RfD).<sup>16</sup> The screening process helped identify the more sensitive  
6 endpoints for different types of effects within each health effect domain (e.g., different target  
7 systems) and provided information on the exposure levels that could contribute to the most  
8 sensitive effects, used for the RfC and RfD, as well as to additional noncancer effects as  
9 exposure increases. These more sensitive endpoints were also used to investigate the impacts of  
10 pharmacokinetic uncertainty and variability.

11           The general process used to derive the RfD and RfC was as follows (see Figure 5-1):

- 12
- 13       (1) Consider all studies described in Section 4 that report adverse noncancer health effects or  
14       markers for such effects and provide quantitative dose-response data<sup>17</sup>.
- 15       (2) Consider for each study/endpoint possible points of departure (PODs) on the basis of  
16       applied dose, with the order of preference being first a benchmark dose (BMD)<sup>18</sup> derived  
17       from empirical modeling of the dose-response data, then a no-observed-adverse-effect  
18       level (NOAEL), and lastly a lowest-observed-adverse-effect level (LOAEL).
- 19       (3) Adjust each POD by endpoint/study-specific “uncertainty factors” (UFs), accounting for  
20       uncertainties and adjustments in the extrapolation from the study conditions to conditions  
21       of human exposure, to derive candidate RfCs (cRfCs) or RfDs (cRfDs) intended to be  
22       protective for each endpoint (individually) on the basis of applied dose.


---

16 In U.S. EPA noncancer health assessments, the RfC (RfD) is an estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation (daily oral) exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious effects during a lifetime. It can be derived from a NOAEL, LOAEL, or benchmark concentration (dose), with uncertainty factors generally applied to reflect limitations of the data used.

17 Adequate dose-response data comprise, at a minimum, one exposure group and an appropriate control group, from which one can derive a LOAEL (or a NOAEL, if evidence of the effect is available from some other comparable study).

18 More precisely, it is the BMDL, i.e., the (one-sided) 95% lower confidence bound on the dose corresponding to the benchmark response for the effect that is used as the POD.

- 1 (4) Array the cRfCs and cRfDs across the following health effect domains: (1) neurotoxic  
 2 effects; (2) systemic (body weight) and organ toxicity (kidney, liver) effects;  
 3 (3) immunotoxic effects; (4) reproductive effects; and (5) developmental effects.



4  
 5 **Figure 5-1. Flow-chart of the process used to derive the RfD and RfC for**  
 6 **noncancer effects.**

- 7  
 8  
 9 (5) Select as candidate critical effects those endpoints with the lowest cRfCs or cRfDs,  
 10 within each of these effect domains, taking into account the confidence in each estimate.  
 11 When there are alternative estimates available for a particular endpoint, preference is  
 12 given to studies whose design characteristics (e.g., species, statistical power, exposure  
 13 level(s) and duration, endpoint measures) are better suited for determining the most  
 14 sensitive human health effects of chronic TCE exposure.
- 15 (6) For each candidate critical effect selected in step 5, use, to the extent possible, the  
 16 physiologically based pharmacokinetic (PBPK) model developed in Section 3.5 to  
 17 calculate an internal dose POD (idPOD) for plausible internal dose-metrics that were  
 18 selected on the basis of what is understood about the role of different TCE metabolites in  
 19 toxicity and the mode of action (MOA) for toxicity.

*This document is a draft for review purposes only and does not constitute Agency policy.*

- 1 (7) For each idPOD for each candidate critical effect, use the PBPK model to estimate  
2 interspecies and within-human pharmacokinetic variability (or just within-human  
3 variability for human-based PODs). The results of this calculation are 99<sup>th</sup> percentile  
4 estimates of the human equivalent concentration and human equivalent dose (HEC<sub>99</sub> and  
5 HED<sub>99</sub>) for each candidate critical effect.<sup>19</sup>
- 6 (8) Adjust each HEC<sub>99</sub> or HED<sub>99</sub> by endpoint/study-specific UFs (which, due to the use of  
7 the PBPK model, may differ from the UFs used in step 3) to derive a PBPK model-based  
8 candidate RfCs (p-cRfC) and RfD (p-cRfD) for each candidate critical effect.
- 9 (9) Characterize the uncertainties in the cRfCs, cRfDs, p-cRfCs, and p-cRfDs, with the  
10 inclusion of quantitative uncertainty analyses of pharmacokinetic uncertainty and  
11 variability as derived from the Bayesian population analysis using the PBPK model.
- 12 (10) Evaluate the most sensitive cRfCs, p-cRfCs, cRfDs, and p-cRfDs, taking into account the  
13 confidence in the estimates, to arrive at an RfC and RfD for TCE.  
14  
15

16 In contrast to the approach used in most assessments, in which the RfC and RfD are each  
17 based on a single critical effect, the final RfC and RfD for TCE were based on multiple critical  
18 effects that resulted in very similar candidate RfC and RfD values at the low end of the full range  
19 of values. This approach was taken here it is considered to provide more robust estimates of the  
20 RfC and RfD and because it highlights the multiple effects that are yielding very similar  
21 candidate values. The results of this process are summarized in the sections below, with  
22 technical details presented in Appendix F.  
23

### 5.1.1. Modeling Approaches and Uncertainty Factors for Developing Candidate Reference Values Based on Applied Dose

24 This section summarizes the general methodology used with all the TCE studies and  
25 endpoints for developing cRfCs and cRfDs on the basis of applied dose. A detailed discussion of  
26 the application of these approaches to the studies and endpoints for each health effect domain  
27 follows in the next section (see Section 5.1.2).

28 Standard adjustments<sup>20</sup> were made to the applied doses to obtain continuous inhalation  
29 exposures and daily average oral doses over the study exposure period (see Appendix F for

---

<sup>19</sup> The choice of the 99th percentile is discussed in Section 5.1.3.2.

<sup>20</sup> Discontinuous exposures (e.g., gavage exposures once a day, 5 days/week, or inhalation exposures for 5 days/week, 6 hours/day) were adjusted to the continuous exposure yielding the same cumulative exposure. For inhalation studies, these adjustments are equivalent to those recommended by U.S. EPA (1994b) for deriving a

1 details), except for effects for which there was sufficient evidence that the effect was more  
2 closely associated with administered exposure level (e.g., changes in visual function). The PODs  
3 based on applied dose in the following sections and in Appendix F are presented in terms of the  
4 adjusted doses (except where noted).

5 As described above, wherever possible,<sup>21</sup> benchmark dose modeling was conducted to  
6 obtain benchmark dose lower bounds (BMDLs) to serve as PODs for the cRfCs and cRfDs.  
7 Note that not all quantitative dose-response data are amenable to benchmark dose modeling. For  
8 example, while nonnumerical data (e.g., data presented in line or bar graphs rather than in tabular  
9 form) were considered for developing LOAELs or NOAELs, they were not used for benchmark  
10 dose modeling. In addition, sometimes the available models used do not provide an adequate fit  
11 to the data. For the benchmark dose modeling for this assessment, the EPA's BenchMark Dose  
12 Software (BMDS), which is freely available at [www.epa.gov/ncea/bmds](http://www.epa.gov/ncea/bmds), was used. For  
13 dichotomous responses, the log-logistic, multistage, and Weibull models were fitted. This subset  
14 of BMDS dichotomous models was used to reduce modeling demands, and these particular  
15 models were selected because, as a group, they have been found to be capable of describing the  
16 great majority of dose-response data sets, and specifically for some TCE data sets (Falk Filipsson  
17 and Victorin, 2003). For continuous responses, the distinct models available in BMDS—the  
18 power, polynomial, and Hill models—were fitted. For some reproductive and developmental  
19 data sets, two nested models (the nested logistic and the Rai and Van Ryzin models in BMDS<sup>22</sup>)  
20 were fitted to examine and account for potential intralitter correlations. Models with  
21 unconstrained power parameters <1 were considered when the dose-response relationship  
22 appeared supralinear, but these models often yield very low BMDL estimates and there was no  
23 situation in which an unconstrained model with a power parameter <1 was selected for the data  
24 sets modeled here. In most cases, a constrained model or the Hill model provided an adequate fit  
25 to such a dose-response relationship. In a few cases, the highest-dose group was dropped to  
26 obtain an improved fit to the lower-dose groups. See Appendix F for further details on model  
27 fitting and parameter constraints.

---

human equivalent concentration for a Category 3 gas for which the blood:air partition coefficient in laboratory animals is greater than that in humans (see Section 3.1 for discussion of the TCE blood:air partition coefficient).

<sup>21</sup>An exception was for the systemic effect of decreased body weight, which was observed in multiple chronic studies. Dose-response data were available, but the resources were not invested into modeling these data because the endpoint appeared a priori to be less sensitive than others and was not expected to be a critical effect.

<sup>22</sup>The BMDS v1.4 module for the National Center for Toxicological Research model failed with the TCE data sets.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 After fitting these models to the data sets, the following procedure for model selection was  
2 applied. First, models were rejected if the  $p$ -value for goodness of fit was  $<0.10$ .<sup>23</sup> Second,  
3 models were rejected if they did not appear to adequately fit the low-dose region of the  
4 dose-response relationship, based on an examination of graphical displays of the data and scaled  
5 residuals. If the BMDL estimates from the remaining models were “sufficiently close” (with a  
6 criterion of within twofold for “sufficiently close”), then the model with the lowest Akaike  
7 Information Criteria (AIC) was selected.<sup>24</sup> If the BMDL estimates from the remaining models  
8 are not sufficiently close, some model dependence is assumed. With no clear biological or  
9 statistical basis to choose among them, the lowest BMDL was chosen as a reasonable  
10 conservative estimate, unless the lowest BMDL appeared to be an outlier, in which case further  
11 judgments were made. Additionally, for continuous models, constant variance models were used  
12 for model parsimony unless the  $p$ -value for the test of homogenous variance was  $<0.10$ , in which  
13 case the modeled variance models were considered.

14 For benchmark response (BMR) selection, statistical and biological considerations were  
15 taken into account. For dichotomous responses, our general approach was to use 10% extra risk  
16 as the BMR for borderline or minimally adverse effects and either 5% or 1% extra risk for  
17 adverse effects, with 1% reserved for the most severe effects. For continuous responses, the  
18 preferred approach for defining the BMR is to use a preestablished cut-point for the minimal  
19 level of change in the endpoint at which the effect is generally considered to become biologically  
20 significant (e.g., there is substantial precedence for using a 10% change in weight for organ and  
21 body weights and a 5% change in weight for fetal weight). In the absence of a well-established  
22 cut-point, a BMR of 1 (control) standard deviation (SD) change from the control mean, or  
23 0.5 SD for effects considered to be more serious, was generally selected. For one neurological  
24 effect (traverse time), a doubling (i.e., twofold change) was selected because the control SD  
25 appeared unusually small.

26 After the PODs were determined for each study/endpoint, UFs were applied to obtain the  
27 cRfCs and cRfDs. Uncertainty factors are used to address differences between study conditions  
28 and conditions of human environmental exposure (U.S. EPA, 2002c). These include  
29

---

23In a few cases in which none of the models fit the data with  $p > 0.10$ , linear models were selected on the basis of an adequate visual fit overall.

24Akaike Information Criteria—a measure of information loss from a dose-response model that can be used to compare a set of models. Among a specified set of models, the model with the lowest AIC is considered the “best.” If two or more models share the lowest AIC, an average of the BMDLs could be used, but averaging was not used in this assessment because for the one occasion in which models shared the lowest AIC, a selection was made based on visual fit.

1  
2 (a) *Extrapolating from laboratory animals to humans:* If a POD is derived from  
3 experimental animal data, it is divided by an UF to reflect pharmacokinetic and  
4 pharmacodynamic differences that may make humans more sensitive than laboratory  
5 animals. For oral exposures, the standard value for the interspecies UF is 10, which  
6 breaks down (approximately) to a factor of three for pharmacokinetic differences  
7 (which is removed if the PBPK model is used) and a factor of three for  
8 pharmacodynamic differences. For inhalation exposures, ppm equivalence across  
9 species is generally assumed, in which case pharmacokinetic differences are  
10 considered to be negligible, and the standard value used for the interspecies UF is 3,  
11 which is ascribed to pharmacodynamic differences. These standard values were used  
12 for all the cRfCs and cRfDs based on laboratory animal data in this assessment.

13 (b) *Human (intraspecies) variability:* RfCs and RfDs apply to the human population,  
14 including sensitive subgroups, but studies rarely examine sensitive humans. Sensitive  
15 humans could be adversely affected at lower exposures than a general study  
16 population; consequently, PODs from general-population studies are divided by an  
17 UF to address sensitive humans. Similarly, the animals used in most laboratory  
18 animal studies are considered to be “typical” or “average” responders, and the human  
19 (intraspecies) variability UF is also applied to PODs from such studies to address  
20 sensitive subgroups. The standard value for the human variability UF is 10, which  
21 breaks down (approximately) to a factor of three for pharmacokinetic variability  
22 (which is removed if the PBPK model is used) and a factor of three for  
23 pharmacodynamic variability. This standard value was used for all the PODs in this  
24 assessment with the exception of the PODs for a few immunological effects that were  
25 based on data from a sensitive (autoimmune-prone) mouse strain; for those PODs, an  
26 UF of 3 was used for human variability.

27 (c) *Uncertainty in extrapolating from subchronic to chronic exposures:*<sup>25</sup> RfCs and RfDs  
28 apply to lifetime exposure, but sometimes the best (or only) available data come from  
29 less-than-lifetime studies. Lifetime exposure can induce effects that may not be  
30 apparent or as large in magnitude in a shorter study; consequently, a dose that elicits a  
31 specific level of response from a lifetime exposure may be less than the dose eliciting  
32 the same level of response from a shorter exposure period. Thus, PODs based on  
33 subchronic exposure data are generally divided by a subchronic-to-chronic UF, which  
34 has a standard value of 10. If there is evidence suggesting that exposure for longer  
35 time periods does not increase the magnitude of an effect, a lower value of three or  
36 one might be used. For some reproductive and developmental effects, chronic  
37 exposure is that which covers a specific window of exposure that is relevant for  
38 eliciting the effect, and subchronic exposure would correspond to an exposure that is  
39 notably less than the full window of exposure.

---

25 Rodent studies exceeding 90 days of exposure are considered chronic, and rodent studies with 4 weeks to 90 days of exposure are considered subchronic (see [http://www.epa.gov/iris/help\\_gloss.htm](http://www.epa.gov/iris/help_gloss.htm)).



1 (d) *Uncertainty in extrapolating from LOAELs to NOAELs:* PODs are intended to be  
2 estimates of exposure levels without appreciable risk under the study conditions so  
3 that, after the application of appropriate UFs for interspecies extrapolation, human  
4 variability, and/or duration extrapolation, the absence of appreciable risk is conveyed  
5 to the RfC or RfD exposure level to address sensitive humans with lifetime exposure.  
6 Under the NOAEL/LOAEL approach to determining a POD, however, adverse  
7 effects are sometimes observed at all study doses. If the POD is a LOAEL, it is  
8 divided by an UF to better estimate a NOAEL. The standard value for the  
9 LOAEL-to-NOAEL UF is 10, although sometimes a value of three is used if the  
10 effect is considered minimally adverse at the response level observed at the LOAEL  
11 or even one if the effect is an early marker for an adverse effect. For one POD in this  
12 assessment, a value of 30 was used for the LOAEL-to-NOAEL UF because the  
13 incidence rate for the adverse effect was  $\geq 90\%$  at the LOAEL.

14 (e) *Additional database uncertainties:* A database UF of 1, 3 or 10 is used to reflect the  
15 potential for deriving an underprotective toxicity value as a result of an incomplete  
16 characterization of the chemical's toxicity. No database UF was used in this  
17 assessment. See Section 5.1.4.1 for additional discussion of the uncertainties  
18 associated with the overall database for TCE.  
19

### 5.1.2. Candidate Critical Effects by Effect Domain

20 A large number of endpoints and studies were considered within each of the five health  
21 effect domains. A comprehensive list of all endpoints/studies that were considered for  
22 developing cRfCs and cRfDs is shown in Tables 5-1–5-5. These tables also summarize the  
23 PODs for the various study endpoints, the UFs applied, and the resulting cRfCs or cRfDs.  
24 Inhalation and oral studies are presented together so that the extent of the available data, as well  
25 as concordance or lack thereof in the responses across routes of exposure, is evident. In addition,  
26 the PBPK model developed in Section 3.5 will be applied to each candidate critical effect to  
27 develop a POD based on internal dose (idPOD); and subsequent extrapolation of the idPOD to  
28 pharmacokinetically sensitive humans is performed for both inhalation and oral human  
29 exposures, regardless of the route of exposure in the original study.

30 The sections below discuss the cRfCs and cRfDs developed from the effects and studies  
31 identified in the hazard characterization (see Section 4) that were suitable for the derivation of  
32 reference values (i.e., that provided quantitative dose-response data). Because the general  
33 approach for applying UFs was discussed above, the sections below only discuss the selection of  
34 particular UFs when there are study characteristics that require additional judgment as to the  
35 appropriate UF values and possible deviations from the standard values usually assigned.  
36

#### **5.1.2.1.1. Candidate Critical Neurological Effects on the Basis of Applied Dose**

1           As summarized in Section 4.11.1.1, both human and experimental animal studies have  
2 associated TCE exposure with effects on several neurological domains. The strongest  
3 neurological evidence of hazard is for changes in trigeminal nerve function or morphology and  
4 impairment of vestibular function. There is also evidence for effects on motor function; changes  
5 in auditory, visual, and cognitive function or performance; structural or functional changes in the  
6 brain; and neurochemical and molecular changes. Studies with numerical dose-response  
7 information are summarized in Table 5-1, with their corresponding cRfCs or cRfDs shown in  
8 Table 5-2. Because impairment of vestibular function occurs at higher exposures, such changes  
9 were not considered candidate critical effects; but the other neurological effect domains are  
10 represented. For trigeminal nerve effects, cRfC estimates based on two human studies are in a

**Table. 5-1 Summary of studies of neurological effects suitable for dose-response assessment**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>Trigeminal nerve effects</b>				<b>Section 4.3.1</b>
Mhiri et al. (2004)	Human phosphate industry workers (23 exposed, 23 controls)	Inhalation: Exposure ranged from 50–150 ppm, for 6 hr/d for at least 2 yr	Increased trigeminal somatosensory evoked potential latency	Table 4-20
Ruijten et al. (1991)	Human mail printing workers (31 exposed, 28 controls)	Inhalation: Mean cumulative exposure: 704 ppm × yrs; mean exposure duration: 16 yr	Increased latency in masseter reflex	Table 4-20
Barret et al. (1992)	Rat, Sprague-Dawley, female, 7/group	Oral: 0, 2,500 mg/kg; 1 dose/d, 5 d/wk, 10 wk	Increased internode length and fiber diameter in class A fibers of the trigeminal nerve observed with TCE treatment; changes in fatty acid composition.	Table 4-21
<b>Auditory effects</b>				<b>Section 4.3.2</b>
Rebert et al. (1991)	Rat, Long Evans, male, 10/group	Inhalation: 0, 1,600, and 3,200 ppm; 12 h/d, 12 wks	Significant decreases in BAER amplitude and an increase in latency of appearance of the initial peak (P1).	Table 4-23
Albee et al. (2006)	Rat, Fischer 344, male and female, 10/sex/group	Inhalation: 0, 250, 800, 2,500 ppm; 6 h/d, 5 d/wk, 13 wk	Mild frequency specific hearing deficits; focal loss of cochlear hair cells.	Table 4-23
Crofton and Zhao (1997)	Rat, Long Evans, male, 8–10/group	Inhalation: 0, 800, 1,600, 2,400, and 3,200 ppm; 6 h/d, 5 d/wk, 13 wk	Increased auditory thresholds as measured by BAERs for the 16 kHz tone	Table 4-23

This document is a draft for review purposes only and does not constitute Agency policy.

**Table. 5-1 Summary of studies of neurological effects suitable for dose-response assessment (continued)**

<b>Effect type</b> Study reference	<b>Species, strain (if appl.), sex, number used for dose-response assessment</b>	<b>Exposure(s) used for dose-response assessment</b>	<b>Endpoint(s) used for dose-response assessment</b>	<b>Section 4 Section/Table</b>
<b>Psychomotor effects</b>				<b>Section 4.3.6</b>
Waseem et al. (2001)	Rat, Wistar, male, 8/group	Inhalation: 0, 376 ppm for up to 180 d; 4 h/d, 5 d/wk	Changes in locomotor activity.	Table 4-31
Nunes et al. (2001)	Rat, Sprague-Dawley, male, 10/group	Oral: 0, 2,000 mg/kg/day; 7 d	Increased foot splay.	Table 4-30
Moser et al. (1995)	Rat, Fischer 344, female, 8/dose	Oral: 0, 150, 500, 1,500, and 5,000 mg/kg, 1 dose	Neuro-muscular impairment	Table 4-30
		0, 50, 150, 500, and 1,500 mg/kg/day, 14 d	Increased rearing activity	Table 4-30
<b>Visual function effects</b>				<b>Section 4.3.4</b>
Blain et al. (1994)	Rabbit, New Zealand albino, male, 6–8/group	Inhalation: 0, 350, 700 ppm; 4 h/d, 4 d/wk, 12 wk	Weekly electroretinograms (ERGs) and oscillatory potentials (OPs).	Table 4-26
<b>Cognitive effects</b>				<b>Section 4.3.5 and 4.3.6</b>
Kulig et al. (1987)	Rat, Wistar, male, 8/dose	Inhalation: 0, 500, 1,000, and 1,500 ppm; 16 h/d, 5 d/wk, 18 wk	Increased time in 2-choice visual discrimination test.	Table 4-31
Isaacson et al. (1990)	Rat, Sprague Dawley, male weanlings, 12/dose	Oral: (1) 0 mg/kg/day, 8 wk (2) 47 mg/kg/day, 4 wk + 0 mg/kg/day, 4 wk (3) 47 mg/kg/day, 4 wk + 0 mg/kg/day, 2 wk + 24 mg/kg/day, 2 wk	Demyelination of hippocampus	Table 4-28

**Table. 5-1 Summary of studies of neurological effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>Mood and sleep disorders</b>				<b>Section 4.3.7</b>
Albee et al. (2006)	Rat, Fischer 344, male and female, 10/sex/group	Inhalation: 0, 250, 800, 2,500 ppm; 6 h/d, 5 d/wk, 13 wk	Increased handling reactivity	Table 4-33
Arito et al. (1994)	Rat, Wistar, male, 5/group	Inhalation: 0,50, 100, and 300 ppm; 8 h/d, 5 d/wk, for 6 wk	Significant decreases in wakefulness.	Table 4-33
<b>Other neurological effects</b>				<b>Section 4.3.9</b>
Kjellstrand et al. (1987)	Rat, Sprague-Dawley, female	0, 300 ppm, 24 h/d, 24 d	Sciatic nerve regeneration was inhibited.	Table 4-36
	Mouse, NMRI, male	0, 150, or 300 ppm, 24 h/d, 24 d	Sciatic nerve regeneration was inhibited.	Table 4-36
Gash et al. (2008)	Rat, Fischer 344, male, 9/group	Oral: 0 and 1,000 mg/kg; 5 d/wk, 6 wk	Degeneration of dopamine-containing neurons in substantia nigra.	Table 4-35

**Table 5-2. Neurological effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
<b>Trigeminal nerve effects</b>												

Mhiri et al. (2004)	Human	LOAEL	40	1	1	10	10	1	100	0.40		Abnormal trigeminal somatosensory evoked potentials; preferred POD based on middle of reported range of 50–150 ppm.
	Human	LOAEL	6	1	1	10	10	1	100	0.06		Alternate POD based on U-TCA and Ikeda et al. (1972).
Ruijten et al. (1991)	Human	LOAEL	14	1	1	10	3	1	30	0.47		Trigeminal nerve effects; POD based on mean cumulative exposure and mean duration, UFl <sub>loael</sub> = 3 due to early marker effect and minimal degree of change.
Barret et al. (1992)	Rat	LOAEL	1,800	10	10	10	10	1	10,000 <sup>c</sup>		0.18	Morphological changes; uncertain adversity; some effects consistent with demyelination.
<b>Auditory effects</b>												
Rebert et al. (1991)	Rat	NOAEL	800	10	3	10	1	1	300	2.7		
Albee et al. (2006)	Rat	NOAEL	140	10	3	10	1	1	300	0.47		
Crofton and Zhao (1997)	Rat	BMDL	274	10	3	10	1	1	300	0.91		Preferred, due to better dose-response data, amenable to BMD modeling. BMR = 10dB absolute change.
<b>Psychomotor effects</b>												
Waseem et al. (2001)	Rat	LOAEL	45	1	3	10	3	1		0.45		Changes in locomotor activity; transient, minimal degree of adversity; no effect reported in same study for oral exposures (210 mg/kg/day).
Nunes et al. (2001)	Rat	LOAEL	2,000	10	10	10	3	1	3,000		0.67	↑ Foot splaying; minimal adversity.

**Table 5-2. Neurological effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs (continued)**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
Moser et al. (1995)	Rat	BMDL	248	3	10	10	1	1	300		0.83	↑ # rears (standing on hindlimbs); BMR = 1 SD change.
	Rat	NOAEL	500	3	10	10	1	1	300		1.7	↑ Severity score for neuromuscular changes.
<b>Visual function effects</b>												
Blain et al. (1994)	Rabbit	LOAEL	350	10	3	10	10	1	3,000	0.12		POD not adjusted to continuous exposure because visual effects more closely associated with administered exposure.
<b>Cognitive effects</b>												
Kulig et al. (1987)	Rat	NOAEL	500	1	3	10	1	1	30	17		↑ time in 2-choice visual discrim. test; test involves multiple systems but largely visual so not adjusted to continuous exposure.
Isaacson et al. (1990)	Rat	LOAEL	47	10	10	10	10	1	10,000 <sup>c</sup>		0.0047	Demyelination in hippocampus.
<b>Mood and sleep disorders</b>												
Albee et al. (2006)	Rat	NOAEL	140	10	3	10	1	1	300	0.47		Hyperactivity.
Arito et al. (1994)	Rat	LOAEL	12	3	3	10	10	1	1,000	0.012		Changes in wakefulness.
<b>Other neurological effects</b>												
Kjellstrand et al. (1987)	Rat	LOAEL	300	10	3	10	10	1	3,000	0.10		↓ regeneration of sciatic nerve.
	Mouse	LOAEL	150	10	3	10	10	1	3,000	0.050		↓ regeneration of sciatic nerve.
Gash et al. (2008)	Rat	LOAEL	710	10	10	10	10	1	10,000 <sup>c</sup>		0.071	Degeneration of dopaminergic neurons.

<sup>a</sup> Adjusted to continuous exposure unless otherwise noted. For inhalation studies, adjustments yield a POD that is a human equivalent concentration as recommended for a Category 3 gas in EPA (1994) in the absence of PBPK modeling. Same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup> Product of individual uncertainty factors.

**Table 5-2. Neurological effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs (continued)**

<sup>c</sup> EPA’s report on the RfC and RfD processes (U.S. EPA, 2002) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the subsequent application of the PBPK model for candidate critical effects will reduce the values of some of the individual UFs.

UF<sub>SC</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF. Shaded studies/endpoints were selected as candidate critical effects/studies.



1 similar range of 0.4–0.5 ppm (Mhiri et al., 2004; Ruijten et al., 1991). There remains some  
2 uncertainty as to the exposure characterization, as shown by the use of an alternative POD for  
3 Mhiri et al. (2004) based on urinary trichloroacetic acid (TCA) resulting in a fivefold smaller  
4 cRfC. However, the overall confidence in these estimates is relatively high because they are  
5 based on humans exposed under chronic or nearly chronic conditions. Other human studies (e.g.,  
6 Barret et al., 1984), while indicative of hazard, did not have adequate exposure information for  
7 quantitative estimates of an inhalation POD. A cRfD of 0.2 mg/kg/day was developed from the  
8 only oral study demonstrating trigeminal nerve changes, a subchronic study in rats (Barret et al.,  
9 1992). This estimate required multiple extrapolations with a composite uncertainty factor of  
10 10,000.<sup>26</sup>

11 For auditory effects, a high confidence cRfC of about 0.7 ppm was developed based on  
12 BMD modeling of data from Crofton and Zhao (1997); and cRfCs developed from two other  
13 auditory studies (Albee et al., 2006; Rebert et al., 1991) were within about fourfold. No oral data  
14 were available for auditory effects. For psychomotor effects, the available human studies (e.g.,  
15 Rasmussen et al., 1983) did not have adequate exposure information for quantitative estimates of  
16 an inhalation POD. However, a relatively high confidence cRfC of 0.5 ppm was developed from  
17 a study in rats (Waseem et al., 2001). Two cRfDs within a narrow range of 0.7–1.7 mg/kg/day  
18 were developed based on two oral studies reporting psychomotor effects (Moser et al., 1995;  
19 Nunes et al., 2001), although varying in degree of confidence.

20 For the other neurological effects, the estimated cRfCs and cRfDs were more uncertain,  
21 as there were fewer studies available for any particular endpoint, and the PODs from several  
22 studies required more adjustment to arrive at a cRfC or cRfD. However, the endpoints in these  
23 studies also tended to be indicative of more sensitive effects and, therefore, they need to be  
24 considered. The lower cRfCs fall in the range 0.01–0.1 ppm and were based on effects on visual  
25 function in rabbits (Blain et al., 1994), wakefulness in rats (Arito et al., 1994), and regeneration  
26 of the sciatic nerve in mice and rats (Kjellstrand et al., 1987). Of these, altered wakefulness  
27 (Arito et al., 1994) has both the lowest POD and the lowest cRfC. There is relatively high  
28 confidence in this study, as it shows a clear dose-response trend, with effects persisting  
29 postexposure. For the subchronic-to-chronic UF, a value of 3 was used because, even though it  
30 was just a 6-week study, there was no evidence of a greater impact on wakefulness following  
31 6 weeks of exposure than there was following 2 weeks of exposure at the LOAEL, although

---

<sup>26</sup>U.S. EPA's report on the RfC and RfD processes ([U.S. EPA, 2002c](#)) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the subsequent application of the PBPK model for candidate critical effects will reduce the values of some of the individual UFs.

1 there was an effect of repeated exposure on the postexposure period impacts of higher exposure  
2 levels. The cRfDs, in the range 0.005–0.07, were based on demyelination in the hippocampus  
3 (Isaacson et al., 1990) and degeneration of dopaminergic neurons (Gash et al., 2008), both in  
4 rats. In both these cases, adjusting for study design characteristics led to a composite uncertainty  
5 factor of 10,000,<sup>27</sup> so the confidence in these cRfDs is lower. However, no other studies of these  
6 effects are available.

7 In summary, although there is high confidence both in the hazard and in the cRfCs and  
8 cRfDs for trigeminal nerve, auditory, or psychomotor effects, the available data suggest that the  
9 more sensitive indicators of TCE neurotoxicity are changes in wakefulness, regeneration of the  
10 sciatic nerve, demyelination in the hippocampus and degeneration of dopaminergic neurons.  
11 Therefore, these more sensitive effects are considered the candidate critical effects for  
12 neurotoxicity, albeit with more uncertainty in the corresponding cRfCs and cRfDs. Of these  
13 more sensitive effects, for the reasons discussed above, there is greater confidence in the changes  
14 in wakefulness reported by Arito et al. (1994). In addition, trigeminal nerve effects are  
15 considered a candidate critical effect because this is the only type of neurological effect for  
16 which human data are available, and the POD for this effect is similar to that from the most  
17 sensitive rodent study (Arito et al., 1994, for changes in wakefulness). Between the two human  
18 studies of trigeminal nerve effects, Ruijten et al. (1991) is preferred for deriving noncancer  
19 reference values because its exposure characterization is considered more reliable.

#### 5.1.2.1.2. Candidate Critical Kidney Effects on the Basis of Applied Dose

21 As summarized in Section 4.11.1.2, multiple lines of evidence support TCE  
22 nephrotoxicity in the form of tubular toxicity, mediated predominantly through the glutathione  
23 (GSH) conjugation product dichlorovinyl cysteine (DCVC). Available human studies, while  
24 providing evidence of hazard, did not have adequate exposure information for quantitative  
25 estimates of PODs. Several studies in rodents, some of chronic duration, have shown  
26 histological changes, nephropathy, or increased kidney/body weight ratios. Studies with  
27 numerical dose-response information are summarized in Table 5-3, with their corresponding  
28 cRfCs or cRfDs shown in Table 5-4.

---

27U.S. EPA's report on the RfC and RfD processes ([U.S. EPA, 2002c](#)) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the subsequent application of the PBPK model for candidate critical effects will reduce the values of some of the individual UFs.

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 5-3. Summary of studies of kidney, liver, and body weight effects suitable for dose-response assessment**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>Histological changes in kidney</b>				<b>Section 4.4.4</b>
Maltoni (1986)	Rat, Sprague-Dawley, M, 116–124/group	Inhalation: 0, 100, 300, 600 ppm, 7 h/d, 5 d/wk, 104 wk exposure, observed for lifespan	Meganeucleocytosis	Table 4-49, Table 4-43
NTP (1990)	Rat, F344/N, male and female, 48–50/group	Oral: 0, 500, 1,000 mg/kg/day, 5 d/wk, 103 wk	Cytomegaly and karyomegaly	Table 4-45, Table 4-44
NCI (1976)	Mouse, B6C3F1, female, 20–50/group	Oral: 0, 869, 1,739 mg/kg/day, 5 d/wk, TWA during exposure period (78 wk), observed for 90 wk	Toxic nephrosis	Table 4-46, Table 4-44
NTP (1988)	Rat, Marshall, F, 44–50/group	Oral: 0, 500, 1,000 mg/kg/day, 5 d/wk, 104 wk	Toxic nephropathy	Table 4-47, Table 4-44
<b>↑ kidney/body weight ratio</b>				<b>Section 4.4.4</b>
Kjellstrand et al., (1983a)	Mouse, NMRI, M, 10–20/group	Inhalation: 0 (air), 37, 75, 150, 225, 300, 450, 900, 1,800, 3,600 ppm; continuous and intermittent exposures for 30–120 d	Increased kidney/body weight ratio	Table 4-43
Woolhiser et al. (2006)	Rat, Sprague-Dawley, F, 16/group	Inhalation: 0, 100, 300, and 1,000 ppm, 6 hr/d, 5 d/wk, for 4 wk	Increased kidney/body weight ratio	Table 4-43

**Table 5-3. Summary of studies of kidney, liver, and body weight effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>↑ liver/body weight ratio</b>				<b>Section 4.5.4.1</b>
Kjellstrand et al. (1983a)	Mouse, NMRI, M, 10–20/group	Inhalation: 0 (air), 37, 75, 150, 225, 300, 450, 900, 1,800, 3,600 ppm; continuous and intermittent exposures for 30–120 d	Increased liver/body weight ratio	Table 4-58
Woolhiser et al. (2006)	Rat, Sprague-Dawley, F, 16/group	Inhalation: 0, 100, 300, and 1,000 ppm, 6 hr/d, 5 d/wk, for 4 wk	Increased liver/body weight ratio	Table 4-58
Buben and O'Flaherty (1985)	Mouse, Swiss-Cox, 12–15/group	Oral: 0, 100, 200, 400, 800, 1,600, 2,400, and 3,200 mg/kg/day, 5 d/wk for 6 wk	Increased liver/body weight ratio	Table 4-54
<b>Decreased body weight</b>				
NTP (1990)	Mouse, B6C3F1, M, 48–50/group	Oral: 0, 1,000 mg/kg/day, 5 d/wk, 103 wk	Decreased body weight.	NA
NCI (1976)	Rat, Osborn-Mendel, M and F, 20–50/group	Oral: 0, 549, 1,097 mg/kg/day, 5 d/wk, TWA during exposure period (78 wk), observed at 110 wk	Decreased body weight.	NA

**Table 5-4. Kidney, liver, and body weight effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
<b>Histological changes in kidney</b>												
Maltoni (1986)	Rat	BMDL	40.2	1	3	10	1	1	30	1.3		meganucleocytosis; BMR = 10% extra risk
	Rat	BMDL	34	1	10	10	1	1	100		0.34	meganucleocytosis; BMR = 10% extra risk
NTP (1990)	Rat	LOAEL	360	1	10	10	10	1	1,000		0.36	cytomegaly and karyomegaly; considered minimally adverse, but UF <sub>loael</sub> = 10 due to high response rate (≥98%) at LOAEL; also in mice, but use NCI (1976) for that species
NCI (1976)	Mouse	LOAEL	620	1	10	10	30	1	3,000		0.21	toxic nephrosis; UF <sub>loael</sub> = 30 due to >90% response at LOAEL for severe effect
NTP (1988)	Rat	BMDL	9.45	1	10	10	1	1	100		0.0945	toxic nephropathy; female Marshall (most sensitive sex/strain); BMR = 5% extra risk
<b>↑ kidney/body weight ratio</b>												
Kjellstrand et al. (1983a)	Mouse	BMDL	34.7	1	3	10	1	1	30	1.2		BMR = 10% increase; 30 d, but 120 d @ 120 ppm not more severe so UF <sub>sc</sub> = 1; results are for males, which were slightly more sensitive, and yielded better fit to variance model
Woolhiser et al. (2006)	Rat	BMDL	15.7	1	3	10	1	1	30	0.52		BMR = 10% increase; UF <sub>sc</sub> = 1 based on Kjellstrand et al. (1983b) result
<b>↑ liver/body weight ratio</b>												
Kjellstrand et al. (1983a)	Mouse	BMDL	21.6	1	3	10	1	1	30	0.72		BMR = 10% increase; UF <sub>sc</sub> = 1 based on not more severe at 4 months
Woolhiser et al. (2006)	Rat	BMDL	25.2	1	3	10	1	1	30	0.84		BMR = 10% increase; UF <sub>sc</sub> = 1 based on Kjellstrand et al. (1983b) result
Buben and O'Flaherty (1985)	Mouse	BMDL	81.5	1	10	10	1	1	100		0.82	BMR = 10% increase; UF <sub>sc</sub> = 1 based on Kjellstrand et al. (1983b) result

This document is a draft for review purposes only and does not constitute Agency policy.

**Table 5-4. Kidney, liver, and body weight effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs (continued)**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
<b>Histological changes in kidney</b>												
NTP (1990)	Mouse	LOAEL	710	1	10	10	10	1	1,000		0.71	
NCI (1976)	Rat	LOAEL	360	1	10	10	10	1	1,000		0.36	Reflects several, but not all, strains/sexes.

<sup>a</sup> Adjusted to continuous exposure unless otherwise noted. For inhalation studies, adjustments yield a POD that is a human equivalent concentration as recommended for a Category 3 gas in U.S. EPA (1994b) in the absence of PBPK modeling. Same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup> Product of individual uncertainty factors.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.  
Shaded studies/endpoints were selected as candidate critical effects/studies.

1 The cRfCs developed from three suitable inhalation studies, one reporting  
2 meganucleocytosis in rats (Maltoni et al., 1986) and two others reporting increased kidney  
3 weights in mice (Kjellstrand et al., 1983a) and rats (Woolhiser et al., 2006),<sup>28</sup> are in a narrow  
4 range of 0.5–1.3 ppm. All three utilized BMD modeling and, thus, take into account statistical  
5 limitations of the Woolhiser et al. (2006) and Kjellstrand et al. (1983a) studies, such as  
6 variability in responses or the use of low numbers of animals in the experiment. The response  
7 used for kidney weight increases was the organ weight as a percentage of body weight, to  
8 account for any commensurate decreases in body weight, although the results did not generally  
9 differ much when absolute weights were used instead. Although the two studies reporting  
10 kidney weight changes were subchronic, longer-term experiments by Kjellstrand et al. (1983a)  
11 did not report increased severity, so no subchronic-to-chronic uncertainty factor was used in the  
12 derivation of the cRfC. The high response level of 73% at the lowest dose for  
13 meganucleocytosis in the chronic study of Maltoni et al. (1986) implies more uncertainty in the  
14 low-dose extrapolation. However, it is the only inhalation study that includes histopathological  
15 analysis, and it uses relatively high numbers of animals per dose group.

16 The suitable oral studies give cRfDs within a narrow range of 0.09–0.4 mg/kg/day, as  
17 shown in Table 5-4, although the degree of confidence in the cRfDs varies considerably. For  
18 cRfDs based on National Toxicology Program (NTP, 1990) and National Cancer Institute (NCI,  
19 1976) chronic studies in rodents, extremely high response rates of >90% precluded BMD  
20 modeling. An UF of 10 was applied for extrapolation from a LOAEL to a NOAEL in the NTP  
21 (1990) study because the effect (cytomegaly and karyomegaly), although minimally adverse, was  
22 observed at such a high incidence. An UF of 30 was applied for extrapolation from a LOAEL to  
23 a NOAEL in the NCI (1976) study because of the high incidence of a clearly adverse effect  
24 (toxic nephrosis). There is more confidence in the cRfDs based on meganucleocytosis reported  
25 in Maltoni et al. (1986) and toxic nephropathy NTP (1988), as BMD modeling was used to  
26 estimate BMDLs. Because these two oral studies measured somewhat different endpoints, but  
27 both were sensitive markers of nephrotoxic responses, they were considered to have similarly  
28 strong weight from a hazard perspective. For meganucleocytosis, a BMR of 10% extra risk was  
29 selected because the effect was considered to be minimally adverse. For toxic nephropathy, a  
30 BMR of 5% extra risk was used because toxic nephropathy is a severe toxic effect. This BMR  
31 required substantial extrapolation below the observed responses (about 60%); however, the  
32 response level seemed warranted for this type of effect and the ratio of the BMD to the BMDL

---

28Woolhiser et al. (2006) is an Organisation for Economic Co-operation and Development guideline immunotoxicity study performed by the Dow Chemical Company, certified by Dow as conforming to Good Laboratory Practices as published by the U.S. EPA for the Toxic Substances Control Act.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 was not large (1.56). Thus, from a dose-response extrapolation perspective, there is more  
2 confidence in Maltoni et al. (1986). However, the effect observed in NTP (1988) is more severe  
3 and therefore also merits consideration.

4 In summary, there is high confidence in the hazard and moderate confidence in the cRfCs  
5 and cRfDs for histopathological and weight changes in the kidney. These effects are considered  
6 to be candidate critical effects for several reasons. First, they appear to be the most sensitive  
7 indicators of toxicity that are available for the kidney. In addition, as discussed in Section 3.5,  
8 some pharmacokinetic data indicate substantially more production of GSH-conjugates thought to  
9 mediate TCE kidney effects in humans relative to rats and mice, although there is uncertainty in  
10 these data due to possible analytic errors. As discussed above, several studies are considered  
11 reliable for developing cRfCs and cRfDs for these endpoints. For histopathological changes, in  
12 general, the most sensitive were selected as candidate critical studies. These were the only  
13 available inhalation study (Maltoni et al., 1986), the Maltoni et al. (1986) and NTP (1988) oral  
14 studies in rats, and the NCI (1976) oral study in mice. For oral studies in rats, Maltoni et al.  
15 (1986) was considered in addition to NTP (1988), despite its having a higher cRfD, because of  
16 the much greater degree of low-dose extrapolation necessary for NTP (1988) and the excessive  
17 mortality present in that study. While the NCI (1976) study has even greater uncertainty, as  
18 discussed above, with a high response incidence at the POD that necessitates greater low-dose  
19 extrapolation, it is included to add a second species to the set of candidate critical effects. For  
20 kidney weight changes, both available studies were chosen as candidate critical studies.

### 5.1.2.1.3. Candidate Critical Liver Effects on the Basis of Applied Dose

22 As summarized in Section 4.11.1.3, while there is only limited epidemiologic evidence of  
23 TCE hepatotoxicity, TCE clearly leads to liver toxicity in laboratory animals, likely through its  
24 oxidative metabolites. Available human studies contribute to the overall weight of evidence of  
25 hazard, but did not have adequate exposure information for quantitative estimates of PODs. In  
26 rodent studies, TCE causes a wide array of hepatotoxic endpoints: increased liver weight, small  
27 transient increases in DNA synthesis, changes in ploidy, cytomegaly, increased nuclear size, and  
28 proliferation of peroxisomes. Increased liver weight (hepatomegaly, or specifically increased  
29 liver/body weight ratio) has been the most studied endpoint across a range of studies in both  
30 sexes of rats and mice, with a variety of exposure routes and durations. Hepatomegaly was  
31 selected as the critical liver effect for multiple reasons. First, it has been consistently reported in  
32 multiple studies in rats and mice following both inhalation and oral routes of exposure. In  
33 addition, it appears to accompany the other hepatic effects at the doses tested, and hence

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 constitutes a hepatotoxicity marker of similar sensitivity to the other effects. Finally, in several  
2 studies, there are good dose-response data for BMD modeling.

3 As shown in Table 5-4, cRfCs for hepatomegaly developed from the two most suitable  
4 subchronic inhalation studies (Kjellstrand et al., 1983a; Woolhiser et al., 2006), while in different  
5 species (rats and mice, respectively), are both based on similar PODs derived from BMD  
6 modeling, have the same composite uncertainty factor of 30, and result in similar cRfC estimates  
7 of about 0.8 ppm. The cRfD for hepatomegaly developed from the oral study of Buben and  
8 O’Flaherty (1985) in mice also was based on a POD derived from BMD modeling and resulted  
9 in a cRfD estimate of 0.8 mg/kg/day. Among the studies reporting liver weight changes  
10 (reviewed in Section 4.5 and Appendix E), this study had by far the most extensive  
11 dose-response data. The response used in each case was the liver weight as a percentage of body  
12 weight, to account for any commensurate decreases in body weight, although the results did not  
13 generally differ much when absolute weights were used instead.

14 There is high confidence in all these candidate reference values. BMD modeling takes  
15 into account statistical limitations such as variability in response or low numbers of animals and  
16 standardizes the response rate at the POD. Although the studies were subchronic, hepatomegaly  
17 occurs rapidly with TCE exposure, and the degree of hepatomegaly does not increase with  
18 chronic exposure (Kjellstrand et al., 1983a), so no subchronic-to-chronic uncertainty factor was  
19 used.

20 In summary, there is high confidence both in the hazard and the cRfCs and cRfDs for  
21 hepatomegaly. Hepatomegaly also appears to be the most sensitive indicator of toxicity that is  
22 available for the liver and is therefore considered a candidate critical effect. As discussed above,  
23 several studies are considered reliable for developing cRfCs and cRfDs for this endpoint, and,  
24 since they all indicated similar sensitivity but represented different species and/or routes of  
25 exposure, were all considered candidate critical studies.

#### 26 **5.1.2.1.4. Candidate Critical Body Weight Effects on the Basis of Applied Dose**

27 The chronic oral bioassays NCI (1976) and NTP (1990) reported decreased body weight  
28 with TCE exposure, as shown in Table 5-4. However, the lowest doses in these studies were  
29 quite high, even on an adjusted basis (see PODs in Table 5-4). These were not considered  
30 critical effects because they are not likely to be the most sensitive noncancer endpoints, and were  
31 not considered candidate critical effects.

#### 5.1.2.1.5. Candidate Critical Immunological Effects on the Basis of Applied Dose

1 As summarized in Section 4.11.1.4, the human and experimental animal studies of TCE  
2 and immune-related effects provide strong evidence for a role of TCE in autoimmune disease  
3 and in a specific type of generalized hypersensitivity syndrome, while there are fewer data  
4 pertaining to immunosuppressive effects. Available human studies, while providing evidence of  
5 hazard, did not have adequate exposure information for quantitative estimates of PODs. Several  
6 studies in rodents were available on autoimmune and immunosuppressive effects that were  
7 adequate for deriving cRfCs and cRfDs. Studies with numerical dose-response information are  
8 summarized in Table 5-5, with their corresponding cRfCs or cRfDs summarized in Table 5-6.

9 For decreased thymus weights, a cRfD from the only suitable study (Keil et al., 2009) is  
10 0.00035 mg/kg/day based on results from nonautoimmune-prone B6C3F1 mice, with a  
11 composite uncertainty factor of 1,000 for a POD that is a LOAEL (the dose-response relationship  
12 is sufficiently supralinear that attempts at BMD modeling did not result in adequate fits to these  
13 data). Thymus weights were not significantly affected in autoimmune prone mice in the same  
14 study, consistent with the results reported by Kaneko et al. (2000) in autoimmune-prone mice. In  
15 addition, Keil et al. (2009) and Peden-Adams et al. (2008) reported that for several  
16 immunotoxicity endpoints associated with TCE, the autoimmune-prone strain appeared to be less  
17 sensitive than the nonautoimmune prone B6C3F1 strain. In rats, Woolhiser et al. (2006) reported  
18 no significant change in thymus weights in the Sprague-Dawley (S-D) strain. These data are  
19 consistent with normal mice being sensitive to this effect as compared to autoimmune-prone  
20 mice or S-D rats, so the results of Keil et al. (2009) are not necessarily discordant with the other  
21 studies.

22 For autoimmune effects, the cRfC from the only suitable inhalation study (Kaneko et al.,  
23 2000) is 0.07 ppm. This study reported changes in immunoreactive organs (i.e., liver and spleen)  
24 in autoimmune-prone mice. BMD modeling was not feasible, so a LOAEL was used as the  
25 POD. The standard value of 10 was used for the LOAEL-to-NOAEL UF because the  
26 inflammation was reported to include sporadic necrosis in the hepatic lobules at the LOAEL, so  
27 this was considered an adverse effect. A value of 3 was used for the human (intraspecies)  
28 variability UF because the effect was induced in autoimmune-prone mice, a sensitive mouse  
29 strain for such an effect. The cRfDs from the oral studies (Cai et al., 2008; Griffin et al., 2000;  
30 Keil et al., 2009) spanned about a 100-fold range from 0.004–0.5 mg/kg/day. Each of the studies  
31 used different markers for autoimmune effects, which may explain the over 100-fold range of  
32 PODs (0.4–60 mg/kg/day). The most sensitive endpoint, reported by Keil et al. (2009), was  
33 increases in anti-dsDNA and anti-ssDNA antibodies, early markers for systemic lupus  
34 erythematosus (SLE), in B6C3F1 mice exposed to the lowest tested dose of 0.35 mg/kg/day,

1 yielding a cRfD of 0.004 mg/kg/day. Therefore, the results of Keil et al. (2009) are not  
2 discordant with the higher PODs and cRfDs derived from the other oral studies that examined  
3 more frank autoimmune effects.

**Table 5-5. Summary of studies of immunological effects suitable for dose-response assessment**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>↓ thymus weight</b>				<b>Section 4.6.2.3</b>
Keil et al. (2009)	Mouse, B6C3F1, Female, 10/group	Oral: 0, 1,400, or 14,000 ppb TCE (0, 0.35, or 3.5 mg/kg/day), 27 wk	Decreased thymus weights; decrease in thymus cellularity	Table 4-78
<b>Autoimmunity</b>				<b>Section 4.6.2.3</b>
Kaneko et al., (2000)	5/group	Inhalation: 0, 500, 1,000, or 2,000 ppm TCE, 4 h/d, 6 d/wk, 8 wk	Liver inflammation, splenomegaly and hyperplasia of lymphatic follicles	Table 4-78
Keil et al. (2009)	Mouse, B6C3F1, Female, 10/group	Oral: 0, 1,400, or 14,000 ppb TCE (0, 0.35, or 3.5 mg/kg/day), 27 wk	Increased anti-dsDNA and anti-ssDNA antibodies	Table 4-78
Griffin et al. (2000)	Mouse, MRL +/+, Female, 8/group	Oral: 0, 21, 100, or 400 mg/kg/day, 32 wk	Various signs of autoimmune hepatitis (serology, ex vivo assays of cultured splenocytes, clinical and histopathologic findings)	Table 4-78
Cai et al. (2008)	Mouse, MRL +/+, Female, 5/group	Oral: 0 or 60 mg/kg/day, 48 wk	Hepatic necrosis; hepatocyte proliferation; leukocyte infiltrate in the liver, lungs, and kidneys;	Table 4-78

**Table 5-5. Summary of studies of immunological effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>Immunosuppression</b>				<b>Section 4.6.2.1</b>
Woolhiser et al. (2006)	Rat, Sprague-Dawley, female, 16/group	Inhalation: 0, 100, 300, or 1,000 ppm, 6 h/d, 5 d/wk, 4-wk	Decreased plaque-forming cell assay response.	Table 4-76
Sanders et al. (1982b)	Mouse, CD-1, Female, 7-25/group	Oral: 0, 0.1, 1.0, 2.5, or 5.0 mg/mL (0, 18, 217, 393, or 660 mg/kg/day, from Tucker et al., 1982), 4 or 6 mo	Decreased humoral immunity, cell-mediated immunity, and bone marrow stem cell colonization.	Table 4-76

**Table 5-6. Immunological effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
<b>↓ thymus weight</b>												
Keil et al. (2009)	Mouse	LOAEL	0.35	1	10	10	10	1	1,000		0.00035	↓ thymus weight; corresponding decrease in total thymic cellularity reported at 10× higher dose
<b>Autoimmunity</b>												
Kaneko et al., (2000)	Mouse (MRL-lpr/ lpr)	LOAEL	70	10	3	3	10	1	1,000	0.070		Changes in immunoreactive organs—liver (incl. sporadic necrosis in hepatic lobules), spleen; UF <sub>h</sub> = 3 due to autoimmune-prone mouse
Keil et al. (2009)	Mouse	LOAEL	0.35	1	10	10	1	1	100		0.0035	↑ anti-dsDNA and anti-ssDNA Abs (early markers for SLE) (B6C3F1 mouse); UF <sub>loael</sub> = 1 due to early marker
Griffin et al. (2000)	Mouse (MRL+/+ )	BMDL	13.4	1	10	3	1	1	30		0.45	Various signs of autoimmune hepatitis; BMR = 10% extra risk for > minimal effects
Cai et al. (2008)	Mouse (MRL+/+ )	LOAEL	60	1	10	3	10	1	300		0.20	Inflammation in liver, kidney, lungs, and pancreas, which may lead to SLE-like disease; UF <sub>h</sub> = 3 due to autoimmune-prone mouse; UF <sub>loael</sub> = 10 since some hepatic necrosis
<b>Immunosuppression</b>												
Woolhiser et al. (2006)	Rat	BMDL	31.2	10	3	10	1	1	300	0.10		↓ PFC response; BMR = 1 SD change
Sanders et al. (1982b)	Mouse	NOAEL	190	1	10	10	1	1	100		1.9	↓ humoral response to sRBC; largely transient during exposure
	Mouse	LOAEL	18	1	10	10	3	1	300		0.060	↓ stem cell bone marrow recolonization (sustained); females more sensitive
	Mouse	LOAEL	18	1	10	10	3	1	300		0.060	↓ cell-mediated response to sRBC (largely transient during exposure); females more sensitive

<sup>a</sup> Adjusted to continuous exposure unless otherwise noted. For inhalation studies, adjustments yield a POD that is a human equivalent concentration as recommended for a Category 3 gas in U.S. EPA (1994b) in the absence of PBPK modeling. Same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup> Product of individual uncertainty factors.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.

Shaded studies/endpoints were selected as candidate critical effects/studies.

1 For immunosuppressive effects, the only suitable inhalation study (Woolhiser et al.,  
2 2006) gave a cRfC of 0.08 ppm. The cRfDs from the only suitable oral study (Sanders et al.,  
3 1982b) ranged from 0.06 mg/kg/day to 2 mg/kg/day, based on different markers for  
4 immunosuppression. Woolhiser et al. (2006) reported decreased plaque-forming cell (PFC)  
5 response in rats. Data from Woolhiser et al. (2006) were amenable to BMD modeling, but there  
6 is notable uncertainty in the modeling. First, it is unclear what should constitute the cut-point for  
7 characterizing the change as minimally biologically significant, so a BMR of 1 control SD  
8 change was used. In addition, the dose-response relationship is supralinear, and the highest  
9 exposure group was dropped to improve the fit to the low-dose data points. Nonetheless, the  
10 uncertainty in the BMD modeling is no greater than the uncertainty inherent in the use of a  
11 LOAEL or NOAEL. The more sensitive endpoints reported by Sanders et al. (1982b), both of  
12 which were in female mice exposed to a LOAEL of 18 mg/kg/day TCE in drinking water for  
13 4 months, were decreased cell-mediated response to sheep red blood cells (sRBC) and decreased  
14 stem cell bone recolonization, a sign of impaired bone marrow function. The cRfD based on  
15 these endpoints is 0.06 mg/kg/day, with a LOAEL-to-NOAEL UF of 3 because, although the  
16 immunosuppressive effects may not be adverse in and of themselves, multiple effects were  
17 observed suggesting potentially less resilience to an insult requiring an immunological response.

18 In summary, there is high qualitative confidence for TCE immunotoxicity and moderate  
19 confidence in the cRfCs and cRfDs that can be derived from the available studies. Decreased  
20 thymus weight reported at relatively low exposures in nonautoimmune-prone mice is a clear  
21 indicator of immunotoxicity (Keil et al., 2009), and is therefore considered a candidate critical  
22 effect. A number of studies have also reported changes in markers of immunotoxicity at  
23 relatively low exposures. Therefore, among markers for autoimmune effects, the more sensitive  
24 measures of autoimmune changes in liver and spleen (Kaneko et al., 2000) and increased  
25 anti-dsDNA and anti-ssDNA antibodies (Keil et al., 2009) are considered the candidate critical  
26 effects. Similarly, for markers of immunosuppression, the more sensitive measures of decreased  
27 PFC response (Woolhiser et al., 2006), decreased stem cell bone marrow recolonization, and  
28 decreased cell-mediated response to sRBC (both from Sanders et al., 1982b) are considered the  
29 candidate critical effects.

30

#### **5.1.2.1.6. Candidate Critical Respiratory Tract Effects on the Basis of Applied Dose**

31 As summarized in Section 4.11.1.5, available data are suggestive of TCE causing  
32 respiratory tract toxicity, based primarily on short-term studies in mice and rats. However, these  
33 studies are generally at high inhalation exposures and over durations of less than 2 weeks. Thus,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 these were not considered critical effects because such data are not necessarily indicators of  
2 longer-term effects at lower exposure and are not likely to be the most sensitive noncancer  
3 endpoints for chronic exposures. Therefore, cRfCs and cRfDs were not developed for them.  
4

#### 5.1.2.1.7. Candidate Critical Reproductive Effects on the Basis of Applied Dose

5 As summarized in Section 4.11.1.6, both human and experimental animal studies have  
6 associated TCE exposure with adverse reproductive effects. The strongest evidence of hazard is  
7 for effects on sperm and male reproductive outcomes, with evidence from multiple human  
8 studies and several experimental animal studies. There is also substantial evidence for effects on  
9 the male reproductive tract and male serum hormone levels, as well as evidence for effects on  
10 male reproductive behavior. There are fewer data and more limited support for effects on female  
11 reproduction. Studies with numerical dose-response information are summarized in Table 5-7,  
12 with their corresponding cRfCs or cRfDs summarized in Table 5-8.  
13

#### 5.1.2.1.8. Male reproductive effects (effects on sperm and reproductive tract)

14 A number of available studies have reported functional and structural changes in sperm  
15 and male reproductive organs and effects on male reproductive outcomes following TCE  
16 exposure (see Table 5-8). A cRfC of 0.014 ppm was derived based on hyperzoospermia reported  
17 in the available human study (Chia et al., 1996), but there is substantial uncertainty in this  
18 estimate due to multiple issues.<sup>29</sup> Among the rodent inhalation studies, the cRfC of 0.2 ppm  
19 based on increased abnormal sperm in the mouse reported by Land et al. (1981) is considered  
20 relatively reliable because it is based on BMD modeling rather than a LOAEL or NOAEL.  
21 However, increased sperm abnormalities do not appear to be the most sensitive effect, as

---

<sup>29</sup>Mean exposure estimates for the exposure groups were limited because they were defined in terms of ranges and because they were based on mean urinary TCA (mg/g creatinine). There is substantial uncertainty in the conversion of urinary TCA to TCE exposure level (see discussion of Mhiri et al. (2004), for neurotoxicity, above). In addition, there was uncertainty about the adversity of the effect being measured. While rodent evidence supports effects of TCE on sperm, and hyperzoospermia has reportedly been associated with infertility, the adversity of the hyperzoospermia (i.e., high sperm density) outcome measured in the Chia et al. (1996) study is unclear. Furthermore, the cut-point used to define hyperzoospermia in this study (i.e., >120 million sperm per mL ejaculate) is lower than some other reported cut-points, such as 200 and 250 million sperm/mL. A BMR of 10% extra risk was used on the assumption that this is a minimally adverse effect, but biological significance of this effect level is unclear.



1 Kumar et al. (2001b; 2000a; 2000b) reported a similar POD to be a LOAEL for reported multiple  
2 effects on sperm and testes, as well as altered testicular enzyme markers in the rat. Although  
3 there are greater uncertainties associated with the cRfC of 0.02 ppm for this effect and a  
4 composite UF of 3,000 was applied to the POD, the uncertainties are generally typical of those  
5 encountered in RfC derivations.

**Table 5-7. Summary of studies of reproductive effects suitable for dose-response assessment**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>Effects on sperm, male reproductive outcomes</b>				<b>Sections 4.8.1.1–4.8.1.2</b>
Chia et al. (1996)	Human, 85 men (37 low exposure, 48 high exposure)	Inhalation: Mean personal air TCE: 29.6 ppm; Mean U-TCA: 22.4 mg/g creatinine	Decreased normal sperm morphology and hyperzoospermia	Table 4-85
Land et al. (1981)	Mouse, C57BlxC3H (F1), M, 5 or 10/group	Inhalation: 0, 200, 2,000 ppm, 4 h/d, 5 d exposure, 23 d rest	Increased percent morphologically abnormal epididymal sperm	Table 4-86
Kan et al. (2007)	Mouse, CD-1, male, 4/group	Inhalation: 0 or 1,000 ppm, 6 h/d, 5 d/wk, 4 wk	Abnormalities of the head and tail in sperm located in the epididymal lumen.	Table 4-86
Xu et al. (2004)	Mouse, CD-1, male, 4–27/group	Inhalation: 0 or 1,000 ppm, 6 h/d, 5 d/wk, 6 wk	Decreased in vitro sperm-oocyte binding and in vivo fertilization.	Table 4-86
Kumar et al. (2000b)	Rat, Wistar, male, 12–13/group	Inhalation: 0 or 376 ppm, 4 h/d, 5 d/wk, 2–10 wk exposed, 2–8 wk unexposed.	Multiple sperm effects.	Table 4-86
Kumar et al., (2001b)	Rat, Wistar, male, 6/group	Inhalation: 0 or 376 ppm, 4 h/d, 5 d/wk, 24 wk	Multiple sperm effects, increasing severity from 12–24 weeks exposure.	Table 4-86

**Table 5-7. Summary of studies of reproductive effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
George et al. (1985)	Mouse, CD-1, male and female, 20 pairs/treatment group; 40 controls/sex	Oral: 0, 173, 362, or 737 mg/kg/day, Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased sperm motility in F0 and F1 males.	Table 4-87
DuTeaux et al., (2004b)	Rat, Sprague-Dawley, male, 3/group, or Simonson albino (UC Davis), male, 3/group	Oral: 0, 143, or 270 mg/kg/day, 14 d	Decreased ability of sperm to fertilize oocytes collected from untreated females. Oxidative damage to sperm membrane in head and mid-piece.	Table 4-87
<b>Male reproductive tract effects</b>				<b>Section 4.8.1.2</b>
Forkert et al. (2002)	Mouse, CD-1, male, 6/group	Inhalation: 0 or 1,000 ppm, 6 h/d, 5 d/wk, 19 d over 4 wk	Sloughing of epididymal epithelial cells.	Table 4-86
Kan et al. (2007)	Mouse, CD-1, male, 4/group	Inhalation: 0 or 1,000 ppm, 6 h/d, 5 d/wk, 1-4 wk	Degeneration and sloughing of epididymal epithelial cells (more severe by 4 wks). Vesiculation in cytoplasm, disintegration of basolateral cell membranes, sloughing of epithelial cells.	Table 4-86
Kumar et al. (2000b)	Rat, Wistar, male, 12-13/group	Inhalation: 0 or 376 ppm, 4 h/d, 5 d/wk, 2-10 wk exposed, 2-8 wk unexposed	Smaller, necrotic spermatogenic tubules.	Table 4-86

**Table 5-7. Summary of studies of reproductive effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
Kumar et al. (2001b)	Rat, Wistar, male, 6/group	Inhalation: 0 or 376 ppm, 4 h/d, 5 d/wk, 24 wk	Decreased testes weight, numbers of spermatogenic cells and spermatids, testes atrophy, smaller tubules devoid of spermatocytes and spermatids, hyperplastic Leydig cells, altered testicular enzyme markers. Increasing severity from 12–24 weeks exposure.	Table 4-86
George et al. (1985)	Mouse, CD-1, male and female, 20 pairs/treatment group; 40 controls/sex	Oral: 0, 173, 362, or 737 mg/kg/day, Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased testes and seminal vesicle weights in F0.	Table 4-87
George et al. (1986)	Rat, F334, males and female, 20 pairs/treatment group, 40 controls/sex	Oral: 0, 72, 186, or 389 mg/kg/day (estimated), Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Increased testes and epididymis weights in F0.	Table 4-87
<b>Female maternal weight gain</b>				<b>Section 4.8.3.2</b>
Carney et al. (2006)	Rat, Sprague-Dawley, females, 27 dams/group	Inhalation: 0, 50, 150, or 600 ppm, 6 h/d; GD 6–20	Decreased BW gain on GD 6–9	Table 4-96
Schwetz et al. (1975)	Rat, Sprague-Dawley, female, 20–35/group	Inhalation: 0 or 300 ppm, 7 h/d; GD 6–15	Decreased BW gain on GD 6–9	Table 4-96

**Table 5-7. Summary of studies of reproductive effects suitable for dose-response assessment (continued)**

<b>Effect type Study reference</b>	<b>Species, strain (if appl.), sex, number used for dose-response assessment</b>	<b>Exposure(s) used for dose-response assessment</b>	<b>Endpoint(s) used for dose-response assessment</b>	<b>Section 4 Section/Table</b>
Narotsky et al. (1995)	Rat, Fischer 344, females, 8–12 dams/group	Oral: 0, 10.1, 32, 101, 320, 475, 633, 844, or 1,125 mg/kg/day, GD 6–15	Decreased BW gain on GD 6–8 and 6–20.	Table 4-98
Manson et al. (1984)	Rat, Long-Evans, female, 23–25/group	Oral: 0, 10, 100, or 1,000 mg/kg/day, 6 wks: 2 wk pre mating, 1 wk mating period, GD 1–21	Decreased gestation BW gain.	Table 4-87
George et al. (1986)	Rat, F334, males and female, 20 pairs/treatment group, 40 controls/sex	Oral: 0, 72, 186, or 389 mg/kg/day (estimated), Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased term and postpartum dam BW in F0 and F1.	Table 4-87
<b>Female reproductive outcomes</b>				<b>Section 4.8.3.2</b>
Narotsky et al. (1995)	Rat, Fischer 344, females, 8–12 dams/group	Oral: 0, 10.1, 32, 101, 320, 475, 633, 844, or 1,125 mg/kg/day, GD 6–15	Delayed parturition.	Table 4-98
<b>Reproductive behavior</b>				<b>Section 4.8.1.2</b>
Zenick et al. (1984)	Rat, Long-Evans, male, 10/group	Oral: 0, 10, 100, or 1,000 mg/kg/day, 5 d/wk, 6 wk exposure, 4 wks recovery	Impaired copulatory performance.	Table 4-87

**Table 5-7. Summary of studies of reproductive effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
George et al. (1986)	Rat, F334, males and female, 20 pairs/treatment group, 40 controls/sex	Oral: 0, 72, 186, or 389 mg/kg/day (estimated), Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased F0 mating in cross-over mating trials.	Table 4-87
<b>Reproductive effects from exposure to both sexes</b>				<b>Section 4.8.1.2</b>
George et al. (1986)	Rat, F334, males and female, 20 pairs/treatment group, 40 controls/sex	Oral: 0, 72, 186, or 389 mg/kg/day (estimated), Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased F0 litters/pair and live F1 pups/litter.	Table 4-87

**Table 5-8. Reproductive effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
<b>Effects on sperm, male reproductive outcomes</b>												
Chia et al. (1996)	Human	BMDL	1.43	10	1	10	1	1	100	0.014		Hyperzoospermia; exposure estimates based on U-TCA from Ikeda et al. (1972); BMR = 10% extra risk
Land et al. (1981)	Mouse	BMDL	46.9	10	3	10	1	1	300	0.16		↑ abnormal sperm; BMR = 0.5 SD
Kan et al. (2007)	Mouse	LOAEL	180	10	3	10	10	1	3,000	0.060		↑ abnormal sperm; Land et al. (1981) cRfC preferred due to BMD modeling
Xu et al. (2004)	Mouse	LOAEL	180	10	3	10	10	1	3,000	0.060		↓ fertilization
Kumar et al. (2001b; 2000b)	Rat	LOAEL	45	10	3	10	10	1	3,000	0.015		Multiple sperm effects, increasing severity from 12–24 weeks
	Rat	LOAEL	45	1	3	10	10	1	300	0.15		Pre- and postimplantation losses; UF <sub>sc</sub> = 1 due to exposure covered time period for sperm development; higher response for preimplantation losses
George et al. (1985)	Mouse	NOAEL	362	1	10	10	1	1	100		3.6	↓ sperm motility
DuTeaux et al., (2004a)	Rat	LOAEL	141	10	10	10	10	1	10,000 <sup>c</sup>		0.014	↓ ability of sperm to fertilize in vitro
<b>Male reproductive tract effects</b>												
Forkert et al. (2002); Kan et al. (2007)	Mouse	LOAEL	180	10	3	10	10	1	3,000	0.060		Effects on epididymis epithelium
Kumar et al. (2001b; 2000b)	Rat	LOAEL	45	10	3	10	10	1	3,000	0.015		Testes effects, altered testicular enzyme markers, increasing severity from 12–24 weeks
George et al. (1985)	Mouse	NOAEL	362	1	10	10	1	1	100		3.6	↓ testis/seminal vesicle weights
George et al. (1986)	Rat	NOAEL	186	1	10	10	1	1	100		1.9	↑ testis/epididymis weights

**Table 5-8. Reproductive effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs (continued)**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
<b>Female maternal weight gain</b>												
Carney et al. (2006)	Rat	BMDL	10.5	1	3	10	1	1	30	0.35		↓ BW gain; BMR = 10% decrease
Schwetz et al. (1975)	Rat	LOAEL	88	1	3	10	10	1	300	0.29		↓ mat BW; Carney et al. (2006) cRfC preferred due to BMD modeling
Narotsky et al. (1995)	Rat	BMDL	108	1	10	10	1	1	100		1.1	↓ BW gain; BMR = 10% decrease
Manson et al. (1984)	Rat	NOAEL	100	1	10	10	1	1	100		1.0	↓ BW gain; Narotsky et al. (1995) preferred due to BMD modeling (different strain)
George et al. (1986)	Rat	NOAEL	186	1	10	10	1	1	100		1.9	↓ postpartum BW; Narotsky et al. (1995) cRfD preferred due to BMD modeling
<b>Female reproductive outcomes</b>												
Narotsky et al. (1995)	Rat	LOAEL	475	1	10	10	10	1	1,000		0.48	Delayed parturition
<b>Reproductive behavior</b>												
Zenick et al. (1984)	Rat	NOAEL	100	1	10	10	1	1	100		1.0	↓ copulatory performance in males
George et al. (1986)	Rat	LOAEL	389	1	10	10	10	1	1,000		0.39	↓ mating (both sexes exposed)
<b>Reproductive effects from exposure to both sexes</b>												
George et al. (1986)	Rat	BMDL	179	1	10	10	1	1	100		1.8	↓ # litters/pair; BMR = 0.5 SD
	Rat	BMDL	152	1	10	10	1	1	100		1.5	↓ live pups/litter; BMR = 0.5 SD

<sup>a</sup> Adjusted to continuous exposure unless otherwise noted. For inhalation studies, adjustments yield a POD that is a human equivalent concentration as recommended for a Category 3 gas in U.S. EPA (1994b) in the absence of PBPK modeling. Same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup> Product of individual UFs.

<sup>c</sup> EPA's report on the RfC and RfD processes (U.S. EPA, 2002c) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the subsequent application of the PBPK model for candidate critical effects will reduce the values of some of the individual UFs.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.



Shaded studies/endpoints were selected as candidate critical effects/studies.

This document is a draft for review purposes only and does not constitute Agency policy.

1 Standard values of 3, 10, and 10 were used for the interspecies UF, the human variability  
2 UF, and the LOAEL-to-NOAEL UF, respectively. In addition, although the study would have  
3 qualified as a chronic exposure study based on its duration of 24 weeks (i.e., >10% of lifetime),  
4 statistically significant decreases in testicular weight and in sperm count and motility were  
5 already observed from subchronic exposure (12 weeks) to the same TCE exposure concentration  
6 and these effects became more severe after 24 weeks of exposure. Moreover, several testicular  
7 enzyme markers associated with spermatogenesis and germ cell maturation had significantly  
8 altered activities after 12 weeks of exposure, with more severe alterations at 24 weeks, and  
9 histological changes were also observed in the testes at 12 weeks, with the testes being severely  
10 deteriorated by 24 weeks. Thus, since the single exposure level used was already a LOAEL from  
11 subchronic exposure, and the testes were even more seriously affected by longer exposures, a  
12 subchronic-to-chronic UF of 10 was applied.<sup>30</sup> Note that for the cRfC derived for pre and  
13 postimplantation losses reported by Kumar et al. (2000b), the subchronic-to-chronic UF was not  
14 applied because the exposure covered the time period for sperm development. This cRfC was  
15 0.2 ppm, similar to that derived from Land et al. (1981) based on BMD modeling of increases in  
16 abnormal sperm.

17 At a higher inhalation POD, Xu et al. (2004) reported decreased fertilization following  
18 exposure in male mice, and Forkert et al. (2002) and Kan et al. (2007) reported effects on the  
19 epididymal epithelium in male mice. Kan et al. (2007) reported degenerative effects on the  
20 epididymis as early as 1 week into exposure that became more severe at 4 weeks of exposure  
21 when the study ended; increases in abnormal sperm were also observed. As with the cRfC  
22 developed from the Kumar et al. (2001b; 2000a; 2000b), a composite UF of 3,000 was applied to  
23 these data, but the uncertainties are again typical of those encountered in RfC derivations.  
24 Standard values of 3 for the interspecies UF, 10 for the human variability UF, 10 for the  
25 LOAEL-to-NOAEL UF, and 10 for the subchronic-to-chronic UF were applied to each of the  
26 study PODs.

27 Among the oral studies, cRfDs derived for decreased sperm motility and changes in  
28 reproductive organ weights in rodents reported by George et al. (George et al., 1985, 1986) were  
29 relatively high (2–4 mg/kg/day), and these effects were not considered candidate critical effects.  
30 The remaining available oral study of male reproductive effects is DuTeaux et al. (2004a), which  
31 reported decreased ability of sperm from TCE-exposed rats to fertilize eggs in vitro. This effect

---

30Alternatively, the value of the LOAEL-to-NOAEL UF could have been increased above 10 to reflect the extreme severity of the effects at the LOAEL after 24 weeks; however, the comparison of the 12-week and 24-week results gives such a clear depiction of the progression of the effects, it was more compelling to frame the issue as a subchronic-to-chronic extrapolation issue.

1 occurred in the absence of changes in combined testes/epididymes weight, sperm concentration  
2 or motility, or histological changes in the testes or epididymes. DuTeaux et al. (2004a)  
3 hypothesize that the effect is due to oxidative damage to the sperm. A LOAEL was used as the  
4 POD, and the standard UF values of 10 were used for each of the UFs, i.e., the  
5 subchronic-to-chronic UF (14-day study; substantially less than the 70-day time period for sperm  
6 development), the interspecies UF for oral exposures, the human variability UF, and the  
7 LOAEL-to-NOAEL UF. The resulting composite UF was 10,000,<sup>31</sup> and this yielded a cRfD of  
8 0.01 mg/kg/day. The excessive magnitude of the composite UF, however, highlights the  
9 uncertainty in this estimate.

10 In summary, there is high qualitative confidence for TCE male reproductive tract toxicity  
11 and lower confidence in the cRfCs and cRfDs that can be derived from the available studies.  
12 Relatively high PODs are derived from several studies reporting less sensitive endpoints (George  
13 et al., 1985, 1986; 1981), and correspondingly higher cRfCs and cRfDs suggest that they are not  
14 likely to be critical effects. The studies reporting more sensitive endpoints also tend to have  
15 greater uncertainty. For the human study by Chia et al. (1996), as discussed above, there are  
16 uncertainties in the characterization of exposure and the adversity of the effect measured in the  
17 study. For the Kumar et al. (2001b; 2000a; 2000b), Forkert et al. (2002) and Kan et al. (2007)  
18 studies, the severity of the sperm and testes effects appears to be continuing to increase with  
19 duration even at the end of the study, so it is plausible that a lower exposure for a longer duration  
20 may elicit similar effects. For the DuTeaux et al. (2004a) study, there is also duration- and  
21 low-dose extrapolation uncertainty due to the short duration of the study in comparison to the  
22 time period for sperm development as well as the lack of a NOAEL at the tested doses. Overall,  
23 even though there are limitations in the quantitative assessment, there remains sufficient  
24 evidence to consider these to be candidate critical effects.

#### 5.1.2.1.9. Other reproductive effects

26 With respect to female reproductive effects, several studies reporting decreased maternal  
27 weight gain were suitable for deriving candidate reference values (see Table 5-8). The cRfCs  
28 from the two inhalation studies (Carney et al., 2006; Schwetz et al., 1975) yielded virtually the  
29 same estimate (0.3–0.4 ppm), although the Carney et al. (2006) result is preferred due to the use

---

31 U.S. EPA's report on the RfC and RfD processes ([U.S. EPA, 2002c](#)) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the subsequent application of the PBPK model for candidate critical effects will reduce the values of some of the individual UFs.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 of BMD modeling, which obviates the need for the 10-fold LOAEL-to-NOAEL UF used for  
2 Schwetz et al. (1975) (the other UFs, with a product of 30, were the same). The cRfDs for this  
3 endpoint from the three oral studies were within twofold of each other (1.1–1.9 mg/kg/day), with  
4 the same composite UFs of 100. The most sensitive estimate of Narotsky et al. (1995) is  
5 preferred due to the use of BMD modeling and the apparent greater sensitivity of the rat strain  
6 used.

7 With respect to other reproductive effects, the most reliable cRfD estimates of about  
8 2 mg/kg/day, derived from BMD modeling with composite UFs of 100, are based on decreased  
9 litters/pair and decreased live pups/litter in rats reported in the continuous breeding study of  
10 George et al. (1986). Both of these effects were considered severe adverse effects, so a BMR of  
11 a 0.5 control SD shift from the control mean was used. Somewhat lower cRfDs of  
12 0.4–1 mg/kg/day were derived based on delayed parturition in females (Narotsky et al., 1995),  
13 decreased copulatory performance in males (Zenick et al., 1984), and decreased mating for both  
14 exposed males and females in cross-over mating trials (George et al., 1986), all with composite  
15 UFs of 100 or 1,000 depending on whether a LOAEL or NOAEL was used.

16 In summary, there is moderate confidence both in the hazard and the cRfCs and cRfDs  
17 for reproductive effects other than the male reproductive effects discussed previously. While  
18 there are multiple studies suggesting decreased maternal body weight with TCE exposure, this  
19 systemic change may not be indicative of more sensitive reproductive effects. None of the  
20 estimates developed from other reproductive effects is particularly uncertain or unreliable.  
21 Therefore, delayed parturition (Narotsky et al., 1995) and decreased mating (George et al.,  
22 1986), which yielded the lowest cRfDs, were considered candidate critical effects. These effects  
23 were also included so that candidate critical reproductive effects from oral studies would not  
24 include only that reported by DuTeaux et al. (2004a), from which deriving the cRfD entailed a  
25 higher degree of uncertainty.

#### 26 **5.1.2.1.10. Candidate Critical Developmental Effects on the Basis of Applied Dose**

27 As summarized in Section 4.11.1.7, both human and experimental animal studies have  
28 associated TCE exposure with adverse developmental effects. Weakly suggestive epidemiologic  
29 data and fairly consistent experimental animal data support TCE exposure posing a hazard for  
30 increased prenatal or postnatal mortality and decreased pre or postnatal growth. In addition,  
31 congenital malformations following maternal TCE exposure have been reported in a number of  
32 epidemiologic and experimental animal studies. There is also some support for TCE effects on  
33 neurological and immunological development. Available human studies, while indicative of

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 hazard, did not have adequate exposure information for quantitative estimates of PODs, so only  
2 experimental animal studies are considered here. Studies with numerical dose-response  
3 information are summarized in Table 5-9, with their corresponding cRfCs or cRfDs summarized  
4 in Table 5-10.

5 For pre and postnatal mortality and growth, a cRfC of 0.06 ppm for resorptions,  
6 decreased fetal weight, and variations in skeletal development indicative of delays in ossification  
7 was developed based on the single available (rat) inhalation study considered (Healy et al., 1982)  
8 and utilizing the composite UF of 300 for an inhalation POD that is a LOAEL. The cRfDs for  
9 pre and postnatal mortality derived from oral studies were within about a 10-fold range of  
10 0.4–5 mg/kg/day, depending on the study and specific endpoint assessed. Of these, the estimate  
11 based on Narotsky et al. (1995) rat data was both the most sensitive and most reliable cRfD. The  
12 dose response for increased full-litter resorptions from this study is based on BMD modeling.  
13 Because of the severe nature of this effect, a BMR of 1% extra risk was used. The ratio of the  
14 resulting BMD to the BMDL was 5.7, which is on the high side, but given the severity of the  
15 effect and the low background response, a judgment was made to use 1% extra risk.

16 Alternatively, a 10% extra risk could have been used, in which case the POD would have been  
17 considered more analogous to a LOAEL than a NOAEL, and a LOAEL-to-NOAEL UF of 10  
18 would have been applied, ultimately resulting in the same cRfD estimate. The cRfDs for altered  
19 pre and postnatal growth developed from the oral studies ranged about 10-fold from  
20 0.8–8 mg/kg/day, all utilizing the composite UFs for the corresponding type of POD. The cRfDs  
21 for decreased fetal weight, both of which were based on NOAELs, were consistent, being about  
22 twofold apart (George et al., 1985; Narotsky et al., 1995). The cRfD based on postnatal growth  
23 at 21 days, reported in George et al. (1986), was lower and is preferred because it was based on  
24 BMD modeling. A BMR of 5% decrease in weight was used for postnatal growth at 21 days  
25 because decreases in weight gain so early in life were considered similar to effects on fetal  
26 weight.

27 For congenital defects, there is relatively high confidence in the cRfD for eye defects in  
28 rats reported in Narotsky et al. (1995), derived using a composite UF of 100 for BMD modeling  
29 in a study of duration that encompasses the full window of eye development. However, the most  
30 sensitive developmental effect by far was heart malformations in the rat reported by  
31 Johnson et al. (2003), yielding a cRfD estimate of 0.0002 mg/kg/day, also with a composite UF  
32 of 100. As discussed in detail in Section 4.8 and summarized in Section 4.11.1.7, although this  
33 study has important limitations, the overall weight of evidence supports an effect of TCE on  
34 cardiac development, and this is the only study of heart malformations available for conducting  
35 dose-response analysis. Individual data were kindly provided by Dr. Johnson (personal

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 communication from Paula Johnson, University of Arizona, to Susan Makris, EPA, 25 August  
2 2008), and, for analyses for which the pup was the unit of measure, BMD modeling was done

**Table 5-9. Summary of studies of developmental effects suitable for dose-response assessment**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>Pre and postnatal mortality</b>				<b>Section 4.8.1.2 and 4.8.3.2</b>
George et al. (1985)	Mouse, CD-1, male and female, 20 pairs/treatment group; 40 controls/sex	Oral: 0, 173, 362, or 737 mg/kg/day, Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Increase perinatal mortality (PND 0-21)	Table 4-87
Narotsky et al. (1995)	Rat, Fischer 344, females, 8-12 dams/group	Oral: 0, 10.1, 32, 101, 320, 475, 633, 844, or 1,125 mg/kg/day, GD 6-15	Increased resorptions, prenatal loss, and postnatal mortality	Table 4-98
Manson et al. (1984)	Rat, Long-Evans, female, 23-25/group	Oral: 0, 10, 100, or 1,000 mg/kg/day, 6 wk: 2 wk pre mating, 1 wk mating period, GD 1-21	Increased neonatal deaths on PND 1, 10, and 14.	Table 4-87
Healy et al. (1982)	Rat, Wistar, females, 31-32 dams/group	Inhalation: 0 or 100 ppm, 4 h/d; GD 8-21	Increased resorptions.	Table 4-96
<b>Pre and postnatal growth</b>				<b>Section 4.8.3.2</b>
Healy et al. (1982)	Rat, Wistar, females, 31-32 dams/group	Inhalation: 0 or 100 ppm, 4 h/d; GD 8-21	Decreased fetal weight, increased bipartite or absent skeletal ossification centers	Table 4-96

**Table 5-9. Summary of studies of developmental effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
Narotsky et al. (1995)	Rat, Fischer 344, females, 8–12 dams/group	Oral: 0, 10.1, 32, 101, 320, 475, 633, 844, or 1,125 mg/kg/day, GD 6–15	Decreased pup BW on PND 1 and 6.	Table 4-98
George et al. (1985)	Mouse, CD-1, male and female, 20 pairs/treatment group; 40 controls/sex	Oral: 0, 173, 362, or 737 mg/kg/day, Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased live birth weights, PND 4 pup body weights.	Table 4-87
George et al. (1986)	Rat, F334, males and female, 20 pairs/treatment group, 40 controls/sex	Oral: 0, 72, 186, or 389 mg/kg/day (estimated), Breeders exposed 1 wk pre mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased F1 BW on PND 4-80.	Table 4-87
<b>Congenital defects</b>				<b>Section 4.8.3.2</b>
Narotsky et al. (1995)	Rat, Fischer 344, females, 8–12 dams/group	Oral: 0, 10.1, 32, 101, 320, 475, 633, 844, or 1,125 mg/kg/day, GD 6–15	Increased incidence of eye defects.	Table 4-98
Johnson et al. (2003)	Rat, Sprague-Dawley, female, 9–13/group, 55 in control group	Oral: 0, 0.00045, 0.048, 0.218, or 129 mg/kg/day, GD 0–22	Increased percentage of abnormal hearts; increased percentage of litters with abnormal hearts.	Table 4-76



**Table 5-9. Summary of studies of developmental effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
<b>Developmental neurotoxicity</b>				<b>Sections 4.3.8.2 and 4.8.3.2</b>
George et al. (1986)	Rat, F334, males and female, 20 pairs/treatment group, 40 controls/sex	Oral: 0, 72, 186, or 389 mg/kg/day (estimated), Breeders exposed 1 wk pre-mating, then for 13 wk; pregnant females exposed throughout gestation (i.e., 18 wk total)	Decreased locomotor, as assessed by increased time required for pups to cross the first grid in open-field testing.	Table 4-34 and Table 4-98
Fredriksson et al. (1993)	Mouse, NMRI, male pups, 12 pups from 3–4 different litters/group	Oral: 0, 50, or 290 mg/kg/day, PND 10–16	Decreased rearing activity on PND 60.	Table 4-34 and Table 4-98
Taylor et al. (1985)	Rat, Sprague-Dawley, females, no. dams/group not reported	Oral: 0, 312, 625, or 1,250 mg/L (0, 45, 80, or 140 mg/kg/day estimated), dams (and pups) exposed from 14 d prior to mating until end of lactation	Increased exploratory behavior in 60- and 90-d old male rats (offspring).	Table 4-34 and Table 4-98
Isaacson and Taylor (1989)	Rat, Sprague-Dawley, females, 6 dams/group	Oral: 0, 4.0, or 8.1 mg/d (0, 15, or 32 mg/kg/day estimated) <sup>a</sup> , dams (and pups) exposed from 14 d prior to mating until end of lactation.	Decreased myelinated fibers in the stratum lacunosum-moleculare of pups; decreased myelin in the hippocampus.	Table 4-34 and Table 4-98
<b>Developmental immunotoxicity</b>				<b>Section 4.8.3.2</b>

**Table 5-9. Summary of studies of developmental effects suitable for dose-response assessment (continued)**

Effect type Study reference	Species, strain (if appl.), sex, number used for dose-response assessment	Exposure(s) used for dose-response assessment	Endpoint(s) used for dose-response assessment	Section 4 Section/Table
Peden-Adams et al. (2006)	Mouse, B6C3F1, dams and both sexes offspring, 5 dams/group; 5–7 pups/group at 3 wks; 4–5 pups/sex/group at 8 wks	Oral: 0, 1,400, or 14,000 ppb in water (0, 0.37, or 3.7 mg/kg/day estimated), parental mice and/or offspring exposed during mating, and from GD 0 thru 3 or 8 wk of age	Suppressed PFC responses in males and in females. Delayed hypersensitivity response increased at 8 wks of age in females. Splenic cell population decreased in 3 wk old pups. Increased thymic T-cells at 8 wks of age. Delayed hypersensitivity response increased at 8 wks of age in males and females	Table 4-98

<sup>a</sup>The Isaacson and Taylor (1989) and Taylor et al. (1985) studies report different doses despite identical study designs and administered concentrations, both studies taking TCE degradation into account. Taylor et al. (1985) report total consumption of 646 mg, 1102 mg, and 1991 mg TCE for rats exposed to 312, 625, and 1250 mg TCE/L drinking water, respectively. Dividing by the 56 days of exposure and the average 250 g per rat for female SD rats of those ages yields estimated doses of roughly 45, 80, and 140 mg/kg/day, respectively. Isaacson and Taylor (1989) report average doses of TCE of 4.0 and 8.1 mg/day corresponding to exposures of 312 and 625 mg TCE/L drinking water, respectively. Dividing by the average 250 g per rat yields estimated doses of 16 and 32 mg/kg/day, respectively. Thus the estimated doses for Taylor et al. (1985) are nearly 3-times higher than those for Isaacson and Taylor (1989), for reasons unknown.

**Table 5-10. Developmental effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
<b>Pre and postnatal mortality</b>												
George et al. (1985)	Mouse	NOAEL	362	1	10	10	1	1	100		3.6	↑ perinatal mortality
Narotsky et al. (1995)	Rat	LOAEL	475	1	10	10	10	1	1,000		0.48	Postnatal mortality; Manson et al. (1984) cRfD preferred for same endpoint due to NOAEL vs. LOAEL
Manson et al. (1984)	Rat	NOAEL	100	1	10	10	1	1	100		1.0	↑ neonatal death
Healy et al. (1982)	Rat	LOAEL	17	1	3	10	10	1	300	0.057		Resorptions
Narotsky et al. (1995)	Rat	BMDL	469	1	10	10	1	1	100		4.7	Prenatal loss; BMR = 1% extra risk
	Rat	BMDL	32.2	1	10	10	1	1	100		0.32	Resorptions; BMR = 1% extra risk
<b>Pre and postnatal growth</b>												
Healy et al. (1982)	Rat	LOAEL	17	1	3	10	10	1	300	0.057		↓ fetal weight; skeletal effects
Narotsky et al. (1995)	Rat	NOAEL	844	1	10	10	1	1	100		8.4	↓ fetal weight
George et al. (1985)	Mouse	NOAEL	362	1	10	10	1	1	100		3.6	↓ fetal weight
George et al. (1986)	Rat	BMDL	79.7	1	10	10	1	1	100		0.80	↓ BW at d21; BMR = 5% decrease
<b><u>Congenital defects</u></b>												
Narotsky et al. (1995)	Rat	BMDL	60	1	10	10	1	1	100		0.60	Eye defects; low BMR (1%), but severe effect and low bkgd. rate (<1%)
Johnson et al. (2003)	Rat	BMDL	0.0146	1	10	10	1	1	100		0.00015	Heart malformations (litters); BMR = 10% extra risk (only ~1/10 from each litter affected); highest-dose group (1,000-fold higher than next highest) dropped to improve model fit.

**Table 5-10. Developmental effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs (continued)**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
	Rat	BMDL	0.0207	1	10	10	1	1	100		0.00021	Heart malformations (pups); BMR = 1% extra risk; preferred due to accounting for intralitter effects via nested model and pups being the unit of measure; highest-dose group (1,000-fold higher than next highest) dropped to improve model fit
<b>Developmental neurotoxicity</b>												
George et al. (1986)	Rat	BMDL	72.6	1	10	10	1	1	100		0.73	↓ locomotor activity; BMR = doubling of traverse time; results from females (males similar with BMDL = 92)
Fredriksson et al. (1993)	Mouse	LOAEL	50	3	10	10	10	1	3,000		0.017	↓ rearing postexposure; pup gavage dose; No effect at tested doses on locomotion behavior; UF <sub>sc</sub> = 3 because exposure only during PND 10–16
Taylor et al. (1985)	Rat	LOAEL	45	1	10	10	10	1	1,000		0.045	↑ exploration postexposure; estimated dam dose; Less sensitive than Isaacson and Taylor (1989), but included because exposure is preweaning, so can utilize PBPK model
Isaacson and Taylor (1989)	Rat	LOAEL	16	1	10	10	10	1	1,000		0.016	↓ myelination in hippocampus; estimated dam dose
<b>Developmental immunotoxicity</b>												

**Table 5-10. Developmental effects in studies suitable for dose-response assessment, and corresponding cRfCs and cRfDs (continued)**

Effect type Supporting studies	Species	POD type	POD <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC (ppm)	cRfD (mg/kg/day)	Effect; comments
Peden-Adams et al. (2006)	Mouse	LOAEL	0.37	1	10	10	10	1	1,000		0.00037	↓ PFC, ↑DTH; POD is estimated dam dose (exposure throughout gestation and lactation + to 3 or 8 wks of age); UF LOAEL = 10 since ↑ DTH and also multiple immuno. effects

<sup>a</sup>Adjusted to continuous exposure unless otherwise noted. For inhalation studies, adjustments yield a POD that is a human equivalent concentration as recommended for a Category 3 gas in U.S. EPA (1994b) in the absence of PBPK modeling. Same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup>Product of individual uncertainty factors.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.

Shaded studies/endpoints were selected as candidate critical effects/studies.

This document is a draft for review purposes only and does not constitute Agency policy.

٤٣      DRAFT      DO NOT CITE OR QUOTE

1 using nested models because accounting for the intralitter correlation improved model fit. For  
2 these latter analyses, a 1% extra risk of a pup having a heart malformation was used as the BMR  
3 because of the severity of the effect, since, for example, some of the types of malformations  
4 observed could have been fatal. The ratio of the resulting BMD to the BMDL was about three.

5 For developmental neurotoxicity, the cRfD estimates based on the four oral studies span a  
6 wide range from 0.02–0.8 mg/kg/day. The most reliable estimate, with a composite UF of 100,  
7 is based on BMD modeling of decreased locomotor activity in rats reported in George et al.  
8 (1986), although a nonstandard BMR of a twofold change was selected because the control SD  
9 appeared unusually small. The cRfDs developed for decreased rearing postexposure in mice  
10 (Fredriksson et al., 1993), increased exploration postexposure in rats (Taylor et al., 1985) and  
11 decreased myelination in the hippocampus of rats (Isaacson and Taylor, 1989), while being more  
12 than 10-fold lower, are all within a threefold range of 0.02–0.05 mg/kg/day. Importantly, there  
13 is some evidence from adult neurotoxicity studies of TCE causing demyelination, so there is  
14 additional biological support for the latter effect. There is greater uncertainty in the Fredriksson  
15 et al. (1993), the cRfD for which utilized a subchronic-to-chronic UF of three rather than one,  
16 because exposure during postnatal day (PND) 10–16 does not cover the full developmental  
17 window (Rice and Barone, 2000). The cRfDs derived from Taylor et al. (1985) and (Isaacson  
18 and Taylor, 1989) used the composite UF of 1,000 for a POD that is a LOAEL. While there is  
19 greater uncertainty in these endpoints, none of the uncertainties is particularly high, and they also  
20 appear to be more sensitive indicators of developmental neurotoxicity than that from George  
21 et al. (1986).

22 A cRfD of 0.0004 mg/kg/day was developed from the study (Peden-Adams et al., 2006)  
23 that reported developmental immunotoxicity. The main effects observed were significantly  
24 decreased PFC response and increased delayed-type hypersensitivity. The data on these effects  
25 were kindly provided by Dr. Peden-Adams (personal communication from Margie  
26 Peden-Adams, Medical University of South Carolina, to Jennifer Jinot, EPA, 26 August 2008);  
27 however, the dose-response relationships were sufficiently supralinear that attempts at BMD  
28 modeling did not result in adequate fits to these data. Thus, the LOAEL was used as the POD.  
29 Although decreased PFC response may not be considered adverse in and of itself, a  
30 LOAEL-to-NOAEL UF of 10 was used because of the increased delayed-type hypersensitivity at  
31 the same dose. While there is uncertainty in this estimate, it is notable that decreased PFC  
32 response was also observed in an immunotoxicity study in adult animals (Woolhiser et al., 2006),  
33 lending biological plausibility to the effect.

34 In summary, there is moderate-to-high confidence both in the hazard and the cRfCs and  
35 cRfDs for developmental effects of TCE. It is also noteworthy that the PODs for the more

1 sensitive developmental effects were similar to or, in most cases, lower than the PODs for the  
2 more sensitive reproductive effects, suggesting that developmental effects are not a result of  
3 paternal or maternal toxicity. Among inhalation studies, cRfCs were only developed for effects  
4 in rats reported in Healy et al. (1982), so the effects of resorptions, decreased fetal weight, and  
5 delayed skeletal ossification were considered candidate critical developmental effects. Because  
6 resorptions were also reported in oral studies, the most sensitive (rat) oral study (and most  
7 reliable for dose-response analysis) of Narotsky et al. (1995) was also selected as a candidate  
8 critical study for this effect. The confidence in the oral studies and candidate reference values  
9 developed for more sensitive endpoints is more moderate, but still sufficient for consideration as  
10 candidate critical effects. The most sensitive endpoints by far are the increased fetal heart  
11 malformations in rats reported by Johnson et al. (2003) and the developmental immunotoxicity in  
12 mice reported by Peden-Adams et al. (2006), and these are both considered candidate critical  
13 effects. Neurodevelopmental effects are a distinct type among developmental effects. Thus, the  
14 next most sensitive endpoints of decreased rearing postexposure in mice (Fredriksson et al.,  
15 1993), increased exploration postexposure in rats (Taylor et al., 1985) and decreased myelination  
16 in the hippocampus of rats (Isaacson and Taylor, 1989) are also considered candidate critical  
17 effects.

18

#### **5.1.2.1.11. Summary of cRfCs, cRfDs, and Candidate Critical Effects**

19 An overall summary of the cRfCs, cRfDs, and candidate critical effects across the health  
20 effect domains is shown in Tables 5-11–5-12. These tables present, for each type of noncancer  
21 effect, the relative ranges of the cRfC and cRfD developed for the different endpoints. The  
22 candidate critical effects selected above for each effect domain are shown in bold. As discussed  
23 above, these effects were generally selected to represent the most sensitive endpoints, across  
24 species where possible. From these candidate critical effects, candidate reference values based  
25 on internal dose-metrics from the PBPK model (p-cRfCs and p-cRfDs) were developed where  
26 possible. Application of the PBPK model is discussed in the next section.

27

#### **5.1.3. Application of Physiologically Based Pharmacokinetic (PBPK) Model to Inter- and Intraspecies Extrapolation for Candidate Critical Effects**

28 For the candidate critical effects, the use of PBPK modeling of internal doses could  
29 justify, where appropriate, replacement of the uncertainty factors for pharmacokinetic inter and  
30 intraspecies extrapolation. For more details on PBPK modeling used to estimate levels of

*This document is a draft for review purposes only and does not constitute Agency policy.*



- 1 dose-metrics corresponding to different exposure scenarios in rodents and humans, as well as a
- 2 qualitative discussion of the uncertainties and limitations of the model, see Section 3.5.

**Table 5-11. Ranges of cRfCs based on applied dose for various noncancer effects associated with inhalation TCE exposure**

cRfC range (ppm)	Neurological	Systemic/organ-specific	Immunological	Reproductive	Developmental
10–100	Impaired visual discrimination (rat)				
1–10		<b>Kidney meganucleocytosis (rat)</b> ↑ <b>kidney weight (mouse)</b>			
0.1–1	Ototoxicity (rat) Hyperactivity (rat) Changes in locomotor activity (rat) <b>Trigeminal nerve effects (human)</b> Impaired visual function (rabbit) ↓ <b>regeneration of sciatic nerve (rat)</b>	↑ <b>liver weight (rat)</b> ↑ <b>liver weight (mouse)</b> ↑ <b>kidney weight (rat)</b>	↓ <b>PFC response (rat)</b>	↓ maternal body weight gain (rat) ↑ abnormal sperm (mouse) pre/postimplantation losses (male rat exp)	
0.01–0.1	↓ <b>regeneration of sciatic nerve (mouse)</b> <b>Disturbed wakefulness (rat)</b>		<b>Autoimmune changes (MRL—lpr/lpr mouse)</b>	<b>Effects on epididymis epithelium (mouse)</b> ↓ <b>fertilization (male mouse exp)</b> <b>Testes and sperm effects (rat)</b> <b>Hyperzoospermia (human)</b>	<b>Resorptions (female rat)</b> ↓ <b>fetal weight (rat)</b> <b>Skeletal effects (rat)</b>

Endpoints in **bold** were selected as candidate critical effects (see Sections 5.1.2.1–5.1.2.8).

**Table 5-12. Ranges of cRfDs based on applied dose for various noncancer effects associated with oral TCE exposure**

cRfD range (mg/kg/day)	Neurological	Systemic/organ-specific	Immunological	Reproductive	Developmental
1–10	↑ neuromuscular changes (rat)	↓ BW (mouse)	↓ humoral response to sRBC (mouse)	↓ testis/seminal vesicle weight (mouse) ↓ sperm motility (mouse) ↑ testis/epididymis weight (rat) ↓ litters/pair (rat) ↓ live pups/litter (rat) ↓ BW gain (rat) ↓ copulatory performance (rat)	↓ fetal weight (rat) Prenatal loss (rat) ↓ fetal weight (mouse) ↑ neonatal mortality (mouse, rat)
0.1–1	↑ # rears (rat) ↑ foot splaying (rat) Trigeminal nerve effect (rat)	↑ <b>liver weight (mouse)</b> ↓ BW (mouse) ↓ BW (rat) <b>Toxic nephropathy</b> (other rat strains/sexes and <b>mouse</b> ) <b>Meganucleocytosis (male Sprague-Dawley rat)</b>	Signs of autoimmune hepatitis (MRL +/- mouse) Inflamm. in various tissues (MRL +/- mouse)	<b>Delayed parturition (rat)</b> ↓ <b>mating (rat)</b>	↓ BW at PND 21 (rat) ↓ locomotor activity (rat) Eye defects (rat) <b>Resorptions (rat)</b>
0.01–0.1	<b>Degeneration of dopaminergic neurons (rat)</b>	<b>Toxic nephropathy (female Marshall rat)</b>	↓ <b>cell-mediated response to sRBC (mouse)</b> ↓ <b>stem cell bone marrow recolonization (mouse)</b>	↓ <b>ability of sperm to fertilize (rat)</b>	↑ <b>exploration (postexp.) (rat)</b> ↓ <b>rearing (postexp.) (mouse)</b> ↓ <b>myelination in hippocampus (rat)</b>
0.001–0.01	<b>Demyelination in hippocampus (rat)</b>		↑ <b>anti-dsDNA and anti-ssDNA Abs (early marker for SLE) (mouse)</b>		
10 <sup>-4</sup> –0.001			↓ <b>thymus weight (mouse)</b>		<b>Immunotox (↓ PFC, ↑ DTH) (B6C3F1 mouse)</b> <b>Heart malformations (rat)</b>

Endpoints in **bold** were selected as candidate critical effects (see Sections 5.1.2.1–5.1.2.8).

1 Quantitative analyses of the PBPK modeling uncertainties and their implications for  
2 dose-response assessment, utilizing the results of the Bayesian analysis of the PBPK model, are  
3 discussed separately in Section 5.1.4.  
4

#### 5.1.3.1.1. Selection of Dose-metrics for Different Endpoints

5 One area of scientific uncertainty in noncancer dose-response assessment is the  
6 appropriate scaling between rodent and human doses for equivalent responses. As discussed  
7 above, the interspecies UF of 10 is usually thought of as a product of two factors of  
8 (approximately) three each for pharmacokinetics and pharmacodynamics. In this assessment,  
9 EPA's cross-species scaling methodology, grounded in general principles of allometric variation  
10 of biologic processes, is used for describing pharmacokinetic equivalence (Allen and Fisher,  
11 1993; Allen et al., 1987; Crump et al., 1989; "Supplementary data for TCE assessment: Rat  
12 population example," 2011; U.S. EPA, 1992, 2005c) U.S. EPA 2011. Briefly, in the absence of  
13 adequate information to the contrary, the methodology determines pharmacokinetic equivalence  
14 across species through equal average lifetime concentrations or AUCs of the toxicant. Thus, in  
15 cases where the PBPK model can predict internal concentrations of the active moiety, equivalent  
16 daily AUCs are assumed to address cross-species pharmacokinetics, and the interspecies UF is  
17 reduced to 3 to account for the remaining pharmacodynamic factor.

18 In the absence of directly estimated AUCs, the cross-species scaling methodology  
19 assumes that, unless there is evidence to the contrary ("Supplementary data for TCE assessment:  
20 Rat population example," 2011; U.S. EPA, 1992, 2005c)

- 21  
22
- 23 (1) The production of the active moiety(ies) is proportional to dose
  - 24 (2) The clearance of the active moiety(ies) scales allometrically by body weight to the  $\frac{3}{4}$   
25 power; and
  - 26 (3) The tissue distribution is equal across species  
27  
28

29 Under these assumptions, for oral exposures, pharmacokinetic equivalence of AUCs between  
30 animals to humans is expressed on the basis of mg/kg<sup>3/4</sup>/d, not mg/kg/day ("body-weight  
31 scaling"). For inhalation exposures, pharmacokinetic equivalence would be on the basis of  
32 equivalent air concentrations, since the alveolar ventilation rate (which determines dose, for a  
33 constant air concentration) scales approximately by body weight to the  $\frac{3}{4}$  power, cancelling out  
34 the assumed scaling dependence of clearance.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1           However, when one or more metabolites are thought to be the toxicologically active  
2 compound(s), it is often the case that a PBPK model can predict the rate of production of the  
3 active moiety(ies) (i.e., the rate of metabolism) but cannot predict AUCs due to lack of data to  
4 inform clearance. In this case, assumption (1) above can be replaced by the PBPK model, while  
5 the other two cross-species scaling methodology assumptions are retained. The resulting  
6 pharmacokinetic equivalence can therefore be expressed on the basis of rate of  
7 metabolism/kg<sup>3/4</sup>/d.<sup>32</sup> Thus, in cases where the PBPK model can predict the rate of production of  
8 the active metabolite(s), equivalent daily amounts metabolized through the appropriate pathway  
9 per unit body weight to the <sup>3/4</sup> power are assumed to address cross-species pharmacokinetics, and  
10 the interspecies UF is reduced to 3 to account for the remaining pharmacodynamic factor.

11           In addition, in some cases when AUCs cannot be estimated, there are data to replace  
12 assumption (2), above, that the clearance of the active moiety(ies) scales allometrically by body  
13 weight to the <sup>3/4</sup> power. Often, this is considered for toxicity associated with local (in situ)  
14 production of “reactive” metabolites whose concentrations cannot be directly measured in the  
15 target tissue. In such a case, an alternative approach of scaling the rate of local metabolism by  
16 target tissue mass, rather than body weight to the <sup>3/4</sup> power, is appropriate if the metabolites are  
17 sufficiently reactive *and* are cleared by “spontaneous” deactivation (i.e., changes in chemical  
18 structure without the need of biological influences). In particular, use of this alternative scaling  
19 approach requires evidence that (1) the active moiety or moieties do not leave the target tissue in  
20 appreciable quantities (i.e., are cleared primarily by in situ transformation to other chemical  
21 species and/or binding to/reactions with cellular components); and (2) the clearance of the active  
22 moieties from the target tissue is governed by biochemical reactions whose rates are independent  
23 of body weight (e.g., purely chemical reactions). If these conditions are met, equivalent daily  
24 amounts metabolized through the appropriate pathway per unit target tissue mass are assumed to  
25 address cross-species pharmacokinetics, and the interspecies UF is reduced to 3 to account for  
26 the remaining pharmacodynamic factor.

---

32 Consider a circulating stable metabolite *X*. Under a one-compartment model, at steady-state, the production of *X* will be equal to the clearance of *X*, so that

$$R_{met} = V_d \times BW \times C_X \times k_{cl}$$

where  $R_{met}$  = rate of production of *X* (mg/time),  $V_d$  = fractional volume of distribution,  $BW$  = body weight,  $C_X$  = concentration of *X* and  $k_{cl}$  = clearance of *X* in units of 1/time. Then, for the concentration  $C_X$  to be equivalent between experimental animals (*A*) and humans (*H*):

$$C_X = [R_{met}/BW \times k_{cl} \times V_d]_H = [R_{met}/BW \times k_{cl} \times V_d]_A.$$

Under the cross-species scaling methodology, it is assumed that  $V_d$  is the same across species, so  $[R_{met}/BW \times k_{cl}]_H = [R_{met}/BW \times k_{cl}]_A$ . Next, under the cross-species scaling methodology,  $k_{cl}$  (with units of 1/time) is assumed to scale according to  $BW^{-1/4}$  (U.S. EPA, 2005c) U.S. EPA 2011, leading to:

$$R_{met(H)}/BW_H^{3/4} = R_{met(A)}/BW_A^{3/4}$$

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Finally, there is the case where local metabolism, rather than systemically delivered  
2 metabolite(s), is thought to be involved in toxicity, but there are inadequate data to determine  
3 either the rate of local metabolism or its clearance. In this case, assumption (1) above can be  
4 replaced by the assumption that local metabolism will be proportional to blood concentration.  
5 Because tissue blood flow approximately scales allometrically by body weight to the  $\frac{3}{4}$  power,  
6 combining this with assumptions (2) and (3) above will lead to the AUC of the parent compound  
7 in blood as an appropriate surrogate for local metabolism. Thus, in this case, equivalent daily  
8 AUCs of the parent compound are assumed to address cross-species pharmacokinetics, and the  
9 interspecies UF is reduced to 3 to account for the remaining pharmacodynamic factor.

10 To summarize, the internal dose-metric for addressing cross-species pharmacokinetics is  
11 based on the Agency's cross-species scaling methodology. The preferred dose-metric under this  
12 methodology is equivalent daily AUC of the active moiety (parent compound or metabolite).  
13 For metabolites, in cases where the rate of production, but not the rate of clearance, of the active  
14 moiety can be estimated, the preferred dose-metric is the rate of metabolism (through the  
15 appropriate pathway) scaled by body weight to the  $\frac{3}{4}$  power. If there are sufficient data to  
16 consider the active metabolite moiety(ies) "reactive" and cleared through nonbiological  
17 processes, then the preferred dose-metric is the rate of metabolism (through the appropriate  
18 pathway) scaled by the tissue mass. Finally, if local metabolism is thought to be involved, but  
19 cannot be estimated with the available data, then the AUC of the parent compound in blood is  
20 considered an appropriate surrogate and thus the preferred dose-metric.

21 These dose-metrics were then also used in addressing the pharmacokinetic component,  
22  $UF_{h-pk}$ , of the UF for human (intraspecies) variability. Because all the dose-metrics used for  
23 TCE were for adults, and the dose-metrics are not very sensitive to the plausible range of adult  
24 body weight, for convenience the body weight  $\frac{3}{4}$  scaling used for interspecies extrapolation was  
25 retained for characterization of human variability. However, it should be emphasized that this  
26 intraspecies characterization is of pharmacokinetics only, and not pharmacodynamics.

27 In general, an attempt was made to use tissue-specific dose-metrics representing  
28 particular pathways or metabolites identified from available data on the role of metabolism in  
29 toxicity for each endpoint (discussed in more detail below). The selection was limited to  
30 dose-metrics for which uncertainty and variability could be adequately characterized by the  
31 PBPK model (see Section 3.5). For most endpoints, sufficient information on the role of  
32 metabolites or MOA was not available to identify likely relevant dose-metrics, and more  
33 "upstream" metrics representing either parent compound or total metabolism had to be used.  
34 The "primary" or "preferred" dose-metric referred to in subsequent tables has the greater  
35 biological support for its involvement in toxicity, whereas "alternative" dose-metrics are those

1 that may also be plausibly involved (discussed further below). A discussion of the dose-metrics  
2 selected for particular noncancer endpoints follows.

#### 5.1.3.1.2. Kidney toxicity (meganucleocytosis, increased kidney weight, toxic nephropathy)

3 As discussed in Sections 4.4.6–4.4.7, there is sufficient evidence to conclude that  
4 TCE-induced kidney toxicity is caused predominantly by GSH conjugation metabolites either  
5 produced in situ in or delivered systemically to the kidney. As discussed in Section 3.3.3.2,  
6 bioactivation of S-dichlorovinyl glutathione (DCVG), DCVC, and  
7 N-acetyl-S-(1,2-dichlorovinyl)-L-cysteine (NAcDCVC) within the kidney, either by beta-lyase,  
8 flavin mono-oxygenase (FMO), or cytochrome P450 (CYP), produces reactive species, any or all  
9 of which may cause nephrotoxicity. Therefore, multiple lines of evidence support the conclusion  
10 that renal bioactivation of DCVC is the preferred basis for internal dose extrapolations for  
11 TCE-induced kidney toxicity. However, uncertainties remain as to the relative contribution from  
12 each bioactivation pathway; and quantitative clearance data necessary to calculate the  
13 concentration of each species are lacking. Moreover, the estimates of the amount bioactivated  
14 are indirect, derived from the difference between overall GSH conjugation flux and NAcDCVC  
15 excretion (see Section 3.5.7.3.1).

16 Under the cross-species scaling methodology, the rate of renal bioactivation of DCVC  
17 would be scaled by body weight to the  $\frac{3}{4}$  power. However, it is necessary to consider whether  
18 there are adequate data to support use of the alternative scaling by target tissue mass. For the  
19 beta-lyase pathway, Dekant et al. (1988) reported in trapping experiments that the postulated  
20 reactive metabolites decompose to stable (unreactive) metabolites in the presence of water.  
21 Moreover, the necessity of a chemical trapping mechanism to detect the reactive metabolites  
22 suggests a very rapid reaction such that it is unlikely that the reactive metabolites leave the site  
23 of production. Therefore, these data support the conclusion that, for this bioactivation pathway,  
24 clearance is chemical in nature and hence species-independent. If this were the only  
25 bioactivation pathway, then scaling by kidney weight would be supported. With respect to the  
26 FMO bioactivation pathway, Sausen and Elfarra (1991) reported that after direct dosing of the  
27 postulated reactive sulfoxide (DCVC sulfoxide), the sulfoxide was detected as an excretion  
28 product in bile. These data suggest that reactivity in the tissue to which the sulfoxide was  
29 delivered (the liver, in this case) is insufficient to rule out a significant role for enzymatic or  
30 other biologically mediated systemic clearance. Therefore, according to the criteria outlined  
31 above, for this bioactivation pathway, the data support scaling the rate of metabolism by body  
32 weight to the  $\frac{3}{4}$  power. For P450-mediated bioactivation producing NAcDCVC sulfoxide, the  
33 only relevant data on clearance are from a study of the structural analogue to DCVC,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether (FDVE; Sheffels et al., 2004), which  
2 reported that the postulated reactive sulfoxide was detected in urine. This suggests that the  
3 sulfoxide is sufficiently stable to be excreted by the kidney and supports the scaling of the rate of  
4 metabolism by body weight to the  $\frac{3}{4}$  power.

5 Therefore, because the contributions to TCE-induced nephrotoxicity from each possible  
6 bioactivation pathway are not clear, and the scaling by body weight to the  $\frac{3}{4}$  power is supported  
7 for two of the identified three bioactivation pathways, it is decided here to scale the DCVC  
8 bioactivation rate by body weight to the  $\frac{3}{4}$  power. The primary internal dose-metric for  
9 TCE-induced kidney toxicity is thus, the weekly rate of DCVC bioactivation per unit body  
10 weight to the  $\frac{3}{4}$  power (**ABioactDCVCBW34 [mg/kg<sup>3/4</sup>/week]**). However, it should be noted  
11 that due to the larger relative kidney weight in rats as compared to humans, scaling by kidney  
12 weight instead of body weight to the  $\frac{3}{4}$  power would only change the quantitative interspecies  
13 extrapolation by about twofold,<sup>33</sup> so the sensitivity of the results to the scaling choice is  
14 relatively small. In addition, quantitative estimates for this dose-metric are only available in rats  
15 and humans, and not in mice. Accordingly, this metric was only used for extrapolating results  
16 from rat toxicity studies.

17 An alternative dose-metric that also involves the GSH conjugation pathway is the amount  
18 of GSH conjugation scaled by the  $\frac{3}{4}$  power of body weight (**AMetGSHBW34 [mg/kg<sup>3/4</sup>/week]**).  
19 This dose-metric uses the total flux of GSH conjugation as the toxicologically-relevant dose,  
20 and, thus, incorporates any direct contributions from DCVG and DCVC, which are not addressed  
21 in the DCVC bioactivation metric. The rationale for scaling by body weight to the  $\frac{3}{4}$  power  
22 rather than target tissue mass is the same as above. Because of the lack of availability of the  
23 DCVC bioactivation dose-metric in mice, the GSH conjugation metric is used as the primary  
24 dose-metric for the nephrotoxicity endpoint in studies of mice.

25 Another alternative dose-metric is the total amount of TCE metabolism (oxidation and  
26 GSH conjugation together) scaled by the  $\frac{3}{4}$  power of body weight (**TotMetabBW34**  
27 **[mg/kg<sup>3/4</sup>/week]**). This dose-metric uses the total flux of TCE metabolism as the toxicologically  
28 relevant dose, and, thus, incorporates the possible involvement of oxidative metabolites, acting  
29 either additively or interactively, in addition to GSH conjugation metabolites in nephrotoxicity  
30 (see Section 4.4.6). However, this dose-metric is given less weight than those involving GSH  
31 conjugation because, as discussed in Sections 4.4.6, the weight of evidence supports the  
32 conclusion that GSH conjugation metabolites play a predominant role in nephrotoxicity. The

---

<sup>33</sup>The range of the difference is 2.1–2.4-fold using the posterior medians for the relative kidney weight in rats and humans from the PBPK model described in Section 3.5 (see Table 3-37), and body weights of 0.3–0.4 kg for rats and 60–70 kg for humans.



1 rationale for scaling by body weight to the  $\frac{3}{4}$  power rather than target tissue mass is the same as  
2 above.  
3

#### 5.1.3.1.3. Liver weight increases (hepatomegaly)

4 As discussed in Section 4.5.6, there is substantial evidence that oxidative metabolism is  
5 involved in TCE hepatotoxicity, based primarily on similarities in noncancer effects with a  
6 number of oxidative metabolites of TCE (e.g., chloral hydrate [CH], TCA, and dichloroacetic  
7 acid [DCA]). While TCA is a stable, circulating metabolite, CH and DCA are relatively  
8 short-lived, although enzymatically cleared (see Section 3.3.3.1). As discussed in Section  
9 4.5.6.2.1, there is substantial evidence that TCA alone does not adequately account for the  
10 hepatomegaly induced by TCE; therefore, unlike in previous dose-response analyses (Barton and  
11 Clewell, 2000; Clewell and Andersen, 2004), the AUC of TCA in plasma or in liver were not  
12 considered as dose-metrics. However, there are inadequate data across species to quantify the  
13 dosimetry of CH and DCA, and other intermediates of oxidative metabolism (such as TCE-oxide  
14 or dichloroacetylchloride) may be involved in hepatomegaly. Thus, due to uncertainties as to the  
15 active moiety(ies), but given the strong evidence associating TCE liver effects with oxidative  
16 metabolism in the liver, hepatic oxidative metabolism is the preferred basis for internal dose  
17 extrapolations of TCE-induced liver weight increases.

18 Under the cross-species scaling methodology, the rate of hepatic oxidative metabolism  
19 would be scaled by body weight to the  $\frac{3}{4}$  power. However, it is necessary to consider whether  
20 there are adequate data to support use of the alternative scaling by target tissue mass. Several of  
21 the oxidative metabolites are stable and systemically available, and several of those that are  
22 cleared rapidly are metabolized enzymatically, so, according to the criteria discussed above,  
23 there are insufficient data to support the conclusions that the active moiety or moieties do not  
24 leave the target tissue in appreciable quantities and are cleared by mechanisms whose rates are  
25 independent of body weight.

26 Therefore, the primary internal dose-metric for TCE-induced liver weight changes is  
27 selected to be the weekly rate of hepatic oxidation per unit body weight to the  $\frac{3}{4}$  power  
28 ( $AMetLiv1BW^{3/4}$  [mg/kg<sup>3/4</sup>/week]). The use of this dose-metric is also supported by the analysis  
29 in Section 4.5.6.2.1 showing much more consistency in the dose-response relationships for  
30 TCE-induced hepatomegaly across studies and routes of exposure using this metric and the total  
31 oxidative metabolism dose-metric (discussed below) as compared to the AUC of TCE in blood.  
32 It should be noted that due to the larger relative liver weight in mice as compared to humans,  
33 scaling by liver weight instead of body weight to the  $\frac{3}{4}$  power would only change the

1 quantitative interspecies extrapolation by about fourfold,<sup>34</sup> so the sensitivity of the results to the  
2 scaling choice is relatively modest.

3 It is also known that the lung has substantial capacity for oxidative metabolism, with  
4 some proportion of the oxidative metabolites produced there entering systemic circulation. Thus,  
5 it is possible that extrahepatic oxidative metabolism can contribute to TCE-induced  
6 hepatomegaly. Therefore, the total amount of oxidative metabolism of TCE scaled by the  
7  $\frac{3}{4}$  power of body weight (**TotOxMetabBW<sup>34</sup> [mg/kg<sup>3/4</sup>/week]**) was selected as an alternative  
8 dose-metric (the rationale for the body weight to the  $\frac{3}{4}$  power scaling is analogous to that for  
9 hepatic oxidative metabolism, above).

10

#### **5.1.3.1.4. Developmental toxicity—heart malformations**

11 As discussed in Section 4.8.3.2.1, several studies have reported that the prenatal exposure  
12 to TCE oxidative metabolites TCA or DCA also induces heart malformations, suggesting that  
13 oxidative metabolism is involved in TCE-induced heart malformations. However, there are  
14 inadequate data across species to quantify the dosimetry of DCA, and it is unclear if other  
15 products of TCE oxidative metabolism are involved. Therefore, the total amount of oxidative  
16 metabolism of TCE scaled by the  $\frac{3}{4}$  power of body weight (TotOxMetabBW<sup>34</sup> [mg/kg<sup>3/4</sup>/week])  
17 was selected as the primary dose-metric. The rationale for the scaling by body weight to the  
18  $\frac{3}{4}$  power is analogous to that for hepatic oxidative metabolism, above.

19 An alternative dose-metric that is considered here is the AUC of TCE in (maternal) blood  
20 (AUCCBld [mg-hour/L/day]). The placenta is a highly perfused tissue, and TCE is known to  
21 cross the placenta to the fetus, with rats showing similar (within twofold) maternal and fetal  
22 blood TCE concentrations (see Section 3.2). This dose-metric accounts for the possible roles  
23 either of local metabolism or of TCE itself.

24

#### **5.1.3.1.5. Reproductive toxicity—decreased ability of sperm to fertilize oocytes**

25 The decreased ability of sperm to fertilize oocytes observed by DuTeaux et al. (2004a)  
26 occurred in the absence of changes in combined testes/epididymes weight, sperm concentration  
27 or motility, or histological changes in the testes or epididymes. However, there was evidence of

---

<sup>34</sup>The range of the difference is 3.5–3.9-fold using the posterior medians for the relative liver weight in mice and humans from the PBPK model described in Section 3.5 (see Table 3-37), and body weights of 0.03–0.04 kg for mice and 60–70 kg for humans.

1 oxidative damage to the sperm, and DuTeaux et al. (2003) previously reported the ability of the  
2 rat epididymis and efferent ducts to metabolize TCE oxidatively. Based on this evidence,  
3 DuTeaux et al. (2004a) hypothesize that the decreased ability to fertilize is due to oxidative  
4 damage to the sperm from local metabolism. Thus, the primary dose-metric for this endpoint is  
5 selected to be the AUC of TCE in blood (AUC<sub>CBld</sub> [mg-hour/L/day]), based on the assumption  
6 that in situ oxidation of systemically-delivered TCE (the flow rate of which scales as body  
7 weight to the  $\frac{3}{4}$  power) is the determinant of toxicity.

8 Because metabolites causing oxidative damage may be delivered systemically to the  
9 target tissue, an alternative dose-metric that is considered here is total oxidative metabolism of  
10 TCE scaled by the  $\frac{3}{4}$  power of body weight (TotOxMetabBW $\frac{3}{4}$  [mg/kg $\frac{3}{4}$ /day]). The rationale  
11 for the scaling by body weight to the  $\frac{3}{4}$  power is analogous to that for hepatic oxidative  
12 metabolism, above. Because oxidative metabolites make up the majority of TCE metabolism,  
13 total metabolism gives very similar results (within 1.2-fold) to total oxidative metabolism and is  
14 therefore not included as a dose-metric.

#### 15 **5.1.3.1.6. Other reproductive and developmental effects and neurological effects and immunologic effects**

16 For all other candidate critical endpoints listed in Tables 5-11–5-12, including  
17 developmental effects other than heart malformations and reproductive effects other than  
18 decreased ability of sperm to fertilize, there is insufficient information for site-specific  
19 determinations of an appropriate dose-metric. While TCE metabolites and/or metabolizing  
20 enzymes have been reported in some of these tissues (e.g., male reproductive tract), their general  
21 roles in toxicity in the respective tissues have not been established. The choice of total  
22 metabolism as the primary dose-metric is based on the observation that, in general, TCE toxicity  
23 is associated with metabolism rather than the parent compound. It is acknowledged that there is  
24 no compelling evidence that definitively establishes one metric as more plausible than the other  
25 in any particular case. Nonetheless, as a general inference in the absence of specific data, total  
26 metabolism is viewed as more likely to be involved in toxicity than the concentration of TCE  
27 itself.

28 Therefore, given that the majority of the toxic and carcinogenic responses in many tissues  
29 to TCE appears to be associated with metabolism, the primary dose-metric is selected to be total  
30 metabolism of TCE scaled by the  $\frac{3}{4}$  power of body weight (TotMetabBW $\frac{3}{4}$  [mg/kg $\frac{3}{4}$ /d]). The  
31 rationale for the scaling by body weight to the  $\frac{3}{4}$  power is analogous to that for the other  
32 metabolism dose-metrics, above. Because oxidative metabolites make up the majority of TCE

1 metabolism, total oxidative metabolism gives very similar results (within 1.2-fold) to total  
2 metabolism and is therefore not included as a dose-metric.

3 An alternative dose-metric that is considered here is the AUC of TCE in blood  
4 (AUCCBld [mg-hour/L/day]). This dose-metric would account for the possible role of local  
5 metabolism, which is determined by TCE delivered in blood via systemic circulation to the target  
6 tissue (the flow rate of which scales as body weight to the  $\frac{3}{4}$  power), and the possible role of  
7 TCE itself. This dose-metric would also be most applicable to tissues that have similar  
8 tissue:blood partition coefficients across and within species.

9 Because the PBPK model described in Section 3.5 did not include a fetal compartment,  
10 the maternal internal dose-metric is taken as a surrogate for developmental effects in which  
11 exposure was before or during pregnancy (Fredriksson et al., 1993; Johnson et al., 2003;  
12 Narotsky et al., 1995; Taylor et al., 1985). This was considered reasonable because TCE and the  
13 major circulating metabolites (TCA and trichloroethanol [TCOH]) appear to cross the placenta  
14 (see Sections 3.2, 3.3, and 4.10 (Fisher et al., 1989; Ghantous et al., 1986)), and maternal  
15 metabolizing capacity is generally greater than that of the fetus (see Section 4.10). In the cases  
16 where exposure continues after birth (Isaacson and Taylor, 1989; Peden-Adams et al., 2006), no  
17 PBPK model-based internal dose was used. Because of the complicated fetus/neonate dosing  
18 that includes transplacental, lactational, and direct (if dosing continues postweaning) exposure,  
19 the maternal internal dose is no more accurate a surrogate than applied dose in this case.

20

#### 5.1.3.1.7. Methods for Inter- and Intraspecies Extrapolation Using Internal Doses<sup>35</sup>


21 As shown in Figures 5-2 and 5-3, the  
22 general approach taken to use the  
23 internal dose-metrics in deriving  
24 HECs and HEDs was to first apply  
25 the rodent PBPK model to get rodent  
26 values for the dose-metrics

---


35An alternative approach (e.g., Clewell et al., 2002) applies the UFs to the internal dose prior to using the human PBPK model to derive a human exposure level. As noted by Barton and Clewell (2000) for previous TCE PBPK models, because the human PBPK model for TCE is linear for all the dose metrics over very broad dose and concentration ranges, essentially identical results would be obtained using this alternative approach. Specifically, for all the primary dose metrics, the difference in the two approaches is less than 2-fold, with the results from the critical studies differing by <0.1%. For some studies using AUCBld as an alternative dose metric, the difference ranged from 3- to 7-fold. Overall, use of the alternative approach would not significantly change the noncancer dose-response assessment of TCE, and the derived RfC and RfD would be identical.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

corresponding to the applied doses in a study reporting noncancer effects. The idPOD is then obtained either directly from the internal dose corresponding to the applied dose LOAEL or NOAEL, or by dose-response modeling of responses with respect to the internal doses to derive a BMDL in terms of internal dose. Separately, the human PBPK model is run for a range of continuous exposures from  $10^{-1}$  to  $2 \times 10^3$  ppm or mg/kg/day to obtain the relationship between human exposure and internal dose for the same dose-metric used for the rodent. The human equivalent exposure (HEC or HED) corresponding to the idPOD is derived by interpolation. It should be noted that median values of dose-metrics were used for rodents, whereas both median and 99<sup>th</sup> percentile values were used for humans. As discussed in Section 3.5, the rodent population model characterizes study-to-study variation, while, within a study, animals with the same sex/species/strain combination were assumed to be identical pharmacokinetically and represented by the group average (typically the only data reported).



1 **Figure 5-2. Flow-chart for dose-response analyses of rodent noncancer**  
 2 **effects using PBPK model-based dose-metrics. Square nodes indicate point**  
 3 **values, circle nodes indicate distributions, and the inverted triangle indicates**  
 4 **a (deterministic) functional relationship.**  
 5



1 **Figure 5-3. Schematic of combined interspecies, intraspecies, and**  
 2 **route-to-route extrapolation from a rodent study LOAEL or NOAEL. In the**  
 3 **case where BMD modeling is performed, the applied dose values are replaced**  
 4 **by the corresponding median internal dose estimate, and the idPOD is the**  
 5 **modeled BMDL in internal dose units.**  
 6

1 Therefore, use of median dose-metric values can be interpreted as assuming that the animals in  
2 the noncancer toxicity study were all “typical” animals and the idPOD is for a rodent that is  
3 pharmacokinetically “typical.” In practice, the use of median or mean internal doses for rodents  
4 did not make much difference except when the uncertainty in the rodent dose-metric was high.  
5 The impact of the uncertainty in the rodent PBPK dose-metrics is analyzed quantitatively in  
6 Section 5.1.4.2.

7 The human population model characterizes individual-to-individual variation, in addition  
8 to its uncertainty. The “median” value for the HEC or HED was calculated as a point of  
9 comparison but was not actually used for derivation of candidate reference values. Because the  
10 RfC and RfD are intended to characterize the dose below which a sensitive individual would  
11 likely not experience adverse effects, the overall 99<sup>th</sup> percentile of the combined uncertainty and  
12 variability distribution was used for deriving the HEC and HED (denoted HEC<sub>99</sub> and HED<sub>99</sub>)  
13 from each idPOD.<sup>36</sup> As shown in Figures 5-2 and 5-3, the HEC<sub>99</sub> or HED<sub>99</sub> replaces the quantity  
14  $POD/(UF_{is-adj} \times UF_{h-pk})$  in the calculation of the RfC or RfD, i.e., the pharmacokinetic  
15 components of the UFs representing interspecies extrapolation and human interindividual  
16 variability.

17 As calculated, the extrapolated HEC<sub>99</sub> and HED<sub>99</sub> can be interpreted as being the dose or  
18 exposure for which there is 99% likelihood that a randomly selected individual will have an  
19 internal dose less than or equal to the idPOD derived from the rodent study. By contrast, the  
20 HEC<sub>50</sub> and HED<sub>50</sub> can be interpreted as being the dose or exposure for which there is 50%  
21 likelihood that a randomly selected individual will have an internal dose less than or equal to the  
22 idPOD derived from the rodent study. Values of HEC<sub>99</sub> or HED<sub>99</sub> are shown for each study and  
23 dose-metric considered in Tables 5-13–5-18. In addition, values of HEC<sub>50</sub> or HED<sub>50</sub> are shown  
24 for comparison, to give a sense of the difference between the median and the 99% confidence  
25 bound for combined uncertainty and variability. The separate contributions of uncertainty and

---

36While for uncertainty, a 95th percentile is often selected by convention, there is no explicit guidance on the selection of the percentile for human toxicokinetic variability. Ideally, all sources of uncertainty and variability would be included, and percentile selected that is more in line with the levels of risk at which cancer dose-response is typically characterized (e.g., 106 to 104) along with a level of confidence. However, only toxicokinetic uncertainty and variability is assessed quantitatively. Because the distribution here incorporates both uncertainty and variability simultaneously, a percentile higher than the 95th (a conventional choice for uncertainty *only*) was selected. However, percentiles greater than the 99th are likely to be progressively less reliable due to the unknown shape of the tail of the input uncertainty and variability distributions for the PBPK model parameters (which were largely assumed to be normal or lognormal), and the fact that only 42 individuals were incorporated in the PBPK model for characterization of uncertainty and inter-individual variability (see Section 3.5). This concern is somewhat ameliorated because the candidate reference values also incorporate use of UFs to account for inter- and intraspecies toxicodynamic sensitivity.



**Table 5-13. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical neurological effects**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Trigeminal nerve effects</b>													
Ruijten et al. (1991)	Human	LOAEL		14	1	1	10	3	1	30	0.47		Trigeminal nerve effects
		HEC	14	5.3	1	1	3	3	1	10	0.53		[TotMetabBW34]
		HEC	14	8.3	1	1	3	3	1	10	0.83		[AUCCBld]
		HED	7.4	7.3	1	1	3	3	1	10		0.73	[TotMetabBW34] (route-to-route)
		HED	59	14	1	1	3	3	1	10		1.4	[AUCCBld] (route-to-route)
<b>Cognitive effects</b>													
Isaacson et al. (1990)	Rat	LOAEL		47	10	10	10	10	1	10,000 <sup>c</sup>		0.0047	demyelination in hippocampus
		HED	9.4	9.2	10	3	3	10	1	1,000		0.0092	[TotMetabBW34]
		HED	31	4.3	10	3	3	10	1	1,000		0.0043	[AUCCBld]
		HEC	18	7.1	10	3	3	10	1	1,000	0.0071		[TotMetabBW34] (route-to-route)
		HEC	3.8	2.3	10	3	3	10	1	1,000	0.0023		[AUCCBld] (route-to-route)
<b>Mood and sleep disorders</b>													
Arito et al. (1994)	Rat	LOAEL		12	3	3	10	10	1	1,000	0.012		Changes in wakefulness
		HEC	13	4.8	3	3	3	10	1	300	0.016		[TotMetabBW34]
		HEC	15	9.0	3	3	3	10	1	300	0.030		[AUCCBld]
		HED	6.6	6.5	3	3	3	10	1	300		0.022	[TotMetabBW34] (route-to-route)
		HED	65	15	3	3	3	10	1	300		0.051	[AUCCBld] (route-to-route)

**Table 5-13. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical neurological effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Other neurological effects</b>													
Kjellstrand et al. (1987)	Rat	LOAEL		300	10	3	10	10	1	3,000	0.10		↓ regeneration of sciatic nerve
		HEC	274	93	10	3	3	10	1	1,000	0.093		[TotMetabBW34]
		HEC	487	257	10	3	3	10	1	1,000	0.26		[AUCCBld]
		HED	110	97	10	3	3	10	1	1,000		0.097	[TotMetabBW34] (route-to-route)
		HED	436	142	10	3	3	10	1	1,000		0.14	[AUCCBld] (route-to-route)
	Mouse	LOAEL		150	10	3	10	10	1	3,000	0.050		↓ regeneration of sciatic nerve
		HEC	378	120	10	3	3	10	1	1,000	0.12		[TotMetabBW34]
		HEC	198	108	10	3	3	10	1	1,000	0.11		[AUCCBld]
		HED	145	120	10	3	3	10	1	1,000		0.12	[TotMetabBW34] (route-to-route)
		HED	237	76	10	3	3	10	1	1,000		0.076	[AUCCBld] (route-to-route)
Gash et al. (2008)	Rat	LOAEL		710	10	10	10	10	1	10,000 <sup>c</sup>		0.071	degeneration of dopaminergic neurons
		HED	56	53	10	3	3	10	1	1,000		0.053	[TotMetabBW34]
		HED	571	192	10	3	3	10	1	1,000		0.19	[AUCCBld]
		HEC	126	47	10	3	3	10	1	1,000	0.047		[TotMetabBW34] (route-to-route)
		HEC	679	363	10	3	3	10	1	1,000	0.36		[AUCCBld] (route-to-route)

<sup>a</sup>Applied dose POD adjusted to continuous exposure unless otherwise noted. POD, HEC<sub>99</sub>, and HED<sub>99</sub> have same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup>Product of individual uncertainty factors, rounded to 3, 10, 30, 100, 300, 1,000, 3,000, or 10,000 [see Footnote c below].

<sup>c</sup>EPA's report on the RfC and RfD processes (U.S. EPA, 2002c) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the application of the PBPK model for candidate critical effects reduces the values of some of the individual UFs for the p-cRfCs and p-cRfDs.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.

Shaded rows represent the p-cRfC or p-cRfD using the preferred PBPK model dose-metric.

**Table 5-14. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical kidney effects**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Histological changes in kidney</b>													
Maltoni (1986) (inhalation)	Rat	BMDL		40.2	1	3	10	1	1	30	1.3		meganucleocytosis; BMR = 10%
		HEC	0.28	0.038	1	3	3	1	1	10	0.0038		[ABioactDCVCBW34]
		HEC	0.45	0.058	1	3	3	1	1	10	0.0058		[AMetGSHBW34]
		HEC	39	15.3	1	3	3	1	1	10	1.5		[TotMetabBW34]
		HED	0.22	0.023	1	3	3	1	1	10		0.0023	[ABioactDCVCBW34] (route-to-route)
		HED	0.35	0.036	1	3	3	1	1	10		0.0036	[AMetGSHBW34] (route-to-route)
		HED	19	19	1	3	3	1	1	10		1.9	[TotMetabBW34] (route-to-route)
NCI (1976)	Mouse	LOAEL		620	1	10	10	30	1	3,000		0.21	toxic nephrosis
		HED	2.9	0.30	1	3	3	30	1	300		0.00101	[AMetGSHBW34]
		HED	51	48	1	3	3	30	1	300		0.160	[TotMetabBW34]
		HEC	3.9	0.50	1	3	3	30	1	300	0.00165		[AMetGSHBW34] (route-to-route)
		HEC	113	42	1	3	3	30	1	300	0.140		[TotMetabBW34] (route-to-route)

**Table 5-14. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical kidney effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>b</sub> <sup>comp</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
NTP (1988)	Rat	BMDL		9.45	1	10	10	1	1	100		0.0945	toxic nephropathy; BMR = 5%; female Marshall (most sensitive sex/strain)
		HED	0.033	0.0034	1	3	3	1	1	10		0.00034	[ABioactDCVCBW34]
		HED	0.053	0.0053	1	3	3	1	1	10		0.00053	[AMetGSHBW34]
		HED	0.75	0.74	1	3	3	1	1	10		0.074	[TotMetabBW34]
		HEC	0.042	0.0056	1	3	3	1	1	10	0.00056		[ABioactDCVCBW34] (route-to-route)
		HEC	0.067	0.0087	1	3	3	1	1	10	0.00087		[AMetGSHBW34] (route-to-route)
		HEC	1.4	0.51	1	3	3	1	1	10	0.051		[TotMetabBW34] (route-to-route)
<b>Histological changes in kidney (continued)</b>													
Maltoni (1986) (oral)	Rat	BMDL		34	1	10	10	1	1	100		0.34	meganucleocytosis; BMR = 10%
		HED	0.15	0.015	1	3	3	1	1	10		0.0015	[ABioactDCVCBW34]
		HED	0.25	0.025	1	3	3	1	1	10		0.0025	[AMetGSHBW34]
		HED	11	11	1	3	3	1	1	10		0.11	[TotMetabBW34]
		HEC	0.19	0.025	1	3	3	1	1	10	0.0025		[ABioactDCVCBW34] (route-to-route)
		HEC	0.31	0.041	1	3	3	1	1	10	0.0041		[AMetGSHBW34] (route-to-route)
		HEC	22	8.5	1	3	3	1	1	10	0.85		[TotMetabBW34] (route-to-route)

**Table 5-14. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical kidney effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sub>b</sub>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>↑ kidney/body weight ratio</b>													
Kjellstrand et al. (1983b)	Mouse	BMDL		34.7	1	3	10	1	1	30	1.2		BMR = 10%
		HEC	0.88	0.12	1	3	3	1	1	10	0.012		[AMetGSHBW34]
		HEC	52	21	1	3	3	1	1	10	2.1		[TotMetabBW34]
		HED	0.69	0.070	1	3	3	1	1	10		0.0070	[AMetGSHBW34] (route-to-route)
		HED	25	25	1	3	3	1	1	10		2.5	[TotMetabBW34] (route-to-route)
Woolhiser et al. (2006)	Rat	BMDL		15.7	1	3	10	1	1	30	0.52		BMR = 10%
		HEC	0.099	0.013	1	3	3	1	1	10	0.0013		[ABioactDCVCBW34]
		HEC	0.17	0.022	1	3	3	1	1	10	0.0022		[AMetGSHBW34]
		HEC	29	11	1	3	3	1	1	10	1.1		[TotMetabBW34]
		HED	0.078	0.0079	1	3	3	1	1	10		0.00079	[ABioactDCVCBW34] (route-to-route)
		HED	0.13	0.013	1	3	3	1	1	10		0.0013	[AMetGSHBW34] (route-to-route)
		HED	14	14	1	3	3	1	1	10		1.4	[TotMetabBW34] (route-to-route)

<sup>a</sup>Applied dose POD adjusted to continuous exposure unless otherwise noted. POD, HEC<sub>99</sub>, and HED<sub>99</sub> have same units as cRfC or cRfD.

<sup>b</sup>Product of individual uncertainty factors, rounded to 3, 10, 30, 100, 300, 1,000, or 3,000.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.  
Shaded rows represent the p-cRfC or p-cRfD using the preferred PBPK model dose-metric.

**Table 5-15. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical liver effects**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>↑ liver/body weight ratio</b>													
Kjellstrand et al. (1983b)	Mouse	BMDL		21.6	1	3	10	1	1	30	0.72		BMR = 10% increase
		HEC	25	9.1	1	3	3	1	1	10	0.91		[AMetLiv1BW34]
		HEC	75	24.9	1	3	3	1	1	10	2.5		[TotOxMetabBW34]
		HED	9.0	7.9	1	3	3	1	1	10		0.79	[AMetLiv1BW34] (route-to-route)
		HED	32	25.7	1	3	3	13	1	10		2.6	[TotOxMetabBW34] (route-to-route)
Woolhiser et al. (2006)	Rat	BMDL		25	1	3	10	1	1	30	0.83		BMR = 10% increase
		HEC	53	19	1	3	3	1	1	10	1.9		[AMetLiv1BW34]
		HEC	46	16	1	3	3	1	1	10	1.6		[TotOxMetabBW34]
		HED	19	16	1	3	3	1	1	10		1.6	[AMetLiv1BW34] (route-to-route)
		HED	20	17	1	3	3	1	1	10		1.7	[TotOxMetabBW34] (route-to-route)
Buben and O'Flaherty (1985)	Mouse	BMDL		82	1	10	10	1	1	100		0.82	BMR = 10% increase
		HED	12	10	1	3	3	1	1	10		1.0	[AMetLiv1BW34]
		HED	15	13	1	3	3	1	1	10		1.3	[TotOxMetabBW34]
		HEC	32	11	1	3	3	1	1	10	1.1		[AMetLiv1BW34] (route-to-route)
		HEC	34	11	1	3	3	1	1	10	1.1		[TotOxMetabBW34] (route-to-route)

<sup>a</sup>Applied dose POD adjusted to continuous exposure unless otherwise noted. POD, HEC<sub>99</sub>, and HED<sub>99</sub> have same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup>Product of individual uncertainty factors, rounded to 3, 10, 30, 100, 300, 1,000, or 3,000.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.

Shaded rows represent the p-cRfC or p-cRfD using the preferred PBPK model dose-metric.

**Table 5-16. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical immunological effects**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>↓ thymus weight</b>													
Keil et al. (2009)	Mouse	LOAEL		0.35	1	10	10	10	1	1,000		0.00035	↓ thymus weight
		HED	0.049	0.048	1	3	3	10	1	100		0.00048	[TotMetabBW34]
		HED	0.20	0.016	1	3	3	10	1	100		0.00016	[AUCCBld]
		HEC	0.092	0.033	1	3	3	10	1	100	0.00033		[TotMetabBW34] (route-to-route)
		HEC	0.014	0.0082	1	3	3	10	1	100	0.000082		[AUCCBld] (route-to-route)
<b>Autoimmunity</b>													
Kaneiko et al. (2000)	Mouse	LOAEL		70	10	3	3	10	1	1,000	0.070		Changes in immunoreactive organs - liver (including sporadic necrosis in hepatic lobules), spleen; UF <sub>h</sub> = 3 due to autoimmune-prone mouse
		HEC	97	37	10	3	1	10	1	300	0.12		[TotMetabBW34]
		HEC	121	69	10	3	1	10	1	300	0.23		[AUCCBld]
		HED	44	42	10	3	1	10	1	300		0.14	[TotMetabBW34] (route-to-route)
		HED	181	57	10	3	1	10	1	300		0.19	[AUCCBld] (route-to-route)
Keil et al. (2009)	Mouse	LOAEL		0.35	1	10	10	1	1	100		0.0035	↑ anti-dsDNA and anti-ssDNA Abs (early markers for SLE)
		HED	0.049	0.048	1	3	3	1	1	10		0.0048	[TotMetabBW34]
		HED	0.20	0.016	1	3	3	1	1	10		0.0016	[AUCCBld]
		HEC	0.092	0.033	1	3	3	1	1	10	0.0033		[TotMetabBW34] (route-to-route)
		HEC	0.014	0.0082	1	3	3	1	1	10	0.00082		[AUCCBld] (route-to-route)

**Table 5-16. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical immunological effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Immunosuppression</b>													
Woolhiser et al. (2006)	Rat	BMDL		24.9	10	3	10	1	1	300	0.083		↓ PFC response; BMR = 1 SD change; dropped highest dose
		HEC	29	11	10	3	3	1	1	100	0.11		[TotMetabBW34]; all does groups
		HEC	263	140	10	3	3	1	1	100	1.4		[AUCCBld] ; all does groups
		HED	14	14	10	3	3	1	1	100		0.14	[TotMetabBW34] (route-to-route) ; all does groups
		HED	282	91	10	3	3	1	1	100		0.91	[AUCCBld] (route-to-route) ; all does groups
Sanders et al. (1982a)	Mouse	LOAEL		18	1	10	10	3	1	300		0.060	↓ stem cell bone marrow recolonization (sustained); ↓ cell-mediated response to sRBC (largely transient during exposure); females more sensitive
		HED	2.5	2.5	1	3	3	3	1	30		0.083	[TotMetabBW34]
		HED	8.8	0.84	1	3	3	3	1	30		0.028	[AUCCBld]
		HEC	4.8	1.7	1	3	3	3	1	30	0.057		[TotMetabBW34] (route-to-route)
		HEC	0.73	0.43	1	3	3	3	1	30	0.014		[AUCCBld] (route-to-route)

<sup>a</sup>Applied use POD adjusted to continuous exposure unless otherwise noted. POD, HEC<sub>99</sub>, and HED<sub>99</sub> have same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup>Product of individual uncertainty factors, rounded to 3, 10, 30, 100, 300, 1,000, or 3,000.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.  
Shaded rows represent the p-cRfC or p-cRfD using the preferred PBPK model dose-metric.



**Table 5-17. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical reproductive effects**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Effects on sperm, male reproductive outcomes</b>													
Chia et al. (1996)	Human	BMDL		1.4	10	1	10	1	1	100	0.014		Hyperzoospermia; BMR = 10% extra risk
		HEC	1.4	0.50	10	1	3	1	1	30	0.0017		[TotMetabBW34]
		HEC	1.4	0.83	10	1	3	1	1	30	0.0028		[AUCCBld]
		HED	0.74	0.73	10	1	3	1	1	30		0.024	[TotMetabBW34] (route-to-route)
		HED	15	1.6	10	1	3	1	1	30		0.053	[AUCCBld] (route-to-route)
Xu et al. (2004)	Mouse	LOAEL		180	10	3	10	10	1	3,000	0.060		↓ fertilization
		HEC	190	67	10	3	3	10	1	1,000	0.067		[TotMetabBW34]
		HEC	321	170	10	3	3	10	1	1,000	0.17		[AUCCBld]
		HED	80	73	10	3	3	10	1	1,000		0.073	[TotMetabBW34] (route-to-route)
		HED	324	104	10	3	3	10	1	1,000		0.10	[AUCCBld] (route-to-route)
Kumar et al (2000b)	Rat	LOAEL		45	10	3	10	10	1	3,000	0.015		Multiple sperm effects, increasing severity from 12–24 weeks
		HEC	32	13	10	3	3	10	1	1,000	0.013		[TotMetabBW34]
		HEC	91	53	10	3	3	10	1	1,000	0.053		[AUCCBld]
		HED	16	16	10	3	3	10	1	1,000		0.016	[TotMetabBW34] (route-to-route)
		HED	157	49	10	3	3	10	1	1,000		0.049	[AUCCBld] (route-to-route)

**Table 5-17. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical reproductive effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
DuTeaux et al. (2004b)	Rat	LOAEL		141	10	10	10	10	1	10,000 <sup>c</sup>		0.014	↓ ability of sperm to fertilize in vitro
		HED	66	16	10	3	3	10	1	1,000		0.016	[AUCCBld]
		HED	65	42	10	3	3	10	1	1,000		0.042	[TotOxMetabBW34]
		HEC	16	9.3	10	3	3	10	1	1,000	0.0093		[AUCCBld] (route-to-route)
		HEC	160	43	10	3	3	10	1	1,000	0.043		[TotOxMetabBW34] (route-to-route)
<b>Male reproductive tract effects</b>													
Forkert et al. (2002); Kan et al. (2007)	Mouse	LOAEL		180	10	3	10	10	1	3,000	0.060		Effects on epididymis epithelium
		HEC	190	67	10	3	3	10	1	1,000	0.067		[TotMetabBW34]
		HEC	321	170	10	3	3	10	1	1,000	0.17		[AUCCBld]
		HED	80	73	10	3	3	10	1	1,000		0.073	[TotMetabBW34] (route-to-route)
		HED	324	104	10	3	3	10	1	1,000		0.10	[AUCCBld] (route-to-route)
Kumar et al. (2000b) (2001a)	Rat	LOAEL		45	10	3	10	10	1	3,000	0.015		Testes effects, testicular enzyme markers, increasing severity from 12–24 weeks
		HEC	32	13	10	3	3	10	1	1,000	0.013		[TotMetabBW34]
		HEC	91	53	10	3	3	10	1	1,000	0.053		[AUCCBld]
		HED	16	16	10	3	3	10	1	1,000		0.016	[TotMetabBW34] (route-to-route)
		HED	157	49	10	3	3	10	1	1,000		0.049	[AUCCBld] (route-to-route)
<b>Female reproductive outcomes</b>													
Narotsky et al. (1995)	Rat	LOAEL		475	1	10	10	10	1	1,000		0.48	Delayed parturition
		HED	47	44	1	3	3	10	1	100		0.44	[TotMetabBW34]
		HED	350	114	1	3	3	10	1	100		1.1	[AUCCBld]
		HEC	98	37	1	3	3	10	1	100	0.37		[TotMetabBW34] (route-to-route)
		HEC	363	190	1	3	3	10	1	100	1.9		[AUCCBld] (route-to-route)

**Table 5-17. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical reproductive effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Reproductive behavior</b>													
George et al. (1986)	Rat	LOAEL		389	1	10	10	10	1	1,000		0.39	↓ mating (both sexes exposed)
		HED	85	77	1	3	3	10	1	100		0.77	[TotMetabBW34]
		HED	167	52	1	3	3	10	1	100		0.52	[AUCCBld]
		HEC	204	71	1	3	3	10	1	100	0.71		[TotMetabBW34] (route-to-route)
		HEC	103	60	1	3	3	10	1	100	0.60		[AUCCBld] (route-to-route)

<sup>a</sup>Applied dose POD adjusted to continuous exposure unless otherwise noted. POD, HEC<sub>99</sub>, and HED<sub>99</sub> have same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup>Product of individual uncertainty factors, rounded to 3, 10, 30, 100, 300, 1,000, 3,000, or 10,000 (see footnote [c] below).

<sup>c</sup>EPA's report on the RfC and RfD processes (U.S. EPA, 2002c) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the application of the PBPK model for candidate critical effects reduces the values of some of the individual UFs for the p-cRfCs and p-cRfDs.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.

Shaded rows represent the p-cRfC or p-cRfD using the preferred PBPK model dose-metric.

**Table 5-18. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical developmental effects**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Pre and postnatal mortality</b>													
Healy et al. (1982)	Rat	LOAEL		17	1	3	10	10	1	300	0.057		Resorptions
		HEC	16	6.2	1	3	3	10	1	100	0.062		[TotMetabBW34]
		HEC	23	14	1	3	3	10	1	100	0.14		[AUCCBld]
		HED	8.7	8.5	1	3	3	10	1	100		0.085	[TotMetabBW34] (route-to-route)
		HED	73	20	1	3	3	10	1	100		0.20	[AUCCBld] (route-to-route)
Narotsky et al. (1995)	Rat	BMDL		32.2	1	10	10	1	1	100		0.32	Resorptions; BMR = 1% extra risk
		HED	29	28	1	3	3	1	1	10		2.8	[TotMetabBW34]
		HED	95	29	1	3	3	1	1	10		2.9	[AUCCBld]
		HEC	57	23	1	3	3	1	1	10	2.3		[TotMetabBW34] (route-to-route)
		HEC	40	24	1	3	3	1	1	10	2.4		[AUCCBld] (route-to-route)
<b>Pre and postnatal growth</b>													
Healy et al. (1982)	Rat	LOAEL		17	1	3	10	10	1	300	0.057		↓ fetal weight; skeletal effects
		HEC	16	6.2	1	3	3	10	1	100	0.062		[TotMetabBW34]
		HEC	23	14	1	3	3	10	1	100	0.14		[AUCCBld]
		HED	8.7	8.5	1	3	3	10	1	100		0.085	[TotMetabBW34] (route-to-route)
		HED	73	20	1	3	3	10	1	100		0.20	[AUCCBld] (route-to-route)

**Table 5-18. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical developmental effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
<b>Congenital defects</b>													
Johnson et al. (2003)	Rat	BMDL		0.0207	1	10	10	1	1	100		0.00021	Heart malformations (pups); BMR = 1% extra risk; highest-dose group (1,000-fold higher than next highest) dropped to improve model fit
		HED	0.0058	0.0052	1	3	3	1	1	10		0.00052	[TotOxMetabBW34]
		HED	0.019	0.0017	1	3	3	1	1	10		0.00017	[AUCCBld]
		HEC	0.012	0.0037	1	3	3	1	1	10	0.00037		[TotOxMetabBW34] (route-to-route)
		HEC	0.0016	0.00093	1	3	3	1	1	10	0.000093		[AUCCBld] (route-to-route)
<b>Developmental neurotoxicity</b>													
Fredriksson et al. (1993)	Mouse	LOAEL		50	3	10	10	10	1	3,000		0.017	↓ rearing postexposure; pup gavage dose
		HED	4.2	4.1	3	3	3	10	1	300		0.014	[TotMetabBW34]
		HED	27	3.5	3	3	3	10	1	300		0.012	[AUCCBld]
		HEC	8.0	3.0	3	3	3	10	1	300	0.010		[TotMetabBW34] (route-to-route)
		HEC	3.1	1.8	3	3	3	10	1	300	0.0061		[AUCCBld] (route-to-route)
Taylor et al. (1985)	Rat	LOAEL		45	1	10	10	10	1	1,000		0.045	↑ exploration postexposure; estimated dam dose
		HED	11	11	1	3	3	10	1	100		0.11	[TotMetabBW34]
		HED	30	4.1	1	3	3	10	1	100		0.041	[AUCCBld]
		HEC	22	8.4	1	3	3	10	1	100	0.084		[TotMetabBW34] (route-to-route)

**Table 5-18. cRfCs and cRfDs (based on applied dose) and p-cRfCs and p-cRfDs (based on PBPK modeled internal dose-metrics) for candidate critical developmental effects (continued)**

Effect type Candidate critical studies	Species	POD type	HEC <sub>50</sub> or HED <sub>50</sub>	POD, HEC <sub>99</sub> , or HED <sub>99</sub> <sup>a</sup>	UF <sub>sc</sub>	UF <sub>is</sub>	UF <sub>h</sub>	UF <sub>loael</sub>	UF <sub>db</sub>	UF <sub>comp</sub> <sup>b</sup>	cRfC or p-cRfC (ppm)	cRfD or p-cRfD (mg/kg/day)	Candidate critical effect; comments [dose-metric]
		HEC	3.7	2.2	1	3	3	10	1	100	0.022		[AUCCBld] (route-to-route)
Isaacson and Taylor (1989)	Rat	LOAEL		16	1	10	10	10	1	1,000		0.016	↓ myelination in hippocampus; estimated dam dose
<b>Developmental immunotoxicity</b>													
Peden-Adams et al. (2006)	Mouse	LOAEL		0.37	1	10	10	10	1	1,000		0.00037	↓ PFC, ↑DTH; POD is estimated dam dose (exposure throughout gestation and lactation + to 3 or 8 wks of age)

<sup>a</sup>Applied dose POD adjusted to continuous exposure unless otherwise noted. POD, HEC<sub>99</sub>, and HED<sub>99</sub> have same units as cRfC (ppm) or cRfD (mg/kg/day).

<sup>b</sup>Product of individual uncertainty factors, rounded to 3, 10, 30, 100, 300, 1,000, or 3,000.

UF<sub>sc</sub> = subchronic-to-chronic UF; UF<sub>is</sub> = interspecies UF; UF<sub>h</sub> = human variability UF; UF<sub>loael</sub> = LOAEL-to-NOAEL UF; UF<sub>db</sub> = database UF.

Shaded rows represent the p-cRfC or p-cRfD using the preferred PBPK model dose-metric or, in the cases where the PBPK model was not used, the cRfD or cRfC based on applied dose.

1 variability in the human PBPK model are analyzed quantitatively, along with the uncertainty in  
2 the rodent PBPK dose-metrics as mentioned above, in Section 5.1.4.2.

3 Because they are derived from rodent internal dose estimates, the HEC and HED are  
4 derived in the same manner independent of the route of administration of the original rodent  
5 study. Therefore, a route-to-route extrapolation from an oral (inhalation) study in rodents to a  
6 HEC (HED) in humans is straight-forward. As shown in Tables 5-13–5-18, route-to-route  
7 extrapolation was performed for a number of endpoints with low cRfCs and cRfDs to derive  
8 p-cRfDs and p-cRfCs.

#### 5.1.3.1.8. Results and Discussion of p-RfCs and p-RfDs for Candidate Critical Effects

10 Tables 5-13–5-18 present the p-cRfCs and p-cRfDs developed using the PBPK internal  
11 dose-metrics, along with the cRfCs and cRfDs based on applied dose for comparison, for each  
12 health effect domain.

13 The greatest impact of using the PBPK model was, as expected, for kidney effects, since  
14 as discussed in Sections 3.3 and 3.5, some toxicokinetic data indicate substantially more GSH  
15 conjugation of TCE and subsequent bioactivation of GSH-conjugates in humans relative to rats  
16 or mice. In addition, as discussed in Sections 3.3 and 3.5, the available in vivo data indicate high  
17 interindividual variability in the amount of TCE conjugated with GSH. The overall impact is  
18 that the p-cRfCs and p-cRfDs based on the preferred dose-metric of bioactivated DCVC are  
19 300- to 400-fold lower than the corresponding cRfCs and cRfDs based on applied dose. As  
20 shown in Figure 3-20 in Section 3.5, for this dose-metric there is about a 30- to 100-fold  
21 difference (depending on exposure route and level) between rats and humans in the “central  
22 estimates” of interspecies differences for the fraction of TCE that is bioactivated as DCVC. The  
23 uncertainty in the human central estimate is only on the order of twofold (in either direction),  
24 while that in the rat central estimate is substantially greater, about 10-fold (in either direction).  
25 In addition, the interindividual variability about the human median estimate is on the order of  
26 10-fold (in either direction). However, as noted in Section 3.3.3.2, there are a number of  
27 discrepancies in estimates for the extent of GSH conjugation that may be related to different  
28 analytical methods, and it is possible that GSH conjugation data to which the PBPK model was  
29 calibrated overestimated the extent of DCVG formation by a substantial amount. Thus, there  
30 remain significant uncertainties in the human estimates of GSH conjugation derived from the  
31 PBPK model. Moreover, the estimates of the amount bioactivated are indirect, derived from the  
32 difference between overall GSH conjugation flux and NAcDCVC excretion (see Section  
33 3.5.7.3.1). Therefore, while there is a high degree of confidence in the nephrotoxic hazard posed

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 by TCE, there is less confidence in the p-cRfCs and p-RfDs derived using GSH conjugation  
2 dose-metrics for these effects.

3 In addition, in two cases in which BMD modeling was employed, using internal  
4 dose-metrics led to a sufficiently different dose-response shape so as to change the resulting  
5 reference value by greater than fivefold. For the Woolhiser et al. (2006) decreased PFC  
6 response, this occurred with the AUC of TCE in blood dose-metric, leading to a p-cRfC 17-fold  
7 higher than the cRfC based on applied dose. However, the model fit for this effect using this  
8 metric was substantially worse than the fit using the preferred metric of Total oxidative  
9 metabolism. Moreover, whereas an adequate fit was obtained with applied dose only with the  
10 highest-dose group dropped, all the dose groups were included when the total oxidative  
11 metabolism dose-metric was used while still resulting in a good model fit. Therefore, it appears  
12 that using this metric resolves some of the low-dose supralinearity in the dose-response curve.  
13 Nonetheless, the overall impact of the preferred metric was minimal, as the p-cRfC based on the  
14 Total oxidative metabolism metric was less than 1.4-fold larger than the cRfC based on applied  
15 dose. The second case in which BMD modeling based on internal doses changed the candidate  
16 reference value by more than fivefold was for resorptions reported by Narotsky et al. (1995).  
17 Here, the p-cRfDs were seven- to eightfold larger than the corresponding cRfD based on applied  
18 dose. However, for applied dose there is substantial uncertainty in the low-dose curvature of the  
19 dose-response curve. This uncertainty persisted with the use of internal dose-metrics, so the  
20 BMD remains somewhat uncertain (see figures in Appendix F). In the remaining cases, which  
21 generally involved the “generic” dose-metrics of total metabolism and AUC of TCE in blood, the  
22 p-cRfCs and p-cRfDs were within fivefold of the corresponding cRfC or cRfD based on applied  
23 dose, with the vast majority within threefold. This suggests that the standard UFs for inter and  
24 intraspecies pharmacokinetic variability are fairly accurate in capturing these differences for  
25 these TCE studies.

#### 26 **5.1.4. Uncertainties in cRfCs and cRfDs**

##### **5.1.4.1.1. Qualitative Uncertainties**

27 An underlying assumption in deriving reference values for noncancer effects is that the  
28 dose-response relationship for these effects has a threshold. Thus, a fundamental uncertainty is  
29 the validity of that assumption. For some effects, in particular effects on very sensitive processes  
30 (e.g., developmental processes) or effects for which there is a nontrivial background level and  
31 even small exposures may contribute to background disease processes in more susceptible

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 people, a practical threshold (i.e., a threshold within the range of environmental exposure levels  
2 of regulatory concern) may not exist.

3         Nonetheless, under the assumption of a threshold, the desired exposure level to have as a  
4 reference value is the maximum level at which there is no appreciable risk for an adverse effect  
5 in (nonnegligible) sensitive subgroups (of humans). However, because it is not possible to know  
6 what this level is, “uncertainty factors” are used to attempt to address quantitatively various  
7 aspects, depending on the data set, of qualitative uncertainty.

8         First there is uncertainty about the “point of departure” for the application of UFs.  
9 Conceptually, the POD should represent the maximum exposure level at which there is no  
10 appreciable risk for an adverse effect in the study population under study conditions (i.e., the  
11 threshold in the dose-response relationship). Then, the application of the relevant UFs is  
12 intended to convey that exposure level to the corresponding exposure level for sensitive human  
13 subgroups exposed continuously for a lifetime. In fact, it is again not possible to know that  
14 exposure level even for a laboratory study because of experimental limitations (e.g., the power to  
15 detect an effect, dose spacing, measurement errors, etc.), and crude approximations like the  
16 NOAEL or a BMDL are used. If a LOAEL is used as the POD, the LOAEL-to-NOAEL UF is  
17 applied as an adjustment factor to get a better approximation of the desired exposure level  
18 (threshold), but the necessary extent of adjustment is unknown.

19         If a BMDL is used as the POD, there are uncertainties regarding the appropriate  
20 dose-response model to apply to the data, but these should be minimal if the modeling is in the  
21 observable range of the data. There are also uncertainties about what BMR to use to best  
22 approximate the desired exposure level (threshold, see above). For continuous endpoints, in  
23 particular, it is often difficult to identify the level of change that constitutes the “cut-point” for an  
24 adverse effect. Sometimes, to better approximate the desired exposure level, a BMR somewhat  
25 below the observable range of the data is selected. In such cases, the model uncertainty is  
26 increased, but this is a trade-off to reduce the uncertainty about the POD not being a good  
27 approximation for the desired exposure level.

28         For each of these types of PODs, there are additional uncertainties pertaining to  
29 adjustments to the administered exposures (doses). Typically, administered exposures (doses)  
30 are converted to equivalent continuous exposures (daily doses) over the study exposure period  
31 under the assumption that the effects are related to concentration  $\times$  time, independent of the daily  
32 (or weekly) exposure regimen (i.e., a daily exposure of 6 hours to 4 ppm is considered equivalent  
33 to 24 hours of exposure to 1 ppm). However, the validity of this assumption is generally  
34 unknown, and, if there are dose-rate effects, the assumption of  $C \times t$  equivalence would tend to  
35 bias the POD downwards. Where there is evidence that administered exposure better correlates

1 to the effect than equivalent continuous exposure averaged over the study exposure period (e.g.,  
2 visual effects), administered exposure was not adjusted. For the PBPK analyses in this  
3 assessment, the actual administered exposures are taken into account in the PBPK modeling, and  
4 equivalent daily values (averaged over the study exposure period) for the dose-metrics are  
5 obtained (see above, Section 5.1.3.2). Additional uncertainties about the PBPK-based estimates  
6 include uncertainties about the appropriate dose-metric for each effect, although for some effects  
7 there was better information about relevant dose-metrics than for others (see Section 5.1.3.1).  
8 Furthermore, as discussed in Section 3.3.3.2, there remains substantial uncertainty in the  
9 extrapolation of GSH conjugation from rodents to humans due to limitations in the available  
10 data.

11 Second, there is uncertainty about the UFs. The human variability UF is to some extent  
12 an adjustment factor because for more sensitive people, the dose-response relationship shifts to  
13 lower exposures. However, there is uncertainty about the extent of the adjustment required, i.e.,  
14 about the distribution of human susceptibility. Therefore, in the absence of data on a more  
15 sensitive population(s) or on the distribution of susceptibility in the general population, an UF of  
16 10 is generally used, in part for pharmacokinetic variability and in part for pharmacodynamic  
17 variability. The PBPK analyses in this assessment attempt to account for the pharmacokinetic  
18 portion of human variability using human data on pharmacokinetic variability. A quantitative  
19 uncertainty analysis of the PBPK-derived dose-metrics used in the assessment is presented in  
20 Section 5.1.4.2 below. There is still uncertainty regarding the susceptible subgroups for TCE  
21 exposure and the extent of pharmacodynamic variability.

22 If the data used to determine a particular POD are from laboratory animals, an  
23 interspecies extrapolation UF is used. This UF is also to some extent an adjustment factor for the  
24 expected scaling for toxicologically-equivalent doses across species (i.e., according to body  
25 weight to the  $3/4$  power for oral exposure). However, there is also uncertainty about the true  
26 extent of interspecies differences for specific noncancer effects from specific chemical  
27 exposures. Often, the “adjustment” component of this UF has been attributed to  
28 pharmacokinetics, while the “uncertainty” component has been attributed to pharmacodynamics,  
29 but as discussed above in Section 5.1.3.1, this is not the only interpretation supported. For oral  
30 exposures, the standard value for the interspecies UF is 10, which can be viewed as breaking  
31 down (approximately) to a factor of three for the “adjustment” (nominally pharmacokinetics) and  
32 a factor of three for the “uncertainty” (nominally pharmacodynamics). For inhalation exposures,  
33 no adjustment across species is generally assumed for fixed air concentrations (ppm  
34 equivalence), and the standard value for the interspecies UF is 3 reflects “uncertainty”  
35 (nominally pharmacodynamics only). The PBPK analyses in this assessment attempt to account

1 for the “adjustment” portion of interspecies extrapolation using rodent pharmacokinetic data to  
2 estimate internal doses for various dose-metrics. With respect to the “uncertainty” component,  
3 quantitative uncertainty analyses of the PBPK-derived dose-metrics used in the assessment are  
4 presented in Section 5.1.4.2 below. However, these only address the pharmacokinetic  
5 uncertainties in a particular dose-metric, and there is still uncertainty regarding the true  
6 dose-metrics. Nor do the PBPK analyses address the uncertainty in either cross-species  
7 pharmacodynamic differences (i.e., about the assumption that equal doses of the appropriate  
8 dose-metric convey equivalent risk across species for a particular endpoint from a specific  
9 chemical exposure) or in cross-species pharmacokinetic differences not accounted for by the  
10 PBPK model dose-metrics (e.g., departures from the assumed interspecies scaling of clearance of  
11 the active moiety, in the cases where only its production is estimated). A value of 3 is typically  
12 used for the “uncertainty” about cross-species differences, and this generally represents true  
13 uncertainty because it is usually unknown, even after adjustments have been made to account for  
14 the expected interspecies differences, whether humans have more or less susceptibility, and to  
15 what degree, than the laboratory species in question.

16 If only subchronic data are available, the subchronic-to-chronic UF is to some extent an  
17 adjustment factor because, if the effect becomes more severe with increasing exposure, then  
18 chronic exposure would shift the dose-response relationship to lower exposures. However, the  
19 true extent of the shift is unknown.

20 Sometimes a database UF is also applied to address limitations or uncertainties in the  
21 database. The overall database for TCE is quite extensive, with studies for many different types  
22 of effects, including 2-generation reproductive studies, as well as neurological, immunological,  
23 and developmental immunological studies. In addition, there were sufficient data to develop a  
24 reliable PBPK model to estimate route-to-route extrapolated doses for some candidate critical  
25 effects for which data were only available for one route of exposure. Thus, there is a high degree  
26 of confidence that the TCE database was sufficient to identify some sensitive endpoints.

27

#### **5.1.4.1.2. Quantitative Uncertainty Analysis of Physiologically Based Pharmacokinetic (PBPK) Model-Based Dose-metrics for Lowest-Observed-Adverse-Effect Level (LOAEL) or No-Observed-Adverse-Effect Level (NOAEL)-Based Points of Departure (PODs)**

28 The Bayesian analysis of the PBPK model for TCE generates distributions of uncertainty  
29 and variability in the internal dose-metrics that can be readily used for characterizing the  
30 uncertainty and variability in the PBPK model-based derivations of the HEC and HED.  
31 However, in the primary analysis, a number of simplifications are made including (1) use of


1 median estimates for rodent internal doses and (2) expressing the “sensitive human” HEC and  
2 HED in terms of combined uncertainty and variability. Therefore, a 2-dimensional quantitative  
3 uncertainty and variability analysis is performed, the objective of which is to characterize the  
4 impact of these assumptions.

5 As shown in Figure 5-4, the overall approach taken for the uncertainty analysis is similar  
6 to that used for the point estimates except for the carrying through of separate uncertainty and  
7 variability distributions throughout the analysis. In particular, to address simplification  
8 (1), above, the distribution of rodent internal dose estimates is carried through; and to address  
9 simplification (2), above, uncertainty and variability distributions in human internal dose  
10 estimates are kept distinct.

11 Because of a lack of tested software and limitations of time and resources, this analysis  
12 was not performed for idPODs based on BMD modeling, and was only performed for idPODs  
13 derived from a LOAEL or NOAEL. However, for those endpoints for which BMD modeling  
14 was performed, for the purposes of this uncertainty analysis, an alternative idPOD was used  
15 based on the study LOAEL or NOAEL.

16 In brief, the methodology involves an iterative process of sampling from three separate  
17 distributions – the uncertainty distribution of rodent PBPK model parameters, the uncertainty  
18 distribution of human population PBPK parameters, and the variability distribution of human  
19 individual PBPK model parameters – the latter two of which are related hierarchically. For a  
20 sample from the rodent parameter distribution, the corresponding idPOD is calculated. Then, an  
21 individual is sampled from a human population distribution, which itself is sampled from the  
22 uncertainty distribution of population parameters. For this individual, a human equivalent  
23 exposure (HEC or HED) corresponding to the idPOD is derived by interpolation. Taking  
24 multiple individuals from this population, a HEC or HED corresponding to the median and  
25 99<sup>th</sup> percentile individuals is then derived. Repeating this process (starting again with a sample  
26 from the rodent distribution) results in two distributions (both reflecting uncertainty): one of  
27 “typical” individuals represented by the distribution of population medians, and one of  
28 “sensitive” individuals represented by the distribution of an upper percentile of the population  
29 (e.g., 99<sup>th</sup> percentile). This uncertainty reflects both uncertainty in the rodent internal dose and  
30 uncertainty in the human population parameters. Thus, for selected quantiles of the population  
31 and level of confidence (e.g., X<sup>th</sup> percentile individual at Y<sup>th</sup>% confidence), the interpretation is  
32 that at the resulting HEC or HED, there is Y% confidence that X% of the population has an  
33 internal dose less than that of the rodent in the toxicity study.

34 As shown in Tables 5-19–5-23, the HEC<sub>99</sub> and HED<sub>99</sub> derived using the rodent median  
35 dose-metrics and the combined uncertainty and variability in human dose-metrics is generally



1 near (within 1.3-fold of) the median confidence level estimate of the HEC and HED for the  
 2 99<sup>th</sup> percentile individual. Therefore, the interpretation is that there is about 50% confidence that  
 3 human exposure at the HEC<sub>99</sub> or HED<sub>99</sub> will, in 99% of the human population, lead to an internal  
 4

1 **Figure 5-4. Flow-chart for uncertainty analysis of HECs and HEDs derived**  
2 **using PBPK model-based dose-metrics. Square nodes indicate point values,**  
3 **circle nodes indicate distributions, and the inverted triangle indicates a**  
4 **(deterministic) functional relationship.**  
5

**Table 5-19. Comparison of “sensitive individual” HECs or HEDs for neurological effects based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the NOAEL or LOAEL**

Candidate critical effect Candidate critical study (species)	POD type	Ratio HEC/D <sub>50</sub> : HEC/D <sub>99</sub>	HEC <sub>x</sub> or HED <sub>x</sub>			[Dose-metric]
			X = 99	X = 99, median	X = 99, 95lcb	
<b>Neurological</b>						
Trigeminal nerve effects Ruijten et al. (1991) (human)	HEC	2.62	5.4	5.4	2.6	[TotMetabBW34]
	HEC	1.68	8.3	8.3	4.9	[AUCCBld]
	HED	1.02	7.3	7.2	3.8	[TotMetabBW34] (rtr)
	HED	4.31	14	16	8.0	[AUCCBld] (rtr)
Demyelination in hippocampus Isaacson et al. (1990) (rat)	HED	1.02	9.21	9.20	7.39	[TotMetabBW34]
	HED	7.20	4.29	5.28	2.52	[AUCCBld]
	HEC	2.59	7.09	6.77	4.94	[TotMetabBW34] (rtr)
	HEC	1.68	2.29	2.42	0.606	[AUCCBld] (rtr)
Changes in wakefulness Arito et al. (1994) (rat)	HEC	2.65	4.79	4.86	2.37	[TotMetabBW34]
	HEC	1.67	9	9.10	4.63	[AUCCBld]
	HED	1.02	6.46	6.50	3.39	[TotMetabBW34] (rtr)
	HED	4.25	15.2	18.0	8.33	[AUCCBld] (rtr)
↓ regeneration of sciatic nerve Kjellstrand et al. (1987) (rat)	HEC	2.94	93.1	93.6	38.6	[TotMetabBW34]
	HEC	1.90	257	266	114	[AUCCBld]
	HED	1.13	97.1	96.8	43.4	[TotMetabBW34] (rtr)
	HED	3.08	142	147	78.0	[AUCCBld] (rtr)
↓ regeneration of sciatic nerve Kjellstrand et al. (1987) (mouse)	HEC	3.16	120	125	48.8	[TotMetabBW34]
	HEC	1.84	108	111	59.7	[AUCCBld]
	HED	1.21	120	121	57.0	[TotMetabBW34] (rtr)
	HED	2.13	75.8	79.1	53.4	[AUCCBld] (rtr)
Degeneration of dopaminergic neurons Gash et al. (2008) (rat)	HED	1.06	53	53.8	17.1	[TotMetabBW34]
	HED	2.98	192	199	94.7	[AUCCBld]
	HEC	2.70	46.8	47.9	14.2	[TotMetabBW34] (rtr)

HEC<sub>99</sub> = the 99<sup>th</sup> percentile of the combined human uncertainty and variability distribution of continuous exposure concentrations that lead to the (fixed) median estimate of the rodent internal dose at the POD.

HEC<sub>99,median</sub> (or HEC<sub>99,95lcb</sub>) = the median (or 95<sup>th</sup> percentile lower confidence bound) estimate of the uncertainty distribution of continuous exposure concentrations for which the 99<sup>th</sup> percentile individual has an internal dose less than the (uncertain) rodent internal dose at the POD.

rtr = route-to-route extrapolation using PBPK model and the specified dose-metric.

Shaded rows denote results for the primary dose-metric.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1  
2  
3  
4

**Table 5-20. Comparison of “sensitive individual” HECs or HEDs for kidney and liver effects based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the NOAEL or LOAEL**

Candidate critical effect Candidate critical study (species)	POD type	Ratio HEC/D <sub>50</sub> : HEC/D <sub>99</sub>	HEC <sub>x</sub> or HED <sub>x</sub>			[Dose-metric]
			X = 99	X = 99, median	X = 99, 95lcb	
<b>Kidney</b>						
Meganucleocytosis [NOAEL]* Maltoni et al. (1986) (rat inhalation)	HEC	7.53	0.0233	0.0260	0.00366	[ABioactDCVCBW34]
	HEC	7.70	0.0364	0.0411	0.00992	[AMetGSHBW34]
	HEC	2.57	8.31	7.97	4.03	[TotMetabBW34]
	HED	9.86	0.0140	0.0156	0.00216	[ABioactDCVCBW34] (rtr)
	HED	9.83	0.0223	0.0242	0.00597	[AMetGSHBW34] (rtr)
	HED	1.02	10.6	10.7	5.75	[TotMetabBW34] (rtr)
Toxic nephrosis NCI (1976) (mouse)	HED	9.51	0.30	0.32	0.044	[AMetGSHBW34]
	HED	1.05	48	48.9	16.2	[TotMetabBW34]
	HEC	7.78	0.50	0.514	0.0703	[AMetGSHBW34] (rtr)
	HEC	2.67	42	43.5	13.7	[TotMetabBW34] (rtr)
Toxic nephropathy [LOAEL]* NTP (1988) (rat)	HED	9.75	0.121	0.126	0.0177	[ABioactDCVCBW34]
	HED	9.64	0.193	0.210	0.0379	[AMetGSHBW34]
	HED	1.03	33.1	33.1	11.1	[TotMetabBW34]
	HEC	7.55	0.201	0.204	0.0269	[ABioactDCVCBW34] (rtr)
	HEC	7.75	0.314	0.353	0.0676	[AMetGSHBW34] (rtr)
	HEC	2.59	28.2	27.2	8.77	[TotMetabBW34] (rtr)
Meganucleocytosis [NOAEL]* Maltoni et al. (1986) (rat oral)	HED	9.85	0.0133	0.0145	0.00158	[ABioactDCVCBW34]
	HED	9.86	0.0214	0.0249	0.00366	[AMetGSHBW34]
	HED	1.02	8.7	8.57	4.95	[TotMetabBW34]
	HEC	7.55	0.022	0.0249	0.00256	[ABioactDCVCBW34] (rtr)
	HEC	7.71	0.0349	0.0424	0.00615	[AMetGSHBW34] (rtr)
	HEC	2.60	6.66	6.31	3.70	[TotMetabBW34] (rtr)
↑ kidney/body weight ratio [NOAEL]* Kjellstrand et al. (1983a) (mouse)	HEC	7.69	0.111	0.103	0.00809	[AMetGSHBW34]
	HEC	2.63	34.5	33.7	13.5	[TotMetabBW34]
	HED	9.78	0.068	0.00641	0.00497	[AMetGSHBW34] (rtr)
	HED	1.03	39.9	39.2	17.9	[TotMetabBW34] (rtr)

5

*This document is a draft for review purposes only and does not constitute Agency policy.*



**Table 5-20. Comparison of “sensitive individual” HECs or HEDs for kidney and liver effects based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the NOAEL or LOAEL (continued)**

Candidate critical effect Candidate critical study (species)	POD type	Ratio HEC/D <sub>50</sub> : HEC/D <sub>99</sub>	HEC <sub>X</sub> or HED <sub>X</sub>			[Dose-metric]
			X = 99	X = 99, median	X = 99, 95lcb	
↑ kidney/body weight ratio [NOAEL] <sup>a</sup> Woolhiser et al. (2006) (rat)	HEC	7.53	0.0438	0.0481	0.00737	[ABioactDCVCBW34]
	HEC	7.70	0.0724	0.0827	0.0179	[AMetGSHBW34]
	HEC	2.54	16.1	15.2	7.56	[TotMetabBW34]
	HED	9.84	0.0264	0.0282	0.00447	[ABioactDCVCBW34] (rtr)
	HED	9.81	0.0444	0.0488	0.0111	[AMetGSHBW34] (rtr)
	HED	1.02	19.5	19.2	10.5	[TotMetabBW34] (rtr)
<b>Liver</b>						
↑ liver/body weight ratio [LOAEL] <sup>a</sup> Kjellstrand et al. (1983a) (mouse)	HEC	2.85	16.2	16.3	6.92	[AMetLiv1BW34]
	HEC	3.63	40.9	38.1	15.0	[TotOxMetabBW34]
	HED	1.16	14.1	14.1	5.85	[AMetLiv1BW34] (rtr)
	HED	1.53	40.1	39.4	17.9	[TotOxMetabBW34] (rtr)
↑ liver/body weight ratio [NOAEL] <sup>a</sup> Woolhiser et al. (2006) (rat)	HEC	2.86	20.7	21.0	11.0	[AMetLiv1BW34]
	HEC	2.94	18.2	17.1	8.20	[TotOxMetabBW34]
	HED	1.20	17.8	17.7	9.94	[AMetLiv1BW34] (rtr)
	HED	1.21	19.6	19.3	10.5	[TotOxMetabBW34] (rtr)
↑ liver/body weight ratio [LOAEL] <sup>a</sup> Buben and O'Flaherty (1985) (mouse)	HED	1.14	8.82	8.95	4.17	[AMetLiv1BW34]
	HED	1.14	9.64	9.78	5.28	[TotOxMetabBW34]
	HEC	2.80	10.1	9.97	4.83	[AMetLiv1BW34] (rtr)
	HEC	3.13	7.83	7.65	4.23	[TotOxMetabBW34] (rtr)

<sup>a</sup>BMDL used for p-cRfC or p-cRfD, but LOAEL or NOAEL (as noted) used for uncertainty analysis.

HEC<sub>99</sub> = the 99<sup>th</sup> percentile of the combined human uncertainty and variability distribution of continuous exposure concentrations that lead to the (fixed) median estimate of the rodent internal dose at the POD.

HEC<sub>99,median</sub> (or HEC<sub>99,95lcb</sub>) = the median (or 95<sup>th</sup> percentile lower confidence bound) estimate of the uncertainty distribution of continuous exposure concentrations for which the 99<sup>th</sup> percentile individual has an internal dose less than the (uncertain) rodent internal dose at the POD.

rtr = route-to-route extrapolation using PBPK model and the specified dose-metric.

Shaded rows denote results for the primary dose-metric.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11

1  
2  
3  
4

**Table 5-21. Comparison of “sensitive individual” HECs or HEDs for immunological effects based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the NOAEL or LOAEL**

Candidate critical effect Candidate critical study (species)	POD type	Ratio HEC/D <sub>50</sub> : HEC/D <sub>99</sub>	HEC <sub>x</sub> or HED <sub>x</sub>			[Dose-metric]
			X = 99	X = 99, median	X = 99, 95lcb	
<b>Immunological</b>						
Changes in immunoreactive organs—liver (including sporadic necrosis in hepatic lobules), spleen Kaneko et al. (2000) (mouse)	HEC	2.65	36.7	38.3	16.0	[TotMetabBW34]
	HEC	1.75	68.9	70.0	37.1	[AUCCBld]
	HED	1.04	42.3	43.3	21.3	[TotMetabBW34] (rtr)
	HED	3.21	56.5	59.0	39.8	[AUCCBld] (rtr)
↑ anti-dsDNA and anti-ssDNA Abs (early markers for SLE); ↓ thymus weight Keil et al. (2009) (mouse)	HED	1.02	0.0482	0.0483	0.0380	[TotMetabBW34]
	HED	12.1	0.0161	0.0189	0.00363	[AUCCBld]
	HEC	2.77	0.0332	0.0337	0.0246	[TotMetabBW34] (rtr)
	HEC	1.69	0.00821	0.00787	0.00199	[AUCCBld] (rtr)
↓ PFC response [NOAEL] <sup>a</sup> Woolhiser et al. (2006) (rat)	HEC	2.54	16.1	15.2	7.56	[TotMetabBW34]
	HEC	1.73	59.6	60.1	26.2	[AUCCBld]
	HED	1.02	19.5	19.2	10.5	[TotMetabBW34] (rtr)
	HED	3.21	52	55.9	33.0	[AUCCBld] (rtr)
↓ stem cell bone marrow recolonization; ↓ cell-mediated response to sRBC Sanders et al. (1982b) (mouse)	HED	1.02	2.48	2.48	1.94	[TotMetabBW34]
	HED	10.5	0.838	0.967	0.187	[AUCCBld]
	HEC	2.77	1.72	1.75	1.28	[TotMetabBW34] (rtr)
	HEC	1.68	0.43	0.412	0.103	[AUCCBld] (rtr)

5  
6  
7  
8  
9  
10  
11  
12  
13  
14

<sup>a</sup>BMDL used for p-cRFc or p-cRFd, but LOAEL or NOAEL (as noted) used for uncertainty analysis.

HEC<sub>99</sub> = the 99<sup>th</sup> percentile of the combined human uncertainty and variability distribution of continuous exposure concentrations that lead to the (fixed) median estimate of the rodent internal dose at the POD.

HEC<sub>99,median</sub> (or HEC<sub>99,95lcb</sub>) = the median (or 95<sup>th</sup> percentile lower confidence bound) estimate of the uncertainty distribution of continuous exposure concentrations for which the 99<sup>th</sup> percentile individual has an internal dose less than the (uncertain) rodent internal dose at the POD.

rtr = route-to-route extrapolation using PBPK model and the specified dose-metric.

Shaded rows denote results for the primary dose-metric.

1  
2  
3  
4

**Table 5-22. Comparison of “sensitive individual” HECs or HEDs for reproductive effects based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the NOAEL or LOAEL**

Candidate critical effect Candidate critical study (species)	POD type	Ratio HEC/D <sub>50</sub> : HEC/D <sub>99</sub>	HEC <sub>x</sub> or HED <sub>x</sub>			[Dose-metric]
			X = 99	X = 99, median	X = 99, 95lcb	
<b>Reproductive</b>						
Hyperzoospermia Chia et al. (1996) (human)	HEC	2.78	0.50	0.53	0.25	[TotMetabBW34]
	HEC	1.68	0.83	0.83	0.49	[AUCCBld]
	HED	1.02	0.73	0.71	0.37	[TotMetabBW34] (rtr)
	HED	9.69	1.6	2.0	0.92	[AUCCBld] (rtr)
↓ fertilization Xu et al. (2004) (mouse)	HEC	2.85	66.6	72.3	26.6	[TotMetabBW34]
	HEC	1.89	170	171	97.1	[AUCCBld]
	HED	1.09	73.3	76.9	32.9	[TotMetabBW34] (rtr)
	HED	3.11	104	109	67.9	[AUCCBld] (rtr)
Multiple sperm effects, testicular enzyme markers Kumar et al. (2001a; 2000b) (rat)	HEC	2.53	12.8	12.2	6.20	[TotMetabBW34]
	HEC	1.72	53.2	54.4	23.2	[AUCCBld]
	HED	1.02	15.8	15.7	8.60	[TotMetabBW34] (rtr)
	HED	3.21	48.8	52.6	30.6	[AUCCBld] (rtr)
↓ ability of sperm to fertilize in vitro DuTeaux et al. (2004b) (rat)	HED	4.20	15.6	18.1	4.07	[AUCCBld]
	HED	1.57	41.7	41.9	32.0	[TotOxMetabBW34]
	HEC	1.67	9.3	10.1	2.09	[AUCCBld] (rtr)
	HEC	3.75	42.5	55.6	39.1	[TotOxMetabBW34] (rtr)
Effects on epididymis epithelium Forkert et al. (2002); Kan et al. (2007) (mouse)	HEC	2.85	66.6	72.3	26.6	[TotMetabBW34]
	HEC	1.89	170	171	97.1	[AUCCBld]
	HED	1.09	73.3	76.9	32.9	[TotMetabBW34] (rtr)
	HED	3.11	104	109	67.9	[AUCCBld] (rtr)
Testes effects Kumar et al. (2001a; 2000b) (rat)	HEC	2.53	12.8	12.2	6.20	[TotMetabBW34]
	HEC	1.72	53.2	54.4	23.2	[AUCCBld]
	HED	1.02	15.8	15.7	8.60	[TotMetabBW34] (rtr)
	HED	3.21	48.8	52.6	30.6	[AUCCBld] (rtr)
Delayed parturition Narotsky et al. (1995) (rat)	HED	1.06	44.3	43.9	15.1	[TotMetabBW34]
	HED	3.07	114	119	47.7	[AUCCBld]
	HEC	2.66	36.9	35.3	11.6	[TotMetabBW34] (rtr)
	HEC	1.91	190	197	48.1	[AUCCBld] (rtr)

5

1 **Table 5-22. Comparison of “sensitive individual” HECs or HEDs for**  
 2 **reproductive effects based on PBPK modeled internal dose-metrics at**  
 3 **different levels of confidence and sensitivity, at the NOAEL or LOAEL**  
 4 **(continued)**  
 5

Candidate critical effect Candidate critical study (species)	POD type	Ratio HEC/D <sub>50</sub> : HEC/D <sub>99</sub>	HEC <sub>x</sub> or HED <sub>x</sub>			[Dose-metric]
			X = 99	X = 99, median	X = 99, 95lcb	
↓ mating (both sexes exposed) George et al. (1986) (rat)	HED	1.10	77.4	77.1	34.2	[TotMetabBW34]
	HED	3.21	51.9	55.8	14.7	[AUCCBld]
	HEC	2.86	71.1	70.0	29.5	[TotMetabBW34] (rtr)
	HEC	1.73	59.5	63.3	8.14	[AUCCBld] (rtr)

6  
 7 HEC<sub>99</sub> = the 99<sup>th</sup> percentile of the combined human uncertainty and variability distribution of continuous exposure  
 8 concentrations that lead to the (fixed) median estimate of the rodent internal dose at the POD.

9 HEC<sub>99,median</sub> (or HEC<sub>99,95lcb</sub>) = the median (or 95<sup>th</sup> percentile lower confidence bound) estimate of the uncertainty  
 10 distribution of continuous exposure concentrations for which the 99<sup>th</sup> percentile individual has an internal dose  
 11 less than the (uncertain) rodent internal dose at the POD.

12 rtr = route-to-route extrapolation using PBPK model and the specified dose-metric.

13 Shaded rows denote results for the primary dose-metric.  
 14

**Table 5-23. Comparison of “sensitive individual” HECs or HEDs for developmental effects based on PBPK modeled internal dose-metrics at different levels of confidence and sensitivity, at the NOAEL or LOAEL**

Candidate critical effect Candidate critical study (species)	POD type	Ratio HEC/D <sub>50</sub> : HEC/D <sub>99</sub>	HEC <sub>X</sub> or HED <sub>X</sub>			[Dose-metric]
			X = 99	X = 95, median	X = 95, 95lcb	
<b>Developmental</b>						
Resorptions Healy et al. (1982) (rat)	HEC	2.58	6.19	6.02	3.13	[TotMetabBW34]
	HEC	1.69	13.7	13.9	7.27	[AUCCBld]
	HED	1.02	8.5	8.50	4.61	[TotMetabBW34] (rtr)
	HED	3.68	19.7	22.4	11.5	[AUCCBld] (rtr)
Resorptions [LOAEL] <sup>a</sup> Narotsky et al. (1995) (rat)	HED	1.06	44.3	43.9	15.1	[TotMetabBW34]
	HED	3.07	114	119	47.7	[AUCCBld]
	HEC	2.66	36.9	35.3	11.6	[TotMetabBW34] (rtr)
	HEC	1.91	190	197	48.1	[AUCCBld] (rtr)
↓ fetal weight; skeletal effects Healy et al. (1982) (rat)	HEC	2.58	6.19	6.02	3.13	[TotMetabBW34]
	HEC	1.69	13.7	13.9	7.27	[AUCCBld]
	HED	1.02	8.5	8.50	4.61	[TotMetabBW34] (rtr)
	HED	3.68	19.7	22.4	11.5	[AUCCBld] (rtr)
Heart malformations (pups) [LOAEL] <sup>a</sup> Johnson et al. (2003) (rat)	HED	1.02	0.012	0.012	0.0102	[TotOxMetabBW34]
	HED	11.6	0.00382	0.00476	0.00112	[AUCCBld]
	HEC	2.75	0.00848	0.00866	0.00632	[TotOxMetabBW34] (rtr)
	HEC	1.70	0.00216	0.00221	0.000578	[AUCCBld] (rtr)
↓ rearing postexposure Fredriksson et al. (1993) (mouse)	HED	1.02	4.13	4.19	2.22	[TotMetabBW34]
	HED	7.69	3.46	4.21	0.592	[AUCCBld]
	HEC	2.71	2.96	2.96	1.48	[TotMetabBW34] (rtr)
	HEC	1.68	1.84	1.81	0.302	[AUCCBld] (rtr)
↑ exploration postexposure Taylor et al. (1985) (rat)	HED	1.02	10.7	10.7	8.86	[TotMetabBW34]
	HED	7.29	4.11	5.08	1.16	[AUCCBld]
	HEC	2.57	8.36	7.94	5.95	[TotMetabBW34] (rtr)
	HEC	1.68	2.19	2.31	0.580	[AUCCBld] (rtr)

<sup>a</sup>BMDL used for p-cRfC or p-cRfD, but LOAEL or NOAEL (as noted) used for uncertainty analysis.

HEC<sub>99</sub> = the 99<sup>th</sup> percentile of the combined human uncertainty and variability distribution of continuous exposure concentrations that lead to the (fixed) median estimate of the rodent internal dose at the POD.

HEC<sub>99,median</sub> (or HEC<sub>99,95lcb</sub>) = the median (or 95<sup>th</sup> percentile lower confidence bound) estimate of the uncertainty distribution of continuous exposure concentrations for which the 99<sup>th</sup> percentile individual has an internal dose less than the (uncertain) rodent internal dose at the POD.

rtr = route-to-route extrapolation using PBPK model and the specified dose-metric.

Shaded rows denote results for the primary dose-metric.

1 dose less than or equal to that in the subjects (rodent or human) exposed at the POD in the  
2 corresponding study.

3 In several cases, the uncertainty, as reflected in the ratio between the 95% and 50%  
4 confidence bounds on the 99<sup>th</sup> percentile individual, was rather high (e.g.,  $\geq 5$ -fold), and reflected  
5 primarily uncertainty in the rodent internal dose estimates, discussed previously in Section 3.5.7.  
6 The largest uncertainties (ratios between 95% to 50% confidence bounds of 8- to 10-fold) were  
7 for kidney effects in mice using the AMetGSHBW34 dose-metric (Kjellstrand et al., 1983a; NCI,  
8 1976). More moderate uncertainties (ratios between 95% to 50% confidence bounds of five- to  
9 eightfold) were evident in some oral studies using the AUCCBld dose-metric (Fredriksson et al.,  
10 1993; George et al., 1986; Keil et al., 2009; Sanders et al., 1982b), as well as in studies reporting  
11 kidney effects in rats in which the ABioactDCVCBW34 or AMetGSHBW34 dose-metrics were  
12 used (Maltoni et al., 1986; NTP, 1988; Woolhiser et al., 2006). Therefore, in these cases, a POD  
13 that is protective of the 99<sup>th</sup> percentile individual at a confidence level higher than 50% could be  
14 as much as an order of magnitude lower.

15 For comparison, Tables 5-19 and 5-23 also show the ratios of the overall 50<sup>th</sup> percentile  
16 to the overall 99<sup>th</sup> percentile HECs and HEDs, reflecting combined human uncertainty and  
17 variability at the median study/endpoint idPOD. The smallest ratios (up to 1.2-fold) are for total,  
18 oxidative, and hepatic oxidative metabolism dose-metrics from oral exposures, due to the large  
19 hepatic first-pass effect resulting in virtually all of the oral intake being metabolized before  
20 systemic circulation. Conversely, the large hepatic first-pass results in high variability in the  
21 blood concentration of TCE following oral exposures, with ratios up to 12-fold at low exposures  
22 (e.g., 90 vs. 99% first-pass would result in amounts metabolized differing by about 10% but TCE  
23 blood concentrations differing by about 10-fold). From inhalation exposures, there is moderate  
24 variability in these metrics, about two- to threefold. For GSH conjugation and bioactivated  
25 DCVC, however, variability is high (eight- to 10-fold) for both exposure routes, which follows  
26 from the incorporation in the PBPK model analysis of the data from Lash et al. (1999a) showing  
27 substantial interindividual variability in GSH conjugation in humans.

28 Finally, it is important to emphasize that this analysis only addresses pharmacokinetic  
29 uncertainty and variability, so other aspects of extrapolation addressed in the UFs (e.g., LOAEL  
30 to NOAEL, subchronic to chronic, and pharmacodynamic differences), discussed above, are not  
31 included in the level of confidence.  
32

## 5.1.5. Summary of Noncancer Reference Values

### 5.1.5.1.1. Preferred Candidate Reference Values (cRfCs, cRfD, p-cRfCs and p-cRfDs) for Candidate Critical Effects

1           The candidate critical effects that yielded the lowest p-cRfC or p-cRfD for each type of  
2 effect, based on the primary dose-metric, are summarized in Tables 5-24 (p-cRfCs) and 5-25  
3 (p-cRfDs). These results are extracted from Tables 5-13–5-18. In cases where a route-to-route  
4 extrapolated p-cRfC (p-cRfD) is lower than the lowest p-cRfC (p-cRfD) from an inhalation  
5 (oral) study, both values are presented in the table. In addition, if there is greater than usual  
6 uncertainty associated with the lowest p-cRfC or p-cRfD for a type of effect, then the endpoint  
7 with the next lowest value is also presented. Furthermore, given those selections, the same sets  
8 of critical effects and studies are displayed across both tables, with the exception of two oral  
9 studies for which route-to-route extrapolation was not performed. Tables 5-24 and 5-25 are  
10 further summarized in Tables 5-26 and 5-27 to present the overall preferred p-cRfC and p-cRfD  
11 for each type of noncancer effect. The purpose of these summary tables is to show the most  
12 sensitive endpoints for each type of effect and the apparent relative sensitivities (based on  
13 reference value estimates) of the different types of effects.

14           For neurological, kidney, immunological, and developmental effects, the lowest p-cRfCs  
15 were derived from oral studies by route-to-route extrapolation. This appears to be a function of  
16 the lack of comparable inhalation studies for many effects studied via the oral exposure route, for  
17 which there is a larger database of studies. For the liver and reproductive effects, inhalation  
18 studies yielded a p-cRfC lower than the lowest route-to-route extrapolated p-cRfC for that type  
19 of effect. Conversely, the lowest p-cRfDs were derived from oral studies with the exception of  
20 reproductive effects, for which route-to-route extrapolation from an inhalation study in humans  
21 also yielded among the lowest p-cRfDs. The only effect for which there were comparable  
22 studies for comparing a p-cRfC from an inhalation study with a p-cRfC estimated by  
23 route-to-route extrapolation from an oral study was increased liver weight in the mouse. The  
24 primary dose-metric of amount of TCE oxidized in the liver yielded similar p-cRfCs of 1.0 and  
25 1.1 ppm for the inhalation result and the route-to-route extrapolated result, respectively (see  
26 Table 5-15).

27           As can be seen in these tables, the most sensitive types of effects (the types with the  
28 lowest p-cRfCs and p-cRfDs) appear to be developmental, kidney, and immunological (adult and  
29 developmental) effects, and then neurological and reproductive effects, in that order. Lastly, the  
30 liver effects have p-cRfC and p-cRfD values that are about 3½ orders of magnitude higher than  
31 those for developmental, kidney, and immunological effects.

1  
2

*This document is a draft for review purposes only and does not constitute Agency policy.*



1  
2

**Table 5-24. Lowest p-cRfCs or cRfCs for different effect domains**

Effect domain Effect type	Candidate critical effect (Species/Critical Study)	p-cRfC or cRfC in ppm (composite uncertainty factor)		
		Preferred dose-metric <sup>a</sup>	Default methodology	Alternative dose-metrics/studies (Tables 5-8–5-13)
<b>Neurologic</b>				
Trigeminal nerve effects	Trigeminal nerve effects (human/Ruijten et al., 1991)	0.54 (10)	0.47 (30)	0.83 (10)
Cognitive effects	Demyelination in hippocampus (rat/Isaacson et al., 1990)	0.0071 (1,000)	– [rtr]	0.0023 (1,000)
Mood/sleep changes	Changes in wakefulness (rat/Arito et al., 1994)	0.016 (300)	0.012 (1,000)	0.030 (300)
<b>Kidney</b>				
Histological changes	<i>Toxic nephropathy</i> (rat/NTP, 1988)	0.00056 (10)	– [rtr]	0.00087–1.3 (10–300)
	Toxic nephrosis (mouse/NCI, 1976)	0.0017 (300)	– [rtr]	
	Meganucleocytosis (rat/Maltoni et al., 1986)	0.0025 (10)	– [rtr]	
↑ kidney weight	↑ kidney weight (rat/Woolhiser et al., 2006)	0.0013 (10)	0.52 (30)	0.0022–2.1 (10–30)
<b>Liver</b>				
↑ liver weight	↑ liver weight (mouse/Kjellstrand et al., 1983a)	0.91 (10)	0.72 (30)	0.83–2.5 (10–30)
<b>Immunologic</b>				
↓ thymus weight	↓ <b>thymus weight</b> (mouse/Keil et al., 2009)	<b>0.00033</b> (100)	– [rtr]	0.000082 (100)
Immuno-suppression	↓ stem cell recolonization (mouse/Sanders et al., 1982b)	0.057 (30)	– [rtr]	0.014–1.4 (30–100)
	Decreased PFC response (rat/Woolhiser et al., 2006)	0.11 (100)	0.083 (300)	
Autoimmunity	↑ anti-dsDNA and anti-ssDNA Abs (mouse/Keil et al., 2009)	0.0033 (10)	– [rtr]	0.00082–0.23 (10–300)
	Autoimmune organ changes mouse/Kaneko et al. (2000)	0.12 (300)	0.070 (1,000)	

3

**Table 5-24. Lowest p-cRfCs or cRfCs for different effect domains (continued)**

Effect domain Effect type	Candidate critical effect (Species/Critical Study)	p-cRfC or cRfC in ppm (composite uncertainty factor)		
		Preferred dose-metric <sup>a</sup>	Default methodology	Alternative dose-metrics/studies (Tables 5-8–5-13)
<b>Reproductive</b>				
Effects on sperm and testes	↓ ability of sperm to fertilize (rat/DuTeaux et al., 2004b)	0.0093 (1,000)	– [rtr]	0.028–0.17 (30–1,000)
	Multiple effects (rat/2001a; Kumar et al., 2000b)	0.013 (1,000)	0.015 (3,000)	
	Hyperzoospermia (human/Chia et al., 1996) <sup>b</sup>	0.017 (30)	0.014 (100)	
<b>Developmental</b>				
Congenital defects	<b>Heart malformations</b> (rat/Johnson et al., 2003)	<b>0.00037</b> (10)	– [rtr]	0.000093 (10)
Develop. neurotox.	↓ rearing postexposure (rat/Fredriksson et al., 1993)	0.028 (300)	– [rtr]	0.0077–0.084 (100–300)
Pre/postnatal mortality/growth	Resorptions/↓ fetal weight/ skeletal effects (rat/Healy et al., 1982)	0.062 (100)	0.057 (300)	0.14–2.4 (10–100)

1  
2 <sup>a</sup>The critical effects/studies and p-cRfCs used to derive the RfC are in **bold**; supporting effects/studies and p-cRfCs  
3 in *italics*.

4 <sup>b</sup>Greater than usual degree of uncertainty (see Section 5.1.2).

5  
6 rtr = route-to-route extrapolated result.

7

1  
2

**Table 5-25. Lowest p-cRfDs or cRfDs for different effect domains**

Effect domain Effect type	Candidate critical effect (Species/Critical Study)	p-cRfD or cRfD in mg/kg/day (composite uncertainty factor)		
		Preferred dose-metric <sup>a</sup>	Default methodology	Alternative dose-metrics/studies (Tables 5-8–5-13)
<b>Neurologic</b>				
Trigeminal nerve effects	Trigeminal nerve effects (human/Ruijten et al., 1991)	0.73 (10)	– [rtr]	1.4 (10)
Cognitive effects	Demyelination in hippocampus (rat/Isaacson et al., 1990)	0.0092 (1,000)	0.0047 (10,000 <sup>b</sup> )	0.0043 (1,000)
Mood/sleep changes	Changes in wakefulness (rat/Arito et al., 1994)	0.022 (300)	– [rtr]	0.051 (300)
<b>Kidney</b>				
Histological changes	<i>Toxic nephropathy</i> (rat/NTP, 1988)	0.00034 (10)	0.0945 (100)	0.00053–1.9 (10–300)
	Toxic nephrosis (mouse/NCI, 1976)	0.0010 (300)		
	Meganucleocytosis (rat/Maltoni et al., 1986)	0.0015 (10)	0.34 (100)	
↑ kidney weight	↑ <i>kidney weight</i> (rat/Woolhiser et al., 2006)	0.00079 (10)	– [rtr]	0.0013–2.5 (10)
<b>Liver</b>				
↑ liver weight	↑ liver weight (mouse/Kjellstrand et al., 1983a)	0.79 (10)	– [rtr]	0.82–2.6 (10–100)
<b>Immunologic</b>				
↓ thymus weight	↓ <b>thymus weight</b> (mouse/Keil et al., 2009)	<b>0.00048</b> (100)	0.00035 (1,000)	0.00016 (100)
Immuno-suppression	↓ stem cell recolonization (mouse/Sanders et al., 1982b)	0.083 (30)	0.060 (300)	0.028–0.91 (30–100)
	Decreased PFC response (rat/Woolhiser et al., 2006)	0.14 (100)	– [rtr]	
Autoimmunity	↑ anti-dsDNA and anti-ssDNA Abs (mouse/Keil et al., 2009)	0.0048 (10)	0.0035 (100)	0.0016–0.19 (10–300)
	Autoimmune organ changes (Kaneko et al., 2000)	0.14 (300)	– [rtr]	

3

**Table 5-25. Lowest p-cRfDs or cRfDs for different effect domains (continued)**

Effect domain Effect type	Candidate critical effect (Species/Critical Study)	p-cRfD or cRfD in mg/kg/day (composite uncertainty factor)		
		Preferred dose-metric <sup>a</sup>	Default methodology	Alternative dose-metrics/studies (Tables 5-8–5-13)
<b>Reproductive</b>				
Effects on sperm and testes	↓ ability of sperm to fertilize (DuTeaux et al., 2004b)	0.016 (1,000)	0.014 (10,000 <sup>b</sup> )	0.042–0.10 (30–1,000)
	Multiple effects (rat/2001a; Kumar et al., 2000b)	0.016 (1,000)	– [rtr]	
	Hyperzoospermia (human/Chia et al., 1996) <sup>c</sup>	0.024 (30)	– [rtr]	
<b>Developmental</b>				
Develop. immunotox.	↓ PFC, ↑ DTH (rat/Peden-Adams et al., 2006) <sup>d</sup>	<b>0.00037</b> (1,000)	Same as preferred	–
Congenital defects	<b>Heart malformations</b> (rat/Johnson et al., 2003)	<b>0.00052</b> (10)	0.00021 (100)	0.00017 (10)
Develop. neurotox.	↓ rearing postexposure (rat/Fredriksson et al., 1993) <sup>d</sup>	0.016 (1,000)	Same as preferred	0.017–0.11 (100–3,000)
Pre/postnatal mortality/growth	Resorptions/↓ fetal weight/ skeletal effects (rat/Healy et al., 1982)	0.085 (100)	[rtr]	0.70–2.9 (10–100)

<sup>a</sup> The critical effects/studies and p-cRfDs or cRfDs used to derive the RfD are in **bold**; supporting effects/studies and p-cRfDs in *italics*.

<sup>b</sup> EPA's report on the RfC and RfD processes (U.S. EPA, 2002c) recommends not deriving reference values with a composite UF of greater than 3,000; however, composite UFs exceeding 3,000 are considered here because the derivation of the cRfCs and cRfDs is part of a screening process and the application of the PBPK model for candidate critical effects reduces the values of some of the individual UFs for the p-cRfCs and p-cRfDs.

<sup>c</sup> Greater than usual degree of uncertainty (see Section 5.1.2).

<sup>d</sup> No PBPK model based analyses were done, so cRfD on the basis of applied dose only.

rtr = route-to-route extrapolated result (no value for default methodology).

**Table 5-26. Lowest p-cRfCs for candidate critical effects for different types of effect based on primary dose-metric**

Type of effect	Effect (primary dose-metric)	p-cRfC (ppm)
Neurological	Demyelination in hippocampus in rats (TotMetabBW34)	0.007 (rtr)
Kidney	Toxic nephropathy in rats (ABioactDCVCBW34)	0.0006 (rtr)
Liver	Increased liver weight in mice (AMetLiv1BW34)	0.9
Immunological	Decreased thymus weight in mice (TotMetabBW34)	0.0003 (rtr)
Reproductive	Decreased ability of rat sperm to fertilize (AUCCBld)	0.009 (rtr) <sup>a</sup>
Developmental	Heart malformations in rats (TotOxMetabBW34)	0.0004 (rtr)

<sup>a</sup>This value is supported by the p-cRfC value of 0.01 ppm for multiple testes and sperm effects from an inhalation study in rats.

rtr = route-to-route extrapolated result.

**Table 5-27. Lowest p-cRfDs for candidate critical effects for different types of effect based on primary dose-metric**

Type of effect	Effect (primary dose-metric)	p-cRfD (mg/kg/day)
Neurological	Demyelination in hippocampus in rats (TotMetabBW34)	0.009
Kidney	Toxic nephropathy in rats (ABioactDCVCBW34)	0.0003
Liver	Increased liver weight in mice (AMetLiv1BW34)	0.8 (rtr)
Immunological	Decreased thymus weight in mice (TotMetabBW34)	0.0005
Reproductive	Decreased ability of rat sperm to fertilize (AUCCBld) and multiple testes and sperm effects (TotMetabBW34) <sup>a</sup>	0.02
Developmental	Heart malformations in rats (TotOxMetabBW34)	0.0005 <sup>b</sup>

<sup>a</sup>Endpoints from two different studies yielded the same p-cRfD value.

1 <sup>b</sup>This value is supported by the cRfD value of 0.0004 mg/kg/day derived for developmental immunotoxicity effects  
2 in mice (Peden-Adams et al., 2006); however, no PBPK analyses were done for this latter effect, so the value of  
3 0.0004 mg/kg/day is based on applied dose.  
4 rtr = route-to-route extrapolated result.

#### 5.1.5.1.2. Reference Concentration

5 The goal is to select an overall RfC that is well supported by the available data (i.e.,  
6 without excessive uncertainty given the extensive database) and protective for all the candidate  
7 critical effects, recognizing that individual candidate RfC values are by nature somewhat  
8 imprecise. The lowest candidate RfC values within each health effect category span a 3,000-fold  
9 range from 0.0003–0.9 ppm (see Table 5-26). One approach to selecting a RfC would be to  
10 select the lowest calculated value of 0.0003 ppm for decreased thymus weight in mice.  
11 However, as can be seen in Table 5-24, three p-cRfCs are in the relatively narrow range of  
12 0.0003–0.0006 ppm at the low end of the overall range. Given the somewhat imprecise nature of  
13 the individual candidate RfC values, and the fact that multiple effects/studies lead to similar  
14 candidate RfC values, the approach taken in this assessment is to select a RfC supported by  
15 multiple effects/studies. The advantages of this approach, which is only possible when there is a  
16 relatively large database of studies/effects and when multiple candidate values happen to fall  
17 within a narrow range at the low end of the overall range, are that it leads to a more robust RfC  
18 (less sensitive to limitations of individual studies) and that it provides the important  
19 characterization that the RfC exposure level is similar for multiple noncancer effects rather than  
20 being based on a sole explicit critical effect.

21 Table 5-28 and Table 5-29 summarize the PODs and UFs for the two critical and one  
22 supporting studies/effects, respectively, corresponding to the p-cRfCs that have been chosen as  
23 the basis of the RfC for TCE noncancer effects. Each of these lowest candidate p-cRfCs, ranging  
24 from 0.0003–0.0006 ppm, for developmental, immunologic, and kidney effects, are values  
25 derived from route-to-route extrapolation using the PBPK model. The lowest p-cRfC estimate  
26 (for a primary dose-metric) from an inhalation studies is 0.001 ppm for kidney effects, which is  
27 higher than the route-to-route extrapolated p-cRfC from the most sensitive oral study. For each  
28 of the candidate RfCs, the PBPK model was used for inter- and intraspecies extrapolation, based  
29 on the preferred dose-metric for each endpoint.

30 There is moderate confidence in the lowest p-cRfC for developmental effects (heart  
31 malformations) (see Section 5.1.2.8) and the lowest p-cRfC estimate for immunological effects  
32 (see Section 5.1.2.5), and these are considered the critical effects used for deriving the RfC. For  
33 developmental effects, although the available study has important limitations, the overall weight  
34 of evidence supports an effect of TCE on cardiac development. For immunological effects, there

1 is high confidence in the evidence for an immunotoxic hazard from TCE, but the available  
2 dose-response data preclude application of BMD modeling.

3 For kidney effects (see Section 5.1.2.2), there is high confidence in the evidence for a  
4 nephrotoxic hazard from TCE. Moreover, the lowest p-cRfC for kidney effects (toxic  
5 nephropathy) is derived from a chronic study and is based on BMD modeling. However, as

6 **Table 5-28. Summary of critical studies, effects, PODs, and UFs used to**  
7 **derive the RfC**

8

Keil et al. (2009)—Decreased thymus weight in female B6C3F1 mice exposed for 30 weeks by drinking water.

- idPOD = 0.139 mg TCE metabolized/kg<sup>3/4</sup>/d, which is the PBPK model-predicted internal dose at the applied dose LOAEL of 0.35 mg/kg/day (continuous) (no BMD modeling due to inadequate model fit caused by supralinear dose-response shape) (see Appendix F, Section F.6.4).
- HEC<sub>99</sub> = 0.033 ppm (lifetime continuous exposure) derived from combined interspecies, intraspecies, and route-to-route extrapolation using PBPK model.
- UF<sub>loael</sub> = 10 because POD is a LOAEL for an adverse effect.
- UF<sub>is</sub> = 3.16 because the PBPK model was used for interspecies extrapolation.
- UF<sub>h</sub> = 3.16 because the PBPK model was used to characterize human toxicokinetic variability.
- p-cRfC = 0.033/100 = 0.00033 ppm (2 µg/m<sup>3</sup>).

Johnson et al. (2003)—fetal heart malformations in S-D rats exposed from GD 1–22 by drinking water.

- idPOD = 0.0142 mg TCE metabolized by oxidation/kg<sup>3/4</sup>/d, which is the BMDL from BMD modeling using PBPK model-predicted internal doses, with highest-dose group (1,000-fold higher than next highest-dose group) dropped, pup as unit of analysis, BMR = 1% (due to severity of defects, some of which could have been fatal), and a nested Log-logistic model to account for intralitter correlation (see Appendix F, Section F.6.5).
- HEC<sub>99</sub> = 0.0037 ppm (lifetime continuous exposure) derived from combined interspecies, intraspecies, and route-to-route extrapolation using PBPK model.
- UF<sub>is</sub> = 3.16 because the PBPK model was used for interspecies extrapolation.
- UF<sub>h</sub> = 3.16 because the PBPK model was used to characterize human toxicokinetic variability.
- p-cRfC = 0.0037/10 = 0.00037 ppm (2 µg/m<sup>3</sup>).

9

10 GD = gestation day.

11

12

13

14

15

**Table 5-29. Summary of supporting studies, effects, PODs, and UFs for the RfC**

NTP (1988)—Toxic nephropathy in female Marshall rats exposed for 104 weeks by oral gavage (5 d/wk).

- idPOD = 0.0132 mg DCVC bioactivated/kg<sup>3/4</sup>/d, which is the BMDL from BMD modeling using PBPK model-predicted internal doses, BMR = 5% (clearly toxic effect), and log-logistic model (see Appendix F, Section F.6.1).
- HEC<sub>99</sub> = 0.0056 ppm (lifetime continuous exposure) derived from combined interspecies, intraspecies, and route-to-route extrapolation using PBPK model.
- UF<sub>is</sub> = 3.16 because the PBPK model was used for interspecies extrapolation.
- UF<sub>h</sub> = 3.16 because the PBPK model was used to characterize human toxicokinetic variability.
- p-cRfC = 0.0056/10 = 0.00056 ppm (3 µg/m<sup>3</sup>).

1  
2  
3 discussed in Section 3.3.3.2, there remains substantial uncertainty in the extrapolation of GSH  
4 conjugation from rodents to humans due to limitations in the available data. In addition, the  
5 p-cRfC for toxic nephropathy had greater dose-response uncertainty since the estimation of its  
6 POD involved extrapolation from high response rates (>60%). Therefore, toxic nephropathy is  
7 considered supportive but is not used as a primary basis for the RfC. The other sensitive  
8 p-cRfCs for kidney effects in Table 5-19 were all within a factor of 5 of that for toxic  
9 nephropathy; however, these values similarly relied on the uncertain interspecies extrapolation of  
10 GSH conjugation.

11 As a whole, the estimates support a RfC of 0.0004 ppm (0.4 ppb or 2 µg/m<sup>3</sup>). This  
12 estimate essentially reflects the midpoint between the similar p-cRfC estimates for the two  
13 critical effects (0.00033 ppm for decreased thymus weight in mice and 0.00037 ppm for heart  
14 malformations in rats), rounded to one significant figure. This estimate is also within a factor of  
15 two of the p-cRfC estimate of 0.00006 ppm for the supporting effect of toxic nephropathy in rats.  
16 Thus, there is robust support for a RfC of 0.0004 ppm provided by estimates for multiple effects  
17 from multiple studies. The estimates are based on PBPK model-based estimates of internal dose  
18 for interspecies, intraspecies, and route-to-route extrapolation, and there is sufficient confidence  
19 in the PBPK model and support from mechanistic data for one of the dose-metrics  
20 (TotOxMetabBW34 for the heart malformations). There is high confidence that  
21 ABioactDCVCBW34 and AMetGSHBW34 would be appropriate dose-metrics for kidney  
22 effects, but there is substantial uncertainty in the PBPK model predictions for these dose-metrics  
23 in humans (see Section 5.1.3.1). Note that there is some human evidence of developmental heart  
24 defects from TCE exposure in community studies (see Section 4.8.3.1.1) and of kidney toxicity  
25 in TCE-exposed workers (see Section 4.4.1).

26 In summary, the RfC is **0.0004 ppm** (0.4 ppb or 2 µg/m<sup>3</sup>) based on route-to-route  
27 extrapolated results from oral studies for the critical effects of heart malformations (rats) and

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 immunotoxicity (mice). This RfC value is further supported by route-to-route extrapolated  
2 results from an oral study of toxic nephropathy (rats).

3

### 5.1.5.1.3. Reference Dose

4 As with the RfC determination above, the goal is to select an overall RfD that is well  
5 supported by the available data (i.e., without excessive uncertainty given the extensive database)  
6 and protective for all the candidate critical effects, recognizing that individual candidate RfD  
7 values are by nature somewhat imprecise. The lowest candidate RfD values within each health  
8 effect category span a nearly 3,000-fold range from 0.0003–0.8 mg/kg/day (see Table 5-26).  
9 One approach to selecting a RfC would be to select the lowest calculated value of 0.0003 ppm  
10 for toxic nephropathy in rats. However, as can be seen in Table 5-25, multiple p-cRfDs or cRfDs  
11 from oral studies are in the relatively narrow range of 0.0003–0.0008 mg/kg/day at the low end  
12 of the overall range. Given the somewhat imprecise nature of the individual candidate RfD  
13 values, and the fact that multiple effects/studies lead to similar candidate RfD values, the  
14 approach taken in this assessment is to select a RfD supported by multiple effects/studies. The  
15 advantages of this approach, which is only possible when there is a relatively large database of  
16 studies/effects and when multiple candidate values happen to fall within a narrow range at the  
17 low end of the overall range, are that it leads to a more robust RfD (less sensitive to limitations  
18 of individual studies) and that it provides the important characterization that the RfD exposure  
19 level is similar for multiple noncancer effects rather than being based on a sole explicit critical  
20 effect.

21 Table 5-30 and Table 5-31 summarize the PODs and UFs for the three critical and  
22 two supporting studies/effects, respectively, corresponding to the p-cRfDs or cRfDs that have  
23 been chosen as the basis of the RfD for TCE noncancer effects. Two of the lowest p-cRfDs for  
24 the primary dose-metrics—0.0008 mg/kg/day for increased kidney weight in rats and 0.0005  
25 mg/kg/day for both heart malformations in rats and decreased thymus weights in mice—are  
26 derived using the PBPK model for inter- and intraspecies extrapolation, and a third—  
27 0.0003 mg/kg/day for increased toxic nephropathy in rats—is derived using the PBPK model for  
28 inter- and intraspecies extrapolation as well as route-to-route extrapolation from an inhalation  
29 study. The other of these lowest values—0.0004 mg/kg/day for developmental immunotoxicity  
30 (decreased PFC response and increased delayed-type hypersensitivity) in mice—is based on  
31 applied dose.

32 There is moderate confidence in the p-cRfDs for decreased thymus weights (see  
33 Section 5.1.2.5) and heart malformations (see Section 5.1.2.8) and the cRfD for developmental

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 immunological effects (see Section 5.1.2.8), and these effects are considered the critical effects  
2 used for deriving the RfD. For developmental effects, although the available study has important  
3 limitations, the overall weight of evidence supports an effect of TCE on cardiac development.  
4 For adult and developmental immunological effects, there is high confidence in the evidence for  
5 an immunotoxic hazard from TCE. However, the available dose-response data for  
6 immunological effects preclude application of BMD modeling.

7 For kidney effects (see Section 5.1.2.2), there is high confidence in the evidence for a  
8 nephrotoxic hazard from TCE. Moreover, the two lowest p-cRfDs for kidney effects (toxic  
9 nephropathy and increased kidney weight) are both based on BMD modeling and one is derived  
10 from a chronic study. However, as discussed in Section 3.3.3.2, there remains substantial  
11 uncertainty in the extrapolation of GSH conjugation from rodents to humans due to limitations in  
12 the available data. In addition, the p-cRfD value for toxic nephropathy had greater  
13 dose-response uncertainty since the estimation of its POD involved extrapolation from high  
14 response rates (>60%). Therefore, kidney effects are considered supportive but are not used as a  
15 primary basis for the RfD.

16  
17 **Table 5-30. Summary of critical studies, effects, PODs, and UFs used to**  
18 **derive the RfD**  
19

Keil et al. (2009)—Decreased thymus weight in female B6C3F1 mice exposed for 30 weeks by drinking water.

- idPOD = 0.139 mg TCE metabolized/kg<sup>3/4</sup>/d, which is the PBPK model-predicted internal dose at the applied dose LOAEL of 0.35 mg/kg/day (continuous) (no BMD modeling due to inadequate model fit caused by supralinear dose-response shape) (see Appendix F, Section F.6.4).
- HED<sub>99</sub> = 0.048 mg/kg/day (lifetime continuous exposure) derived from combined interspecies and intraspecies extrapolation using PBPK model.
- UF<sub>loael</sub> = 10 because POD is a LOAEL for an adverse effect.
- UF<sub>is</sub> = 3.16 because the PBPK model was used for interspecies extrapolation.
- UF<sub>h</sub> = 3.16 because the PBPK model was used to characterize human toxicokinetic variability.
- p-cRfD = 0.048/100 = 0.00048 mg/kg/day.

Peden-Adams et al. (2006)—Decreased PFC response (3 and 8 weeks), increased delayed-type hypersensitivity (8 weeks) in pups exposed from GD 0–3- or 8-weeks-of-age through drinking water (placental and lactational transfer, and pup ingestion).

- $POD = 0.37$  mg/kg/day is the applied dose LOAEL (estimated daily dam dose) (no BMD modeling due to inadequate model fit caused by supralinear dose-response shape). No PBPK modeling was attempted due to lack of appropriate models/parameters to account for complicated fetal/pup exposure pattern (see Appendix F, Section F.6.6).
- $UF_{loael} = 10$  because POD is a LOAEL for multiple adverse effects.
- $UF_{is} = 10$  for interspecies extrapolation because PBPK model was not used.
- $UF_h = 10$  for human variability because PBPK model was not used.
- $cRfD = 0.37/1,000 = 0.00037$  mg/kg/day.

Johnson et al. (2003)—fetal heart malformations in S-D rats exposed from GD 1–22 by drinking water

- $idPOD = 0.0142$  mg TCE metabolized by oxidation/kg<sup>3/4</sup>/d, which is the BMDL from BMD modeling using PBPK model-predicted internal doses, with highest-dose group (1,000-fold higher than next highest-dose group) dropped, pup as unit of analysis, BMR = 1% (due to severity of defects, some of which could have been fatal), and a nested Log-logistic model to account for intralitter correlation (see Appendix F, Section F.6.5).
- $HED_{99} = 0.0051$  mg/kg/day (lifetime continuous exposure) derived from combined interspecies and intraspecies extrapolation using PBPK model.
- $UF_{is} = 3.16$  because the PBPK model was used for interspecies extrapolation.
- $UF_h = 3.16$  because the PBPK model was used to characterize human toxicokinetic variability.
- $p-cRfD = 0.0051/10 = 0.00051$  mg/kg/day.

1  
2  
3

GD = gestation day.

**Table 5-31. Summary of supporting studies, effects, PODs, and UFs for the RfD**

NTP (1988)—Toxic nephropathy in female Marshall rats exposed for 104 weeks by oral gavage (5 d/wk).

- idPOD = 0.0132 mg DCVC bioactivated/kg<sup>3/4</sup>/d, which is the BMDL from BMD modeling using PBPK model-predicted internal doses, BMR = 5% (clearly toxic effect), and Log-logistic model (see Appendix F, Section F.6.1).
- HED<sub>99</sub> = 0.0034 mg/kg/day (lifetime continuous exposure) derived from combined interspecies and intraspecies extrapolation using PBPK model.
- UF<sub>is</sub> = 3.16 because the PBPK model was used for interspecies extrapolation.
- UF<sub>h</sub> = 3.16 because the PBPK model was used to characterize human toxicokinetic variability.
- p-cRfD = 0.0034/10 = 0.00034 mg/kg/day.

Woolhiser et al. (2006)—Increased kidney weight in female S-D rats exposed for 4 weeks by inhalation (6 h/d, 5 d/wk).

- idPOD = 0.0309 mg DCVC bioactivated/kg<sup>3/4</sup>/d, which is the BMDL from BMD modeling using PBPK model-predicted internal doses, BMR = 10%, and Hill model with constant variance (see Appendix F, Section F.6.3).
- HED<sub>99</sub> = 0.0079 mg/kg/day (lifetime continuous exposure) derived from combined interspecies and intraspecies extrapolation using PBPK model.
- UF<sub>sc</sub> = 1 because Kjellstrand et al. (1983a) reported that in mice, kidney effects after exposure for 120 d was no more severe than those after 30 d exposure.
- UF<sub>is</sub> = 3.16 because the PBPK model was used for interspecies extrapolation.
- UF<sub>h</sub> = 3.16 because the PBPK model was used to characterize human toxicokinetic variability.
- p-cRfC = 0.0079/10 = 0.00079 mg/kg/day.

1  
2  
3 As a whole, the estimates support a RfD of 0.0005 mg/kg/day. This estimate is within  
4 20% of the estimates for the critical effects—0.0004 mg/kg/day for developmental  
5 immunotoxicity (decreased PFC and increased delayed-type hypersensitivity) in mice, and  
6 0.0005 mg/kg/day both for heart malformations in rats and for decreased thymus weights in  
7 mice. This estimate is also within approximately a factor of two of the supporting effect  
8 estimates of 0.0003 mg/kg/day for toxic nephropathy in rats and 0.0008 mg/kg/day for increased  
9 kidney weight in rats. Thus, there is strong, robust support for a RfD of 0.0005 mg/kg/day  
10 provided by the concordance of estimates derived from multiple effects from multiple studies.  
11 The estimates for kidney effects, thymus effects, and developmental heart malformations are  
12 based on PBPK model-based estimates of internal dose for interspecies and intraspecies  
13 extrapolation, and there is sufficient confidence in the PBPK model and support from  
14 mechanistic data for one of the dose-metrics (TotOxMetabBW34 for the heart malformations).

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 There is high confidence that ABioactDCVCBW34 would be an appropriate dose-metric for  
2 kidney effects, but there is substantial uncertainty in the PBPK model predictions for this  
3 dose-metric in humans (see Section 5.1.3.1). Note that there is some human evidence of  
4 developmental heart defects from TCE exposure in community studies (see Section 4.8.3.1.1)  
5 and of kidney toxicity in TCE-exposed workers (see Section 4.4.1).

6 In summary, the RfD is **0.0005 mg/kg/day** based on the critical effects of heart  
7 malformations (rats), adult immunological effects (mice), and developmental immunotoxicity  
8 (mice), all from oral studies. This RfD value is further supported by results from an oral study  
9 for the effect of toxic nephropathy (rats) and route-to-route extrapolated results from an  
10 inhalation study for the effect of increased kidney weight (rats).

## 11 **5.2. DOSE-RESPONSE ANALYSIS FOR CANCER ENDPOINTS**

12 This section describes the dose-response analysis for cancer endpoints. Section 5.2.1  
13 discusses the analyses of data from chronic rodent bioassays. Section 5.2.2 discusses the  
14 analyses of human epidemiologic data. Section 5.2.3 discusses the choice of the preferred  
15 inhalation unit risk and oral slope factor estimates, as well as the application of age-dependent  
16 adjustment factors to the slope factor and unit risk estimates.

### 17 **5.2.1. Dose-Response Analyses: Rodent Bioassays**

18 This section describes the calculation of cancer slope factor and unit risk estimates based  
19 on rodent bioassays. First, all the available studies (i.e., chronic rodent bioassays) were  
20 considered, and those suitable for dose-response modeling were selected for analysis (see  
21 Section 5.2.1.1). Then dose-response modeling using the linearized multistage model was  
22 performed using applied doses (default dosimetry) as well as PBPK model-based internal doses  
23 (see Section 5.2.1.2). Bioassays for which time-to-tumor data were available were analyzed  
24 using poly-3 adjustment techniques and using a Multistage Weibull model. In addition, a cancer  
25 potency estimate for different tumor types combined was derived from bioassays in which there  
26 was more than one type of tumor response in the same sex and species. Slope factor and unit  
27 risk estimates based on PBPK model-estimated internal doses were then extrapolated to human  
28 population slope factor and unit risk estimates using the human PBPK model. From these results  
29 (see Section 5.2.1.3), estimates from the most sensitive bioassay (i.e., that with the greatest slope  
30 factor or unit risk estimate) for each combination of administration route, sex, and species, based  
31 on the PBPK model-estimated internal doses, were considered as candidate slope factor or unit

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 risk estimates for TCE. Uncertainties in the rodent-based dose-response analyses are described  
 2 in Section 5.2.1.4.

3

**5.2.1.1.1. Rodent Dose-Response Analyses: Studies and Modeling Approaches**

4 The rodent cancer bioassays that were identified for consideration for dose-response  
 5 analysis are listed in Tables 5-32 (inhalation bioassays) and 5-33 (oral bioassays) for each  
 6 sex/species combination. The bioassays selected for dose-response analysis are marked with an  
 7 asterisk; rationales for rejecting the bioassays that were not selected are provided in the  
 8 “Comments” columns of the tables. For the selected bioassays, the tissues/organs that exhibited  
 9 a TCE-associated carcinogenic response and for which dose-response modeling was performed  
 10 are listed in the “Tissue/Organ” columns.

11

**Table 5-32. Inhalation bioassays**

12

13

Study	Strain	Tissue/Organ	Comments
<b>Female mice</b>			
Fukuda et al. (1983) <sup>a</sup>	Crj:CD-1 (ICR)	Lung	
Henschler et al. (1980) <sup>a</sup>	Han:NMRI	Lymphoma	
Maltoni et al. (1986) <sup>a</sup>	B6C3F1	Liver, Lung	
Maltoni et al. (1986)	Swiss	–	No dose-response
<b>Male mice</b>			
Henschler et al. (1980)	Han:NMRI	–	No dose-response
Maltoni et al. (1986)	B6C3F1	Liver	Exp #BT306: excessive fighting
Maltoni et al. (1986)	B6C3F1	Liver	Exp #BT306bis. Results similar to Swiss mice
Maltoni et al. (1986) <sup>a</sup>	Swiss	Liver	
<b>Female rats</b>			
Fukuda et al. (1983)	Sprague-Dawley	–	No dose-response
Henschler et al. (1980)	Wistar	–	No dose-response
Maltoni et al. (1986)	Sprague-Dawley	–	No dose-response
<b>Male rats</b>			

Henschler et al. (1980)	Wistar	–	No dose-response
Maltoni et al. (1986) <sup>a</sup>	Sprague-Dawley	Kidney, Leydig cell, Leukemia	

1  
2  
3  
4  
5

<sup>a</sup>Selected for dose-response analysis.

“No dose-response” = no tumor incidence data suitable for dose-response modeling.

1  
2

**Table 5-33. Oral bioassays**

<b>Study</b>	<b>Strain</b>	<b>Tissue/organ</b>	<b>Comments</b>
<b>Female mice</b>			
Henschler et al. (1984)	Han:NMRI	–	Toxicity, no dose-response
NCI (1976) <sup>a</sup>	B6C3F1	Liver, lung, sarcomas and lymphomas	
NTP (1990)	B6C3F1	Liver, lung, lymphomas	Single dose
Van Duuren et al. (1979)	Swiss	Liver	Single dose, no dose-response
<b>Male mice</b>			
Anna et al. (1994)	B6C3F1	Liver	Single dose
Bull et al. (2002)	B6C3F1	Liver	Single dose
Henschler et al. (1984)	Han:NMRI	–	Toxicity, no dose-response
NCI (1976) <sup>a</sup>	B6C3F1	Liver	
NTP (1990)	B6C3F1	Liver	Single dose
Van Duuren et al. (1979)	Swiss	–	Single dose, no dose-response
<b>Female rats</b>			
NCI (1976)	Osborne-Mendel	–	Toxicity, no dose-response
NTP (1988)	ACI	–	No dose-response
NTP (1988) <sup>a</sup>	August	Leukemia	
NTP (1988)	Marshall	–	No dose-response
NTP (1988)	Osborne-Mendel	Adrenal cortex	Adenomas only
NTP (1990)	F344/N	–	No dose-response
<b>Male rats</b>			
NCI (1976)	Osborne-Mendel	–	Toxicity, no dose-response
NTP (1988)	ACI	–	No dose-response
NTP (1988) <sup>a</sup>	August	Subcutaneous tissue sarcomas	
NTP (1988) <sup>a</sup>	Marshall	Testes	

*This document is a draft for review purposes only and does not constitute Agency policy.*



NTP (1988) <sup>a</sup>	Osborne-Mendel	Kidney	
NTP (1990) <sup>a</sup>	F344/N	Kidney	

<sup>a</sup> Selected for dose-response analysis.

“No dose-response” = no tumor incidence data suitable for dose-response modeling.

The general approach used was to model each sex/species/bioassay tumor response to determine the most sensitive bioassay response (in terms of human equivalent exposure or dose) for each sex/species combination. The various modeling approaches, model selection, and slope factor and unit risk derivation are discussed below. Modeling was done using the applied dose or exposure (default dosimetry) and several internal dose-metrics. The dose-metrics used in the dose-response modeling are discussed in Section 5.2.1.2. Because of the large volume of analyses and results, detailed discussions about how the data were modeled using the various dosimetry and modeling approaches and results for individual data sets are provided in Appendix G. The overall results are summarized and discussed in Section 5.2.1.3.

Most tumor responses were modeled using the multistage model in EPA’s BMDS ([www.epa.gov/ncea/bmds](http://www.epa.gov/ncea/bmds)). The multistage model is a flexible model, capable of fitting most cancer bioassay data, and it is EPA’s long-standing model for the modeling of such cancer data. The multistage model has the general form

$$P(d) = 1 - \exp\left[-(q_0 + q_1d + q_2d^2 + \dots + q_kd^k)\right] \quad (\text{Eq. 5-1})$$

where  $P(d)$  represents the lifetime risk (probability) of cancer at dose  $d$ , and parameters  $q_i \geq 0$ , for  $i = 0, 1, \dots, k$ . For each data set, the multistage model was evaluated for one stage and  $(n - 1)$  stages, where  $n$  is the number of dose groups in the bioassay. A detailed description of how the data were modeled, as well as tables of the dose-response input data and figures of the multistage modeling results, is provided in Appendix G.

Only models with acceptable fit ( $p > 0.05$ ) were considered.<sup>37</sup> If 1-parameter and 2-parameter models were both acceptable (in no case was there a 3-parameter model), the more parsimonious model (i.e., the 1-parameter model) was selected unless the inclusion of the

<sup>37</sup> When considering multiple types of model for noncancer effects,  $p > 0.10$  is used. For cancer, there is a prior preference for the multistage model, thus the  $p > 0.05$  (which increases the probability of accepting the preferred model).

1 2<sup>nd</sup> parameter resulted in a statistically significant<sup>38</sup> improvement in fit. If two different  
2 1-parameter models were available (e.g., a 1-stage model and a 3-stage model with  $\beta_1$  and  $\beta_2$   
3 both equal to 0), the one with the best fit, as indicated by the lowest AIC value, was selected. If  
4 the AIC values were the same (to three significant figures), then the lower-stage model was  
5 selected. Visual fit and scaled chi-square residuals were also considered for confirmation in  
6 model selection. For two data sets, the highest-dose group was dropped to improve the fit in the  
7 lower dose range.

8 From the selected model for each data set, the maximum likelihood estimate (MLE) for  
9 the dose corresponding to a specified level of risk (i.e., the benchmark dose, or BMD) and its  
10 95% lower confidence bound (BMDL) were estimated.<sup>39</sup> In most cases, the risk level, or BMR,  
11 was 10% extra risk;<sup>40</sup> however, in a few cases with low response rates, a BMR of 5%, or even  
12 1%, extra risk was used to avoid extrapolation above the range of the data. As discussed in  
13 Section 4.4, there is sufficient evidence to conclude that a mutagenic MOA is operative for  
14 TCE-induced kidney tumors, so linear extrapolation from the BMDL to the origin was used to  
15 derive slope factor and unit risk estimates for this site. For all other tumor types, the available  
16 evidence supports the conclusion that the MOA(s) for TCE-induced rodent tumors is unknown,  
17 as discussed in Sections 4.5–4.10 and summarized in Section 4.11.2.3. Therefore, linear  
18 extrapolation was also used based on the general principles outlined in EPA’s *Guidelines for*  
19 *Carcinogen Risk Assessment* (U.S. EPA, 2005c) and reviewed below in Section 5.2.1.4.1. Thus,  
20 for all TCE-associated rodent tumors, slope factor and unit risk estimates are equal to  
21 BMR/BMDL (e.g., 0.10/BMDL<sub>10</sub> for a BMR of 10%). See Section 5.2.1.3 for a summary of the  
22 slope factor and unit risk estimates for each sex/species/bioassay/tumor type.

23 Some of the bioassays exhibited differential early mortality across the dose groups, and,  
24 for three such male rat studies (identified with checkmarks in the “Time-to-tumor” column of  
25 Table 5-34), analyses that take individual animal survival times into account were performed.  
26 (For bioassays with differential early mortality occurring primarily before the time of the  
27 1<sup>st</sup> tumor [or 52 weeks, whichever came first], the effects of early mortality were largely  
28 accounted for by adjusting the tumor incidence for animals at risk, as described in Appendix G,  
29 and the dose-response data were modeled using the regular multistage model, as discussed  
30 above, rather than approaches that account for individual animal survival times.)

---

38Using a standard criterion for nested models, that the difference in  $-2 \times \log$ -likelihood exceeds 3.84 (the 95th percentile of  $\chi^2$  [1]).

39BMDS estimates confidence intervals using the profile likelihood method.

40Extra risk over the background tumor rate is defined as  $[P(d) - P(0)] / [1 - P(0)]$ , where  $P(d)$  represents the lifetime risk (probability) of cancer at dose  $d$ .

1           Two approaches were used to take individual survival times into account. First, EPA's  
2 Multistage Weibull (MSW) software<sup>41</sup> was used for time-to-tumor modeling. The Multistage  
3 Weibull time-to-tumor model has the general form  
4

---

<sup>41</sup>This software has been thoroughly tested and externally reviewed. In February 2009, it will become available on U.S. EPA's Web site.

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 5-34. Specific dose-response analyses performed and dose-metrics used**

Bioassay	Strain	Endpoint	Applied dose	PBPK-based—primary dose-metric	PBPK-based—alternative dose-metric(s)	Time-to-tumor
<b>INHALATION</b>						
<b>Female mice</b>						
Fukuda et al. (1983)	Crj:CD-1 (ICR)	Lung adenomas and carcinomas	√	AMetLngBW34	TotOxMetabBW34 AUCCBld	
Henschler et al. (1980)	Han:NMRI	Lymphoma	√	TotMetabBW34	AUCCBld	
Maltoni et al. (1986)	B6C3F1	Liver hepatomas	√	AMetLiv1BW34	TotOxMetabBW34	
		Lung adenomas and carcinomas	√	AMetLngBW34	TotOxMetabBW34 AUCCBld	
		Combined risk	√			
<b>Male mice</b>						
Maltoni et al. (1986)	Swiss	Liver hepatomas	√	AMetLiv1BW34	TotOxMetabBW34	
<b>Female rats</b>						
None selected						
<b>Male rats</b>						
Maltoni et al. (1986)	Sprague-Dawley	Kidney adenomas and carcinomas	√	ABioactDCVCBW34	AMetGSHBW34 TotMetabBW34	
		Leydig cell tumors	√	TotMetabBW34	AUCCBld	
		Leukemias	√	TotMetabBW34	AUCCBld	
		Combined risk	√			

**Table 5-34. Specific dose-response analyses performed and dose-metrics used (continued)**

Bioassay	Strain	Endpoint	Applied dose	PBPK-based—primary dose-metric	PBPK-based—alternative dose-metric(s)	Time-to-tumor
<b>ORAL</b>						
<b>Female mice</b>						
NCI (1976)	B6C3F1	Liver carcinomas	√	AMetLiv1BW34	TotOxMetabBW34	
		Lung adenomas and carcinomas	√	AMetLngBW34	TotOxMetabBW34 AUCCBld	
		Multiple sarcomas/lymphomas	√	TotMetabBW34	AUCCBld	
		Combined risk	√			
<b>Male mice</b>						
NCI (1976)	B6C3F1	Liver carcinomas	√	AMetLiv1BW34	TotOxMetabBW34	
<b>Female rats</b>						
NTP (1988)	August	Leukemia	√	TotMetabBW34	AUCCBld	
<b>Male rats</b>						
NTP (1988)	August	Subcutaneous tissue sarcomas	√	TotMetabBW34	AUCCBld	
NTP (1988)	Marshall	Testicular interstitial cell tumors	√	TotMetabBW34	AUCCBld	√
NTP (1988)	Osborne-Mendel	Kidney adenomas and carcinomas	√	ABioactDCVCBW34	AMetGSHBW34 TotMetabBW34	√
NTP (1990)	F344/N	Kidney adenomas and carcinomas	√	ABioactDCVCBW34	AMetGSHBW34 TotMetabBW34	√

**PBPK-based dose-metric abbreviations:**

ABioactDCVCBW34 = Amount of DCVC bioactivated in the kidney per unit body weight<sup>3/4</sup> (mg DCVC/kg<sup>3/4</sup>/week).

AMetGSHBW34 = Amount of TCE conjugated with GSH per unit body weight<sup>3/4</sup> (mg TCE/kg<sup>3/4</sup>/week).

AMetLiv1BW34 = Amount of TCE oxidized per unit body weight<sup>3/4</sup> (mg TCE/kg<sup>3/4</sup>/week).

AMetLngBW34 = Amount of TCE oxidized in the respiratory tract per unit body weight<sup>3/4</sup> (mg TCE/kg<sup>3/4</sup>/week).

AUCCBld = Area under the curve of the venous blood concentration of TCE (mg-hour/L/week).

TotMetabBW34 = Total amount of TCE metabolized per unit body weight<sup>3/4</sup> (mg TCE/kg<sup>3/4</sup>/week).

TotOxMetabBW34 = Total amount of TCE oxidized per unit body weight<sup>3/4</sup> (mg TCE/kg<sup>3/4</sup>/week).

$$P(d,t) = 1 - \exp\left[-(q_0 + q_1d + q_2d^2 + \dots + q_kd^k) \times (t-t_0)^z\right] \quad (\text{Eq. 5-2})$$

where  $P(d,t)$  represents the probability of a tumor by age  $t$  for dose  $d$ , and parameters  $z \geq 1$ ,  $t_0 \geq 0$ , and  $q_i \geq 0$  for  $i = 0, 1, \dots, k$ , where  $k =$  the number of dose groups; the parameter  $t_0$  represents the time between when a potentially fatal tumor becomes observable and when it causes death. (All of our analyses used the model for incidental tumors, which has no  $t_0$  term.) Although the fit of the MSW model can be assessed visually using the plot feature of the MSW software, because there is no applicable goodness-of-fit statistic with a well-defined asymptotic distribution, an alternative survival-adjustment technique, “poly-3 adjustment,” was also applied (Portier and Bailer, 1989). This technique was used to adjust the tumor incidence denominators based on the individual animal survival times.<sup>42</sup> The adjusted incidence data then served as inputs for EPA’s BMDS multistage model, and model (i.e., stage) selection was conducted as already described above. Under both survival-adjustment approaches, BMDs and BMDLs were obtained and slope factor and unit risks derived as discussed above for the standard multistage model approach. See Appendix G for a more detailed description of the MSW modeling and for the results of both the MSW and poly-3 approaches for the individual data sets. A comparison of the results for the three different data sets and the various dose-metrics used is presented in Section 5.2.1.3.

For bioassays that exhibited more than one type of tumor response in the same sex and species (these studies have a row for “combined risk” in the “Endpoint” column of Table 5-34), the cancer potency for the different tumor types combined was estimated. The combined tumor risk estimate describes the risk of developing tumors for *any* (not all together) of the tumor types that exhibited a TCE-associated tumor response; this estimate then represents the total excess cancer risk. The model for the combined tumor risk is also multistage, with the sum of the stage-specific multistage coefficients from the individual tumor models serving as the stage-specific coefficients for the combined risk model (i.e., for each  $q_i$ ,  $q_{i[\text{combined}]} = q_{i1} + q_{i2} + \dots + q_{ik}$ , where the  $q_i$ s are the coefficients for the powers of dose and  $k$  is the number of tumor types being combined) (Bogen, 1990; NRC, 1994). This model assumes that the occurrences of two or more tumor types are independent. Although the resulting model equation can be readily solved for a given BMR to obtain an MLE (BMD) for the combined risk,

---

<sup>42</sup>Each tumorless animal is weighted by its fractional survival time (number of days on study divided by 728 days, the typical number of days in a two-year bioassay) raised to the power of three to reflect the fact that animals are at greater risk of cancer at older ages. Animals with tumors are given a weight of one. The sum of the weights of all the animals in an exposure group yields the effective survival-adjusted denominator.

1 the confidence bounds for the combined risk estimate are not calculated by available modeling  
2 software. Therefore, the confidence bounds on the combined BMD were estimated using a  
3 Bayesian approach, computed using Markov chain Monte Carlo techniques and implemented  
4 using the freely available WinBugs software (Spiegelhalter et al., 2003). Use of WinBugs for  
5 derivation of a distribution of BMDs for a single multistage model has been demonstrated by  
6 Kopylev et al. (2007), and this approach can be straightforwardly generalized to derive the  
7 distribution of BMDs for the combined tumor load. For further details on the implementation of  
8 this approach and for the results of the analyses, see Appendix G.  
9

#### 5.2.1.1.2. Rodent Dose-Response Analyses: Dosimetry

10 In modeling the applied doses (or exposures), default dosimetry procedures were applied  
11 to convert applied rodent doses to human equivalent doses. Essentially, for inhalation exposures,  
12 “ppm equivalence” across species was assumed. For oral doses,  $\frac{3}{4}$ -power body-weight scaling  
13 was used, with a default average human body weight of 70 kg. See Appendix G for more details  
14 on the default dosimetry procedures.

15 In addition to applied doses, several internal dose-metrics were used in the dose-response  
16 modeling for each tumor type. Use of internal dose-metrics in dose-response modeling is  
17 described here briefly. For more details on the PBPK modeling used to estimate the levels of the  
18 dose-metrics corresponding to different exposure scenarios in rodents and humans, as well as a  
19 qualitative discussion of the uncertainties and limitations of the model, see Section 3.5; for a  
20 more detailed discussion of how the dose-metrics were used in dose-response modeling, see  
21 Appendix G. Quantitative analyses of the uncertainties and their implications for dose-response  
22 assessment, utilizing the results of the Bayesian analysis of the PBPK model, are discussed  
23 separately in Section 5.2.1.4.2.  
24

#### 5.2.1.1.3. Selection of dose-metrics for different tumor types

25 One area of scientific uncertainty in cancer dose-response assessment is the appropriate  
26 scaling between rodent and human doses for equivalent responses. As discussed above, for  
27 applied dose, the standard dosimetry assumptions for equal lifetime carcinogenic risk are, for  
28 inhalation exposure, the same lifetime exposure concentration in air, and, for oral exposure, the  
29 same lifetime daily dose scaled by body weight to the  $\frac{3}{4}$  power. In this assessment, the  
30 cross-species scaling methodology, grounded in the principles of allometric variation of biologic  
31 processes, is used for describing pharmacokinetic equivalence (Allen and Fisher, 1993; Allen et

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 al., 1987; Crump et al., 1989; "Supplementary data for TCE assessment: Rat population  
2 example," 2011; U.S. EPA, 1992, 2005c) U.S. EPA 2011. Briefly, in the absence of adequate  
3 information to the contrary, the methodology determines pharmacokinetic equivalence across  
4 species through equal average lifetime concentrations or AUCs of the toxicant. Thus, in cases  
5 where the PBPK model can predict internal concentrations of the active moiety, equivalent daily  
6 AUCs are assumed to address cross-species pharmacokinetics. For cancer assessments, there is  
7 currently no adjustment for pharmacodynamic differences.

8 More detailed discussion of the cross-species scaling methodology, and its implications  
9 for dose-metric selection, was presented for the noncancer dose-response analyses in Section  
10 5.1.3.1, and those details are not repeated here.

11 To summarize, the preferred dose-metric under this methodology is equivalent daily  
12 AUC of the active moiety (parent compound or metabolite). For metabolites, in cases where the  
13 rate of production, but not the rate of clearance, of the active moiety can be estimated, the  
14 preferred dose-metric is the rate of metabolism (through the appropriate pathway) scaled by body  
15 weight to the  $\frac{3}{4}$  power. If there are sufficient data to consider the active metabolite moiety(ies)  
16 "reactive" and cleared through nonbiological processes, then the preferred dose-metric is the rate  
17 of metabolism (through the appropriate pathway) scaled by the tissue mass. Finally, if local  
18 metabolism is thought to be involved but cannot be estimated with the available data, then the  
19 AUC of the parent compound in blood is considered an appropriate surrogate and thus the  
20 preferred dose-metric.

21 Generally, an attempt was made to use tissue-specific dose-metrics representing  
22 particular pathways or metabolites identified from available data as having a likely role in the  
23 induction of a tissue-specific cancer. Where insufficient information was available to establish  
24 particular metabolites or pathways of likely relevance to a tissue-specific cancer, more general  
25 "upstream" metrics representing either parent compound or total metabolism had to be used. In  
26 addition, the selection of dose-metrics was limited to metrics that could be adequately estimated  
27 by the PBPK model (see Section 3.5). The (PBPK-based) dose-metrics used for the different  
28 tumor types are listed in Table 5-34. For each tumor type, the "primary" dose-metric referred to  
29 in Table 5-34 is the metric representing the particular metabolite or pathway whose involvement  
30 in carcinogenicity has the greatest biological support, whereas "alternative" dose-metrics  
31 represent upstream metabolic pathways (or TCE distribution, in the case of AUCCBld) that may  
32 be more generally involved.



#### 5.2.1.1.3.1. **Kidney**

1 As discussed in Sections 4.4.6–4.4.7, there is sufficient evidence to conclude that  
2 TCE-induced kidney tumors in rats are primarily caused by GSH-conjugation metabolites either  
3 produced in situ in or delivered systemically to the kidney. As discussed in Section 3.3.3.2,  
4 bioactivation of these metabolites within the kidney, either by beta-lyase, FMO, or P450s,  
5 produces reactive species. Therefore, multiple lines of evidence support the conclusion that  
6 renal bioactivation of DCVC is the preferred basis for internal dose extrapolations of  
7 TCE-induced kidney tumors. However, uncertainties remain as to the relative contributions from  
8 each bioactivation pathway, and quantitative clearance data necessary to calculate the  
9 concentration of each species are lacking. Moreover, the estimates of the amount bioactivated  
10 are indirect, derived from the difference between overall GSH conjugation flux and NAcDCVC  
11 excretion (see Section 3.5.7.3.1).

12 The rationales for the dose-metrics for kidney tumors are the same as for kidney  
13 noncancer toxicity, discussed above in Section 5.1.3.1.1, and not repeated here. The primary  
14 internal dose-metric for TCE-induced kidney tumors is the weekly rate of DCVC bioactivation  
15 per unit body weight to the  $\frac{3}{4}$  power (**ABioactDCVCBW $\frac{3}{4}$**  [mg/kg $\frac{3}{4}$ /week]). Due to the larger  
16 relative kidney weight in rats as compared to humans, using the alternative scaling by kidney  
17 weight instead of body weight to the  $\frac{3}{4}$  power would only change the quantitative interspecies  
18 extrapolation by about twofold,<sup>43</sup> so the sensitivity of the results to the scaling choice is  
19 relatively small. An alternative dose-metric that also involves the GSH conjugation pathway is  
20 the amount of GSH conjugation scaled by the  $\frac{3}{4}$  power of body weight (**AMetGSHBW $\frac{3}{4}$**   
21 [mg/kg $\frac{3}{4}$ /week]). This dose-metric uses the total flux of GSH conjugation as the  
22 toxicologically-relevant dose, and, thus, incorporates any direct contributions from DCVG and  
23 DCVC, which are not addressed in the DCVC bioactivation metric. Another alternative  
24 dose-metric is the total amount of TCE metabolism (oxidation and GSH conjugation together)  
25 scaled by the  $\frac{3}{4}$  power of body weight (**TotMetabBW $\frac{3}{4}$**  [mg/kg $\frac{3}{4}$ /week]). This dose-metric  
26 uses the total flux of TCE metabolism as the toxicologically relevant dose, and, thus,  
27 incorporates the possible involvement of oxidative metabolites, acting either additively or  
28 interactively, in addition to GSH conjugation metabolites in nephrocarcinogenicity (see  
29 Section 4.4.6). While there is no evidence that TCE oxidative metabolites can on their own  
30 induce kidney cancer, some nephrotoxic effects attributable to oxidative metabolites (e.g.,  
31 peroxisome proliferation) may modulate the nephrocarcinogenic potency of GSH metabolites.

---

<sup>43</sup>The range of the difference is 2.1–2.4-fold using the posterior medians for the relative kidney weight in rats and humans from the PBPK model described in Section 3.5 (see Table 3-36) and body weights of 0.3–0.4 kg for rats and 60–70 kg for humans.

1 However, this dose-metric is given less weight than those involving GSH conjugation because,  
2 as discussed in Sections 4.4.6 and 4.4.7, the weight of evidence supports the conclusion that  
3 GSH conjugation metabolites play a predominant role in nephrocarcinogenicity.  
4

#### 5.2.1.1.3.2. Liver

5 As discussed in Section 4.5.6, there is substantial evidence that oxidative metabolism is  
6 involved in TCE hepatocarcinogenicity, based primarily on noncancer and cancer effects similar  
7 to those observed with TCE being observed with a number of oxidative metabolites of TCE (e.g.,  
8 CH, TCA, and DCA). While TCA is a stable, circulating metabolite, CH and DCA are relatively  
9 short-lived, although enzymatically cleared (see Section 3.3.3.1). As discussed in Sections 4.5.6  
10 and 4.5.7, there is now substantial evidence that TCA does not adequately account for the  
11 hepatocarcinogenicity of TCE; therefore, unlike in previous dose-response analyses (Clewell and  
12 Andersen, 2004; Rhomberg, 2000), the AUC of TCA in plasma and in liver were not considered  
13 as dose-metrics. However, there are inadequate data across species to quantify the dosimetry of  
14 CH and DCA, and other intermediates of oxidative metabolism (such as TCE-oxide or  
15 dichloroacetylchloride) also may be involved in carcinogenicity. Thus, due to uncertainties as to  
16 the active moiety(ies), but the strong evidence associating TCE liver effects with oxidative  
17 metabolism in the liver, hepatic oxidative metabolism is the preferred basis for internal dose  
18 extrapolations of TCE-induced liver tumors.

19 The rationales for the dose-metrics for liver tumors are the same as for liver noncancer  
20 toxicity, discussed above in Section 5.1.3.1.2, and not repeated here. The primary internal  
21 dose-metric for TCE-induced liver tumors is selected to be the weekly rate of hepatic oxidation  
22 per unit body weight to the  $\frac{3}{4}$  power (**AMetLiv1BW<sup>3/4</sup> [mg/kg<sup>3/4</sup>/week]**). Due to the larger  
23 relative liver weight in mice as compared to humans, scaling by liver weight instead of body  
24 weight to the  $\frac{3}{4}$  power would only change the quantitative interspecies extrapolation by about  
25 fourfold,<sup>44</sup> so the sensitivity of the results to the scaling choice is relatively modest. The total  
26 amount of oxidative metabolism of TCE scaled by the  $\frac{3}{4}$  power of body weight  
27 (**TotOxMetabBW<sup>3/4</sup> [mg/kg<sup>3/4</sup>/week]**) was selected as an alternative dose-metric (the  
28 justification for the body weight to the  $\frac{3}{4}$  power scaling is analogous to that for hepatic oxidative  
29 metabolism, above). This dose-metric accounts for the possible additional contributions of  
30 systemically delivered products of lung oxidation.

---

<sup>44</sup>The range of the difference is 3.5–3.9-fold using the posterior medians for the relative liver weight in mice and humans from the PBPK model described in Section 3.5 (see Table 3-37) and body weights of 0.03–0.04 kg for mice and 60–70 kg for humans.

### 5.2.1.1.3.3.Lung

2 As discussed in Section 4.7.3, in situ oxidative metabolism in the respiratory tract may be  
3 more important to lung toxicity than systemically delivered metabolites, at least as evidenced by  
4 acute pulmonary toxicity. While chloral was originally implicated as the active metabolite,  
5 based on either acute toxicity or mutagenicity of chloral and/or chloral hydrate, more recent  
6 evidence suggests that other oxidative metabolites may also contribute to lung toxicity. These  
7 data include the identification of dichloroacetyl lysine adducts in Clara cells (Forkert et al.,  
8 2006), and the induction of pulmonary toxicity by TCE in CYP2E1-null mice, which may  
9 generate a different spectrum of oxidative metabolites as compared to wild-type mice  
10 (respiratory tract tissue also contains P450s from the CYP2F family). Overall, the weight of  
11 evidence supports the selection of respiratory tract oxidation of TCE as the preferred basis for  
12 internal dose extrapolations of TCE-induced lung tumors. However, uncertainties remain as to  
13 the relative contributions from different oxidative metabolites, and quantitative clearance data  
14 necessary to calculate the concentration of each species are lacking.

15 Under the cross-species scaling methodology, the rate of respiratory tract oxidation  
16 would be scaled by body weight to the  $\frac{3}{4}$  power. For chloral, as discussed in Section 4.7.3, the  
17 reporting of substantial TCOH but no detectable chloral hydrate in blood following TCE  
18 exposure from experiments in isolated, perfused lungs (Dalbey and Bingham, 1978) support the  
19 conclusion that chloral does not leave the target tissue in substantial quantities, but that there is  
20 substantial clearance by enzyme-mediated biotransformation. Dichloroacetyl chloride is a  
21 relatively-short-lived intermediate from aqueous (nonenzymatic) decomposition of TCE-oxide  
22 that can be trapped with lysine or degrade further to form DCA, among other products (Cai and  
23 Guengerich, 1999). Cai and Guengerich (1999) reported a half-life of TCE-oxide under aqueous  
24 conditions of 12 s at 23EC, a time-scale that would be shorter at physiological conditions (37EC)  
25 and that includes formation of dichloroacetyl chloride as well as its decomposition. Therefore,  
26 evidence for this metabolite suggests its clearance both is sufficiently rapid so that it would  
27 remain at the site of formation and is nonenzymatically mediated so that its rate would be  
28 independent of body weight. Other oxidative metabolites may also play a role, but, because they  
29 have not been identified, no inferences can be made as to their clearance.

30 Therefore, because it is not clear what the contributions to TCE-induced lung tumors are  
31 from different oxidative metabolites produced in situ and the scaling by body weight to the  
32  $\frac{3}{4}$  power is supported for at least one of the possible active moieties, it was decided here to scale  
33 the rate of respiratory tract tissue oxidation of TCE by body weight to the  $\frac{3}{4}$  power. The primary

1 internal dose-metric for TCE-induced lung tumors is, thus, the weekly rate of respiratory tract  
2 oxidation per unit body weight to the  $\frac{3}{4}$  power (**AMetLngBW34 [mg/kg<sup>3/4</sup>/week]**). It should be  
3 noted that, due to the larger relative respiratory tract tissue weight in mice as compared to  
4 humans, scaling by tissue weight instead of body weight to the  $\frac{3}{4}$  power would change the  
5 quantitative interspecies extrapolation by less than twofold,<sup>45</sup> so the sensitivity of the results to  
6 the scaling choice is relatively small.

7 While there is substantial evidence that acute pulmonary toxicity is related to pulmonary  
8 oxidative metabolism, for carcinogenicity, it is possible that, in addition to locally produced  
9 metabolites, systemically-delivered oxidative metabolites also play a role. Therefore, total  
10 oxidative metabolism scaled by the  $\frac{3}{4}$  power of body weight (**TotOxMetabBW34**  
11 **[mg/kg<sup>3/4</sup>/week]**) was selected as an alternative dose-metric (the justification for the body weight  
12 to the  $\frac{3}{4}$  power scaling is analogous to that for respiratory tract oxidative metabolism, above).

13 Another alternative dose-metric considered here is the AUC of TCE in blood (**AUCCBld**  
14 **[mg-hour/L/week]**). This dose-metric would account for the possibility that local metabolism is  
15 determined primarily by TCE delivered in blood via systemic circulation to pulmonary tissue  
16 (the flow rate of which scales as body weight to the  $\frac{3}{4}$  power), as assumed in previous PBPK  
17 models, rather than TCE delivered in air via diffusion to the respiratory tract, as is assumed in  
18 the PBPK model described in Section 3.5. However, as discussed in Section 3.5 and  
19 Appendix A, the available pharmacokinetic data provide greater support for the updated model  
20 structure. This dose-metric also accounts for the possible role of TCE itself in pulmonary  
21 carcinogenicity (consistent with the assumption that the same average concentration of TCE in  
22 blood will lead to a similar lifetime cancer risk across species).

#### 5.2.1.1.3.4. Other sites

24 For all other sites listed in Table 5-34, there is insufficient information for site-specific  
25 determinations of appropriate dose-metrics. While TCE metabolites and/or metabolizing  
26 enzymes have been reported in some of these tissues (e.g., male reproductive tract), their roles in  
27 carcinogenicity for these specific sites have not been established. Although “primary” and  
28 “alternative” dose-metrics are defined, they do not differ appreciably in their degrees of  
29 plausibility.

---

<sup>45</sup>The range of the difference is 1.6–1.8-fold using the posterior medians for the relative respiratory tract tissue weight in mice and humans from the PBPK model described in Section 3.5 (see Table 3-37), and body weights of 0.03–0.04 kg for mice and 60–70 kg for humans.

1           Given that the majority of the toxic and carcinogenic responses to TCE appear to be  
2 associated with metabolism, total metabolism of TCE scaled by the  $\frac{3}{4}$  power of body weight was  
3 selected as the primary dose-metric (**TotMetabBW<sup>3/4</sup> [mg/kg<sup>3/4</sup>/week]**). This dose-metric uses  
4 the total flux of TCE metabolism as the toxicologically-relevant dose, and, thus, incorporates the  
5 possible involvement of any TCE metabolite in carcinogenicity.

6           An alternative dose-metric considered here is the AUC of TCE in blood. This  
7 dose-metric would account for the possibility that the determinant of carcinogenicity is local  
8 metabolism, governed primarily by TCE delivered in blood via systemic circulation to the target  
9 tissue (the flow rate of which scales as body weight to the  $\frac{3}{4}$  power). This dose-metric also  
10 accounts for the possible role of TCE itself in carcinogenicity (consistent with the assumption  
11 that the same average concentration of TCE in blood will lead to a similar lifetime cancer risk  
12 across species).


#### 5.2.1.1.4. Methods for dose-response analyses using internal dose-metrics

14           As shown in Figure 5-5, the general approach taken for the use of internal dose-metrics in  
15 dose-response modeling was to first apply the rodent PBPK model to obtain rodent values for the  
16 dose-metrics corresponding to the applied doses in a bioassay. Then, dose-response modeling  
17 for a tumor response was performed using the internal dose-metrics and the multistage model or  
18 the survival-adjusted modeling approaches described above to obtain a BMD and BMDL in  
19 terms of the dose-metric. On an internal dose basis, humans and rodents are presumed to have  
20 similar lifetime cancer risks, and the relationship between human internal and external doses is  
21 essentially linear at low doses up to 0.1 mg/kg/day or 0.1 ppm, and nearly linear up to  
22 10 mg/kg/day or 10 ppm. Therefore, the BMD and BMDL were then converted human  
23 equivalent doses (or exposures) using conversion ratios estimated from the human PBPK model  
24 at 0.001 mg/kg/day or 0.001 ppm (see Table 5-35). Because the male and female conversions  
25 differed by less than 11%, the human BMDLs were derived using the mean of the sex-specific  
26 conversion factors (except for testicular tumors, for which only male conversion factors were  
27 used). Finally, a slope factor or unit risk estimate for that tumor response was derived from the  
28 human “BMDLs” as described above (i.e., BMR/BMDL). Note that the converted “BMDs” and  
29 “BMDLs” are not actually human equivalent BMDs and BMDLs corresponding to the BMR  
30 because the conversion was not made in the dose range of the BMD; the converted BMDs and  
31 BMDLs are merely intermediaries to obtain a converted slope factor or unit risk estimate. In  
32 addition, it should be noted that median values of dose-metrics were used for rodents, whereas  
33 mean values were used for humans. Because the rodent population model characterizes

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 study-to-study variation, animals of the same sex/species/strain combination within a study were  
2 assumed to be identical. Therefore, use of median dose-metric values for rodents can be  
3 interpreted as assuming that the animals in the bioassay were all “typical” animals and the  
4 dose-response model is estimating a “risk to the typical rodent.” In practice, the use of median  
5 or mean internal doses for rodents did not make much difference except when the uncertainty in  
6 the dose-metric was high (e.g., AMetLungBW34 dose-metric in mice). A quantitative analysis  
7 of the impact of the uncertainty in the rodent PBPK dose-metrics is included in Section 5.2.1.4.2.  
8 On the other hand, the human population model characterizes individual-to-individual variation.  
9 Because the quantity of interest is the human population mean risk, the expected value  
10 (averaging over the uncertainty) of the population mean (averaging over the variability)  
11 dose-metric was used for the conversion  
12

13 **Figure 5-5. Flow-chart for dose-response analyses of rodent bioassays using**  
14 **PBPK model-based dose-metrics. Square nodes indicate point values,**  
15 **circular nodes indicate distributions, and the inverted triangles indicate a**  
16 **(deterministic) functional relationship.**



1  
2  
3  
4  
5

**Table 5-35. Mean PBPK model predictions for weekly internal dose in humans exposed continuously to low levels of TCE via inhalation (ppm) or orally (mg/kg/day)**

*This document is a draft for review purposes only and does not constitute Agency policy.*

Dose-metric	0.001 ppm		0.001 mg/kg/day	
	Female	Male	Female	Male
ABioactDCVCBW34	0.00324	0.00324	0.00493	0.00515
AMetGSHBW34	0.00200	0.00200	0.00304	0.00318
AMetLiv1BW34	0.00703	0.00683	0.0157	0.0164
AMetLngBW34	0.00281	0.00287	$6.60 \times 10^{-5}$	$6.08 \times 10^{-5}$
AUCCBld	0.00288	0.00298	0.000411	0.000372
TotMetabBW34	0.0118	0.0117	0.0188	0.0196
TotOxMetabBW34	0.00984	0.00970	0.0157	0.0164

1  
2 See note to Table 5-34 for dose-metric abbreviations. Values represent the mean of the (uncertainty) distribution of  
3 population means for each sex and exposure scenario, generated from Monte Carlo simulation of 500 populations of  
4 500 individuals each.

5  
6  
7 to human slope factor or unit risks. Therefore, the extrapolated slope factor or unit risk estimates  
8 can be interpreted as the expected “average risk” across the population based on rodent  
9 bioassays.

#### 10 11 **5.2.1.1.5. Rodent Dose-Response Analyses: Results**

12 A summary of the PODs and slope factor and unit risk estimates for each  
13 sex/species/bioassay/tumor type is presented in Tables 5-36 (inhalation studies) and 5-37 (oral  
14 studies). The PODs for individual tumor types were extracted from the modeling results in the  
15 figures in Appendix G. For the applied dose (default dosimetry) analyses, the POD is the BMDL  
16 from the male human (“M”) BMDL entry at the top of the figure for the selected model; male  
17 results were extracted because the default weight for males in the PBPK modeling is 70 kg,  
18 which is the overall human weight in EPA’s default dosimetry methods (for inhalation, male and  
19 female results are identical). As described in Section 5.2.1.2 above, for internal dose-metrics,  
20 male and female results were averaged, and the converted human “BMDLs” are not true BMDLs  
21 because they were converted outside the linear range of the PBPK models. It can be seen in  
22 Appendix G that the male and female results were similar for all the dose-metrics.

23 For two data sets, the highest dose (exposure) group was dropped to get a better fit when  
24 using applied doses. This technique can improve the fit when the response tends to plateau with  
increasing dose. Plateauing typically occurs when metabolic saturation alters the pattern of



Table 5-36. Summary of PODs and unit risk estimates for each sex/species/bioassay/tumor type (inhalation)

Study	Tumor type	BMR	PODs (ppm, in human equivalent exposures) <sup>a</sup>							
			Applied dose	AUC CBld	TotMetab BW34	TotOxMetab BW34	AMetLng BW34	AMetLiv1 BW34	AMetGSH BW34	ABioact DCVCBW34
<b>Female mouse</b>										
Fukuda	Lung AD + CARC	0.1	26.3	55.5		31.3	38.8			
Henschler	Lymphoma	0.1	11.0 <sup>b</sup>	-- <sup>b</sup>	9.84					
Maltoni	Lung AD + CARC	0.1	44.6	96.6		51.4	55.7			
	Liver	0.05	37.1			45.8		41.9		
	Combined	0.05	15.7			20.7				
<b>Male mouse</b>										
Maltoni	Liver	0.1	34.3			51		37.9		
<b>Male rat</b>										
Maltoni	Leukemia	0.05	28.2 <sup>c</sup>	-- <sup>b</sup>	28.3					
	Kidney AD + CARC	0.01	22.7		13.7				0.197	0.121
	Leydig cell	0.1	18.6 <sup>c</sup>	-- <sup>d</sup>	18.1					
	Combined	0.01	1.44		1.37					
<b>Unit risk estimate (ppm<sup>-1</sup>)<sup>e</sup></b>										
Study	Tumor type	Applied dose	AUC CBld	TotMetab BW34	TotOxMetab BW34	AMetLng BW34	AMetLiv1 BW34	AMetGSH BW34	ABioact DCVCBW34	
<b>Female mouse</b>										
Fukuda	Lung AD + CARC	$3.8 \times 10^{-3}$	$1.8 \times 10^{-3}$		$3.2 \times 10^{-3}$	$2.6 \times 10^{-3}$				
Henschler	Lymphoma	$9.1 \times 10^{-3}$		$1.0 \times 10^{-2}$						
Maltoni	Lung AD + CARC	$2.2 \times 10^{-3}$	$1.0 \times 10^{-3}$		$1.9 \times 10^{-3}$	$1.8 \times 10^{-3}$				
	Liver	$1.3 \times 10^{-3}$			$1.1 \times 10^{-3}$		$1.2 \times 10^{-3}$			
	Combined	$3.2 \times 10^{-3}$			$2.4 \times 10^{-3}$					

**Table 5-36. Summary of PODs and unit risk estimates for each sex/species/bioassay/tumor type (inhalation) (continued)**

Study	Tumor type	Unit risk estimate (ppm <sup>-1</sup> ) <sup>e</sup>							
		Applied dose	AUC CBld	TotMetab BW34	TotOxMetab BW34	AMetLng BW34	AMetLiv1 BW34	AMetGSH BW34	ABioact DCVCBW34
<b>Male mouse</b>									
Maltoni	Liver	$2.9 \times 10^{-3}$			$2.0 \times 10^{-3}$		$2.6 \times 10^{-3}$		
<b>Male rat</b>									
Maltoni	Leukemia	$1.8 \times 10^{-3}$		<b><math>1.8 \times 10^{-3}</math></b>					
	Kidney AD + CARC	$4.4 \times 10^{-4}$		$7.3 \times 10^{-4}$				$5.1 \times 10^{-2}$	<b><math>8.3 \times 10^{-2}</math></b>
	Leydig cell	$5.4 \times 10^{-3}$		<b><math>5.5 \times 10^{-3}</math></b>					
	Combined	$7.0 \times 10^{-3}$		$7.3 \times 10^{-3}$					

<sup>a</sup>For the applied doses, the PODs are BMDLs. However, for the internal dose-metrics, the PODs are not actually human equivalent BMDLs corresponding to the BMR because the interspecies conversion does not apply to the dose range of the BMDL; the converted BMDLs are merely intermediaries to obtain a converted unit risk estimate. The calculation that was done is equivalent to using linear extrapolation from the BMDLs in terms of the internal dose-metric to get a unit risk estimate for low-dose risk in terms of the internal dose-metric and then converting that estimate to a unit risk estimate in terms of human equivalent exposures. The PODs reported here are what one would get if one then used the unit risk estimate to calculate the human exposure level corresponding to a 10% extra risk, but the unit risk estimate is not intended to be extrapolated upward out of the low-dose range, e.g., above  $10^{-4}$  risk. In addition, for the internal dose-metrics, the PODs are the average of the male and female human "BMDL" results presented in Appendix G.

<sup>b</sup>Inadequate fit to control group, but the primary metric, TotMetabBW34, fits adequately.

<sup>c</sup>Dropped highest-dose group to improve model fit.

<sup>d</sup>Inadequate overall fit.

<sup>e</sup>Unit risk estimate = BMR/POD. Results for the primary dose-metric are in bold.

AD = adenoma, CARC = carcinoma.

Table 5-37. Summary of PODs and slope factor estimates for each sex/species/bioassay/tumor type (oral)

Study	Tumor type	BMR	PODs (mg/kg/day, in human equivalent doses) <sup>a</sup>							
			Applied dose	AUC CB1d	TotMetab BW34	TotOxMetab BW34	AMetLng BW34	AMetLiv1 BW34	AMetGSH BW34	ABioact DCVCBW34
<b>Female mouse</b>										
NCI	Liver carc	0.1	26.5			17.6		14.1		
	Lung AD + CARC	0.1	41.1	682		24.7	757			
	Leukemias + sarcomas	0.1	43.1	733	20.6					
	Combined	0.05	7.43			5.38				
<b>Male mouse</b>										
NCI	Liver carc	0.1	8.23			4.34		3.45		
<b>Female rat</b>										
NTP (1988)	Leukemia	0.05	72.3	3,220	21.7					
<b>Male rat</b>										
NTP (1990) <sup>c</sup>	Kidney AD + CARC	0.1	32		11.5				0.471	0.292
NTP (1988)										
Marshall <sup>d</sup>	Testicular	0.1	3.95	167	1.41					
August	Subcut sarcoma	0.05	60.2	2,560	21.5					
Osborne-Mendel <sup>c</sup>	Kidney AD + CARC	0.1	41.5		14.3				0.648	0.402
<b>Female mouse</b>										
NCI	Liver carc		$3.8 \times 10^{-3}$			$5.7 \times 10^{-3}$		$7.1 \times 10^{-3}$		
	Lung AD + CARC		$2.4 \times 10^{-3}$	$1.5 \times 10^{-4}$		$4.0 \times 10^{-3}$	$1.3 \times 10^{-4}$			
	Leukemias + sarcomas		$2.3 \times 10^{-3}$	$1.4 \times 10^{-4}$	$4.9 \times 10^{-3}$					
	Combined		$6.7 \times 10^{-3}$			$9.3 \times 10^{-3}$				

**Table 5-37. Summary of PODs and slope factor estimates for each sex/species/bioassay/tumor type (oral)  
(continued)**

Study	Tumor type	Slope factor estimate (mg/kg/day) <sup>-1</sup> <sup>b</sup>							
		Applied dose	AUC CBId	TotMetab BW34	TotOxMetab BW34	AMetLng BW34	AMetLiv1 BW34	AMetGSH BW34	ABioact DCVCBW34
<b>Male mouse</b>									
NCI	Liver carc	$1.2 \times 10^{-2}$			$2.3 \times 10^{-2}$		$2.9 \times 10^{-2}$		
<b>Female rat</b>									
NTP (1988)	Leukemia	$6.9 \times 10^{-4}$	$1.6 \times 10^{-5}$	<b><math>2.3 \times 10^{-3}</math></b>					
<b>Male rat</b>									
NTP (1990) <sup>c</sup>	Kidney AD + CARC	$1.6 \times 10^{-3}$		$4.3 \times 10^{-3}$				$1.1 \times 10^{-1}$	<b><math>1.7 \times 10^{-1}</math></b>
NTP (1988)									
Marshall <sup>d</sup>	Testicular	$2.5 \times 10^{-2}$	$6.0 \times 10^{-4}$	<b><math>7.1 \times 10^{-2}</math></b>					
August	Subcut sarcoma	$8.3 \times 10^{-4}$	$2.0 \times 10^{-5}$	<b><math>2.3 \times 10^{-3}</math></b>					
Osborne-Mendel <sup>c</sup>	Kidney AD + CARC	$2.4 \times 10^{-3}$		$7.0 \times 10^{-3}$				$1.5 \times 10^{-1}$	<b><math>2.5 \times 10^{-1}</math></b>

<sup>a</sup>For the applied doses, the PODs are BMDLs. However, for the internal dose-metrics, the PODs are not actually human equivalent BMDLs corresponding to the BMR because the interspecies conversion does not apply to the dose range of the BMDL; the converted BMDLs are merely intermediaries to obtain a converted slope factor estimate. The calculation that was done is equivalent to using linear extrapolation from the BMDLs in terms of the internal dose-metric to get a slope factor estimate for low-dose risk in terms of the internal dose-metric and then converting that estimate to a slope factor estimate in terms of human equivalent doses. The PODs reported here are what one would get if one then used the slope factor estimate to calculate the human dose level corresponding to a 10% extra risk, but the slope factor estimate is not intended to be extrapolated upward out of the low-dose range, e.g., above  $10^{-4}$  risk. In addition, for the internal dose-metrics, the PODs are the average of the male and female human “BMDL” results presented in Appendix G.

<sup>b</sup>Slope factor estimate = BMR/POD. Results for the primary dose-metric are in bold.

<sup>c</sup>Using MSW adjusted incidences (see text and Table 5-38).

<sup>d</sup>Using poly-3 adjusted incidences (see text and Table 5-38).

AD = adenoma, CARC = carcinoma.

1 metabolite formation or when survival is impacted at higher doses, and it is assumed that these  
2 high-dose responses are less relevant to low-dose risk. The highest-dose group was not dropped  
3 to improve the fit for any of the internal dose-metrics because it was felt that if the dose-metric  
4 was an appropriate reflection of internal dose of the reactive metabolite(s), then use of the  
5 dose-metric should have ameliorated the plateauing in the dose-response relationship (note that  
6 survival-impacted data sets were addressed using survival adjustment techniques). For a 3<sup>rd</sup> data  
7 set (Henschler lymphomas), it might have helped to drop the highest exposure group, but there  
8 were only two exposure groups, so this was not done. As a result, the selected model, although it  
9 had an adequate fit overall, did not fit the control group very well (the model estimated a higher  
10 background response than was observed); thus, the BMD and BMDL were likely overestimated  
11 and the risk underestimated. The estimates from the NCI (1976) oral male mouse liver cancer  
12 data set are also somewhat more uncertain because the response rate was extrapolated down from  
13 a response rate of about 50% extra risk to the BMR of 10% extra risk.

14 Some general patterns can be observed in Tables 5-36 and 5-37. For inhalation, the unit  
15 risk estimates for different dose-metrics were generally similar (within about 2.5-fold) for most  
16 tumor types. The exception was for kidney cancer, where the estimates varied by over 2 orders  
17 of magnitude, with the AMetGSHBW34 and ABioactDCVCBW34 metrics yielding the highest  
18 estimates. This occurs because pharmacokinetic data indicate, and the PBPK model predicts,  
19 substantially more GSH conjugation (as a fraction of intake), and hence subsequent  
20 bioactivation, in humans relative to rats. The range of the risk estimates for individual tumor  
21 types overall (across tumor types and dose-metrics) was encompassed by the range of estimates  
22 across the dose-metrics for kidney cancer in the male rat, which was from  $4.4 \times 10^{-4}$  per ppm  
23 (applied dose) to  $8.3 \times 10^{-2}$  per ppm (ABioactDCVCBW34).

24 For oral exposure, the slope factor estimates are more variable across dose-metrics  
25 because of first-pass effects in the liver (median estimates for the fraction of TCE metabolized in  
26 *one* pass through the liver in mice, rats, and humans are >0.8). Here, the exception is for the risk  
27 estimates for cancer of the liver itself, which are also within about a 2.5-fold range, because the  
28 liver gets the full dose of all the metrics during that “first pass.” For the other tumor types, the  
29 range of estimates across dose-metrics varies from about 30-fold to over two orders of  
30 magnitude, with the estimates based on AUCCBld and AMetLngBW34 being at the low end and  
31 those based on AMetGSHBW34 and ABioactDCVCBW34 again being at the high end. For  
32 AUCCBld, the PBPK model predicted the blood concentrations to scale more closely to body  
33 weight rather than the  $\frac{3}{4}$  power of body weight, so the extrapolated human unit risks using this  
34 dose-metric are smaller than those obtained by applied dose or other dose-metrics that included  
35  $\frac{3}{4}$  power body weight scaling. For AMetLngBW34, pharmacokinetic data indicate, and the

1 PBPK model predicts, that the human respiratory tract metabolizes a lower fraction of total TCE  
2 intake than the mouse respiratory tract, so the extrapolated risk to humans based on this metric is  
3 lower than that obtained using applied dose or other dose-metrics. Overall, the oral slope factor  
4 estimates for individual tumor types ranged from  $1.6 \times 10^{-5}$  per mg/kg/day (female rat leukemia,  
5 AUCCB1d) to  $2.5 \times 10^{-1}$  per mg/kg/day (male Osborne-Mendel rat kidney,  
6 ABioactDCVCBW34), a range of over 4 orders of magnitude. It must be recognized, however,  
7 that not all dose-metrics are equally credible, and, as will be presented below, the slope factor  
8 estimates for total cancer risk for the most sensitive bioassay response for each sex/species  
9 combination using the primary (preferred) dose-metrics fall within a very narrow range.

10 Results for survival-adjusted analyses are summarized in Table 5-38. For the  
11 time-independent (BMDS) multistage model, the risk estimates using poly-3 adjustment are  
12 higher than those without poly-3 adjustment. This is to be expected because the  
13 poly-3 adjustment decreases denominators when accounting for early mortality, and, for these  
14 data sets, the higher-dose groups had greater early mortality. The difference was fairly modest  
15 for the kidney cancer data sets (about 30% higher) but somewhat larger for the testicular cancer  
16 data set (about 150% higher).

17 In addition, the MSW time-to-tumor model generated higher risk estimates than the  
18 poly-3 adjustment technique. The MSW results were about 40% higher for the NTP F344 rat  
19 kidney cancer data sets and about 60% higher for the NTP Osborne-Mendel rat kidney cancer  
20 data sets. For the NTP Marshall rat testicular cancer data set, the discrepancies were greater; the  
21 results ranged from about 100% to 180% higher for the different dose-metrics. As discussed in  
22 Section 5.2.1.1, these two approaches differ in the way they take early mortality into account.  
23 The poly-3 technique merely adjusts the tumor incidence denominators, using a constant power  
24 3 of time, to reflect the fact that animals are at greater risk of cancer at older ages. The MSW  
25 model estimates risk as a function of time (and dose), and it estimates the power (of time)  
26 parameter for each data set.<sup>46</sup> For the NTP F344 rat kidney cancer and NTP Marshall rat  
27 testicular cancer data sets, the estimated power parameter was close to 3 in each case, ranging  
28 from 3.0–3.7; for the NTP Osborne-Mendel rat kidney cancer data sets, however, the estimated  
29 power parameter was about 10 for each of the dose-metrics, presumably reflecting the fact that  
30 these were late-occurring tumors (the earliest occurred at 92 weeks). Using a higher power

---

46Conceptually, the approaches differ most when different tumor contexts (incidental or fatal) are considered, because the poly-3 technique only accounts for time of death, while the MSW model can account for the tumor context and attempt to estimate an induction time (t0), although this was not done for any of the data sets in this assessment.

1 parameter than 3 in the poly-3 adjustment would give even less weight to nontumor-bearing  
 2 animals that die early and would, thus, increase the adjusted incidence even more in the

3 **Table 5-38. Comparison of survival-adjusted results for 3 oral male rat data**  
 4 **sets<sup>a</sup>**  
 5

Dose-metric	Adjustment method	BMR	POD (mg/kg/day)	BMD:BMDL	Slope factor estimate (per mg/kg/day)
<b>NTP (1990) F344 rat kidney AD + CARC</b>					
Applied dose	unadj BMDS	0.05	56.9	1.9	$8.8 \times 10^{-4}$
	poly-3 BMDS	0.1	89.2	1.9	$1.1 \times 10^{-3}$
	MSW	0.05	32.0	2.6	$1.6 \times 10^{-3}$
TotMetabBW34	unadj BMDS	0.05	20.2	2.1	$2.5 \times 10^{-3}$
	poly-3 BMDS	0.1	31.8	1.7	$3.1 \times 10^{-3}$
	MSW	0.05	11.5	3.1	$4.3 \times 10^{-3}$
AMetGSHBW34	unadj BMDS	0.05	0.841	1.9	$5.9 \times 10^{-2}$
	poly-3 BMDS	0.1	1.32	1.9	$7.6 \times 10^{-2}$
	MSW	0.05	0.471	2.4	$1.1 \times 10^{-1}$
ABioactDCVCBW34	unadj BMDS	0.05	0.522	1.9	$9.6 \times 10^{-2}$
	poly-3 BMDS	0.1	0.817	1.9	$1.2 \times 10^{-1}$
	MSW	0.05	0.292	2.4	$1.7 \times 10^{-1}$
<b>NTP (1988) Osborne-Mendel rat kidney AD + CARC</b>					
Applied dose	unadj BMDS	0.1	86.6	1.7	$1.2 \times 10^{-3}$
	poly-3 BMDS	0.1	65.9	1.7	$1.5 \times 10^{-3}$
	MSW	0.1	41.5	2.0	$2.4 \times 10^{-3}$
TotMetabBW34	unadj BMDS	0.1	30.4	1.7	$3.3 \times 10^{-3}$
	poly-3 BMDS	0.1	23.1	1.7	$4.3 \times 10^{-3}$
	MSW	0.1	14.3	2.0	$7.0 \times 10^{-3}$
AMetGSHBW34	unadj BMDS	0.1	1.35	1.7	$7.4 \times 10^{-2}$
	poly-3 BMDS	0.1	1.03	1.7	$9.7 \times 10^{-2}$
	MSW	0.1	0.648	2.0	$1.5 \times 10^{-1}$
ABioactDCVCBW34	unadj BMDS	0.1	0.835	1.7	$1.2 \times 10^{-1}$
	poly-3 BMDS	0.1	0.636	1.7	$1.6 \times 10^{-1}$
	MSW	0.1	0.402	2.0	$2.5 \times 10^{-1}$
<b>NTP (1988) Marshall rat testicular tumors</b>					
Applied dose	unadj BMDS	0.1	9.94	1.4	$1.0 \times 10^{-2}$
	poly-3 BMDS	0.1	3.95	1.5	$2.5 \times 10^{-2}$
	MSW	0.1	1.64	5.2	$6.1 \times 10^{-2}$
AUCCBld	unadj BMDS	0.1	427	1.4	$2.3 \times 10^{-4}$
	poly-3 BMDS	0.1	167	1.6	$6.0 \times 10^{-4}$
	MSW	0.1	60.4	2.6	$1.7 \times 10^{-3}$
TotMetabBW34	unadj BMDS	0.1	3.53	4.3	$2.8 \times 10^{-2}$
	poly-3 BMDS	0.1	1.41	1.5	$7.1 \times 10^{-2}$
	MSW	0.1	0.73	9.4	$1.4 \times 10^{-1}$

6  
 7 <sup>a</sup>For the applied doses, the PODs are BMDLs. However, for the internal dose-metrics, the PODs are not actually  
 8 human equivalent BMDLs corresponding to the BMR because the interspecies conversion does not apply to the

1 dose range of the BMDL; the converted BMDLs are merely intermediaries to obtain a converted slope factor  
2 estimate. Results for the primary dose-metric are in bold.

3

4 AD = adenoma, CARC = carcinoma.

5



1 highest-dose groups where the early mortality is most pronounced, increasing the slope factor  
2 estimate. Nonetheless, as noted above, the MSW results were only about 60% higher for the  
3 NTP Osborne-Mendel rat kidney cancer data sets for which MSW estimated a power parameter  
4 of about 10.

5 In general, the risk estimates from the MSW model would be preferred because, as  
6 discussed above, this model incorporates more information (e.g., tumor context) and estimates  
7 the power parameter rather than using a constant value of three. From Table 5-38, it can be seen  
8 that the results from MSW yielded higher BMD:BMDL ratios than the results from the  
9 poly-3 technique. These ratios were only slightly higher and not unusually large for MSW  
10 model analyses of the NTP (1988, 1990) kidney tumor estimates, and this, along with the  
11 adequate fit (assessed visually) of the MSW model, supports using the slope factor estimates  
12 from the MSW modeling of rat kidney tumor incidence. On the other hand, the BMD:BMDL  
13 ratio was relatively large for the applied dose analysis and, in particular, for the preferred  
14 dose-metric analysis (9.4-fold) of the NTP Marshall rat testicular tumor data set. Therefore, for  
15 this endpoint, the poly-3-adjusted results were used, although they may underestimate risk  
16 somewhat as compared to the MSW model.

17 In addition to the results from dose-response modeling of individual tumor types, the  
18 results of the combined tumor risk analyses for the three bioassays in which the rodents exhibited  
19 increased risks at multiple sites are also presented in Tables 5-36 and 5-37, in the rows labeled  
20 “combined” under the column heading “Tumor Type.” These results were extracted from the  
21 detailed results in Appendix G. Note that, because of the computational complexity of the  
22 combined tumor analyses, dose-response modeling was only done using applied dose and a  
23 common upstream internal dose-metric, rather than using the different preferred dose-metrics for  
24 each tumor type within a combined tumor analysis.

25 For the Maltoni female mouse inhalation bioassay, the combined tumor risk estimates are  
26 bounded by the highest individual tumor risk estimates and the sums of the individual tumor  
27 risks estimates (the risk estimates are upper bounds, so the combined risk estimate, i.e., the upper  
28 bound on the sum of the individual central tendency estimates, should be less than the sum of the  
29 individual upper bound estimates), as one would expect. The common upstream internal  
30 dose-metric used for the combined analysis was TotOxMetabBW34, which is not the primary  
31 metric for either of the individual tumor types. For the liver tumors, the primary metric was  
32 AMetLiv1BW34, but as can be seen in Table 5-36, it yields results similar to those for  
33 TotOxMetabBW34. Likewise, for the lung tumors, the primary metric was AMetLngBW34,  
34 which yields a unit risk estimate slightly smaller than for TotOxMetabBW34. Thus, the results of

1 the combined analysis using TotOxMetabBW34 as a common metric is not likely to substantially  
2 over- or underestimate the combined risk based on preferred metrics for each of the tumor types.

3 For the Maltoni male rat inhalation bioassay, the combined risk estimates are also  
4 reasonably bounded, as expected. The common upstream internal dose-metric used for the  
5 combined analysis was TotMetabBW34, which is the primary metric for two of the  
6 three individual tumor types. However, as can be seen in Table 5-36, the risk estimate for the  
7 preferred dose-metric for the third tumor type, ABioactDCVCBW34 for the kidney tumors, is  
8 substantially higher than the risk estimates for the primary dose-metrics for the other two tumor  
9 types and would dominate a combined tumor risk estimate across primary dose-metrics; thus, the  
10 ABioactDCVCBW34-based kidney tumor risk estimate alone can reasonably be used to  
11 represent the total cancer risk for the bioassay using preferred internal dose-metrics, although it  
12 would underestimate the combined risk to some extent (e.g., the kidney-based estimate is  
13  $8.3 \times 10^{-2}$  per ppm; the combined estimate would be about  $9 \times 10^{-2}$  per ppm, rounded to  
14 one significant figure).

15 For the third bioassay (NCI female mouse oral bioassay), the combined tumor risk  
16 estimates are once again reasonably bounded. The common upstream internal dose-metric used  
17 for the combined analysis was TotOxMetabBW34, which is not the primary metric for any of the  
18 three individual tumor types but was considered to be the most suitable metric to apply as a basis  
19 for combining risk across these different tumor types. The slope factor estimate for the lung  
20 based on the primary dose-metric for that site becomes negligible compared to the estimates for  
21 the other two tumor types (see Table 5-37). However, the slope factor estimates for the  
22 remaining two tumor types are both somewhat underestimated using the TotOxMetabBW34  
23 metric rather than the primary metrics for those tumors (the TotOxMetabBW34-based estimate  
24 for leukemias + sarcomas, which is not presented in Table 5-30 because, in the absence of better  
25 mechanistic information, more upstream metrics were used for that individual tumor type, is  
26  $4.1 \times 10^{-3}$  per mg/kg/day). Thus, overall, the combined estimate based on TotOxMetabBW34 is  
27 probably a reasonable estimate for the total tumor risk in this bioassay, although it might  
28 overestimate risk slightly.

29 The most sensitive sex/species results are extracted from Tables 5-29 and 5-30 and  
30 presented in Tables 5-39 (inhalation) and 5-40 (oral) below. The BMD:BMDL ratios for all the  
31 results corresponding to the slope factor and unit risk estimates based on the preferred  
32 dose-metrics ranged from 1.3–2.1. For inhalation, the most sensitive bioassay responses based  
33 on the preferred dose-metrics ranged from  $2.6 \times 10^{-3}$  per ppm to  $8.3 \times 10^{-2}$  per ppm across the  
34 sex/species combinations (with the exception of the female rat, which exhibited no apparent

1 TCE-associated response in the 3 available bioassays). For oral exposure, the most sensitive  
 2 bioassay responses

3  
 4  
 5  
 6

**Table 5-39. Inhalation: most sensitive bioassay for each sex/species combination<sup>a</sup>**

Sex/species	Endpoint (study)	Unit risk per ppm		
		Preferred dose-metric	Default methodology	Alternative dose-metrics, studies, or endpoints
Female mouse	Lymphoma (Henschler et al., 1980)	$1.0 \times 10^{-2}$	$9.1 \times 10^{-3}$	$1 \times 10^{-3} \sim 4 \times 10^{-3}$
Male mouse	Liver hepatoma (Maltoni et al., 1986)	$2.6 \times 10^{-3}$	$2.9 \times 10^{-3}$	$2 \times 10^{-3}$
Female rat	–	–	–	–
Male rat	Leukemia+ Kidney AD and CARC+ Leydig cell tumors (Maltoni et al., 1986)	$8.3 \times 10^{-2}$	$7.0 \times 10^{-3}$	$4 \times 10^{-4} \sim 5 \times 10^{-2}$ [individual site results]

7  
 8 <sup>a</sup>Results extracted from Table 5-36.  
 9 AD = adenoma, CARC = carcinoma.

10  
 11  
 12  
 13

**Table 5-40. Oral: most sensitive bioassay for each sex/species combination<sup>a</sup>**

Sex/species	Endpoint (Study)	Unit risk per mg/kg/day		
		Preferred dose-metric	Default methodology	Alternative dose-metrics, studies, or endpoints
Female mouse	Liver CARC + lung AD and CARC+ sarcomas + leukemias (NCI, 1976)	$9.3 \times 10^{-3}$	$6.7 \times 10^{-3}$	$1 \times 10^{-4} \sim 7 \times 10^{-3}$ [individual site results]
Male mouse	Liver CARC (NCI, 1976)	$2.9 \times 10^{-2}$	$1.2 \times 10^{-2}$	$2 \times 10^{-2}$
Female rat	Leukemia (NTP, 1988)	$2.3 \times 10^{-3}$	$6.9 \times 10^{-4}$	$2 \times 10^{-5}$

*This document is a draft for review purposes only and does not constitute Agency policy.*

Male rat	Kidney AD + CARC (NTP, 1988, Osborne-Mendel)	$2.5 \times 10^{-1}$	$2.4 \times 10^{-3}{}^b$	$2 \times 10^{-5} \sim 2 \times 10^{-1}$
----------	---	----------------------	--------------------------	--

<sup>a</sup>Results extracted from Table 5-37.

<sup>b</sup>Most sensitive male rat result using default methodology is  $2.5 \times 10^{-2}$  per mg/kg/day for NTP (1988) Marshall rat testicular tumors.

AD = adenoma, CARC = carcinoma.

based on the preferred dose-metrics ranged from  $2.3 \times 10^{-3}$  per mg/kg/day to  $2.5 \times 10^{-1}$  per mg/kg/day across the sex/species combinations. For both routes of exposure, the most sensitive sex/species response was (or was dominated by, in the case of the combined tumors in the male rat by inhalation) male rat kidney cancer based on the preferred dose-metric of ABioactDCVCBW34.

#### 5.2.1.1.6. Uncertainties in Dose-Response Analyses of Rodent Bioassays

#### 5.2.1.1.7. Qualitative discussion of uncertainties

All risk assessments involve uncertainty, as study data are extrapolated to make inferences about potential effects in humans from environmental exposure. The largest sources of uncertainty in the TCE rodent-based cancer risk estimates are interspecies extrapolation and low-dose extrapolation. Some limited human (occupational) data from which to estimate human cancer risk are available, and cancer risk estimates based on these data are developed in Section 5.2.2 below. In addition, some quantitative uncertainty analyses of the interspecies differences in pharmacokinetics were conducted and are presented in Section 5.2.1.4.2.

The rodent bioassay data offer conclusive evidence of carcinogenicity in both rats and mice, and the available epidemiologic and mechanistic data support the relevance to humans of the TCE-induced carcinogenicity observed in rodents. The epidemiologic data provide sufficient evidence that TCE is “carcinogenic to humans” (see Section 4.11). There is even some evidence of site concordance with the rodent findings, although site concordance is not essential to human relevance and, in fact, is not observed across TCE-exposed rats and mice. The strongest evidence in humans is for TCE-induced kidney tumors, with fairly strong evidence for lymphomas and some lesser support for liver tumors; each of these tumor types has also been observed in TCE rodent bioassays. Furthermore, the mechanistic data are supportive of human relevance because, while the exact reactive species associated with TCE-induced tumors are not known, the metabolic pathways for TCE are qualitatively similar for rats, mice, and humans (see

1 Section 3.3). The impact of uncertainties with respect to quantitative differences in TCE  
2 metabolism is discussed in Section 5.2.1.4.2.

3 Typically, the cancer risk estimated is for the total cancer burden from all sites that demonstrate  
4 an increased tumor incidence for the most sensitive experimental species and sex. It is expected  
5 that this approach is protective of the human population, which is more diverse but is exposed to  
6 lower exposure levels.

7 For the inhalation unit risk estimates, the preferred estimate from the most sensitive  
8 species and sex was the estimate of  $8.3 \times 10^{-2}$  per ppm for the male rat, which was based on  
9 multiple tumors observed in this sex/species but was dominated by the kidney tumor risk  
10 estimated with the dose-metric for bioactivated DCVC. This estimate was the high end of the  
11 range of estimates (see Table 5-39) but was within an order of magnitude of other estimates,  
12 such as the preferred estimate for the female mouse and the male rat kidney estimate based on  
13 the GSH conjugation dose-metric, which provide additional support for an estimate of this  
14 magnitude. The preferred estimate for the male mouse was about an order of magnitude and a  
15 half lower. The female rat showed no apparent TCE-associated tumor response in the three  
16 available inhalation bioassays; however, this apparent absence of response is inconsistent with  
17 the observations of increased cancer risk in occupationally exposed humans and in female rats in  
18 oral bioassays. In Section 5.2.2.2, an inhalation unit risk estimate based on the human data is  
19 derived and can be compared to the rodent-based estimate.

20 For the oral slope factor estimate, the preferred estimate from the most sensitive species  
21 and sex was the estimate of  $2.5 \times 10^{-1}$  per mg/kg/day, again for the male rat, based on the kidney  
22 tumor risk estimated with the dose-metric for bioactivated DCVC. This estimate was at the high  
23 end of the range of estimates (see Table 5-40) but was within an order of magnitude of other  
24 estimates, such as the preferred male mouse estimate and the male rat kidney estimate based on  
25 the GSH conjugation dose-metric, which provide additional support for an estimate of this  
26 magnitude. The preferred estimates for the female mouse and the female rat were about another  
27 order of magnitude lower. Some of the oral slope factor estimates based on the alternative  
28 dose-metric of AUC for TCE in the blood were as much as three orders of magnitude lower, but  
29 these estimates were considered less credible than those based on the preferred dose-metrics. In  
30 Section 5.2.2.3, an oral slope factor estimate based on the human (inhalation) data is derived  
31 using the PBPK model for route-to-route extrapolation; this estimate can be compared to the  
32 rodent-based estimate.

33 Furthermore, the male rat kidney tumor estimates from the inhalation (Maltoni et al., 1986) and  
34 oral (NTP, 1988) studies were consistent on the basis of internal dose using the dose-metric for  
35 bioactivated DCVC. In particular, the linearly extrapolated slope (i.e., the BMR/BMDL) per unit

1 of internal dose derived from Maltoni et al. (1986) male rat kidney tumor data was  $2.4 \times 10^{-1}$  per  
2 weekly mg DCVC bioactivated per unit body weight<sup>3/4</sup>, while the analogous slope derived from  
3 NTP (1988) male rat kidney tumor data was  $9.3 \times 10^{-2}$  per weekly mg DCVC bioactivated per  
4 unit body weight<sup>3/4</sup> (MSW-modeled results), a difference of less than

5  
6  
7 threefold.<sup>47</sup> These results also suggest that differences between routes of administration are  
8 adequately accounted for by the PBPK model using this dose-metric.

9 Regarding low-dose extrapolation, a key consideration in determining what extrapolation  
10 approach to use is the MOA(s). However, MOA data are lacking or limited for each of the  
11 cancer responses associated with TCE exposure, with the exception of the kidney tumors (see  
12 Section 4.11). For the kidney tumors, the weight of the available evidence supports the  
13 conclusion that a mutagenic MOA is operative (see Section 4.4); this MOA supports linear  
14 low-dose extrapolation. For the other TCE-induced tumors, the MOA(s) is unknown. When the  
15 MOA(s) cannot be clearly defined, EPA generally uses a linear approach to estimate low-dose  
16 risk (U.S. EPA, 2005c), based on the following general principles:

- 17  
18
- 19 • A chemical's carcinogenic effects may act additively to ongoing biological processes,  
20 given that diverse human populations are already exposed to other agents and have  
21 substantial background incidences of various cancers.
  - 22 • A broadening of the dose-response curve (i.e., less rapid fall-off of response with  
23 decreasing dose) in diverse human populations and, accordingly, a greater potential for  
24 risks from low-dose exposures (!!! INVALID CITATION !!!) is expected for two reasons:  
25 First, even if there is a “threshold” concentration for effects at the cellular level, that  
26 threshold is expected to differ across individuals. Second, greater variability in response  
27 to exposures would be anticipated in heterogeneous populations than in inbred laboratory  
28 species under controlled conditions (due to, e.g., genetic variability, disease status, age,  
29 nutrition, and smoking status).

---

47For the Maltoni et al. (1986) male rat kidney tumors, the unit risk estimate of  $8.3 \times 10^{-2}$  per ppm using the ABioactDCVCBW34 dose metric, from Table 5-36, is divided by the average male and female internal doses at 0.001 ppm, (0.0034/0.001), from Table 5-35, to yield a unit risk in internal dose units of  $2.4 \times 10^{-2}$ . For the NTP (1988) male rat kidney tumors, the unit risk estimate of  $2.5 \times 10^{-1}$  per mg/kg/day using the ABioactDCVCBW34 dose metric, from Table 5-37, is divided by the average male and female internal doses at 0.001 mg/kg/day, (0.0027/0.001), from Table 5-35, to yield a unit risk in internal dose units of  $9.3 \times 10^{-2}$ . Note that the original BMDLs and unit risks from BMD modeling were in internal dose units that were then converted to applied dose units using the values in Table 5-35, so this calculation reverses that conversion.

- The general use of linear extrapolation provides reasonable upper-bound estimates that are believed to be health-protective (U.S. EPA, 2005c) and also provides consistency across assessments.

Additional uncertainties arise from the specific dosimetry assumptions, the model structures and parameter estimates in the PBPK models, the dose-response modeling of data in the observable range, and the application of the results to potentially sensitive human populations. As discussed in Section 5.2.1.2.1, one uncertainty in the tissue-specific dose-metrics used here is whether to scale the rate of metabolism by tissue mass or body weight to the  $\frac{3}{4}$  in the absence of specific data on clearance; however, in the cases where this is an issue (the lung, liver, and kidney), the impact of this choice is relatively modest (less than twofold to about fourfold). An additional dosimetry assumption inherent in this analysis is that equal concentrations of the active moiety over a lifetime yield equivalent lifetime risk of cancer across species, and the extent to which this is true for TCE is unknown. Furthermore, it should be noted that use of tissue-specific dosimetry inherently presumes site concordance of tumors across species.

With respect to uncertainties in the estimates of internal dose themselves, a quantitative analysis of the uncertainty and variability in the PBPK model-predicted dose-metric estimates and their impacts on cancer risk estimates is presented in Section 5.2.1.4.2. Additional uncertainties in the PBPK model were discussed in Section 3.5. Furthermore, this assessment examined a variety of dose-metrics for the different tumor types using PBPK models for rats, mice, and humans, so the impact of dose-metric selection can be assessed. As discussed in Section 5.2.1.2.1, there is strong support for the primary dose-metrics selected for kidney, liver, and, to a lesser extent, lung. For the other tumor sites, there is more uncertainty about dose-metric selection. The cancer slope factor and unit risk estimates obtained using the preferred dose-metrics were generally similar (within about threefold) to those derived using default dosimetry assumptions (e.g., equal risks result from equal cumulative equivalent exposures or doses), with the exception of the bioactivated DCVC dose-metric for rat kidney tumors and the metric for the amount of TCE oxidized in the respiratory tract for mouse lung tumors occurring from oral exposure (see Tables 5-39 and 5-40). The higher risk estimates for kidney tumors based on the bioactivated DCVC dose-metric are to be expected because pharmacokinetic data indicate, and the PBPK model predicts, substantially more GSH conjugation (as a fraction of intake), and hence subsequent bioactivation, in humans relative to rats. Nonetheless, there is substantial uncertainty in the quantitative extrapolation of GSH conjugation from rodents to humans due to limitations in the available data. The lower risk estimates for lung tumors from oral TCE exposure based on the metric for the amount of TCE

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 oxidized in the respiratory tract are because there is a greater first-pass effect in human liver  
2 relative to mouse liver following oral exposure and because the gavage dosing used in rodent  
3 studies leads to a large bolus dose that potentially overwhelms liver metabolism to a greater  
4 extent than a more graded oral exposure. Both of these effects result in relatively more TCE  
5 being available for metabolism in the lung for mice than for humans. In addition, mice have  
6 greater respiratory metabolism relative to humans. However, because oxidative metabolites  
7 produced in the liver may contribute to respiratory tract effects, using respiratory tract  
8 metabolism alone as a dose-metric may underestimate lung tumor risk. The slope factor or unit  
9 risk estimates obtained using the alternative dose-metrics were also generally similar to those  
10 derived using default dosimetry assumptions, with the exception of the metric for the amount of  
11 TCE conjugated with GSH for rat kidney tumors, again because humans have greater GSH  
12 conjugation, and the AUC of TCE in blood for all the tumor types resulting from oral exposure,  
13 again because of first-pass effects.


14 With respect to uncertainties in the dose-response modeling, the two-step approach of  
15 modeling only in the observable range, as put forth in EPA's cancer assessment guidelines (U.S.  
16 EPA, 2005c), is designed in part to minimize model dependence. The ratios of the BMDs to the  
17 BMDLs give some indication of the statistical uncertainties in the dose-response modeling.  
18 These ratios did not exceed a value of 2.5 for all the primary analyses used in this assessment.  
19 Thus, overall, modeling uncertainties in the observable range are considered to be minimal.  
20 Some additional uncertainty is conveyed by uncertainties in the survival adjustments made to  
21 some of the bioassay data; however, their impact is also believed to be minimal relative to the  
22 uncertainties already discussed (i.e., interspecies and low-dose extrapolations).

23 Regarding the cancer risks to potentially sensitive human populations or life stages,  
24 pharmacokinetic data on 42 individuals were used in the Bayesian population analysis of the  
25 PBPK model discussed in Section 3.5. The impacts of these data on the predicted population  
26 mean are incorporated in the quantitative uncertainty analyses presented in Section 5.2.1.4.2.  
27 These data do not, however, reflect the full range of metabolic variability in the human  
28 population (they are all from healthy, mostly male, human volunteers) and do not address  
29 specific potentially sensitive subgroups (see Section 4.10). Moreover, there is inadequate  
30 information about disease status, coexposures, and other factors that make humans vary in their  
31 responses to TCE. It will be a challenge for future research to quantify the differential risk  
32 indicated by different risk factors or exposure scenarios.

33



**5.2.1.1.8. Quantitative uncertainty analysis of physiologically based pharmacokinetic (PBPK) model-based dose-metrics**



1           The Bayesian analysis of the PBPK model for TCE generates distributions of uncertainty  
2 and variability in the internal dose-metrics that can be readily fed into dose-response analysis.  
3 As shown in Figure 5-6, the overall approach taken for the uncertainty analysis is similar to that  
4 used for the point estimates except that distributions are carried through the analysis rather than  
5 median or expected values. In particular, the PBPK model-based rodent internal doses are  
6 carried through to a distribution of BMDs (which also includes sampling variance from the  
7 number of responding and at risk animals in the bioassay). This distribution of BMDs generates  
8 a distribution of cancer slope factors based on internal dose, which then is combined with the

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 **Figure 5-6. Flow-chart for uncertainty analysis of dose-response analyses of**  
2 **rodent bioassays using PBPK model-based dose-metrics. Square nodes**  
3 **indicate point values, circular nodes indicate distributions, and the inverted**  
4 **triangles indicate a (deterministic) functional relationship.**  
5

1 (uncertainty) distribution of the human population mean conversion to applied dose or exposure.  
2 The resulting distribution for the human population mean risk per unit dose or exposure accounts  
3 for uncertainty in the PBPK model parameters (rodent and human) and the binomial sampling  
4 error in the bioassays. These distributions can then be compared with the point estimates, based  
5 on median rodent dose-metrics and mean human population dose-metrics, reported in  
6 Tables 5-36 and 5-37. Details of the implementation of this uncertainty analysis, which used the  
7 WinBugs software in conjunction with the R statistical package, are reported in Appendix G.

8 Overall, as shown in Tables 5-41 and 5-42, the 95% confidence upper bound of the  
9 distributions for the linearly extrapolated risk per unit dose or exposure ranged from one- to  
10 eightfold higher than the point slope factors and unit risks derived using the BMDLs reported in  
11 Tables 5-36 and 5-37. The largest differences, up to 4-fold, for rat kidney tumors and 8-fold for  
12 mouse lung tumors, primarily reflect the substantial uncertainty in the internal dose-metrics for  
13 rat kidney DCVC and GSH conjugation and for mouse lung oxidation (see Section 3.5).  
14 Additionally, despite the differences in the degree of uncertainty due to the PBPK model across  
15 endpoints and dose-metrics, the only case where the choice of the most sensitive bioassay for  
16 each sex/speciescombination would change based on the 95% confidence upper bounds reported  
17 in Tables 5-41 and 5-42 would be for female mouse inhalation bioassays. Even in this case, the  
18 difference between slope factor or unit risk estimate for the most sensitive and next most  
19 sensitive study/endpoint was only twofold.

### 5.2.2. Dose-Response Analyses: Human Epidemiologic Data

20 Of the epidemiological studies of TCE and cancer, only two had sufficient  
21 exposure-response information for potential dose-response analysis. The two studies, Charbotel  
22 et al. (2006) and Moore et al. (2010), were both case-control studies of TCE and kidney cancer,  
23 and both had quantitative cumulative exposure estimates for the individual subjects. In the study  
24 by Moore et al. (2010), however, the cumulative exposure estimates were assessed by experts  
25 based on categorical metrics for frequency and intensity of exposure and not continuous  
26 measures. Moore et al. (2010) also used a categorical confidence-of-exposure metric to classify  
27 different jobs because of the potential for exposure misclassification from this approach. While  
28 the detailed approach used by Moore et al. should be fairly reliable for general rankings, the  
29 resulting estimates are not expected to be as quantitatively accurate as those in the Charbotel  
30 et al. (2006) study, which relied on a task-exposure matrix based on decades of measurements  
31 from the Arve Valley workshops (Fevotte et al., 2006; see also Section 4.4 for more discussion  
32 of the exposure assessments). Thus, the Charbotel et al. (2006) study was selected as the sole  
33 basis for the derivation of an inhalation unit risk estimate for kidney cancer (see Section 5.2.2.1).

*This document is a draft for review purposes only and does not constitute Agency policy.*

**Table 5-41. Summary of PBPK model-based uncertainty analysis of unit risk estimates for each sex/species/bioassay/tumor type (inhalation)**

Study	Tumor Type	BMR	Dose-metric	Unit risk estimates (ppm) <sup>-1</sup>				
				From	Summary statistics of unit risk distribution			
				Table 5-36	Mean	5% lower bound	Median	95% upper bound
<b>Female mouse</b>								
Fukuda	Lung AD + CARC <sup>a</sup>	0.1	<b>AMetLngBW34</b>	<b>2.6 × 10<sup>-3</sup></b>	5.65 × 10 <sup>-3</sup>	2.34 × 10 <sup>-4</sup>	1.49 × 10 <sup>-3</sup>	2.18 × 10 <sup>-2</sup>
			TotOxMetabBW34	3.2 × 10 <sup>-3</sup>	1.88 × 10 <sup>-3</sup>	3.27 × 10 <sup>-4</sup>	1.52 × 10 <sup>-3</sup>	4.59 × 10 <sup>-3</sup>
			AUCCBld	1.8 × 10 <sup>-3</sup>	1.01 × 10 <sup>-3</sup>	1.54 × 10 <sup>-4</sup>	8.36 × 10 <sup>-4</sup>	2.44 × 10 <sup>-3</sup>
Henschler	Lymphoma <sup>b</sup>	0.1	<b>TotMetabBW34</b>	<b>1.0 × 10<sup>-2</sup></b>	4.38 × 10 <sup>-3</sup>	6.06 × 10 <sup>-4</sup>	3.49 × 10 <sup>-3</sup>	1.11 × 10 <sup>-2</sup>
Maltoni	Lung AD + CARC <sup>a</sup>	0.1	<b>AMetLngBW34</b>	<b>1.8 × 10<sup>-3</sup></b>	3.88 × 10 <sup>-3</sup>	1.48 × 10 <sup>-4</sup>	1.04 × 10 <sup>-3</sup>	1.52 × 10 <sup>-2</sup>
			TotOxMetabBW34	1.9 × 10 <sup>-3</sup>	1.10 × 10 <sup>-3</sup>	3.73 × 10 <sup>-4</sup>	9.52 × 10 <sup>-4</sup>	2.32 × 10 <sup>-3</sup>
			AUCCBld	1.0 × 10 <sup>-3</sup>	5.25 × 10 <sup>-4</sup>	1.63 × 10 <sup>-4</sup>	4.64 × 10 <sup>-4</sup>	1.10 × 10 <sup>-3</sup>
	Liver	0.05	<b>AMetLiv1BW34</b>	<b>1.2 × 10<sup>-3</sup></b>	6.27 × 10 <sup>-4</sup>	2.18 × 10 <sup>-4</sup>	5.39 × 10 <sup>-4</sup>	1.32 × 10 <sup>-3</sup>
			TotOxMetabBW34	1.1 × 10 <sup>-3</sup>	5.98 × 10 <sup>-4</sup>	1.81 × 10 <sup>-4</sup>	5.07 × 10 <sup>-4</sup>	1.31 × 10 <sup>-3</sup>
<b>Male mouse</b>								
Maltoni	Liver	0.1	<b>AMetLiv1BW34</b>	<b>2.6 × 10<sup>-3</sup></b>	1.35 × 10 <sup>-3</sup>	4.28 × 10 <sup>-4</sup>	1.16 × 10 <sup>-3</sup>	2.93 × 10 <sup>-3</sup>
			TotOxMetabBW34	2.0 × 10 <sup>-3</sup>	1.23 × 10 <sup>-3</sup>	4.24 × 10 <sup>-4</sup>	1.06 × 10 <sup>-3</sup>	2.60 × 10 <sup>-3</sup>
<b>Male rat</b>								
Maltoni	Leukemia <sup>b</sup>	0.05	<b>TotMetabBW34</b>	<b>1.8 × 10<sup>-3</sup></b>	9.38 × 10 <sup>-4</sup>	1.26 × 10 <sup>-4</sup>	7.86 × 10 <sup>-4</sup>	2.25 × 10 <sup>-3</sup>
	Kidney AD + CARC	0.01	<b>ABioactDCVCBW34</b>	<b>8.3 × 10<sup>-2</sup></b>	9.07 × 10 <sup>-2</sup>	3.66 × 10 <sup>-3</sup>	3.64 × 10 <sup>-2</sup>	3.21 × 10 <sup>-1</sup>
			AMetGSHBW34	5.1 × 10 <sup>-2</sup>	3.90 × 10 <sup>-2</sup>	2.71 × 10 <sup>-3</sup>	2.20 × 10 <sup>-2</sup>	1.30 × 10 <sup>-1</sup>
			TotMetabBW34	7.3 × 10 <sup>-4</sup>	3.94 × 10 <sup>-4</sup>	8.74 × 10 <sup>-5</sup>	3.42 × 10 <sup>-4</sup>	8.74 × 10 <sup>-4</sup>
	Leydig cell <sup>b</sup>	0.1	<b>TotMetabBW34</b>	<b>5.5 × 10<sup>-3</sup></b>	4.34 × 10 <sup>-3</sup>	1.99 × 10 <sup>-3</sup>	3.98 × 10 <sup>-3</sup>	7.87 × 10 <sup>-3</sup>

<sup>a</sup>WinBUGS dose-response analyses did not adequately converge for the AMetLngBW34 dose-metric using the 3<sup>rd</sup>-order multistage model (used for results in Table 5-36), but did converge when the 2<sup>nd</sup>-order model was used. Summary statistics reflect results of 2<sup>nd</sup>-order model calculations.

<sup>b</sup>Poor dose-response fits in point estimates for AUCCBld, so not included in uncertainty analysis.

AD = adenoma, CARC = carcinoma.

**Table 5-42. Summary of PBPK model-based uncertainty analysis of slope factor estimates for each sex/species/bioassay/tumor type (oral)**

Study	Tumor type	BMR	Dose-metric	Slope factor estimates (mg/kg/day) <sup>-1</sup>				
				From	Summary statistics of slope factor distribution			
				Table 5-37 or 5-38	Mean	5% lower bound	Median	95% upper bound
<b>Female mouse</b>								
NCI	Liver CARC	0.1	<b>AMetLiv1BW34</b>	<b>7.1 × 10<sup>-3</sup></b>	3.26 × 10 <sup>-3</sup>	9.35 × 10 <sup>-4</sup>	2.44 × 10 <sup>-3</sup>	8.35 × 10 <sup>-3</sup>
			TotOxMetabBW34	5.7 × 10 <sup>-3</sup>	2.63 × 10 <sup>-3</sup>	8.76 × 10 <sup>-4</sup>	2.01 × 10 <sup>-3</sup>	6.60 × 10 <sup>-3</sup>
	Lung AD + CARC <sup>a</sup>	0.1	<b>AMetLngBW34</b>	<b>1.3 × 10<sup>-4</sup></b>	1.28 × 10 <sup>-4</sup>	6.73 × 10 <sup>-6</sup>	4.12 × 10 <sup>-5</sup>	4.62 × 10 <sup>-4</sup>
			TotOxMetabBW34	4.0 × 10 <sup>-3</sup>	1.84 × 10 <sup>-3</sup>	5.29 × 10 <sup>-4</sup>	1.39 × 10 <sup>-3</sup>	4.73 × 10 <sup>-3</sup>
			AUCCBld	1.5 × 10 <sup>-4</sup>	7.16 × 10 <sup>-5</sup>	4.40 × 10 <sup>-6</sup>	3.39 × 10 <sup>-5</sup>	2.18 × 10 <sup>-4</sup>
	Leukemias + sarcomas	0.1	<b>TotMetabBW34</b>	<b>4.9 × 10<sup>-3</sup></b>	1.60 × 10 <sup>-3</sup>	1.42 × 10 <sup>-4</sup>	1.13 × 10 <sup>-3</sup>	4.65 × 10 <sup>-3</sup>
AUCCBld			1.4 × 10 <sup>-4</sup>	6.36 × 10 <sup>-5</sup>	3.10 × 10 <sup>-6</sup>	2.90 × 10 <sup>-5</sup>	1.94 × 10 <sup>-4</sup>	
<b>Male mouse</b>								
NCI	Liver CARC	0.1	<b>AMetLiv1BW34</b>	<b>2.9 × 10<sup>-2</sup></b>	1.65 × 10 <sup>-2</sup>	4.70 × 10 <sup>-3</sup>	1.25 × 10 <sup>-2</sup>	4.25 × 10 <sup>-2</sup>
			TotOxMetabBW34	2.3 × 10 <sup>-2</sup>	1.32 × 10 <sup>-2</sup>	4.41 × 10 <sup>-3</sup>	1.01 × 10 <sup>-2</sup>	3.29 × 10 <sup>-2</sup>
<b>Female rat</b>								
NTP (1988)	Leukemia	0.05	<b>TotMetabBW34</b>	<b>2.3 × 10<sup>-3</sup></b>	1.89 × 10 <sup>-3</sup>	5.09 × 10 <sup>-4</sup>	1.43 × 10 <sup>-3</sup>	4.69 × 10 <sup>-3</sup>
			AUCCBld	1.6 × 10 <sup>-5</sup>	1.56 × 10 <sup>-5</sup>	3.39 × 10 <sup>-6</sup>	1.07 × 10 <sup>-5</sup>	3.98 × 10 <sup>-5</sup>
<b>Male rat</b>								
NTP (1990)	Kidney AD + CARC <sup>b</sup>	0.1	<b>ABioactDCVCBW34</b>	<b>1.2 × 10<sup>-1</sup></b>	1.40 × 10 <sup>-1</sup>	5.69 × 10 <sup>-3</sup>	5.24 × 10 <sup>-2</sup>	5.18 × 10 <sup>-1</sup>
			AMetGSHBW34	7.6 × 10 <sup>-2</sup>	6.18 × 10 <sup>-2</sup>	4.00 × 10 <sup>-3</sup>	3.27 × 10 <sup>-2</sup>	2.11 × 10 <sup>-1</sup>
			TotMetabBW34	3.1 × 10 <sup>-3</sup>	2.49 × 10 <sup>-3</sup>	7.14 × 10 <sup>-4</sup>	1.96 × 10 <sup>-3</sup>	5.96 × 10 <sup>-3</sup>

**Table 5-42. Summary of PBPK model-based uncertainty analysis of slope factor estimates for each sex/species/bioassay/tumor type (oral) (continued)**

Study	Tumor type	BMR	Dose-metric	Slope factor estimates (mg/kg/day) <sup>-1</sup>				
				From	Summary statistics of slope factor distribution			
				Table 5-37 or 5-38	Mean	5% lower bound	Median	95% upper bound
NTP (1988)								
Marshall	Testicular <sup>b</sup>	0.1	<b>TotMetabBW34</b>	<b>7.1 × 10<sup>-2</sup></b>	6.18 × 10 <sup>-2</sup>	1.92 × 10 <sup>-2</sup>	4.89 × 10 <sup>-2</sup>	1.45 × 10 <sup>-1</sup>
			AUCCBld	6.0 × 10 <sup>-4</sup>	5.45 × 10 <sup>-4</sup>	1.18 × 10 <sup>-4</sup>	3.70 × 10 <sup>-4</sup>	1.44 × 10 <sup>-3</sup>
August	Subcut sarcoma	0.05	<b>TotMetabBW34</b>	<b>2.3 × 10<sup>-3</sup></b>	1.65 × 10 <sup>-3</sup>	4.58 × 10 <sup>-4</sup>	1.27 × 10 <sup>-3</sup>	4.04 × 10 <sup>-3</sup>
			AUCCBld	2.0 × 10 <sup>-5</sup>	1.35 × 10 <sup>-5</sup>	1.53 × 10 <sup>-6</sup>	8.34 × 10 <sup>-6</sup>	3.73 × 10 <sup>-5</sup>
Osborne-Mendel	Kidney AD + CARC <sup>b</sup>	0.1	<b>ABioactDCVCBW34</b>	<b>1.6 × 10<sup>-1</sup></b>	1.61 × 10 <sup>-1</sup>	5.45 × 10 <sup>-3</sup>	6.35 × 10 <sup>-2</sup>	6.02 × 10 <sup>-1</sup>
			AMetGSHBW34	9.7 × 10 <sup>-2</sup>	7.47 × 10 <sup>-2</sup>	3.90 × 10 <sup>-3</sup>	3.85 × 10 <sup>-2</sup>	2.54 × 10 <sup>-1</sup>
			TotMetabBW34	4.3 × 10 <sup>-3</sup>	2.73 × 10 <sup>-3</sup>	5.40 × 10 <sup>-4</sup>	2.10 × 10 <sup>-3</sup>	6.89 × 10 <sup>-3</sup>

<sup>a</sup>WinBUGS dose-response analyses did not adequately converge for AMetLngBW34 dose-metric using the 3<sup>rd</sup>-order multistage model (used for results in Table 5-37), but did converge when the 2<sup>nd</sup>-order model was used. Summary statistics reflect results of 2<sup>nd</sup>-order model calculations.

<sup>b</sup>Using poly-3 adjusted incidences from Table 5-38 (software for WinBUGS-based analyses using the MSW model was not developed).

AD = adenoma, CARC = carcinoma.

1 Other epidemiological studies were used in Section 5.2.2.2 below to provide information for a  
2 comparison of relative risk (RR) estimates across cancer types. These epidemiologic data were  
3 used to derive an adjusted inhalation unit risk estimate for the combined risk of developing  
4 kidney cancer, non-Hodgkin lymphoma (NHL), or liver cancer. The human PBPK model was  
5 then used to perform route-to-route extrapolation to derive an oral slope factor estimate for the  
6 combined risk of kidney cancer, NHL, or liver cancer (see Section 5.2.2.3).  
7

#### **5.2.2.1.1. Inhalation Unit Risk Estimate for Renal Cell Carcinoma Derived from Charbotel et al. (2006) Data**

8 The Charbotel et al. (2006) case-control study of 86 incident renal cell carcinoma (RCC)  
9 cases and 316 age-and sex-matched controls, with individual cumulative exposure estimates for  
10 TCE for each subject, provides a sufficient human data set for deriving quantitative cancer risk  
11 estimates for RCC in humans. The study is a high-quality study that used a detailed exposure  
12 assessment (Fevotte et al., 2006) and took numerous potential confounding factors, including  
13 exposure to other chemicals, into account (see Section 4.4). A significant dose-response  
14 relationship was reported for cumulative TCE exposure and RCC (Charbotel et al., 2006).  
15

16 The derivation of an inhalation unit risk estimate, defined as the plausible upper bound  
17 lifetime risk of cancer from chronic inhalation of TCE per unit of air concentration, for RCC  
18 incidence in the U.S. population, based on results of the Charbotel et al. (2006) case-control  
19 study, is presented in the following subsections.

#### **5.2.2.1.2. Renal cell carcinoma (RCC) results from the Charbotel et al. (2006) study**

20 Charbotel et al. (2006) analyzed their data using conditional logistic regression, matching  
21 on sex and age, and reported results (odds ratios [ORs]) for cumulative TCE exposure categories,  
22 adjusted for tobacco smoking and body mass index (Charbotel et al., 2006, Table 6). The  
23 exposure categories were constructed as tertiles based on the cumulative exposure levels in the  
24 exposed control subjects. The results are summarized in Table 5-43, with mean exposure levels  
25 kindly provided by Dr. Charbotel (personal communication from Barbara Charbotel, University  
26 of Lyon, to Cheryl Scott, U.S. EPA, 11 April 2008).

27 For additional details and discussion of the Charbotel et al. (2006) study, see Section 4.4  
28 and Appendix B.  
29



### **5.2.2.1.3. Prediction of lifetime extra risk of renal cell carcinoma (RCC) incidence from trichloroethylene (TCE) exposure**

1           The categorical results summarized in Table 5-43 were used for predicting the extra risk  
2 of RCC incidence from continuous environmental exposure to TCE. Extra risk is defined as  
3

1 **Table 5-43. Results from Charbotel et al. (2006) on relationship between**  
 2 **TCE exposure and RCC**  
 3

Cumulative exposure category	Mean Cumulative exposure (ppm × yrs)	Adjusted OR (95% CI)
Nonexposed		1
Low	62.4	1.62 (0.75, 3.47)
Medium	253.2	1.15 (0.47, 2.77)
High	925.0	2.16 (1.02, 4.60)

4  
 5 CI = confidence interval.  
 6  
 7

$$8 \quad \text{Extra risk} = (R_x - R_o)/(1 - R_o), \quad (\text{Eq. 5-3})$$

9  
 10 where  $R_x$  is the lifetime risk in the exposed population and  $R_o$  is the lifetime risk in an  
 11 unexposed population (i.e., the background risk). Because kidney cancer is a rare event, the ORs  
 12 in Table 5-43 can be used as estimates of the relative risk ratio,  $RR = R_x/R_o$  (Rothman and  
 13 Greenland, 1998). A weighted linear regression model was used to model the dose-response data  
 14 in Table 5-43 to obtain a slope estimate (regression coefficient) for RR of RCC versus  
 15 cumulative exposure, under the commonly employed assumption that exposure was measured  
 16 without error. Use of a linear model in the observable range of the data is often a good general  
 17 approach for epidemiological data because such data are frequently too limited (i.e., imprecise),  
 18 as is the case here, to clearly identify an alternate model (U.S. EPA, 2005c). This linear  
 19 dose-response function was then used to calculate lifetime extra risks in an actuarial program  
 20 (life-table analysis) that accounts for age-specific rates of death and background disease, under  
 21 the common assumption that the RR is independent of age.<sup>48</sup> In addition, it is generally assumed  
 22 that RR estimates transfer across populations, independent of background disease rates—in this  
 23 case, the RR estimates based on the Charbotel et al. (2006) study, which was conducted in  
 24 France, are assumed to apply to the U.S. population.<sup>49</sup>

---

48This program is an adaptation of the approach previously used by the Committee on the Biological Effects of Ionizing Radiation (NRC, 1988). The same methodology was also used in U.S. EPA's 1,3-butadiene health risk assessment (U.S. EPA, 2002a). A spreadsheet illustrating the extra risk calculation for the derivation of the LEC<sub>01</sub> for RCC incidence is presented in Appendix H.

49 In any event, background kidney cancer rates between the U.S. and France are similar, with estimated age-adjusted incidence rates of 14.1 per 100,000 in the U.S. (Surveillance, Epidemiology, and End Results: <http://seer.cancer.gov/statfacts/html/kidrp.html>) and 10.4 per 100,000 in France (European Cancer Observatory: <http://eu-cancer.iarc.fr/cancer-19-kidney.html,en>).

1 For the weighted linear regression, the weights used for the RR estimates were the  
2 inverses of the variances, which were calculated from the confidence intervals. Using this  
3 approach,<sup>50</sup> a linear regression coefficient of 0.001205 per ppm × year  
4 (standard error = 0.0008195 per ppm × year) was obtained from the categorical results.

5 For the life-table analysis, U.S. age-specific all-cause mortality rates for 2004 for both  
6 sexes and all race groups combined (CDC, 2007) were used to specify the all-cause background  
7 mortality rates in the actuarial program. Because the goal is to estimate the unit risk for extra  
8 risk of cancer incidence, not mortality, and because the Charbotel et al. (2006) data are incidence  
9 data, RCC incidence rates were used for the cause-specific background “mortality” rates in the  
10 life-table analysis.<sup>51</sup> Surveillance, Epidemiology, and End Results (SEER) 2001–2005  
11 cause-specific background incidence rates for RCC were obtained from the SEER public-use  
12 database.<sup>52</sup> SEER collects good-quality cancer incidence data from a variety of geographical  
13 areas in the United States. The incidence data used here are from SEER 17, a registry of  
14 17 states, cities, or regions covering about 26% of the United States population  
15 (<http://seer.cancer.gov>). The risks were computed up to age 85 years for continuous exposures to  
16 TCE.<sup>53</sup> Conversions between occupational TCE exposures and continuous environmental  
17 exposures were made to account for differences in the number of days exposed per year (240 vs.  
18 365 days) and in the amount of air inhaled per day (10 vs. 20 m<sup>3</sup>; U.S. EPA, 1994b). The  
19 standard error for the regression coefficient from the weighted linear regression calculation  
20 described above was used to compute the 95% upper confidence limit (UCL) for the slope  
21 estimate, and this value was used to derive 95% UCLs for risk estimates (or 95% lower  
22 confidence limits [LCLs] for corresponding exposure estimates), based on a normal  
23 approximation.

24 Point estimates and one-sided 95% UCLs for the extra risk of RCC incidence associated  
25 with varying levels of environmental exposure to TCE based on linear regression of the  
26 Charbotel et al. (2006) categorical results were determined by the actuarial program; the results

---

<sup>50</sup>Equations for this weighted linear regression approach are presented in Rothman (1986) and summarized in Appendix H.

<sup>51</sup>No adjustment was made for using RCC incidence rates rather than mortality rates to represent cause-specific mortality in the actuarial program because the RCC incidence rates are negligible in comparison to the all-cause mortality rates. Otherwise, all-cause mortality rates for each age interval would have been adjusted to reflect people dying of a cause other than RCC or being diagnosed with RCC.

<sup>52</sup>In accordance with the “SEER Program Coding and Staging Manual 2007”

([http://seer.cancer.gov/manuals/2007/SPCSM\\_2007\\_AppendixC\\_p6.pdf](http://seer.cancer.gov/manuals/2007/SPCSM_2007_AppendixC_p6.pdf)), pages C-831 to C-833, RCC was specified as ICD-0-3 histological types coded 8312, 8260, 8310, 8316-8320, 8510, 8959, and 8255 (mixed types).

<sup>53</sup>Rates above age 85 years are not included because cause-specific disease rates are less stable for those ages. Note that 85 years is not employed here as an average lifespan but, rather, as a cut-off point for the life-table analysis, which uses actual age-specific mortality rates.

1 are presented in Section 5.2.13. The models based on cumulative exposure yield extra risk  
 2 estimates that are fairly linear for exposures up to 1 ppm or so.

3 Consistent with EPA’s *Guidelines for Carcinogen Risk Assessment* (U.S. EPA, 2005c),  
 4 the same data and methodology were also used to estimate the exposure level ( $EC_x$ : “effective  
 5 concentration corresponding to an extra risk of  $x\%$ ”) and the associated 95% lower confidence  
 6 limit of the effective concentration corresponding to an extra risk of 1% ( $LEC_x$  [lowest effective  
 7 concentration],  $x = 0.01$ ). A 1% extra risk level is commonly used for the determination of the  
 8 POD for epidemiological data. Use of a 1% extra risk level for these data is supported by the  
 9 fact that, based on the actuarial program, the risk ratio (i.e.,  $R_x/R_o$ ) for an extra risk of 1% for  
 10 RCC incidence is 1.9, which is in the range of the ORs reported by Charbotel et al. (see  
 11 Table 5-43). Thus, 1% extra risk was selected for determination of the POD, and, consistent  
 12 with the *Guidelines for Carcinogen Risk Assessment*, the LEC value corresponding to that risk  
 13 level was used as the actual POD. For the linear model that was selected, the unit risk is  
 14 independent of the benchmark risk level used to determine the POD (at low exposures/risk  
 15 levels; see Table 5-44); however, selection of a benchmark risk level is generally useful for  
 16 comparisons across models.

17 As discussed in Section 4.4, there is sufficient evidence to conclude that a mutagenic  
 18 MOA is operative for TCE-induced kidney tumors, which supports the use of linear low-dose  
 19 extrapolation from the POD. The  $EC_{01}$ ,  $LEC_{01}$ , and inhalation unit risk estimates for RCC  
 20 incidence using the linear cumulative exposure model are presented in Table 5-45. Converting  
 21 the units,  $5.49 \times 10^{-3}$  per ppm corresponds to a unit risk of  $1.02 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$  for RCC  
 22 incidence.

23  
 24 **Table 5-44. Extra risk estimates for RCC incidence from various levels of**  
 25 **lifetime exposure to TCE, using linear cumulative exposure model**  
 26

Exposure concentration (ppm)	MLE of extra risk	95% UCL on extra risk
0.001	$2.603 \times 10^{-6}$	$5.514 \times 10^{-6}$
0.01	$2.603 \times 10^{-5}$	$5.514 \times 10^{-5}$
0.1	$2.602 \times 10^{-4}$	$5.512 \times 10^{-4}$
1.0	$2.598 \times 10^{-3}$	$5.496 \times 10^{-3}$
10.0	$2.562 \times 10^{-2}$	$5.333 \times 10^{-2}$

1 **Table 5-45. EC<sub>01</sub>, LEC<sub>01</sub>, and unit risk estimates for RCC incidence, using**  
 2 **linear cumulative exposure model**  
 3

EC <sub>01</sub> (ppm)	LEC <sub>01</sub> (ppm)	unit risk (per ppm)*
3.87	1.82	$5.49 \times 10^{-3}$

4 \*Unit risk = 0.01/LEC<sub>01</sub>.  
 5  
 6  
 7

#### 5.2.2.1.4. Uncertainties in the renal cell carcinoma (RCC) unit risk estimate

8 The two major sources of uncertainty in quantitative cancer risk estimates are generally  
 9 interspecies extrapolation and high-dose to low-dose extrapolation. The unit risk estimate for  
 10 RCC incidence derived from the Charbotel et al. (2006) results is not subject to interspecies  
 11 uncertainty because it is based on human data. A major uncertainty remains in the extrapolation  
 12 from occupational exposures to lower environmental exposures. There was some evidence of a  
 13 contribution to increased RCC risk from peak exposures; however, there remained an apparent  
 14 dose-response relationship for RCC risk with increasing cumulative exposure without peaks, and  
 15 the OR for exposure with peaks compared to exposure without peaks was not significantly  
 16 elevated (Charbotel et al., 2006). Although the actual exposure-response relationship at low  
 17 exposure levels is unknown, the conclusion that a mutagenic MOA is operative for TCE-induced  
 18 kidney tumors supports the linear low-dose extrapolation that was used (U.S. EPA, 2005c).

19 Another notable source of uncertainty in the cancer unit risk estimate is the dose-response  
 20 model used to model the study data to estimate the POD. A weighted linear regression across the  
 21 categorical ORs was used to obtain a slope estimate; use of a linear model in the observable  
 22 range of the data is often a good general approach for human data because epidemiological data  
 23 are frequently too limited (i.e., imprecise) to clearly identify an alternate model (U.S. EPA,  
 24 2005c). The Charbotel et al. (2006) study is a relatively small case-control study, with only 86  
 25 RCC cases, 37 of which had TCE exposure; thus, the dose-response data upon which to specify a  
 26 model are indeed limited.

27 In accordance with EPA's *Guidelines for Carcinogen Risk Assessment*, the lower bound  
 28 on the EC<sub>01</sub> is used as the POD; this acknowledges some of the uncertainty in estimating the  
 29 POD from the available dose-response data. In this case, the statistical uncertainty associated  
 30 with the EC<sub>01</sub> is relatively small, as the ratio between the EC<sub>01</sub> and the LEC<sub>01</sub> is about twofold.  
 31 The inhalation unit risk estimate of  $5.49 \times 10^{-3}$  per ppm presented above, which is calculated  
 32 based on a linear extrapolation from the POD (LEC<sub>01</sub>), is expected to provide an upper bound on

1 the risk of cancer incidence. However, for certain applications, such as benefit-cost analyses,  
2 estimates of “central tendency” for the risk below the POD are desired. Because a linear  
3 dose-response model was used in the observable range of the human data and the POD was  
4 within the low-dose linear range for extra risk as a function of exposure, linear extrapolation  
5 below the  $LEC_{01}$  has virtually the same slope as the 95% UCL on the actual (linear)  
6 dose-response model in the low-dose range (i.e., below the POD). This is illustrated in  
7 Table 5-44, where the 95% UCL on extra risk for RCC incidence predicted by the dose-response  
8 model is about  $5.51 \times 10^{-3}$  per ppm for exposures at or below about 0.1 ppm, which is virtually  
9 equivalent to the unit risk estimate of  $5.49 \times 10^{-3}$  per ppm derived from the  $LEC_{01}$  (see  
10 Table 5-45). The same holds for the central tendency (weighted least squares) estimates of the  
11 extra risk from the (linear) dose-response model (i.e., the dose-response model prediction of  
12  $2.60 \times 10^{-3}$  per ppm from Table 5-44 is virtually identical to the value of  $2.58 \times 10^{-3}$  per ppm  
13 obtained from linear extrapolation below the  $EC_{01}$ , i.e., by dividing 0.01 extra risk by the  $EC_{01}$  of  
14 3.87 from Table 5-45). In other words, because the dose-response model that was used to model  
15 the data in the observable range is already low-dose linear near the POD, if one assumes that the  
16 same linear model is valid for the low-dose range, one can use the central tendency (weighted  
17 least squares) estimate from the model to derive a statistical “best estimate” of the slope rather  
18 than relying on an extrapolated risk estimate ( $0.01/EC_{01}$ ). [The extrapolated risk estimates are  
19 not generally central tendency estimates in any statistical sense because once risk is extrapolated  
20 below the  $EC_{01}$  using the formulation  $0.01/EC_{01}$ , it is no longer a function of the original model  
21 which generated the  $EC_{01}$ s and the  $LEC_{01}$ s.]

22 An important source of uncertainty in the underlying Charbotel et al. (2006) study is the  
23 retrospective estimation of TCE exposures in the study subjects. This case-control study was  
24 conducted in the Arve Valley in France, a region with a high concentration of workshops  
25 devoted to screw cutting, which involves the use of TCE and other degreasing agents. Since the  
26 1960s, occupational physicians of the region have collected a large quantity of well-documented  
27 measurements, including TCE air concentrations and urinary metabolite levels (Fevotte et al.,  
28 2006). The study investigators conducted a comprehensive exposure assessment to estimate  
29 cumulative TCE exposures for the individual study subjects, using a detailed occupational  
30 questionnaire with a customized task-exposure matrix for the screw-cutting workers and a more  
31 general occupational questionnaire for workers exposed to TCE in other industries (Fevotte et  
32 al., 2006). The exposure assessment even attempted to take dermal exposure from hand-dipping  
33 practices into account by equating it with an equivalent airborne concentration based on  
34 biological monitoring data. Despite the appreciable effort of the investigators, considerable  
35 uncertainty associated with any retrospective exposure assessment is inevitable, and some  
36 exposure misclassification is unavoidable. Such exposure misclassification was most likely for

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 the 19 deceased cases and their matched controls, for which proxy respondents were used, and  
2 for exposures outside the screw-cutting industry (295 of 1,486 identified job periods involved  
3 TCE exposure; 120 of these were not in the screw-cutting industry).

4 Although the exposure estimates from Moore et al. (2010) were not considered to be as  
5 quantitatively accurate as those of Charbotel et al. (2006), as discussed at the beginning of  
6 Section 5.2.2, it is worth noting, in the context of uncertainty in the exposure assessment, that the  
7 exposure estimates in Moore et al. (2010) are substantially lower than those of Charbotel et al.  
8 (2006) for comparable OR estimates. For example, for all subjects and high-confidence  
9 assessments only, respectively, Moore et al. (2010) report OR estimates of 1.19 and 1.77 for  
10 cumulative exposures  $< 1.58 \text{ ppm} \times \text{years}$  and 2.02 and 2.23 for cumulative  
11 exposures  $\geq 1.58 \text{ ppm} \times \text{years}$ . Charbotel et al. (2006), on the other hand, report OR estimates  
12 for all subjects of 1.62, 1.15, and 2.16 for mean cumulative exposures of 62.4, 253.2, and  
13 925.0  $\text{ppm} \times \text{years}$ , respectively. If the exposure estimates for Charbotel et al. (2006) are  
14 overestimated, as suggested by the exposure estimates from Moore et al. (2010), the slope of the  
15 linear regression model, and hence the unit risk estimate, would be correspondingly  
16 underestimated.

17 Another noteworthy source of uncertainty in the Charbotel et al. (2006) study is the  
18 possible influence of potential confounding or modifying factors. This study population, with a  
19 high prevalence of metal-working, also had relatively high prevalences of exposure to petroleum  
20 oils, cadmium, petroleum solvents, welding fumes, and asbestos (Fevotte et al., 2006). Other  
21 exposures assessed included other solvents (including other chlorinated solvents), lead, and  
22 ionizing radiation. None of these exposures was found to be significantly associated with RCC  
23 at a  $p = 0.05$  significance level. Cutting fluids and other petroleum oils were associated with  
24 RCC at a  $p = 0.1$  significance level; however, further modeling suggested no association with  
25 RCC when other significant factors were taken into account (Charbotel et al., 2006). Moreover,  
26 a review of other studies suggested that potential confounding from cutting fluids and other  
27 petroleum oils is of minimal concern (see Section 4.4.2.3). Nonetheless, a sensitivity analysis  
28 was conducted using the OR estimates further adjusted for cutting fluids and other petroleum oils  
29 from the unpublished report by Charbotel et al. (2005), and an essentially identical unit risk  
30 estimate of  $5.46 \times 10^{-3}$  per ppm was obtained.<sup>54</sup> In addition, the medical questionnaire included

---

54 The OR estimates further adjusted for cutting fluids and other petroleum oils were 1.52 (95% CI: 0.66, 3.49), 1.07 (0.39, 2.88), and 1.96 (0.71, 5.37) for the low, medium, and high cumulative exposure groups, respectively (Charbotel et al., 2005). For the linear regression model, these OR estimates yielded a shallower slope estimate of 0.0009475 per  $\text{ppm} \times \text{year}$  but a larger SE of 0.0009709 per  $\text{ppm} \times \text{year}$ . In the lifetable analysis, these latter estimates in turn yielded a slightly higher EC01 estimate (4.92 ppm versus 3.87 ppm), because of the shallower slope estimate, but an essentially identical LEC01, because of the larger SE.

1 familial kidney disease and medical history, such as kidney stones, infection, chronic dialysis,  
2 hypertension, and use of antihypertensive drugs, diuretics, and analgesics. Body mass index  
3 (BMI) was also calculated, and lifestyle information such as smoking habits and coffee  
4 consumption was collected. Univariate analyses found high levels of smoking and BMI to be  
5 associated with increased odds of RCC, and these two variables were included in the conditional  
6 logistic regressions. Thus, although impacts of other factors are possible, this study took great  
7 pains to attempt to account for potential confounding or modifying factors.

8 Some other sources of uncertainty associated with the epidemiological data are the  
9 dose-metric and lag period. As discussed above, there was some evidence of a contribution to  
10 increased RCC risk from peak TCE exposures; however, there appeared to be an independent  
11 effect of cumulative exposure without peaks. Cumulative exposure is considered a good  
12 measure of total exposure because it integrates exposure (levels) over time. If there is a  
13 contributing effect of peak exposures, not already taken into account in the cumulative exposure  
14 metric, the linear slope may be overestimated to some extent. Sometimes cancer data are  
15 modeled with the inclusion of a lag period to discount more recent exposures not likely to have  
16 contributed to the onset of cancer. In an unpublished report (Charbotel et al., 2005), Charbotel  
17 et al. also present the results of a conditional logistic regression with a 10-year lag period, and  
18 these results are very similar to the unlagged results reported in their published paper, suggesting  
19 that the lag period might not be an important factor in this study.

20 Some additional sources of uncertainty are not so much inherent in the exposure-response  
21 modeling or in the epidemiologic data themselves but, rather, arise in the process of obtaining  
22 more general Agency risk estimates from the epidemiologic results. EPA cancer risk estimates  
23 are typically derived to represent an upper bound on increased risk of cancer incidence for all  
24 sites affected by an agent for the general population. From experimental animal studies, this is  
25 accomplished by using tumor incidence data and summing across all the tumor sites that  
26 demonstrate significantly increased incidences, customarily for the most sensitive sex and  
27 species, to attempt to be protective of the general human population. However, in estimating  
28 comparable risks from the Charbotel et al. (2006) epidemiologic data, certain limitations are  
29 encountered. For one thing, these epidemiology data represent a geographically limited (Arve  
30 Valley, France) and likely not very diverse population of working adults. Thus, there is  
31 uncertainty about the applicability of the results to a more diverse general population.  
32 Additionally, the Charbotel et al. (2006) study was a study of RCC only, and so the risk estimate  
33 derived from it does not represent all the tumor sites that may be affected by TCE. The issue of  
34 cancer risk at other sites is addressed in the next section (see Section 5.2.2.2).



#### 5.2.2.1.5. Conclusions regarding the renal cell carcinoma (RCC) unit risk estimate

1 An EC<sub>01</sub> of 3.9 ppm was calculated using a life-table analysis and linear modeling of the  
2 categorical conditional logistic regression results for RCC incidence reported in a high-quality  
3 case-control study. Linear low-dose extrapolation from the LEC<sub>01</sub> yielded a lifetime extra RCC  
4 incidence unit risk estimate of  $5.5 \times 10^{-3}$  per ppm ( $1.0 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ) of continuous TCE  
5 exposure. The assumption of low-dose linearity is supported by the conclusion that a mutagenic  
6 MOA is operative for TCE-induced kidney tumors. The inhalation unit risk estimate is expected  
7 to provide an upper bound on the risk of RCC incidence; however, this is just the risk estimate  
8 for RCC. A risk estimate for total cancer risk to humans would need to include the risk for other  
9 potential TCE-associated cancers.

#### 5.2.2.1.6. Adjustment of the Inhalation Unit Risk Estimate for Multiple Sites

11 Human data on TCE exposure and cancer risk sufficient for dose-response modeling are  
12 only available for RCC, yet human and rodent data suggest that TCE exposure increases the risk  
13 of cancer at other sites as well. In particular, there is evidence from human (and rodent) studies  
14 for increased risks of NHL and liver cancer (see Section 4.11). Therefore, the inhalation unit  
15 risk estimate derived from human data for RCC incidence was adjusted to account for potential  
16 increased risk of those tumor types. To make this adjustment, a factor accounting for the relative  
17 contributions to the extra risk for cancer incidence from TCE exposure for these three tumor  
18 types combined versus the extra risk for RCC alone was estimated, and this factor was applied to  
19 the unit risk estimate for RCC to obtain a unit risk estimate for the three tumor types combined  
20 (i.e., lifetime extra risk for developing *any* of the three types of tumor). This estimate is  
21 considered a better estimate of total cancer risk from TCE exposure than the estimate for RCC  
22 alone.

23 Although only the Charbotel et al. (2006) study was found adequate for direct estimation  
24 of inhalation unit risks, the available epidemiologic data provide sufficient information for  
25 estimating the *relative* potency of TCE across tumor sites. In particular, the relative  
26 contributions to extra risk (for cancer incidence) were calculated from two different data sets to  
27 derive the adjustment factor for adjusting the unit risk estimate for RCC to a unit risk estimate  
28 for the three types of cancers (RCC, NHL, and liver) combined. The first calculation is based on  
29 the results of the meta-analyses of human epidemiologic data for the three tumor types (see  
30 Appendix C); the second calculation is based on the results of the Raaschou-Nielsen et al. (2003)  
31 study, the largest single human epidemiologic study by far with RR estimates for all three tumor  
32 types. The approach for each calculation was to use the RR estimates and estimates of the

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 lifetime background risk in an unexposed population,  $R_o$ , to calculate the lifetime risk in the  
2 exposed population,  $R_x$ , where  $R_x = RR \times R_o$ , for each tumor type. Then, the extra risk from  
3 TCE exposure for each tumor type could be calculated using the equation in Section 5.2.2.1.2.  
4 Finally, the extra risks were summed across the three tumor types and the ratio of the sum of the  
5 extra risks to the extra risk for RCC was derived. For the first calculation, the summary relative  
6 risk estimates (RRm's) from the meta-analyses for NHL, kidney cancer, and liver (and biliary)  
7 cancer were used as the RR estimates. For the second calculation, the SIR estimates from the  
8 Raaschou-Nielsen et al. (2003) study were used. For both calculations,  $R_o$  for RCC was taken  
9 from the life-table analysis described in Section 5.2.2.1.2 and presented in Appendix H, which  
10 estimated a lifetime risk for RCC incidence up to age 85 years. For  $R_o$  values for the other  
11 two sites, SEER statistics for the lifetime risk of developing cancer were used  
12 (<http://seer.cancer.gov/statfacts/html/nhl.html> and  
13 <http://seer.cancer.gov/statfacts/html/livibd.html>).

14 In both cases, an underlying assumption in deriving the relative potencies is that the  
15 relative values of the age-specific background incidence risks for the person-years from the  
16 epidemiologic studies for each tumor type approximate the relative values of the lifetime  
17 background incidence risks for those tumor types. In other words, at least on a proportional  
18 basis, the lifetime background incidence risks (for the United States population) for each site  
19 approximate the age-specific background incidence risks for the study populations. A further  
20 assumption is that the lifetime risk of RCC up to 85 years is an adequate approximation to the  
21 full lifetime risk, which is what was used for the other two tumor types. The first calculation,  
22 based on the results of the meta-analyses for the three tumor types, has the advantage of being  
23 based on a large data set, incorporating data from many different studies. However, this  
24 calculation relies on a number of additional assumptions. First, it is assumed that the RRm's  
25 from the meta-analyses, which are based on different groups of studies, reflect similar overall  
26 TCE exposures, i.e., that the overall TCE exposures are similar across the different groups of  
27 studies that went into the different meta-analyses for the three tumor types. Second, it is  
28 assumed that the RRm's, which incorporate RR estimates for both mortality and incidence,  
29 represent good estimates for cancer incidence risk from TCE exposure. In addition, it is assumed  
30 that the RRm for kidney cancer, for which RCC estimates from individual studies were used  
31 when available, is a good estimate for the overall RR for RCC and that the RRm estimate for  
32 NHL, for which different studies used different classification schemes, is a good estimate for the  
33 overall RR for NHL. The second calculation, based on the results of the Raaschou-Nielsen et al.  
34 (2003) study, the largest single study with RR estimates for all three tumor types, has the  
35 advantage of having RR estimates that are directly comparable. In addition, the

1 Raaschou-Nielsen et al. study provided data for the precise tumor types of interest for the  
2 calculation, i.e., RCC, NHL, and liver (and biliary) cancer.

3 The input data and results of the calculations are presented in Table 5-46. The value for  
4 the ratio of the sum of the extra risks to the extra risk for RCC alone was 3.28 in calculation #1  
5 and 4.36 in calculation #2, which together suggest that 4 is a reasonable factor to use to adjust  
6 the inhalation unit risk estimate based on RCC for multiple sites to obtain a total cancer unit risk  
7 estimate.<sup>55</sup> Using this factor to adjust the unit risk estimate based on RCCs entails the further  
8 fundamental assumption that the dose-response relationships for the other two tumor types (NHL  
9 and liver cancer) are similarly linear, i.e., that the relative potencies are roughly maintained at  
10 lower exposure levels. This assumption is consistent with EPA's *Guidelines for Carcinogen*  
11 *Risk Assessment* (U.S. EPA, 2005c), which recommends low-dose linear extrapolation in the  
12 absence of sufficient evidence to support a nonlinear MOA.

13 Applying the factor of 4 to the  
14 lifetime extra RCC incidence unit  
15 risk estimate of  $5.49 \times 10^{-3}$  per ppm  
16 ( $1.0 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ) of continuous  
17 TCE exposure yields a cancer unit  
18 risk estimate of  $2.2 \times 10^{-2}$  per ppm  
19 ( $4.1 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ). Table 5-46  
20 also presents calculations for just  
21 kidney and NHL extra risks  
22 combined, because the strongest  
23 human evidence is for those two  
24 tumor types. For those two tumor  
25 types, the calculations support a  
26 factor of 3.56 Applying this factor to  
27 the RCC unit risk estimate yields an  
28 estimate of  $1.6 \times 10^{-2}$  per ppm,  
29 which results in the same estimate as  
30 for the three tumor types combined  
31 when finally rounded to one

---

55 Both the geometric and arithmetic means of the two values for the ratio are 3.8, which rounds to 4, in keeping with the imprecise nature of the adjustment factor. The factor of 4 is within 25% of either calculated ratio.

56 The geometric and mean of the two values for the ratio, 2.62 and 3.29, is 2.96, and the arithmetic mean is 2.94, which both round to 3, in keeping with the imprecise nature of the adjustment factor. The factor of 3 is within 15% of either calculated ratio.

1 significant figure, i.e.,  $2 \times 10^{-2}$  per  
2 ppm (or  $3 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ , which is  
3 still similar to the three-tumor-type  
4 estimate in those units).

5 In addition to the uncertainties in the underlying RCC estimate, there are uncertainties  
6 related to the assumptions inherent in these calculations for adjusting to multiple sites, as  
7 detailed above. Nonetheless, the fact that the calculations based on two different data sets  
8 yielded comparable values for the adjustment factor (both within 25% of the selected factor of 4)  
9 provides more robust support for the use of the factor of 4. Additional uncertainties pertain to  
10 the weight of evidence supporting the association of TCE exposure with increased risk of cancer  
11 for the three cancer types. As discussed in Section 4.11.2, it was found that the weight of  
12 evidence for kidney cancer was sufficient to classify TCE as “carcinogenic to humans.” It was  
13 also concluded that there was strong evidence that TCE causes NHL as well, although the  
14 evidence for liver cancer was more limited. In addition, the rodent studies demonstrate clear  
15

1 **Table 5-46. Relative contributions to extra risk for cancer incidence from**  
 2 **TCE exposure for multiple tumor types**  
 3

	<b>RR</b>	<b>Ro</b>	<b>Rx</b>	<b>Extra risk</b>	<b>Ratio to kidney value</b>
<b>Calculation #1: using RR estimates from the meta-analyses</b>					
Kidney (RCC)	1.27	0.0107	0.01359	0.002920	1
NHL	1.23	0.0202	0.02485	0.004742	1.62
Liver (and biliary) cancer	1.29	0.0066	0.008514	0.001927	0.66
			<b>sum</b>	0.009589	<b>3.28</b>
Kidney + NHL only			<b>sum</b>	0.007662	2.62
<b>Calculation #2: using RR estimates from Rasschou-Nielsen et al. (2003)</b>					
Kidney (RCC)	1.20	0.0107	0.01284	0.002163	1
NHL	1.24	0.0202	0.02505	0.004948	2.29
Liver (and biliary) cancer	1.35	0.0066	0.008910	0.002325	1.07
			<b>sum</b>	0.009436	<b>4.36</b>
Kidney + NHL only			<b>sum</b>	0.007111	3.29

4  
 5  
 6 evidence of multisite carcinogenicity, with tumor types including those for which associations  
 7 with TCE exposure are observed in human studies, i.e., liver and kidney cancers and NHLs.  
 8 Overall, the evidence was found to be sufficiently persuasive to support the use of the adjustment  
 9 factor of 4 based on these three cancer types, resulting in a cancer inhalation unit risk estimate of  
 10  $2.2 \times 10^{-2}$  per ppm ( $4.1 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ). Alternatively, if one were to use the factor based  
 11 only on the two cancer types with the strongest human evidence, the cancer inhalation unit risk  
 12 estimate would be only slightly reduced (25%).  
 13

#### 5.2.2.1.7. Route-to-Route Extrapolation Using Physiologically Based Pharmacokinetic (PBPK) Model

14 Route-to-route extrapolation of the inhalation unit risk estimate was performed using the  
 15 PBPK model described in Section 3.5. The (partial) unit risk estimates for NHL and liver cancer  
 16 were derived as for the total cancer inhalation unit risk estimate in Section 5.2.2.2 above, except

1 that the ratios of extra risk for the individual tumor types relative to kidney cancer were used as  
2 adjustment factors rather than the ratio of the sum. As presented in Table 5-46, for NHL, the  
3 ratios from the two different calculations were 1.62 and 2.29, so a factor of 2 was used; for liver  
4 cancer, the ratios were 0.66 and 1.07, so a factor of 1 was used. (With the ratio of one for kidney  
5 cancer itself, the combined adjustment factor is 4, reproducing the factor of 4 used to estimate  
6 the total cancer unit risk from the multiple sites in Section 5.2.2.2.)

7 Because different internal dose-metrics are preferred for each target tissue site, a separate  
8 route-to-route extrapolation was performed for each site-specific unit risk estimate calculated in  
9 Sections 5.2.2.1 and 5.2.2.2. As shown in Figure 5-7, the approach taken to apply the human  
10 PBPK model in the low-dose range where external and internal doses are linearly related to  
11 derive a conversion that is the ratio of internal dose per mg/kg/day to internal dose per ppm. The  
12 expected value of the population mean for this conversion factor (in ppm per mg/kg/day) was  
13 used to extrapolate each inhalation unit risk in units of risk per ppm to an oral slope factor in  
14 units of risk per mg/kg/day. Note that this conversion is the *mean of the ratio* of internal dose  
15 predictions, and is not the same as taking the *ratio of the mean* of internal dose predictions in  
16 Table 5-35.<sup>57</sup>


17 Table 5-47 shows the results of this  
18 route-to-route extrapolation for the  
19 “primary” and “alternative”  
20 dose-metrics. For reference,  
21 route-to-route extrapolation based on  
22 total intake (i.e., ventilation rate  $\times$  air  
23 concentration = oral dose  $\times$  BW)  
24 using the parameters in the PBPK  
25 model would yield an expected  
26 population average conversion of  
27 0.95 ppm per mg/kg/day. For  
28 TotMetabBW34,  
29 TotOxMetabBW34, and  
30 AMetLiv1BW34, the conversion is  
31 2.0–2.8 ppm per mg/kg/day, greater  
32 than that based on intake. This is

---

<sup>57</sup>For route-to-route extrapolation based on dose-response analysis performed on internal dose, as is the case for rodent bioassays, it would be appropriate to use the values in Table 5-35 to first “unconvert” the unit risk based on one route, and then reconvert to a unit risk based on the other route.

1 because of the greater metabolic first  
2 pass in the liver, which leads to a  
3 higher percentage of intake being  
4 metabolized via oral exposure  
5 relative to inhalation exposure for  
6 the same intake. Conversely, for the  
7 AUC in blood, the conversion is 0.14  
8 ppm per mg/kg/day, less than that  
9 based on intake—the greater first  
10 pass in the liver means lower blood  
11 levels of parent compound via oral  
12 exposure relative to inhalation for  
13 the same intake. The conversion for  
14 the primary dose-metric for the  
15 kidney, ABioactDCVCBW34, is 1.7  
16 ppm per mg/kg/day, less than that for  
17 total, oxidative, or liver oxidative  
18 metabolism. This is because the  
19 majority of metabolism in first pass  
20 through the liver is via oxidation,  
21 whereas with inhalation exposure,  
22 more parent compound reaches the  
23 kidney, in which metabolism is via  
24 GSH conjugation.

25 When one sums the oral slope factor estimates based on the primary (preferred)  
26 dose-metrics for the three individual tumor types shown in Table 5-47, the resulting total cancer  
27 oral slope factor estimate is  **$4.64 \times 10^{-2}$  per mg/kg/day**. In the case of the oral  
28 route-extrapolated results, the ratio of the risk estimate for the three tumor types combined to the  
29 risk estimate for kidney cancer alone is 5.0. This value differs from the factor of 4 used for the



1



1 **Figure 5-7. Flow-chart for route-to-route extrapolation of human**  
2 **site-specific cancer inhalation unit risks to oral slope factors. Square nodes**  
3 **indicate point values, circle nodes indicate distributions, and the inverted**  
4 **triangle indicates a (deterministic) functional relationship.**  
5

**Table 5-47. Route-to-route extrapolation of site-specific inhalation unit risks to oral slope factors**

	<b>Kidney</b>	<b>NHL</b>	<b>Liver</b>
Inhalation unit risk (risk per ppm)	$5.49 \times 10^{-3}$	$1.10 \times 10^{-2}$	$5.49 \times 10^{-3}$
Primary dose-metric	ABioactDCVCBW34 <sup>a</sup>	TotMetabBW34	AMetLiv1BW34
ppm per mg/kg/day <sup>b</sup>	1.70	1.97	2.82
Oral slope factor (risk per mg/kg/day)	$9.33 \times 10^{-3}$	$2.16 \times 10^{-2}$	$1.55 \times 10^{-2}$
Alternative dose-metric	TotMetabBW34	AUCCBld	TotOxMetabBW34
ppm per mg/kg/day <sup>b</sup>	1.97	0.137	2.04
Oral slope factor (risk per mg/kg/day)	$1.08 \times 10^{-2}$	$1.50 \times 10^{-3}$	$1.12 \times 10^{-2}$

<sup>a</sup>The AMetGSHBW34 dose-metric gives the same route-to-route conversion because there is no route dependence in the pathway between GSH conjugation and DCVC bioactivation.

<sup>b</sup>Average of expected population mean of males and females. Male and female estimates differed by <1% for ABioactDCVCBW34; TotMetabBW34, AMetLiv1BW34, and TotOxMetabBW34, and <15% for AUCCBld. Uncertainty on the population mean route-to-route conversion, expressed as the ratio between the 97.5% quantile the 2.5% quantile, is about 2.6-fold for ABioactDCVCBW34, 1.5-fold for TotMetabBW34, AMetLiv1BW34, and TotOxMetabBW34, and about 3.4-fold for AUCCBld.

total cancer inhalation unit risk estimate because of the different dose-metrics used for the different tumor types when the route-to-route extrapolation is performed. If only the kidney cancer and NHL results, for which the evidence is strongest, were combined, the resulting total cancer oral slope factor estimate would be  $3.09 \times 10^{-2}$  per mg/kg/day, and the ratio of this risk estimate to that for kidney cancer alone would be 3.3.

If one were to use some of the risk estimates based on alternative dose-metrics in Table 5-40, the total cancer risk estimate would vary depending on for which tumor type(s) an alternative metric was used. The most extreme difference would occur when the alternative metric is used for NHL and liver tumors; in that case, the resulting total cancer oral slope factor estimate would be  $2.20 \times 10^{-2}$  per mg/kg/day, and the ratio of this risk estimate to that for kidney cancer alone (based on the primary dose-metric of ABioactDCVCBW34) would be 2.4.

The uncertainties in these conversions are relatively modest. As discussed in the note to Table 5-47, the 95% confidence range for the route-to-route conversions at its greatest spans 3.4-fold. The greatest uncertainty is in the selection of the dose-metric for NHL, since the use of the alternative dose-metric of AUCCBld yields a converted oral slope factor that is 14-fold lower

1 than that using the primary dose-metric of TotMetabBW34. However, for the other two tumor  
2 sites, the range of conversions is tighter, and lies within threefold of the conversion based solely  
3 on intake.  
4

### 5.2.3. Summary of Unit Risk Estimates

#### 5.2.3.1.1. Inhalation Unit Risk Estimate

5 The inhalation unit risk for TCE is defined as a plausible upper bound lifetime extra risk  
6 of cancer from chronic inhalation of TCE per unit of air concentration. The preferred estimate of  
7 the inhalation unit risk for TCE is  $2.20 \times 10^{-2}$  per ppm ( **$2 \times 10^{-2}$  per ppm [ $4 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ]**  
8 rounded to one significant figure), based on human kidney cancer risks reported by Charbotel  
9 et al. (2006) and adjusted for potential risk for tumors at multiple sites. This estimate is based on  
10 good-quality human data, thus avoiding the uncertainties inherent in interspecies extrapolation.

11 This value is supported by inhalation unit risk estimates from multiple rodent bioassays,  
12 the most sensitive of which range from  **$1 \times 10^{-2}$  to  $2 \times 10^{-1}$  per ppm [ $2 \times 10^{-6}$  to**  
13  **$3 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ]**. From the inhalation bioassays selected for analysis in Section 5.2.1.1, and  
14 using the preferred PBPK model-based dose-metrics, the inhalation unit risk estimate for the  
15 most sensitive sex/species is  $8 \times 10^{-2}$  per ppm [ $2 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ], based on kidney adenomas  
16 and carcinomas reported by Maltoni et al. (1986) for male Sprague-Dawley rats. Leukemias and  
17 Leydig cell tumors were also increased in these rats, and, although a combined analysis for these  
18 tumor types that incorporated the different site-specific preferred dose-metrics was not  
19 performed, the result of such an analysis is expected to be similar, about  $9 \times 10^{-2}$  per ppm  
20 [ $2 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ]. The next most sensitive sex/species from the inhalation bioassays is the  
21 female mouse, for which lymphomas were reported by Henschler et al. (1980); these data yield a  
22 unit risk estimate of  $1.0 \times 10^{-2}$  per ppm [ $2 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ]. In addition, the 90% confidence  
23 intervals reported in Table 5-41 for male rat kidney tumors from Maltoni et al. (1986) and female  
24 mouse lymphomas from Henschler et al. (1980), derived from the quantitative analysis of PBPK  
25 model uncertainty, both included the estimate based on human data of  $2 \times 10^{-2}$  per ppm.  
26 Furthermore, PBPK model-based route-to-route extrapolation of the results for the most sensitive  
27 sex/species from the oral bioassays, kidney tumors in male Osborne-Mendel rats and testicular  
28 tumors in Marshall rats (NTP, 1988), leads to inhalation unit risk estimates of  $2 \times 10^{-1}$  per ppm  
29 [ $3 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ] and  $4 \times 10^{-2}$  per ppm [ $8 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ], respectively, with the preferred  
30 estimate based on human data falling within the route-to-route extrapolation of the 90%

1 confidence intervals reported in Table 5-42.58 Finally, for all these estimates, the ratios of  
2 BMDs to the BMDLs did not exceed a value of three, indicating that the uncertainties in the  
3 dose-response modeling for determining the POD in the observable range are small.

4 Although there are uncertainties in these various estimates, as discussed in  
5 Sections 5.2.1.4, 5.2.2.1.3, and 5.2.2.2, confidence in the proposed inhalation unit risk estimate  
6 of  $2 \times 10^{-2}$  per ppm [ $4 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ], based on human kidney cancer risks reported by  
7 Charbotel et al. (2006) and adjusted for potential risk for tumors at multiple sites (as discussed in  
8 Section 5.2.2.2), is further increased by the similarity of this estimate to estimates based on  
9 multiple rodent data sets.

#### 5.2.3.1.2. Oral Slope Factor Estimate

11 The oral slope factor for TCE is defined as a plausible upper bound lifetime extra risk of  
12 cancer from chronic ingestion of TCE per mg/kg/day oral dose. The preferred estimate of the  
13 oral slope factor is  $4.64 \times 10^{-2}$  per mg/kg/day ( **$5 \times 10^{-2}$  per mg/kg/day** rounded to  
14 one significant figure), resulting from PBPK model-based route-to-route extrapolation of the  
15 inhalation unit risk estimate based on the human kidney cancer risks reported in Charbotel et al.  
16 (2006) and adjusted for potential risk for tumors at multiple sites. This estimate is based on  
17 good-quality human data, thus avoiding uncertainties inherent in interspecies extrapolation. In  
18 addition, uncertainty in the PBPK model-based route-to-route extrapolation is relatively low  
19 ([Chiu, 2006](#); Chiu and White, 2006). In this particular case, extrapolation using different  
20 dose-metrics yielded expected population mean risks within about a twofold range, and, for any  
21 particular dose-metric, the 95% confidence interval for the extrapolated population mean risks  
22 for each site spanned a range of no more than about threefold.

23 This value is supported by oral slope factor estimates from multiple rodent bioassays, the  
24 most sensitive of which range from  **$3 \times 10^{-2}$  to  $3 \times 10^{-1}$  per mg/kg/day**. From the oral bioassays  
25 selected for analysis in Section 5.2.1.1, and using the preferred PBPK model-based dose-metrics,

---

58For oral-to-inhalation extrapolation of NTP ([1988](#)) male rat kidney tumors, the unit risk estimate of  $2.5 \times 10^{-1}$  per mg/kg/day using the ABioactDCVCBW34 dose metric, from Table 5-37, is divided by the average male and female internal doses at 0.001 mg/kg/day, (0.00504/0.001), and then multiplied by the average male and female internal doses at 0.001 ppm, (0.00324/0.001), both from Table 5-35, to yield a unit risk of  $1.6 \times 10^{-1}$  [ $3.0 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ]. For oral-to-inhalation extrapolation of NTP ([1988](#)) male rat testicular tumors, the unit risk estimate of  $7.1 \times 10^{-2}$  per mg/kg/day using the TotMetabBW34 dose metric, from Table 5-37, is divided by the male internal dose at 0.001 mg/kg/day, (0.0192/0.001), and then multiplied by the male internal doses at 0.001 ppm, (0.0118/0.001), both from Table 5-35, to yield a unit risk of  $4.4 \times 10^{-2}$  [ $8.1 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ].

1 the oral slope factor estimate for the most sensitive sex/species is  $3 \times 10^{-1}$  per mg/kg/day, based  
2 on kidney tumors in male Osborne-Mendel rats (NTP, 1988). The oral slope factor estimate for  
3 testicular tumors in male Marshall rats (NTP, 1988) is somewhat lower at  $7 \times 10^{-2}$  per  
4 mg/kg/day. The next most sensitive sex/species result from the oral studies is for male mouse  
5 liver tumors (NCI, 1976), with an oral slope factor estimate of  $3 \times 10^{-2}$  per mg/kg/day. In  
6 addition, the 90% confidence intervals reported in Table 5-42 for male Osborne-Mendel rat  
7 kidney tumors (NTP, 1988), male F344 rat kidney tumors (NTP, 1990), and male Marshall rat  
8 testicular tumors (NTP, 1988), derived from the quantitative analysis of PBPK model  
9 uncertainty, all included the estimate based on human data of  $5 \times 10^{-2}$  per mg/kg/day, while the  
10 upper 95% confidence bound for male mouse liver tumors from NCI (1976) was slightly below  
11 this value at  $4 \times 10^{-2}$  per mg/kg/day. Furthermore, PBPK model-based route-to-route  
12 extrapolation of the most sensitive endpoint from the inhalation bioassays, male rat kidney  
13 tumors from Maltoni et al. (1986), leads to an oral slope factor estimate of  $1 \times 10^{-1}$  per  
14 mg/kg/day, with the preferred estimate based on human data falling within the route-to-route  
15 extrapolation of the 90% confidence interval reported in Table 5-41.<sup>59</sup> Finally, for all these  
16 estimates, the ratios of BMDs to the BMDLs did not exceed a value of three, indicating that the  
17 uncertainties in the dose-response modeling for determining the POD in the observable range are  
18 small.

19 Although there are uncertainties in these various estimates, as discussed in  
20 Sections 5.2.1.4, 5.2.2.1.3, 5.2.2.2, and 5.2.2.3, confidence in the proposed oral slope factor  
21 estimate of  $5 \times 10^{-2}$  per mg/kg/day, resulting from PBPK model-based route-to-route  
22 extrapolation of the inhalation unit risk estimate based on the human kidney cancer risks reported  
23 in Charbotel et al. (2006) and adjusted for potential risk for tumors at multiple sites (as discussed  
24 in Section 5.2.2.2), is further increased by the similarity of this estimate to estimates based on  
25 multiple rodent data sets.

26

### 5.2.3.1.3. Application of Age-Dependent Adjustment Factors

27 When there is sufficient weight of evidence to conclude that a carcinogen operates  
28 through a mutagenic MOA, and in the absence of chemical-specific data on age-specific

---

<sup>59</sup>For the Maltoni et al. (1986) male rat kidney tumors, the unit risk estimate of  $8.3 \times 10^{-2}$  per ppm using the ABioactDCVCBW34 dose metric, from Table 5-36, is divided by the average male and female internal doses at 0.001 ppm, (0.00324/0.001) and then multiplied by the average male and female internal doses at 0.001 mg/kg/day, (0.00504/0.001), both from Table 5-35, to yield a unit risk of  $1.3 \times 10^{-1}$  per mg/kg/day.

1 susceptibility, EPA's *Supplemental Guidance for Assessing Susceptibility from Early-Life*  
2 *Exposure to Carcinogens* (U.S. EPA, 2005d) advises that increased early-life susceptibility be  
3 assumed and recommends that default age-dependent adjustment factors (ADAFs) be applied to  
4 adjust for this potential increased susceptibility from early-life exposure. As discussed in  
5 Section 4.4, there is sufficient evidence to conclude that a mutagenic MOA is operative for  
6 TCE-induced kidney tumors. In addition, as described in Section 4.10, TCE-specific data are  
7 inadequate for quantification of early-life susceptibility to TCE carcinogenicity. Therefore, as  
8 recommended in the *Supplemental Guidance*, the default ADAFs are applied.

9 See the *Supplemental Guidance* for detailed information on the general application of  
10 these adjustment factors. In brief, the *Supplemental Guidance* establishes ADAFs for  
11 three specific age groups. The current ADAFs and their age groupings are 10 for <2 years,  
12 three for 2 to <16 years, and one for 16 years and above (U.S. EPA, 2005d). For risk  
13 assessments based on specific exposure assessments, the 10-fold and threefold adjustments to the  
14 slope factor or unit risk estimates are to be combined with age-specific exposure estimates when  
15 estimating cancer risks from early-life (<16-years-of-age) exposure. Currently, due to lack of  
16 appropriate data, no ADAFs are used for other life-stages, such as the elderly. However, the  
17 ADAFs and their age groups may be revised over time. The most current information on the  
18 application of ADAFs for cancer risk assessment can be found at  
19 [www.epa.gov/cancerguidelines](http://www.epa.gov/cancerguidelines).

20 In the case of TCE, the inhalation unit risk and oral slope factor estimates reflect lifetime  
21 risk for cancer at multiple sites, and a mutagenic MOA has been established for one of these  
22 sites, the kidney. The following subsections illustrate how one might apply the default ADAFs  
23 to the *kidney-cancer component* of the inhalation unit risk and oral slope factor estimates for  
24 TCE. These are **sample calculations**, and individual risk assessors should use exposure-related  
25 parameters (e.g., age-specific water ingestion rates) that are appropriate for their particular risk  
26 assessment applications.

27 In addition to the uncertainties discussed above for the inhalation and oral total cancer  
28 unit risk or slope factor estimates, there are uncertainties in the application of ADAFs to adjust  
29 for potential increased early-life susceptibility. For one thing, the adjustment is made only for  
30 the kidney-cancer component of total cancer risk because that is the tumor type for which the  
31 weight of evidence was sufficient to conclude that TCE-induced carcinogenesis operates through  
32 a mutagenic MOA. However, it may be that TCE operates through a mutagenic MOA for other  
33 tumor types as well or that it operates through other MOAs that might also convey increased  
34 early-life susceptibility. Additionally, the ADAFs are general default factors, and it is uncertain

1 to what extent they reflect increased early-life susceptibility for exposure to TCE, if increased  
2 early-life susceptibility occurs.

3 Furthermore, the assumption of increased early-life susceptibility, invoked by the finding  
4 of a mutagenic MOA for kidney cancer, is in contradiction to the assumption that RR is  
5 independent of age that was used to derive the unit risk estimates in the life-table analysis. In  
6 some other assessments faced with a similar situation, a small modification has been made to the  
7 derivation of the unit risk estimate to avoid the contradictory assumptions (by calculating an  
8 adult-exposure-only unit risk estimate for the application of ADAFs). This has the effect of  
9 slightly reducing the unit risk estimate to which the ADAFs are applied. Because there are  
10 multiple cancer types for TCE but the finding of a mutagenic MOA applies to only one of them,  
11 and because under these circumstances application of the ADAFs already has a minimal impact  
12 on the total risk for most exposure scenarios, as discussed with respect to the examples in  
13 Sections 5.2.3.3.1 and 5.2.3.3.2 below, no attempt was made to modify the kidney cancer unit  
14 risk estimate for this assessment. Such a modification would have substantially increased the  
15 complexity of the calculations, which are already more elaborate than the standard ADAF  
16 applications, without having much quantitative impact on the final risk estimates.

#### 5.2.3.1.4. Example application of age-dependent adjustment factors (ADAFs) for inhalation exposures.

18 A calculation template for application of the ADAFs is provided in Table 5-48. In the  
19 example provided, it is assumed that an individual is exposed to  $1 \mu\text{g}/\text{m}^3$  in air from birth  
20 through age 70 years. Using the template, risk estimates for different exposure scenarios can be  
21 obtained by changing the exposure concentrations (including possibly zero for some age groups).  
22 The steps in the calculation are as follows:

- 23  
24
- 25 (1) Separate the kidney cancer contribution from the NHL + liver cancer contribution to the  
26 inhalation unit risk estimate. From Section 5.2.2.1.4, the kidney lifetime unit risk is  
27  $1.0 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$  in air. Subtracting this from the total lifetime unit risk of  $4.1 \times 10^{-6}$   
28 per  $\mu\text{g}/\text{m}^3$  from Section 5.2.2.2 results in the estimated contribution of NHL + liver  
29 cancer being  $3.1 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ .
  - 30 (2) Assign a lifetime unit risk estimate for each age group. The template shows the  
31 recommended age groupings from U.S. EPA (2005b) in Column A, along with the age  
32 group duration (Column D), and the fraction of lifetime each age group represents  
33 (Column E; used as a duration adjustment). For each age group, the (unadjusted) lifetime

1 unit risk estimates for kidney cancer, total cancer, and NHL + liver cancer are shown in  
2 columns Column F, I, and J, respectively.

3 (3) For each age group, the kidney cancer inhalation unit risk estimate (Column F) is  
4 multiplied by the risk per  $\mu\text{g}/\text{m}^3$  equivalence (Column B), the exposure concentration  
5 (Column C), the duration adjustment (Column E), and the ADAF (Column G), to obtain  
6 the partial risk from exposure during those ages (Column H). For inhalation exposures, a  
7 “risk per  $\mu\text{g}/\text{m}^3$  equivalence” of 1 is assumed across age groups (i.e., equivalent risk from  
8 equivalent exposure levels in air, independent of body size), as shown in Column B. In  
9 this calculation, a unit lifetime exposure of  $1 \mu\text{g}/\text{m}^3$  is assumed, as shown in Column C.

10



**Table 5-48. Sample calculation for total lifetime cancer risk based on the kidney unit risk estimate, potential risk for NHL and liver cancer, and potential increased early-life susceptibility, assuming a constant lifetime exposure to 1 µg/m<sup>3</sup> of TCE in air**

Col A	Col B	Col C	Col D	Col E	Col F	Col G	Col H	Col I	Col J	Col K	Col L	
	Exposure scenario parameters				Dose-response assessment calculations							
<b>Units:</b>		(µg/m <sup>3</sup> )	year	-	(µg/m <sup>3</sup> ) <sup>-1</sup>	-		(µg/m <sup>3</sup> ) <sup>-1</sup>	(µg/m <sup>3</sup> ) <sup>-1</sup>			
<b>Age group</b>	Risk per µg/m <sup>3</sup> air equivalence	Exposure concentration	Age group duration	Duration adjustment (Col D/Sum(Col D))	Kidney cancer unadjusted lifetime unit risk (p 5-137 [5.2.2.1.4])	Default ADAF	<b>Kidney cancer ADAF-adjusted partial risk (Col B × Col C × Col E × Col F × Col G)</b>	Kidney cancer+NHL+ liver cancer unadjusted lifetime unit risk (p 5-139 [5.2.2.2])	NHL+ liver cancer lifetime unit risk (Col I – Col F)	<b>NHL and liver cancer partial risk (Col B × Col C × Col E × Col J)</b>	<b>Total partial risk (Col H + Col K)</b>	
Birth to <1 month	1	1.000	0.083	0.0012	1.0E-06	10	<b>1.2E-08</b>	4.1E-06	3.1E-06	<b>3.7E-09</b>	<b>1.6E-08</b>	
1 to <3 months	1	1.000	0.167	0.0024	1.0E-06	10	<b>2.4E-08</b>	4.1E-06	3.1E-06	<b>7.4E-09</b>	<b>3.1E-08</b>	
3 to <6 months	1	1.000	0.250	0.0036	1.0E-06	10	<b>3.6E-08</b>	4.1E-06	3.1E-06	<b>1.1E-08</b>	<b>4.7E-08</b>	
6 to <12 months	1	1.000	0.500	0.0071	1.0E-06	10	<b>7.1E-08</b>	4.1E-06	3.1E-06	<b>2.2E-08</b>	<b>9.4E-08</b>	
1 to <2 years	1	1.000	1.000	0.0143	1.0E-06	10	<b>1.4E-07</b>	4.1E-06	3.1E-06	<b>4.4E-08</b>	<b>1.9E-07</b>	
2 to <3 years	1	1.000	1.000	0.0143	1.0E-06	3	<b>4.3E-08</b>	4.1E-06	3.1E-06	<b>4.4E-08</b>	<b>8.7E-08</b>	
3 to <6 years	1	1.000	3.000	0.0429	1.0E-06	3	<b>1.3E-07</b>	4.1E-06	3.1E-06	<b>1.3E-07</b>	<b>2.6E-07</b>	
6 to <11 years	1	1.000	5.000	0.0714	1.0E-06	3	<b>2.1E-07</b>	4.1E-06	3.1E-06	<b>2.2E-07</b>	<b>4.4E-07</b>	
11 to <16 years	1	1.000	5.000	0.0714	1.0E-06	3	<b>2.1E-07</b>	4.1E-06	3.1E-06	<b>2.2E-07</b>	<b>4.4E-07</b>	
16 to <21	1	1.000	5.000	0.0714	1.0E-06	1	<b>7.1E-08</b>	4.1E-06	3.1E-06	<b>2.2E-07</b>	<b>2.9E-07</b>	
21-70	1	1.000	49.000	0.7000	1.0E-06	1	<b>7.0E-07</b>	4.1E-06	3.1E-06	<b>2.2E-06</b>	<b>2.9E-06</b>	
		<b>Total exposure duration:</b>	<b>70</b>							<b>Total unit risk:</b>	<b>4.8E-06</b>	

- 1 (4) For each age group, the NHL + liver cancer unit risk estimate (Column J) is multiplied by  
2 the risk per  $\mu\text{g}/\text{m}^3$  equivalence (Column B), the exposure concentration (Column C), and  
3 the duration adjustment (Column E), to obtain the partial risk from exposure during those  
4 ages (Column K).
- 5 (5) For each age group, the ADAF-adjusted partial risk for kidney cancer (Column H) is  
6 added to the partial risk for NHL + liver cancer (Column K), resulting in the total partial  
7 risk (Column L).
- 8 (6) The age-group-specific partial risks are added together to obtain the estimated total  
9 lifetime risk (bottom of Column L).

10  
11  
12 From the example calculation, based on continuous exposure to  $1 \mu\text{g}/\text{m}^3$  from birth to age 70, the  
13 estimated total lifetime risk is  $4.8 \times 10^{-6}$ , which corresponds to a lifetime unit risk estimate of  
14  $4.8 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ . The risk-specific air concentrations at risk levels of  $10^{-6}$ ,  $10^{-5}$ , and  $10^{-4}$   
15 are 0.21, 2.1, and  $21 \mu\text{g}/\text{m}^3$ , respectively.

16 This total cancer unit risk estimate of  $4.8 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$  ( $2.6 \times 10^{-2}$  per ppm), adjusted  
17 for potential increased early-life susceptibility, is only minimally (17.5%) increased over the  
18 unadjusted total cancer unit risk estimate because the kidney cancer risk estimate that gets  
19 adjusted for potential increased early-life susceptibility is only part of the total cancer risk  
20 estimate. Thus, foregoing the ADAF adjustment in the case of full lifetime calculations will not  
21 seriously impact the resulting risk estimate. For less-than-lifetime exposure calculations, the  
22 impact of applying the ADAFs will increase as the proportion of time at older ages decreases.  
23 The maximum impact will be when exposure is for only the first 2 years of life, in which case the  
24 partial lifetime total cancer risk estimate for exposure to  $1 \mu\text{g}/\text{m}^3$  adjusted for potential increased  
25 early-life susceptibility is  $10 \times (1 \mu\text{g}/\text{m}^3) \times (1.0 \times 10^{-6} \text{ per } \mu\text{g}/\text{m}^3) \times (2 / 70)$  for the kidney cancer  
26 risk +  $(1 \mu\text{g}/\text{m}^3) \times (3.1 \times 10^{-6} \text{ per } \mu\text{g}/\text{m}^3) \times (2 / 70)$  for the NHL and liver cancer, or  $3.7 \times 10^{-7}$ ,  
27 which is over three times greater than the unadjusted partial lifetime total cancer risk estimate for  
28 exposure to  $1 \mu\text{g}/\text{m}^3$  of  $(1 \mu\text{g}/\text{m}^3) \times (4.1 \times 10^{-6} \text{ per } \mu\text{g}/\text{m}^3) \times (2 / 70)$ , or  $1.2 \times 10^{-7}$ .

#### 5.2.3.1.5. Example application of age-dependent adjustment factors (ADAFs) for oral drinking water exposures

30 For oral exposures, the calculation of risk estimates adjusted for potential increased  
31 early-life susceptibility is complicated by the fact that for a constant exposure level, e.g., a  
32 constant concentration of TCE in drinking water, doses will vary by age because of different  
33 age-specific uptake rates, e.g., drinking water consumption rates. Different EPA Program or

1 Regional Offices may have different default age-specific uptake rates that they use for risk  
2 assessments for specific exposure scenarios, and the calculations presented below are merely to  
3 illustrate the general approach to applying ADAFs for oral TCE exposures, using exposure to  
4 1 µg/L of TCE in drinking water from birth through age 70 years as an example. Using the  
5 template, risk estimates for different exposure scenarios can be obtained by changing the intake  
6 rates and exposure concentrations (including possibly zero for some age groups). The steps in  
7 the calculation, illustrated in the template in Table 5-49, are as follows:

8  
9  
10 (1) Separate the kidney cancer contribution from the NHL + liver cancer contribution to the  
11 oral slope factor estimate. From Section 5.2.2.3, the kidney lifetime oral slope factor is  
12  $9.3 \times 10^{-3}$  per mg/kg/day. Subtracting this from the total lifetime oral slope factor of  
13  $4.6 \times 10^{-2}$  per mg/kg/day from Section 5.2.2.3 results in an estimated contribution from  
14 NHL + liver cancer of  $3.7 \times 10^{-2}$  per mg/kg/day.

15 (2) Assign a lifetime oral slope factor estimate for each age group. The template shows the  
16 recommended age groupings from U.S. EPA (2005b) in Column A, along with the age  
17 group duration (Column D), and the fraction of lifetime each age group represents  
18 (Column E; used as a duration adjustment). For each age group, the (unadjusted) lifetime  
19 oral slope factor estimates for kidney cancer, total cancer, and NHL + liver cancer are  
20 shown in columns Column F, I, and J, respectively.

21 (3) For each age group, the kidney cancer oral slope factor estimate (Column F) is multiplied  
22 by the drinking water ingestion rate (Column B), the exposure concentration (Column C),  
23 the duration adjustment (Column E), and the ADAF (Column G), to obtain the partial risk  
24 from exposure during those ages (Column H). Age-specific water ingestion rates in  
25 L/kg/day, taken from EPA's *Child-Specific Exposure Factors Handbook* (U.S. EPA,  
26 2008b), are shown in Column B.<sup>60</sup> In this calculation, a lifetime unit exposure of 1 µg/L  
27 is assumed, as shown in Column C.

28 (4) For each age group, the NHL + liver cancer oral slope factor estimate (Column J) is  
29 multiplied by the drinking water ingestion rate (Column B), the exposure concentration

---

60 Values for the 90th percentile were taken from Table 3-19 of U.S. EPA (2008c) (consumers-only estimates of combined direct and indirect water ingestion from community water). The 90th percentile was based on the policy in the U.S. EPA Office of Water for determining risk through direct and indirect consumption of drinking water. Community water was used in the illustration because U.S. EPA only regulates community water sources and not private wells and cisterns or bottled water. Data for "consumers only" (i.e., excluding individuals who did not ingest community water) were used because formula-fed infants (as opposed to breast-fed infants, who consume very little community water), children, and young adolescents are often the population of concern with respect to water consumption. For the 16-21 and 21+ age groups, the standard default rate for adults was used (i.e., 2 L/day ÷ 70 kg, or 0.029 L/kg/day) (U.S. EPA, 1997b, page 3-1), which is identical to the 90th percentile for the 18 to <21 age group.

1 (Column C), and the duration adjustment (Column E), to obtain the partial risk from  
2 exposure during those ages (Column K).

3

**Table 5-49. Sample calculation for total lifetime cancer risk based on the kidney cancer slope factor estimate, potential risk for NHL and liver cancer, and potential increased early-life susceptibility, assuming a constant lifetime exposure to 1 µg/L of TCE in drinking water**

Col A	Col B	Col C	Col D	Col E	Col F	Col G	Col H	Col I	Col J	Col K	Col L	
	Exposure scenario parameters				Dose-response assessment calculations							
<b>Units:</b>	L water/kg/d	mg/L water	year	-	(mg/kg/day) <sup>-1</sup>	-	-	(mg/kg/day) <sup>-1</sup>	(mg/kg/day) <sup>-1</sup>	-	-	
<b>Age group</b>	ingestion rate	Exposure concentration	Age group duration	Duration adjustment (Col D/Sum(Col D))	Kidney cancer unadjusted lifetime slope factor (p 5–144 [Table 5-40])	Default ADAF	<b>Kidney cancer ADAF-adjusted partial risk (Col B × Col C × Col E × Col F × Col G)</b>	Kidney cancer+NHL+ liver cancer unadjusted lifetime slope factor (p 5–143 [5.2.2.3])	NHL+ liver cancer lifetime slope factor (Col I – Col F)	<b>NHL and liver cancer partial risk (Col B × Col C × Col E × Col J)</b>	<b>Total partial risk (Col H + Col K)</b>	
<b>Birth to &lt;1 month</b>	0.238	0.001	0.083	0.0012	9.3E-03	10	<b>2.6E-08</b>	4.6E-02	3.7E-02	<b>1.0E-08</b>	<b>3.7E-08</b>	
<b>1 to &lt;3 months</b>	0.228	0.001	0.167	0.0024	9.3E-03	10	<b>5.0E-08</b>	4.6E-02	3.7E-02	<b>2.0E-08</b>	<b>7.0E-08</b>	
<b>3 to &lt;6 months</b>	0.148	0.001	0.250	0.0036	9.3E-03	10	<b>4.9E-08</b>	4.6E-02	3.7E-02	<b>1.9E-08</b>	<b>6.9E-08</b>	
<b>6 to &lt;12 months</b>	0.112	0.001	0.500	0.0071	9.3E-03	10	<b>7.4E-08</b>	4.6E-02	3.7E-02	<b>2.9E-08</b>	<b>1.0E-07</b>	
<b>1 to &lt;2 years</b>	0.056	0.001	1.000	0.0143	9.3E-03	10	<b>7.4E-08</b>	4.6E-02	3.7E-02	<b>2.9E-08</b>	<b>1.0E-07</b>	
<b>2 to &lt;3 years</b>	0.052	0.001	1.000	0.0143	9.3E-03	3	<b>2.1E-08</b>	4.6E-02	3.7E-02	<b>2.7E-08</b>	<b>4.8E-08</b>	
<b>3 to &lt;6 years</b>	0.049	0.001	3.000	0.0429	9.3E-03	3	<b>5.9E-08</b>	4.6E-02	3.7E-02	<b>7.7E-08</b>	<b>1.4E-07</b>	
<b>6 to &lt;11 years</b>	0.035	0.001	5.000	0.0714	9.3E-03	3	<b>7.0E-08</b>	4.6E-02	3.7E-02	<b>9.2E-08</b>	<b>1.6E-07</b>	
<b>11 to &lt;16 years</b>	0.026	0.001	5.000	0.0714	9.3E-03	3	<b>5.2E-08</b>	4.6E-02	3.7E-02	<b>6.8E-08</b>	<b>1.2E-07</b>	
<b>16 to &lt;21</b>	0.029	0.001	5.000	0.0714	9.3E-03	1	<b>1.9E-08</b>	4.6E-02	3.7E-02	<b>7.6E-08</b>	<b>9.5E-08</b>	
<b>21–70</b>	0.029	0.001	49.000	0.7000	9.3E-03	1	<b>1.9E-07</b>	4.6E-02	3.7E-02	<b>7.5E-07</b>	<b>9.3E-07</b>	
		<b>Total exposure duration:</b>	<b>70</b>							<b>Total unit risk:</b>	<b>1.9E-06</b>	

1 (5) For each age group, the ADAF-adjusted partial risk for kidney cancer (Column H) is  
2 added to the partial risk for NHL + liver cancer (Column K), resulting in the total partial  
3 risk (Column L).

4 (6) The age-group-specific partial risks are added together to obtain the estimated total  
5 lifetime risk (bottom of Column L).  
6  
7

8 Because the TCE intake is not constant across age groups, one does not calculate a  
9 lifetime unit risk estimate in terms of risk per mg/kg/day adjusted for potential increased  
10 early-life susceptibility. One could calculate a unit risk estimate for TCE in drinking water in  
11 terms of  $\mu\text{g/L}$  from the result in Table 5-49, but this is dependent on the water ingestion rates  
12 used. Based on the example calculation assuming continuous exposure to  $1 \mu\text{g/L}$  of TCE in  
13 drinking water from birth to age 70 years and using the drinking water intake rates shown,  
14 estimated total lifetime risk is  $1.9 \times 10^{-6}$ , which corresponds to a lifetime drinking water unit risk  
15 estimate of  $1.9 \times 10^{-6}$  per  $\mu\text{g/L}$ . The corresponding risk-specific drinking water concentrations  
16 at risk levels of  $10^{-6}$ ,  $10^{-5}$ , and  $10^{-4}$  are 0.53, 5.3, and 53  $\mu\text{g/L}$ , respectively. For different  
17 exposure and intake parameters, the risk-specific drinking water concentrations would need to be  
18 recalculated.

19 As with the adjusted inhalation risk estimate in Section 5.2.3.3.1, the lifetime total cancer  
20 risk estimate of  $1.9 \times 10^{-6}$  calculated for lifetime exposure to  $1 \mu\text{g/L}$  of TCE in drinking water  
21 adjusted for potential increased early-life susceptibility is only minimally (25%) increased over  
22 the unadjusted total cancer unit risk estimate. (This calculation is not shown, but if one omits the  
23 ADAFs for each of the age groups in Table 5-49, the resulting total lifetime risk estimate is  
24  $1.5 \times 10^{-6}$ .) Unlike with inhalation exposure under the assumption of ppm equivalence, the oral  
25 intake rates are higher in the potentially more susceptible younger age groups. This would tend  
26 to yield a larger relative impact of adjusting for potential increased early-life susceptibility for  
27 oral risk estimates compared to inhalation risk estimates. In the case of TCE, however, this  
28 impact is partially offset by the lesser proportion of the total oral cancer risk that is accounted for  
29 by the kidney cancer risk, which is the component of total risk that is being adjusted for potential  
30 increased early-life susceptibility, based on the primary dose-metrics (1/5 vs. 1/4 for inhalation).  
31 Thus, as with lifetime inhalation risk, foregoing the ADAF adjustment in the case of full lifetime  
32 calculations will not seriously impact the resulting risk estimate. For less-than-lifetime exposure  
33 calculations, the impact of applying the ADAFs will increase as the proportion of time at older  
34 ages decreases. The maximum impact will be when exposure is for only the first 2 years of life,  
35 in which case the partial lifetime total cancer risk estimate for exposure to  $1 \mu\text{g/L}$  adjusted for  
36 potential increased early-life susceptibility is  $3.8 \times 10^{-7}$  (adding partial risks from Table 5-49 for

1 the appropriate ages groups), which is almost three times greater than the unadjusted partial  
2 lifetime total cancer risk estimate for exposure to 1 µg/L of  $5 \times (0.001 \text{ mg/L}) \times (0.103 \text{ L/kg/day})$   
3  $\times (9.33 \times 10^{-3} \text{ per mg/kg/day}) \times (2/70)$ , or  $1.4 \times 10^{-7}$ , where 5 is the factor for the multiple  
4 cancer types for oral exposure, 0.103 L/kg/day is the time-weighted ingestion rate for the  
5 1<sup>st</sup> two years of life using the rates in Table 5-49,  $9.33 \times 10^{-3} \text{ per mg/kg/day}$  is the unadjusted  
6 oral slope factor estimate for kidney cancer, and 2/70 is the duration adjustment.  
7

### 5.3. KEY RESEARCH NEEDS FOR TCE DOSE-RESPONSE ANALYSES

8 For noncancer dose-response assessment, key research that would substantially improve  
9 the accuracy or utility of TCE noncancer risk estimates includes:

- 10  
11  
12 • Research to obtain toxicokinetic data to better quantify the amount of bioactivation of  
13 DCVC to toxic moiety(ies) in rats and humans, including data on human variability in  
14 DCVC bioactivation.
- 15 • Research to obtain mechanistic data that would identify the active moiety(ies) for  
16 TCE-induced immunological effects and developmental cardiac defects. As a  
17 corollary, data on human variability pharmacokinetics of the active moiety after TCE  
18 exposure would also be informative.
- 19 • Research to obtain mechanistic data that would quantitatively inform the  
20 pharmacodynamic factors that would make individuals more or less susceptible to  
21 kidney, immunological, and developmental cardiac defects induced by TCE.
- 22 • Research to obtain TCE dose-response data on kidney effects, immunological effects,  
23 and developmental cardiac defects at a larger number of doses at and below the  
24 current LOAELs, so as to better describe the dose-response shape at low effect levels.  
25 Ideally, studies would be based on human epidemiologic data with good quantitative  
26 exposure assessment. Studies in laboratory animals would need to address the  
27 limitations in the currently available studies. For example, studies of cardiac defects  
28 would need to address limitations of the Johnson et al. (2003) study described in  
29 Section 4.8.3.3.2.
- 30 • Development of a probabilistic approach to noncancer dose-response analysis that  
31 would enable calculation of a risk-specific dose for noncancer effects, while capturing  
32 uncertainty and variability quantitatively.

1 For cancer dose-response assessment, key research that would substantially improve the  
2 accuracy or utility of TCE cancer risk estimates includes:

- 3 • Research to obtain toxicokinetic data to better quantify the amount of bioactivation of  
4 DCVC to toxic moiety(ies) in humans, including data on human variability in DCVC  
5 bioactivation.
- 6 • Research to obtain mechanistic data that would identify the active moiety(ies) for  
7 TCE-induced liver tumors and NHL. As a corollary, data on human variability  
8 pharmacokinetics of the active moiety after TCE exposure would also be informative.
- 9 • Research to obtain mechanistic data that would quantitatively inform the  
10 pharmacodynamic factors that would make individuals more or less susceptible to  
11 kidney tumors, liver tumors, and NHL induced by TCE. This includes data on  
12 life-stage-specific susceptibility that would replace the default ADAFs for kidney  
13 tumors and the assumption of no life-stage-specific susceptibility for liver tumors and  
14 NHL.
- 15 • Research to obtain human epidemiologic dose-response data on TCE-induced kidney  
16 tumors, liver tumors, and NHL with good quantitative exposure assessment.
- 17 • Research to obtain additional human epidemiologic data on TCE exposure and other  
18 tumors, so as to better estimate the total risk of cancer from TCE exposure.
- 19 • Development of a probabilistic approach to cancer dose-response analysis that would  
20 enable calculation of a differential susceptibility to carcinogenic effects, while  
21 capturing uncertainty and variability quantitatively

1



## 6. MAJOR CONCLUSIONS IN THE CHARACTERIZATION OF HAZARD AND DOSE RESPONSE

### 6.1. HUMAN HAZARD POTENTIAL

1 This section summarizes the human hazard potential for trichloroethylene (TCE). For extensive  
2 discussions and references, see Section 2 for Exposure, Section 3 for toxicokinetics and  
3 physiologically based pharmacokinetic (PBPK) modeling, and Sections 4.1–4.9 for the  
4 epidemiologic and experimental studies of TCE noncancer and cancer toxicity. Section 4.10  
5 summarizes information on susceptibility, and Section 4.11 provides a more detailed summary  
6 and references for noncancer toxicity and carcinogenicity.

#### 6.1.1. Exposure (see Section 2)

8 TCE is a volatile compound with moderate water solubility. Most TCE produced today is used  
9 for metal degreasing. The highest environmental releases are to the air. Ambient air monitoring  
10 data suggest that mean levels have remained fairly constant since 1999 at about  $0.3 \mu\text{g}/\text{m}^3$   
11 ( $0.06 \text{ ppb}$ ). As discussed in Section 2, in 2006, ambient air monitors ( $n = 258$ ) had annual means  
12 ranging from  $0.03\text{--}7.73 \mu\text{g}/\text{m}^3$  with a median of  $0.13$  and an overall average of  $0.23 \mu\text{g}/\text{m}^3$ .  
13 Indoor levels are commonly three or more times higher than outdoor levels due to releases from  
14 building materials and consumer products. Vapor intrusion is a likely significant source in  
15 situations where residences are located near soils or groundwater with high contamination levels  
16 and sparse indoor air sampling had detected TCE levels ranging from  $1\text{--}140 \mu\text{g}/\text{m}^3$ . TCE is  
17 among the most common groundwater contaminants and the one present in the highest  
18 concentration in a summary of ground water analyses reported in 1982. The median level of  
19 TCE in groundwater, based on a large survey by the U.S. Geological Survey for 1985–2001, is  
20  $0.15 \mu\text{g}/\text{L}$ . It has also been detected in a wide variety of foods in the  $1\text{--}100 \mu\text{g}/\text{kg}$  range. None  
21 of the environmental sampling has been done using statistically based national surveys.  
22 However, a substantial amount of air and groundwater data have been collected allowing  
23 reasonably well supported estimates of typical daily intakes by the general population:  
24 inhalation— $13 \mu\text{g}/\text{day}$  and water ingestion— $0.2 \mu\text{g}/\text{day}$ . The limited food data suggest an intake  
25 of about  $5 \mu\text{g}/\text{day}$ , but this must be considered preliminary. Higher exposures have occurred to  
26 various occupational groups, particularly with vapor degreasing that has the highest potential for  
27 exposure because vapors can escape into the work place. For example, past studies of aircraft  
28 workers have shown short term peak exposures in the hundreds of ppm ( $>500,000 \mu\text{g}/\text{m}^3$ ) and  
29 long term exposures in the low tens of ppm ( $>50,000 \mu\text{g}/\text{m}^3$ ). Occupational exposures have

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 likely decreased in recent years due to better release controls, improvements in worker  
2 protection, and substituting other solvents for TCE.  
3 Exposure to a variety of TCE related compounds, which include metabolites of TCE and other  
4 parent compounds that produce similar metabolites, can alter or enhance TCE metabolism and  
5 toxicity by generating higher internal metabolite concentrations than would result from TCE  
6 exposure by itself. Available estimates suggest that exposures to most of these TCE-related  
7 compounds are comparable to or greater than TCE itself.

8

**6.1.2. Toxicokinetics and Physiologically Based Pharmacokinetic Modeling (see Section 3 and Appendix A)**

9 TCE is a lipophilic compound that readily crosses biological membranes. Exposures may occur  
10 via the oral, dermal, and inhalation route, with evidence for systemic availability from each  
11 route. TCE can also be transferred transplacentally and through breast milk ingestion. TCE is  
12 rapidly and nearly completely absorbed from the gut following oral administration, and animal  
13 studies indicate that exposure vehicle may impact the time course of absorption: oily vehicles  
14 may delay absorption whereas aqueous vehicles result in a more rapid increase in blood  
15 concentrations. See Section 3.1 for additional discussion of TCE absorption.

16 Following absorption to the systemic circulation, TCE distributes from blood to solid tissues by  
17 each organ's solubility. This process is mainly determined by the blood:tissue partition  
18 coefficients, which are largely determined by tissue lipid content. Adipose partitioning is high,  
19 so adipose tissue may serve as a reservoir for TCE, and accumulation into adipose tissue may  
20 prolong internal exposures. TCE attains high concentrations relative to blood in the brain,  
21 kidney, and liver—all of which are important target organs of toxicity. TCE is cleared via  
22 metabolism mainly in three organs: the kidney, liver, and lungs. See Section 3.2 for additional  
23 discussion of TCE distribution.

24 The metabolism of TCE is an important determinant of its toxicity. Metabolites are generally  
25 thought to be responsible for toxicity—especially for the liver and kidney. Initially, TCE may be  
26 oxidized via cytochrome P450 (CYP) isoforms or conjugated with glutathione (GSH) by  
27 glutathione S-transferase enzymes. While CYP2E1 is generally accepted to be the CYP isoform  
28 most responsible for TCE oxidation, others forms may also contribute. There are conflicting  
29 data as to which glutathione-S-transferase (GST) isoforms are responsible for TCE conjugation,  
30 with one rat study indicating alpha-class GSTs and another rat study indicating mu and pi-class  
31 GST. The balance between oxidative and conjugative metabolites generally favors the oxidative  
32 pathway, especially at lower concentrations, and inhibition of CYP-dependent oxidation in vitro

1 increases glutathione conjugation in renal preparations. However, different investigators have  
2 reported considerably different rates for TCE conjugation in human liver and kidney cell  
3 fractions, perhaps due to different analytical methods. The inferred flux through the GSH  
4 pathway differs by more than four orders of magnitude across data sets. While the available data  
5 are consistent with the higher values being overestimates, the degree of overestimation is  
6 unclear, and differing results may be attributable to true interindividual variation. Overall, there  
7 remains significant uncertainty in the quantitative estimation of TCE GSH conjugation. See  
8 Section 3.3 for additional discussion of TCE metabolism.

9 Once absorbed, TCE is excreted primarily either in breath as unchanged TCE or carbon dioxide  
10 [CO<sub>2</sub>], or in urine as metabolites. Minor pathways of elimination include excretion of  
11 metabolites in saliva, sweat, and feces. Following oral administration or upon cessation of  
12 inhalation exposure, exhalation of unmetabolized TCE is a major elimination pathway. Initially,  
13 elimination of TCE upon cessation of inhalation exposure demonstrates a steep concentration-  
14 time profile: TCE is rapidly eliminated in the minutes and hours postexposure, and then the rate  
15 of elimination via exhalation decreases. Following oral or inhalation exposure, urinary  
16 elimination of parent TCE is minimal, with urinary elimination of the metabolites trichloroacetic  
17 acid and trichloroethanol accounting for the bulk of the absorbed dose of TCE. See Section 3.4  
18 for additional discussion of TCE excretion.

19 As part of this assessment, a comprehensive Bayesian PBPK model-based analysis of the  
20 population toxicokinetics of TCE and its metabolites was developed in mice, rats, and humans  
21 (also reported in Chiu et al., 2009). This analysis considered a wider range of physiological,  
22 chemical, in vitro, and in vivo data than any previously published analysis of TCE. The  
23 toxicokinetics of the “population average,” its population variability, and their uncertainties are  
24 characterized and estimates of experimental variability and uncertainty are included in this  
25 analysis. The experimental database included separate sets for model calibration and evaluation  
26 for rats and humans; fewer data were available in mice, and were all used for model calibration.  
27 The total combination of these approaches and PBPK analysis substantially supports the model  
28 predictions. In addition, the approach employed yields an accurate characterization of the  
29 uncertainty in metabolic pathways for which available data were sparse or relatively indirect,  
30 such as GSH conjugation and respiratory tract metabolism. Key conclusions from the model  
31 predictions include (1) as expected, TCE is substantially metabolized, primarily by oxidation at  
32 doses below saturation; (2) GSH conjugation and subsequent bioactivation in humans appears to  
33 be 10- to 100-fold greater than previously estimated; and (3) mice had the greatest rate of  
34 respiratory tract oxidative metabolism compared to rats and humans. However, there are  
35 uncertainties as to the accuracy of the analytical method used for some of the available in vivo

1 data on GSH conjugation. Because these data are highly influential, the PBPK modeling results  
2 for the flux of GSH conjugation should be interpreted with caution. Thus, there is lower  
3 confidence in the accuracy of GSH conjugation predictions as compared to other dose-metrics,  
4 such as those related to the parent compound, total metabolism, or oxidative metabolites. The  
5 predictions of the PBPK model are subsequently used in noncancer and cancer dose-response  
6 analyses for inter and intraspecies extrapolation of toxicokinetics (see Section 6.2, below). See  
7 Section 3.5 and Appendix A for additional discussion of and details about PBPK modeling of  
8 TCE and metabolites.

9

### 6.1.3. Noncancer Toxicity

10 This section summarizes the weight of evidence for TCE noncancer toxicity. Based on the  
11 available human epidemiologic data and experimental and mechanistic studies, it is concluded  
12 that TCE poses a potential human health hazard for noncancer toxicity to the central nervous  
13 system, the kidney, the liver, the immune system, the male reproductive system, and the  
14 developing fetus. The evidence is more limited for TCE toxicity to the respiratory tract and  
15 female reproductive system. The conclusions pertaining to specific endpoints within these  
16 tissues and systems are summarized below.

17

#### 6.1.3.1.1. Neurological Effects (see Sections 4.3 and 4.11.1.1 and Appendix D)

18 Both human and animal studies have associated TCE exposure with effects on several  
19 neurological domains. Multiple epidemiologic studies in different populations have reported  
20 abnormalities in trigeminal nerve function in association with TCE exposure. Two small studies  
21 did not report an association between TCE exposure and trigeminal nerve function. However,  
22 statistical power was limited, exposure misclassification was possible, and, in one case, methods  
23 for assessing trigeminal nerve function were not available. As a result, these studies do not  
24 provide substantial evidence against a causal relationship between TCE exposure and trigeminal  
25 nerve impairment. Laboratory animal studies have also demonstrated TCE-induced changes in  
26 the morphology of the trigeminal nerve following short-term exposures in rats. However, one  
27 study reported no significant changes in trigeminal somatosensory evoked potential in rats  
28 exposed to TCE for 13 weeks. See Section 4.3.1 for additional discussion of studies of  
29 alterations in nerve conduction and trigeminal nerve effects. Human chamber, occupational, and  
30 geographic based/drinking water studies have consistently reported subjective symptoms such as  
31 headaches, dizziness, and nausea which are suggestive of vestibular system impairments. One

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 study reported changes in nystagmus threshold (a measure of vestibular system function)  
2 following an acute TCE exposure. There are only a few laboratory animal studies relevant to  
3 this neurological domain, with reports of changes in nystagmus, balance, and handling reactivity.  
4 See Section 4.3.3 for additional discussion of TCE effects on vestibular function. Fewer and  
5 more limited epidemiologic studies are suggestive of TCE exposure being associated with  
6 delayed motor function, and changes in auditory, visual, and cognitive function or performance  
7 (see Sections 4.3.2, 4.3.4, 4.3.5, and 4.3.6). Acute and subchronic animal studies show  
8 disruption of the auditory system, changes in visual evoked responses to patterns or flash  
9 stimulus, and neurochemical and molecular changes. Animal studies suggest that while the  
10 effects on the auditory system lead to permanent function impairments and histopathology,  
11 effects on the visual system may be reversible with termination of exposure. Additional acute  
12 studies reported structural or functional changes in hippocampus, such as decreased myelination  
13 or decreased excitability of hippocampal CA1 neurons, although the relationship of these effects  
14 to overall cognitive function is not established (see Section 4.3.9). An association between TCE  
15 exposure and sleep changes has also been demonstrated in rats (see Section 4.3.7). Some  
16 evidence exists for motor-related changes in rats/mice exposed acutely/subchronically to TCE,  
17 but these effects have not been reported consistently across all studies (see Section 4.3.6).  
18 Gestational exposure to TCE in humans has been reported to be associated with  
19 neurodevelopmental abnormalities including neural tube defects, encephalopathy, impaired  
20 cognition, aggressive behavior, and speech and hearing impairment. Developmental  
21 neurotoxicological changes have also been observed in animals including aggressive behaviors  
22 following an in utero exposure to TCE and a suggestion of impaired cognition as noted by  
23 decreased myelination in the CA1 hippocampal region of the brain. See Section 4.3.8 for  
24 additional discussion of developmental neurological effects of TCE. Therefore, overall, the  
25 strongest neurological evidence of human toxicological hazard is for changes in trigeminal nerve  
26 function or morphology and impairment of vestibular function, based on both human and  
27 experimental studies, while fewer and more limited evidence exists for delayed motor function,  
28 changes in auditory, visual, and cognitive function or performance, and neurodevelopmental  
29 outcomes.

30

#### **6.1.3.1.2. Kidney Effects (see Sections 4.4.1, 4.4.4, 4.4.6, and 4.11.1.2)**

31 Kidney toxicity has also been associated with TCE exposure in both human and animal  
32 studies. There are few human data pertaining to TCE-related noncancer kidney toxicity;  
33 however, several available studies reported elevated excretion of urinary proteins, considered

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 nonspecific markers of nephrotoxicity, among TCE-exposed subjects compared to unexposed  
2 controls. While some of these studies include subjects previously diagnosed with kidney cancer,  
3 other studies report similar results in subjects that are disease free. Some additional support for  
4 TCE nephrotoxicity in humans is provided by two studies of end-stage renal disease; a study  
5 reporting a greater incidence of end-stage renal disease in TCE-exposed workers as compared to  
6 unexposed controls and a second study reporting a greater risk for progression from IgA or  
7 membranous nephropathy glomerulonephritis to end-stage renal disease and TCE-exposure. See  
8 Section 4.4.1 for additional discussion of human data on the noncancer kidney effects of TCE.  
9 Laboratory animal and in vitro data provide additional support for TCE nephrotoxicity. TCE  
10 causes renal toxicity in the form of cytomegaly and karyomegaly of the renal tubules in male and  
11 female rats and mice following either oral or inhalation exposure. In rats, the pathology of TCE-  
12 induced nephrotoxicity appears distinct from age-related nephropathy. Increased kidney weights  
13 have also been reported in some rodent studies. See Section 4.4.4 for additional discussion of  
14 laboratory animal data on the noncancer kidney effects of TCE. Further studies with TCE  
15 metabolites have demonstrated a potential role for dichlorovinyl cysteine (DCVC),  
16 trichloroethanol, and trichloroacetic acid (TCA) in TCE-induced nephrotoxicity. Of these,  
17 available data suggest that DCVC induced renal effects are most similar to those of TCE and that  
18 DCVC is formed in sufficient amounts following TCE exposure to account for these effects.  
19 TCE or DCVC have also been shown to be cytotoxic to primary cultures of rat and human renal  
20 tubular cells. See Section 4.4.6 for additional discussion on the role of metabolism in the  
21 noncancer kidney effects of TCE. Overall, multiple lines of evidence support the conclusion that  
22 TCE causes nephrotoxicity in the form of tubular toxicity, mediated predominantly through the  
23 TCE GSH conjugation product DCVC.

24

#### **6.1.3.1.3. Liver Effects (see Sections 4.5.1, 4.5.3, 4.5.4, 4.5.6, and 4.11.1.3, and Appendix E)**

25 Liver toxicity has also been associated with TCE exposure in both human and animal  
26 studies. Although there are few human studies on liver toxicity and TCE exposure, several  
27 available studies have reported TCE exposure to be associated with significant changes in serum  
28 liver function tests, widely used in clinical settings in part to identify patients with liver disease,  
29 or changes in plasma or serum bile acids. Additional, more limited human evidence for TCE  
30 induced liver toxicity includes reports suggesting an association between TCE exposure and liver  
31 disorders, and case reports of liver toxicity including hepatitis accompanying immune-related  
32 generalized skin diseases, jaundice, hepatomegaly, hepatosplenomegaly, and liver failure in  
33 TCE-exposed workers. Cohort studies examining cirrhosis mortality and either TCE exposure or

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 solvent exposure are generally null, but these studies cannot rule out an association with TCE  
2 because of their use of death certificates where there is a high degree (up to 50%) of  
3 underreporting. Overall, while some evidence exists of liver toxicity as assessed from liver  
4 function tests, the data are inadequate for making conclusions regarding causality. See  
5 Section 4.5.1 for additional discussion of human data on the noncancer liver effects of TCE. In  
6 rats and mice, TCE exposure causes hepatomegaly without concurrent cytotoxicity. Like  
7 humans, laboratory animals exposed to TCE have been observed to have increased serum bile  
8 acids, although the toxicological importance of this effect is unclear. Other effects in the rodent  
9 liver include small transient increases in DNA synthesis, cytomegaly in the form of “swollen” or  
10 enlarged hepatocytes, increased nuclear size probably reflecting polyploidization, and  
11 proliferation of peroxisomes. Available data also suggest that TCE does not induce substantial  
12 cytotoxicity, necrosis, or regenerative hyperplasia, as only isolated, focal necroses and mild to  
13 moderate changes in serum and liver enzyme toxicity markers having been reported. These  
14 effects are consistently observed across rodent species and strains, although the degree of  
15 response at a given mg/kg-day dose appears to be highly variable across strains, with mice on  
16 average appearing to be more sensitive. See Sections 4.5.3 and 4.5.4 for additional discussion of  
17 laboratory animal data on the noncancer liver effects of TCE. While it is likely that oxidative  
18 metabolism is necessary for TCE-induced effects in the liver, the specific metabolite or  
19 metabolites responsible is less clear. However, the available data are strongly inconsistent with  
20 TCA being the sole or predominant active moiety for TCE-induced liver effects, particularly  
21 with respect to hepatomegaly. See Section 4.5.6 for additional discussion on the role of  
22 metabolism in the noncancer liver effects of TCE. Overall, TCE, likely through its oxidative  
23 metabolites, clearly leads to liver toxicity in laboratory animals, with mice appearing to be more  
24 sensitive than other laboratory animal species, but there is only limited epidemiologic evidence  
25 of hepatotoxicity being associated with TCE exposure.

26

#### **6.1.3.1.4. Immunological Effects (see Sections 4.6.1.1, 4.6.2, and 4.11.1.4)**

27 Effects related the immune system have also been associated with TCE exposure in both  
28 human and animal studies. A relationship between systemic autoimmune diseases, such as  
29 scleroderma, and occupational exposure to TCE has been reported in several recent studies, and a  
30 meta-analysis of scleroderma studies resulted in a statistically significant combined odds ratio for  
31 any exposure in men (odds ratio [OR]: 2.5, 95% confidence interval [CI]: 1.1, 5.4), with a lower  
32 relative risk seen in women (OR: 1.2, 95% CI: 0.58, 2.6). The human data at this time do not  
33 allow a determination of whether the difference in effect estimates between men and women

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 reflects the relatively low background risk of scleroderma in men, gender-related differences in  
2 exposure prevalence or in the reliability of exposure assessment, a gender-related difference in  
3 susceptibility to the effects of TCE, or chance. Additional human evidence for the  
4 immunological effects of TCE includes studies reporting TCE-associated changes in levels of  
5 inflammatory cytokines in occupationally-exposed workers and infants exposed via indoor air at  
6 air concentrations typical of such exposure scenarios (see Section 6.1.1, above); a large number  
7 of case reports (mentioned above) of a severe hypersensitivity skin disorder, distinct from  
8 contact dermatitis and often accompanied by hepatitis; and a reported association between  
9 increased history of infections and exposure to TCE contaminated drinking water. See  
10 Section 4.6.1.1 for additional discussion of human data on the immunological effects of TCE.  
11 Immunotoxicity has also been reported in experimental rodent studies of TCE. Numerous  
12 studies have demonstrated accelerated autoimmune responses in autoimmune-prone mice,  
13 including changes in cytokine levels similar to those reported in human studies, with more severe  
14 effects, including autoimmune hepatitis, inflammatory skin lesions, and alopecia, manifesting at  
15 longer exposure periods. Immunotoxic effects have been also reported in B6C3F1 mice, which  
16 do not have a known particular susceptibility to autoimmune disease. Developmental  
17 immunotoxicity in the form of hypersensitivity responses have been reported in TCE-treated  
18 guinea pigs and mice via drinking water pre- and postnatally. Evidence of localized  
19 immunosuppression has also been reported in mice and rats. See Section 4.6.2 for additional  
20 discussion of laboratory animal data on the immunological effects of TCE. Overall, the human  
21 and animal studies of TCE and immune-related effects provide strong evidence for a role of TCE  
22 in autoimmune disease and in a specific type of generalized hypersensitivity syndrome, while  
23 there are less data pertaining to immunosuppressive effects.  
24

#### **6.1.3.1.5. Respiratory Tract Effects (see Sections 4.7.1.1, 4.7.2.1, 4.7.3, and 4.11.1.5)**

25 The very few human data on TCE and pulmonary toxicity are too limited for drawing  
26 conclusions (see Section 4.7.1.1), but laboratory studies in mice and rats have shown toxicity in  
27 the bronchial epithelium, primarily in Clara cells, following acute exposures to TCE (see  
28 Section 4.7.2.1). A few studies of longer duration have reported more generalized toxicity, such  
29 as pulmonary fibrosis in mice and pulmonary vasculitis in rats. However, respiratory tract  
30 effects were not reported in other longer-term studies. Acute pulmonary toxicity appears to be  
31 dependent on oxidative metabolism, although the particular active moiety is not known. While  
32 earlier studies implicated chloral produced in situ by CYP enzymes in respiratory tract tissue in  
33 toxicity, the evidence is inconsistent and several other possibilities are viable. Although humans

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 appear to have lower overall capacity for enzymatic oxidation in the lung relative to mice, CYP  
2 enzymes do reside in human respiratory tract tissue, suggesting that, qualitatively, the respiratory  
3 tract toxicity observed in rodents is biologically plausible in humans. See Section 4.7.3 for  
4 additional discussion of the role of metabolism in the noncancer respiratory tract toxicity of  
5 TCE. Therefore, overall, data are suggestive of TCE causing respiratory tract toxicity, based  
6 primarily on short-term studies in mice and rats, with available human data too few and limited  
7 to add to the weight of evidence for pulmonary toxicity.  
8

#### **6.1.3.1.6. Reproductive Effects (see Sections 4.8.1 and 4.11.1.6)**

9 A number of human and laboratory animal studies suggest that TCE exposure has the  
10 potential for male reproductive toxicity, with a more limited number of studies examining female  
11 reproductive toxicity. Human studies have reported TCE exposure to be associated (in all but  
12 one case statistically-significantly) with increased sperm density and decreased sperm quality,  
13 altered sexual drive or function, or altered serum endocrine levels. Measures of male fertility,  
14 however, were either not reported or reported to be unchanged with TCE exposure, though the  
15 statistical power of the available studies is quite limited. Epidemiologic studies have identified  
16 possible associations of TCE exposure with effects on female fertility and with menstrual cycle  
17 disturbances, but these data are fewer than those available for male reproductive toxicity. See  
18 Section 4.8.1.1 for additional discussion of human data on the reproductive effects of TCE.  
19 Evidence of similar effects, particularly for male reproductive toxicity, is provided by several  
20 laboratory animal studies that reported effects on sperm, libido/copulatory behavior, and serum  
21 hormone levels, although some studies that assessed sperm measures did not report treatment-  
22 related alterations. Additional adverse effects on male reproduction have also been reported,  
23 including histopathological lesions in the testes or epididymides and altered in vitro sperm-  
24 oocyte binding or in vivo fertilization due to TCE or metabolites. While reduced fertility in  
25 rodents was only observed in one study, this is not surprising given the redundancy and  
26 efficiency of rodent reproductive capabilities. In addition, although the reduced fertility  
27 observed in the rodent study was originally attributed to systemic toxicity, the database as a  
28 whole suggests that TCE does induce reproductive toxicity independent of systemic effects.  
29 Fewer data are available in rodents on female reproductive toxicity. While in vitro oocyte  
30 fertilizability has been reported to be reduced as a result of TCE exposure in rats, a number of  
31 other laboratory animal studies did not report adverse effects on female reproductive function.  
32 See Section 4.8.1.2 for additional discussion of laboratory animal data on the reproductive  
33 effects of TCE. Very limited data are available to elucidate the mode of action (MOA) for these

1 effects, though some aspects of a putative MOA (e.g., perturbations in testosterone biosynthesis)  
2 appear to have some commonalities between humans and animals (see Section 4.8.1.3.2).  
3 Together, the human and laboratory animal data support the conclusion that TCE exposure poses  
4 a potential hazard to the male reproductive system, but are more limited with regard to the  
5 potential hazard to the female reproductive system.  
6

#### **6.1.3.1.7. Developmental Effects (see Sections 4.8.3 and 4.11.1.7)**

7 The relationship between TCE exposure (direct or parental) and developmental toxicity  
8 has been investigated in a number of epidemiologic and laboratory animal studies. Postnatal  
9 developmental outcomes examined include developmental neurotoxicity (addressed above with  
10 neurotoxicity), developmental immunotoxicity (addressed above with immunotoxicity), and  
11 childhood cancers. Prenatal effects examined include death (spontaneous abortion, perinatal  
12 death, pre- or postimplantation loss, resorptions), decreased growth (low birth weight, small for  
13 gestational age, intrauterine growth restriction, decreased postnatal growth), and congenital  
14 malformations, in particular cardiac defects. Some epidemiological studies have reported  
15 associations between parental exposure to TCE and spontaneous abortion or perinatal death, and  
16 decreased birth weight or small for gestational age, although other studies reported mixed or null  
17 findings. While comprising both occupational and environmental exposures, these studies are  
18 overall not highly informative due to the small numbers of cases and limited exposure  
19 characterization or to the fact that exposures were to a mixture of solvents. See Section 4.8.3.1  
20 for additional discussion of human data on the developmental effects of TCE. However,  
21 multiple well conducted studies in rats and mice show analogous effects of TCE exposure: pre-  
22 or postimplantation losses, increased resorptions, perinatal death, and decreased birth weight.  
23 Interestingly, the rat studies reporting these effects used Fischer 344 or Wistar rats, while several  
24 other studies, all of which used Sprague-Dawley rats, reported no increased risk in these  
25 developmental measures, suggesting a strain difference in susceptibility. See Section 4.8.3.2 for  
26 additional discussion of laboratory animal data on the developmental effects of TCE. Therefore,  
27 overall, based on weakly suggestive epidemiologic data and fairly consistent laboratory animal  
28 data, it can be concluded that TCE exposure poses a potential hazard for prenatal losses and  
29 decreased growth or birth weight of offspring.

30 With respect to congenital malformations, epidemiology and experimental animal studies  
31 of TCE have reported increases in total birth defects, central nervous system defects, oral cleft  
32 defects, eye/ear defects, kidney/urinary tract disorders, musculoskeletal birth anomalies,  
33 lung/respiratory tract disorders, skeletal defects, and cardiac defects. Human occupational cohort

1 studies, while not consistently reporting positive results, are generally limited by the small  
2 number of observed or expected cases of birth defects. While only one of the epidemiological  
3 studies specifically reported observations of eye anomalies, studies in rats have identified  
4 increases in the incidence of fetal eye defects following oral exposures during the period of  
5 organogenesis with TCE or its oxidative metabolites dichloroacetic acid (DCA) and TCA. The  
6 epidemiological studies, while individually limited, as a whole show relatively consistent  
7 elevations, some of which were statistically significant, in the incidence of cardiac defects in  
8 TCE-exposed populations compared to reference groups. In laboratory animal models, avian  
9 studies were the first to identify adverse effects of TCE exposure on cardiac development, and  
10 the initial findings have been confirmed multiple times. Additionally, administration of TCE and  
11 its metabolites TCA and DCA in maternal drinking water during gestation has been reported to  
12 induce cardiac malformations in rat fetuses. It is notable that a number of other studies, several  
13 of which were well-conducted, did not report induction of cardiac defects in rats, mice, or rabbits  
14 in which TCE was administered by inhalation or gavage. However, many of these studies used a  
15 traditional free-hand section technique on fixed fetal specimens, and a fresh dissection technique  
16 that can enhance detection of anomalies was used in the positive studies by Dawson et al. (1993)  
17 and Johnson et al. (2003)(2005). Nonetheless, two studies that used the same or similar fresh  
18 dissection technique did not report cardiac anomalies. Differences in other aspects of  
19 experimental design may have been contributing factors to the differences in observed response.  
20 In addition, mechanistic studies, such as the treatment-related alterations in endothelial cushion  
21 development observed in avian in ovo and in vitro studies, provide a plausible mechanistic basis  
22 for defects in septal and valvular morphogenesis observed in rodents, and consequently support  
23 the plausibility of cardiac defects induced by TCE in humans. Therefore, while the studies by  
24 Dawson et al. (1993) and Johnson et al. (2003)(2005) have significant limitations, including the  
25 lack of clear dose-response relationship for the incidence of any specific cardiac anomaly and the  
26 pooling of data collected over an extended period, there is insufficient reason to dismiss their  
27 findings. See Section 4.8.3.3.2 for additional discussion of the conclusions with respect to  
28 TCE-induced cardiac malformations. Therefore, overall, based on weakly suggestive, but  
29 overall consistent, epidemiologic data, in combination with evidence from experimental animal  
30 and mechanistic studies, it can be concluded that TCE exposure poses a potential hazard for  
31 congenital malformations, including cardiac defects, in offspring.

32

**6.1.4. Carcinogenicity (see Sections 4.1, 4.2, 4.4.2, 4.4.5, 4.4.7, 4.5.2, 4.5.5, 4.5.6, 4.5.7, 4.6.1.2, 4.6.2.4, 4.7.1.2, 4.7.2.2, 4.7.4, 4.8.2, 4.9, and 4.11.2, and Appendices B and C)**

1 Following EPA (2005c) *Guidelines for Carcinogen Risk Assessment*, based on the  
2 available data as of 2010, TCE is characterized as “*carcinogenic to humans*” by all routes of  
3 exposure. This conclusion is based on convincing evidence of a causal association between TCE  
4 exposure in humans and kidney cancer. The consistency of increased kidney cancer relative risk  
5 estimates across a large number of independent studies of different designs and populations from  
6 different countries and industries provides compelling evidence given the difficulty, a priori, in  
7 detecting effects in epidemiologic studies when the relative risks are modest, the cancers are  
8 relatively rare, and therefore, individual studies have limited statistical power. This strong  
9 consistency of the epidemiologic data on TCE and kidney cancer argues against chance, bias,  
10 and confounding as explanations for the elevated kidney cancer risks. In addition, statistically  
11 significant exposure-response trends are observed in high-quality studies. These studies were  
12 conducted in populations with high TCE exposure intensity or had the ability to identify  
13 TCE-exposed subjects with high confidence. These studies addressed important potential  
14 confounders and biases, further supporting the observed associations with kidney cancer as  
15 causal. See Section 4.4.2 for additional discussion of the human epidemiologic data on TCE  
16 exposure and kidney cancer. In a meta-analysis of 15 studies with high exposure potential, a  
17 statistically significant summary relative risk estimate was observed for overall TCE exposure  
18 (RRm: 1.27 [95% CI: 1.13, 1.43]). The summary relative risk estimate was greater for the  
19 highest TCE exposure groups (RRm: 1.58 [95% CI: 1.28, 1.96];  $n = 13$  studies). Meta-analyses  
20 investigating the influence of individual studies and the sensitivity of the results to alternate  
21 relative risk estimate selections found the summary relative risk estimates to be highly robust.  
22 Furthermore, there was no indication of publication bias or significant heterogeneity across the  
23 15 studies. It would require a substantial amount of negative data from informative studies (i.e.,  
24 studies having a high likelihood of TCE exposure in individual study subjects and which meet, to  
25 a sufficient degree, the standards of epidemiologic design and analysis in a systematic review) to  
26 contradict this observed association. See Section 4.4.2.5 and Appendix C for additional  
27 discussion of the kidney cancer meta-analysis.

28 The human evidence of carcinogenicity from epidemiologic studies of TCE exposure is  
29 compelling for non-Hodgkin lymphoma (NHL) but less convincing than for kidney cancer. High  
30 quality studies generally reported excess relative risk estimates, with statistically significant  
31 increases in three studies with overall TCE exposure, and a statistically significant increase in the  
32 high TCE exposure group and statistically significant trend in a fourth study (see  
33 Section 4.6.1.2). The consistency of the association between TCE exposure and NHL is further  
34 supported by the results of meta-analyses (see Section 4.6.1.2.2 and Appendix C). A statistically

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 significant summary relative risk estimate was observed for overall TCE exposure (RRm: 1.23  
2 [95% CI: 1.07, 1.42];  $n = 17$  studies), and, as with kidney cancer, the summary relative risk  
3 estimate was greater for the highest TCE exposure groups (RRm: 1.43 [95% CI: 1.13, 1.82];  $n =$   
4 13 studies) than for overall TCE exposure. Sensitivity analyses indicated that these results and  
5 their statistical significance were not overly influenced by any single study or choice of  
6 individual (study-specific) risk estimates, and in all of the influence and sensitivity analyses, the  
7 RRm estimate was statistically significantly increased. Some heterogeneity was observed,  
8 particularly between cohort and case-control studies, but it was not statistically significant. In  
9 addition, there was some evidence of potential publication bias. Thus, while the evidence is  
10 strong for NHL, issues of study heterogeneity, potential publication bias, and weaker exposure-  
11 response results contribute greater uncertainty.

12 The evidence is more limited for liver and biliary tract cancer mainly because only cohort studies  
13 are available and most of these studies have small numbers of cases due the comparative rarity of  
14 liver and biliary tract cancer. While most high quality studies reported excess relative risk  
15 estimates, they were generally based on small numbers of cases or deaths, with the result of wide  
16 confidence intervals on the estimates. The low number of liver cancer cases in the available  
17 studies made assessing exposure-response relationships difficult. See Section 4.5.2 for  
18 additional discussion of the human epidemiologic data on TCE exposure and liver cancer. A  
19 consistency of the association between TCE exposure and liver cancer is supported by the results  
20 of meta-analyses (see Section 4.5.2 and Appendix C). These meta-analyses found a statistically  
21 significant increased summary relative risk estimate for liver and biliary tract cancer of 1.29  
22 (95% CI: 1.07, 1.56;  $n = 9$  studies) with overall TCE exposure; but the meta-analyses using only  
23 the highest exposure groups yielded a lower, and nonstatistically significant, summary estimate  
24 for primary liver cancer (1.28 [95% CI: 0.93, 1.77],  $n = 8$  studies). Although there was no  
25 evidence of heterogeneity or publication bias and the summary estimates were fairly insensitive  
26 to the use of alternative relative risk estimates, the statistical significance of the summary  
27 estimates depends heavily on the one large study by Raaschou-Nielsen et al. (2003). There were  
28 fewer adequate studies with high exposure potential available for meta-analysis of liver cancer  
29 (9 versus 17 for NHL and 15 for kidney), leading to lower statistical power, even with pooling.  
30 Thus, while there is epidemiologic evidence of an association between TCE exposure and liver  
31 cancer, the much more limited database, both in terms of number of available studies and  
32 number of cases upon which the studies are based, contributes to greater uncertainty as compared  
33 to the evidence for kidney cancer or lymphoma.

34 In addition to the body of evidence pertaining to kidney cancer, NHL, and liver cancer, the  
35 available epidemiologic studies also provide more limited evidence of an association between

1 TCE exposure and other types of cancer, including bladder, esophageal, prostate, cervical, breast,  
2 and childhood leukemia. Differences between these sets of data and the data for kidney cancer,  
3 NHL, and liver cancer are observations from fewer numbers of studies, a mixed pattern of  
4 observed risk estimates, and the general absence of exposure-response data from the studies  
5 using a quantitative TCE-specific exposure measure.

6 There are several other lines of supporting evidence for TCE carcinogenicity in humans by all  
7 routes of exposure. First, multiple chronic bioassays in rats and mice have reported increased  
8 incidences of tumors with TCE treatment via inhalation and oral gavage, including tumors in the  
9 kidney, liver, and lymphoid tissues-target tissues of TCE carcinogenicity also seen in  
10 epidemiological studies. Of particular note is the site-concordant finding of low, but biologically  
11 and sometimes statistically significant, increases in the incidence of kidney tumors in multiple  
12 strains of rats treated with TCE by either inhalation or corn oil gavage (see Section 4.4.5). The  
13 increased incidences were only detected at the highest tested doses, and were greater in male  
14 than female rats; although, notably, pooled incidences in females from five rat strains tested by  
15 NTP (1988, 1990) resulted in a statistically significant trend. Although these studies have shown  
16 limited increases in kidney tumors, and several individual studies have a number of limitations,  
17 given the rarity of these tumors as assessed by historical controls and the repeatability of this  
18 result across studies and strains, these are considered biologically significant. Therefore, while  
19 individual studies provide only suggestive evidence of renal carcinogenicity, the database as a  
20 whole supports the conclusion that TCE is a kidney carcinogen in rats, with males being more  
21 sensitive than females. No other tested laboratory species (i.e., mice and hamsters) have  
22 exhibited increased kidney tumors, with no adequate explanation for these species differences  
23 (particularly with mice, which have been extensively tested). With respect to the liver, TCE and  
24 its oxidative metabolites chloral hydrate (CH), TCA, and DCA are clearly carcinogenic in mice,  
25 with strain and sex differences in potency that appear to parallel, qualitatively, differences in  
26 background tumor incidence. Data in other laboratory animal species are limited; thus, except  
27 for DCA which is carcinogenic in rats, inadequate evidence exists to evaluate the  
28 hepatocarcinogenicity of these compounds in rats or hamsters. However, to the extent that there  
29 is hepatocarcinogenic potential in rats, TCE is clearly less potent in the strains tested in this  
30 species than in B6C3F1 and Swiss mice. See Section 4.5.5 for additional discussion of  
31 laboratory animal data on TCE-induced liver tumors. Additionally, there is more limited  
32 evidence for TCE-induced lymphatic cancers in rats and mice, lung tumors in mice, and  
33 testicular tumors in rats. With respect to the lymphatic cancers, two studies in mice reported  
34 increased incidences of lymphomas in females of two different strains, and two studies in rats  
35 reported leukemias in males of one strain and females of another. However, these tumors had

1 relatively modest increases in incidence with treatment, and were not reported to be increased in  
2 other studies. See Section 4.6.2.4 for additional discussion of laboratory animal data on  
3 TCE-induced lymphatic tumors. With respect to lung tumors, rodent bioassays have  
4 demonstrated a statistically significant increase in pulmonary tumors in mice following chronic  
5 inhalation exposure to TCE, and nonstatistically significant increases in mice exposed orally; but  
6 pulmonary tumors were not reported in other species tested (i.e., rats and hamsters) (see  
7 Section 4.7.2.2). Finally, increased testicular (interstitial or Leydig cell) tumors have been  
8 observed in multiple studies of rats exposed by inhalation and gavage, although in some cases  
9 high (> 75%) control rates of testicular tumors in rats limited the ability to detect a treatment  
10 effect. See Section 4.8.2.2 for additional discussion of laboratory animal data on TCE-induced  
11 tumors of the reproductive system. Overall, TCE is clearly carcinogenic in rats and mice. The  
12 apparent lack of site concordance across laboratory animal studies may be due to limitations in  
13 design or conduct in a number of rat bioassays and/or genuine interspecies differences in  
14 qualitative or quantitative sensitivity (i.e., potency). Nonetheless, these studies have shown  
15 carcinogenic effects across different strains, sexes, and routes of exposure, and site-concordance  
16 is not necessarily expected for carcinogens. Of greater import is the finding that there is site-  
17 concordance between the main cancers observed in TCE-exposed humans and those observed in  
18 rodent studies—in particular, cancers of the kidney, liver, and lymphoid tissues.

19 A second line of supporting evidence for TCE carcinogenicity in humans consists of  
20 toxicokinetic data indicating that TCE is well absorbed by all routes of exposure, and that TCE  
21 absorption, distribution, metabolism, and excretion are qualitatively similar in humans and  
22 rodents. As summarized above, there is evidence that TCE is systemically available, distributes  
23 to organs and tissues, and undergoes systemic metabolism from all routes of exposure.

24 Therefore, although the strongest evidence from epidemiologic studies largely involves  
25 inhalation exposures, the evidence supports TCE carcinogenicity being applicable to all routes of  
26 exposure. In addition, there is no evidence of major qualitative differences across species in  
27 TCE absorption, distribution, metabolism, and excretion. Extensive in vivo and in vitro data  
28 show that mice, rats, and humans all metabolize TCE via two primary pathways: oxidation by  
29 CYPs and conjugation with glutathione via GSTs. Several metabolites and excretion products  
30 from both pathways have been detected in blood and urine from exposed humans as well as from  
31 at least one rodent species. In addition, the subsequent distribution, metabolism, and excretion of  
32 TCE metabolites are qualitatively similar among species. Therefore, humans possess the  
33 metabolic pathways that produce the TCE metabolites thought to be involved in the induction of  
34 rat kidney and mouse liver tumors, and internal target tissues of both humans and rodents  
35 experience a similar mix of TCE and metabolites. See Sections 3.1–3.4 for additional discussion

1 of TCE toxicokinetics. Quantitative interspecies differences in toxicokinetics do exist, and are  
2 addressed through PBPK modeling (see Section 3.5 and Appendix A). Importantly, these  
3 quantitative differences affect only interspecies extrapolations of carcinogenic potency, and do  
4 not affect inferences as to the carcinogenic hazard for TCE.

5 Finally, available mechanistic data do not suggest a lack of human carcinogenic hazard from  
6 TCE exposure. In particular, these data do not suggest qualitative differences between humans  
7 and test animals that would preclude any of the hypothesized key events in the carcinogenic  
8 MOA in rodents from occurring in humans. For the kidney, the predominance of positive  
9 genotoxicity data in the database of available studies of TCE metabolites derived from GSH  
10 conjugation (in particular DCVC), together with toxicokinetic data consistent with their systemic  
11 delivery to and in situ formation in the kidney, supports the conclusion that a mutagenic MOA is  
12 operative in TCE-induced kidney tumors. While supporting the biological plausibility of this  
13 hypothesized MOA, available data on the von Hippel-Lindau (VHL) gene in humans or  
14 transgenic animals do not conclusively elucidate the role of VHL mutation in TCE-induced renal  
15 carcinogenesis. Cytotoxicity and compensatory cell proliferation, similarly presumed to be  
16 mediated through metabolites formed after GSH-conjugation of TCE, have also been suggested  
17 to play a role in the MOA for renal carcinogenesis, as high incidences of nephrotoxicity have  
18 been observed in animals at doses that induce kidney tumors. Human studies have reported  
19 markers for nephrotoxicity at current occupational exposures, although data are lacking at lower  
20 exposures. Nephrotoxicity is observed in both mice and rats, in some cases with nearly  
21 100% incidence in all dose groups, but kidney tumors are only observed at low incidences in rats  
22 at the highest tested doses. Therefore, nephrotoxicity alone appears to be insufficient, or at least  
23 not rate-limiting, for rodent renal carcinogenesis, since maximal levels of toxicity are reached  
24 before the onset of tumors. In addition, nephrotoxicity has not been shown to be necessary for  
25 kidney tumor induction by TCE in rodents. In particular, there is a lack of experimental support  
26 for causal links, such as compensatory cellular proliferation or clonal expansion of initiated cells,  
27 between nephrotoxicity and kidney tumors induced by TCE. Furthermore, it is not clear if  
28 nephrotoxicity is one of several key events in a MOA, if it is a marker for an “upstream” key  
29 event (such as oxidative stress) that may contribute independently to both nephrotoxicity and  
30 renal carcinogenesis, or if it is incidental to kidney tumor induction. Moreover, while  
31 toxicokinetic differences in the GSH conjugation pathway along with their uncertainty are  
32 addressed through PBPK modeling, no data suggest that any of the proposed key events for  
33 TCE-induced kidney tumors in rats are precluded in humans. See Section 4.4.7 for additional  
34 discussion of the MOA for TCE-induced kidney tumors. Therefore, TCE-induced rat kidney



1 tumors provide additional support for the convincing human evidence of TCE-induced kidney  
2 cancer, with mechanistic data supportive of a mutagenic MOA.  
3 With respect to other tumor sites, data are insufficient to conclude that any of the other  
4 hypothesized MOAs are operant. In the liver, a mutagenic MOA mediated by CH, which has  
5 evidence for genotoxic effects, or some other oxidative metabolite of TCE cannot be ruled out,  
6 but data are insufficient to conclude it is operant. A second MOA hypothesis for TCE-induced  
7 liver tumors involves activation of the peroxisome proliferator activated receptor alpha (PPAR $\alpha$ )  
8 receptor. Clearly, in vivo administration of TCE leads to activation of PPAR $\alpha$  in rodents and  
9 likely does so in humans as well. However, the evidence as a whole does not support the view  
10 that PPAR $\alpha$  is the sole operant MOA mediating TCE hepatocarcinogenesis. Rather, there is  
11 evidential support for multiple TCE metabolites and multiple toxicity pathways contributing to  
12 TCE-induced liver tumors. Furthermore, recent experiments have demonstrated that PPAR $\alpha$   
13 activation and the sequence of key events in the hypothesized MOA are not sufficient to induce  
14 hepatocarcinogenesis (Yang et al., 2007). Moreover, the demonstration that the PPAR $\alpha$  agonist  
15 di(2-ethylhexyl) phthalate induces tumors in PPAR $\alpha$ -null mice supports the view that the events  
16 comprising the hypothesized PPAR $\alpha$  activation MOA are not necessary for liver tumor induction  
17 in mice by this PPAR $\alpha$  agonist (Ito et al., 2007). See Section 4.5.7 for additional discussion of  
18 the MOA for TCE-induced liver tumors. For mouse lung tumors, as with the liver, a mutagenic  
19 MOA involving CH has also been hypothesized, but there are insufficient data to conclude that it  
20 is operant. A second MOA hypothesis for mouse lung tumors has been posited involving other  
21 effects of oxidative metabolites including cytotoxicity and regenerative cell proliferation, but  
22 experimental support remains limited, with no data on proposed key events in experiments of  
23 duration two weeks or longer. See Section 4.7.4 for additional discussion of the MOA for  
24 TCE-induced lung tumors. A MOA subsequent to in situ oxidative metabolism, whether  
25 involving mutagenicity, cytotoxicity, or other key events, may also be relevant to other tissues  
26 where TCE would undergo CYP metabolism. For instance, CYP2E1, oxidative metabolites, and  
27 protein adducts have been reported in the testes of rats exposed to TCE, and, in some rat  
28 bioassays, TCE exposure increased the incidence of rat testicular tumors. However, inadequate  
29 data exist to adequately define a MOA hypothesis for this tumor site (see Section 4.8.2.3 for  
30 additional discussion of the MOA for TCE-induced testicular tumors).

31

#### **6.1.5. Susceptibility (see Sections 4.10 and 4.11.3)**

32 There is some evidence that certain populations may be more susceptible to exposure to TCE.  
33 Factors affecting susceptibility examined include lifestage, gender, genetic polymorphisms,

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 race/ethnicity, preexisting health status, and lifestyle factors and nutrition status. Factors that  
2 affect early lifestage susceptibility include exposures such as transplacental transfer and breast  
3 milk ingestion, early lifestage-specific toxicokinetics, and differential outcomes in early  
4 lifestages such as developmental cardiac defects (see Section 4.10.1). Because the weight of  
5 evidence supports a mutagenic MOA being operative for TCE carcinogenicity in the kidney (see  
6 Section 4.4.7), and there is an absence of chemical-specific data to evaluate differences in  
7 carcinogenic susceptibility, early-life susceptibility should be assumed and the age-dependent  
8 adjustment factors (ADAFs) should be applied, in accordance with the Supplemental Guidance  
9 (see summary below in Section 6.2.2.5). Fewer data are available on later lifestages, although  
10 there is suggestive evidence to indicate that older adults may experience increased adverse  
11 effects than younger adults due to greater tissue distribution of TCE. In general, more studies  
12 specifically designed to evaluate effects in early and later lifestages are needed in order to more  
13 fully characterize potential life stage-related TCE toxicity. Gender-specific (see  
14 Section 4.10.2.1) differences also exist in toxicokinetics (e.g., cardiac outputs, percent body fat,  
15 expression of metabolizing enzymes) and susceptibility to toxic endpoints (e.g., gender-specific  
16 effects on the reproductive system, gender differences in baseline risks to endpoints such as  
17 scleroderma or liver cancer). Genetic variation (see Section 4.10.2.2) likely has an effect on the  
18 toxicokinetics of TCE. Increased CYP2E1 activity and GST polymorphisms may influence  
19 susceptibility of TCE due to effects on production of toxic metabolites or may play a role in  
20 variability in toxic response. Differences in genetic polymorphisms related to the metabolism of  
21 TCE have also been observed among various race/ethnic groups (see Section 4.10.2.3).  
22 Preexisting diminished health status (see Section 4.10.2.4) may alter the response to TCE  
23 exposure. Individuals with increased body mass may have an altered toxicokinetic response due  
24 to the increased uptake of TCE into fat. Other conditions that may alter the response to TCE  
25 exposure include diabetes and hypertension, and lifestyle and nutrition factors (see  
26 Section 4.10.2.5) such as alcohol consumption, tobacco smoking, nutritional status, physical  
27 activity, and socioeconomic status. Alcohol intake has been associated with inhibition of TCE  
28 metabolism in both humans and experimental animals. In addition, such conditions have been  
29 associated with increased baseline risks for health effects also associated with TCE, such as  
30 kidney cancer and liver cancer. However, the interaction between TCE and known risk factors  
31 for human diseases is not known, and further evaluation of the effects due to these factors is  
32 needed.

33 In sum, there is some evidence that certain populations may be more susceptible to exposure to  
34 TCE. Factors affecting susceptibility examined include lifestage, gender, genetic  
35 polymorphisms, race/ethnicity, preexisting health status, and lifestyle factors and nutrition status.

1 However, except in the case of toxicokinetic variability characterized using the PBPK model  
2 described in Section 3.5, there are inadequate chemical-specific data to quantify the degree of  
3 differential susceptibility due to such factors.  
4

## 6.2. DOSE-RESPONSE ASSESSMENT

5 This section summarizes the major conclusions of the dose-response analysis for TCE noncancer  
6 effects and carcinogenicity, with more detailed discussions in Section 5.  
7

### 6.2.1. Noncancer Effects (see Section 5.1)

#### 6.2.1.1.1. Background and Methods

8 As summarized above, based on the available human epidemiologic data and experimental and  
9 mechanistic studies, it is concluded that TCE poses a potential human health hazard for  
10 noncancer toxicity to the central nervous system, the kidney, the liver, the immune system, the  
11 male reproductive system, and the developing fetus. The evidence is more limited for TCE  
12 toxicity to the respiratory tract and female reproductive system.

13 Dose-response analysis for a noncancer endpoint generally involves two steps: (1) the  
14 determination of a point of departure (POD) derived from a benchmark dose (BMD),<sup>61</sup> a  
15 no-observed-adverse-effect level (NOAEL), or a lowest-observed-adverse-effect level (LOAEL);  
16 and (2) adjustment of the POD by endpoint/study-specific “uncertainty factors” (UFs),  
17 accounting for adjustments and uncertainties in the extrapolation from the study conditions to  
18 conditions of human exposure.

19 Because of the large number of noncancer health effects associated with TCE exposure and the  
20 large number of studies reporting on these effects, in contrast to toxicological reviews for  
21 chemicals with smaller databases of studies, a formal, quantitative screening process (see  
22 Section 5.1) was used to reduce the number of endpoints and studies to those that would best  
23 inform the selection of the *critical effects* for the inhalation reference concentration (RfC) and  
24 oral reference dose (RfD).<sup>62</sup> As described in Section 5.1, for all studies described in Section 4

---

61 More precisely, it is the benchmark dose lower bound (BMDL), i.e., the (one-sided) 95% lower confidence bound on the dose corresponding to the benchmark response (BMR) for the effect, that is used as the POD.

62 In EPA noncancer health assessments, the RfC [RfD] is an estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation [daily oral] exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious effects during a lifetime. It can be derived

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 which report adverse noncancer health effects and provided quantitative dose-response data,  
2 PODs on the basis of applied dose, adjusted by endpoint/study-specific UFs, were used to  
3 develop candidate RfCs (cRfCs) and candidate RfDs (cRfDs) intended to be protective for each  
4 endpoint individually. Candidate critical effects—those with the lowest cRfCs and cRfDs taking  
5 into account the confidence in each estimate—were selected within each of the following health  
6 effect domains: (1) neurological, (2) systemic/organ system; (3) immunological;  
7 (4) reproductive; and (5) developmental. For each of these candidate critical effects, the PBPK  
8 model developed in Section 3.5 was used for interspecies, intraspecies, and route-to-route  
9 extrapolation on the basis of internal dose to develop PBPK model-based PODs. Plausible  
10 internal dose-metrics were selected based on what is understood about the role of different TCE  
11 metabolites in toxicity and the MOA for toxicity. These PODs were then adjusted by  
12 endpoint/study-specific UFs, taking into account the use of the PBPK model, to develop PBPK  
13 model-based candidate RfCs (p-cRfCs) and candidate RfDs (p-cRfDs). The most sensitive  
14 cRfCs, p-cRfCs, cRfDs, and p-cRfDs were then evaluated, taking into account the confidence in  
15 each estimate, to arrive at overall candidate RfCs and RfDs for each health effect type. Then, the  
16 RfC and RfD for TCE were selected so as to be protective of the most sensitive effects. In  
17 contrast to the approach used in most assessments, in which the RfC and RfD are each based on  
18 a single critical effect, the final RfC and RfD for TCE were based on multiple critical effects that  
19 resulted in very similar candidate RfC and RfD values at the low end of the full range of values.  
20 This approach was taken here because it provides robust estimates of the RfC and RfD and  
21 because it highlights the multiple effects that are all yielding very similar candidate values.  
22

#### **6.2.1.1.2. Uncertainties and Application of Uncertainty Factors (UFs) (see Section 5.1.1 and 5.1.4)**

23 An underlying assumption in deriving reference values for noncancer effects is that the dose-  
24 response relationship for these effects has a threshold. Thus, a fundamental uncertainty is the  
25 validity of that assumption. For some effects, in particular effects on very sensitive processes  
26 (e.g., developmental processes) or effects for which there is a nontrivial background level and  
27 even small exposures may contribute to background disease processes in more susceptible  
28 people, a practical threshold (i.e., a threshold within the range of environmental exposure levels  
29 of regulatory concern) may not exist.

---

from a NOAEL, LOAEL, or benchmark concentration [dose], with uncertainty factors generally applied to reflect limitations of the data used.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Nonetheless, under the assumption of a threshold, the desired exposure level to have as a  
2 reference value is the maximum level at which there is no appreciable risk for an adverse effect  
3 in sensitive subgroups (of humans). However, because it is not possible to know what this level  
4 is, “uncertainty factors” are used to attempt to address quantitatively various aspects, depending  
5 on the data set, of qualitative uncertainty.

6 First there is uncertainty about the “point of departure” for the application of UFs. Conceptually,  
7 the POD should represent the maximum exposure level at which there is no appreciable risk for  
8 an adverse effect in the study population under study conditions (i.e., the threshold in the dose-  
9 response relationship). Then, the application of the relevant UFs is intended to convey that  
10 exposure level to the corresponding exposure level for sensitive human subgroups exposed  
11 continuously for a lifetime. In fact, it is again not possible to know that exposure level even for a  
12 laboratory study because of experimental limitations (e.g., the power to detect an effect, dose  
13 spacing, measurement errors, etc.), and crude approximations like the NOAEL or a BMDL are  
14 used. If a LOAEL is used as the POD, the LOAEL-to-NOAEL UF is applied as an adjustment  
15 factor to better approximate the desired exposure level (threshold), although the necessary extent  
16 of adjustment is unknown. The standard value for the LOAEL-to-NOAEL UF is 10, although  
17 sometimes a value of 3 is used if the effect is considered minimally adverse at the response level  
18 observed at the LOAEL or even one if the effect is an early marker for an adverse effect. For  
19 one POD in this assessment, a value of 30 was used for the LOAEL-to-NOAEL UF because the  
20 incidence rate for the adverse effect was  $\geq 90\%$  at the LOAEL.

21 If a BMDL is used as the POD, there are uncertainties regarding the appropriate dose-response  
22 model to apply to the data, but these should be minimal if the modeling is in the observable  
23 range of the data. There are also uncertainties about what BMR to use to best approximate the  
24 desired exposure level (threshold, see above). For continuous endpoints, in particular, it is often  
25 difficult to identify the level of change that constitutes the “cut-point” for an adverse effect.  
26 Sometimes, to better approximate the desired exposure level, a BMR somewhat below the  
27 observable range of the data is selected. In such cases, the model uncertainty is increased, but  
28 this is a trade-off to reduce the uncertainty about the POD not being a good approximation for  
29 the desired exposure level.

30 For each of these types of PODs, there are additional uncertainties pertaining to adjustments to  
31 the administered exposures (doses). Typically, administered exposures (doses) are converted to  
32 equivalent continuous exposures (daily doses) over the study exposure period under the  
33 assumption that the effects are related to concentration  $\times$  time, independent of the daily (or  
34 weekly) exposure regimen (i.e., a daily exposure of 6 hours to 4 ppm is considered equivalent to  
35 24 hours of exposure to 1 ppm). However, the validity of this assumption is generally unknown,

1 and, if there are dose-rate effects, the assumption of concentration times time ( $C \times t$ ) equivalence  
2 would tend to bias the POD downwards. Where there is evidence that administered exposure  
3 better correlates to the effect than equivalent continuous exposure averaged over the study  
4 exposure period (e.g., visual effects), administered exposure was not adjusted. For the PBPK  
5 analyses in this assessment, the actual administered exposures are taken into account in the  
6 PBPK modeling, and equivalent daily values (averaged over the study exposure period) for the  
7 dose-metrics are obtained (see above, Section 5.1.3.2). Additional uncertainties about the  
8 PBPK-based estimates include uncertainties about the appropriate dose-metric for each effect,  
9 although for some effects there was better information about relevant dose-metrics than for  
10 others, and uncertainties in the PBPK model predictions for the dose-metrics in humans,  
11 particularly for GSH conjugation (see Section 5.1.3.1).

12 There is also uncertainty about the other UFs. The human variability UF is to some extent an  
13 adjustment factor because for more sensitive people, the dose-response relationship shifts to  
14 lower exposures. But there is uncertainty about the extent of the adjustment required, i.e., about  
15 the distribution of human susceptibility. Therefore, in the absence of data on a susceptible  
16 population(s) or on the distribution of susceptibility in the general population, an UF of 10 is  
17 generally used, which breaks down (approximately) to a factor of 3 for pharmacokinetic  
18 variability and a factor of 3 for pharmacodynamic variability. This standard value was used for  
19 all the PODs based on applied dose in this assessment with the exception of the PODs for a few  
20 immunological effects that were based on data from a sensitive (autoimmune-prone) mouse  
21 strain. For those PODs, an UF of 3 (reflecting pharmacokinetics only) was used for human  
22 variability. The PBPK analyses in this assessment attempt to account for the pharmacokinetic  
23 portion of human variability using human data on pharmacokinetic variability. For PBPK  
24 model-based candidate reference values, the pharmacokinetic component of this UF was omitted.

25 A quantitative uncertainty analysis of the PBPK derived dose-metrics used in the assessment is  
26 presented in Section 5.1.4.2 in Section 5. There is still uncertainty regarding the susceptible  
27 subgroups for TCE exposure and the extent of pharmacodynamic variability.

28 If the data used to determine a particular POD are from laboratory animals, an interspecies  
29 extrapolation UF is used. This UF is also to some extent an adjustment factor for the expected  
30 scaling for toxicologically equivalent doses across species (i.e., according to body weight to the  
31  $3/4$  power for oral exposures). However, there is also uncertainty about the true extent of  
32 interspecies differences for specific noncancer effects from specific chemical exposures. For  
33 oral exposures, the standard value for the interspecies UF is 10, which can be viewed as breaking  
34 down (approximately) to a factor of 3 for the “adjustment” (nominally pharmacokinetics) and a  
35 factor of 3 for the “uncertainty” (nominally pharmacodynamics). For inhalation exposures for

1 systemic toxicants such as TCE, no adjustment across species is generally assumed for fixed air  
2 concentrations (ppm equivalence), and the standard value for the interspecies UF is 3 reflects  
3 “uncertainty” (nominally pharmacodynamics only). The PBPK analyses in this assessment  
4 attempt to account for the “adjustment” portion of interspecies extrapolation using rodent  
5 pharmacokinetic data to estimate internal doses for various dose-metrics. Equal doses of these  
6 dose-metrics, appropriately scaled, are then assumed to convey equivalent risk across species.  
7 For PBPK model-based candidate reference values, the “adjustment” component of this UF was  
8 omitted. With respect to the “uncertainty” component, quantitative uncertainty analyses of the  
9 PBPK-derived dose-metrics used in the assessment are presented in Section 5.1.4.2 in Section 5.  
10 However, these only address the pharmacokinetic uncertainties in a particular dose-metric, and  
11 there is still uncertainty regarding the true dose-metrics. Nor do the PBPK analyses address the  
12 uncertainty in either cross-species pharmacodynamic differences (i.e., about the assumption that  
13 equal doses of the appropriate dose-metric convey equivalent risk across species for a particular  
14 endpoint from a specific chemical exposure) or in cross-species pharmacokinetic differences not  
15 accounted for by the PBPK model dose-metrics (e.g., departures from the assumed interspecies  
16 scaling of clearance of the active moiety, in the cases where only its production is estimated). A  
17 value of 3 is typically used for the “uncertainty” about cross-species differences, and this  
18 generally represents true uncertainty because it is usually unknown, even after adjustments have  
19 been made to account for the expected interspecies differences, whether humans have more or  
20 less susceptibility, and to what degree, than the laboratory species in question.  
21 RfCs and RfDs apply to lifetime exposure, but sometimes the best (or only) available data come  
22 from less-than-lifetime studies. Lifetime exposure can induce effects that may not be apparent or  
23 as large in magnitude in a shorter study; consequently, a dose that elicits a specific level of  
24 response from a lifetime exposure may be less than the dose eliciting the same level of response  
25 from a shorter exposure period. If the effect becomes more severe with increasing exposure,  
26 then chronic exposure would shift the dose-response relationship to lower exposures, although  
27 the true extent of the shift is unknown. PODs based on subchronic exposure data are generally  
28 divided by a subchronic-to-chronic UF, which has a standard value of 10. If there is evidence  
29 suggesting that exposure for longer time periods does not increase the magnitude of an effect, a  
30 lower value of 3 or 1 might be used. For some reproductive and developmental effects, chronic  
31 exposure is that which covers a specific window of exposure that is relevant for eliciting the  
32 effect, and subchronic exposure would correspond to an exposure that is notably less than the full  
33 window of exposure.  
34 Sometimes a database UF is also applied to address limitations or uncertainties in the database.  
35 The overall database for TCE is quite extensive, with studies for many different types of effects,

1 including 2-generation reproductive studies, as well as neurological and immunological studies.  
2 In addition, there were sufficient data to develop a reliable PBPK model to estimate route-to-  
3 route extrapolated doses for some candidate critical effects for which data were only available  
4 for one route of exposure. Thus, there is a high degree of confidence that the TCE database was  
5 sufficient to identify some sensitive endpoints, and no database UF was used in this assessment.  
6

#### **6.2.1.1.3. Candidate Critical Effects and Reference Values (see Sections 5.1.2 and 5.1.3)**

7 A large number of endpoints and studies were considered within each health effect  
8 domain. Section 5 contains a comprehensive discussion of all endpoints/studies which were  
9 considered for developing candidate reference values (cRfCs, cRfDs, p-cRfCs, and p-cRfDs),  
10 their PODs, and the UFs applied. The summary below reviews the selection of candidate critical  
11 effects for each health effect domain, the confidence in the reference values, the selection of  
12 PBPK model-based dose-metrics, and the impact of PBPK modeling on the candidate reference  
13 values.  
14

#### **6.2.1.1.4. Neurological effects**

15 Candidate reference values were developed for several neurological domains for which there was  
16 evidence of hazard (see Tables 5-2 and 5-13). There is higher confidence in the candidate  
17 reference values for trigeminal nerve, auditory, or psychomotor effects, but the available data  
18 suggest that the more sensitive indicators of TCE neurotoxicity are changes in wakefulness,  
19 regeneration of the sciatic nerve, demyelination in the hippocampus and degeneration of  
20 dopaminergic neurons. Therefore, these more sensitive effects are considered the candidate  
21 critical effects for neurotoxicity, albeit with more uncertainty in the corresponding candidate  
22 reference values. Of these more sensitive effects, there is greater confidence in the changes in  
23 wakefulness reported by Arito et al. (1994). In addition, trigeminal nerve effects are considered  
24 a candidate critical effect because this is the only type of neurological effect for which human  
25 data are available, and the POD for this effect is similar to that from the most sensitive rodent  
26 study (Arito et al., 1994, for changes in wakefulness). Between the two human studies of  
27 trigeminal nerve effects, Ruijten et al. (1991) is preferred for deriving noncancer reference  
28 values because its exposure characterization is considered more reliable.  
29 Because of the lack of specific data as to the metabolites involved and the MOA for the  
30 candidate critical neurologic effects, PBPK model predictions of total metabolism (scaled by  
31 body weight to the  $\frac{3}{4}$  power) were selected as the preferred dose-metric based on the general

*This document is a draft for review purposes only and does not constitute Agency policy.*



1 observation that TCE toxicity is associated with metabolism. The area-under-the-curve (AUC)  
2 of TCE in blood was used as an alternative dose-metric. With these dose-metrics, the candidate  
3 reference values derived using the PBPK model were only modestly (~threefold or less) different  
4 than those derived on the basis of applied dose.  
5

#### 6.2.1.1.5. Kidney effects

6 Candidate reference values were developed for histopathological and weight changes in the  
7 kidney (see Tables 5-4 and 5-14), and these are considered to be candidate critical effects for  
8 several reasons. First, they appear to be the most sensitive indicators of toxicity that are  
9 available for the kidney. In addition, as discussed in Sections 3.3 and 3.5, both in vitro and in  
10 vivo pharmacokinetic data indicate substantially more production of GSH-conjugates thought to  
11 mediate TCE kidney effects in humans relative to rats and mice. Several studies are considered  
12 reliable for developing candidate reference values for these endpoints. For histopathological  
13 changes, these were the only available inhalation study (the rat study of Maltoni et al., 1986), the  
14 NTP (1988) study in rats, and the National Cancer Institute (NCI, 1976) study in mice. For  
15 kidney weight changes, both available studies (Kjellstrand et al., 1983a; Woolhiser et al., 2006)  
16 were chosen as candidate critical studies.

17 Due to the substantial evidence supporting the role of GSH conjugation metabolites in TCE-  
18 induced nephrotoxicity, the preferred PBPK model dose-metrics for kidney effects were the  
19 amount of DCVC bioactivated in the kidney for rat studies and the amount of GSH conjugation  
20 (both scaled by body weight to the  $3/4$  power) for mouse studies (inadequate toxicokinetic data are  
21 available in mice for predicting the amount of DCVC bioactivation). With these dose-metrics,  
22 the candidate reference values derived using the PBPK model were 300- to 400-fold lower than  
23 those derived on the basis of applied dose. As discussed above and in Section 3, this is due to  
24 the available in vivo and in vitro data supporting not only substantially more GSH conjugation in  
25 humans than in rodents, but also substantial interindividual toxicokinetic variability. Overall,  
26 there is high confidence in the nephrotoxic hazard from TCE exposure and in the appropriateness  
27 of the dose-metrics discussed above; however, there is substantial uncertainty in the  
28 extrapolation of GSH conjugation from rodents to humans due to limitations in the available data  
29 (see Section 3.3.3.2).  
30

#### 6.2.1.1.6. Liver effects

1 Hepatomegaly appears to be the most sensitive indicator of toxicity that is available for the liver  
2 and is therefore, considered a candidate critical effect. Several studies are considered reliable for  
3 developing high confidence candidate reference values for this endpoint. Since they all indicated  
4 similar sensitivity but represented different species and/or routes of exposure, they were all  
5 considered candidate critical studies (see Tables 5-4 and 5-14).  
6 Due to the substantial evidence supporting the role of oxidative metabolism in TCE-induced  
7 hepatomegaly (and evidence against TCA being the sole mediator of TCE-induced hepatomegaly  
8 (Evans et al., 2009)), the preferred PBPK model dose-metric for liver effects was the amount of  
9 hepatic oxidative metabolism (scaled by body weight to the  $\frac{3}{4}$  power). Total (hepatic and  
10 extrahepatic) oxidative metabolism (scaled by body weight to the  $\frac{3}{4}$  power) was used as an  
11 alternative dose-metric. With these dose-metrics, the candidate reference values derived using  
12 the PBPK model were only modestly (~threefold or less) different than those derived on the  
13 basis of applied dose.  
14

#### 6.2.1.1.7. Immunological effects

15 There is high qualitative confidence for TCE immunotoxicity and moderate confidence in the  
16 candidate reference values that can be derived from the available studies (see Tables 5-6  
17 and 5-16). Decreased thymus weight reported at relatively low exposures in nonautoimmune-  
18 prone mice is a clear indicator of immunotoxicity (Keil et al., 2009), and is therefore, considered  
19 a candidate critical effect. A number of studies have also reported changes in markers of  
20 immunotoxicity at relatively low exposures. Among markers for autoimmune effects, the more  
21 sensitive measures of autoimmune changes in liver and spleen (Kaneko et al., 2000) and  
22 increased anti-dsDNA and anti-ssDNA antibodies (early markers for systemic lupus  
23 erythematosus) (Keil et al., 2009) are considered the candidate critical effects. For markers of  
24 immunosuppression, the more sensitive measures of decreased PFC response (Woolhiser et al.,  
25 2006), decreased stem cell bone marrow recolonization, and decreased cell-mediated response to  
26 sRBC (both from Sanders et al., 1982b) are considered the candidate critical effects.  
27 Developmental immunological effects are discussed below as part of the summary of  
28 developmental effects (see Section 6.2.1.3.6).  
29 Because of the lack of specific data as to the metabolites involved and the MOA for the  
30 candidate critical immunologic effects, PBPK model predictions of total metabolism (scaled by  
31 body weight to the  $\frac{3}{4}$  power) was selected as the preferred dose-metric based on the general  
32 observation that TCE toxicity is associated with metabolism. The AUC of TCE in blood was

1 used as an alternative dose-metric. With these dose-metrics, the candidate reference values  
2 derived using the PBPK model were, with one exception, only modestly (~threefold or less)  
3 different than those derived on the basis of applied dose. For the Woolhiser et al. (2006)  
4 decreased PFC response, with the alternative dose-metric of AUC of TCE in blood, BMD  
5 modeling based on internal doses changed the candidate reference value by 17-fold higher than  
6 the cRfC based on applied dose. However, the dose-response model fit for this effect using this  
7 metric was substantially worse than the fit using the preferred metric of total oxidative  
8 metabolism, with which the change in candidate reference value was only 1.3-fold.  
9

#### 6.2.1.1.8. Reproductive effects

10 While there is high qualitative confidence in the male reproductive hazard posed by TCE, there  
11 is lower confidence in the reference values that can be derived from the available studies of these  
12 effects (see Tables 5-8 and 5-17). Relatively high PODs are derived from several studies  
13 reporting less sensitive endpoints (George et al., 1985, 1986; Land et al., 1981), and  
14 correspondingly higher cRfCs and cRfDs suggest that they are not likely to be critical effects.  
15 The studies reporting more sensitive endpoints also tend to have greater uncertainty. For the  
16 human study by Chia et al. (1996), there are uncertainties in the characterization of exposure and  
17 the adversity of the effect measured in the study. For the Kumar et al. (2001b; 2000a; 2000b),  
18 Forkert et al. (2002) and Kan et al. (2007) studies, the severity of the sperm and testes effects  
19 appears to be continuing to increase with duration even at the end of the study, so it is plausible  
20 that a lower exposure for a longer duration may elicit similar effects. For the DuTeaux et al.  
21 (2004a) study, there is also duration- and low-dose extrapolation uncertainty due to the short  
22 duration of the study in comparison to the time period for sperm development as well as the lack  
23 of a NOAEL at the tested doses. Overall, even though there are limitations in the quantitative  
24 assessment, there remains sufficient evidence to consider these to be candidate critical effects.  
25 There is moderate confidence both in the hazard and the candidate reference values for  
26 reproductive effects other than male reproductive effects. While there are multiple studies  
27 suggesting decreased maternal body weight with TCE exposure, this systemic change may not be  
28 indicative of more sensitive reproductive effects. None of the estimates developed from other  
29 reproductive effects is particularly uncertain or unreliable. Therefore, delayed parturition  
30 (Narotsky et al., 1995) and decreased mating (George et al., 1986), which yielded the lowest  
31 cRfDs, were considered candidate critical effects. These effects were also included so that  
32 candidate critical reproductive effects from oral studies would not include only that reported by  
33 DuTeaux et al. (2004a), from which deriving the cRfD entailed a higher degree of uncertainty.

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 Because of the general lack of specific data as to the metabolites involved and the MOA for the  
2 candidate critical reproductive effects, PBPK model predictions of total metabolism (scaled by  
3 body weight to the  $\frac{3}{4}$  power) was selected as the preferred dose-metric based on the general  
4 observation that TCE toxicity is associated with metabolism. The AUC of TCE in blood was  
5 used as an alternative dose-metric. The only exception to this was for the DuTeaux et al. (2004a)  
6 study, which suggested that local oxidative metabolism of TCE in the male reproductive tract  
7 was involved in the effects reported. Therefore, in this case, AUC of TCE in blood was  
8 considered the preferred dose-metric, while total oxidative metabolism (scaled by body weight to  
9 the  $\frac{3}{4}$  power) was considered the alternative metric. With these dose-metrics, the candidate  
10 reference values derived using the PBPK model were only modestly (~3.5-fold or less) different  
11 than those derived on the basis of applied dose.  
12

#### 6.2.1.1.9. Developmental effects

13 There is moderate-to-high confidence both in the hazard and the candidate reference values for  
14 developmental effects of TCE (see Tables 5-10 and 5-18). It is also noteworthy that the PODs  
15 for the more sensitive developmental effects were similar to or, in most cases, lower than the  
16 PODs for the more sensitive reproductive effects, suggesting that developmental effects are not a  
17 result of paternal or maternal toxicity. Among inhalation studies, candidate reference values  
18 were only developed for effects in rats reported in Healy et al. (1982), of resorptions, decreased  
19 fetal weight, and delayed skeletal ossification. These were all considered candidate critical  
20 developmental effects. Because resorptions were also reported in oral studies, the most sensitive  
21 (rat) oral study for this effect (and most reliable for dose-response analysis) of Narotsky et al.  
22 (1995) was also selected as a candidate critical study. The confidence in the oral studies and  
23 candidate reference values developed for more sensitive endpoints is more moderate, but still  
24 sufficient for consideration as candidate critical effects. The most sensitive endpoints by far are  
25 the increased fetal heart malformations in rats reported by Johnson et al. (2003) and the  
26 developmental immunotoxicity in mice reported by Peden-Adams et al. (2006), and these are  
27 both considered candidate critical effects. Neurodevelopmental effects are a distinct type among  
28 developmental effects. Thus, the next most sensitive endpoints of decreased rearing  
29 postexposure in mice (Fredriksson et al., 1993), increased exploration postexposure in rats  
30 (Taylor et al., 1985) and decreased myelination in the hippocampus of rats (Isaacson and Taylor,  
31 1989) are also considered candidate critical effects.  
32 Because of the general lack of specific data as to the metabolites involved and the MOA for the  
33 candidate critical reproductive effects, PBPK model predictions of total metabolism (scaled by

1 body weight to the  $\frac{3}{4}$  power) was selected as the preferred dose-metric based on the general  
2 observation that TCE toxicity is associated with metabolism. The AUC of TCE in blood was  
3 used as an alternative dose-metric. The only exception to this was for the Johnson et al. (2003)  
4 study, which suggested that oxidative metabolites were involved in the effects reported based on  
5 similar effects being reported from TCA and DCA exposure. Therefore, in this case, total  
6 oxidative metabolism (scaled by body weight to the  $\frac{3}{4}$  power) was considered the preferred dose-  
7 metric, while AUC of TCE in blood was considered the alternative metric. With these dose-  
8 metrics, the candidate reference values derived using the PBPK model were, with one exception,  
9 only modestly (~threefold or less) different than those derived on the basis of applied dose. For  
10 resorptions reported by Narotsky et al. (1995), BMD modeling based on internal doses changed  
11 the candidate reference value by seven to eightfold larger than the corresponding cRfD based on  
12 applied dose. However, there is substantial uncertainty in the low-dose curvature of the  
13 dose-response curve for modeling both with applied and internal dose, so the BMD remains  
14 somewhat uncertain for this endpoint/study. Finally, for two studies (Isaacson and Taylor, 1989;  
15 Peden-Adams et al., 2006), PBPK modeling of internal doses was not performed due to the  
16 inability to model the complicated exposure pattern (in utero, followed by lactational transfer,  
17 followed by drinking water postweaning).  
18

#### **6.2.1.1.10. Summary of most sensitive candidate reference values**

19 As shown in Section 5.1.3 and 5.1.5, the most sensitive candidate reference values are for the  
20 developmental effect of heart malformations in rats (candidate RfC of 0.0004 ppm and candidate  
21 RfD of 0.0005 mg/kg-day), developmental immunotoxicity in mice exposed pre and postnatally  
22 (candidate RfD of 0.0004 mg/kg-day), immunological effects in mice (lowest candidate RfCs of  
23 0.0003–0.003 ppm and lowest candidate RfDs of 0.0005–0.005 mg/kg-day), and kidney effects  
24 in rats and mice (candidate RfCs of 0.0006–0.002 ppm and candidate RfDs of  
25 0.0003–0.001 mg/kg-day). The most sensitive candidate reference values also generally have  
26 low composite uncertainty factors (with the exception of some mouse immunological and kidney  
27 effects), so they are expected to be reflective of the most sensitive effects as well. Thus, the  
28 most sensitive candidate references values for multiple effects span about an order of magnitude  
29 for both inhalation (0.0003–0.003 ppm [0.002–0.02 mg/m<sup>3</sup>]) and oral (0.0004–0.005 mg/kg-day)  
30 exposures. The most sensitive candidate references values for neurological and reproductive  
31 effects are about an order of magnitude higher (lowest candidate RfCs of 0.007–0.02 ppm  
32 [0.04–0.1 mg/m<sup>3</sup>] and lowest candidate RfDs of 0.009–0.02 mg/kg-day). Lastly, the liver effects

1 have candidate reference values that are another two orders of magnitude higher (candidate RfCs  
2 of 1–2 ppm [6–10 mg/m<sup>3</sup>] and candidate RfDs of 0.9–2 mg/kg-day).  
3

#### 6.2.1.1.11. Noncancer Reference Values (see Section 5.1.5)

##### *Reference concentration*

4 The goal is to select an overall RfC that is well supported by the available data (i.e., without  
5 excessive uncertainty given the extensive database) and protective for all the candidate critical  
6 effects, recognizing that individual candidate RfC values are by nature somewhat imprecise. As  
7 discussed in Section 5.1 in Section 5, the lowest candidate RfC values within each health effect  
8 category span a 3,000-fold range from 0.0003–0.9 ppm (see Table 5-26). One approach to  
9 selecting a RfC would be to select the lowest calculated value of 0.0003 ppm for decreased  
10 thymus weight in mice. However, three candidate RfCs (cRfCs and p-cRfCs) are in the  
11 relatively narrow range of 0.0003–0.0006 ppm at the low end of the overall range (see  
12 Table 5-24). Given the somewhat imprecise nature of the individual candidate RfC values, and  
13 the fact that multiple effects/studies lead to similar candidate RfC values, the approach taken in  
14 this assessment is to select a RfC supported by multiple effects/studies. The advantages of this  
15 approach, which is only possible when there is a relatively large database of studies/effects and  
16 when multiple candidate values happen to fall within a narrow range at the low end of the overall  
17 range, are that it leads to a more robust RfC (less sensitive to limitations of individual studies)  
18 and that it provides the important characterization that the RfC exposure level is similar for  
19 multiple noncancer effects rather than being based on a sole explicit critical effect.  
20 Therefore, two critical and one supporting studies/effects were chosen as the basis of the RfC for  
21 TCE noncancer effects (see Tables 5-28 and 5-29). These lowest candidate RfCs, ranging from  
22 0.0003–0.0006 ppm for developmental, kidney, and immunologic effects, are values derived  
23 from route-to-route extrapolation using the PBPK model. The lowest candidate RfC estimate  
24 from an inhalation study is 0.001 ppm for kidney effects, which is higher than the route-to-route  
25 extrapolated candidate RfC estimate from the most sensitive oral study. For all of the candidate  
26 RfCs, the PBPK model was used for inter and intraspecies extrapolation, based on the preferred  
27 dose-metric for each endpoint. There is moderate-to-high confidence in the lowest candidate  
28 RfC for immunological effects (see Section 5.1.2.5), and moderate confidence in the lowest  
29 candidate RfC for developmental effects (heart malformations) (see Section 5.1.2.8); these are  
30 considered the critical effects for deriving the RfC. For kidney effects (toxic nephropathy), there  
31 is high confidence in the nephrotoxic hazard from TCE exposure and in the appropriateness of

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 the selected dose-metric; however, as discussed in Section 3.3.3.2, there remains substantial  
2 uncertainty in the extrapolation of GSH conjugation from rodents to humans due to limitations in  
3 the available data, and thus toxic nephropathy is considered a supporting effect.  
4 As a whole, the estimates support a RfC of 0.0004 ppm (0.4 ppb or 2  $\mu\text{g}/\text{m}^3$ ). This estimate  
5 essentially reflects the midpoint between the similar candidate RfC estimates for the two critical  
6 effects (0.00033 ppm for decreased thymus weight in mice and 0.00037 ppm for heart  
7 malformations in rats), rounded to one significant figure. This estimate is also within a factor of  
8 2 of the candidate RfC estimate of 0.00006 ppm for the supporting effect of toxic nephropathy in  
9 rats. Thus, this assessment does not rely on a single estimate alone; rather, each estimate is  
10 supported by estimates of similar magnitude from other effects. In other words, there is robust  
11 support for an RfC of 0.0004 ppm provided by estimates for multiple effects from multiple  
12 studies. The estimates are based on PBPK model-based estimates of internal dose for  
13 interspecies, intraspecies, and route-to-route extrapolation, and there is sufficient confidence in  
14 the PBPK model and support from mechanistic data for one of the dose-metrics (total oxidative  
15 metabolism for the heart malformations). There is high confidence that bioactivation of DCVC  
16 and total GSH metabolism would be appropriate dose-metrics for toxic nephropathy, but there is  
17 substantial uncertainty in the PBPK model predictions for these dose-metrics in humans (see  
18 Section 5.1.3.1). Note that there is some human evidence of developmental heart defects from  
19 TCE exposure in community studies (see Section 4.8.3.1.1) and of kidney toxicity in TCE-  
20 exposed workers (see Section 4.4.1).  
21 In summary, the RfC is **0.0004 ppm** (0.4 ppb or 2  $\mu\text{g}/\text{m}^3$ ) based on route-to-route extrapolated  
22 results from oral studies for the critical effects of heart malformations (rats) and immunotoxicity  
23 (mice). This RfC value is further supported by route-to-route extrapolated results from an oral  
24 study of toxic nephropathy (rats).  
25

#### 6.2.1.1.12. Reference dose

26 As with the RfC determination above, the goal is to select an overall RfD that is well supported  
27 by the available data (i.e., without excessive uncertainty given the extensive database) and  
28 protective for all the candidate critical effects, recognizing that individual candidate RfD values  
29 are by nature somewhat imprecise. As discussed in Section 5.1 in Section 5, the lowest  
30 candidate RfD values (cRfDs and p-cRfDs) within each health effect category span a nearly  
31 3,000-fold range from 0.0003–0.8 mg/kg-day (see Table 5-26). However, multiple candidate  
32 RfDs are in the relatively narrow range of 0.0003–0.0008 mg/kg-day at the low end of the  
33 overall range. Given the somewhat imprecise nature of the individual candidate RfD values, and

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 the fact that multiple effects/studies lead to similar candidate RfD values, the approach taken in  
2 this assessment is to select a RfD supported by multiple effects/studies. The advantages of this  
3 approach, which is only possible when there is a relatively large database of studies/effects and  
4 when multiple candidate values happen to fall within a narrow range at the low end of the overall  
5 range, are that it leads to a more robust RfD (less sensitive to limitations of individual studies)  
6 and that it provides the important characterization that the RfD exposure level is similar for  
7 multiple noncancer effects rather than being based on a sole explicit critical effect.  
8 Therefore, three critical and two supporting studies/effects were chosen as the basis of the RfD  
9 for TCE noncancer effects (see Tables 5-30 and 5-31). All but one of the lowest candidate RfD  
10 values—0.0008 mg/kg-day for increased kidney weight in rats, 0.0005 mg/kg-day for both heart  
11 malformations in rats and decreased thymus weights in mice, and 0.0003 mg/kg-day for  
12 increased toxic nephropathy in rats—are derived using the PBPK model for inter and  
13 intraspecies extrapolation, based on the preferred dose-metric for each endpoint, and the latter  
14 value is derived also using the PBPK model for route-to-route extrapolation from an inhalation  
15 study. The other of these lowest candidate RfDs—0.0004 mg/kg-day for developmental  
16 immunotoxicity (decreased PFC response and increased delayed-type hypersensitivity) in  
17 mice—is based on applied dose. There is moderate-to-high confidence in the candidate RfDs for  
18 decreased thymus weights (see Section 5.1.2.5) and developmental immunological effects, and  
19 moderate confidence in that for heart malformations (see Section 5.1.2.8); these are considered  
20 the critical effects for deriving the RfC. For kidney effects, there is high confidence in the  
21 nephrotoxic hazard from TCE exposure and in the appropriateness of the selected dose-metric;  
22 however, as discussed in Section 3.3.3.2, there remains substantial uncertainty in the  
23 extrapolation of GSH conjugation from rodents to humans due to limitations in the available  
24 data, and thus these effects are considered supporting effects.  
25 As a whole, the estimates support a RfD of 0.0005 mg/kg-day. This estimate is within 20% of  
26 the estimates for the critical effects—0.0004 mg/kg-day for developmental immunotoxicity  
27 (decreased PFC and increased delayed-type hypersensitivity) in mice and 0.0005 mg/kg-day for  
28 both heart malformations in rats and decreased thymus weights in mice. This estimate is also  
29 within approximately a factor of 2 of the supporting effect estimates of 0.0003 mg/kg-day for  
30 toxic nephropathy in rats and 0.0008 mg/kg-day for increased kidney weight in rats. Thus, this  
31 assessment does not rely on any single estimate alone; rather, each estimate is supported by  
32 estimates of similar magnitude from other effects. In other words, there is strong, robust support  
33 for an RfD of 0.0005 mg/kg-day provided by the concordance of estimates derived from multiple  
34 effects from multiple studies. The estimates for kidney effects, thymus effects, and  
35 developmental heart malformations are based on PBPK model-based estimates of internal dose



1 for interspecies and intraspecies extrapolation, and there is sufficient confidence in the PBPK  
2 model and support from mechanistic data for one of the dose-metrics (total oxidative metabolism  
3 for the heart malformations). There is high confidence that bioactivation of DCVC would be an  
4 appropriate dose-metric for toxic nephropathy, but there is substantial uncertainty in the PBPK  
5 model predictions for this dose-metric in humans (see Section 5.1.3.1). Note that there is some  
6 human evidence of developmental heart defects from TCE exposure in community studies (see  
7 Section 4.8.3.1.1) and of kidney toxicity in TCE-exposed workers (see Section 4.4.1).  
8 In summary, the RfD is **0.0005 mg/kg-day** based on the critical effects of heart malformations  
9 (rats), adult immunological effects (mice), and developmental immunotoxicity (mice), and toxic  
10 nephropathy (rats), all from oral studies. This RfD value is further supported by results from an  
11 oral study for the effect of toxic nephropathy (rats) and route-to-route extrapolated results from  
12 an inhalation study for the effect of increased kidney weight (rats).  
13

## 6.2.2. Cancer (see Section 5.2)

### 6.2.2.1.1. Background and Methods (rodent: see Section 5.2.1.1; human: see Section 5.2.2.1)

14 As summarized above, following EPA (2005c) *Guidelines for Carcinogen Risk Assessment*, TCE  
15 is characterized as “*carcinogenic to humans*” by all routes of exposure, based on convincing  
16 evidence of a causal association between TCE exposure in humans and kidney cancer, but there  
17 is also human evidence of TCE carcinogenicity in the liver and lymphoid tissues. This  
18 conclusion is further supported by rodent bioassay data indicating carcinogenicity of TCE in rats  
19 and mice at tumor sites that include those identified in human epidemiologic studies. Therefore,  
20 both human epidemiologic studies as well as rodent bioassays were considered for deriving  
21 PODs for dose-response assessment of cancer endpoints. For PODs derived from rodent  
22 bioassays, default dosimetry procedures were applied to convert applied rodent doses to human  
23 equivalent doses. Essentially, for inhalation exposures, “ppm equivalence” across species was  
24 assumed. For oral doses,  $\frac{3}{4}$ -power body-weight scaling was used, with a default average human  
25 body weight of 70 kg. In addition to applied doses, several internal dose-metrics estimated using  
26 a PBPK model for TCE and its metabolites were used in the dose-response modeling for each  
27 tumor type. In general, an attempt was made to use tissue-specific dose-metrics representing  
28 particular pathways or metabolites identified from available data as having a likely role in the  
29 induction of a tissue-specific cancer. Where insufficient information was available to establish  
30 particular metabolites or pathways of likely relevance to a tissue-specific cancer, more general

1 “upstream” metrics had to be used. In addition, the selection of dose-metrics was limited to  
2 metrics that could be adequately estimated by the PBPK model.

3 Regarding low-dose extrapolation, a key consideration in determining what extrapolation  
4 approach to use is the MOA(s). However, MOA data are lacking or limited for each of the  
5 cancer responses associated with TCE exposure, with the exception of the kidney tumors. For  
6 the kidney tumors, the weight of the available evidence supports the conclusion that a mutagenic  
7 MOA is operative; this MOA supports linear low-dose extrapolation. For the other TCE-induced  
8 tumors, the MOA(s) is unknown. When the MOA(s) cannot be clearly defined, EPA generally  
9 uses a linear approach to estimate low-dose risk (2005c), based on the following general  
10 principles:

- 11
- 12
- 13 • A chemical’s carcinogenic effects may act additively to ongoing biological processes,  
14 given that diverse human populations are already exposed to other agents and have  
15 substantial background incidences of various cancers.
- 16 • A broadening of the dose-response curve (i.e., less rapid fall-off of response with  
17 decreasing dose) in diverse human populations and, accordingly, a greater potential for  
18 risks from low-dose exposures (Lutz et al., 2005; Zeise et al., 1987) is expected for two  
19 reasons: First, even if there is a “threshold” concentration for effects at the cellular level,  
20 that threshold is expected to differ across individuals. Second, greater variability in  
21 response to exposures would be anticipated in heterogeneous populations than in inbred  
22 laboratory species under controlled conditions (due to, e.g., genetic variability, disease  
23 status, age, nutrition, and smoking status).
- 24 • The general use of linear extrapolation provides reasonable upper-bound estimates that  
25 are believed to be health-protective (U.S. EPA, 2005c) and also provides consistency  
26 across assessments.

#### 27

**6.2.2.1.2. Inhalation Unit Risk Estimate (rodent: see Section 5.2.1.3; human: see  
Section 5.2.2.1 and 5.2.2.2)**

28 The inhalation unit risk for TCE is defined as a plausible upper bound lifetime extra risk of  
29 cancer from chronic inhalation of TCE per unit of air concentration. The inhalation unit risk for  
30 TCE is  $2.20 \times 10^{-2}$  per ppm ( $2 \times 10^{-2}$  per ppm [ $4 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ]) rounded to one significant  
31 figure), based on human kidney cancer risks reported by Charbotel et al. (2006) and adjusted for  
32 potential risk for tumors at multiple sites. This estimate is based on good-quality human data,  
33 thus avoiding the uncertainties inherent in interspecies extrapolation. The Charbotel et al. (2006)  
34 case-control study of 86 incident renal cell carcinoma (RCC) cases and 316 age- and sex-  
35 matched controls, with individual cumulative exposure estimates for TCE inhalation for each

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 subject, provides a sufficient human data set for deriving quantitative cancer risk estimates for  
2 RCC in humans. The study is a high-quality study which used a detailed exposure assessment  
3 (Fevotte et al., 2006) and took numerous potential confounding factors, including exposure to  
4 other chemicals, into account. A significant dose-response relationship was reported for  
5 cumulative TCE exposure and RCC (Charbotel et al., 2006). Human data on TCE exposure and  
6 cancer risk sufficient for dose-response modeling are only available for RCC, yet human and  
7 rodent data suggest that TCE exposure increases the risk of cancer at other sites as well. In  
8 particular, there is evidence from human (and rodent) studies for increased risks of lymphoma  
9 and liver cancer. Therefore, the inhalation unit risk estimate derived from human data for RCC  
10 incidence was adjusted to account for potential increased risk of those tumor types. To make this  
11 adjustment, a factor accounting for the relative contributions to the extra risk for cancer  
12 incidence from TCE exposure for these three tumor types combined versus the extra risk for  
13 RCC alone was estimated, and this factor was applied to the unit risk estimate for RCC to obtain  
14 a unit risk estimate for the three tumor types combined (i.e., lifetime extra risk for developing  
15 *any* of the three types of tumor). This estimate is considered a better estimate of total cancer risk  
16 from TCE exposure than the estimate for RCC alone. Although only the Charbotel et al. (2006)  
17 study was found adequate for direct estimation of inhalation unit risks, the available  
18 epidemiologic data provide sufficient information for estimating the *relative* potency of TCE  
19 across tumor sites. In particular, the relative contributions to extra risk (for cancer incidence)  
20 were calculated from two different data sets to derive the adjustment factor for adjusting the unit  
21 risk estimate for RCC to a unit risk estimate for the three types of cancers (RCC, lymphoma, and  
22 liver) combined. The first calculation is based on the results of the meta-analyses of human  
23 epidemiologic data for the three tumor types; the second calculation is based on the results of the  
24 Raaschou-Nielsen et al. (2003) study, the largest single human epidemiologic study by far with  
25 RR estimates for all three tumor types. These calculations support an adjustment factor of .  
26 The inhalation unit risk based on human epidemiologic data is supported by inhalation unit risk  
27 estimates from multiple rodent bioassays, the most sensitive of which range from  $1 \times 10^{-2}$  to  $2 \times$   
28  $10^{-1}$  per ppm [ $2 \times 10^{-6}$  to  $3 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ]. From the inhalation bioassays selected for  
29 analysis in Section 5.2.1.1, and using the preferred PBPK model-based dose-metrics, the  
30 inhalation unit risk estimate for the most sensitive sex/species is  $8 \times 10^{-2}$  per ppm [ $2 \times 10^{-5}$  per  
31  $\mu\text{g}/\text{m}^3$ ], based on kidney adenomas and carcinomas reported by Maltoni et al. (1986) for male  
32 Sprague-Dawley rats. Leukemias and Leydig cell tumors were also increased in these rats, and,  
33 although a combined analysis for these tumor types which incorporated the different site-specific  
34 preferred dose-metrics was not performed, the result of such an analysis is expected to be  
35 similar, about  $9 \times 10^{-2}$  per ppm [ $2 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ]. The next most sensitive sex/species from

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 the inhalation bioassays is the female mouse, for which lymphomas were reported by Henschler  
2 et al. (1980); these data yield a unit risk estimate of  $1.0 \times 10^{-2}$  per ppm [ $2 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ]. In  
3 addition, the 90% confidence intervals (i.e., 5 to 95% bounds) reported in Table 5-41 for male rat  
4 kidney tumors from Maltoni et al. (1986) and female mouse lymphomas from Henschler et al.  
5 (1980), derived from the quantitative analysis of PBPK model uncertainty, both included the  
6 estimate based on human data of  $2 \times 10^{-2}$  per ppm. Furthermore, PBPK model-based route-to-  
7 route extrapolation of the results for the most sensitive sex/species from the oral bioassays,  
8 kidney tumors in male Osborne-Mendel rats and testicular tumors in Marshall rats (NTP, 1988),  
9 leads to inhalation unit risk estimates of  $2 \times 10^{-1}$  per ppm [ $3 \times 10^{-5}$  per  $\mu\text{g}/\text{m}^3$ ] and  $4 \times 10^{-2}$  per  
10 ppm [ $8 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ], respectively, with the preferred estimate based on human data falling  
11 within the route-to-route extrapolation of the 90% confidence intervals reported in Table 5-42.  
12 Finally, for all these estimates, the ratios of BMDs to the BMDLs did not exceed a value of 3,  
13 indicating that the uncertainties in the dose-response modeling for determining the POD in the  
14 observable range are small.

15 Although there are uncertainties in these various estimates, confidence in the proposed inhalation  
16 unit risk estimate of  $2 \times 10^{-2}$  per ppm [ $4 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ ], based on human kidney cancer risks  
17 reported by Charbotel et al. (2006) and adjusted for potential risk for tumors at multiple sites (as  
18 summarized above in Section 6.1.4), is further increased by the similarity of this estimate to  
19 estimates based on multiple rodent data sets. Application of the ADAFs for the kidney cancer  
20 risks, due to the weight of evidence supporting a mutagenic MOA for this endpoint, is  
21 summarized below in Section 6.2.2.5.

22

### **6.2.2.1.3. Oral Slope Factor Estimate (rodent: see Section 5.2.1.3; human: see Section 5.2.2.3)**

23 The oral slope factor for TCE is defined as a plausible upper bound lifetime extra risk of cancer  
24 from chronic ingestion of TCE per mg/kg-day oral dose. The oral slope factor is  $4.64 \times 10^{-2}$  per  
25 mg/kg-day ( **$5 \times 10^{-2}$  per mg/kg-day** rounded to one significant figure), resulting from PBPK  
26 model-based route-to-route extrapolation of the inhalation unit risk estimate based on the human  
27 kidney cancer risks reported in Charbotel et al. (2006) and adjusted for potential risk for tumors  
28 at multiple sites. This estimate is based on good-quality human data, thus avoiding uncertainties  
29 inherent in interspecies extrapolation. In addition, uncertainty in the PBPK model-based route-  
30 to-route extrapolation is relatively low (Chiu, 2006; Chiu and White, 2006). In this particular  
31 case, extrapolation using different dose-metrics yielded expected population mean risks within  
32 about a twofold range, and, for any particular dose-metric, the 95% confidence interval for the

*This document is a draft for review purposes only and does not constitute Agency policy.*

1 extrapolated population mean risks for each site spanned a range of no more than about  
2 threefold.

3 This value is supported by oral slope factor estimates from multiple rodent bioassays, the most  
4 sensitive of which range from  $3 \times 10^{-2}$  to  $3 \times 10^{-1}$  per mg/kg-day. From the oral bioassays  
5 selected for analysis in Section 5.2.1.1, and using the preferred PBPK model-based dose-metrics,  
6 the oral slope factor estimate for the most sensitive sex/species is  $3 \times 10^{-1}$  per mg/kg-day, based  
7 on kidney tumors in male Osborne-Mendel rats (NTP, 1988). The oral slope factor estimate for  
8 testicular tumors in male Marshall rats (NTP, 1988) is somewhat lower at  $7 \times 10^{-2}$  per mg/kg-  
9 day. The next most sensitive sex/species result from the oral studies is for male mouse liver  
10 tumors (NCI, 1976), with an oral slope factor estimate of  $3 \times 10^{-2}$  per mg/kg-day. In addition,  
11 the 90% confidence intervals reported in Table 5-42 for male Osborne-Mendel rat kidney tumors  
12 (NTP, 1988), male F344 rat kidney tumors (NTP, 1990), and male Marshall rat testicular tumors  
13 (NTP, 1988), derived from the quantitative analysis of PBPK model uncertainty, all included the  
14 estimate based on human data of  $5 \times 10^{-2}$  per mg/kg-day, while the upper 95% confidence bound  
15 for male mouse liver tumors from NCI (1976) was slightly below this value at  $4 \times 10^{-2}$  per  
16 mg/kg-day. Furthermore, PBPK model-based route-to-route extrapolation of the most sensitive  
17 endpoint from the inhalation bioassays, male rat kidney tumors from Maltoni et al. (1986), leads  
18 to an oral slope factor estimate of  $1 \times 10^{-1}$  per mg/kg-day, with the preferred estimate based on  
19 human data falling within the route-to-route extrapolation of the 90% confidence interval  
20 reported in Table 5-41. Finally, for all these estimates, the ratios of BMDs to the BMDLs did  
21 not exceed a value of three, indicating that the uncertainties in the dose-response modeling for  
22 determining the POD in the observable range are small.

23 Although there are uncertainties in these various estimates, confidence in the proposed oral slope  
24 factor estimate of  $5 \times 10^{-2}$  per mg/kg-day, resulting from PBPK model-based route-to-route  
25 extrapolation of the inhalation unit risk estimate based on the human kidney cancer risks reported  
26 in Charbotel et al. (2006) and adjusted for potential risk for tumors at multiple sites (as  
27 summarized above), is further increased by the similarity of this estimate to estimates based on  
28 multiple rodent data sets. Application of the ADAFs for the kidney cancer risks, due to the  
29 weight of evidence supporting a mutagenic MOA for this endpoint, is summarized below in  
30 Section 6.2.2.5.

31

#### 6.2.2.1.4. Uncertainties in Cancer Dose-Response Assessment

#### 6.2.2.1.5. Uncertainties in estimates based on human epidemiologic data (see Section 5.2.2.1.3)

1 All risk assessments involve uncertainty, as study data are extrapolated to make inferences about  
2 potential effects in humans from environmental exposure. The values for the slope factor and  
3 unit risk estimates are based on good quality human data, which avoids interspecies  
4 extrapolation, one of the major sources of uncertainty in quantitative cancer risk estimates.

5 A remaining major uncertainty in the unit risk estimate for RCC incidence derived from  
6 the Charbotel et al. (2006) study is the extrapolation from occupational exposures to lower  
7 environmental exposures. There was some evidence of a contribution to increased RCC risk  
8 from peak exposures; however, there remained an apparent dose-response relationship for RCC  
9 risk with increasing cumulative exposure without peaks, and the OR for exposure with peaks  
10 compared to exposure without peaks was not significantly elevated (Charbotel et al., 2006).  
11 Although the actual exposure-response relationship at low exposure levels is unknown, the  
12 conclusion that a mutagenic MOA is operative for TCE-induced kidney tumors supports the  
13 linear low-dose extrapolation that was used (U.S. EPA, 2005c). Additional support for use of  
14 linear extrapolation is discussed above in Section 6.2.2.1.

15 Another source of uncertainty is the dose-response model used to model the study data to  
16 estimate the POD. A weighted linear regression across the categorical ORs was used to obtain a  
17 slope estimate; use of a linear model in the observable range of the data is often a good general  
18 approach for human data because epidemiological data are frequently too limited (the Charbotel  
19 et al. [(2006)] study had 86 RCC cases, 37 of which had TCE exposure) to clearly identify an  
20 alternate model (U.S. EPA, 2005c). The ratio of the maximum likelihood estimate of the  
21 effective concentration for a 1% response ( $EC_{01}$ ) to the  $LEC_{01}$ , which gives some indication of  
22 the statistical uncertainties in the dose-response modeling, was about a factor of 2.

23 A further source of uncertainty is the retrospective estimation of TCE exposures in the Charbotel  
24 et al. (2006) study. This case-control study was conducted in the Arve Valley in France, a region  
25 with a high concentration of screw cutting workshops using TCE and other degreasing agents.  
26 Since the 1960s, occupational physicians of the region have collected a large quantity of well-  
27 documented measurements, including TCE air concentrations and urinary metabolite levels  
28 (Fevotte et al., 2006). The study investigators conducted a comprehensive exposure assessment  
29 to estimate cumulative TCE exposures for the individual study subjects, using a detailed  
30 occupational questionnaire with a customized task-exposure matrix for the screw-cutting workers  
31 and a more general occupational questionnaire for workers exposed to TCE in other industries

1 (Fevotte et al., 2006). The exposure assessment also attempted to take dermal exposure from  
2 hand-dipping practices into account by equating it with an equivalent airborne concentration  
3 based on biological monitoring data. Despite the appreciable effort of the investigators,  
4 considerable uncertainty associated with any retrospective exposure assessment is inevitable, and  
5 some exposure misclassification is unavoidable. Such exposure misclassification was most  
6 likely for the 19 deceased cases and their matched controls, for which proxy respondents were  
7 used, and for exposures outside the screw-cutting industry. The exposure estimates from the  
8 RCC study of Moore et al. (2010) were not considered to be as quantitatively accurate as those of  
9 Charbotel et al. (2006) and so were not used for derivation of a unit risk estimate (see Section  
10 5.2.2); nonetheless, it should be noted that these exposure estimates are substantially lower than  
11 those of Charbotel et al. (2006) for comparable OR estimates. If the exposure estimates for  
12 Charbotel et al. (2006) are overestimated, as suggested by the exposure estimates from Moore et  
13 al. (2010), the slope of the linear regression model, and hence the unit risk estimate, would be  
14 correspondingly underestimated.

15 Another source of uncertainty in the Charbotel et al. (2006) study is the possible influence of  
16 potential confounding or modifying factors. This study population, with a high prevalence of  
17 metal-working, also had relatively high prevalences of exposure to petroleum oils, cadmium,  
18 petroleum solvents, welding fumes, and asbestos (Fevotte et al., 2006). Other exposures  
19 assessed included other solvents (including other chlorinated solvents), lead, and ionizing  
20 radiation. None of these exposures was found to be significantly associated with RCC at a  
21  $p = 0.05$  significance level. Cutting fluids and other petroleum oils were associated with RCC at  
22 a  $p = 0.1$  significance level; however, further modeling suggested no association with RCC when  
23 other significant factors were taken into account (Charbotel et al., 2006). Moreover, a review of  
24 other studies suggested that potential confounding from cutting fluids and other petroleum oils is  
25 of minimal concern (see Section 4.4.2.3). Nonetheless, a sensitivity analysis was conducted  
26 using the OR estimates further adjusted for cutting fluids and other petroleum oils from the  
27 unpublished report by Charbotel et al. (2005), and an essentially identical unit risk estimate of  
28  $5.46 \times 10^{-3}$  per ppm was obtained. In addition, the medical questionnaire included familial  
29 kidney disease and medical history, such as kidney stones, infection, chronic dialysis,  
30 hypertension, and use of antihypertensive drugs, diuretics, and analgesics. Body mass index  
31 (BMI) was also calculated, and lifestyle information such as smoking habits and coffee  
32 consumption was collected. Univariate analyses found high levels of smoking and BMI to be  
33 associated with increased odds of RCC, and these two variables were included in the conditional  
34 logistic regressions. Thus, although impacts of other factors are possible, this study took great  
35 pains to attempt to account for potential confounding or modifying factors.

1 Some other sources of uncertainty associated with the epidemiological data are the dose-metric  
2 and lag period. As discussed above, there was some evidence of a contribution to increased RCC  
3 risk from peak TCE exposures; however, there appeared to be an independent effect of  
4 cumulative exposure without peaks. Cumulative exposure is considered a good measure of total  
5 exposure because it integrates exposure (levels) over time. If there is a contributing effect of  
6 peak exposures, not already taken into account in the cumulative exposure metric, the linear  
7 slope may be overestimated to some extent. Sometimes cancer data are modeled with the  
8 inclusion of a lag period to discount more recent exposures not likely to have contributed to the  
9 onset of cancer. In an unpublished report, Charbotel et al. (2005) also present the results of a  
10 conditional logistic regression with a 10-year lag period, and these results are very similar to the  
11 unlagged results reported in their published paper, suggesting that the lag period might not be an  
12 important factor in this study.

13 Some additional sources of uncertainty are not so much inherent in the exposure-response  
14 modeling or in the epidemiologic data themselves but, rather, arise in the process of obtaining  
15 more general Agency risk estimates from the epidemiologic results. EPA cancer risk estimates  
16 are typically derived to represent an upper bound on increased risk of cancer incidence for all  
17 sites affected by an agent for the general population. From experimental animal studies, this is  
18 accomplished by using tumor incidence data and summing across all the tumor sites that  
19 demonstrate significantly increased incidences, customarily for the most sensitive sex and  
20 species, to attempt to be protective of the general human population. However, in estimating  
21 comparable risks from the Charbotel et al. (2006) epidemiologic data, certain limitations are  
22 encountered. For one thing, these epidemiology data represent a geographically limited (Arve  
23 Valley, France) and likely not very diverse population of working adults. Thus, there is  
24 uncertainty about the applicability of the results to a more diverse general population.  
25 Additionally, the Charbotel et al. (2006) study was a study of RCC only, and so the risk estimate  
26 derived from it does not represent all the tumor sites that may be affected by TCE. This  
27 uncertainty was addressed by adjusting the RCC estimate to multiple sites, but there are also  
28 uncertainties related to the assumptions inherent in the calculations for this adjustment. As  
29 discussed in Section 5.2.2.2, adequate quantitative dose-response data were only available for  
30 one cancer type in humans, so other human data were used to adjust the estimate derived for  
31 RCC to include risk for other cancers with substantial human evidence of hazard (lymphoma and  
32 liver cancer). The relative contributions to extra risk (for cancer incidence) were calculated from  
33 two different data sets to derive an adjustment factor. The first calculation is based on the results  
34 of the meta-analyses for the three tumor types; the second calculation is based on the results of  
35 the Raaschou-Nielsen et al. (2003) study, the largest single study by far with RR estimates for all



1 three tumor types. The fact that the calculations based on two different data sets yielded  
2 comparable values for the adjustment factor (both within 25% of the selected factor of 4)  
3 provides more robust support for the use of the factor of 4. Additional uncertainties pertain to  
4 the weight of evidence supporting the association of TCE exposure with increased risk of cancer  
5 for the three cancer types. As discussed in Section 4.11.2, it is concluded that the weight of  
6 evidence for kidney cancer is sufficient to classify TCE as “carcinogenic to humans.” It is also  
7 concluded that there is strong evidence that TCE causes lymphoma as well, although the  
8 evidence for liver cancer was more limited. In addition, the rodent studies demonstrate clear  
9 evidence of multisite carcinogenicity, with tumor types including those for which associations  
10 with TCE exposure are observed in human studies, i.e., liver and kidney cancers and lymphomas.  
11 Overall, the evidence is sufficiently persuasive to support the use of the adjustment factor of 4  
12 based on these three cancer types. Alternatively, if one were to use the factor based only on the  
13 two cancer types with the strongest human evidence, the cancer inhalation unit risk estimate  
14 would be only slightly reduced (25%).  
15 Finally, the value for the oral slope factor estimate was based on route-to-route extrapolation of  
16 the inhalation unit risk based on human data using predictions from the PBPK model. Because  
17 different internal dose-metrics are preferred for each target tissue site, a separate route-to-route  
18 extrapolation was performed for each site-specific slope factor estimate. As discussed above,  
19 uncertainty in the PBPK model-based route-to-route extrapolation is relatively low (Chiu, 2006;  
20 Chiu and White, 2006). In this particular case, extrapolation using different dose-metrics yielded  
21 expected population mean risks within about a twofold range, and, for any particular dose-  
22 metric, the 95% confidence interval for the extrapolated population mean risks for each site  
23 spanned a range of no more than about threefold.

24

#### **6.2.2.1.6. Uncertainties in estimates based on rodent bioassays (see Section 5.2.1.4)**

25 With respect to rodent-based cancer risk estimates, the cancer risk is typically estimated from the  
26 total cancer burden from all sites that demonstrate an increased tumor incidence for the most  
27 sensitive experimental species and sex. It is expected that this approach is protective of the  
28 human population, which is more diverse but is exposed to lower exposure levels. In the case of  
29 TCE, the impact of selection of the bioassay is limited, since, as discussed in Sections 5.2.1.3  
30 and 5.2.3, estimates based on the two or three most sensitive bioassays are within an order of  
31 magnitude of each other, and are consistent across routes of exposure when extrapolated using  
32 the PBPK model.

1 Another source of uncertainty in the TCE rodent-based cancer risk estimates is interspecies  
2 extrapolation. Several plausible PBPK model-based dose-metrics were used for extrapolation of  
3 toxicokinetics, but the cancer slope factor and unit risk estimates obtained using the preferred  
4 dose-metrics were generally similar (within about threefold) to those derived using default  
5 dosimetry assumptions, with the exception of the bioactivated DCVC dose-metric for rat kidney  
6 tumors and the metric for the amount of TCE oxidized in the respiratory tract for mouse lung  
7 tumors occurring from oral exposure. However, there is greater biological support for these  
8 selected dose-metrics. The uncertainty in the PBPK model predictions themselves was analyzed  
9 quantitatively through an analysis of the impact of parameter uncertainties in the PBPK model.  
10 The 95% lower bounds on the BMD including parameter uncertainties in the PBPK model were  
11 no more than fourfold lower than those based on central estimates of the PBPK model  
12 predictions. The greatest uncertainty was for slope factors and unit risks derived from rat kidney  
13 tumors, primarily reflecting the substantial uncertainty in the rat internal dose and in the  
14 extrapolation of GSH conjugation from rodents to humans.

15 Regarding low-dose extrapolation, a key consideration in determining what extrapolation  
16 approach to use is the MOA(s). However, MOA data are lacking or limited for each of the  
17 cancer responses associated with TCE exposure, with the exception of the kidney tumors. For  
18 the kidney tumors, the weight of the available evidence supports the conclusion that a mutagenic  
19 MOA is operative; this MOA supports linear low-dose extrapolation. For the other TCE-induced  
20 tumors, the data either support a complex MOA or are inadequate to specify the key events and  
21 MOAs involved. When the MOA(s) cannot be clearly defined, EPA generally uses a linear  
22 approach to estimate low-dose risk (U.S. EPA, 2005c), based on the general principles discussed  
23 above.

24 With respect to uncertainties in the dose-response modeling, the two-step approach of modeling  
25 only in the observable range, as put forth in EPA's *Guidelines for Carcinogen Risk Assessment*  
26 (U.S. EPA, 2005c), is designed in part to minimize model dependence. The ratios of the BMDs  
27 to the BMDLs, which give some indication of the statistical uncertainties in the dose-response  
28 modeling, did not exceed a value of 2.5 for all the primary analyses used in this assessment.  
29 Thus, overall, modeling uncertainties in the observable range are considered to be minimal.  
30 Some additional uncertainty is conveyed by uncertainties in the survival adjustments made to  
31 some of the bioassay data; however, a comparison of the results of two different survival  
32 adjustment methods suggest that their impact is minimal relative to the uncertainties already  
33 discussed.

#### 6.2.2.1.7. Application of Age-Dependent Adjustment Factors (see Section 5.2.3.3)

1 When there is sufficient weight of evidence to conclude that a carcinogen operates through a  
2 mutagenic MOA, and in the absence of chemical-specific data on age-specific susceptibility,  
3 EPA's *Supplemental Guidance for Assessing Susceptibility from Early-Life Exposure to*  
4 *Carcinogens* (U.S. EPA, 2005c) recommends the application of default ADAFs to adjust for  
5 potential increased susceptibility from early-life exposure. See the *Supplemental Guidance* for  
6 detailed information on the general application of these adjustment factors. In brief, the  
7 *Supplemental Guidance* establishes ADAFs for three specific age groups. The current ADAFs  
8 and their age groupings are 10 for <2 years, 3 for 2 to <16 years, and 1 for 16 years and above  
9 (U.S. EPA, 2005c). For risk assessments based on specific exposure assessments, the 10-fold  
10 and threefold adjustments to the slope factor or unit risk estimates are to be combined with age-  
11 specific exposure estimates when estimating cancer risks from early-life (<16 years age)  
12 exposure.

13 In the case of TCE, the inhalation unit risk and oral slope factor estimates reflect lifetime risk for  
14 cancer at multiple sites, and a mutagenic MOA has been established for one of these sites, the  
15 kidney. In addition, as discussed in Section 4.10, inadequate TCE-specific data exists to quantify  
16 early-life susceptibility to TCE carcinogenicity; therefore, as recommended in the *Supplemental*  
17 *Guidance*, the default ADAFs are used. As illustrated in the example calculations in  
18 Sections 5.2.3.3.1 and 5.2.3.3.2, application of the default ADAFs to the kidney cancer  
19 inhalation unit risk and oral slope factor estimates for TCE is likely to have minimal impact on  
20 the total cancer risk except when exposure is primarily during early life.

21 In addition to the uncertainties discussed above for the inhalation and oral total cancer unit risk  
22 and slope factor estimates, there are uncertainties in the application of ADAFs to adjust for  
23 potential increased early-life susceptibility. The adjustment is made only for the kidney cancer  
24 component of total cancer risk because that is the tumor type for which the weight of evidence  
25 was sufficient to conclude that TCE-induced carcinogenesis operates through a mutagenic MOA.  
26 However, it may be that TCE operates through a mutagenic MOA for other tumor types as well  
27 or that it operates through other MOAs that might also convey increased early-life susceptibility.  
28 Additionally, the ADAFs from the 2005 Supplemental Guidance are not specific to TCE, and it  
29 is uncertain to what extent they reflect increased early-life susceptibility to kidney cancer from  
30 exposure to TCE, if increased early-life susceptibility occurs.

31

### 6.3. OVERALL CHARACTERIZATION OF TCE HAZARD AND DOSE RESPONSE

1           There is substantial potential for human exposure to TCE, as it has a widespread presence  
2 in ambient air, indoor air, soil, and groundwater. At the same time, humans are likely to be  
3 exposed to a variety of compounds that are either metabolites of TCE or which have common  
4 metabolites or targets of toxicity. Once exposed, humans, as well as laboratory animal species,  
5 rapidly absorb TCE, which is then distributed to tissues via systemic circulation, extensively  
6 metabolized, and then excreted primarily in breath as unchanged TCE or CO<sub>2</sub>, or in urine as  
7 metabolites.

8           Based on the available human epidemiologic data and experimental and mechanistic  
9 studies, it is concluded that TCE poses a potential human health hazard for noncancer toxicity to  
10 the central nervous system, the kidney, the liver, the immune system, the male reproductive  
11 system, and the developing fetus. The evidence is more limited for TCE toxicity to the  
12 respiratory tract and female reproductive system. Following EPA (2005c) *Guidelines for*  
13 *Carcinogen Risk Assessment*, TCE is characterized as “*carcinogenic to humans*” by all routes of  
14 exposure. This conclusion is based on convincing evidence of a causal association between TCE  
15 exposure in humans and kidney cancer. The human evidence of carcinogenicity from  
16 epidemiologic studies of TCE exposure is compelling for lymphoma, but less convincing than  
17 for kidney cancer, and more limited for liver and biliary tract cancer. Less human evidence is  
18 found for an association between TCE exposure and other types of cancer, including bladder,  
19 esophageal, prostate, cervical, breast, and childhood leukemia, breast. Further support for the  
20 characterization of TCE as “*carcinogenic to humans*” by all routes of exposure is derived from  
21 positive results in multiple rodent cancer bioassays in rats and mice of both sexes, similar  
22 toxicokinetics between rodents and humans, mechanistic data supporting a mutagenic MOA for  
23 kidney tumors, and the lack of mechanistic data supporting the conclusion that any of the  
24 MOA(s) for TCE-induced rodent tumors are irrelevant to humans.

25           As TCE toxicity and carcinogenicity are generally associated with TCE metabolism,  
26 susceptibility to TCE health effects may be modulated by factors affecting toxicokinetics,  
27 including lifestage, gender, genetic polymorphisms, race/ethnicity, preexisting health status,  
28 lifestyle, and nutrition status. In addition, while some of these factors are known risk factors for  
29 effects associated with TCE exposure, it is not known how TCE interacts with known risk factors  
30 for human diseases.

31           For noncancer effects, the most sensitive types of effects, based either on human  
32 equivalent concentrations/doses or on candidate RfCs/RfDs, appear to be developmental, kidney,  
33 and immunological (adult and developmental) effects. The neurological and reproductive effects  
34 appear to be about an order of magnitude less sensitive, with liver effects another two orders of

1 magnitude less sensitive. The RfC of **0.0004 ppm** (0.4 ppb or 2  $\mu\text{g}/\text{m}^3$ ) is based on route-to-  
2 route extrapolated results from oral studies for the critical effects of heart malformations (rats)  
3 and immunotoxicity (mice). This RfC value is further supported by route-to-route extrapolated  
4 results from an oral study of toxic nephropathy (rats). Similarly, the RfD for noncancer effects  
5 of **0.0005 mg/kg-day** is based on the critical effects of heart malformations (rats), adult  
6 immunological effects (mice), and developmental immunotoxicity (mice), all from oral studies.  
7 This RfD value is further supported by results from an oral study for the effect of toxic  
8 nephropathy (rats) and route-to-route extrapolated results from an inhalation study for the effect  
9 of increased kidney weight (rats). There is high confidence in these noncancer reference values,  
10 as they are supported by moderate- to high-confidence estimates for multiple effects from  
11 multiple studies.

12 For cancer, the inhalation unit risk is  **$2 \times 10^{-2}$  per ppm** [ **$4 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$** ], based on  
13 human kidney cancer risks reported by Charbotel et al. (2006) and adjusted, using human  
14 epidemiologic data, for potential risk for tumors at multiple sites. The oral slope factor for  
15 cancer is  **$5 \times 10^{-2}$  per mg/kg-day**, resulting from PBPK model-based route-to-route extrapolation  
16 of the inhalation unit risk estimate based on the human kidney cancer risks reported in Charbotel  
17 et al. (2006) and adjusted, using human epidemiologic data, for potential risk for tumors at  
18 multiple sites. There is high confidence in these unit risks for cancer, as they are based on good  
19 quality human data, as well as being similar to unit risk estimates based on multiple rodent  
20 bioassays. There is both sufficient weight of evidence to conclude that TCE operates through a  
21 mutagenic MOA for kidney tumors and a lack of TCE-specific quantitative data on early-life  
22 susceptibility. Generally, the application of age-dependent adjustment factors (ADAFs) is  
23 recommended when assessing cancer risks for a carcinogen with a mutagenic MOA. However,  
24 because the ADAF adjustment applies only to the kidney cancer component of the total risk  
25 estimate, it is likely to have a minimal impact on the total cancer risk except when exposures are  
26 primarily during early life.

*This document is a draft for review purposes only and does not constitute Agency policy.*

## 7. REFERENCES

- 1 Abbas, R. and Fisher, J. W. (1997). A physiologically based pharmacokinetic model for  
2 trichloroethylene and its metabolites, chloral hydrate, trichloroacetate, dichloroacetate,  
3 trichloroethanol, and trichloroethanol glucuronide in B6C3F1 mice. *Toxicol Appl*  
4 *Pharmacol*, 147, 15-30. <http://dx.doi.org/10.1006/taap.1997.8190>
- 5 Abbas, R. R.; Seckel, C. S.; Kidney, J. K.; Fisher, J. W. (1996). Pharmacokinetic analysis of  
6 chloral hydrate and its metabolism in B6C3F1 mice. *Drug Metab Dispos*, 24, 1340-1346.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/8971140>
- 8 Abbas, R.; Seckel, C. S.; MacMahon, K. L.; Fisher, J. W. (1997). Determination of kinetic rate  
9 constants for chloral hydrate, trichloroethanol, trichloroacetic acid and dichloroacetic  
10 acid: A physiologically based modeling approach. *Toxicologist*, 36, 32-33.
- 11 Abbate, C.; Giorgianni, C.; Munao, F.; Brecciaroli, R. (1993). Neurotoxicity induced by  
12 exposure to toluene: an electrophysiologic study. *Int Arch Occup Environ Health*, 64,  
13 389-392.
- 14 Abraham, D.; Patel, P.; Cooper, A. (1995). Isolation from rat kidney of a cytosolic high  
15 molecular weight cysteine-S-conjugate beta-lyase with activity toward leukotriene E4. *J*  
16 *Biol Chem*, 270, 180-188. <http://www.ncbi.nlm.nih.gov/pubmed/7814371>
- 17 Abraham, D.; Thomas, R.; Cooper, A. (1995). Glutamine transaminase K is not a major cysteine  
18 S-conjugate beta-lyase of rat kidney mitochondria: evidence that a high-molecular weight  
19 enzyme fulfills this role. *Mol Pharmacol*, 48, 855-860.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/7476916>
- 21 Adachi, H.; Kubota, I.; Okamura, N.; Iwata, H.; Tsujimoto, M.; Nakazato, H., . . . Noguchi, T.  
22 (1989). Purification and characterization of human microsomal dipeptidase. *J Biochem*,  
23 105, 957-961. <http://www.ncbi.nlm.nih.gov/pubmed/2768222>
- 24 Adami, H. O.; Chow, W. H.; Nyrén, O.; Berne, C.; Linet, M. S.; Ekblom, A., . . . Fraumeni, J. F.  
25 (1996). Excess risk of primary liver cancer in patients with diabetes mellitus. *J Natl*  
26 *Cancer Inst*, 88, 1472-1477. <http://www.ncbi.nlm.nih.gov/pubmed/8841022>
- 27 Adams, C.; Keil, D.; Meyers, K.; Eudaly, A.; Smythe, J.; Eudaly, J., . . . Peden-Adams, M. M.  
28 (2003). Lifetime exposure to trichloroethylene (TCE) modulates immune function.  
29 *Toxicologist*, 72, 375.
- 30 Adams, L. A. and Lindor, K. D. (2007). Nonalcoholic fatty liver disease. *Ann Epidemiol*, 17,  
31 863-869. <http://dx.doi.org/10.1016/j.annepidem.2007.05.013>
- 32 Adgate, J. L.; Church, T. R.; Ryan, A. D.; Ramachandran, G.; Fredrickson, A. L.; Stock, T. H., . . .  
33 . Sexton, K. (2004). Outdoor, indoor, and personal exposure to VOCs in children.  
34 *Environ Health Perspect*, 112, 1386-1392. <http://dx.doi.org/10.1289/ehp.7107>
- 35 Adgate, J.; Eberly, L.; Stroebel, C.; Pellizzari, E.; Sexton, K. (2004). Personal, indoor, and  
36 outdoor VOC exposures in a probability sample of children. *J Expo Anal Environ*  
37 *Epidemiol*, 14, S4-S13. <http://dx.doi.org/10.1038/sj.jea.7500353>
- 38 ADHS. (Arizona Department of Health Services). (1990). Incidence study of childhood cancer in  
39 Maricopa County: 1965-1986. Phoenix, AZ: Author.
- 40 ADHS. (Arizona Department of Health Services). (1995). Update of the incidence of childhood  
41 cancers and testicular cancer in Southwest Tucson: 1987-1991. Phoenix, AZ: Author.

DRAFT– DO NOT QUOTE OR CITE

- 1 Adler, I. D. (1993). Synopsis of the in vivo results obtained with the 10 known or suspected  
2 aneugens tested in the CEC collaborative study. *DNA Repair*, 287, 131-137.  
3 [http://dx.doi.org/10.1016/0027-5107\(93\)90152-6](http://dx.doi.org/10.1016/0027-5107(93)90152-6)
- 4 Adli, M. and Baldwin, A. S. (2006). IKK-i/IKKepsilon controls constitutive, cancer cell-  
5 associated NF-kappaB activity via regulation of Ser-536 p65/RelA phosphorylation. *J*  
6 *Biol Chem*, 281, 26976-26984. <http://dx.doi.org/10.1074/jbc.M603133200>
- 7 Aickin, M. (2004). Bayes without priors. *J Clin Epidemiol*, 57, 4-13.  
8 [http://dx.doi.org/10.1016/S0895-4356\(03\)00251-8](http://dx.doi.org/10.1016/S0895-4356(03)00251-8)
- 9 Aickin, M.; Chapin, C. A.; Flood, T. J.; Englender, S. J.; Caldwell, G. G. (1992). Assessment of  
10 the spatial occurrence of childhood leukaemia mortality using standardized rate ratios  
11 with a simple linear Poisson model. *Int J Epidemiol*, 21, 649-655.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/1521967>
- 13 Aikata, H.; Takaishi, H.; Kawakami, Y.; Takahashi, S.; Kitamoto, M.; Nakanishi, T., . . . Ide, T.  
14 (2000). Telomere reduction in human liver tissues with age and chronic inflammation.  
15 *Exp Cell Res*, 256, 578-582. <http://dx.doi.org/10.1006/excr.2000.4862>
- 16 Akira, S. and Takeda, K. (2004). Toll-like receptor signalling. *Nat Rev Immunol*, 4, 499-511.  
17 <http://dx.doi.org/10.1038/nri1391>
- 18 Akiyama, T. E.; Nicol, C. J.; Fievet, C.; Staels, B.; Ward, J. M.; Auwerx, J., . . . Peters, J. M.  
19 (2001). Peroxisome proliferator-activated receptor-alpha regulates lipid homeostasis, but  
20 is not associated with obesity: studies with congenic mouse lines. *J Biol Chem*, 276,  
21 39088-39093. <http://dx.doi.org/10.1074/jbc.M107073200>
- 22 Albanese, R.; Banks, H.; Evans, M.; Potter, L. (2002). Physiologically based pharmacokinetic  
23 models for the transport of trichloroethylene in adipose tissue. *Bull Math Biol*, 64, 97-  
24 131. <http://dx.doi.org/10.1006/bulm.2001.0268>
- 25 Albee, R.; Nitschke, K.; Mattsson, J.; Stebbins, K. (1997). Dichloroacetylene: effects on the rat  
26 trigeminal nerve somatosensory evoked potential. *Neurotoxicol Teratol*, 19, 27-37.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/9088008>
- 28 Albee, R.; Spencer, P.; Johnson, K.; Bradley, G.; Marable, B.; Wilmer, J.; Mattsson, J. (2006).  
29 Lack of trigeminal nerve toxicity in rats exposed to trichloroethylene vapor for 13 weeks.  
30 *Int J Toxicol*, 25, 531-540. <http://dx.doi.org/10.1080/10915810600972678>
- 31 Alberati-Giani, D.; Malherbe, P.; Kohler, C.; Lang, G.; Kiefer, V.; Lahm, H. W.; Cesura, A. M.  
32 (1995). Cloning and characterization of a soluble kynurenine aminotransferase from rat  
33 brain: Identity with kidney cysteine conjugate [beta]-lyase. *J Neurochem*, 64, 1448-1455.  
34 <http://dx.doi.org/10.1046/j.1471-4159.1995.64041448.x>
- 35 Albers, J. W.; Wald, J. J.; Werner, R. A.; Franzblau, A.; Berent, S. (1999). Absence of  
36 polyneuropathy among workers previously diagnosed with solvent-induced toxic  
37 encephalopathy. *J Occup Environ Med*, 41, 500-509.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/10390702>
- 39 Albert, D.; Albert, A.; Vernace, M.; Sebastian, J.; Hsia, E. (2005). Analysis of a cluster of cases  
40 of Wegener granulomatosis. *J Clin Rheumatol*, 11, 188-193.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/16357755>
- 42 Albertini, S. (1990). Analysis of nine known or suspected spindle poisons for mitotic  
43 chromosome malsegregation using *Saccharomyces cerevisiae* D61.M. *Mutagenesis*, 5,  
44 453-459. <http://www.ncbi.nlm.nih.gov/pubmed/2263203>



DRAFT– DO NOT QUOTE OR CITE

- 1 Alexander, D. D.; Kelsh, M. A.; Mink, P. J.; Mandel, J. H.; Basu, R.; Weingart, M. (2007). A  
2 meta-analysis of occupational trichloroethylene exposure and liver cancer. *Int Arch*  
3 *Occup Environ Health*, 81, 127-143. <http://dx.doi.org/10.1007/s00420-007-0201-4>
- 4 Alexander, D. D.; Mink, P. J.; Adami, H. O.; Chang, E. T.; Cole, P.; Mandel, J. S.; Trichopoulos,  
5 D. (2007). The non-Hodgkin lymphomas: a review of the epidemiologic literature. *Int J*  
6 *Cancer*, 120 Suppl 12, 1-39. <http://dx.doi.org/10.1002/ijc.22719>
- 7 Alexander, D. D.; Mink, P. J.; Mandel, J. H.; Kelsh, M. A. (2006). A meta-analysis of  
8 occupational trichloroethylene exposure and multiple myeloma or leukaemia. *Occup Med*  
9 (Lond), 56, 485-493. <http://dx.doi.org/10.1093/occmed/kql083>
- 10 Alimov, A.; Kost-Alimova, M.; Liu, J.; Li, C.; Bergerheim, U.; Imreh, S., . . . Zabarovsky, E. R.  
11 (2000). Combined LOH/CGH analysis proves the existence of interstitial 3p deletions in  
12 renal cell carcinoma. *Oncogene*, 19, 1392-1399. <http://dx.doi.org/10.1038/sj.onc.1203449>
- 13 Allen, B. C. and Fisher, J. W. (1993). Pharmacokinetic modeling of trichloroethylene and  
14 trichloroacetic acid in humans. *Risk Anal*, 13, 71-86.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/8451462>
- 16 Allen, B. C.; Shipp, A. M.; Crump, K. S.; Kilian, B.; Hogg, M. L.; Tudor, J.; Keller, B. (1987).  
17 Investigation of cancer risk assessment methods (3 volumes plus summary report).  
18 Washington, D.C.: U.S. Environmental Protection Agency.
- 19 Allen, D.; Pearse, G.; Haseman, J.; Maronpot, R. (2004). Prediction of rodent carcinogenesis: an  
20 evaluation of prechronic liver lesions as forecasters of liver tumors in NTP  
21 carcinogenicity studies. *Toxicol Pathol*, 32, 393-401.  
22 <http://dx.doi.org/10.1080/01926230490440934>
- 23 Allen, J. W.; Collins, B. W.; Evansky, P. A. (1994). Spermatid micronucleus analyses of  
24 trichloroethylene and chloral hydrate effects in mice. *Mutat Res Lett*, 323, 81-88.  
25 [http://dx.doi.org/10.1016/0165-7992\(94\)90049-3](http://dx.doi.org/10.1016/0165-7992(94)90049-3)
- 26 Altmann, L.; Welge, P.; Mensing, T.; Lilienthal, H.; Voss, B.; Wilhelm, M. (2002). Chronic  
27 exposure to trichloroethylene affects neuronal plasticity in rat hippocampal slices.  
28 *Environ Toxicol Pharmacol*, 12, 157-167.
- 29 Altuntas, T.; Park, S.; Kharasch, E. (2004). Sulfoxidation of cysteine and mercapturic acid  
30 conjugates of the sevoflurane degradation product fluoromethyl-2,2-difluoro-1-  
31 (trifluoromethyl)vinyl ether (compound A). *Chem Res Toxicol*, 17, 435-445.  
32 <http://dx.doi.org/10.1021/tx034254k>
- 33 Amacher, D. E. and Zelljadt, I. (1983). The morphological transformation of Syrian hamster  
34 embryo cells by chemicals reportedly nonmutagenic to *Salmonella typhimurium*.  
35 *Carcinogenesis*, 4, 291-296.
- 36 Amet, Y.; Berthou, F.; Fournier, G.; Dreano, Y.; Bardou, L.; Cledes, J.; Menez, J. F. (1997).  
37 Cytochrome P450 4A and 2E1 expression in human kidney microsomes. *Biochem*  
38 *Pharmacol*, 53, 765-771. [http://dx.doi.org/10.1016/S0006-2952\(96\)00821-0](http://dx.doi.org/10.1016/S0006-2952(96)00821-0)
- 39 Ammini, C.; Fernandez-Canon, J.; Shroads, A.; Cornett, R.; Cheung, J.; James, M., . . .  
40 Stacpoole, P. (2003). Pharmacologic or genetic ablation of maleylacetoacetate isomerase  
41 increases levels of toxic tyrosine catabolites in rodents. *Biochem Pharmacol*, 66, 2029-  
42 2038. <http://www.ncbi.nlm.nih.gov/pubmed/14599561>
- 43 Andelman, J. B. (1985). Human exposures to volatile halogenated organic chemicals in indoor  
44 and outdoor air. *Environ Health Perspect*, 62, 313-318.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/4085436>

DRAFT– DO NOT QUOTE OR CITE

- 1 Anders, M. W. and Jakobson, I. (1985). Biotransformation of halogenated solvents. *Scand J*  
2 *Work Environ Health*, 11, 23-32. <http://www.ncbi.nlm.nih.gov/pubmed/3906866>
- 3 Andersen, M. E.; 3rd, C. H.; Gargas, M. L.; Smith, F. A.; Reitz, R. H. (1987). Physiologically  
4 based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol*  
5 *Appl Pharmacol*, 87, 185-205. [http://dx.doi.org/10.1016/0041-008X\(87\)90281-X](http://dx.doi.org/10.1016/0041-008X(87)90281-X)
- 6 Andersen, M. E.; Gargas, M. L.; 3rd, C. H.; Severyn, K. M. (1987). Quantitative evaluation of  
7 the metabolic interactions between trichloroethylene and 1,1-dichloroethylene in vivo  
8 using gas uptake methods. *Toxicol Appl Pharmacol*, 89, 149-157.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/3603553>
- 10 Anderson, B. J.; McKee, A. D.; Holford, N. H. (1997). Size, myths and the clinical  
11 pharmacokinetics of analgesia in paediatric patients. *Clin Pharmacokinet*, 33, 313-327.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/9391745>
- 13 Anderson, S. P.; Howroyd, P.; Liu, J.; Qian, X.; Bahnemann, R.; Swanson, C., . . . Corton, J. C.  
14 (2004). The transcriptional response to a peroxisome proliferator-activated receptor alpha  
15 agonist includes increased expression of proteome maintenance genes. *J Biol Chem*, 279,  
16 52390-52398. <http://dx.doi.org/10.1074/jbc.M409347200>
- 17 Andersson, L.; Haraldsson, B.; Johansson, C.; Barregard, L. (2008). Methodological issues on  
18 the use of urinary alpha-1-microglobuline in epidemiological studies. *Nephrol Dial*  
19 *Transplant*, 23, 1252-1256. <http://dx.doi.org/10.1093/ndt/gfm729>
- 20 Anna, C. H.; Maronpot, R. R.; Pereira, M. A.; Foley, J. F.; Malarkey, D. E.; Anderson, M. W.  
21 (1994). ras proto-oncogene activation in dichloroacetic acid-, trichloroethylene- and  
22 tetrachloroethylene-induced liver tumors in B6C3F1 mice. *Carcinogenesis*, 15, 2255-  
23 2261. <http://dx.doi.org/10.1093/carcin/15.10.2255>
- 24 Anti, M.; Marra, G.; Rapaccini, G. L.; Rumi, C.; Bussa, S.; Fadda, G., . . . Pompili, M. (1994).  
25 DNA ploidy pattern in human chronic liver diseases and hepatic nodular lesions. *Flow*  
26 *cytometric analysis on echo-guided needle liver biopsy. Cancer*, 73, 281-288.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/8293389>
- 28 Anttila, A.; Pukkala, E.; Sallmen, M.; Hernberg, S.; Hemminki, K. (1995). Cancer incidence  
29 among Finnish workers exposed to halogenated hydrocarbons. *J Occup Environ Med*, 37,  
30 797-806. <http://www.ncbi.nlm.nih.gov/pubmed/7552463>
- 31 Antti-Poika, M. (1982). Overall prognosis of patients with diagnosed chronic organic solvent  
32 intoxication. *Int Arch Occup Environ Health*, 51, 127-138.  
33 <http://dx.doi.org/10.1007/BF00378157>
- 34 Anway, M. D.; Cupp, A. S.; Uzumcu, M.; Skinner, M. K. (2005). Epigenetic transgenerational  
35 actions of endocrine disruptors and male fertility. *Science*, 308, 1466-1469.  
36 <http://dx.doi.org/10.1126/science.1108190>
- 37 Aranyi, C.; O'Shea, W. J.; Graham, J. A.; Miller, F. J. (1986). The effects of inhalation of  
38 organic chemical air contaminants on murine lung host defenses. *Fundam Appl Toxicol*,  
39 6, 713-720. [http://dx.doi.org/10.1016/0272-0590\(86\)90184-3](http://dx.doi.org/10.1016/0272-0590(86)90184-3)
- 40 Aratani, J.; Suzuki, H.; Hashimoto, K. (1993). Measurement of vibratory perception threshold  
41 (VPT) in workers exposed to organic solvents. *Environ Res*, 61, 357-361.  
42 <http://dx.doi.org/10.1006/enrs.1993.1080>
- 43 Arif, A. A. and Shah, S. M. (2007). Association between personal exposure to volatile organic  
44 compounds and asthma among US adult population. *Int Arch Occup Environ Health*, 80,  
45 711-719. <http://dx.doi.org/10.1007/s00420-007-0183-2>

DRAFT– DO NOT QUOTE OR CITE

- 1 Arito, H.; Takahashi, M.; Ishikawa, T. (1994). Effect of subchronic inhalation exposure to low-  
2 level trichloroethylene on heart rate and wakefulness-sleep in freely moving rats. *Sangyo*  
3 *Eiseigaku Zasshi*, 36, 1-8. <http://www.ncbi.nlm.nih.gov/pubmed/8126932>
- 4 Aronson, K.; Siemiatycki, J.; Dewar, R.; Gérin, M. (1996). Occupational risk factors for prostate  
5 cancer: Results from a case-control study in Montréal, Québec, Canada. *Am J Epidemiol*,  
6 143, 363-373. <http://www.ncbi.nlm.nih.gov/pubmed/8633620>
- 7 Arp, E. W.; Wolf, P. H.; Checkoway, H. (1983). Lymphocytic leukemia and exposures to  
8 benzene and other solvents in the rubber industry. *J Occup Med*, 25, 598-602.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/6886869>
- 10 Aryal, B. K.; Khuder, S. A.; Schaub, E. A. (2001). Meta-analysis of systemic sclerosis and  
11 exposure to solvents. *Am J Ind Med*, 40, 271-274. <http://dx.doi.org/10.1002/ajim.1098>
- 12 Asal, N. R.; Geyer, J. R.; Risser, D. R.; Lee, E. T.; Kadamani, S.; Cherng, N. (1988). Risk  
13 factors in renal cell carcinoma. II. Medical history, occupation, multivariate analysis, and  
14 conclusions. *Cancer Detect Prev*, 13, 263-279.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/3266567>
- 16 Asal, N. R.; Risser, D. R.; Kadamani, S.; Geyer, J. R.; Lee, E. T.; Cherng, N. (1988). Risk  
17 factors in renal cell carcinoma: I. Methodology, demographics, tobacco, beverage use,  
18 and obesity. *Cancer Detect Prev*, 11, 359-377.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/3390857>
- 20 Ashby, J.; Brady, A.; Elcombe, C. R.; Elliott, B. M.; Ishmael, J.; Odum, J., . . . Purchase, I. F.  
21 (1994). Mechanistically-based human hazard assessment of peroxisome proliferator-  
22 induced hepatocarcinogenesis. *Hum Exp Toxicol*, 13, S1-S117.  
23 <http://dx.doi.org/10.1177/096032719401300201>
- 24 Ashley, D. L.; Bonin, M. A.; Cardinali, F. L.; McCraw, J. M.; Wooten, J. V. (1994). Blood  
25 concentrations of volatile organic compounds in a nonoccupationally exposed US  
26 population and in groups with suspected exposure. *Clin Chem*, 40, 1401-1404.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/8013127>
- 28 Astrand, I. (1975). Uptake of solvents in the blood and tissues of man. A review. *Scand J Work*  
29 *Environ Health*, 1, 199-218. <http://www.ncbi.nlm.nih.gov/pubmed/779027>
- 30 Astrand, I. and Ovrum, P. (1976). Exposure to trichloroethylene I Uptake and distribution in  
31 man. *Scand J Work Environ Health*, 4, 199-211.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/1019594>
- 33 ATSDR. (1990). Toxicological profile for cis-, trans-1,2-dichloroethene.
- 34 ATSDR. (1996). Toxicological profile for 1,2-dichloroethene. Retrieved from  
35 <http://www.atsdr.cdc.gov/ToxProfiles/tp.asp?id=464&tid=82>.
- 36 ATSDR. (1997). Toxicological profile for tetrachloroethylene. Retrieved from  
37 <http://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=265&tid=48>.
- 38 ATSDR. (1997). Toxicological profile for trichloroethylene. Retrieved from  
39 <http://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=265&tid=48>.
- 40 ATSDR. (1998). Volatile organic compounds in drinking water and adverse pregnancy  
41 outcomes: United States Marine Corps Base, Camp Lejeune, North Carolina. Retrieved  
42 from <http://www.atsdr.cdc.gov/hs/lejeune/>.
- 43 ATSDR. (2003). Impact of trichloroethylene exposure on oral motor, speech, and hearing in  
44 children. Retrieved from file://L:\Lab\NrCEA\TCE Documents\TCE PDFs4\ATSDR2003  
45 TCE\_OralMotor\_Speech\_Hearing\_Children FinalReport.pdf.

DRAFT– DO NOT QUOTE OR CITE

- 1 ATSDR. (2003). Progress report: Survey of specific childhood cancers and birth defects among  
2 children whose mothers were pregnant while living at U.S. Marine Corps Base Camp  
3 Lejeune, North Carolina, 1968-1985. Retrieved from  
4 <http://www.atsdr.cdc.gov/sites/lejeune/lejeunehealthsurvey.pdf>.
- 5 ATSDR. (Agency for Toxic Substances and Disease Registry). (1995). Toxicological profile for  
6 1,1,1-trichloroethane. Atlanta, GA: U.S. Department of Health and Human Services.
- 7 ATSDR. (Agency for Toxic Substances and Disease Registry). (1996). Biennial report to  
8 congress (1991 and 1992). Atlanta, GA: U.S. Department of Health and Human  
9 Services.
- 10 ATSDR. (Agency for Toxic Substances and Disease Registry). (1997). Toxicological profile for  
11 trichloroethylene (update). (PB98-101165). Atlanta, GA: U.S. Department of Health and  
12 Human Services, Public Health Service, Agency for Toxic Substances and Disease  
13 Registry. Retrieved from <http://www.atsdr.cdc.gov/toxprofiles/tp19-p.pdf>.
- 14 ATSDR. (Agency for Toxic Substances and Disease Registry). (1998). Health consultation:  
15 Evaluation of health impacts from potential past exposure to tetrachloroethylene in the  
16 Natick public water supply. Atlanta, GA: Agency for Toxic Substances & Disease  
17 Registry. Retrieved from <http://www.atsdr.cdc.gov/hac/pha/pha.asp?docid=1270&pg=0>.
- 18 ATSDR. (Agency for Toxic Substances and Disease Registry). (2000). Public Health  
19 Assessment for Tucson International Airport Area (a/k/a El Vado Residential Properties)  
20 Tucson, Pima County, Arizona, Region 9. CERCLIS No. AZD980737530. Office of  
21 Environmental Health, Environmental Health Consultation Services: U.S. Department of  
22 Health and Human Services.
- 23 ATSDR. (Agency for Toxic Substances and Disease Registry). (2001). Final Report: Evaluation  
24 of priority health conditions in a community with historical contamination by  
25 trichloroethylene. Atlanta, GA: U.S. Department of Health and Human Services.
- 26 ATSDR. (Agency for Toxic Substances and Disease Registry). (2003). Health consultation.  
27 View-master factory supply well (a.k.a. Mattel Portland Operations) Beaverton,  
28 Washington County, Oregon. Atlanta, GA: U.S. Department of Health and Human  
29 Services. Retrieved from <http://www.atsdr.cdc.gov/hac/pha/pha.asp?docid=293&pg=0>.
- 30 ATSDR. (Agency for Toxic Substances and Disease Registry). (2004). Feasibility investigation  
31 of worker exposure to trichloroethylene at the view-master factory in Beaverton, Oregon.  
32 Atlanta, GA: Author. Retrieved from  
33 <http://www.ntis.gov/search/product.aspx?ABBR=PB2005102173>.
- 34 ATSDR. (Agency for Toxic Substances and Disease Registry). (2004). Interaction profile: 1,1,1-  
35 trichloroethane, 1,1-dichloroethane, trichloroethylene, and tetrachloroethylene. Atlanta,  
36 GA: Author. Retrieved from <http://www.atsdr.cdc.gov/interactionprofiles/ip02.html>.
- 37 ATSDR. (Agency for Toxic Substances and Disease Registry). (2006). Health Consultation:  
38 Public health implications of exposures to low-level volatile organic compounds in public  
39 drinking water: Endicott area investigation Broome County, New York. Atlanta, GA: U.S.  
40 Department of Health and Human Services.
- 41 ATSDR. (Agency for Toxic Substances and Disease Registry). (2006). Health statistics review:  
42 Cancer and birth outcome analysis: Endicott area investigation: Endicott area, town of  
43 Union, Broome County, New York. Atlanta, GA: U.S. Department of Health and  
44 Human Services. Retrieved from

DRAFT– DO NOT QUOTE OR CITE

- 1 <http://www.atsdr.cdc.gov/HAC/pha/EndicottAreaInvestigation/EndicottHealthStatsReviewHC052606.pdf>.
- 2
- 3 ATSDR. (Agency for Toxic Substances and Disease Registry). (2006). Toxicological profile for  
4 1,1,1 trichloroethane. Atlanta, GA: Author. Retrieved from  
5 <http://www.atsdr.cdc.gov/toxpro2.html>.
- 6 ATSDR. (Agency for Toxic Substances and Disease Registry). (2007). Analyses of groundwater  
7 flow, contaminant fate and transport and distribution of drinking water at Tarawa Terrace  
8 and vicinity, U.S. Marine Corps Base Camp Lejeune, North Carolina: Historical  
9 reconstruction and present-day conditions. Atlanta, GA: U.S. Department of Health and  
10 Human Services, Agency for Toxic Substances and Disease Registry. Retrieved from  
11 <http://www.atsdr.cdc.gov/sites/lejeune/tarawaterrace.html>.
- 12 ATSDR. (Agency for Toxic Substances and Disease Registry). (2008). Health consultation.  
13 Cancer and birth outcome analysis, Endicott area, Town of Union, Broome County, New  
14 York. Health statistics review follow-up. Atlanta, GA: U.S. Department of Health and  
15 Human Services. Retrieved from  
16 <http://www.atsdr.cdc.gov/hac/pha/EndicottAreaInvestigationFollowUp/EndicottAreaHC051508.pdf>.
- 17
- 18 ATSDR. (Agency for Toxic Substances and Disease Registry). (2009). Study on birth defects  
19 and childhood cancers. Atlanta, GA: U.S. Department of Health and Human Services,  
20 Public Health Services. Retrieved from  
21 <http://www.atsdr.cdc.gov/sites/lejeune/update.html>.
- 22 Attallah, A. M.; Tabll, A. A.; Salem, S. F.; El-Sadany, M.; Ibrahim, T. A.; Osman, S.; El-  
23 Dosoky, I. M. (1999). DNA ploidy of liver biopsies from patients with liver cirrhosis and  
24 hepatocellular carcinoma: a flow cytometric analysis. *Cancer Lett*, 142, 65-69.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/10424782>
- 26 Auperin, A.; Benhamou, S.; Ory-Paoletti, C.; Flamant, R. (1994). Occupational risk factors for  
27 renal cell carcinoma: A case-control study. *Occup Environ Med*, 51, 426-428.  
28 <http://dx.doi.org/10.1136/oem.51.6.426>
- 29 Austin, E. W.; Parrish, J. M.; Kinder, D. H.; Bull, R. J. (1996). Lipid peroxidation and formation  
30 of 8-hydroxydeoxyguanosine from acute doses of halogenated acetic acids. *Fundam Appl*  
31 *Toxicol*, 31, 77-82. <http://dx.doi.org/10.1006/faat.1996.0078>
- 32 Austin, H.; Delzell, E.; Grufferman, S.; Levine, R.; Morrison, A. S.; Stolley, P. D.; Cole, P.  
33 (1987). Case-control study of hepatocellular carcinoma, occupation, and chemical  
34 exposures. *J Occup Med*, 29, 665-669. <http://www.ncbi.nlm.nih.gov/pubmed/2821204>
- 35 Auwerx, J.; Schoonjans, K.; Fruchart, J. C.; Staels, B. (1996). Transcriptional control of  
36 triglyceride metabolism: fibrates and fatty acids change the expression of the LPL and  
37 apo C-III genes by activating the nuclear receptor PPAR. *Atherosclerosis*, 124 Suppl,  
38 S29-S37. <http://www.ncbi.nlm.nih.gov/pubmed/8831913>
- 39 Axelson, O.; Seldén, A.; Andersson, K.; Hogstedt, C. (1994). Updated and expanded Swedish  
40 cohort study on trichloroethylene and cancer risk. *J Occup Med*, 36, 556-562.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/8027881>
- 42 Baden, J. M.; Kelley, M.; Mazze, R. I.; Simmon, V. F. (1979). Mutagenicity of inhalation  
43 anaesthetics: trichloroethylene, divinyl ether, nitrous oxide and cyclopropane. *Br J*  
44 *Anaesth*, 51, 417-421.

DRAFT– DO NOT QUOTE OR CITE

- 1 Badger, T.; Ronis, M.; Seitz, H.; Albano, E.; Ingelman-Sundberg, M.; Lieber, C. (2003). Alcohol  
2 metabolism: role in toxicity and carcinogenesis. *Alcohol Clin Exp Res*, 27, 336-347.  
3 <http://dx.doi.org/10.1097/01.ALC.0000052583.87673.37>
- 4 Bai, C. L.; Canfield, P. J.; Stacey, N. H. (1992). Effects of hexachloro-1,3-butadiene and 1,1,2,2-  
5 tetrachloroethylene on individual serum bile acids. *Toxicol Ind Health*, 8, 191-203.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/1502697>
- 7 Bai, C.; Canfield, P. J.; Stacey, N. H. (1992). Individual serum bile acids as early indicators of  
8 carbon tetrachloride- and chloroform-induced liver injury. *Toxicology*, 75, 221-234.
- 9 Bai, C.-L. and Stacey, N. H. (1993). Mechanism of trichloroethylene-induced elevation of  
10 individual serum bile acids: II in vitro and in vivo interference by trichloroethylene with  
11 bile acid transport in isolated rat hepatocytes. *Toxicol Appl Pharmacol*, 121, 296-302.  
12 <http://dx.doi.org/10.1006/taap.1993.1157>
- 13 Bailly, M.; Bain, C.; Favrot, M. C.; Ozturk, M. (1995). Somatic mutations of von Hippel-Lindau  
14 (VHL) tumor-suppressor gene in European kidney cancers. *Int J Cancer*, 63, 660-664.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/7591282>
- 16 Baker, V. A.; Harries, H. M.; Waring, J. F.; Duggan, C. M.; Ni, H. A.; Jolly, R. A., . . . Rockett,  
17 J. C. (2004). Clofibrate-induced gene expression changes in rat liver: a cross-laboratory  
18 analysis using membrane cDNA arrays. *Environ Health Perspect*, 112, 428-438.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/15033592>
- 20 Bakke, B.; Stewart, P.; Waters, M. (2007). Uses of and exposure to trichloroethylene in U.S.  
21 industry: a systematic literature review. *J Occup Environ Hyg*, 4, 375-390.  
22 <http://dx.doi.org/10.1080/15459620701301763>
- 23 Balkwill, F. and Coussens, L. M. (2004). Cancer: an inflammatory link. *Nature*, 431, 405-406.  
24 <http://dx.doi.org/10.1038/431405a>
- 25 Balkwill, F.; Charles, K. A.; Mantovani, A. (2005). Smoldering and polarized inflammation in  
26 the initiation and promotion of malignant disease. *Cancer Cell*, 7, 211-217.  
27 <http://dx.doi.org/10.1016/j.ccr.2005.02.013>
- 28 Ballestar, E. and Esteller, M. (2002). The impact of chromatin in human cancer: linking DNA  
29 methylation to gene silencing. *Carcinogenesis*, 23, 1103-1109.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/12117766>
- 31 Balluz, L.; Philen, R.; Ortega, L.; Rosales, C.; Brock, J.; Barr, D.; Kieszak, S. (2001).  
32 Investigation of systemic lupus erythematosus in Nogales, Arizona. *Am J Epidemiol*,  
33 154, 1029-1036. <http://www.ncbi.nlm.nih.gov/pubmed/11724719>
- 34 Band, P. R.; Le, N. D.; Fang, R.; Deschamps, M.; Gallagher, R. P.; Yang, P. (2000).  
35 Identification of occupational cancer risks in British Columbia: A population-based case-  
36 control study of 995 incident breast cancer cases by menopausal status, controlling for  
37 confounding factors. *J Occup Environ Med*, 42, 284-310.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/10738708>
- 39 Banerjee, S. and Van Duuren, B. (1978). Covalent binding of the carcinogen trichloroethylene to  
40 hepatic microsomal proteins and to exogenous DNA in vitro. *Cancer Res*, 38, 776-780.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/626981>
- 42 Banks, P. M. (1992). Changes in diagnosis of non-Hodgkin's lymphomas over time. *Cancer Res*,  
43 52, 5453s-5455s. <http://www.ncbi.nlm.nih.gov/pubmed/1394153>
- 44 Bannasch, P. (1996). Pathogenesis of hepatocellular carcinoma: sequential cellular, molecular,  
45 and metabolic changes. 14, 161-197. <http://www.ncbi.nlm.nih.gov/pubmed/9055578>

- 1 Bannasch, P. (2001). Hormonal and hormone-like effects eliciting hepatocarcinogenesis. *Folia*  
2 *Histochem Cytobiol*, 39 Suppl 2, 28-29. <http://www.ncbi.nlm.nih.gov/pubmed/11820616>
- 3 Bannasch, P.; Haertel, T.; Su, Q. (2003). Significance of hepatic preneoplasia in risk  
4 identification and early detection of neoplasia. *Toxicol Pathol*, 31, 134-139.  
5 <http://dx.doi.org/10.1080/01926230390173923>
- 6 Bannasch, P.; Nehrbass, D.; Kopp-Schneider, A. (2001). Significance of hepatic preneoplasia for  
7 cancer chemoprevention. *IARC Sci Publ*, 154, 223-240.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/11220662>
- 9 Barak, Y.; Sadovsky, Y.; Shalom-Barak, T. (2008). PPAR Signaling in Placental Development  
10 and Function. *PPAR Research*, 2008, 142082. <http://dx.doi.org/10.1155/2008/142082>
- 11 Baraona, E.; Shoichet, L.; Navder, K.; Lieber, C. (2002). Mediation by nitric oxide of the  
12 stimulatory effects of ethanol on blood flow. *Life Sci*, 70, 2987-2995.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/12138012>
- 14 Baraona, E.; Zeballos, G.; Shoichet, L.; Mak, K.; Lieber, C. (2002). Ethanol consumption  
15 increases nitric oxide production in rats, and its peroxynitrite-mediated toxicity is  
16 attenuated by polyenylphosphatidylcholine. *Alcohol Clin Exp Res*, 26, 883-889.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/12068258>
- 18 Bardodej, Z. and Vyskocil, J. (1956). The problem of trichloroethylene in occupational  
19 medicine: trichloroethylene metabolism and its effect on the nervous system evaluated as  
20 a means of hygienic control. *Arch Environ Occup Health*, 13, 581-592.
- 21 Barrass, N. C.; Price, R. J.; Lake, B. G.; Orton, T. C. (1993). Comparison of the acute and  
22 chronic mitogenic effects of the peroxisome proliferators methylclofenapate and clofibric  
23 acid in rat liver. *Carcinogenesis*, 14, 1451-1456.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/8330363>
- 25 Barret, L.; Arsac, P.; Vincent, M.; Faure, J.; Garrel, S.; Reymond, F. (1982). Evoked trigeminal  
26 nerve potential in chronic trichloroethylene intoxication. *Clin Toxicol*, 19, 419-423.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/7143527>
- 28 Barret, L.; Faure, J.; Guillard, B.; Chomat, D.; Didier, B.; Debru, J. (1984). Trichloroethylene  
29 occupational exposure: elements for better prevention. *Int Arch Occup Environ Health*,  
30 53, 283-289. <http://www.ncbi.nlm.nih.gov/pubmed/6715052>
- 31 Barret, L.; Garrel, S.; Danel, V.; Debru, J. L. (1987). Chronic trichloroethylene intoxication: a  
32 new approach by trigeminal-evoked potentials? *Arch Environ Occup Health*, 42, 297-  
33 302. <http://www.ncbi.nlm.nih.gov/pubmed/3452297>
- 34 Barret, L.; Torch, S.; Leray, C.; Sarliève, L.; Saxod, R. (1992). Morphometric and biochemical  
35 studies in trigeminal nerve of rat after trichloroethylene or dichloroacetylene oral  
36 administration. *Neurotoxicology*, 13, 601-614.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/1282229>
- 38 Barret, L.; Torch, S.; Usson, Y.; Gonthier, B.; Saxod, R. (1991). A morphometric evaluation of  
39 the effects of trichloroethylene and dichloroacetylene on the rat mental nerve Preliminary  
40 results. *Neurosci Lett*, 131, 141-144. <http://www.ncbi.nlm.nih.gov/pubmed/1762688>
- 41 Barter, Z. E.; Bayliss, M. K.; Beaune, P. H.; Boobis, A. R.; Carlile, D. J.; Edwards, R. J., . . .  
42 Rostami-Hodjegan, A. (2007). Scaling factors for the extrapolation of in vivo metabolic  
43 drug clearance from in vitro data: reaching a consensus on values of human microsomal  
44 protein and hepatocellularity per gram of liver. *Curr Drug Metab*, 8, 33-45.

DRAFT– DO NOT QUOTE OR CITE

- 1 Barton, H. A. and Clewell, H. J., III. (2000). Evaluating noncancer effects of trichloroethylene:  
2 Dosimetry, mode of action, and risk assessment. *Environ Health Perspect*, 108, 323-334.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/10807562>
- 4 Barton, H. A. and Das, S. (1996). Alternatives for a risk assessment on chronic noncancer effects  
5 from oral exposure to trichloroethylene. *Regul Toxicol Pharmacol*, 24, 269-285.  
6 <http://dx.doi.org/10.1006/rtph.1996.0140>
- 7 Barton, H. A.; Chiu, W. A.; Setzer, R. W.; Andersen, M. E.; Bailer, A. J.; Bois, F. Y., . . . Tan,  
8 Y. M. (2007). Characterizing uncertainty and variability in physiologically-based  
9 pharmacokinetic (PBPK) models: State of the science and needs for research and  
10 implementation. *Toxicol Sci*, 99, 395-402. <http://dx.doi.org/10.1093/toxsci/kfm100>
- 11 Barton, H. A.; Creech, J. R.; Godin, C. S.; Randall, G. M.; Seckel, C. S. (1995). Chloroethylene  
12 mixtures: pharmacokinetic modeling and in vitro metabolism of vinyl chloride,  
13 trichloroethylene, and trans-1,2-dichloroethylene in rat. *Toxicol Appl Pharmacol*, 130,  
14 237-247. <http://dx.doi.org/10.1006/taap.1995.1029>
- 15 Barton, H. A.; Flemming, C. D.; Lipscomb, J. C. (1996). Evaluating human variability in  
16 chemical risk assessment: hazard identification and dose-response assessment for  
17 noncancer oral toxicity of trichloroethylene. *Toxicology*, 111, 271-287.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/8711742>
- 19 Barton, H.; Bull, R.; Schultz, I.; Andersen, M. (1999). Dichloroacetate (DCA) dosimetry:  
20 interpreting DCA-induced liver cancer dose response and the potential for DCA to  
21 contribute to trichloroethylene-induced liver cancer. *Toxicol Lett*, 106, 9-21.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/10378446>
- 23 Bartonicek, V. (1962). Metabolism and excretion of trichloroethylene after inhalation by human  
24 subjects. *Occup Environ Med*, 19, 134-141.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/13865497>
- 26 Bartonicek, V. and Teisinger, J. (1962). Effect of tetraethyl thiuram disulphide (disulfiram) on  
27 metabolism of trichloroethylene in man. *Occup Environ Med*, 19, 216-221.  
28 <http://dx.doi.org/10.1136/oem.19.3.216>
- 29 Bartosiewicz, M.; Penn, S.; Buckpitt, A. (2001). Applications of gene arrays in environmental  
30 toxicology: fingerprints of gene regulation associated with cadmium chloride,  
31 benzo(a)pyrene, and trichloroethylene. *Environ Health Perspect*, 109, 71-74.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/11171528>
- 33 Bartsch, H.; Malaveille, C.; Barbin, A.; Planche, G. (1979). Mutagenic and alkylating  
34 metabolites of halo-ethylenes, chlorobutadienes and dichlorobutenes produced by rodent  
35 or human liver tissues: Evidence for oxirane formation by P450-linked microsomal  
36 mono-oxygenases. *Arch Toxicol*, 41, 249-277. <http://dx.doi.org/10.1007/BF00296896>
- 37 Beaudreuil, S.; Lasfargues, G.; Laueriere, L.; El Ghoul, Z.; Fourquet, F.; Longuet, C., . . .  
38 Buchler, M. (2005). Occupational exposure in ANCA-positive patients: A case-control  
39 study. *Kidney Int*, 67, 1961-1966. <http://dx.doi.org/10.1111/j.1523-1755.2005.00295.x>
- 40 Beaumont, J. J. and Breslow, N. E. (1981). Power considerations in epidemiologic studies of  
41 vinyl chloride workers. *Am J Epidemiol*, 114, 725-734.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/7030065>
- 43 Becker, N.; Deeg, E.; Nieters, A. (2004). Population-based study of lymphoma in Germany:  
44 rationale, study design and first results. *Leuk Res*, 28, 713-724.  
45 <http://dx.doi.org/10.1016/j.leukres.2003.11.010>



DRAFT– DO NOT QUOTE OR CITE

- 1 Beckstead, M. J.; Weiner, J. L.; Eger, E. I.; Gong, D. H.; Mihic, S. J. (2000). Glycine and  
2 [gamma]-aminobutyric acid(A) receptor function is enhanced by inhaled drugs of abuse.  
3 *Mol Pharmacol*, 57, 1199-1205. <http://www.ncbi.nlm.nih.gov/pubmed/10825391>
- 4 Beekman, J. M.; Boess, F.; Hildebrand, H.; Kalkuhl, A.; Suter, L. (2006). Gene expression  
5 analysis of the hepatotoxicant methapyrilene in primary rat hepatocytes: an  
6 interlaboratory study. *Environ Health Perspect*, 114, 92-99.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/16393664>
- 8 Begg, C. B. and Mazumdar, M. (1994). Operating characteristics of a rank correlation test for  
9 publication bias. *Biometrics*, 50, 1088-1101.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/7786990>
- 11 Beland, F. (1999). NTP technical report on the toxicity and metabolism studies of chloral hydrate  
12 (CAS No. 302-17-0). Administered by gavage to F344/N rats and B6C3F1 mice. U.S.  
13 Dept. of Health and Human Services, National Institutes of Health: Rockville, MD.
- 14 Benedetti, M. S.; Whomsley, R.; Canning, M. (2007). Drug metabolism in the paediatric  
15 population and in the elderly. *Drug Discov Today*, 12, 599-610.  
16 <http://dx.doi.org/10.1016/j.drudis.2007.06.011>
- 17 Benichou, J.; Chow, W. H.; McLaughlin, J. K.; Mandel, J. S.; Fraumeni, J. F. (1998). Population  
18 attributable risk of renal cell cancer in Minnesota. *Am J Epidemiol*, 148, 424-430.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/9737554>
- 20 Benson, R. (2004). [Correspondence from Robert Benson (U. S. EPA) to Dr. P. D. Johnson  
21 (University of Arizona)].
- 22 Beppu, K. (1968). Transmission of the anesthetic agents through the placenta in painless delivery  
23 and their effects on newborn infants. *Keio J Med*, 17, 81-107.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/5727834>
- 25 Berger, J. and Daxenbichler, G. (2002). DNA methylation of nuclear receptor genes--possible  
26 role in malignancy. *J Steroid Biochem Mol Biol*, 80, 1-11.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/11867259>
- 28 Berger, T. and Horner, C. M. (2003). In vivo exposure of female rats to toxicants may affect  
29 oocyte quality. *Reprod Toxicol*, 17, 273-281. [http://dx.doi.org/10.1016/S0890-](http://dx.doi.org/10.1016/S0890-6238(03)00009-1)  
30 [6238\(03\)00009-1](http://dx.doi.org/10.1016/S0890-6238(03)00009-1)
- 31 Bergeron, A.; Jorquera, R.; Tanguay, R. (2003). [Hereditary tyrosinemia: an endoplasmic  
32 reticulum stress disorder?]. *Med Sci (Paris)*, 19, 976-980.  
33 <http://dx.doi.org/10.1051/medsci/20031910976>
- 34 Bergman, K. (1983). Interactions of trichloroethylene with DNA in vitro and with RNA and  
35 DNA of various mouse tissues in vivo. *Arch Toxicol*, 54, 181-193.  
36 <http://www.ncbi.nlm.nih.gov/pubmed/6197950>
- 37 Berman, E.; Schlicht, M.; Moser, V. C.; MacPhail, R. C. (1995). A multidisciplinary approach to  
38 toxicological screening: I. Systemic toxicity. *J Toxicol Environ Health*, 45, 127-143.  
39 <http://dx.doi.org/10.1080/15287399509531986>
- 40 Bernad, P. G.; Newell, S.; Spyker, D. A. (1987). Neurotoxicity and behavior abnormalities in a  
41 cohort chronically exposed to trichloroethylene. *Vet Hum Toxicol*, 29, 475.
- 42 Bernauer, U.; Birner, G.; Dekant, W.; Henschler, D. (1996). Biotransformation of  
43 trichloroethene: dose-dependent excretion of 2,2,2-trichloro-metabolites and mercapturic  
44 acids in rats and humans after inhalation. *Arch Toxicol*, 70, 338-346.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/8975632>

DRAFT– DO NOT QUOTE OR CITE

- 1 Bernillon, P. and Bois, F. Y. (2000). Statistical issues in toxicokinetic modeling: a bayesian  
2 perspective. *Environ Health Perspect*, 108, 883-893.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/11035998>
- 4 Béroud, C.; Joly, D.; Gallou, C.; Staroz, F.; Orfanelli, M. T.; Junien, C. (1998). Software and  
5 database for the analysis of mutations in the VHL gene. *Nucleic Acids Res*, 26, 256-258.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/9399847>
- 7 Besson, H.; Brennan, P.; Becker, N.; Nieters, A.; De Sanjosé, S.; Font, R., . . . Boffetta, P.  
8 (2006). Tobacco smoking, alcohol drinking and non-Hodgkin's lymphoma: A European  
9 multicenter case-control study (Epilymph). *Int J Cancer*, 119, 901-908.  
10 <http://dx.doi.org/10.1002/ijc.21913>
- 11 Best, D. H. and Coleman, W. B. (2007). Treatment with 2-AAF blocks the small hepatocyte-like  
12 progenitor cell response in retrorsine-exposed rats. *J Hepatol*, 46, 1055-1063.  
13 <http://dx.doi.org/10.1016/j.jhep.2007.01.040>
- 14 Beyer, R. P.; Fry, R. C.; Lasarev, M. R.; McConnachie, L. A.; Meira, L. B.; Palmer, V. S., . . .  
15 Consortium, M. o. t. T. R. (2007). Multicenter study of acetaminophen hepatotoxicity  
16 reveals the importance of biological endpoints in genomic analyses. *Toxicol Sci*, 99, 326-  
17 337. <http://dx.doi.org/10.1093/toxsci/kfm150>
- 18 Bhunya, S. P. and Behera, B. C. (1987). Relative genotoxicity of trichloroacetic acid (TCA) as  
19 revealed by different cytogenetic assays: Bone marrow chromosome aberration,  
20 micronucleus and sperm-head abnormality in the mouse. *Mutat Res Genet Toxicol*  
21 *Environ Mutagen*, 188, 215-221. [http://dx.doi.org/10.1016/0165-1218\(87\)90092-9](http://dx.doi.org/10.1016/0165-1218(87)90092-9)
- 22 Bhunya, S. P. and Jena, G. B. (1996). The evaluation of clastogenic potential of trichloroacetic  
23 acid (TCA) in chick in vivo test system. *Mutat Res Genet Toxicol Environ Mutagen*, 367,  
24 253-259. [http://dx.doi.org/10.1016/S0165-1218\(96\)90085-3](http://dx.doi.org/10.1016/S0165-1218(96)90085-3)
- 25 Bignami, M.; Conti, G.; Conti, L.; Crebelli, R.; Misuraca, F.; Puglia, A. M., . . . Carere, A.  
26 (1980). Mutagenicity of halogenated aliphatic hydrocarbons in *Salmonella typhimurium*,  
27 *Streptomyces coelicolor* and *Aspergillus nidulans*. *Chem Biol Interact*, 30, 9-23.  
28 [http://dx.doi.org/10.1016/0009-2797\(80\)90110-6](http://dx.doi.org/10.1016/0009-2797(80)90110-6)
- 29 Binaschi, S. and Cantu, L. (1983). Vestibular Function In Solvent Exposure: Clinical Criteria.  
30 46, 205-210.
- 31 Birner, G.; Bernauer, U.; Werner, M.; Dekant, W. (1997). Biotransformation, excretion and  
32 nephrotoxicity of haloalkene-derived cysteine S-conjugates. *Arch Toxicol*, 72, 1-8.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/9458184>
- 34 Birner, G.; Richling, C.; Henschler, D.; Anders, M. W.; Dekant, W. (1994). Metabolism of  
35 tetrachloroethene in rats: Identification of N[epsilon]-(dichloroacetyl)-L-lysine and  
36 N[epsilon]-(trichloroacetyl)-L-lysine as protein adducts. *Chem Res Toxicol*, 7, 724-732.  
37 <http://dx.doi.org/10.1021/tx00042a003>
- 38 Birner, G.; Vamvakas, S.; Dekant, W.; Henschler, D. (1993). Nephrotoxic and genotoxic N-  
39 acetyl-S-dichlorovinyl-L-cysteine is a urinary metabolite after occupational 1,1,2-  
40 trichloroethene exposure in humans: implications for the risk of trichloroethene exposure.  
41 *Environ Health Perspect*, 99, 281-284. <http://www.ncbi.nlm.nih.gov/pubmed/8319644>
- 42 Birner, G.; Werner, M.; Rosner, E.; Mehler, C.; Dekant, W. (1998). Biotransformation,  
43 excretion, and nephrotoxicity of the hexachlorobutadiene metabolite (E)-N-acetyl-S-  
44 (1,2,3,4, 4-pentachlorobutadienyl)-L-cysteine sulfoxide. *Chem Res Toxicol*, 11, 750-757.  
45 <http://dx.doi.org/10.1021/tx970216n>

DRAFT– DO NOT QUOTE OR CITE

- 1 Bjornsson, H. T.; Sigurdsson, M. I.; Fallin, M. D.; Irizarry, R. A.; Aspelund, T.; Cui, H., . . .  
2 Feinberg, A. P. (2008). Intra-individual change over time in DNA methylation with  
3 familial clustering. *JAMA*, 299, 2877-2883. <http://dx.doi.org/10.1001/jama.299.24.2877>
- 4 Blackburn, A.; Coggan, M.; Tzeng, H.; Lantum, H.; Polekhina, G.; Parker, M., . . . Board, P.  
5 (2001). GSTZ1d: a new allele of glutathione transferase zeta and maleylacetoacetate  
6 isomerase. *Pharmacogenetics*, 11, 671-678.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/11692075>
- 8 Blackburn, A.; Tzeng, H.; Anders, M.; Board, P. (2000). Discovery of a functional  
9 polymorphism in human glutathione transferase zeta by expressed sequence tag database  
10 analysis. *Pharmacogenetics*, 10, 49-57. <http://www.ncbi.nlm.nih.gov/pubmed/10739172>
- 11 Blackshaw, A. W. (1970). Histochemical localization of testicular enzyme. In A. D. Johnson, W.  
12 R. Gomes & N. L. VanDemark (Eds.), *The testis* (pp. 73-173). New York, NY: Academic  
13 Press.
- 14 Blain, L.; Lachapelle, P.; Molotchnikoff, S. (1994). Electroretinal responses are modified by  
15 chronic exposure to trichloroethylene. *Neurotoxicology*, 15, 627-631.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/7854598>
- 17 Blair, A.; Haas, T.; Prosser, R.; Morrissette, M.; Blackman, K.; Grauman, D., . . . Moran, F.  
18 (1989). Mortality among United States Coast Guard marine inspectors. *Arch Environ*  
19 *Health*, 44, 150 - 156.
- 20 Blair, A.; Hartge, P.; Stewart, P. A.; McAdams, M.; Lubin, J. (1998). Mortality and cancer  
21 incidence of aircraft maintenance workers exposed to trichloroethylene and other organic  
22 solvents and chemicals: Extended follow-up. *Occup Environ Med*, 55, 161-171.  
23 <http://dx.doi.org/10.1136/oem.55.3.161>
- 24 Blair, A.; Linos, A.; Stewart, P. A.; Burmeister, L. F.; Gibson, R.; Everett, G., . . . Cantor, K. P.  
25 (1993). Evaluation of risks for non-Hodgkin's lymphoma by occupation and industry  
26 exposures from a case-control study. *Am J Ind Med*, 23, 301-312.  
27 <http://dx.doi.org/10.1002/ajim.4700230207>
- 28 Blake, J. E.; Compton, K. V.; Schmidt, W.; Orrego, H. (1988). Accuracy of death certificates in  
29 the diagnosis of alcoholic liver cirrhosis. *Alcohol Clin Exp Res*, 12, 168-172.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/3279851>
- 31 Blake, M. J.; Castro, L.; Leeder, J. S.; Kearns, G. L. (2005). Ontogeny of drug metabolizing  
32 enzymes in the neonate. *Semin Fetal Neonatal Med*, 10, 123-138.  
33 <http://dx.doi.org/10.1016/j.siny.2004.11.001>
- 34 Blatter, B. M.; Roeleveld, N.; Zielhuis, G. A.; Verbeek, A. L. (1997). Assessment of  
35 occupational exposure in a population based case-control study: comparing postal  
36 questionnaires with personal interviews. *Occup Environ Med*, 54, 54-59.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/9072035>
- 38 Blechacz, B. R. and Gores, G. J. (2008). Cholangiocarcinoma. *Clin Liver Dis*, 12, 131-150, ix.  
39 <http://dx.doi.org/10.1016/j.cld.2007.11.003>
- 40 Bloemen, L. and Tomenson, J. (1995). Increased incidence of renal cell tumours in a cohort of  
41 cardboard workers exposed to trichloroethylene. *Arch Toxicol*, 70, 129-133.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/8773187>
- 43 Bloemen, L.; Monster, A.; Kezic, S.; Commandeur, J.; Veulemans, H.; Vermeulen, N.; Wilmer,  
44 J. (2001). Study on the cytochrome P-450- and glutathione-dependent biotransformation

DRAFT– DO NOT QUOTE OR CITE

- 1 of trichloroethylene in humans. *Int Arch Occup Environ Health*, 74, 102-108.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/11317702>
- 3 Blossom, S. J. and Doss, J. C. (2007). Trichloroethylene alters central and peripheral immune  
4 function in autoimmune-prone MRL(+/+) mice following continuous developmental and  
5 early life exposure. *J Immunotoxicol*, 4, 129-141.  
6 <http://dx.doi.org/10.1080/15476910701337035>
- 7 Blossom, S. J. and Gilbert, K. M. (2006). Exposure to a metabolite of the environmental  
8 toxicant, trichloroethylene, attenuates CD4+ T cell activation-induced cell death by  
9 metalloproteinase-dependent FasL shedding. *Toxicol Sci*, 92, 103-114.  
10 <http://dx.doi.org/10.1093/toxsci/kfj212>
- 11 Blossom, S. J.; Doss, J. C.; Gilbert, K. M. (2007). Chronic exposure to a trichloroethylene  
12 metabolite in autoimmune-prone MRL+/+ mice promotes immune modulation and  
13 alopecia. *Toxicol Sci*, 95, 401-411. <http://dx.doi.org/10.1093/toxsci/kfl149>
- 14 Blossom, S. J.; Doss, J. C.; Hennings, L. J.; Jernigan, S.; Melnyk, S.; James, S. J. (2008).  
15 Developmental exposure to trichloroethylene promotes CD4+ T cell differentiation and  
16 hyperactivity in association with oxidative stress and neurobehavioral deficits in MRL+/+  
17 mice. *Toxicol Appl Pharmacol*, 231, 344-353.  
18 <http://dx.doi.org/10.1016/j.taap.2008.05.009>
- 19 Blossom, S. J.; Pumford, N. R.; Gilbert, K. M. (2004). Activation and attenuation of apoptosis of  
20 CD4+ T cells following in vivo exposure to two common environmental toxicants,  
21 trichloroacetaldehyde hydrate and trichloroacetic acid. *J Autoimmun*, 23, 211-220.  
22 <http://dx.doi.org/10.1016/j.jaut.2004.06.007>
- 23 Board, P.; Chelvanayagam, G.; Jermin, L.; Tetlow, N.; Tzeng, H.; Anders, M.; Blackburn, A.  
24 (2001). Identification of novel glutathione transferases and polymorphic variants by  
25 expressed sequence tag database analysis. *Drug Metab Dispos*, 29, 544-547.  
26 <http://www.ncbi.nlm.nih.gov/pubmed/11259348>
- 27 Boerrigter, M. E. (2004). Mutagenicity of the peroxisome proliferators clofibrate, Wyeth 14,643  
28 and di-2-ethylhexyl phthalate in the lacZ plasmid-based transgenic mouse mutation  
29 assay. *Journal of Carcinogenesis*, 3, 7. <http://dx.doi.org/10.1186/1477-3163-3-7>
- 30 Boffetta, P. and de Vocht, F. (2007). Occupation and the risk of non-Hodgkin lymphoma. *Cancer*  
31 *Epidemiol Biomarkers Prev*, 16, 369-372. <http://dx.doi.org/10.1158/1055-9965.EPI-06-1055>
- 32
- 33 Bogen, K. T. (1990). *Uncertainty in Environmental Health Risk Assessment*. New York: Garland  
34 Publishing Company.
- 35 Bogen, K. T.; BW Jr, C.; Machicao, L. K. (1992). Dermal absorption of dilute aqueous  
36 chloroform, trichloroethylene, and tetrachloroethylene in hairless guinea pigs. *Toxicol*  
37 *Sci*, 18, 30-39. [http://dx.doi.org/10.1016/0272-0590\(92\)90192-K](http://dx.doi.org/10.1016/0272-0590(92)90192-K)
- 38 Boice, J. D.; Cohen, S. S.; Mumma, M. T.; Dupree Ellis, E.; Eckerman, K. F.; Leggett, R. W., . . .  
39 . Henderson, B. E. (2006). Mortality among radiation workers at Rocketdyne (Atomics  
40 International), 1948-1999. *Radiat Res*, 166, 98-115. <http://dx.doi.org/10.1667/RR3582.1>
- 41 Boice, J. D.; Marano, D. E.; Cohen, S. S.; Mumma, M. T.; Blot, W. J.; Brill, A. B., . . .  
42 McLaughlin, J. K. (2006). Mortality among Rocketdyne workers who tested rocket  
43 engines, 1948-1999. *J Occup Environ Med*, 48, 1070-1092.  
44 <http://dx.doi.org/10.1097/01.jom.0000240661.33413.b5>

DRAFT– DO NOT QUOTE OR CITE

- 1 Boice, J.; Marano, D.; Fryzek, J.; Sadler, C.; McLaughlin, J. (1999). Mortality among aircraft  
2 manufacturing workers. *Occup Environ Med*, 56, 581-597.  
3 <http://dx.doi.org/10.1136/oem.56.9.581>
- 4 Bois, F. (2000). Statistical analysis of Clewell et al. PBPK model of trichloroethylene kinetics.  
5 *Environ Health Perspect*, 108, 307-316. <http://www.ncbi.nlm.nih.gov/pubmed/10807560>
- 6 Bois, F. (2000). Statistical analysis of Fisher et al. PBPK model of trichloroethylene kinetics.  
7 *Environ Health Perspect*, 108 Suppl 2, 275-282.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/10807558>
- 9 Bolm-Audorff, U.; Ahrens, W.; Jockel, K.-H.; Molik, B.; Greiser, E.; Timm, J. (1988).  
10 Experience with supplementary questionnaires in a lung cancer case-referent study. In D.  
11 Henon & M. Goldberg (Eds.), *Proceedings of the workshop on methodology of*  
12 *assessment of occupational exposures in the context of epidemiological detection of*  
13 *cancer risks* (pp. 117-168). Luxembourg
- 14 Bolt, H. M. and Filser, J. G. (1977). Irreversible binding of chlorinated ethylenes to  
15 macromolecules. *Environ Health Perspect*, 21, 107-112.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/612434>
- 17 Bolt, H.; Lammert, M.; Selinski, S.; Brüning, T. (2004). Urinary alpha1-microglobulin excretion  
18 as biomarker of renal toxicity in trichloroethylene-exposed persons. *Int Arch Occup*  
19 *Environ Health*, 77, 186-190. <http://dx.doi.org/10.1007/s00420-003-0500-3>
- 20 Bombail, V.; Moggs, J. G.; Orphanides, G. (2004). Perturbation of epigenetic status by toxicants.  
21 *Toxicol Lett*, 149, 51-58. <http://dx.doi.org/10.1016/j.toxlet.2004.01.003>
- 22 Bonatti, S.; Cavalieri, Z.; Viaggi, S.; Abbondandolo, A. (1992). The analysis of 10 potential  
23 spindle poisons for their ability to induce CREST-positive micronuclei in human diploid  
24 fibroblasts. *Mutagenesis*, 7, 111-114. <http://www.ncbi.nlm.nih.gov/pubmed/1579065>
- 25 Boorman, G. A.; Dellarco, V.; Dunnick, J. K.; Chapin, R. E.; Hunter, S.; Hauchman, F., . . . Sills,  
26 R. C. (1999). Drinking water disinfection byproducts: review and approach to toxicity  
27 evaluation. *Environ Health Perspect*, 107, 207-217.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/10229719>
- 29 Borzio, M.; Fargion, S.; Borzio, F.; Fracanzani, A. L.; Croce, A. M.; Stroffolini, T., . . . Roncalli,  
30 M. (2003). Impact of large regenerative, low grade and high grade dysplastic nodules in  
31 hepatocellular carcinoma development. *J Hepatol*, 39, 208-214.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/12873817>
- 33 Boublik, T.; Fried, V.; Hala, E. (1984). The vapour pressures of pure substances: selected values  
34 of the temperature dependence of the vapour pressures of some pure substances in the  
35 normal and low pressure region. Amsterdam, The Netherlands: Elsevier.
- 36 Bove, F. J. (1996). Public drinking water contamination and birthweight, prematurity, fetal  
37 deaths, and birth defects. *Toxicol Ind Health*, 12, 255-266.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/8794538>
- 39 Bove, F. J.; Fulcomer, M. C.; Klotz, J. B.; Esmart, J.; Dufficy, E. M.; Savrin, J. E. (1995). Public  
40 drinking water contamination and birth outcomes. *Am J Epidemiol*, 141, 850-862.
- 41 Bovenzi, M.; Barbone, F.; Pisa, F.; Betta, A.; Romeo, L.; Tonello, A., . . . Caramaschi, P. (2004).  
42 A case-control study of occupational exposures and systemic sclerosis. *Int Arch Occup*  
43 *Environ Health*, 77, 10-16. <http://dx.doi.org/10.1007/s00420-003-0462-5>
- 44 Bovo, R.; Aimoni, C.; Martini, A. (2006). Immune-mediated inner ear disease. *Acta Otolaryngol*,  
45 126, 1012-1021. <http://dx.doi.org/10.1080/00016480600606723>

DRAFT– DO NOT QUOTE OR CITE

- 1 Bowler, R. M.; Mergler, D.; Huel, G.; Harrison, R.; Cone, J. (1991). Neuropsychological  
2 impairment among former microelectronics workers. *Neurotoxicology*, 12, 87-103.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/2014071>
- 4 Box, G. E. P.; Hunter, W. G.; Hunter, J. S. (1978). *Statistics for experimenters: an introduction*  
5 *to design, data analysis, and model building*. New York, NY: Wiley.
- 6 Boyault, S.; Rickman, D. S.; de Reynies, A.; Balabaud, C.; Rebouissou, S.; Jeannot, E., . . .  
7 Zucman-Rossi, J. (2007). Transcriptome classification of HCC is related to gene  
8 alterations and to new therapeutic targets. *Hepatology*, 45, 42-52.  
9 <http://dx.doi.org/10.1002/hep.21467>
- 10 Boyer, A.; Finch, W.; Runyan, R. (2000). Trichloroethylene inhibits development of embryonic  
11 heart valve precursors in vitro. *Toxicol Sci*, 53, 109-117.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/10653528>
- 13 Boyes, W. K.; Bercegeay, M.; Krantz, T.; Evans, M.; Benignus, V.; Simmons, J. E. (2005).  
14 Momentary brain concentration of trichloroethylene predicts the effects on rat visual  
15 function. *Toxicol Sci*, 87, 187-196. <http://dx.doi.org/10.1093/toxsci/kfi242>
- 16 Boyes, W. K.; Bushnell, P. J.; Crofton, K. M.; Evans, M.; Simmons, J. E. (2000). Neurotoxic and  
17 pharmacokinetic responses to trichloroethylene as a function of exposure scenario.  
18 *Environ Health Perspect*, 108, 317-322. <http://www.ncbi.nlm.nih.gov/pubmed/10807561>
- 19 Boyes, W.; Bercegeay, M.; Ali, J.; Krantz, T.; McGee, J.; Evans, M., . . . Simmons, J. (2003).  
20 Dose-based duration adjustments for the effects of inhaled trichloroethylene on rat visual  
21 function. *Toxicol Sci*, 76, 121-130. <http://dx.doi.org/10.1093/toxsci/kfg213>
- 22 Boyle, C. A. and Brann, E. A. (1992). Proxy respondents and the validity of occupational and  
23 other exposure data. The Selected Cancers Cooperative Study Group. *Am J Epidemiol*,  
24 136, 712-721. <http://www.ncbi.nlm.nih.gov/pubmed/1442737>
- 25 Bradford, B. U.; Kono, H.; Isayama, F.; Kosyk, O.; Wheeler, M. D.; Akiyama, T. E., . . . Rusyn,  
26 I. (2005). Cytochrome P450 CYP2E1, but not nicotinamide adenine dinucleotide  
27 phosphate oxidase, is required for ethanol-induced oxidative DNA damage in rodent  
28 liver. *Hepatology*, 41, 336-344. <http://dx.doi.org/10.1002/hep.20532>
- 29 Braissant, O. and Wahli, W. (1998). Differential expression of peroxisome proliferator-activated  
30 receptor-alpha, -beta, and -gamma during rat embryonic development. *Endocrinology*,  
31 139, 2748-2754. <http://www.ncbi.nlm.nih.gov/pubmed/9607781>
- 32 Branda, M. and Wands, J. R. (2006). Signal transduction cascades and hepatitis B and C related  
33 hepatocellular carcinoma. *Hepatology*, 43, 891-902. <http://dx.doi.org/10.1002/hep.21196>
- 34 Brauch, H.; Weirich, G.; Hornauer, M.; Störkel, S.; Wöhl, T.; Brüning, T. (1999).  
35 Trichloroethylene exposure and specific somatic mutations in patients with renal cell  
36 carcinoma. *J Natl Cancer Inst*, 91, 854-861.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/10340905>
- 38 Brauch, H.; Weirich, G.; Klein, B.; Rabstein, S.; Bolt, H.; Brüning, T. (2004). VHL mutations in  
39 renal cell cancer: does occupational exposure to trichloroethylene make a difference.  
40 *Toxicol Lett*, 151, 301-310. <http://dx.doi.org/10.1016/j.toxlet.2003.12.074>
- 41 Brautbar, N. and Williams, J. (2002). Industrial solvents and liver toxicity: risk assessment, risk  
42 factors and mechanisms. *Int J Hyg Environ Health*, 205, 479-491.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/12455270>

DRAFT– DO NOT QUOTE OR CITE

- 1 Brenner, H. and Gefeller, O. (1993). Use of the positive predictive value to correct for disease  
2 misclassification in epidemiologic studies. *Am J Epidemiol*, 138, 1007-1015.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/8256775>
- 4 Breslow, N. E. and Day, N. E. (1980). *Statistical methods in cancer research: Volume 1: The*  
5 *analysis of case-control studies (Vol. 32)*. Lyon, France: International Agency for  
6 Research on Cancer.
- 7 Breslow, N. E. and Day, N. E. (1987). *Statistical methods in cancer research, volume 2 - the*  
8 *design and analysis of cohort studies*. Oxford, UK: World Health Organization,  
9 International Agency for Research on Cancer.
- 10 Bringmann, G.; God, R.; Feineis, D.; Wesemann, W.; Riederer, P.; Rausch, W., . . . Sontag, K.  
11 (1995). The TaClo concept: 1-trichloromethyl-1,2,3,4-tetrahydro-beta-carboline (TaClo),  
12 a new toxin for dopaminergic neurons. *J Neural Transm Suppl*, 46, 235-244.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/8821060>
- 14 Briving, C.; Jacobson, I.; Hamberger, A.; Kjellstrand, P.; Haglid, K. G.; Rosengren, L. E. (1986).  
15 Chronic effects of perchloroethylene and trichloroethylene on the gerbil brain amino  
16 acids and glutathione. *Neurotoxicology*, 7, 101-108.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/2872637>
- 18 Brodsky, W. Y. and Uryvaeva, I. V. (1977). Cell polyploidy: its relation to tissue growth and  
19 function. *Int Rev Cytol*, 50, 275-332. <http://www.ncbi.nlm.nih.gov/pubmed/332657>
- 20 Bronley-DeLancey, A.; McMillan, D.; McMillan, J.; Jollow, D.; Mohr, L.; Hoel, D. (2006).  
21 Application of cryopreserved human hepatocytes in trichloroethylene risk assessment:  
22 relative disposition of chloral hydrate to trichloroacetate and trichloroethanol. *Environ*  
23 *Health Perspect*, 114, 1237-1242. <http://www.ncbi.nlm.nih.gov/pubmed/16882532>
- 24 Bronzetti, G.; Corsi, C.; Nieri, R. (1980). [Use of the D7 strain of *S. cerevisiae* in the  
25 determination of environmental risk. 1. Genetic effects of trichloroethylene]. *Boll Soc Ital*  
26 *Biol Sper*, 56, 1315-1321. <http://www.ncbi.nlm.nih.gov/pubmed/7006634>
- 27 Bronzetti, G.; Galli, A.; Corsi, C.; Cundari, E.; Del Carratore, R.; Nieri, R.; Paolini, M. (1984).  
28 Genetic and biochemical investigation on chloral hydrate in vitro and in vivo. *Mutat Res*,  
29 141, 19-22. <http://www.ncbi.nlm.nih.gov/pubmed/6384771>
- 30 Bross, G.; DiFranceisco, D.; Desmond, M. E. (1983). The effects of low dosages of  
31 trichloroethylene on chick development. *Toxicology*, 28, 283-294.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/6648978>
- 33 Brown, H. S.; Bishop, D. R.; Rowan, C. A. (1984). The role of skin absorption as a route of  
34 exposure for volatile organic compounds (VOCs) in drinking water. *Am J Public Health*,  
35 74, 479-484. <http://www.ncbi.nlm.nih.gov/pubmed/6711723>
- 36 Brown, R. P.; Delp, M. D.; Lindstedt, S. L.; Rhomberg, L. R.; Beliles, R. P. (1997).  
37 Physiological parameter values for physiologically based pharmacokinetic models.  
38 *Toxicol Ind Health*, 13, 407-484. <http://dx.doi.org/10.1177/074823379701300401>
- 39 Brownson, R. C. (1988). A case-control study of renal cell carcinoma in relation to occupation,  
40 smoking, and alcohol consumption. *Arch Environ Health*, 43, 238-241.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/3382249>
- 42 Brüning, T.; Lammert, M.; Kempkes, M.; Thier, R.; Golka, K.; Bolt, H. (1997). Influence of  
43 polymorphisms of GSTM1 and GSTT1 for risk of renal cell cancer in workers with long-  
44 term high occupational exposure to trichloroethene. *Arch Toxicol*, 71, 596-599.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/9285043>

DRAFT– DO NOT QUOTE OR CITE

- 1 Brüning, T.; Mann, H.; Melzer, H.; Sundberg, A.; Bolt, H. (1999). Pathological excretion  
2 patterns of urinary proteins in renal cell cancer patients exposed to trichloroethylene.  
3 *Occup Med (Lond)*, 49, 299-305. <http://www.ncbi.nlm.nih.gov/pubmed/10628055>
- 4 Brüning, T.; Pesch, B.; Wiesenhütter, B.; Rabstein, S.; Lammert, M.; Baumüller, A.; Bolt, H.  
5 (2003). Renal cell cancer risk and occupational exposure to trichloroethylene: results of a  
6 consecutive case-control study in Arnsberg, Germany. *Am J Ind Med*, 43, 274-285.  
7 <http://dx.doi.org/10.1002/ajim.10185>
- 8 Brüning, T.; Sundberg, A.; Birner, G.; Lammert, M.; Bolt, H.; Appelkvist, E., . . . Dallner, G.  
9 (1999). Glutathione transferase alpha as a marker for tubular damage after  
10 trichloroethylene exposure. *Arch Toxicol*, 73, 246-254.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/10463390>
- 12 Brüning, T.; Vamvakas, S.; Makropoulos, V.; Birner, G. (1998). Acute intoxication with  
13 trichloroethene: clinical symptoms, toxicokinetics, metabolism, and development of  
14 biochemical parameters for renal damage. *Toxicol Sci*, 41, 157-165.  
15 <http://dx.doi.org/10.1006/toxs.1997.2401>
- 16 Brüning, T.; Weirich, G.; Hornauer, M.; Höfler, H.; Brauch, H. (1997). Renal cell carcinomas in  
17 trichloroethene (TRI) exposed persons are associated with somatic mutations in the von  
18 Hippel-Lindau (VHL) tumour suppressor gene. *Arch Toxicol*, 71, 332-335.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/9137812>
- 20 Brzezinski, M. R.; Boutelet-Bochan, H.; Person, R. E.; Fantel, A. G.; Juchau, M. R. (1999).  
21 Catalytic activity and quantitation of cytochrome P-450 2E1 in prenatal human brain. *J*  
22 *Pharmacol Exp Ther*, 289, 1648-1653. <http://www.ncbi.nlm.nih.gov/pubmed/10336564>
- 23 Buben, J. A. and O'Flaherty, E. J. (1985). Delineation of the role of metabolism in the  
24 hepatotoxicity of trichloroethylene and perchloroethylene: A dose-effect study. *Toxicol*  
25 *Appl Pharmacol*, 78, 105-122. <http://www.ncbi.nlm.nih.gov/pubmed/2994252>
- 26 Buckpitt, A.; Chang, A.-M.; Weir, A.; Van Winkle, L.; Duan, X.; Philpot, R.; Plopper, C. (1995).  
27 Relationship of cytochrome P450 activity to Clara cell cytotoxicity IV Metabolism of  
28 naphthalene and naphthalene oxide in microdissected airways from mice, rats, and  
29 hamsters. *Mol Pharmacol*, 47, 74-81. <http://www.ncbi.nlm.nih.gov/pubmed/7838135>
- 30 Budavari, S., (Ed.). (1996). *The Merck Index: An Encyclopedia of Chemicals, Drugs, and*  
31 *Biologicals* (12th ed.). Rahway, NJ: Merck & Co., Inc.
- 32 Bull, R. J. (2000). Mode of action of liver tumor induction by trichloroethylene and its  
33 metabolites, trichloroacetate and dichloroacetate. *Environ Health Perspect*, 108, 241-259.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/10807555>
- 35 Bull, R. J. (2004). Trichloroethylene and liver tumors in mice. *Symposium on New Scientific*  
36 *Research Related to Health Effects of TCE*. Washington, DC.
- 37 Bull, R. J.; Orner, G. A.; Cheng, R. S.; Stillwell, L.; Stauber, A. J.; Sasser, L. B., . . . Thrall, B.  
38 D. (2002). Contribution of dichloroacetate and trichloroacetate to liver tumor induction in  
39 mice by trichloroethylene. *Toxicol Appl Pharmacol*, 182, 55-65.  
40 <http://dx.doi.org/10.1006/taap.2002.9427>
- 41 Bull, R. J.; Sanchez, I. M.; Nelson, M. A.; Larson, J. L.; Lansing, A. J. (1990). Liver tumor  
42 induction in B6C3F1 mice by dichloroacetate and trichloroacetate. *Toxicology*, 63, 341-  
43 359. [http://dx.doi.org/10.1016/0300-483X\(90\)90195-M](http://dx.doi.org/10.1016/0300-483X(90)90195-M)



DRAFT– DO NOT QUOTE OR CITE

- 1 Bull, R. J.; Sasser, L. B.; Lei, X. C. (2004). Interactions in the tumor-promoting activity of  
2 carbon tetrachloride, trichloroacetate, and dichloroacetate in the liver of male B6C3F1  
3 mice. *Toxicology*, 199, 169-183. <http://dx.doi.org/10.1016/j.tox.2004.02.018>
- 4 Bunce, N. J. and Schneider, U. A. (1994). Chemical lifetimes of chlorinated aliphatic priority  
5 pollutants in the Canadian troposphere. *J Photochem Photobiol A*, 81, 93 - 101.
- 6 Burg, J. and Gist, G. (1997). The potential impact on women from environmental exposures. *J*  
7 *Wom Health Gend Base Med*, 6, 159-161.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/9140851>
- 9 Burg, J. and Gist, G. (1999). Health effects of environmental contaminant exposure: an intrafile  
10 comparison of the Trichloroethylene Subregistry. *Arch Environ Health*, 54, 231-241.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/10433181>
- 12 Burg, J. R.; Gist, G.; Allred, S. L.; Rudtke, T. M.; Pallos, L. L.; Cusack, C. D. (1995). The  
13 national exposure registry - morbidity analyses of noncancer outcomes from the  
14 trichloroethylene subregistry baseline data. 4, 237 - 257.
- 15 Burmaster, D. E. (1982). The new pollution: groundwater contamination. *Environment*, 24, 6-  
16 13;33-36.
- 17 Bushnell, P. (1997). Concentration-time relationships for the effects of inhaled trichloroethylene  
18 on signal detection behavior in rats. *Fundam Appl Toxicol*, 36, 30-38.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/9073464>
- 20 Bushnell, P. and Oshiro, W. (2000). Behavioral components of tolerance to repeated inhalation  
21 of trichloroethylene (TCE) in rats. *Neurotoxicol Teratol*, 22, 221-229.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/10758351>
- 23 Butler, T. C. (1949). Metabolic transformations of trichloroethylene. *J Pharmacol Exp Ther*, 97,  
24 84-92.
- 25 Buxton, P. and Hayward, M. (1967). Polyneuritis cranialis associated with industrial  
26 trichloroethylene poisoning. *J Neurol Neurosurg Psychiatry*, 30, 511-518.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/4170824>
- 28 Byers, V. S.; Levin, A. S.; Ozonoff, D. M.; Baldwin, R. W. (1988). Association between clinical  
29 symptoms and lymphocyte abnormalities in a population with chronic domestic exposure  
30 to industrial solvent-contaminated domestic water supply and a high incidence of  
31 leukaemia. *Cancer Immunol Immunother*, 27, 77-81.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/3260823>
- 33 Cai, H. and Guengerich, F. (2001). Reaction of trichloroethylene and trichloroethylene oxide  
34 with cytochrome P450 enzymes: Inactivation and sites of modification. *Chem Res*  
35 *Toxicol*, 14, 451-458. <http://www.ncbi.nlm.nih.gov/pubmed/11304134>
- 36 Cai, H. and Guengerich, F. (2001). Reaction of trichloroethylene oxide with proteins and dna:  
37 instability of adducts and modulation of functions. *Chem Res Toxicol*, 14, 54-61.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/11170508>
- 39 Cai, H. and Guengerich, F. P. (1999). Mechanisms of aqueous decomposition of  
40 trichloroethylene oxide. 121, 11656-11663.
- 41 Cai, P.; König, R.; Boor, P. J.; Kondraganti, S.; Kaphalia, B. S.; Khan, M. F.; Ansari, G. A. S.  
42 (2008). Chronic exposure to trichloroethene causes early onset of SLE-like disease in  
43 female MRL +/+ mice. *Toxicol Appl Pharmacol*, 228, 68-75.  
44 <http://dx.doi.org/10.1016/j.taap.2007.11.031>

DRAFT– DO NOT QUOTE OR CITE

- 1 Cai, P.; König, R.; Khan, M. F.; Kaphalia, B. S.; Ansari, G. A. (2007). Differential immune  
2 responses to albumin adducts of reactive intermediates of trichloroethene in MRL+/  
3 mice. *Toxicol Appl Pharmacol*, 220, 278-283.  
4 <http://dx.doi.org/10.1016/j.taap.2007.01.020>
- 5 Cai, P.; König, R.; Khan, M. F.; Qiu, S.; Kaphalia, B. S.; Ansari, G. A. (2006). Autoimmune  
6 response in MRL+/  
7 mice following treatment with dichloroacetyl chloride or  
8 dichloroacetic anhydride. *Toxicol Appl Pharmacol*, 216, 248-255.  
9 <http://dx.doi.org/10.1016/j.taap.2006.05.010>
- 10 Cakmak, A.; Ekici, A.; Ekici, M.; Arslan, M.; Iteginli, A.; Kurtipek, E.; Kara, T. (2004).  
11 Respiratory findings in gun factory workers exposed to solvents. *Respir Med*, 98, 52-56.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/14959814>
- 13 Caldwell, J. and Keshava, N. (2006). Key issues in the modes of action and effects of  
14 trichloroethylene metabolites for liver and kidney tumorigenesis. *Environ Health*  
15 *Perspect*, 114, 1457-1463. <http://www.ncbi.nlm.nih.gov/pubmed/16966105>
- 16 Caldwell, J. C.; Keshava, N.; Evans, M. V. (2008). Difficulty of mode of action determination  
17 for trichloroethylene: An example of complex interactions of metabolites and other  
18 chemical exposures. *Environ Mol Mutagen*, 49, 142-154.  
19 <http://dx.doi.org/10.1002/em.20350>
- 20 Caldwell, P. T.; Manziello, A.; Howard, J.; Palbykin, B.; Runyan, R. B.; Selmin, O. (2010).  
21 Gene expression profiling in the fetal cardiac tissue after folate and low-dose  
22 trichloroethylene exposure. *Birth Defects Res A Clin Mol Teratol*, 88, 111-127.  
23 <http://dx.doi.org/10.1002/bdra.20631>
- 24 Caldwell, P. T.; Thorne, P. A.; Johnson, P. D.; Boitano, S.; Runyan, R. B.; Selmin, O. (2008).  
25 Trichloroethylene disrupts cardiac gene expression and calcium homeostasis in rat  
26 myocytes. *Toxicol Sci*, 104, 135-143. <http://dx.doi.org/10.1093/toxsci/kfn078>
- 27 Callen, D. F.; Wolf, C. R.; Philpot, R. M. (1980). Cytochrome P-450 mediated genetic activity  
28 and cytotoxicity of seven halogenated aliphatic hydrocarbons in *Saccharomyces*  
29 *cerevisiae*. *Mutat Res Genet Toxicol Environ Mutagen*, 77, 55-63.  
30 [http://dx.doi.org/10.1016/0165-1218\(80\)90120-2](http://dx.doi.org/10.1016/0165-1218(80)90120-2)
- 31 Calvisi, D. F.; Factor, V. M.; Ladu, S.; Conner, E. A.; Thorgeirsson, S. S. (2004). Disruption of  
32 beta-catenin pathway or genomic instability define two distinct categories of liver cancer  
33 in transgenic mice. *Gastroenterology*, 126, 1374-1386.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/15131798>
- 35 Calvisi, D. F.; Ladu, S.; Conner, E. A.; Factor, V. M.; Thorgeirsson, S. S. (2004). Disregulation  
36 of E-cadherin in transgenic mouse models of liver cancer. *Lab Invest*, 84, 1137-1147.  
37 <http://dx.doi.org/10.1038/labinvest.3700147>
- 38 Calvisi, D. F.; Ladu, S.; Gorden, A.; Farina, M.; Conner, E. A.; Lee, J. S., . . . Thorgeirsson, S. S.  
39 (2006). Ubiquitous activation of Ras and Jak/Stat pathways in human HCC.  
40 *Gastroenterology*, 130, 1117-1128. <http://dx.doi.org/10.1053/j.gastro.2006.01.006>
- 41 Calvisi, D. F.; Ladu, S.; Gorden, A.; Farina, M.; Lee, J. S.; Conner, E. A., . . . Thorgeirsson, S. S.  
42 (2007). Mechanistic and prognostic significance of aberrant methylation in the molecular  
43 pathogenesis of human hepatocellular carcinoma. *J Clin Invest*, 117, 2713-2722.  
<http://dx.doi.org/10.1172/JCI31457>

DRAFT– DO NOT QUOTE OR CITE

- 1 Campo, P.; Blachere, V.; Payan, J. P.; Cossec, B.; Ducos, P. (2006). Toluene toxicokinetics and  
2 metabolism parameters in the rat and guinea pig. *Environ Toxicol Pharmacol*, 21, 276-  
3 282.
- 4 Campo, P.; Loquet, G.; Blachère, V.; Roure, M. (1999). Toluene and styrene intoxication route  
5 in the rat cochlea. *Neurotoxicol Teratol*, 21, 427-434.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/10440486>
- 7 Cao, K.; Hollenbach, J.; Shi, X.; Shi, W.; Chopek, M.; Fernández-Viña, M. A. (2001). Analysis  
8 of the frequencies of HLA-A, B, and C alleles and haplotypes in the five major ethnic  
9 groups of the United States reveals high levels of diversity in these loci and contrasting  
10 distribution patterns in these populations. *Hum Immunol*, 62, 1009-1030.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/11543903>
- 12 Cappaert, N. L. M.; Klis, S. F. L.; Baretta, A. B.; Muijser, H.; Smoorenburg, G. F. (2000). Ethyl  
13 benzene-induced ototoxicity in rats: A dose-dependent mid-frequency hearing loss. *J*  
14 *Assoc Res Otolaryngol*, 1, 292-299. <http://dx.doi.org/10.1007/s101620010050>
- 15 Cappaert, N. L. M.; Klis, S. F. L.; Muijser, H.; de Groot, J. C. M. J.; Kulig, B. M.; Smoorenburg,  
16 G. F. (1999). The ototoxic effects of ethyl benzene in rats. *Hear Res*, 137, 91-102.  
17 [http://dx.doi.org/10.1016/S0378-5955\(99\)00141-0](http://dx.doi.org/10.1016/S0378-5955(99)00141-0)
- 18 Carney, E. W.; Thorsrud, B. A.; Dugard, P. H.; Zablony, C. L. (2006). Developmental toxicity  
19 studies in Crl:CD (SD) rats following inhalation exposure to trichloroethylene and  
20 perchloroethylene. *Birth Defects Res B Dev Reprod Toxicol*, 77, 405-412.  
21 <http://dx.doi.org/10.1002/bdrb.20091>
- 22 Carpenter, S. P.; Lasker, J. M.; Raucy, J. L. (1996). Expression, induction, and catalytic activity  
23 of the ethanol-inducible cytochrome P450 (CYP2E1) in human fetal liver and  
24 hepatocytes. *Mol Pharmacol*, 49, 260-268.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/8632758>
- 26 Carrieri, M.; Magosso, D.; Piccoli, P.; Zanetti, E.; Trevisan, A.; Bartolucci, G. B. (2007). Acute,  
27 nonfatal intoxication with trichloroethylene. *Arch Toxicol*, 81, 529-532.  
28 <http://dx.doi.org/10.1007/s00204-007-0180-y>
- 29 Carter, J. H.; Carter, H. W.; DeAngelo, A. B. (1995). Biochemical, pathologic and morphometric  
30 alterations induced in male B6C3F1 mouse liver by short-term exposure to dichloroacetic  
31 acid. *Toxicol Lett*, 81, 55-71. [http://dx.doi.org/10.1016/0378-4274\(95\)03409-9](http://dx.doi.org/10.1016/0378-4274(95)03409-9)
- 32 Carter, J. H.; Carter, H. W.; Deddens, J. A.; Hurst, B. M.; George, M. H.; DeAngelo, A. B.  
33 (2003). A 2-year dose-response study of lesion sequences during hepatocellular  
34 carcinogenesis in the male B6C3F(1) mouse given the drinking water chemical  
35 dichloroacetic acid. *Environ Health Perspect*, 111, 53-64.  
36 <http://dx.doi.org/10.1289/ehp.5442>
- 37 CDC. (Centers for Disease Control and Prevention). (2007). National Vital Statistics Reports:  
38 Deaths: Final data for 2004. Hyattsville, MD. Retrieved from  
39 <http://www.cdc.gov/nchs/products/nvsr.htm>.
- 40 CDHS. (1986). Organic chemical contamination of large public water systems in California.
- 41 Chakrabarti, S. K. and Tuchweber, B. (1988). Studies of acute nephrotoxic potential of  
42 trichloroethylene in Fischer 344 rats. *J Toxicol Environ Health*, 23, 147-158.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/3343692>
- 44 CHALUPA, B.; SYNKOVA, J.; SEVCIK, M. (1960). The assessment of  
45 electroencephalographic changes and memory disturbances in acute intoxications with

- 1 industrial poisons. *Br J Ind Med*, 17, 238-241.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/13692202>
- 3 Chang, L. W.; Daniel, F. B.; DeAngelo, A. B. (1992). Analysis of DNA strand breaks induced in  
4 rodent liver in vivo, hepatocytes in primary culture, and a human cell line by chlorinated  
5 acetic acids and chlorinated acetaldehydes. *Environ Mol Mutagen*, 20, 277-288.  
6 <http://dx.doi.org/10.1002/em.2850200406>
- 7 Chang, Y.; Tai, C.; Lin, R.; Yang, S.; Chen, C.; Shih, T.; Liou, S. (2003). A proportionate cancer  
8 morbidity ratio study of workers exposed to chlorinated organic solvents in Taiwan. *Ind*  
9 *Health*, 41, 77-87. <http://www.ncbi.nlm.nih.gov/pubmed/12725467>
- 10 Chang, Y.; Tai, C.; Yang, S.; Chen, C.; Shih, T.; Lin, R.; Liou, S. (2003). A cohort mortality  
11 study of workers exposed to chlorinated organic solvents in Taiwan. *Ann Epidemiol*, 13,  
12 652-660. [http://dx.doi.org/10.1016/S1047-2797\(03\)00038-3](http://dx.doi.org/10.1016/S1047-2797(03)00038-3)
- 13 Chang, Y.; Tai, C.; Yang, S.; Lin, R.; Sung, F.; Shih, T.; Liou, S. (2005). Cancer incidence  
14 among workers potentially exposed to chlorinated solvents in an electronics factory. *J*  
15 *Occup Health*, 47, 171-180. <http://www.ncbi.nlm.nih.gov/pubmed/15824483>
- 16 Channel, S. R.; Latendresse, J. R.; Kidney, J. K.; Grabau, J. H.; Lane, J. W.; Steel-Goodwin, L.;  
17 Gothaus, M. C. (1998). A subchronic exposure to trichloroethylene causes lipid  
18 peroxidation and hepatocellular proliferation in male B6C3F1 mouse liver. *Toxicol Sci*,  
19 43, 145-154. <http://dx.doi.org/10.1093/toxsci/43.2.145>
- 20 Chapin, R. E. and Sloane, R. A. (1997). Reproductive assessment by continuous breeding:  
21 evolving study design and summaries of ninety studies. *Environ Health Perspect*, 105  
22 Suppl 1, 199-205. <http://www.ncbi.nlm.nih.gov/pubmed/9114287>
- 23 Chapin, R. E.; Norton, R. M.; Popp, J. A.; Bus, J. S. (1982). The effects of 2,5-hexanedione on  
24 reproductive hormones and testicular enzyme activities in the F-344 rat. *Toxicol Appl*  
25 *Pharmacol*, 62, 262-272. <http://www.ncbi.nlm.nih.gov/pubmed/6120587>
- 26 Charbonneau, M.; Couture, J.; Plaa, G. L. (1991). Inhalation versus oral administration of  
27 acetone: effect of the vehicle on the potentiation of CCl4-induced liver injury. *Toxicol*  
28 *Lett*, 57, 47-54.
- 29 Charbotel, B.; Fevotte, J.; Hours, M.; Martin, J. L.; Bergeret, A. (2005). Case-control study on  
30 renal cell cancer and occupational trichloroethylene exposure in the Arve Valley  
31 (France). Lyon, France: Institut Universitaire de Médecine du Travail.
- 32 Charbotel, B.; Fevotte, J.; Hours, M.; Martin, J. L.; Bergeret, A. (2006). Case-control study on  
33 renal cell cancer and occupational exposure to trichloroethylene. Part II: Epidemiological  
34 aspects. *Ann Occup Hyg*, 50, 777-787. <http://dx.doi.org/10.1093/annhyg/mel039>
- 35 Charbotel, B.; Fevotte, J.; Martin, J.-L.; Bergeret, A. (2009). Cancer du rein et expositions au  
36 trichloroéthylène : les valeurs limites d'exposition professionnelle françaises en vigueur  
37 sont-elles adaptées ? *Rev Epidemiol Sante Publique*, 57, 41-47.  
38 <http://dx.doi.org/10.1016/j.respe.2008.09.008>
- 39 Charbotel, B.; Gad, S.; Caïola, D.; Bérout, C.; Fevotte, J.; Bergeret, A., . . . Richard, S. (2007).  
40 Trichloroethylene exposure and somatic mutations of the VHL gene in patients with  
41 Renal Cell Carcinoma. *J Occup Med Toxicol*, 2, 13. [http://dx.doi.org/10.1186/1745-](http://dx.doi.org/10.1186/1745-6673-2-13)  
42 [6673-2-13](http://dx.doi.org/10.1186/1745-6673-2-13)
- 43 Charles, J. M. and Leeming, N. M. (1998). Chronic dietary toxicity/oncogenicity studies on 2,4-  
44 dichlorophenoxybutyric acid in rodents. *Toxicol Sci*, 46, 21-30.  
45 <http://dx.doi.org/10.1006/toxs.1998.2502>

DRAFT– DO NOT QUOTE OR CITE

- 1 Checkoway, H.; Pearce, N.; Crawford-Brown, D. J. (1989). Research methods in occupational  
2 epidemiology. New York: Oxford University Press.
- 3 Chen, C. W. (2000). Biologically based dose-response model for liver tumors induced by  
4 trichloroethylene. *Environ Health Perspect*, 108 Suppl 2, 335-342.  
5 <http://www.ncbi.nlm.nih.gov/pubmed/10807563>
- 6 Chen, C.; Hennig, G.; Whiteley, H.; Manautou, J. (2002). Protection against acetaminophen  
7 hepatotoxicity by clofibrate pretreatment: role of catalase induction. *J Biochem Mol*  
8 *Toxicol*, 16, 227-234. <http://dx.doi.org/10.1002/jbt.10043>
- 9 Chen, X.; Cheung, S. T.; So, S.; Fan, S. T.; Barry, C.; Higgins, J., . . . Brown, P. O. (2002). Gene  
10 expression patterns in human liver cancers. *Mol Biol Cell*, 13, 1929-1939.  
11 <http://dx.doi.org/10.1091/mbc.02-02-0023>.
- 12 Chen, Y.; Cai, J.; Anders, M.; Stevens, J.; Jones, D. (2001). Role of mitochondrial dysfunction in  
13 S-(1,2-dichlorovinyl)-l-cysteine-induced apoptosis. *Toxicol Appl Pharmacol*, 170, 172-  
14 180. <http://dx.doi.org/10.1006/taap.2000.9107>
- 15 Chen, Z. Y.; White, C. C.; He, C. Y.; Liu, Y. F.; Eaton, D. L. (1995). Zonal differences in DNA  
16 synthesis activity and cytochrome P450 gene expression in livers of male F344 rats  
17 treated with five nongenotoxic carcinogens. *J Environ Pathol Toxicol Oncol*, 14, 83-99.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/9372837>
- 19 Cherrie, J.; Kromhout, H.; Semple, S. (2001). The importance of reliable exposure estimates in  
20 deciding whether trichloroethylene can cause kidney cancer. *J Cancer Res Clin Oncol*,  
21 127, 400-404. <http://www.ncbi.nlm.nih.gov/pubmed/11414201>
- 22 Cheung, C.; Akiyama, T. E.; Ward, J. M.; Nicol, C. J.; Feigenbaum, L.; Vinson, C.; Gonzalez, F.  
23 J. (2004). Diminished hepatocellular proliferation in mice humanized for the nuclear  
24 receptor peroxisome proliferator-activated receptor alpha. *Cancer Res*, 64, 3849-3854.  
25 <http://dx.doi.org/10.1158/0008-5472.CAN-04-0322>
- 26 Chia, S. E.; Goh, V. H.; Ong, C. N. (1997). Endocrine profiles of male workers with exposure to  
27 trichloroethylene. *Am J Ind Med*, 32, 217-222. [http://dx.doi.org/10.1002/\(SICI\)1097-0274\(199709\)32:3<217::AID-AJIM6>3.0.CO;2-U](http://dx.doi.org/10.1002/(SICI)1097-0274(199709)32:3<217::AID-AJIM6>3.0.CO;2-U)
- 28 Chia, S.-E.; Ong, C.-N.; Tsakok, M. F.; Ho, A. (1996). Semen parameters in workers exposed to  
29 trichloroethylene. *Reprod Toxicol*, 10, 295-299. [http://dx.doi.org/10.1016/0890-6238\(96\)00058-5](http://dx.doi.org/10.1016/0890-6238(96)00058-5)
- 30  
31
- 32 Chieco, P.; Moslen, M. T.; Reynolds, E. S. (1981). Effect of administrative vehicle on oral 1,1-  
33 dichloroethylene toxicity. *Toxicol Appl Pharmacol*, 57, 146-155.
- 34 Chiesa, R.; Donato, F.; Tagger, A.; Favret, M.; Ribero, M. L.; Nardi, G., . . . Callea, F. (2000).  
35 Etiology of hepatocellular carcinoma in Italian patients with and without cirrhosis.  
36 *Cancer Epidemiol Biomarkers Prev*, 9, 213-216.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/10698484>
- 38 Chiu, B. C. and Weisenburger, D. D. (2003). An update of the epidemiology of non-Hodgkin's  
39 lymphoma. 4, 161-168. <http://www.ncbi.nlm.nih.gov/pubmed/14715098>
- 40 Chiu, W. A. (2006). Statistical issues in physiologically based pharmacokinetic modeling  
41 Toxicokinetics And Risk Assessment (pp. 269). New York: Informa Healthcare.
- 42 Chiu, W. A. (In Press). Trichloroacetic acid: Updated estimates of its bioavailability and its  
43 contribution to trichloroethylene-induced mouse hepatomegaly.  
44 <http://dx.doi.org/10.1016/j.tox.2011.04.009>

- 1 Chiu, W. A. and Bois, F. Y. (2007). An Approximate Method for Population Toxicokinetic  
2 Analysis With Aggregated Data. *Journal of Agricultural, Biological, and Environmental*  
3 *Statistics*, 12, 346-363. <http://dx.doi.org/10.1198/108571107X229340>
- 4 Chiu, W. A. and White, P. (2004). *PBPK Model-Based Route Extrapolation: Exact and*  
5 *Approximate Solutions for Use in Risk Assessment*.
- 6 Chiu, W. A. and White, P. (2006). Steady-state solutions to PBPK models and their applications  
7 to risk assessment I: Route-to-route extrapolation of volatile chemicals. *Risk Anal*, 26,  
8 769-780. <http://dx.doi.org/10.1111/j.1539-6924.2006.00762.x>
- 9 Chiu, W. A.; Micallef, S.; Monster, A. C.; Bois, F. Y. (2007). Toxicokinetics of inhaled  
10 trichloroethylene and tetrachloroethylene in humans at 1 ppm: empirical results and  
11 comparisons with previous studies. *Toxicol Sci*, 95, 23-36.  
12 <http://dx.doi.org/10.1093/toxsci/kfl129>
- 13 Chiu, W. A.; Okino, M. S.; Evans, M. V. (2009). Characterizing uncertainty and population  
14 variability in the toxicokinetics of trichloroethylene and metabolites in mice, rats, and  
15 humans using an updated database, physiologically based pharmacokinetic (PBPK)  
16 model, and Bayesian approach. *Toxicol Appl Pharmacol*, 241, 36-60.  
17 <http://dx.doi.org/10.1016/j.taap.2009.07.032>
- 18 Chiu, W. A.; Okino, M. S.; Lipscomb, J. C.; Evans, M. V. (2006). Issues in the pharmacokinetics  
19 of trichloroethylene and its metabolites. *Environ Health Perspect*, 114, 1450-1456.  
20 <http://dx.doi.org/10.1289/ehp.8691>
- 21 Chiu, W.; Caldwell, J.; Keshava, N.; Scott, C. (2006). Key scientific issues in the health risk  
22 assessment of trichloroethylene. *Environ Health Perspect*, 114, 1445-1449.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/16966103>
- 24 Cho, W. C. (2007). OncomiRs: the discovery and progress of microRNAs in cancers. *Mol*  
25 *Cancer*, 6, 60. <http://dx.doi.org/10.1186/1476-4598-6-60>
- 26 Chow, W. H. and Devesa, S. S. (2008). Contemporary epidemiology of renal cell cancer. *Cancer*  
27 *J*, 14, 288-301. <http://dx.doi.org/10.1097/PPO.0b013e3181867628>
- 28 Chow, W.; Gridley, G.; McLaughlin, J.; Mandel, J.; Wacholder, S.; Blot, W., . . . Fraumeni, J.  
29 (1994). Protein intake and risk of renal cell cancer. *J Natl Cancer Inst*, 86, 1131-1139.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/8028035>
- 31 Cicmanec, J.; Condie, L.; Olson, G.; Wang, S. (1991). 90-Day toxicity study of dichloroacetate  
32 in dogs. *Fundam Appl Toxicol*, 17, 376-389.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/1765225>
- 34 Clapp, R. W. and Hoffman, K. (2008). Cancer mortality in IBM Endicott plant workers, 1969-  
35 2001: an update on a NY production plant. *Environ Health*, 7, 13.  
36 <http://dx.doi.org/10.1186/1476-069X-7-13>
- 37 Clay, P. (2008). Assessment of the genotoxicity of trichloroethylene and its metabolite, S-(1,2-  
38 dichlorovinyl)-L-cysteine (DCVC), in the comet assay in rat kidney. *Mutagenesis*, 23,  
39 27-33. <http://dx.doi.org/10.1093/mutage/gem034>
- 40 Clegg, E. D.; Cook, J. C.; Chapin, R. E.; Foster, P. M. D.; Daston, G. P. (1997). Leydig cell  
41 hyperplasia and adenoma formation: Mechanisms and relevance to humans. *Reprod*  
42 *Toxicol*, 11, 107-121. [http://dx.doi.org/10.1016/S0890-6238\(96\)00203-1](http://dx.doi.org/10.1016/S0890-6238(96)00203-1)
- 43 Clewell, H. J. and Andersen, M. E. (2004). Applying mode-of-action and pharmacokinetic  
44 considerations in contemporary cancer risk assessments: An example with

- 1 trichloroethylene. *Crit Rev Toxicol*, 34, 385-445.  
2 <http://dx.doi.org/10.1080/10408440490500795>
- 3 Clewell, H. J., III; Gentry, P. R.; Covington, T. R.; Gearhart, J. M. (2000). Development of a  
4 physiologically based pharmacokinetic model of trichloroethylene and its metabolites for  
5 use in risk assessment. *Environ Health Perspect*, 108, 283-305.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/10807559>
- 7 Clewell, H.; Andersen, M.; Barton, H. (2002). A consistent approach for the application of  
8 pharmacokinetic modeling in cancer and noncancer risk assessment. *Environ Health*  
9 *Perspect*, 110, 85-93. <http://www.ncbi.nlm.nih.gov/pubmed/11781169>
- 10 Coberly, S.; Oudiz, D. J.; Overstreet, J. W.; Wiley, L. M. (1992). Effects of maternal exposure to  
11 trichloroethylene (TCE) on cell proliferation in the mouse preimplantation embryo.  
12 *Reprod Toxicol*, 6, 241-245. <http://www.ncbi.nlm.nih.gov/pubmed/1591481>
- 13 Cocco, P.; t'Mannetje, A.; Fadda, D.; Melis, M.; Becker, N.; de Sanjosé, S., . . . Boffetta, P.  
14 (2010). Occupational exposure to solvents and risk of lymphoma subtypes: results from  
15 the Epilymph case-control study. *Occup Environ Med*, 67, 341-347.  
16 <http://dx.doi.org/10.1136/oem.2009.046839>
- 17 Cogger, V. C.; Warren, A.; Fraser, R.; Ngu, M.; McLean, A. J.; Le Couteur, D. G. (2003).  
18 Hepatic sinusoidal pseudocapillarization with aging in the non-human primate. *Exp*  
19 *Gerontol*, 38, 1101-1107. <http://www.ncbi.nlm.nih.gov/pubmed/14580863>
- 20 Cohn, P.; Fagliano, J.; Klotz, J. (1994). Assessing human health effects from chemical  
21 contaminants in drinking water. 91, 719-722.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/7800286>
- 23 Cohn, P.; Klotz, J.; Bove, F.; Berkowitz, M.; Fagliano, J. (1994). Drinking water contamination  
24 and the incidence of leukemia and non-Hodgkin's lymphoma. *Environ Health Perspect*,  
25 102, 556-561. <http://www.ncbi.nlm.nih.gov/pubmed/9679115>
- 26 Cojocel, C.; Beuter, W.; Müller, W.; Mayer, D. (1989). Lipid peroxidation: A possible  
27 mechanism of trichloroethylene-induced nephrotoxicity. *Toxicology*, 55, 131-141.  
28 [http://dx.doi.org/10.1016/0300-483X\(89\)90180-7](http://dx.doi.org/10.1016/0300-483X(89)90180-7)
- 29 Coleridge, H. M.; Coleridge, J. C. G.; Luck, J. C.; Norman, J. (1968). The effect of four volatile  
30 anaesthetic agents on the impulse activity of two types of pulmonary receptor. *Br J*  
31 *Anaesth*, 40, 484-492.
- 32 Collier, J.; Selmin, O.; Johnson, P.; Runyan, R. (2003). Trichloroethylene effects on gene  
33 expression during cardiac development. *Birth Defects Res A Clin Mol Teratol*, 67, 488-  
34 495. <http://dx.doi.org/10.1002/bdra.10073>
- 35 Colt, J. and Blair, A. (1998). Parental occupational exposures and risk of childhood cancer.  
36 *Environ Health Perspect*, 106 Suppl 3, 909-925.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/9646055>
- 38 Columbano, A. and Ledda-Columbano, G. M. (2003). Mitogenesis by ligands of nuclear  
39 receptors: an attractive model for the study of the molecular mechanisms implicated in  
40 liver growth. *Cell Death Differ*, 10 Suppl 1, S19-S21.  
41 <http://dx.doi.org/10.1038/sj.cdd.4401113>
- 42 Columbano, A.; Ledda-Columbano, G. M.; Coni, P. P.; Faa, G.; Liguori, C.; Santa Cruz, G.;  
43 Pani, P. (1985). Occurrence of cell death (apoptosis) during the involution of liver  
44 hyperplasia. *Lab Invest*, 52, 670-675. <http://www.ncbi.nlm.nih.gov/pubmed/4010225>

- 1 Columbano, A.; Ledda-Columbano, G. M.; Ennas, M. G.; Curto, M.; Chelo, A.; Pani, P. (1990).  
2 Cell proliferation and promotion of rat liver carcinogenesis: Different effect of hepatic  
3 regeneration and mitogen induced hyperplasia on the development of enzyme-altered  
4 foci. *Carcinogenesis*, 11, 771-776. <http://dx.doi.org/10.1093/carcin/11.5.771>
- 5 Columbano, A.; Ledda-Columbano, G. M.; Lee, G.; Rajalakshmi, S.; Sarma, D. S. (1987).  
6 Inability of mitogen-induced liver hyperplasia to support the induction of enzyme-altered  
7 islands induced by liver carcinogens. *Cancer Res*, 47, 5557-5559.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/2889525>
- 9 Colvin, M.; Myers, J.; Nell, V.; Rees, D.; Cronje, R. (1993). A cross-sectional survey of  
10 neurobehavioral effects of chronic solvent exposure on workers in a paint manufacturing  
11 plant. *Environ Res*, 63, 122-132. <http://dx.doi.org/10.1006/enrs.1993.1134>
- 12 Commandeur, J. and Vermeulen, N. (1990). Identification of N-acetyl(2,2-dichlorovinyl)- and N-  
13 acetyl(1,2-dichlorovinyl)-L-cysteine as two regioisomeric mercapturic acids of  
14 trichloroethylene in the rat. *Chem Res Toxicol*, 3, 212-218.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/1966701>
- 16 Coni, P.; Simbula, G.; De Prati, A. C.; Menegazzi, M.; Suzuki, H.; Sarma, D. S. R., . . .  
17 Columbano, A. (1993). Differences in the steady-state levels of c-fos, c-jun and c-myc  
18 messenger RNA during mitogen-induced liver growth and compensatory regeneration.  
19 *Hepatology*, 17, 1109-1116.
- 20 Conner, E. A.; Lemmer, E. R.; Omori, M.; Wirth, P. J.; Factor, V. M.; Thorgeirsson, S. S.  
21 (2000). Dual functions of E2F-1 in a transgenic mouse model of liver carcinogenesis.  
22 *Oncogene*, 19, 5054-5062. <http://dx.doi.org/10.1038/sj.onc.1203885>
- 23 Conner, E. A.; Lemmer, E. R.; Sánchez, A.; Factor, V. M.; Thorgeirsson, S. S. (2003). E2F1  
24 blocks and c-Myc accelerates hepatic ploidy in transgenic mouse models. *Biochem*  
25 *Biophys Res Commun*, 302, 114-120. <http://www.ncbi.nlm.nih.gov/pubmed/12593856>
- 26 Contrera, J. F.; Jacobs, A. C.; DeGeorge, J. J. (1997). Carcinogenicity testing and the evaluation  
27 of regulatory requirements for pharmaceuticals. *Regul Toxicol Pharmacol*, 25, 130-145.  
28 <http://dx.doi.org/10.1006/rtph.1997.1085>
- 29 Cook, J. C.; Klinefelter, G. R.; Hardisty, J. F.; Sharpe, R. M.; Foster, P. M. D. (1999). Rodent  
30 leydig cell tumorigenesis: A review of the physiology, pathology, mechanisms and  
31 relevance to humans. *Crit Rev Toxicol*, 29, 169-261.
- 32 Cooper, A. (2004). The role of glutamine transaminase K (GTK) in sulfur and alpha-keto acid  
33 metabolism in the brain, and in the possible bioactivation of neurotoxicants. *Neurochem*  
34 *Int*, 44, 557-577. <http://dx.doi.org/10.1016/j.neuint.2003.12.002>
- 35 Cooper, G. and Stroehla, B. (2003). The epidemiology of autoimmune diseases. *Autoimmun*  
36 *Rev*, 2, 119-125. <http://www.ncbi.nlm.nih.gov/pubmed/12848952>
- 37 Cooper, G. S.; Makris, S. L.; Nietert, P. J.; Jinot, J. (2009). Evidence of autoimmune-related  
38 effects of trichloroethylene exposure from studies in mice and humans. *Environ Health*  
39 *Perspect*, 117, 696-702. <http://dx.doi.org/10.1289/ehp.11782>
- 40 Cooper, G. S.; Parks, C. G.; Treadwell, E. L.; St Clair, E. W.; Gilkeson, G. S.; Dooley, M. A.  
41 (2004). Occupational risk factors for the development of systemic lupus erythematosus. *J*  
42 *Rheumatol*, 31, 1928-1933. <http://www.ncbi.nlm.nih.gov/pubmed/15468355>
- 43 Cooper, G.; Miller, F.; Pandey, J. (1999). The role of genetic factors in autoimmune disease:  
44 implications for environmental research. *Environ Health Perspect*, 107 Suppl 5, 693-700.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/10502533>



DRAFT– DO NOT QUOTE OR CITE

- 1 Coopman, V.; Cordonnier, J.; De Letter, E.; Piette, M. (2003). Tissue distribution of  
2 trichloroethylene in a case of accidental acute intoxication by inhalation. *Forensic Sci Int*,  
3 134, 115-119. <http://www.ncbi.nlm.nih.gov/pubmed/12850404>
- 4 Copeland, K. T.; Checkoway, H.; McMichael, A. J.; Holbrook, R. H. (1977). Bias due to  
5 misclassification in the estimation of relative risk. *Am J Epidemiol*, 105, 488-495.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/871121>
- 7 Cornett, R.; James, M.; Henderson, G.; Cheung, J.; Shroads, A.; Stacpoole, P. (1999). Inhibition  
8 of glutathione S-transferase zeta and tyrosine metabolism by dichloroacetate: A potential  
9 unifying mechanism for its altered biotransformation and toxicity. *Biochem Biophys Res*  
10 *Commun*, 262, 752-756. <http://dx.doi.org/10.1006/bbrc.1999.1287>
- 11 Corton, J. C. (2008). Evaluation of the role of peroxisome proliferator-activated receptor alpha  
12 (PPARalpha) in mouse liver tumor induction by trichloroethylene and metabolites. *Crit*  
13 *Rev Toxicol*, 38, 857-875. <http://dx.doi.org/10.1080/10408440802209796>
- 14 Corton, J. C. and Lapinskas, P. J. (2005). Peroxisome proliferator-activated receptors: Mediators  
15 of phthalate ester-induced effects in the male reproductive tract? *Toxicol Sci*, 83, 4-17.  
16 <http://dx.doi.org/10.1093/toxsci/kfi011>
- 17 Cosby, N. C. and Dukelow, W. R. (1992). Toxicology of maternally ingested trichloroethylene  
18 (TCE) on embryonal and fetal development in mice and of TCE metabolites on in vitro  
19 fertilization. *Toxicol Sci*, 19, 268-274. <http://www.ncbi.nlm.nih.gov/pubmed/1516784>
- 20 Costa, A. K. and Ivanetich, K. M. (1984). Chlorinated ethylenes: their metabolism and effect on  
21 DNA repair in rat hepatocytes. *Carcinogenesis*, 5, 1629-1636.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/6499114>
- 23 Costa, G.; Merletti, F.; Segnan, N. (1989). A mortality cohort study in a north Italian aircraft  
24 factory. *Br J Ind Med*, 46, 738-743. <http://www.ncbi.nlm.nih.gov/pubmed/2818961>
- 25 Costantini, A. S.; Benvenuti, A.; Vineis, P.; Kriebel, D.; Tumino, R.; Ramazzotti, V., . . . Miligi,  
26 L. (2008). Risk of leukemia and multiple myeloma associated with exposure to benzene  
27 and other organic solvents: Evidence from the Italian Multicenter Case-control study. *Am*  
28 *J Ind Med*, 51, 803-811. <http://dx.doi.org/10.1002/ajim.20592>
- 29 Costas, K.; Knorr, R. S.; Condon, S. K. (2002). A case-control study of childhood leukemia in  
30 Woburn, Massachusetts: The relationship between leukemia incidence and exposure to  
31 public drinking water. *Sci Total Environ*, 300, 23-35. [http://dx.doi.org/10.1016/S0048-](http://dx.doi.org/10.1016/S0048-9697(02)00169-9)  
32 [9697\(02\)00169-9](http://dx.doi.org/10.1016/S0048-9697(02)00169-9)
- 33 Costet, P.; Legendre, C.; Moré, J.; Edgar, A.; Galtier, P.; Pineau, T. (1998). Peroxisome  
34 proliferator-activated receptor alpha-isoform deficiency leads to progressive dyslipidemia  
35 with sexually dimorphic obesity and steatosis. *J Biol Chem*, 273, 29577-29585.  
36 <http://www.ncbi.nlm.nih.gov/pubmed/9792666>
- 37 Counts, J.; Sarmiento, J.; Harbison, M.; Downing, J.; McClain, R.; Goodman, J. (1996). Cell  
38 proliferation and global methylation status changes in mouse liver after phenobarbital  
39 and/or choline-devoid, methionine-deficient diet administration. *Carcinogenesis*, 17,  
40 1251-1257. <http://www.ncbi.nlm.nih.gov/pubmed/8681439>
- 41 Coyle, Y. M.; Hynan, L. S.; Euhus, D. M.; Minhajuddin, A. T. (2005). An ecological study of the  
42 association of environmental chemicals on breast cancer incidence in Texas. *Breast*  
43 *Cancer Res Treat*, 92, 107-114. <http://dx.doi.org/10.1007/s10549-004-8268-z>
- 44 Crabb, D.; Pinaire, J.; Chou, W.; Sissom, S.; Peters, J.; Harris, R.; Stewart, M. (2001).  
45 Peroxisome proliferator-activated receptors (PPAR) and the mitochondrial aldehyde

DRAFT– DO NOT QUOTE OR CITE

- 1 dehydrogenase (ALDH2) promoter in vitro and in vivo. *Alcohol Clin Exp Res*, 25, 945-  
2 952. <http://www.ncbi.nlm.nih.gov/pubmed/11505017>
- 3 Crebelli, R. and Carere, A. (1989). Genetic toxicology of 1,1,2-trichloroethylene. *Mutat Res*,  
4 221, 11-37. [http://dx.doi.org/10.1016/0165-1110\(89\)90043-2](http://dx.doi.org/10.1016/0165-1110(89)90043-2)
- 5 Crebelli, R.; Bignami, M.; Conti, L.; Carere, A. (1982). Mutagenicity of trichloroethylene in  
6 *Salmonella typhimurium* TA100. *Ann Ist Super Sanita*, 18, 117-121.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/6763484>
- 8 Crebelli, R.; Conti, G.; Conti, L.; Carere, A. (1985). Mutagenicity of trichloroethylene,  
9 trichloroethanol and chloral hydrate in *Aspergillus nidulans*. *Mutat Res Genet Toxicol*  
10 *Environ Mutagen*, 155, 105-111. [http://dx.doi.org/10.1016/0165-1218\(85\)90126-0](http://dx.doi.org/10.1016/0165-1218(85)90126-0)
- 11 Crebelli, R.; Conti, G.; Conti, L.; Carere, A. (1991). In vitro studies with nine known or  
12 suspected spindle poisons: results in tests for chromosome malsegregation in *Aspergillus*  
13 *nidulans*. *Mutagenesis*, 6, 131-136. <http://www.ncbi.nlm.nih.gov/pubmed/2056914>
- 14 Crofton, K. and Zhao, X. (1997). The ototoxicity of trichloroethylene: extrapolation and  
15 relevance of high-concentration, short-duration animal exposure data. *Fundam Appl*  
16 *Toxicol*, 38, 101-106. <http://www.ncbi.nlm.nih.gov/pubmed/9268609>
- 17 Crofton, K. M. and Zhao, X. (1993). Mid-frequency hearing loss in rats following inhalation  
18 exposure to trichloroethylene: evidence from reflex modification audiometry.  
19 *Neurotoxicol Teratol*, 15, 413-423. <http://www.ncbi.nlm.nih.gov/pubmed/8302243>
- 20 Crofton, K. M.; Lassiter, T. L.; Rebert, C. S. (1994). Solvent-induced ototoxicity in rats: an  
21 atypical selective mid-frequency hearing deficit. *Hear Res*, 80, 25-30.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/7852200>
- 23 Crump, K.; Allen, B.; Shipp, A. (1989). Choice of dose measure for extrapolating carcinogenic  
24 risk from animals to humans: An empirical investigation of 23 chemicals. *Health Phys*,  
25 57, 387-393. <http://www.ncbi.nlm.nih.gov/pubmed/2606698>
- 26 Cruzan, G.; Bus, J.; Banton, M.; Gingell, R.; Carlson, G. (2009). Mouse specific lung tumors  
27 from CYP2F2-mediated cytotoxic metabolism: an endpoint/toxic response where data  
28 from multiple chemicals converge to support a mode of action. *Regul Toxicol Pharmacol*,  
29 55, 205-218. <http://dx.doi.org/10.1016/j.yrtph.2009.07.002>
- 30 Cummings, B. S. and Lash, L. H. (2000). Metabolism and toxicity of trichloroethylene and S-  
31 (1,2-dichlorovinyl)-L-cysteine in freshly isolated human proximal tubular cells. *Toxicol*  
32 *Sci*, 53, 458-466. <http://www.ncbi.nlm.nih.gov/pubmed/10696794>
- 33 Cummings, B. S.; Lasker, J. M.; Lash, L. H. (2000). Expression of glutathione-dependent  
34 enzymes and cytochrome P450s in freshly isolated and primary cultures of proximal  
35 tubular cells from human kidney. *J Pharmacol Exp Ther*, 293, 677-685.  
36 <http://www.ncbi.nlm.nih.gov/pubmed/10773044>
- 37 Cummings, B. S.; Parker, J. C.; Lash, L. H. (2000). Role of cytochrome P450 and glutathione S-  
38 transferase alpha in the metabolism and cytotoxicity of trichloroethylene in rat kidney.  
39 *Biochem Pharmacol*, 59, 531-543. [http://dx.doi.org/10.1016/S0006-2952\(99\)00374-3](http://dx.doi.org/10.1016/S0006-2952(99)00374-3)
- 40 Cummings, B. S.; Parker, J. C.; Lash, L. H. (2001). Cytochrome p450-dependent metabolism of  
41 trichloroethylene in rat kidney. *Toxicol Sci*, 60, 11-19.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/11222868>
- 43 Cummings, B. S.; Zangar, R. C.; Novak, R. F.; Lash, L. H. (1999). Cellular distribution of  
44 cytochromes P-450 in the rat kidney. *Drug Metab Dispos*, 27, 542-548.

DRAFT– DO NOT QUOTE OR CITE

- 1 Cummings, B.; Zangar, R.; Novak, R.; Lash, L. (2000). Cytotoxicity of trichloroethylene and S-  
2 (1, 2-dichlorovinyl)-L-cysteine in primary cultures of rat renal proximal tubular and distal  
3 tubular cells. *Toxicology*, 150, 83-98. <http://www.ncbi.nlm.nih.gov/pubmed/10996665>
- 4 Cutler, J.; Parker, G.; Rosen, S.; Prenney, B.; Healey, R.; Caldwell, G. (1986). Childhood  
5 leukemia in Woburn, Massachusetts. *Public Health Rep*, 101, 201-205.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/3083476>
- 7 Dahlgren, J.; Takhar, H.; Anderson-Mahoney, P.; Kotlerman, J.; Tarr, J.; Warshaw, R. (2007).  
8 Cluster of systemic lupus erythematosus (SLE) associated with an oil field waste site: a  
9 cross sectional study. *Environ Health*, 6, 8. <http://dx.doi.org/10.1186/1476-069X-6-8>
- 10 Dai, X.; De Souza, A. T.; Dai, H.; Lewis, D. L.; Lee, C. K.; Spencer, A. G., . . . He, Y. D.  
11 (2007). PPARalpha siRNA-treated expression profiles uncover the causal sufficiency  
12 network for compound-induced liver hypertrophy. 3, e30.  
13 <http://dx.doi.org/10.1371/journal.pcbi.0030030>
- 14 Dai, Y.; Leng, S.; Li, L.; Niu, Y.; Huang, H.; Cheng, J.; Zheng, Y. (2004). Genetic  
15 polymorphisms of cytokine genes and risk for trichloroethylene-induced severe  
16 generalized dermatitis: a case-control study. *Biomarkers*, 9, 470-478.  
17 <http://dx.doi.org/10.1080/13547500400026920>
- 18 Dajee, M.; Lazarov, M.; Zhang, J. Y.; Cai, T.; Green, C. L.; Russell, A. J., . . . Khavari, P. A.  
19 (2003). NF-kappaB blockade and oncogenic Ras trigger invasive human epidermal  
20 neoplasia. *Nature*, 421, 639-643. <http://dx.doi.org/10.1038/nature01283>
- 21 Dalbey, W. and Bingham, E. (1978). Metabolism of trichloroethylene by the isolated perfused  
22 lung. *Toxicol Appl Pharmacol*, 43, 267-277.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/635916>
- 24 Dallas, C. E. (1991). Validation and application of pharmacokinetic models for interspecies  
25 extrapolations in toxicity risk assessments of volatile organics.
- 26 Dallas, C.; Gallo, J.; Ramanathan, R.; Muralidhara, S.; Bruckner, J. (1991). Physiological  
27 pharmacokinetic modeling of inhaled trichloroethylene in rats. *Toxicol Appl Pharmacol*,  
28 110, 303-314. <http://www.ncbi.nlm.nih.gov/pubmed/1891776>
- 29 Daniel, F. B.; DeAngelo, A. B.; Stober, J. A.; Olson, G. R.; Page, N. P. (1992).  
30 Hepatocarcinogenicity of chloral hydrate, 2-chloroacetaldehyde, and dichloroacetic acid  
31 in the male B6C3F1 mouse. *Fundam Appl Toxicol*, 19, 159-168.  
32 [http://dx.doi.org/10.1016/0272-0590\(92\)90147-A](http://dx.doi.org/10.1016/0272-0590(92)90147-A)
- 33 Daniell, W. E.; Claypoole, K. H.; Checkoway, H.; Smith-Weller, T.; Dager, S. R.; Townes, B.  
34 D.; Rosenstock, L. (1999). Neuropsychological function in retired workers with previous  
35 long-term occupational exposure to solvents. *Occup Environ Med*, 56, 93-105.  
36 <http://www.ncbi.nlm.nih.gov/pubmed/10448313>
- 37 Darnerud, P.; Brandt, I.; Feil, V.; Bakke, J. (1989). Dichlorovinyl cysteine (DCVC) in the mouse  
38 kidney: tissue-binding and toxicity after glutathione depletion and probenecid treatment.  
39 *Arch Toxicol*, 63, 345-350. <http://www.ncbi.nlm.nih.gov/pubmed/2818197>
- 40 Das, R. M. and Scott, J. E. (1994). Trichloroethylene-induced pneumotoxicity in fetal and  
41 neonatal mice. *Toxicol Lett*, 73, 227-239. <http://www.ncbi.nlm.nih.gov/pubmed/8091431>
- 42 David, R. M.; Clewell, H. J.; Gentry, P. R.; Covington, T. R.; Morgott, D. A.; Marino, D. J.  
43 (2006). Revised assessment of cancer risk to dichloromethane II. Application of  
44 probabilistic methods to cancer risk determinations. *Regul Toxicol Pharmacol*, 45, 55-65.  
45 <http://dx.doi.org/10.1016/j.yrtph.2005.12.003>

DRAFT– DO NOT QUOTE OR CITE

- 1 David, R. M.; Moore, M. R.; Cifone, M. A.; Finney, D. C.; Guest, D. (1999). Chronic  
2 peroxisome proliferation and hepatomegaly associated with the hepatocellular  
3 tumorigenesis of di(2-ethylhexyl)phthalate and the effects of recovery. *Toxicol Sci*, 50,  
4 195-205. <http://www.ncbi.nlm.nih.gov/pubmed/10478855>
- 5 Davidson, I. W. F. and Beliles, R. P. (1991). Consideration of the target organ toxicity of  
6 trichloroethylene in terms of metabolite toxicity and pharmacokinetics. *Drug Metab Rev*,  
7 23, 493-599. <http://dx.doi.org/10.3109/03602539109029772>
- 8 Davis, R. R.; Murphy, W. J.; Snawder, J. E.; Striley, C. A.; Henderson, D.; Khan, A.; Krieg, E.  
9 F. (2002). Susceptibility to the ototoxic properties of toluene is species specific. *Hear*  
10 *Res*, 166, 24-32. <http://www.ncbi.nlm.nih.gov/pubmed/12062755>
- 11 Davis, S.; Laszlo Pallos, L.; Wu, J.; Sapp, J.; Cusack, C. (2005). ATSDR's trichloroethylene  
12 subregistry methods and results: 1989-2000. *Arch Environ Occup Health*, 60, 130-139.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/17153085>
- 14 Dawson, B. V.; Johnson, P. D.; Goldberg, S. J.; Ulreich, J. B. (1990). Cardiac teratogenesis of  
15 trichloroethylene and dichloroethylene in a mammalian model. *J Am Coll Cardiol*, 16,  
16 1304-1309.
- 17 Dawson, B.; Johnson, P.; Goldberg, S.; Ulreich, J. (1993). Cardiac teratogenesis of halogenated  
18 hydrocarbon-contaminated drinking water. *J Am Coll Cardiol*, 21, 1466-1472.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/8473657>
- 20 De Baere, S.; Meyer, E.; Dirinck, I.; Lambert, W.; Piette, M.; Van Peteghem, C.; De Leenheer,  
21 A. (1997). Tissue distribution of trichloroethylene and its metabolites in a forensic case. *J*  
22 *Anal Toxicol*, 21, 223-227. <http://www.ncbi.nlm.nih.gov/pubmed/9171208>
- 23 de Jong, P. and Brenner, B. (2004). From secondary to primary prevention of progressive renal  
24 disease: the case for screening for albuminuria. *Kidney Int*, 66, 2109-2118.  
25 <http://dx.doi.org/10.1111/j.1523-1755.2004.66001.x>
- 26 De Roos, A. J.; Olshan, A. F.; Teschke, K.; Poole, C.; Savitz, D. A.; Blatt, J., . . . Pollock, B. H.  
27 (2001). Parental occupational exposures to chemicals and incidence of neuroblastoma in  
28 offspring. *Am J Epidemiol*, 154, 106-114.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/11447042>
- 30 De Roos, A.; Olshan, A.; Teschke, K.; Poole, C.; Savitz, D.; Blatt, J., . . . Pollock, B. (2001).  
31 Parental occupational exposures to chemicals and incidence of neuroblastoma in  
32 offspring. *Am J Epidemiol*, 154, 106-114. <http://dx.doi.org/10.1093/aje/154.2.106>
- 33 Deane, M.; Swan, S.; Harris, J.; Epstein, D.; Neutra, R. (1989). Adverse pregnancy outcomes in  
34 relation to water contamination, Santa Clara County, California, 1980-1981. *Am J*  
35 *Epidemiol*, 129, 894-904. <http://www.ncbi.nlm.nih.gov/pubmed/2705433>
- 36 DeAngelo, A. B.; Daniel, F. B.; McMillan, L.; Wernsing, P.; Savage, R. E. J. (1989). Species  
37 and strain sensitivity to the induction of peroxisome proliferation by chloroacetic acids.  
38 *Toxicol Appl Pharmacol*, 101, 285-298. [http://dx.doi.org/10.1016/0041-008X\(89\)90277-](http://dx.doi.org/10.1016/0041-008X(89)90277-9)  
39 9
- 40 DeAngelo, A. B.; Daniel, F. B.; Most, B. M.; Olson, G. R. (1996). The carcinogenicity of  
41 dichloroacetic acid in the male fischer 344 rat. *Toxicology*, 114, 207-221.  
42 [http://dx.doi.org/10.1016/S0300-483X\(96\)03510-X](http://dx.doi.org/10.1016/S0300-483X(96)03510-X)
- 43 DeAngelo, A. B.; Daniel, F. B.; Most, B. M.; Olson, G. R. (1997). Failure of monochloroacetic  
44 acid and trichloroacetic acid administered in the drinking water to produce liver cancer in

DRAFT– DO NOT QUOTE OR CITE

- 1 male F344/N rats. *J Toxicol Environ Health*, 52, 425-445.  
2 <http://dx.doi.org/10.1080/00984109708984074>
- 3 DeAngelo, A. B.; Daniel, F. B.; Wong, D. M.; George, M. H. (2008). The induction of  
4 hepatocellular neoplasia by trichloroacetic acid administered in the drinking water of the  
5 male B6C3F1 mouse. *J Toxicol Environ Health A*, 71, 1056-1068.  
6 <http://dx.doi.org/10.1080/15287390802111952>
- 7 DeAngelo, A. B.; George, M. H.; House, D. E. (1999). Hepatocarcinogenicity in the male  
8 B6C3F1 mouse following a lifetime exposure to dichloroacetic acid in the drinking  
9 water: Dose-response determination and modes of action. *J Toxicol Environ Health A*,  
10 58, 485-507. <http://www.ncbi.nlm.nih.gov/pubmed/10632141>
- 11 Deerberg, F. and Müller-Peddinghaus, R. (1970). [Comparative studies on the occurrence of  
12 spontaneous diseases and spontaneous deaths in conventional and specific germ-free  
13 female NMRI-HAN mice]. *Z Versuchstierkd*, 12, 341-347.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/5508977>
- 15 Deerberg, F.; Pittermann, W.; Rapp, K. (1974). [Lung tumor in the mouse: a progressive tumor].  
16 *Vet Pathol*, 11, 430-441. <http://www.ncbi.nlm.nih.gov/pubmed/4462259>
- 17 Dees, C. and Travis, C. (1993). The mitogenic potential of trichloroethylene in B6C3F1 mice.  
18 *Toxicol Lett*, 69, 129-137. <http://www.ncbi.nlm.nih.gov/pubmed/8212055>
- 19 Dees, C. and Travis, C. (1994). Trichloroacetate stimulation of liver DNA synthesis in male and  
20 female mice. *Toxicol Lett*, 70, 343-355. [http://dx.doi.org/10.1016/0378-4274\(94\)90129-5](http://dx.doi.org/10.1016/0378-4274(94)90129-5)
- 21 Degrassi, F. and Tanzarella, C. (1988). Immunofluorescent staining of kinetochores in  
22 micronuclei: A new assay for the detection of aneuploidy. *Mutat Res Genet Toxicol*  
23 *Environ Mutagen*, 203, 339-345. [http://dx.doi.org/10.1016/0165-1161\(88\)90030-1](http://dx.doi.org/10.1016/0165-1161(88)90030-1)
- 24 Dehon, B.; Humbert, L.; Devisme, L.; Stievenart, M.; Mathieu, D.; Houdret, N.; Lhermitte, M.  
25 (2000). Tetrachloroethylene and trichloroethylene fatality: case report and simple  
26 headspace SPME-capillary gas chromatographic determination in tissues. *J Anal Toxicol*,  
27 24, 22-26. <http://www.ncbi.nlm.nih.gov/pubmed/10654565>
- 28 Dekant, W. (1990). Bioactivation of halogenated alkenes by glutathione conjugation a  
29 mechanistic explanation for their renal carcinogenicity. *J Cancer Res Clin Oncol*, 116  
30 (Suppl. Part 2), 1063.
- 31 Dekant, W.; Berthold, K.; Vamvakas, S.; Henschler, D.; Anders, M. W. (1988). Thioacylating  
32 intermediates as metabolites of S-(1,2-dichlorovinyl)-L-cysteine and S-(1,2,2-  
33 trichlorovinyl)-L-cysteine formed by cysteine conjugate beta-lyase. *Chem Res Toxicol*, 1,  
34 175-178. <http://dx.doi.org/10.1021/tx00003a008>
- 35 Dekant, W.; Metzler, M.; Henschler, D. (1984). Novel metabolites of trichloroethylene through  
36 dechlorination reactions in rats, mice and humans. *Biochem Pharmacol*, 33, 2021-2027.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/6743352>
- 38 Dekant, W.; Metzler, M.; Henschler, D. (1986). Identification of S-1,2-dichlorovinyl-N-acetyl-  
39 cysteine as a urinary metabolite of trichloroethylene: A possible explanation for its  
40 nephrocarcinogenicity in male rats. *Biochem Pharmacol*, 35, 2455-2458.  
41 [http://dx.doi.org/10.1016/0006-2952\(86\)90039-0](http://dx.doi.org/10.1016/0006-2952(86)90039-0)
- 42 Dekant, W.; Schulz, A.; Metzler, M.; Henschler, D. (1986). Absorption, elimination, and  
43 metabolism of trichloroethylene: a quantitative comparison between rats and mice.  
44 *Xenobiotica*, 16, 143-152.

DRAFT– DO NOT QUOTE OR CITE

- 1 Dekant, W.; Vamvakas, S.; Berthold, K.; Schmidt, S.; Wild, D.; Henschler, D. (1986). Bacterial  
2 beta-lyase mediated cleavage and mutagenicity of cysteine conjugates derived from the  
3 nephrocarcinogenic alkenes trichloroethylene, tetrachloroethylene and  
4 hexachlorobutadiene. *Chem Biol Interact*, 60, 31-45. [http://dx.doi.org/10.1016/0009-  
5 2797\(86\)90015-3](http://dx.doi.org/10.1016/0009-2797(86)90015-3)
- 6 Delahunt, B.; Bethwaite, P. B.; Nacey, J. N. (1995). Occupational risk for renal cell carcinoma.  
7 A case-control study based on the New Zealand Cancer Registry. *Br J Urol*, 75, 578-582.  
8 <http://dx.doi.org/10.1111/j.1464-410X.1995.tb07410.x>
- 9 DeMarini, D. M.; Perry, E.; Shelton, M. L. (1994). Dichloroacetic acid and related compounds:  
10 Induction of prophage in *E. coli* and mutagenicity and mutation spectra in *Salmonella*  
11 TA100. *Mutagenesis*, 9, 429-437. <http://dx.doi.org/10.1093/mutage/9.5.429>
- 12 DerSimonian, R. and Laird, N. (1986). Meta-analysis in clinical trials. *Contemp Clin Trials*, 7,  
13 177-188. <http://www.ncbi.nlm.nih.gov/pubmed/3802833>
- 14 Dewar, R.; Siemiatyckia, J.; Geacuterin, M. (1994). Loss of statistical power associated with the  
15 use of a job-exposure matrix in occupational case-control studies. *Appl Occup Environ*  
16 *Hyg*, 6, 508-515. <http://dx.doi.org/10.1080/1047322X.1991.10387920>
- 17 Ding, R.; Shen, T.; Zhu, Q. X. (2006). [Effects of TCE and PCE on cultured human keratinocyte  
18 lipid peroxidation and protective effect of vitamin E on it]. *Zhonghua Lao Dong Wei*  
19 *Sheng Zhi Ye Bing Za Zhi*, 24, 662-665. <http://www.ncbi.nlm.nih.gov/pubmed/17181943>
- 20 Diot, E.; Lesire, V.; Guilmot, J.; Metzger, M.; Pilore, R.; Rogier, S., . . . Lasfargues, G. (2002).  
21 Systemic sclerosis and occupational risk factors: a case-control study. *Occup Environ*  
22 *Med*, 59, 545-549. <http://www.ncbi.nlm.nih.gov/pubmed/12151611>
- 23 DiRenzo, A. B.; Gandolfi, A. J.; Sipes, I. G. (1982). Microsomal bioactivation and covalent  
24 binding of aliphatic halides to DNA. *Toxicol Lett*, 11, 243-252.
- 25 Dix, K. J.; Kedderis, G. L.; Borghoff, S. J. (1997). Vehicle-dependent oral absorption and target  
26 tissue dosimetry of chloroform in male rats and female mice. *Toxicol Lett*, 91, 197-209.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/9217240>
- 28 Djouadi, F.; Weinheimer, C. J.; Saffitz, J. E.; Pitchford, C.; Bastin, J.; Gonzalez, F. J.; Kelly, D.  
29 P. (1998). A gender-related defect in lipid metabolism and glucose homeostasis in  
30 peroxisome proliferator- activated receptor alpha- deficient mice. *J Clin Invest*, 102,  
31 1083-1091. <http://dx.doi.org/10.1172/JCI3949>
- 32 Dobrev, I. D.; Andersen, M. E.; Yang, R. S. (2001). Assessing interaction thresholds for  
33 trichloroethylene in combination with tetrachloroethylene and 1,1,1-trichloroethane using  
34 gas uptake studies and PBPK modeling. *Arch Toxicol*, 75, 134-144.  
35 <http://dx.doi.org/10.1007/s002040100216>
- 36 Dobrev, I. D.; Andersen, M. E.; Yang, R. S. (2002). In silico toxicology: Simulating interaction  
37 thresholds for human exposure to mixtures of trichloroethylene, tetrachloroethylene, and  
38 1,1,1-trichloroethane. *Environ Health Perspect*, 110, 1031-1039.  
39 <http://dx.doi.org/12361929>
- 40 Dohn, D. R. and Anders, M. W. (1982). Assay of cysteine conjugate beta-lyase activity with S-  
41 (2-benzothiazolyl)cysteine as the substrate. *Anal Biochem*, 120, 379-386.  
42 [http://dx.doi.org/10.1016/0003-2697\(82\)90361-X](http://dx.doi.org/10.1016/0003-2697(82)90361-X)
- 43 Dolinoy, D. C.; Weidman, J. R.; Jirtle, R. L. (2007). Epigenetic gene regulation: linking early  
44 developmental environment to adult disease. *Reprod Toxicol*, 23, 297-307.  
45 <http://dx.doi.org/10.1016/j.reprotox.2006.08.012>

DRAFT– DO NOT QUOTE OR CITE

- 1 Donoghue, A. M.; Dryson, E. W.; Wynn-Williams, G. (1995). Contrast sensitivity in organic-  
2 solvent-induced chronic toxic encephalopathy. *J Occup Environ Med*, 37, 1357-1363.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/8749741>
- 4 Dorfmueller, M. A.; Henne, S. P.; York, R. G.; Bornschein, R. L.; Manson, J. M. (1979).  
5 Evaluation of teratogenicity and behavioral toxicity with inhalation exposure of maternal  
6 rats to trichloroethylene. *Toxicology*, 14, 153-166.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/538767>
- 8 Dorne, J. L.; Walton, K.; Renwick, A. G. (2005). Human variability in xenobiotic metabolism  
9 and pathway-related uncertainty factors for chemical risk assessment: a review. *Food*  
10 *Chem Toxicol*, 43, 203-216. <http://dx.doi.org/10.1016/j.fct.2004.05.011>
- 11 Dosemeci, M.; Cocco, P.; Chow, W. H. (1999). Gender differences in risk of renal cell  
12 carcinoma and occupational exposures to chlorinated aliphatic hydrocarbons. *Am J Ind*  
13 *Med*, 36, 54-59. <http://www.ncbi.nlm.nih.gov/pubmed/10361587>
- 14 Dosemeci, M.; Cocco, P.; Gómez, M.; Stewart, P. A.; Heineman, E. F. (1994). Effects of three  
15 features of a job-exposure matrix on risk estimates. *Epidemiology*, 5, 124-127.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/8117771>
- 17 Dosemeci, M.; Wacholder, S.; Lubin, J. H. (1990). Does nondifferential misclassification of  
18 exposure always bias a true effect toward the null value? *Am J Epidemiol*, 132, 746-748.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/2403115>
- 20 Døssing, M.; Petersen, K. T.; Vyberg, M.; Olsen, J. H. (1997). Liver cancer among employees in  
21 Denmark. *Am J Ind Med*, 32, 248-254. <http://www.ncbi.nlm.nih.gov/pubmed/9219654>
- 22 Douglas, G.; Gingerich, J.; Soper, L.; Potvin, M.; Bjarnason, S. (1999). Evidence for the lack of  
23 base-change and small-deletion mutation induction by trichloroethylene in lacZ  
24 transgenic mice. *Environ Mol Mutagen*, 34, 190-194.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/10529743>
- 26 Dow, J. and Green, T. (2000). Trichloroethylene induced vitamin B(12) and folate deficiency  
27 leads to increased formic acid excretion in the rat. *Toxicology*, 146, 123-136.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/10814845>
- 29 Dowsley, T.; Ulreich, J.; Bolton, J.; Park, S.; Forkert, P. (1996). CYP2E1-dependent  
30 bioactivation of 1,1-dichloroethylene in murine lung: formation of reactive intermediates  
31 and glutathione conjugates. *Toxicol Appl Pharmacol*, 139, 42-48.  
32 <http://dx.doi.org/10.1006/taap.1996.0141>
- 33 Drake, V. J.; Koprowski, S. L.; Hu, N.; Smith, S. M.; Lough, J. (2006). Cardiogenic effects of  
34 trichloroethylene and trichloroacetic acid following exposure during heart specification  
35 of avian development. *Toxicol Sci*, 94, 153-162. <http://dx.doi.org/10.1093/toxsci/kfl083>
- 36 Drake, V.; Koprowski, S.; Lough, J.; Hu, N.; Smith, S. (2006). Trichloroethylene exposure  
37 during cardiac valvuloseptal morphogenesis alters cushion formation and cardiac  
38 hemodynamics in the avian embryo. *Environ Health Perspect*, 114, 842-847.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/16759982>
- 40 Driscoll, T.; Hamdan, H.; Wang, G.; Wright, P.; Stacey, N. (1992). Concentrations of individual  
41 serum or plasma bile acids in workers exposed to chlorinated aliphatic hydrocarbons. *Br J*  
42 *Ind Med*, 49, 700-705. <http://www.ncbi.nlm.nih.gov/pubmed/1419858>
- 43 Dryver, E.; Brandt, L.; Kauppinen, T.; Olsson, H. (2004). Occupational exposures and non-  
44 Hodgkin's lymphoma in Southern Sweden. *Int J Occup Environ Health*, 10, 13-21.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/15070021>

DRAFT– DO NOT QUOTE OR CITE

- 1 D'Souza, R.; Bruckner, J.; Feldman, S. (1985). Oral and intravenous trichloroethylene  
2 pharmacokinetics in the rat. *J Toxicol Environ Health*, 15, 587-601.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/4046066>
- 4 DT Jr, L. and Calo, J. M. (1981). Distribution coefficients of chlorinated hydrocarbons in dilute  
5 air-water systems for groundwater contamination applications. *Journal of Chemical and*  
6 *Engineering Data*, 26, 382-585.
- 7 Du, C. L. and Wang, J. D. (1998). Increased morbidity odds ratio of primary liver cancer and  
8 cirrhosis of the liver among vinyl chloride monomer workers. *Occup Environ Med*, 55,  
9 528-532. <http://www.ncbi.nlm.nih.gov/pubmed/9849539>
- 10 Duffel, M. W. and Jakoby, W. B. (1982). Cysteine S-conjugate N-acetyltransferase from rat  
11 kidney microsomes. *Mol Pharmacol*, 21, 444-448.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/6892478>
- 13 Dugard, P. (2000). Effects of trichloroethylene (TCE) on an in vitro chick atrioventricular canal  
14 culture. *Toxicol Sci*, 56, 437-438. <http://www.ncbi.nlm.nih.gov/pubmed/10960254>
- 15 Dumas, S.; Parent, M. E.; Siemiatycki, J.; Brisson, J. (2000). Rectal cancer and occupational risk  
16 factors: a hypothesis-generating, exposure-based case-control study. *Int J Cancer*, 87,  
17 874-879.
- 18 DuTeaux, S. B.; Berger, T.; Hess, R. A.; Sartini, B. L.; Miller, M. G. (2004). Male reproductive  
19 toxicity of trichloroethylene: sperm protein oxidation and decreased fertilizing ability.  
20 *Biol Reprod*, 70, 1518-1526. <http://dx.doi.org/10.1095/biolreprod.103.022210>
- 21 DuTeaux, S.; Hengel, M.; DeGroot, D.; Jelks, K.; Miller, M. (2003). Evidence for  
22 trichloroethylene bioactivation and adduct formation in the rat epididymis and efferent  
23 ducts. *Biol Reprod*, 69, 771-779. <http://dx.doi.org/10.1095/biolreprod.102.014845>
- 24 DuTeaux, S.; Newman, J.; Morisseau, C.; Fairbairn, E.; Jelks, K.; Hammock, B.; Miller, M.  
25 (2004). Epoxide hydrolases in the rat epididymis: possible roles in xenobiotic and  
26 endogenous fatty acid metabolism. *Toxicol Sci*, 78, 187-195.  
27 <http://dx.doi.org/10.1093/toxsci/kfh066>
- 28 Eacho, P. I.; Lanier, T. L.; Brodhecker, C. A. (1991). Hepatocellular DNA synthesis in rats given  
29 peroxisome proliferating agents: Comparison of WY-14,643 to clofibrac acid, nafenopin  
30 and LY171883. *Carcinogenesis*, 12, 1557-1561.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/1893515>
- 32 Eastin, W. C.; Mennear, J. H.; Tennant, R. W.; Stoll, R. E.; Branstetter, D. G.; Bucher, J. R., . . .  
33 Mahler, J. F. (2001). Tg.AC genetically altered mouse: assay working group overview of  
34 available data. *Toxicol Pathol*, 29 Suppl, 60-80.  
35 <http://dx.doi.org/10.1080/019262301753178483>
- 36 Eastmond, D. A. and Tucker, J. D. (1989). Identification of aneuploidy-inducing agents using  
37 cytokinesis-blocked human lymphocytes and an antikinetochores antibody. *Environ Mol*  
38 *Mutagen*, 13, 34-43. <http://www.ncbi.nlm.nih.gov/pubmed/2783409>
- 39 Eastmond, D. A.; Hartwig, A.; Anderson, D.; Anwar, W. A.; Cimino, M. C.; Dobrev, I., . . .  
40 Vickers, C. (2009). Mutagenicity testing for chemical risk assessment: Update of the  
41 WHO/IPCS Harmonized Scheme. *Mutagenesis*, 24, 341-349.  
42 <http://dx.doi.org/10.1093/mutage/geb014>
- 43 EFSA. (European Food Safety Authority). (2006). Opinion of the scientific panel on food  
44 additives, flavourings, processing aids and materials in contact with food (AFC) on a



DRAFT– DO NOT QUOTE OR CITE

- 1 request from the commission related to a new long-term carcinogenicity study on  
2 aspartame. *EFSA J*, 4, 1-44. <http://dx.doi.org/10.2903/j.efsa.2006.356>
- 3 Egger, M.; Smith, G. D.; Schneider, M.; Minder, C. (1997). Bias in meta-analysis detected by a  
4 simple, graphical test. *Br Med J*, 315, 629-634.
- 5 EHS. (Environmental Health Strategies). (1997). Final Report. Correspondence from Paul A.  
6 Cammer, Ph.D., Trichloroethylene Issues Group, to Cheryl Siegel Scott, US  
7 Environmental Protection Agency. December 22, 1997. Menlo Park, CA: Author.
- 8 Eichenlaub-Ritter, U.; Baart, E.; Yin, H.; Betzendahl, I. (1996). Mechanisms of spontaneous and  
9 chemically-induced aneuploidy in mammalian oogenesis: basis of sex-specific  
10 differences in response to aneugens and the necessity for further tests. *Mutat Res*, 372,  
11 279-294. <http://www.ncbi.nlm.nih.gov/pubmed/9015146>
- 12 El Ghawabi, S. M.; Mansoor, M. B.; El Gamel, M. S.; El Saharti, A. A.; El Enany, F. F. (1973).  
13 Chronic trichloroethylene exposure. 56, 715-724.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/4618847>
- 15 Elcombe, C. R. (1985). Species differences in carcinogenicity and peroxisome proliferation due  
16 to trichloroethylene: a biochemical human hazard assessment. *J Ind Hyg Toxicol*, 8, 6-17.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/3868383>
- 18 Elcombe, C. R.; Rose, M. S.; Pratt, I. S. (1985). Biochemical, histological, and ultrastructural  
19 changes in rat and mouse liver following the administration of trichloroethylene: Possible  
20 relevance to species differences in hepatocarcinogenicity. *Toxicol Appl Pharmacol*, 79,  
21 365-376. [http://dx.doi.org/10.1016/0041-008X\(85\)90135-8](http://dx.doi.org/10.1016/0041-008X(85)90135-8)
- 22 Elfarra, A. A. and Anders, M. W. (1984). Renal processing of glutathione conjugates role in  
23 nephrotoxicity. *Biochem Pharmacol*, 33, 3729-3732.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/6508831>
- 25 Elfarra, A. A.; Jakobson, I.; Anders, M. W. (1986). Mechanism of S-(1,2-  
26 dichlorovinyl)glutathione-induced nephrotoxicity. *Biochem Pharmacol*, 35, 283-288.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/2867768>
- 28 Elfarra, A. and Hwang, I. (1990). In vivo metabolites of S-(2-benzothiazolyl)-L-cysteine as  
29 markers of in vivo cysteine conjugate beta-lyase and thiol glucuronosyl transferase  
30 activities. *Drug Metab Dispos*, 18, 917-922.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/1981537>
- 32 Elfarra, A.; Krause, R.; Last, A.; Lash, L.; Parker, J. (1998). Species- and sex-related differences  
33 in metabolism of trichloroethylene to yield chloral and trichloroethanol in mouse, rat, and  
34 human liver microsomes. *Drug Metab Dispos*, 26, 779-785.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/9698293>
- 36 Elfarra, A.; Lash, L.; Anders, M. (1987). Alpha-ketoacids stimulate rat renal cysteine conjugate  
37 beta-lyase activity and potentiate the cytotoxicity of S-(1,2-dichlorovinyl)-L-cysteine.  
38 *Mol Pharmacol*, 31, 208-212. <http://www.ncbi.nlm.nih.gov/pubmed/3807895>
- 39 Elizur, A.; Adair-Kirk, T. L.; Kelley, D. G.; Griffin, G. L.; Demello, D. E.; Senior, R. M. (2008).  
40 Tumor necrosis factor-alpha from macrophages enhances LPS-induced clara cell  
41 expression of keratinocyte-derived chemokine. *Am J Respir Cell Mol Biol*, 38, 8-15.  
42 <http://dx.doi.org/10.1165/rcmb.2007-0203OC>
- 43 Elofsson, S. A.; Gamberale, F.; Hindmarsh, T. (1980). Exposure to organic solvents. A cross-  
44 sectional epidemiologic investigation on occupationally exposed car and industrial spray

- 1 painters with special reference to the nervous system. *Scand J Work Environ Health*, 6,  
2 239 - 273.
- 3 Elovaara, E.; Hemminki, K.; Vainio, H. (1979). Effects of methylene chloride, trichloroethane,  
4 trichloroethylene, tetrachloroethylene and toluene on the development of chick embryos.  
5 *Toxicology*, 12, 111-119. [http://dx.doi.org/10.1016/0300-483X\(79\)90037-4](http://dx.doi.org/10.1016/0300-483X(79)90037-4)
- 6 El-Serag, H. B. (2004). Hepatocellular carcinoma: recent trends in the United States.  
7 *Gastroenterology*, 127, S27-S34. <http://www.ncbi.nlm.nih.gov/pubmed/15508094>
- 8 El-Serag, H. B. (2007). Epidemiology of hepatocellular carcinoma in USA. *Hepatology*  
9 *Research*, 37 Suppl 2, S88-S94. <http://dx.doi.org/10.1111/j.1872-034X.2007.00168.x>
- 10 El-Serag, H. B. and Rudolph, K. L. (2007). Hepatocellular carcinoma: epidemiology and  
11 molecular carcinogenesis. *Gastroenterology*, 132, 2557-2576.  
12 <http://dx.doi.org/10.1053/j.gastro.2007.04.061>
- 13 Emmert, B.; Bünger, J.; Keuch, K.; Müller, M.; Emmert, S.; Hallier, E.; Westphal, G. A. (2006).  
14 Mutagenicity of cytochrome P450 2E1 substrates in the Ames test with the metabolic  
15 competent *S. typhimurium* strain YG7108pin3ERb5. *Toxicology*, 228, 66-76.  
16 <http://dx.doi.org/10.1016/j.tox.2006.08.013>
- 17 Engel, L. S.; Vaughan, T. L.; Gammon, M. D.; Chow, W. H.; Risch, H. A.; Dubrow, R., . . .  
18 Fraumeni, J. F. (2002). Occupation and risk of esophageal and gastric cardia  
19 adenocarcinoma. *Am J Ind Med*, 42, 11-22. <http://dx.doi.org/10.1002/ajim.10077>
- 20 Epstein, D. L.; Nolen, G. A.; Randall, J. L.; Christ, S. A.; Read, E. J.; Stober, J. A.; Smith, M. K.  
21 (1992). Cardiopathic effects of dichloroacetate in the fetal Long-Evans rat. *Teratology*,  
22 46, 225-235. <http://dx.doi.org/10.1002/tera.1420460306>
- 23 Ertle, T.; Henschler, D.; Mueller, G.; Spassowski, M. (1972). Metabolism of trichloroethylene in  
24 man I The significance of trichloroethanol in long-term exposure conditions. *Arch*  
25 *Toxicol*, 29, 171-188. <http://www.ncbi.nlm.nih.gov/pubmed/4642755>
- 26 Evans, M. V.; Chiu, W. A.; Okino, M. S.; Caldwell, J. C. (2009). Development of an updated  
27 PBPK model for trichloroethylene and metabolites in mice, and its application to discern  
28 the role of oxidative metabolism in TCE-induced hepatomegaly. *Toxicol Appl*  
29 *Pharmacol*, 236, 329-340. <http://dx.doi.org/10.1016/j.taap.2009.02.013>
- 30 Evert, M.; Schneider-Stock, R.; Dombrowski, F. (2003). Apolipoprotein A-IV mRNA  
31 overexpression in early preneoplastic hepatic foci induced by low-number pancreatic islet  
32 transplants in streptozotocin-diabetic rats. *Pathol Res Pract*, 199, 373-379.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/12924437>
- 34 Eyre, R.; Stevens, D.; Parker, J.; Bull, R. (1995). Acid-labile adducts to protein can be used as  
35 indicators of the cysteine S-conjugate pathway of trichloroethene metabolism. *J Toxicol*  
36 *Environ Health*, 46, 443-464. <http://www.ncbi.nlm.nih.gov/pubmed/8523471>
- 37 Eyre, R.; Stevens, D.; Parker, J.; Bull, R. (1995). Renal activation of trichloroethene and S-(1,2-  
38 dichlorovinyl)-L-cysteine and cell proliferative responses in the kidneys of F344 rats and  
39 B6C3F1 mice. *J Toxicol Environ Health*, 46, 465-481.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/8523472>
- 41 Fagliano, J.; Berry, M.; Bove, F.; Burke, T. (1990). Drinking water contamination and the  
42 incidence of leukemia: An ecologic study. *Am J Public Health*, 80, 1209-1212.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/2400032>
- 44 Fahrig, R.; Madle, S.; Baumann, H. (1995). Genetic toxicology of trichloroethylene (TCE).  
45 *Mutat Res*, 340, 1-36. <http://www.ncbi.nlm.nih.gov/pubmed/8524349>

DRAFT– DO NOT QUOTE OR CITE

- 1 Fairman, M. J.; Scott, B. B.; Toothill, C.; Losowsky, M. S. (1977). Jejunal mucosal gamma  
2 glutamyl transferase activity in coeliac disease. *Gut*, 18, 484-487.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/17564>
- 4 Falk Filipsson, A. and Victorin, K. (2003). Comparison of available benchmark dose softwares  
5 and models using trichloroethylene as a model substance. *Regul Toxicol Pharmacol*, 37,  
6 343-355. <http://www.ncbi.nlm.nih.gov/pubmed/12758215>
- 7 Fan, A. (1988). Trichloroethylene: water contamination and health risk assessment. *Rev Environ*  
8 *Contam Toxicol*, 101, 55-92. <http://www.ncbi.nlm.nih.gov/pubmed/3275994>
- 9 Fang, H. L.; Strom, S. C.; Cai, H.; Falany, C. N.; Kocarek, T. A.; Runge-Morris, M. (2005).  
10 Regulation of human hepatic hydroxysteroid sulfotransferase gene expression by the  
11 peroxisome proliferator-activated receptor alpha transcription factor. *Mol Pharmacol*, 67,  
12 1257-1267. <http://dx.doi.org/10.1124/mol.104.005389>
- 13 Farazi, P. A. and Depinho, R. A. (2006). Hepatocellular carcinoma pathogenesis: from genes to  
14 environment. *Nat Rev Cancer*, 6, 674-687. <http://dx.doi.org/10.1038/nrc1934>
- 15 Fattovich, G.; Stroffolini, T.; Zagni, I.; Donato, F. (2004). Hepatocellular carcinoma in cirrhosis:  
16 incidence and risk factors. *Gastroenterology*, 127, S35-S50.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/15508101>
- 18 Fechter, L. D.; Gearhart, C.; Fulton, S.; Campbell, J.; Fisher, J.; Na, K., . . . Pouyatos, B. (2007).  
19 Promotion of noise-induced cochlear injury by toluene and ethylbenzene in the rat.  
20 *Toxicol Sci*, 98, 542-551.
- 21 Fechter, L. D.; Liu, Y.; Herr, D. W.; Crofton, K. M. (1998). Trichloroethylene ototoxicity:  
22 Evidence for a cochlear origin. *Toxicol Sci*, 42, 28-35.  
23 <http://dx.doi.org/10.1093/toxsci/42.1.28>
- 24 Feinberg, A. P.; Ohlsson, R.; Henikoff, S. (2006). The epigenetic progenitor origin of human  
25 cancer. *Nat Rev Genet*, 7, 21-33. <http://dx.doi.org/10.1038/nrg1748>
- 26 Feitelson, M. A.; Sun, B.; Satiroglu Tufan, N. L.; Liu, J.; Pan, J.; Lian, Z. (2002). Genetic  
27 mechanisms of hepatocarcinogenesis. *Oncogene*, 21, 2593-2604.  
28 <http://dx.doi.org/10.1038/sj.onc.1205434>
- 29 Feldman, R.; Chirico-Post, J.; Proctor, S. (1988). Blink reflex latency after exposure to  
30 trichloroethylene in well water. *Arch Environ Health*, 43, 143-148.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/3377549>
- 32 Feldman, R.; Niles, C.; Proctor, S.; Jabre, J. (1992). Blink reflex measurement of effects of  
33 trichloroethylene exposure on the trigeminal nerve. *Muscle Nerve*, 15, 490-495.  
34 <http://dx.doi.org/10.1002/mus.880150413>
- 35 Ferlay, J.; Bray, F.; Pisani, P.; Parkin, D. M. (2004). GLOBOCAN 2002: Cancer incidence,  
36 mortality and prevalence worldwide.
- 37 Fernandez, J. G.; Droz, P. O.; Humbert, B. E.; Caperos, J. R. (1977). Trichloroethylene  
38 exposure: simulation of uptake, excretion, and metabolism using a mathematical model.  
39 *Occup Environ Med*, 34, 43-55. <http://www.ncbi.nlm.nih.gov/pubmed/843463>
- 40 Ferreira-Gonzalez, A.; DeAngelo, A. B.; Nasim, S.; Garrett, C. T. (1995). Ras oncogene  
41 activation during hepatocarcinogenesis in B6C3F1 male mice by dichloroacetic and  
42 trichloroacetic acids. *Carcinogenesis*, 16, 495-500.  
43 <http://dx.doi.org/10.1093/carcin/16.3.495>
- 44 Fevotte, J.; Charbotel, B.; Muller-Beauté, P.; Martin, J. L.; Hours, M.; Bergeret, A. (2006). Case-  
45 control study on renal cell cancer and occupational exposure to trichloroethylene. Part I:

- 1 Exposure assessment. *Ann Occup Hyg*, 50, 765-775.  
2 <http://dx.doi.org/10.1093/annhyg/mel040>
- 3 Figgs, L.; Dosemeci, M.; Blair, A. (1995). United States non-Hodgkin's lymphoma surveillance  
4 by occupation 1984-1989: a twenty-four state death certificate study. *Am J Ind Med*, 27,  
5 817-835. <http://www.ncbi.nlm.nih.gov/pubmed/7645576>
- 6 Filser, J. G. and Bolt, H. M. (1979). Pharmacokinetics of halogenated ethylenes in rats. *Arch*  
7 *Toxicol*, 42, 123-136. <http://dx.doi.org/10.1007/BF00316492>
- 8 Finckh, A.; Cooper, G. S.; Chibnik, L. B.; Costenbader, K. H.; Watts, J.; Pankey, H., . . .  
9 Karlson, E. W. (2006). Occupational silica and solvent exposures and risk of systemic  
10 lupus erythematosus in urban women. *Arthritis Rheum*, 54, 3648-3654.  
11 <http://dx.doi.org/10.1002/art.22210>
- 12 Fiserova-Bergerova, V.; Tichy, M.; Di Carlo, F. J. (1984). Effects of biosolubility on pulmonary  
13 uptake and disposition of gases and vapors of lipophilic chemicals. *Drug Metab Rev*, 15,  
14 1033-1070.
- 15 Fisher, J. (2000). Physiologically based pharmacokinetic models for trichloroethylene and its  
16 oxidative metabolites. *Environ Health Perspect*, 108, 265-273.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/10807557>
- 18 Fisher, J. W. and Allen, B. C. (1993). Evaluating the risk of liver cancer in humans exposed to  
19 trichloroethylene using physiological models. *Risk Anal*, 13, 87-95.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/8451463>
- 21 Fisher, J. W.; Gargas, M. L.; Allen, B. C.; Andersen, M. E. (1991). Physiologically based  
22 pharmacokinetic modeling with trichloroethylene and its metabolite, trichloroacetic acid,  
23 in the rat and mouse. *Toxicol Appl Pharmacol*, 109, 183-195.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/2068722>
- 25 Fisher, J. W.; Mahle, D.; Abbas, R. (1998). A human physiologically based pharmacokinetic  
26 model for trichloroethylene and its metabolites, trichloroacetic acid and free  
27 trichloroethanol. *Toxicol Appl Pharmacol*, 152, 339-359.  
28 <http://dx.doi.org/10.1006/taap.1998.8486>
- 29 Fisher, J. W.; Whittaker, T. A.; Taylor, D. H.; 3rd, C. H.; Andersen, M. E. (1989).  
30 Physiologically based pharmacokinetic modeling of the pregnant rat: a multiroute  
31 exposure model for trichloroethylene and its metabolite, trichloroacetic acid. *Toxicol*  
32 *Appl Pharmacol*, 99, 395-414. [http://dx.doi.org/10.1016/0041-008X\(89\)90149-X](http://dx.doi.org/10.1016/0041-008X(89)90149-X)
- 33 Fisher, J. W.; Whittaker, T. A.; Taylor, D. H.; Clewell, H. J., III; Andersen, M. E. (1990).  
34 Physiologically based pharmacokinetic modeling of the lactating rat and nursing pup: a  
35 multiroute exposure model for trichloroethylene and its metabolite, trichloroacetic acid.  
36 *Toxicol Appl Pharmacol*, 102, 497-513. <http://www.ncbi.nlm.nih.gov/pubmed/2315918>
- 37 Fisher, J.; Channel, S.; Eggers, J.; Johnson, P.; MacMahon, K.; Goodyear, C., . . . Graeter, L.  
38 (2001). Trichloroethylene, trichloroacetic acid, and dichloroacetic acid: Do they affect  
39 fetal rat heart development. *Int J Toxicol*, 20, 257-267.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/11766123>
- 41 Fisher, J.; Mahle, D.; Bankston, L.; Greene, R.; Gearhart, J. (1997). Lactational transfer of  
42 volatile chemicals in breast milk. *Am Ind Hyg Assoc J*, 58, 425-431.  
43 <http://dx.doi.org/10.1080/15428119791012667>
- 44 Fitzgerald, J. E.; Sanyer, J. L.; Schardein, J. L.; Lake, R. S.; McGuire, E. J.; de la Iglesia, F. A.  
45 (1981). Carcinogen bioassay and mutagenicity studies with the hypolipidemic agent

DRAFT– DO NOT QUOTE OR CITE

- 1 gemfibrozil. *J Natl Cancer Inst*, 67, 1105-1116.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/7029098>
- 3 Fleming-Jones, M. E. and Smith, R. E. (2003). Volatile organic compounds in foods: A five year  
4 study. *J Agric Food Chem*, 51, 8120-8127.
- 5 Flodin, U.; Fredriksson, M.; Persson, B. (1987). Multiple myeloma and engine exhausts, fresh  
6 wood, and creosote: a case-referent study. *Am J Ind Med*, 12, 519-529.
- 7 Flood, T. (1988). Report on mortality in Maricopa County.
- 8 Flood, T. J. (1997). Case-referent study of childhood leukemia in Maricopa County, Arizona:  
9 1965-1990. Phoenix, AZ: Arizona Dept. of Health Services, Bureau of Epidemiology  
10 and Disease Control Services, Office of Chronic Disease Epidemiology, Office of  
11 Environmental Health.
- 12 Forand, S. P. (2004). Leukaemia incidence among workers in the shoe and boot manufacturing  
13 industry: a case-control study. *Environ Health*, 3, 7. [http://dx.doi.org/10.1186/1476-](http://dx.doi.org/10.1186/1476-069X-3-7)  
14 [069X-3-7](http://dx.doi.org/10.1186/1476-069X-3-7)
- 15 Ford, E.; Rhodes, S.; McDiarmid, M.; Schwartz, S.; Brown, J. (1995). Deaths from acute  
16 exposure to trichloroethylene. *J Occup Environ Med*, 37, 749-754.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/7670923>
- 18 Forkert, P. (1999). 1,1-Dichloroethylene-induced Clara cell damage is associated with in situ  
19 formation of the reactive epoxide. Immunohistochemical detection of its glutathione  
20 conjugate. *Am J Respir Cell Mol Biol*, 20, 1310-1318.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/10340951>
- 22 Forkert, P. (1999). In vivo formation and localization of 1,1-dichloroethylene epoxide in murine  
23 liver: identification of its glutathione conjugate 2-S-glutathionyl acetate. *J Pharmacol Exp*  
24 *Ther*, 290, 1299-1306. <http://www.ncbi.nlm.nih.gov/pubmed/10454507>
- 25 Forkert, P. G. and Birch, D. W. (1989). Pulmonary toxicity of trichloroethylene in mice: covalent  
26 binding and morphological manifestations. *Drug Metab Dispos*, 17, 106-113.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/2566461>
- 28 Forkert, P. G. and Forkert, L. (1994). Trichloroethylene induces pulmonary fibrosis in mice. *Can*  
29 *J Physiol Pharmacol*, 72, 205-210. <http://www.ncbi.nlm.nih.gov/pubmed/8069766>
- 30 Forkert, P. G.; Sylvestre, P. L.; Poland, J. S. (1985). Lung injury induced by trichloroethylene.  
31 *Toxicology*, 35, 143-160. <http://www.ncbi.nlm.nih.gov/pubmed/4002240>
- 32 Forkert, P.; Baldwin, R.; Millen, B.; Lash, L.; Putt, D.; Shultz, M.; Collins, K. (2005).  
33 Pulmonary bioactivation of trichloroethylene to chloral hydrate: relative contributions of  
34 CYP2E1, CYP2F, and CYP2B1. *Drug Metab Dispos*, 33, 1429-1437.  
35 <http://dx.doi.org/10.1124/dmd.105.005074>
- 36 Forkert, P.; Lash, L.; Nadeau, V.; Tardif, R.; Simmonds, A. (2002). Metabolism and toxicity of  
37 trichloroethylene in epididymis and testis. *Toxicol Appl Pharmacol*, 182, 244-254.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/12183104>
- 39 Forkert, P.; Lash, L.; Tardif, R.; Tanphaichitr, N.; Vandervoort, C.; Moussa, M. (2003).  
40 Identification of trichloroethylene and its metabolites in human seminal fluid of workers  
41 exposed to trichloroethylene. *Drug Metab Dispos*, 31, 306-311.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/12584157>
- 43 Forkert, P.; Malkinson, A.; Rice, P.; Moussa, M. (1999). Diminished CYP2E1 expression and  
44 formation of 2-S-glutathionyl acetate, a glutathione conjugate derived from 1,1-

- 1 dichloroethylene epoxide, in murine lung tumors. *Drug Metab Dispos*, 27, 68-73.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/9884311>
- 3 Forkert, P.; Millen, B.; Lash, L.; Putt, D.; Ghanayem, B. (2006). Pulmonary bronchiolar  
4 cytotoxicity and formation of dichloroacetyl lysine protein adducts in mice treated with  
5 trichloroethylene. *J Pharmacol Exp Ther*, 316, 520-529.  
6 <http://dx.doi.org/10.1124/jpet.105.093062>
- 7 Fort, D. J.; Stover, E. L.; Rayburn, J. R.; Hull, M.; Bantle, J. A. (1993). Evaluation of the  
8 developmental toxicity of trichloroethylene and detoxification metabolites using  
9 *Xenopus*. *Birth Defects Res B Dev Reprod Toxicol*, 13, 35-45.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/8100651>
- 11 Fort, D.; Rayburn, J.; DeYoung, D.; Bantle, J. (1991). Assessing the efficacy of an Aroclor 1254-  
12 induced exogenous metabolic activation system for FETAX. *Drug Chem Toxicol*, 14,  
13 143-160. <http://dx.doi.org/10.3109/01480549109017873>
- 14 Fox, A.; Sullivan, B.; Buffini, J.; Neichin, M.; Nicora, R.; Hoehler, F., . . . Stoltz, R. (1996).  
15 Reduction of serum lactate by sodium dichloroacetate, and human pharmacokinetic-  
16 pharmacodynamic relationships. *J Pharmacol Exp Ther*, 279, 686-693.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/8930172>
- 18 Fox, A.; Yang, X.; Murli, H.; Lawlor, T.; Cifone, M.; Reno, F. (1996). Absence of mutagenic  
19 effects of sodium dichloroacetate. *Fundam Appl Toxicol*, 32, 87-95.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/8812237>
- 21 Fox, T. R.; Schumann, A. M.; Watanabe, P. G.; Yano, B. L.; Maher, V. M.; McCormick, J. J.  
22 (1990). Mutational analysis of the H-ras oncogene in spontaneous C57BL/6 x C3H/He  
23 mouse liver tumors and tumors induced with genotoxic and nongenotoxic  
24 hepatocarcinogens. *Cancer Res*, 50, 4014-4019.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/2191770>
- 26 Fraga, M. F.; Ballestar, E.; Villar-Garea, A.; Boix-Chornet, M.; Espada, J.; Schotta, G., . . .  
27 Esteller, M. (2005). Loss of acetylation at Lys16 and trimethylation at Lys20 of histone  
28 H4 is a common hallmark of human cancer. *Nat Genet*, 37, 391-400.  
29 <http://dx.doi.org/10.1038/ng1531>
- 30 Fraga, M. F.; Herranz, M.; Espada, J.; Ballestar, E.; Paz, M. F.; Roper, S., . . . Esteller, M.  
31 (2004). A mouse skin multistage carcinogenesis model reflects the aberrant DNA  
32 methylation patterns of human tumors. *Cancer Res*, 64, 5527-5534.  
33 <http://dx.doi.org/10.1158/0008-5472.CAN-03-4061>
- 34 Franco, A.; Costoya, M. A.; Roca, E. (2007). Estimating risk during showering exposure to  
35 VOCs of workers in a metal-degreasing facility. *J Toxicol Environ Health A*, 70, 627-  
36 637. <http://dx.doi.org/10.1080/15287390600974551>
- 37 Franco, J. L.; Teixeira, A.; Meotti, F. C.; Ribas, C. M.; Stringari, J.; Garcia Pomblum, S. C., . . .  
38 Farina, M. (2006). Cerebellar thiol status and motor deficit after lactational exposure to  
39 methylmercury. *Environ Res*, 102, 22-28. <http://dx.doi.org/10.1016/j.envres.2006.02.003>
- 40 Fredriksson, A.; Danielsson, B. R. G.; Eriksson, P. (1993). Altered behaviour in adult mice  
41 orally exposed to tri- and tetrachloroethylene as neonates. *Toxicol Lett*, 66, 13-19.  
42 [http://dx.doi.org/10.1016/0378-4274\(93\)90074-8](http://dx.doi.org/10.1016/0378-4274(93)90074-8)
- 43 Fredriksson, M.; Bengtsson, N. O.; Hardell, L.; Axelson, O. (1989). Colon cancer, physical  
44 activity, and occupational exposures. A case-control study. *Cancer*, 63, 1838-1842.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/2702592>

DRAFT– DO NOT QUOTE OR CITE

- 1 Fritschi, L. and Siemiatycki, J. (1996). Lymphoma, myeloma and occupation: results of a case-  
2 control study. *Int J Cancer*, 67, 498-503. [http://dx.doi.org/10.1002/\(SICI\)1097-](http://dx.doi.org/10.1002/(SICI)1097-)  
3 [0215\(19960807\)67:4<498::AID-IJC6>3.0.CO;2-N](http://dx.doi.org/10.1002/(SICI)1097-0215(19960807)67:4<498::AID-IJC6>3.0.CO;2-N)
- 4 Fritschi, L. and Siemiatycki, J. (1996). Melanoma and occupation: results of a case-control  
5 study. *Occup Environ Med*, 53, 168-173. <http://www.ncbi.nlm.nih.gov/pubmed/8704857>
- 6 Fritschi, L.; Nadon, L.; Benke, G.; Lakhani, R.; Latreille, B.; Parent, M. E.; Siemiatycki, J.  
7 (2003). Validation of expert assessment of occupational exposures. *Am J Ind Med*, 43,  
8 519-522. <http://dx.doi.org/10.1002/ajim.10208>
- 9 Fritschi, L.; Siemiatycki, J.; Richardson, L. (1996). Self-assessed versus expert-assessed  
10 occupational exposures. *Am J Epidemiol*, 144, 521-527.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/8781468>
- 12 Frøyland, L.; Madsen, L.; Vaagenes, H.; Totland, G. K.; Auwerx, J.; Kryvi, H., . . . Berge, R. K.  
13 (1997). Mitochondrion is the principal target for nutritional and pharmacological control  
14 of triglyceride metabolism. *J Lipid Res*, 38, 1851-1858.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/9323594>
- 16 Fujita, J.; Marino, M. W.; Wada, H.; Jungbluth, A. A.; Mackrell, P. J.; Rivadeneira, D. E., . . .  
17 Daly, J. M. (2001). Effect of TNF gene depletion on liver regeneration after partial  
18 hepatectomy in mice. *Surgery*, 129, 48-54.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/11150033>
- 20 Fukuda, K.; Takemoto, K.; Tsuruta, H. (1983). Inhalation carcinogenicity of trichloroethylene in  
21 mice and rats. *Ind Health*, 21, 243-254. <http://www.ncbi.nlm.nih.gov/pubmed/6654707>
- 22 Furge, K. A.; Dykema, K.; Petillo, D.; Westphal, M.; Zhang, Z. F.; Kort, E. J.; Teh, B. T. (2007).  
23 Combining differential expression, chromosomal and pathway analyses for the molecular  
24 characterization of renal cell carcinoma. 1, S21-S27.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/18542781>
- 26 Furnus, C. C.; Ulrich, M. A.; Terreros, M. C.; Dulout, F. N. (1990). The induction of aneuploidy  
27 in cultured Chinese hamster cells by propionaldehyde and chloral hydrate. *Mutagenesis*,  
28 5, 323-326. <http://www.ncbi.nlm.nih.gov/pubmed/2398816>
- 29 Fürstenberger, G. and Senn, H. J. (2002). Insulin-like growth factors and cancer. *Lancet Oncol*,  
30 3, 298-302. <http://www.ncbi.nlm.nih.gov/pubmed/12067807>
- 31 Fuscoe, J. C.; Afshari, A. J.; George, M. H.; DeAngelo, A. B.; Tice, R. R.; Salman, T.; Allen, J.  
32 W. (1996). In vivo genotoxicity of dichloroacetic acid: Evaluation with the mouse  
33 peripheral blood micronucleus assay and the single cell gel assay. *Environ Mol Mutagen*,  
34 27, 1-9. [http://dx.doi.org/10.1002/\(SICI\)1098-2280\(1996\)27:1<1::AID-EM1>3.0.CO;2-L](http://dx.doi.org/10.1002/(SICI)1098-2280(1996)27:1<1::AID-EM1>3.0.CO;2-L)
- 35 Gaglio, P. J.; Liu, H.; Dash, S.; Cheng, S.; Dunne, B.; Ratterree, M., . . . LaBrecque, D. (2002).  
36 Liver regeneration investigated in a non-human primate model (*Macaca mulatta*). *J*  
37 *Hepatol*, 37, 625-632. <http://www.ncbi.nlm.nih.gov/pubmed/12399229>
- 38 Gagnaire, F.; Marignac, B.; Langlais, C.; Bonnet, P. (2001). Ototoxicity in rats exposed to ortho-  
39 , meta- and para-xylene vapours for 13 weeks. *Pharmacol Toxicol*, 89, 6-14.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/11484912>
- 41 Gallou, C.; Longueaux, S.; Deloménie, C.; Méjean, A.; Martin, N.; Martinet, S., . . . Dupret, J.  
42 (2001). Association of GSTT1 non-null and NAT1 slow/rapid genotypes with von  
43 Hippel-Lindau tumour suppressor gene transversions in sporadic renal cell carcinoma.  
44 *Pharmacogenetics*, 11, 521-535. <http://www.ncbi.nlm.nih.gov/pubmed/11505222>

DRAFT– DO NOT QUOTE OR CITE

- 1 Galloway, S. M.; Armstrong, M. J.; Reuben, C.; Colman, S.; Brown, B.; Cannon, C., . . . Zeiger,  
2 E. (1987). Chromosome aberrations and sister chromatid exchanges in Chinese hamster  
3 ovary cells: Evaluations of 108 chemicals. *Environ Mol Mutagen*, 10, 1-175.  
4 <http://dx.doi.org/10.1002/em.2850100502>
- 5 Gama, C. and Meira, J. B. (1978). Occupational acro-osteolysis. *J Bone Joint Surg Am*, 60, 86-  
6 90.
- 7 Gamberale, G.; Annwall, G.; Olson, B. A. (1976). Exposure to trichloroethylene: III  
8 psychological functions. *Scand J Work Environ Health*, 4, 220-224.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/1019596>
- 10 Gandhi, M.; Aweeka, F.; Greenblatt, R. M.; Blaschke, T. F. (2004). Sex differences in  
11 pharmacokinetics and pharmacodynamics. *Annu Rev Pharmacol Toxicol*, 44, 499-523.  
12 <http://dx.doi.org/10.1146/annurev.pharmtox.44.101802.121453>
- 13 Gandillet, A.; Alexandre, E.; Royer, C.; Cinqualbre, J.; Jaeck, D.; Richert, L. (2003). Hepatocyte  
14 ploidy in regenerating livers after partial hepatectomy, drug-induced necrosis, and  
15 cirrhosis. *Eur Surg Res*, 35, 148-160. <http://dx.doi.org/10.1159/000070044>
- 16 Garabrant, D. H.; Lacey J. V., J.; Laing, T. J.; Gillespie, B. W.; Mayes, M. D.; Cooper, B. C.;  
17 Schottenfeld, D. (2003). Scleroderma and solvent exposure among women. *Am J*  
18 *Epidemiol*, 157, 493-500. <http://dx.doi.org/10.1093/aje/kwf223>
- 19 Garabrant, D.; Held, J.; Langholz, B.; Bernstein, L. (1988). Mortality of aircraft manufacturing  
20 workers in southern California. *Am J Ind Med*, 13, 683-693.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/3389363>
- 22 Garfinkel, L. and Stellman, S. D. (1988). Smoking and lung cancer in women: findings in a  
23 prospective study. *Cancer Res*, 48, 6951-6955.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/3180103>
- 25 Gargas, M. L.; Burgess, R. J.; Voisard, D. E.; Cason, G. H.; Andersen, M. E. (1989). Partition  
26 coefficients of low-molecular-weight volatile chemicals in various liquids and tissues.  
27 *Toxicol Appl Pharmacol*, 98, 87-99. <http://www.ncbi.nlm.nih.gov/pubmed/2929023>
- 28 Gargas, M. L.; Seybold, P. G.; Andersen, M. E. (1988). Modeling the tissue solubilities and  
29 metabolic rate constant (Vmax) of halogenated methanes, ethanes, and ethylenes. *Toxicol*  
30 *Lett*, 43, 235-256. [http://dx.doi.org/10.1016/0378-4274\(88\)90031-8](http://dx.doi.org/10.1016/0378-4274(88)90031-8)
- 31 Gart, J. J.; Chu, K. C.; Tarone, R. E. (1979). Statistical issues in interpretation of chronic  
32 bioassay tests for carcinogenicity. *J Natl Cancer Inst*, 62, 957-974.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/285297>
- 34 Garte, S.; Gaspari, L.; Alexandrie, A. K.; Ambrosone, C.; Autrup, H.; Autrup, J. L., . . . Taioli, E.  
35 (2001). Metabolic gene polymorphism frequencies in control populations. *Cancer*  
36 *Epidemiol Biomarkers Prev*, 10, 1239-1248.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/11751440>
- 38 Gash, D.; Rutland, K.; Hudson, N.; Sullivan, P.; Bing, G.; Cass, W., . . . Prince, T. (2008).  
39 Trichloroethylene: Parkinsonism and complex 1 mitochondrial neurotoxicity. *Ann*  
40 *Neurol*, 63, 184-192. <http://dx.doi.org/10.1002/ana.21288>
- 41 Ge, R.; Tao, L.; Kramer, P. M.; Cunningham, M. L.; Pereira, M. A. (2002). Effect of peroxisome  
42 proliferators on the methylation and protein level of the c-myc protooncogene in B6C3F1  
43 mice liver. *J Biochem Mol Toxicol*, 16, 41-47. <http://dx.doi.org/10.1002/jbt.10019>



DRAFT– DO NOT QUOTE OR CITE

- 1 Ge, R.; Wang, W.; Kramer, P. M.; Yang, S.; Tao, L.; Pereira, M. A. (2001). Wy-14,643-induced  
2 hypomethylation of the c-myc gene in mouse liver. *Toxicol Sci*, 62, 28-35.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/11399790>
- 4 Ge, R.; Yang, S.; Kramer, P. M.; Tao, L.; Pereira, M. A. (2001). The effect of dichloroacetic acid  
5 and trichloroacetic acid on DNA methylation and cell proliferation in B6C3F1 mice. *J*  
6 *Biochem Mol Toxicol*, 15, 100-106. <http://dx.doi.org/10.1002/jbt.5>
- 7 Gelderblom, W. C.; Galendo, D.; Abel, S.; Swanevelder, S.; Marasas, W. F.; Wild, C. P. (2001).  
8 Cancer initiation by fumonisin B(1) in rat liver--role of cell proliferation. *Cancer Lett*,  
9 169, 127-137. <http://www.ncbi.nlm.nih.gov/pubmed/11431101>
- 10 Gelfand, A. E. and Smith, A. F. M. (1990). Sampling based approaches to calculating marginal  
11 densities. *J Am Stat Assoc*, 85, 398 - 409.
- 12 Gelman, A.; Bois, F.; Jiang, J. (1996). Physiological pharmacokinetic analysis using population  
13 modeling and informative prior distributions. *J Am Stat Assoc*, 91, 1400-1412.
- 14 Gelman, A.; Carlin, J. B.; Stern, H. S.; Rubin, D. B. (2004). *Bayesian data analysis: Chapman*  
15 *and Hall*.
- 16 George, J. D.; Reel, J. R.; Myers, C. B.; Lawton, A. D. (1985). Trichloroethylene: Reproduction  
17 and fertility assessment in CD-1 mice when administered in the feed. (RTI-152).  
18 Research Triangle Park, NC: NIHEHS.
- 19 George, J. D.; Reel, J. R.; Myers, C. B.; Lawton, A. D. (1986). Trichloroethylene: Reproduction  
20 and fertility assessment in F344 rats when administered in the feed. Retrieved from  
21 <file://L:\Lab\NCEA\TCE Documents\TCE PDFs4\ George et al rats 1986 Part 1.pdf>.
- 22 George, J.; Byth, K.; Farrell, G. C. (1995). Age but not gender selectively affects expression of  
23 individual cytochrome P450 proteins in human liver. *Biochem Pharmacol*, 50, 727-730.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/7669077>
- 25 George, M. H.; Moore, T.; Kilburn, S.; Olson, G. R.; DeAngelo, A. B. (2000). Carcinogenicity  
26 of chloral hydrate administered in drinking water to the male F344/N rat and male  
27 B6C3F1 mouse. *Toxicol Pathol*, 28, 610-618.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/10930049>
- 29 Ghantous, H.; Danielsson, B. R. G.; Dencker, L.; Gorczak, J.; Vesterberg, O. (1986).  
30 Trichloroacetic acid accumulates in murine amniotic fluid after tri- and  
31 tetrachloroethylene inhalation. *Basic Clin Pharmacol Toxicol*, 58, 105-114.  
32 <http://dx.doi.org/10.1111/j.1600-0773.1986.tb00078.x>
- 33 Ghoshal, A. K. and Farber, E. (1984). The induction of liver cancer by dietary deficiency of  
34 choline and methionine without added carcinogens. *Carcinogenesis*, 5, 1367-1370.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/6488458>
- 36 Giardino, N. J. and Andelman, J. B. (1996). Characterization of the emissions of  
37 trichloroethylene, chloroform, and 1,2-dibromo-3-chloropropane in a full-size,  
38 experimental shower. *J Expo Sci Environ Epidemiol*, 6, 413-423.
- 39 Gibson, D.; Aardema, M.; Kerckaert, G.; Carr, G.; Brauning, R.; LeBoeuf, R. (1995).  
40 Detection of aneuploidy-inducing carcinogens in the Syrian hamster embryo (SHE) cell  
41 transformation assay. *Mutat Res*, 343, 7-24.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/7753109>
- 43 Gilbert, K. M.; Whitlow, A. B.; Pumford, N. R. (2004). Environmental contaminant and  
44 disinfection by-product trichloroacetaldehyde stimulates T cells in vitro. *Int*  
45 *Immunopharmacol*, 4, 25-36. <http://dx.doi.org/10.1016/j.intimp.2003.10.001>

DRAFT– DO NOT QUOTE OR CITE

- 1 Gilbert, K.; Griffin, J.; Pumford, N. (1999). Trichloroethylene activates CD4+ T cells: potential  
2 role in an autoimmune response. *Drug Metab Rev*, 31, 901-916.  
3 <http://dx.doi.org/10.1081/DMR-100101945>
- 4 Gilbert, K.; Pumford, N.; Blossom, S. (2006). Environmental contaminant trichloroethylene  
5 promotes autoimmune disease and inhibits T-cell apoptosis in MRL(+/+) mice. *J*  
6 *Immunotoxicol*, 3, 263-267. <http://dx.doi.org/10.1080/15476910601023578>
- 7 Gilkeson, G. S.; Keil, D.; Peden-Adams, M. B. J. D. B. T. I. T. t. s. b. o. r. a. (2004). Immune  
8 effects of trichloroethylene on autoimmune disease in mice. 87 - 98 PUBL- The Medical  
9 University of South Carolina Press.
- 10 Gilks, W. R.; Richardson, S.; Spiegelhalter, D. J. (1996). *Markov Chain Monte Carlo in Practice*.  
11 Boca Raton, FL: Chapman & Hall/CRC.
- 12 Giller, S.; Le Curieux, F.; Gauthier, L.; Erb, F.; Marzin, D. (1995). Genotoxicity assay of chloral  
13 hydrate and chloropicrine. *Mutat Res*, 348, 147-152.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/8544866>
- 15 Giller, S.; Le Curieux, F.; Erb, F.; Marzin, D. (1997). Comparative genotoxicity of halogenated  
16 acetic acids found in drinking water. *Mutagenesis*, 12, 321-328.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/9379909>
- 18 Ginsberg, G.; Hattis, D.; Miller, R.; Sonawane, B. (2004). Pediatric pharmacokinetic data:  
19 Implications for environmental risk assessment for children. *Pediatrics*, 113, 973-983.
- 20 Ginsberg, G.; Hattis, D.; Russ, A.; Sonawane, B. (2005). Pharmacokinetic and pharmacodynamic  
21 factors that can affect sensitivity to neurotoxic sequelae in elderly individuals. *Environ*  
22 *Health Perspect*, 113, 1243-1249. <http://dx.doi.org/10.1289/ehp.7568>
- 23 Ginsberg, G.; Hattis, D.; Sonawane, B. (2004). Incorporating pharmacokinetic differences  
24 between children and adults in assessing children's risks to environmental toxicants.  
25 *Toxicol Appl Pharmacol*, 198, 164-183. <http://dx.doi.org/10.1016/j.taap.2003.10.010>
- 26 Ginsberg, G.; Hattis, D.; Sonawane, B.; Russ, A.; Banati, P.; Kozlak, M., . . . Goble, R. (2002).  
27 Evaluation of child/adult pharmacokinetic differences from a database derived from the  
28 therapeutic drug literature. *Toxicol Sci*, 66, 185-200.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/11896285>
- 30 Gochfeld, M. (2007). Framework for gender differences in human and animal toxicology.  
31 *Environ Res*, 104, 4-21. <http://dx.doi.org/10.1016/j.envres.2005.12.005>
- 32 Goel, S.; Rao, G.; Pandya, K.; Shanker, R. (1992). Trichloroethylene toxicity in mice: A  
33 biochemical, hematological and pathological assessment. *Indian J Exp Biol*, 30, 402-406.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/1459618>
- 35 Goeptar, A. R.; Commandeur, J. N. M.; van Ommen, B.; van Bladeren, P. J.; Vermeulen, N. P.  
36 E. (1995). Metabolism and kinetics of trichloroethylene in relation to toxicity and  
37 carcinogenicity Relevance of the mercapturic acid pathway. *Chem Res Toxicol*, 8, 3-21.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/7703363>
- 39 Goh, V. H.; Chia, S. E.; Ong, C. N. (1998). Effects of chronic exposure to low doses of  
40 trichloroethylene on steroid hormone and insulin levels in normal men. *Environ Health*  
41 *Perspect*, 106, 41-44. <http://www.ncbi.nlm.nih.gov/pubmed/9417767>
- 42 Gold, L. S. and E, Z. (Eds.). (1997). *Handbook of carcinogenic potency and genotoxicity*  
43 *databases*. Boca Raton: CRC Press.
- 44 Gold, L. S.; Stewart, P. A.; Milliken, K.; Purdue, M.; Severson, R.; Seixas, N., . . . De Roos, A.  
45 J. (In Press). The relationship between multiple myeloma and occupational exposure to

- 1 six chlorinated solvents. *Occup Environ Med*.  
2 <http://dx.doi.org/10.1136/oem.2009.054809>
- 3 Goldberg, M.; Parent, M.; Siemiatycki, J.; Désy, M.; Nadon, L.; Richardson, L., . . . Valois, M.  
4 (2001). A case-control study of the relationship between the risk of colon cancer in men  
5 and exposures to occupational agents. *Am J Ind Med*, 39, 531-546.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/11385637>
- 7 Goldberg, S.; Lebowitz, M.; Graver, E.; Hicks, S. (1990). An association of human congenital  
8 cardiac malformations and drinking water contaminants. *J Am Coll Cardiol*, 16, 155-164.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/2358589>
- 10 Goldsworthy, T. L. and Popp, J. A. (1987). Chlorinated hydrocarbon-induced peroxisomal  
11 enzyme activity in relation to species and organ carcinogenicity. *Toxicol Appl*  
12 *Pharmacol*, 88, 225-233. [http://dx.doi.org/10.1016/0041-008X\(87\)90008-1](http://dx.doi.org/10.1016/0041-008X(87)90008-1)
- 13 Goldsworthy, T. L.; Lyght, O.; Burnett, V. L.; Popp, J. A. (1988). Potential role of [alpha]-  
14 2[mu]-globulin, protein droplet accumulation, and cell replication in the renal  
15 carcinogenicity of rats exposed to trichloroethylene, perchloroethylene, and  
16 pentachloroethane. *Toxicol Appl Pharmacol*, 96, 367-379.  
17 [http://dx.doi.org/10.1016/0041-008X\(88\)90095-6](http://dx.doi.org/10.1016/0041-008X(88)90095-6)
- 18 Gomez, M.; Cocco, P.; Dosemeci, M.; Stewart, P. (1994). Occupational exposure to chlorinated  
19 aliphatic hydrocarbons: Job exposure matrix. *Am J Ind Med*, 26, 171-183.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/7977394>
- 21 Gonzalez-Leon, A.; Merdink, J.; Bull, R.; Schultz, I. (1999). Effect of pre-treatment with  
22 dichloroacetic or trichloroacetic acid in drinking water on the pharmacokinetics of a  
23 subsequent challenge dose in B6C3F1 mice. *Chem Biol Interact*, 123, 239-253.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/10654841>
- 25 Gorla, G. R.; Malhi, H.; Gupta, S. (2001). Polyploidy associated with oxidative injury attenuates  
26 proliferative potential of cells. *J Cell Sci*, 114, 2943-2951.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/11686298>
- 28 Graham, D. J.; Staffa, J. A.; Shatin, D.; Andrade, S. E.; Schech, S. D.; La Grenade, L., . . . Platt,  
29 R. (2004). Incidence of hospitalized rhabdomyolysis in patients treated with lipid-  
30 lowering drugs. *JAMA*, 292, 2585-2590. <http://dx.doi.org/10.1001/jama.292.21.2585>
- 31 Gramantieri, L.; Melchiorri, C.; Chieco, P.; Gaiani, S.; Stecca, B.; Casali, A.; Bolondi, L. (1996).  
32 Alteration of DNA ploidy and cell nuclearity in human hepatocellular carcinoma  
33 associated with HBV infection. *J Hepatol*, 25, 848-853.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/9007712>
- 35 Grandjean, E.; Munchinger, R.; Turrian, V.; Haas, P. A.; Knoepfel, H.-K.; Rosenmund, H.  
36 (1955). Investigations into the effects of exposure to trichlorethylene in mechanical  
37 engineering. *Occup Environ Med*, 12, 131-142.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/14363593>
- 39 Grasl-Kraupp, B.; Waldhör, T.; Huber, W.; Schulte-Hermann, R. (1993). Glutathione S-  
40 transferase isoenzyme patterns in different subtypes of enzyme-altered rat liver foci  
41 treated with the peroxisome proliferator nafenopin or with phenobarbital. *Carcinogenesis*,  
42 14, 2407-2412. <http://www.ncbi.nlm.nih.gov/pubmed/8242872>
- 43 Grawé, J.; Nüsse, M.; Adler, I. (1997). Quantitative and qualitative studies of micronucleus  
44 induction in mouse erythrocytes using flow cytometry. I. Measurement of micronucleus  
45 induction in peripheral blood polychromatic erythrocytes by chemicals with known and

DRAFT– DO NOT QUOTE OR CITE

- 1 suspected genotoxicity. *Mutagenesis*, 12, 1-8.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/9025090>
- 3 Green, L. and Lash, T. (1999). Re: "Renal cell cancer correlated with occupational  
4 exposure to trichloroethylene". *J Cancer Res Clin Oncol*, 125, 430-432.  
5 <http://www.ncbi.nlm.nih.gov/pubmed/10394965>
- 6 Green, T. (2000). Pulmonary toxicity and carcinogenicity of trichloroethylene: species  
7 differences and modes of action. *Environ Health Perspect*, 108 Suppl 2, 261-264.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/10807556>
- 9 Green, T. and Prout, M. S. (1985). Species differences in response to trichloroethylene: II  
10 Biotransformation in rats and mice. *Toxicol Appl Pharmacol*, 79, 401-411.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/3929429>
- 12 Green, T.; Dow, J.; Ellis, M. K.; Foster, J. R.; Odum, J. (1997). The role of glutathione  
13 conjugation in the development of kidney tumours in rats exposed to trichloroethylene.  
14 *Chem Biol Interact*, 105, 99-117. <http://www.ncbi.nlm.nih.gov/pubmed/9251723>
- 15 Green, T.; Dow, J.; Foster, J. (2003). Increased formic acid excretion and the development of  
16 kidney toxicity in rats following chronic dosing with trichloroethanol, a major metabolite  
17 of trichloroethylene. *Toxicology*, 191, 109-119.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/12965114>
- 19 Green, T.; Dow, J.; Foster, J.; Hext, P. (1998). Formic acid excretion in rats exposed to  
20 trichloroethylene: a possible explanation for renal toxicity in long-term studies.  
21 *Toxicology*, 127, 39-47. <http://www.ncbi.nlm.nih.gov/pubmed/9699792>
- 22 Green, T.; Dow, J.; Ong, C.; Ng, V.; Ong, H.; Zhuang, Z., . . . Bloemen, L. (2004). Biological  
23 monitoring of kidney function among workers occupationally exposed to  
24 trichloroethylene. *Occup Environ Med*, 61, 312-317.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/15031388>
- 26 Green, T.; Mainwaring, G. W.; Foster, J. R. (1997). Trichloroethylene-induced mouse lung  
27 tumors: studies of the mode of action and comparisons between species. *Fundam Appl*  
28 *Toxicol*, 37, 125-130. <http://dx.doi.org/10.1006/faat.1997.2312>
- 29 Green, T.; Odum, J.; Nash, J. A.; Foster, J. R. (1990). Perchloroethylene-induced rat kidney  
30 tumors: An investigation of the mechanisms involved and their relevance to humans.  
31 *Toxicol Appl Pharmacol*, 103, 77-89. [http://dx.doi.org/10.1016/0041-008X\(90\)90264-U](http://dx.doi.org/10.1016/0041-008X(90)90264-U)
- 32 Greenberg, M.; Burton, G.; Fisher, J. (1999). Physiologically based pharmacokinetic modeling of  
33 inhaled trichloroethylene and its oxidative metabolites in B6C3F1 mice. *Toxicol Appl*  
34 *Pharmacol*, 154, 264-278. <http://dx.doi.org/10.1006/taap.1998.8594>
- 35 Greene, A. K.; Wiener, S.; Puder, M.; Yoshida, A.; Shi, B.; Perez-Atayde, A. R., . . . O'Reilly,  
36 M. S. (2003). Endothelial-directed hepatic regeneration after partial hepatectomy. *Ann*  
37 *Surg*, 237, 530-535. <http://dx.doi.org/10.1097/01.SLA.0000059986.96051.EA>
- 38 Greenland, S. (1992). A semi-Bayes approach to the analysis of correlated multiple associations,  
39 with an application to an occupational cancer-mortality study. *Stat Med*, 11, 219-230.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/1579760>
- 41 Greenland, S.; Salvan, A.; Wegman, D. H.; Hallock, M. F.; Smith, T. J. (1994). A case-control  
42 study of cancer mortality at a transformer-assembly facility. *Int Arch Occup Environ*  
43 *Health*, 66, 49-54. <http://dx.doi.org/10.1007/BF00386579>

DRAFT– DO NOT QUOTE OR CITE

- 1 Gregersen, P. (1988). Neurotoxic effects of organic solvents in exposed workers: two controlled  
2 follow-up studies after 55 and 106 years. *Am J Ind Med*, 14, 681-701.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/2852897>
- 4 Greim, H.; Bonse, G.; Radwan, Z.; Reichert, D.; Henschler, D. (1975). Mutagenicity in vitro and  
5 potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane  
6 formation. *Biochem Pharmacol*, 24, 2013-2017. [http://dx.doi.org/10.1016/0006-  
7 2952\(75\)90396-2](http://dx.doi.org/10.1016/0006-2952(75)90396-2)
- 8 Greten, F. R.; Eckmann, L.; Greten, T. F.; Park, J. M.; Li, Z. W.; Egan, L. J., . . . Karin, M.  
9 (2004). IKKbeta links inflammation and tumorigenesis in a mouse model of colitis-  
10 associated cancer. *Cell*, 118, 285-296. <http://dx.doi.org/10.1016/j.cell.2004.07.013>
- 11 Griffin, J. M.; Blossom, S. J.; Jackson, S. K.; Gilbert, K. M.; Pumford, N. R. (2000).  
12 Trichloroethylene accelerates an autoimmune response by Th1 T cell activation in MRL  
13 +/+ mice. *Immunopharmacology*, 46, 123-137. [http://dx.doi.org/10.1016/S0162-  
14 3109\(99\)00164-2](http://dx.doi.org/10.1016/S0162-3109(99)00164-2)
- 15 Griffin, J. M.; Gilbert, K. M.; Lamps, L. W.; Pumford, N. R. (2000). CD4(+) T-cell activation  
16 and induction of autoimmune hepatitis following trichloroethylene treatment in MRL+/+  
17 mice. *Toxicol Sci*, 57, 345-352. <http://www.ncbi.nlm.nih.gov/pubmed/11006364>
- 18 Griffin, J. M.; Gilbert, K. M.; Pumford, N. R. (2000). Inhibition of CYP2E1 reverses CD4+ T-  
19 cell alterations in trichloroethylene-treated MRL+/+ mice. *Toxicol Sci*, 54, 384-389.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/10774820>
- 21 Grosse, Y.; Baan, R.; Secretan-lauby, B.; El ghissassi, F.; Bouvard, V.; Benbrahim-tallaa, L., . . .  
22 Straif, K. (2011). Carcinogenicity of chemicals in industrial and consumer products, food  
23 contaminants and flavourings, and water chlorination byproducts. *Lancet Oncol*, 12, 328-  
24 329. [http://dx.doi.org/10.1016/S1470-2045\(11\)70088-2](http://dx.doi.org/10.1016/S1470-2045(11)70088-2)
- 25 Groupwater Vapor Project, Endicott, New York:,. (2006). Summary of findings, working draft.  
26 Cited in ATSDR. Endicott Area, Town of Union, Broome County, New York: Health  
27 Statistics Reviews, Cancer and Birth Outcome Analysis.
- 28 Groves, C.; Hayden, P.; Lock, E.; Schnellmann, R. (1993). Differential cellular effects in the  
29 toxicity of haloalkene and haloalkane cysteine conjugates to rabbit renal proximal  
30 tubules. *J Biochem Toxicol*, 8, 49-56. <http://www.ncbi.nlm.nih.gov/pubmed/8492303>
- 31 Gu, Z. W.; Sele, B.; Chmara, D.; Jalbert, P.; Vincent, M.; Vincent, F., . . . Faure, J. (1981).  
32 [Effects of trichlorethylene and its metabolites on the rate of sister chromatid exchange.  
33 In vivo and in vitro study on the human lymphocytes]. *Ann Genet*, 24, 105-106.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/6977287>
- 35 Gu, Z. W.; Sele, B.; Jalbert, P.; Vincent, M.; Vin. (1981). Induction d'echanges entre les  
36 chromatides soeurs (SCE) par le trichloroethylene et ses metabolites [Induction of sister  
37 chromatid exchange by trichloroethylene and its metabolites]. *Toxicol Eur Res*, 3, 63-67.
- 38 Gualandi, G. (1987). Use of alpha- and beta-tubulin mutants for the study of spontaneous and  
39 induced chromosomal mis-distribution in *Aspergillus nidulans*. *Mutat Res*, 178, 33-41.
- 40 Guehl, D.; Bezar, E.; Dovero, S.; Boraud, T.; Bioulac, B.; Gross, C. (1999). Trichloroethylene  
41 and parkinsonism: A human and experimental observation. *Eur J Neurol*, 6, 609-611.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/10457397>
- 43 Guengerich, F. P. (2004). Metabolism of trichlorethylene and covalent binding of reaction  
44 products. *Proceedings of symposium on new scientific research related to the health  
45 effects of trichlorethylene*. Washington, DC.

DRAFT– DO NOT QUOTE OR CITE

- 1 Guengerich, F. P. and Shimada, T. (1991). Oxidation of toxic and carcinogenic chemicals by  
2 human cytochrome P-450 enzymes. *Chem Res Toxicol*, 4, 391-407.  
3 <http://dx.doi.org/10.1021/tx00022a001>
- 4 Guengerich, F. P.; Crawford, W. M. J.; Watanabe, P. G. (1979). Activation of vinyl chloride to  
5 covalently bound metabolites: roles of 2-chloroethylene oxide and 2-chloroacetaldehyde.  
6 *Biochemistry*, 18, 5177-5182.
- 7 Guengerich, F. P.; Kim, D.-H.; Iwasaki, M. (1991). Role of human cytochrome P-450 IIE1 in the  
8 oxidation of many low molecular weight cancer suspects. *Chem Res Toxicol*, 4, 168-179.  
9 <http://dx.doi.org/10.1021/tx00020a008>
- 10 Gun, R. T.; Grygorcewicz, C.; Nettelbeck, T. J. (1978). Choice reaction time in workers using  
11 trichloroethylene. *Med J Aust*, 1, 535-536. <http://www.ncbi.nlm.nih.gov/pubmed/683058>
- 12 Gupta, S. (2000). Hepatic polyploidy and liver growth control. *Semin Cancer Biol*, 10, 161-171.  
13 <http://dx.doi.org/10.1006/scbi.2000.0317>
- 14 Gustavsson, P.; Jakobsson, R.; Johansson, H.; Lewin, F.; Norell, S.; Rutkvist, L. E. (1998).  
15 Occupational exposures and squamous cell carcinoma of the oral cavity, pharynx, larynx,  
16 and oesophagus: A case-control study in Sweden. *Occup Environ Med*, 55, 393-400.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/9764099>
- 18 Guyton, K. Z.; Barone, S.; Brown, R. C.; Euling, S. Y.; Jinot, J.; Makris, S. (2008). Mode of  
19 action frameworks: a critical analysis. *J Toxicol Environ Health B Crit Rev*, 11, 16-31.  
20 <http://dx.doi.org/10.1080/10937400701600321>
- 21 Guyton, K. Z.; Chiu, W. A.; Bateson, T. F.; Jinot, J.; Scott, C. S.; Brown, R. C.; Caldwell, J. C.  
22 (2009). A reexamination of the PPAR-alpha activation mode of action as a basis for  
23 assessing human cancer risks of environmental contaminants. *Environ Health Perspect*,  
24 117, 1664-1672. <http://dx.doi.org/10.1289/ehp.0900758>
- 25 Gyorke, S. and Terentyev, D. (2008). Modulation of ryanodine receptor by luminal calcium and  
26 accessory proteins in health and cardiac disease. *Cardiovasc Res*, 77, 245-255.  
27 <http://dx.doi.org/10.1093/cvr/cvm038>
- 28 Habib, G. M.; Barrios, R.; Shi, Z. Z.; Lieberman, M. W. (1996). Four distinct membrane-bound  
29 dipeptidase RNAs are differentially expressed and show discordant regulation with  
30 gamma-glutamyl transpeptidase. *J Biol Chem*, 271, 16273-16280.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/8663190>
- 32 Hack, C. E.; Chiu, W. A.; Jay Zhao, Q.; Clewell, H. J. (2006). Bayesian population analysis of a  
33 harmonized physiologically based pharmacokinetic model of trichloroethylene and its  
34 metabolites. *Regul Toxicol Pharmacol*, 46, 63-83.  
35 <http://dx.doi.org/10.1016/j.yrtph.2006.05.012>
- 36 Haddad, S.; Charest-Tardif, G.; K, K. (2000). Physiologically based modeling of the maximal  
37 effect of metabolic interactions on the kinetics of components of complex chemical  
38 mixtures. *J Toxicol Environ Health A*, 61, 209-223.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/11036509>
- 40 Haglid, K. G.; Briving, C.; Hansson, H.-A.; Rosengren, L.; Kjellstrand, P.; Stavron, D., . . .  
41 Wronski, A. (1981). Trichloroethylene: long-lasting changes in the brain after  
42 rehabilitation. *Neurotoxicology*, 2, 659-673.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/7200581>
- 44 Hakkola, J.; Pasanen, M.; Hukkanen, J.; Pelkonen, O.; Maenpaa, J.; Edwards, R. J., . . . Raunio,  
45 H. (1996). Expression of xenobiotic-metabolizing cytochrome P450 forms in human full-

- 1 term placenta. *Biochem Pharmacol*, 51, 403-411. <http://dx.doi.org/10.1016/0006->  
2 2952(95)02184-1
- 3 Hakkola, J.; Pelkonen, O.; Pasanen, M.; Raunio, H. (1998). Xenobiotic-metabolizing cytochrome  
4 P450 enzymes in the human fetoplacental unit: Role in intrauterine toxicity. *Crit Rev*  
5 *Toxicol*, 28, 35-72. <http://dx.doi.org/10.1080/10408449891344173>
- 6 Hakkola, J.; Raunio, H.; Purkunen, R.; Pelkonen, O.; Saarikoski, S.; Cresteil, T.; Pasanen, M.  
7 (1996). Detection of cytochrome P450 gene expression in human placenta in first  
8 trimester of pregnancy. *Biochem Pharmacol*, 52, 379-383.  
9 [http://dx.doi.org/10.1016/0006-2952\(96\)00216-X](http://dx.doi.org/10.1016/0006-2952(96)00216-X)
- 10 Hakkola, J.; Tanaka, E.; Pelkonen, O. (1998). Developmental expression of cytochrome P450  
11 enzymes in human liver. *Pharmacol Toxicol*, 82, 209-217.  
12 <http://dx.doi.org/10.1111/j.1600-0773.1998.tb01427.x>
- 13 Halmes, N.; Perkins, E.; McMillan, D.; Pumford, N. (1997). Detection of trichloroethylene-  
14 protein adducts in rat liver and plasma. *Toxicol Lett*, 92, 187-194.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/9334829>
- 16 Hamada, T. and Tanaka, H. (1995). Transfer of methyl chloroform, trichloroethylene and  
17 tetrachloroethylene to milk, tissues and expired air following intraruminal or oral  
18 administration in lactating goats and milk-fed kids. *Environ Pollut*, 87, 313-318.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/15091581>
- 20 Hamdan, H. and Stacey, N. H. (1993). Mechanism of trichloroethylene-induced elevation of  
21 individual serum bile acids: I correlation of trichloroethylene concentrations to bile acids  
22 in rat serum. *Toxicol Appl Pharmacol*, 121, 291-295.  
23 <http://dx.doi.org/10.1006/taap.1993.1156>
- 24 Hanahan, D. and Weinberg, R. A. (2000). The hallmarks of cancer. *Cell*, 100, 57-70.  
25 [http://dx.doi.org/10.1016/S0092-8674\(00\)81683-9](http://dx.doi.org/10.1016/S0092-8674(00)81683-9)
- 26 Hansch, C.; Leo, A.; Hoekman, D. (1995). Exploring QSAR: Hydrophobic, electronic, and steric  
27 constants. Washington, DC: American Chemical Society.
- 28 Hansen, J.; Raaschou-Nielsen, O.; Christensen, J. M.; Johansen, I.; McLaughlin, J. K.; Lipworth,  
29 L., . . . Olsen, J. H. (2001). Cancer incidence among Danish workers exposed to  
30 trichloroethylene. *J Occup Environ Med*, 43, 133-139.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/11227631>
- 32 Hardell, L.; Bengtsson, N. O.; Jonsson, U.; Eriksson, S.; Larsson, L. G. (1984). Aetiological  
33 aspects on primary liver cancer with special regard to alcohol, organic solvents and acute  
34 intermittent porphyria - An epidemiological investigation. *Br J Cancer*, 50, 389-397.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/6087869>
- 36 Hardell, L.; Eriksson, M.; Degerman, A. (1994). Exposure to phenoxyacetic acids,  
37 chlorophenols, or organic solvents in relation to histopathology, stage, and anatomical  
38 localization of non-Hodgkin's lymphoma. *Cancer Res*, 54, 2386-2389.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/8162585>
- 40 Hardell, L.; Eriksson, M.; Lenner, P.; Lundgren, E. (1981). Malignant lymphoma and exposure  
41 to chemicals, especially organic solvents, chlorophenols and phenoxy acids: A case-  
42 control study. *Br J Cancer*, 43, 169-176. <http://www.ncbi.nlm.nih.gov/pubmed/7470379>
- 43 Hardin, B. D.; Bond, G. P.; Sikov, M. R.; Andrew, F. D.; Beliles, R. P.; Niemeier, R. W. (1981).  
44 Testing of selected workplace chemicals for teratogenic potential. *Scand J Work Environ*  
45 *Health*, 7, 66-75. <http://www.ncbi.nlm.nih.gov/pubmed/7330632>

DRAFT– DO NOT QUOTE OR CITE

- 1 Hardin, B. D.; Kelman, B. J.; Brent, R. L. (2004). Trichloroethylene and cardiac malformations.  
2 Environ Health Perspect, 112, A607-608; author reply A608-609.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/15289177>
- 4 Hardin, B.; Kelman, B.; Brent, R. (2005). Trichloroethylene and dichloroethylene: a critical  
5 review of teratogenicity. Birth Defects Res A Clin Mol Teratol, 73, 931-955.  
6 <http://dx.doi.org/10.1002/bdra.20192>
- 7 Harrington, J. M.; Whitby, H.; Gray, C. N.; Reid, F. J.; Aw, T. C.; Waterhouse, J. A. (1989).  
8 Renal disease and occupational exposure to organic solvents: A case referent approach.  
9 Br J Ind Med, 46, 643-650. <http://dx.doi.org/10.1136/oem.46.9.643>
- 10 Harrington-Brock, K.; Doerr, C. L.; Moore, M. M. (1998). Mutagenicity of three disinfection by-  
11 products: Di- and trichloroacetic acid and chloral hydrate in L5178Y(+/-) --3.7.2C mouse  
12 lymphoma cells. Mutat Res Genet Toxicol Environ Mutagen, 413, 265-276.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/9651541>
- 14 Harris, N. L.; Jaffe, E. S.; Diebold, J.; Flandrin, G.; Muller-Hermelink, H. K.; Vardiman, J.  
15 (2000). Lymphoma classification--from controversy to consensus: The R.E.A.L. and  
16 WHO Classification of lymphoid neoplasms. Ann Oncol, 11, S3-S10.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/10707771>
- 18 Haseman, J. K.; Huff, J.; Boorman, G. A. (1984). Use of historical control data in  
19 carcinogenicity studies in rodents. Toxicol Pathol, 12, 126-135.  
20 <http://dx.doi.org/10.1177/019262338401200203>
- 21 Hasmall, S. C. and Roberts, R. A. (2000). The nongenotoxic hepatocarcinogens  
22 diethylhexylphthalate and methylclofenapate induce DNA synthesis preferentially in  
23 octoploid rat hepatocytes. Toxicol Pathol, 28, 503-509.  
24 <http://dx.doi.org/10.1177/019262330002800401>
- 25 Hassall, C.; Brendel, K.; Gandolfi, A. (1983). Regulation of a S(trans-1,2-dichlorovinyl)-L-  
26 cysteine-induced renal tubular toxicity by glutathione. J Appl Toxicol, 3, 321-325.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/6677655>
- 28 Hassan, M. M.; Frome, A.; Patt, Y. Z.; El-Serag, H. B. (2002). Rising prevalence of hepatitis C  
29 virus infection among patients recently diagnosed with hepatocellular carcinoma in the  
30 United States. J Clin Gastroenterol, 35, 266-269.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/12192205>
- 32 Hattis, D.; Ginsberg, G.; Sonawane, B.; Smolenski, S.; Russ, A.; Kozlak, M.; Goble, R. (2003).  
33 Differences in pharmacokinetics between children and adults: II. Childrens variability in  
34 drug elimination half-lives and in some parameters needed for physiologically-based  
35 pharmacokinetic modeling. Risk Anal, 23, 117-142. <http://dx.doi.org/10.1111/1539-6924.00295>
- 36  
37 Haworth, S.; Lawlor, T.; Mortelmans, K.; Speck, W.; Zeiger, E. (1983). Salmonella mutagenicity  
38 test results for 250 chemicals. Environ Mutagen, 5, 3-142.  
39 <http://dx.doi.org/10.1002/em.2860050703>
- 40 Hayden, M. S. and Ghosh, S. (2004). Signaling to NF-kappaB. Genes Dev, 18, 2195-2224.  
41 <http://dx.doi.org/10.1101/gad.1228704>
- 42 Hays, T.; Rusyn, I.; Burns, A. M.; Kennett, M. J.; Ward, J. M.; Gonzalez, F. J.; Peters, J. M.  
43 (2005). Role of peroxisome proliferator-activated receptor-alpha (PPARalpha) in  
44 bezafibrate-induced hepatocarcinogenesis and cholestasis. Carcinogenesis, 26, 219-227.  
45 <http://dx.doi.org/10.1093/carcin/bgh285>



DRAFT– DO NOT QUOTE OR CITE

- 1 Healy, T. E. J.; Poole, T. R.; Hopper, A. (1982). Rat fetal development and maternal exposure to  
2 trichloroethylene 100 ppm. *Br J Anaesth*, 54, 337-341.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/7066129>
- 4 Heath, C. W.; Lally, C. A.; Calle, E. E.; McLaughlin, J. K.; Thun, M. J. (1997). Hypertension,  
5 diuretics, and antihypertensive medications as possible risk factors for renal cell cancer.  
6 *Am J Epidemiol*, 145, 607-613. <http://www.ncbi.nlm.nih.gov/pubmed/9098177>
- 7 Hegi, M. E.; Fox, T. R.; Belinsky, S. A.; Devereux, T. R.; Anderson, M. W. (1993). Analysis of  
8 activated protooncogenes in B6C3F1 mouse liver tumors induced by ciprofibrate, a  
9 potent peroxisome proliferator. *Carcinogenesis*, 14, 145-149.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/8425263>
- 11 Hein, M. J.; Waters, M. A.; Ruder, A. M.; Stenzel, M. R.; Blair, A.; Stewart, P. A. (2010).  
12 Statistical modeling of occupational chlorinated solvent exposures for case-control  
13 studies using a literature-based database. *Ann Occup Hyg*, 54, 459-472.  
14 <http://dx.doi.org/10.1093/annhyg/meq027>
- 15 Heineman, E. F.; Cocco, P.; Gomez, M. R.; Dosemeci, M.; Stewart, P. A.; Hayes, R. B., . . .  
16 Blair, A. (1994). Occupational exposure to chlorinated aliphatic hydrocarbons and risk of  
17 astrocytic brain cancer. *Am J Ind Med*, 26, 155-169.  
18 <http://dx.doi.org/10.1002/ajim.4700260203>
- 19 Heinemann, K.; Willich, S.; Heinemann, L.; DoMinh, T.; Möhner, M.; Heuchert, G. (2000).  
20 Occupational exposure and liver cancer in women: results of the Multicentre  
21 International Liver Tumour Study (MILTS). *Occup Med (Lond)*, 50, 422-429.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/10994245>
- 23 Hejtmancik, M. R.; Trela, B. A.; Kurtz, P. J.; Persing, R. L.; Ryan, M. J.; Yarrington, J. T.;  
24 Chhabra, R. S. (2002). Comparative gavage subchronic toxicity studies of o-chloroaniline  
25 and m-chloroaniline in F344 rats and B6C3F1 mice. *Toxicol Sci*, 69, 234-243.
- 26 Helliwell, P. J. and Hutton, A. M. (1950). Trichlorethylene anesthesia. *Anaesthesia*, 5, 4-13.  
27 <http://dx.doi.org/10.1111/j.1365-2044.1950.tb13680.x>
- 28 Henderson, G.; Yan, Z.; James, M.; Davydova, N.; Stacpoole, P. (1997). Kinetics and  
29 metabolism of chloral hydrate in children: identification of dichloroacetate as a  
30 metabolite. *Biochem Biophys Res Commun*, 235, 695-698.  
31 <http://dx.doi.org/10.1006/bbrc.1997.6868>
- 32 Hendler, A. H. and Crow, W. L. (1992). Paper No. 92-75.05.
- 33 Hendriksen, P. J.; Freidig, A. P.; Jonker, D.; Thissen, U.; Bogaards, J. J.; Mumtaz, M. M., . . .  
34 Stierum, R. H. (2007). Transcriptomics analysis of interactive effects of benzene,  
35 trichloroethylene and methyl mercury within binary and ternary mixtures on the liver and  
36 kidney following subchronic exposure in the rat. *Toxicol Appl Pharmacol*, 225, 171-188.  
37 <http://dx.doi.org/10.1016/j.taap.2007.08.017>
- 38 Hennig, U. G.; Rudd, N. L.; Hoar, D. I. (1988). Kinetochore immunofluorescence in  
39 micronuclei: a rapid method for the in situ detection of aneuploidy and chromosome  
40 breakage in human fibroblasts. *Mutat Res*, 203, 405-414.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/3057373>
- 42 Henning, S. M. and Swendseid, M. E. (1996). The role of folate, choline, and methionine in  
43 carcinogenesis induced by methyl-deficient diets. *Adv Exp Med Biol*, 399, 143-155.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/8937554>

DRAFT– DO NOT QUOTE OR CITE

- 1 Henschler, D.; Eder, E.; Neudecker, T.; Metzler, M. (1977). Carcinogenicity of  
2 trichloroethylene: fact or artifact? *Arch Toxicol*, 37, 233-236.
- 3 Henschler, D.; Elsaesser, H.; Romen, W.; Eder, E. (1984). Carcinogenicity study of  
4 trichloroethylene, with and without epoxide stabilizers, in mice. *J Cancer Res Clin Oncol*,  
5 107, 149-156. <http://www.ncbi.nlm.nih.gov/pubmed/6736101>
- 6 Henschler, D.; Romen, W.; Elsaesser, H. M.; Reichert, D.; Eder, E.; Radwan, Z. (1980).  
7 Carcinogenicity study of trichloroethylene by longterm inhalation in three animal species.  
8 *Arch Toxicol*, 43, 237-248.
- 9 Henschler, D.; Vamvakas, S.; Lammert, M.; Dekant, W.; Kraus, B.; Thomas, B.; Ulm, K. (1995).  
10 Increased incidence of renal cell tumors in a cohort of cardboard workers exposed to  
11 trichloroethene. *Arch Toxicol*, 69, 291-299.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/7654132>
- 13 Herbert, V.; Gardner, A.; Colman, N. (1980). Mutagenicity of dichloroacetate, an ingredient of  
14 some formulations of pangamic acid (trade-named &quot;vitamin B15&quot;). *Am J*  
15 *Clin Nutr*, 33, 1179-1182. <http://www.ncbi.nlm.nih.gov/pubmed/6992558>
- 16 Herman, J.; Umar, A.; Polyak, K.; Graff, J.; Ahuja, N.; Issa, J., . . . Baylin, S. (1998). Incidence  
17 and functional consequences of hMLH1 promoter hypermethylation in colorectal  
18 carcinoma. *PNAS*, 95, 6870-6875. <http://www.ncbi.nlm.nih.gov/pubmed/9618505>
- 19 Hernberg, S.; Kauppinen, T.; Riala, R.; Korkala, M. L.; Asikainen, U. (1988). Increased risk for  
20 primary liver cancer among women exposed to solvents. *Scand J Work Environ Health*,  
21 14, 356-365. <http://www.ncbi.nlm.nih.gov/pubmed/3212412>
- 22 Hernberg, S.; Korkala, M. L.; Asikainen, U.; Riala, R. (1984). Primary liver cancer and exposure  
23 to solvents. *Int Arch Occup Environ Health*, 54, 147-153.  
24 <http://dx.doi.org/10.1007/BF00378517>
- 25 Herren-Freund, S. L.; Pereira, M. A.; Khoury, M. D.; Olson, G. (1987). The carcinogenicity of  
26 trichloroethylene and its metabolites, trichloroacetic acid and dichloroacetic acid, in  
27 mouse liver. *Toxicol Appl Pharmacol*, 90, 183-189. [http://dx.doi.org/10.1016/0041-](http://dx.doi.org/10.1016/0041-008X(87)90325-5)  
28 [008X\(87\)90325-5](http://dx.doi.org/10.1016/0041-008X(87)90325-5)
- 29 Herrera, E.; Samper, E.; Martín-Caballero, J.; Flores, J. M.; Lee, H. W.; Blasco, M. A. (1999).  
30 Disease states associated with telomerase deficiency appear earlier in mice with short  
31 telomeres. *EMBO J*, 18, 2950-2960. <http://dx.doi.org/10.1093/emboj/18.11.2950>
- 32 Hers, I.; Zapf-Gilje, R.; Li, L.; Atwater, J. (2001). The use of indoor air measurements to  
33 evaluate intrusion of subsurface VOC vapors into buildings. *J Air Waste Manag Assoc*,  
34 51, 1318-1331. <http://www.ncbi.nlm.nih.gov/pubmed/11575885>
- 35 Higgins, J. P.; Thompson, S. G.; Deeks, J. J.; Altman, D. G. (2003). Measuring inconsistency in  
36 meta-analyses. *Br Med J Int Ed*, 327, 557-560.  
37 <http://dx.doi.org/10.1136/bmj.327.7414.557>
- 38 Hill, A. B. (1965). The environment and disease: Association or causation? *Proc R Soc Med*, 58,  
39 295-300. <http://www.ncbi.nlm.nih.gov/pubmed/14283879>
- 40 Hinchman, C. A. and Ballatori, N. (1990). Glutathione-degrading capacities of liver and kidney  
41 in different species. *Biochem Pharmacol*, 40, 1131-1135. [http://dx.doi.org/10.1016/0006-](http://dx.doi.org/10.1016/0006-2952(90)90503-D)  
42 [2952\(90\)90503-D](http://dx.doi.org/10.1016/0006-2952(90)90503-D)
- 43 Hines, R. N. and McCarver, D. G. (2002). The ontogeny of human drug-metabolizing enzymes:  
44 Phase I oxidative enzymes. *J Pharmacol Exp Ther*, 300, 355-360.  
45 <http://dx.doi.org/10.1124/jpet.300.2.355>

DRAFT– DO NOT QUOTE OR CITE

- 1 Hirsch, A. R.; Rankin, K. M.; Panelli, P. P. (1996). Trichloroethylene exposure and headache. 7,  
2 126 - 138.
- 3 Hissink, E. M.; Bogaards, J. J. P.; Freidig, A. P.; Commandeur, J. N. M.; Vermeulen, N. P. E.;  
4 Bladeren, P. J. v. (2002). The use of in vitro metabolic parameters and physiologically  
5 based pharmacokinetic (PBPK) modeling to explore the risk assessment of  
6 trichloroethylene. *Environ Toxicol Pharmacol*, 11, 259 - 271.
- 7 Hobara, T.; Kobayashi, H.; Higashihara, E.; Kawamoto, T.; Sakai, T. (1984). Acute effects of  
8 1,1,1-trichloroethane, trichloroethylene, and toluene on the hematologic parameters in  
9 dogs. *Arch Environ Contam Toxicol*, 13, 589-593.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/6486885>
- 11 Hodgen, G. D. and Sherins, R. J. (1973). Enzymes as markers of testicular growth and  
12 development in the rat. *Endocrinology*, 93, 985-989.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/4728225>
- 14 Hoel, D. G. and Walburg, H. E. (1972). Statistical analysis of survival experiments. *J Natl*  
15 *Cancer Inst*, 49, 361-372. <http://www.ncbi.nlm.nih.gov/pubmed/5076821>
- 16 Hoffman, S.; Mishima, N.; Krug, E. L. (2004). An improved model for evaluating  
17 trichloroethylene and its metabolites as cardiac specific teratogens. In D. Jollow (Ed.),  
18 *Trichloroethylene: the scientific basis of risk assessment: The Medical University of*  
19 *South Carolina Press*.
- 20 Hoivik, D. J.; Qualls, C. W.; Mirabile, R. C.; Cariello, N. F.; Kimbrough, C. L.; Colton, H. M., . .  
21 . Miller, R. T. (2004). Fibrates induce hepatic peroxisome and mitochondrial proliferation  
22 without overt evidence of cellular proliferation and oxidative stress in cynomolgus  
23 monkeys. *Carcinogenesis*, 25, 1757-1769. <http://dx.doi.org/10.1093/carcin/bgh182>
- 24 Hooper, N. M.; Hesp, R. J.; Tiekou, S. (1994). Metabolism of aspartame by human and pig  
25 intestinal microvillar peptidases. *Biochem J*, 298 Pt 3, 635-639.  
26 <http://www.ncbi.nlm.nih.gov/pubmed/8141778>
- 27 Hoover, R. N. and Strasser, P. H. (1980). Artificial sweeteners and human bladder cancer.  
28 Preliminary results. *Lancet*, 1, 837-840. <http://www.ncbi.nlm.nih.gov/pubmed/6103207>
- 29 Horesovsky, G.; Ginsler, J.; Everitt, J.; Walker, C. (1994). Increased susceptibility to in vitro  
30 transformation of cells carrying the Eker tumor susceptibility mutation. *Carcinogenesis*,  
31 15, 2183-2187. <http://www.ncbi.nlm.nih.gov/pubmed/7955051>
- 32 Hornung, R. W.; Pinney, S. M.; Lodwick, J.; Killough, G. G.; Brewer, D. E.; Nasuta, J. (2008).  
33 Estimation of radon exposures to workers at the Fernald Feed Materials Production  
34 Center 1952-1988. *J Expo Sci Environ Epidemiol*, 18, 512-523.  
35 <http://dx.doi.org/10.1038/sj.jes.7500645>
- 36 Hortelano, S.; Dewez, B.; Genaro, A. M.; Díaz-Guerra, M. J.; Boscá, L. (1995). Nitric oxide is  
37 released in regenerating liver after partial hepatectomy. *Hepatology*, 21, 776-786.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/7533126>
- 39 Horvath, A. L.; Getzen, F. W.; Maczynska, Z. (1999). IUPAC-NIST Solubility Data Series 67.  
40 Halogenated Ethanes and Ethenes with Water. *J Phys Chem Ref Data*, 28, 395.  
41 <http://dx.doi.org/10.1063/1.556039>
- 42 Howard, P. H.; Boethling, R. S.; Jarvis, W. F.; Meylan, W. M.; Michalenko, E. M. (1991).  
43 *Handbook of environmental degradation rates*. Chelsea, MI: Lewis Publishers.
- 44 Howroyd, P.; Swanson, C.; Dunn, C.; Cattley, R. C.; Corton, J. C. (2004). Decreased longevity  
45 and enhancement of age-dependent lesions in mice lacking the nuclear receptor

DRAFT– DO NOT QUOTE OR CITE

- 1 peroxisome proliferator-activated receptor alpha (PPARalpha). *Toxicol Pathol*, 32, 591-  
2 599. <http://www.ncbi.nlm.nih.gov/pubmed/15603543>
- 3 Hrelia, P.; Maffei, F.; Vigagni, F.; Fimognari, C.; Flori, P.; Stanzani, R.; Forti, G. C. (1994).  
4 Interactive effects between trichloroethylene and pesticides at metabolic and genetic level  
5 in mice. *Environ Health Perspect*, 102, 31-34.
- 6 HSDB. (Hazardous Substances Data Bank). (1996). Trichloroethylene. Bethesda, MD: National  
7 Library of Medicine. Retrieved from [http://toxnet.nlm.nih.gov/cgi-](http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB)  
8 [bin/sis/htmlgen?HSDB](http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB)
- 9 HSDB. (Hazardous Substances Data Bank). (2002). Trichloroethylene. Bethesda, MD: National  
10 Library of Medicine. Retrieved from [http://toxnet.nlm.nih.gov/cgi-](http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB)  
11 [bin/sis/htmlgen?HSDB](http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB)
- 12 Hsia, C. Y.; Huo, T. I.; Chiang, S. Y.; Lu, M. F.; Sun, C. L.; Wu, J. C., . . . Lee, S. D. (2007).  
13 Evaluation of interleukin-6, interleukin-10 and human hepatocyte growth factor as tumor  
14 markers for hepatocellular carcinoma. *Eur J Surg Oncol*, 33, 208-212.  
15 <http://dx.doi.org/10.1016/j.ejso.2006.10.036>
- 16 Hu, C.; Jiang, L.; Geng, C.; Zhang, X.; Cao, J.; Zhong, L. (2008). Possible involvement of  
17 oxidative stress in trichloroethylene-induced genotoxicity in human HepG2 cells. *Mutat*  
18 *Res*, 652, 88-94. <http://dx.doi.org/10.1016/j.mrgentox.2008.01.002>
- 19 Hu, Y.; Hakkola, J.; Oscarson, M.; Ingelman-Sundberg, M. (1999). Structural and functional  
20 characterization of the 5'-flanking region of the rat and human cytochrome P450 2E1  
21 genes: Identification of a polymorphic repeat in the human gene. *Biochem Biophys Res*  
22 *Commun*, 263, 286-293. <http://dx.doi.org/10.1006/bbrc.1999.1362>
- 23 Huang, H. L.; Li, L. Y.; Chen, B. J.; Huang, J. X.; Kuang, S. R. (2002). New problems caused by  
24 occupational trichloroethylene exposure. *Int J Immunopathol Pharmacol*, 15, 30-32.
- 25 Hunter, E.; Rogers, E.; Schmid, J.; Richard, A. (1996). Comparative effects of haloacetic acids in  
26 whole embryo culture. *Teratology*, 54, 57-64. [http://dx.doi.org/10.1002/\(SICI\)1096-](http://dx.doi.org/10.1002/(SICI)1096-9926(199606)54:2<57::AID-TERA1>3.0.CO;2-1)  
27 [9926\(199606\)54:2<57::AID-TERA1>3.0.CO;2-1](http://dx.doi.org/10.1002/(SICI)1096-9926(199606)54:2<57::AID-TERA1>3.0.CO;2-1)
- 28 IARC. (International Agency for Research on Cancer). (1995). Dry cleaning, some chlorinated  
29 solvents and other industrial chemicals: Summary of data reported and evaluation. (IARC  
30 Monographs on the Evaluation of Carcinogenic Risks to Humans, Volume 63). Lyon,  
31 France: International Agency for Research on Cancer, World Health Organization.
- 32 IARC. (International Agency for Research on Cancer). (1995). Dry cleaning, some chlorinated  
33 solvents and other industrial chemicals: views and expert opinions of an IARC Working  
34 Group on the Evaluation of Carcinogenic Risks of Humans; February; Lyon, France.  
35 Lyon, France: Author.
- 36 IARC. (International Agency for Research on Cancer). (1995). Tetrachloroethylene. Lyon,  
37 France: Author. Retrieved from  
38 <http://monographs.iarc.fr/ENG/Monographs/vol63/index.php>.
- 39 IARC. (International Agency for Research on Cancer). (2004). Tobacco smoke and involuntary  
40 smoking. Lyon, France: Author.
- 41 Iavicoli, I.; Marinaccio, A.; Carelli, G. (2005). Effects of occupational trichloroethylene  
42 exposure on cytokine levels in workers. *J Occup Environ Med*, 47, 453-457.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/15891523>
- 44 ICRP. (International Commission of Radiological Protection). (2003). Basic anatomical and  
45 physiological data for use in radiological protection: reference values. A report of age-

- 1 and gender-related differences in the anatomical and physiological characteristics of  
2 reference individuals. ICRP Publication 89. (89). New York: International Commission  
3 On Radiological Protection.
- 4 IEI. (International Epidemiology Institute). (2005). Rocketdyne worker health study: Final  
5 report: Executive summary & appendices. Rockville, MD: Author. Retrieved from  
6 [http://www.boeing.com/aboutus/environment/santa\\_susana/healthstudy.html](http://www.boeing.com/aboutus/environment/santa_susana/healthstudy.html).
- 7 Ikbal, M.; Tastekin, A.; Dogan, H.; Pirim, I.; Ors, R. (2004). The assessment of genotoxic effects  
8 in lymphocyte cultures of infants treated with chloral hydrate. *Mutat Res*, 564, 159-164.  
9 <http://dx.doi.org/10.1016/j.mrgentox.2004.08.007>
- 10 Ikeda, M. (1977). Metabolism of trichloroethylene and tetrachloroethylene in human subjects.  
11 *Environ Health Perspect*, 21, 239-245. <http://www.ncbi.nlm.nih.gov/pubmed/612449>
- 12 Ikeda, M. and Imamura, T. (1973). Biological half-life of trichloroethylene and  
13 tetrachloroethylene in human subjects. *Int Arch Occup Environ Health*, 31, 209-224.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/4593976>
- 15 Ikeda, M.; Ohtsuji, H.; Imamura, T.; Komoike, Y. (1972). Urinary excretion of total trichloro-  
16 compounds, trichloroethanol, and trichloroacetic acid as a measure of exposure to  
17 trichloroethylene and tetrachloroethylene. *Br J Ind Med*, 29, 328-333.  
18 <http://dx.doi.org/10.1136/oem.29.3.328>
- 19 Ikeda, M.; Otsuji, H.; Kawai, H.; Kuniyoshi, M. (1971). Excretion kinetics of urinary metabolites  
20 in a patient addicted to trichloroethylene. *Br J Ind Med*, 28, 203-206.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/5572691>
- 22 ILSI. (International Life Sciences Institute). (1992). Similarities and differences between  
23 children and adults: Implications for risk assessment. New York: ILSI Press.
- 24 Inoue, O.; Seiji, K.; Kawai, T.; Jin, C.; Liu, Y.-T.; Chen, Z., . . . Ikeda, M. (1989). Relationship  
25 between vapor exposure and urinary metabolite excretion among workers exposed to  
26 trichloroethylene. *Am J Ind Med*, 15, 103-110.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/2784624>
- 28 IOM. (Institute of Medicine). (2003). Gulf war and health. Washington, DC: National  
29 Academies Press.
- 30 Isaacs, K.; Evans, M.; Harris, T. (2004). Visualization-based analysis for a mixed-inhibition  
31 binary PBPK model: determination of inhibition mechanism. *J Pharmacokinetic*  
32 *Pharmacodyn*, 31, 215-242. <http://www.ncbi.nlm.nih.gov/pubmed/15518245>
- 33 Isaacson, L. and Taylor, D. (1989). Maternal exposure to 1,1,2-trichloroethylene affects myelin  
34 in the hippocampal formation of the developing rat. *Brain Res*, 488, 403-407.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/2743138>
- 36 Isaacson, L. G.; Spohler, S. A.; Taylor, D. H. (1990). Trichloroethylene affects learning and  
37 decreases myelin in the rat hippocampus. *Neurotoxicol Teratol*, 12, 375-381.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/2392097>
- 39 Isaacson, P.; Bean, J. A.; Splinter, R.; Olson, D. B.; Kohler, J. (1985). Drinking water and cancer  
40 incidence in Iowa. III. Association of cancer with indices of contamination. *Am J*  
41 *Epidemiol*, 121, 856-869. <http://www.ncbi.nlm.nih.gov/pubmed/4014178>
- 42 Ito, Y.; Yamanoshita, O.; Asaeda, N.; Tagawa, Y.; Lee, C. H.; Aoyama, T., . . . Nakajima, T.  
43 (2007). Di(2-ethylhexyl)phthalate induces hepatic tumorigenesis through a peroxisome  
44 proliferator-activated receptor alpha-independent pathway. *J Occup Health*, 49, 172-182.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/17575397>

DRAFT– DO NOT QUOTE OR CITE

- 1 Iyer, R.; Frink, E.; Ebert, T.; Anders, M. (1998). Cysteine conjugate beta-lyase-dependent  
2 metabolism of compound A (2-[fluoromethoxy]-1,1,3,3,3-pentafluoro-1-propene) in  
3 human subjects anesthetized with sevoflurane and in rats given compound A.  
4 *Anesthesiology*, 88, 611-618. <http://www.ncbi.nlm.nih.gov/pubmed/9523802>
- 5 Jacob, S.; Hery, M.; Protois, J. C.; Rossert, J.; Stengel, B. (2007). New insight into solvent-  
6 related end-stage renal disease: Occupations, products and types of solvents at risk.  
7 *Occup Environ Med*, 64, 843–848. <http://dx.doi.org/10.1136/oem.2007.032482>
- 8 Jaffe, D.; Gandolfi, A.; Nagle, R. (1984). Chronic toxicity of S-(trans-1,2-dichlorovinyl)-L-  
9 cysteine in mice. *J Appl Toxicol*, 4, 315-319.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/6520320>
- 11 Jaffe, D.; Hassall, C.; Gandolfi, A.; Brendel, K. (1985). Production of DNA single strand breaks  
12 in rabbit renal tissue after exposure to 1,2-dichlorovinylcysteine. *Toxicology*, 35, 25-33.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/4002236>
- 14 Jaffe, E. S.; N.L., H.; Stein, H.; Vardiman, J. W. (2001). IARC WHO classification of tumours,  
15 no 3: Pathology and genetics of tumours of haematopoietic and lymphoid tissues. Lyon,  
16 France: International Agency for Research on Cancer. Retrieved from  
17 <http://apps.who.int/bookorders/anglais/detart1.jsp?sesslan=1&codlan=1&codcol=70&codcch=3>.
- 18
- 19 Jakobson, I.; Holmberg, B.; Ekner, A. (1986). Venous blood levels of inhaled trichloroethylene  
20 in female rats and changes induced by interacting agents. *Acta Pharmacol Toxicol*, 59,  
21 135-143. <http://www.ncbi.nlm.nih.gov/pubmed/3776551>
- 22 Jakobson, I.; Wahlberg, J. E.; Holmberg, B.; Johansson, G. (1982). Uptake via the blood and  
23 elimination of 10 organic solvents following epicutaneous exposure of anesthetized  
24 guinea pigs. *Toxicol Appl Pharmacol*, 63, 181-187. [http://dx.doi.org/10.1016/0041-](http://dx.doi.org/10.1016/0041-008X(82)90039-4)  
25 [008X\(82\)90039-4](http://dx.doi.org/10.1016/0041-008X(82)90039-4)
- 26 Jakubowski, M. and Wieczorek, H. (1988). The effects of physical effort on pulmonary uptake of  
27 selected organic compound vapours. *Pol J Occup Med*, 1, 62-71.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/2980152>
- 29 James, M.; Cornett, R.; Yan, Z.; Henderson, G.; Stacpoole, P. (1997). Glutathione-dependent  
30 conversion to glyoxylate, a major pathway of dichloroacetate biotransformation in  
31 hepatic cytosol from humans and rats, is reduced in dichloroacetate-treated rats. *Drug*  
32 *Metab Dispos*, 25, 1223-1227. <http://www.ncbi.nlm.nih.gov/pubmed/9351896>
- 33 Jansson, C.; Johansson, A. L.; Bergdahl, I. A.; Dickman, P. W.; Plato, N.; Adami, J., . . .  
34 Lagergren, J. (2005). Occupational exposures and risk of esophageal and gastric cardia  
35 cancers among male Swedish construction workers. *Cancer Causes Control*, 16, 755-764.  
36 <http://dx.doi.org/10.1007/s10552-005-1723-2>
- 37 Jansson, C.; Plato, N.; Johansson, A. L.; Nyrén, O.; Lagergren, J. (2006). Airborne occupational  
38 exposures and risk of oesophageal and cardia adenocarcinoma. *Occup Environ Med*, 63,  
39 107-112. <http://dx.doi.org/10.1136/oem.2005.022467>
- 40 Jasińska, M. (1965). [General anesthesia with trichloroethylene in children with congenital  
41 developmental defects of the jaw and face]. 18, 935-940.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/5214455>
- 43 Jaspers, R. M. A.; Muijser, H.; Lammers, J. H. C. M.; Kulig, B. M. (1993). Mid-frequency  
44 hearing loss and reduction of acoustic startle responding in rats following

- 1 trichloroethylene exposure. *Neurotoxicol Teratol*, 15, 407-412.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/8302242>
- 3 Jemal, A.; Siegel, R.; Ward, E.; Hao, Y.; Xu, J.; Murray, T.; Thun, M. J. (2008). Cancer  
4 statistics, 2008. *CA Cancer J Clin*, 58, 71-96. <http://dx.doi.org/10.3322/CA.2007.0010>
- 5 Jensen, O. M.; Knudsen, J. B.; McLaughlin, J. K.; Sorensen, B. L. (1988). The Copenhagen case-  
6 control study of renal pelvis and ureter cancer: role of smoking and occupational  
7 exposures. *Int J Cancer*, 41, 557-561. <http://www.ncbi.nlm.nih.gov/pubmed/3356489>
- 8 Ji, B. T.; Blair, A.; Shu, X. O.; Chow, W. H.; Hauptmann, M.; Dosemeci, M., . . . Zheng, W.  
9 (2008). Occupation and breast cancer risk among Shanghai women in a population-based  
10 cohort study. *Am J Ind Med*, 51, 100-110. <http://dx.doi.org/10.1002/ajim.20507>
- 11 Ji, J. and Hemminki, K. (2005). Variation in the risk for liver and gallbladder cancers in  
12 socioeconomic and occupational groups in Sweden with etiological implications. *Int*  
13 *Arch Occup Environ Health*, 78, 641-649. <http://dx.doi.org/10.1007/s00420-005-0015-1>
- 14 Jia, H. L.; Ye, Q. H.; Qin, L. X.; Budhu, A.; Forgues, M.; Chen, Y., . . . Wang, X. W. (2007).  
15 Gene expression profiling reveals potential biomarkers of human hepatocellular  
16 carcinoma. *Clin Cancer Res*, 13, 1133-1139. <http://dx.doi.org/10.1158/1078-0432.CCR-06-1025>
- 17  
18 Jia, M.; Coats, B.; Chadha, M.; Frentzen, B.; Perez-Rodriguez, J.; Chadik, P., . . . Stacpoole, P.  
19 (2006). Human kinetics of orally and intravenously administered low-dose 1,2-(13)C-  
20 dichloroacetate. *J Clin Pharmacol*, 46, 1449-1459.  
21 <http://dx.doi.org/10.1177/0091270006292627>
- 22 Jirtle, R. L.; Meyer, S. A.; Brockenbrough, J. S. (1991). Liver tumor promoter phenobarbital: A  
23 biphasic modulator of hepatocyte proliferation. *Prog Clin Biol Res*, 369, 209-216.
- 24 Johnson, A. D.; VanDemark, W. L.; Gomes, W. R. (1967). The effect of antispermatogenic and  
25 hormone treatments on testicular cholesterol. *Fed Proc*, 26, 645.
- 26 Johnson, P. D.; Dawson, B. V.; Goldberg, S. J. (1998). Cardiac teratogenicity of  
27 trichloroethylene metabolites. *J Am Coll Cardiol*, 32, 540-545.  
28 [http://dx.doi.org/10.1016/S0735-1097\(98\)00232-0](http://dx.doi.org/10.1016/S0735-1097(98)00232-0)
- 29 Johnson, P. D.; Dawson, B. V.; Goldberg, S. J.; Mays, M. Z. (2004). Trichloroethylene: Johnson  
30 et al.'s Response. *Environ Health Perspect*, 112, a608-a609.  
31 <http://dx.doi.org/10.1289/ehp.112-a608>
- 32 Johnson, P.; Dawson, B.; Goldberg, S. (1998). A review: trichloroethylene metabolites: potential  
33 cardiac teratogens. *Environ Health Perspect*, 106 Suppl 4, 995-999.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/9703484>
- 35 Johnson, P.; Goldberg, S.; Mays, M.; Dawson, B. (2003). Threshold of trichloroethylene  
36 contamination in maternal drinking waters affecting fetal heart development in the rat.  
37 *Environ Health Perspect*, 111, 289-292. <http://www.ncbi.nlm.nih.gov/pubmed/12611656>
- 38 Johnsrud, E. K.; Koukouritaki, S. B.; Divakaran, K.; Brunengraber, L. L.; Hines, R. N.; p  
39 McCarver, D. G. (2003). Human hepatic CYP2E1 expression during development. *J*  
40 *Pharmacol Exp Ther*, 307, 402-407. <http://dx.doi.org/10.1124/jpet.102.053124>
- 41 Jones, T.; Qin, C.; Schaeffer, V.; Stevens, J. (1988). Immunohistochemical localization of  
42 glutamine transaminase K, a rat kidney cysteine conjugate beta-lyase, and the  
43 relationship to the segment specificity of cysteine conjugate nephrotoxicity. *Mol*  
44 *Pharmacol*, 34, 621-627. <http://www.ncbi.nlm.nih.gov/pubmed/3193955>

DRAFT– DO NOT QUOTE OR CITE

- 1 Jorquera, R. and Tanguay, R. (2001). Fumarylacetoacetate, the metabolite accumulating in  
2 hereditary tyrosinemia, activates the ERK pathway and induces mitotic abnormalities and  
3 genomic instability. *Hum Mol Genet*, 10, 1741-1752.  
4 <http://www.ncbi.nlm.nih.gov/pubmed/11532983>
- 5 Juntunen, J.; Antti-Poika, M.; Tola, S.; Partanen, T. (1982). Clinical prognosis of patients with  
6 diagnosed chronic solvent intoxication. *Acta Neurol Scand*, 65, 488-503.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/7113659>
- 8 Juntunen, J.; Hupli, V.; Hernberg, S.; Luisto, M. (1980). Neurological picture of organic solvent  
9 poisoning in industry: a retrospective clinical study of 37 patients. *Int Arch Occup*  
10 *Environ Health*, 46, 219-231. <http://www.ncbi.nlm.nih.gov/pubmed/7450887>
- 11 Juran, B. D. and Lazaridis, K. N. (2006). Genomics and complex liver disease: Challenges and  
12 opportunities. *Hepatology*, 44, 1380-1390. <http://dx.doi.org/10.1002/hep.21453>
- 13 Kadry, A.; Turkall, R.; Skowronski, G.; Abdel-Rahman, M. (1991). Soil adsorption alters  
14 kinetics and bioavailability of trichloroethylene in orally exposed female rats. *Toxicol*  
15 *Lett*, 58, 337-346. <http://www.ncbi.nlm.nih.gov/pubmed/1957329>
- 16 Kafer, E. (1986). Tests which distinguish induced crossing-over and aneuploidy from secondary  
17 segregation in *Aspergillus* treated with chloral hydrate or gamma-rays. *Mutat Res*, 164,  
18 145-166. [http://dx.doi.org/10.1016/0165-1161\(86\)90006-3](http://dx.doi.org/10.1016/0165-1161(86)90006-3)
- 19 Kamijima, M.; Hisanaga, N.; Wang, H.; Nakajima, T. (2007). Occupational trichloroethylene  
20 exposure as a cause of idiosyncratic generalized skin disorders and accompanying  
21 hepatitis similar to drug hypersensitivities. *Int Arch Occup Environ Health*, 80, 357-370.  
22 <http://dx.doi.org/10.1007/s00420-006-0147-y>
- 23 Kamijima, M.; Wang, H.; Huang, H.; Li, L.; Shibata, E.; Lin, B., . . . Nakajima, T. (2008).  
24 Trichloroethylene causes generalized hypersensitivity skin disorders complicated by  
25 hepatitis. *J Occup Health*, 50, 328-338. <http://www.ncbi.nlm.nih.gov/pubmed/18540116>
- 26 Kamijo, Y.; Hora, K.; Nakajima, T.; Kono, K.; Takahashi, K.; Ito, Y., . . . Aoyama, T. (2007).  
27 Peroxisome proliferator-activated receptor alpha protects against glomerulonephritis  
28 induced by long-term exposure to the plasticizer di-(2-ethylhexyl)phthalate. *J Am Soc*  
29 *Nephrol*, 18, 176-188. <http://dx.doi.org/10.1681/asn.2006060597>
- 30 Kan, F.; Forkert, P.; Wade, M. (2007). Trichloroethylene exposure elicits damage in epididymal  
31 epithelium and spermatozoa in mice. *Histol Histopathol*, 22, 977-988.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/17523075>
- 33 Kaneko, T.; Saegusa, M.; Tasaka, K.; Sato, A. (2000). Immunotoxicity of trichloroethylene: a  
34 study with MRL-lpr/lpr mice. *J Appl Toxicol*, 20, 471-475.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/11180269>
- 36 Kaneko, T.; Wang, P.-Y.; Sato, A. (1994). Enzymes induced by ethanol differently affect the  
37 pharmacokinetics of trichloroethylene and 1,1,1-trichloroethane. *Occup Environ Med*, 51,  
38 113-119. <http://www.ncbi.nlm.nih.gov/pubmed/8111458>
- 39 Kaneko, T.; Wang, P.-Y.; Sato, A. (1994). Partition coefficients of some acetate esters and  
40 alcohols in water, blood, olive oil, and rat tissues. *Occup Environ Med*, 51, 68-72.
- 41 Kanisawa, M. and Suzuki, S. (1978). Induction of renal and hepatic tumors in mice by  
42 ochratoxin A, a mycotoxin. *Gann*, 69, 599-600.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/710809>



- 1 Kappas, A. (1989). On the mechanisms of induced aneuploidy in *Aspergillus nidulans* and  
2 validation of tests for genomic mutations. *Prog Clin Biol Res*, 318, 377-384.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/2696976>
- 4 Karami, S.; Boffetta, P.; Brennan, P.; Stewart, P. A.; Zaridze, D.; Matveev, V., . . . Moore, L. E.  
5 (2011). Renal cancer risk and occupational exposure to polycyclic aromatic hydrocarbons  
6 and plastics. *J Occup Environ Med*, 53, 218-223.  
7 <http://dx.doi.org/10.1097/JOM.0b013e31820a40a3>
- 8 Kargalioglu, Y.; McMillan, B. J.; Minear, R. A.; Plewa, M. J. (2002). Analysis of the  
9 cytotoxicity and mutagenicity of drinking water disinfection by-products in *Salmonella*  
10 *typhimurium*. *Teratog Carcinog Mutagen*, 22, 113-128.  
11 <http://dx.doi.org/10.1002/tcm.10010>
- 12 Karpinets, T. V. and Foy, B. D. (2005). Tumorigenesis: the adaptation of mammalian cells to  
13 sustained stress environment by epigenetic alterations and succeeding matched mutations.  
14 *Carcinogenesis*, 26, 1323-1334. <http://dx.doi.org/10.1093/carcin/bgi079>
- 15 Karunanayake, C. P.; McDuffie, H. H.; Dosman, J. A.; Spinelli, J. J.; Pahwa, P. (2008).  
16 Occupational exposures and non-Hodgkin's lymphoma: Canadian case-control study.  
17 *Environ Health*, 7, 44. <http://dx.doi.org/10.1186/1476-069X-7-44>
- 18 Kato-Weinstein, J.; Lingohr, M. K.; Orner, G. A.; Thrall, B. D.; Bull, R. J. (1998). Effects of  
19 dichloroacetate on glycogen metabolism in B6C3F1 mice. *Toxicology*, 130, 141-154.  
20 [http://dx.doi.org/10.1016/S0300-483X\(98\)00106-1](http://dx.doi.org/10.1016/S0300-483X(98)00106-1)
- 21 Kato-Weinstein, J.; Stauber, A. J.; Orner, G. A.; Thrall, B. D.; Bull, R. J. (2001). Differential  
22 effects of dihalogenated and trihalogenated acetates in the liver of B6C3F1 mice. *J Appl*  
23 *Toxicol*, 21, 81-89. <http://dx.doi.org/10.1002/jat.717>
- 24 Kauffmann, B.; White, K.; Sanders, V.; Douglas, K.; Sain, L.; Borzelleca, J.; Munson, A. (1982).  
25 Humoral and cell-mediated immune status in mice exposed to chloral hydrate. *Environ*  
26 *Health Perspect*, 44, 147-151. <http://www.ncbi.nlm.nih.gov/pubmed/6979478>
- 27 Kautiainen, A.; Vogel, J.; Turteltaub, K. (1997). Dose-dependent binding of trichloroethylene to  
28 hepatic DNA and protein at low doses in mice. *Chem Biol Interact*, 106, 109-121.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/9366897>
- 30 Kawamoto, T.; Hobara, T.; Aoki, T.; Kobayashi, H.; Iwamoto, S.; Sakai, T., . . . Ogino, K.  
31 (1988). Metabolism of chloral hydrate in the anoxic perfused liver. *Toxicol Lett*, 40, 225-  
32 231. <http://www.ncbi.nlm.nih.gov/pubmed/3354006>
- 33 Kawamoto, T.; Hobara, T.; Nakamura, K.; Imamura, A.; Ogino, K.; Kobayashi, H., . . . Sakai, T.  
34 (1988). Induction of cytochrome P-450, cytochrome b-5, NADPH-cytochrome c  
35 reductase and change of cytochrome P-450 isozymes with long-term trichloroethylene  
36 treatment. *Toxicology*, 53, 239-249. <http://www.ncbi.nlm.nih.gov/pubmed/3145630>
- 37 Keech, A.; Simes, J.; Barter, P.; Best, J.; Scott, R.; Taskinen, M. R.; Committee, F. M. (2006).  
38 Correction to the FIELD study report. *Lancet*, 368, 1415.  
39 [http://dx.doi.org/10.1016/S0140-6736\(06\)69594-9](http://dx.doi.org/10.1016/S0140-6736(06)69594-9)
- 40 Keech, A.; Simes, R. J.; Barter, P.; Best, J.; Scott, R.; Taskinen, M. R., . . . investigators, F. s.  
41 (2005). Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people  
42 with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet*, 366,  
43 1849-1861. [http://dx.doi.org/10.1016/S0140-6736\(05\)67667-2](http://dx.doi.org/10.1016/S0140-6736(05)67667-2)
- 44 Keil, D. E.; Peden-Adams, M. M.; Wallace, S.; Ruiz, P.; Gilkeson, G. S. (2009). Assessment of  
45 trichloroethylene (TCE) exposure in murine strains genetically-prone and non-prone to

- 1 develop autoimmune disease. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 44,  
2 443-453. <http://dx.doi.org/10.1080/10934520902719738>
- 3 Keller, D. A. and Hd'A, H. (1988). Mechanistic studies on chloral toxicity: Relationship to  
4 trichloroethylene carcinogenesis. *Toxicol Lett*, 42, 183-191.  
5 [http://dx.doi.org/10.1016/0378-4274\(88\)90076-8](http://dx.doi.org/10.1016/0378-4274(88)90076-8)
- 6 Kelsh, M. A.; Alexander, D. D.; Mink, P. J.; Mandel, J. H. (2010). Occupational  
7 trichloroethylene exposure and kidney cancer: a meta-analysis. *Epidemiology*, 21, 95-  
8 102. <http://dx.doi.org/10.1097/EDE.0b013e3181c30e92>
- 9 Kenck, C.; Wilhelm, M.; Bugert, P.; Staehler, G.; Kovacs, G. (1996). Mutation of the VHL gene  
10 is associated exclusively with the development of non-papillary renal cell carcinomas. *J*  
11 *Pathol*, 179, 157-161. [http://dx.doi.org/10.1002/\(SICI\)1096-  
12 9896\(199606\)179:2<157::AID-PATH557>3.0.CO;2-S](http://dx.doi.org/10.1002/(SICI)1096-9896(199606)179:2<157::AID-PATH557>3.0.CO;2-S)
- 13 Kernan, G. J.; Ji, B. T.; Dosemeci, M.; Silverman, D. T.; Balbus, J.; Zahm, S. H. (1999).  
14 Occupational risk factors for pancreatic cancer: A case-control study based on death  
15 certificates from 24 U.S. States. *Am J Ind Med*, 36, 260-270.  
16 [http://dx.doi.org/10.1002/\(SICI\)1097-0274\(199908\)36:2<260::AID-AJIM5>3.0.CO;2-P](http://dx.doi.org/10.1002/(SICI)1097-0274(199908)36:2<260::AID-AJIM5>3.0.CO;2-P)
- 17 Keshava, N. and Caldwell, J. (2006). Key issues in the role of peroxisome proliferator-activated  
18 receptor agonism and cell signaling in trichloroethylene toxicity. *Environ Health*  
19 *Perspect*, 114, 1464-1470. <http://www.ncbi.nlm.nih.gov/pubmed/16966106>
- 20 Keshava, N.; Chiu, W. A.; Caldwell, J. C. (2007). PPARalpha and TCE: Keshava et al. respond.  
21 *Environ Health Perspect*, 115, A15-A16. <http://dx.doi.org/10.1289/ehp.115-a15>
- 22 Kessler, W.; Numtip, W.; Grote, K.; Csanady, G. A.; Chahoud, I.; Filser, J. G. (2004). Blood  
23 burden of di(2-ethylhexyl) phthalate and its primary metabolite mono(2-ethylhexyl)  
24 phthalate in pregnant and nonpregnant rats and marmosets. *Toxicol Appl Pharmacol*, 195,  
25 142-153. <http://dx.doi.org/10.1016/j.taap.2003.11.014>
- 26 Ketcha, M. M.; Stevens, D. K.; Warren, D. A.; Bishop, C. T.; Brashear, W. T. (1996).  
27 Conversion of trichloroacetic acid to dichloroacetic acid in biological samples. *J Anal*  
28 *Toxicol*, 20, 236-241. <http://www.ncbi.nlm.nih.gov/pubmed/8835661>
- 29 Keys, D. A.; Schultz, I. R.; Mahle, D. A.; Fisher, J. W. (2004). A quantitative description of  
30 suicide inhibition of dichloroacetic acid in rats and mice. *Toxicol Sci*, 82, 381-393.  
31 <http://dx.doi.org/10.1093/toxsci/kfh276>
- 32 Keys, D.; Bruckner, J.; Muralidhara, S.; Fisher, J. (2003). Tissue dosimetry expansion and cross-  
33 validation of rat and mouse physiologically based pharmacokinetic models for  
34 trichloroethylene. *Toxicol Sci*, 76, 35-50. <http://dx.doi.org/10.1093/toxsci/kfg212>
- 35 Kezic, S.; Monster, A.; Krüse, J.; Verberk, M. (2000). Skin absorption of some vaporous  
36 solvents in volunteers. *Int Arch Occup Environ Health*, 73, 415-422.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/11007346>
- 38 Kezic, S.; Monster, A.; van de Gevel, I.; Krüse, J.; Opdam, J.; Verberk, M. (2001). Dermal  
39 absorption of neat liquid solvents on brief exposures in volunteers. 62, 12-18.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/11258863>
- 41 Khan, M.; Kaphalia, B.; Prabhakar, B.; Kanz, M.; Ansari, G. (1995). Trichloroethene-induced  
42 autoimmune response in female MRL +/+ mice. *Toxicol Appl Pharmacol*, 134, 155-160.  
43 <http://dx.doi.org/10.1006/taap.1995.1179>
- 44 Khan, M.; Wu, X.; Ansari, G. (2001). Anti-malondialdehyde antibodies in MRL+/+ mice treated  
45 with trichloroethene and dichloroacetyl chloride: possible role of lipid peroxidation in

- 1 autoimmunity. *Toxicol Appl Pharmacol*, 170, 88-92.  
2 <http://dx.doi.org/10.1006/taap.2000.9086>
- 3 Khandwala, H. M.; McCutcheon, I. E.; Flyvbjerg, A.; Friend, K. E. (2000). The effects of  
4 insulin-like growth factors on tumorigenesis and neoplastic growth. *Endocr Rev*, 21, 215-  
5 244. <http://www.ncbi.nlm.nih.gov/pubmed/10857553>
- 6 Kilburn, K. (2002). Do duration of exposure, proximity to electronic manufacturing plants, and  
7 involvement in a lawsuit affect chlorinated solvent toxicity. *Arch Environ Health*, 57,  
8 121-126. <http://www.ncbi.nlm.nih.gov/pubmed/12194156>
- 9 Kilburn, K. (2002). Is neurotoxicity associated with environmental trichloroethylene (TCE).  
10 *Arch Environ Health*, 57, 113-120. <http://www.ncbi.nlm.nih.gov/pubmed/12194155>
- 11 Kilburn, K. and Thornton, J. (1996). Prediction equations for simple and visual two-choice  
12 reactions times in environmental neurotoxicology. *Arch Environ Health*, 51, 439-444.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/9012322>
- 14 Kilburn, K. H. and Warshaw, R. H. (1992). Neurobehavioral effects of formaldehyde and  
15 solvents on histology technicians: repeated testing across time. *Environ Res*, 58, 134-146.  
16 [http://dx.doi.org/10.1016/S0013-9351\(05\)80210-5](http://dx.doi.org/10.1016/S0013-9351(05)80210-5)
- 17 Kilburn, K. H. and Warshaw, R. H. (1992). Prevalence of symptoms of systemic lupus  
18 erythematosus (SLE) and of fluorescent antinuclear antibodies associated with chronic  
19 exposure to trichloroethylene and other chemicals in well water. *Environ Res*, 57, 1-9.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/1740091>
- 21 Kilburn, K. H. and Warshaw, R. H. (1993). Effects on neurobehavioral performance of chronic  
22 exposure to chemically contaminated well water. *Toxicol Ind Health*, 9, 391-404.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/8367882>
- 24 Kilburn, K. H. and Warshaw, R. H. (1993). Neurobehavioral testing of subjects exposed  
25 residentially to groundwater contaminated from an aluminum die-casting plant and local  
26 referents. *J Toxicol Environ Health*, 39, 483-496.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/8345533>
- 28 Kilburn, K. H.; Warshaw, R.; Thornton, J. C. (1987). Formaldehyde impairs memory,  
29 equilibrium, and dexterity in histology technicians: Effects which persist for days after  
30 exposure. *Arch Environ Occup Health*, 42, 117-120.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/3579365>
- 32 Kim, C. F.; Jackson, E. L.; Woolfenden, A. E.; Lawrence, S.; Babar, I.; Vogel, S., . . . Jacks, T.  
33 (2005). Identification of bronchioalveolar stem cells in normal lung and lung cancer.  
34 *Cell*, 121, 823-835. <http://dx.doi.org/10.1016/j.cell.2005.03.032>
- 35 Kim, D. and Ghanayem, B. (2006). Comparative metabolism and disposition of trichloroethylene  
36 in Cyp2e1<sup>-/-</sup> and wild-type mice. *Drug Metab Dispos*, 34, 2020-2027.  
37 <http://dx.doi.org/10.1124/dmd.106.010538>
- 38 Kim, H. J.; Bruckner, J. V.; Dallas, C. E.; Gallo, J. M. (1990). Effect of dosing vehicles on the  
39 pharmacokinetics of orally administered carbon tetrachloride in rats. *Toxicol Appl*  
40 *Pharmacol*, 102, 50-60.
- 41 Kim, H. J.; Odend'hal, S.; Bruckner, J. V. (1990). Effect of oral dosing vehicles on the acute  
42 hepatotoxicity of carbon tetrachloride in rats. *Toxicol Appl Pharmacol*, 102, 34-49.
- 43 Kim, S.; Kim, D.; Pollack, G. M.; Collins, L. B.; Rusyn, I. (2009). Pharmacokinetic analysis of  
44 trichloroethylene metabolism in male B6C3F1 mice: Formation and disposition of  
45 trichloroacetic acid, dichloroacetic acid, S-(1,2-dichlorovinyl)glutathione and S-(1,2-

DRAFT– DO NOT QUOTE OR CITE

- 1 dichlorovinyl)-L-cysteine. *Toxicol Appl Pharmacol*, 238, 90-99.  
2 <http://dx.doi.org/10.1016/j.taap.2009.04.019>
- 3 Kim, S.; Kupke, K.; Ierardi-Curto, L.; Holme, E.; Greter, J.; Tanguay, R., . . . Levy, H. (2000).  
4 Hepatocellular carcinoma despite long-term survival in chronic tyrosinaemia I. *J Inherit*  
5 *Metab Dis*, 23, 791-804. <http://www.ncbi.nlm.nih.gov/pubmed/11196105>
- 6 Kimmerle, G. and Eben, A. (1973). Metabolism, excretion and toxicology of trichloroethylene  
7 after inhalation: 1 experimental exposure on rats. *Arch Toxicol*, 30, 115-126.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/4685263>
- 9 Kimmerle, G. and Eben, A. (1973). Metabolism, excretion and toxicology of trichloroethylene  
10 after inhalation: 2 experimental human exposure. *Arch Toxicol*, 30, 127-138.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/4685264>
- 12 King, G. S.; Smialek, J. E.; Troutman, W. G. (1985). Sudden death in adolescents resulting from  
13 the inhalation of typewriter correction fluid. *JAMA*, 253, 1604-1606.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/3974043>
- 15 Kioski, C.; Heubach, S.; Petersen, N. (1990). The incidence of childhood leukemia and testicular  
16 cancer in Pima County: 1970-1986.
- 17 Kioski, C.; Petersen, N.; Heubach, H.; Chapin, C. (1990). Cancer incidence and mortality in an  
18 East Phoenix area overlaying groundwater contaminated with volatile organic  
19 compounds.
- 20 Kiran, S.; Cocco, P.; Mannetje, A.; Satta, G.; D'Andrea, I.; Becker, N., . . . Boffetta, P. (2010).  
21 Occupational exposure to ethylene oxide and risk of lymphoma. *Epidemiology*, 21, 905-  
22 910. <http://dx.doi.org/10.1097/EDE.0b013e3181f4cc0f>
- 23 Kishi, R.; Harabuchi, I.; Ikeda, T.; Katakura, Y.; Miyake, H. (1993). Acute effects of  
24 trichloroethylene on blood concentrations and performance decrements in rats and their  
25 relevance to humans. *Occup Environ Med*, 50, 470-480.  
26 <http://www.ncbi.nlm.nih.gov/pubmed/8507600>
- 27 Kitada, T.; Seki, S.; Kawakita, N.; Kuroki, T.; Monna, T. (1995). Telomere shortening in chronic  
28 liver diseases. *Biochem Biophys Res Commun*, 211, 33-39.  
29 <http://dx.doi.org/10.1006/bbrc.1995.1774>
- 30 Kjellstrand, P.; Bjerkemo, M.; Mortensen, I.; Mansson, L. (1981). Effects of long-term exposure  
31 to trichloroethylene on the behavior of Mongolian gerbils (*Meriones unguiculatus*). *J*  
32 *Toxicol Environ Health*, 8, 787-793.
- 33 Kjellstrand, P.; Holmquist, B.; Alm, P.; Kanje, M.; Romare, S.; Jonsson, I., . . . Bjerkemo, M.  
34 (1983). Trichloroethylene: further studies of the effects on body and organ weights and  
35 plasma butyrylcholinesterase activity in mice. *Basic Clin Pharmacol Toxicol*, 53, 375-  
36 384. <http://www.ncbi.nlm.nih.gov/pubmed/6659967>
- 37 Kjellstrand, P.; Holmquist, B.; Mandahl, N.; Bjerkemo, M. (1983). Effects of continuous  
38 trichloroethylene inhalation on different strains of mice. *Basic Clin Pharmacol Toxicol*,  
39 53, 369-374. <http://www.ncbi.nlm.nih.gov/pubmed/6659966>
- 40 Kjellstrand, P.; Kanje, M.; Bjerkemo, M. (1987). Regeneration of the sciatic nerve in mice and  
41 rats exposed to trichloroethylene. *Toxicol Lett*, 38, 187-191.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/3629629>
- 43 Kjellstrand, P.; Kanje, M.; Mansson, L.; Bjerkemo, M.; Mortensen, I.; Lanke, J.; Holmquist, B.  
44 (1981). Trichloroethylene: effects on body and organ weights in mice, rats and gerbils.  
45 *Toxicology*, 21, 105-115. <http://www.ncbi.nlm.nih.gov/pubmed/7281199>

DRAFT– DO NOT QUOTE OR CITE

- 1 Kjellstrand, P.; Lanke, J.; Bjerkemo, M.; Zetterqvist, L.; Mansson, L. (1980). Irreversible effects  
2 of trichloroethylene exposure on the central nervous system. *Scand J Work Environ*  
3 *Health*, 6, 40-47. <http://www.ncbi.nlm.nih.gov/pubmed/7384766>
- 4 Klaunig, J. E.; Babich, M. A.; Baetcke, K. P.; Cook, J. C.; Corton, J. C.; David, R. M., . . .  
5 Fenner-Crisp, P. A. (2003). PPARalpha agonist-induced rodent tumors: Modes of action  
6 and human relevance. *Crit Rev Toxicol*, 33, 655-780.  
7 <http://dx.doi.org/10.1080/713608372>
- 8 Kligerman, A. D.; Bryant, M. F.; Doerr, C. L.; Erexson, G. L.; Evansky, P. A.; Kwanyuen, P.;  
9 McGee, J. K. (1994). Inhalation studies of the genotoxicity of trichloroethylene to  
10 rodents. *DNA Repair*, 322, 87-96.
- 11 Kobayashi, M.; Ikeda, K.; Hosaka, T.; Sezaki, H.; Someya, T.; Akuta, N., . . . Kumada, H.  
12 (2006). Dysplastic nodules frequently develop into hepatocellular carcinoma in patients  
13 with chronic viral hepatitis and cirrhosis. *Cancer*, 106, 636-647.  
14 <http://dx.doi.org/10.1002/cncr.21607>
- 15 Koch, R.; Schlegelmilch, R.; Wolf, H. U. (1988). Genetic effects of chlorinated ethylenes in the  
16 yeast *Saccharomyces cerevisiae*. *Mutat Res*, 206, 209-216.  
17 [http://dx.doi.org/10.1016/0165-1218\(88\)90162-0](http://dx.doi.org/10.1016/0165-1218(88)90162-0)
- 18 Kochen, W.; Kohlmüller, D.; De Biasi, P.; Ramsay, R. (2003). The endogeneous formation of  
19 highly chlorinated tetrahydro-beta-carbolines as a possible causative mechanism in  
20 idiopathic Parkinson's disease. *Adv Exp Med Biol*, 527, 253-263.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/15206739>
- 22 Koizumi, A. (1989). Potential of physiologically based pharmacokinetics to amalgamate kinetic  
23 data of trichloroethylene and tetrachloroethylene obtained in rats and man. *Occup*  
24 *Environ Med*, 46, 239-249. <http://www.ncbi.nlm.nih.gov/pubmed/2713280>
- 25 Konietzko, H.; Elster, I.; Bencsath, A.; Drysch, K.; Weichardt, H. (1975). [EEG changes under  
26 standardized trichloroethylene load (author's transl)]. *Int Arch Occup Environ Health*, 35,  
27 257-264. <http://www.ncbi.nlm.nih.gov/pubmed/1176181>
- 28 Kopylev, L.; Chen, C.; White, P. (2007). Towards quantitative uncertainty assessment for cancer  
29 risks: central estimates and probability distributions of risk in dose-response modeling.  
30 *Regul Toxicol Pharmacol*, 49, 203-207.
- 31 Korrapati, M.; Lock, E.; Mehendale, H. (2005). Molecular mechanisms of enhanced renal cell  
32 division in protection against S-1,2-dichlorovinyl-L-cysteine-induced acute renal failure  
33 and death. *Am J Physiol Renal Physiol*, 289, F175-F185.  
34 <http://dx.doi.org/10.1152/ajprenal.00418.2004>
- 35 Kozak, E. M. and Tate, S. S. (1982). Glutathione-degrading enzymes of microvillus membranes.  
36 *J Biol Chem*, 257, 6322-6327. <http://www.ncbi.nlm.nih.gov/pubmed/6122685>
- 37 Krasowski, M. and Harrison, N. (2000). The actions of ether, alcohol and alkane general  
38 anaesthetics on GABAA and glycine receptors and the effects of TM2 and TM3  
39 mutations. *Br J Pharmacol*, 129, 731-743. <http://dx.doi.org/10.1038/sj.bjp.0703087>
- 40 Kraupp-Grasl, B.; Huber, W.; Putz, B.; Gerbracht, U.; Schulte-Hermann, R. (1990). Tumor  
41 promotion by the peroxisome proliferator nafenopin involving a specific subtype of  
42 altered foci in rat liver. *Cancer Res*, 50, 3701-3708.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/1971196>
- 44 Krause, R. J.; Lash, L. H.; Elfarra, A. A. (2003). Human kidney flavin-containing  
45 monooxygenases and their potential roles in cysteine s-conjugate metabolism and

- 1 nephrotoxicity. *J Pharmacol Exp Ther*, 304, 185-191.  
2 <http://dx.doi.org/10.1124/jpet.102.042911>
- 3 Krejci, M.; Ridgewell, R.; Koechel, D. (1991). Acute effects of the D-isomer of S-(1,2-  
4 dichlorovinyl)cysteine on renal function and ultrastructure in the pentobarbital-  
5 anesthetized dog: site-specific toxicity involving the S1 and S2 cells of the proximal  
6 tubule. *Toxicology*, 69, 151-164. <http://www.ncbi.nlm.nih.gov/pubmed/1949044>
- 7 Krishnadasan, A.; Kennedy, N.; Zhao, Y.; Morgenstern, H.; Ritz, B. (2007). Nested case-control  
8 study of occupational chemical exposures and prostate cancer in aerospace and radiation  
9 workers. *Am J Ind Med*, 50, 383-390. <http://dx.doi.org/10.1002/ajim.20458>
- 10 Krishnadasan, A.; Kennedy, N.; Zhao, Y.; Morgenstern, H.; Ritz, B. (2008). Nested case-control  
11 study of occupational physical activity and prostate cancer among workers using a job  
12 exposure matrix. *Cancer Causes Control*, 19, 107-114. [http://dx.doi.org/10.1007/s10552-](http://dx.doi.org/10.1007/s10552-007-9076-7)  
13 [007-9076-7](http://dx.doi.org/10.1007/s10552-007-9076-7)
- 14 Kroneld, R. (1989). Volatile pollutants in the environment and human tissues. *Bull Environ*  
15 *Contam Toxicol*, 42, 873-877. <http://www.ncbi.nlm.nih.gov/pubmed/2743020>
- 16 Kubo, Y.; Mitani, H.; Hino, O. (1994). Allelic loss at the predisposing gene locus in spontaneous  
17 and chemically induced renal cell carcinomas in the Eker rat. *Cancer Res*, 54, 2633-2635.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/8168090>
- 19 Kudryavtsev, B. N.; Kudryavtseva, M. V.; Sakuta, G. A.; Stein, G. I. (1993). Human hepatocyte  
20 polyploidization kinetics in the course of life cycle. *Virchows Arch B Cell Pathol Incl*  
21 *Mol Pathol*, 64, 387-393. <http://www.ncbi.nlm.nih.gov/pubmed/8148960>
- 22 Kulig, B. M. (1987). The effects of chronic trichloroethylene exposure on neurobehavioral  
23 functioning in the rat. *Neurotoxicol Teratol*, 9, 171-178.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/3657753>
- 25 Kuller, L. H.; Ockene, J. K.; Meilahn, E.; Wentworth, D. N.; Svendsen, K. H.; Neaton, J. D.  
26 (1991). Cigarette smoking and mortality. MRFIT Research Group. *Prev Med*, 20, 638-  
27 654. <http://www.ncbi.nlm.nih.gov/pubmed/1758843>
- 28 Kumar, M.; Kumar, R.; Hissar, S. S.; Saraswat, M. K.; Sharma, B. C.; Sakhuja, P.; Sarin, S. K.  
29 (2007). Risk factors analysis for hepatocellular carcinoma in patients with and without  
30 cirrhosis: a case-control study of 213 hepatocellular carcinoma patients from India. *J*  
31 *Gastroenterol Hepatol*, 22, 1104-1111. [http://dx.doi.org/10.1111/j.1440-](http://dx.doi.org/10.1111/j.1440-1746.2007.04908.x)  
32 [1746.2007.04908.x](http://dx.doi.org/10.1111/j.1440-1746.2007.04908.x)
- 33 Kumar, M.; Tewari, S.; Sharma, P.; Verma, V. K.; Chauhan, L. K.; Agarwal, S. K., . . . Goel, S.  
34 K. (2009). Study of genetic polymorphism in solvent exposed population and its  
35 correlation to in vitro effect of trichloroethylene on lymphocytes. *J Environ Biol*, 30, 685-  
36 691. <http://www.ncbi.nlm.nih.gov/pubmed/20136049>
- 37 Kumar, P.; Prasad, A. K.; Dutta, K. K. (2000). Steroidogenic alterations in testes and sera of rats  
38 exposed to trichloroethylene (TCE) by inhalation. *Hum Exp Toxicol*, 19, 117-121.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/10773841>
- 40 Kumar, P.; Prasad, A. K.; Saxena, D. K.; Manu, U.; Maji, B. K.; Dutta, K. K. (2000). Fertility  
41 and general reproduction studies in trichloroethylene exposed rats. 43, 117-126.
- 42 Kumar, P.; Prasad, A.; Maji, B.; Mani, U.; Dutta, K. (2001). Hepatotoxic alterations induced by  
43 inhalation of trichloroethylene (TCE) in rats. *Biomed Environ Sci*, 14, 325-332.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/11862613>

DRAFT– DO NOT QUOTE OR CITE

- 1 Kumar, P.; Prasad, A.; Mani, U.; Maji, B.; Dutta, K. (2001). Trichloroethylene induced testicular  
2 toxicity in rats exposed by inhalation. *Hum Exp Toxicol*, 20, 585-589.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/11926613>
- 4 Kung, H.-C.; Hoyert, D. L.; Xu, J.; Murphy, S. L. (2008). Deaths: Final data for 2005, *National*  
5 *Vital Statistics Reports*. Retrieved from  
6 [http://www.cdc.gov/nchs/data/nvsr/nvsr56/nvsr56\\_10.pdf](http://www.cdc.gov/nchs/data/nvsr/nvsr56/nvsr56_10.pdf)
- 7 Kurasawa, K. (1988). Selective damage of pulmonary nonciliated bronchiolar epithelial (Clara)  
8 cells by trichloroethylene in rats. *Sangyo Eiseigaku Zasshi*, 30, 121-133.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/3418956>
- 10 Kvam, B. M.; Romundstad, P. R.; Boffetta, P.; Andersen, A. (2005). Cancer in the Norwegian  
11 printing industry. *Scand J Work Environ Health*, 31, 36-43.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/15751617>
- 13 Kylin, B.; Axell, K.; Samuel, H. E.; Lindborg, A. (1967). Effect of inhaled trichloroethylene on  
14 the CNS: as measured by optokinetic nystagmus. *Arch Environ Occup Health*, 15, 48-52.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/4951693>
- 16 Kyrklund, T. and Haglid, K. (1991). Brain lipid composition in guinea pigs after intrauterine  
17 exposure to perchloroethylene. *Pharmacol Toxicol*, 68, 146-148.  
18 <http://dx.doi.org/10.1111/j.1600-0773.1991.tb02054.x>
- 19 Kyrklund, T.; Kjellstrand, P.; Haglid, K. G. (1986). Fatty acid changes in rat brain ethanolamine  
20 phosphoglycerides during and following chronic exposure to trichloroethylene. *Toxicol*  
21 *Appl Pharmacol*, 85, 145-153. <http://www.ncbi.nlm.nih.gov/pubmed/3764901>
- 22 Kyrklund, T.; Kjellstrand, P.; Haglid, K. G. (1988). Effects of exposure to Freon 11, 1,1,1-  
23 trichloroethane or perchloroethylene on the lipid and fatty-acid composition of rat  
24 cerebral cortex. *Scand J Work Environ Health*, 14, 91-94.
- 25 La Vecchia, C.; Negri, E.; Franceschi, S.; D'Avanzo, B.; Boyle, P. (1994). A case-control study  
26 of diabetes mellitus and cancer risk. *Br J Cancer*, 70, 950-953.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/7947103>
- 28 Lacey, J. V. J.; Garabrant, D. H.; Laing, T. J.; Gillespie, B. W.; Mayes, M. D.; Cooper, B. C.;  
29 Schottenfeld, D. (1999). Petroleum distillate solvents as risk factors for undifferentiated  
30 connective tissue disease (UCTD). *Am J Epidemiol*, 149, 761-770.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/10206626>
- 32 Lagakos, S. W.; Wessen, B. J.; Zelen, M. (1986). An analysis of contaminated well water and  
33 health effects in Woburn, Massachusetts. *J Am Stat Assoc*, 81, 583-596.
- 34 Laham, S. (1970). Studies on placental transfer: Trichlorethylene. *Ind Med Surg*, 39, 46-49.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/5263053>
- 36 Lake, B. G.; Evans, J. G.; Cunninghame, M. E.; Price, R. J. (1993). Comparison of the hepatic  
37 effects of nafenopin and WY-14,643 on peroxisome proliferation and cell replication in  
38 the rat and Syrian hamster. *Environ Health Perspect*, 101, 241-247.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/8013414>
- 40 Lalwani, N. D.; Dethloff, L. A.; Haskins, J. R.; Robertson, D. G.; de la Iglesia, F. A. (1997).  
41 Increased nuclear ploidy, not cell proliferation, is sustained in the peroxisome  
42 proliferator-treated rat liver. *Toxicol Pathol*, 25, 165-176.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/9125775>

DRAFT– DO NOT QUOTE OR CITE

- 1 Lamb, J. C. and Hentz, K. L. (2006). Toxicological review of male reproductive effects and  
2 trichloroethylene exposure: assessing the relevance to human male reproductive health.  
3 *Reprod Toxicol*, 22, 557-563. <http://dx.doi.org/10.1016/j.reprotox.2006.07.001>
- 4 Land, P. C.; Owen, E. L.; Linde, H. W. (1981). Morphologic changes in mouse spermatozoa  
5 after exposure to inhalational anesthetics during early spermatogenesis. *Anesthesiology*,  
6 54, 53-56. <http://www.ncbi.nlm.nih.gov/pubmed/6109470>
- 7 Landrigan, P. J.; Stein, G. F.; Kominsky, J. R.; Ruhe, R. L.; Watanabe, A. S. (1987). Common-  
8 source community and industrial exposure to trichloroethylene. *Arch Environ Occup*  
9 *Health*, 42, 327-332. <http://www.ncbi.nlm.nih.gov/pubmed/3439809>
- 10 Lane, S. E.; Watts, R. A.; Bentham, G.; Innes, N. J.; Scott, D. G. I. (2003). Are environmental  
11 factors important in primary systemic vasculitis? A case-control study. *Arthritis Rheum*,  
12 48, 814-823. <http://dx.doi.org/10.1002/art.10830>
- 13 Langholz, B. and Goldstein, L. (1996). <http://dx.doi.org/10.1214/ss/1032209663>
- 14 Laparé, S.; Tardif, R.; Brodeur, J. (1995). Effect of various exposure scenarios on the biological  
15 monitoring of organic solvents in alveolar air. II. 1,1,1-Trichloroethane and  
16 trichloroethylene. *Int Arch Occup Environ Health*, 67, 375-394.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/8567088>
- 18 Lapis, K.; Zalatnai, A.; Timár, F.; Thorgeirsson, U. P. (1995). Quantitative evaluation of  
19 lysozyme- and CD68-positive Kupffer cells in diethylnitrosamine-induced hepatocellular  
20 carcinomas in monkeys. *Carcinogenesis*, 16, 3083-3085.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/8603489>
- 22 Larsen, G. L. and Stevens, J. L. (1986). Cysteine conjugate beta-lyase in the gastrointestinal  
23 bacterium *Eubacterium limosum*. *Mol Pharmacol*, 29, 97-103.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/3945231>
- 25 Larson, J. and Bull, R. (1992). Species differences in the metabolism of trichloroethylene to the  
26 carcinogenic metabolites trichloroacetate and dichloroacetate. *Toxicol Appl Pharmacol*,  
27 115, 278-285. <http://www.ncbi.nlm.nih.gov/pubmed/1641861>
- 28 Larson, J. L. and Bull, R. J. (1989). Effect of ethanol on the metabolism of trichloroethylene. *J*  
29 *Toxicol Environ Health*, 28, 395-406. <http://dx.doi.org/10.1080/15287398909531359>
- 30 Larson, J. L. and Bull, R. J. (1992). Metabolism and lipoperoxidative activity of trichloroacetate  
31 and dichloroacetate in rats and mice. *Toxicol Appl Pharmacol*, 115, 268-277.  
32 [http://dx.doi.org/10.1016/0041-008X\(92\)90332-M](http://dx.doi.org/10.1016/0041-008X(92)90332-M)
- 33 Lash, L. H.; Fisher, J. W.; Lipscomb, J. C.; Parker, J. C. (2000). Metabolism of  
34 trichloroethylene. *Environ Health Perspect*, 108, 177-200.
- 35 Lash, L. H.; Jones, D. P.; Anders, M. W. (1988). Glutathione homeostasis and glutathione S-  
36 conjugate toxicity in kidney. *Rev Biochem Toxicol*, 9, 29-67.
- 37 Lash, L. H.; Lipscomb, J. C.; Putt, D. A.; Parker, J. C. (1999). Glutathione conjugation of  
38 trichloroethylene in human liver and kidney: Kinetics and individual variation. *Drug*  
39 *Metab Dispos*, 27, 351-359. <http://www.ncbi.nlm.nih.gov/pubmed/10064565>
- 40 Lash, L. H.; Nelson, R. M.; Van Dyke, R. A.; Anders, M. W. (1990). Purification and  
41 characterization of human kidney cytosolic cysteine conjugate beta-lyase activity. *Drug*  
42 *Metab Dispos*, 18, 50-54. <http://www.ncbi.nlm.nih.gov/pubmed/2139845>
- 43 Lash, L. H.; Parker, J. C.; Scott, C. S. (2000). Modes of action of trichloroethylene for kidney  
44 tumorigenesis. *Environ Health Perspect*, 108, 225-240.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/10807554>



DRAFT– DO NOT QUOTE OR CITE

- 1 Lash, L. H.; Putt, D. A.; Huang, P.; Hueni, S. E.; Parker, J. C. (2007). Modulation of hepatic and  
2 renal metabolism and toxicity of trichloroethylene and perchloroethylene by alterations in  
3 status of cytochrome P450 and glutathione. *Toxicology*, 235, 11-26.  
4 <http://dx.doi.org/10.1016/j.tox.2007.03.001>
- 5 Lash, L. H.; Qian, W.; Putt, D. A.; Hueni, S. E.; Elfarra, A. A.; Krause, R. J.; Parker, J. C.  
6 (2001). Renal and hepatic toxicity of trichloroethylene and its glutathione-derived  
7 metabolites in rats and mice: sex-, species-, and tissue-dependent differences. *J*  
8 *Pharmacol Exp Ther*, 297, 155-164. <http://www.ncbi.nlm.nih.gov/pubmed/11259540>
- 9 Lash, L. H.; Qian, W.; Putt, D. A.; Jacobs, K.; Elfarra, A. A.; Krause, R. J.; Parker, J. C. (1998).  
10 Glutathione conjugation of trichloroethylene in rats and mice: Sex-, species-, and tissue-  
11 dependent differences. *Drug Metab Dispos*, 26, 12-19.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/9443846>
- 13 Lash, L. H.; Visarius, T. M.; Sall, J. M.; Qian, W.; Tokarz, J. J. (1998). Cellular and subcellular  
14 heterogeneity of glutathione metabolism and transport in rat kidney cells. *Toxicology*,  
15 130, 1-15. <http://www.ncbi.nlm.nih.gov/pubmed/9846992>
- 16 Lash, L. H.; Xu, Y.; Elfarra, A. A.; Duescher, R. J.; Parker, J. C. (1995). Glutathione-dependent  
17 metabolism of trichloroethylene in isolated liver and kidney cells of rats and its role in  
18 mitochondrial and cellular toxicity. *Drug Metab Dispos*, 23, 846-853.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/7493552>
- 20 Lash, L.; Elfarra, A.; Anders, M. (1986). Renal cysteine conjugate beta-lyase. Bioactivation of  
21 nephrotoxic cysteine S-conjugates in mitochondrial outer membrane. *J Biol Chem*, 261,  
22 5930-5935. <http://www.ncbi.nlm.nih.gov/pubmed/3700378>
- 23 Lash, L.; Elfarra, A.; Rakiewicz-Nemeth, D.; Anders, M. (1990). Bioactivation mechanism of  
24 cytotoxic homocysteine S-conjugates. *Arch Biochem Biophys*, 276, 322-330.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/2154944>
- 26 Lash, L.; Hueni, S.; Putt, D. (2001). Apoptosis, necrosis, and cell proliferation induced by S-  
27 (1,2-dichlorovinyl)-L-cysteine in primary cultures of human proximal tubular cells.  
28 *Toxicol Appl Pharmacol*, 177, 1-16. <http://dx.doi.org/10.1006/taap.2001.9295>
- 29 Lash, L.; Putt, D.; Brashear, W.; Abbas, R.; Parker, J.; Fisher, J. (1999). Identification of S-(1,2-  
30 dichlorovinyl)glutathione in the blood of human volunteers exposed to trichloroethylene.  
31 *J Toxicol Environ Health A*, 56, 1-21. <http://www.ncbi.nlm.nih.gov/pubmed/9923751>
- 32 Lash, L.; Putt, D.; Hueni, S.; Krause, R.; Elfarra, A. (2003). Roles of necrosis, Apoptosis, and  
33 mitochondrial dysfunction in S-(1,2-dichlorovinyl)-L-cysteine sulfoxide-induced  
34 cytotoxicity in primary cultures of human renal proximal tubular cells. *J Pharmacol Exp*  
35 *Ther*, 305, 1163-1172. <http://dx.doi.org/10.1124/jpet.102.046185>
- 36 Lash, L.; Putt, D.; Parker, J. (2006). Metabolism and tissue distribution of orally administered  
37 trichloroethylene in male and female rats: identification of glutathione- and cytochrome  
38 P-450-derived metabolites in liver, kidney, blood, and urine. *J Toxicol Environ Health A*,  
39 69, 1285-1309. <http://dx.doi.org/10.1080/15287390500360133>
- 40 Lash, L.; Sausen, P.; Duescher, R.; Cooley, A.; Elfarra, A. (1994). Roles of cysteine conjugate  
41 beta-lyase and S-oxidase in nephrotoxicity: studies with S-(1,2-dichlorovinyl)-L-cysteine  
42 and S-(1,2-dichlorovinyl)-L-cysteine sulfoxide. *J Pharmacol Exp Ther*, 269, 374-383.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/8169843>
- 44 Lash, T. L. and Green, L. C. (1993). Blink reflex measurement of effects of trichloroethylene  
45 exposure on the trigeminal nerve [letter]. *Muscle Nerve*, 16, 217-220.

DRAFT– DO NOT QUOTE OR CITE

- 1 Laslo-Baker, D.; Barrera, M.; Knittel-Keren, D.; Kozer, E.; Wolpin, J.; Khattak, S., . . . Koren,  
2 G. (2004). Child neurodevelopmental outcome and maternal occupational exposure to  
3 solvents. *Arch Pediatr Adolesc Med*, 158, 956-961.  
4 <http://dx.doi.org/10.1001/archpedi.158.10.956>
- 5 Latendresse, J. R. and Pereira, M. A. (1997). Dissimilar characteristics of n-methyl-n-  
6 nitrosourea-initiated foci and tumors promoted by dichloroacetic acid or trichloroacetic  
7 acid in the liver of female B6C3F1 mice. *Toxicol Pathol*, 25, 433-440.  
8 <http://dx.doi.org/10.1177/019262339702500501>
- 9 Lau, S. S.; Kleiner, H. E.; Monks, T. J. (1995). Metabolism as a determinant of species  
10 susceptibility to 2,3,5-(triglutathion-S-yl)hydroquinone-mediated nephrotoxicity. The  
11 role of N-acetylation and N-deacetylation. *Drug Metab Dispos*, 23, 1136-1142.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/8654203>
- 13 Laughter, A. R.; Dunn, C. S.; Swanson, C. L.; Howroyd, P.; Cattley, R. C.; Corton, J. C. (2004).  
14 Role of the peroxisome proliferator-activated receptor alpha (PPARalpha) in responses to  
15 trichloroethylene and metabolites, trichloroacetate and dichloroacetate in mouse liver.  
16 *Toxicology*, 203, 83-98. <http://dx.doi.org/10.1016/j.tox.2004.06.014>
- 17 Laureno, R. (1993). Trichloroethylene does not cause trigeminal neuropathy [letter]. *Muscle*  
18 *Nerve*, 16, 217.
- 19 Lawrence, T.; Hageman, T.; Balkwill, F. (2007). Cancer. Sex, cytokines, and cancer. *Science*,  
20 317, 51-52. <http://dx.doi.org/10.1126/science.1146052>
- 21 Le Mesurier, S. M.; Lykke, A. W. J.; Stewart, B. W. (1980). Reduced yield of pulmonary  
22 surfactant: patterns of response following administration of chemicals to rats by  
23 inhalation. *Toxicol Lett*, 5, 89-93. <http://www.ncbi.nlm.nih.gov/pubmed/6892858>
- 24 Leakey, J. E.; Seng, J. E.; Allaben, W. T. (2003). Body weight considerations in the B6C3F1  
25 mouse and the use of dietary control to standardize background tumor incidence in  
26 chronic bioassays. *Toxicol Appl Pharmacol*, 193, 237-265.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/14644626>
- 28 Leakey, J.; Seng, J.; Latendresse, J.; Hussain, N.; Allen, L.; Allaben, W. (2003). Dietary  
29 controlled carcinogenicity study of chloral hydrate in male B6C3F1 mice. *Toxicol Appl*  
30 *Pharmacol*, 193, 266-280. <http://www.ncbi.nlm.nih.gov/pubmed/14644627>
- 31 Leandri, M.; Schizzi, R.; Scielzo, C.; Favale, E. (1995). Electrophysiological evidence of  
32 trigeminal root damage after trichloroethylene exposure. *Muscle Nerve*, 18, 467-468.  
33 <http://dx.doi.org/10.1002/mus.880180416>
- 34 Leavitt, S. A.; DeAngelo, A. B.; George, M. H.; Ross, J. A. (1997). Assessment of the  
35 mutagenicity of dichloroacetic acid in lacI transgenic B6C3F1 mouse liver.  
36 *Carcinogenesis*, 18, 2101-2106. <http://www.ncbi.nlm.nih.gov/pubmed/9395208>
- 37 Ledda-Columbano, G. M.; Columbano, A.; Curto, M.; Ennas, M. G.; Coni, P.; Sarma, D. S.;  
38 Pani, P. (1989). Further evidence that mitogen-induced cell proliferation does not support  
39 the formation of enzyme-altered islands in rat liver by carcinogens. *Carcinogenesis*, 10,  
40 847-850. <http://www.ncbi.nlm.nih.gov/pubmed/2565171>
- 41 Ledda-Columbano, G. M.; Coni, P.; Simbula, G.; Zedda, I.; Columbano, A. (1993).  
42 Compensatory regeneration, mitogen-induced liver growth, and multistage chemical  
43 carcinogenesis. *Environ Health Perspect*, 101 Suppl 5, 163-168.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/8013404>

DRAFT– DO NOT QUOTE OR CITE

- 1 Ledda-Columbano, G. M.; Curto, M.; Piga, R.; Zedda, A. I.; Menegazzi, M.; Sartori, C., . . .  
2 Columbano, A. (1998). In vivo hepatocyte proliferation is inducible through a TNF and  
3 IL-6-independent pathway. *Oncogene*, 17, 1039-1044.  
4 <http://dx.doi.org/10.1038/sj.onc.1202018>
- 5 Ledda-Columbano, G. M.; Perra, A.; Concas, D.; Cossu, C.; Molotzu, F.; Sartori, C., . . .  
6 Columbano, A. (2003). Different effects of the liver mitogens triiodo-thyronine and  
7 ciprofibrate on the development of rat hepatocellular carcinoma. *Toxicol Pathol*, 31, 113-  
8 120. <http://www.ncbi.nlm.nih.gov/pubmed/12597455>
- 9 Lee, D.; Park, I.; Kim, J.; Lee, Y.; Kim, D.; Kang, S. (1998). Neurobehavioral changes in shoe  
10 manufacturing workers. *Neurotoxicol Teratol*, 20, 259-263.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/9638683>
- 12 Lee, J. S.; Chu, I. S.; Heo, J.; Calvisi, D. F.; Sun, Z.; Roskams, T., . . . Thorgeirsson, S. S.  
13 (2004). Classification and prediction of survival in hepatocellular carcinoma by gene  
14 expression profiling. *Hepatology*, 40, 667-676. <http://dx.doi.org/10.1002/hep.20375>
- 15 Lee, J. S.; Chu, I. S.; Mikaelyan, A.; Calvisi, D. F.; Heo, J.; Reddy, J. K.; Thorgeirsson, S. S.  
16 (2004). Application of comparative functional genomics to identify best-fit mouse  
17 models to study human cancer. *Nat Genet*, 36, 1306-1311.  
18 <http://dx.doi.org/10.1038/ng1481>
- 19 Lee, K.; Bruckner, J.; Muralidhara, S.; Gallo, J. (1996). Characterization of presystemic  
20 elimination of trichloroethylene and its nonlinear kinetics in rats. *Toxicol Appl*  
21 *Pharmacol*, 139, 262-271. <http://dx.doi.org/10.1006/taap.1996.0165>
- 22 Lee, K.; Muralidhara, S.; Schnellmann, R.; Bruckner, J. (2000). Contribution of direct solvent  
23 injury to the dose-dependent kinetics of trichloroethylene: portal vein administration to  
24 rats. *Toxicol Appl Pharmacol*, 164, 46-54. <http://dx.doi.org/10.1006/taap.2000.8891>
- 25 Lee, K.; Muralidhara, S.; White, C.; Bruckner, J. (2000). Mechanisms of the dose-dependent  
26 kinetics of trichloroethylene: oral bolus dosing of rats. *Toxicol Appl Pharmacol*, 164, 55-  
27 64. <http://dx.doi.org/10.1006/taap.2000.8892>
- 28 Lee, L. J.-H.; Chan, C.-C.; Chung, C.-W.; Ma, Y.-C.; Wang, G.-S.; Wang, J.-D. (2002). Health  
29 risk assessment on residents exposed to chlorinated hydrocarbons contaminated in  
30 groundwater of a hazardous waste site. *J Toxicol Environ Health A*, 65, 219-235.  
31 <http://dx.doi.org/10.1080/15287390252800828>
- 32 Lee, L.; Chung, C.; Ma, Y.; Wang, G.; Chen, P.; Hwang, Y.; Wang, J. (2003). Increased  
33 mortality odds ratio of male liver cancer in a community contaminated by chlorinated  
34 hydrocarbons in groundwater. *Occup Environ Med*, 60, 364-369.  
35 <http://dx.doi.org/10.1136/oem.60.5.364>
- 36 Lehmann, I.; Rehwagen, M.; Diez, U.; Seiffart, A.; Rolle-Kampczyk, U.; Richter, M., . . .  
37 Herbarth, O. (2001). Enhanced in vivo IgE production and T cell polarization toward the  
38 type 2 phenotype in association with indoor exposure to VOC: Results of the LARS  
39 study. *Int J Hyg Environ Health*, 204, 211-221. <http://dx.doi.org/10.1078/1438-4639-00100>
- 40  
41 Lehmann, I.; Thoenke, A.; Rehwagen, M.; Rolle-Kampczyk, U.; Schlink, U.; Schulz, R., . . .  
42 Herbarth, O. (2002). The influence of maternal exposure to volatile organic compounds  
43 on the cytokine secretion profile of neonatal T cells. *Environ Toxicol*, 17, 203-210.  
44 <http://dx.doi.org/10.1002/tox.10055>

DRAFT– DO NOT QUOTE OR CITE

- 1 Lehnart, S. E.; Mongillo, M.; Bellinger, A.; Lindegger, N.; Bi-Xing, C.; Hsueh, W., . . . Marks,  
2 A. R. (2008). Leaky Ca<sup>2+</sup> release channel/ryanodine receptor 2 causes  
3 seizures and sudden cardiac death in mice. *J Clin Invest*, 118, 2230-2245.
- 4 Leigh, J. P. and Jiang, W. Y. (1993). Liver cirrhosis deaths within occupations and industries in  
5 the California occupational mortality study. *Addiction*, 88, 767-779.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/8329968>
- 7 Lemke, L. E. and McQueen, C. A. (1995). Acetylation and its role in the mutagenicity of the  
8 antihypertensive agent hydralazine. *Drug Metab Dispos*, 23, 559-565.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/7587931>
- 10 Leopardi, P.; Zijno, A.; Bassani, B.; Pacchierotti, F. (1993). In vivo studies on chemically  
11 induced aneuploidy in mouse somatic and germinal cells. *Mutat Res*, 287, 119-130.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/7683378>
- 13 Leuschner, J. and Beuscher, N. (1998). Studies on the mutagenic and carcinogenic potential of  
14 chloral hydrate. *Arzneimittelforschung*, 48, 961-968.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/9825112>
- 16 Leuschner, J. and Leuschner, F. (1991). Evaluation of the mutagenicity of chloral hydrate in  
17 vitro and in vivo. *Arzneimittelforschung*, 41, 1101-1103.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/1799393>
- 19 Levy, D.; Lipton, R.; Holzman, P. (1981). Smooth pursuit eye movements: effects of alcohol and  
20 chloral hydrate. *J Psychiatr Res*, 16, 1-11. <http://www.ncbi.nlm.nih.gov/pubmed/7205696>
- 21 Lewis, G. D.; Reynolds, R. C.; Johnson, A. R. (1984). Some effects of trichloroethylene on  
22 mouse lungs and livers. *Vascul Pharmacol*, 15, 139-144.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/6425111>
- 24 Lewis, R. J., Sr. (Ed.). (2001). *Hawley's Condensed Chemical Dictionary* (14 ed.). New York,  
25 NY: John Wiley & Sons, Inc.
- 26 Lewitt, M.; Brismar, K.; Wang, J.; Wivall-Helleryd, I.; Sindelar, P.; Gonzalez, F., . . . Bobek, G.  
27 (2001). Responses of insulin-like growth factor (IGF)-I and IGF-binding proteins to  
28 nutritional status in peroxisome proliferator-activated receptor-alpha knockout mice.  
29 *Growth Horm IGF Res*, 11, 303-313. <http://dx.doi.org/10.1054/ghir.2001.0247>
- 30 Li, H.; Dai, Y.; Huang, H.; Li, L.; Leng, S.; Cheng, J., . . . Zheng, Y. (2007). HLA-B\*1301 as a  
31 biomarker for genetic susceptibility to hypersensitivity dermatitis induced by  
32 trichloroethylene among workers in China. *Environ Health Perspect*, 115, 1553-1556.  
33 <http://dx.doi.org/10.1289/ehp.10325>
- 34 Li, T.; Schultz, I.; Keys, D. A.; Campbell, J. L.; Fisher, J. W. (2008). Quantitative evaluation of  
35 dichloroacetic acid kinetics in human--a physiologically based pharmacokinetic modeling  
36 investigation. *Toxicology*, 245, 35-48. <http://dx.doi.org/10.1016/j.tox.2007.12.010>
- 37 Liang, J. C. and Brinkley, B. R. (1985). Chemical probes and possible targets for the induction of  
38 aneuploidy. *Basic Life Sci*, 36, 491-505. <http://www.ncbi.nlm.nih.gov/pubmed/4096703>
- 39 Liang, J. C. and Pacchierotti, F. (1988). Cytogenetic investigation of chemically-induced  
40 aneuploidy in mouse spermatocytes. *Mutat Res-Fundam Mol Mech Mutagen*, 201, 325-  
41 335. [http://dx.doi.org/10.1016/0027-5107\(88\)90021-8](http://dx.doi.org/10.1016/0027-5107(88)90021-8)
- 42 Liangpunsakul, S.; Kolwankar, D.; Pinto, A.; Gorski, J.; Hall, S.; Chalasani, N. (2005). Activity  
43 of CYP2E1 and CYP3A enzymes in adults with moderate alcohol consumption: a  
44 comparison with nonalcoholics. *Hepatology*, 41, 1144-1150.  
45 <http://dx.doi.org/10.1002/hep.20673>

- 1 Libbrecht, L. (2006). Hepatic progenitor cells in human liver tumor development. *World J*  
2 *Gastroenterol*, 12, 6261-6265. <http://www.ncbi.nlm.nih.gov/pubmed/17072946>
- 3 Lide, D. R. (Ed.). (1998). *handbook of chemistry and physics (79th ed.)*. Boca Raton, FL: CRC  
4 Press.
- 5 Lieber, C. (2004). The discovery of the microsomal ethanol oxidizing system and its physiologic  
6 and pathologic role. *Drug Metab Rev*, 36, 511-529. [http://dx.doi.org/10.1081/DMR-](http://dx.doi.org/10.1081/DMR-200033441)  
7 200033441
- 8 Ligocki, M. P.; Leuenberger, C.; Pankow, J. F. (1985). Trace organic-compounds in rain: 2. Gas  
9 scavenging of neutral organic-compounds. *Atmos Environ*, 19, 1609-1617.
- 10 Lilly, P. D.; Simmons, J. E.; Pegram, R. A. (1994). Dose-dependent vehicle differences in the  
11 acute toxicity of bromodichloromethane. *Fundam Appl Toxicol*, 23, 132-140.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/7958557>
- 13 Lin, L. W.; Lin, X. Y.; He, Y. M.; Gao, S. D.; Lin, X. D. (2003). Biological characteristics of  
14 HCC by ultrasound-guided aspiration biopsy and its clinical application. *World J*  
15 *Gastroenterol*, 9, 941-945. <http://www.ncbi.nlm.nih.gov/pubmed/12717834>
- 16 Lin, Y. Z.; Brunt, E. M.; Bowling, W.; Hafenrichter, D. G.; Kennedy, S. C.; Flye, M. W.;  
17 Ponder, K. P. (1995). Ras-transduced diethylnitrosamine-treated hepatocytes develop into  
18 cancers of mixed phenotype in vivo. *Cancer Res*, 55, 5242-5250.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/7585583>
- 20 Lindbohm, M. L.; Sallmén, M.; Kyyrönen, P.; Kauppinen, T.; Pukkala, E. (2009). Risk of liver  
21 cancer and exposure to organic solvents and gasoline vapors among Finnish workers. *Int*  
22 *J Cancer*, 124, 2954-2959. <http://dx.doi.org/10.1002/ijc.24309>
- 23 Lindbohm, M. L.; Taskinen, H.; Sallmen, M.; Hemminki, K. (1990). Spontaneous abortions  
24 among women exposed to organic solvents. *Am J Ind Med*, 17, 449-463.  
25 <http://dx.doi.org/10.1002/ajim.4700170404>
- 26 Lindstrom, K. (1973). Psychological performances of workers exposed to various solvents.  
27 *Scand J Work Environ Health*, 10, 151-155.
- 28 Lindstrom, K. (1980). Changes in psychological performances of solvent-poisoned and solvent  
29 exposed workers. *Am J Ind Med*, 1, 69-84.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/7342756>
- 31 Lindström, K.; Antti-Poika, M.; Tola, S.; Hyytiäinen, A. (1982). Psychological prognosis of  
32 diagnosed chronic organic solvent intoxication. *Neurobehav Toxicol Teratol*, 4, 581-588.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/7177310>
- 34 Lingohr, M.; Bull, R.; Kato-Weinstein, J.; Thrall, B. (2002). Dichloroacetate stimulates glycogen  
35 accumulation in primary hepatocytes through an insulin-independent mechanism. *Toxicol*  
36 *Sci*, 68, 508-515. <http://www.ncbi.nlm.nih.gov/pubmed/12151648>
- 37 Lingohr, M.; Thrall, B.; Bull, R. (2001). Effects of dichloroacetate (DCA) on serum insulin  
38 levels and insulin-controlled signaling proteins in livers of male B6C3F1 mice. *Toxicol*  
39 *Sci*, 59, 178-184. <http://www.ncbi.nlm.nih.gov/pubmed/11134557>
- 40 Lipscomb, J. C.; Confer, P. D.; Miller, M. R.; Stamm, S. C.; Snawder, J. E.; Bandiera, S. M.  
41 (1998). Metabolism of trichloroethylene and chloral hydrate by the Japanese medaka  
42 (*Oryzias latipes*) in vitro. *Environ Toxicol Chem*, 17, 325-332.
- 43 Lipscomb, J. C.; Fisher, J. W.; Confer, P. D.; Byczkowski, J. Z. (1998). In vitro to in vivo  
44 extrapolation for trichloroethylene metabolism in humans. *Toxicol Appl Pharmacol*, 152,  
45 376-387. <http://dx.doi.org/10.1006/taap.1998.8485>

DRAFT– DO NOT QUOTE OR CITE

- 1 Lipscomb, J. C.; Garrett, C. M.; Snawder, J. E. (1997). Cytochrome P450-dependent metabolism  
2 of trichloroethylene: interindividual differences in humans. *Toxicol Appl Pharmacol*,  
3 142, 311-318. <http://dx.doi.org/10.1006/taap.1996.8040>
- 4 Lipscomb, J. C.; Garrett, C. M.; Snawder, J. E. (1998). Use of kinetic and mechanistic data in  
5 species extrapolation of bioactivation: cytochrome P-450 dependent trichloroethylene  
6 metabolism at occupationally relevant concentrations. *J Occup Health*, 40, 110-117.
- 7 Lipscomb, J. C.; Mahle, D. A.; Brashear, W. T.; Barton, H. A. (1995). Dichloroacetic acid:  
8 Metabolism in cytosol. *Drug Metab Dispos*, 23, 1202-1205.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/8591719>
- 10 Lipscomb, J. C.; Mahle, D. A.; Brashear, W. T.; Garrett, C. M. (1996). A species comparison of  
11 chloral hydrate metabolism in blood and liver. *Biochem Biophys Res Commun*, 227,  
12 340-350. <http://dx.doi.org/10.1006/bbrc.1996.1511>
- 13 Lipscomb, J. C.; Teuschler, L. K.; Swartout, J.; Popken, D.; Cox, T.; Kedderis, G. L. (2003). The  
14 impact of cytochrome P450 2E1-dependent metabolic variance on a risk-relevant  
15 pharmacokinetic outcome in humans. *Risk Anal*, 23, 1221-1238.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/14641897>
- 17 Little, J. C.; Daisey, J. M.; Nazaroff, W. W. (1992). Transport of subsurface contaminants into  
18 buildings: an exposure pathway for volatile organics. *Environ Sci Technol*, 26, 2058-  
19 2066.
- 20 Liu, E.; Jelinek, J.; Pastore, Y. D.; Guan, Y.; Prchal, J. F.; Prchal, J. T. (2003). Discrimination of  
21 polycythemias and thrombocytoses by novel, simple, accurate clonality assays and  
22 comparison with PRV-1 expression and BFU-E response to erythropoietin. *Blood*, 101,  
23 3294-3301. <http://dx.doi.org/10.1182/blood-2002-07-2287>
- 24 Liu, Y.; Bartlett, M. G.; White, C. A.; Muralidhara, S.; Bruckner, J. V. (2009). Presystemic  
25 elimination of trichloroethylene in rats following environmentally relevant oral  
26 exposures. *Drug Metab Dispos*, 37, 1994-1998.  
27 <http://dx.doi.org/10.1124/dmd.109.028100>
- 28 Liu, Y.-T.; Jin, C.; Chen, Z.; Cai, S.-X.; Yin, S.-N.; Li, G.-L., . . . Ikeda, M. (1988). Increased  
29 subjective symptom prevalence among workers exposed to trichloroethylene at sub-OEL  
30 levels. *Tohoku J Exp Med*, 155, 183-195. <http://www.ncbi.nlm.nih.gov/pubmed/3212780>
- 31 Lock, E.; Reed, C.; McMillan, J.; Oatis, J.; Schnellmann, R. (2007). Lack of formic acid  
32 production in rat hepatocytes and human renal proximal tubule cells exposed to chloral  
33 hydrate or trichloroacetic acid. *Toxicology*, 230, 234-243.  
34 <http://dx.doi.org/10.1016/j.tox.2006.11.055>
- 35 Lockey, J.; Kelly, C.; Cannon, G.; Colby, T.; Aldrich, V.; Livingston, G. (1987). Progressive  
36 systemic sclerosis associated with exposure to trichloroethylene. *J Occup Med*, 29, 493-  
37 496. <http://www.ncbi.nlm.nih.gov/pubmed/3612322>
- 38 Loeber, C. and Runyan, R. (1990). A comparison of fibronectin, laminin, and  
39 galactosyltransferase adhesion mechanisms during embryonic cardiac mesenchymal cell  
40 migration in vitro. *Dev Biol*, 140, 401-412.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/2142656>
- 42 Loeber, C.; Hendrix, M.; Diez De Pinos, S.; Goldberg, S. (1988). Trichloroethylene: a cardiac  
43 teratogen in developing chick embryos. *Pediatr Res*, 24, 740-744.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/3205631>

DRAFT– DO NOT QUOTE OR CITE

- 1 Lopreato, G.; Phelan, R.; Borghese, C.; Beckstead, M.; Mihic, S. (2003). Inhaled drugs of abuse  
2 enhance serotonin-3 receptor function. *Drug Alcohol Depend*, 70, 11-15.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/12681521>
- 4 Lorente, C.; Cordier, S.; Bergeret, A.; De Walle, H.; Goujard, J.; Aymé, S., . . . Bianchi, F.  
5 (2000). Maternal occupational risk factors for oral clefts. *Occupational Exposure and*  
6 *Congenital Malformation Working Group. Scand J Work Environ Health*, 26, 137-145.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/10817379>
- 8 Lorimore, S.; Coates, P.; Wright, E. (2003). Radiation-induced genomic instability and bystander  
9 effects: inter-related nontargeted effects of exposure to ionizing radiation. *Oncogene*, 22,  
10 7058-7069. <http://dx.doi.org/10.1038/sj.onc.1207044>
- 11 Lowengart, R. A.; Peters, J. M.; Cicioni, C.; Buckley, J.; Bernstein, L.; Preston-Martin, S.;  
12 Rappaport, E. (1987). Childhood leukemia and parents' occupational and home  
13 exposures. *J Natl Cancer Inst*, 79, 39-46. <http://www.ncbi.nlm.nih.gov/pubmed/3474448>
- 14 Luebke, R. W.; Chen, D. H.; Dietert, R.; Yang, Y.; King, M.; Luster, M. I. (2006). The  
15 comparative immunotoxicity of five selected compounds following developmental or  
16 adult exposure. *J Toxicol Environ Health B Crit Rev*, 9, 1-26.  
17 <http://dx.doi.org/10.1080/15287390500194326>
- 18 Luedde, T.; Beraza, N.; Kotsikoris, V.; van Loo, G.; Nenci, A.; De Vos, R., . . . Pasparakis, M.  
19 (2007). Deletion of NEMO/IKKgamma in liver parenchymal cells causes steatohepatitis  
20 and hepatocellular carcinoma. *Cancer Cell*, 11, 119-132.  
21 <http://dx.doi.org/10.1016/j.ccr.2006.12.016>
- 22 Lumpkin, M. H.; Bruckner, J. V.; Campbell, J. L.; Dallas, C. E.; White, C. A.; Fisher, J. W.  
23 (2003). Plasma binding of trichloroacetic acid in mice, rats, and humans under cancer  
24 bioassay and environmental exposure conditions. *Drug Metab Dispos*, 31, 1203-1207.  
25 <http://dx.doi.org/10.1124/dmd.31.10.1203>
- 26 Lundberg, I.; Alfredsson, L.; Plato, N.; Sverdrup, B.; Klareskog, L.; Kleinau, S. (1994).  
27 Occupation, occupational exposure to chemicals and rheumatological disease: A register  
28 based cohort study. *Scand J Rheumatol*, 23, 305-310.  
29 <http://dx.doi.org/10.3109/03009749409099278>
- 30 Lundgren, B. and DePierre, J. W. (1989). Proliferation of peroxisomes and induction of cytosolic  
31 and microsomal epoxide hydrolases in different strains of mice and rats after dietary  
32 treatment with clofibrate. *Xenobiotica*, 19, 867-881.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/2815829>
- 34 Lundgren, B.; Meijer, J.; DePierre, J. W. (1987). Induction of cytosolic and microsomal epoxide  
35 hydrolases and proliferation of peroxisomes and mitochondria in mouse liver after dietary  
36 exposure to p-chlorophenoxyacetic acid, 2,4-dichlorophenoxyacetic acid and 2,4,5-  
37 trichlorophenoxyacetic acid. *Biochem Pharmacol*, 36, 815-821.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/3032197>
- 39 Lutz, W. K.; Gaylor, D. W.; Conolly, R. B.; Lutz, R. W. (2005). Nonlinearity and thresholds in  
40 dose-response relationships for carcinogenicity due to sampling variation, logarithmic  
41 dose scaling, or small differences in individual susceptibility. *Toxicol Appl Pharmacol*,  
42 207, S565-S569. <http://dx.doi.org/10.1016/j.taap.2005.01.038>
- 43 Lybarger, J. A.; Lichtveld, M. Y.; Amler, R. W. (1999). Biomedical testing of the kidney for  
44 persons exposed to hazardous substances in the environment. *Ren Fail*, 21, 263-274.

DRAFT– DO NOT QUOTE OR CITE

- 1 Lynch, A. and Parry, J. (1993). The cytochalasin-B micronucleus/kinetochore assay in vitro:  
2 studies with 10 suspected aneugens. *Mutat Res*, 287, 71-86.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/7683386>
- 4 Lyngge, E.; Anttila, A.; Hemminki, K. (1997). Organic solvents and cancer. *Cancer Causes*  
5 *Control*, 8, 406-419. <http://dx.doi.org/10.1023/A:1018461406120>
- 6 Macchiarini, P. (2006). Primary tracheal tumours. *Lancet Oncol*, 7, 83-91.  
7 [http://dx.doi.org/10.1016/S1470-2045\(05\)70541-6](http://dx.doi.org/10.1016/S1470-2045(05)70541-6)
- 8 Macdonald, N.; Holden, P. R.; Roberts, R. A. (1999). Addition of peroxisome proliferator-  
9 activated receptor alpha to guinea pig hepatocytes confers increased responsiveness to  
10 peroxisome proliferators. *Cancer Res*, 59, 4776-4780.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/10519382>
- 12 Mackay, J.; Fox, V.; Griffiths, K.; Fox, D.; Howard, C.; Coutts, C., . . . Styles, J. (1995).  
13 Trichloroacetic acid: investigation into the mechanism of chromosomal damage in the in  
14 vitro human lymphocyte cytogenetic assay and the mouse bone marrow micronucleus  
15 test. *Carcinogenesis*, 16, 1127-1133. <http://www.ncbi.nlm.nih.gov/pubmed/7767975>
- 16 Maeda, S.; Kamata, H.; Luo, J. L.; Leffert, H.; Karin, M. (2005). IKKbeta couples hepatocyte  
17 death to cytokine-driven compensatory proliferation that promotes chemical  
18 hepatocarcinogenesis. *Cell*, 121, 977-990. <http://dx.doi.org/10.1016/j.cell.2005.04.014>
- 19 Magnusson, B. and Kligman, A. M. (1969). The identification of contact allergens by animal  
20 assay. The guinea pig maximization test. *J Invest Dermatol*, 52, 268-276.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/5774356>
- 22 Mahle, D. A.; Gearhart, J. M.; Grigsby, C. C.; Mattie, D. R.; Barton, H. A.; Lipscomb, J. C.;  
23 Cook, R. S. (2007). Age-dependent partition coefficients for a mixture of volatile organic  
24 solvents in Sprague-Dawley rats and humans. *J Toxicol Environ Health A*, 70, 1745-  
25 1751. <http://dx.doi.org/10.1080/15287390701458991>
- 26 Mahr, A. D.; Neogi, T.; Merkel, P. A. (2006). Epidemiology of Wegener's granulomatosis:  
27 Lessons from descriptive studies and analyses of genetic and environmental risk  
28 determinants. *Clin Exp Rheumatol*, 24, S82-S91.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/16859601>
- 30 Mailhes, J.; Aardema, M.; Marchetti, F. (1993). Investigation of aneuploidy induction in mouse  
31 oocytes following exposure to vinblastine-sulfate, pyrimethamine, diethylstilbestrol  
32 diphosphate, or chloral hydrate. *Environ Mol Mutagen*, 22, 107-114.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/8359152>
- 34 Maitre, A.; Hours, M.; Bonnetterre, V.; Arnaud, J.; Arslan, M. T.; Carpentier, P., . . . de  
35 Gaudemaris, R. (2004). Systemic sclerosis and occupational risk factors: role of solvents  
36 and cleaning products. *J Rheumatol*, 31, 2395-2401.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/15570640>
- 38 Makwana, O.; King, N. M.; Ahles, L.; Selmin, O.; Granzier, H. L.; Runyan, R. B. (2010).  
39 Exposure to low-dose trichloroethylene alters shear stress gene expression and function  
40 in the developing chick heart. *Cardiovasc Toxicol*, 10, 100-107.  
41 <http://dx.doi.org/10.1007/s12012-010-9066-y>
- 42 Malarkey, D. E.; Johnson, K.; Ryan, L.; Boorman, G.; Maronpot, R. R. (2005). New insights into  
43 functional aspects of liver morphology. *Toxicol Pathol*, 33, 27-34.  
44 <http://dx.doi.org/10.1080/01926230590881826>



DRAFT– DO NOT QUOTE OR CITE

- 1 Malherbe, P.; Alberati-Giani, D.; Kohler, C.; Cesura, A. M. (1995). Identification of a  
2 mitochondrial form of kynurenine aminotransferase/glutamine transaminase K from rat  
3 brain. *FEBS Lett*, 367, 141-144. <http://www.ncbi.nlm.nih.gov/pubmed/7796908>
- 4 Mallin, K. (1990). Investigation of a bladder cancer cluster in northwestern Illinois. *Am J*  
5 *Epidemiol*, 132, S96-106. <http://www.ncbi.nlm.nih.gov/pubmed/2356842>
- 6 Mally, A.; Walker, C.; Everitt, J.; Dekant, W.; Vamvakas, S. (2006). Analysis of renal cell  
7 transformation following exposure to trichloroethene in vivo and its metabolite S-  
8 (dichlorovinyl)-L-cysteine in vitro. *Toxicology*, 224, 108-118.  
9 <http://dx.doi.org/10.1016/j.tox.2006.04.036>
- 10 Maloney, E. K. and Waxman, D. J. (1999). trans-Activation of PPARalpha and PPARgamma by  
11 structurally diverse environmental chemicals. *Toxicol Appl Pharmacol*, 161, 209-218.  
12 <http://dx.doi.org/10.1006/taap.1999.8809>
- 13 Maltoni, C.; Lefemine, G.; Cotti, G.; Perino, G. (1988). Long-term carcinogenicity bioassays on  
14 trichloroethylene administered by inhalation to Sprague-Dawley rats and Swiss and  
15 B6C3F1 mice. *Ann N Y Acad Sci*, 534, 316-342.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/3389663>
- 17 Maltoni, C.; Lefemine, G.; G, C. (1986). Experimental research on trichloroethylene  
18 carcinogenesis. In M. M. C Maltoni (Ed.), *Archives of Research on Industrial*  
19 *Carcinogenesis* (Vol. 5, pp. ). Princeton, NJ: Princeton Scientific Publishing.
- 20 Mandel, J. H.; Kelsh, M. A.; Mink, P. J.; Alexander, D. D.; Kalmes, R. M.; Weingart, M., . . .  
21 Goodman, M. (2006). Occupational trichloroethylene exposure and non-Hodgkin's  
22 lymphoma: a meta-analysis and review. *Occup Environ Med*, 63, 597-607.  
23 <http://dx.doi.org/10.1136/oem.2005.022418>
- 24 Mandel, J. S. (2001). Renal cell cancer correlated with occupational exposure to  
25 trichloroethylene. *J Cancer Res Clin Oncol*, 127, 265-268.
- 26 Mandel, J. S.; McLaughlin, J. K.; Schlehofer, B.; Mellempgaard, A.; Helmert, U.; Lindblad, P., . . .  
27 . Adami, H.-O. (1995). International renal-cell cancer study. IV. Occupation. *Int J*  
28 *Cancer*, 61, 601-605. <http://dx.doi.org/10.1002/ijc.2910610503>
- 29 Manson, J.; Murphy, M.; Richdale, N.; Smith, M. (1984). Effects of oral exposure to  
30 trichloroethylene on female reproductive function. *Toxicology*, 32, 229-242.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/6474485>
- 32 Marano, D.; Boice, J.; Fryzek, J.; Morrison, J.; Sadler, C.; McLaughlin, J. (2000). Exposure  
33 assessment for a large epidemiological study of aircraft manufacturing workers. *Appl*  
34 *Occup Environ Hyg*, 15, 644-656. <http://dx.doi.org/10.1080/10473220050075653>
- 35 Markwald, R. R.; Runyan, R. B.; Kitten, G. T.; Funderburg, F. M.; Bernanke, D. H.; Brayer, P.  
36 R. (1984). Use of collagen gel cultures to study heart development: preteoglycan and  
37 glycoprotein interactions during formation of endocardial cushion tissue. In R. L.  
38 Trelstad (Ed.), *The role of extracellular matrix in development* (pp. 323-350): Alan R.  
39 Liss.
- 40 Markwald, R.; Eisenberg, C.; Eisenberg, L.; Trusk, T.; Sugi, Y. (1996). Epithelial-mesenchymal  
41 transformations in early avian heart development. *Acta Anat*, 156, 173-186.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/9124035>
- 43 Maronpot, R. R. (2007). *Biological basis of differential susceptibility to hepatocarcinogenesis*  
44 *among mouse strains*. Paper presented at the 3rd Workshop of the Standing Committee  
45 on Risk Analysis Issues and Reviews: Mouse liver tumors: Benefits and constraints on

- 1 use in human health risk assessment, qualitative and quantitative aspects, Washington,  
2 DC.
- 3 Maronpot, R. R.; Flake, G.; Huff, J. (2004). Relevance of animal carcinogenesis findings to  
4 human cancer predictions and prevention. *Toxicol Pathol*, 32, 40-48.
- 5 Maronpot, R. R.; Fox, T.; Malarkey, D. E.; Goldsworthy, T. L. (1995). Mutations in the ras  
6 proto-oncogene: clues to etiology and molecular pathogenesis of mouse liver tumors.  
7 *Toxicology*, 101, 125-156. [http://dx.doi.org/10.1016/0300-483X\(95\)03112-S](http://dx.doi.org/10.1016/0300-483X(95)03112-S)
- 8 Maronpot, R.; Anna, C.; Devereux, T.; Lucier, G.; Butterworth, B.; Anderson, M. (1995).  
9 Considerations concerning the murine hepatocarcinogenicity of selected chlorinated  
10 hydrocarbons. *Prog Clin Biol Res*, 391, 305-323.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/8532724>
- 12 Marrazzini, A.; Betti, C.; Bernacchi, F.; Barrai, I.; Barale, R. (1994). Micronucleus test and  
13 metaphase analyses in mice exposed to known and suspected spindle poisons.  
14 *Mutagenesis*, 9, 505-515. <http://www.ncbi.nlm.nih.gov/pubmed/7854141>
- 15 Marshall, E.; Gensburg, L.; Deres, D.; Geary, N.; Cayo, M. (1997). Maternal residential  
16 exposure to hazardous wastes and risk of central nervous system and musculoskeletal  
17 birth defects. *Arch Environ Health*, 52, 416-425.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/9541362>
- 19 Marsman, D. S.; Cattley, R. C.; Conway, J. G.; Popp, J. A. (1988). Relationship of hepatic  
20 peroxisome proliferation and replicative DNA synthesis to the hepatocarcinogenicity of  
21 the peroxisome proliferators di(2-ethylhexyl)phthalate and [4-chloro-6-(2,3-xylidino)-2-  
22 pyrimidinylthio]acetic acid (Wy-14,643) in rats. *Cancer Res*, 48, 6739-6744.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/3180084>
- 24 Marsman, D. S.; Goldsworthy, T. L.; Popp, J. A. (1992). Contrasting hepatocytic peroxisome  
25 proliferation, lipofuscin accumulation and cell turnover for the hepatocarcinogens Wy-  
26 14,643 and clofibric acid. *Carcinogenesis*, 13, 1011-1017.  
27 <http://dx.doi.org/10.1093/carcin/13.6.1011>
- 28 Martinez-Larrañaga, M. R.; Anadón, A.; Diaz, M. J.; Fernandez, R.; Sevil, B.; Fernandez-Cruz,  
29 M. L., . . . Anton, R. (1996). Induction of cytochrome P4501A1 and P4504A1 activities  
30 and peroxisomal proliferation by fumonisin B1. *Toxicol Appl Pharmacol*, 141, 185-194.  
31 <http://dx.doi.org/10.1006/taap.1996.0275>
- 32 Mather, G.; Exon, J.; Koller, L. (1990). Subchronic 90 day toxicity of dichloroacetic and  
33 trichloroacetic acid in rats. *Toxicology*, 64, 71-80.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/2219134>
- 35 Mattioli, S.; Truffelli, D.; Baldasseroni, A.; Risi, A.; Marchesini, B.; Giacomini, C., . . . Buiatti,  
36 E. (2002). Occupational risk factors for renal cell cancer: a case--control study in  
37 northern Italy. *J Occup Environ Med*, 44, 1028-1036.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/12448354>
- 39 Mazzullo, M.; Bartoli, S.; Bonora, B.; Colacci, A.; Lattanzi, G.; Niero, A., . . . Grilli, S. (1992).  
40 In vivo and in vitro interaction of trichloroethylene with macromolecules from various  
41 organs of rat and mouse. *Res Commun Mol Pathol Pharmacol*, 76, 192-208.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/1376481>
- 43 McCarthy, T. B. and Jones, R. D. (1983). Industrial gassing poisonings due to trichloroethylene,  
44 perchlorethylene, and 1,1,1-trichloroethane, 1961-80. *Occup Environ Med*, 40, 450-455.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/6626474>

- 1 McCarver, D. G. and Hines, R. N. (2002). The ontogeny of human drug-metabolizing enzymes:  
2 Phase II conjugation enzymes and regulatory mechanisms. *J Pharmacol Exp Ther*, 300,  
3 361-366. <http://dx.doi.org/10.1124/jpet.300.2.361>
- 4 McCarver, D. G.; Byun, R.; Hines, R. N.; Hichme, M.; Wegenek, W. (1998). A genetic  
5 polymorphism in the regulatory sequences of human CYP2E1: Association with  
6 increased chlorzoxazone hydroxylation in the presence of obesity and ethanol intake.  
7 *Toxicol Appl Pharmacol*, 152, 276-281. <http://dx.doi.org/10.1006/taap.1998.8532>
- 8 McConnell, G.; Ferguson, D. M.; Pearson, C. R. (1975). Chlorinated hydrocarbons and the  
9 environment. *Endeavour*, 34, 13-18. <http://www.ncbi.nlm.nih.gov/pubmed/54249>
- 10 McCredie, M. and Stewart, J. H. (1993). Risk factors for kidney cancer in New South Wales. IV.  
11 Occupation. *Br J Ind Med*, 50, 349-354. <http://www.ncbi.nlm.nih.gov/pubmed/8494775>
- 12 McCunney, R. J. (1988). Diverse manifestations of trichloroethylene. *Br J Ind Med*, 45, 122-126.  
13 <http://dx.doi.org/10.1136/oem.45.2.122>
- 14 McDaniel, T.; Martin, P.; Ross, N.; Brown, S.; Lesage, S.; Pauli, B. (2004). Effects of  
15 chlorinated solvents on four species of North American amphibians. *Arch Environ*  
16 *Contam Toxicol*, 47, 101-109. <http://www.ncbi.nlm.nih.gov/pubmed/15346783>
- 17 McEntee, G. P.; Batts, K. A.; Katzmann, J. A.; Ilstrup, D. M.; Nagorney, D. M. (1992).  
18 Relationship of nuclear DNA content to clinical and pathologic findings in patients with  
19 primary hepatic malignancy. *Surgery*, 111, 376-379.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/1313603>
- 21 McGregor, D.; Reynolds, D.; Zeiger, E. (1989). Conditions affecting the mutagenicity of  
22 trichloroethylene in *Salmonella*. *Environ Mol Mutagen*, 13, 197-202.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/2651116>
- 24 McGuire, V.; Nelson, L. M.; Koepsell, T. D.; Checkoway, H.; Longstreth, W. T.; Jr. (1998).  
25 Assessment of occupational exposures in community-based case-control studies. *Annu*  
26 *Rev Public Health*, 19, 35-53.
- 27 McKinney, P. A.; Alexander, F. E.; Cartwright, R. A.; Parker, L. (1991). Parental occupations of  
28 children with leukaemia in west Cumbria, north Humberside, and Gateshead. *Br Med J*,  
29 302, 681-687. <http://dx.doi.org/10.1136/bmj.302.6778.681>
- 30 McLaughlin, J. and Blot, W. (1997). A critical review of epidemiology studies of  
31 trichloroethylene and perchloroethylene and risk of renal-cell cancer. *Int Arch Occup*  
32 *Environ Health*, 70, 222-231. <http://www.ncbi.nlm.nih.gov/pubmed/9342621>
- 33 McLaughlin, J. K.; Hrubec, Z.; Blot, W. J.; Fraumeni, J. F. (1995). Smoking and cancer  
34 mortality among U.S. veterans: a 26-year follow-up. *Int J Cancer*, 60, 190-193.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/7829214>
- 36 McLaughlin, J. K.; Lipworth, L.; Tarone, R. E. (2006). Epidemiologic aspects of renal cell  
37 carcinoma. *Semin Oncol*, 33, 527-533.  
38 <http://dx.doi.org/10.1053/j.seminoncol.2006.06.010>
- 39 McLean, A. J.; Cogger, V. C.; Chong, G. C.; Warren, A.; Markus, A. M.; Dahlstrom, J. E.; Le  
40 Couteur, D. G. (2003). Age-related pseudocapillarization of the human liver. *J Pathol*,  
41 200, 112-117. <http://dx.doi.org/10.1002/path.1328>
- 42 McMichael, A. J.; Spirtas, R.; Gamble, J. F.; Tousey, P. M. (1976). Mortality among rubber  
43 workers: relationship to specific jobs. *J Occup Environ Med*, 18, 178-185.
- 44 McNutt, L. A. (2003). Estimating the relative risk in cohort studies and clinical trials of common  
45 outcomes. *Am J Epidemiol*, 157, 940-943. <http://dx.doi.org/10.1093/aje/kwg074>

DRAFT– DO NOT QUOTE OR CITE

- 1 MDPH. (Massachusetts Department of Public Health), (1997). Woburn childhood leukemia  
2 follow-up study: Information booklet. Boston, MA: Massachusetts Department of Public  
3 Health, Bureau of Environmental Health Assessment.
- 4 MDPH. (Massachusetts Department of Public Health). (1997). Woburn childhood leukemia  
5 follow-up study: Volume I: Analyses. Boston, MA: Massachusetts Department of Public  
6 Health, Bureau of Environmental Health Assessment. Retrieved from  
7 [http://www.mass.gov/Eeohhs2/docs/dph/environmental/investigations/woburn\\_cancer\\_leukemia\\_follow\\_up\\_study\\_1997.pdf](http://www.mass.gov/Eeohhs2/docs/dph/environmental/investigations/woburn_cancer_leukemia_follow_up_study_1997.pdf).  
8
- 9 MDPH. (Massachusetts Department of Public Health). (1997). Woburn childhood leukemia  
10 follow-up study: Volume II: Appendix. Boston, MA: Massachusetts Department of  
11 Public Health, Bureau of Environmental Health Assessment.
- 12 Mehendale, H. (2000). PPAR-alpha: a key to the mechanism of hepatoprotection by clofibrate.  
13 *Toxicol Sci*, 57, 187-190. <http://www.ncbi.nlm.nih.gov/pubmed/11006348>
- 14 Melchiorri, C.; Bolondi, L.; Chieco, P.; Pagnoni, M.; Gramantieri, L.; Barbara, L. (1994).  
15 Diagnostic and prognostic value of DNA ploidy and cell nuclearity in ultrasound-guided  
16 liver biopsies. *Cancer*, 74, 1713-1719. <http://www.ncbi.nlm.nih.gov/pubmed/8082072>
- 17 Melchiorri, C.; Chieco, P.; Zedda, A. I.; Coni, P.; Ledda-Columbano, G. M.; Columbano, A.  
18 (1993). Ploidy and nuclearity of rat hepatocytes after compensatory regeneration or  
19 mitogen-induced liver growth. *Carcinogenesis*, 14, 1825-1830.
- 20 Mellemegaard, A.; Engholm, G.; McLaughlin, J. K.; Olsen, J. H. (1994). Occupational risk factors  
21 for renal-cell carcinoma in Denmark. *Scand J Work Environ Health*, 20, 160-165.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/7973487>
- 23 Melnick, R. L.; Huff, J.; Barrett, J. C.; Maronpot, R. R.; Lucier, G.; Portier, C. J. (1993). Cell  
24 proliferation and chemical carcinogenesis: symposium overview. *Environ Health*  
25 *Perspect*, 101 Suppl 5, 3-7. <http://www.ncbi.nlm.nih.gov/pubmed/7912189>
- 26 Melnick, R. L.; Kohn, M. C.; Portier, C. J. (1996). Implications for risk assessment of suggested  
27 nongenotoxic mechanisms of chemical carcinogenesis. *Environ Health Perspect*, 104  
28 Suppl 1, 123-134. <http://www.ncbi.nlm.nih.gov/pubmed/8722116>
- 29 Melnick, R. L.; Thayer, K. A.; Bucker, J. R. (2007). Conflicting views on chemical  
30 carcinogenesis arising from the design and evaluation of rodent carcinogenicity studies.  
31 *Environ Health Perspect*, 116, 130-135. <http://dx.doi.org/10.1289/ehp.9989>
- 32 Melnick, R.; Jameson, C.; Goehl, T.; Maronpot, R.; Collins, B.; Greenwell, A., . . . Agarwal, D.  
33 (1987). Application of microencapsulation for toxicology studies. II. Toxicity of  
34 microencapsulated trichloroethylene in Fischer 344 rats. *Fundam Appl Toxicol*, 8, 432-  
35 442. <http://www.ncbi.nlm.nih.gov/pubmed/3111923>
- 36 Menegazzi, M.; Carcereri-De Prati, A.; Suzuki, H.; Shinozuka, H.; Pibiri, M.; Piga, R., . . .  
37 Ledda-Columbano, G. M. (1997). Liver cell proliferation induced by nafenopin and  
38 cyproterone acetate is not associated with increases in activation of transcription factors  
39 NF-kappaB and AP-1 or with expression of tumor necrosis factor alpha. *Hepatology*, 25,  
40 585-592. <http://dx.doi.org/10.1002/hep.510250316>
- 41 Mensing, T.; Welge, P.; Voss, B.; Fels, L.; Fricke, H.; Brüning, T.; Wilhelm, M. (2002). Renal  
42 toxicity after chronic inhalation exposure of rats to trichloroethylene. *Toxicol Lett*, 128,  
43 243-247. <http://www.ncbi.nlm.nih.gov/pubmed/11869834>
- 44 Merdink, J. L.; Bull, R. J.; Schultz, I. R. (2000). Trapping and identification of the  
45 dichloroacetate radical from the reductive dehalogenation of trichloroacetate by mouse

DRAFT– DO NOT QUOTE OR CITE

- 1 and rat liver microsomes. *Free Radic Biol Med*, 29, 125-130.  
2 [http://dx.doi.org/10.1016/S0891-5849\(00\)00330-0](http://dx.doi.org/10.1016/S0891-5849(00)00330-0)
- 3 Merdink, J. L.; Robison, L. M.; Stevens, D. K.; Hu, M.; Parker, J. C.; Bull, R. J. (2008). Kinetics  
4 of chloral hydrate and its metabolites in male human volunteers. *Toxicology*, 245, 130-  
5 140. <http://dx.doi.org/10.1016/j.tox.2007.12.018>
- 6 Merdink, J.; Gonzalez-Leon, A.; Bull, R.; Schultz, I. (1998). The extent of dichloroacetate  
7 formation from trichloroethylene, chloral hydrate, trichloroacetate, and trichloroethanol  
8 in B6C3F1 mice. *Toxicol Sci*, 45, 33-41. <http://dx.doi.org/10.1006/toxs.1998.2500>
- 9 Merdink, J.; Stenner, R.; Stevens, D.; Parker, J.; Bull, R. (1999). Effect of enterohepatic  
10 circulation on the pharmacokinetics of chloral hydrate and its metabolites in F344 rats. *J*  
11 *Toxicol Environ Health A*, 57, 357-368. <http://www.ncbi.nlm.nih.gov/pubmed/10405189>
- 12 Mergler, D.; Huel, G.; Bowler, R.; Frenette, B.; Cone, J. (1991). Visual dysfunction among  
13 former microelectronics assembly workers. *Arch Environ Health*, 46, 326-334.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/1772256>
- 15 Merrick, B. A.; Robinson, M.; Condie, L. W. (1989). Differing hepatotoxicity and lethality after  
16 subacute trichloroethylene exposure in aqueous or corn oil gavage vehicles in B6C3F1  
17 mice. *J Appl Toxicol*, 9, 15-21. <http://www.ncbi.nlm.nih.gov/pubmed/2926094>
- 18 Messing, K.; Punnett, L.; Bond, M.; Alexanderson, K.; Pyle, J.; Zahm, S., . . . de Grosbois, S.  
19 (2003). Be the fairest of them all: challenges and recommendations for the treatment of  
20 gender in occupational health research. *Am J Ind Med*, 43, 618-629.  
21 <http://dx.doi.org/10.1002/ajim.10225>
- 22 Mester, B.; Nieters, A.; Deeg, E.; Elsner, G.; Becker, N.; Seidler, A. (2006). Occupation and  
23 malignant lymphoma: A population based case control study in Germany. *Occup Environ*  
24 *Med*, 63, 17-26. <http://dx.doi.org/10.1136/oem.2005.020453>
- 25 Meyer, K.; Lee, J.-S.; Dyck, P. A.; Cao, W.-Q.; Rao, M. S.; Thorgeirsson, S. S.; Reddy, J. K.  
26 (2003). Molecular profiling of hepatocellular carcinomas developing spontaneously in  
27 acyl-CoA oxidase deficient mice: Comparison with liver tumors induced in wild-type  
28 mice by a peroxisome proliferator and a genotoxic carcinogen. *Carcinogenesis*, 24, 975-  
29 984. <http://dx.doi.org/10.1093/carcin/bgg040>
- 30 Mhiri, C.; Choyakh, F.; Ben, H. M.; Feki, I.; Ben, M. M.; Zouari, N. (2004). Trigeminal  
31 somatosensory evoked potentials in trichloroethylene-exposed workers. 9, 102-107.
- 32 Michalik, L.; Desvergne, B.; Dreyer, C.; Gavillet, M.; Laurini, R.; Wahli, W. (2002). PPAR  
33 expression and function during vertebrate development. *Int J Dev Biol*, 46, 105-114.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/11902671>
- 35 Michel, C.; Desdouets, C.; Slaoui, M.; Isaacs, K. R.; Roberts, R. A.; Boitier, E. (2007).  
36 Diethylnitrosamine initiation does not alter clofibrac acid-induced hepatocarcinogenesis  
37 in the rat. *Toxicol Sci*, 99, 58-69. <http://dx.doi.org/10.1093/toxsci/kfm168>
- 38 Mikol, Y. B.; Hoover, K. L.; Creasia, D.; Poirier, L. A. (1983). Hepatocarcinogenesis in rats fed  
39 methyl-deficient, amino acid-defined diets. *Carcinogenesis*, 4, 1619-1629.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/6317218>
- 41 Miligi, L.; Costantini, A. S.; Benvenuti, A.; Kriebel, D.; Bolejack, V.; Tumino, R., . . . Vineis, P.  
42 (2006). Occupational exposure to solvents and the risk of lymphomas. *Epidemiology*, 17,  
43 552-561. <http://dx.doi.org/10.1097/01.ede.0000231279.30988.4d>
- 44 Miller, B. M. and Adler, I.-D. (1992). Aneuploidy induction in mouse spermatocytes.  
45 *Mutagenesis*, 7, 69-76.

DRAFT– DO NOT QUOTE OR CITE

- 1 Miller, R. E. and Guengerich, F. P. (1982). Oxidation of trichloroethylene by liver microsomal  
2 cytochrome P-450: evidence for chlorine migration in a transition state not involving  
3 trichloroethylene oxide. *Biochemistry*, 21, 1090-1097.  
4 <http://dx.doi.org/10.1021/bi00534a041>
- 5 Miller, R. E. and Guengerich, F. P. (1983). Metabolism of trichloroethylene in isolated  
6 hepatocytes, microsomes, and reconstituted enzyme systems containing cytochrome P-  
7 450. *Cancer Res*, 43, 1145-1152. <http://www.ncbi.nlm.nih.gov/pubmed/6825087>
- 8 Miller, R. T.; Shah, R. S.; Cattley, R. C.; Popp, J. A. (1996). The peroxisome proliferations WY-  
9 14,643 and methylclofenapate induce hepatocyte ploidy alterations and ploidy-specific  
10 DNA synthesis in F344 rats. *Toxicol Appl Pharmacol*, 138, 317-323.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/8658533>
- 12 Mills, N. C. and Means, A. R. (1972). Sorbitol dehydrogenase of rat testis: changes of activity  
13 during development, after hypophysectomy and following gonadotrophic hormone  
14 administration. *Endocrinology*, 91, 147-156.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/4337211>
- 16 Minamishima, Y. A.; Nakayama, K.; Nakayama, K.-I. (2002). Recovery of liver mass without  
17 proliferation of hepatocytes after partial hepatectomy in Skp2-deficient mice. *Cancer Res*,  
18 62, 995-999. <http://www.ncbi.nlm.nih.gov/pubmed/11861371>
- 19 Mirer, F. E. (2010). New evidence on the health hazards and control of metalworking fluids  
20 since completion of the OSHA advisory committee report. *Am J Ind Med*, 53, 792-801.  
21 <http://dx.doi.org/10.1002/ajim.20853>
- 22 Mirsalis, J. C.; Tyson, C. K.; Steinmetz, K. L.; Loh, E. K.; Hamilton, C. M.; Bakke, J. P.;  
23 Spalding, J. W. (1989). Measurement of unscheduled DNA synthesis and S-phase  
24 synthesis in rodent hepatocytes following in vivo treatment: Testing of 24 compounds.  
25 *Environ Mol Mutagen*, 14, 155-164. <http://dx.doi.org/10.1002/em.2850140305>
- 26 Mishima, N.; Hoffman, S.; Hill, E. G.; Krug, E. L. (2006). Chick embryos exposed to  
27 trichloroethylene in an ex ovo culture model show selective defects in early endocardial  
28 cushion tissue formation. *Birth Defects Res A Clin Mol Teratol*, 76, 517-527.  
29 <http://dx.doi.org/10.1002/bdra.20283>
- 30 Mitchell, A. B. S. and Parsons-Smith, B. G. (1969). Trichloroethylene neuropathy. *Br Med J*, 1,  
31 422-423.
- 32 Miura, Y.; Anzai, K.; Takahashi, S.; TBTIFL, O. (1997). A novel lipophilic spin probe for the  
33 measurement of radiation damage in mouse brain using in vivo electron spin resonance  
34 (ESR). *FEBS Lett*, 419, 99-102. <http://www.ncbi.nlm.nih.gov/pubmed/9426228>
- 35 Mjaatvedt, C.; Krug, E.; Markwald, R. (1991). An antiserum (ES1) against a particulate form of  
36 extracellular matrix blocks the transition of cardiac endothelium into mesenchyme in  
37 culture. *Dev Biol*, 145, 219-230. <http://www.ncbi.nlm.nih.gov/pubmed/2040370>
- 38 Mjaatvedt, C.; Lepera, R.; Markwald, R. (1987). Myocardial specificity for initiating endothelial-  
39 mesenchymal cell transition in embryonic chick heart correlates with a particulate  
40 distribution of fibronectin. *Dev Biol*, 119, 59-67.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/3539667>
- 42 Moghaddam, A. P.; Abbas, R.; Fisher, J. W.; Lipscomb, J. C. (1997). The role of mouse  
43 intestinal microflora in the metabolism of trichloroethylene, an in vivo study. *Hum Exp*  
44 *Toxicol*, 16, 629-635. <http://dx.doi.org/10.1177/096032719701601101>

DRAFT– DO NOT QUOTE OR CITE

- 1 Moghaddam, A. P.; Abbas, R.; Fisher, J. W.; Stavrou, S.; Lipscomb, J. C. (1996). Formation of  
2 dichloroacetic acid by rat and mouse gut microflora, an in vitro study. *Biochem Biophys*  
3 *Res Commun*, 228, 639-645. <http://dx.doi.org/10.1006/bbrc.1996.1709>
- 4 Moinzadeh, P.; Breuhahn, K.; Stützer, H.; Schirmacher, P. (2005). Chromosome alterations in  
5 human hepatocellular carcinomas correlate with aetiology and histological grade--results  
6 of an explorative CGH meta-analysis. *Br J Cancer*, 92, 935-941.  
7 <http://dx.doi.org/10.1038/sj.bjc.6602448>
- 8 Monster, A. C. (1979). Difference in uptake, elimination, and metabolism in exposure to  
9 trichloroethylene, 1,1,1-trichloroethane, and tetrachloroethylene. *Int Arch Occup Environ*  
10 *Health*, 42, 311-317. <http://www.ncbi.nlm.nih.gov/pubmed/422272>
- 11 Monster, A. C. and Houtkooper, J. M. (1979). Estimation of individual uptake of  
12 trichloroethylene, 1,1,1-trichloroethane and tetrachloroethylene from biological  
13 parameters. *Int Arch Occup Environ Health*, 42, 319-323.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/422273>
- 15 Monster, A. C.; Boersma, G.; Duba, W. C. (1976). Pharmacokinetics of trichloroethylene in  
16 volunteers, influence of workload and exposure concentration. *Int Arch Occup Environ*  
17 *Health*, 38, 87-102. <http://www.ncbi.nlm.nih.gov/pubmed/1002309>
- 18 Monster, A. C.; Boersma, G.; Duba, W. C. (1979). Kinetics of trichloroethylene in repeated  
19 exposure of volunteers. *Int Arch Occup Environ Health*, 42, 283-292.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/422269>
- 21 Moore, L. E.; Boffetta, P.; Karami, S.; Brennan, P.; Stewart, P. S.; Hung, R., . . . Rothman, N.  
22 (2010). Occupational trichloroethylene exposure and renal carcinoma risk: Evidence of  
23 genetic susceptibility by reductive metabolism gene variants. *Cancer Res*, 70, 6527-6536.  
24 <http://dx.doi.org/10.1158/0008-5472.CAN-09-4167>
- 25 Moore, L. E.; Wilson, R. T.; Campleman, S. L. (2005). Lifestyle factors, exposures, genetic  
26 susceptibility, and renal cell cancer risk: A review. *Cancer Invest*, 23, 240-255.  
27 <http://dx.doi.org/10.1081/cnv-200055962>
- 28 Moore, M. M. and Harrington-Brock, K. (2000). Mutagenicity of trichloroethylene and its  
29 metabolites: Implications for the risk assessment of trichloroethylene. *Environ Health*  
30 *Perspect*, 108, 215-223. <http://www.ncbi.nlm.nih.gov/pubmed/10807553>
- 31 Morata, T. C.; Fiorini, A. C.; Fischer, F. M.; Colacioppo, S.; Wallingford, K. M.; Krieg, E. F., . .  
32 . Cesar, C. L. (1997). Toluene-induced hearing loss among rotogravure printing workers.  
33 *Scand J Work Environ Health*, 23, 289-298.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/9322820>
- 35 Morata, T. C.; Johnson, A. C.; Nysten, P.; Svensson, E. B.; Cheng, J.; Krieg, E. F., . . . Franks, J.  
36 (2002). Audiometric findings in workers exposed to low levels of styrene and noise. *J*  
37 *Occup Environ Med*, 44, 806-814.  
38 <http://dx.doi.org/10.1097/01.jom.0000026645.83602.60>
- 39 Morgan, J. and Cassady, R. (2002). Community cancer assessment in response to long-time  
40 exposure to perchlorate and trichloroethylene in drinking water. *J Occup Environ Med*,  
41 44, 616-621. <http://www.ncbi.nlm.nih.gov/pubmed/12134524>
- 42 Morgan, R. W.; Kelsh, M. A.; Zhao, K.; et al. (2000). Erratum. *Epidemiology*, 11, 360.
- 43 Morgan, R. W.; Kelsh, M. A.; Zhao, K.; Heringer, S. (1998). Mortality of aerospace workers  
44 exposed to Trichloroethylene. *Epidemiology*, 9, 424-431.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/9647907>

DRAFT– DO NOT QUOTE OR CITE

- 1 Morgan, R. W.; Kelsh, M. A.; Zhao, K.; Heringer, S. (2000). Mortality of aerospace workers  
2 exposed to trichloroethylene. *Epidemiology*, 11, 360.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/9647907>
- 4 Morgenstern, H. (1998). Ecologic studies. In S. Greenland (Ed.), *Modern epidemiology* (pp. 459-  
5 480): Lippincott -Raven.
- 6 Morgenstern, H.; Froines, J.; Ritz, B.; Young, B. (1997). Epidemiologic study to determine  
7 possible adverse effects to Rocketdyne/Atomics International workers from exposure to  
8 ionizing radiation. Berkeley, CA: Public Health Institute. Retrieved from  
9 <http://www.mapcruzin.com/rocketdyne-cancer/radiation.pdf>.
- 10 Morgenstern, H.; Froines, J.; Ritz, B.; Young, B. (1999). Epidemiologic study to determine  
11 possible adverse effects to Rocketdyne/Atomics International workers from exposure to  
12 selected chemicals. Berkeley, CA: Public Health Institute. Retrieved from  
13 [http://www.ssflpanel.org/files/UCLA\\_chem.pdf](http://www.ssflpanel.org/files/UCLA_chem.pdf).
- 14 Morimura, K.; Cheung, C.; Ward, J. M.; Reddy, J. K.; Gonzalez, F. J. (2006). Differential  
15 susceptibility of mice humanized for peroxisome proliferator-activated receptor alpha to  
16 Wy-14,643-induced liver tumorigenesis. *Carcinogenesis*, 27, 1074-1080.  
17 <http://dx.doi.org/10.1093/carcin/bgi329>
- 18 Morioka, I.; Miyai, N.; Yamamoto, H.; Miyashita, K. (2000). Evaluation of combined effect of  
19 organic solvents and noise by the upper limit of hearing. *Ind Health*, 38, 252-257.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/10812851>
- 21 Moriya, M.; Ohta, T.; Watanabe, K.; Miyazawa, T.; Kato, K.; Shirasu, Y. (1983). Further  
22 mutagenicity studies on pesticides in bacterial reversion assay systems. *Mutat Res*, 116,  
23 185-216. <http://www.ncbi.nlm.nih.gov/pubmed/6339892>
- 24 Morrow, L.; Ryan, C.; Goldstein, G.; Hodgson, M. (1989). A distinct pattern of personality  
25 disturbance following exposure to mixtures of organic solvents. *J Occup Med*, 31, 743-  
26 746. <http://www.ncbi.nlm.nih.gov/pubmed/2795250>
- 27 Morrow, L.; Steinhauer, S.; Hodgson, M. (1992). Delay in P300 latency in patients with organic  
28 solvent exposure. *Arch Neurol*, 49, 315-320.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/1536637>
- 30 Mortelmans, K.; Haworth, S.; Lawlor, T.; Speck, W.; Tainer, B.; Zeiger, E. (1986). Salmonella  
31 mutagenicity tests: II. Results from the testing of 270 chemicals. *Environ Mutagen*, 8, 1-  
32 119. <http://dx.doi.org/10.1002/em.2860080702>
- 33 Morton, L. D.; Youssef, A. F.; Lloyd, E.; Kiorpes, A. L.; Goldsworthy, T. L.; Fort, F. L. (2002).  
34 Evaluation of carcinogenic responses in the Eker rat following short-term exposure to  
35 selected nephrotoxins and carcinogens. *Toxicol Pathol*, 30, 559-564.  
36 <http://www.ncbi.nlm.nih.gov/pubmed/12371664>
- 37 Morton, L. M.; Hartge, P.; Holford, T. R.; Holly, E. A.; Chiu, B. C.; Vineis, P., . . . Zheng, T.  
38 (2005). Cigarette smoking and risk of non-Hodgkin lymphoma: A pooled analysis from  
39 the International Lymphoma Epidemiology Consortium (interlymph). *Cancer Epidemiol*  
40 *Biomarkers Prev*, 14, 925-933. <http://dx.doi.org/10.1158/1055-9965.EPI-04-0693>
- 41 Morton, L. M.; Holford, T. R.; Leaderer, B.; Zhang, Y.; Zahm, S. H.; Boyle, P., . . . Zheng, T.  
42 (2003). Alcohol use and risk of non-Hodgkin's lymphoma among Connecticut women  
43 (United States). *Cancer Causes Control*, 14, 687-694.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/14575367>



DRAFT– DO NOT QUOTE OR CITE

- 1 Morton, L. M.; Turner, J. J.; Cerhan, J. R.; Linet, M. S.; Treseler, P. A.; Clarke, C. A., . . .  
2 Weisenburger, D. D. (2007). Proposed classification of lymphoid neoplasms for  
3 epidemiologic research from the Pathology Working Group of the International  
4 Lymphoma Epidemiology Consortium (InterLymph). *Blood*, 110, 695-708.  
5 <http://dx.doi.org/10.1182/blood-2006-11-051672>
- 6 Moser, V. C.; Cheek, B. M.; MacPhail, R. C. (1995). A multidisciplinary approach to  
7 toxicological screening III Neurobehavioral toxicity. *J Toxicol Environ Health*, 45, 173-  
8 210. <http://www.ncbi.nlm.nih.gov/pubmed/7783252>
- 9 Moser, V. C.; MacPhail, R. C.; Gennings, C. (2003). Neurobehavioral evaluations of mixtures of  
10 trichloroethylene, heptachlor, and di(2-ethylhexyl)phthalate in a full-factorial design.  
11 *Toxicology*, 188, 125-137. <http://www.ncbi.nlm.nih.gov/pubmed/12767685>
- 12 Moser, V. C.; Phillips, P. M.; McDaniel, K. L.; MacPhail, R. C. (1999). Behavioral evaluation of  
13 the neurotoxicity produced by dichloroacetic acid in rats. *Neurotoxicol Teratol*, 21, 719-  
14 731. <http://www.ncbi.nlm.nih.gov/pubmed/10560779>
- 15 Moslen, M. T.; Reynolds, E. S.; Boor, P. J.; Bailey, K.; Szabo, S. (1977). Trichloroethylene-  
16 induced deactivation of cytochrome P-450 and loss of liver glutathione in vivo. *Res*  
17 *Commun Mol Pathol Pharmacol*, 16, 109-120.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/841173>
- 19 Motohashi, Y. and Miyazaki, Y. (1990). Time-dependent effects of trichloroethylene on motor  
20 activity in rats. *Chronobiol Int*, 7, 193-201.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/2268881>
- 22 Mueller, W. F.; Coulston, F.; Korte, F. (1982). Comparative metabolism of 14C-  
23 trichloroethylene in chimpanzees, baboons, and rhesus monkeys. *Chemosphere*, 11, 215-  
24 218.
- 25 Muijser, H.; Lammers, J.; Kullig, B. (2000). Effects of exposure to trichloroethylene and noise  
26 on hearing in rats. *Noise Health*, 2, 57-66.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/12689479>
- 28 Muller, G.; Spassovski, M.; Henschler, D. (1972). Trichloroethylene exposure and  
29 trichloroethylene metabolites in urine and blood. *Arch Toxicol*, 29, 335-340.
- 30 Muller, G.; Spassovski, M.; Henschler, D. (1974). Metabolism of trichloroethylene in man II  
31 Pharmacokinetics of metabolites. *Arch Toxicol*, 32, 283-295.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/4480028>
- 33 Muller, G.; Spassowski, M.; Henschler, D. (1975). Metabolism of trichloroethylene in man III  
34 Interaction of trichloroethylene and ethanol. *Arch Toxicol*, 33, 173-189.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/1173750>
- 36 Murphy, J. P. (1990). Exposure to wells G and H in Woburn, Massachusetts. Boston, MA:  
37 Massachusetts Department of Public Health. Retrieved from  
38 [http://www.mass.gov/Eeohhs2/docs/dph/environmental/investigations/woburn\\_exposure\\_](http://www.mass.gov/Eeohhs2/docs/dph/environmental/investigations/woburn_exposure_wells.pdf)  
39 [wells.pdf](http://www.mass.gov/Eeohhs2/docs/dph/environmental/investigations/woburn_exposure_wells.pdf).
- 40 Nagaya, T.; Ishikawa, N.; Hata, H. (1989). Sister-chromatid exchanges in lymphocytes of  
41 workers exposed to trichloroethylene. *Mutat Res*, 222, 279-282.
- 42 Nagaya, T.; Ishikawa, N.; Hata, H. (1989). Urinary total protein and "beta"-2-microglobulin in  
43 workers exposed to trichloroethylene. *Environ Res*, 50, 86-92.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/2676510>

DRAFT– DO NOT QUOTE OR CITE

- 1 Nagaya, T.; Ishikawa, N.; Hata, H. (1990). No change in serum dopamine-"beta"-hydroxylase  
2 activity in workers exposed to trichloroethylene. *Toxicol Lett*, 54, 221-227.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/2260120>
- 4 Nagaya, T.; Ishikawa, N.; Hata, H.; Otobe, T. (1993). Subclinical and reversible hepatic effects  
5 of occupational exposure to trichloroethylene. *Int Arch Occup Environ Health*, 64, 561-  
6 563. <http://dx.doi.org/10.1007/BF00517701>
- 7 Nakai, J. S.; Stathopoulos, P. B.; Campbell, G. L.; Chu, I.; Li-Muller, A.; Aucoin, R. (1999).  
8 Penetration of chloroform, trichloroethylene, and tetrachloroethylene through human  
9 skin. *J Toxicol Environ Health A*, 58, 157-170.  
10 <http://dx.doi.org/10.1080/009841099157368>
- 11 Nakajima, T. and Aoyama, T. (2000). Polymorphism of drug-metabolizing enzymes in relation  
12 to individual susceptibility to industrial chemicals. *Ind Health*, 38, 143-152.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/10812837>
- 14 Nakajima, T.; Kamijo, Y.; Usuda, N.; Liang, Y.; Fukushima, Y.; Kametani, K., . . . Aoyama, T.  
15 (2000). Sex-dependent regulation of hepatic peroxisome proliferation in mice by  
16 trichloroethylene via peroxisome proliferator-activated receptor alpha (PPARalpha).  
17 *Carcinogenesis*, 21, 677-682. <http://www.ncbi.nlm.nih.gov/pubmed/10753203>
- 18 Nakajima, T.; Koyama, Y.; Sato, A. (1982). Dietary modification of metabolism and toxicity of  
19 chemical substances - with special reference to carbohydrate. *Biochem Pharmacol*, 31,  
20 1005-1011. <http://www.ncbi.nlm.nih.gov/pubmed/7082357>
- 21 Nakajima, T.; Moriguchi, M.; Mitsumoto, Y.; Sekoguchi, S.; Nishikawa, T.; Takashima, H., . . .  
22 Kagawa, K. (2004). Centrosome aberration accompanied with p53 mutation can induce  
23 genetic instability in hepatocellular carcinoma. *Mod Pathol*, 17, 722-727.  
24 <http://dx.doi.org/10.1038/modpathol.3800115>
- 25 Nakajima, T.; Okino, T.; Okuyama, S.; Kaneko, T.; Yonekura, I.; Sato, A. (1988). Ethanol-  
26 induced enhancement of trichloroethylene metabolism and hepatotoxicity: difference  
27 from the effect of phenobarbital. *Toxicol Appl Pharmacol*, 94, 227-237.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/3388420>
- 29 Nakajima, T.; Wang, R. S.; Katakura, Y.; Kishi, R.; Elovaara, E.; Park, S. S., . . . Vainio, H.  
30 (1992). Sex-, age- and pregnancy-induced changes in the metabolism of toluene and  
31 trichloroethylene in rat liver in relation to the regulation of cytochrome P450IIE1 and  
32 P450IIC11 content. *J Pharmacol Exp Ther*, 261, 869-874.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/1602390>
- 34 Nakajima, T.; Wang, R.-S.; Elovaara, E.; Park, S. S.; Gelboin, H. V.; Vainio, H. (1992). A  
35 comparative study on the contribution of cytochrome P450 isozymes to metabolism of  
36 benzene, toluene and trichloroethylene in rat liver. *Biochem Pharmacol*, 43, 251-257.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/1739413>
- 38 Nakajima, T.; Wang, R.-S.; Elovaara, E.; Park, S. S.; Gelboin, H. V.; Vainio, H. (1993).  
39 Cytochrome P450-related differences between rats and mice in the metabolism of  
40 benzene, toluene and trichloroethylene in liver microsomes. *Biochem Pharmacol*, 45,  
41 1079-1085. [http://dx.doi.org/10.1016/0006-2952\(93\)90252-R](http://dx.doi.org/10.1016/0006-2952(93)90252-R)
- 42 Nakajima, T.; Wang, R.-S.; Murayama, N.; Sato, A. (1990). Three forms of trichloroethylene-  
43 metabolizing enzymes in rat liver induced by ethanol, phenobarbital, and 3-  
44 methylcholanthrene. *Toxicol Appl Pharmacol*, 102, 546-552.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/2315921>

DRAFT– DO NOT QUOTE OR CITE

- 1 Nakajima, T.; Yamanoshita, O.; Kamijima, M.; Kishi, R.; Ichihara, G. (2003). Generalized skin  
2 reactions in relation to trichloroethylene exposure: a review from the viewpoint of drug-  
3 metabolizing enzymes. *J Occup Health*, 45, 8-14.  
4 <http://www.ncbi.nlm.nih.gov/pubmed/14605423>
- 5 Nakajima, Y.; Mironov, V.; Yamagishi, T.; Nakamura, H.; Markwald, R. (1997). Expression of  
6 smooth muscle alpha-actin in mesenchymal cells during formation of avian endocardial  
7 cushion tissue: a role for transforming growth factor beta3. *Dev Dyn*, 209, 296-309.  
8 [http://dx.doi.org/10.1002/\(SICI\)1097-0177\(199707\)209:3<296::AID-  
9 AJA5>3.0.CO;2-D](http://dx.doi.org/10.1002/(SICI)1097-0177(199707)209:3<296::AID-AJA5>3.0.CO;2-D)
- 10 Nakanishi, S.; Shiohara, E.; Tsukada, M.; Yamazaki, Y.; Okumura, K. (1978). Acetaldehyde  
11 level in the blood and liver aldehyde dehydrogenase activities in trichloroethylene-treated  
12 rats. *Arch Toxicol*, 41, 207-214. <http://www.ncbi.nlm.nih.gov/pubmed/736791>
- 13 Narotsky, M. G. and Kavlock, R. J. (1995). A multidisciplinary approach to toxicological  
14 screening: II. Developmental toxicity. *J Toxicol Environ Health*, 45, 145-171.  
15 <http://dx.doi.org/10.1080/15287399509531987>
- 16 Narotsky, M. G.; Weller, E. A.; Chinchilli, V. M.; Kavlock, R. J. (1995). Nonadditive  
17 developmental toxicity in mixtures of trichloroethylene, Di(2-ethylhexyl) phthalate, and  
18 heptachlor in a 5 x 5 x 5 design. *Fundam Appl Toxicol*, 27, 203-216.  
19 <http://dx.doi.org/10.1093/toxsci/27.2.203>
- 20 Natarajan, A.; Duivenvoorden, W.; Meijers, M.; Zwanenburg, T. (1993). Induction of mitotic  
21 aneuploidy using Chinese hamster primary embryonic cells. Test results of 10 chemicals.  
22 *Mutat Res*, 287, 47-56. <http://www.ncbi.nlm.nih.gov/pubmed/7683384>
- 23 Naugler, W. E.; Sakurai, T.; Kim, S.; Maeda, S.; Kim, K.; Elsharkawy, A. M.; Karin, M. (2007).  
24 Gender disparity in liver cancer due to sex differences in MyD88-dependent IL-6  
25 production. *Science*, 317, 121-124. <http://dx.doi.org/10.1126/science.1140485>
- 26 NCI. (National Institutes of Health, National Cancer Institute). (1976). Carcinogenesis bioassay  
27 of trichloroethylene. (NCI-CG-TR-2). Bethesda, MD: National Cancer Institute.  
28 Retrieved from [http://ntp.niehs.nih.gov/ntp/htdocs/LT\\_rpts/tr002.pdf](http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr002.pdf).
- 29 NCI. (National Institutes of Health, National Cancer Institute). (2008). SEER cancer statistics  
30 review, 1975-2005. Bethesda, MD. Retrieved from [http://seer.cancer.gov/csr/1975\\_2005](http://seer.cancer.gov/csr/1975_2005).
- 31 Neghab, M. and Stacey, N. (1997). Toluene-induced elevation of serum bile acids: relationship  
32 to bile acid transport. *J Toxicol Environ Health*, 52, 249-268.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/9316646>
- 34 Neghab, M.; Qu, S.; Bai, C. L.; Caples, J.; Stacey, N. H. (1997). Raised concentration of serum  
35 bile acids following occupational exposure to halogenated solvents, 1,1,2-trichloro-1,2,2-  
36 trifluoroethane and trichloroethylene. *Int Arch Occup Environ Health*, 70, 187-194.  
37 <http://dx.doi.org/10.1007/s004200050205>
- 38 Nelson, G. M.; Swank, A. E.; Brooks, L. R.; Bailey, K. C.; George, S. E. (2001). Metabolism,  
39 microflora effects, and genotoxicity in haloacetic acid-treated cultures of rat cecal  
40 microbiota. *Toxicol Sci*, 60, 232-241. <http://www.ncbi.nlm.nih.gov/pubmed/11248134>
- 41 Nelson, H. H.; Wiencke, J. K.; Christiani, D. C.; Cheng, T. J.; Zuo, Z.-F.; Schwartz, B. S., . . .  
42 Kelsey, K. T. (1995). Ethnic differences in the prevalence of the homozygous deleted  
43 genotype of glutathione S-transferase theta. *Carcinogenesis*, 16, 1243-1245.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/7767992>

DRAFT– DO NOT QUOTE OR CITE

- 1 Nelson, J. and Zenick, H. (1986). The effect of trichloroethylene on male sexual behavior:  
2 possible opioid role. *Neurobehav Toxicol Teratol*, 8, 441-445.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/3785506>
- 4 Nelson, L.; Longstreth, W.; Koepsell, T.; Checkoway, H.; van Belle, G. (1994). Completeness  
5 and accuracy of interview data from proxy respondents: demographic, medical, and life-  
6 style factors. *Epidemiology*, 5, 204-217. <http://www.ncbi.nlm.nih.gov/pubmed/8172996>
- 7 Nelson, M. A. and Bull, R. J. (1988). Induction of strand breaks in DNA by trichloroethylene  
8 and metabolites in rat and mouse liver in vivo. *Toxicol Appl Pharmacol*, 94, 45-54.  
9 [http://dx.doi.org/10.1016/0041-008X\(88\)90335-3](http://dx.doi.org/10.1016/0041-008X(88)90335-3)
- 10 Nelson, M. A.; Lansing, A. J.; Sanchez, I. M.; Bull, R. J.; Springer, D. L. (1989). Dichloroacetic  
11 acid and trichloroacetic acid-induced DNA strand breaks are independent of peroxisome  
12 proliferation. *Toxicology*, 58, 239-248. [X](http://dx.doi.org/10.1016/0300-483X(89)90139-)  
13
- 14 Nelson, M. A.; Sanchez, I. M.; Bull, R. J.; Sylvester, S. R. (1990). Increased expression of c-myc  
15 and c-H-ras in dichloroacetate and trichloroacetate-induced liver tumors in B6C3F1 mice.  
16 *Toxicology*, 64, 47-57.
- 17 Nesslany, F. and Marzin, D. (1999). A micromethod for the in vitro micronucleus assay.  
18 *Mutagenesis*, 14, 403-410. <http://www.ncbi.nlm.nih.gov/pubmed/10390508>
- 19 Nestmann, E. R.; Chu, I.; Kowbel, D. J.; Matula, T. I. (1980). Short-lived mutagen in Salmonella  
20 produced by reaction of trichloroacetic acid and dimethyl sulphoxide. *Can J Genet Cytol*,  
21 22, 35-40. <http://dx.doi.org/10.1139/g80-006>
- 22 Ni, Y. C.; Kadlubar, F. F.; Fu, P. P. (1995). Formation of malondialdehyde-modified 2'-  
23 deoxyguanosinyl adduct from metabolism of chloral hydrate by mouse liver microsomes.  
24 *Biochem Biophys Res Commun*, 216, 1110-1117.  
25 <http://dx.doi.org/10.1006/bbrc.1995.2735>
- 26 Ni, Y. C.; Wong, T. Y.; Lloyd, R. V.; Heinze, T. M.; Shelton, S.; Casciano, D., . . . Fu, P. P.  
27 (1996). Mouse liver microsomal metabolism of chloral hydrate, trichloroacetic acid, and  
28 trichloroethanol leading to induction of lipid peroxidation via a free radical mechanism.  
29 *Drug Metab Dispos*, 24, 81-90. <http://www.ncbi.nlm.nih.gov/pubmed/8825194>
- 30 Ni, Y.; Wong, T.; Kadlubar, F.; Fu, P. (1994). Hepatic metabolism of chloral hydrate to free  
31 radical(s) and induction of lipid peroxidation. *Biochem Biophys Res Commun*, 204, 937-  
32 943. <http://dx.doi.org/10.1006/bbrc.1994.2550>
- 33 Nickerson, M. L.; Jaeger, E.; Shi, Y.; Durocher, J. A.; Mahurkar, S.; Zaridze, D., . . . Moore, L.  
34 E. (2008). Improved identification of von Hippel-Lindau gene alterations in clear cell  
35 renal tumors. *Clin Cancer Res*, 14, 4726-4734. [0432.CCR-07-4921](http://dx.doi.org/10.1158/1078-)  
36
- 37 Nickoloff, B. J.; Ben-Neriah, Y.; Pikarsky, E. (2005). Inflammation and cancer: is the link as  
38 simple as we think. *J Invest Dermatol*, 124, x-xiv. [202X.2005.23724.x](http://dx.doi.org/10.1111/j.0022-)  
39
- 40 Niederlehner, B.; Cairns, J.; Smith, E. (1998). Modeling acute and chronic toxicity of nonpolar  
41 narcotic chemicals and mixtures to *Ceriodaphnia dubia*. *Ecotoxicol Environ Saf*, 39, 136-  
42 146. <http://dx.doi.org/10.1006/eesa.1997.1621>
- 43 NIEHS. (National Institutes of Health, National Institute for Environmental Health Sciences).  
44 (1998). Tumor incidence in control animals by route and vehicle of administration,  
45 B6C3F1 mice. Research Triangle Park, NC.

DRAFT– DO NOT QUOTE OR CITE

- 1 Nietert, P. J.; Sutherland, S. E.; Silver, R. M.; Pandey, J. P.; Knapp, R. G.; Hoel, D. G.;  
2 Dosemeci, M. (1998). Is occupational organic solvent exposure a risk factor for  
3 scleroderma? *Arthritis Rheum*, 41, 1111-1118. [http://dx.doi.org/10.1002/1529-](http://dx.doi.org/10.1002/1529-0131(199806)41:6<1111::AID-ART19>3.0.CO;2-J)  
4 [0131\(199806\)41:6<1111::AID-ART19>3.0.CO;2-J](http://dx.doi.org/10.1002/1529-0131(199806)41:6<1111::AID-ART19>3.0.CO;2-J)
- 5 Niklasson, M.; Tham, R.; Larsby, B.; Eriksson, B. (1993). Effects of toluene, styrene,  
6 trichloroethylene, and trichloroethane on the vestibulo- and opto-oculo motor system in  
7 rats. *Neurotoxicol Teratol*, 15, 327-334. <http://www.ncbi.nlm.nih.gov/pubmed/8277926>
- 8 NIOSH. (National Institute for Occupational Safety and Health). (1992). Health Hazard  
9 Evaluation Report HETA 86-469-2189, James River Corporation, Newnan, Georgia.  
10 (HETA 86-469-2189). Springfield, VA.
- 11 NIOSH. (National Institute for Occupational Safety and Health). (1998). Criteria for a  
12 recommended standard: Occupational exposure to metalworking fluids. (98-102).  
13 Atlanta, GA: Author. Retrieved from <http://www.cdc.gov/niosh/98-102.html>.
- 14 Nissen, S. E.; Nicholls, S. J.; Wolski, K.; Howey, D. C.; McErlean, E.; Wang, M. D., . . . Russo,  
15 J. M. (2007). Effects of a potent and selective PPAR-alpha agonist in patients with  
16 atherogenic dyslipidemia or hypercholesterolemia: two randomized controlled trials.  
17 *JAMA*, 297, 1362-1373. <http://dx.doi.org/10.1001/jama.297.12.1362>
- 18 Nitcheva, D. K.; Piegorsch, W. W.; West, R. W. (2007). On use of the multistage dose-response  
19 model for assessing laboratory animal carcinogenicity. *Regul Toxicol Pharmacol*, 48,  
20 135-147. <http://dx.doi.org/10.1016/j.yrtph.2007.03.002>
- 21 Nöhammer, C.; Brunner, F.; Wölkart, G.; Staber, P.; Steyrer, E.; Gonzalez, F., . . . Hoefler, G.  
22 (2003). Myocardial dysfunction and male mortality in peroxisome proliferator-activated  
23 receptor alpha knockout mice overexpressing lipoprotein lipase in muscle. *Lab Invest*, 83,  
24 259-269. <http://www.ncbi.nlm.nih.gov/pubmed/12594240>
- 25 Noland-Gerbec, E. A.; Pfohl, R. J.; Taylor, D. H.; Bull, R. J. (1986). 2-Deoxyglucose uptake in  
26 the developing rat brain upon pre- and postnatal exposure to trichloroethylene.  
27 *Neurotoxicology*, 7, 157-164. <http://www.ncbi.nlm.nih.gov/pubmed/3822256>
- 28 Nomiyama, H. and Nomiyama, K. (1979). Pathway and rate of metabolism of trichloroethylene  
29 in rats and rabbits. *Ind Health*, 17, 29-37.
- 30 Nomiyama, K. and Nomiyama, H. (1971). Metabolism of trichloroethylene in human: sex  
31 difference in urinary excretion of trichloroacetic acid and trichloroethanol. *Int Arch*  
32 *Occup Environ Health*, 28, 37-48. <http://www.ncbi.nlm.nih.gov/pubmed/5088555>
- 33 Nordlund, L. A.; Carstensen, J. M.; Pershagen, G. (1997). Cancer incidence in female smokers: a  
34 26-year follow-up. *Int J Cancer*, 73, 625-628.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/9398036>
- 36 Nordlund, L. A.; Carstensen, J. M.; Pershagen, G. (1999). Are male and female smokers at equal  
37 risk of smoking-related cancer: evidence from a Swedish prospective study. *Scand J*  
38 *Public Health*, 27, 56-62. <http://www.ncbi.nlm.nih.gov/pubmed/10847673>
- 39 Nordström, M.; Hardell, L.; Magnuson, A.; Hagberg, H.; Rask-Andersen, A. (1998).  
40 Occupational exposures, animal exposure and smoking as risk factors for hairy cell  
41 leukaemia evaluated in a case-control study. *Br J Cancer*, 77, 2048-2052.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/9667691>
- 43 NRC. (National Research Council). (1983). Risk assessment in the federal government:  
44 Managing the process. Washington, DC: National Academy Press.

DRAFT– DO NOT QUOTE OR CITE

- 1 NRC. (National Research Council). (1988). Health risks of radon and other internally deposited  
2 alpha-emitters: BEIR IV. (PB88-225594). Washington, DC: National Research Council;  
3 National Academy Press. Retrieved from  
4 <http://www.nap.edu/openbook.php?isbn=0309037972>.
- 5 NRC. (National Research Council). (1993). Pesticides in the diets of infants and children.  
6 Washington, DC: National Academy Press.
- 7 NRC. (National Research Council). (1994). Science and judgment in risk assessment.  
8 Washington, DC: National Academy Press.
- 9 NRC. (National Research Council). (1997). Environmental epidemiology, volume 2: Use of the  
10 gray literature and other data in environmental epidemiology (Vol. 2). Washington, DC:  
11 National Academies Press.
- 12 NRC. (National Research Council). (2006). Assessing the human health risks of  
13 trichloroethylene: Key scientific issues. Washington, DC: The National Academies  
14 Press. Retrieved from [http://nae.edu/nae/naepcms.nsf/weblinks/MKEZ-  
15 6SSHPD?OpenDocument](http://nae.edu/nae/naepcms.nsf/weblinks/MKEZ-6SSHPD?OpenDocument).
- 16 NRC. (National Research Council). (2009). Contaminated water supplies at Camp Lejeune:  
17 Assessing potential health effects. Washington, DC: National Academies Press.
- 18 NTP. (National Toxicology Program). (1988). Toxicology and carcinogenesis studies of  
19 trichloroethylene (CAS No. 79-01-6) in four strains of rats (ACI, August, Marshall,  
20 Osborne-Mendel)(gavage studies). Research Triangle Park, NC: Public Health Service,  
21 U.S. Department of Health and Human Services. Retrieved from  
22 [http://ntp.niehs.nih.gov/ntp/htdocs/LT\\_rpts/tr273.pdf](http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr273.pdf).
- 23 NTP. (National Toxicology Program). (1990). Carcinogenesis studies of trichloroethylene  
24 (without epichlorohydrin) (CAS no 79-01-6) in F344/N rats and B6C3F mice (gavage  
25 studies). Research Triangle Park, NC: Author.
- 26 Nunes, J.; Ehrich, M.; Robertson, J. (2001). Toxicosis associated with dual oral exposure of rats  
27 to lead and trichloroethylene. *Toxicol Pathol*, 29, 451-457.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/11560250>
- 29 Nutley, E.; Tcheong, A.; Allen, J.; Collins, B.; Ma, M.; Lowe, X., . . . Wyrobek, A. (1996).  
30 Micronuclei induced in round spermatids of mice after stem-cell treatment with chloral  
31 hydrate: evaluations with centromeric DNA probes and kinetochore antibodies. *Environ*  
32 *Mol Mutagen*, 28, 80-89. [http://dx.doi.org/10.1002/\(SICI\)1098-  
33 2280\(1996\)28:2<80::AID-EM3>3.0.CO;2-I](http://dx.doi.org/10.1002/(SICI)1098-2280(1996)28:2<80::AID-EM3>3.0.CO;2-I)
- 34 NYSDEC. (New York State Department of Environmental Conservation). (2007). Project  
35 information summary: Solvents, contamination and cleanup at Endicott. Albany, NY.  
36 Retrieved from <http://www.dec.ny.gov/chemical/24887.html>.
- 37 Nysdoh. (2005). Trichloroethene air criteria document. Draft report. Retrieved from  
38 [file:///L:/Lab/NCEA/TCE Documents/TCE PDFs4\NYS DH 2005 Draft Trichloroethene  
39 Air Criteria.pdf](file:///L:/Lab/NCEA/TCE Documents/TCE PDFs4/NYS DH 2005 Draft Trichloroethene Air Criteria.pdf).
- 40 NYSDOH. (New York State Department of Health). (2006). Trichloroethene air criteria  
41 document: Final report. Troy, NY.
- 42 O'Donnell, L.; McLachlan, R. I.; Wreford, N. G.; Robertson, D. M. (1994). Testosterone  
43 promotes the conversion of round spermatids between stages VII and VIII of the rat  
44 spermatogenic cycle. *Endocrinology*, 135, 2608-2614.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/7988449>

- 1 Odum, J.; Foster, J. R.; Green, T. (1992). A mechanism for the development of Clara cell lesions  
2 in the mouse lung after exposure to trichloroethylene. *Chem Biol Interact*, 83, 135-153.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/1505057>
- 4 Ogata, M.; Takatsuka, Y.; Tomokuni, K. (1971). Excretion of organic chlorine compounds in the  
5 urine of persons exposed to vapours of trichloroethylene and tetrachloroethylene. *Occup*  
6 *Environ Med*, 28, 386-391. <http://dx.doi.org/10.1136/oem.28.4.386>
- 7 Ohta, M.; Saito, T.; Saito, K.; Kurasaki, M.; Hosokawa, T. (2001). Effect of trichloroethylene on  
8 spatiotemporal pattern of LTP in mouse hippocampal slices. *Int J Neurosci*, 111, 257-  
9 271. <http://www.ncbi.nlm.nih.gov/pubmed/11912680>
- 10 Okino, T.; Nakajima, T.; Nakano, M. (1991). Morphological and biochemical analyses of  
11 trichloroethylene hepatotoxicity: differences in ethanol- and phenobarbital-pretreated  
12 rats. *Toxicol Appl Pharmacol*, 108, 379-389.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/2020965>
- 14 Olivier, P.; Plancke, M. O.; Marzin, D.; Clavey, V.; Sauzies, J.; Fruchart, J. C. (1988). Effects  
15 of fenofibrate, gemfibrozil and nicotinic acid on plasma lipoprotein levels in normal and  
16 hyperlipidemic mice. A proposed model for drug screening. *Atherosclerosis*, 70, 107-  
17 114. <http://www.ncbi.nlm.nih.gov/pubmed/3162679>
- 18 Olshan, A. F.; De Roos, A. J.; Teschke, K.; Neglia, J. P.; Stram, D. O.; Pollock, B. H.;  
19 Castleberry, R. P. (1999). Neuroblastoma and parental occupation. *Cancer Causes*  
20 *Control*, 10, 539-549. <http://www.ncbi.nlm.nih.gov/pubmed/10616823>
- 21 Olsson, A.; Skogh, T.; Axelson, O.; Wingren, G. (2004). Occupations and exposures in the work  
22 environment as determinants for rheumatoid arthritis. *Occup Environ Med*, 61, 233-238.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/14985518>
- 24 Olsson, A.; Skogh, T.; Wingren, G. (2000). Occupational determinants for rheumatoid arthritis.  
25 *Scand J Work Environ Health*, 26, 243-249.  
26 <http://www.ncbi.nlm.nih.gov/pubmed/10901117>
- 27 Oneta, C.; Lieber, C.; Li, J.; Rüttimann, S.; Schmid, B.; Lattmann, J., . . . Seitz, H. (2002).  
28 Dynamics of cytochrome P4502E1 activity in man: induction by ethanol and  
29 disappearance during withdrawal phase. *J Hepatol*, 36, 47-52.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/11804663>
- 31 Ono, Y.; Somiya, I.; Kawamura, M. (1991). The evaluation of genotoxicity using DNA repairing  
32 test for chemicals produced in chlorination and ozonation processes. *Water Sci Technol*,  
33 23, 329-338.
- 34 Opdam, J. J. (1989). Intra and interindividual variability in the kinetics of a poorly and highly  
35 metabolising solvent. *Br J Ind Med*, 46, 831-845.  
36 <http://www.ncbi.nlm.nih.gov/pubmed/2611156>
- 37 Opdam, J. J. (1989). Respiratory input in inhalation experiments. *Occup Environ Med*, 46, 145-  
38 156. <http://dx.doi.org/10.1136/oem.46.3.145>
- 39 Orsi, L.; Monnereau, A.; Dananche, B.; Berthou, C.; Fenaux, P.; Marit, G., . . . Clavel, J. (2010).  
40 Occupational exposure to organic solvents and lymphoid neoplasms in men: Results of a  
41 French case-control study. *Occup Environ Med*, 67, 664-672.  
42 <http://dx.doi.org/10.1136/oem.2009.049460>
- 43 Oshiro, W.; Krantz, Q.; Bushnell, P. (2004). A search for residual behavioral effects of  
44 trichloroethylene (TCE) in rats exposed as young adults. *Neurotoxicol Teratol*, 26, 239-  
45 251. <http://dx.doi.org/10.1016/j.ntt.2003.12.005>

DRAFT– DO NOT QUOTE OR CITE

- 1 Ou, J.; Ou, Z.; McCarver, D.; Hines, R.; Oldham, K.; Ackerman, A.; Pritchard, K. (2003).  
2 Trichloroethylene decreases heat shock protein 90 interactions with endothelial nitric  
3 oxide synthase: implications for endothelial cell proliferation. *Toxicol Sci*, 73, 90-97.  
4 <http://dx.doi.org/10.1093/toxsci/kfg062>
- 5 Owens, A. H. and Marshall, E. K. (1955). Further studies on the metabolic rate of chloral hydrate  
6 and trichloroethanol. *Johns Hopkins Med J*, 97, 320-326.
- 7 Palmer, C. N.; Hsu, M. H.; Griffin, K. J.; Raucy, J. L.; Johnson, E. F. (1998). Peroxisome  
8 proliferator activated receptor-alpha expression in human liver. *Mol Pharmacol*, 53, 14-  
9 22. <http://www.ncbi.nlm.nih.gov/pubmed/9443928>
- 10 Pannett, B.; Coggon, D.; Acheson, E. (1985). A job-exposure matrix for use in population based  
11 studies in England and Wales. *Br J Ind Med*, 42, 777-783.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/4063222>
- 13 Parent, M. E.; Hua, Y.; Siemiatycki, J. (2000). Occupational risk factors for renal cell carcinoma  
14 in Montreal. *Am J Ind Med*, 38, 609-618.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/11071683>
- 16 Parent, M. E.; Siemiatycki, J.; Fritschi, L. (2000). Workplace exposures and oesophageal cancer.  
17 *Occup Environ Med*, 57, 325-334. <http://www.ncbi.nlm.nih.gov/pubmed/10769298>
- 18 Paris, France: Commission of the European Communities.
- 19 Park, P. A.; Gilmour, M. I.; Selgrade, M. K. (1993). Pulmonary defenses to streptococcal  
20 infection following acute exposure to trichloroethylene (TCE) or chloroform (CHC13).  
21 *Toxicologist*, 13, 107.
- 22 Park, S.; Osterloh, J.; Vamvakas, S.; Hashmi, M.; Anders, M.; Cashman, J. (1992). Flavin-  
23 containing monooxygenase-dependent stereoselective S-oxygenation and cytotoxicity of  
24 cysteine S-conjugates and mercapturates. *Chem Res Toxicol*, 5, 193-201.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/1643249>
- 26 Parker, M. J.; Tarlow, M. J.; Milne Anderson, J. (1984). Glue sniffing and cerebral infarction.  
27 *Arch Dis Child*, 59, 675-677. <http://www.ncbi.nlm.nih.gov/pubmed/6465942>
- 28 Parkin, D. M.; Whelan, S. L.; Ferlay, J.; Teppo, L.; Thomas, D. B. (Eds.). (2002). Cancer  
29 incidence in five continents: Volume VIII. Lyon, France: International Agency for  
30 Research on Cancer.
- 31 Parkinson, A.; Mudra, D. R.; Johnson, C.; Dwyer, A.; Carroll, K. M. (2004). The effects of  
32 gender, age, ethnicity, and liver cirrhosis on cytochrome P450 enzyme activity in human  
33 liver microsomes and inducibility in cultured human hepatocytes. *Toxicol Appl*  
34 *Pharmacol*, 199, 193-209. <http://dx.doi.org/10.1016/j.taap.2004.01.010>
- 35 Parnell, M.; Koller, L.; Exon, J.; Arnzen, J. (1986). Trichloroacetic acid effects on rat liver  
36 peroxisomes and enzyme-altered foci. *Environ Health Perspect*, 69, 73-79.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/2880713>
- 38 Parrish, J. M.; Austin, E. W.; Stevens, D. K.; Kinder, D. H.; Bull, R. J. (1996). Haloacetate-  
39 induced oxidative damage to DNA in the liver of male B6C3F1 mice. *Toxicology*, 110,  
40 103-111. [http://dx.doi.org/10.1016/0300-483X\(96\)03342-2](http://dx.doi.org/10.1016/0300-483X(96)03342-2)
- 41 Parry, E. M.; Hague, A.; Parry, J. M. (1990). A study of mitotic division fidelity and numerical  
42 chromosome changes in ageing Syrian hamster dermal cells. *Mutat Res*, 237, 83-93.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/2366802>



DRAFT– DO NOT QUOTE OR CITE

- 1 Parry, J. M.; Parry, E. M.; Bourner, R.; Doherty, A.; Ellard, S.; O'Donovan, J., . . . Hess, P.  
2 (1996). The detection and evaluation of aneugenic chemicals. *Mutat Res*, 353, 11-46.  
3 [http://dx.doi.org/10.1016/0027-5107\(95\)00242-1](http://dx.doi.org/10.1016/0027-5107(95)00242-1)
- 4 Partanen, T.; Heikkila, P.; Hernberg, S.; Kauppinen, T.; Moneta, G.; Ojajarvi, A. (1991). Renal  
5 cell cancer and occupational exposure to chemical agents. *Scand J Work Environ Health*,  
6 17, 231-239. <http://www.ncbi.nlm.nih.gov/pubmed/1925434>
- 7 Pastino, G. M.; Yap, W. Y.; Carroquino, M. (2000). Human variability and susceptibility to  
8 trichloroethylene. *Environ Health Perspect*, 108 Suppl 2, 201-214.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/10807552>
- 10 Paykoc, Z. V. and Powell, J. F. (1945). The excretion of sodium trichloroacetate. *J Pharmacol*  
11 *Exp Ther*, 85, 289-293.
- 12 Peden-Adams, M. M.; Eudaly, J. G.; Lee, A. M.; Miller, J.; Keil, D. E.; Gilkeson, G. S. (2008).  
13 Lifetime exposure to trichloroethylene (TCE) does not accelerate autoimmune disease in  
14 MRL +/+ mice. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 43, 1402-1409.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/18780217>
- 16 Peden-Adams, M.; Eudaly, J.; Heesemann, L.; Smythe, J.; Miller, J.; Gilkeson, G.; Keil, D.  
17 (2006). Developmental immunotoxicity of trichloroethylene (TCE): studies in B6C3F1  
18 mice. *J Environ Sci Health A Tox Hazard Subst Environ Eng*, 41, 249-271.  
19 <http://dx.doi.org/10.1080/10934520500455289>
- 20 Pelekis, M.; Gephart, L. A.; Lerman, S. E. (2001). Physiological-model-based derivation of the  
21 adult and child pharmacokinetic intraspecies uncertainty factors for volatile organic  
22 compounds. *Regul Toxicol Pharmacol*, 33, 12-20.  
23 <http://dx.doi.org/10.1006/rtph.2000.1436>
- 24 Pellizzari, E. D.; Hartwell, T. D.; 3rd, H. B.; Waddell, R. D.; Whitaker, D. A.; Erickson, M. D.  
25 (1982). Purgeable organic compounds in mother's milk. *Bull Environ Contam Toxicol*,  
26 28, 322-328. <http://www.ncbi.nlm.nih.gov/pubmed/7082873>
- 27 Perbellini, L.; Olivato, D.; Zedde, A.; Miglioranza, R. (1991). Acute trichloroethylene poisoning  
28 by ingestion: clinical and pharmacokinetic aspects. *Intensive Care Med*, 17, 234-235.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/1744309>
- 30 Percy, C.; 3rd, S. E.; Gloeckler, L. (1981). Accuracy of cancer death certificates and its effect on  
31 cancer mortality statistics. *Am J Public Health*, 71, 242-250.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/7468855>
- 33 Percy, C.; Ries, L. G.; Van Holten, V. D. (1990). The accuracy of liver cancer as the underlying  
34 cause of death on death certificates. *Public Health Rep*, 105, 361-367.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/2116637>
- 36 Pereira, M. A. (1996). Carcinogenic activity of dichloroacetic acid and trichloroacetic acid in the  
37 liver of female B6C3F1 mice. *Fundam Appl Toxicol*, 31, 192-199.  
38 <http://dx.doi.org/10.1006/faat.1996.0091>
- 39 Pereira, M. A. and Phelps, J. B. (1996). Promotion by dichloroacetic acid and trichloroacetic acid  
40 of N-methyl-N-nitrosourea-initiated cancer in the liver of female B6C3F1 mice. *Cancer*  
41 *Lett*, 102, 133-141. [http://dx.doi.org/10.1016/0304-3835\(96\)04156-0](http://dx.doi.org/10.1016/0304-3835(96)04156-0)
- 42 Pereira, M. A.; Kramer, P. M.; Conran, P. B.; Tao, L. (2001). Effect of chloroform on  
43 dichloroacetic acid and trichloroacetic acid-induced hypomethylation and expression of  
44 the c-myc gene and on their promotion of liver and kidney tumors in mice.  
45 *Carcinogenesis*, 22, 1511-1519. <http://www.ncbi.nlm.nih.gov/pubmed/11532874>

DRAFT– DO NOT QUOTE OR CITE

- 1 Pereira, M. A.; Li, K.; Kramer, P. M. (1997). Promotion by mixtures of dichloroacetic acid and  
2 trichloroacetic acid of N-methyl-N-nitrosourea-initiated cancer in the liver of female  
3 B6C3F1 mice. *Cancer Lett*, 115, 15-23. [http://dx.doi.org/10.1016/S0304-3835\(97\)04699-](http://dx.doi.org/10.1016/S0304-3835(97)04699-5)  
4 5
- 5 Pereira, M. A.; Wang, W.; Kramer, P. M.; Tao, L. (2004). Prevention by methionine of  
6 dichloroacetic acid-induced liver cancer and DNA hypomethylation in mice. *Toxicol Sci*,  
7 77, 243-248. <http://dx.doi.org/10.1093/toxsci/kfh031>
- 8 Pereira, M.; Wang, W.; Kramer, P.; Tao, L. (2004). DNA hypomethylation induced by non-  
9 genotoxic carcinogens in mouse and rat colon. *Cancer Lett*, 212, 145-151.  
10 <http://dx.doi.org/10.1016/j.canlet.2004.03.024>
- 11 Perkins, N. D. (2004). NF-kappaB: tumor promoter or suppressor. *Trends Cell Biol*, 14, 64-69.  
12 <http://dx.doi.org/10.1016/j.tcb.2003.12.004>
- 13 Perkins, N. D. and Gilmore, T. D. (2006). Good cop, bad cop: the different faces of NF-kappaB.  
14 *Cell Death Differ*, 13, 759-772. <http://dx.doi.org/10.1038/sj.cdd.4401838>
- 15 Perocco, P. and Prodi, G. (1981). DNA damage by haloalkanes in human lymphocytes cultured  
16 in vitro. *Cancer Lett*, 13, 213-218. [http://dx.doi.org/10.1016/0304-3835\(81\)90020-3](http://dx.doi.org/10.1016/0304-3835(81)90020-3)
- 17 Perrot, N.; Nalpas, B.; Yang, C. S.; Beaune, P. H. (1989). Modulation of cytochrome P450  
18 isozymes in human liver, by ethanol and drug intake. *Eur J Clin Invest*, 19, 549-555.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/2515975>
- 20 Perry, S.; Schofield, M.; MacFarlane, M.; Lock, E.; King, L.; Gibson, G.; Goldfarb, P. (1993).  
21 Isolation and expression of a cDNA coding for rat kidney cytosolic cysteine conjugate  
22 beta-lyase. *Mol Pharmacol*, 43, 660-665. <http://www.ncbi.nlm.nih.gov/pubmed/8502223>
- 23 Persson, B. and Fredrikson, M. (1999). Some risk factors for non-Hodgkin's lymphoma. *Int J*  
24 *Occup Med Environ Health*, 12, 135-142.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/10465904>
- 26 Persson, B.; Dahlander, A. M.; Fredriksson, M.; Brage, H. N.; Ohlson, C. G.; Axelson, O.  
27 (1989). Malignant lymphomas and occupational exposures. *Br J Ind Med*, 46, 516-520.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/2775671>
- 29 Persson, B.; Fredriksson, M.; Olsen, K.; Boeryd, B.; Axelson, O. (1993). Some occupational  
30 exposures as risk factors for malignant lymphomas. *Cancer*, 72, 1773-1778.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/8348507>
- 32 Pesch, B.; Haerting, J.; Ranft, U.; Klimpel, A.; Oelschlagel, B.; Schill, W. (2000). Occupational  
33 risk factors for urothelial carcinoma: agent-specific results from a case-control study in  
34 Germany. MURC Study Group. Multicenter Urothelial and Renal Cancer. *Int J*  
35 *Epidemiol*, 29, 238-247.
- 36 Pesch, B.; Haerting, J.; Ranft, U.; Klimpel, A.; Oelschlagel, B.; Schill, W. (2000). Occupational  
37 risk factors for renal cell carcinoma: Agent-specific results from a case-control study in  
38 Germany. *Int J Epidemiol*, 29, 1014-1024. <http://dx.doi.org/10.1093/ije/29.6.1014>
- 39 Peter, R.; Böcker, R.; Beaune, P.; Iwasaki, M.; Guengerich, F.; Yang, C. (1990). Hydroxylation  
40 of chlorzoxazone as a specific probe for human liver cytochrome P-450IIE1. *Chem Res*  
41 *Toxicol*, 3, 566-573. <http://www.ncbi.nlm.nih.gov/pubmed/2103328>
- 42 Peters, J. M.; Cattley, R. C.; Gonzalez, F. J. (1997). Role of PPAR alpha in the mechanism of  
43 action of the nongenotoxic carcinogen and peroxisome proliferator Wy-14,643.  
44 *Carcinogenesis*, 18, 2029-2033.

DRAFT– DO NOT QUOTE OR CITE

- 1 Peters, J. M.; Preston-Martin, S.; Yu, M. C. (1981). Brain tumors of children and occupational  
2 exposures of parents. *Science*, 213, 235-237.
- 3 Peters, J.; Garabrant, D.; Wright, W.; Bernstein, L.; Mack, T. (1985). Uses of a cancer registry in  
4 the assessment of occupational cancer risks. 69, 157-161.  
5 <http://www.ncbi.nlm.nih.gov/pubmed/3834325>
- 6 Pfaffenberger, C. D.; Peoples, A. J.; Enos, H. F. (1980). Distribution of volatile halogenated  
7 organic compounds between rat blood serum and adipose tissue. *Int J Environ Anal*  
8 *Chem*, 8, 55-65. <http://www.ncbi.nlm.nih.gov/pubmed/7399799>
- 9 Phillips, T. and Macdonald, R. (1971). Comparative effect of pethidine, trichloroethylene, and  
10 Entonox on fetal and neonatal acid-base and PO<sub>2</sub>. *Br Med J*, 3, 558-560.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/5571451>
- 12 Piegorsch, W. W. and Bailer, A. J. (1997). *Statistics for environmental biology and toxicology*.  
13 London, England: Chapman & Hall.
- 14 Pikarsky, E.; Porat, R. M.; Stein, I.; Abramovitch, R.; Amit, S.; Kasem, S., . . . Ben-Neriah, Y.  
15 (2004). NF-kappaB functions as a tumour promoter in inflammation-associated cancer.  
16 *Nature*, 431, 461-466. <http://dx.doi.org/10.1038/nature02924>
- 17 Plewa, M. J.; Kargalioglu, Y.; Vanker, D.; Minear, R. A.; Wagner, E. D. (2002). Mammalian  
18 cell cytotoxicity and genotoxicity analysis of drinking water disinfection by-products.  
19 *Environ Mol Mutagen*, 40, 134-142. <http://dx.doi.org/10.1002/em.10092>
- 20 Poet, T.; Corley, R.; Thrall, K.; Edwards, J.; Tanojo, H.; Weitz, K., . . . Wester, R. (2000).  
21 Assessment of the percutaneous absorption of trichloroethylene in rats and humans using  
22 MS/MS real-time breath analysis and physiologically based pharmacokinetic modeling.  
23 *Toxicol Sci*, 56, 61-72. <http://www.ncbi.nlm.nih.gov/pubmed/10869454>
- 24 Pogribny, I. P.; Tryndyak, V. P.; Woods, C. G.; Witt, S. E.; Rusyn, I. (2007). Epigenetic effects  
25 of the continuous exposure to peroxisome proliferator WY-14,643 in mouse liver are  
26 dependent upon peroxisome proliferator activated receptor alpha. *Mutat Res-Fundam*  
27 *Mol Mech Mutagen*, 625, 62-71. <http://dx.doi.org/10.1016/j.mrfmmm.2007.05.004>
- 28 Pohl, H. R.; Roney, N.; Wilbur, S.; Hansen, H.; De Rosa, C. T. (2003). Six interaction profiles  
29 for simple mixtures. *Chemosphere*, 53, 183-197.
- 30 Pollack, G. M.; Li, R. C.; Ermer, J. C.; Shen, D. D. (1985). Effects of route of administration and  
31 repetitive dosing on the disposition kinetics of di(2-ethylhexyl) phthalate and its mono-  
32 de-esterified metabolite in rats. *Toxicol Appl Pharmacol*, 79, 246-256.  
33 [http://dx.doi.org/10.1016/0041-008X\(85\)90346-1](http://dx.doi.org/10.1016/0041-008X(85)90346-1)
- 34 Popper, H. (1986). Aging and the liver. In H. Popper & F. Schaffner (Eds.), *Progress in liver*  
35 *diseases: Volume VIII* (pp. 659–683). Orlando, FL: Grune and Stratton.
- 36 Porru, S.; Placidi, D.; Carta, A.; Gelatti, U.; Ribero, M. L.; Tagger, A., . . . Donato, F. (2001).  
37 Primary liver cancer and occupation in men: a case-control study in a high-incidence area  
38 in Northern Italy. *Int J Cancer*, 94, 878-883.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/11745492>
- 40 Porter, R. (1993). Follow-up of childhood leukemia incidence rate in Maricopa County  
41 standardized incidence ratio of area F to area R for 1987.
- 42 Portier, C. J. and Bailer, A. J. (1989). Testing for increased carcinogenicity using a survival-  
43 adjusted quantal response test. *Toxicol Sci*, 12, 731-737.

DRAFT– DO NOT QUOTE OR CITE

- 1 Portier, C. J.; Hedges, J. C.; Hoel, D. G. (1986). Age-specific models of mortality and tumor  
2 onset for historical control animals in the National Toxicology Program's carcinogenicity  
3 experiments. *Cancer Res*, 46, 4372-4378.
- 4 Povey, A.; Guppy, M.; Wood, M.; Knight, C.; Black, C.; Silman, A. (2001). Cytochrome P2  
5 polymorphisms and susceptibility to scleroderma following exposure to organic solvents.  
6 *Arthritis Rheum*, 44, 662-665. [http://dx.doi.org/10.1002/1529-  
7 0131\(200103\)44:3<662::AID-ANR115>3.0.CO;2-P](http://dx.doi.org/10.1002/1529-0131(200103)44:3<662::AID-ANR115>3.0.CO;2-P)
- 8 Powell, J. F. (1945). Trichlorethylene: absorption, elimination and metabolism. *Occup Environ*  
9 *Med*, 2, 142-145.
- 10 Prasanna, H. R.; Hart, R. W.; Magee, P. N. (1989). Effect of food restriction on the metabolism  
11 of dimethylnitrosamine (NDMA) in rats. *Anticancer Res*, 9, 93-96.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/2705761>
- 13 Pravecek, T. L.; Channel, S. R.; Schmidt, W. J.; Kidney, J. K. (1996). Cytotoxicity and  
14 metabolism of dichloroacetic acid and trichloroacetic acid in B6C3F1 mouse liver tissue.  
15 *Arthritis Rheum*, 39, 261-269.
- 16 Prendergast, J. A.; Jones, R. A.; Jenkins, L. J. J.; Siegel, J. (1967). Effects on experimental  
17 animals of long-term inhalation of trichloroethylene, carbon tetrachloride, 1,1,1-  
18 trichloroethane, dichlorodifluoromethane, and 1,1-dichloroethylene. *Toxicol Appl*  
19 *Pharmacol*, 10, 270-289. <http://www.ncbi.nlm.nih.gov/pubmed/4962252>
- 20 Price, P. J.; Hassett, C. M.; Mansfield, J. I. (1978). Transforming activities of trichloroethylene  
21 and proposed industrial alternatives. *In Vitro Cell Dev Biol Anim*, 14, 290-293.
- 22 Price, R. G.; Taylor, S. A.; Chivers, I.; Arce-Tomas, M.; Crutcher, E.; Franchini, I., . . .  
23 Eisenberger, U. (1996). Development and validation of new screening tests for  
24 nephrotoxic effects. *Hum Exp Toxicol*, 1, S10-S19.
- 25 Price, R. G.; Wedeen, R.; Lichtveld, M. Y.; Lybarger, J. A.; Porter, G. A.; Fels, L., . . . Finn, W.  
26 F. (1999). Urinary biomarkers: Roles in risk assessment to environmental and  
27 occupational nephrotoxins: Monitoring of effects and evaluation of mechanisms of  
28 toxicity. *Ren Fail*, 21, xiii-xviii.
- 29 Prout, M. S.; Provan, W. M.; Green, T. (1985). Species differences in response to  
30 trichloroethylene: I pharmacokinetics in rats and mice. *Toxicol Appl Pharmacol*, 79, 389-  
31 400. <http://www.ncbi.nlm.nih.gov/pubmed/4035686>
- 32 Pryor, G. T.; Dickinson, J.; Howd, R. A.; Rebert, C. S. (1983). Transient cognitive deficits and  
33 high-frequency hearing loss in weanling rats exposed to toluene. *Neurotoxicol Teratol*, 5,  
34 53-57.
- 35 Pryor, G. T.; Rebert, C. S.; Howd, R. A. (1987). Hearing loss in rats caused by inhalation of  
36 mixed xylenes and styrene. *J Appl Toxicol*, 7, 55-61.
- 37 Purdue, M. P.; Severson, R. K.; Colt, J. S.; Stewart, P.; De Roos, A. J.; Cerhan, J. R., . . . Blair,  
38 A. (2009). Degreasing and risk of non-Hodgkin lymphoma. *Occup Environ Med*, 66,  
39 557-560. <http://dx.doi.org/10.1136/oem.2008.040386>
- 40 Purdue, M.; Bakke, B.; Stewart, P.; De Roos, A.; Schenk, M.; Lynch, C., . . . Colt, J. (2011). A  
41 case-control study of occupational exposure to trichloroethylene and non-Hodgkin  
42 lymphoma. *Environ Health Perspect*, 119, 232-238.  
43 <http://dx.doi.org/10.1289/ehp.1002106>
- 44 Raaschou-Nielsen, O.; Hansen, J.; Christensen, J.; Blot, W.; McLaughlin, J.; Olsen, J. (2001).  
45 Urinary concentrations of trichloroacetic acid in Danish workers exposed to

- 1 trichloroethylene, 1947-1985. *Am J Ind Med*, 39, 320-327.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/11241565>
- 3 Raaschou-Nielsen, O.; Hansen, J.; McLaughlin, J.; Kolstad, H.; Christensen, J.; Tarone, R.;  
4 Olsen, J. (2003). Cancer risk among workers at Danish companies using  
5 trichloroethylene: A cohort study. *Am J Epidemiol*, 158, 1182-1192.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/14652303>
- 7 Raaschou-Nielsen, O.; Hansen, J.; Thomsen, B.; Johansen, I.; Lipworth, L.; McLaughlin, J.;  
8 Olsen, J. (2002). Exposure of Danish workers to trichloroethylene, 1947-1989. *Appl*  
9 *Occup Environ Hyg*, 17, 693-703. <http://dx.doi.org/10.1080/10473220290096186>
- 10 Radican, L.; Blair, A.; Stewart, P.; Wartenberg, D. (2008). Mortality of aircraft maintenance  
11 workers exposed to trichloroethylene and other hydrocarbons and chemicals: extended  
12 follow-up. *J Occup Environ Med*, 50, 1306-1319.  
13 <http://dx.doi.org/10.1097/JOM.0b013e3181845f7f>
- 14 Radican, L.; Wartenberg, D.; Rhoads, G.; Schneider, D.; Wedeen, R.; Stewart, P.; Blair, A.  
15 (2006). A retrospective occupational cohort study of end-stage renal disease in aircraft  
16 workers exposed to trichloroethylene and other hydrocarbons. *J Occup Environ Med*, 48,  
17 1-12. <http://www.ncbi.nlm.nih.gov/pubmed/16404204>
- 18 Raijmakers, M. T. M.; Steegers, E. A. P.; Peters, W. H. M. (2001). Glutathione S-transferases  
19 and thiol concentrations in embryonic and early fetal tissues. *Hum Reprod*, 16, 2445-  
20 2450. <http://www.ncbi.nlm.nih.gov/pubmed/11679536>
- 21 Rake, J. P.; Visser, G.; Labrune, P.; Leonard, J. V.; Ullrich, K.; Smit, G. P. (2002). Glycogen  
22 storage disease type I: diagnosis, management, clinical course and outcome. Results of  
23 the European Study on Glycogen Storage Disease Type I (ESGSD I). *Eur J Pediatr*, 161  
24 Suppl 1, S20-S34. <http://dx.doi.org/10.1007/s00431-002-0999-4>
- 25 Rakoff-Nahoum, S. and Medzhitov, R. (2007). Regulation of spontaneous intestinal  
26 tumorigenesis through the adaptor protein MyD88. *Science*, 317, 124-127.  
27 <http://dx.doi.org/10.1126/science.1140488>
- 28 Rakyan, V. K.; Blewitt, M. E.; Druker, R.; Preis, J. I.; Whitelaw, E. (2002). Metastable epialleles  
29 in mammals. *Trends in Genetics*, 18, 348-351.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/12127774>
- 31 Rakyan, V. K.; Chong, S.; Champ, M. E.; Cuthbert, P. C.; Morgan, H. D.; Luu, K. V.; Whitelaw,  
32 E. (2003). Transgenerational inheritance of epigenetic states at the murine Axin(Fu)  
33 allele occurs after maternal and paternal transmission. *PNAS*, 100, 2538-2543.  
34 <http://dx.doi.org/10.1073/pnas.0436776100>
- 35 Ramanakumar, A. V.; Nadon, L.; Siemiatycki, J. (2008). Exposures in painting related  
36 occupations and risk of selected cancers: results from a case-control study in Montreal.  
37 *Am J Ind Med*, 51, 419-427. <http://dx.doi.org/10.1002/ajim.20564>
- 38 Ramdhan, D. H.; Kamijima, M.; Wang, D.; Ito, Y.; Naito, H.; Yanagiba, Y., . . . Nakajima, T.  
39 (2010). Differential response to trichloroethylene-induced hepatosteatosis in wild-type  
40 and PPARalpha-humanized mice. *Environ Health Perspect*, 118, 1557-1563.  
41 <http://dx.doi.org/10.1289/ehp.1001928>
- 42 Ramdhan, D. H.; Kamijima, M.; Yamada, N.; Ito, Y.; Yanagiba, Y.; Nakamura, D., . . .  
43 Nakajima, T. (2008). Molecular mechanism of trichloroethylene-induced hepatotoxicity  
44 mediated by CYP2E1. *Toxicol Appl Pharmacol*, 231, 300-307.  
45 <http://dx.doi.org/10.1016/j.taap.2008.04.020>

DRAFT– DO NOT QUOTE OR CITE

- 1 Ramsdell, A. and Markwald, R. (1997). Induction of endocardial cushion tissue in the avian  
2 heart is regulated, in part, by TGFbeta-3-mediated autocrine signaling. *Dev Biol*, 188, 64-  
3 74. <http://dx.doi.org/10.1006/dbio.1997.8637>
- 4 Ramsey, J. C. and Andersen, M. E. (1984). A physiologically based description of the inhalation  
5 pharmacokinetics of styrene in rats and humans. *Toxicol Appl Pharmacol*, 73, 159-175.  
6 [http://dx.doi.org/10.1016/0041-008X\(84\)90064-4](http://dx.doi.org/10.1016/0041-008X(84)90064-4)
- 7 Rao, M. S.; Tatematsu, M.; Subbarao, V.; Ito, N.; Reddy, J. K. (1986). Analysis of peroxisome  
8 proliferator-induced preneoplastic and neoplastic lesions of rat liver for placental form of  
9 glutathione S-transferase and gamma-glutamyltranspeptidase. *Cancer Res*, 46, 5287-  
10 5290. <http://www.ncbi.nlm.nih.gov/pubmed/2875789>
- 11 Rapson, W. H.; Nazar, M. A.; Butsky, V. V. (1980). Mutagenicity produced by aqueous  
12 chlorination of organic compounds. *Bull Environ Contam Toxicol*, 24, 590-596.  
13 <http://dx.doi.org/10.1007/BF01608160>
- 14 Rasmussen, K. and Sabroe, S. (1986). Neuropsychological symptoms among metal workers  
15 exposed to halogenated hydrocarbons. *Scand J Public Health*, 14, 161-168.  
16 <http://dx.doi.org/10.1177/140349488601400309>
- 17 Rasmussen, K.; Arlien-Søborg, P.; Sabroe, S. (1993). Clinical neurological findings among metal  
18 degreasers exposed to chlorinated solvents. *Acta Neurol Scand*, 87, 200-204.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/8475690>
- 20 Rasmussen, K.; Brogren, C.-H.; Sabroe, S. (1993). Subclinical affection of liver and kidney  
21 function and solvent exposure. *Int Arch Occup Environ Health*, 64, 445-448.  
22 <http://dx.doi.org/10.1007/BF00517951>
- 23 Rasmussen, K.; Jeppesen, H. J.; Sabroe, S. (1993). Solvent-induced chronic toxic  
24 encephalopathy. *Am J Ind Med*, 23, 779-792. <http://dx.doi.org/10.1002/ajim.4700230511>
- 25 Rasmussen, K.; Jeppesen, H.; Sabroe, S. (1993). Psychometric tests for assessment of brain  
26 function after solvent exposure. *Am J Ind Med*, 24, 553-565.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/8266931>
- 28 Rasmussen, K.; Sabroe, S.; Wohler, M.; Ingerslev, H. J.; Kappel, B.; Nielsen, J. (1988). A  
29 genotoxic study of metal workers exposed to trichloroethylene Sperm parameters and  
30 chromosome aberrations in lymphocytes. *Int Arch Occup Environ Health*, 60, 419-423.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/3410552>
- 32 Raunio, H.; Hakkola, J.; Hukkanen, J.; Pelkonen, O.; Edwards, R.; Boobis, A.; Anttila, S. (1998).  
33 Expression of xenobiotic metabolising cytochrome P450's in human pulmonary tissue.  
34 *Arch Toxicol Suppl*, 20, 465-469.
- 35 Ray, W. A.; Griffin, M. R.; Schaffner, W. (1987). Psychotropic drug use and the risk of hip  
36 fracture. *316/7* (363-369).
- 37 Reamon-Buettner, S. M. and Borlak, J. (2007). A new paradigm in toxicology and teratology:  
38 altering gene activity in the absence of DNA sequence variation. *Reprod Toxicol*, 24, 20-  
39 30. <http://dx.doi.org/10.1016/j.reprotox.2007.05.002>
- 40 Rebert, C. S.; Boyes, W. K.; Pryor, G. T.; Svendsgaard, D. J.; Kassay, K. M.; Gordon, G. R.;  
41 Shinsky, N. (1993). Combined effects of solvents on the rat's auditory system: styrene  
42 and trichloroethylene. *Int J Psychophysiol*, 14, 49-59.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/8432679>

DRAFT– DO NOT QUOTE OR CITE

- 1 Rebert, C. S.; Day, V. L.; Matteucci, M. J.; Pryor, G. T. (1991). Sensory-evoked potentials in  
2 rats chronically exposed to trichloroethylene: predominant auditory dysfunction.  
3 *Neurotoxicol Teratol*, 13, 83-90. <http://www.ncbi.nlm.nih.gov/pubmed/2046630>
- 4 Rebert, C. S.; Schwartz, R. W.; Svendsgaard, D. J.; Pryor, G. T.; Boyes, W. K. (1995).  
5 Combined effects of paired solvents on the rat's auditory system. *Toxicology*, 105, 345-  
6 354. <http://www.ncbi.nlm.nih.gov/pubmed/8571371>
- 7 Reif, J.; Burch, J.; Nuckols, J.; Metzger, L.; Ellington, D.; Anger, W. (2003). Neurobehavioral  
8 effects of exposure to trichloroethylene through a municipal water supply. *Environ Res*,  
9 93, 248-258. <http://www.ncbi.nlm.nih.gov/pubmed/14615234>
- 10 Rennix, C. P.; Quinn, M. M.; Amoroso, P. J.; Eisen, E. A.; Wegman, D. H. (2005). Risk of breast  
11 cancer among enlisted Army women occupationally exposed to volatile organic  
12 compounds. *Am J Ind Med*, 48, 157-167.
- 13 Reznik, G.; Ward, J. M.; Hardisty, J. F.; Russfield, A. (1979). Renal carcinogenic and  
14 nephrotoxic effects of the flame retardant tris(2,3-dibromopropyl) phosphate in F344 rats  
15 and (C57BL/6N X C3H/HeN)F1 mice. *J Natl Cancer Inst*, 63, 205-212.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/286830>
- 17 Rhee, I.; Bachman, K.; Park, B.; Jair, K.; Yen, R.; Schuebel, K., . . . Vogelstein, B. (2002).  
18 DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature*, 416,  
19 552-556. <http://dx.doi.org/10.1038/416552a>
- 20 Rhomberg, L. R. (2000). Dose-response analyses of the carcinogenic effects of trichloroethylene  
21 in experimental animals. *Environ Health Perspect*, 108, 343-358.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/10807564>
- 23 Rice, D. C. and Barone, S., Jr. (2000). Critical periods of vulnerability for the developing  
24 nervous system: Evidence from humans and animal models. *Environ Health Perspect*,  
25 108, 511-533. <http://www.ncbi.nlm.nih.gov/pubmed/10852851>
- 26 Richard, A. and Hunter, E. (1996). Quantitative structure-activity relationships for the  
27 developmental toxicity of haloacetic acids in mammalian whole embryo culture.  
28 *Teratology*, 53, 352-360. [http://dx.doi.org/10.1002/\(SICI\)1096-  
29 9926\(199606\)53:6<352::AID-TERA6>3.0.CO;2-1](http://dx.doi.org/10.1002/(SICI)1096-9926(199606)53:6<352::AID-TERA6>3.0.CO;2-1)
- 30 Richardson, D. B.; Terschüren, C.; Hoffmann, W. (2008). Occupational risk factors for non-  
31 Hodgkin's lymphoma: a population-based case-control study in Northern Germany. *Am J*  
32 *Ind Med*, 51, 258-268. <http://dx.doi.org/10.1002/ajim.20552>
- 33 Richardson, F. C.; Boucheron, J. A.; Dyroff, M. C.; Popp, J. A.; Swenberg, J. A. (1986).  
34 Biochemical and morphologic studies of heterogeneous lobe responses in  
35 hepatocarcinogenesis. *Carcinogenesis*, 7, 247-251.  
36 <http://www.ncbi.nlm.nih.gov/pubmed/2868806>
- 37 Richmond, R. E.; Carter, J. H.; Carter, H. W.; Daniel, F. B.; DeAngelo, A. B. (1995).  
38 Immunohistochemical analysis of dichloroacetic acid (DCA)-induced  
39 hepatocarcinogenesis in male Fischer (F344) rats. *Cancer Lett*, 92, 67-76.  
40 [http://dx.doi.org/10.1016/0304-3835\(94\)03756-9](http://dx.doi.org/10.1016/0304-3835(94)03756-9)
- 41 Riederer, F.; Luborzewski, A.; God, R.; Bringmann, G.; Scholz, J.; Feineis, D.; Moser, A.  
42 (2002). Modification of tyrosine hydroxylase activity by chloral derived beta-carbolines  
43 in vitro. *J Neurochem*, 81, 814-819. <http://www.ncbi.nlm.nih.gov/pubmed/12065640>

DRAFT– DO NOT QUOTE OR CITE

- 1 Ripp, S. L.; Itagaki, K.; Philpot, R. M.; Elfarra, A. A. (1999). Species and sex differences in  
2 expression of flavin-containing monooxygenase form 3 in liver and kidney microsomes.  
3 *Drug Metab Dispos*, 27, 46-52. <http://www.ncbi.nlm.nih.gov/pubmed/9884308>
- 4 Ripp, S. L.; Overby, L. H.; Philpot, R. M.; Elfarra, A. A. (1997). Oxidation of cysteine S-  
5 conjugates by rabbit liver microsomes and cDNA-expressed flavin-containing mono-  
6 oxygenases: studies with S-(1,2-dichlorovinyl)-L-cysteine, S-(1,2,2-trichlorovinyl)-L-  
7 cysteine, S-allyl-L-cysteine, and S-benzyl-L-cysteine. *Mol Pharmacol*, 51, 507-515.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/9058607>
- 9 Ristoff, E. and Larsson, A. (2007). Inborn errors in the metabolism of glutathione. *Orphanet J*  
10 *Rare Dis*, 2, 16. <http://dx.doi.org/10.1186/1750-1172-2-16>
- 11 Ritz, B. (1999). Cancer mortality among workers exposed to chemicals during uranium  
12 processing. *J Occup Environ Med*, 41, 556-566.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/10412097>
- 14 Ritz, B. (1999). Radiation exposure and cancer mortality in uranium processing workers.  
15 *Epidemiology*, 10, 531-538. <http://www.ncbi.nlm.nih.gov/pubmed/10468427>
- 16 Ritz, B.; Morgenstern, H.; Crawford-Brown, D.; Young, B. (2000). The effects of internal  
17 radiation exposure on cancer mortality in nuclear workers at Rocketdyne/Atomics  
18 International. *Environ Health Perspect*, 108, 743-751.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/10964795>
- 20 Ritz, B.; Morgenstern, H.; Froines, J.; Young, B. B. (1999). Effects of exposure to external  
21 ionizing radiation on cancer mortality in nuclear workers monitored for radiation at  
22 Rocketdyne/Atomics International. *Am J Ind Med*, 35, 21-31.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/9884742>
- 24 Robbiano, L.; Baroni, D.; Carrozzino, R.; Mereto, E.; Brambilla, G. (2004). DNA damage and  
25 micronuclei induced in rat and human kidney cells by six chemicals carcinogenic to the  
26 rat kidney. *Toxicology*, 204, 187-195. <http://dx.doi.org/10.1016/j.tox.2004.06.057>
- 27 Robbiano, L.; Mereto, E.; Migliazzi Morando, A.; Pastore, P.; Brambilla, G. (1998). Increased  
28 frequency of micronucleated kidney cells in rats exposed to halogenated anaesthetics.  
29 *Mutat Res*, 413, 1-6. <http://www.ncbi.nlm.nih.gov/pubmed/9602852>
- 30 Robins, J. (1987). A graphical approach to the identification and estimation of causal parameters  
31 in mortality studies with sustained exposure periods. *J Clin Epidemiol*, 2, 139S-161S.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/3667861>
- 33 Rodenbeck, S.; Sanderson, L.; Rene, A. (2000). Maternal exposure to trichloroethylene in  
34 drinking water and birth-weight outcomes. *Arch Environ Health*, 55, 188-194.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/10908102>
- 36 Rodriguez, C. E.; Mahle, D. A.; Gearhart, J. M.; Mattie, D. R.; Lipscomb, J. C.; Cook, R. S.;  
37 Barton, H. A. (2007). Predicting age-appropriate pharmacokinetics of six volatile organic  
38 compounds in the rat utilizing physiologically based pharmacokinetic modeling. *Toxicol*  
39 *Sci*, 98, 43-56. <http://dx.doi.org/10.1093/toxsci/kfm082>
- 40 Rosenbaum, A. (2005). The HAPEM5 user's guide: Hazardous air pollutant exposure model,  
41 Version 5. San Francisco, CA: ICF Consulting.
- 42 Roskams, T. A.; Theise, N. D.; Balabaud, C.; Bhagat, G.; Bhathal, P. S.; Bioulac-Sage, P., . . .  
43 West, A. B. (2004). Nomenclature of the finer branches of the biliary tree: canals,  
44 ductules, and ductular reactions in human livers. *Hepatology*, 39, 1739-1745.  
45 <http://dx.doi.org/10.1002/hep.20130>



DRAFT– DO NOT QUOTE OR CITE

- 1 Roskams, T.; Yang, S. Q.; Koteish, A.; Durnez, A.; DeVos, R.; Huang, X., . . . Diehl, A. M.  
2 (2003). Oxidative stress and oval cell accumulation in mice and humans with alcoholic  
3 and nonalcoholic fatty liver disease. *Am J Pathol*, 163, 1301-1311.  
4 [http://dx.doi.org/10.1016/S0002-9440\(10\)63489-X](http://dx.doi.org/10.1016/S0002-9440(10)63489-X)
- 5 Rossi, A. M.; Migliore, L.; Barale, R.; Loprieno, N. (1983). In vivo and in vitro mutagenicity  
6 studies of a possible carcinogen, trichloroethylene, and its two stabilizers,  
7 epichlorohydrin and 1,2-epoxybutane. *Birth Defects Res B Dev Reprod Toxicol*, 3, 75-  
8 87.
- 9 Rothman, K. J. (1986). *Modern epidemiology*. New York, NY: Little Brown & Co.
- 10 Rothman, K. J. and Greenland, S. (1998). *Modern epidemiology* (2nd ed.). Philadelphia, PA:  
11 Lippincott-Raven Publishers.
- 12 Rothman, K. J.; Greenland, S.; Lash, T. L. (2008). Case-control studies, chapter 8 *Modern*  
13 *epidemiology* (3rd ed., pp. 111-127). Philadelphia, PA: Lippincott Williams & Wilkins.
- 14 Ruckenstein, M. J. (2004). Autoimmune inner ear disease. *Curr Opin Otolaryngol Head Neck*  
15 *Surg*, 12, 426-430.
- 16 Rufer, E. S.; Hacker, T. A.; Flentke, G. R.; Drake, V. J.; Brody, M. J.; Lough, J.; Smith, S. M.  
17 (2010). Altered cardiac function and ventricular septal defect in avian embryos exposed  
18 to low-dose trichloroethylene. *Toxicol Sci*, 113, 444-452.  
19 <http://dx.doi.org/10.1093/toxsci/kfp269>
- 20 Rufer, E. S.; Hacker, T.; Lough, J.; Smith, S. M. (2008). Low-dose trichloroethylene exposure  
21 during valvuloseptal morphogenesis causes ventricular septal defects in hatched chicks.  
22 *Toxicologist*, 102, 314.
- 23 Ruijten, M. W. M. M.; Verberk, M. M.; Salle, H. J. A. (1991). Nerve function in workers with  
24 long term exposure to trichloroethene. *Occup Environ Med*, 48, 87-92.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/1998613>
- 26 Russo, A. and Levis, A. (1992). Detection of aneuploidy in male germ cells of mice by means of  
27 a meiotic micronucleus assay. *Mutat Res*, 281, 187-191.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/1371841>
- 29 Russo, A. and Levis, A. G. (1992). Further evidence for the aneuploidogenic properties of  
30 chelating agents: induction of micronuclei in mouse male germ cells by EDTA. *Environ*  
31 *Mol Mutagen*, 19, 125-131. <http://www.ncbi.nlm.nih.gov/pubmed/1541253>
- 32 Russo, A.; Pacchierotti, F.; Metalli, P. (1984). Nondisjunction induced in mouse spermatogenesis  
33 by chloral hydrate, a metabolite of trichloroethylene. *Environ Mutagen*, 6, 695-703.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/6479114>
- 35 Russo, A.; Stocco, A.; Majone, F. (1992). Identification of kinetochore-containing (CREST+)  
36 micronuclei in mouse bone marrow erythrocytes. *Mutagenesis*, 7, 195-197.
- 37 Rusyn, I.; Bradham, C. A.; Cohn, L.; Schoonhoven, R.; Swenberg, J. A.; Brenner, D. A.;  
38 Thurman, R. G. (1999). Corn oil rapidly activates nuclear factor-kappaB in hepatic  
39 Kupffer cells by oxidant-dependent mechanisms. *Carcinogenesis*, 20, 2095-2100.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/10545411>
- 41 Rusyn, I.; Peters, J. M.; Cunningham, M. L. (2006). Modes of action and species-specific effects  
42 of di-(2-ethylhexyl)phthalate in the liver. *Crit Rev Toxicol*, 36, 459-479.  
43 <http://dx.doi.org/10.1080/10408440600779065>

DRAFT– DO NOT QUOTE OR CITE

- 1 Sabel, G. V. and Clark, T. P. (1984). Volatile organic compounds as indicators of municipal  
2 solid waste leachate contamination. 2, 119-130. (as cited in U.S. EPA 2001a in NRC  
3 2006).
- 4 Sagawa, K.; Nishitani, H.; Kawai, H.; Kuge, Y.; Ikeda, M. (1973). Transverse lesion of spinal  
5 cord after accidental exposure to trichloroethylene. *Int Arch Arbeitsmed*, 31, 257-264.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/4767703>
- 7 Saghir, S. and Schultz, I. (2002). Low-dose pharmacokinetics and oral bioavailability of  
8 dichloroacetate in naive and GST-zeta-depleted rats. *Environ Health Perspect*, 110, 757-  
9 763. <http://www.ncbi.nlm.nih.gov/pubmed/12153755>
- 10 Saihan, E.; Burton, J.; Heaton, K. (1978). A new syndrome with pigmentation, scleroderma,  
11 gynaecomastia, Raynaud's phenomenon and peripheral neuropathy. *Br J Dermatol*, 99,  
12 437-440. <http://www.ncbi.nlm.nih.gov/pubmed/213095>
- 13 Saillenfait, A. M.; Langonne, I.; Sabate, J. P. (1995). Developmental toxicity of  
14 trichloroethylene, tetrachloroethylene and four of their metabolites in rat whole embryo  
15 culture. *Arch Toxicol*, 70, 71-82. <http://dx.doi.org/10.1007/BF02733666>
- 16 Sakurai, T.; Maeda, S.; Chang, L.; Karin, M. (2006). Loss of hepatic NF-kappa B activity  
17 enhances chemical hepatocarcinogenesis through sustained c-Jun N-terminal kinase 1  
18 activation. *PNAS*, 103, 10544-10551. <http://dx.doi.org/10.1073/pnas.0603499103>
- 19 Sallmen, M.; Lindbohm, M. L.; Anttila, A.; Kyyronen, P.; Taskinen, H.; Nykyri, E.; Hemminki,  
20 K. (1998). Time to pregnancy among the wives of men exposed to organic solvents.  
21 *Occup Environ Med*, 55, 24-30. <http://www.ncbi.nlm.nih.gov/pubmed/9536159>
- 22 Sallmén, M.; Lindbohm, M. L.; Kyyrönen, P.; Nykyri, E.; Anttila, A.; Taskinen, H.; Hemminki,  
23 K. (1995). Reduced fertility among women exposed to organic solvents. *Am J Ind Med*,  
24 27, 699-713. <http://dx.doi.org/10.1002/ajim.4700270506>
- 25 Salvan, A. (1990). Occupational exposure and cancer mortality at an electrical manufacturing  
26 plant: A case-control study. University of California, Los Angeles, CA.
- 27 Salvan, A.; Stayner, L.; Steenland, K.; Smith, R. (1995). Selecting an exposure lag period.  
28 *Epidemiology*, 6, 387-390. <http://www.ncbi.nlm.nih.gov/pubmed/7548346>
- 29 Salvini, M.; Binaschi, S.; Riva, M. (1971). Evaluation of the psychophysiological functions in  
30 humans exposed to trichloroethylene. *Occup Environ Med*, 28, 293-295.  
31 <http://www.ncbi.nlm.nih.gov/pubmed/5557850>
- 32 Sanchez, I. M. and Bull, R. J. (1990). Early induction of reparative hyperplasia in the liver of  
33 B6C3F1 mice treated with dichloroacetate and trichloroacetate. *Toxicology*, 64, 33-46.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/2219131>
- 35 Sanders, V. M.; Tucker, A. N.; White, K. L. J.; Kauffmann, B. M.; Hallett, P.; Carchman, R. A., .  
36 . . Munson, A. E. (1982). Humoral and cell-mediated immune status in mice exposed to  
37 trichloroethylene in the drinking water. *Toxicol Appl Pharmacol*, 62, 358-368.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/7071855>
- 39 Sanders, V.; Kauffmann, B.; White, K.; Douglas, K.; Barnes, D.; Sain, L., . . . Munson, A.  
40 (1982). Toxicology of chloral hydrate in the mouse. *Environ Health Perspect*, 44, 137-  
41 146. <http://www.ncbi.nlm.nih.gov/pubmed/7084146>
- 42 Santibanez, M.; Vioque, J.; Alguacil, J.; Barber, X.; de la Hera, G.; Kauppinen, T. (2008).  
43 Occupational exposures and risk of oesophageal cancer by histological type: a case-  
44 control study in eastern Spain. *Occup Environ Med*, 65, 774-781.  
45 <http://dx.doi.org/10.1136/oem.2007.037929>

DRAFT– DO NOT QUOTE OR CITE

- 1 Sanz, N.; Díez-Fernández, C.; Cascales, M. (1996). Variations of hepatic antioxidant systems  
2 and DNA ploidy in rats aged 2 to 8 months. *Biochim Biophys Acta*, 1315, 123-130.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/8608169>
- 4 Sapkota, A.; Williams, D.; Buckley, T. J. (2005). Tollbooth workers and mobile source-related  
5 hazardous air pollutants: how protective is the indoor environment? *Environ Sci Technol*,  
6 39, 2936-2943. <http://www.ncbi.nlm.nih.gov/pubmed/15926536>
- 7 Sarangapani, R.; Gentry, P. R.; Covington, T. R.; Teegarden, J. G.; 3rd, C. H. (2003).  
8 Evaluation of the potential impact of age- and gender-specific lung morphology and  
9 ventilation rate on the dosimetry of vapors. *Inhal Toxicol*, 15, 987-1016.  
10 <http://dx.doi.org/10.1080/713857276>
- 11 Sargent, L. M.; Zhou, X.; Keck, C. L.; Sanderson, N. D.; Zimonjic, D. B.; Popescu, N. C.;  
12 Thorgeirsson, S. S. (1999). Nonrandom cytogenetic alterations in hepatocellular  
13 carcinoma from transgenic mice overexpressing c-Myc and transforming growth factor-  
14 alpha in the liver. *Am J Pathol*, 154, 1047-1055.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/10233843>
- 16 Sato, A. (1993). Confounding factors in biological monitoring of exposure to organic solvents.  
17 *Int Arch Occup Environ Health*, 65, S61-S67.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/8406940>
- 19 Sato, A. and Nakajima, T. (1978). Differences following skin or inhalation exposure in the  
20 absorption and excretion kinetics of trichloroethylene and toluene. *Occup Environ Med*,  
21 35, 43-49. <http://www.ncbi.nlm.nih.gov/pubmed/629888>
- 22 Sato, A. and Nakajima, T. (1979). A vial-equilibration method to evaluate the drug-metabolizing  
23 enzyme activity for volatile hydrocarbons. *Toxicol Appl Pharmacol*, 47, 41-46.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/34248>
- 25 Sato, A. and Nakajima, T. (1985). Enhanced metabolism of volatile hydrocarbons in rat liver  
26 following food deprivation, restricted carbohydrate intake and administration of ethanol,  
27 phenobarbital, polychlorinated biphenyl and 3-methylcholanthrene: A comparative study.  
28 *Xenobiotica*, 15, 67. <http://dx.doi.org/10.3109/00498258509045336>
- 29 Sato, A.; Endoh, K.; Johanson, G. (1991). The Use of Models to Investigate the Toxicokinetic  
30 Behavior of Organic Solvents. Lewis Publishers, Chelsea, Michigan, 131-152.
- 31 Sato, A.; Endoh, K.; Kaneko, T.; Johanson, G. (1991). A simulation study of physiological  
32 factors affecting pharmacokinetic behaviour of organic solvent vapours. *Occup Environ*  
33 *Med*, 48, 342-347. <http://www.ncbi.nlm.nih.gov/pubmed/2039747>
- 34 Sato, A.; Nakajima, T.; Fujiwara, Y.; Murayama, N. (1977). A pharmacokinetic model to study  
35 the excretion of trichloroethylene and its metabolites after an inhalation exposure. *Occup*  
36 *Environ Med*, 34, 56-63. <http://www.ncbi.nlm.nih.gov/pubmed/843464>
- 37 Sato, A.; Nakajima, T.; Koyama, Y. (1980). Effects of chronic ethanol consumption on hepatic  
38 metabolism of aromatic and chlorinated hydrocarbons in rats. *Br J Ind Med*, 37, 382-386.  
39 <http://dx.doi.org/10.1136/oem.37.4.382>
- 40 Sato, A.; Nakajima, T.; Koyama, Y. (1981). Dose-related effects of a single dose of ethanol on  
41 the metabolism in rat liver of some aromatic and chlorinated hydrocarbons. *Toxicol Appl*  
42 *Pharmacol*, 60, 8-15. [http://dx.doi.org/10.1016/0041-008X\(81\)90129-0](http://dx.doi.org/10.1016/0041-008X(81)90129-0)
- 43 Sato, A.; Nakajima, T.; Koyama, Y. (1983). Interaction between ethanol and carbohydrate on the  
44 metabolism in rat liver of aromatic and chlorinated hydrocarbons. *Toxicol Appl*  
45 *Pharmacol*, 68, 242-249. [http://dx.doi.org/10.1016/0041-008X\(83\)90008-X](http://dx.doi.org/10.1016/0041-008X(83)90008-X)

DRAFT– DO NOT QUOTE OR CITE

- 1 Saunders, R. A. (1967). A new hazard in closed environmental atmospheres. *Arch Environ*  
2 *Health*, 14, 380-384. <http://www.ncbi.nlm.nih.gov/pubmed/4951670>
- 3 Sausen, P. and Elfarra, A. (1991). Reactivity of cysteine S-conjugate sulfoxides: formation of S-  
4 [1-chloro-2-(S-glutathionyl)vinyl]-L-cysteine sulfoxide by the reaction of S-(1,2-  
5 dichlorovinyl)-L-cysteine sulfoxide with glutathione. *Chem Res Toxicol*, 4, 655-660.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/1807449>
- 7 Sausen, P. J. and Elfarra, A. A. (1990). Cysteine conjugate S-oxidase. Characterization of a  
8 novel enzymatic activity in rat hepatic and renal microsomes. *J Biol Chem*, 265, 6139-  
9 6145. <http://www.ncbi.nlm.nih.gov/pubmed/2318851>
- 10 Savolainen, H.; Pfaffli, P.; Tengen, M.; Vainio, H. (1977). Trichloroethylene and 1,1,1-  
11 trichloroethane: effects on brain and liver after five days intermittent inhalation. *Arch*  
12 *Toxicol*, 38, 229-237. <http://www.ncbi.nlm.nih.gov/pubmed/578726>
- 13 Saxena, D. K.; Srivastava, R. S.; Lal, B.; Chandra, S. V. (1987). The effect of lead exposure on  
14 the testis of growing rats. *Exp Toxicol Pathol*, 31, 249-252.
- 15 Saygun, M.; Cakmak, A.; Ekici, A.; Pinar, T.; Bulcun, E.; Ulu, N.; Ekici, M. (2007). Five annual  
16 observations of respiratory findings in gun factory workers exposed to solvents. *J Occup*  
17 *Environ Med*, 49, 909-912. <http://dx.doi.org/10.1097/JOM.0b013e318124a8f3>
- 18 Sbrana, I.; Di Sibio, A.; Lomi, A.; Scarcelli, V. (1993). C-mitosis and numerical chromosome  
19 aberration analyses in human lymphocytes: 10 known or suspected spindle poisons.  
20 *Mutat Res*, 287, 57-70. <http://www.ncbi.nlm.nih.gov/pubmed/7683385>
- 21 Scharf, J. G.; Dombrowski, F.; Ramadori, G. (2001). The IGF axis and hepatocarcinogenesis. *J*  
22 *Clin Pathol*, 54, 138-144. <http://www.ncbi.nlm.nih.gov/pubmed/11376124>
- 23 Schatten, H. and Chakrabarti, A. (1998). Centrosome structure and function is altered by chloral  
24 hydrate and diazepam during the first reproductive cell cycles in sea urchin eggs. *Eur J*  
25 *Cell Biol*, 75, 9-20.
- 26 Scheff, P. A. and Wadden, R. A. (1993). Receptor modeling of volatile organic compounds 1  
27 Emission inventory and validation. *Environ Sci Technol*, 27, 617-625.  
28 <http://dx.doi.org/10.1021/es00041a005>
- 29 Schenk, M.; Purdue, M.; Colt, J.; Hartge, P.; Blair, A.; Stewart, P., . . . Severson, R. (2009).  
30 Occupation/industry and risk of non-Hodgkin's lymphoma in the United States. *Occup*  
31 *Environ Med*, 66, 23-31. <http://dx.doi.org/10.1136/oem.2007.036723>
- 32 Schlehofer, B.; Heuer, C.; Blettner, M.; Niehoff, D.; Wahrendorf, J. (1995). Occupation,  
33 smoking and demographic factors, and renal cell carcinoma in Germany. *Int J Epidemiol*,  
34 24, 51-57. <http://www.ncbi.nlm.nih.gov/pubmed/7797356>
- 35 Schlesselman, J. J. and Stolley, P. D. (1982). Case control studies: design, conduct, analysis.  
36 New York: Oxford University Press.
- 37 Schlipkötter, H. W.; Fodor, G. G.; Winneke, G. (1974). Zum Einfluss neurotoxischer  
38 Industriegifte auf unfallgenerierendes Fehlverhalten bei Kontroll- und Steuertätigkeiten.  
39 Wilhelmshaven, Dortmund: Wirtschaftsverlag Nordwest.
- 40 Schmucker, D. L. (2001). Liver function and phase I drug metabolism in the elderly: a paradox.  
41 *Drugs Aging*, 18, 837-851. <http://www.ncbi.nlm.nih.gov/pubmed/11772124>
- 42 Schmucker, D. L. (2005). Age-related changes in liver structure and function: Implications for  
43 disease. *Exp Gerontol*, 40, 650-659. <http://dx.doi.org/10.1016/j.exger.2005.06.009>

DRAFT– DO NOT QUOTE OR CITE

- 1 Schmucker, D. L.; Mooney, J. S.; Jones, A. L. (1978). Stereological analysis of hepatic fine  
2 structure in the Fischer 344 rat. Influence of sublobular location and animal age. *J Cell*  
3 *Biol*, 78, 319-337. <http://www.ncbi.nlm.nih.gov/pubmed/690169>
- 4 Schraml, P.; Zhaou, M.; Richter, J.; Brüning, T.; Pommer, M.; Sauter, G., . . . Moch, H. (1999).  
5 [Analysis of kidney tumors in trichloroethylene exposed workers by comparative  
6 genomic hybridization and DNA sequence analysis]. *Verh Dtsch Ges Pathol*, 83, 218-  
7 224. <http://www.ncbi.nlm.nih.gov/pubmed/10714214>
- 8 Schulte-Hermann, R.; Timmermann-Trosiener, I.; Schuppler, J. (1986). Facilitated expression of  
9 adaptive responses to phenobarbital in putative pre-stages of liver cancer. *Carcinogenesis*,  
10 7, 1651-1655. <http://www.ncbi.nlm.nih.gov/pubmed/2875809>
- 11 Schultz, I. R.; Merdink, J. L.; Gonzalez-Leon, A.; Bull, R. J. (1999). Comparative toxicokinetics  
12 of chlorinated and brominated haloacetates in F344 rats. *Toxicol Appl Pharmacol*, 158,  
13 103-114. <http://dx.doi.org/10.1006/taap.1999.869>
- 14 Schultz, I. R.; Merdink, J. L.; Gonzalez-Leon, A.; Bull, R. J. (2002). Dichloroacetate  
15 toxicokinetics and disruption of tyrosine catabolism in B6C3F1 mice: Dose-response  
16 relationships and age as a modifying factor. *Toxicology*, 173, 229-247.  
17 [http://dx.doi.org/10.1016/S0300-483X\(02\)00034-3](http://dx.doi.org/10.1016/S0300-483X(02)00034-3)
- 18 Schumucker, D. L.; Mooney, J. S.; Jones, A. L. (1977). Age-related changes in the hepatic  
19 endoplasmic reticulum: a quantitative analysis. *Science*, 197, 1005-1008.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/887935>
- 21 Schwartz-Arad, D.; Zajicek, G.; Bartfeld, E. (1989). The streaming liver IV: DNA content of the  
22 hepatocyte increases with its age. 9, 93-99.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/2709954>
- 24 Schwetz, B. A.; Leong, B. K. J.; Gehring, P. J. (1975). The effect of maternally inhaled  
25 trichloroethylene, perchloroethylene, methyl chloroform, and methylene chloride on  
26 embryonal and fetal development in mice and rats. *Toxicol Appl Pharmacol*, 32, 84-96.  
27 [http://dx.doi.org/10.1016/0041-008X\(75\)90197-0](http://dx.doi.org/10.1016/0041-008X(75)90197-0)
- 28 Scott, C. and Chiu, W. (2006). Trichloroethylene cancer epidemiology: a consideration of select  
29 issues. *Environ Health Perspect*, 114, 1471-1478.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/16966107>
- 31 Scott, J. E.; Forkert, P. G.; Oulton, M.; Rasmusson, M.-G.; Temple, S.; Fraser, M. O.;  
32 Whitefield, S. (1988). Pulmonary toxicity of trichloroethylene: induction of changes in  
33 surfactant phospholipids and phospholipase A2 activity in the mouse lung. *Exp Mol*  
34 *Pathol*, 49, 141-150. <http://www.ncbi.nlm.nih.gov/pubmed/3396665>
- 35 Seelbach, A.; Fissler, B.; Madle, S. (1993). Further evaluation of a modified micronucleus assay  
36 with V79 cells for detection of aneugenic effects. *Mutat Res*, 303, 163-169.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/7694134>
- 38 Seidler, A.; Mohner, M.; Berger, J.; Mester, B.; Deeg, E.; Elsner, G., . . . Becker, N. (2007).  
39 Solvent exposure and malignant lymphoma: A population-based case-control study in  
40 Germany. *J Occup Med Toxicol*, 2, 2. <http://dx.doi.org/10.1186/1745-6673-2-2>
- 41 Seitz, C. S.; Lin, Q.; Deng, H.; Khavari, P. A. (1998). Alterations in NF-kappaB function in  
42 transgenic epithelial tissue demonstrate a growth inhibitory role for NF-kappaB. *PNAS*,  
43 95, 2307-2312. <http://www.ncbi.nlm.nih.gov/pubmed/9482881>

DRAFT– DO NOT QUOTE OR CITE

- 1 Seitz, H. K. and Stickel, F. (2006). Risk factors and mechanisms of hepatocarcinogenesis with  
2 special emphasis on alcohol and oxidative stress. *Biol Chem*, 387, 349-360.  
3 <http://dx.doi.org/10.1515/bc.2006.047>
- 4 Seldén, A. and Ahlberg, G. (1991). Mortality and cancer morbidity after exposure to military  
5 aircraft fuel. *Aviat Space Environ Med*, 62, 789-794.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/1930063>
- 7 Selden, A.; Hultberg, B.; Ulander, A.; Ahlberg, G. J. (1993). Trichloroethylene exposure in  
8 vapour degreasing and the urinary excretion of N-acetyl-"beta"-D-glucosaminidase. *Arch*  
9 *Toxicol*, 67, 224-226. <http://www.ncbi.nlm.nih.gov/pubmed/8494503>
- 10 Selgrade, M. K. and Gilmour, M. I. (2010). Suppression of pulmonary host defenses and  
11 enhanced susceptibility to respiratory bacterial infection in mice following inhalation  
12 exposure to trichloroethylene and chloroform. *J Immunotoxicol*, 7, 350-356.  
13 <http://dx.doi.org/10.3109/1547691X.2010.520139>
- 14 Sellers, E. M.; Lang, M.; Koch-Weser, J.; LeBlanc, E.; Kalant, H. (1972). Interaction of chloral  
15 hydrate and ethanol in man I Metabolism. *Clin Pharmacol Ther*, 13, 37-49.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/4550322>
- 17 Sellers, E.; Lang-Sellers, M.; Koch-Weser, J. (1978). Comparative metabolism of chloral hydrate  
18 and triclofos. *J Clin Pharmacol*, 18, 457-461.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/711926>
- 20 Selmin, O. I.; Thorne, P. A.; Caldwell, P. T.; Taylor, M. R. (2008). Trichloroethylene and  
21 trichloroacetic acid regulate calcium signaling pathways in murine embryonal carcinoma  
22 cells p19. *Cardiovasc Toxicol*, 8, 47-56. <http://dx.doi.org/10.1007/s12012-008-9014-2>
- 23 Seng, J.; Agrawal, N.; Horsley, E.; Leakey, T.; Scherer, E.; Xia, S., . . . Leakey, J. (2003).  
24 Toxicokinetics of chloral hydrate in ad libitum-fed, dietary-controlled, and calorically  
25 restricted male B6C3F1 mice following short-term exposure. *Toxicol Appl Pharmacol*,  
26 193, 281-292. <http://www.ncbi.nlm.nih.gov/pubmed/14644628>
- 27 Seo, M.; Ikeda, K.; Okamura, T.; Kida, K.; Satoh, M.; Inagaki, N., . . . Nagase, H. (2008).  
28 Enhancing effect of chlorinated organic solvents on histamine release and inflammatory  
29 mediator production. *Toxicology*, 243, 75-83. <http://dx.doi.org/10.1016/j.tox.2007.09.024>
- 30 Seppalainen, A. M. and Antti-Poika, M. (1983). Time course of electrophysiological findings for  
31 patients with solvent poisoning. A descriptive study. *Scand J Work Environ Health*, 9,  
32 15-24. <http://www.ncbi.nlm.nih.gov/pubmed/6857184>
- 33 Sexton, K.; Adgate, J. L.; Church, T. R.; Ashley, D. L.; Needham, L. L.; Ramachandran, G., . . .  
34 Ryan, A. D. (2005). Children's exposure to volatile organic compounds as determined by  
35 longitudinal measurements in blood. *Environ Health Perspect*, 113, 342-349.  
36 <http://dx.doi.org/10.1289/ehp.7412>
- 37 Sexton, K.; Mongin, S. J.; Adgate, J. L.; Pratt, G. C.; Ramachandran, G.; Stock, T. H.; Morandi,  
38 M. T. (2007). Estimating volatile organic compound concentrations in selected  
39 microenvironments using time-activity and personal exposure data. *J Toxicol Environ*  
40 *Health A*, 70, 465-476. <http://dx.doi.org/10.1080/15287390600870858>
- 41 Shafer, T. J.; Bushnell, P. J.; Benignus, V. A.; Woodward, J. J. (2005). Perturbation of voltage-  
42 sensitive Ca<sup>2+</sup> channel function by volatile organic solvents. *J Pharmacol Exp Ther*, 315,  
43 1109-1118. <http://dx.doi.org/10.1124/jpet.105.090027>

- 1 Shah, J. J. and Singh, H. B. (1988). Distribution of volatile organic chemicals in outdoor and  
2 indoor air: a national VOCs data base. *Environ Sci Technol*, 22, 1381-1388.  
3 <http://dx.doi.org/10.1021/es00177a001>
- 4 Shah, Y. M.; Morimura, K.; Yang, Q.; Tanabe, T.; Takagi, M.; Gonzalez, F. J. (2007).  
5 Peroxisome proliferator-activated receptor alpha regulates a microRNA-mediated  
6 signaling cascade responsible for hepatocellular proliferation. *Mol Cell Biol*, 27, 4238-  
7 4247. <http://dx.doi.org/10.1128/MCB.00317-07>
- 8 Shankar, K.; Vaidya, V.; Corton, J.; Bucci, T.; Liu, J.; Waalkes, M.; Mehendale, H. (2003).  
9 Activation of PPAR-alpha in streptozotocin-induced diabetes is essential for resistance  
10 against acetaminophen toxicity. *FASEB J*, 17, 1748-1750.  
11 <http://dx.doi.org/10.1096/fj.02-1186fje>
- 12 Shannon, H.; Haines, T.; Bernholz, C.; Julian, J.; Verma, D.; Jamieson, E.; Walsh, C. (1988).  
13 Cancer morbidity in lamp manufacturing workers. *Am J Ind Med*, 14, 281-290.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/3189346>
- 15 Shao, J.; Stapleton, P. L.; Lin, Y. S.; Gallagher, E. P. (2007). Cytochrome p450 and glutathione  
16 s-transferase mRNA expression in human fetal liver hematopoietic stem cells. *Drug*  
17 *Metab Dispos*, 35, 168-175. <http://dx.doi.org/10.1124/dmd.106.012757>
- 18 Sharpe, C. R.; Rochon, J. E.; Adam, J. M.; Suissa, S. (1989). Case-control study of hydrocarbon  
19 exposures in patients with renal cell carcinoma. *CMAJ*, 140, 1309-1318.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/2720514>
- 21 Shearer, B. G. and Hoekstra, W. J. (2003). Recent advances in peroxisome proliferator-activated  
22 receptor science. *Curr Med Chem*, 10, 267-280.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/12570700>
- 24 Sheffels, P.; Schroeder, J.; Altuntas, T.; Liggitt, H.; Kharasch, E. (2004). Role of cytochrome  
25 P4503A in cysteine S-conjugates sulfoxidation and the nephrotoxicity of the sevoflurane  
26 degradation product fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether (compound  
27 A) in rats. *Chem Res Toxicol*, 17, 1177-1189. <http://dx.doi.org/10.1021/tx049899e>
- 28 Sher, T.; Yi, H. F.; McBride, O. W.; Gonzalez, F. J. (1993). cDNA cloning, chromosomal  
29 mapping, and functional characterization of the human peroxisome proliferator activated  
30 receptor. *Biochemistry*, 32, 5598-5604.
- 31 Sherins, R. J. and Hodgen, G. D. (1976). Testicular gamma glutamyl-transpeptidase: an index of  
32 Sertoli cell function in man. *Journal of Reproduction and Fertility*, 48, 191-193.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/9507>
- 34 Shiao, Y.; Kamata, S.; Li, L.; Hooth, M.; DeAngelo, A.; Anderson, L.; Wolf, D. (2002).  
35 Mutations in the VHL gene from potassium bromate-induced rat clear cell renal tumors.  
36 *Cancer Lett*, 187, 207-214. <http://www.ncbi.nlm.nih.gov/pubmed/12359370>
- 37 Shiao, Y.; Rice, J.; Anderson, L.; Diwan, B.; Hard, G. (1998). von Hippel-Lindau gene  
38 mutations in N-nitrosodimethylamine-induced rat renal epithelial tumors. *J Natl Cancer*  
39 *Inst*, 90, 1720-1723. <http://www.ncbi.nlm.nih.gov/pubmed/9827526>
- 40 Shih, C.; Chen, H.; Chiu, T. (2001). Acute exposure to trichloroethylene differentially alters the  
41 susceptibility to chemoconvulsants in mice. *Toxicology*, 162, 35-42.  
42 <http://www.ncbi.nlm.nih.gov/pubmed/11311456>
- 43 Shimada, T.; Swanson, A. F.; Leber, P.; Williams, G. M. (1985). Activities of chlorinated ethane  
44 and ethylene compounds in the Salmonella/rat microsome mutagenesis and rat

- 1 hepatocyte/DNA repair assays under vapor phase exposure conditions. *Cell Biol Toxicol*,  
2 1, 159-179. <http://dx.doi.org/10.1007/BF00120162>
- 3 Shimada, T.; Yamazaki, H.; Mimura, M.; Inui, Y.; Guengerich, F. P. (1994). Interindividual  
4 variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs,  
5 carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30  
6 Caucasians. *J Pharmacol Exp Ther*, 270, 414-423.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/8035341>
- 8 Shindell, S. and Ulrich, S. (1985). A cohort study of employees of a manufacturing plant using  
9 trichlorethylene. *J Occup Environ Med*, 27, 577-579.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/4032094>
- 11 Shirasu, Y.; Moriya, M.; Kato, K.; Furuhashi, A.; Kada, T. (1976). Mutagenicity screening of  
12 pesticides in the microbial system. *Mutat Res*, 40, 19-30. [http://dx.doi.org/10.1016/0165-](http://dx.doi.org/10.1016/0165-1218(76)90018-5)  
13 [1218\(76\)90018-5](http://dx.doi.org/10.1016/0165-1218(76)90018-5)
- 14 Short, B. G. (1993). Cell proliferation and renal carcinogenesis. *Environ Health Perspect*, 101  
15 Suppl 5, 115-120. <http://www.ncbi.nlm.nih.gov/pubmed/7516872>
- 16 Shroads, A. L.; Guo, X.; Dixit, V.; Liu, H. P.; James, M. O.; Stacpoole, P. W. (2008). Age-  
17 dependent kinetics and metabolism of dichloroacetate: possible relevance to toxicity. *J*  
18 *Pharmacol Exp Ther*, 324, 1163-1171. <http://dx.doi.org/10.1124/jpet.107.134593>
- 19 Shu, X. O.; Perentesis, J. P.; Wen, W.; Buckley, J. D.; Boyle, E.; Ross, J. A., . . . Group, C. s. O.  
20 (2004). Parental exposure to medications and hydrocarbons and ras mutations in children  
21 with acute lymphoblastic leukemia: a report from the Children's Oncology Group. *Cancer*  
22 *Epidemiol Biomarkers Prev*, 13, 1230-1235.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/15247135>
- 24 Shu, X. O.; Stewart, P.; Wen, W. Q.; Han, D.; Potter, J. D.; Buckley, J. D., . . . Robison, L. L.  
25 (1999). Parental occupational exposure to hydrocarbons and risk of acute lymphocytic  
26 leukemia in offspring. *Cancer Epidemiol Biomarkers Prev*, 8, 783-791.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/10498397>
- 28 Shultz, J. and Weiner, H. (1979). Alteration of the enzymology of chloral hydrate reduction in  
29 the presence of ethanol. *Biochem Pharmacol*, 28, 3379-3384.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/43731>
- 31 Siegel, J. H. and Kasmin, F. E. (1997). Biliary tract diseases in the elderly: management and  
32 outcomes. *Gut*, 41, 433-435. <http://www.ncbi.nlm.nih.gov/pubmed/9391238>
- 33 Siemiatycki, J. (1991). Risk factors for cancer in the workplace. Boca Raton, FL: CRC Press.
- 34 Siemiatycki, J.; Dewar, R.; Nadon, L.; Gérin, M. (1994). Occupational risk factors for bladder  
35 cancer: results from a case-control study in Montreal, Quebec, Canada. *Am J Epidemiol*,  
36 140, 1061-1080. <http://www.ncbi.nlm.nih.gov/pubmed/7998589>
- 37 Siemiatycki, J.; Fritschi, L.; Nadon, L.; Gérin, M. (1997). Reliability of an expert rating  
38 procedure for retrospective assessment of occupational exposures in community-based  
39 case-control studies. *Am J Ind Med*, 31, 280-286.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/9055950>
- 41 Siemiatycki, J.; Wacholder, S.; Richardson, L.; Dewar, R.; Gérin, M. (1987). Discovering  
42 carcinogens in the occupational environment. Methods of data collection and analysis of  
43 a large case-referent monitoring system. *Scand J Work Environ Health*, 13, 486-492.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/3433050>



DRAFT– DO NOT QUOTE OR CITE

- 1 Sigal, S. H.; Rajvanshi, P.; Gorla, G. R.; Sokhi, R. P.; Saxena, R.; Gebhard, D. R., . . . Gupta, S.  
2 (1999). Partial hepatectomy-induced polyploidy attenuates hepatocyte replication and  
3 activates cell aging events. *Am J Physiol*, 276, G1260-G1272.  
4 <http://www.ncbi.nlm.nih.gov/pubmed/10330018>
- 5 Simmon, V. F.; Kauhanen, K.; Tardiff, R. G. (1977). Mutagenic activity of chemicals identified  
6 in drinking water. In B. B. F. S. D Scott (Ed.), *Progress in Genetic Toxicology* (Vol. 2,  
7 pp. 249-268). New York; Amsterdam: Elsevier/North Holland Press.
- 8 Simmons, J. E.; Boyes, W. K.; Bushnell, P. J.; Raymer, J. H.; Limsakun, T.; McDonald, A., . . .  
9 Evans, M. V. (2002). A physiologically based pharmacokinetic model for  
10 trichloroethylene in the male long-evans rat. *Toxicol Sci*, 69, 3-15.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/12215655>
- 12 Singh, M. and Sinha, U. (1976). Chloral hydrate induced haploidization in *Aspergillus nidulans*.  
13 *Experientia*, 32, 1144-1145.
- 14 Singh, M. and Sinha, U. (1979). Mitotic haploidization and growth of *Aspergillus nidulans* on  
15 media containing chloral hydrate. 14, 1-4.
- 16 Sinks, T.; Lushniak, B.; Haussler, B.; Sniezek, J.; Deng, J.; Roper, P., . . . Coates, R. (1992).  
17 Renal cell cancer among paperboard printing workers. *Epidemiology*, 3, 483-489.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/1420513>
- 19 Skender, L. J.; Karacic, V.; Prpic-Majic, D. (1991). A comparative study of human levels of  
20 trichloroethylene and tetrachloroethylene after occupational exposure. *Arch Environ*  
21 *Occup Health*, 46, 174-178. <http://www.ncbi.nlm.nih.gov/pubmed/2039273>
- 22 Slikker, J. W.; Andersen, M. E.; Bogdanffy, M. S.; Bus, J. S.; Cohen, S. D.; Conolly, R. B., . . .  
23 Wallace, K. (2004). Dose-dependent transitions in mechanisms of toxicity: Case studies.  
24 *Toxicol Appl Pharmacol*, 201, 226-294. <http://dx.doi.org/10.1016/j.taap.2004.06.027>
- 25 Slikker, J. W.; Andersen, M. E.; Bogdanffy, M. S.; Bus, J. S.; Cohen, S. D.; Conolly, R. B., . . .  
26 Wallace, K. (2004). Dose-dependent transitions in mechanisms of toxicity. *Toxicol Appl*  
27 *Pharmacol*, 201, 203-225. <http://dx.doi.org/10.1016/j.taap.2004.06.019>
- 28 Sliwińska-Kowalska, M.; Jedlińska, U.; Rządzińska, A.; Rajkowska, E. (1999). [Hair cell  
29 regeneration in a chicken's inner ear after damage due to exposure to industrial noise].  
30 *Otolaryngol Pol*, 53, 469-477. <http://www.ncbi.nlm.nih.gov/pubmed/10581958>
- 31 Smith, G. (1970). The investigation of the mental effects of trichlorethylene. *Ergonomics*, 13,  
32 580-586. <http://www.ncbi.nlm.nih.gov/pubmed/5494866>
- 33 Smith, M. K.; Randall, J. L.; Read, E. J.; Stober, J. A. (1989). Teratogenic activity of  
34 trichloroacetic acid in the rat. *Teratology*, 40, 445-451.  
35 <http://dx.doi.org/10.1002/tera.1420400506>
- 36 Smith, M.; Randall, J.; Read, E.; Stober, J. (1992). Developmental toxicity of dichloroacetate in  
37 the rat. *Teratology*, 46, 217-223. <http://dx.doi.org/10.1002/tera.1420460305>
- 38 Snedcor, G. W. and Cochran, W. G. (1980). *Statistical methods* (7th ed.). Ames, IA: Iowa State  
39 University Press.
- 40 Snyder, R.; Pullman, J.; Carter, J.; Carter, H.; DeAngelo, A. (1995). In vivo administration of  
41 dichloroacetic acid suppresses spontaneous apoptosis in murine hepatocytes. *Cancer Res*,  
42 55, 3702-3705. <http://www.ncbi.nlm.nih.gov/pubmed/7641179>
- 43 Song, B. J.; Veech, R. L.; Saenger, P. (1990). Cytochrome P450IIE1 is elevated in lymphocytes  
44 from poorly controlled insulin-dependent diabetics. *J Clin Endocrinol Metab*, 71, 1036-  
45 1040. <http://www.ncbi.nlm.nih.gov/pubmed/2205622>

DRAFT– DO NOT QUOTE OR CITE

- 1 Soni, M.; Mangipudy, R.; Mumtaz, M.; Mehendale, H. (1998). Tissue repair response as a  
2 function of dose during trichloroethylene hepatotoxicity. *Toxicol Sci*, 42, 158-165.  
3 <http://dx.doi.org/10.1006/toxs.1998.2427>
- 4 Soni, M.; Ramaiah, S.; Mumtaz, M.; Clewell, H.; Mehendale, H. (1999). Toxicant-inflicted  
5 injury and stimulated tissue repair are opposing toxicodynamic forces in predictive  
6 toxicology. *Regul Toxicol Pharmacol*, 29, 165-174.  
7 <http://dx.doi.org/10.1006/rtph.1998.1280>
- 8 Sonnenfeld, N.; Hertz-Picciotto, I.; Kaye, W. E. (2001). Tetrachloroethylene in drinking water  
9 and birth outcomes at the US Marine Corps Base at Camp Lejeune, North Carolina. *Am J*  
10 *Epidemiol*, 154, 902-908. <http://www.ncbi.nlm.nih.gov/pubmed/11700244>
- 11 Sora, S. and Agostini Carbone, M. L. (1987). Chloral hydrate, methylmercury hydroxide and  
12 ethidium bromide affect chromosomal segregation during meiosis of *Saccharomyces*  
13 *cerevisiae*. *Mutat Res Lett*, 190, 13-17. [http://dx.doi.org/10.1016/0165-7992\(87\)90075-3](http://dx.doi.org/10.1016/0165-7992(87)90075-3)
- 14 Soresi, M.; Giannitrapani, L.; D'Antona, F.; Florena, A. M.; La Spada, E.; Terranova, A., . . .  
15 Montalto, G. (2006). Interleukin-6 and its soluble receptor in patients with liver cirrhosis  
16 and hepatocellular carcinoma. *World J Gastroenterol*, 12, 2563-2568.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/16688802>
- 18 Soucek, B. and Vlachova, D. (1960). Excretion of trichloroethylene metabolites in human urine.  
19 *Occup Environ Med*, 17, 60-64. <http://www.ncbi.nlm.nih.gov/pubmed/13832993>
- 20 Spiegelhalter, D.; Thomas, A.; Best, N. (2003). WinBugs version 1.4 user manual. Cambridge,  
21 UK: MRC Biostatistics Unit.
- 22 Spielberg, S. P. (1996). N-acetyltransferases: pharmacogenetics and clinical consequences of  
23 polymorphic drug metabolism. *J Pharmacokinet Biopharm*, 24, 509-519.  
24 <http://www.ncbi.nlm.nih.gov/pubmed/9131488>
- 25 Spirtas, R.; Stewart, P. A.; Lee, J. S.; Marano, D. E.; Forbes, C. D.; Grauman, D. J., . . . Cohen, J.  
26 L. (1991). Retrospective cohort mortality study of workers at an aircraft maintenance  
27 facility. I. Epidemiological results. *Br J Ind Med*, 48, 515-530.  
28 <http://dx.doi.org/10.1136/oem.48.8.515>
- 29 Squillace, P. J.; Moran, M. J.; Price, C. V. (2004). VOCs in shallow groundwater in new  
30 residential/commercial areas of the United States. *Environ Sci Technol*, 38, 5327-5338.
- 31 SRI International. (1992). 1992 directory of chemical producers United States of America.  
32 Menlo Park, CA: Author.
- 33 Srivastava, S. and Srivastava, S. (1991). Effect of di(2-ethylhexyl)phthalate on 17 $\beta$ -  
34 hydroxysteroid dehydrogenase activity in testis of rat. *Toxicol Lett*, 57, 235-239.  
35 [http://dx.doi.org/10.1016/0378-4274\(91\)90150-5](http://dx.doi.org/10.1016/0378-4274(91)90150-5)
- 36 Staats, D. A.; Fisher, J. W.; Connolly, R. B. (1991). Gastrointestinal absorption of xenobiotics in  
37 physiologically based pharmacokinetic models: A two-compartment description. *Drug*  
38 *Metab Dispos*, 19, 144-148. <http://www.ncbi.nlm.nih.gov/pubmed/1673388>
- 39 Stadler, J. C. and Kennedy, G. L. J. (1996). Evaluation of the sensory irritation potential of  
40 volatile organic chemicals from carpets--alone and in combination. *Food Chem Toxicol*,  
41 34, 1125-1130. <http://www.ncbi.nlm.nih.gov/pubmed/9119325>
- 42 Stanley, L. A.; Blackburn, D. R.; Devereaux, S.; Foley, J.; Lord, P. G.; Maronpot, R. R., . . .  
43 Anderson, M. W. (1994). Ras mutations in methylclofenapate-induced B6C3F1 and  
44 C57BL/10J mouse liver tumours. *Carcinogenesis*, 15, 1125-1131.  
45 <http://www.ncbi.nlm.nih.gov/pubmed/8020144>

- 1 Staples, R. E. (1974). Detection of visceral alterations in mammalian fetuses. *Teratology*, 9,  
2 A37–A38.
- 3 Stauber, A. J. and Bull, R. J. (1997). Differences in phenotype and cell replicative behavior of  
4 hepatic tumors induced by dichloroacetate (DCA) and trichloroacetate (TCA). *Toxicol*  
5 *Appl Pharmacol*, 144, 235-246. <http://dx.doi.org/10.1006/taap.1997.8159>
- 6 Stauber, A. J.; Bull, R. J.; Thrall, B. D. (1998). Dichloroacetate and trichloroacetate promote  
7 clonal expansion of anchorage-independent hepatocytes in vivo and in vitro. *Toxicol*  
8 *Appl Pharmacol*, 150, 287-294. <http://dx.doi.org/10.1006/taap.1998.8417>
- 9 Steinberg, A. (1993). Should chloral hydrate be banned. *Pediatrics*, 92, 442-446.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/8361800>
- 11 Stemhagen, A.; Slade, J.; Altman, R.; Bill, J. (1983). Occupational risk factors and liver cancer:  
12 A retrospective case-control study of primary liver cancer in New Jersey. *Am J*  
13 *Epidemiol*, 117, 443-454. <http://www.ncbi.nlm.nih.gov/pubmed/6837558>
- 14 Stemmer, K.; Ellinger-Ziegelbauer, H.; Ahr, H. J.; Dietrich, D. R. (2007). Carcinogen-specific  
15 gene expression profiles in short-term treated Eker and wild-type rats indicative of  
16 pathways involved in renal tumorigenesis. *Cancer Res*, 67, 4052-4068.  
17 <http://dx.doi.org/10.1158/0008-5472.CAN-06-3587>
- 18 Stenner, R.; Merdink, J.; Stevens, D.; Springer, D.; Bull, R. (1997). Enterohepatic recirculation  
19 of trichloroethanol glucuronide as a significant source of trichloroacetic acid. *Metabolites*  
20 *of trichloroethylene*. *Drug Metab Dispos*, 25, 529-535.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/9152590>
- 22 Stephens, E. A.; Taylor, J. A.; Kaplan, N.; Yang, C. H.; Hsieh, L. L.; Lucier, G. W.; Bell, D. A.  
23 (1994). Ethnic variation in the CYP2E1 gene: polymorphism analysis of 695 African-  
24 Americans, European-Americans and Taiwanese. *Pharmacogenetics*, 4, 185-192.
- 25 Stephenson, R. P. (1956). A modification of receptor theory. *Br J Pharmacol*, 11, 379–393.
- 26 Steup, D. R.; Hall, P.; McMillan, D. A.; Sipes, I. G. (1993). Time course of hepatic injury and  
27 recovery following coadministration of carbon tetrachloride and trichloroethylene in  
28 Fischer-344 rats. *Toxicol Pathol*, 21, 327-334.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/8248722>
- 30 Steup, D. R.; Wiersma, D.; McMillan, D. A.; Sipes, I. G. (1991). Pretreatment with drinking  
31 water solutions containing trichloroethylene or chloroform enhances the hepatotoxicity of  
32 carbon tetrachloride in Fischer 344 rats. *Toxicol Sci*, 16, 798-809.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/1884917>
- 34 Stevens, J. (1985). Cysteine conjugate beta-lyase activities in rat kidney cortex: subcellular  
35 localization and relationship to the hepatic enzyme. *Biochem Biophys Res Commun*, 129,  
36 499-504. <http://www.ncbi.nlm.nih.gov/pubmed/4015643>
- 37 Stevens, J. and Jakoby, W. B. (1983). Cysteine conjugate beta-lyase. *Mol Pharmacol*, 23, 761-  
38 765.
- 39 Stevens, J.; Ayoubi, N.; Robbins, J. (1988). The role of mitochondrial matrix enzymes in the  
40 metabolism and toxicity of cysteine conjugates. *J Biol Chem*, 263, 3395-3401.  
41 <http://www.ncbi.nlm.nih.gov/pubmed/3343250>
- 42 Stevens, J.; Hatzinger, P.; Hayden, P. (1989). Quantitation of multiple pathways for the  
43 metabolism of nephrotoxic cysteine conjugates using selective inhibitors of L-alpha-  
44 hydroxy acid oxidase (L-amino acid oxidase) and cysteine conjugate beta-lyase. *Drug*  
45 *Metab Dispos*, 17, 297-303. <http://www.ncbi.nlm.nih.gov/pubmed/2568912>

DRAFT– DO NOT QUOTE OR CITE

- 1 Stevens, J.; Robbins, J.; Byrd, R. (1986). A purified cysteine conjugate beta-lyase from rat  
2 kidney cytosol. Requirement for an alpha-keto acid or an amino acid oxidase for activity  
3 and identity with soluble glutamine transaminase K. *J Biol Chem*, 261, 15529-15537.  
4 <http://www.ncbi.nlm.nih.gov/pubmed/3782077>
- 5 Stewart, B. W.; Le Mesurier, S. M.; Lykke, A. W. J. (1979). Correlation of biochemical and  
6 morphological changes induced by chemical injury to the lung. *Chem Biol Interact*, 26,  
7 321-338. <http://www.ncbi.nlm.nih.gov/pubmed/509695>
- 8 Stewart, P. A.; Lee, J. S.; Marano, D. E.; Spirtas, R.; Forbes, C. D.; Blair, A. (1991).  
9 Retrospective cohort mortality study of workers at an aircraft maintenance facility II  
10 Exposures and their assessment. *Occup Environ Med*, 48, 531-537.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/1878309>
- 12 Stewart, P. A.; Lees, P. S.; Francis, M. (1996). Quantification of historical exposures in  
13 occupational cohort studies. *Scand J Work Environ Health*, 22, 405-414.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/9000307>
- 15 Stewart, P. and Dosemeci, M. (2005). Exposure estimation in occupational epidemiology studies.  
16 *Third Meeting on Assessing Human Health Risks of Trichloroethylene*. Irvine, CA.
- 17 Stewart, R. D.; Dodd, H. C.; Gay, H. H.; Erley, D. S. (1970). Experimental human exposure to  
18 trichloroethylene. *Arch Environ Health*, 20, 64-71.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/5409662>
- 20 Stewart, R. D.; Hake, C. L.; Peterson, J. E. (1974). "Degreaser's flush". *Arch Environ Health*, 29,  
21 1-5.
- 22 Stewart, W. F. and Correa-Villaseor, A. (1991). False positive exposure errors and low exposure  
23 prevalence in community-based case-control studies. *Appl Occup Environ Hyg*, 6, 534-  
24 540.
- 25 Stopps, G. J. and McLaughlin, M. (1967). Psychophysiological testing of human subjects  
26 exposed to solvent vapors. *Am Ind Hyg Assoc J*, 28, 43-50.  
27 <http://www.ncbi.nlm.nih.gov/pubmed/6032934>
- 28 Stott, W. T.; Quast, J. F.; Watanabe, P. G. (1982). The pharmacokinetics and macromolecular  
29 interactions of trichloroethylene in mice and rats. *Toxicol Appl Pharmacol*, 62, 137-151.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/7064149>
- 31 Stuckhardt, J. L. and Poppe, S. M. (1984). Fresh visceral examination of rat and rabbit fetuses  
32 used in teratogenicity testing. *Teratog Carcinog Mutagen*, 4, 181-188.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/6145223>
- 34 Styles, J. A.; Kelly, M.; Pritchard, N. R.; Ekcombe, C. R. (1988). A species comparison of acute  
35 hyperplasia induced by the peroxisome proliferator methylclofenapate: Involvement of  
36 the binucleated hepatocyte. *Carcinogenesis*, 9, 1647-1655.  
37 <http://dx.doi.org/10.1093/carcin/9.9.1647>
- 38 Styles, J. A.; Wyatt, I.; Coutts, C. (1991). Trichloroacetic acid: studies on uptake and effects on  
39 hepatic DNA and liver growth in mouse. *Carcinogenesis*, 12, 1715-1719.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/1893532>
- 41 Su, Q. and Bannasch, P. (2003). Relevance of hepatic preneoplasia for human  
42 hepatocarcinogenesis. *Toxicol Pathol*, 31, 126-133.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/12597457>

DRAFT– DO NOT QUOTE OR CITE

- 1 Subramoniam, A.; Goel, S. K.; Pandya, K. P.; Seth, P. K. (1989). Influence of trichloroethylene  
2 treatment on phosphoinositides in rat brain. *Toxicol Lett*, 49, 55-60.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/2554538>
- 4 Sung, T.; Chen, P.; Jyuhn-Hsiarn Lee, L.; Lin, Y.; Hsieh, G.; Wang, J. (2007). Increased  
5 standardized incidence ratio of breast cancer in female electronics workers. *BMC Public*  
6 *Health*, 7, 102. <http://dx.doi.org/10.1186/1471-2458-7-102>
- 7 Sung, T.; Wang, J.; Chen, P. (2008). Increased risk of cancer in the offspring of female  
8 electronics workers. *Reprod Toxicol*, 25, 115-119.  
9 <http://dx.doi.org/10.1016/j.reprotox.2007.08.004>
- 10 Supplementary data for TCE assessment: Cancer rodents input data. (2011).  
11 Supplementary data for TCE assessment: Cancer rodents model selections. (2011).  
12 Supplementary data for TCE assessment: Cancer rodents plots. (2011).  
13 Supplementary data for TCE assessment: Cancer rodents results. (2011).  
14 Supplementary data for TCE assessment: Cancer rodents uncertainty analysis. (2011).  
15 Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-inhalation histograms,  
16 inhalation bioassays. (2011).  
17 Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-inhalation historams,  
18 oral bioassays. (2011).  
19 Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-oral histograms,  
20 inhalation bioassay. (2011).  
21 Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-oral histograms, oral  
22 bioassays. (2011).  
23 Supplementary data for TCE assessment: Hack mouse population calibration evaluation. (2011).  
24 Supplementary data for TCE assessment: Hack mouse subject calibration. (2011).  
25 Supplementary data for TCE assessment: Human posteriors by subject. (2011).  
26 Supplementary data for TCE assessment: Mouse posteriors by subject. (2011).  
27 Supplementary data for TCE assessment: Non-cancer HECs plots from human inhalation studies.  
28 (2011).  
29 Supplementary data for TCE assessment: Non-cancer HECs plots from rodent inhalation studies.  
30 (2011).  
31 Supplementary data for TCE assessment: Non-cancer HECs plots from rodent oral studies.  
32 (2011).  
33 Supplementary data for TCE assessment: Non-cancer HEDs altPOD plots from rodent inhalation  
34 studies. (2011).  
35 Supplementary data for TCE assessment: Non-cancer HEDs plots from human inhalation studies.  
36 (2011).  
37 Supplementary data for TCE assessment: Non-cancer HEDs plots from rodent inhalation studies.  
38 (2011).  
39 Supplementary data for TCE assessment: Non-cancer HEDs plots from rodent oral studies.  
40 (2011).  
41 Supplementary data for TCE assessment: Non-cancer input data contin. (2011).  
42 Supplementary data for TCE assessment: Non-cancer input data dichot. (2011).  
43 Supplementary data for TCE assessment: Non-cancer plots contin. (2011).  
44 Supplementary data for TCE assessment: Non-cancer plots dichot. (2011).  
45 Supplementary data for TCE assessment: Non-cancer results contin. (2011).

DRAFT– DO NOT QUOTE OR CITE

- 1 Supplementary data for TCE assessment: Non-cancer results dichot. (2011).  
2 Supplementary data for TCE assessment: Rat population example. (2011).  
3 Supplementary data for TCE assessment: Rat posterior by subject. (2011).  
4 Supplementary data for TCE assessment: Rodents time to tumor results. (2011).  
5 Sverdrup, B.; Kallberg, H.; Bengtsson, C.; Lundberg, I.; Padyukov, L.; Alfredsson, L., . . .  
6 Group, E. I. o. R. A. S. (2005). Association between occupational exposure to mineral oil  
7 and rheumatoid arthritis: results from the Swedish EIRA case-control study. *Arthritis Res*  
8 *Ther*, 7, R1296-R1303. <http://dx.doi.org/10.1186/ar1824>  
9 Swaen, G. M. (1995). Increased incidence of renal cell tumours in a cohort of cardboard workers  
10 exposed to trichloroethylene [letter to the editor] [see Henschler, 1995, for a reply]. *Arch*  
11 *Toxicol*, 70, 127-128.  
12 Swan, S.; Shaw, G.; Harris, J.; Neutra, R. (1989). Congenital cardiac anomalies in relation to  
13 water contamination, Santa Clara County, California, 1981-1983. *Am J Epidemiol*, 129,  
14 885-893. <http://www.ncbi.nlm.nih.gov/pubmed/2784935>  
15 Sweeney, C.; Farrow, D.; Schwartz, S.; Eaton, D.; Checkoway, H.; Vaughan, T. (2000).  
16 Glutathione S-transferase M1, T1, and P1 polymorphisms as risk factors for renal cell  
17 carcinoma: A case-control study. *Cancer Epidemiol Biomarkers Prev*, 9, 449-454.  
18 <http://www.ncbi.nlm.nih.gov/pubmed/10794492>  
19 Sweeney, L. M.; Kirman, C. R.; Gargas, M. L.; Dugard, P. H. (2009). Contribution of  
20 trichloroacetic acid to liver tumors observed in perchloroethylene (perc)-exposed mice.  
21 *Toxicology*, 260, 77-83. <http://dx.doi.org/10.1016/j.tox.2009.03.008>  
22 Sweet, C. W. and Vermette, S. J. (1992). Toxic volatile organic compounds in urban air in  
23 Illinois. *Environ Sci Technol*, 26, 165-173. <http://dx.doi.org/10.1021/es00025a020>  
24 Szyf, M. (2007). The dynamic epigenome and its implications in toxicology. *Toxicol Sci*, 100, 7-  
25 23. <http://dx.doi.org/10.1093/toxsci/kfm177>  
26 Szymańska, K.; Moore, L. E.; Rothman, N.; Chow, W. H.; Waldman, F.; Jaeger, E., . . . Brennan,  
27 P. (2010). TP53, EGFR, and KRAS mutations in relation to VHL inactivation and  
28 lifestyle risk factors in renal-cell carcinoma from central and eastern Europe. 293, 92-98.  
29 <http://dx.doi.org/10.1016/j.canlet.2009.11.024>  
30 't Mannetje, A.; Dryson, E.; Walls, C.; McLean, D.; McKenzie, F.; Maule, M., . . . Pearce, N.  
31 (2008). High risk occupations for non-Hodgkin's lymphoma in New Zealand: case-  
32 control study. *Occup Environ Med*, 65, 354-363.  
33 <http://dx.doi.org/10.1136/oem.2007.035014>  
34 't Mannetje, A.; Steenland, K.; Attfield, M.; Boffetta, P.; Checkoway, H.; DeKlerk, N.; Koskela,  
35 R. (2002). Exposure-response analysis and risk assessment for silica and silicosis  
36 mortality in a pooled analysis of six cohorts. *Occup Environ Med*, 59, 723-728.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/12409529>  
38 Takashima, K.; Ito, Y.; Gonzalez, F. J.; Nakajima, T. (2008). Different mechanisms of DEHP-  
39 induced hepatocellular adenoma tumorigenesis in wild-type and Ppar alpha-null mice. *J*  
40 *Occup Health*, 50, 169-180. <http://www.ncbi.nlm.nih.gov/pubmed/18403868>  
41 Takeda, S.; Kondo, M.; Kumada, T.; Koshikawa, T.; Ueda, R.; Nishio, M., . . . Nakao, A. (1996).  
42 Allelic-expression imbalance of the insulin-like growth factor 2 gene in hepatocellular  
43 carcinoma and underlying disease. *Oncogene*, 12, 1589-1592.  
44 <http://www.ncbi.nlm.nih.gov/pubmed/8622877>

- 1 Takubo, K. and Kaminishi, M. (2001). [Diseases of the digestive tract and telomere lengths:  
2 significance and problems of telomere measurement]. *Nippon Shokakibyo Gakkai*  
3 *Zasshi*, 98, 144-150. <http://www.ncbi.nlm.nih.gov/pubmed/11235188>
- 4 Tanaka, K.; Smith, P. F.; Stromberg, P. C.; Eydeloth, R. S.; Herold, E. G.; Grossman, S. J., . . .  
5 Keenan, K. P. (1992). Studies of early hepatocellular proliferation and peroxisomal  
6 proliferation in Sprague-Dawley rats treated with tumorigenic doses of clofibrate.  
7 *Toxicol Appl Pharmacol*, 116, 71-77. <http://www.ncbi.nlm.nih.gov/pubmed/1529455>
- 8 Tang, X.; Li, L.; Huang, J.; Deng, Y. (2002). Guinea pig maximization test for trichloroethylene  
9 and its metabolites. *Biomed Environ Sci*, 15, 113-118.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/12244752>
- 11 Tang, X.; Que, B.; Song, X.; Li, S.; Yang, X.; Wang, H., . . . Li, L. (2008). Characterization of  
12 liver injury associated with hypersensitive skin reactions induced by trichloroethylene in  
13 the guinea pig maximization test. *J Occup Health*, 50, 114-121.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/18403861>
- 15 Tanguay, R.; Jorquera, R.; Poudrier, J.; St-Louis, M. (1996). Tyrosine and its catabolites: from  
16 disease to cancer. *Acta Biochim Pol*, 43, 209-216.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/8790725>
- 18 Tao, L.; Kramer, P. M.; Ge, R.; Pereira, M. A. (1998). Effect of dichloroacetic acid and  
19 trichloroacetic acid on DNA methylation in liver and tumors of female B6C3F1 mice.  
20 *Toxicol Sci*, 43, 139-144. <http://dx.doi.org/10.1006/toxs.1998.2449>
- 21 Tao, L.; Li, Y.; Kramer, P.; Wang, W.; Pereira, M. (2004). Hypomethylation of DNA and the  
22 insulin-like growth factor-II gene in dichloroacetic and trichloroacetic acid-promoted  
23 mouse liver tumors. *Toxicology*, 196, 127-136.  
24 <http://dx.doi.org/10.1016/j.tox.2003.11.011>
- 25 Tao, L.; Wang, W.; Li, L.; Kramer, P.; Pereira, M. (2004). Effect of dibromoacetic acid on DNA  
26 methylation, glycogen accumulation, and peroxisome proliferation in mouse and rat liver.  
27 *Toxicol Sci*, 82, 62-69. <http://dx.doi.org/10.1093/toxsci/kfh266>
- 28 Tao, L.; Yang, S.; Xie, M.; Kramer, P. M.; Pereira, M. A. (2000). Effect of trichloroethylene and  
29 its metabolites, dichloroacetic acid and trichloroacetic acid, on the methylation and  
30 expression of c-jun and c-myc protooncogenes in mouse liver: Prevention by methionine.  
31 *Toxicol Sci*, 54, 399-407. <http://www.ncbi.nlm.nih.gov/pubmed/10774822>
- 32 Tao, L.; Yang, S.; Xie, M.; Kramer, P. M.; Pereira, M. A. (2000). Hypomethylation and  
33 overexpression of c-jun and c-myc protooncogenes and increased DNA methyltransferase  
34 activity in dichloroacetic and trichloroacetic acid-promoted mouse liver tumors. *Cancer*  
35 *Lett*, 158, 185-193. [http://dx.doi.org/10.1016/S0304-3835\(00\)00518-8](http://dx.doi.org/10.1016/S0304-3835(00)00518-8)
- 36 Taskinen, H.; Anttila, A.; Lindbohm, M.-L.; Sallmén, M.; Hemminki, K. (1989). Spontaneous  
37 abortions and congenital malformations among the wives of men occupationally exposed  
38 to organic solvents. *Scand J Work Environ Health*, 15, 345-352.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/2799322>
- 40 Taskinen, H.; Kyyronen, P.; Hemminki, K. (1994). Laboratory work and pregnancy outcome. *J*  
41 *Occup Med*, 36, 311-319. <http://www.ncbi.nlm.nih.gov/pubmed/8195901>
- 42 Tateishi, M.; Suzuki, S.; Shimizu, H. (1978). Cysteine conjugate beta-lyase in rat liver. A novel  
43 enzyme catalyzing formation of thiol-containing metabolites of drugs. *J Biol Chem*, 253,  
44 8854-8859. <http://www.ncbi.nlm.nih.gov/pubmed/721818>

DRAFT– DO NOT QUOTE OR CITE

- 1 Tateishi, T.; Nakura, H.; Asoh, M.; Watanabe, M.; Tanaka, M.; Kumai, T., . . . Kobayashi, S.  
2 (1997). A comparison of hepatic cytochrome P450 protein expression between infancy  
3 and postinfancy. *Life Sci*, 61, 2567-2574. <http://dx.doi.org/10.1016/S0024->  
4 [3205\(97\)01011-4](http://dx.doi.org/10.1016/S0024-3205(97)01011-4)
- 5 Tatham, L.; Tolbert, P.; Kjeldsberg, C. (1997). Occupational risk factors for subgroups of non-  
6 Hodgkin's lymphoma. *Epidemiology*, 8, 551-558.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/9270958>
- 8 Taylor, D. H.; Lagory, K. E.; Zaccaro, D. J.; Pfohl, R. J.; Laurie, R. D. (1985). Effect of  
9 trichloroethylene on the exploratory and locomotor activity of rats exposed during  
10 development. *Sci Total Environ*, 47, 415-420.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/4089608>
- 12 Templin, M. V.; Parker, J. C.; Bull, R. J. (1993). Relative formation of dichloroacetate and  
13 trichloroacetate from trichloroethylene in male B6C3F1 mice. *Toxicol Appl Pharmacol*,  
14 123, 1-8. <http://dx.doi.org/10.1006/taap.1993.1214>
- 15 Templin, M. V.; Parker, J. C.; Bull, R. J. (1995). Erratum for: relative formation of  
16 dichloroacetate and trichloroacetate from trichloroethylene in male B6C3F1 mice.  
17 *Toxicol Appl Pharmacol*, 133, 177.
- 18 Templin, M. V.; Stevens, D. K.; Stenner, R. D.; Bonate, P. L.; Tuman, D.; Bull, R. J. (1995).  
19 Factors affecting species differences in the kinetics of metabolites of trichloroethylene. *J*  
20 *Toxicol Environ Health*, 44, 435-447. <http://www.ncbi.nlm.nih.gov/pubmed/7723076>
- 21 Tennant, R. W.; Elwell, M. R.; Spalding, J. W.; Griesemer, R. A. (1991). Evidence that toxic  
22 injury is not always associated with induction of chemical carcinogenesis. *Mol Carcinog*,  
23 4, 420-440. <http://dx.doi.org/10.1002/mc.2940040604>
- 24 TERRACINI, B. and PARKER, V. (1965). A PATHOLOGICAL STUDY ON THE TOXICITY  
25 OF S-DICHLOROVINYLL-CYSTEINE. *Food Cosmet Toxicol*, 3, 67-74.  
26 <http://www.ncbi.nlm.nih.gov/pubmed/14342797>
- 27 Teschke, K.; Olshan, A. F.; Daniels, J. L.; De Roos, A. J.; Parks, C. G.; Schulz, M.; Vaughan, T.  
28 L. (2002). Occupational exposure assessment in case-control studies: opportunities for  
29 improvement. *Occup Environ Med*, 59, 575-593; discussion 594.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/12205230>
- 31 Thai, S.; Allen, J.; DeAngelo, A.; George, M.; Fuscoe, J. (2001). Detection of early gene  
32 expression changes by differential display in the livers of mice exposed to dichloroacetic  
33 acid. *Carcinogenesis*, 22, 1317-1322. <http://www.ncbi.nlm.nih.gov/pubmed/11470764>
- 34 Thai, S.-F.; Allen, J. W.; DeAngelo, A. B.; George, M. H.; Fuscoe, J. C. (2003). Altered gene  
35 expression in mouse livers after dichloroacetic acid exposure. *Mutat Res Rev Mutat Res*,  
36 543, 167-180. [http://dx.doi.org/10.1016/S1383-5742\(03\)00014-0](http://dx.doi.org/10.1016/S1383-5742(03)00014-0)
- 37 Tham, R.; Bunnfors, I.; Eriksson, B.; Larsby, B.; Lindgren, S.; Odkvist, L. M. (1984). Vestibulo-  
38 ocular disturbances in rats exposed to organic solvents. *Acta Pharmacol Toxicol*, 54, 58-  
39 63. <http://dx.doi.org/10.1111/j.1600-0773.1984.tb01895.x>
- 40 Tham, R.; Larsby, B.; Odkvist, L. M.; Norlander, B.; Hyden, D.; Aschan, G.; Bertler, A. (1979).  
41 The influence of trichloroethylene and related drugs on the vestibular system. *Basic Clin*  
42 *Pharmacol Toxicol*, 44, 336-342. <http://www.ncbi.nlm.nih.gov/pubmed/314225>
- 43 Tharappel, J. B.; Cunningham, M. L.; Spear, B. T.; Glauert, H. P. (2001). Differential activation  
44 of hepatic NF- $\kappa$ B in rats and hamsters by the peroxisome proliferators Wy-14,643,  
45 gemfibrozil, and dibutyl phthalate. *Toxicol Sci*, 62, 20-27.



- 1 Thiele, D. L.; Eigenbrodt, E. H.; Ware, A. J. (1982). Cirrhosis after repeated trichloroethylene  
2 and 1,1,1-trichloroethane exposure. *Gastroenterology*, 83, 926-929.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/7106523>
- 4 Thorgeirsson, S. S. (2006). Genomic decoding of hepatocellular carcinoma. *Gastroenterology*,  
5 131, 1344-1346. <http://dx.doi.org/10.1053/j.gastro.2006.08.060>
- 6 Thrall, K. and Poet, T. (2000). Determination of biokinetic interactions in chemical mixtures  
7 using real-time breath analysis and physiologically based pharmacokinetic modeling. *J*  
8 *Toxicol Environ Health A*, 59, 653-670. <http://www.ncbi.nlm.nih.gov/pubmed/10839498>
- 9 Till, C.; Koren, G.; Rovet, J. F. (2001). Prenatal exposure to organic solvents and child  
10 neurobehavioral performance. *Neurotoxicol Teratol*, 23, 235-245.  
11 [http://dx.doi.org/10.1016/S0892-0362\(01\)00141-6](http://dx.doi.org/10.1016/S0892-0362(01)00141-6)
- 12 Till, C.; Westall, C. A.; Koren, G.; Nulman, I.; Rovet, J. F. (2005). Vision abnormalities in  
13 young children exposed prenatally to organic solvents. *Neurotoxicology*, 26, 599-613.  
14 <http://dx.doi.org/10.1016/j.neuro.2005.05.011>
- 15 Till, C.; Westall, C. A.; Rovet, J. F.; Koren, G. (2001). Effects of maternal occupational exposure  
16 to organic solvents on offspring visual functioning: A prospective controlled study.  
17 *Teratology*, 64, 134-141. <http://dx.doi.org/10.1002/tera.1056>
- 18 Tola, S.; Vilhunen, R.; Jarvinen, E.; Korkala, M.-L. (1980). A cohort study on workers exposed  
19 to trichloroethylene. *J Occup Environ Med*, 22, 737-740.
- 20 Toma, M.; Grosser, M.; Herr, A.; Aust, D.; Meye, A.; Hoefling, C., . . . Baretton, G. (2008). Loss  
21 of heterozygosity and copy number abnormality in clear cell renal cell carcinoma  
22 discovered by high-density affymetrix 10K single nucleotide polymorphism mapping  
23 array. *Neoplasia*, 10, 634-642. <http://www.ncbi.nlm.nih.gov/pubmed/18592004>
- 24 Tomisawa, H.; Ichihara, S.; Fukazawa, H.; Ichimoto, N.; Tateishi, M.; Yamamoto, I. (1986).  
25 Purification and characterization of human hepatic cysteine-conjugate beta-lyase.  
26 *Biochem J*, 235, 569-575. <http://www.ncbi.nlm.nih.gov/pubmed/3741406>
- 27 Tomisawa, H.; Suzuki, S.; Ichihara, S.; Fukazawa, H.; Tateishi, M. (1984). Purification and  
28 characterization of C-S lyase from *Fusobacterium varium*. A C-S cleavage enzyme of  
29 cysteine conjugates and some S-containing amino acids. *J Biol Chem*, 259, 2588-2593.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/6698982>
- 31 Tong, Z.; Board, P. G.; Anders, M. W. (1998). Glutathione transferase zeta catalyses the  
32 oxygenation of the carcinogen dichloroacetic acid to glyoxylic acid. *Biochem J*, 331 ( Pt  
33 2), 371-374. <http://www.ncbi.nlm.nih.gov/pubmed/9531472>
- 34 Toraason, M.; Clark, J.; Dankovic, D.; Mathias, P.; Skaggs, S.; Walker, C.; Werren, D. (1999).  
35 Oxidative stress and DNA damage in Fischer rats following acute exposure to  
36 trichloroethylene or perchloroethylene. *Toxicology*, 138, 43-53.  
37 [http://dx.doi.org/10.1016/S0300-483X\(99\)00083-9](http://dx.doi.org/10.1016/S0300-483X(99)00083-9)
- 38 Toth, B.; Hornung, D.; Scholz, C.; Djalali, S.; Friese, K.; Jeschke, U. (2007). Peroxisome  
39 proliferator-activated receptors: new players in the field of reproduction. *Am J Reprod*  
40 *Immunol*, 58, 289-310. <http://dx.doi.org/10.1111/j.1600-0897.2007.00514.x>
- 41 Triebig, G.; Bestler, W.; Baumeister, P.; Valentin, H. (1983). Investigations on neurotoxicity of  
42 chemical substances at the workplace: 4. Determination of the motor and sensory nerve  
43 conduction velocity in persons occupationally exposed to a mixture of organic solvents.  
44 *Int Arch Occup Environ Health*, 52, 139-150.

DRAFT– DO NOT QUOTE OR CITE

- 1 Triebig, G.; Essing, H.; Schaller, K.; Valentin, H. (1976). [Biochemical and psychological  
2 examinations of trichloroethylene exposed volunteers (author's transl)]. Zentralbl  
3 Bakteriologie, Parasitenkunde, Infektionskrankheiten Hygiene, Abteilung 1: Originalarbeiten, Reihe B, 163, 383-416.  
4 <http://www.ncbi.nlm.nih.gov/pubmed/1020533>
- 5 Triebig, G.; Lehl, S.; Kinzel, W.; Erzigkeit, H.; Galster, J.; Schaller, K. (1977).  
6 [Psychopathometrische Ergebnisse von Nachuntersuchungen von trichloroethylen-exponierten Personen  
7 (author's transl)]. Zentralbl Bakteriologie, Parasitenkunde, Infektionskrankheiten Hygiene, Abteilung 1: Originalarbeiten,  
8 Reihe B, 164, 314-377. <http://www.ncbi.nlm.nih.gov/pubmed/329621>
- 9 Triebig, G.; Schaller, K.; Erzigkeit, H.; Valentin, H. (1977). [Biochemical investigations and  
10 psychological studies of persons chronically exposed to trichloroethylene with regard to  
11 non-exposure intervals (author's transl)]. Int Arch Occup Environ Health, 38, 149-162.  
12 <http://www.ncbi.nlm.nih.gov/pubmed/832911>
- 13 Triebig, G.; Trautner, P.; Weltle, D.; Saure, E.; Valentin, H. (1982). [Investigations on  
14 neurotoxicity of chemical substances at the workplace. III. Determination of the motor  
15 and sensory nerve conduction velocity in persons occupationally exposed to  
16 trichloroethylene]. Int Arch Occup Environ Health, 51, 25-34.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/7152699>
- 18 Troster, A. I. and Ruff, R. M. (1990). Neuropsychological sequelae of exposure to the  
19 chlorinated hydrocarbon solvents trichloroethylene and trichloroethane. Arch Clin  
20 Neuropsychol, 5, 31-47. <http://www.ncbi.nlm.nih.gov/pubmed/14589542>
- 21 Truhaut, R. (1951). [Contribution to the study of toxicology of trichloroethylene; establishment  
22 of a method of determination applicable to atmosphere and biological media]. Ann Pharm  
23 Fr, 9, 175-187. <http://www.ncbi.nlm.nih.gov/pubmed/14838437>
- 24 Tryndyak, V. P.; Muskhelishvili, L.; Kovalchuk, O.; Rodriguez-Juarez, R.; Montgomery, B.;  
25 Churchwell, M. I., . . . Pogribny, I. P. (2006). Effect of long-term tamoxifen exposure on  
26 genotoxic and epigenetic changes in rat liver: implications for tamoxifen-induced  
27 hepatocarcinogenesis. Carcinogenesis, 27, 1713-1720.  
28 <http://dx.doi.org/10.1093/carcin/bgl050>
- 29 Tu, A. S.; Murray, T. A.; Hatch, K. M.; Sivak, A.; Milman, H. A. (1985). In vitro transformation  
30 of BALB/c-3T3 cells by chlorinated ethanes and ethylenes. Cancer Lett, 28, 85-92.  
31 [http://dx.doi.org/10.1016/0304-3835\(85\)90096-5](http://dx.doi.org/10.1016/0304-3835(85)90096-5)
- 32 Tucker, A. N.; Sanders, V. M.; Barnes, D. W.; Bradshaw, T. J.; White, K. L. J.; Sain, L. E., . . .  
33 Munson, A. E. (1982). Toxicology of trichloroethylene in the mouse. Toxicol Appl  
34 Pharmacol, 62, 351-357. <http://www.ncbi.nlm.nih.gov/pubmed/7071854>
- 35 Tugwood, J. D.; Aldridge, T. C.; Lambe, K. G.; Macdonald, N.; Woodyatt, N. J. (1996).  
36 Peroxisome proliferator-activated receptors: structures and function. Ann N Y Acad Sci,  
37 804, 252-265.
- 38 Tzeng, H. F.; Blackburn, A. C.; Board, P. G.; Anders, M. W. (2000). Polymorphism- and  
39 species-dependent inactivation of glutathione transferase zeta by dichloroacetate. Chem  
40 Res Toxicol, 13, 231-236. <http://dx.doi.org/10.1021/tx990175q>
- 41 U.S. EPA. (U.S. Environmental Protection Agency). (1980). Ambient water quality criteria for  
42 trichloroethylene. (EPA/440/5-80/077). Washington, DC: U.S. Environmental Protection  
43 Agency, Office of Water Regulations and Standards. Retrieved from  
44 <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=2000M4KP.txt>.

DRAFT– DO NOT QUOTE OR CITE

- 1 U.S. EPA. (U.S. Environmental Protection Agency). (1984). Health assessment document for  
2 1,1,1-trichloroethane (methyl chloroform) [final report].
- 3 U.S. EPA. (U.S. Environmental Protection Agency). (1985). Health assessment document for  
4 trichloroethylene [final report]. (EPA/600/8-82/006F). Research Triangle Park, NC: U.S.  
5 Environmental Protection Agency, Office of Research and Development. Retrieved from  
6 <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=2000BFCV.txt>.
- 7 U.S. EPA. (U.S. Environmental Protection Agency). (1987). Addendum to the health assessment  
8 document for trichloroethylene: updated carcinogenicity assessment for trichloroethylene  
9 [review draft]. (EPA/600/8-82/006FA). Washington, DC: U.S. Environmental Protection  
10 Agency, Office of Research and Development. Retrieved from  
11 <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=300016WQ.txt>.
- 12 U.S. EPA. (U.S. Environmental Protection Agency). (1992). Dermal exposure assessment:  
13 Principles and applications. (EPA/600/8-91/011B). Washington, DC: U.S. Environmental  
14 Protection Agency, Office of Health and Environmental Assessment. Retrieved from  
15 <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=12188>.
- 16 U.S. EPA. (U.S. Environmental Protection Agency). (1994). Interim policy for particle size and  
17 limit concentration issues in inhalation toxicity studies. Washington, DC: U.S.  
18 Environmental Protection Agency, Office of Pesticide Products, Health Effects Division.  
19 Retrieved from <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=186068>.
- 20 U.S. EPA. (U.S. Environmental Protection Agency). (1994). Methods for derivation of inhalation  
21 reference concentrations and application of inhalation dosimetry. (EPA/600/8-90/066F).  
22 Washington, DC: U.S. Environmental Protection Agency, Office of Research and  
23 Development. Retrieved from  
24 <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=71993>.
- 25 U.S. EPA. (U.S. Environmental Protection Agency). (1995). National public water system  
26 supervision program: FY 1994 compliance report. (EPA 812-R-95-001). Washington,  
27 DC: Author. Retrieved from  
28 <http://www.ntis.gov/search/product.aspx?ABBR=PB95270534>.
- 29 U.S. EPA. (U.S. Environmental Protection Agency). (1997). Exposure factors handbook.  
30 (EPA/600/P-95/002FA-C). Washington, DC: U.S. Environmental Protection Agency,  
31 Office of Research and Development. Retrieved from  
32 <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=12464>.
- 33 U.S. EPA. (U.S. Environmental Protection Agency). (1997). Exposure factors handbook: Vol 1 -  
34 General factors. (EPA/600/P-95/002FA). Washington, DC: U.S. Environmental  
35 Protection Agency, Office of Research and Development. Retrieved from [http://epa-  
36 prgs.ornl.gov/radionuclides/EFH\\_Final\\_1997\\_EPA600P95002Fa.pdf](http://epa-prgs.ornl.gov/radionuclides/EFH_Final_1997_EPA600P95002Fa.pdf).
- 37 U.S. EPA. (U.S. Environmental Protection Agency). (1998). EPA data file: Office of  
38 groundwater and drinking water. Washington, DC: Author.
- 39 U.S. EPA. (U.S. Environmental Protection Agency). (2000). Benchmark dose technical guidance  
40 document [external review draft]. (EPA/630/R-00/001). Washington, DC: U.S.  
41 Environmental Protection Agency, Risk Assessment Forum. Retrieved from  
42 <http://www.epa.gov/raf/publications/benchmark-dose-doc-draft.htm>.
- 43 U.S. EPA. (U.S. Environmental Protection Agency). (2000). Science policy council handbook:  
44 Risk characterization. (EPA 100-B-00-002). Washington, D.C.: U.S. Environmental

DRAFT– DO NOT QUOTE OR CITE

- 1 Protection Agency, Office of Research and Development, Office of Science Policy.  
2 Retrieved from <http://www.epa.gov/osa/spc/pdfs/rchandbk.pdf>.
- 3 U.S. EPA. (U.S. Environmental Protection Agency). (2001). Sources, emission and exposure for  
4 trichloroethylene (TCE) and related chemicals. (EPA/600/R-00/009). Washington, DC:  
5 U.S. Environmental Protection Agency, Office of Research and Development.
- 6 U.S. EPA. (U.S. Environmental Protection Agency). (2002). A review of the reference dose and  
7 reference concentration processes. (EPA/630/P-02/0002F). Washington, DC: U.S.  
8 Environmental Protection Agency, Risk Assessment Forum. Retrieved from  
9 <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=51717>.
- 10 U.S. EPA. (U.S. Environmental Protection Agency). (2002). Health assessment of 1,3-butadiene.  
11 (EPA/600/P-98/001F). Washington, DC: U.S. Environmental Protection Agency, Office  
12 of Research and Development.
- 13 U.S. EPA. (U.S. Environmental Protection Agency). (2002). OSWER draft guidance for  
14 evaluating the vapor intrusion to indoor air pathway from groundwater and soils  
15 (subsurface vapor intrusion guidance). (EPA 530-D-02-004). Washington, DC: Author.  
16 Retrieved from <http://epa.gov/osw/hazard/correctiveaction/eis/vapor/guidance.pdf>.
- 17 U.S. EPA. (U.S. Environmental Protection Agency). (2003). Framework for cumulative risk  
18 assessment. (EPA/630/P-02/001F). Washington, DC: U.S. Environmental Protection  
19 Agency, Risk Assessment Forum. Retrieved from  
20 <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=30004TJH.txt>.
- 21 U.S. EPA. (U.S. Environmental Protection Agency). (2003). Occurrence estimation methodology  
22 and occurrence findings report of the six-year review of existing National Primary  
23 Drinking Water Regulations. (EPA-815/R-03-006). Washington, D.C.: Author.
- 24 U.S. EPA. (U.S. Environmental Protection Agency). (2003). Toxicological review of  
25 dichloroacetic acid (CAS No. 79-43-6). (EPA 635/R-03/007). Washington, DC: Author.  
26 Retrieved from <http://www.epa.gov/iris/subst/0654.htm>.
- 27 U.S. EPA. (U.S. Environmental Protection Agency). (2004). Estimated per capita water ingestion  
28 and body weight in the United States: An update. (EPA-822/R-00-001). Washington,  
29 D.C.: Author. Retrieved from  
30 <http://www.epa.gov/waterscience/criteria/drinking/percapita/2004.pdf>.
- 31 U.S. EPA. (U.S. Environmental Protection Agency). (2005). Documentation of environmental  
32 indicator determination: RCRA corrective action. IBM-Endicott. New York, NY:  
33 Author. Retrieved from [http://www.epa.gov/Region2/waste/ibm\\_e725.pdf](http://www.epa.gov/Region2/waste/ibm_e725.pdf).
- 34 U.S. EPA. (U.S. Environmental Protection Agency). (2005). Guidance on selecting age groups  
35 for monitoring and assessing childhood exposures to environmental contaminants (Final).  
36 (EPA/630/P-03/003F). Washington, DC: U.S. Environmental Protection Agency, Risk  
37 Assessment Forum. Retrieved from  
38 <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=146583>.
- 39 U.S. EPA. (U.S. Environmental Protection Agency). (2005). Guidelines for carcinogen risk  
40 assessment. (EPA/630/P-03/001F). Washington, DC: U.S. Environmental Protection  
41 Agency, Risk Assessment Forum. Retrieved from <http://www.epa.gov/cancerguidelines/>.
- 42 U.S. EPA. (U.S. Environmental Protection Agency). (2005). Supplemental guidance for  
43 assessing susceptibility from early-life exposure to carcinogens. (EPA/630/R-03/003F).  
44 Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum.

DRAFT– DO NOT QUOTE OR CITE

- 1 Retrieved from [http://www.epa.gov/cancerguidelines/guidelines-carcinogen-](http://www.epa.gov/cancerguidelines/guidelines-carcinogen-supplement.htm)  
2 supplement.htm.
- 3 U.S. EPA. (U.S. Environmental Protection Agency). (2006). Peer review handbook (3rd edition).  
4 (EPA/100/B-06/002). Washington, DC: U.S. Environmental Protection Agency, Science  
5 Policy Council. Retrieved from <http://www.epa.gov/peerreview/>.
- 6 U.S. EPA. (U.S. Environmental Protection Agency). (2006). Technology transfer network 1999  
7 national-scale air toxics assessment, from  
8 <http://www.epa.gov/ttn/atw/nata1999/index.html>
- 9 U.S. EPA. (U.S. Environmental Protection Agency). (2007). AirData: Access to Air Pollution  
10 Data (Version 2007). Research Triangle Park, NC: Office of Air Quality Planning and  
11 Standards. Retrieved from <http://www.epa.gov/air/data/index.html>
- 12 U.S. EPA. (U.S. Environmental Protection Agency). (2008). 1,1,2-Trichloro-1,2,2-  
13 trifluoroethane. Washington, D.C.: Author. Retrieved from  
14 <http://www.epa.gov/IRIS/subst/0123.htm>.
- 15 U.S. EPA. (U.S. Environmental Protection Agency). (2008). Child-specific exposure factors  
16 handbook. (EPA/600/R-06/096F). Washington, DC: U.S. Environmental Protection  
17 Agency, National Center for Environmental Assessment. Retrieved from  
18 <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=199243>.
- 19 U.S. EPA. (U.S. Environmental Protection Agency). (2008). Factoids: Drinking water and  
20 ground water statistics for 2008. (EPA-816/K-08-004). Washington, D.C.: Author.  
21 Retrieved from [http://www.epa.gov/ogwdw000/databases/pdfs/data\\_factoids\\_2008.pdf](http://www.epa.gov/ogwdw000/databases/pdfs/data_factoids_2008.pdf).
- 22 U.S. GAO. (U.S. Government Accountability Office). (2007). Testimony before the  
23 subcommittee on oversight and investigations, committee on energy and commerce,  
24 house of representatives: Defense health care: Issues related to past drinking water  
25 contamination at marine corps base Camp Lejeune. Washington, DC: Author.
- 26 U.S. GAO. (U.S. Government Accountability Office). Report to congressional committees:  
27 Defense health care: Activities related to past drinking water contamination at marine  
28 corps base Camp Lejeune. Washington, DC: Author.
- 29 Ulrich, R. G. (2003). The toxicogenomics of nuclear receptor agonists. *Curr Opin Chem Biol*, 7,  
30 505-510. <http://www.ncbi.nlm.nih.gov/pubmed/12941426>
- 31 Ulrich, R. G.; Rockett, J. C.; Gibson, G. G.; Pettit, S. D. (2004). Overview of an interlaboratory  
32 collaboration on evaluating the effects of model hepatotoxicants on hepatic gene  
33 expression. *Environ Health Perspect*, 112, 423-427.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/15033591>
- 35 Umezu, T.; Yonemoto, J.; Soma, Y.; Miura, T. (1997). Behavioral effects of trichloroethylene  
36 and tetrachloroethylene in mice. *Pharmacol Biochem Behav*, 58, 665-671.  
37 [http://dx.doi.org/10.1016/S0091-3057\(97\)00046-4](http://dx.doi.org/10.1016/S0091-3057(97)00046-4)
- 38 Urabe, Y.; Nouse, K.; Higashi, T.; Nakatsukasa, H.; Hino, N.; Ashida, K., . . . Tsuji, T. (1996).  
39 Telomere length in human liver diseases. 16, 293-297.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/8938628>
- 41 USAF. (U.S. Air Force). (2004). Determination of partition coefficients for a mixture of volatile  
42 organic compounds in rats and humans at different life stages. (AFRL-HE-WP-TR-2005-  
43 0012). Wright-Patterson Air Force Base, OH: U.S. Air Force Research Laboratory.

DRAFT– DO NOT QUOTE OR CITE

- 1 USGS. (U.S. Geological Survey). (2006). Volatile organic compounds in the nation's ground  
2 water and drinking water supply wells. Reston, VA: Author. Retrieved from  
3 <http://pubs.usgs.gov/circ/circ1292/pdf/circular1292.pdf>.
- 4 Vagnarelli, P.; De Sario, A.; De Carli, L. (1990). Aneuploidy induced by chloral hydrate  
5 detected in human lymphocytes with the Y97 probe. *Mutagenesis*, 5, 591-592.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/2263216>
- 7 Vaidya, V.; Shankar, K.; Lock, E.; Bucci, T.; Mehendale, H. (2003). Renal injury and repair  
8 following S-1, 2 dichlorovinyl-L-cysteine administration to mice. *Toxicol Appl*  
9 *Pharmacol*, 188, 110-121. <http://www.ncbi.nlm.nih.gov/pubmed/12691729>
- 10 Vaidya, V.; Shankar, K.; Lock, E.; Bucci, T.; Mehendale, H. (2003). Role of tissue repair in  
11 survival from s-(1,2-dichlorovinyl)-L-cysteine-induced acute renal tubular necrosis in the  
12 mouse. *Toxicol Sci*, 74, 215-227. <http://dx.doi.org/10.1093/toxsci/kfg111>
- 13 Vaidya, V.; Shankar, K.; Lock, E.; Dixon, D.; Mehendale, H. (2003). Molecular mechanisms of  
14 renal tissue repair in survival from acute renal tubule necrosis: role of ERK1/2 pathway.  
15 *Toxicol Pathol*, 31, 604-618. <http://www.ncbi.nlm.nih.gov/pubmed/14585728>
- 16 Vainio, H.; Linnainmaa, K.; Kähönen, M.; Nickels, J.; Hietanen, E.; Marniemi, J.; Peltonen, P.  
17 (1983). Hypolipidemia and peroxisome proliferation induced by phenoxyacetic acid  
18 herbicides in rats. *Biochem Pharmacol*, 32, 2775-2779.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/6626247>
- 20 Vainio, H.; Nickels, J.; Linnainmaa, K. (1982). Phenoxy acid herbicides cause peroxisome  
21 proliferation in Chinese hamsters. *Scand J Work Environ Health*, 8, 70-73.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/7134925>
- 23 Valic, E.; Waldhor, T.; Konnaris, C.; Michitsch, A.; Wolf, C. (1997). Acquired dyschromatopsia  
24 in combined exposure to solvents and alcohol. *Int Arch Occup Environ Health*, 70, 403-  
25 406. <http://www.ncbi.nlm.nih.gov/pubmed/9439987>
- 26 Vamvakas, S.; Bittner, D.; Köster, U. (1993). Enhanced expression of the protooncogenes c-myc  
27 and c-fos in normal and malignant renal growth. *Toxicol Lett*, 67, 161-172.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/8451757>
- 29 Vamvakas, S.; Bruning, T.; Thomasson, B.; Lammert, M.; Baumuller, A.; Bolt, H. M., . . . Ulm,  
30 K. (1998). Renal cell cancer correlated with occupational exposure to trichloroethene. *J*  
31 *Cancer Res Clin Oncol*, 124, 374-382.
- 32 Vamvakas, S.; Dekant, W.; Berthold, K.; Schmidt, S.; Wild, D.; Henschler, D. (1987).  
33 Enzymatic transformation of mercapturic acids derived from halogenated alkenes to  
34 reactive and mutagenic intermediates. *Biochem Pharmacol*, 36, 2741-2748.  
35 [http://dx.doi.org/10.1016/0006-2952\(87\)90258-9](http://dx.doi.org/10.1016/0006-2952(87)90258-9)
- 36 Vamvakas, S.; Dekant, W.; Henschler, D. (1989). Assessment of unscheduled DNA synthesis in  
37 a cultured line of renal epithelial cells exposed to cysteine S-conjugates of haloalkenes  
38 and haloalkanes. *Mutat Res*, 222, 329-335. [http://dx.doi.org/10.1016/0165-](http://dx.doi.org/10.1016/0165-1218(89)90108-0)  
39 [1218\(89\)90108-0](http://dx.doi.org/10.1016/0165-1218(89)90108-0)
- 40 Vamvakas, S.; Dekant, W.; Schiffmann, D.; Henschler, D. (1988). Induction of unscheduled  
41 DNA synthesis and micronucleus formation in Syrian hamster embryo fibroblasts treated  
42 with cysteine S-conjugates of chlorinated hydrocarbons. *Cell Biol Toxicol*, 4, 393-403.  
43 <http://dx.doi.org/10.1007/BF00117768>

DRAFT– DO NOT QUOTE OR CITE

- 1 Vamvakas, S.; Elfarra, A.; Dekant, W.; Henschler, D.; Anders, M. (1988). Mutagenicity of  
2 amino acid and glutathione S-conjugates in the Ames test. *Mutat Res*, 206, 83-90.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/2901035>
- 4 Vamvakas, S.; Richter, H.; Bittner, D. (1996). Induction of dedifferentiated clones of LLC-PK1  
5 cells upon long-term exposure to dichlorovinylcysteine. *Toxicology*, 106, 65-74.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/8571403>
- 7 van Bree, L.; Commandeur, J.; Lamberts, B.; Cornelissen, M.; van Roon, M.; Laterveer, H.; de  
8 Vries, J. (1990). Induction of drug metabolism enzymes by dihalogenated biphenyls. *J*  
9 *Biochem Toxicol*, 5, 57-63. <http://dx.doi.org/10.1002/jbt.2570050109>
- 10 Van Hummelen, P. and Kirsch-Volders, M. (1992). Analysis of eight known or suspected  
11 aneugens by the in vitro human lymphocyte micronucleus test. *Mutagenesis*, 7, 447-455.  
12 <http://dx.doi.org/10.1093/mutage/7.6.447>
- 13 van Lieshout, E. M. and Peters, W. H. (1998). Age and gender dependent levels of glutathione  
14 and glutathione S-transferases in human lymphocytes. *Carcinogenesis*, 19, 1873-1875.  
15 <http://www.ncbi.nlm.nih.gov/pubmed/9806172>
- 16 van Lieshout, E.; Knapen, M.; Lange, W.; Steegers, E.; Peters, W. (1998). Localization of  
17 glutathione S-transferases alpha and pi in human embryonic tissues at 8 weeks gestational  
18 age. *Hum Reprod*, 13, 1380-1386. <http://www.ncbi.nlm.nih.gov/pubmed/9647576>
- 19 Van Duuren, B. L.; Goldschmidt, B. M.; Loewengart, G.; Smith, A. C.; Melchionne, S.;  
20 Seidman, I.; Roth, D. (1979). Carcinogenicity of halogenated olefinic and aliphatic  
21 hydrocarbons in mice. *J Natl Cancer Inst*, 63, 1433-1439.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/292813>
- 23 Vartiainen, T.; Pukkala, E.; Rienoja, T.; Strandman, T.; Kaksonen, K. (1993). Population  
24 exposure to tri- and tetrachloroethene and cancer risk: Two cases of drinking water  
25 pollution. *Chemosphere*, 27, 1171-1181. [http://dx.doi.org/10.1016/0045-6535\(93\)90165-](http://dx.doi.org/10.1016/0045-6535(93)90165-2)  
26 [2](http://dx.doi.org/10.1016/0045-6535(93)90165-2)
- 27 Veeramachaneni, D.; Palmer, J.; Amann, R. (2001). Long-term effects on male reproduction of  
28 early exposure to common chemical contaminants in drinking water. *Hum Reprod*, 16,  
29 979-987. <http://www.ncbi.nlm.nih.gov/pubmed/11331649>
- 30 Verma, Y. and Rana, S. (2003). Gender differences in the metabolism of benzene, toluene and  
31 trichloroethylene in rat with special reference to certain biochemical parameters. *J*  
32 *Environ Biol*, 24, 135-140. <http://www.ncbi.nlm.nih.gov/pubmed/12974453>
- 33 Vernon, R. J. and Ferguson, R. K. (1969). Effects of trichloroethylene on visual-motor  
34 performance. *Arch Environ Occup Health*, 18, 894-900.  
35 <http://www.ncbi.nlm.nih.gov/pubmed/5770690>
- 36 Vesterberg, O. and Astrand, I. (1976). Exposure to trichloroethylene monitored by analysis of  
37 metabolites in blood and urine. *J Occup Med*, 18, 224-226.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/1271124>
- 39 Vesterberg, O.; Gorczak, J.; Krasts, M. (1976). Exposure to trichloroethylene II Metabolites in  
40 blood and urine. *Scand J Work Environ Health*, 4, 212-219.
- 41 Vickers, A. E. and Lucier, G. W. (1996). Estrogen receptor levels and occupancy in hepatic  
42 sinusoidal endothelial and Kupffer cells are enhanced by initiation with  
43 diethylnitrosamine and promotion with 17alpha-ethinylestradiol in rats. *Carcinogenesis*,  
44 17, 1235-1242. <http://www.ncbi.nlm.nih.gov/pubmed/8681437>

- 1 Vieira, I.; Sonnier, M.; Cresteil, T. (1996). Developmental expression of CYP2E1 in the human  
2 liver: Hypermethylation control of gene expression during the neonatal period. *Eur J*  
3 *Biochem*, 238, 476-483. <http://dx.doi.org/10.1111/j.1432-1033.1996.0476z.x>
- 4 Vielhauer, V.; Sarafoff, M.; Gais, P.; Rabes, H. M. (2001). Cell type-specific induction of O6-  
5 alkylguanine-DNA alkyltransferase mRNA expression in rat liver during regeneration,  
6 inflammation and preneoplasia. *J Cancer Res Clin Oncol*, 127, 591-602.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/11599795>
- 8 Vig, P.; Russo, F. P.; Edwards, R. J.; Tadrous, P. J.; Wright, N. A.; Thomas, H. C., . . . Forbes, S.  
9 J. (2006). The sources of parenchymal regeneration after chronic hepatocellular liver  
10 injury in mice. *Hepatology*, 43, 316-324. <http://dx.doi.org/10.1002/hep.21018>
- 11 Villaschi, S.; Giovanetti, A.; Lombardi, C. C.; Nicolai, G.; Garbati, M.; Androzzzi, U. (1991).  
12 Damage and repair of mouse bronchial epithelium following acute inhalation of  
13 trichloroethylene. *Exp Lung Res*, 17, 601-614.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/1860457>
- 15 Vineis, P.; Miligi, L.; Costantini, A. (2007). Exposure to solvents and risk of non-Hodgkin  
16 lymphoma: clues on putative mechanisms. *Cancer Epidemiol Biomarkers Prev*, 16, 381-  
17 384. <http://dx.doi.org/10.1158/1055-9965.EPI-07-0124>
- 18 Vogelstein, B. and Kinzler, K. W. (2004). Cancer genes and the pathways they control. *Nat Med*,  
19 10, 789-799. <http://dx.doi.org/10.1038/nm1087>
- 20 Vollmar, B.; Pradarutti, S.; Richter, S.; Menger, M. D. (2002). In vivo quantification of ageing  
21 changes in the rat liver from early juvenile to senescent life. 22, 330-341.  
22 <http://www.ncbi.nlm.nih.gov/pubmed/12296967>
- 23 Von Tungeln, L. S.; Yi, P.; Bucci, T. J.; Samokyszyn, V. M.; Chou, M. W.; Kadlubar, F. F.; Fu,  
24 P. P. (2002). Tumorigenicity of chloral hydrate, trichloroacetic acid, trichloroethanol,  
25 malondialdehyde, 4-hydroxy-2-nonenal, crotonaldehyde, and acrolein in the B6C3F1  
26 neonatal mouse. *Cancer Lett*, 185, 13-19. [http://dx.doi.org/10.1016/S0304-  
27 3835\(02\)00231-8](http://dx.doi.org/10.1016/S0304-3835(02)00231-8)
- 28 Voss, C.; Zerban, H.; Bannasch, P.; Berger, M. R. (2005). Lifelong exposure to di-(2-  
29 ethylhexyl)-phthalate induces tumors in liver and testes of Sprague-Dawley rats.  
30 *Toxicology*, 206, 359-371. <http://dx.doi.org/10.1016/j.tox.2004.07.016>
- 31 Voss, K. A.; Riley, R.; Dunn, C.; Christopher Corton, J. (2006). The role of tumor necrosis factor  
32 alpha and the peroxisome proliferator-activated receptor alpha in modulating the effects  
33 of fumonisin in mouse liver. *Toxicology*, 222, 165-174.  
34 <http://dx.doi.org/10.1016/j.tox.2006.02.012>
- 35 Wada, N.; Marsman, D. S.; Popp, J. A. (1992). Dose-related effects of the hepatocarcinogen,  
36 Wy-14,643, on peroxisomes and cell replication. *Fundam Appl Toxicol*, 18, 149-154.  
37 <http://www.ncbi.nlm.nih.gov/pubmed/1601204>
- 38 Wainfan, E. and Poirier, L. A. (1992). Methyl groups in carcinogenesis: effects on DNA  
39 methylation and gene expression. *Cancer Res*, 52, 2071s-2077s.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/1544143>
- 41 Walgren, J. E.; Kurtz, D. T.; McMillan, J. M. (2000). Expression of PPAR(alpha) in human  
42 hepatocytes and activation by trichloroacetate and dichloroacetate. *Res Commun Mol*  
43 *Pathol Pharmacol*, 108, 116-132. <http://www.ncbi.nlm.nih.gov/pubmed/11758968>
- 44 Walgren, J. E.; Kurtz, D. T.; McMillan, J. M. (2000). The effect of the trichloroethylene  
45 metabolites trichloroacetate and dichloroacetate on peroxisome proliferation and DNA



- 1 synthesis in cultured human hepatocytes. *Cell Biol Toxicol*, 16, 257-273.  
2 <http://dx.doi.org/10.1023/A:1007638227821>
- 3 Wallace, L. A.; Pellizzari, E.; Hartwell, T.; Rosenzweig, M.; Erickson, M.; Sparacino, C.; Zelon,  
4 H. (1984). Personal exposure to volatile organic compounds: 1. Direct measurements in  
5 breathing-zone air, drinking water, food and exhaled breath. *Environ Res*, 35, 293-319.
- 6 Walles, S. A. S. (1986). Induction of single-strand breaks in dna of mice by trichloroethylene and  
7 tetrachloroethylene. *Toxicol Lett*, 31, 31-35. [http://dx.doi.org/10.1016/0378-](http://dx.doi.org/10.1016/0378-4274(86)90191-8)  
8 [4274\(86\)90191-8](http://dx.doi.org/10.1016/0378-4274(86)90191-8)
- 9 Wang, G. and Stacey, N. H. (1990). Elevation of individual serum bile acids on exposure to  
10 trichloroethylene or "alpha"-naphthylisothiocyanate. *Toxicol Appl Pharmacol*, 105, 209-  
11 215. <http://www.ncbi.nlm.nih.gov/pubmed/2219116>
- 12 Wang, G.; Ansari, G. A.; Khan, M. F. (2007). Involvement of lipid peroxidation-derived  
13 aldehyde-protein adducts in autoimmunity mediated by trichloroethene. *J Toxicol*  
14 *Environ Health A*, 70, 1977-1985. <http://dx.doi.org/10.1080/15287390701550888>
- 15 Wang, G.; Cai, P.; Ansari, G. A. S.; Khan, M. F. (2007). Oxidative and nitrosative stress in  
16 trichloroethene-mediated autoimmune response. *Toxicology*, 229, 186-193.  
17 <http://dx.doi.org/10.1016/j.tox.2006.10.014>
- 18 Wang, G.; König, R.; Ansari, G. A.; Khan, M. F. (2008). Lipid peroxidation-derived aldehyde-  
19 protein adducts contribute to trichloroethene-mediated autoimmunity via activation of  
20 CD4+ T cells. *Free Radic Biol Med*, 44, 1475-1482.  
21 <http://dx.doi.org/10.1016/j.freeradbiomed.2008.01.012>
- 22 Wang, G.; Takano, T.; Tomita, K.; Nakata, K.; Nakamura, K. (1993). [Interaction of  
23 trichloroethylene and isopropyl alcohol in the perfused rat liver]. *Nippon Eiseigaku*  
24 *Zasshi*, 48, 1000-1005. <http://www.ncbi.nlm.nih.gov/pubmed/8107291>
- 25 Wang, J.-D. (2004). Increased mortality odds ratio of male liver cancer in a community  
26 contaminated by chlorinated hydrocarbons in groundwater. *Symposium on New Scientific*  
27 *Research Related to the Health Effects of Trichloroethylene*. Washington, DC.
- 28 Wang, J.-L.; Chen, W.-L.; Tsai, S.-Y.; Sung, P.-Y.; Huang, R.-N. (2001). An in vitro model for  
29 evaluation of vaporous toxicity of trichloroethylene and tetrachloroethylene to CHO-K1  
30 cells. *Chem Biol Interact*, 137, 139-154. [http://dx.doi.org/10.1016/S0009-](http://dx.doi.org/10.1016/S0009-2797(01)00226-5)  
31 [2797\(01\)00226-5](http://dx.doi.org/10.1016/S0009-2797(01)00226-5)
- 32 Wang, R.; Nakajima, T.; Honma, T. (1999). Trichloroethylene inhibits aldehyde dehydrogenase  
33 only for aliphatic aldehydes of short chains in rats. *Toxicology*, 132, 9-18.  
34 <http://www.ncbi.nlm.nih.gov/pubmed/10199577>
- 35 Wang, R.; Zhang, Y.; Lan, Q.; Holford, T. R.; Leaderer, B.; Zahm, S. H., . . . Zheng, T. (2009).  
36 Occupational exposure to solvents and risk of non-Hodgkin lymphoma in Connecticut  
37 women. *Am J Epidemiol*, 169, 176-185. <http://dx.doi.org/10.1093/aje/kwn300>
- 38 Wang, Z.; Hall, S. D.; Maya, J. F.; Li, L.; Asghar, A.; Gorski, J. C. (2003). Diabetes mellitus  
39 increases the in vivo activity of cytochrome P450 2E1 in humans. *Br J Clin Pharmacol*,  
40 55, 77-85. <http://www.ncbi.nlm.nih.gov/pubmed/12534643>
- 41 Wanson, J.-C.; Bernaert, D.; Penasse, W.; Mosselmans, R.; Bannasch, P. (1980). Separation in  
42 distinct subpopulations by elutriation of liver cells following exposure of rats to N-  
43 nitrosomorpholine. *Cancer Res*, 40, 459-471.
- 44 Ward, E. M.; Thun, M. J.; Hannan, L. M.; Jemal, A. (2006). Interpreting cancer trends. *Ann N Y*  
45 *Acad Sci*, 1076, 29-53. <http://dx.doi.org/10.1196/annals.1371.048>

DRAFT– DO NOT QUOTE OR CITE

- 1 Ward, J. M.; Peters, J. M.; Perella, C. M.; Gonzalez, F. J. (1998). Receptor and nonreceptor-  
2 mediated organ-specific toxicity of di(2-ethylhexyl)phthalate (DEHP) in peroxisome  
3 proliferator-activated receptor alpha-null mice. *Toxicol Pathol*, 26, 240-246.
- 4 Warr, T.; Parry, E.; Parry, J. (1993). A comparison of two in vitro mammalian cell cytogenetic  
5 assays for the detection of mitotic aneuploidy using 10 known or suspected aneugens.  
6 *Mutat Res*, 287, 29-46. <http://www.ncbi.nlm.nih.gov/pubmed/7683382>
- 7 Warren, D. A.; Graeter, L. J.; Channel, S. R.; Eggers, J. S.; Goodyear, C. D.; Macmahon, K. L., .  
8 . . Baker, W. H. (2006). Trichloroethylene, trichloroacetic acid, and dichloroacetic acid:  
9 do they affect eye development in the Sprague-Dawley rat. *Int J Toxicol*, 25, 279-284.  
10 <http://dx.doi.org/10.1080/10915810600745975>
- 11 Wartenberg, D.; Reyner, D.; Scott, C. S. (2000). Trichloroethylene and cancer: Epidemiologic  
12 evidence. *Environ Health Perspect*, 108, 161-176.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/10807550>
- 14 Waseem, M.; Ali, M.; Dogra, S.; Dutta, K.; Kaw, J. (2001). Toxicity of trichloroethylene  
15 following inhalation and drinking contaminated water. *J Appl Toxicol*, 21, 441-444.  
16 <http://www.ncbi.nlm.nih.gov/pubmed/11746189>
- 17 Waskell, L. (1978). A study of the mutagenicity of anesthetics and their metabolites. *DNA*  
18 *Repair*, 57, 141-153. <http://www.ncbi.nlm.nih.gov/pubmed/351387>
- 19 Watanabe, S. and Fukui, T. (2000). Suppressive effect of curcumin on trichloroethylene-induced  
20 oxidative stress. *J Nutr Sci Vitaminol*, 46, 230-234.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/11234915>
- 22 Watanabe, T.; Shimada, H.; Tanaka, Y. (1978). Human hepatocytes and aging: a  
23 cytophotometrical analysis in 35 sudden-death cases. *Virchows Arch B Cell Pathol Incl*  
24 *Mol Pathol*, 27, 307-316. <http://www.ncbi.nlm.nih.gov/pubmed/98902>
- 25 Watson, R. E. (2002). Effects of Phenobarbital on DNA Methylation in GC-Rich Regions of  
26 Hepatic DNA from Mice That Exhibit Different Levels of Susceptibility to Liver  
27 Tumorigenesis. *Toxicol Sci*, 68, 51-58. <http://dx.doi.org/10.1093/toxsci/68.1.51>
- 28 Watson, R. E.; Jacobson, C. F.; Williams, A. L.; Howard, W. B.; DeSesso, J. M. (2006).  
29 Trichloroethylene-contaminated drinking water and congenital heart defects: a critical  
30 analysis of the literature. *Reprod Toxicol*, 21, 117-147.  
31 <http://dx.doi.org/10.1016/j.reprotox.2005.07.013>
- 32 Waxman, S. and Wurbach, E. (2007). De-regulation of common housekeeping genes in  
33 hepatocellular carcinoma. *BMC Genomics*, 8, 243. [http://dx.doi.org/10.1186/1471-2164-](http://dx.doi.org/10.1186/1471-2164-8-243)  
34 [8-243](http://dx.doi.org/10.1186/1471-2164-8-243)
- 35 Weiderpass, E.; Pukkala, E.; Kauppinen, T.; Mutanen, P.; Paakkulainen, H.; Vasama-Neuvonen,  
36 K., . . . Partanen, T. (1999). Breast cancer and occupational exposures in women in  
37 Finland. *Am J Ind Med*, 36, 48-53. <http://www.ncbi.nlm.nih.gov/pubmed/10361586>
- 38 Weiderpass, E.; Vainio, H.; Kauppinen, T.; Vasama-Neuvonen, K.; Partanen, T.; Pukkala, E.  
39 (2003). Occupational exposures and gastrointestinal cancers among Finnish women. *J*  
40 *Occup Environ Med*, 45, 305-315. <http://www.ncbi.nlm.nih.gov/pubmed/12661188>
- 41 Weidman, J. R.; Dolinoy, D. C.; Murphy, S. K.; Jirtle, R. L. (2007). Cancer susceptibility:  
42 epigenetic manifestation of environmental exposures. *Cancer J*, 13, 9-16.  
43 <http://dx.doi.org/10.1097/PPO.0b013e31803c71f2>
- 44 Weinberg, H. S.; Krasner, S. W.; Richardson, S. D.; Thruston, A. D. J. (2002). The occurrence of  
45 disinfection by-products (DBPs) of health concern in drinking water: results of a

DRAFT– DO NOT QUOTE OR CITE

- 1 nationwide DBP occurrence study. (EPA/600/R-02/068). Athens, GA: U.S.  
2 Environmental Protection Agency, National Exposure Research Laboratory.
- 3 Weisel, C. P. and Jo, W.-K. (1996). Ingestion, inhalation, and dermal exposures to chloroform  
4 and trichloroethene from tap water. *Environ Health Perspect*, 104, 48-51.
- 5 Weisel, C. P.; Kim, H.; Haltmeier, P.; Klotz, J. B. (1999). Exposure estimates to disinfection by-  
6 products of chlorinated drinking water. *Environ Health Perspect*, 107, 103-110.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/9924004>
- 8 Weisenburger, D. D. (1992). Pathological classification of non-Hodgkin's lymphoma for  
9 epidemiological studies. *Cancer Res*, 52, 5456s-5462s.
- 10 Weiss, J. R.; Moysich, K. B.; Swede, H. (2005). Epidemiology of male breast cancer. *Cancer*  
11 *Epidemiol Biomarkers Prev*, 14, 20-26. <http://www.ncbi.nlm.nih.gov/pubmed/15668471>
- 12 Weiss, P. A. (1971). A cell is not an island entire of itself. *Perspect Biol Med*, 14, 182-205.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/4925931>
- 14 Weistenhofer, W.; Blaszkewicz, M.; Bolt, H. M.; Golka, K. (2008). N-acetyltransferase-2 and  
15 medical history in bladder cancer cases with a suspected occupational disease (BK 1301)  
16 in Germany. *J Toxicol Environ Health A*, 71, 906-910.  
17 <http://dx.doi.org/10.1080/15287390801988681>
- 18 Wells, G. M.; Schroth, W.; Brauch, H.; Ross, E. A. (2009). Bilateral renal-cell carcinoma  
19 associated with an acquired VHL mutation and long-term trichloroethylene exposure.  
20 *Clin Nephrol*, 71, 708-713. <http://www.ncbi.nlm.nih.gov/pubmed/19473641>
- 21 Wendling, O.; Chambon, P.; Mark, M. (1999). Retinoid X receptors are essential for early mouse  
22 development and placentogenesis. *PNAS*, 96, 547-551.  
23 <http://www.ncbi.nlm.nih.gov/pubmed/9892670>
- 24 Werner, H. and Le Roith, D. (2000). New concepts in regulation and function of the insulin-like  
25 growth factors: implications for understanding normal growth and neoplasia. *Cell Mol*  
26 *Life Sci*, 57, 932-942. <http://www.ncbi.nlm.nih.gov/pubmed/10950308>
- 27 Werner, M.; Birner, G.; Dekant, W. (1995). The role of cytochrome P450A1/2 in the sex-  
28 specific sulfoxidation of the hexachlorobutadiene metabolite, N-acetyl-S-  
29 (pentachlorobutadienyl)-L-cysteine in rats. *Drug Metab Dispos*, 23, 861-868.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/7493554>
- 31 Werner, M.; Birner, G.; Dekant, W. (1996). Sulfoxidation of mercapturic acids derived from tri-  
32 and tetrachloroethene by cytochromes P450 3A: A bioactivation reaction in addition to  
33 deacetylation and cysteine conjugate beta-lyase mediated cleavage. *Chem Res Toxicol*, 9,  
34 41-49. <http://dx.doi.org/10.1021/tx950075u>
- 35 Werner, M.; Guo, Z.; Birner, G.; Dekant, W.; Guengerich, F. (1995). The sulfoxidation of the  
36 hexachlorobutadiene metabolite N-acetyl-S-(1,2,3,4,4-pentachlorobutadienyl)-L-cysteine  
37 is catalyzed by human cytochrome P450 3A enzymes. *Chem Res Toxicol*, 8, 917-923.  
38 <http://www.ncbi.nlm.nih.gov/pubmed/8555406>
- 39 West, G. B.; Woodruff, W. H.; Brown, J. H. (2002). Allometric scaling of metabolic rate from  
40 molecules and mitochondria to cells and mammals. *PNAS*, 99, 2473-2478.
- 41 Westergren, I.; Kjellstrand, P.; Linder, L.; Johansson, B. (1984). Reduction of brain specific  
42 gravity in mice prenatally exposed to trichloroethylene. *Toxicol Lett*, 23, 223-226.  
43 <http://www.ncbi.nlm.nih.gov/pubmed/6506097>

DRAFT– DO NOT QUOTE OR CITE

- 1 Wheeler, C. W. and Guenther, T. M. (1990). Spectroscopic quantitation of cytochrome P-450 in  
2 human lung microsomes. *J Biochem Toxicol*, 5, 269-272.  
3 <http://www.ncbi.nlm.nih.gov/pubmed/2096222>
- 4 Wheeler, C. W.; Wrighton, S. A.; Guenther, T. M. (1992). Detection of human lung  
5 cytochromes P-450 that are immunochemically related to cytochrome P450III<sub>E1</sub> and  
6 cytochrome P-450III<sub>A</sub>. *Biochem Pharmacol*, 44, 183-186.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/1632834>
- 8 White, A. E.; Takehisa, S.; Eger EI, I. I.; Wolff, S.; Stevens, W. C. (1979). Sister chromatid  
9 exchanges induced by inhaled anesthetics. 50, 426-430.
- 10 White, J. F. and Carlson, G. P. (1981). Epinephrine-induced cardiac arrhythmias in rabbits  
11 exposed to trichloroethylene: potentiation by ethanol. *Toxicol Appl Pharmacol*, 60, 466-  
12 471.
- 13 White, J. F. and Carlson, G. P. (1981). Epinephrine-induced cardiac arrhythmias in rabbits  
14 exposed to trichloroethylene: role of trichloroethylene metabolites. *Toxicol Appl*  
15 *Pharmacol*, 60, 458-465.
- 16 White, K.; David, D.; Butterworth, L.; Klykken, P. (2000). Assessment of autoimmunity-  
17 inducing potential using the brown Norway rat challenge model. *Toxicol Lett*, 112-113,  
18 443-451. <http://www.ncbi.nlm.nih.gov/pubmed/10720764>
- 19 White, R.; Feldman, R.; Eviator, I.; Jabre, J.; Niles, C. (1997). Hazardous waste and  
20 neurobehavioral effects: a developmental perspective. *Environ Res*, 73, 113-124.  
21 <http://dx.doi.org/10.1006/enrs.1997.3699>
- 22 WHO. (World Health Organization). (1977). Manual of the international statistical classification  
23 of diseases, injuries, and causes of death. Geneva, Switzerland: Author.
- 24 WHO. (World Health Organization). (1985). Environmental health criteria: Trichloroethylene.  
25 Geneva, Switzerland: Author.
- 26 Wideroff, L.; Gridley, G.; Mellemejaer, L.; Chow, W. H.; Linet, M.; Keehn, S., . . . Olsen, J. H.  
27 (1997). Cancer incidence in a population-based cohort of patients hospitalized with  
28 diabetes mellitus in Denmark. *J Natl Cancer Inst*, 89, 1360-1365.  
29 <http://www.ncbi.nlm.nih.gov/pubmed/9308706>
- 30 Wiesenhütter, B.; Selinski, S.; Golka, K.; Brüning, T.; Bolt, H. M. (2007). Re-assessment of the  
31 influence of polymorphisms of phase-II metabolic enzymes on renal cell cancer risk of  
32 trichloroethylene-exposed workers. *Int Arch Occup Environ Health*, 81, 247-251.  
33 <http://dx.doi.org/10.1007/s00420-007-0200-5>
- 34 Wigle, D. T.; Turner, M. C.; Gomes, J.; Parent, M. E. (2008). Role of hormonal and other factors  
35 in human prostate cancer. *J Toxicol Environ Health B Crit Rev*, 11, 242-259.  
36 <http://dx.doi.org/10.1080/10937400701873548>
- 37 Wilcosky, T. C.; Checkoway, H.; Marshall, E. G.; Tyroler, H. A. (1984). Cancer mortality and  
38 solvent exposures in the rubber industry. *AIHA J*, 45, 809-811.  
39 <http://www.ncbi.nlm.nih.gov/pubmed/6549104>
- 40 Wilkens, L.; Flemming, P.; Gebel, M.; Bleck, J.; Terkamp, C.; Wingen, L., . . . Schlegelberger,  
41 B. (2004). Induction of aneuploidy by increasing chromosomal instability during  
42 dedifferentiation of hepatocellular carcinoma. *PNAS*, 101, 1309-1314.  
43 <http://dx.doi.org/10.1073/pnas.0305817101>

- 1 Williams, A. L. and DeSesso, J. M. (2008). Trichloroethylene and ocular malformations:  
2 analysis of extant literature. *Int J Toxicol*, 27, 81-95.  
3 <http://dx.doi.org/10.1080/10915810701876760>
- 4 Williams, F.; Meenagh, A.; Darke, C.; Acosta, A.; Daar, A. S.; Gorodezky, C., . . . Middleton, D.  
5 (2001). Analysis of the distribution of HLA-B alleles in populations from five continents.  
6 *Hum Immunol*, 62, 645-650. <http://www.ncbi.nlm.nih.gov/pubmed/11390040>
- 7 Williams, P.; Benton, L.; Warmerdam, J.; Sheehan, P. (2002). Comparative risk analysis of six  
8 volatile organic compounds in California drinking water. *Environ Sci Technol*, 36, 4721-  
9 4728. <http://www.ncbi.nlm.nih.gov/pubmed/12487291>
- 10 Wilson, J. G. (1965). Methods for administering agents and detecting malformations in  
11 experimental animals. In J. Warkany (Ed.), *Teratology: Principles and techniques* (8 ed.,  
12 pp. 262-267). Chicago, IL: University of Chicago Press.
- 13 Wilson, R.; Donahue, M.; Gridley, G.; Adami, J.; El Ghormli, L.; Dosemeci, M. (2008). Shared  
14 occupational risks for transitional cell cancer of the bladder and renal pelvis among men  
15 and women in Sweden. *Am J Ind Med*, 51, 83-99. <http://dx.doi.org/10.1002/ajim.20522>
- 16 Windemuller, F. J. B. and Ettema, J. H. (1978). Effects of combined exposure to  
17 trichloroethylene and alcohol on mental capacity. *Int Arch Occup Environ Health*, 41, 77-  
18 85. <http://www.ncbi.nlm.nih.gov/pubmed/631909>
- 19 Windham, G. C.; Shusterman, D.; Swan, S. H.; Fenster, L.; Eskenazi, B. (1991). Exposure to  
20 organic solvents and adverse pregnancy outcome. *Am J Ind Med*, 20, 241-259.  
21 <http://dx.doi.org/10.1002/ajim.4700200210>
- 22 Windham, G. C.; Zhang, L.; Gunier, R.; Croen, L. A.; Grether, J. K. (2006). Autism spectrum  
23 disorders in relation to distribution of hazardous air pollutants in the San Francisco Bay  
24 area. *Environ Health Perspect*, 114, 1438-1444.  
25 <http://www.ncbi.nlm.nih.gov/pubmed/16966102>
- 26 Winneke, G. (1982). Acute behavioral effects of exposure to some organic solvents -  
27 psychophysiological aspects. *Acta Neurol Scand*, 92, 117-129.  
28 <http://www.ncbi.nlm.nih.gov/pubmed/6962647>
- 29 Winneke, G.; Fodor, G.; Schlipkoter, H. (1978). Carbon monoxide, trichloroethylene, and  
30 alcohol: reliability and validity of neurobehavioral effects.
- 31 Winneke, G.; Kramer, U.; Kastka, J. (1976). Zur Beeinflussung psychomotor Leistungen Durch  
32 Alkohol und durch verschiedene Lösungsmitteldämpfe. In M. Horvath (Ed.), *Adverse*  
33 *effects of environmental chemicals and psychotropic drugs* (Vol. 2, pp. 99-110).  
34 Amsterdam, The Netherlands: Elsevier.
- 35 Withey, J. R. and Karpinski, K. (1985). The fetal distribution of some aliphatic chlorinated  
36 hydrocarbons in the rat after vapor phase exposure. *Biol Res Pregnancy Perinatol*, 6, 79-  
37 88. <http://www.ncbi.nlm.nih.gov/pubmed/4016168>
- 38 Withey, J. R.; Collins, B. T.; Collins, P. G. (1983). Effect of vehicle on the pharmacokinetics and  
39 uptake of four halogenated hydrocarbons from the gastrointestinal tract of the rat. *J Appl*  
40 *Toxicol*, 3, 249-253. <http://dx.doi.org/10.1002/jat.2550030506>
- 41 Wolf, D. C.; Goldsworthy, T. L.; Janszen, D. B.; Harden, R.; Donner, E. M.; David, C. S.;  
42 Everitt, J. I. (2000). Promotion by sodium barbital induces early development but does  
43 not increase the multiplicity of hereditary renal tumors in Eker rats. *Carcinogenesis*, 21,  
44 1553-1558. <http://www.ncbi.nlm.nih.gov/pubmed/10910958>

DRAFT– DO NOT QUOTE OR CITE

- 1 Wolff, D. L. and Siegmund, R. (1978). The circadian dependent effect of trichloroethylene on  
2 spontaneous locomotor activity and of tetrachloroethane on mortality in mice. 97, 345-  
3 352.
- 4 Wolfgang, G.; Gandolfi, A.; Stevens, J.; Brendel, K. (1989). In vitro and in vivo nephrotoxicity  
5 of the L and D isomers of S-(1,2-dichlorovinyl)-cysteine. *Toxicology*, 58, 33-42.  
6 <http://www.ncbi.nlm.nih.gov/pubmed/2815092>
- 7 Wolfgang, G.; Gandolfi, A.; Stevens, J.; Brendel, K. (1989). N-acetyl S-(1,2-dichlorovinyl)-L-  
8 cysteine produces a similar toxicity to S-(1,2-dichlorovinyl)-L-cysteine in rabbit renal  
9 slices: differential transport and metabolism. *Toxicol Appl Pharmacol*, 101, 205-219.  
10 <http://www.ncbi.nlm.nih.gov/pubmed/2815079>
- 11 Wood, L. D.; Parsons, D. W.; Jones, S.; Lin, J.; Sjoblom, T.; Leary, R. J., . . . Vogelstein, B.  
12 (2007). The genomic landscapes of human breast and colorectal cancers. *Science*, 318,  
13 1108-1113. <http://dx.doi.org/10.1126/science.1145720>
- 14 Woodruff, T. J.; Axelrad, D. A.; Caldwell, J.; Morello-Frosch, R.; Rosenbaum, A. (1998). Public  
15 health implications of 1990 air toxics concentrations across the United States. *Environ*  
16 *Health Perspect*, 106, 245-251. <http://www.ncbi.nlm.nih.gov/pubmed/9518474>
- 17 Woods, C. G.; Burns, A. M.; Bradford, B. U.; Ross, P. K.; Kosyk, O.; Swenberg, J. A., . . .  
18 Rusyn, I. (2007). WY-14,643-induced cell proliferation and oxidative stress in mouse  
19 liver are independent of NADPH oxidase. *Toxicol Sci*, 98, 366-374.  
20 <http://www.ncbi.nlm.nih.gov/pubmed/17483499>
- 21 Woods, C. G.; Heuvel, J. P.; Rusyn, I. (2007). Genomic profiling in nuclear receptor-mediated  
22 toxicity. *Toxicol Pathol*, 35, 474-494. <http://dx.doi.org/10.1080/01926230701311351>
- 23 WOOLF, B. (1955). On estimating the relation between blood group and disease. *Ann Hum*  
24 *Genet*, 19, 251-253. <http://www.ncbi.nlm.nih.gov/pubmed/14388528>
- 25 Woolhiser, M. R.; Krieger, S. M.; Thomas, J.; Hotchkiss, J. A. (2006). Trichloroethylene (TCE):  
26 Immunotoxicity potential in CD rats following a 4-week vapor inhalation exposure.  
27 Midland, MI.
- 28 Wright, P. F. A.; Thomas, W. D.; Stacey, N. H. (1991). Effects of trichloroethylene on hepatic  
29 and splenic lymphocytotoxic activities in rodents. *Toxicology*, 70, 231-242.  
30 <http://www.ncbi.nlm.nih.gov/pubmed/1763417>
- 31 Wu, C. and Schaum, J. (2000). Exposure assessment of trichloroethylene. *Environ Health*  
32 *Perspect*, 108(Suppl 2), 359-363.
- 33 Wu, C. and Schaum, J. (2001). Sources, emissions and exposures for trichloroethylene (TCE)  
34 and related chemicals. (EPA/600/R-00/099). Washington, DC: U.S. Environmental  
35 Protection Agency.
- 36 Wu, K. L. and Berger, T. (2007). Trichloroethylene metabolism in the rat ovary reduces oocyte  
37 fertilizability. *Chem Biol Interact*, 170, 20-30.  
38 <http://dx.doi.org/10.1016/j.cbi.2007.06.038>
- 39 Wu, K. L. and Berger, T. (2008). Ovarian gene expression is stable after exposure to  
40 trichloroethylene. *Toxicol Lett*, 177, 59-65. <http://dx.doi.org/10.1016/j.toxlet.2007.12.008>
- 41 Wu, X. Z. and Chen, D. (2006). Origin of hepatocellular carcinoma: role of stem cells. *J*  
42 *Gastroenterol Hepatol*, 21, 1093-1098. [http://dx.doi.org/10.1111/j.1440-](http://dx.doi.org/10.1111/j.1440-1746.2006.04485.x)  
43 [1746.2006.04485.x](http://dx.doi.org/10.1111/j.1440-1746.2006.04485.x)
- 44 Xu, H.; Tanphaichitr, N.; Forkert, P.; Anupriwan, A.; Weerachatanukul, W.; Vincent, R., . . .  
45 Wade, M. (2004). Exposure to trichloroethylene and its metabolites causes impairment of

- 1 sperm fertilizing ability in mice. *Toxicol Sci*, 82, 590-597.  
2 <http://dx.doi.org/10.1093/toxsci/kfh277>
- 3 Xu, W. and Adler, I. (1990). Clastogenic effects of known and suspect spindle poisons studied  
4 by chromosome analysis in mouse bone marrow cells. *Mutagenesis*, 5, 371-374.  
5 <http://www.ncbi.nlm.nih.gov/pubmed/2398818>
- 6 Xu, X.; Yang, R.; Wu, N.; Zhong, P.; Ke, Y.; Zhou, L., . . . Wu, B. (2009). Severe  
7 hypersensitivity dermatitis and liver dysfunction induced by occupational exposure to  
8 trichloroethylene. *Ind Health*, 47, 107-112.  
9 <http://www.ncbi.nlm.nih.gov/pubmed/19367038>
- 10 Yamamura, K.; Ikeda, T.; Sadmoto, T.; Maehara, N.; Harabuchi, I.; Takashima, H.; Kiyosawa,  
11 H. (1983). Effects of trichloroethylene exposure on hearing: an investigation of cochlear  
12 microphonics and action potential of the guinea pig. *Eur J Appl Physiol*, 52, 47-50.  
13 <http://www.ncbi.nlm.nih.gov/pubmed/6686127>
- 14 Yang, Q.; Ito, S.; Gonzalez, F. J. (2007). Hepatocyte-restricted constitutive activation of PPAR  
15 alpha induces hepatoproliferation but not hepatocarcinogenesis. *Carcinogenesis*, 28,  
16 1171-1177. <http://dx.doi.org/10.1093/carcin/bgm046>
- 17 Yano, M.; Yamamoto, T.; Kobayashi, S.; Ikeda, Y.; Matsuzaki, M. (2008). Defective Ca<sup>2+</sup>  
18 cycling as a key pathogenic mechanism of heart failure. *Circ J*, 72 Suppl A, A22-A30.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/18772523>
- 20 Yauck, J.; Malloy, M.; Blair, K.; Simpson, P.; McCarver, D. (2004). Proximity of residence to  
21 trichloroethylene-emitting sites and increased risk of offspring congenital heart defects  
22 among older women. *Birth Defects Res A Clin Mol Teratol*, 70, 808-814.  
23 <http://dx.doi.org/10.1002/bdra.20060>
- 24 Yeh, M. M.; Larson, A. M.; Campbell, J. S.; Fausto, N.; Rulyak, S. J.; Swanson, P. E. (2007).  
25 The expression of transforming growth factor-alpha in cirrhosis, dysplastic nodules, and  
26 hepatocellular carcinoma: an immunohistochemical study of 70 cases. *Am J Surg Pathol*,  
27 31, 681-689. <http://dx.doi.org/10.1097/PAS.0b013e31802ff7aa>
- 28 Yeldandi, A. V.; Milano, M.; Subbarao, V.; Reddy, J. K.; Rao, M. S. (1989). Evaluation of liver  
29 cell proliferation during ciprofibrate-induced hepatocarcinogenesis. *Cancer Lett*, 47, 21-  
30 27. <http://www.ncbi.nlm.nih.gov/pubmed/2636030>
- 31 Yeung, R. S.; Xiao, G. H.; Everitt, J. I.; Jin, F.; Walker, C. L. (1995). Allelic loss at the tuberous  
32 sclerosis 2 locus in spontaneous tumors in the Eker rat. *Mol Carcinog*, 14, 28-36.  
33 <http://www.ncbi.nlm.nih.gov/pubmed/7546222>
- 34 Yoon, H. S.; Monks, T. J.; Walker, C. L.; Lau, S. S. (2001). Transformation of kidney epithelial  
35 cells by a quinol thioether via inactivation of the tuberous sclerosis-2 tumor suppressor  
36 gene. *Mol Carcinog*, 31, 37-45. <http://www.ncbi.nlm.nih.gov/pubmed/11398196>
- 37 Yoon, J. S.; Mason, J. M.; Valencia, R.; Woodruff, R. C.; Zimmering, S. (1985). Chemical  
38 mutagenesis testing in *Drosophila*. IV. Results of 45 coded compounds tested for the  
39 National Toxicology Program. *Environ Mutagen*, 7, 349-367.  
40 <http://dx.doi.org/10.1002/em.2860070310>
- 41 Yoon, M.; Madden, M. C.; Barton, H. A. (2007). Extrahepatic metabolism by CYP2E1 in PBPK  
42 modeling of lipophilic volatile organic chemicals: Impacts on metabolic parameter  
43 estimation and prediction of dose metrics. *J Toxicol Environ Health A*, 70, 1527-1541.
- 44 Yoshida, M.; Fukabori, S.; Hara, K.; Yuasa, H.; Nakaaki, K.; Yamamura, Y.; Yoshida, K.  
45 (1996). Concentrations of trichloroethylene and its metabolites in blood and urine after

DRAFT– DO NOT QUOTE OR CITE

- 1 acute poisoning by ingestion. *Hum Exp Toxicol*, 15, 254-258.  
2 <http://www.ncbi.nlm.nih.gov/pubmed/8839215>
- 3 Yoshioka, T.; Krauser, J. A.; Guengerich, F. P. (2002). Microsomal oxidation of  
4 tribromoethylene and reactions of tribromoethylene oxide. *Chem Res Toxicol*, 15, 1414-  
5 1420. <http://www.ncbi.nlm.nih.gov/pubmed/12437332>
- 6 Yu, K. O.; Barton, H. A.; Mahle, D. A.; Frazier, J. M. (2000). In vivo kinetics of trichloroacetate  
7 in male Fischer 344 rats. *Toxicol Sci*, 54, 302-311.  
8 <http://www.ncbi.nlm.nih.gov/pubmed/10774812>
- 9 Yu, M. C.; Garabrant, D. H.; Peters, J. M.; Mack, T. M. (1988). Tobacco, alcohol, diet,  
10 occupation, and carcinoma of the esophagus. *Cancer Res*, 48, 3843-3848.  
11 <http://www.ncbi.nlm.nih.gov/pubmed/3378219>
- 12 Yu, S.; Rao, S.; Reddy, J. (2003). Peroxisome proliferator-activated receptors, fatty acid  
13 oxidation, steatohepatitis and hepatocarcinogenesis. *Curr Mol Med*, 3, 561-572.  
14 <http://www.ncbi.nlm.nih.gov/pubmed/14527087>
- 15 Yuan, J. M.; Castelao, J. E.; Gago-Dominguez, M.; Yu, M. C.; Ross, R. K. (1998). Tobacco use  
16 in relation to renal cell carcinoma. *Cancer Epidemiol Biomarkers Prev*, 7, 429-433.  
17 <http://www.ncbi.nlm.nih.gov/pubmed/9610793>
- 18 Yue, F.; Huang, H. L.; Huang, J. X.; Liang, L. Y.; Huang, Z. L.; Wei, Q. Y.; Chen, X. M. (2007).  
19 HLA-DM polymorphism and risk of trichloroethylene induced medicamentosa-like  
20 dermatitis. *Biomed Environ Sci*, 20, 506-511.  
21 <http://www.ncbi.nlm.nih.gov/pubmed/18348411>
- 22 Zack, M.; Cannon, S.; Loyd, D.; Heath, C. W.; Falletta, J. M.; Jones, B., . . . Crowley, S. (1980).  
23 Cancer in children of parents exposed to hydrocarbon-related industries and occupations.  
24 *Am J Epidemiol*, 111, 329-336. <http://www.ncbi.nlm.nih.gov/pubmed/7361757>
- 25 Zahm, S. H. (1992). Computerized occupational referent population system (CORPS): Study  
26 documentation. Rockville, MD: National Cancer Institute & National Institute for  
27 Occupational Safety and Health.
- 28 Zajicek, G. and Schwartz-Arad, D. (1990). Streaming liver. VII: DNA turnover in acinus zone-3.  
29 10, 137-140. <http://www.ncbi.nlm.nih.gov/pubmed/2385154>
- 30 Zajicek, G.; Arber, N.; Schwartz-Arad, D. (1991). Streaming liver. VIII: Cell production rates  
31 following partial hepatectomy. 11, 347-351.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/1779714>
- 33 Zajicek, G.; Schwartz-Arad, D.; Bartfeld, E. (1989). The streaming liver. V: Time and age-  
34 dependent changes of hepatocyte DNA content, following partial hepatectomy. 9, 164-  
35 171. <http://www.ncbi.nlm.nih.gov/pubmed/2747441>
- 36 Zandi-Nejad, K.; Eddy, A.; Glasscock, R.; Brenner, B. (2004). Why is proteinuria an ominous  
37 biomarker of progressive kidney disease. *Kidney Int Suppl* S76-S89.  
38 <http://dx.doi.org/10.1111/j.1523-1755.2004.09220.x>
- 39 Zanelli, U.; Puccini, P.; Acerbi, D.; Ventura, P.; Gervasi, P. G. (1996). Induction of peroxisomal  
40 beta-oxidation and P-450 4A-dependent activities by pivalic and trichloroacetic acid in  
41 rat liver and kidney. *Arch Toxicol*, 70, 145-149.
- 42 Zeise, L.; Wilson, R.; Crouch, E. A. C. (1987). Dose-response relationships for carcinogens: a  
43 review. *Environ Health Perspect*, 73, 259-308.



DRAFT– DO NOT QUOTE OR CITE

- 1 Zenick, H.; Blackburn, K.; Hope, E.; Richdale, N.; Smith, M. K. (1984). Effects of  
2 trichloroethylene exposure on male reproductive function in rats. *Toxicology*, 31, 237-  
3 250. <http://www.ncbi.nlm.nih.gov/pubmed/6740699>
- 4 Zeppa, P.; Benincasa, G.; Troncone, G.; Lucariello, A.; Zabatta, A.; Cochand-Priollet, B., . . .  
5 Palombini, L. (1998). Retrospective evaluation of DNA ploidy of hepatocarcinoma on  
6 cytologic samples. *Diagn Cytopathol*, 19, 323-329.  
7 <http://www.ncbi.nlm.nih.gov/pubmed/9812223>
- 8 Zhang, B.; Pan, X.; Cobb, G. P.; Anderson, T. A. (2007). microRNAs as oncogenes and tumor  
9 suppressors. *Dev Biol*, 302, 1-12. <http://dx.doi.org/10.1016/j.ydbio.2006.08.028>
- 10 Zhang, G. and Stevens, J. (1989). Transport and activation of S-(1,2-dichlorovinyl)-L-cysteine  
11 and N-acetyl-S-(1,2-dichlorovinyl)-L-cysteine in rat kidney proximal tubules. *Toxicol*  
12 *Appl Pharmacol*, 100, 51-61. <http://www.ncbi.nlm.nih.gov/pubmed/2763302>
- 13 Zhang, S. H.; Miao, D. Y.; Liu, A. L.; Zhang, L.; Wei, W.; Xie, H.; Lu, W. Q. (2010).  
14 Assessment of the cytotoxicity and genotoxicity of haloacetic acids using microplate-  
15 based cytotoxicity test and CHO/HGPRT gene mutation assay. *Mutat Res*, 703, 174-179.  
16 <http://dx.doi.org/10.1016/j.mrgentox.2010.08.014>
- 17 Zhang, Y.; Cantor, K. P.; Lynch, C. F.; Zheng, T. (2004). A population-based case-control study  
18 of occupation and renal cell carcinoma risk in Iowa. *J Occup Environ Med*, 46, 235-240.  
19 <http://www.ncbi.nlm.nih.gov/pubmed/15091286>
- 20 Zhao, Y.; Krishnadasan, A.; Kennedy, N.; Morgenstern, H.; Ritz, B. (2005). Estimated effects of  
21 solvents and mineral oils on cancer incidence and mortality in a cohort of aerospace  
22 workers. *Am J Ind Med*, 48, 249-258. <http://dx.doi.org/10.1002/ajim.20216>
- 23 Zhou, H.; Randers-Pehrson, G.; Geard, C. R.; Brenner, D. J.; Hall, E. J.; Hei, T. K. (2003).  
24 Interaction between radiation-induced adaptive response and bystander mutagenesis in  
25 mammalian cells. *Radiat Res*, 160, 512-516.  
26 <http://www.ncbi.nlm.nih.gov/pubmed/14565832>
- 27 Zhou, Y. C. and Waxman, D. J. (1998). Activation of peroxisome proliferator-activated receptors  
28 by chlorinated hydrocarbons and endogenous steroids. *Environ Health Perspect*, 106  
29 Suppl 4, 983-988. <http://www.ncbi.nlm.nih.gov/pubmed/9703482>
- 30 Zhu, J.; Newhook, R.; Marro, L.; Chan, C. (2005). Selected volatile organic compounds in  
31 residential air in the city of Ottawa, Canada. *Environ Sci Technol*, 39, 3964-3971.  
32 <http://www.ncbi.nlm.nih.gov/pubmed/15984771>
- 33 Zielinska-Psuja, B.; Orłowski, J.; Kowalowska, J.; Zieba-Proc, B. (1998). Metabolic implications  
34 of benzene and trichloroethylene. Part I. The effect trichloroethylene on phenol  
35 elimination in rats. *Bromatol Chem Toksykol*, 31, 13-20.
- 36 Zielinski, A. (1973). General health state of women professionally exposed to trichloroethylene  
37 vapours. *Med Pr*, 24, 263-271.
- 38 Zordan, M.; Osti, M.; Pesce, M.; Costa, R. (1994). Chloral hydrate is recombinogenic in the  
39 wing spot test in *Drosophila melanogaster*. *Mutat Res*, 322, 111-116.  
40 <http://www.ncbi.nlm.nih.gov/pubmed/7519317>

41

42

43