

AR 201-12863B

Robust Summaries

RECEIVED
DPT/NCIS
2000 DEC 27 AM 8:17

Melting Point

Type	Melting Point
Test Substance	1,3,5-Trioxane CAS Number: 11 0-88-3

Method

- Guideline
- Test Type Melting Point
- GLP No
- Year Unknown

Result

- Melting Point 64 deg C
- Remarks Field for Results Handbook data

Conclusions

Remarks field Melting point is 64° C.

Data Quality

- Reliability Klimisch Code 2. A reliability code of 2 is assigned to data from reference handbooks.

References

1. The Merck Index. 9th ed. Rahway, New Jersey: Merck & Co., Inc., 1976. 1249

Boiling Point

Type	Boiling Point
Test Substance	1,3,5-Trioxane CAS Number: 11 0-88-3

Method

- Guideline
- Test Type Boiling Point
- GLP No
- Year unknown

Result

- Boiling Point 114.5 deg C @ 759 mm Hg

Remarks Field for Results Handbook data

Conclusions

Remarks field 114.5 deg C @ 759 mm Hg

Data Quality

- Reliability Klimisch Code 2. A reliability code of 2 is assigned to data from reference handbooks.

References

1. The Merck Index. 9th ed. Rahway, New Jersey: Merck & Co., Inc., 1976. 1249

Vapor Pressure

Type Vapor Pressure
Test Substance 1,3,5-Trioxane
CAS Number: 11 0-88-3

Method

- Guideline
- Test Type Vapor Pressure
- GLP No
- Year unknown

Result

- Vapor Pressure 10 mm Hg @ 20 deg C

Remarks Field for Determined value
Results

Conclusions

Remarks field This material is a volatile solid at normal room temperatures. VP values for solids are not generally reported but, in this case, it is of value in risk assessment.

Data Quality

- Reliability Klimisch Code 2. Physical measurement conducted by reliable laboratory, considered reliable although not conducted under GLP conditions.

References

Determination by Celanese Chemicals at the Corpus Christi Technical Center.

Other

This result is supported by the MPBPWIN v1.40 modeling program which uses three methods to estimate the VP based on the experimentally determined melting point and boiling point (both found in the EPIWIN data base) to estimate values of 11.8, 10.5 and 11.1 mm Hg @ 25° C. with 10.5 mm selected by the program as the preferred estimated value.

Reference for Supporting Studies MPBPWIN v1.40 as found in EPIWIN v3.05, Syracuse Research Corporation, Syracuse NY (April 2000).

Partition Coefficient

Type	Partition Coefficient
Test Substance	1,3,5-Trioxane CAS Number: 11 0-88-3

Method

- Method Shake flask
- Test Type Partition Coefficient
- GLP No
- 1989

Remarks Field for Test Conditions	◇ Replicates	One replicate per test concentration
	◇ Concentration	55, 102, 551 mg test material
	◇ Volumes	Water 25 to 35 ml n-Octanol 25 ml
	◇ Analysis	Triplicate GC analysis of each phase of each replicate
	◇ Variation From Current OECD Guideline	This study used single replicates of three test material concentration in relatively fixed volumes of Octanol/water. The current (1995) OECD 107 guideline calls for duplicate runs at three solvent ratios and possibly test substance quantities. As the partition coefficient is near unity this deviation is considered to have minimal effect on the outcome.

Result

- Partition Coefficient Log $K_{o/w}$ = -0.47
- Temperature 25°C

Remarks Field for Results	All three concentration of test material gave similar partition coefficients, they were $P_{o/w}$ of 0.36, 0.32 and 0.34 for test material quantities of 55, 102, 551 mg, respectively.
---------------------------	---

Conclusions

Remarks field	Log $K_{o/w}$ was determined to be -0.47 using the shake-flask method
---------------	---

Data Quality

- Reliability Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although it could not be established that the study was conducted in accord with GLP standards. Study design and reporting meets current EPA/OECD guidelines with minor exceptions.

References

Reprint Of A Test Report "Determination of partition coefficient n-octanol/water of 1,3,5-Trioxane by shake flask method." Analytical report Number 107243/01 Test period: May 1989

Other

This result is supported by a literature value of -0.43' and by the the Log Kow v1.66 modeling program which estimates a value of -0.56 for Log **K_o/w**

Reference for
Supporting Studies

1. Hansch. C., A. Leo and D. Hoekman. 1995. Exploring QSAR. Hydrophobic, Electronic, and Steric Constants. ACS Professional Reference Book. Washington, DC: American Chemical Society as cited in Documentation for KOWIN estimation program version 4.00.5000, Syracuse Research Corporation, Syracuse, NY (April 2000)
2. Log Kow v1.66 as found in EPIWIN v3.05, Syracuse Research Corporation, Syracuse NY (April 2000).

Water Solubility

Type Water Solubility

Test Substance 1,3,5-Trioxane
CAS Number: 11 0-88-3

Method

- Guideline
- Test Type Water Solubility
- GLP No
-  

Result

- Solubility 0 17.2 g/100 ml @ 18° C,
0 21.2 g/100 ml @ 25° C

Remarks Field for Results Handbook data

Conclusions

Remarks field Water solubility approximately 200 g/l at room temperature.

Data Quality

- Reliability Klimisch Code 2. A reliability code of 2 is assigned to data from reference handbooks.

References

1. The Merck Index. 9th ed. Rahway, New Jersey: Merck & Co., Inc., 1976. 1249

Fate in the Environment

Photodegradation

Type	Photodegradation
Test Substance	1,3,5-Trioxane CAS Number: 110-88-3

Method

- **Guideline** Estimated using version 1.90 of the Atmospheric Oxidation Program for Microsoft Windows (AOPWIN) which estimates the rate constant for the atmospheric, gas-phase reaction between photochemically produced hydroxyl radicals and organic chemicals. The rate constants estimated by the program are then used to calculate atmospheric half-lives for organic compounds based upon average atmospheric concentrations of hydroxyl radical.
- **Test Type** Photodegradation Estimate
- **GLP** No
- **☑** 2000

Result

- Experimental •OH rate constant $6.2 \times 10^{-12} \text{ cm}^3/\text{molecule-sec}$ (2)
- APOWIN estimated •OH rate constant $10.2 \times 10^{-12} \text{ cm}^3/\text{molecule-sec}$

Remarks Field for Results APOWIN estimated the reaction rate but lists a measured value for this material in the reference **materials**.² Based on the measured rate constant and using the defaults in APOWIN for atmospheric hydroxyl radical concentration, the estimated half-life is approximately 25 hours. Using the APOWIN estimated rate constant the estimated half-life is approximately 12.5 hours.

Conclusions

Remarks field The atmospheric half-life of **1,2,3-Trioxane** in the atmosphere is estimated to be in the range of 25 hours.

Data Quality

- **Reliability** Klimisch Code 1. A reliability code of 2 is assigned a result using an accepted method of estimation. Since there is a literature value for the reaction rate of this material with hydroxyl radical, and since it is similar to the APOWIN estimated value the estimate is considered to have a higher reliability and is assigned a code of 1.

References

1. Syracuse Research Corporation, Syracuse, NY (April 2000)
2. Atkinson, R. 1989. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. J. Phys. Chem. Ref. Data Monograph No. 1. NY: Amer. Inst. Physics & Amer. Chem Soc.

Water Stability**Type****Water Stability****Test Substance****1,3,5-Trioxane**

CAS Number: 11 0-88-3

Purity > 99 % by GC

Source: Fluka (ZAX A 2205/02/GC_73 1)

Method

- Guideline OECD 111 (1981)
- Test Type Hydrolysis as a Function of **pH**
- GLP Yes
- Year 2000

Remarks Field for 0 Duration
Test Conditions

Six days (preliminary test only)

0 Analytical
Method

Direct injection into GC using flame ionization detector.

0 Buffers

Target PH	Buffer System	Measured pH
4.0	Citric acid/sodium chloride, sodium hydroxide	3.96
7.0	Potassium dihydrogen phosphate/disodium hydrogenphosphate	7.03
9.0	Sodium borate/hydrochloric acid	9.04

0 Vessels

One-liter water-jacketed glass vessels with screw caps.

0 Replicates

One at **pH** 4 and 7, two at **pH** 9.

0 Temperature

49.8 – 50.0 °C

0 Sampling

About 12 samplings in six days.

0 Additional
Testing

Not conducted since material showed less than ten percent degradation at 50° in four days.

Results

- Nominal Target Concentration 875 mg/l

- Measured Concentrations

Conditions		Measured Concentration (mg/l)		Percent Degradation 6 days
PH	T (deg C)	Initial	Final	
4.06 (±0.05)	49.8 (±0.1)	905	867	4.2
6.98 (±0.05)	50.0 (±0.1)	893	843	5.6
8.84 (±0.05)	49.9 (±0.1)	918	909	1.0
8.85 (±0.05)	50.0 (±0.1)	874	860	1.6


- Percent Degradation Less than 10% at 50" in six days
- Breakdown Products None

Remarks Field for Results

pH 4.06		pH 6.98		pH 8.84		pH 8.85	
Time	Conc.	Time	Conc.	Time	Conc.	Time	Conc.
0.0	905	0.0	893	0.0	918	0.0	874
1.0	899	1.1	883	1.0	920	1.0	880
4.1	898	3.0	905	3.0	922	3.0	890
5.0	911	5.0	887	5.0	934	5.0	869
22.0	894	22.0	868	-*	-	-	-
24.0	891	24.1	877	-	-	-	-
26.6	900	27.5	870	48.0	901	48.4	847
29.0	910	29.0	878	50.0	923	50.0	864
47.9	886	48.2	888	52.1	922	52.1	863
52.0	877	52.0	871	72.6	916	72.3	858
72.0	883	72.0	859	76.6	913	76.4	858
76.0	873	145.1	851	141.6	918	141.5	859
145.1	851	146.3	845	144.3	907	144.3	859
146.1	867	147.4	843	148.5	909	148.5	860

* Sampling in the 24-hour time range not conducted for pH 9 determinations

Remarks for Results, continued



Conclusions

Remarks field

. Trioxane is stable at **pH** 4, 7 or 9 for six days at 50° C under the conditions specified in the OECD 111 Guideline.

. Trioxane is estimated to have the following half-life (**t_{1/2}**) at 25 ° C.

Experiment	t _{1/2} , 25°C (days)
pH 4	780
PH7	840
pH 9	3200
pH 9 (2nd test)	2000

Data Quality

- Reliability

Klimisch Code 1. May be used without restriction.

Reference

Physico-chemical properties of Trioxane (Hydrolysis as a Function of **pH**.) ZAX Analytik. Study No. OOLO0453, BASF AG. Ludwigshafen November 2000.

Other

The hydrolysis-modeling program found in EPIWIN has no valid model for ethers.

Reference for supporting study

1. HYDROWIN modeling program, version 1.67, as found in EPIWIN v 3.05, Syracuse Research Corporation, Syracuse NY (April 2000).

Theoretical Distribution (Fugacity)**Type** **Theoretical Distribution (Fugacity)**

Test Substance 1,3,5-Trioxane

Method

- **Guideline** Estimated using the **MacKay** model with standard defaults contained in **EPIWIN v 3.05.**
- **Test Type** Level III Fugacity Model
- **GLP** No
- **Year** **2000**

Result

- **Distribution** 0 Air 4.9 %
 0 Water **53.4 %**
 0 Soil 41.6 %
 0 Sediment 0.09 %

Remarks Field for Results 0 This is the currently accepted model for theoretical distribution estimation.
 0 Experimentally determined vapor pressure and water solubility were used to improve the modeling.
 0 Calculated K_{oc} is 0.152

Conclusions

Remarks field This material is expected to environmentally distribute primarily in water and soil

Data Quality

- **Reliability** Klimisch Code 2. A reliability code of 2 is assigned a result using an accepted method of estimation.

References

Syracuse. . . Research . . . Corporation, Syracuse., NY (**April 2000**)

Biodegradation**Type** Ready Biodegradation**Test Substance** 1,3,5-Trioxane
CAS Number: 11 O-88-3**Method**

- **Guideline** MITI Test
- **Test Type** Ready Biodegradation
- **GLP** No
- **Year** 1981
- **Contact Time** 28 days
- **Innoculum** Activated sludge from **BASF's** waste water treatment plant

Result

- **Result** % Biodegradation in 28 days 2 %

Remarks Field for Results Not readily biodegradable

Conclusions

Remarks field Not readily biodegradable

Data Quality

- **Reliability** Klimisch Code 2 Study design, conduct and reporting are considered reliable to address the test endpoint although it could not be established that the study was conducted in accord with GLP standards.

References

Weitere Untersuchungen vom Trioxan, **1,3-Dioxepam** und Dibutylformal. Respirometrische Tests Über 28 Tage, TUU/W – K 210 -1130, BASF AG, Ludwigshafen, April 198 1.

Other

- This study is supported by an earlier study, sponsored by Celanese, in which Dioxolane was tested for biodegradation using a municipal secondary effluent and measuring oxygen consumption with a manometric respirometer. After 15 days Dioxolane showed 4.8 % of the THOD.¹
- Other short-term BOD studies conducted by BASF.
- The BIOWIN **V4.0** model found in EPIWIN. Two of the three models included predict that Trioxane will not rapidly **biodegrade**.²

References for
Supporting Studies

1. Report to Celanese Chemical Company Inc. on Toxicology and Fate of Selected Industrial Chemicals in Aquatic Ecosystems. J.R. Walton and E.M. Davis, University of Texas at Houston. December 1980.
2. EPIWIN v 3.05, Syracuse Research Corporation, Syracuse NY (April 2000).

Biodegradation, Inherent

Type Inherent Biodegradation

Test Substance 1,3,5-Trioxane
CAS Number: 110-88-3

Method

- Guideline OECD 302B
- Test Type Inherent Biodegradation, Zahn-Wellens method with aeration
- GLP No
- Year 1984
- Contact Time 23 days
- Inoculum Activated sludge from **BASF's** waste water treatment plant

Remarks Field for Test Conditions

- ◇ Conditions Trioxane is known to volatilize directly from water. A control was run with aeration only to determine the potential extent of "air stripping"
- ◇ Acclimation Waste-water treatment plant receives some Trioxane; **thus** it is considered pre-acclimated.


Result

- Degradation Cannot be determined
% Biodegradation in 23 days
- Results
 - 79% loss of DOC from control due to air stripping.
 - 98 % loss of DOC from inoculated sample due to air stripping and biodegradation
- Breakdown Products Not identified

Remarks Field for \diamond DOC level at Each Time

Day of Test	DOC (mg/l)	
	Aeration only	Aeration plus Bacteria
0	388	386
1	360	378
2	323	297
3	292	267
4	276	219
5	245	183
6	241	156
7	231	120
8	212	108
9	200	118
10	177	109
11	169	93.0
14	134	75.0
16	109	62.0
18	100	60.0
21	83.0	49.0
23	70.0	25.0

0 DOC Loss Curves



Conclusions

Remarks field

Trioxane was effectively removed from solution by aeration in the presence of a bacterial inoculum; however, the relative contributions of biodegradation and volatilization could not be determined.

Data Quality

- Reliability

Klimisch Code 2 Study design, conduct and reporting are considered reliable to address the test endpoint although it could not be established that the study was conducted in accord with GLP standards.

References Reprint Of A Test Report: Determination of the Ultimate and Inherent Biodegradation and the Elimination from Water of Trioxan in a Batch Test with Activated Sludge BASF Aktiengesellschaft, Ecological Studies D-67056 Ludwigshafen, January 1984

Other This result is supported by the **BIOWIN V4.0** model found in **EPIWIN** that estimates a time of “weeks” for ultimate biodegradation.

References for Supporting Studies Syracuse Research Corporation, Syracuse, NY (April 2000).

Biodegradation, Inherent**Type** **Inherent** **Biodegradation**

Test Substance **1,3,5-Trioxane**
 CAS Number: 110-88-3

Method

- Guideline Modified OECD 302B
- Test Type Inherent Biodegradation, Modified Zahn-Wellens method without aeration using Nitrate as oxygen source
- GLP No
- Year 1984
- Contact Time 23 days
- Inoculum Activated sludge from BASF's waste water treatment plant

Remarks Field for
 Test Conditions

- ◇ Conditions Trioxane is known to volatilize directly from water. A previous test showed that loss to air with aeration was significant. This study used a closed system with the addition of 2 g/l Sodium nitrate as an oxygen source. The control was test material with Sodium nitrate without bacteria.
- ◇ Acclimation The wastewater treatment plant receives some Trioxane; thus, it is considered **preacclimated**.


Result

- Degradation ◇ % Biodegradation in 18 days > 90%
- Results Inherently biodegradable
- Breakdown Products Not identified

Remarks Field for Results ◇ DOC level at Each Time

Day of Test	DOC (mg/l)	
	Nitrate only	Nitrate plus Bacteria
0	384	381
1	377	381
2	371	369
5	370	318
7	333	252
9	343	228
12	321	168
14	320	131
16	294	72.0
17	322	28.0
18	314	8.0

◇ DOC Loss Curves



Conclusions

Remarks field

Trioxane was effectively removed from solution under these conditions where volatilization was not possible. It is assumed that essentially complete biodegradation occurred under these conditions.

Data Quality

- Reliability

Klimisch Code 2 Study design, conduct and reporting are considered reliable to address the test endpoint although it could not be established that the study was conducted in accord with GLP standards.

References

Reprint Of A Test Report: Determination of the Inherent and Ultimate Biodegradability of Trioxan in a Modified Batch Test with Activated Sludge Modification: No Aeration but Use of Nitrate as Oxygen Supply BASF Aktiengesellschaft, Ecological Studies D-67056 Ludwigshafen March 1984

Other

This result is supported by the BIOWIN V4.0 model found in EPIWIN that estimates a time of “weeks” for ultimate biodegradation.

References for Supporting Studies

Syracuse Research Corporation, Syracuse, NY (April 2000).

- Dilution water Reconstituted freshwater according to DIN 38 412, Part L1, October 1982.
Demineralized water resalted with 294.0 mg/l $\text{CaCl}_2 \times 2 \text{H}_2\text{O}$; 123.3 mg/l $\text{MgSO}_4 \times 7 \text{H}_2\text{O}$; 63.0 mg/l NaHCO_3 ; 5.5 mg/l KCl
- . Water Conditions
 - Total hardness: **2.5 mmol/l**
 - Acid capacity: **0.8 mmol/l**
 - Ratio Ca/Mg ions: 4:1
 - Ratio Na/K ions: 10:1
 - pH about 8.0
- . Photoperiod 16 hours light and 8 hours dark
- Water Temp 20- 21 deg C at all test times

0 Solvent	Test water
0 Vessel	All-glass aquaria, 30 X 22 X 24 cm, containing 10 liters
0 Fish per group	Ten, one replicate per concentration
0 Fish per vessel	Ten
0 Exposure period	96 hours
0 Observation times	1,4, 24, 48, 72 and 96 hours
0 Solution pH range	7.4 -7.8 at all concentrations and at 1, 24, 28, 72 and 96 hours
0 Dissolved oxygen	Above 7.8 mg/l for all solutions at 0 and 48 hours

Results

- Nominal Concentrations 0, 1000, 2150, 4640 and 10000 mg/l
- Units mg./l.
- LC_{50} 4030 (96, 72 and 48 hour), 5340 (24 hour)
- LC_0 2150 (24 to 96 hours)

Remarks Field for 0 Mortality Results

Nominal Conc (mg/ml)	Number Dead Fish After						
	0 hr	1 hr	4 hr	24 hr	48 hr	72 hr	96 hr
1,000	10	0	0	0	0	0	0
2,150	10	0	0	0	0	0	0
4,640	10	2	3	3	7	7	7
10,000	10	10	10	10	10	10	10

- ◇ Biological Observations Restricted to mortality
- ◇ Control Mortality Zero

Conclusions

Remarks field

- ◇ The LC_{50} was found to be 4030 mg/l at 96, 72 and 48 hours; it was 5340 at 24 hours.
- ◇ The LC_0 was found to be 2150 mg/l at all time intervals.
- ◇ The NOEC was 2 150 mg/l. The lowest concentration causing 100% mortality was 10000 mg/ml.
- ◇ The study closely followed the OECD 203 guideline with the exception of the recommended species of fish.

Data Quality

- Reliability Klimisch Code 2 . Study design, conduct and reporting are considered reliable to address the test endpoint although it could not be established that the study was conducted in accord with GLP standards. Study design and reporting meets current EPA/OECD guidelines with minor exceptions

References

Bericht über die Prüfung der akuten Toxizität an der Goldorfe (*Leuciscus idus* L., Goldvariante); englische Fassung Prüfung zur Einstufung in Wassergefährdungsklasse. Substance 87/525, 1,3,5-Trioxane. BASF Department of Toxicology July 1988.

Other

- 0 This study is supported by an earlier study, sponsored by Celanese, in which Sheepshead minnows (*Cyprinodon variegatus*, 5 per group) were exposed to Trioxane at concentrations of **10000, 20000** and 30000 **mg/l**. In this study, the 96 **hour-LC₅₀** was reported to be 16350 **mg/l**. and the **96-hour LCO** was 10000 **mg/l**.¹
- 0 The EPA ECOSAR Modeling Program found in EPIWIN, estimates the **96-hour LC₅₀** for fish to be 17000 **mg/l**.²

References for supporting studies

1. Report to Celanese Chemical Company Inc. on Toxicology and Fate of Selected Industrial Chemicals in Aquatic Ecosystems. J.R. Walton and E.M. Davis, University of Texas at Houston. December 1980.
2. ECOSAR modeling program, version **0.99f**, as found in EPIWIN v 3.05, Syracuse Research Corporation, Syracuse NY (April 2000).

Acute Toxicity to Aquatic Invertebrates**Type** Acute Toxicity to Aquatic Invertebrates

Test Substance 1,3,5-Trioxane
 CAS Number: 11 0-88-3
 Purity not specified

Method

- Guideline OECD 202 (1984)
- Test Type Daphnia, acute immobilization
- GLP No
- Year 1989
- Analytical Procedures None
- Species/Strain *Daphnia magna* STRAUS
- Test Details Static
- Statistical Methods None necessary

Remarks	Field for	◇ Age at study initiation	Four to 24 hours
Test Conditions	0	Test conditions	As specified in OECD 202.
	0	Solvent	Test water
	0	Vessel	50 ml glass beaker with 20 ml solution
	0	Daphnids per group	Twenty
	0	Daphnids per vessel	Five
	0	Exposure period	48 hours
	0	Observation times	3, 6, 24, 48 hours
	0	Solution pH range	7.72 to 7.80 at all concentrations at 0 and 48 hours
	0	Dissolved oxygen	Above 8 mg/l for all solutions at 0 and 48 hours

Results

- Nominal Concentrations 0, 0.1, 1, 10, 100 and 1000 mg/l
- Units mg./l.
- EC₅₀ > 1000 at 24 and 48 hours
- EC₀ > 1000 at 24 and 48 hours

Remarks Field for V Immobilization o No animal was immobilized at any observation time.
 Results o No other adverse effects were reported
 0 Controls were normal

Conclusions

Remarks field The **EC₅₀** (48 hour) and **EC₀** (48 hour) were greater than 1000 **mg/l** under these conditions. The **study** closely followed the OECD 202 guideline.

Data Quality

- Reliability Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although it could not be established that the study was conducted in accord with GLP standards. Study design and reporting meets current **EPA/OECD** guidelines with minor exceptions.

References

Fraunhofer-Institute für Umweltchemie und Ökotoxikologie, Daphnia, Acute Immobilisation, Test-Substanz: **1,3,5-Trioxane**. 27.4.1989 IUCT-Nr.: BALU1

Other

This study is supported by an earlier study, sponsored by Celanese, in which Trioxane was tested at **5000, 10000, 15000, 20000** and 30000 **mg/l**. In this study the 48 **hour-EC₅₀** was reported to be 15200 **mg/l**.¹
 The EPA ECOSAR Modeling Program found in **EPIWIN**, estimates the 48-hour **LC₅₀** for daphnia to be 15000 **mg/l**.²

- References for supporting studies
2. Report to Celanese Chemical Company **Inc.** on Toxicology and Fate of Selected Industrial Chemicals in Aquatic Ecosystems. J.R. Walton and E.M. Davis, University of Texas at Houston, December 1980.
 3. ECOSAR modeling program, version **0.99f**, as found in **EPIWIN** v 3.05, Syracuse Research Corporation, Syracuse NY (April 2000).

Toxicity to Aquatic Plants**Type** **Toxicity to Aquatic Plants**

Test Substance **1,3,5-Trioxane**
 CAS Number: 110-88-3
 Purity not specified

Method

- Guideline OECD 201
- Test Type Algae Growth Inhibition
- GLP No
- ॐ 1990
- Species/Strain *Scenedesmus subspicatus* strain SAG 86.81
- Element Basis Cell growth
- Exposure Period 72 hours
- Analytical Monitoring **No**
- Statistical Methods None necessary due to lack of significant inhibition

Remarks Field for Test Conditions

0 Test Temperature Range Stated as 20° C	
0 Growth Medium Chemistry	OECD specified medium
0 Exposure Vessel	Erlenmeyer flask (250 ml) containing 100 ml media
0 Water Chemistry	pH remained between 7.9 and 9.7 with all test concentration similar to control.
0 Light Level and Quality	Not specified in report
0 Test Design	o Replicates: Four replicate determinations o Concentrations: 0, 3.9, 7.8, 15.625, 31.25, 62.5, 125, 250 and 500 mg/l
0 Method of calculating mean	Arithmetic
0 Exposure period	24, 48 and 72 hours

- 0 Deviations from OECD Guideline The study report does not specifically state that it complied with the OECD guideline; however, except for the temperature being one degree lower than recommended by the guideline, all other criteria were met.

Results

- Nominal Concentrations 0, 3.9, 7.8, 15.625, 31.25, 62.5, 125, 250 and 500 mg/l
- Units mg./l.
- EC₂₀ > 500
- EC₅₀ > 500
- EC₉₀ > 500
- NOEC 500

Remarks Field for 0 Biological Observations None noted

0 Cell Density in Each Flask at Each Time Point

Conc (mg/l)	Cell Growth (Fluorescence Units)		
	24 hrs	48 hrs	72 hrs
Control	262	812	2103
3.9	259	909	2368
7.8	253	825	2194
15.625	263	810	2157
31.25	246	792	2100
62.5	228	716	2023
125	227	705	1988
250	203	674	1766
500	212	675	1769

Conclusions

Remarks field The EC₂₀ (48 hour) and EC₅₀ and EC₉₀ (72 hour) were greater than 500 mg/l under these conditions.

Data Quality

Reliability

Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although it could not be established that the study was conducted in accord with GLP standards. Study design and reporting meets current OECD guidelines with minor exceptions.

References

Determination of the inhibitory effect of **1,3,5-Trioxane** on the cell multiplication of algae. BASF AG, Ludwigshafen/RheinGermany. Project Number 2/w569/89/t0, 2/w569/89/t24, 2/w569/89/t48 and 2/w569/89/t730. Translation of a report originally dated 23 Feb 1990.

Other

This study is supported by an earlier study, sponsored by Celanese, in which Trioxane was tested for growth inhibition of *Selenastrum cupricornutum*. In this study, algae growth was measured out to 14 days of exposure at Trioxane concentrations of 1000, 5000 or 10000 mg/l. Significant inhibition was seen only at 10000 mg/l and 5000 mg/l was determined to be the NOEC. ¹

This study is also supported by the EPA ECOSAR Modeling Program found in EPIWIN², which estimates the 96-hour EC₅₀ for green algae to be 8245 mg/l.

References for supporting studies

1. Report to Celanese Chemical Company Inc. on Toxicology and Fate of Selected Industrial Chemicals in Aquatic Ecosystems. J.R. Walton and E.M. Davis, University of Texas at Houston. December 1980.
2. ECOSAR modeling program, version 0.99f, as found in EPIWIN v 3.05, Syracuse Research Corporation, Syracuse NY (April 2000).

Acute Health Effects**Acute Oral Toxicity**

Type	Acute Oral Toxicity
Test Substance	Trioxane CAS Number: 110-88-3 Clear colorless crystalline material

Method

- Guideline Federal Hazardous Substance Labeling Act provisional guidelines FR 8/12/1961
- GLP No
- Year 1962
- Species Rat
- Strain Albino, unspecified strain
- Route of administration Oral Gavage
- Doses 2520, 5000 and 10000 mg/kg
- Sex Male
- Number of Animals/group Five
- Vehicle Water

Remarks Field for Test Conditions	0 Age at Study Initiation	Unknown, weight between 229 and 281 g
	0 Doses	2520, 5000 and 10000 mg/kg
	0 Volume administered	1 to 12 ml aqueous solution
	0 Post-dose observation period	14 Days
	0 Other	The high dose was split into two administrations in a single 8-hour period. Animals were not fasted.

Results

- **LD₅₀** 8190 **mg/kg** (95% confidence limits of 6210 to 10080 **mg/kg**)

	Dose	<u>Mortality</u>
• Number of deaths at each dose level	2520 mg/kg	0/5
	5000 mg/kg	0/5
	10000 mg/kg	4/5

Remarks Field for Results **O** Time of death Not stated

O Clinical Signs Not stated

◇ Body Weights

Dose (mg/kg)	Initial weight	Weight gain	Control gain
2520	263	+24	+60
5000	249	+39	+60
10000	248	+23	+60

O Necropsy Findings None

O Target Organs None identified

Conclusions

Remarks field Study documentation is good for the time. Results are consistent with other data for the material, including a literature study. The study appears to have been well conducted.

Data Quality

- Reliability Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards.

References

Range Finding Tests on Trioxane for Celanese Corporation of America. Industrial Hygiene Foundation of America, Inc, Mellon Institute, October November 1962.

Other

This study is supported by a study¹ in the literature that reports the acute oral LD₅₀ in the rat to be 8,500 **mg/kg**

- References for supporting studies
1. Czajkowska, T, Krysiak, B and Popińska, E. Experimental studies of toxic effects of 1,3,5-trioxane and 1,3-dioxolane. I. Acute toxic effect. Med Pr; Vol 38, 1987, P184-90.

Acute Inhalation Toxicity**Type** Acute Inhalation Toxicity

Test Substance 1,3,5-Trioxane
 CAS Number: 110-88-3
 C-235
 Lot41115

Method

- **Guideline** None specified
- **GLP** Yes
- **Year** 1986
- **Species** Rat
- **Strain** Charles River CD@
- **Route of administration** Whole-body inhalation as vapor
- **Doses** 8,370 and 10,643 ppm (maximum attainable concentration at high dose)
- **Sex** Males and Females
- **Exposure Period** Four hours
- **Number of Animals/group** Five animals of each sex per dose level
- **Vehicle** Air

Remarks Field for Test Conditions

0	Age at Study Initiation	Males: 9 weeks Females: 12 weeks
0	Doses	Initial dose was 8,370 ppm (measured), as mortality was not observed, a second group was exposed at the maximum attainable vapor concentration, which was measured at 10,643 ppm. Particle size distribution analysis showed no particles were present, thus it is assumed that this was a vapor exposure.
0	Post-dose observation period	14 Days

Results

- LC₅₀ >10,643 ppm
- Number of deaths at each dose level No deaths reported

Remarks Field for Results

Clinical Signs
8,370 ppm

Sign	Onset	Duration	# Animals
Lacrimation	4 hours	14 days	Few
Shallow breathing	4 hours	< 24 hours	6/10
Irregular breathing	4 hours	2 days	9/10
Reduced activity	45 minutes	< 24 hours	Most
Nasal discharge	4 hours	14 days	About half

Clinical Signs
10,643 ppm

Sign	Onset	Duration	# Animals
Lacrimation	15 minutes	13 days	many
Shallow breathing	4 hours	< 24 hours	half
Irregular breathing:	30 minutes	5 days	All
Reduced activity	15 minutes	< 24 hours	All
Nasal discharge	4 hours	14 days	Few to most

Mean Body Weights
8,370 ppm

Day	0	2	3	5	8	15
Males	338	310	319	325	342	364
Females	230	210	218	221	233	239
Males	329	293	292	310	330	352
Females	226	210	216	216	224	235

10,643 ppm

Necropsy Findings
8370 ppm

Encrusted ear	1/10
Dilated kidney pelvis	1/10
Discolored lymph node	1/10

Necropsy	Encrusted ear	2/10
Findings	Dilated kidney pelvis	1/10
10,643 ppm	Calculi in kidney pelvis	1/10
	Discolored lymph node	1/10
	Ureter distended	1/10
	Urinary bladder thickened	1/10
	Urinary bladder calculi	1/10

Other Necropsy findings considered unremarkable. No target organs were identified at necropsy. Response of males and females was similar. There was a compound-related reduction in body weight following exposure. Animals gained body weight normally during the second week of the observation period

Conclusions

Remarks field

All test animals survived until termination. There was a slight diminution in body weight for each sex in each group following exposure; however, body weights during the second week were unremarkable. Other signs of treatment included increased secretory response, respiratory distress, and general signs of poor condition. Gross postmortem findings were considered unremarkable.

Data Quality

- Reliability

Klimisch Code 1 Although not a guideline study, the study was substantially similar, well documented, conducted under **GLPs** and measured concentrations of test material were used.

References

An Acute Inhalation Toxicity Study of Trioxane (C-235) in the Rat. **Bio/dynamics** Inc. Project 85-7832, submitted to Celanese Corporation, Feb 27 1986.

Other

This study is supported by a study in the literature that reports the acute inhalation **LC₅₀** to be greater than 26000 **mg/m³** (**>6500 ppm**).'

References for supporting data

1. Czajkowska, T, **Krysiak**, B and **Popińska**, E. Experimental studies of toxic effects of **1,3,5-trioxane** and **1,3-dioxolane**. I. Acute toxic effect. Med Pr; Vol 38, 1987, **P184-90**.

Acute Dermal Toxicity**Type** Acute Dermal Toxicity

Test Substance Trioxane
 CAS Number: 110-88-3
 Clear colorless crystalline material

Method

- Guideline Federal Hazardous Substance Labeling Act provisional guidelines FR 8/12/1961
- GLP No
- Year 1962
- Species Rabbit
- Strain Albino, unspecified strain
- Route of administration Dermal administration
- Doses 3,980 mg/kg
- Sex Male
- Exposure Period Twenty-four hours
- Number of Animals/group Four
- Vehicle None

Remarks	Field for	0	Age at Study Initiation	Unknown
Test Conditions		0	Doses	Single dose level of 3,980 mg/kg.
		0	Post-dose observation period	14 Days
		0	Other	Two animals had skin abraded prior to exposure

Results

- LD₅₀ >3,980 mg/kg
- Number of deaths at each dose level No deaths reported

Remarks Field for Clinical Signs The only reported clinical sign was very slight to moderate erythema but no appreciable edema of the skin after removal of the wrap. This disappeared by study termination

Results 3,980 mg/kg

Sex	Day	
	0	14
Males	2520 g	2570 g

Mean Body Weights

Necropsy Findings None

Conclusions

Remarks field

All test animals survived until termination. Two of the four rabbits showed a reduction in body weight gain as compared to controls. Gross postmortem findings were unremarkable. The material produced slight to moderate skin irritation.

Data Quality

- Reliability

Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards.

References

Range Finding Tests on Trioxane for Celanese Corporation of America. Industrial Hygiene Foundation of America, Inc, Mellon Institute, **October-November** 1962.

Other

This study is supported by a study in the literature that reports the acute dermal LD₅₀ in rabbits to be greater than 15000 g/kg (1).

References for supporting data

1. Czajkowska, T, Krysiak, B and Popińska, E. Experimental studies of toxic effects of **1,3,5-trioxane** and **1,3-dioxolane**. I. Acute toxic effect. Med Pr; Vol 38, 1987, **P184-90**.

Repeated Dose Toxicity, 28-Day Oral

Type	Repeated Dose Toxicity, 28-Day Oral
Test Substance	Trioxane CAS Number: 11 O-88-3 99.1% purity (0.8% water, 0.1% Triethanolamine)
Method	
• Guideline	OECD 407 Repeat Dose Oral Toxicity (198 1)
• GLP	Yes
• Year	1990
• Species	Rat
• Strain	Wistar [Hoe: WISKf(SPF7 1)] Source: Hoechst AG, Kastengrund
• Route of administration	Oral gavage
• Duration of Test	29 days
• Doses	0, 40, 200, or 1000 mg/kg/day
• Sex	Male and Female
• Exposure Period	Once daily
• Frequency of Treatment	0 Seven days a week 0 Twenty-eight doses in twenty-nine days
• Number of Animals/group	Five of each sex
• Vehicle	Deionized water
• Control Group and Treatment	Five animals of each sex dosed with vehicle (water)
• Post-Exposure Observation Period	Twenty-four hours after last dose.
• Statistical Methods	Referenced established methods

Remarks Field for Test Conditions	0	Age at study initiation	Six weeks
	0	Number of animals per Sex per dose	Five of each sex per concentration
	0	Satellite groups	None
	0	Housing	Housed 5 to a group on soft wood granules
	0	Test Material Preparation	Trioxane was prepared fresh each day of dosing and was not analyzed for concentration or stability. It is known to be relatively stable in water solution.
	0	Rationale for dose selection	The high-dose of 1000 mg/kg day is the maximum dose level recommended in the OECD Guideline. The fivefold spacing was selected to assure that a NOEL would be established.
	0	Clinical observations performed and frequency	<ul style="list-style-type: none"> ■ Mortality and gross signs: Twice daily except once daily on weekends and holidays. ■ Body weights were measured twice a week ■ Possible neurological disturbances were evaluated weekly ■ Weekly examinations were conducted for the following: eyes for signs of opacity, the mucus membranes of the mouth for damage, and the teeth for unusual growth.
	0	Terminal observation	<ul style="list-style-type: none"> ■ Blood taken for hematology and clinical chemistry (Na, K, P04, uric acid, creatinin, glucose, BUN, Ca, Cl, SGOT, SGPT, AP, GGT, total protein, albumin, globulin (calculated) A/G ratio). ■ Serum electrophoresis. ■ Urinalysis for a 16 hour sample ■ Gross postmortem examination including external surfaces, all orifices, eyes, teeth, and inner organs of all animals. ■ Organ weights for heart, lungs, liver, kidneys, spleen, testes and adrenals. ■ Microscopic examination for several tissues.
	0	Histopathology	The following tissues were examined from all animals: Heart, lungs, liver, kidneys, spleen, testes and adrenals, thymus, jejunum, colon, bone marrow and stomach.
	0	Differences from current guideline	The current OECD 407 guideline (adopted in 1995) recommends additional tissues for gross and histopathological examination.

Results

. NOAEL 200 mg/kg /day

. LOEAL 1000 mg/kg/day

- Mortality All animals survived the duration of the study.

Toxic Responses

Dose	Toxic Responses
40 mg/kg	None
200 mg/kg	None
1000 mg/kg	<ul style="list-style-type: none"> ▪ Males: Significant decrease in leucocyte count, increase in gamma-glutamyl-transpeptidase. No changes in body or organ weight nor any significant histopathological findings. The study director did not consider the finding of a single animal reported to show testicular atrophy significant. ▪ Females: Significant decrease in leucocyte count, increase in gamma-glutamyl-transpeptidase, increase in SGPT and SGOT, decrease in serum protein and glucose levels. No changes in body or organ weight nor any histopathological findings

Remarks Field for Results

◇ Body Weights (Males)	Dose (mg/kg)	Initial weight	Final weight	Weight gain
	0	124±7	282±31	158±25
40	122±7	282±15	160±13	
200	121±3	281±15	161±05	
1000	122±5	257±22	135±18	

0 Body Weights (Females)	Dose (mg/kg)	Initial weight	Final weight	Weight gain
	0	112±4	195±15	83±10
40	112±2	195±10	83±12	
200	110±6	194±14	83±16	
1000	113±4	188±07	75±04	

0 Food/water	Dose	Food Consumption (g/ 00g body wt/day)		Water Consumption (g/100g bodywt/16 hr)	
		Males	Females	Males	Females
	0	10.0	9.7	13.1	12.4
	40	10.2	10.1	12.5	12.3
	200	10.4	10.1	12.6	14.8
	1000	9.7	9.9	11.9	10.3

0 Clinical Signs No clinical signs of toxicity were observed.

Dose (mg/kg)	Leukocytes ($10^9/L$)	
	Males	Females
0	6.1±0.6	4.3±0.8
40	6.3±1.4	5.5±1.2
200	4.8±0.1	4.7±2.2
1000	3.6±0.6*	2.6±0.5*

Dose (mg/kg)	Selected Clinical Chemistry Values - Males				
	GGT (U/L)	SGOT (U/L)	SGPT (U/L)	Protein (g/L)	Glucose (mmol/L)
0	0±0	77±11	43±12	57±4	10.8±6.2
40	0±0	68±06	36±03	56±3	9.6±0.8
200	1±1	72±09	43±03	57±2	10.5±0.8
1000	1±1	75±12	58±14	56±1	10.4±1.1

Dose (mg/kg)	Selected Clinical Chemistry Values - Females				
	GGT (U/L)	SGOT (U/L)	SGPT (U/L)	Protein (g/L)	Glucose (mmol/L)
0	1±1	64±4	31±3	59±4	16.6±3.4
40	0±0	71±16	31±3	57±3	11.9±1.3
200	0±1	61±9	33±5	56±2	12.0±5.9
1000	1±1	84±9*	45±2*	53±2*	7.5±0.9*

Other chemistry parameters showed significant changes from controls but were considered not compound-related as they were within the range of historical controls for this strain.

0 Necropsy Findings None reported

0 organ Weight The absolute spleen weights of high-dose males were significantly reduced compared to controls, relative spleen weights of males were not significantly affected. In females, the mid-dose mean spleen weight was significantly higher than control but this was disregarded due to lack of dose response.

Dose (mg/kg)	Selected Final Organ Weights, males (grams)			
	Spleen	Kidney	Testes	Liver
0	0.61±0.12	1.85±0.19	2.99±0.33	11.61±1.50
40	0.60±0.06	1.89±0.10	2.95±0.28	11.05±1.26
200	0.63±0.07	1.85±0.12	3.19±0.18	11.30±0.94
1000	0.47±0.06*	1.65±0.25	2.48±1.02	11.40±1.66

Dose (mg/kg)	Selected Final Organ Weights, females (grams)		
	Spleen	Kidney	Liver
0	0.42±0.03	1.45±0.14	8.44±0.83
40	0.47±0.03	1.35±0.07	7.84±0.62
200	0.52±0.06*	1.45±0.09	8.47±0.69
1000	0.39±0.05	1.29±0.09	7.74±0.56

- 0** Histopath The examining pathologist considered no finding significant. Few findings were reported and treated animals were similar to controls. Of interest for this material is the testis, in which one high-dose male (of five) was reported to show testicular atrophy.
- 0** Target Organs Blood **leucocytes**, liver
- 0** Other Serum electrophoresis results were similar for treated and control groups.

Conclusions

Remarks field

The NOEL was determined to be 200 **mg/kg/day**. Adverse clinical effects were not observed at any dose level. Body weight gain was unaffected; food and water consumption were similar for all groups. Leukocyte levels were significantly depressed at the high-dose level. Clinical chemistry parameters indicative of hepatic damage were elevated in high-dose females but no corresponding histopathologic effects were observed.

This is a well-documented guideline study conducted under **GLPs**

Data Quality

- Reliability

Klimisch Code 1. Reliable without restriction, study meets GLP standards and/or most requirements.

References

Trioxane. Subakute **orale** Toxizität (28 Application in 29 Tagen) an **SPF-Wistar-Ratten**. Hoechst AG Central Toxicology Report number 90.05 13 22 May 1990.

* = p < 0.05

Other

This study is supported by a gavage study of duration four or seven months, depending on the dose level, conducted using male rats. In this study, groups of 8-10 rats were administered Trioxane as a water solution, 5 days a week. The high dose was 850 **mg/kg** and the duration of dosing was 4 months. Doses of 213 or 106 **mg/kg** were administered for 7 months. Body weight gain, as compared to controls, was slightly reduced at 850 and at 213 **mg/kg** and slightly increased at 106 **mg/kg**. At the end of the exposure period, blood was taken and evaluated for several hematology and clinical chemistry parameters. Heart, liver, lung, kidney, adrenals and spleen were weighed at necropsy and examined microscopically. No specific organ effects were reported and it was concluded that the mortality rate did not show unexpectedly high mortality or other toxicity with longer-term exposure. Examination of the limited data presented suggest that 213 **mg/kg** was near the NOAEL for Trioxane after 7-months of administration. It could not be determined if the same parameters were examined in high-dose animals, which were sacrificed **after** only 4 months of exposure, or if a concurrent control group was sacrificed with the high-dose group. It is not known if there were any effects on spleen or **WBC's** as the information available for this study lacks sufficient documentation for thorough evaluation and comparison with the repeated-dose studies.

Reference for
supporting study

Experimental studies of the toxic effects of **1,3,5-trioxane** and **1,3-dioxolane**. II. Cumulation of toxic effect. Czajkowska T; Krysiak B Med Pr; VOL 38, 1987, P244-9

Repeated Dose Toxicity, Two-Week Inhalation**Type** **Repeated Dose Toxicity, Two-Week Inhalation**

Test Substance **1,3,5-Trioxane**
 CAS Number: 110-88-3
 White crystalline solid, C-235

Method

- Guideline None Specified
- GLP No
- Year 1983
- Species Rat
- Strain CD (Sprague-Dawley derived)
 Charles River, Wilmington MA
- Route of administration Whole body inhalation of vapor
- Duration of Test 12 days
- Doses 0, 103,984 or 4940 ppm (Mean, measured concentrations)
- Sex Male and Female
- Exposure Period Six hours per day
- Frequency of Treatment Five days a week
 Ten exposures in twelve days
- Number of Animals/group Five of each sex
- Control Group and Treatment Five animals of each sex exposed only to air under the same chamber conditions
- Post-Exposure Observation Period None, animals sacrificed immediately after last exposure

- Statistical Methods
 - Body weight data, hematology and clinical chemistry parameters, organ weights and organ/body weight ratios were analyzed. Mean values of the treated groups were compared to control at each time interval.
 - Statistical evaluation of equality of means was made by the appropriate one-way analysis of variance technique, followed by a multiple comparison procedure if needed. First, Bartlett's test was performed to determine if groups had equal variance. If the variances were equal, parametric procedures were used; if not, nonparametric procedures were used. The parametric procedures were the standard one-way **ANOVA** using the F distribution to assess significance. If significant differences among the means were indicated, Dunnett's test was used to determine which means were significantly different from the control. If a nonparametric procedure for testing equality of means was needed, the **Kruskal-Wallis** test was used, and if differences were indicated a summed rank test (Dunn) was used to determine which treatments differed from control.
 - A statistical test for trend in the dose levels was also performed. In the parametric case, (i.e. equal variance) standard regression techniques with a test for trend and lack of fit were used. In the nonparametric case, Jonckheere's test for monotonic trend was used.
 - The test for equal variance (Bartlett's) was conducted at the 1%, two-sided risk level. All other statistical tests were conducted at the 5% and 1%, two-sided risk level.

- Remarks Field for Test Conditions
 - Age at study initiation
 - . Males: 41 days
 - . Females: 58 days
 - Number of animals per Sex per dose
 - . Five of each sex per concentration
 - ◇ Satellite groups
 - . None
 - ◇ Housing
 - . Individually housed in stainless steel cages

 - ◇ Clinical observations performed and frequency
 - . Mortality and gross signs: Twice daily
 - . Abnormal signs: Daily
 - . Detailed physical examination: Twice weekly

 - Terminal observations
 - . Blood taken for hematology and clinical chemistry (sgpt, bun, glucose, total protein, albumin, globulin (calculated) **A/G** ratio).
 - . Complete gross postmortem examination including external surfaces, all orifices, the cranial cavity, carcass, the external surface of the brain and spinal cord, the thoracic, abdominal and pelvic cavities and their viscera and the cervical tissues and organs were examined for all animals.

- 0 Histopathology The following tissues were examined from all control and high-exposure animals only. Bone marrow smear, right kidney, liver, lungs, peribronchial lymph nodes, nasal turbinates and thymus.

Results

- . NOAEL 0 Males: Not determined (< 103 ppm)
0 Females: 103 ppm
- . LOEAL 0 Males 103 ppm
0 Females: 984 ppm
- Mortality All animals survived the duration of the study.

- Toxic Responses

Dose	Toxic Response
103 ppm	A decrease in mean absolute and relative spleen weights was observed for males only. All other findings occurred sporadically and were not considered to be related to exposure
984 ppm	A decrease in mean absolute and relative spleen weights was observed for males and females. All other findings occurred sporadically and were not considered to be related to exposure
4940 ppm	Exposure-related effects were noted in each sex of this group. These included increased secretory responses, reduced righting reflex and grip strength, persistent pupillary constriction and decreased mean body weights throughout most of the study. In addition, hemoglobin, hematocrit, erythrocyte counts and lymphocyte counts were elevated, while total leukocytes and segmented neutrophils were decreased. Slight to significant increases in mean serum glutamic pyruvic transaminase, total protein and albumin values, and a decrease in glucose values were also noted. In the absence of supportive microscopic findings in the liver and kidneys of these animals, the significance of the clinical pathology is unclear. Mean absolute and relative spleen weights were decreased for the high-dose animals. Increased mean relative weights for all other weighed organs were associated with significantly decreased mean terminal body weights for this group. The mucosa of the anterior nasal cavity from the high-exposure males and females showed squamous metaplasia with necrosis and desquamation. Acute rhinitis with neutrophil exudation into the nasal cavity was a concomitant change.

- Remarks Field for \diamond Body Weights Results

	Test Day and Time					
Males	Pretest	1 (AM)	1 (PM)	5 (AM)	5 (PM)	12 (AM)
Control	162	202	189	224	214	276
103 ppm	163	203	190	226	215	278
984 ppm	162	202	187	223	212	268
4940 ppm	163	207	189	202**	193**	239**
Females						
Control	183	201	191	198	193	228
103 ppm	182	197	184	205	195	219
984 ppm	183	199	185	203	192	213
4940 ppm	183	199	183	189	179	200**

0 Clinical Signs

Clinical Signs		Day (# Animals/10)			
		2	5	9	12
0 ppm	Lacrimation	0	0	0	1
	Mucoid nasal discharge	0	0	0	1
	Swollen conjunctiva	0	0	0	4
103 ppm	Lacrimation	2	2	1	5
	Mucoid nasal discharge	8	8	1	0
	Dry rales	2	0	0	1
	Swollen conjunctiva	0	0	0	4
984 ppm	Lacrimation	9	5	0	7
	Mucoid nasal discharge	10	9	3	1
	Red nasal discharge	0	0	0	1
	Salivation	1	1	0	0
	Dry rales	1	0	0	0
	Swollen conjunctiva	0	0	0	1
4940 ppm	Lacrimation	10	1	2	6
	Mucoid nasal discharge	9	10	7	6
	Red nasal discharge	0	0	0	1
	Dry rales	0	0	0	4
	Moist rales	0	0	2	1
	Reduced righting reflex	10	6	10	9
	Reduced grip	9	1	9	5
	Persistent pupil constriction	8	0	8	1
	Swollen conjunctiva	0	0	0	3
	Yellow fur	0	0	4	7
	Negative toe pinch reflex	1	0	0	0
	Yellow ano-genital fur	1	1	2	1

0 Hematology

PPM	HGB	HCT	RBC	Platelets	MCV	MCH	MCHC	Clot T	WBC
Males	g/dl	%	10 ⁶ /mm	10 ⁵ /mm	μ ³ /cell	μg	g/dl	min	103/mm
0	13.6	40	6.20	15.22	65	22	33.9	1.4	13.9
103	14.0	41	6.28	13.43	66	22.3	33.9	1.6	13.8
984	13.9	41	6.11	14.16	67	22.7	34.1	1.4	13.3
4940	15.2**	45**	6.93**	15.14	64	22	34.2	1.7	6.2**
Females									
0	14.1	42	6.45	10.77	65	21.9	33.7	1.6	11.0
103	14.5	43	6.76	13.24	64	21.5	33.3	1.5	11.0
984	14.8	44	6.79	13.26	65	21.8	33.8	1.7	10.8
4940	15.4**	46**	7.13*	11.46	64	21.7	33.7	1.5	8.5

- 0** Necropsy findings Gross postmortem examinations revealed no differences among groups which were considered to represent an effect of exposure to the test substance
- 0** Organ weights Mean absolute and relative spleen weights were decreased for the 103 ppm-group males and the mid and high-dose males and females; differences were indicative of an exposure-response relationship. Increased mean relative weights for all other weighed organs (brain, heart, kidneys, liver, testes/ovaries) were noted for the high-dose males and/or females; however, these differences were associated with the significantly decreased mean terminal body weights exhibited by these animals.
- 0** Histopathology Significant changes in the mucosa of the anterior nasal cavity were noted in the high-dose males and females. The columnar ciliated epithelium was replaced by a less specialized squamous epithelium. The metaplasia was focal, and the epithelium showed various degrees of degeneration, necrosis and desquamation. Concomitant with this change was an acute inflammation of the nasal mucosa characterized by collections of neutrophils in the lamina propria and mucosa. Exudation of neutrophils produced an accumulation of purulent exudate in the nasal cavity. In a few cases, a small layer of packed neutrophils was adherent to the mucosa. Proteinaceous material, usually with enmeshed neutrophils, was seen predominantly in the high-dose males. Microscopic changes in other organs occurred sporadically in both the control and high-exposure animals and did not indicate any exposure related effects.

Findings in the Nasal Turbinate (NI section)	Exposure Concentration (ppm)			
	0	103	984	4940
Acute Rhinitis	0/10	3/10	6/10	7/10
Mucosa:				
-Squamous Metaplasia	0/10	0/10	8/10	10/10
-Erosion	0/10	0/10	7/10	10/10
-Hyperplasia	0/10	0/10	2/10	0/10

Conclusions

Remarks field

Two-weeks of inhalation exposures at concentrations of 984 ppm and above produced a variety of toxic sequelae including a decrease in weight gain, splenic atrophy and squamous metaplasia in the mucosa of the anterior nasal passage. The LOAEL was 103 ppm for males showing a decrease in mean absolute and relative spleen weights. 103 ppm was considered a NOAEL for females.

Study documentation is good. Results are consistent with other data in the literature and the study appears to have been well conducted.

Data Quality

- Reliability

Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards.

References

Bio/Dynamics Inc. A Two-Week Inhalation Toxicity Study of C-235 in the Rat. Project Number 82-7572 1983 Submitted to Celanese Corporation.

Other

This study is supported by a **12-month** inhalation study in the literature where concentrations of 50 500 and 2500 mg/m^3 were found to be toxic to rats or guinea pigs, and a NOEL concentration of 50 mg/m^3 was reported. (50 mg/m^3 is about 12 ppm) The 500 mg/m^3 groups and higher showed adverse pathological changes of the kidneys and respiratory epithelium. Few details are available. Considering the extended length of this study and that the next higher dose level was 500 mg/m^3 (about 120 ppm), which was apparently a LOAEL; the 103 ppm NOAEL/LOAEL from the 14-day inhalation study is in accord with the longer study. It is not known if there were any specific adverse effects on spleen or **WBC's** in this long-term study since the available report lacks sufficient documentation for adequate evaluation and comparison with the repeated-dose study.

References for supporting studies

1. Indulski et al. MAC Values for trioxane and dioxolane at the work place proposed on the basis of animal studies. Fourteenth International Congress on Occupational Health in the Chemical Industry (MEDICHEM), pp 548-556 (1986). As summarized in Toxikologische Bewertung. Heidelberg, Berufsgenossenschaft der chemischen Industrie *Trioxane* Nr. 185 (1992)

** = $p < 0.01$

Genetic Toxicology *in vitro*

Reverse mutation assay, Ames Test

Type Reverse mutation assay - *S. typhimurium*

Test Substance 1,3,5-Trioxane
 CAS Number: 110-88-3
 Substance-Nr.:88/164

Method

- **Guideline** None specified
- **GLP** No
- **Year** 1988
- **Species**
 - 0 *S. typhimurium*: TA1535, **TA100**, TA 1537, TA98,
 - 0 Metabolic activation. Tested with and without
 - 0 Rat liver S-9
 - 0 0.5 ml S-9 per 100 ml agar plate when used
 - 0 Triple plate
 - 0 Independent repeat using pre-incubation method for repeat
- **Concentrations tested**
 - Concentrations tested (micrograms per plate)
 - First Experiment (no preincubation) and second experiment (preincubation)
 - 0 TA98 0, 20, 100, 500, 2500 and 5000
 - 0 **TA100** 0, 20, 100, 500, 2500 and 5000
 - 0 TA1535 0, 20, 100, 500, 2500 and 5000
 - 0 TA1537 0, 20, 100, 500, 2500 and 5000
- **Statistical Methods** Not specified
- Remarks Field for Test Conditions**
 - 0 Positive controls
 - 0 Without activation
 - o **AAC**: TA1537
 - o **NPD**: TA98,
 - o **MNNG**: TA1535, TA100
 - 0 With activation
 - 0 2-Aminoanthracene: all strains
 - 0 Solvent: aqua dest.
 - 0 Study generally follows OECD 47 1 differences include only four strains of bacterial used where five is recommended and AAC used as sole positive control under activation conditions.

Results

- **Result** No dose-dependent increase in the number of revertants in any bacterial strain in the presence or absence of metabolic activation. Positive controls demonstrated the sensitivity of the test system
- **Cytotoxic concentration** No cytotoxicity observed at any level
- **Genotoxic Effects** Not genotoxic under these conditions

Remarks Field for Results No visible precipitation was observed at any concentration. No positive responses (doubling of the control mutation rate) were observed. The material was water soluble

Conclusions

Remarks field No genotoxic activity. Study well conducted. Documentation is fair.

Data Quality

- **Reliability** Klimisch Code 2 Reliable with restrictions. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards. Supporting studies increase the reliability of the conclusion.

References

Report on the Study of **1,3,5-Trioxane** (ZST Test Substance No.: **88/164**) in the Ames Test. Project 40M0 164/884027. BASF AG. Department of Toxicology, Z 470, **Ludwigshafen/Rhein**, Germany (1988)

Other

This study is supported by a single-plate test from another laboratory using five strains of bacteria (TA98, **TA100**, **TA1535**, TA 1537 and TA1538) exposed with and without S-9 activation at eight concentrations of Trioxane from 0.5 to 5000 micrograms per plate. No mutagenic response was recorded in this study. ¹

Additional support comes from two additional published studies in prokaryotes which produced negative results. In a triple plate test, with preincubation in the presence and absence of S-9 with strains TA97, TA98, **TA100** and **TA1535** at concentrations from 100 to 10000 micrograms per plate, Trioxane was reported to be without mutagenic activity.² The mutagenic activity of Trioxane was investigated in **five** strains of *S. typhimurium*: TA1535, TA1537, **TA1538**, TA98 and **TA100** with and without activation by liver microsomes (induced with Aroclor 1254). It displayed no mutagenic activity under any of these conditions.³

References for
Supporting Studies

1. Mutagenicity Evaluation of C- 120 in the Ames **Salmonella/Microsome Plate Test**. Litton Bionetics Project # 20988. Sponsored by Celanese Corporation. 1980.
2. Zeiger, E, Anderson, B, **Haworth, S**, Lawlor, T, Mortelmans, K. Salmonella mutagenicity tests. 4 Results from the testing of 300 chemicals. *Environ Mol Mutagen* **11(Suppl 12):1-158,1988**.
3. Kowalski, Z, Spiechowicz, E, And Baranski, B. Absence of mutagenicity of trioxane and dioxolane in Salmonella typhimurium. *Mutat Res*; 136 (3). 1984. 169-172

Mammalian Cell Transformation Assay, *in vitro*

Type Mammalian Cell Transformation Assay, *in vitro*

Test Substance 1,3,5-Trioxane
CAS Number: 11 O-88-3
C-235

Method

- **Guideline** None specified
- **GLP** No
- **Year** 1981
- **Cell Type**
 - 0 C3H 10T-1/2 clone 8
 - 0 Six replicates at each concentration in 60 mm dishes, 11 -day incubation
 - 0 Six replicates at each concentration in T-25 flasks, 11 -day incubation
 - 0 Six replicates at each concentration in 60 mm dishes, **38-day** incubation
 - 0 Six replicates at each concentration in T-25 flasks, **38-day** incubation
- **Concentrations tested** Concentrations tested (micrograms per ml)
 - 0 11 -day incubation: 0, **1, 10, 100, 500, 1000, 5000, 10000** and 20000
 - 0 **38-day** incubation: 0, 1, 10, 100, 500, **1000, 5000, 10000** and 20000
 - 0 **24-** ~~hour~~ exposure to test material
 - 0 60 mm dishes sealed in jar to reduce loss of test material
 - 0 Vehicle: water or culture media
- **Statistical Methods** Not specified
- **Remarks Field for Test Conditions**
 - 0 Positive control: Benzpyrene
 - 0 Cultures re-fed on regular schedule
 - 0 Colonies examined macroscopically and microscopically.
 - 0 300 cells per flask or dish
 - 0 Colony-forming potential determined at 11 -day interval
 - 0 Foci transformation potential determined at **38-day** interval

Results

- **Result** No increase in the number of transformed colonies or transformed foci was noted at any concentration of test material. Positive controls demonstrated the sensitivity of the test system
- **Cytotoxic Concentration** Severe cytotoxicity observed above 10000 micrograms/ml, slight **dose-**dependent cytotoxicity observed at 2000 microgram per ml and above.
- **Genotoxic Effects** Not genotoxic under these conditions

Remarks Field for Results The test material was water-soluble; precipitation was not recorded.

Conclusions

Remarks field No genotoxic activity. Study was well conducted. Documentation is good.

Data Quality

- **Reliability** **Klimisch** Code 2. Reliable with restrictions, study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards.

References

Mammalian Cell Transformation Assay, University of Minnesota Experimental Pathology Laboratory, submitted to Celanese Corporation, dated 8/27/8 1.

Mouse lymphoma forward mutation assay**Type** Mouse lymphoma forward mutation assay**Test Substance** 1,3,5-Trioxane
CAS Number: 110-88-3**Method**

- **Guideline** None specified
- **GLP** No
- **System of Testing** Non-bacterial
- **Year** 1980
- **Species/Strain** Mouse lymphoma L5 178Y TK+/-
- **Metabolic activation**
 - Tested with and without
 - 0 Rat liver S-9
 - 0 Aroclor 1245 induced
 - 0 Final concentration 5% S-9 in cell suspensions
- **Concentrations tested** 0, 0.313, 2.5, 6.25, 12.5 and 15 **mg/ml** for both non-activation and activation conditions in first trial; **0, 0.156, 0.625, 2.5, 5.0 and 7.5 mg/ml** in the second trial under activation conditions only.
- **Statistical Methods** Simple ratio criteria

Remarks Field for Test Conditions

- 0 Independent repeat of activation conditions only
- 0 Triplicate plates for counting mutant colonies
- 0 Solvent, water
- 0 Negative control, water
- 0 Positive controls
 - o Without activation - EMS
 - o With activation - DMN

Results

- Result No dose-dependent increase in the number of mutants was observed in the absence of metabolic activation; however, the number of mutants was found to increase in the presence of metabolic activation. Positive controls demonstrated the sensitivity of the test system.
- Cytotoxic Concentration No cytotoxicity observed under non-activation conditions. Under activation conditions, cytotoxicity was dose-dependent **with** moderate cytotoxicity observed at the lowest dose and extensive cytotoxicity reported for the highest concentrations.
- Genotoxic Effects
 - 0 No genotoxic activity under non-activation conditions
 - 0 Concentration-dependent mutational activity under activation conditions.

Remarks Field for Results

- 0 Material was soluble in water
- 0 Under activation conditions, 4 fold increases in mutant frequency were observed at the moderately-toxic 0.156 **mg/ml** and the **frequence** increased to the 10 to 20 fold range at highly toxic concentration of 5 and 7.5 **mg/ml**.
- 0 The positive control (**DMN**) under activation gave lower than expected mutational frequencies in the first trial but normal frequencies in the second trial indicating that the S9 activity was adequate.

Conclusions

Remarks field

- ◇ No genotoxic activity under non-activation conditions
- ◇ Concentration-dependent mutational activity under activation conditions.
- ◇ Mutational activity appeared to correlate with cytotoxicity suggesting metabolism by S-9 to active material.
- ◇ Study was well conducted, although there was no GLP certification the study appears to have been conducted using a GLP-quality protocol in a GLP compliant laboratory.

Data Quality

- Reliability Klimisch Code 2. Reliable with restrictions. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards.

References

Mutagenic Evaluation of C- 120 in the Mouse Lymphoma Forward Mutation Assay. Submitted to Celanese Corporation. Litton Bionetics, Inc, Kensington Maryland. **LBI** Project No. 20989, November 1980.

Genetic Toxicology *in vivo***Mouse micronucleus assay****Type** Mouse micronucleus assay

Test Substance **1,3,5-Trioxane**
 CAS Number: 11 0-88-3
 Source: Nitrogen Plant in Tamow, Poland

Method

- **Guideline** None specified
- **GLP** No
- **Year** 1983
- **Species/Strain** Mice/ **BALB/c**
- **Sex** Male
- **Route of Administration** **IP** Injection in water
- **Doses** **0, 2125 and 4250 mg/kg** split into two doses given 24 hours apart
- **Exposure Period** Approximately 30 hours
- **Statistical Methods** The **Wilcoxon** test was used for statistical analysis of results. A positive response was defined as consisting of an increase in the incidence of polychromatic erythrocytes with micronuclei to at least twice the negative control value and a significance of at least $p \leq 0.05$.

- Remarks Field for Test Conditions**
- ① **Age at Study Initiation** Seven to eight weeks
 - ① **Number of animals per dose** Four per group. The dose was split into 2 and injected at **24-hour** intervals. Animals were sacrificed 6 hours after the second dose of test material. Positive control groups contained three animals each.
 - ① **Rationale for Animals** The OECD guideline specifies rats of each sex unless it has been shown in other studies that there is no sex difference in toxicity. The toxicity of Trioxane is known to be similar in males and females

- 0** Control Groups and Treatment . Negative Control: Water injection
 . Positive Control: Mitomycin C in water. In a range of 0.65-5.2 mg/kg
- 0** Clinical Observations . None reported
- 0** Criteria for Evaluating Results . See statistical methods.
- 0** Criteria For Selection of MTD . Not specifically stated but the total high dose was half the “approximate lethal dose”.
- 0** Differences from Current OECD Guideline . This study varies from the current OECD 474 guideline. The significant variations are:
 - The guideline specifies 5 animals/group
 - The criteria of bone marrow toxicity at the high dose may not have been met; however, the “limit test” dosing of 2000 mg/kg was exceeded. The suggested number of dose levels is also irrelevant in light of the limit dose being met.

Results

- Genotoxic Effects None indicated.

Remarks Field for Results

◇ Induction of Micronucleated Cells

Trioxane			
Dose (mg/kg)	n	Number PCE examined	Incidence of micronuclei in PCE % ± sem
0	4	8000	0.42 ± 0.20
2125	4	8000	0.39 ± 0.20
4250	4	7550	0.47 ± 0.25
<i>Positive Control Mitomycin C</i>			
0.65	3	5650	1.73 ± 0.57
1.30	3	5660	3.00 ± 0.18
2.60	3	2800	3.47 ± 0.34
5.20	3	2300	2.24 ± 0.74

0 Mortality No mortality reported

Conclusions

Remarks field

0 There was no induction of micronucleated polychromatic cells associated with administration of Trioxane.

0 The high dose of dioxolane was stated to be toxic to the animals by the authors.

Data Quality

- Reliability

Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint.

Reference

Przybojewska B, Dziubautowska E, Kowalski Z. Genotoxic effects of dioxolane and trioxane in mice evaluated by the micronucleus test. **Toxicol Lett** **21:349-52** (1984)

In Vivo Unscheduled DNA Synthesis (UDS) Assay**Type** In Vivo Unscheduled DNA Synthesis (UDS) Assay

Test Substance 1,3,5-Trioxane
 CAS Number: 110-88-3
 99.9%
 Batch 65-3997

Method

- **Guideline** OECD 486 (using revised draft guideline dated March 1996)
- **GLP** Yes
- **Reference** 1997
- **Species** Rat
- **Strain/Sex** Wistar (Chbb : THOM; SPF) male
- **Route** Oral gavage
- **Vehicle** Water
- **Dose Levels** 0, 250, 5000, 1000 and 2000 mg/kg
- **Dosing Duration** Single dose
- **Cell Harvest Time** 4 and 18 hours after dosing
- **Statistical Methods** Not conducted since negative result

Remarks Field for Test Conditions

- ◇ Concentration of test substance determined analytically in vehicle
- ◇ High dose is maximum recommended under OECD 486
- ◇ Rats weighed mean of 258 grams at time of dosing, age not specified
- ◇ Three animals per group (dose level and harvest time)
- ◇ Positive control 50 mg/kg 2-acetylaminofluorene (in corn oil)
- ◇ Hepatocytes harvested by collagenase perfusion 4 and 18 hours after dosing
- ◇ After attachment, cells incubated 4-hours with tritiated-thymidine
- ◇ Cells washed and incubated 14-hours longer with cold thymidine (chase)
- ◇ Cells coated with photo emulsion, exposed 19 hours and developed
- V 100 random cells per animal counted for nuclear grains

Results

- Result No treatment-related increase in the number of cells under repair
- Cytotoxic Concentration > 2000 mg/kg
- Genotoxic Effects Not genotoxic under these conditions

Remarks Field for
Results

Dose	Viability (percent)	
	4-Hour Harvest	18-Hour Harvest
0	84	84
250	87	84
500	86	86
1000	82	84
2000	81	88
Positive Control	89	79

Cell
viability

DNA
Repair
Activity

Dose	Percent cells in repair	
	4-Hour Harvest	18-Hour Harvest
0	8.67	7.33
250	Not examined	Not examined
500	10.0	12.0
1000	8.67	11.7
2000	10.0	6.33
Positive Control	78.7	84.7

Conclusions

Remarks field

No genotoxic activity. Study was well conducted. Documentation is very good. Study exceeds requirements of current OECD 486 in that determination of cells in repair was further segregated into 1). Percentage cells in repair with net nuclear grain count greater than or equal to zero. and 2). Percentage cells in repair with net nuclear grain count greater than or equal to five.

Data Quality

- Reliability Klimisch Code 1. Reliable without restriction, study meets GLP standards and/or most requirements

References

In Vivo Unscheduled DNA Synthesis (UDS) Assay with 1,3,5-Trioxane in Rat Hepatocytes, Single Oral Administration. Department of Toxicology, BASG AG, Ludwigshafen. 12 March 1996.

Other

Male Rat Dominant Lethal, Oral

This study is the same as found in the Reproductive toxicology section. There it is listed as Dominant lethal, oral dosing and is found on page 87 of this document. The design of this study gives both genetic toxicity and reproductive toxicity information; thus, it is listed in both sections.

Male Rat Dominant Lethal, Inhalation

This study is the same as found in the Reproductive toxicology section. There it is listed as Dominant lethal, inhalation dosing and is found on page 90 of this document. The design of this study gives both genetic toxicity and reproductive toxicity information.

Reproductive Toxicology

Dominant Lethal, Oral Dosing

Type Dominant Lethal, Oral Dosing

Test Substance 1,3,5-Trioxane
CAS Number: 11 0-88-3

Method

- Guideline None, basically in accord with OECD 478
- G L P No
- Year 1984
- Species Rat
- Strain Albino, Wistar
- Route of administration Oral Gavage
- Doses 0, 850, 1700 mg/kg
- Sex Male
- Number of Animals/group 10
- Vehicle Water

Remarks	Field for	0	Age at Study Initiation	3 ½ to 4 months; weight 300 to 320 g
Test Conditions		◇	Doses	0,850 and 1700 mg/kg/day
		◇	Dosing	Males only
		◇	Dosing Schedule	Five days a week
		◇	Dosing Duration	Eight weeks
		◇	Mating Interval	Weekly
		◇	Mating Ratio	2: 1 females:males
		◇	Variations from OECD 478 Protocol Guideline	Guideline suggests that the number of males should be sufficient that 30-50 pregnant rats be evaluated at each time interval. In this study, 14 to 20 pregnant females were evaluated at each time; however, more time periods than typical were evaluated in this study. Current guideline suggests three dose levels; only two were used in this study.

Results

- Result No evidence of dominant lethal effect

- Dominant Lethal Rate

Dose	Dominant Lethals per Female (by week)							
	1	2	3	4	5	6	7	8
0 mg/kg	0.84	0.83	1.45	1.29	0.89	1.11	0.93	0.67
850 mg/kg	1.11	1.06	0.39	0.26	0.35	1.17	0.44	0.37
1700 mg/kg	0.44	1.56	0.85	0.61	0.85	0.95	0.71	0.88

Number of deaths at each dose level

Dose	<u>Mortality</u>
0 mg/kg	0/10
850 mg/kg	0/10
1700 mg/kg	0/10

Remarks Field for Results

◊ Time of death No deaths

0 Clinical Signs Behavior and appearance did not differ to any significant degree from controls

◊ Body Weights	Dose (mg/kg)	Body Weight gain (percent control gain)
	850	50%
1700	42%	

0 Organ Weights Absolute and relative weights of liver, kidney and spermatic vesicle were increased at both dose levels

0 Necropsy and Microscopic Findings Necropsy finding are not discussed. Histopathology was performed on the testes. Focal necrosis of the seminiferous epithelium was reported in 1/10 control, 3/10 850 mg/kg males and an unspecified number of high dose males. It was reported that the testicular lesions were bilateral in 3/10 high-dose males. Severity was dose dependent.

- 0** Target Organs Testes by histopathology but no change in fertility. Other organ weights were altered but there is no histopathology for confirmation. No changes in testes weight resulted from dosing.
- 0** Criteria for Dominant Lethal Effect Investigators used early resorptions as a measure of dominant lethal effect; however, neither implants per female, live fetuses per female nor preimplantation loss per female was affected by treatment.

Conclusions

Remarks field

Oral exposure of male rats to 859 or 1700 mg/kg Trioxane per day did not have any significant effects on the fertility rate as measured by the number of preimplantation losses, dead implants or live fetuses. Trioxane did not produce a dominant lethal effect by oral administration under these conditions

The study appears to have been well conducted and is similar to current OECD guideline. Based on reduction in body weight gains, organ changes and histopathology the high-dose level produced signs of toxicity indicating the study is valid regarding achieving a systemically toxic dose.

Data Quality

- Reliability

Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards. Study design and reporting meets current EPA/OECD guidelines with minor exceptions.

References

Barański, B; Stetkiewicz, J; Czajkowska, T; Sitarek, K; Szymczak W. Mutagenic and gonadotoxic properties of trioxane and dioxolane. *Medycyna Pracy* 5: 245-255 (1984)

Other

These results are supported by an inhalation study that is reported in the same publication

Dominant Lethal, Inhalation Dosing**Type** **Dominant Lethal, Inhalation Dosing**

Test Substance **1,3,5-Trioxane**
 CAS Number: 110-88-3

Method

- Guideline None, basically in accord with OECD 478
- GLP No
- Year 1984
- Species Rat
- Strain Albino, Wistar
- Route of administration Inhalation
- Doses 0, 2500 mg/m³
- Sex Male
- Number of Animals/group 14
- Duration of Dosing 12 Months
- Vehicle Air
- Statistical Methods **Kruskal-Wallis** Test followed by non-parametric tests for groups with and without normal distribution.

Remarks Field for Test Conditions	0 Age at Study Initiation 3 ½ to 4 months; weight 300 to 320 g ◇ Doses 0 and 2,500 mg/m ³ ◇ Dosing Males only ◇ Dosing Schedule Five hours a day for five days a week ◇ Dosing Duration 12 Months ◇ Mating Interval At end of study for 1 week duration ◇ Mating Ratio 2: 1 femalesmales ◇ Variations from OECD 478 Protocol Guideline Guideline suggests that the number of males should be sufficient that 30-50 pregnant rats be evaluated at each time interval. In this study, 20 to 22 pregnant females were evaluated. Current guideline suggests three dose levels; only one was used in this study
-----------------------------------	---

0 Other

Concentration of test material measured by chromatography using a published procedure.

2500 mg/m³ is equal to 580 mg/kg/day assuming a 270 ml/min minute volume, a 350 gram rat and 100% absorption.

Results

- Dominant Lethal Rate

Dose	Effect (per female)					
Trioxane (mg/m ³)	Number Pregnant	Live Fetuses	Dead Implants	Number Implants	Corpora Lutea	Preimplant Loss
0	20	11.4	0.6	12	13.7	1.8
2500	22	12.2	0.64	12.8	14.0	1.1

- Number of deaths at each dose level

Dose	<u>Mortality</u>
0 mg/m ³	0/14
2500 mg/m ³	0/14

Remarks Field for Results

- 0 Time of death: No deaths
- 0 Clinical Signs: Not reported
- 0 Body Weights: Not reported
- 0 Organ Weights: Not reported
- 0 Necropsy and Microscopic Findings: Testicular histopathologic effects were similar in control and treated group with regard to leydig cell pathology. Seminiferous tubule pathology was not reported for rats exposed to Trioxane by inhalation.
- 0 Target Organs: None reported

Conclusions

Remarks field

Prolonged exposure of male rats to 2500 mg/m^3 Trioxane by inhalation did not have any significant effects on the fertility rate of these animals, with regard to the measured parameters which were the number of pregnant females, the number of females mating with males, the average litter size, average implantation numbers, pre-implantation losses, and corpus **lutea**. Trioxane did not produce a dominant lethal effect by inhalation at this concentration.

Results are consistent with other data for the material. Since investigation of adverse testicular effects was a primary consideration of the study, and adverse effects on the testes were not reported, it is likely that 2500 mg/m^3 represents a NOAEL for testicular histopathology and for functional effects.

Data Quality

- Reliability

Klimisch Code 2. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards. Study design and reporting meets current EPA/OECD guidelines with minor exceptions.

References

Baranski, B; Stetkiewicz, J. Evaluation of Mutagenic and Gonadotoxic Properties of Trioxane and Dioxolane. *Medycyna Pracy*:35 245-255 (1984)

Other

These results are supported by an oral study reported in the same publication.

Repeated Dose Toxicity, 7-Week Oral with Estrous Cycle Determination**Type** Repeated Dose Toxicity, 7-week Oral**Test Substance** Trioxane
CAS Number: 110-88-3**Method**

- Guideline None
- GLP No
- Year 1990
- Species Rat
- Strain **Wistar/Nofer** Institute Breeding Colony
- Route of administration Oral gavage
- Duration of Test Seven weeks dosing, 13 weeks total
- Doses 0, 190,580, or 1160 **mg/kg/day**
- Sex Female
- Exposure Period Once daily
- Frequency of Treatment Five days a week
Seven weeks
- Number of Animals/group Ten or twelve
- Control Group and Treatment Twelve animals dosed with vehicle
- Vehicle Distilled water
- Post-Exposure Observation Period 5 weeks
- Statistical Methods One-way or two-way analysis of variance and Scheffe's test for multiple comparisons.

Remarks Field for 0 Age at study initiation Four **months**

Test Conditions 0 Number of animals per Sex per dose • Ten per group at 190 and 580 **mg/kg**
 ▪ Twelve animals in the group at 1160 mg/kg

0 Satellite groups None

0 Housing Not reported

0 Clinical observations performed and frequency ▪ Body weights were measured weekly.
 . General health and behavior, interval not reported
 ▪ Vaginal smears for the first 14 days of exposure, for the 14 days of the sixth and seventh week of exposure, and for the 14 days of the fourth and fifth week after ending exposure (11th and 1 2th weeks of study).

0 Terminal observation Body weights, general health

0 Histopathology None reported

Results

- NOAEL 0 580 **mg/kg/day** for estrous cycle effects
 0 Not found for body weight gain
- . LOAEL 0 1160 **mg/kg/day** for effects on estrous cycle
 0 190 **mg/kg/day** for body weight gain
- Mortality No deaths reported.

- Toxic Response

Dose	Effects
190 mg/kg	Significant reduction in body weight gain only at end of dosing period
580 mg/kg	Significant reduction in body weight gain only at week 6, 7 and 8 of study.

1160 mg/kg	<p>Significant decrease in body weight gain reported at weeks 4 to 8, with recovery of body weight gain within two weeks after cessation of dosing.</p> <p>Statistically significant increase in the length of the estrous cycle after 6 to 7 weeks of exposure. Length of estrous cycle returned to normal after dosing was stopped.</p> <p>Behavioral changes were noted in this group it is stated in the publication: “The animals from the 1.16 gm/kg group at the end of the exposure, exhibited changes in appearance and behavior. They sat curled up in the cage corner with ruffled hair coats, they squealed when taken to hand by the experimenter and had sanguineous discharge from the nose. After cessation of exposure these disorders disappeared during the subsequent two weeks and their body weight leveled up to that of the control animals”</p>
------------	--

Remarks Field for Results



Body Weights

0 Provided as weekly bar chart in publication

		Cycle Length (days)			
0	Estrous Cycle Length	Dose (mg/kg)	Week 1-2	Week 6-7	Week 11-12
		0	4.6 ±1.4	4.3±0.4	4.8±1.2
	190	4.3±0.5	4.4±0.9	5.0±1.6	
	580	5.1±1.6	5.4±1.2	4.5±0.6	
	1160	5.2±2.1	6.1±2.5*	5.1±2.8	

0 Necropsy Findings

None Reported

Conclusions

Remarks field

- 0 The authors conclude: “Based on the findings of this study, it seems unlikely that the occupational exposure of women to Trioxane at concentrations not inducing systemic intoxication, can produce alteration in their ovarian functions”
- 0 The study design and conduct appear sound and the systemic toxicity produced is similar to that found in other studies on this material.

Data Quality

- Reliability

Klimisch Code 2 Reliable with restrictions. Study design, conduct and reporting are considered reliable to address the test endpoint although not conducted in accord with GLP standards. The similarity of the results with supporting studies confirms reliability.

References

The effect of oral exposure to trioxane on the oestrous cycle in rats.
Sitarek K; **Barański** B. *Pol J Occup Med*; VOL 3,1990, P209-13

Other

This study is supported by

- ◇ A 2%day gavage study conducted by Hoechst AG in male and female rats where a low **degree** of systemic toxicity was found at 1000 **mg/kg** after 28 days exposure and 200 **mg/kg** was found to be a NOEL.'
- ◇ A 4 or 7-month long oral gavage study in male rats. In this study groups of 8-10 rats were administered Trioxane as a water solution, 5 days a week. The high dose was 850 **mg/kg** and the duration of dosing was 4 months. Doses of 213 or 106 **mg/kg** were administered for 7 months. Body weight gain was similar for all dosed groups as compared to controls. The 850 **mg/kg** (four month duration for this group) did not affect body weigh gain after 4 months of administration. No specific organ effects were reported and it was concluded that the mortality rate did not show accumulated toxic effects of Trioxane. Examination of the limited data presented suggest that 213 **mg/kg** was (or is near) a NOAEL for Trioxane after 7-months of administration.'

References for supporting studies

1. Trioxane. Subakute **orale** Toxizitat (28 Application in 29 **Tagen**) an **SPF**-Wistar-Ratten. Hoechst AG Central Toxicology Report number 90.0513 22 May 1990
2. Experimental studies of the toxic effects of **1,3,5-trioxane** and **1,3-dioxolane**. II. Cumulation of toxic effect. Czajkowska T; Krysiak B *Med Pr*; VOL 38, 1987, P244-9

Developmental Toxicology

Developmental Toxicology, Oral

Type Developmental Toxicology, Oral

Test Substance Trioxane
99.77 % Trioxane as a 20% aqueous solution
Stability data provided and dosage solutions analyzed

Method

- **Guideline** OECD 414 "Teratology" dated 12 May 1981
EC 88/302/EWG
US EPA OPPTS Series 83-3 "Teratology Study" November 1984
Japanese MAFF 4200, 28 January 1985
- **GLP** Yes
- **Year** 1998
- **Species** Rat
- **Strain** Wistar, Hoe: WISKf(SPF71)
- **Route of administration** Oral gavage, as 20% solution in water
- **Doses** 0, 100, 315, 1000 mg/kg
- **Sex** Female, pregnant
- **Exposure Period** Days 7 to 20 of pregnancy
- **Frequency of treatment** Daily
- **Control Group** Water only
- **Duration of test** 14 days
- **Statistical Methods** Two-sided comparison of high dose group followed by one-sided comparison of low-dose group. Fetal data analyzed using multivariate statistics to select the relevant dose groups followed by sequential comparisons with the high-dose group. Feed data analyzed using **Wilcoxon** rank sum test. Body weights of dams used univariate evaluation by t-tests. Fetal mean litter values were compared using the test statistic of Wilks for multivariate comparisons, and the t-test for univariate comparisons. The number of **corporea** leuta, implantation sites and live fetuses, and quotas of dead embryonic primordia undergoing resorption in the animals were analyzed using one-sided **Wilcoxon** tests. Fisher's Exact Test was used to analyze findings at autopsy, body cross-section and skeletal examination; separate analyses were conducted for the individual fetal data and for the litter data at significance levels of 5% and 1%.

Remarks Field for Test Conditions	0	Age at Study Initiation	X-10 weeks at start of in-house breeding procedure
	0	Number of animals per group	23 mated females per dose group
	0	Vehicle	Deionized water
	0	Clinical Observation Performed and Frequency	Behavior and health observed twice daily. Body weights determined on days 1,4, 7, 14, 17, 19 and 2 1 of gestation. Food consumption determined between days 1-4, 4-7, 7-10, 10-14, 14-17, 17-19, and 19-21 of gestation.
	0	Mating Procedures	Virgin females in pre-estrus or estrus phase were mated 1: 1 with males overnight. After detection of sperm in vaginal smears, females were housed individually and presumed pregnant. The day of sperm detection was defined as day-1 of gestation.
	0	Maternal Parameters Assessed During Study	Body weight, feed consumption and clinical signs, corpora lutea , implantations, uterus weight
	0	Fetal Parameters Assessed During Study	Litter size, placental weight, gross malformations, fetal crown-rump length, fetal body weight, sex ratio, body cross sections, skeletal examination
	0	Organs Examined at Necropsy	Full list not given, findings included - ovary, cervix, kidney and uterus
	0	Dose Selection	A dose-range-finding study was conducted using groups of 4 mated Wistar rats dosed at 500 or 1000 mg/kg per day from day 7 - 20 of pregnancy and were killed on day 21. The uterus was opened and the number of live and dead fetuses and the number of conceptuses undergoing resorption were determined. The fetuses were examined for gross major anomalies. Fetal body weight and crown-rump length were recorded. Two animals from the 1000 mg/kg group showed slight loss of body weight from day 7 - 10 and exhibited increased numbers of retarded fetuses. No compound-related effects were observed in the other animals. Based on these results the dose levels of 0, 100, 3 15 and 1000 mg/kg body weight per day were selected for the present study. Testing of dose levels greater than 1000 mg/kg body weight is not necessary according to current OECD Guidelines.

Results

- NOAEL & LOEL for Maternal Toxicity
NOAEL = not established
LOEL = 100 mg/kg/day body weight gain
- NOAEL & LOEL for Developmental Toxicity
NOAEL = 100 mg/kg/day
LOEL = 3 15 mg/kg/day retarded ossification
- Actual Doses Received
0, 100, 315, 1000 mg/kg/day
- Maternal data
Body weight gain and food consumption was significantly decreased in **high**-dose animals from day-10 to the end of the study. Mean corrected body weight gain (day-2 1 body weight minus gravid uterus weight) was decreased with statistically significance $p \leq 0.05$ for all dosed groups. Control 33.6 g., low dose 27.13 g., intermediate dose 28.96 g, and high dose 18.88 g. Although the dose dependency is not clear, there appears to have been reduction in maternal weight gains at all dose levels indicating some maternal toxicity at even the low dose. No compound-related effects were observed at necropsy. Gravid uterus weights were comparable in all groups. One animal in the high dose and one animal in the low dose group showed only implantation sites at **cesarean** section. Based on historical data and lack of dose-dependency, this was not considered compound related. Five dead fetuses were observed in the high-dose group and this was statistically significant compared to the control group. A compound-related effect cannot be ruled out for fetal death at the high dose.
- Fetal data
 - 0 Cesarean Data
Litter size was comparable in all groups. In the high-dose group, fetal body weights and crown-rump length were decreased, and placental weights were increased. Sex ratio was unaffected by treatment.
 - 0 External and visceral effects
The incidence of retarded fetuses was **6/123**, **6/117**, **8/139** and **29/131** for control to high dose respectively. This was statistically significant in the high-dose group compared to control. Two high-dose fetuses showed tail aplasia that may be compound related. Other findings were within the historical control range.
 - 0 Major skeletal defects
Two high-dose fetuses showed tail aplasia, these same two also showed aplasia of the sacral vertebral arch, sacral vertebrae centers and first and second **caudal** vertebrae centers.

One fetus from the low-dose showed fusion of the exoccipital bone with the first cervical vertebrae and dysplasia of the exoccipital bone. One fetus from the high-dose group showed dysplasia of the exoccipital bone. The incidence of these defects was within the historical control range of the rat strain.

• Fetal data

◊ Minor skeletal defects

Effect	Dose Group			
	0	100	315	1000
Longitudinally displaced, fused or fragmented sternbrae	12/123	3/117	1/139	17/131*
Wavy and or thickened ribs	12/123	14/117	56/139*	84/131*
Bent or shortened scapula	0/123	0/117	2/139	12/131*
Bent, shortened or dysplastic humerus	0/123	0/117	1/139	12/131*
Bent or shortened radius	0/123	0/117	0/139	5/131*
Fragmented thoracic vertebral centers	0/123	0/117	2/139	4/131

* = statistically significant

◊ Retardations

Effect	Dose Group			
	0	100	315	1000
Slight or non-ossification of skull bone	42/123	50/117	74/139*	80/131*
Weakly or non-ossified cervical vertebral arch	0/123	0/117	3/139	13/131*
Weakly ossified lumbar vertebral arch	0/123	0/117	2/139	13/131*
Weakly or non-ossified sacral vertebral arch	3/123	3/117	2/139	14/131*
Ossification of less than two caudal vertebral centers	26/123	50/117*	85/139*	98/131*
Weakly ossified ribs	0/123	1/117	3/139	9/131*
Weakly ossified metacarpal 2	0/123	0/117	0/139	5/131*
Non-ossified metacarpal 5	58/123	61/117	92/139*	86/131*
Non-ossified metatarsal 5	3/123	2/117	1/139	11/131*
Weakly or non-ossified sternbrae	18/123	42/117*	45/139*	70/131*
Non-ossified phalanx 3 of the 1 st to 5 th hindpaw toe	1/123	7/117*	3/139	18/131*

* = Statistically significant

◊ External and visceral effects

Retarded fetuses observed at body cross section

0	100	315	1000
---	-----	-----	------

0/116	2/109	1/125	17/121*
-------	-------	-------	---------

* = Statistically significant

- Statistical Results Fetal data statistical determination by Fisher's Exact test (* indicates $p < 0.05$)

Remarks Field for Results Low-dose retardations were not considered compound related by the study director due to lack of dose-response or historical range.

Parameter	Dose Group			
	<u>0</u>	<u>100</u>	<u>315</u>	<u>1000</u>
Pregnancies	20/23	21/23	21/23	21/23
Females with abortion	0	1	0	1
Corpora lueta (total)	280	287	316	298
Implantations (total)	246	245	284	276
Pre-implantation loss, mean percent	12.78	14.10	10.08	7.35
Post-implantation loss, mean percent	2.62	7.65	7.22	8.36
Live fetuses (total)	239	226	264	252
Total intrauterine deaths	7	19	20	24
Males %	48.1	59.7	55.7	50.8
Body weight (mean grams)	3.3	3.3	3.2	2.8*
Crown-rump length (mean mm)	35.1	35.0	34.5	32.8*
Placental weight (mean g.)	0.46	0.47	0.49	0.52
Uterus weight (mean g.)	60.40	57.57	62.57	59.47

Conclusions

Remarks field

Based on the results of this study, Trioxane is not considered a specific developmental toxin. The developmental NOEL was found to be 100 mg/kg/day while a maternal NOEL was not defined since corrected dam body weight gain was affected in all treated groups.

Neither deaths nor clinical signs were observed in any of the animals.

Gravid uterus weights, litter size and fetal sex ratios were not affected. Fetal body weights and crown-rump lengths were decreased in the high-dose group whereas placental weights were increased. Early resorption was not affected by Trioxane administration. Five dead fetuses were observed in five litters at the high dose.

In the high-dose group, two cases of major defects were observed. The incidence of fetuses with minor defects was increased. Retarded ossification was observed in **numerous** bones.

In the mid-dose group, the incidence of minor defects was increased and retarded ossification of individual skull bones and **caudal** vertebral centers were observed.

In the low-dose group, no compound-related effects were observed by morphological observation.

Data Quality

- Reliability Klimisch Code 1. . Reliable without restriction, study meets GLP standards and/or most requirements

References

Hofmann Th. Trioxan, Rat Oral Developmental Toxicity (Teratogenicity) study. Hoechst Marion Roussel, HMR Deutschland **GmbH**, Global Preclinical Development, Drug Safety, Study Number 97.079 1 (1998).

Other

Supporting data comes from previous published developmental toxicity studies. In a developmental toxicity study of Trioxane, pregnant Wistar rats were given 770, 1550, or 3870 **mg/kg** Trioxane orally every other day from day 8 to 20 of gestation. Other rats included in this protocol were given 190 **mg/kg** Trioxane or 20 **mg/kg** Formaldehyde daily during gestational days 8 to 20. Dams were killed on gestational day 21 and necropsied. Placentas were examined for histopathological changes. The numbers of resorptions and live and dead fetuses were recorded. The fetuses were weighed and examined for malformations. Trioxane did not cause any deaths, but caused significant maternal toxicity as evidenced by decreases in body weight gain, feed intake, absolute liver and placenta weights and increases in relative kidney and adrenal weights. The 3890 **mg/kg** dose was associated with hydropic liver in dams. Trioxane induced dose dependent increases in the number of resorptions and decreases in the number of live fetuses and fetal body weight and length. The 770 to 3870 **mg/kg** every-other-day doses induced malformations in the brain, kidneys, and skeleton. The 770-**mg/kg** dose was associated with developmental effects without reported maternal toxicity. The daily 190-**mg/kg** dose did not cause any developmental effects. Trioxane administration was associated with fibrin deposits, inflammatory infiltration, and focal necrosis in the placentas. The authors conclude that Trioxane at sufficiently high concentrations causes fetal lethality, retards fetal development, and induces congenital malformations'.

A study designed to examine the postnatal effects of Trioxane using the every-other-day dosing regime was conducted at 190,580 or 1160 **mg/kg/day** from day 2 to 20 of gestation. In this study, the high dose was associated with high post-natal mortality and reduced maternal instincts. The mid-dose was reported to be associated with reduced active-avoidance acquisition of offspring at 5 months. The low-dose, 190 **mg/kg/day**, was reported to be a NOEL for behavioral developmental effects in the pups'.

References for supporting studies

1. Sitarek, **K**, Baranski, **B**, Stetkiewicz, **J**, Stetkiewicz, **I**, Teratogenicity, Fetal and Placental Toxicity of 1,3,5-Trioxane Administered to Pregnant Female Rats. Polish Journal of Occupational Medicine 1:5 1-6 1, (1988).
2. Sitarek, **K**, **Barański**, **B**. Effects of maternal exposure to trioxane on postnatal development in rats. Pol J Occup Med 3:285-92 (1990)