

**The Flavor and Fragrance High Production Volume Consortia  
Robust Summaries for Terpenoid Tertiary Alcohols and Related  
Esters**

**FFHPVC Terpene Consortium Registration Number**

The evaluation of the quality of the following data uses a systematic approach described by Klimisch [Klimisch et al., 1996]. Based on criteria relating to international testing standards for categorizing data reliability, four reliability categories have been established. The following categories are:

- Reliability code 1.      Reliable without restrictions
- Reliability code 2.      Reliable with restrictions
- Reliability code 3.      Not reliable
- Reliability code 4.      Not assignable

2001 JAN 29 AM 7:20

RECEIVED  
OPT M10

**1 Chemical and Physical Properties**

**1.1 Melting Point**

<b>Substance Name</b>	Linalool (3,7-Dimethyl-1,6-octadien-1-ol)
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Not specified
<b>Melting Point</b>	<20 °C
<b>Conclusion Remarks</b>	Consistent with calculated weighted MP determination (SRC)
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Givaudan Roure (1991a) Determination of ready biodegradability of linalool. Unpublished report.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Method/guideline</b>	Mean or weighted MP
<b>Melting Point</b>	-11.8 °C
<b>Conclusion Remarks</b>	Calculated melting point

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.  
**References** Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	Mean or weighted MP
<b>Melting Point</b>	-13.4 °C
<b>Conclusion Remarks</b>	Calculated melting point
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	Mean or weighted MP
<b>Melting Point</b>	-13.4 °C
<b>Conclusion Remarks</b>	Calculated melting point
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Mean or weighted MP
<b>Melting Point</b>	-2.1 °C
<b>Conclusion Remarks</b>	Calculated melting point
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5

<b>Method/guideline</b>	Mean or weighted MP
<b>Melting Point</b>	12.4 °C
<b>Conclusion Remarks</b>	Calculated melting point, measured solidification temperature for <i>l</i> isomer = 36.4 °C and <i>d</i> isomer 31 .0 °C (Merck, 1997)
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Mean or weighted MP
<b>Melting Point</b>	21.4 °C
<b>Conclusion Remarks</b>	Calculated melting point
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	2-Pinanol
<b>CAS No.</b>	473-54-1
<b>Method/guideline</b>	Mean or weighted MP
<b>Melting Point</b>	22.9 °C
<b>Conclusion Remarks</b>	Calculated melting point
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

## 1.2 Boiling Point

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Boiling Point</b>	205 °C
<b>Pressure</b>	760

**Pressure Unit** mm Hg  
**Data Qualities Reliabilities** Reliability code 1. Reliable without restrictions.  
**References** Fragrance Materials Association (FMA)

<b>Substance Name</b>	2-Pinanol
<b>CAS No.</b>	473-54-1
<b>Boiling Point</b>	205 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Boiling Point</b>	205206 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Boiling Point</b>	200-220 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
-----------------------	---

<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	Using differential scanning calorimetry, ASTM E 537-86 and Method A2 of Commission Directive 92/69/EEC
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Boiling Point</b>	492 +/- 0.5K (219 °C)
<b>Pressure</b>	101.82
<b>Pressure Unit</b>	kPa
<b>Remarks for Results</b>	Boiling Temperature: 492 +/- 0.5 K
<b>Conclusion Remarks</b>	The boiling temperature of plinyl acetate was determined to be 492 +/- 0.5K at a pressure of 101.82 kPa.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Reliabilities</b>	The study was performed in compliance with the UK GLP and these Principles are in accordance with GPL standard published as OECD Environment Monograph No. 45 (OCDE/GD(92)32).
<b>References</b>	Lumsden A. M. (1998) Plinyl Acetate: Determination of general physico-chemical properties. Safeparm Laboratories, Inc. SPL Project Number: 1044/001.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Boiling Point</b>	213 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Boiling Point</b>	80 °C
<b>Pressure</b>	213
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	<b>78-70-6</b>
<b>Boiling Point</b>	<b>198 °C</b>
<b>Pressure</b>	<b>760</b>
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	<b>78-70-6</b>
<b>Boiling Point</b>	<b>198 °C</b>
<b>Pressure</b>	<b>760</b>
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Food and Agricultural Organization of the United States (FAO) (1997) Compendium of food additive specifications addendum 5.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	<b>115-95-7</b>
<b>Boiling Point</b>	<b>213 °C</b>
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restriction.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Boiling Point</b>	230°C
<b>Pressure</b>	760

**Pressure Unit** mm Hg  
**Data Qualities Reliabilities** Reliability code 1. Reliable without restrictions,  
**References** Fragrance Materials Association (FMA)

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Boiling Point</b>	218 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Boiling Point</b>	206-207 °C
<b>Pressure</b>	731
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Merck Index (1997) Merck & Co., Inc. Publishers: Merck Research Laboratories, 12th edition, Whitehouse Station, NJ.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Boiling Point</b>	217 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Food and Agricultural Organization of the US (FAO) (1997) Compendium of food additive specifications addendum 5.

<b>Substance Name</b>	alpha-Terpinyl acetate
-----------------------	------------------------

<b>CAS No.</b>	80-26-2
<b>Boiling Point</b>	220 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Food and Agricultural Organization of the US (FAO) (1997) Compendium of food additive specifications addendum 5.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Boiling Point</b>	220 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Merck Index (1997) Merck & Co., Inc. Publishers: Merck Research Laboratories, 12th edition, Whitehouse Station, NJ.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3. 1. 1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Boiling Point</b>	196 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Lumsden A. M. (1998) Plinyl Acetate: Determination of general physico-chemical properties. Safepharm Laboratories, Inc. SPL Project Number: 1044/001.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Boiling Point</b>	198 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg

**Data Qualities Reliabilities** Reliability code 1. Reliable without restrictions.

**References** Merck Index (1997) Merck & Co., Inc. Publishers: Merck Research Laboratories, 12th edition, Whitehouse Station, NJ.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Stein and Brown Method
<b>Boiling Point</b>	204 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Conclusion Remarks</b>	Calculated
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	Stein and Brown Method
<b>Boiling Point</b>	191 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Conclusion Remarks</b>	Calculated
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Stein and Brown Method
<b>Boiling Point</b>	229 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Conclusion Remarks</b>	Calculated
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	Stein and Brown Method
<b>Boiling Point</b>	200.8 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Conclusion Remarks</b>	Calculated
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Method/guideline</b>	Stein and Brown Method
<b>Boiling Point</b>	214 °C
<b>Pressure</b>	760
<b>Pressure Unit</b>	mm Hg
<b>Conclusion Remarks</b>	Calculated
<b>References</b>	Syracuse Research Corporation (SRC)

### 1.3 Vapor Pressure

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.07 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions,
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.05 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	2-Pinanol
<b>CAS No.</b>	473-54-1
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.05 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.05 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.07 mm Hg

**Temperature** 20 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.

**References** Fragrance Materials Association (FMA)

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	Testing was conducted according to method A4 specified in Commission Directive 92/69/EEC (which constitutes Annex V of Council Directive 67/548/EEC).
<b>Remarks for Test Conditions</b>	The vapor pressure was determined using an isoteniscope system in which sample's vapor pressure was measured using mercury in a glass manometer. The temperature of the sample was regulated by use of a silicone oil bath.
<b>Vapor Pressure</b>	108 Pa = 0.81 mm Hg
<b>Temperature</b>	25 °C
<b>Conclusion Remarks</b>	The vapor pressure was determined to be 108 Pa at 25 C.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was performed in compliance with the UK Principles of GLP and these principles are in accordance with GLP standards published as OECD Environment Monograph No. 45 (OECD/GD(92)32).
<b>General Remarks</b>	The vapor pressure was measured over a range of temperature to enable extrapolation to 298.15 K
<b>References</b>	Tremain S. P. (1997). Plinyl Acetate: Determination of Vapor Pressure. Safepharm Laboratories Ltd. SPL Project No. 1044/003.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.05 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Dihydromyrcenol
-----------------------	-----------------

<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.12 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Calculated
<b>Vapor Pressure</b>	0.08 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Fragrance Materials Association (FMA)

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Measured
<b>Vapor Pressure</b>	0.05027 mm Hg
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Vuilleumeier C., Flament I. and P. Sauvegrain (1995) Headspace analysis study of evaporation rate of perfume ingredients applied onto skin. International Journal of Cosmetic Science, 17(1), 61-76.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Measured
<b>Vapor Pressure</b>	0.15 mm Hg
<b>Temperature</b>	20 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.

**References** Givaudan Roure (1991a) Determination of ready biodegradability of linalool. Unpublished report.

<b>Substance Name</b>	Linalyl acetate
-----------------------	-----------------

**CAS No.** 115-95-7

**Method/guideline** Calculated

**Vapor Pressure** 0.07 mm Hg

**Temperature** 20 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.

**References** Fragrance Materials Association (FMA)

<b>Substance Name</b>	Tetrahydrolinalool
-----------------------	--------------------

**CAS No.** 78-69-3

**Method/guideline** Calculated

**Vapor Pressure** 0.04 mm Hg

**Temperature** 20 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.

**References** Fragrance Materials Association (FMA)

<b>Substance Name</b>	alpha-Terpineol
-----------------------	-----------------

**CAS No.** 98-55-5

**Method/guideline** Calculated

**Vapor Pressure** 0.02 mm Hg

**Temperature** 20 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.

**References** Fragrance Materials Association (FMA)

<b>Substance Name</b>	alpha-Terpineol
-----------------------	-----------------

**CAS No.** 98-55-5

**Year** 1995

**Vapor Pressure** 0.04 mm Hg

**Temperature** 22-25 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.

**References** Misra G., Pavlostathis S.G., Li J., and Perdue E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>GLP</b>	NA
<b>Year</b>	1995
<b>Vapor Pressure</b>	0.16 mm Hg
<b>Temperature</b>	22-25 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Misra G., Pavlostathis S.G., Li J., and Perdue E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (data for component alcohol - linalol, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>GLP</b>	NA
<b>Year</b>	1995
<b>Vapor Pressure</b>	0.14 mm Hg
<b>Temperature</b>	22-25 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Misra G., Pavlostathis S.G., Li J., and Perdue E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

## 1.4 Octanol/Water Partition Coefficients

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The determination was carried out using the HPLC method, Method A8 of Commission Directive 92/69/EEC (which constitutes Annex V of Council Directive 67/548/EEC).
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Remarks for Test Conditions</b>	A preliminary assessment of the partition coefficient was made based on the approximate solubilities of the test material in n-octanol and water. This was performed by visual assessment.
<b>Log Pow</b>	4.09
<b>Temperature</b>	21.5 °C
<b>Conclusion Remarks</b>	The partition coefficient of Plinyl acetate was determined to be $1.23 \times 10^4$ , Log <sub>10</sub> Pow = 4.09
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was performed in compliance with the UK GLP and these Principles are in accordance with GLP standards published as OECD Environment Monograph No. 45 (OECD/GD(92)/32).
<b>General Remarks</b>	Substance having a log <sub>10</sub> Pow of greater than 3 are regard as having the potential of bioaccumulation in the environment.
<b>References</b>	Lumsden A. M. (1998) Plinyl Acetate: Determination of general physico-chemical properties. Safepharm Laboratories, Inc. SPL Project Number: 1044/001.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	3.46
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8

**Method/guideline** Calculated  
**Log Pow** 3.47  
**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.  
**References** Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	3.38
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions,
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Measured Rp-TLC
<b>Log Pow</b>	2.9
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Givaudan Roure (1991 b) Determination of the partition coefficient ( <b>octanol/water</b> ) of linalool. Unpublished report.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	3.46
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Measured

**Log Pow** 2.9  
**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.  
**References** Proctor & Gamble (1996) Submission to EPA. Unpublished report.

---

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	3.6
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	3.33
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Measured RP-TLC
<b>Log Pow</b>	4.3
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Reference</b>	Givaudan Roure (1996a) Partition coefficient n-octanol/water of alpha-terpineol acetate. Unpublished report.

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Calculated

**Log Pow** 4.34  
**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.  
**References** Syracuse Research Corporation (SRC)

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - pinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	2.87
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	2.98
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	2-Pinanol
<b>CAS No.</b>	473-54-1
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	2.98
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	2.85

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions,  
**References** Syracuse Research Corporation (SRC)

<b>Substance Name</b>	2-Pinanol hydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Method/guideline</b>	Calculated based on degradation to 2-pinanol
<b>Log Pow</b>	<b>2.85</b>
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>General Remarks</b>	Based on degradation to 2-Pinanol
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Method/guideline</b>	Calculated based on principal constituents
<b>Log Pow</b>	<b>2.9</b>
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>General Remarks</b>	Calculation based on principle components
<b>References</b>	Syracuse Research Corporation (SRC)

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	<b>78-70-6</b>
<b>Method/guideline</b>	Measured (Karickhoff and Brown method, 1979)
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1995
<b>Log Pow</b>	<b>2.97</b>
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Misra G., Pavlostathis S.G., Li J., and Perdue E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Measured (Karickhoff and Brown method, 1979)
<b>GLP</b>	<b>NG</b>
<b>Year</b>	<b>1995</b>
<b>Log Pow</b>	2.98
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Misra G., Pavlostathis S.G., Li J., and Perdue E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl alcohol)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	Measured (Karickhoff and Brown method, 1979)
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1995
<b>Log Pow</b>	2.87
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Misra G., Pavlostathis S.G., Li J., and Perdue E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

## 1.5 Water Solubility

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The determination was carried out using the flask method, Method A6 of Commission Directive 92/69/EEC (which constitutes Annex V of Council Directive 67/548/EEC).
<b>GLP</b>	<b>Yes</b>
<b>Year</b>	1997

**Remarks for Test Conditions** An aliquot (0.2674) of test material was diluted with glass double distilled waster (100 ml). After shaking at 30 °C for approximately 25-1/2 hours and standing at 20 °C for approximately 24 1/2 hours, solution was filtered and analyzed.

**Value (mg/L) at Temperature** 60.5 mg/l at 20.0 +/- 0.5 °C

**pH value and Concentration** 6.3

**Conclusion Remarks** The water solubility of Plinyl Acetate was determined to be 6.05 X 10E-2 g/l of solution at 20.0 +/- 0.5 °C.

**Data Qualities Reliabilities** Reliability code 1. Reliable without restrictions.

**Data Reliabilities Remarks** The study was performed in compliance with the UK GLP and these Principles are in accordance with GLP standard published as OECD Environment Monograph No. 45 (OECD/GD(92)32).

**References** Lumsden A. M. (1998) Plinyl Acetate: Determination of general physico-chemical properties. Safepharm Laboratories, Inc. SPL Project Number: 1044/001.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>GLP</b>	NA
<b>Value (mg/L) at Temperature</b>	1450 mg/l at 20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions,
<b>References</b>	Givaudan Roure (1991a) Determination of ready biodegradability of linalool. Unpublished report.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>GLP</b>	NA
<b>Value (mg/L) at Temperature</b>	30 mg/l (no temperature assigned)
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Flavor and Extract Manufacturers' Association (FEMA) (1998) Report on the metabolism of linalyl acetate in rat tissue & intestinal homogenates. Unpublished report.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>GLP</b>	NA
<b>Value (mg/L) at Temperature</b>	140 mg/L @ 20 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions.

**References** Givaudan Roure (1991c) Determination of the ready biodegradability of linalyl acetate synthetic. Unpublished report.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>GLP</b>	NA
<b>Value (mg/L) at Temperature</b>	700 mg/l (no temperature assigned)
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Bush Boake Allen (BBA) (1990). Unpublished report to RIFM.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>GLP</b>	NA
<b>Value (mg/L) at Temperature</b>	29,000 mg/l water at 100 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Merck Index (1997) Merck & Co., inc. Publishers: Merck Research Laboratories, 12th edition, Whitehouse Station, NJ.

<b>Substance Name</b>	Pine oil (62.7% alpha-terpineol, 83.4% total terpene alcohol content)
<b>CAS No.</b>	8002-09-3
<b>GLP</b>	NA
<b>Value (mg/L) at Temperature</b>	Insoluble in water
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Graves, W. C. and Swigert, J. P (1994a). Pine Oil: 96-Hours Flow-through Acute Toxicity Test with Bluegill ( <i>Lepomis macrochirus</i> ). Wildlife International Ltd Report to The CSMA Pine Oil Joint Venture. Lab Project ID/Study Number 274A-101.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>GLP</b>	NA
<b>Year</b>	1995

**Remarks for Test Conditions** Water solubility determined at two temperature ranges.

**Value (mg/L) at Temperature** 560 mg/l at 4-8 °C and 867 mg/l at 22-25 °C

**Data Qualities Reliabilities** Reliability code 2. Reliable with restrictions,

**References** Misra G., Pavlostathis S.G., Li J., and **Perdue** E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>GLP</b>	NA
<b>Year</b>	1995
<b>Remarks for Test Conditions</b>	Water solubility determined at two temperature ranges.
<b>Value (mg/L) at Temperature</b>	341 mg/l at 4-8 °C and 716 mg/l at 22-25 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Misra G., Pavlostathis S.G., Li J., and <b>Perdue</b> E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>GLP</b>	NA
<b>Year</b>	1995
<b>Remarks for Test Conditions</b>	Water solubility determined at two temperature ranges.
<b>Value (mg/L) at Temperature</b>	818 mg/l at 4-8 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Misra G., Pavlostathis S.G., Li J., and <b>Perdue</b> E. M. (1995) Characterization and Biodegradation of Selected Monoterpenes. 68th Annual Conference & Exposition, Water Environment Federation; Miami Beach, FL.

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1

**Method/guideline** Calculated at log **Kow=2.85** (ESPKOW)  
**Value (mg/L) at temperature** **863 mg/L** at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	alpha-Terpineol
-----------------------	-----------------

**CAS No.** 98-55-5  
**Method/guideline** Calculated at log **Kow=2.98** (ESPKOW)  
**Value (mg/L) at temperature** **670 mg/L** at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	alpha-Terpineol acetate
-----------------------	-------------------------

**CAS No.** 98-55-5  
**Method/guideline** Calculated at log **Kow=4.34** (ESPKOW)  
**Value (mg/L) at temperature** **9.03 mg/L** at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	Dihydromyrcenol
-----------------------	-----------------

**CAS No.** 18479-58-8  
**Method/guideline** Calculated at log **Kow=3.47** (ESPKOW)  
**Value (mg/L) at temperature** **252 mg/L** at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	Linalool
-----------------------	----------

**CAS No.** 78-70-6  
**Method/guideline** Calculated at log **Kow=2.97** (ESPKOW)

**Value (mg/L) at temperature** 684 mg/L at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	Linalyl acetate
-----------------------	-----------------

**CAS No.** 115-95-7  
**Method/guideline** Calculated at log  $K_{ow}$ =4.39 (ESPKOW)  
**Value (mg/L) at temperature** 8.20 mg/L at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	Myrcenol
-----------------------	----------

**CAS No.** 543-39-5  
**Method/guideline** Calculated at log  $K_{ow}$ =3.46 (ESPKOW)  
**Value (mg/L) at temperature** 261 mg/L at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	Pinane hydroperoxide
-----------------------	----------------------

**CAS No.** 28324-52-9  
**Method/guideline** Calculated at log  $K_{ow}$ =3.02 (ESPKOW)  
**Value (mg/L) at temperature** 160 mg/L at 25 °C  
**Remarks for Data Reliability** Comparable to guidelines/standards  
**References** ESPOW

<b>Substance Name</b>	Tetrahydrolinalool
-----------------------	--------------------

**CAS No.** 78-69-3  
**Method/guideline** Calculated at log  $K_{ow}$ =3.60 (ESPKOW)  
**Value (mg/L) at temperature** 189 mg/L at 25 °C

**Remarks for Data Reliability**    Comparable to guidelines/standards

**References**                            ESPOW

## 2 Environmental Fate and Pathways

### 2.1 Photodegradation

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t<sub>1/2</sub></b>	9.08 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t<sub>1/2</sub></b>	1.24 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t<sub>1/2</sub></b>	1.35 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t<sub>1/2</sub></b>	3.45 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t<sub>1/2</sub></b>	1.07 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t<sub>1/2</sub></b>	1.10 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	Calculation

<b>Test Type</b>	AOPWIN
<b>Half-life t1/2</b>	1.14 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	Pinane hydroperoxide
CAS No.	28324-52-9
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t1/2</b>	6.80 hrs (NOTE: these data are for 2-Pinanol)
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

<b>Substance Name</b>	Tetrahydrolinalool
CAS No.	78-69-3
<b>Method/guideline</b>	Calculation
<b>Test Type</b>	AOPWIN
<b>Half-life t1/2</b>	8.99 hrs
<b>Data Reliabilities Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>References</b>	AOPWIN

## 2.2 Stability in Water

<b>Substance Name</b>	Linalyl acetate
CAS No.	115-95-7
<b>Method/guideline</b>	Hydrolysis in artificial gastric juice
<b>GLP</b>	No

<b>Year</b>	1999
<b>Duration (days)</b>	3 hrs
<b>Analytical procedures</b>	At 37 C, 50 ml of artificial gastric juice adjusted to pH 1.2 or 7.0 was incubated with 50uM linalyl acetate in DMSO with DMSO 0.2%(v/v) of the final mixture. Experiments in acidic gastric juice were performed with and without peptic enzymes. Sample were withdrawn at intervals beginning at 0.1 minutes and ending at 180 minutes and analyzed by Chromopack CP-9003 gas chromatograph.
<b>Temperature</b>	37 °C
<b>Rate constant</b>	Rate constant k=0.006 min <sup>-1</sup> at pH=7.0; rate constant k=>3.5 min <sup>-1</sup> at pH=1.2 without enzyme; rate constant k=>3.5 min <sup>-1</sup> at pH=1.2 with enzyme.
<b>Halflife t1/2</b>	t(1/2)=112.3 min. at pH=7.0: t(1/2)=<2.5 min. at pH=1.2 ,no enzyme; t(1/2)=<2.5 min. at pH=1.2 ,with enzyme
<b>Breakdown products</b>	Linalool and acetic acid formed under all conditions. At different pH with or without enzyme, linalool reacted to form alpha-terpineol in approximately equal amount. Small amounts of geraniol also formed.
<b>Conclusion remarks</b>	Linalyl acetate readily hydrolyzes in neutral and acidic media. The hydrolysis product linalool subsequent partially rearranges to alpha-terpineol and to a lesser extent, geraniol.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Remarks for Reliabilities</b>	Experiments performed with all necessary internal and external standards. Study evaluated the hydrolysis of at least 20 substrates.
<b>References</b>	Buck N. R. and Renwick A.G. (1999) Hydrolysis of tertiary terpenoid esters in artificial gastric juice and intestinal fluid. University of Southampton. Private Communication to FEMA. Unpublished Report.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Hydrolysis in artificial intestinal fluid
<b>GLP</b>	<b>No</b>
<b>Year</b>	1999
<b>Duration (days)</b>	3 hrs
<b>Analytical procedures</b>	At 37 C, 50 ml of artificial intestinal fluid adjusted to pH 7.5 was incubated with 50uM linalyl acetate in DMSO with DMSO 0.2%(v/v) of the final mixture. Experiments in intestinal fluid were performed with and without pancreatin enzymes. Sample were withdrawn at intervals beginning at 0.1 minutes and ending at 180 minutes and analyzed by Chromopack CP gas chromatography.
<b>Temperature</b>	37 °C

<b>Rate constant</b>	Rate constant $k=0.007$ min <sup>-1</sup> at pH=7.5 without enzyme; rate constant $k=0.008$ min <sup>-1</sup> at pH=7.5 with enzyme;
<b>Halflife t<sub>1/2</sub></b>	t(1/2)=96.8 min. at pH=7.5 ,no pancreatin enzyme; t(1/2)=83.1 min. at pH=7.5 ,with enzyme.
<b>Breakdown products</b>	Linalool and acetic acid formed under all conditions. At different pH with or without enzyme, linalool reacted to form alpha-terpineol and smaller amounts of geraniol and nerol.
<b>Conclusion remarks</b>	Linalyl acetate readily hydrolyzes in neutral media. The hydrolysis product linalool subsequent partially rearranges to alpha-terpineol and to a lesser extent, geraniol and geraniol.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Remarks for Reliabilities</b>	Experiments performed with all necessary internal and external standards. Study evaluated the hydrolysis of at least 20 substrates.
<b>References</b>	Buck N. R. and Renwick A.G. (1999) Hydrolysis of tertiary terpenoid esters in artificial gastric juice and intestinal fluid. University of Southampton. Private Communication to FEMA. Unpublished Report.

<b>Substance Name</b>	<i>alpha</i> Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Hydrolysis in artificial gastric juice
<b>GLP</b>	No
<b>Year</b>	1999
<b>Duration (days)</b>	3hrs
<b>Analytical procedures</b>	At 37 C, 50 ml of artificial gastric juice adjusted to pH 1.2 or 7.0 was incubated with 50uM alpha-terpineol acetate in DMSO with DMSO 0.2%(v/v) of the final mixture. Experiments in acidic gastric juice were performed with and without peptic enzymes. Sample were withdrawn at intervals beginning at 0.1 minutes and ending at 180 minutes and analyzed by Chromopack CP-9003 gas chromatograph.
<b>Temperature</b>	37 °C
<b>Rate constant</b>	Rate constant $k=0.00081$ min <sup>-1</sup> at pH=7.0; rate constant $k=0.038$ min <sup>-1</sup> at pH=1.2 without enzyme; rate constant $k=0.042$ min <sup>-1</sup> at pH=1.2 with enzyme.
<b>Halflife t<sub>1/2</sub></b>	t(1/2)=859.7 min. at pH=7.0; t(1/2)=18.3 min. at pH=1.2 ,no enzyme; t(1/2)=16.4 min. at pH=1.2 ,with enzyme
<b>Breakdown products</b>	alpha-Terpineol acetate is hydrolyzed to alpha-terpineol and acetic acid.
<b>Conclusion remarks</b>	alpha-Terpineol readily hydrolyzes to alpha-terpineol and acetic acid in acidic and neutral media.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Remarks for Reliabilities</b>	Experiments performed with all necessary internal and external standards. Study evaluated the hydrolysis of at least 20 substrates.

**References**

Buck N. R. and Renwick A.G. (1999) Hydrolysis of tertiary terpenoid esters in artificial gastric juice and intestinal fluid. University of Southampton. Private Communication to FEMA. Unpublished Report.

<b>Substance Name</b>	alpha Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Hydrolysis in artificial intestinal fluid
<b>GLP</b>	No
<b>Year</b>	1999
<b>Duration (days)</b>	3hrs
<b>Analytical procedures</b>	At 37 C, 50 ml of artificial intestinal fluid adjusted to pH 7.5 was incubated with 50uM alpha-terpineol acetate in DMSO with DMSO 0.2%(v/v) of the final mixture. Experiments in intestinal fluid were performed with and without pancreatin enzymes. Sample were withdrawn at intervals beginning at 0.1 minutes and ending at 180 minutes and analyzed by Chromopack CP gas chromatograph.
<b>Temperature</b>	37 °C
<b>Rate constant</b>	Rate constant $k=0.002 \text{ min}^{-1}$ at pH=7.5 without enzyme; rate constant $k=0.002 \text{ min}^{-1}$ at pH=7.5 with enzyme;
<b>Halflife t1/2</b>	t(1/2)=358.2 min. at pH=7.5 ,no pancreatin enzyme; t(1/2)=371.5 min. at pH=7.5 ,with enzyme.
<b>Breakdown products</b>	alpha-Terpineol acetate is hydrolyzed to alpha-terpineol and acetic acid.
<b>Conclusion remarks</b>	alpha-Terpineol readily hydrolyzes to alpha-terpineol and acetic acid in neutral media.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Remarks for Reliabilities</b>	Experiments performed with all necessary internal and external standards. Study evaluated the hydrolysis of at least 20 substrates.
<b>References</b>	Buck N. R. and Renwick A.G. (1999) Hydrolysis of tertiary terpenoid esters in artificial gastric juice and intestinal fluid. University of Southampton. Private Communication to FEMA. Unpublished Report.

**2.3 Biodegradation**

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The method followed that described in the OECD Guidelines for testing of Chemicals (1992) No. 301 F, "Ready Biodegradability";

<b>Test Type</b>	testing of Chemicals (1992) No. 301F, "Ready Biodegradability; Monometric Respiratory Test" referenced as Method C.4D of Commission Directive 92/69/EEC. Monometric Respiratory Test
<b>GLP</b>	<b>Yes</b>
<b>Year</b>	1997
<b>Contact Time</b>	<b>28</b> days
<b>Innoculum</b>	Activated sludge
<b>Remarks for Test Conditions</b>	100 mg DOC/l at 21 C for 28 days
<b>Degradation % After Time</b>	<b>6%</b> at 28 days
<b>Results</b>	<b>6%</b> after 28 days
<b>10 day Window Criteria</b>	<b>No</b>
<b>Total degradation</b>	<b>No</b>
<b>Conclusion Remarks</b>	Cannot be considered to be readily biodegradable under the terms and conditions of OECD guideline No. 301 F.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The Study was performed in compliance with UK GLP standards. These regulations are in accordance with GLP standard published as OECD monograph No. 45.
<b>Reference</b>	Mead C. (1997b) Plinyl acetate: Assessment of Ready Biodegradability; Manometric Respiratory Test. Safepharm Laboratories, Ltd. SPL Project No. 1044/015.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	<b>125252-49-5</b>
<b>Method/guideline</b>	The method followed that described in the OECD Guidelines for Testing of Chemicals (1984) No. 209 "Activated Sludge Respiration Inhibition Test" and EEC Commission Directive 87/302/EEC.
<b>Test Type</b>	Activated Sludge Respiration inhibition Test.
<b>GLP</b>	<b>Yes</b>
<b>Year</b>	<b>1997</b>
<b>Contact Time</b>	3 hours
<b>Innoculum</b>	Activated Sludge from sewage treatment plant in UK
<b>Remarks for Test Conditions</b>	The test material was aerated for a period of 3 hours at 21 C in the presence of activated sewage sludge with the addition of a synthetic sewage a respiratory substrate. The rate of respiration was determined after 30 min and 3 hours contact time.

<b>Results</b>	The effect of the test material on the respiration of activated sludge gave a 3-hour EC50 of 360 mg/l. The NOEC was determined to be 100 mg/l
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was performed in compliance with UK GLP standard. These regulations are in accordance with GLP Standards published as OECD Monograph No. 45 (OCDE/GD(92)32).
<b>Reference</b>	Mead C. (1997c) <i>Plinyl acetate: Assessment of the Inhibitory Effect on the Respiration of Activated Sewage Sludge</i> . Safepharm Laboratories. SPL Project No. 1044/018.

<b>Substance Name</b>	Linalool Synthetic, 97.6% pure (GLC)
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Ready biodegradability was determined by modified MITI test, OECD guideline
<b>Test Type</b>	Modified MITI test
<b>GLP</b>	NG
<b>Year</b>	1991
<b>Contact Time</b>	28 days
<b>Innoculum</b>	Activated sludge from city waste water treatment plant
<b>Remarks for Test Conditions</b>	Activated sludge was prepared by mixing sludge from biological waste water treatment plant & soil sampled on the bank of the river. Reference Substance: Aniline. Activated sludge conc. 30 mg/l, test substance 100 mg/l, reference substance 100 mg/l.
<b>Degradation % After Time</b>	80% at 28 days for test chemical, for aniline: 79%
<b>Results</b>	80% biodegradation in 28 days.
<b>Total degradation</b>	No
<b>Conclusion Remarks</b>	Linalool synthetic is readily biodegradable under the test conditions.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study followed OECD guidelines.
<b>Reference</b>	Rudio J. (1991) Test Report Nr 90-57/B. Linalool Synthetic. Biodegradation unpublished report.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	The sealed vessel test is a CO2 production test based on OECD Guideline 301 B(1)
<b>Test Type</b>	Sealed Vessel Test

<b>GLP</b>	<b>Yes</b>
<b>Year</b>	<b>1990</b>
<b>Contact Time</b>	28 days
<b>Innoculum</b>	Secondary effluent from an unacclimatized activated sludge plant at URL North
<b>Remarks for Test Conditions</b>	Test is conducted in a 160 ml vessel containing 100 ml mineral salt medium inoculated with secondary effluent and the respective test or reference material. The sealed vessels are incubated at 20 C on a rotary shaker for 28 days.
<b>Degradation % After Time</b>	97.1%
<b>Results</b>	97.1% (95% confidence interval 85.0 - 109.3)
<b>Time required for 10% Degradation</b>	< 4 days
<b>10 day Window Criteria</b>	Yes
<b>Conclusion Remarks</b>	The test material was readily and ultimately biodegradable.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with GLP and OECD guideline.
<b>Reference</b>	Quest International Ltd. (1994d) Biodegradability test of linalool in a sealed vessel test. Unpublished report.

<b>Substance Name</b>	Pine oil (disinfectant) (mixture contains 50-60% alpha-terpineol and 10-l 5% other terpineol isomers)
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Method/guideline</b>	Biodegradability and biochemical oxygen demand.
<b>Test Type</b>	Biodegradability
<b>GLP</b>	NG
<b>Year</b>	1966
<b>Contact Time</b>	20 days
<b>Innoculum</b>	Activated sludge
<b>Degradation % After Time</b>	Not given
<b>Results</b>	Biological Oxygen demand (5 days)= 0.8 mg/lml; COD=1.60 mg/lml; chemical oxygen demand of organic feed to activated sludge = 428 mg/lml; BOD 5-day of feed to activated sludge units= 255 mg/lml; loading on activated sludge units= 47.8.
<b>Data Qualities Reliabilities</b>	Biodegradability cannot be determined from the limited data in this report.
<b>Reference</b>	Ryckman, Edgerley, Burbank and Associates, Inc. (1966) Unpublished report to Hercules Powder Company.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	OECD closed bottle shake flask activated sludge biodegradation in soil.
<b>Test Type</b>	20 day biochemical oxygen demand
<b>GLP</b>	NG(62.7% alpha-terpineol, 83.4% total terpene alcohol content)
<b>Year</b>	1986
<b>Innoculum</b>	Activated sludge
<b>Remarks for Test Conditions</b>	Myrcenol was added by syringe, injected directly into BOD bottle due to limited solubility.
<b>Degradation % After Time</b>	<b>66%</b> in 20 days
<b>Time required for 10% Degradation</b>	<10 days
<b>10 day Window Criteria</b>	Yes
<b>Conclusion Remarks</b>	Myrcenol is biodegradable.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with OECD guideline.
<b>Reference</b>	UCC Business Confidential (1986) Ecological Fate and Effects Testing of UCC Products and Wastewaters During 1985.

<b>Substance Name</b>	alpha-Terpineol acetate (100%)
<b>CAS No.</b>	<b>80-26-2</b>
<b>Method/guideline</b>	The method followed that described in the OECD Guidelines for testing of Chemicals (1992) No. 301F, "Ready Biodegradability; Monometric Respiratory Test" referenced as Method C.4-D of Commission Directive 92/69/EEC.
<b>Test Type</b>	Monometric Respiratory Test
<b>GLP</b>	<b>Yes</b>
<b>Year</b>	<b>1996</b>
<b>Contact Time</b>	<b>28</b> days
<b>Innoculum</b>	Activated sludge
<b>Remarks for Test Conditions</b>	30 mg/L activated sludge, 100 mg/L test substance or reference (aniline) material were incubated for 28 days at 20 °C.
<b>Degradation % After Time</b>	<b>63%</b> after 28 days
<b>Results</b>	63% of the test substance was biodegraded in 28 day whereas 79% of reference material (aniline) was biodegraded during the same period.

<b>10 day Window Criteria</b>	<b>No</b>
<b>Conclusion Remarks</b>	alpha-Terpinyl acetate was biodegradable under the test conditions.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with OECD guideline.
<b>Reference</b>	Givaudan Roure (1996b) Ready biodegradability of <i>alpha</i> -terpineol acetate. Unpublished report.

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	The sealed vessel test is a CO2 production test based on OECD Guideline 301 B(1)
<b>Test Type</b>	Sealed Vessel Test
<b>GLP</b>	Yes
<b>Year</b>	1994
<b>Contact Time</b>	28 days
<b>Innoculum</b>	Secondary effluent from an unacclimatized activated sludge plant at URL North
<b>Remarks for Test Conditions</b>	Test is conducted in a 160 ml vessel containing 100 ml mineral salt medium inoculated with secondary effluent and the respective test or reference material. The sealed vessels are incubated at 20 °C on a rotary shaker for 28 days.
<b>Degradation % After Time</b>	87.3% after 28 days
<b>Results</b>	87.3% (95% confidence interval 85.889%)
<b>10 day Window Criteria</b>	Yes
<b>Conclusion Remarks</b>	The test material was readily and ultimately biodegradable.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with GLP and OECD guideline.
<b>Reference</b>	Quest International Ltd. (1994a) Biodegradability test of terpinyl acetate in a sealed vessel test. Private communication to RIFM.

<b>Substance Name</b>	Tetrahydrolinalool (98%)
<b>CAS No.</b>	<b>78-69-3</b>
<b>Method/guideline</b>	The sealed vessel test is a CO2 production test based on OECD Guideline 301 B(1).
<b>Test Type</b>	Sealed Vessel Test
<b>GLP</b>	<b>Yes</b>

<b>Year</b>	1994
<b>Contact Time</b>	28 days
<b>Innoculum</b>	Secondary effluent from an unacclimatized activated sludge plant at URL North
<b>Remarks for Test Conditions</b>	Test is conducted in a 160 ml vessel containing 100 ml mineral salt medium inoculated with secondary effluent and the respective test or reference material. The sealed vessels are incubated at 17-22 °C on a rotary shaker for 28 days.
<b>Degradation % After Time</b>	103.1 %after 28 days
<b>Results</b>	103.1% (95% confidence interval 986107.7%)
<b>10 day Window Criteria</b>	Yes
<b>Conclusion Remarks</b>	The test material was readily and ultimately biodegradable.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with GLP and OECD guideline.
<b>Reference</b>	Quest International Ltd. (1994b) The ultimate biodegradability of tetrahydrolinalool in the sealed vessel test. Private communication to RIFM.

<b>Substance Name</b>	Linalyl acetate (96.0%)
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Ready Biodegradability was determined by modified MITI test, OECD guideline.
<b>Test Type</b>	Modified MITI test
<b>GLP</b>	Yes
<b>Year</b>	1991
<b>Contact Time</b>	28 days
<b>Innoculum</b>	Activated sludge from city waste water treatment plant.
<b>Remarks for Test Conditions</b>	Activated sludge was prepared by mixing sludge from biological waste water treatment plant & soil sampled on the bank of the river. Reference Substance: Aniline. Activated sludge conc. 30 mg/l, test substance 100 mg/l, ref. substance (aniline) 100 mg/l.
<b>Degradation % After Time</b>	75% after 28 days
<b>Results</b>	75% biodegradation in 28 days.
<b>Conclusion Remarks</b>	The test material was biodegradable.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study followed OECD guidelines.

**Reference** Quest International Ltd. (1991) Biodegradability study of linalyl acetate in a sealed vessel test. Private communication to RIFM.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	The sealed vessel test is a CO <sub>2</sub> production test based on OECD Guideline 301 B(1).
<b>Test Type</b>	Sealed Vessel Test
<b>GLP</b>	Yes
<b>Year</b>	1991
<b>Contact Time</b>	28 days
<b>Innoculum</b>	Secondary effluent from an unacclimatized activated sludge plant at URL North.
<b>Remarks for Test Conditions</b>	Test is conducted in a 160 ml vessel containing 100 ml mineral salt medium inoculated with secondary effluent and the respective test or reference material. The sealed vessels are incubated at 20 C on a rotary shaker for 28 days.
<b>Degradation % After Time</b>	72.1% after 28 days
<b>Results</b>	72.1% (95% confidence interval 68.8-75.4%)
<b>Conclusion Remarks</b>	The test material was biodegradable.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with GLP and OECD guideline.
<b>Reference</b>	Quest International Ltd. (1994c) Biodegradability study of dimyrcetol in a sealed vessel test. Private comm. to RIFM.

## 2.4 Fugacity

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP

**Media** Air-Water Partition Coefficient

**Absorption coefficient** 0.0029

**Data Qualities Reliabilities** Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	15.6
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay

<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	31.3
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2. 11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	97.7
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs

<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Estimated Distribution and Media Concentration</b>	39.7
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
CAS No.	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	89600
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.
<b>Substance Name</b>	Linalool

<b>CAS No.</b>	<b>78-70-6</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	<b>0.455</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	0.317
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	0.223
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2. 11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.005</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.00015</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000013
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.00000082
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air-Water Partition Coefficient

**Absorption coefficient** 0.0016

**Data Qualities Reliabilities** Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	56.7
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I

<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	113.5
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	354.7
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model

<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	144.2
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
-----------------------	----------

<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	895000
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
-----------------------	----------

**CAS No.** 543-39-5

<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.181
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	0.227
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	<b>0.579</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.0129</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D.

(1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration Data Qualities Reliabilities</b>	0.00042 Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration Data Qualities Reliabilities</b>	0.000033 Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or

metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.  
 Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

**References**

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration Data Qualities Reliabilities</b>	0.0000032 Reliable with restriction.

**Data Reliability Remarks**

The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	
<b>Absorption coefficient</b>	

**Estimated Distribution and Media Concentration Data Qualities Reliabilities**

Reliable with restriction.

**Data Reliability Remarks**

The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
-----------------------	-----------------

<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	0.0017
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
-----------------------	-----------------

<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I

<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	58.1
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	116
<b>Data Qualities Reliabilities</b>	Reliable with restriction
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model

<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	363
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	148
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	895000
<b>Data Qualities Reliabilities</b>	Reliable with restriction,
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.186
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	0.222
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	0.579
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D.

(1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	0.0129
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	2.5 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	0.0004
<b>Data Qualities Reliabilities</b>	Reliable with restriction
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or

**References**

metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.  
Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000033
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks**

The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, VP, water solubility, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.000033

**Data Qualities Reliabilities** Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>0.0062</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	<b>25 °C</b> , 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP

<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>483</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>966</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay

<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>3020</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	<b>115-95-7</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	<b>EQC V 2.11 Level I</b>
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	1230
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs

<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	545000
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.118
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
-----------------------	-----------------

<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	0.038
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	0.825
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
CAS No.	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	0.0183
<b>Data Qualities Reliabilities</b>	Reliable with restriction
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Linalyl Acetate
CAS No.	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	0.00057
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000047
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Linalyl Acetate
<b>CAS No.</b>	115-95-7
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, VP, water solubility, estimated log Kow, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.000013
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	0.00048
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient

**Absorption coefficient** 78.3

**Data Qualities Reliabilities** Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	157
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I

<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>490</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydroinalool
<b>CAS No.</b>	<b>78-69-3</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	<b>EQC V 2.11 Level I</b>
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	199
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydroinalool
<b>CAS No.</b>	<b>78-69-3</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model

<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	1130000
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrofuralol
<b>CAS No.</b>	78-69-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.0499
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.
<b>Substance Name</b>	Tetrahydrofuralol
<b>CAS No.</b>	78-69-3

<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	<b>0.206</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	<b>78-69-3</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	<b>0.727</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	<b>78-69-3</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	0.0162
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	<b>78-69-3</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.00051</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D.

(1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000041
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, estimated log Kow, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.000011
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or

**References**

metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.  
Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	0.00046
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	18.8

**Data Qualities Reliabilities** Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
-----------------------	-----------------

**CAS No.** 98-55-5

**Model Conditions** 25 °C, 100,000 lbs

**Test Type** Environmental Equilibrium Partitioning Model

**Method** Mackay

**Model Used** EQC V 2.11 Level I

**input Parameters** MW, water solubility, log Kow, VP, MP

**Media** Sediment-Water Partition Coefficient

**Absorption coefficient** 37.6

**Data Qualities Reliabilities** Reliable with restriction,

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
-----------------------	-----------------

**CAS No.** 98-55-5

**Model Conditions** 25 °C, 100,000 lbs

**Test Type** Environmental Equilibrium Partitioning Model

**Method** Mackay

**Model Used** EQC V 2.11 Level I

**Input Parameters** MW, water solubility, log Kow, VP, MP

<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	117
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	47.8
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay

<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	<b>1130000</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.11
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs

<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	<b>0.477</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	<b>98-55-5</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	<b>0.404</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.
<b>Substance Name</b>	alpha-Terpineol

<b>CAS No.</b>	<b>98-55-5</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration Data Qualities Reliabilities</b>	<b>0.009</b> Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
-----------------------	-----------------

<b>CAS No.</b>	<b>98-55-5</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration Data Qualities Reliabilities</b>	<b>0.00028</b> Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	<b>0.000023</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, water solubility, log Kow, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	<b>0.0000025</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	0.141
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	393
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	785
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient

<b>Absorption coefficient</b>	<b>2450</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	<b>80-26-2</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>998</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	<b>80-26-2</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I

<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	<b>347000</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	<b>0.787</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model

<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	0.0112
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
-----------------------	------------------------

<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	0.197
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
-----------------------	------------------------

**CAS No.** 80-26-2

<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	0.0044
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	0.00014
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	<b>0.000011</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction,
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, estimated water solubility, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.0000055
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D.

(1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	0.0038
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.1 1 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	43
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	86.1
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP

**Media** Suspended Sediment-Water Partition Coefficient

**Absorption coefficient** 269

**Data Qualities Reliabilities** Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	109
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model

<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	<b>645000</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	<b>125252-49-5</b>
<b>Model Conditions</b>	<b>25 °C</b> , 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	<b>0.39</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
-----------------------	---

<b>CAS No.</b>	125252494
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	0.205
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance ▪ plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	0.397
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity

approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	12525249-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	0.0088
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	0.00028
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000022
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, BocaRaton, FL.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, log Kow, water solubility, VP, MP

<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.000005
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	0.0067
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay

<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>13.9</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	<b>4948-28-1</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>27.9</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	<b>4948-28-1</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs

<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	87.1
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	35.4
<b>Absorption coefficient</b>	Fish-Water Partition Coefficient
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.
<b>Substance Name</b>	<i>cis</i> -2-Pinanol

<b>CAS No.</b>	<b>4948-28-1</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	<b>645000</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	<b>4948-28-1</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.17
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	<b>0.506</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	<b>4948-28-1</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	0.317
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.007</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.00022</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, **Boca Raton, FL.**

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	<b>EQC V 2.11 Level I</b>
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000018
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, <b>Boca Raton, FL.</b>

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol

<b>Estimated Distribution and Media Concentration</b>	0.000022
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption coefficient</b>	0.0067
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I

<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>13.9</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	<b>27.9</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model

<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption coefficient</b>	87.1
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
-----------------------	-------------------------

<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption coefficient</b>	35.4
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
-----------------------	-------------------------

<b>CAS No.</b>	4948-29-2
----------------	-----------

<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption coefficient</b>	645000
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans-2-Pinanol</i>
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.17
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	<b>0.506</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	<b>0.317</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D.

(1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.007</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.00022</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or

**References**

metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.  
Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000018
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<i>trans</i> -2-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.000022

<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	<b>28324-52-9</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption Coefficient</b>	<b>0.00336</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	<b>28324-52-9</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP

Media	Soil-Water Partition Coefficient
<b>Absorption Coefficient</b>	<b>20.6</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption Coefficient</b>	41.2
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	<b>28324-52-9</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay

<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption Coefficient</b>	129
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1, Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption Coefficient</b>	52.4
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Model Conditions</b>	25 °C, 100,000 lbs

<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption Coefficient</b>	544000
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.1 1, Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-g
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.463
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL
<b>Substance Name</b>	2-Pinanolhydroperoxide

<b>CAS No.</b>	<b>28324-52-9</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	<b>0.276</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	<b>2-Pinanolhydroperoxide</b>
<b>CAS No.</b>	<b>28324-52-9</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	<b>0.256</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction,
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1992) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	0.0057
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment

<b>Absorption Coefficient</b>	
<b>Estimated Distribution and Media Concentration</b>	0.00018
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or

**References**

metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.  
Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000014
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	2-Pinanolhydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.000005

**Data Qualities Reliabilities** Reliable with restriction.

**Data Reliability Remarks** The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References** Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air-Water Partition Coefficient
<b>Absorption Coefficient</b>	<b>0.00083</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP

<b>Media</b>	Soil-Water Partition Coefficient
<b>Absorption Coefficient</b>	15.6
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Absorption Coefficient</b>	31.3
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay

<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment-Water Partition Coefficient
<b>Absorption Coefficient</b>	97.7
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish-Water Partition Coefficient
<b>Absorption Coefficient</b>	39.7
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Model Conditions</b>	25 °C, 100,000 lbs

<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol-Air Partition Coefficient
<b>Absorption Coefficient</b>	645000
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	0.194
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
-----------------------	----------

<b>CAS No.</b>	<b>8002-09-3</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	<b>0.469</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Soil
<b>Estimated Distribution and Media Concentration</b>	<b>0.33</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.0073</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Model Conditions</b>	<b>25 °C, 100,000 lbs</b>
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Suspended Sediment
<b>Estimated Distribution and Media Concentration</b>	<b>0.00023</b>
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Fish
<b>Estimated Distribution and Media Concentration</b>	0.000019
<b>Data Qualities Reliabilities</b>	Reliable with restriction.
<b>Data Reliability Remarks</b>	The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.
<b>References</b>	Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Model Conditions</b>	25 °C, 100,000 lbs
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Mackay
<b>Model Used</b>	EQC V 2.11 Level I
<b>Input Parameters</b>	MW, estimated log Kow, water solubility, VP, MP
<b>Media</b>	Aerosol
<b>Estimated Distribution and Media Concentration</b>	0.0000025
<b>Data Qualities Reliabilities</b>	Reliable with restriction.

**Data Reliability Remarks**

The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism. Reliable with restriction because this method does not allow for biodegradation or metabolism.

**References**

Level 1 Fugacity-based Environmental Equilibrium Partitioning Model Version 2.11. Based on Mackay, D. (1991) Multimedia environmental models: The fugacity approach. Lewis Publications, CRC Press, Boca Raton, FL.

### 3 Ecotoxicity

#### 3.1 Acute Toxicity to Fish

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - <b>plinyl</b> acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	Followed method in the OECD Guidelines for Testing Chemicals (1992) No. 203, "Fish Acute Toxicity Test" referenced as Method C.I of Commission Directive 92/69/EEC.
<b>Test Type</b>	Acute Fish Toxicity
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/Strain/Supplier</b>	Juvenile rainbow trout
<b>Analytical Monitoring</b>	Moving average method of Thompson
<b>Exposure Period</b>	96 hr
<b>Remarks for Test Conditions</b>	Based on the results of the range-finding study, the following test concentrations were assigned to the definite study: 1.5, 2.7, 4.8, 8.5 & 15 mg/l. Test material was dissolved in and controls were exposed to dimethylformamide.
<b>Observations</b>	8.5 mg/L: Highest concentration resulting in 0% mortality; 15 mg/L: lowest concentration resulting in 100% mortality; No observed effect concentration (NOEC): 2.7 mg/L; sub-lethal effects were observed at concentration of 4.8 mg/L and above.
<b>Nominal Concentrations as mg/L</b>	1.5, 2.7, 4.8, 8.5 & 15 mg/L
<b>Measured Concentrations as mg/L</b>	1.47, 2.55, 4.53, 13.9 mg/l
<b>Remarks for Results</b>	Sub-lethal effects observed were increased pigmentation, swimming at the bottom of the test vessels with increased pigmentation, swimming at the surface with increased pigmentation and the presence of moribund fish.
<b>Conclusion Remarks</b>	The acute toxicity of the test material to the fresh water fish rainbow trout has been investigated and gave a 96-hr LC50 value of 11 mg/L with 95% confidence limit of 8.515 mg/l. No-Observed-Effect Concentration was 2.7 mg/L.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was performed in compliance with U.K. GLP standard. These regulations are in accordance with GLP standards published as OECD monograph No. 45 (OECD/GD/(92)32) and are in conformity with, and implement, the requirement Directive 87/18/EEC.
<b>Reference</b>	Wetton P. M. (1997a). Plinyl Acetate: Acute Toxicity to Rainbow Trout ( <i>Oncorhynchus mykiss</i> ). Safepharm Laboratories Ltd. SPL Project Number: 10441012.

<b>Substance Name</b>	alpha-Terpineol (90% + 10% emulsifier)
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Method used was in accordance with Protocol outlined by Wan et al. (1990, 1991) and by Environment Canada 91990a, 1990b)
<b>Test Type</b>	Acute Fish Toxicity
<b>GLP</b>	NG
<b>Year</b>	1998
<b>Species/Strain/Supplier</b>	Coho salmon and Rainbow trout
<b>Exposure Period</b>	96 hr
<b>Remarks for Test Conditions</b>	10 fishes were exposed to the test substance and observed for 24, 48, 72 and 96 hour. LC50 values were calculated using the "Lethal Computer Program" developed by Stephen (1977, revised in 1983)
<b>Nominal Concentrations as mg/L</b>	5 nominal concentration below 100 ppm were selected
<b>Remarks for Results</b>	LC50 for Coho Salmon =6.8 ppm; LC50 for Rainbow Trout = 6.7 ppm
<b>Conclusion Remarks</b>	The LC50 for Coho Salmon was determined to be 6.8 mg/l and for Rainbow Trout was 6.7 mg/l.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions
<b>Data Reliability Remarks</b>	Article was published in Bull. Environ. Contam. Toxicol.
<b>Reference</b>	Stroh J., Wan, M. T., Isman, M. B. and Moul, D. J. (1998) Evaluation of the acute toxicity to juvenile pacific coho salmon and rainbow trout of some plant essential oils, a formulated product, and the carrier. Bull Enviro. Contam. Toxicol. 60: 923-930

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	The test was conducted by EPA-ASTH procedure using 10 fathead minnows per test concentrations in a total volume of 10 liters. Temperature, fish survival, pH and dissolved oxygen levels were monitored during the test period of 96 hours.
<b>Test Type</b>	Acute Fish Toxicity
<b>GLP</b>	NG
<b>Year</b>	1986
<b>Species/Strain/Supplier</b>	Fathead minnows
<b>Analytical Monitoring</b>	Temperature, pH, fish survival and dissolved oxygen levels

<b>Exposure Period</b>	96 hr
<b>Remarks for Test Conditions</b>	Minimal aeration was supplied when the dissolved oxygen was below 4 mg/L. All samples were tested within six hours after the collection without pH adjustment or solids removal.
<b>Observations</b>	Median lethal concentration (LC50)=3.7 mg/L
<b>Conclusion Remarks</b>	Median lethal concentration (LC50)=3.7 mg/L
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Reference</b>	UCC Business Confidential (1986). Ecological Fate and Effects Testing of UCC Products and Wastewaters During 1985. Unpublished Report.

<b>Substance Name</b>	Pine oil (62.7% alpha-terpineol, 83.4% total terpene alcohol content)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Bioassays were carried out according to the procedures outlined in APHA 1975. Standard Method for examination of water & waste water. APHA, American Water Works Association, Water Pollution Control Federation. 14th Ed. Wash. DC 1193p.
<b>Test Type</b>	Acute Fish Toxicity (range finding test)
<b>GLP</b>	NG
<b>Year</b>	1994
<b>Species/Strain/Supplier</b>	Juvenile Rainbow trout ( <i>Salma, gairdneri</i> ).
<b>Exposure Period</b>	96 hrs
<b>Remarks for Test Conditions</b>	Concentrations tested: 25, 50, 75, 100, 200 & 1000 mg/L. The percentage of survival of the fish in the solution was observed at 24, 48, 72 & 96 hours. LC50 values and their 95% confidence limits were calculated according to Stephan (1977). 96 hour LC50 was calculated to be 71 mg/L (95% CI=50-100)
<b>Conclusion Remarks</b>	
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with APHA procedure.
<b>Reference</b>	Graves, W. C. and Swigert, J. P (1994b). Pine Oil: 96-Hours Flow-through Acute Toxicity Test with the Rainbow Trout ( <i>Oncorhynchus mykiss</i> ). Wildlife International Ltd Report to The CSMA Pine Oil Joint Venture. Lab Project ID/Study Number 274A-102.

<b>Substance Name</b>	Pine oil Disinfectant (80:10:10) (mixture contains 50-60% alpha-terpineol and 10-1 5% terpineol isomers)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	The procedure used was that described in the 1 lth edition of Standard Methods for Examination of Waters and Waste

<b>Test Type</b>	Waters & the Sept 8, 1964 "Dept of Interior Guide for Evaluation of Wildlife. Acute Fish Toxicity
<b>GLP</b>	NG
<b>Year</b>	1965
<b>Species/Strain/Supplier</b>	Bluegill fingerlings ( <i>Lepomis macrochirus</i> )
<b>Exposure Period</b>	24 & 48 hrs
<b>Observations</b>	LC50 at 24 h= 59 mg/L; LC50 at 48 h= 58 mg/L
<b>Conclusion Remarks</b>	The LC50 of the test material was calculated to be 58-59 mg/l
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with Dept of Interior Guideline.
<b>Reference</b>	Ryckman, Edgerley, Burbank and Associates (1965) Investigation of treatability and fish toxicity of certain manufacturing products and wastes and their decay under stream conditions. Unpublished report to Hercules Powder Co.

<b>Substance Name</b>	The study was conducted using structurally related Iso-Linalool (2,4-Dimethyl 2,7-Octadien-4-ol).
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	The protocol followed the standard procedures described in the EPA/OTS guidelines for testing the effects of chemicals on fish and meet the TSCA guidelines as specified in the appropriate Registration Standard.
<b>Test Type</b>	Acute Fish Toxicity
<b>GLP</b>	Yes
<b>Year</b>	1990
<b>Species/Strain/Supplier</b>	Rainbow trout ( <i>Oncorhynchus mykiss</i> )
<b>Analytical Monitoring</b>	All samples analyzed by HPLC
<b>Exposure Period</b>	24, 48, 72 & 96 hours.
<b>Nominal Concentrations as mg/L</b>	100, 60, 36, 22 and 13 mg active ingredient/L
<b>Measured Concentrations as mg/L</b>	77, 44, 25, 17 and 10 mg active ingredient/L
<b>Unit</b>	Mg active ingredient/L
<b>Remarks for Results</b>	The 96-hour LC50 value was estimated by non-linear interpolation to be 33 mg/L; 95% confidence interval was calculated by binomial probability to be 25-44 mg/L. The No Observed Effect Concentration established for this study was 10 mg A.I./L.

**Conclusion Remarks** The 96-hr LC50 value was estimated to 33 mg/L; 95% confidence interval was calculated to be 25-44 mg/L. The No Observed Effect Concentration established for this study was 10 mg A.I./L.

**Data Qualities Reliabilities** Reliability code 1. Reliable without restrictions,

**Data Reliability Remarks** The study was conducted in accordance to GLP.

**Reference** LeLievre, M. (1990a) 2,4-Dimethyl 2,7-octadien-4-ol)-acute toxicity to rainbow trout (*Oncorhynchus mykiss*) under static conditions. Springborn Lab. Inc. Report Submitted to SCM Glidco Organics. SLI Report # 90-g-3476.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	EPA ECOSAR Model
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hr
<b>Observations</b>	96 Hour LC50= 0.60 mg/l
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hr
<b>Observations</b>	96 Hour LC50 = 4.85 mg/l
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity

**Exposure Period** 96 hrs  
**Observations** 96 Hour LC50 = 4.81 mg/l  
**Unit** mg/l  
**Reference** EPA ECOSAR

<b>Substance Name</b>	Linalyl acetate
CAS No.	115-95-7
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hrs
<b>Observations</b>	96 Hour LC50 = 4.9 mg/l using the calculated Kow of 3.46 but is 9.8 if you use the measured log Kow of 2.9
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hr
<b>Observations</b>	96 Hour LC50 = 3.68 mg/l
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hrs
<b>Observations</b>	96 Hour LC50 = 13.7 mg/l

Unit mg/l  
Reference EPA ECOSAR

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hrs
<b>Observations</b>	96 Hour LC50 = 1.75 mg/l based on log Kow = 4.3
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hrs
<b>Observations</b>	96 Hour LC50 = 17.4 mg/l
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	trans-Pinanol
<b>CAS No.</b>	4948-29-2
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hrs
<b>Observations</b>	96 Hour LC50 = 18.2 mg/l
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hrs
<b>Observations</b>	96 Hour LC50 = 18.2 mg/l
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	Pinanol hydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Method/guideline</b>	EPA <b>ECOSAR</b> – calculated as the decomposition product 2-pinanol)
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hrs
<b>Observations</b>	96 Hour LC50 = 18.2 mg/l
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	Pine oil
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	EPA ECOSAR
<b>Test Type</b>	Acute Fish Toxicity
<b>Exposure Period</b>	96 hr
<b>Observations</b>	96 Hour LC50 = 14 mg/l calculated as alpha-Terpineol
<b>Unit</b>	mg/l
<b>Reference</b>	EPA ECOSAR

<b>Substance Name</b>	Pine oil (62.7% alpha-terpineol, 83.4% total terpene alcohol content)
-----------------------	---

<b>CAS No.</b>	<b>8002-09-3</b>
<b>Method/guideline</b>	Bluegills were exposed to a geometric series of 5 test concentrations, solvent control, and a negative (well water) control. Two replicate test chambers were maintained in each treatment and control group, with 10 bluegills in each test chamber.
<b>Test Type</b>	Acute Fish Toxicity
<b>GLP</b>	Yes
<b>Year</b>	1994
<b>Species/Strain/Supplier</b>	Bluegill ( <i>Lepomis macrochirus</i> )
<b>Analytical Monitoring</b>	Quantification of Pine oil in exposure solution was based upon the analysis of alpha terpineol.
<b>Exposure Period</b>	96 hrs
<b>Remarks for Test Conditions</b>	Nominal test concentrations were based upon the results of a range finding toxicity test.
<b>Nominal Concentrations as mg/L</b>	7.8, 13, 22, 36, and 60 mg/L
<b>Measured Concentrations as mg/L</b>	6.4, 11, 21, 36 and 58 mg/L
<b>Remarks for Results</b>	96-Hour LC50: 53 mg/L; 95% Confidence Limit: >36 mg/L; No mortality concentration: 36 mg/L. At 36 mg/L, all fish became lethargic after 72 hrs; at 58 mg/L, all fish became lethargic within 5 hrs; by termination, 65% of fish at 58 mg/L were dead.
<b>Conclusion Remarks</b>	The 96-hour LC50 value for bluegill exposed to Pine oil blend was 53 mg/L. At the 95% confidence level, the LC50 value is above 36 mg/L.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted to conform with Good Laboratory Practice standards published by EPA and OECD.
<b>Reference</b>	Graves W. C. and Swigert, J. P (1994a) Pine Oil: 96-Hours Flow-through Acute Toxicity Test with Bluegill ( <i>Lepomis macrochirus</i> ). Wildlife International Ltd Report to The CSMA Pine Oil Joint Venture. Lab Project ID/Study Number 274A-101

<b>Substance Name</b>	Pine oil (62.7% alpha-terpineol, 83.4% total terpene alcohol content)
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Method/guideline</b>	Rainbow Trout were exposed to a geometric series of 5 test concentrations, solvent control, and a negative (well water) control. Two replicate test chambers were maintained in each treatment and control group, with 10 trout in each test chamber.
<b>Test Type</b>	Acute Fish Toxicity
<b>GLP</b>	Yes
<b>Year</b>	1994

<b>Species/Strain/Supplier</b>	Rainbow Trout ( <i>Oncorhynchus mykiss</i> )
<b>Analytical Monitoring</b>	Measured concentrations of alpha-terpineol were used to calculate the concentration of Pine oil in solution.
<b>Exposure Period</b>	96 hrs
<b>Remarks for Test Conditions</b>	Nominal test concentrations were based upon the results of a range finding toxicity test. Observations were made approximately 3, 24, 48, 72 and 96 hours after test initiation.
<b>Nominal Concentrations as mg/L</b>	6.5, 11, 18, 30 and 50 mg/L
<b>Measured Concentrations as mg/L</b>	6.2, 10, 17, 28 and 48 mg/L
<b>Remarks for Results</b>	96-Hour LC50: 18 mg/L; 95% Confidence limits: 16 & 21 mg/L; No mortality concentration: 10 mg/L.
<b>Conclusion Remarks</b>	The 96-hour LC50 value for Rainbow Trout exposed to Pine Oil Blend was 18 mg/L. The 95% confidence limits were 16 & 21 mg/L and the slope of the concentration-response curve was 9.4. The 96-hrs no mortality concentration was 10 mg/L.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted to conform to GLP published by US EPA and OECD.
<b>Reference</b>	Graves, W. C. and Swigert, J. P (1994b) Pine Oil: 96-Hours Flow-through Acute Toxicity Test with the Rainbow Trout ( <i>Oncorhynchus mykiss</i> ). Wildlife International Ltd Report to The CSMA Pine Oil Joint Venture. Lab Project ID/Study Number 274A-102.

### 3.2 Acute Toxicity to Aquatic Invertebrates

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - pinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	OECD guideline for testing of Chemicals (1984) No. 202, "Daphnia sp., Acute Immobilization test" referenced as Method C.2 of Commission Directive 92/69/EEC.
<b>Test Type</b>	Acute Immobilization Test
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	Following a preliminary range-finding study, 20 daphnids were exposed to an aqueous dispersion of the test material at CONC. of 0.15, 0.27, 0.48, 0.85, 1.5, 2.7, 4.8, 8.5, & 15 mg/l for 48 hrs under static test conditions.

<b>Remarks for Test Conditions</b>	The number of immobilized Daphnia were recorded after 24 and 48 hours.
<b>Nominal Concentrations as mg/L</b>	0.15, 0.27, 0.48, 0.85, 1.5, 2.7, 4.8, 8.5 & 15
<b>Measured concentrations as mg/L</b>	In excess of the required 80% of nominal conc.
<b>EC50, EL50, LC50 at 24, 48 hours</b>	48 hour EC50 = 7.0 mg/ml with 95% CI-6.3-7.6 mg/l.
<b>Remarks for Results</b>	The No Observed Effect Concentration was 4.8 mg/L
<b>Conclusion Remarks</b>	The acute toxicity of <i>Plinyl</i> acetate to Daphnia magna has been investigated and gave a 48-hours EC50 of 7.0 mg/l with 95% CI of 6.3-7.6 mg/l. The No Observed Effect Concentration at 48 hours was 4.8 mg/l.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was in compliance with the UK Principles of Good Laboratory Practice. These Principles are in accordance with GLP standards published as OECD Environment Monograph No. 45 (OECD/GD(92)32).
<b>Reference</b>	Wetton P. M. (1997b) <i>Plinyl</i> Acetate: Acute Toxicity to Daphnia Magna. Safepharm Laboratories, Inc. SPL Project Number: 1044/013.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	The test procedure closely followed those recommended by the EPA Committee on Methods for Toxicity Tests with Aquatic organisms except replicate concentrations were not routinely used.
<b>Year</b>	1985
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	The young Daphnia were randomly added to the test solution. Dissolved oxygen and pH were determined initially and at 48 hours for all test concentration & controls. Mortalities were recorded at 24 & 48 hours.
<b>EC50, EL50, LC50 at 24, 48 hours</b>	LC50=36 mg/L at 48 hours
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with EPA recommended method.
<b>Reference</b>	UCC Business Confidential (1986). Ecological Fate and Effects Testing of UCC Products and Wastewaters During 1985. Unpublished Report.

<b>Substance Name</b>	Linalool (data for structurally related substance, isolinalool (2,4-dimethyl-2,7-octadien-4-ol) clear liquid, 93.73% pure)
<b>CAS No.</b>	78-70-6

<b>Method/guideline</b>	The protocol followed the standard procedures described in the EPA/OTS guidelines for testing the effects of chemicals on daphnids, and meets the TSCA guidelines as specified in the appropriate Registration Standard.
<b>GLP</b>	Yes
<b>Year</b>	1990
<b>Analytical Procedures</b>	High pressure liquid chromatography (HPLC)
<b>Species/Strain/Supplier</b>	Daphnia pulex
<b>Nominal Concentrations as mg/L</b>	180, 110, 65, 40, 23, and 14 mg A.I./L
<b>Measured concentrations as mg/L</b>	110, 63, 37, 24, 14, and 9.3 mg A.I./L
<b>Unit</b>	mg active ingredient/L
<b>EC50, EL50, LC50 at 24, 48 hours</b>	EC50=47 mg/L (95% CI: 42-53) @ 48 h
<b>Biological Observations</b>	At 48 hour, 100% immobilization in 110 mg A.I./L; 90% and 15% immobilization at 63 & 37 mg A.I./L, respectively; sub lethal effects (lethargy) at 63, 37, & 24 mg/L; no immobilization at 24, 14 or 9.3 mg/L;
<b>Control Response</b>	Yes
<b>Statistical Evaluations</b>	EC50 value was calculated by probit analysis
<b>Conclusion Remarks</b>	The EC/50 was 68 (95% CI 37-110) and 47 (95% CI 42-53) mg active ingredient/L at 24 and 48 hours respectively.
<b>Data Qualities Reliabilities</b>	Reliability code 1, Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted in accordance with GLP guidelines
<b>Reference</b>	LeLievre M. (1990b) 2,4-Dimethyl 2,7-Octadien-4-ol - Acute Toxicity to Daphnids (Daphnia pulex) under static conditions. Springborn Lab. Inc. Submitted to SCM Glidco Organics, Inc. SLI Report # 90-g-3494.

<b>Substance Name</b>	Pine oil blend (62.7% alpha-terpineol, 83.4% total terpene alcohol content)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Daphnids were exposed to a geometric series of 5 test concentrations, a solvent control, and a negative (well water) control. Two replicate test chambers were maintained in each treatment and control group, with 10 daphnids in each test chambers.
<b>Test Type</b>	Acute toxicity test with the Daphnia magna
<b>GLP</b>	Yes
<b>Year</b>	1994
<b>Analytical Procedures</b>	Measured concentrations of alpha terpineol were used to calculate the concentrations of Pine Oil Blend in the test

	solutions.
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	Daphnids were impartially assigned to exposure chambers at test initiation. Observations of mortality and other clinical signs were made approximately 19, 24 and 48 hours after the initiation.
<b>Remarks for Test Conditions</b>	Cumulative percent mortality and immobility observed in the treatment groups were used to calculate EC50 values at 24 and 48 hours. The no observed effect concentration was determined by examination of the mortality and clinical observation data.
<b>Nominal Concentrations as mg/L</b>	3.9, 6.5, 11, 18 and 30 mg/L
<b>Measured concentrations as mg/L</b>	3.7, 7.3, 11, 17 and 28 mg/L
<b>EC50, EL50, LC50 at 24, 48 hours</b>	24 mg/L at 48 hours
<b>Biological Observations</b>	<b>48-hour EC50 = 24mg/L</b> ; 95% confidence limits: 17 and 28 mg/L; No observed effect concentration: 11 mg/L.
<b>Control Response</b>	Yes
<b>Statistical Evaluations</b>	Binomial method
<b>Conclusion Remarks</b>	The <b>48-hour</b> EC50 value for daphnids exposed to Pine oil blend was 24 mg/L. The 95% confidence limits were 17 and 28 mg/L. The <b>48-hr</b> no observed effect concentration was 11 mg/L.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was conducted to conform to GLP by the US EPA and OECD.
<b>Reference</b>	Graves, W. C. and Swigert, J. P (1994c) Pine Oil: 48 Hours Flow-Through Acute Toxicity Test with the Cladoceran (Daphnia magna). Wildlife International Ltd. Report to The CSMA Pine Oil Joint Venture. Lab Project ID/No. 274A-103.

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 20.64 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 15.72 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 1.16 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 5.70 mg/l

**Data Reliability Remarks** The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.  
**Reference** ECOSAR

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 1.04 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 5.75 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Pinane hydroperoxide
<b>CAS No.</b>	28324-52-9
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna

<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 20.64 mg/l (calculated as 2-Pinanol)
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Tetrahydrolinalool
CAS No.	78-69-3
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated based on measured Kow
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hrs
<b>Remarks for Results</b>	LC50 = 4.40 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

### 3.3 Acute Toxicity to Aquatic Plants

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The method followed that described in the OECD Guidelines for Testing of Chemicals (1984) No. 201, "Alga, Growth Inhibition Test" referenced as Method C.3 of Commission Directive 92/69/EEC (which constitutes Annex V of Council directive 67/548/EEC).
<b>Test Type</b>	Algae Growth Inhibition test
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/Strain/Supplier</b>	Green Algal, Scenedesmus subspicatus
<b>Exposure Period</b>	72 hr
<b>Analytical Monitoring</b>	Mean cell density was measured spectrophotometrically at 665 nm.

<b>Nominal Concentrations mg/L</b>	15 mg/l
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50>15 mg/l; NOEC>15 mg/l
<b>Control Response</b>	Yes
<b>Statistical Evaluations</b>	A student t-test was carried out on the AUC @ 72h.
<b>Conclusion Remarks</b>	The EC50 & the NOEC was greater than 15 mg/l
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliability Remarks</b>	The study was performed in compliance with UK GLP standard. These regulations are in accordance with GLP standards published as OECD Monograph No. 45 (OCDE/GD(92)32).
<b>Reference</b>	Mead C. (1997a) <i>Plinyl</i> Acetate: Algal Inhibition Test. Safepharm Laboratories, Ltd. SPL Project Number 10441014.

<b>Substance Name</b>	Linalool (data for structurally related isomer, iso-linalool (2,4-dimethyl 2,7-octadien-4-ol) 93.7% pure)
<b>CAS No.</b>	78-70-6
<b>Test Type</b>	Acute toxicity to green alga
<b>GLP</b>	Yes
<b>Year</b>	1990
<b>Species/Strain/Supplier</b>	Freshwater green alga ( <i>Selenastrum capricornutum</i> ).
<b>Endpoint Value</b>	Reduction in cell density after 24, 48, 72 and 96
<b>Exposure Period</b>	96 hr
<b>Remarks for Test Conditions</b>	Test conditions: 96-hours duration, 24-28 C, constant illumination (4,000-5,000 lux), shaking at 100 rpm.
<b>Nominal Concentrations mg/L</b>	32, 16, 8.0, 4.0, 2.0 and 1.0 mg/L
<b>Measured concentrations as mg/L</b>	28, 14, 6.9, 3.3, 1.6 and 0.82 mg/L
<b>Unit</b>	mg/L
<b>NOEC, LOEC or NOEL, LOEL</b>	NOEC = 3.3 mg/L
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Reference</b>	Giddings J. (1990). (2,4-Dimethyl 2,7-octadien-4-ol)-acute toxicity to freshwater green alga ( <i>Selenastrum capricornutum</i> ) under static conditions. Springborn Lab. Inc. Report Submitted to SCM Glidco Organics. SLI Report #390-10-3497.

<b>Substance Name</b>	<i>cis</i> -2-Pinanol
<b>CAS No.</b>	4948-28-1

<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 13.55 mg/l
<b>Conclusion Remarks</b>	EC50 = 13.55 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 10.40 mg/l
<b>Conclusion Remarks</b>	EC50 = 10.40 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 0.14 mg/l
<b>Conclusion Remarks</b>	EC50 = 0.14 mg/l

**Data Reliability Remarks** The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.  
**Reference** ECOSAR

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 3.88 mg/l
<b>Conclusion Remarks</b>	EC50 = 3.88 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 3.93mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR

<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 0.14 81 mg/l [Using Kow = 2.9]
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 0.14 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 3.91 mg/l
<b>Conclusion Remarks</b>	EC50 = 3.91 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	ECOSAR
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hr
<b>NOEC, LOEC or NOEL, LOEL</b>	EC50 = 3.02 mg/l
<b>Conclusion Remarks</b>	EC50 = 3.02 mg/l
<b>Data Reliability Remarks</b>	The data are obtained by a recognized SAR calculation method and are consistent with chemical structure. Data are considered reliable.
<b>Reference</b>	ECOSAR

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Plate Inhibition Assay [Ikawa, 1985]
<b>Test Type</b>	Algal Growth Inhibition Test
<b>GLP</b>	NA
<b>Year</b>	1992
<b>Species/Strain/Supplier</b>	Green algae/Chlorella pyrenoidosa
<b>Exposure Period</b>	48 hr
<b>Analytical Monitoring</b>	Net diameter of inhibition zone=Total diameter -disk diameter.
<b>Remarks for Test Conditions</b>	Plates under fluorescent light and zones of inhibition read after 2 days. Net diameter of inhibition was mean of three disks per plate run on 2 separate occasions. Lightening or total wipe out of color in yellow green Chlorella lawn.
<b>Nominal Concentrations mg/L</b>	10, 1, and 0.1 mg/ml in ethanol
<b>Unit</b>	mg/ml
<b>NOEC, LOEC or NOEL, LOEL</b>	NOEC = 1 mg/ml (1000 mg/L)
<b>Biological Observations</b>	Lightening of lawn color at 10 mg/ml concentration compared to controls
<b>Control Response</b>	Yes
<b>Statistical Evaluations</b>	None
<b>Conclusion Remarks</b>	NOEC = 1.0 mg/ml (1000 mg/L)
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions
<b>Data Reliability Remarks</b>	Data was reported in a peer-reviewed journal.
<b>General Remarks</b>	Authors noted that stored samples produced effects. Newly purchased samples were inactive. Inhibition occurred through vapor phase and not through diffusion in agar medium.
<b>Reference</b>	Ikawa M., Mosley S., and Barbero B. (1992) Inhibitory effects of terpene alcohols and aldehydes on the growth of green alga Chlorella pyrenoidosa. Journal of Chemical Ecology 18( 10): 1755-1 760.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Plate Inhibition Assay [Ikawa, 1985]

<b>Test Type</b>	Algal Growth Inhibition Test
<b>GLP</b>	NA
<b>Year</b>	1992
<b>Species/Strain/Supplier</b>	Green algae/ <i>Chlorella pyrenoidosa</i>
<b>Exposure Period</b>	48 hr
<b>Analytical Monitoring</b>	Net diameter of inhibition <del>zone=Total</del> diameter -disk diameter.
<b>Remarks for Test Conditions</b>	Plates under fluorescent light and zones of inhibition read after 2 days. Net diameter of inhibition was mean of three disks per plate run on 2 separate occasions. Lightening or total wipe out of color in yellow green <i>Chlorella</i> lawn.
<b>Nominal Concentrations mg/L</b>	10, 1, and 0.1 mg/ml in ethanol
<b>Unit</b>	mg/ml
<b>NOEC, LOEC or NOEL, LOEL</b>	NOEC = 10 mg/ml (10,000 mg/L)
<b>Biological Observations</b>	No signs of inhibition of growth of <i>Chlorella</i> p. at any concentration tested.
<b>Control Response</b>	Yes
<b>Statistical Evaluations</b>	None
<b>Conclusion Remarks</b>	No effects on growth of <i>Chlorella pyrenoidosa</i> . NOEC = 10.0 mg/ml (10,000 mg/L)
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions
<b>Data Reliability Remarks</b>	Data was reported in a peer-reviewed.
<b>Reference</b>	Ikawa M., Mosley S., and Barbero B. (1992) Inhibitory effects of terpene alcohols and aldehydes on the growth of green alga <i>Chlorella pyrenoidosa</i> . <i>Journal of Chemical Ecology</i> . <b>18</b> (10): 1755-1760.

## 4 Human Health Data

### 4.1 Acute Toxicity

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	Acute dermal toxicity was determined in male and female New Zealand white rabbits.
<b>Test Type</b>	Acute Dermal LD50 study
<b>GLP</b>	NG
<b>Year</b>	1979
<b>Species/strain</b>	White New Zealand rabbit
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	2M and 1F
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	Two male and one female rabbit were intact and 1 male and 2 females were abraded. Material applied in a close batch test. Animals were observed at 1, 3, 6, and 24 hours, daily for 14 days. All rats were given a complete necropsy.
<b>Value LD50 or LC50 with confidence limits</b>	Dermal LD50 = >5000 mg/kg
<b>Number of deaths at each dose level</b>	No deaths reported.
<b>Remarks for Results</b>	No remarkable necropsy findings.
<b>Conclusion Remarks</b>	Acute dermal LD50 > 5000 mg/kg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Palanker A. L. and Lewis C. A. (1979) Acute toxicity studies in rats and rabbits. Unpublished report to RIFM.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Acute dermal toxicity was determined in albino rabbits
<b>Test Type</b>	Acute Dermal LD50 study
<b>GLP</b>	NG
<b>Year</b>	1970

<b>Species/strain</b>	Albino rabbit
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	3
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	Material was applied to shaved abdomen of rabbits in 24-hour closed-patch test. Animals were observed for 7 days. On day 5, hematology and clinical chemistry were performed. On day 7, animals were sacrificed as and necropsied.
<b>Value LD50 or LC50 with confidence limits</b>	Dermal LD50 = 5610 mg/kg (95% C.I. 3580-8370 mg/kg)
<b>Number of deaths at each dose level</b>	0/3 at 2500 mg/kg, 1/3 at 5000 mg/kg, 3/3 at 10,000 mg/kg.
<b>Remarks for Results</b>	Symptomatology: depression
<b>Conclusion Remarks</b>	Dermal LD50 was determined to be 5610 mg/kg.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Fogelman R.W. (1970) Acute dermal toxicity study in rabbits. Unpublished report to RIFM.

<b>Substance Name</b>	cis-2-Pinanol
<b>CAS No.</b>	4948-28-1
<b>Method/guideline</b>	Acute oral toxicity was determined in male and female Wistar albino rats
<b>Test Type</b>	Acute Oral LD50 study
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1979
<b>Species/strain</b>	Wistar rat
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Vehicle</b>	Corn oil
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	Rats fasted for 18 hours were given the test material at 25% in corn oil. Animals were observed at 1, 3, 6, and 24 hours, daily for 14 days All rats were given a complete necropsy.
<b>Value LD50 or LC50 with confidence limits</b>	Oral LD50= 2050 mg/kg (95% C. I., 1630-2580 mg/kg)
<b>Number of deaths at each dose level</b>	1260 mg/kg, 2/5 female deaths; 2,000 mg/kg, 2/5 male and 3/5 female deaths; 2520 mg/kg, 2/5 male and 4/5 female deaths; 3140 mg/kg, 4/5 male and 3/5 female deaths
<b>Remarks for Results</b>	Depression in 1 male and 5 females at 2000 mg/kg. Depression in 4 male and 5 females at 2520 mg/kg. Depression in all at

3140 mg/kg. No remarkable necropsy findings

**Conclusion Remarks** Acute oral LD50 =2050 mg/kg

**Data Qualities Reliabilities** Reliability code 1. Reliable without restrictions.

**References** Palanker A. L. and Lewis C. A. (1979) Acute toxicity studies in rats and rabbits. Unpublished report to RIFM.

<b>Substance Name</b>	Pine oil • Hercules Fortified Pine-Odor Disinfectant-transparent yellow liquid (6060% terpineol isomers, mainly alpha-terpineol)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Acute Oral toxicity study in rats The technique of Weil, Thompson and Thompson and Weil was employed.
<b>Test Type</b>	Acute Oral toxicity study
<b>GLP</b>	NG
<b>Year</b>	1976
<b>Species/strain</b>	Sprague-Dawley rats
<b>Sex</b>	Female
<b># of animals per sex per dose</b>	4
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	Doses tested: 1350, 4556, 6834, 10250 & 15380 mg/kg
<b>Value LD50 or LC50 with confidence limits</b>	Acute LD50 = 10250 mg/kg +/- 1.475 mg/kg
<b>Number of deaths at each dose level</b>	1350 mg/kg-0/4; 4556 mg/kg-0/4; 6834 mg/kg-1/4; 10250 mg/kg-1/4 and 15380 mg/kg-4/4
<b>Remarks for Results</b>	At necropsy- red discolored lungs, gastritis and pale, discolored kidneys. In addition, mottled livers were noted in 2 rats. Examination of the survivors did not reveal any gross pathologic alterations.
<b>Conclusion Remarks</b>	The oral LD50 of Hercules Fortified Pine-Odor Disinfectant was calculated to be 10,250 mg/kg in female Sprague-Dawley rats, and the substance was considered to be practically nontoxic.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Johnson M. C. and Lewis, R.F. (1976) Report to Hercules Inc. on Acute Toxicity Studies with Hercules Fortified Pine-Odor Disinfectant by Industrial Bio-Test Laboratories inc. Project No. 76-05. June 8, 1976.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	1847958-8
<b>Method/guideline</b>	Acute oral toxicity was determined in rats

<b>Test Type</b>	Acute Oral LD50 study
<b>GLP</b>	NG
<b>Year</b>	1973
<b>Species/strain</b>	Wistar rat
<b>Sex</b>	Male
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	No other details given.
<b>Value LD50 or LC50 with confidence limits</b>	Oral LD50 = 3600 mg/kg (C. I., 3000-4200 mg/kg)
<b>Number of deaths at each dose level</b>	2560 mg/kg, 1/10 deaths; 3200 mg/kg, 5/10 deaths; 4000 mg/kg, 4/10 deaths; 5000 mg/kg, 10/10 deaths.
<b>Remarks for Results</b>	Lethargy occurred in all groups
<b>Conclusion Remarks</b>	Acute oral LD50 = 3600 mg/kg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Moreno O. M. (1973) Dihydromyrcenol. Acute toxicity studies in rats and rabbits. Unpublished report to RIFM.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	543-39-5
<b>Method/guideline</b>	Acute oral toxicity was determined in rats
<b>Test Type</b>	Acute Oral LD50 study
<b>GLP</b>	Ng
<b>Year</b>	1972
<b>Species/strain</b>	Wistar rat
<b>Sex</b>	Male
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	No other details given.
<b>Value LD50 or LC50 with confidence limits</b>	Oral LD50 = 5300 mg/kg (C. I., 4500-6100 mg/kg)
<b>Number of deaths at each dose level</b>	4000 mg/kg, 2/10 deaths; 5,000 mg/kg, 4/10 deaths; 6250 mg/kg, 7/10 deaths; 7,800 mg/kg, 10/10 deaths.
<b>Conclusion Remarks</b>	Acute LD50 study

**Data Qualities Reliabilities** Reliability code 1. Reliable without restrictions.

**References** Moreno O. M. (1972b) Acute toxicity studies in rats and rabbits. Unpublished report to RIFM.

<b>Substance Name</b>	Hercules 70L Pine Oil Disinfectant, transparent yellow liquid
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Acute oral Lethal Dose (LD50) was determined using the techniques of Weil, Thompson, and Thompson and Weil.
<b>Test Type</b>	Acute Oral Lethal Dose
<b>GLP</b>	NG
<b>Year</b>	1976
<b>Species/strain</b>	Albino Sprague Dawley rats
<b>Sex</b>	Female
<b># of animals per sex per dose</b>	4
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	Doses tested 1350, 2025, 3038, 4556, 15380 mg/kg. All doses were administered directly into the stomachs of the rats using a hypodermic syringe equipped with a ball-tipped intubating needle.
<b>Value LD50 or LC50 with confidence limits</b>	Acute Oral LD50 = 4118 mg/kg +/- 418.1 mg/kg
<b>Number of deaths at each dose level</b>	At 1350 mg/kg-0/4; at 2025 mg/kg-0/4; at 3038 mg/kg 0/4; at 4556 mg/kg-3/4, at 15380 mg/kg-4/4.
<b>Remarks for Results</b>	No gross pathological alterations. Hemorrhage in stomach lining at 4556 mg/kg; white mucoid substance in GI tract at 15380 mg/kg.
<b>Conclusion Remarks</b>	The acute oral LD50 value of Hercules 70L Pine oil Disinfectant in the female Sprague Dawley rats was calculated to be 4118 mg/kg +/- 418.1 mg/kg. The substance was classified as slightly toxic.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Baker R. G., Mastri, C. W., Kinoshita, F. K and Keplinger, M. L. (1976) Report to Hercules Inc. on Acute Toxicity Studies with Hercules 70L Pine Oil Disinfectant by Industrial Bio-Test Laboratories, Inc. Project No. 76-05. Dated June 8, 1976.

<b>Substance Name</b>	Riser Pine oil (5060% alpha-terpineol, IO-I 5% other terpineol isomers, 7-15% terpene secondary alcohols and IO-20% terpene hydrocarbons)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Approximate Lethal Dose (A.L.D.) was studied by the Deichman and LaBlanc method.

<b>Test Type</b>	Approximate Lethal Dose Test
<b>GLP</b>	Pre GLP
<b>Year</b>	1961
<b>Species/strain</b>	Rats
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	3
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	Test material was administered undiluted by stomach tube in a single oral dose (ranging from 0.94 g/kg to 7.1 g/kg). The surviving animals were scarified after a period of 2 weeks. A post mortem examination was performed on each of the rats.
<b>Value LD50 or LC50 with confidence limits</b>	The Approximate Lethal Oral dose of Riser Pine oil in the rat was found to be 2.1 g/kg. This dose killed in 54 hours.
<b>Number of deaths at each dose level</b>	Dose 0.94 & 1.4 g/kg-survived; at 2.1 g/kg died in 54 hrs; at 3.2 g/kg died in 36 hrs; at 4.7 & 7.1 died in 12 hours.
<b>Remarks for Results</b>	Signs of intoxication: loss of equilibrium, decreased sensitivity to pain, increased salivation and muscular weakness, increased secretion of lacrimal fluid, occasional clonic convulsions, loss of body wt., hyperemic lungs, hemorrhagic atelectasis, etc.
<b>Conclusion Remarks</b>	The Approximate Lethal Oral dose of Riser Pine oil in the rat was found to be 2.1 g/kg. This dose killed in 54 hours.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Deichmann, W. B. (1961) Report on the Determination of the Approximate Lethal Oral Dose in the Rat of Compounds Submitted by the Hercules Powder Co.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Acute dermal toxicity in the rabbit
<b>Test Type</b>	Acute Dermal LD50 study
<b>GLP</b>	<b>NG</b>
<b>Year</b>	<b>1972</b>
<b>Species/strain</b>	Rabbit
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	3
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	No other details given.

<b>Value LD50 or LC50 with confidence limits</b>	Acute Dermal LD50 was determined to be greater than 5,000 mg/kg
<b>Number of deaths at each dose level</b>	No deaths at 5000 mg/kg dose.
<b>Conclusion Remarks</b>	Dermal LD50 was determined to be >5000 mg/kg.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Moreno O. M. (1972a) Linalyl acetate. Acute toxicity studies in rats and rabbits. Unpublished report to RIFM.

<b>Substance Name</b>	Myrcenol
<b>CAS No.</b>	<b>543-39-5</b>
<b>Method/guideline</b>	Acute dermal toxicity in the rabbit
<b>Test Type</b>	Acute Dermal <b>LD50</b> study
<b>GLP</b>	NG
<b>Year</b>	1972
<b>Species/strain</b>	White New Zealand rabbit
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	No other details given.
<b>Value LD50 or LC50 with confidence limits</b>	Acute dermal LD50 was determined to be greater than 5,000 mg/kg
<b>Number of deaths at each dose level</b>	No deaths at highest dose.
<b>Conclusion Remarks</b>	Dermal LD50 was determined to be >5,000 mg/kg.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Moreno O. M. (1972b) Acute toxicity studies in rats and rabbits, Unpublished report to RIFM.

<b>Substance Name</b>	Dihydromyrcenol
<b>CAS No.</b>	18479-58-8
<b>Method/guideline</b>	Acute dermal toxicity in the rabbit
<b>Test Type</b>	Acute dermal LD50 study
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1973

<b>Species/strain</b>	White New Zealand rabbit
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	No other details given.
<b>Value LD50 or LC50 with confidence limits</b>	Acute Dermal LD50 was determined to be greater than 5,000 mg/kg
<b>Number of deaths at each dose level</b>	No deaths at highest dose.
<b>Conclusion Remarks</b>	Dermal LD50 was determined to be >5000 mg/kg.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Moreno O. M. (1973) Dihydromyrcenol. Acute toxicity studies in rats and rabbits. Unpublished report to RIFM.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The study was performed to assess the acute oral toxicity of Plinyl acetate in Sprague-Dawley CD strain rats. The method followed the OECD Guidelines for Testing of chemicals No 401 "Acute Oral Toxicity" and Method B1 of Commission Directive 92/69/EEC
<b>Test Type</b>	Acute Oral LD50
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/strain</b>	Sprague-Dawley CD Strain rats
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	5 male and 5 female rats were dosed (2000 mg/kg; dose volume 2.11 ml/kg) the animals were observed for deaths or overt signs of toxicity for 112, 1, 2 & 4 hours after dosing and subsequently once daily for 14 days.
<b>Value LD50 or LC50 with confidence limits</b>	>2000mg/kg
<b>Number of deaths at each dose level</b>	No deaths reported
<b>Remarks for Results</b>	There were no deaths or clinical signs of systemic toxicity. All animals showed an expected body weight gain and no abnormalities were noted at necropsy

<b>Conclusion Remarks</b>	The acute oral median lethal dose (LD50) of Plinyl acetate in the Sprague-Dawley CD strain rat was found to be greater than 2000mg/kg.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliabilities Remarks</b>	Study was conducted in compliance with UK GLP and methodology recommended in the OECD Guidelines for testing of Chemicals No.401 "Acute Oral Toxicity" and Method B1 of Commission Directive 92/69/EEC.
<b>References</b>	Sanders A. (1997). Plinyl Acetate: Acute Oral Toxicity (Limit Test) in the Rats. Safepharm Laboratories Ltd. SPL Project No. 1044/004.

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Litchfield-Wilcoxon, 1949 (FDA study)
<b>Test Type</b>	Acute Oral LD50 study
<b>GLP</b>	NG
<b>Year</b>	1964
<b>Species/strain</b>	Osborne-Mendel rat
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Route of Administration</b>	Intubation
<b>Remarks for Test Conditions</b>	Five male and five female young adult Osborne-Mendel rats were fasted for 18 hours prior to treatment. Two weeks thereafter, animals were observed.
<b>Value LD50 or LC50 with confidence limits</b>	Oral LD50= 5075 mg/kg (C. I., 4160-6190 mg/kg)
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	Animals exhibited depression, scrawny appearance, and porphyrin-like deposits around eyes and nose
<b>Conclusion Remarks</b>	Oral LD50 was determined to be 5075 mg/kg for rats.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliabilities Remarks</b>	Data appeared in peer-reviewed journal.
<b>References</b>	Jenner P. M., Hagan, E.C., Taylor, J.M., Cook, E.L. and Fitzhugh, O.G. (1964) Food Flavorings and Compounds of Related Structure I. Acute Oral Toxicity. Food and Cosmetics Toxicology 2:327-343.
<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)

<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The method used test the Acute Inhalation Toxicity Potential of the Plinyl acetate followed the recommendation of the OECD Guidelines for Testing Chemicals No. 403 "Acute Inhalation Toxicity of Chemicals" and method B2 of Commission Directive 92/69/EEC.
<b>Test Type</b>	Acute Inhalation Toxicity
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/strain</b>	Sprague-Dawley CD rats
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Route of Administration</b>	Inhalation (nose only)
<b>Remarks for Test Conditions</b>	Each rat was individually held in a tapered polycarbonate restraining tube fitted onto a single tier of the exposure chamber. Only the nose of each animal was exposed (5 mg/l) to the test atmosphere for 4 hours
<b>Value LD50 or LC50 with confidence limits</b>	LC50 > 5.03 mg/l
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	No deaths, wet fur, hunched posture and pilo-erection, occasional labored &/or noisy respiration, ptosis, tiptoe gait and red/brown staining of the fur and around the eyes snout and head was observed. Dark areas or dark foci on the lungs were seen.
<b>Conclusion Remarks</b>	The acute inhalation median lethal concentration of the Plinyl acetate in the Sprague-Dawley CD rat was > 5.03 mg/l.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliabilities Remarks</b>	Study was performed in compliance with UK GLP standard. These regulations are also in accordance with GLP standards published by OECD monograph No. 45 (OCDE/GD(92)32).
<b>References</b>	Blagden S. M. (1997). Plinyl Acetate: Acute Inhalation Toxicity (Nose Only) Study in the Rat. Safepharm Laboratories Ltd. SPL Project No. 10441017.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Litchfield-Wilcoxon, 1949 (FDA study)
<b>Test Type</b>	Acute Oral LD50 study
<b>GLP</b>	NG
<b>Year</b>	1964

<b>Species/strain</b>	Osborne-Mendel rat
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Route of Administration</b>	Intubation
<b>Remarks for Test Conditions</b>	Five male and five female young adult Osborne-Mendel rats were fasted for 18 hours prior to treatment. Two weeks thereafter, animals were observed.
<b>Value LD50 or LC50 with confidence limits</b>	Oral LD50 = 2790 mg/kg (C. I., 2440-3180 mg/kg)
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	Animals expired between 4 and 18 hours after dose.
<b>Conclusion Remarks</b>	Oral LD50 was determined to be 2790 mg/kg for rats.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliabilities Remarks</b>	Data appeared in peer-reviewed journal.
<b>References</b>	Jenner, P.M., Hagan, E.C., Taylor, J.M, Cook, E.L. and Fitzhugh, O.G. (1964). Food Flavorings and Compounds of Related Structure I. Acute Oral Toxicity. Food and Cosmetics Toxicology 2:327-343.

<b>Substance Name</b>	Tetrahydrolinalool
<b>CAS No.</b>	78-69-3
<b>Method/guideline</b>	Acute oral toxicity was determined in rats
<b>Test Type</b>	Acute LD50 study
<b>GLP</b>	NG
<b>Year</b>	1976
<b>Species/strain</b>	Rats
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	No other details given.
<b>Value LD50 or LC50 with confidence limits</b>	Oral LD50 was found to be greater than 5.0 g/kg
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	Symptomatology: lethargy

<b>Conclusion Remarks</b>	Oral LD50 was found to be greater than 5.0 g/kg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Moreno O.M. (1976) Acute toxicity studies in rats, mice, rabbits and guinea pigs. Unpublished report to RIFM.

<b>Substance Name</b>	Tetrahydrolinalool
CAS No.	78-69-3
<b>Method/guideline</b>	Acute Dermal Toxicity in rabbit was determined.
<b>Test Type</b>	Acute Dermal LD50 study
<b>GLP</b>	NG
<b>Year</b>	1976
<b>Species/strain</b>	Rabbits
Sex	Not reported
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	No other details given
<b>Value LD50 or LC50 with confidence limits</b>	Dermal LD50 was found to be greater than 5 g/kg
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	9/10 animals had moderate redness and moderate edema. 1/10 had slight redness and slight edema.
<b>Conclusion Remarks</b>	Dermal LD50 was found to be greater than 5 g/kg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Moreno O. M. (1976) Acute toxicity studies in rats, mice, rabbits and guinea pigs. Unpublished report to RIFM.

<b>Substance Name</b>	Linalyl acetate
CAS No.	115-95-7
<b>Method/guideline</b>	Litchfield-Wilcoxon, 1949 (FDA study)
<b>Test Type</b>	Acute Oral LD50 study
<b>GLP</b>	NG
<b>Year</b>	164
<b>Species/strain</b>	Osborne-Mendel rat

Sex	Male and Female
# of animals per sex per dose	5
Route of Administration	Intubation
Remarks for Test Conditions	Five male and five female young adult Osborne-Mendel rats were fasted for 18 hours prior to treatment. Two weeks thereafter, animals were observed.
Value LD50 or LC50 with confidence limits	Oral LD50= 14,550 mg/kg (C. I., 12,300-17,170 mg/kg)
Number of deaths at each dose level	Not given
Remarks for Results	Animals showed depression soon after treatment, coma and wet posterior.
Conclusion Remarks	Oral LD50 was determined to be 14,550 mg/kg for rats.
Data Qualities Reliabilities	Reliability code 1. Reliable without restrictions.
Data Reliabilities Remarks	Data appeared in peer-reviewed journal.
References	Jenner P.M., Hagan, EC., Taylor, J.M. Cook, E.L. and Fitzhugh, O.G. (1964) Food Flavorings and Compounds of Related Structure I. Acute Oral Toxicity. Food and Cosmetics Toxicology 2:327-343.

<b>Substance Name</b>	Tetrahydrolinalool
<b>Test Type</b>	Acute Dermal Toxicity study
<b>GLP</b>	NG
<b>Year</b>	1976
<b>Species/strain</b>	Rabbits
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Dermal
<b>Value LD50 or LC50 with confidence limits</b>	Acute Dermal LD50 was determined to be greater than 5.00 g/kg
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	Moderate redness and edema was noted in 9 out of 10 rabbits. Slight redness and edema was noted in one rabbit.
<b>Conclusion Remarks</b>	Acute dermal LD50 of tetrahydrolinalool was determined to be greater than 5.00 g/kg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Moreno. O. M. (1976). Report to RIFM by M.B. Research Laboratories, Inc.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Acute oral toxicity was determined in male Cd-I mice.
<b>Test Type</b>	Acute Oral LD50 study
<b>GLP</b>	NG
<b>Year</b>	1985
<b>Species/strain</b>	CD-I mouse
<b>Sex</b>	Male
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	Animals were assessed for 7 days post-treatment,
<b>Value LD50 or LC50 with confidence limits</b>	Oral LD50 = 2830 mg/kg (95% C. I., 2290-3497 mg/kg)
<b>Number of deaths at each dose level</b>	Not given
<b>Conclusion Remarks</b>	Oral LD50 = 2830 mg/kg
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Yamahara J., Kimura H., Kobayashi M., Okamoto T., Sawada T., Fujimura H., and Chistaka T. (1985) Chologogic action and characteristics of (+/-)-alpha-terpineol-beta-D-O-glucopyranoside, a new monoterpenoid glucoside. Chem. and Pharm. Bulletin. 33(4) 1669.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Approximate Lethal Dose (A.L.D.) was studied by the Deichman and LaBlanc method.
<b>Test Type</b>	Approximate Lethal Dose Test
<b>GLP</b>	NG
<b>Year</b>	1961
<b>Species/strain</b>	Rats
<b>Sex</b>	Not reported
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	Test material was administered undiluted by stomach tube in a single oral dose. The surviving animals were scarified after a period of 2 wks. Post mortem examination were performed.

<b>Value LD50 or LC50 with confidence limits</b>	The Approximate Lethal Oral dose of alpha-Terpineol in the rat was found to be 3.2 g/kg.
<b>Number of deaths at each dose level</b>	Not given
<b>Conclusion Remarks</b>	The Approximate Lethal Oral dose of alpha-Terpineol in the rat was found to be 3.2 g/kg.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Deichmann W. B. (1961) Report on the determination of the approximate lethal oral dose in the rat of compounds. Submitted by the Hercules Powder Co.

<b>Substance Name</b>	20% Pine Oil disinfectant (pine oil contained 76.7% alpha- and gamma-terpineol)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	The study was designed to comply with the standards set forth by EPA proposed Guidelines for Test Procedures (1982), Subsection F. Series 81-1.
<b>Test Type</b>	Acute Oral toxicity study
<b>GLP</b>	Yes
<b>Year</b>	1985
<b>Species/strain</b>	Wistar Albino rats
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	5 male & 5 female rats were dosed by gavage with 20% Pine oil disinfectant in water at 5.0 g/kg and observed for 1, 2, & 4 hrs post dose and twice daily for 14 days.
<b>Value LD50 or LC50 with confidence limits</b>	LD50 was calculated to be greater than 5.0 g/kg (LD50 > 1g/kg for pine oil undiluted.).
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	Symptoms: Lethargy, ataxia, piloerection, chromorhinorrhea, chromodacryorrhea, ptosis and brown staining of anogenital area. Body weight /necropsy normal.
<b>Conclusion Remarks</b>	According to EPA criteria, 20% Pine oil Disinfectant was considered to be Category IV.
<b>Data Qualities Reliabilities</b>	Reliability code 1, Reliable without restrictions.
<b>Data Reliabilities Remarks</b>	Study was conducted in accordance with the GLP regulations of the FDA effective 06/20/79 and EPA effective 11/29/83.
<b>References</b>	Cerven B., Moreno, O. M. and Althenbach, E. (1985a) Single Dose Oral Toxicity in Rats (20% Pine Oil Disinfectant. Report to Hercules Inc. by M. B. Research Laboratories. Project No. MB 85-7546 A.

<b>Substance Name</b>	20% Pine Oil Disinfectant, Pale Yellow Liquid (pine oil contained 76.7% alpha- and gamma-terpineol)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	The study was designed to comply with the standards set forth by EPA proposed Guidelines for Test Procedures (1982), subsection F. Series 81-2.
<b>Test Type</b>	Acute Dermal LD50 study
<b>GLP</b>	Yes
<b>Year</b>	1985
<b>Species/strain</b>	Albino New Zealand rabbits
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	2.0 g/kg of the test material was applied to the shaved dermal sites on the back of health rabbits (5 per sex). The test material was kept in contact with the skin for 24 hours. Dermal responses were recorded on day 1, 7, and 14.
<b>Value LD50 or LC50 with confidence limits</b>	The Dermal LD50 was determined to be greater than 2.0 g/kg (LD50>0.4 g/kg as pine oil undiluted.)
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	Dermal reaction was moderate on day 1, severe on Day 7 and absent to slight on Day 14. An instance of diarrhea decreased fecal output and yellow nasal discharge were noted.
<b>Conclusion Remarks</b>	The Dermal LD50 was determined to be greater than 2.0 g/kg. According to EPA criteria, 20% Pine oil disinfectant was considered to be Category III.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliabilities Remarks</b>	Study was conducted in accordance with the GLP regulations of the FDA effective 06/20/79 and EPA effective 11/29/83.
<b>References</b>	Cerven B., Moreno, O. M. and Althenbach, E. (1985b). Acute Dermal Toxicity in Rabbits (20% Pine Oil Disinfectant. Report to Hercules Inc. by M. B. Research Laboratories. Project No. MB 85-7546 B.

<b>Substance Name</b>	Hercules 70L Pine Oil Disinfectant, transparent yellow liquid
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Acute dermal Lethal Dose (LD50) was determined using the techniques of Weil, Thompson, and Thompson and Weil.
<b>Test Type</b>	Acute Dermal Toxicity
<b>GLP</b>	Ng
<b>Year</b>	1976

<b>Species/strain</b>	White New Zealand rabbits
<b>Sex</b>	Male
<b># of animals per sex per dose</b>	4
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	Test material (3000 mg/kg) was applied to the shaved area on the back of rabbits and occluded for 24 hrs. After 24 hrs, test material was removed and animals were observed for local skin reactions for 14 days.
<b>Value LD50 or LC50 with confidence limits</b>	The Dermal LD50 value was calculated to be greater than 3000 mg/kg.
<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	Severely irritating to the skin of rabbits, beet red erythema, severe edema and 2nd degree bum. Escharosis was observed at the test skin sites at 7 & 14 days. Hemorrhaged lungs and gastroenteritis was also seen in one rabbit each.
<b>Conclusion Remarks</b>	The Dermal LD50 value for Hercules 70L Pine Oil Disinfectant was calculated to be greater than 3000 mg/kg. The test material was considered to be practically nontoxic.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Baker R. G., Mastri, C. W., Kinoshita, F. K and Keplinger, M. L. (1976) Report to Hercules Inc. on Acute Toxicity Studies with Hercules 70L Pine Oil Disinfectant by Industrial Bio-Test Laboratories, Inc. Project No. 76-05. Dated June 8, 1976.

<b>Substance Name</b>	Pine Oil (100%), clear liquid (62.7% alpha-terpineol, 83.4% total terpene alcohol content)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	The test material was tested for acute inhalation toxicity in conformity with the EPA Guidelines. Five male and five female rats (Sprague-Dawley) were exposed for 4 hours to dynamic atmosphere containing a nominal concentration of 9.2-mg/L air.
<b>Test Type</b>	Acute Inhalation toxicity
<b>GLP</b>	Yes
<b>Year</b>	1994
<b>Species/strain</b>	Sprague Dawley rat
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	5
<b>Route of Administration</b>	Inhalation
<b>Remarks for Test Conditions</b>	The actual concentration measured in the breathing zone thrice during the exposure was 3.98 mg/l, 3.92 mg/l and 3.48 mg/l.
<b>Value LD50 or LC50 with confidence limits</b>	The LC50 > 3.79 mg/L.

<b>Number of deaths at each dose level</b>	Not given
<b>Remarks for Results</b>	After exposure, all rats appeared wet and common signs were depressed activity, yellow abdominal and perineal staining (urine), and frequent chromorhinorrhea, one female showed ataxia and also had nasal discharge.
<b>Conclusion Remarks</b>	The LC50 was greater than 3.79 mg/L. The EPA toxicity indicator for Pine oil is best described as Category III/IV, No mortality at 3.79 mg/L.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Data Reliabilities Remarks</b>	Study was conducted in accordance with EPA Guidelines.
<b>References</b>	Robbins G. R. (1994) Acute Inhalation Toxicity Study in Rats. Pine Oil. Cosmopolitan Safety Evaluation, Inc. Report to CSMA Pine Oil Joint Venture. Lab Project ID/Study Number C3371.

## 4.2 Genetic Toxicity *In Vitro*

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Linalyl acetate was tested in the rat hepatocytes for unscheduled DNA synthesis
<b>Test Type</b>	Unscheduled DNA Synthesis
<b>System of Testing</b>	Rat hepatocytes
<b>GLP</b>	NG
<b>Year</b>	1989
<b>Species/Strain</b>	Sprague-Dawley or Fischer Rat
<b>Metabolic Activation</b>	None
<b>Doses/Concentration</b>	300 ug
<b>Remarks for Test Conditions</b>	The Hepatocytes were isolated from adult male rats and incubated for 18-20 hours with the test material in serum-free WME medium containing 5-10uCi/ml 3h-TdR. UDS was measured by electronically counting nuclear grains and subtracting the average number of grains in 3 adjacent nuclear-sized cytoplasmic areas.
<b>Results</b>	Linalyl acetate gave a negative response.
<b>Genotoxic effects</b>	None
<b>Conclusion Remarks</b>	Linalyl acetate was not shown to be clastogenic under the test conditions.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.

**Remarks for Data Reliability** The study was published in a peer-reviewed journal The Toxicologist.

**References** Heck J.D., Vollmuth, T.A., Cifone, M.A, Jagannath, D.R., Myhr, B. and Curren, R.D. (1989) An Evaluation of Food Flavoring Ingredients in a Genetic Toxicity Screening. The Toxicologist Vol 9 NO. 1.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	<b>78-70-6</b>
<b>Method/guideline</b>	The Mouse Lymphoma L5178y TK +/- cell Mutagenesis (MLY) assay was performed according to the procedures described by Clive et al. (1979).
<b>Test Type</b>	Mouse Lymphoma Forward Mutation Assay (MLY).
<b>System of Testing</b>	L5178y Mouse Lymphoma cell line
<b>GLP</b>	NG
<b>Year</b>	1989
<b>Species/Strain</b>	Mice
<b>Metabolic Activation</b>	Aeroclor 1254-induced rat liver S-9 activation system.
<b>Doses/Concentration</b>	3.9 nl/ml -200 nl/ml +S-9: 3.9150 nl/ml, -S-9
<b>Results</b>	Linalool was weakly positive in the presence of rat liver S-9 activation system.
<b>Remarks for Results</b>	At concentration range of 100 - 300 nl/ml linalool treatment resulted in 2.3-4.1 fold increase.
<b>Conclusion Remarks</b>	Linalool was weakly positive in the absence of rat liver S-9 activation system. However, these finding should be viewed with caution in light of the known effect of nonphysiological medium conditions on the outcome of such assays.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal The Toxicologist.
<b>References</b>	Heck J.D., Vollmuth, T.A., Cifone, M.A., Jagannath, D.R., Myhr, B. and Curren, R.D. (1989) An Evaluation of Food Flavoring Ingredients in a Genetic Toxicity Screening. The Toxicologist Vol 9 NO. 1.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	<b>125252-49-5</b>
<b>Method/guideline</b>	Ames
<b>Test Type</b>	Reverse mutation
<b>System of Testing</b>	Bacterial

<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/Strain</b>	Salmonella typhimurium TA 100, TA 98, TA 1535, TA 1537 & TA 1538.
<b>Metabolic Activation</b>	Rat liver <b>microsome</b> fraction <b>S9</b> from Aroclor induced rats.
<b>Doses/Concentration</b>	5 -5000 microliters per plate.
<b>Statistical Methods</b>	All data were analyzed by method recommended by the UKEMS and normally <b>Dunnett's</b> method of linear regression was used.
<b>Remarks for Test Conditions</b>	After 48-hour incubation at 37 °C, each assay plate was counted. Routine positive control plates were prepared: <b>N-ethyl-N-nitro-N-nitrosoguanidine</b> for TA100 & TA1535; <b>9-Aminoacridine</b> for TA1537; <b>4-Nitro-o-phenylenediamine</b> for TA 98; <b>4-Nitroquinoline-1-oxide</b> for TA98 & <b>2-Aminoanthracene</b> which is non-mutagenic in the absence of <b>S9</b> was used.
<b>Results</b>	No biologically significant increase in revertant colonies was observed with any bacterial strains and any dose of the test material either with or without metabolic activation.
<b>Cytotoxic concentration</b>	500 ug/plate
<b>Genotoxic effects</b>	None
<b>Statistical results</b>	NG
<b>Remarks for Results</b>	A slight increase in revertant colonies was observed with tester strain TA98 with metabolic activation only, but the response did not achieve a <b>2-fold</b> increase over the concurrent solvent controls and non-reproducible in three experiments, of which included a pre-incubation method and a tightened test material doserange.
<b>Conclusion Remarks</b>	No evidence of mutagenicity.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions. Data were acquired prior to GLP or OECD guidelines but were obtained by standard methodology and published in a peer reviewed journal.
<b>Remarks for Data Reliability</b>	The study was performed in compliance with UK GLP standard and is in accordance with GLP standards published as OECD Monograph No. 45 (OCDE/GD(92)32).
<b>References</b>	Thompson P. W. (1997) Plinyl Acetate: Reverse Mutation Assay "Ames Test" Using Salmonella Typhimurium. Safepharm Laboratories Ltd. SPL Project Number 1044/010.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance • plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The method used followed that described in the OECD Guidelines for testing Chemicals (1981) No. 473 "Genetic toxicology: Chromosome Aberration Test" and Method B10 of Commission Directive 92/69/EEC.

<b>Test Type</b>	Chromosomal Aberration
<b>System of Testing</b>	In vitro human lymphocytes
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/Strain</b>	Human volunteer whole blood
<b>Metabolic Activation</b>	S9 Fraction prepared from male rat liver treated with single (500 mg/kg, ip) Aroclor 1254 injection
<b>Doses/Concentration</b>	15.31, 30.63, 61.25, 122.5, 245, 490, 980, & 1960 ug/ml
<b>Statistical Methods</b>	Frequency of cells with aberration and frequency of polyploid was compared with the concurrent vehicle control using Fischer's Exact test or Chi-squared test.
<b>Remarks for Test Conditions</b>	Four treatment conditions were used in 4 hrs exposures with the addition of S9 from rat liver at 10% in standard cofactors with cell harvest after 16 & 40 hr expression periods and a 20 & 44 hr continuous exposure in the absence of activation. In experiment 1 the dose range for evaluation was selected from series of 8 dose levels on the basis of toxicity.
<b>Results</b>	All vehicle controls gave frequencies of cells with aberration within the range expected for normal human lymphocytes. All the positive control treatments gave significant increases in the frequency of cells with aberration indicating the satisfactory performance of the test and of the activity of the metabolizing system.
<b>Conclusion Remarks</b>	Plinyl acetate was shown to be non-clastogenic to human lymphocytes in vitro, in presence or absence of liver enzyme metabolizing system.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was performed in compliance with UK GLP standard and is in accordance with GLP standards published as OECD Monograph No. 45 (OCDE/GD(92)32).
<b>References</b>	Wright N. P. (1997) Plinyl Acetate: chromosome aberration test in human lymphocytes in vitro. Safepharm Laboratories Ltd. SPL Project No. 1044101 1.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Test material was assayed for mutagenicity toward Salmonella typhimurium (TA 98, TA 100, TA 1535, & TA 1537) in the presence or absence of Liver S9 fraction from Aroclor 1250 treated Sprague-Dawley rats
<b>Test Type</b>	Reverse mutation (Ames test)
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG

<b>Year</b>	1980
<b>Species/Strain</b>	Bacteria; Salmonella typhimurium Strain TA98, TA1 00, TA1 535 & TA1537
<b>Metabolic Activation</b>	Liver S9 fraction from Aroclor 1250 treated Sprague-Dawley rats
<b>Doses/Concentration</b>	0.03, 0.3, 3.0 & 30.0 umol/plate.
<b>Results</b>	alpha-Terpineol was not found to be mutagenic.
<b>Cytotoxic concentration</b>	NA
<b>Genotoxic effects</b>	None
<b>Conclusion Remarks</b>	alpha-Terpineol was not found to be mutagenic.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal The Toxicologist.
<b>References</b>	Florin I., Rutberg, L, Curvall, M, and Enzell, C. R. (1980) Screening of Tobacco Smoke Constituents for Mutagenicity Using the Ames' Test. Toxicology, 18, 219-232.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Reverse mutation assays using <i>S. typhimurium</i> strains: TA92, TA1535, TA 100, TA 1537 & TA 98 were used according to the method of Ames, The S-9 was prepared from the Fischer rat liver treated with polychlorinated biphenyl (500 mg/kg).
<b>Test Type</b>	Reverse mutation (Ames test)
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1984
<b>Species/Strain</b>	Bacteria: <i>Salmonella typhimurium</i> strains: TA92, TA1 535, TA 100, TA 1537 & TA 98
<b>Metabolic Activation</b>	The S-9 was prepared from the Fischer rat liver treated with polychlorinated biphenyl (500 mg/kg).
<b>Doses/Concentration</b>	Maximum concentration 1.0 mg/plate
<b>Remarks for Test Conditions</b>	The number of revertant (his+) colonies was scored after incubation at 37 C for 2 days. Vehicle: DMSO.
<b>Results</b>	No significant increase in the number of revertant colonies was observed at maximum concentration.
<b>Cytotoxic concentration</b>	NG
<b>Genotoxic effects</b>	No genotoxic effect
<b>Conclusion Remarks</b>	No genotoxic effects of Linalool were observed at maximum concentration of 1.0 mg/plate in Reverse mutation assay (Ames Test)

**Data Qualities Reliabilities** Reliability code 1, Reliable without restrictions.

**Remarks for Data Reliability** The study was published in a peer-reviewed journal Food. Chem. Toxic.

**References** Ishidate M, Sofuni, K., Yoshikawa, M., Nohmi, T., **Sawada, M** and Matsuoka, A. (1984) Primary Mutagenicity Screening of Food Additives Currently Used in Japan. FD. Chem. Tox. **22(8)**, 623-636.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Chromosomal Aberration test was carried out using a Chinese hamster fibroblast cell line. The cells were exposed to Linalool at three different concentrations for 24 & 48 hrs. No metabolic activation system was applied.
<b>Test Type</b>	Chromosomal Aberration test
<b>System of Testing</b>	Chinese hamster fibroblast cell line
<b>GLP</b>	NG
<b>Year</b>	1984
<b>Species/Strain</b>	Hamster: Chinese hamster fibroblast cell line.
<b>Remarks for Test Conditions</b>	The incidence of polyploid cells as well as of structural chromosomal aberrations such as chromatid or chromosomal gaps, breaks, exchanges, ring formation, fragmentations and others, was recorded on each plate. Solvent treated/untreated cells served as negative control. Vehicle: DMSO
<b>Results</b>	Negative chromosomal test.
<b>Statistical results</b>	Results negative if the incidence was < 4.9%
<b>Remarks for Results</b>	The incidence of polyploid cells as well as of structural chromosomal aberrations such as chromatid or chromosomal gaps, breaks, exchanges, ring formation, fragmentations and others was less than 4.9%.
<b>Conclusion Remarks</b>	Linalool is not considered to be clastogenic under the test conditions.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal Fd. Chem. Toxic.
<b>References</b>	Ishidate M, Sofuni, K., Yoshikawa, M., Nohmi, T., <b>Sawada, M</b> and Matsuoka, A. (1984) Primary Mutagenicity Screening of Food Additives Currently Used in Japan. FD. Chem. Tox. <b>22(8)</b> , 623-636.
<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6

<b>Method/guideline</b>	Mutagenic effects of Linalool was tested in 5 tester strains of Salmonella typhimurium TA 1535, TA 1537, TA1538, TA 98 & TA 100 in the presence or absence of S-9 fraction of aroclor 1254 induced male Sprague-Dawley rat liver.
<b>Test Type</b>	Reverse mutation (Ames test)
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1989
<b>Species/Strain</b>	Bacteria: Salmonella typhimurium TA 1535, TA 1537, TA1538, TA 98 & TA 100
<b>Metabolic Activation</b>	S-9 fraction of Aroclor 1254 induced male Sprague-Dawley rat liver.
<b>Doses/Concentration</b>	10,000 ug/plate
<b>Results</b>	Linalool was inactive in Salmonella strains TA 1535, TA 1537, TA 1538, TA 98 & TA 100 both in the presence and absence of metabolic activation system.
<b>Genotoxic effects</b>	No genotoxic effect
<b>Conclusion Remarks</b>	Linalool was not found to be mutagenic in Reverse mutation assay (Ames' test) under the condition of this study.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal The Toxicologist.
<b>References</b>	Heck J.D., Vollmuth, T.A., Cifone, M.A., Jagannath, D.R., Myhr, B. and Curren, R.D. (1989) An Evaluation of Food Flavoring Ingredients in a Genetic Toxicity Screening. The Toxicologist Vo19No.1.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Mutagenic effects of Linalyl acetate was tested in 5 tester strains of Salmonella typhimurium TA 1535, TA 1537, TA1538, TA 98 & TA 100 in the presence or absence of S-9 fraction of Aroclor 1254 induced male Sprague-Dawley rat liver.
<b>Test Type</b>	Reverse mutation (Ames test)
<b>System of Testing</b>	Bacterial
<b>GLP</b>	<b>NG</b>
<b>Year</b>	<b>1989</b>
<b>Species/Strain</b>	Bacteria: Salmonella typhimurium TA 1535, TA 1537, TA1538, TA 98 & TA 100
<b>Metabolic Activation</b>	S-9 fraction of Aroclor 1254 induced male Sprague-Dawley rat liver.
<b>Doses/Concentration</b>	25,000 ug/plate

<b>Results</b>	Linalyl acetate was inactive in Salmonella strains TA 1535, TA 1537, TA 1538, TA 98 & TA 100 both in the presence and absence of metabolic activation system.
<b>Genotoxic effects</b>	No genotoxic effect
<b>Conclusion Remarks</b>	Linalyl acetate was not found to be mutagenic under the test condition.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal The Toxicologist.
<b>References</b>	Heck J.D., Vollmuth, T.A., Cifone, M.A., Jagannath, D.R., Myhr, B. and Curren, R.D. (1989) An Evaluation of Food Flavoring Ingredients in a Genetic Toxicity Screening. The Toxicologist Vol 9 NO. 1.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Linalool was tested in the rat hepatocytes for unscheduled DNA synthesis
<b>Test Type</b>	Unscheduled DNA Synthesis
<b>System of Testing</b>	Rat hepatocytes
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1989
<b>Species/Strain</b>	Sprague-Dawley Rat or Fischer Rat
<b>Metabolic Activation</b>	None
<b>Doses/Concentration</b>	<b>5oug</b>
<b>Remarks for Test Conditions</b>	The Hepatocytes were isolated from adult male rats and incubated for 18-20 hours with the test material in serum-free WME medium containing 5-10uCi/ml 3h-TdR. UDS was measured by electronically counting nuclear grains and subtracting the average number of grains in 3 adjacent nuclear-sized cytoplasmic areas.
<b>Results</b>	Linalool gave a negative response.
<b>Genotoxic effects</b>	None
<b>Conclusion Remarks</b>	Linalool was not shown to be clastogenic under the test conditions.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal The Toxicologist.
<b>References</b>	Heck J.D., Vollmuth, T.A., Cifone, M.A., Jagannath, D.R., Myhr, B. and Curren, R.D. (1989) An Evaluation of Food Flavoring Ingredients in a Genetic Toxicity Screening. The Toxicologist Vol 9 NO. 1.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	The Mouse Lymphoma L5178y TK +/- cell Mutagenesis (MLY) assay was performed according to the procedures described by Clive et al. (1979).
<b>Test Type</b>	Mouse Lymphoma Forward Mutation Assay (MLY).
<b>System of Testing</b>	L5178y Mouse Lymphoma cell line
<b>GLP</b>	NG
<b>Year</b>	1989
<b>Species/Strain</b>	Mice
<b>Metabolic Activation</b>	Aroclor 1254-induced rat liver S-9 activation system.
<b>Doses/Concentration</b>	250-300 nl/ml
<b>Results</b>	alpha-Terpineol was found to be negative in the presence or absence of rat liver S-9 activation system.
<b>Conclusion Remarks</b>	alpha-terpineol was found to be negative in the presence or absence of rat liver S-9 activation system.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal.
<b>References</b>	Heck J.D., Vollmuth, T.A., Cifone, M.A., Jagannath, D.R., Myhr, B. and Curren, R.D. (1989) An Evaluation of Food Flavoring Ingredients in a Genetic Toxicity Screening. The Toxicologist Vol 9 NO. 1.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Mutagenic effects of alpha terpineol was tested in 5 tester strains of Salmonella typhimurium TA 1535, TA 1537, TA1538, TA 98 & TA 100 in the presence or absence of S-9 fraction of Aroclor 1254 induced male Sprague-Dawley rat liver.
<b>Test Type</b>	Reverse mutation (Ames test)
<b>System of Testing</b>	Bacterial
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1989
<b>Species/Strain</b>	Bacteria: Salmonella typhimurium TA 1535, TA 1537, TA1538, TA 98 & TA 100
<b>Metabolic Activation</b>	S-9 fraction of Aroclor 1254 induced male Sprague-Dawley rat liver.

<b>Doses/Concentration</b>	10,000 ug/plate
<b>Results</b>	alpha-Terpineol was inactive in Salmonella strains TA 1535, TA 1537, TA 1538, TA 98 & TA 100 both in the presence and absence of metabolic activation system.
<b>Genotoxic effects</b>	No genotoxic effect
<b>Conclusion Remarks</b>	alpha-Terpineol was not found to be mutagenic under the test condition
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal.
<b>References</b>	Heck J.D., Vollmuth, T.A., Cifone, M.A., Jagannath, D.R., Myhr, B. and Curren, R.D. (1989) An Evaluation of Food Flavoring Ingredients in a Genetic Toxicity Screening. The Toxicologist Vol 9 NO. 1.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Mutagenic effects of linalool were tested in Salmonella typhimurium TA 98 & TA 100 in the presence or absence of S-9 fraction from rat liver.
<b>Test Type</b>	Reverse mutation (Ames test)
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1979
<b>Species/Strain</b>	Bacteria: Salmonella typhimurium TA 98 & TA 100
<b>Metabolic Activation</b>	S-9 fraction from rat liver.
<b>Remarks for Test Conditions</b>	Rats were fed linalool and the ether extract and aqueous phase of 24-hour urine was assayed for mutagenicity. Direct urine samples were assayed with TA 100 and TA 98 either in the presence or absence of betaglucuronidase.
<b>Results</b>	No significant mutagenic activity was detected in either the ether phase or aqueous phase of urine.
<b>Conclusion Remarks</b>	Urine from linalool fed rats was shown to be non mutagenic.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions
<b>References</b>	Rockwell P. and Raw, I. (1979) A mutagenic screening of various herbs, spices, and food additives. Nutrition and Cancer 1(4): 10-15.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6

<b>Method/guideline</b>	0.5 ml of Linalool was administered directly through a gastric tube to 2 rats. Urine was collected on ice for 24 hr period.
<b>Test Type</b>	Reverse mutation (Ames test)
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1979
<b>Species/Strain</b>	Bacteria: Salmonella typhimurium TA 98 & TA 100 & Sprague-Dawley rats
<b>Metabolic Activation</b>	Urine hydrolysis with beta-glucuronidase
<b>Doses/Concentration</b>	172 mg/kg
<b>Remarks for Test Conditions</b>	A screen for mutagenic activity was performed with the direct urine sample, either urine extract or the aqueous fraction. Assay was conducted with Salmonella typhimurium strain TA98 and TA100
<b>Results</b>	No mutagenic activity was detected when the direct urine samples or aqueous fraction were assayed either in presence or absence of beta-glucuronidase. Author had previously tested linalool and found it to be negative.
<b>Genotoxic effects</b>	None
<b>Conclusion Remarks</b>	No mutagenic activity was detected in the urine from rats given Linalool orally.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal.
<b>References</b>	Rockwell P. and Raw, I. (1979) A Mutagenic Screening of Various Herbs, Spices, and Food Additives. Nutrition and Cancer 1: 10-15.

<b>Substance Name</b>	Linalool (substance tested was isomer, isolinalool, colorless liquid)
<b>CAS No.</b>	<b>78-70-6</b>
<b>Method/guideline</b>	Ames test: This method conforms to the guidelines for bacterial mutagenicity testing published by OECD. OECD Guidelines for Testing of Chemicals No. 471 "Reverse Mutation Study," Method B14 of Common Directive 92/69/EEC.
<b>Test Type</b>	Reverse mutation
<b>System of Testing</b>	Bacterial
<b>GLP</b>	<b>Yes</b>
<b>Year</b>	<b>1999</b>
<b>Species/Strain</b>	Salmonella typhimurium strains TA1535, TA1537, TA98 and TA100; E. coli WP2uvrA-
<b>Metabolic Activation</b>	Rat liver homogenate

<b>Doses/Concentration</b>	15-5000 ug/plate
<b>Statistical Methods</b>	Dunnett's method of linear regression
<b>Remarks for Test Conditions</b>	Salmonella typhimurium strains TA1 535, TA1 537, TA98 and TA100 and E. coli WP2uvrA-were treated with test material using the Ames plate incorporation method at up to 6 dose levels, in triplicate, both with and without the addition of a rat liver homogenate metabolizing system. The vehicle DMSO control plates gave counts of revertant colonies within the normal range
<b>Results</b>	Resulted in visible reduction in the growth of the bacterial lawn in all tester strains at 5000 ug/plate; statistically significant increases in revertant colony frequency were observed in tester strains TA100 & TA1535 (without S9 mix only).; statistically significant increases & clear doserelated response in TA1535 with biggest increases at 3000 ug/plate.
<b>Cytotoxic concentration</b>	5000 ug/plate
<b>Remarks for Results</b>	The test material was considered to be mutagenic under the conditions of the test.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was conducted in accordance with GLP.
<b>References</b>	LeLievre, M. (1990a) 2,CDimethyl 2,7-octadien-4-ol)-acute toxicity to rainbow trout (Oncorhynchus mykiss) under static conditions. Springborn Lab. Inc. Report Submitted to SCM Glidco Organics. SLI Report # 90-g-3476.

<b>Substance Name</b>	alpha-Terpineol
<b>CAS No.</b>	98-55-5
<b>Method/guideline</b>	Test substance was assayed for mutagenicity toward 4 histidine-requiring mutants of Salmonella typhimurium (TA98, TA100, TA1535 and TA1537). It was tested qualitatively both with and without metabolic activation using a liver fraction (S-9).
<b>Test Type</b>	Reverse mutation assay
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1980
<b>Species/Strain</b>	Salmonella typhimurium (TA98, TA100, TA1535 and TA1537)
<b>Metabolic Activation</b>	Liver microsomal fraction from aroclor treated rats.
<b>Doses/Concentration</b>	3 umol/plate
<b>Results</b>	No mutagenic activity was detected.
<b>Genotoxic effects</b>	None

<b>Conclusion Remarks</b>	The test material was found to be non mutagenic under the test conditions
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions,
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal.
<b>References</b>	Florin I., Rutberg, L., Curvall, M., and Enzell, CR. (1980) Screening of tobacco smoke constituents for mutagenicity using the Ames test. Toxicology 18: 219-232.

<b>Substance Name</b>	Pine oil blend, clear colorless liquid (76.7% alpha- and <b>gamma</b> -terpineol)
<b>CAS No.</b>	<b>8002-09-3</b>
<b>Method/guideline</b>	Unscheduled DNA synthesis was carried out using rat primary hepatocytes.
<b>Test Type</b>	Unscheduled DNA synthesis
<b>System of Testing</b>	Primary Rat hepatocytes
<b>GLP</b>	Yes
<b>Year</b>	1987
<b>Species/Strain</b>	Male Sprague Dawley rats
<b>Doses/Concentration</b>	Test article was tested at ten dose levels ranging from 0.0003 to 10 ul/ml and was fully evaluated at 5 dose levels of 0.0003, 0.001, 0.003, 0.01 and 0.03 in the initial assay.
<b>Statistical Methods</b>	Method of Casciano and Gaylor ( <i>Mutat Res</i> 122: 81-86, 1983) were used.
<b>Remarks for Test Conditions</b>	Primary rat hepatocytes derived from the liver of adult male Sprague-Dawley rats were used in this study. The test article was dissolved and diluted in DMSO to make up the stock solution. The positive control cpd, <b>7,12</b> -dimethylbenzanthracene, was also dissolved in DMSO. All test article and control treatments were done under subdued yellow lights to avoid possible problems of photoinactivation. In addition to above mentioned doses, 9 dose levels were tested in the confirmatory assay ranging from 0.0001 to 1.0 ul/ml and the same 5 concentrations to test article were fully evaluated.
<b>Cytotoxic concentration</b>	3.0 and 10 ul/ml
<b>Genotoxic effects</b>	None
<b>Statistical results</b>	Under the test conditions, the test article did not cause a significant increase in the Unscheduled DNA synthesis as measured by the mean number of net nuclear grain count at any dose level.
<b>Conclusion Remarks</b>	Pine oil blend is considered negative in Unscheduled DNA synthesis test
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The protocol of this study was designed to comply with all EPA, FDA and OECD test guidelines. This study was also performed in accordance with GLP guidelines.

**References** Curren R. D. (1987) *Unscheduled DNA Synthesis in Rat Primary Hepatocytes with a Confirmatory Assay.* Microbiological Associate, Inc. Report to Chemical Specialties Manufacturers Association, Lab. Study No. T5366.380016.

<b>Substance Name</b>	Pine oil blend, clear colorless liquid
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	The test article was tested in the CHO/HGPRT mutation assay in the absence and presence of an Aroclor-induced rat liver S-9 activation system.
<b>Test Type</b>	CHO/HGPRT MUTATION ASSAY
<b>System of Testing</b>	CHO-K1-BH4 cells
<b>GLP</b>	Yes
<b>Year</b>	1987
<b>Species/Strain</b>	Chinese hamster ovary cells
<b>Metabolic Activation</b>	Aroclor induced male Fischer rat liver S-9 activation system.
<b>Doses/Concentration</b>	The first and the confirmatory assays were conducted at dose levels of 250, 200, 170, 130 and 100 $\mu\text{g/ml}$ in the non-activated study and at 400, 300, 200, 100 and 50 $\mu\text{g/ml}$ in the presence of s-9.
<b>Remarks for Test Conditions</b>	Ethyl methanesulfonate was used as positive control.
<b>Results</b>	The mutant frequencies of the test article treated groups were not increased significantly above the control, whereas EMS, positive control, induced a total of 180 mutants to yield mutant frequency of 271 .1mutants/10E6 clonable cells.
<b>Genotoxic effects</b>	None
<b>Conclusion Remarks</b>	Under the conditions of the assays, Pine Oil Blend was negative in the CHO/HPRT mutation assay.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The protocol of this study was designed to comply with all EPA, FDA and OECD test guidelines. This study was also performed in accordance with GLP guidelines.
<b>References</b>	Yang L. L. (1987) <i>CHO/HGPRT Mutation Assay.</i> Pine Oil Blend. Microbiological Associates Inc Report to Chemical Specialties Manufacturers Association. Lab. Study No. T5366.332001.

<b>Substance Name</b>	Tetrahydroinanolol
<b>CAS No.</b>	78-69-3
<b>Method/guideline</b>	Ames test was performed on five tester strains of Salmonella typhimurium (TA 1535, TA 100, TA 1537, TA 1538, TA 98).
<b>Test Type</b>	Reverse Mutation

<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1983
<b>Species/Strain</b>	Salmonella typhimurium (TA 1535, TA 100, TA 1537, TA 1538, TA 98.
<b>Metabolic Activation</b>	S-9 liver fraction was prepared from Aroclor-pretreated rats (Aroclor 1254, 500 mg/kg, ip).
<b>Doses/Concentration</b>	Up to 3600 ug/plate
<b>Statistical Methods</b>	Statistical significance was determined according to the methods of Kastenbaum and Bowman (1970).
<b>Remarks for Test Conditions</b>	Positive controls were run in each experiment with the reference mutagens sodium azide and benzo[a]pyrene.
<b>Results</b>	No mutagenic activity was detected with any of the Salmonella strains tested.
<b>Cytotoxic concentration</b>	NG
<b>Genotoxic effects</b>	None
<b>Conclusion Remarks</b>	No mutagenic activity was detected with any of the Salmonella strains tested.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed.
<b>References</b>	Wild D., King, M. T., Gocke, E. and Eckhardt. (1983). Study of Artificial Flavouring Substances for Mutagenicity in the Salmonella/Microsome, BASC and Micronucleus tests, Food and Chemical Toxicology 21(6), 707-719.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Ret-assay and mutation test
<b>Test Type</b>	Reverse Mutation
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1986
<b>Species/Strain</b>	Bacteria: E.Coli WP2 uvrA
<b>Metabolic Activation</b>	None
<b>Doses/Concentration</b>	125-1000 ug/plate
<b>Remarks for Test Conditions</b>	Positive for reverse mutation if maximum revertants/spontaneous revertants>2.0.
<b>Results</b>	No mutagenic activity at any dose level.

<b>Cytotoxic concentration</b>	>1000 ug/plate
<b>Genotoxic effects</b>	None
<b>Conclusion Remarks</b>	No mutagenic effects in E. Coli.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Yoo Y.S. (1986) Mutagenic and antimutagenic activities of flavoring agents used in foodstuffs. Journal of the Osaka City Medical Center 34, 267-288

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	The DNA-repair test with Bacillus subtilis was performed as described by Kada et al. (1980).
<b>Test Type</b>	DNA-Repair test
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1978
<b>Species/Strain</b>	Bacillus subtilis H17 Rec+ or M45 Rec-
<b>Metabolic Activation</b>	S9 fraction was prepared from the PCB-treated male Sprague-Dawley rats.
<b>Doses/Concentration</b>	17 ug/disk
<b>Statistical Methods</b>	NG
<b>Results</b>	No difference in zones of inhibition for H17 and M45.
<b>Genotoxic effects</b>	None
<b>Statistical results</b>	NG
<b>Conclusion Remarks</b>	No evidence of mutagenicity was detected under the test conditions.
<b>Data Qualities Reliabilities</b>	Reliability code 3. Data not reliable. Data in Japanese (not translated).
<b>References</b>	Oda Y., Hamano Y., Inoue K., Yamamoto H., Niihara T., and Kunita N. (1978) Mutagenicity of food flavors in bacteria. Shokuhin Eisei Hen 9, 177-181.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Spore plate rec assay
<b>Test Type</b>	DNA-Repair test

<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1986
<b>Species/Strain</b>	Bacillus subtilis H17 Rec+ or M45 Rec-
<b>Metabolic Activation</b>	S9 fraction was prepared from the PCB-treated male Sprague-Dawley rats.
<b>Doses/Concentration</b>	10 ug/disk
<b>Statistical Methods</b>	NG
<b>Results</b>	No difference in zones of inhibition for H17 and M45.
<b>Genotoxic effects</b>	None
<b>Statistical results</b>	NG
<b>Conclusion Remarks</b>	Mutagenicity at 10 nl/disk.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>References</b>	Yoo Y.S. (1986) Mutagenic and antimutagenic activities of flavoring agents used in foodstuffs. Journal of the Osaka City Medical Center 34, 267-288.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	In vitro and In vivo-in vitro Mutagenicity Assay
<b>Test Type</b>	In vivo-in vitro Ames assay
<b>GLP</b>	NG
<b>Year</b>	1979
<b>Species/Strain</b>	Sprague-Dawley rats
<b>Sex</b>	Not given
<b>Route of Administration</b>	Oral gavage
<b>Doses/Concentration</b>	500 ul/rat (approximately 452 mg/rat)
<b>Exposure Period</b>	Urine collected over 24 hrs
<b>Remarks for test conditions</b>	In the in vitro assay, 0.05 to 100 ul/plate of linalool was incubated with strain TA 98 and TA 100 of <i>Salmonella typh.</i> with S-9 prepared from Aroclor 1254-treated rats. In the <i>in vivo-in vitro</i> experiment, rats (2) were given a 0.5 ml (452 mg) oral dose of linalool by oral intubation and the urine was collected over the next 24 hours. Fifty (50) to 300 ul of ether extracts of the urine, the aqueous fraction of the ether extracts, or un-extracted (direct) urine samples were then incubated with

	strains TA 98 and TA 100 for 48 hours. The direct urine mutagenicity assays were performed either in the presence of absence of <b>beta-glucuronidase</b> . Positive controls were sodium azide and picrolic acid for TAIOO and TA98, respectively.
<b>Genotoxic effects</b>	None
<b>NOEL (C)/ LOEL (C)</b>	452 mg/kg
<b>Statistical Evaluation</b>	A positive response in TAIOO was defined as any deviation above the upper limit of 99.9% confidence limits of the mean control values.
<b>Remarks for Results</b>	A standard Ames assay with Salmonella typhimurium strain TA98 and TAIOO was performed with the linalool in the presence or S-9, direct urine sample in the presence or absence of beta-glucuronidase, urine ether extract, or the aqueous fractions of ether extracts without S-9 activation.
<b>Conclusion Remarks</b>	There was no evidence of mutagenicity for linalool or linalool urinary metabolites in rats given 452 mg of linalool.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Remarks for Data Reliability</b>	Study was reported in Journal of Cancer and Nutrition.
<b>References</b>	Rockwell P. and Raw, I. (1979) A mutagenic screening of various herbs, spices, in food additives. Nutrition and Cancer 1: IO-15

<b>Substance Name</b>	Linalyl acetate
CAS No.	115-95-7
<b>Method/guideline</b>	The DNA-repair test with Bacillus subtilis was performed as described by Kada et al. (1980).
<b>Test Type</b>	DNA-Repair test
<b>System of Testing</b>	Bacterial
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1978
<b>Species/Strain</b>	Bacillus subtilis H17 Rec+ or M45 Rec-
<b>Metabolic Activation</b>	S9 fraction was prepared from the PCB-treated male Sprague-Dawley rats.
<b>Doses/Concentration</b>	18 ug/disk
<b>Statistical Methods</b>	<b>NG</b>
<b>Results</b>	No difference in zones of inhibition for H17 and M45.
<b>Genotoxic effects</b>	None
<b>Statistical results</b>	<b>NG</b>
<b>Conclusion Remarks</b>	No evidence of mutagenicity was detected under the test conditions.

**Data Qualities Reliabilities** Reliability code 3. Data not reliable. Data in Japanese (not translated).

**References** Oda Y., Hamono Y., Inoue K., Yamamoto H., Niihara T., and Kunita N. (1978) Mutagenicity of food flavors in bacteria. Shokuhin Eisei Hen 9, 177-181.

<b>Substance Name</b>	alpha-Terpineol acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	The DNA-repair test with Bacillus subtilis was performed as described by Kada et al. (1980).
<b>Test Type</b>	DNA-Repair test
<b>System of Testing</b>	Bacterial
<b>GLP</b>	NG
<b>Year</b>	1978
<b>Species/Strain</b>	Bacillus subtilis HI 7 Rec+ or M45 Rec-
<b>Metabolic Activation</b>	S9 fraction was prepared from the PCB-treated male Sprague-Dawley rats.
<b>Doses/Concentration</b>	19 ug/disk
<b>Statistical Methods</b>	NG
<b>Results</b>	No difference in zones of inhibition for H17 and M45.
<b>Genotoxic effects</b>	None
<b>Statistical results</b>	NG
<b>Conclusion Remarks</b>	No evidence of mutagenicity was detected under the test conditions.
<b>Data Qualities Reliabilities</b>	Reliability code 3. Data not reliable. Data in Japanese (not translated).
<b>References</b>	Oda Y., Hamono Y., Inoue K., Yamamoto H., Niihara T., and Kunita N. (1978) Mutagenicity of food flavors in bacteria. Shokuhin Eisei Hen 9, 177-181.

#### 4.3 Genetic Toxicity *In Vivo*

<b>Substance Name</b>	Pine Oil Blend, clear, colorless liquid (76.7% alpha-and gamma-terpineol)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	Mice were exposed to test article by single IP injection. The high dose <b>was</b> selected to represent 80% of the LD50/7. Bone marrow polychromatic erythrocytes were collected 24, 48 and 72 hours after the treatment. Bone Marrow polychromatic

	erythrocytes were hen examined microscopically for micronuclei.
<b>Test Type</b>	Micronucleus Cy-togenetic Assay
<b>GLP</b>	Yes
<b>Year</b>	1987
<b>Species/Strain</b>	Male and female CD-1 mice
<b>Sex</b>	Male and female
<b>Route of Administration</b>	Intraperitoneal
<b>Doses/Concentration</b>	1155, 578, 116 mg/kg
<b>Exposure Period</b>	24, 48 and 72 hrs
<b>Genotoxic effects</b>	None
<b>NOEL (C)/ LOEL (C)</b>	1155 mg/kg
<b>Statistical Evaluation</b>	Kastenbaum-Bowman tables
<b>Remarks for Results</b>	The number of micronucleated erythrocytes was not significantly increased in test article treated groups, regardless of sex or sacrifice time point. TEM induced a significant increase in micronucleated polychromatic erythrocytes in male and female mice relative to the vehicle control.
<b>Conclusion Remarks</b>	Under the conditions of the assay, Pine Oil Blend didn't increase the incidence of micronucleated polychromatic erythrocytes in bone marrow and was concluded to be negative in the micronucleus test using male and female CD-1 mice.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was performed in accordance with FDA GLP regulation, EPA GLPs and OECD guidelines.
<b>References</b>	Putman D. L. (1987) Micronucleus Cytogenetic Assay in Mice. Pine oil Blend. Microbiological Associates, Inc. Report to Chemical Specialties Manufacturers Association. Lab. Study Number T5366.122.

#### 4.4 Repeat Dose Toxicity

<b>Substance Name</b>	alpha-Terpinyl acetate
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Method used by U.S. Food and Drug Administration.
<b>GLP</b>	P re-GLP
<b>Year</b>	1967

<b>Species/strain</b>	Osborne-Mendel rats
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Diet
<b>Doses/concentration Levels</b>	1000, 2500 or 10000 ppm
<b>Exposure Period</b>	20 weeks
<b>Frequency of Treatment</b>	Continuously in diet
<b>Control Group</b>	Yes
<b>Post Exposure Observation Period</b>	NG
<b>Remarks for Test Conditions</b>	Groups of ten male and ten female Osborne-Mendel rats were provided alpha-terpinyl acetate in the diet at concentrations of 0, 1000, 2500 or 10,000 ppm for 20 weeks. No vehicle was used. The diet was prepared and analyzed weekly. Measurements of body weight, food intake and general condition were recorded weekly. Hematological examinations (white cell counts, red cell counts, hemoglobin and hematocrit) were performed at the termination of the study. Macroscopic examination of all tissues was performed. Histopathological examination was performed on the liver, kidneys, spleen, heart, and testes of 6-8 animals (evenly divided by sex) from the high dose and controls.
<b>NOAEL (NOEL)</b>	10000 ppm
<b>LOAEL (LOEL)</b>	No adverse effects at highest dose
<b>Actual Dose Received by Dose Level and Sex</b>	NG
<b>Toxic Response/effects by Dose Level</b>	None
<b>Statistical Evaluation</b>	NG
<b>Remarks for Results</b>	Determination of the dietary concentration of alpha-terpinyl acetate revealed a weekly loss of 7%. Measurements of body weight, food intake and general condition recorded weekly showed no significant differences between test and control animals at any intake level. At termination, hematological examinations revealed no difference from controls. At necropsy, no differences were reported between major organ weights of test and control animals. Gross examination of tissue of all animals was unremarkable and histopathological examination revealed no treatment-related lesions.
<b>Conclusion Remarks</b>	This study demonstrates a NOAEL in rats of at least 10,000 ppm
<b>Data Qualities Reliabilities</b>	Reliability code 1, Reliable without restrictions. This study was performed by the FDA prior to establishment of GLP & OECD.
<b>References</b>	Hagan E.C., Hansen W.H., Fitzhugh O.G., Jenner P.M., Jones W.I., Taylor J.M., Long E.L., Nelson A.A., and Brouwer J.B. (1967) Food Flavourings and Compounds of Related Structure. II. Subacute and Chronic Toxicity. Food Cosm Toxicology, 5, 141-157.

<b>Substance Name</b>	Linalool
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	A rapid screening protocol incorporating key elements of the US NTP's immunotoxicity tier testing strategy was used to evaluate the effect of the test substance on humoral and cell-mediated immune responses.
<b>GLP</b>	NG
<b>Year</b>	1994
<b>Species/strain</b>	Mice: B6C3F1 mice
<b>Sex</b>	Female
<b>Route of Administration</b>	Intragastric
<b>Doses/concentration Levels</b>	375, 188 & 94 mg/kg/day in methyl cellulose
<b>Exposure Period</b>	5 days
<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	1% Methyl cellulose
<b>Post Exposure Observation Period</b>	10 days
<b>Remarks for Test Conditions</b>	Test material was administered intragastrically on a daily basis for 5 days at 3 dose levels to B6C3F1 mice. A host-resistance assay (Listeria monocytogenes bacterial challenge) was conducted to assess cell-mediated immunity. Humoral immunity was measured by the antibody Plaque-Forming cell (PFC) response to sheep erythrocytes. Cyclophosphamide served as an immunosuppressive positive control agent.
<b>NOAEL (NOEL)</b>	375 mg/kg/day
<b>LOAEL (LOEL)</b>	No adverse effects at highest dose
<b>Toxic Response/effects by Dose Level</b>	None
<b>Remarks for Results</b>	No increase in the mortality or significant alteration to the Plaque-Forming cell response was noted
<b>Conclusion Remarks</b>	Linalool did not mediate the cell-mediated or humoral immune response.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Gaworsky CL., Vollmuth, T.A., Dozier, M.M., Heck, J.D., Dunn, L.T., Ratajczak, H.V. and Thomas, P.T. (1994) An Immunotoxicity Assessment of Food Flavouring Ingredients. <i>Fd Chem. Toxic</i> 32(5), 409-415.
<b>Substance Name</b>	Linalyl acetate

<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	A rapid screening protocol incorporating key elements of the US NTP's immunotoxicity tier testing strategy was used to evaluate the effect of the test substance on humoral and cell-mediated immune responses.
<b>GLP</b>	NG
<b>Year</b>	1994
<b>Species/strain</b>	Mice: CD1 mice
<b>Sex</b>	Female
<b>Route of Administration</b>	Intragastric
<b>Doses/concentration Levels</b>	3000, 1500, 750 mg/kg/day in corn oil
<b>Exposure Period</b>	5 days
<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	Corn oil
<b>Post Exposure Observation Period</b>	10 days
<b>Remarks for Test Conditions</b>	Test material was administered intragastrically on a daily basis for 5 days at 3 dose levels to B6C3F1 mice. A host-resistance assay (Listeria monocytogenes bacterial challenge) was conducted to assess cell-mediate immunity. Humoral immunity was measured by the antibody Plaque-Forming cell (PFC) response to sheep erythrocytes. Cyclophosphamide served as an immunosuppressive positive control agent.
<b>LOAEL (LOEL)</b>	750 mg/kg/day
<b>Toxic Response/effects by Dose Level</b>	Increased mortality following challenge with L. monocytogenes was noted at 750 mg/kg. No significant alteration in the PFC response was noted.
<b>Remarks for Results</b>	The increased mortality observed following challenge with L. monocytogenes was not dose-related and therefore considered to be incidental.
<b>Conclusion Remarks</b>	Linalyl acetate did not mediate the cell-mediated or humoral immune response.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Gaworsky CL., Vollmuth, T.A., Dozier, M.M., Heck, J.D., Dunn, L.T., Ratajczak, H.V. and Thomas, P.T. (1994) An Immunotoxicity Assessment of Food Flavouring Ingredients. <i>Fd Chem. Toxic</i> 32(5), 409-415.
<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	This study was designed to investigate the Systemic toxicity of plinyl acetate and complied with the testing method described

	plinyl acetate and complied with the testing method described in Commission Directive 92/69/EEC (Method B7)
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/Strain</b>	Sprague-Dawley Cri-CD Br Strain Rats
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Gavage
<b>Doses/concentration Levels</b>	15, 150 & 1000 mg/kg/day
<b>Exposure Period</b>	28 days
<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	Arachis oil BP
<b>Post Exposure Observation Period</b>	24 hours
<b>Remarks for Test Conditions</b>	The test material was administered to 3 groups of 5 male and 5 female rats for 28 days. A control group of 5 male and 5 female received the vehicle only.
<b>NOAEL (NOEL)</b>	15 mg/kg/day
<b>LOAEL (LOEL)</b>	150 mg/kg/day
<b>Toxic Response/effects by Dose Level</b>	Increased liver and kidney wt., speckled kidney, microscopic treatment related changes in liver and kidney, accumulation of globular eosinophilic material in male kidney, centrilobular hepatocyte enlargement
<b>Statistical Evaluation</b>	Linear regression analysis, Followed by one way ANOVA incorporating Leven's test for homogeneity.
<b>Remarks for Results</b>	No deaths, no clinically observable symptoms, no effect on body weight, food consumption, hematology and clinical chemistry.
<b>Conclusion Remarks</b>	Oral administration of the test material to rats for a period of 28 consecutive days at the dose levels of up to 1000 mg/kg/day resulted in minor treatment-related changes at 1000 and 150 mg/kg/day. These changes were centrilobular-hepatocyte enlargement, speckled kidney and globular eosinophilic material in renal tubular epithelium.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions. Data are considered to be reliable.
<b>Data Reliabilities Remarks</b>	The study was performed in compliance with UK GLP and these regulations are in accordance with GLP standards published as OECD Monograph No. 45.
<b>References</b>	Thomas O. N. (1997) Plinyl Acetate: Twenty-Eight day Repeated Dose Oral (Gavage) Toxicity Study in the Rat. Safepharm Laboratories, Ltd. SPL Project No. 1044/009.

<b>Substance Name</b>	2,6,6-Trimethylbicyclo[3.1.1]heptan-2-ol thermal rearrangement products (structurally related substance - plinyl acetate, clear colorless non-viscous liquid)
-----------------------	---

<b>CAS No.</b>	125252-49-5
<b>Method/guideline</b>	The study was designed to comply with the testing method described in Commission Directive 92/69/EEC (Method B7)
<b>GLP</b>	Yes
<b>Year</b>	1997
<b>Species/strain</b>	Sprague-Dawley CrI-CD Br Strain Rats
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Gavage
<b>Doses/concentration Levels</b>	0, 150, 400 & 1000 mg/kg/day
<b>Exposure Period</b>	28 days
<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	Arachis oil
<b>Post Exposure Observation Period</b>	24 hours
<b>Remarks for Test Conditions</b>	The study was designed to ascertain Maximum Tolerated Dose Level for 28 days study.
<b>Actual Dose Received by Dose Level and Sex</b>	150, 400, 1000 mg/kg/day
<b>Toxic Response/effects by Dose Level</b>	At 1000 mg/kg: Increased salivation dosing day 2, fur loss after 5 days, hunched posture due to dosing stress; At 400 or 150 mg/kg: increased salivation after Day 4 (due to impalatability &/or local irritant nature of formulation.
<b>Remarks for Results</b>	No deaths, no clinically observable signs of toxicity, no effect on body wt, no macroscopic abnormalities were observed.
<b>Conclusion Remarks</b>	Dose levels of 1000, 100 and 15 mg/kg/day were chosen for the main study.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions. Data are considered to be reliable.
<b>Data Reliabilities Remarks</b>	The study was performed in compliance with UK GLP and these regulations are in accordance with GLP standards published as OECD Monograph No. 45 (OCDE/GD(92)32).
<b>References</b>	Thomas O. N. (1997) (1997) Plinyl Acetate: Twenty-Eight day Repeated Dose Oral (Gavage) Toxicity Study in the Rat. Safepharm Laboratories, Ltd. SPL Project No. 1044/009.

<b>Substance Name</b>	Mixture of Linalool and Citronellol
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	The test mixture was incorporated in the ration at a level designed to provide daily in the food 100 mg of the flavor blend per kg of body wt. The unsupplemented diet was fed to the controls. The rats were fed for 12 weeks.
<b>GLP</b>	Pre GLP
<b>Year</b>	1958

<b>Species/strain</b>	Rat (Unspecified strain)
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Feed
<b>Doses/concentration Levels</b>	100 mg/kg of mixture per day (50 mg/kg bw/day of linalool)
<b>Exposure Period</b>	12 weeks
<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	Feeding tests with a mixture of equal parts of citronellol and linalool fed at a level 100 times the estimated use level in the human diet disclosed no adverse effect on efficiency of food utilization or other observable criteria.
<b>Remarks for Test Conditions</b>	After 12 weeks on test, the urine of 3 rats of each sex per group was examined for the presence of sugar and albumin, blood hemoglobin levels were determined and autopsies were performed on all animals.
<b>Toxic Response/effects by Dose Level</b>	No adverse effects on efficiency of food utilization or other observable physiological criteria were noted.
<b>Remarks for Results</b>	The depression in the growth and food intake of the male rats was attributed to impalatability of the test material at the level administered
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Food and Drug Research Laboratories (1958) Toxicological Screening of Components of Food Flavors Class VI. Citronellol and Linalool.

<b>Substance Name</b>	Linalyl acetate
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Cocarcinogenicity study
<b>GLP</b>	NG
<b>Year</b>	1971
<b>Species/strain</b>	Mice/ICR/Ha Swiss
<b>Sex</b>	Female
<b>Route of Administration</b>	Dermal
<b>Doses/concentration Levels</b>	3 mg/mouse
<b>Exposure Period</b>	67 weeks
<b>Frequency of Treatment</b>	3 times weekly
<b>Control Group</b>	Control groups given solvent only or untreated.
<b>Post Exposure Observation Period</b>	NG

<b>Remarks for Test Conditions</b>	The test material was applied as a solution in acetone to the backs of groups of 20 mice, 3 times per week for 67 weeks.
<b>NOAEL (NOEL)</b>	No carcinogenic activity at 3 mg dose
<b>LOAEL (LOEL)</b>	No effects at single dose level
<b>Toxic Response/effects by Dose Level</b>	None
<b>Remarks for Results</b>	Mice showing tumors for 30 days or more and greater than 1 mm in size were counted. Tumors were confirmed histologically.
<b>Conclusion Remarks</b>	No carcinogenic effects observed.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Data Reliabilities Remarks</b>	Study was a cocarcinogenicity study using linalyl acetate either alone or with benzo(a)pyrene as a cocarcinogen.
<b>References</b>	Van Duuren B., Blazej T., Goldschmidt B. Katz C., Melchionne S., and Sivak A. (1971) Cocarcinogenesis studies on mouse skin and inhibition of tumor induction. Journal of the National Cancer Institute, <b>45(5)</b> , 1039-1044.

<b>Substance Name</b>	Linalyl acetate (data on structurally related ester, linalyl isobutyrate)
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Method used by U.S. Food and Drug Administration.
<b>GLP</b>	Na
<b>Year</b>	1967
<b>Species/strain</b>	Osborne-Mendel rats
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Diet
<b>Doses/concentration Levels</b>	1000, 2500 or 10000 ppm
<b>Actual Dose Received by Dose Level and Sex</b>	NG
<b>Exposure Period</b>	<b>18</b> weeks
<b>Frequency of Treatment</b>	Continuously in diet
<b>Control Group</b>	Yes
<b>Post Exposure Observation Period</b>	NG
<b>Remarks for Test Conditions</b>	Groups of ten male and ten female Osborne-Mendel rats were provided linalyl isobutyrate in the diet at concentrations of 0, 1000, 2500 or 10,000 ppm for 17 weeks. No vehicle was used. The diet was prepared and analyzed weekly. Measurements of body weight, food intake and general condition were recorded weekly. Hematological examinations (white cell counts, red cell counts, hemoglobins and hematocrits) were performed at the

<b>NOAEL (NOEL)</b>	termination of the study. Macroscopic examination of all tissues was performed. Histopathological examination was performed on the liver, kidneys, spleen, heart, and testes of 6-8 animals (evenly divided by sex) from the high dose and controls. 10000 ppm
<b>LOAEL (LOEL)</b>	No adverse effects at highest dose
<b>Toxic Response/effects by Dose Level</b>	None
<b>Remarks for Results</b>	Determination of the dietary concentration of alpha-terpinyl acetate revealed a 7% loss of test material from diet during a 7-day period. Measurements of body weight, food intake and general condition recorded weekly showed no significant differences between test and control animals at any intake level. At termination, hematological examinations revealed no difference from controls. At necropsy, no differences were reported between major organ weights of test and control. At necropsy, no differences were reported between major organ weights of test and control animals. Gross examination of tissue of all animals was unremarkable and histopathological examination revealed no treatment-related lesions.
<b>Conclusion Remarks</b>	This study demonstrates a NOAEL in rats of at least 10,000 ppm.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions. This study was performed by Food and Drug Administration prior to the establishment of GLP and OECD.
<b>References</b>	Hagan E.C., Hansen W.H., Fitzhugh O.G., Jenner P.M., Jones W.I., Taylor J.M., Long E.L., Nelson A. A., and Brouwer, J.S. (1967) Food Flavourings and Compounds of Related Structure. II, Subacute and Chronic Toxicity. Food Cosmetic Toxicology 5, 141-157.

<b>Substance Name</b>	Linalyl acetate (mixture of linalyl isobutyrate & geranyl acetate)
<b>CAS No.</b>	115-95-7
<b>Method/guideline</b>	Method used by FDA. The test mixture was incorporated in the ration at a level designed to provide daily in the food 100 mg of the flavor blend/kg of body wt. The unsupplemented diet was fed to the controls. The rats were fed for 12 weeks.
<b>GLP</b>	Pre GLP
<b>Year</b>	1958
<b>Species/strain</b>	Rat (Unspecified strain)
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Feed
<b>Doses/concentration Levels</b>	Target dose, 100 mg/kg per day of mixture. Measured dose, 112 mg/kg per day (24 mg/kg bw/day of linalyl acetate)
<b>Exposure Period</b>	12 weeks

<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	Male and female rats were maintained on a basal diet for 12 weeks.
<b>Remarks for Test Conditions</b>	After 12 wks on test, the urine of 3 rats/sex/group was examined for the presence of sugar and albumin, blood hemoglobin levels were determined and autopsies were performed on all animals.
<b>Toxic Response/effects by Dose Level</b>	A slight decrease in growth and efficiency of food utilization in females was concluded by the authors not to be an adverse effect.
<b>Remarks for Results</b>	The depression in the growth and food intake of the female rats was attributed to impalatability of the test material at the level administered
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions.
<b>References</b>	Trubeck Laboratories (1958) Toxicological Screening of Components of Linalyl acetate, linalyl isobutyrate, and geranyl acetate in rats. Class VIII. Unpublished Report

#### 4.5 Reproductive Toxicity

<b>Substance Name</b>	Linalool (the study was conducted using natural linalool (coriander oil) 72.9% linalool)
<b>CAS No.</b>	78-70-6
<b>Method/guideline</b>	Virgin female Sprague-Dawley rats (1 O/group) were orally administered a vehicle or the test material at three dosages from one week prior to a cohabitation period through gestation, parturition and a Cday postpartum period.
<b>Test Type</b>	Reproductive/developmental study
<b>GLP</b>	NG
<b>Year</b>	1990
<b>Species/Strain</b>	Sprague-Dawley rat
<b>Sex</b>	Female
<b>Route of Administration</b>	Oral/gavage
<b>Duration of Test</b>	39 days
<b>Doses/Concentration</b>	250, 500 & 1000 mg/kg/day
<b>Premating Exposure period for males</b>	7 days
<b>Frequency of Treatment</b>	Daily
<b>Control Group and Treatment</b>	1% methylcellulose
<b>Remarks for Test Conditions</b>	Clinical signs, body wt & food consumption were recorded throughout the study. Mating performance, fertility, duration of

	gestation/parturition, maternal behavior, litter sized, dystocia, number of implantation sites, and gross lesions at necropsy were also examined. Offspring were examined for viability, sex, external morphology, and body wt at birth and on day 4 postpartum.
<b>NOAEL(NOEL)</b>	250 mg/kg/d (maternal NOAEL)
<b>LOAEL(LOEL)</b>	500 mg/kg/d (maternal)
<b>Parental data and F1 as Appropriate</b>	250 mg/kg/d: <u>increased</u> body weight and food consumption; 500 mg/kg/d: non-statistically significant decrease in body weight & food consumption, decreased gestation index and decreased length of gestation; 1000 mg/kg: statistically significant decrease in body weight & food consumption, statistically significant decrease in gestation index, length of gestation, and litter size
<b>Offspring Toxicity F1 and F2</b>	1000 mg/kg: increased number of pups dying on Days 1-4 postpartum.
<b>Statistical Evaluation</b>	ANOVA followed by Dunnett's test
<b>Remarks for Results</b>	Increased body weights were accompanied by increased food consumption. Effects were not considered adverse effects at 250 mg/kg. No effects on fertility or offspring.
<b>Conclusion Remarks</b>	Dose levels greater than or equal to 500 mg/kg/d produced toxicity in adult female rats and, at 1000 mg/kg/d, decreased viability of fetuses.
<b>Data Reliabilities Qualities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal Teratology.
<b>References</b>	Vollmuth T.A., Bennett, M.B., Hoberman, A.M. and Christian, MS. (1995) An Evaluation of Food Flavoring Ingredients Using an In Vivo Reproductive and Developmental Toxicity Screening Test. Teratology 41(5): 597.

<b>Substance Name</b>	alpha-Terpineol acetate ( The study was performed on cardamon oil, a natural mixture containing >65% tertiary terpenoid alcohol and related esters including alpha-terpineol acetate (51%) alpha terpineol, linalool, linalyl acetate, and terpinen-4-ol)
<b>CAS No.</b>	80-26-2
<b>Method/guideline</b>	Virgin female Sprague-Dawley rats (10/group) were orally administered a vehicle or the test material at three dosages from one week prior to a cohabitation period through gestation, parturition and a 4 day postpartum period.
<b>Test Type</b>	Reproductive/developmental study
<b>GLP</b>	<b>NG</b>
<b>Year</b>	1990
<b>Species/Strain</b>	Sprague-Dawley rat
<b>Sex</b>	Female

<b>Route of Administration</b>	Oral/gavage
<b>Duration of Test</b>	39 days
<b>Doses/Concentration</b>	375, 750 & 1500 mg/kg/day
<b>Premating Exposure period for males</b>	7 days
<b>Frequency of Treatment</b>	Daily
<b>Control Group and Treatment</b>	1% methylcellulose
<b>Remarks for Test Conditions</b>	Clinical signs, body wt & food consumption were recorded throughout the study. Mating performance, fertility, duration of gestation/parturition, maternal behavior, litter sized, dystocia, number of implantation sites, and gross lesions at necropsy were also examined. Offspring were examined for viability, sex, external morphology, and body wt at birth and on day 4.
<b>NOAEL(NOEL)</b>	<375 mg/kg/d (maternal NOAEL) LOAEL at 375 mg/kg/d was based on decreased body weight gain and food consumption). A NOAEL for developmental toxicity in pups was 375 mg/kg/d.
<b>LOAEL(LOEL)</b>	375 mg/kg/d (maternal)
<b>Parental data and FI as Appropriate</b>	375 mg/kg/d: non-statistically significant decrease in body weight gain and food consumption; 750 mg/kg/d: mortality, clinical signs, statistically significant decrease in body weight gain & food consumption, gross lesions at necropsy; 1500 mg/kg: mortality, clinical signs, statistically significant decrease in body weight gain & food consumption, gross lesions at necropsy. Small decreased in average pup weights at 750 and 1500 mg/kg/d and significant increase ( $P<0.01$ ) in pup mortality at 1500 mg/kg/d
<b>Offspring Toxicity F1 and F2</b>	750 mg/kg; decreased body weights: 1500 mg/kg/d increased mortality and decreased body weight gain
<b>Statistical Evaluation</b>	ANOVA followed by Dunnett's test
<b>Remarks for Results</b>	Decreased body weight gain and food consumption were not statistically significant at the lowest dose.
<b>Conclusion Remarks</b>	Dose levels equal to 375 mg/kg/d resulted in depressed body weight gain and food consumption while 750 and 1500 mg/kg/d produced mortality and toxicity (liver weight increase and liver pathology) in adult female rats. At 750 mg/kg/d in offspring, decreased body weight gain during days 1-4 postpartum; at 1500 mg/kg/d, increased mortality and decreased body weight gain of pups. The authors concluded that the test material was not uniquely toxic to the reproductive performance of female rats or the development of their offspring.
<b>Data Reliabilities Qualities</b>	Reliability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	The study was published in a peer-reviewed journal Teratology.
<b>References</b>	Vollmuth T.A., Bennett, M.B., Hoberman, A.M. and Christian, M.S. (1995) An Evaluation of Food Flavoring Ingredients Using an In Vivo Reproductive and Developmental Toxicity Screening Test. Teratology 41(5):597.

## 4.6 Developmental/Teratogenicity Toxicity

<b>Substance Name</b>	Pine oil blend (blend compose of five commercial pine oil samples. Range of composition was <b>35-</b> 76.7% <i>alpha</i> - and gamma-terpineol, I-I 3.1% other terpineol isomers, O-14.6% secondary terpene alcohols, O.I-10% terpene hydrocarbons)
<b>CAS No.</b>	8002-09-3
<b>Method/guideline</b>	The test substance was administered on days 6 through 20 of gestation to pregnant rats at the dosage of 0, 50, 100, 500, 750, or 1000 mg/kg/day at a volume of 10 ml/kg. On day 20 of presumed gestation, all rats were terminated by asphyxiation with CO <sub>2</sub> .
<b>Test Type</b>	Dose range finding study of Teratology study
<b>GLP</b>	Yes
<b>Year</b>	1988
<b>Species/strain</b>	CrI:CD(SD)BR pregnant rats
<b>Sex</b>	Female
<b>Route of Administration</b>	Gavage
<b>Duration of Test</b>	14 days
<b>Doses/concentration Levels</b>	0, 100, 500, 750 or 1000 mg/kg/day
<b>Exposure Period</b>	14 days
<b>Control Group and Treatment</b>	Corn oil
<b>Remarks for Test Conditions</b>	Laparotomies were performed, corpora <b>lutea</b> were counted, and the uterus of each rat was removed, weighed and then examined for number, placement and viability of implantations. Live fetuses were weighed, sexed and gross external alternations were identified. Soft tissue and skeletal evaluations were not performed for this dosage-range (pilot study.)
<b>Maternal Data with Dose Level</b>	There were no deaths or abortions during the course of this study. Necropsy revealed no gross lesions. Excess salivation occurred within 30 min at dosage 100 and above, local alopecia, inhibition of average maternal body weight, decrease in feed consumption for 3 highest doses. At 750 & 1000 mg/kg, reduced average gravid uterine weight,
<b>Fetal Data with Dose Level</b>	Decreased fetal body weight in dosages 100 mg/kg and above. At 500 & 750 mg/kg, slight increase in average number of resorptions/litter.
<b>Remarks for Results</b>	On the basis of this study, dosages of 0, 50, 600 and 1200 mg/kg/day of Pine oil blend were recommended for definitive developmental toxicity study.
<b>Conclusion Remarks</b>	On the basis of this study, dosages of 0, 50, 600 and 1200 mg/kg/day of Pine oil blend were recommended for definitive developmental toxicity study.

<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restrictions.
<b>Data Reliabilities Remarks</b>	The study was conducted in accordance with GLP guidelines
<b>References</b>	Dearlove G. E. (1987) Dosage-Range Developmental Toxicity Study of Pine Oil Blend 1687 Administered Orally Via Gavage to CrI:CD(SD)BR Presumed Pregnant Rats (Pilot Study). Appendix IV to Parent, R.A. (1988). Teratogenicity Study in Albino Rats with Pine Oil Blend. A Report Submitted by Pine Oil Joint Venture to Chemical Specialties Manufacturers Association,

<b>Substance Name</b>	Pine oil blend (blend compose of five commercial pine oil samples. Range of composition was 35- 76.7% alpha- and gamma-terpineol, l-l 3.1% other terpineol isomers, 0-14.6% secondary terpene alcohols, 0.1-10% terpene hydrocarbons)
CAS No.	8002-09-3
<b>Method/guideline</b>	Pine oil Blend was administered by gavage at the doses of 0, 50, 600 or 1200 mg/kg/day to pregnant rats on gestation days 6 through 20. On day 20 of presumed gestation, rats were sacrificed.
<b>Test Type</b>	Teratology study
<b>GLP</b>	Yes
<b>Year</b>	1988
<b>Species/strain</b>	CrI:CD(SD)BR pregnant rats
<b>Sex</b>	Female
<b>Route of Administration</b>	Gavage
<b>Duration of Test</b>	14 days (Gestation day 6 through 20)
<b>Doses/concentration Levels</b>	0, 50, 600 or 1200 mg/kg/day
<b>Exposure Period</b>	14 days
<b>Frequency of Treatment</b>	Daily
<b>Control Group and Treatment</b>	Corn oil
<b>Remarks for Test Conditions</b>	The rats were examined for pregnancy, number and placement of implantations, early and late resorptions, live and dead fetuses, and number of corpora lutea. Fetuses were subsequently weighed and evaluated for both soft and hard tissue abnormalities.
<b>NOAEL Maternal Toxicity</b>	>50 mg/kg
<b>LOAEL Maternal Toxicity</b>	<600 mg/kg
<b>Maternal Data with Dose Level</b>	At 600 & 1200 mg/kg/day: maternal toxicity, decreased food consumption, body weight gains, excess salivation, alopecia, ungroomed coat, ataxia, decreased motor activity, impaired righting reflex and urine stained abdominal fur. Six of the 25

<b>Fetal Data with Dose Level</b>	rats in 1200 mg/kg died and necropsies revealed that adrenal weights were significantly increased in these rats. Increased incidences of delayed fetal ossification, delayed brain development, decreased fetal weights, increased embryofetal mortality, and sunken eyes bulge with associated soft and hard tissue findings at 1200 mg/kg, a dosage that also resulted in maternal death and a low incidence of embryo-fetal death (resorption).
<b>Statistical Evaluation</b>	Bartlett, Kruskal-Walks, Dunn's test, Dunnett's
<b>Conclusion Remarks</b>	Pine oil blend is not expected to produce adverse effects on embryofetal development except at doses that are also maternally toxic. The maternal and developmental NOEL for Pine Oil under the conditions of this study was greater than 50 mg/kg/day but less than 600 mg/kg/day.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restrictions
<b>Data Reliabilities Remarks</b>	The study was conducted in accordance to GLP guidelines.
<b>References</b>	Parent R.A. (1988) Teratogenicity Study in Albino Rats with Pine Oil Blend. A Report Submitted by Pine Oil Joint Venture to Chemical Specialties Manufacturers Association.