

201-15924B

**The Flavor and Fragrance High Production Volume  
Consortia**

**The Terpene Consortium**

**Robust Summaries for Aromatic Terpene Hydrocarbons**

RECEIVED  
OPPT/CRIC  
2005 JUN 16 AM 8:43

*p*-Cymene

CAS No. 99-87-6

**FFHPVC Terpene Consortium Registration Number**

100110

Submitted to the EPA under the HPV Challenge Program by:  
**The Flavor and Fragrance High Production Volume Chemical Consortia**  
1620 I Street, NW, Suite 925  
Washington, DC 20006  
Phone: 202-331-2325  
Fax: 202-463-8998

## **List of Member Companies**

**ARIZONA CHEMICAL**

**BASF CORPORATION**

**BEDOUKIAN RESEARCH, INC.**

**BOISE CASCADE CORPORATION**

**CHAMPION INTERNATIONAL CORPORATION**

**CITRUS AND ALLIED ESSENCES, LTD.**

**DRAGOCO**

**FRAGRANCE RESOURCES, INC.**

**GIVAUDAN**

**HERCULES INCORPORATED**

**INTERNATIONAL FLAVORS & FRAGRANCES, INC.**

**J. MANHEIMER, INC.**

**KURARAY CO., LTD.**

**MEAD CORPORATION**

**MILLENNIUM CHEMICALS**

**POLAROME INTERNATIONAL INCORPORATED**

**QUEST INTERNATIONAL**

**SENSIENT FLAVORS**

**TECNAL CORPORATION**

**THE PROCTOR AND GAMBLE CO.**

**UNILEVER-HPC**

# Table of Contents

<b>1</b>	<b>CHEMICAL AND PHYSICAL PROPERTIES .....</b>	<b>1</b>
1.1	MELTING POINT .....	1
1.2	BOILING POINT .....	3
1.3	VAPOR PRESSURE .....	4
1.4	N-OCTANOL/WATER PARTITION COEFFICIENT .....	5
1.5	WATER SOLUBILITY.....	7
<b>2</b>	<b>ENVIRONMENTAL FATE AND PATHWAYS.....</b>	<b>10</b>
2.1	PHOTODEGRADATION .....	10
2.2	BIODEGRADATION .....	10
2.3	FUGACITY .....	14
<b>3</b>	<b>ECOTOXICITY.....</b>	<b>17</b>
3.1	ACUTE TOXICITY TO FISH .....	17
3.2	ACUTE TOXICITY TO AQUATIC INVERTEBRATES.....	22
3.3	ACUTE TOXICITY TO AQUATIC PLANTS .....	27
<b>4</b>	<b>HUMAN HEALTH TOXICITY.....</b>	<b>30</b>
4.1	ACUTE TOXICITY .....	30
4.2	GENETIC TOXICITY .....	39
4.2.1	<i>In vitro Genotoxicity</i> .....	39
4.2.2	<i>In vivo Genotoxicity</i> .....	47
4.3	REPEAT DOSE TOXICITY.....	51
4.4	REPRODUCTIVE TOXICITY.....	73
4.5	DEVELOPMENTAL/TERATOGENICITY TOXICITY .....	74

# The Flavor and Fragrance High Production Volume Consortia

## Robust Summaries for Aromatic Terpene Hydrocarbons

The evaluation of the quality of the following data uses a systematic approach described by Klimisch [Klimisch *et al.*, 1996]. Based on criteria relating to international testing standards for categorizing data reliability, four reliability categories have been established. The following categories are:

- Reliability code 1. Reliable without restrictions
- Reliability code 2. Reliable with restrictions
- Reliability code 3. Not reliable
- Reliability code 4. Not assignable

## 1 Chemical and Physical Properties

### 1.1 Melting Point

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Melting Point</b>	-67.94 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Merck Index (1996) 12th edition, Susan Budavari, editor, Merck & Co. Inc., Whitehouse Station, NJ.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Melting Point</b>	-68 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	International Programme on Chemical Safety & The Commission of the European Communities (1993) <i>p</i> -Cymene. www.inchem.org.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Melting Point</b>	-67.9 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	CRC Handbook of Chemistry and Physics (1986) 67th edition, Robert C. Weast, editor, The Chemical Rubber Co. Press, Inc. Boca Raton, Florida.

## 1.2 Boiling Point

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Year</b>	1997
<b>Boiling Point</b>	177.1 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Merck Index (1996) 12th edition, Susan Budavari, editor, Merck & Co. Inc., Whitehouse Station, NJ.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Year</b>	1958
<b>Boiling Point</b>	177 °C
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Only short abstract available.
<b>References</b>	Furnas D.W. and Hine, C.H. (1958) Neurotoxicity of some selected hydrocarbons. AMA Arch Ind Health, 18, 9-15.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Boiling Point</b>	177 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability** Code 2. Basic data given: comparable to guidelines/standards.

**References** International Programme on Chemical Safety & The Commission of the European Communities (1993) p-Cymene. [www.inchem.org](http://www.inchem.org).

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Boiling Point</b>	176 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Fragrance Materials Association (FMA) Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Boiling Point</b>	177.1 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	CRC Handbook of Chemistry and Physics (1986) 67th edition, Robert C. Weast, editor, The Chemical Rubber Co Press, Inc., Boca Raton, Florida.

### 1.3 Vapor Pressure

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>Vapor Pressure</b>	1.46 mm Hg (194.6 Pa)
<b>Temperature</b>	25 °C

**Data Qualities Reliabilities** Reliability code 1. Reliable without restriction.

**Remarks for Data Reliability** Code 1. Comparable to guideline study.

**References** Mackay D. Bobra A., Chan D., and Shiu W. (1982) Vapor pressure correlations for low volatility environmental chemicals. Environ. Sci. Technology, 16, 645-649.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Vapor Pressure</b>	1.50 mm Hg (200 Pa)
<b>Temperature</b>	20 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	International Programme on Chemical Safety & The Commission of the European Communities (1993) <i>p</i> -Cymene. www.inchem.org.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Calculated/Antoine & Grain method
<b>Vapor Pressure</b>	1.11 mm Hg (148 Pa)
<b>Temperature</b>	25 °C
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>References</b>	MPBPVP EPI Suite (2000) U S Environmental Protection Agency.

#### 1.4 n-Octanol/Water Partition Coefficient

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured

<b>GLP</b>	No
<b>Year</b>	1980
<b>Log Pow</b>	4.1
<b>Temperature</b>	23 +/-1.5 °C
<b>Remarks for Test Conditions</b>	At a temperature of 23 +/-1.5 °C, a mixture of purified octanol and water was shaken for 30 minutes and separated by centrifugation (10,000 rpm, 30 minutes). <i>p</i> -Cymene was dissolved in the water-saturated octanol and then added to a steel tube which was then sealed and the contents were equilibrated by shaking for 4-5 minute intervals, 10 minutes apart. Afterwards, the tube was centrifuged (10,000 rpm, 30 minutes) and the octanol and water layers were sampled and analyzed by GC. The octanol sample was diluted with methanol prior to analysis. The test was conducted in duplicate.
<b>Remarks for Results</b>	Results given as K = 1.26E4 (6% standard deviation)
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Banerjee S., Yalkowsky, S., and Valvani, S.C. (1980) Water solubility and octanol/water partition coefficients of organics. Limitations of the solubility-partition coefficient correlation. Environmental Science and Technology, 14(10), 1227-1229.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	4.1
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>References</b>	KOWWIN EPI Suite (2000) U S Environmental Protection Agency, (Hansch C. et al., 1995).

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	4.19
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.

## References

Interactive Analysis LogP and LogW Predictor: Database contributed by Syracuse Research Corporation, SciVision, Albany Molecular Research, Inc., eduSoft LC, Cambridge Soft. www.logp.com.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method/guideline</b>	Calculated
<b>Log Pow</b>	3.63
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>References</b>	Mackay D., Bobra, A., Shiu, W.Y., and Yalkowsky, S.H. (1980) Relationships between aqueous solubility and octanol-water partition coefficients. Chemosphere, 9(11), 701-711.

## 1.5 Water Solubility

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>GLP</b>	No
<b>Year</b>	1980
<b>Value (mg/L) at Temperature</b>	23.35 mg/L at 25 °C
<b>Remarks for Test Conditions</b>	Distilled water was mixed with an excess of <i>p</i> -cymene by constant or intermittent shaking in a sealed stainless steel centrifuge tube and allowed to equilibrate (usually within 1 week). Afterwards, the tube was centrifuged (10,000 ppm, 60 minutes) and water samples were taken and analyzed by GC. The test was conducted at least twice and the analysis of samples was conducted in duplicate.
<b>Remarks for Results</b>	Results reported as 174 uM.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Banerjee S., Yalkowsky, S., and Valvani, S.C. (1980) Water solubility and octanol/water partition coefficients of organics. Limitations of the solubility-partition coefficient correlation. Environ. Sci. Technol., 14(10), 1227-1229.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Measured
<b>GLP</b>	No
<b>Value (mg/L) at Temperature</b>	20 mg/L at 25 °C
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	International Programme on Chemical Safety & The Commission of the European Communities (1993) <i>p</i> -Cymene. www.inchem.org.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1974
<b>Value (mg/L) at Temperature</b>	500 mg/L at 25 °C in synthetic seawater
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Price K.S., Waggy, G.T., and Conway, R.A. (1974) Brine shrimp bioassay and seawater BOD of petrochemicals. J Water Pollution Control Fed, 46(1), 63-77.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Calculated
<b>Value (mg/L) at Temperature</b>	11.675 mg/L
<b>Remarks for Results</b>	Reported as LogW = -4.06 W = 0.000087 mol/L W = 0.011675 g/L Lipinski Number: 4
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>References</b>	Interactive Analysis LogP and LogW Predictor: Database contributed by Syracuse Research Corporation, SciVision, Albany Molecular Research, Inc., eduSoft LC, Cambridge Soft. www.logp.com.



## 2 Environmental Fate and Pathways

### 2.1 Photodegradation

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Calculated
<b>Test Type</b>	AOPWIN
<b>Half-life t<sub>1/2</sub></b>	15.03 hours
<b>Remarks for Test Conditions</b>	The data are obtained by a recognized SAR method and are based upon measured OH, ozone and NO <sub>3</sub> rate constants.
<b>Remarks for Results</b>	Reaction with hydroxyl radicals. Experimental rate constant=15 x 10 <sup>-12</sup> cm <sup>3</sup> /molecule.sec
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>References</b>	AOPWIN EPI Suite (2000) U S Environmental Protection Agency.

### 2.2 Stability in Water

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method</b>	Aqueous Base/acid-catalyzed hydrolysis (25 °C) (HYDROWIN v1.67)
<b>Test Type</b>	Calculated
<b>Remarks Results</b>	Rate constant cannot be estimated for this structure. Terpene aromatic hydrocarbons contain no functional group capable of hydrolysis.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite v 3.10 (2000) US Environmental Protection Agency

### 2.3 Biodegradation

<b>Substance Name</b>	<i>p</i> -Cymene
-----------------------	------------------

<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Assay: >95%
<b>Method/guideline</b>	MITI Standard BOD procedure (synthetic seawater)
<b>Test Type</b>	Aerobic
<b>GLP</b>	No
<b>Year</b>	1997
<b>Contact Time</b>	28 days
<b>Innoculum</b>	Activated sludge
<b>Remarks for Test Conditions</b>	Test substance (100 mg/L) is inoculated and incubated with 30 mg/L of sludge. Final day biological oxygen demand (BOD) was reported as the percentage of ultimate degradation. BOD is measured from day 0 continuously to day 28. In the MITI test evidence of biodegradation was a final day BOD>5%.
<b>Degradation % After Time</b>	88% BOD after 28 days
<b>Results</b>	Biodegradation of the homologous series of substituted aromatic hydrocarbons in the MITI test was 88, 39, and 100 % for p-cymene, ethylbenzene, and methylbenzene, respectively
<b>Conclusion Remarks</b>	The authors concluded that p-cymene was biodegradable after 28 days..
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>Reference</b>	Klopman G. and Tu M. (1997) Structure-biodegradability study and computer automated prediction of aerobic biodegradation of chemicals. <i>Environmental Toxicology and Chemistry</i> <b>16</b> (9), 1829-1835.
<b>Substance Name</b>	p-Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	
<b>Method</b>	Rapid Biodegradation Model ((BIOWIN v4.00)
<b>Test Type</b>	Aerobic
<b>GLP</b>	NA
<b>Year</b>	NA
<b>Remarks Results</b>	Expert Survey Biodegradation Results-2.75 weeks by ultimate survey model and 3.51 by primary survey model. Probability of rapid biodegradation by linear model, 0.79 and by non-linear model, 0.90. Readily Biodegradation Probability by linear model, 0.38, and by non-linear model 0.44

<b>Conclusion Remarks</b>	Biodegradation rate similar to that of cumene in freshwater.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite v 3.10 (2000) US Environmental Protection Agency

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method</b>	Modified standard BOD procedure (freshwater)
<b>Test Type</b>	Aerobic
<b>GLP</b>	No
<b>Year</b>	1974
<b>Contact Time</b>	20 days
<b>Innoculum</b>	Domestic wastewater
<b>Remarks for Test Conditions</b>	Domestic wastewater was filtered through glass wool and added (3 ml/bottle) to BOD bottles. Aerated dilution water was added to fill the bottles halfway. Cumene was added to provide concentrations of 3, 7, or 10 mg/L providing a potential oxygen demand of 3-30 mg/L over 20 days. A minimum of 2 concentrations was tested in duplicate. Dissolved oxygen (DO) was monitored periodically and if it dropped below 4.0 mg/L, the bottle contents were re-aerated until a DO level of 7 mg/L was reached. Routine analysis for nitrates and nitrites was performed using the methods described by APHA (1971) with principal modifications as follows: sulfanilic acid was omitted from the color reagent and adjustments to the procedure to allow for smaller sample sizes. Results were recorded as "percent bio-oxidation" which was defined as the difference between the cumulative oxygen uptake for oxidation of the carbonaceous material in the test sample bottle from day 0 to the day of interest in mg/L and the cumulative oxygen uptake in a blank, containing the same amount and type of microbial seed as the test sample bottle, from day 0 to the day of interest in mg/L divided by the initial concentration of the test compound in mg/L times the theoretical oxygen demand or the weight ratio of oxygen required per mg of compound for complete conversion of the compound to CO <sub>2</sub> and water.
<b>Degradation % After Time</b>	40% after 5 days; 62% after 10 days; 63% after 15 days; 70% after 20 days
<b>Time required for 10% degradation</b>	Less than 5 days
<b>Remarks Results</b>	Theoretical oxygen demand = 3.50 mg/mg; measured chemical oxygen demand = 1.13 mg/mg
<b>Conclusion Remarks</b>	In freshwater, cumene showed a 70% bio-oxidation within 20 days. The authors concluded that cumene was biodegradable in freshwater.

**Data Qualities Reliabilities** Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability** Code 2. Basic data given: comparable to guidelines/standards.

**Reference** Price K.S., Waggy, G.T., and Conway, R.A. (1974) Brine shrimp bioassay and seawater BOD of petrochemicals. J Water Pollution Control Fed, 46(1), 63-77.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method/guideline</b>	Modified standard BOD procedure (synthetic seawater)
<b>Test Type</b>	Aerobic
<b>GLP</b>	No
<b>Year</b>	1974
<b>Contact Time</b>	20 days
<b>Innoculum</b>	Settled raw wastewater added to seawater
<b>Remarks for Test Conditions</b>	Wastewater was filtered through glass wool and added (3 ml/bottle) to BOD bottles. Aerated dilution water was added to fill the bottles halfway. Cumene was added to provide concentrations of 3, 7, or 10 mg/L providing a potential oxygen demand of 3-30 mg/L over 20 days. A minimum of 2 concentrations was tested in duplicate. Dissolved oxygen (DO) was monitored periodically and if it dropped below 4.0 mg/L, the bottle contents were re-aerated until a DO level of 7 mg/L was reached. Routine analysis for nitrates and nitrites was performed using the methods described by APHA (1971) with principal modifications as follows: sulfanilic acid was omitted from the color reagent and adjustments to the procedure to allow for smaller sample sizes. Results were recorded as "percent bio-oxidation" which was defined as the difference between the cumulative oxygen uptake for oxidation of the carbonaceous material in the test sample bottle from day 0 to the day of interest in mg/L and the cumulative oxygen uptake in a blank, containing the same amount and type of microbial seed as the test sample bottle, from day 0 to the day of interest in mg/L divided by the initial concentration of the test compound in mg/L times the theoretical oxygen demand or the weight ratio of oxygen required per mg of compound for complete conversion of the compound to CO <sub>2</sub> and water.
<b>Degradation % After Time</b>	3% after 10 days; 3% after 15 days; 2% after 20 days
<b>Results</b>	Theoretical oxygen demand = 3.50 mg/mg; measured chemical oxygen demand = 1.13 mg/mg
<b>Conclusion Remarks</b>	In synthetic seawater, cumene showed a virtually no bio-oxidation (2%) after 20 days. The authors concluded that cumene showed no biodegradation in synthetic seawater.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability** Code 2. Basic data given: comparable to guidelines/standards.

**Reference** Price K.S., Waggy, G.T., and Conway, R.A. (1974) Brine shrimp bioassay and seawater BOD of petrochemicals. J Water Pollut Control Fed, 46(1), 63-77.

## 2.4 Fugacity

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Model Conditions</b>	25 °C, 1,000 kg
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Level III Fugacity Model
<b>Model Used</b>	EQC Model Level III, Mackay, 1996a, 1996b
<b>Input Parameters</b>	MW, VP, log Kow, MP, water solubility
<b>Year</b>	1996
<b>Media</b>	Air
<b>Estimated Distribution and Media Concentration</b>	4.73% into air
<b>Model data and results</b>	Mass amount, half-life and emission rate
<b>Remarks</b>	At emission rate of 1000 kg/hr, half -life in air is 17 hours.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated. The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism
<b>References</b>	Mackay D., A.DiGuardo, S.Paterson, G.Kicsi and C.E.Cowan (1996a) Assessing the fate of new and existing chemicals: a five-stage process. Environ. Toxicol. Chem. 15(9): 1618-1626. Mackay D., A.DiGuardo, S.Paterson and C.E.Cowan (1996b) Evaluating the fate of a variety of of types of chemicals using the EQC model. Environmental Toxicology and Chemistry, 15(9), 1627-1637.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Model Conditions</b>	25 °C, 1,000 kg
<b>Test Type</b>	Environmental Equilibrium Partitioning Model

<b>Method</b>	Level III Fugacity Model
<b>Model Used</b>	EQC Model Level III, Mackay, 1996a, 1996b
<b>Input Parameters</b>	MW, VP, log Kow, MP, water solubility
<b>Year</b>	1996
<b>Media</b>	Water
<b>Estimated Distribution and Media Concentration</b>	27.7% into water
<b>Model data and results</b>	Mass amount, half-life and emission rate
<b>Remarks</b>	At emission rate of 1000 kg/hr, half -life in air is 360 hours.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated. The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism
<b>References</b>	Mackay D., A.DiGuardo, S.Paterson, G.Kicsi and C.E.Cowan (1996a) Assessing the fate of new and existing chemicals: a five-stage process. Environ. Toxicol. Chem. 15(9): 1618-1626. Mackay D., A.DiGuardo, S.Paterson and C.E.Cowan (1996b) Evaluating the fate of a variety of of types of chemicals using the EQC model. Environmental Toxicology and Chemistry, 15(9), 1627-1637.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Model Conditions</b>	25 °C, 1,000 kg
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Level III Fugacity Model
<b>Model Used</b>	EQC Model Level III, Mackay, 1996a, 1996b
<b>Input Parameters</b>	MW, VP, log Kow, MP, water solubility
<b>Year</b>	1996
<b>Media</b>	Soil-Water Partition Coefficient
<b>Estimated Distribution and Media Concentration</b>	65.3% into soil
<b>Model data and results</b>	Mass amount, half-life and emission rate
<b>Remarks</b>	At emission rate of 1000 kg/hr, half -life in air is 360 hours.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.

**Remarks for Data Reliability** Code 4. Calculated. The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism

**References** Mackay D., A.DiGuardo, S.Paterson, G.Kicsi and C.E.Cowan (1996a) Assessing the fate of new and existing chemicals: a five-stage process. Environ. Toxicol. Chem. 15(9): 1618-1626.  
Mackay D., A.DiGuardo, S.Paterson and C.E.Cowan (1996b) Evaluating the fate of a variety of types of chemicals using the EQC model. Environmental Toxicology and Chemistry, 15(9), 1627-1637.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Model Conditions</b>	25 °C, 1,000 kg
<b>Test Type</b>	Environmental Equilibrium Partitioning Model
<b>Method</b>	Level III Fugacity Model
<b>Model Used</b>	EQC Model Level III, Mackay, 1996a, 1996b
<b>Input Parameters</b>	MW, VP, log Kow, MP, water solubility
<b>Year</b>	1996
<b>Media</b>	Sediment-Water Partition Coefficient
<b>Estimated Distribution and Media Concentration</b>	2.22% into sediment
<b>Model data and results</b>	Mass amount, half-life and emission rate
<b>Remarks</b>	At emission rate of 0 kg/hr, half -life in air is 1440 hours.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated. The data are obtained by a recognized fugacity calculation method. Data are considered reliable with restriction because this method does not allow for biodegradation or metabolism
<b>References</b>	Mackay D., A.DiGuardo, S.Paterson, G.Kicsi and C.E.Cowan (1996a) Assessing the fate of new and existing chemicals: a five-stage process. Environ. Toxicol. Chem. 15(9): 1618-1626. Mackay D., A.DiGuardo, S.Paterson and C.E.Cowan (1996b) Evaluating the fate of a variety of types of chemicals using the EQC model. Environmental Toxicology and Chemistry, 15(9), 1627-1637.

### 3 Ecotoxicity

#### 3.1 Acute Toxicity to Fish

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Minimum purity of 80%
<b>Method/guideline</b>	"Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians" (EPA, 1975)
<b>Test Type</b>	Experimental
<b>GLP</b>	Ambiguous
<b>Year</b>	1981
<b>Species/Strain/Supplier</b>	Sheepshead minnow ( <i>Cyprinodon variegatus</i> , 8-15 mm)
<b>Exposure Period</b>	96 hour
<b>Analytical monitoring</b>	Not described.
<b>Remarks for Test Conditions</b>	Groups of 10 sheepshead minnows were used in a 96-hour static test to evaluate the potential toxicity of <i>p</i> -cymene. The test vessels were either 4-L glass jars filled with 3 L of test water (filtered [5 µm] natural seawater) or 19-L glass jars filled with 15 L test solution. No aeration was used. The use of a solvent for <i>p</i> -cymene was not described. Dissolved oxygen was measured at the beginning of the test and daily thereafter. pH was measured in the low and high concentration groups at the beginning and end of the test. Specific nominal concentrations and/or measured concentrations were not reported. LC50s at 24, 48, 72, and 96 hours were calculated with a computer program (Stephan, 1977) that determined the most appropriate statistical method (moving average angle analysis, probit analysis, or binomial probability) to apply.
<b>Endpoint value</b>	24 hour LC50 = 56 (32-100, 95% c.i.) ppm; 48 hour LC50 = 50 (38-68, 95% c.i.) ppm; 72 hour LC50 = 48 (36-64, 95% c.i.) ppm; 96 hour LC50 = 48 (36-64, 95% c.i.) ppm; NOEC=10 ppm
<b>Unit</b>	mg/L
<b>Conclusion Remarks</b>	The authors concluded that substances tested with a 96-hour LC50 ranging from 10-500 ppm were slightly toxic to practically non-toxic.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>Reference</b>	Heitmuller P.T., Hollister, T.A., and Parrish, P.R. (1981). Acute toxicity of 54 industrial chemicals to sheepshead minnows ( <i>Cyprinodon variegatus</i> ). Bull Environm Contam Toxicol., 27, 596-604.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity of 99.4%
<b>Method/guideline</b>	Acute toxicity test using guidelines from 40 CFR 797.1400 (EPA, 1985; 1987)
<b>Test Type</b>	Experimental
<b>GLP</b>	Yes
<b>Year</b>	1995
<b>Species/Strain/Supplier</b>	Rainbow trout ( <i>Oncorhynchus mykiss</i> , 31-46 mm)
<b>Exposure Period</b>	96 hour
<b>Analytical monitoring</b>	HPLC
<b>Remarks for Test Conditions</b>	Tests were conducted using a temperature-controlled water bath with a flow-through system that allowed 13 volume replacements per day. Ten organisms per test vessel were used. Six nominal test concentrations of 8.7, 13, 21, 32, 49, or 75 mg/L with a control group were run in duplicate. Filtered (0.45 um) and unfiltered water from the test vessels was sampled at the beginning, midpoint and end of each test and tested for cumene concentration. In addition, dissolved oxygen, temperature, hardness, and pH were measured daily. The photoperiod was 16:8. LC50s were calculated using a computer program (Stephan, 1983) that used mean measured concentrations and corresponding mortality data (the program used binomial interpolation, moving averages or probit depending on the data).
<b>Nominal concentrations as mg/L</b>	0, 8.7, 13, 21, 32, 49, or 75
<b>Measured concentrations as mg/L</b>	ND (less than 0.27), 0.87, 1.2, 1.9, 2.8, 4.9, or 6.4
<b>Endpoint value</b>	24 hour LC50 = 6.4 (5.5-9.3, 95% c.i.) mg/L; 48 hour LC50 = 5.8 (5.1-6.9, 95% c.i.) mg/L; 72 hour LC50 = 5.2 (4.5-6.2, 95% c.i.) mg/L; 96 hour LC50 = 4.8 (4.2-5.5, 95% c.i.) mg/L; NOEC = 1.9 mg/L
<b>Unit</b>	mg/L
<b>Remarks fields for results</b>	The measured concentrations were approximately 10% of the nominal concentrations. Water hardness, pH and temperature were 30-36 mg/L as CaCO <sub>3</sub> , 7.0, and 12 C, respectively.
<b>Conclusion Remarks</b>	The authors concluded that cumene is moderately toxic to fish but cumene's high volatility would limit its toxicological impact to an aquatic environment.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restriction.
<b>Remarks for Data Reliability</b>	Code 1. Guideline study.
<b>Reference</b>	Glickman A.H., Alexander, H.C., Buccafusco, R.J., Morris, C.R., Francis, B.O., Surprenant, D.C., and Ward, T.J. (1995) An evaluation of the aquatic hazard of cumene (isopropyl benzene). <i>Ecotoxicol Environ Saf.</i> , 31(3), 287-289.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method/guideline</b>	OECD Guideline 203
<b>Test Type</b>	Experimental
<b>GLP</b>	Ambiguous
<b>Year</b>	1993
<b>Species/Strain/Supplier</b>	Red killifish ( <i>Oryzias latipes</i> )
<b>Exposure Period</b>	96 hour
<b>Remarks for Test Conditions</b>	Groups of 10 red killifish were exposed to 5 concentrations of cumene in 2 liters of test solution at 20 °C under semi-static conditions. Specific nominal and measured concentrations were not reported. DMSO and/or dispersant (HCO-40 from Nikkou Chemicals Co.) were used if a solvent was necessary (not reported if these was necessary for cumene).
<b>Endpoint value</b>	96 hour LC50 = 18 mg/L
<b>Unit</b>	mg/L
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>Reference</b>	Yoshioka Y., and Ose, Y. (1993) A quantitative structure-activity relationship and ecotoxicological risk quotient for the protection from chemical pollutants. Environ Toxicol Water Qual., 8, 87-101.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity of 99.4%
<b>Method/guideline</b>	Acute toxicity test using guidelines from 40 CFR 797.1400 (EPA, 1985; 1987)
<b>Test Type</b>	Experimental
<b>GLP</b>	Yes
<b>Year</b>	1995
<b>Species/Strain/Supplier</b>	Sheepshead minnow ( <i>Cyprinodon varigatus</i> , 22-35 mm)
<b>Exposure Period</b>	96 hour
<b>Analytical monitoring</b>	HPLC

<b>Remarks for Test Conditions</b>	Tests were conducted using a temperature-controlled water bath with a flow-through system that allowed 13 volume replacements per day. Ten organisms per test vessel were used. Six nominal test concentrations of 23, 36, 55, 84, 130, or 200 mg/L with a control group were run in duplicate. Filtered (0.45 um) and unfiltered water from the test vessels was sampled at the beginning, midpoint and end of each test and tested for cumene concentration. In addition, dissolved oxygen, temperature, salinity, and pH were measured daily. The photoperiod was 16:8. LC50s were calculated using a computer program (Stephan, 1983) that used mean measured concentrations and corresponding mortality data (the program used binomial interpolation, moving averages or probit depending on the data).
<b>Endpoint value</b>	24 hour LC50 = 8.1 (5.6-14, 95% c.i.) mg/L; 48 hour LC50 = 5.7 (4.8-8.1, 95% c.i.) mg/L; 72 hour LC50 = 4.8 (4.5-5.2, 95% c.i.) mg/L; 96 hour LC50 = 4.7 (4.3-5.6, 95% c.i.) mg/L; NOEC = less than 2.9 mg/L
<b>Unit</b>	mg/L
<b>Nominal concentrations as mg/L</b>	0, 23, 36, 55, 84, 130, or 200
<b>Measured concentrations as mg/L</b>	ND (less than 0.16), 2.9, 4.3, 5.6, 8.1, 14, or 17
<b>Conclusion Remarks</b>	The authors concluded that cumene is moderately toxic to fish but cumene's high volatility would limit its toxicological impact to an aquatic environment.
<b>Remarks for Results</b>	The measured concentrations were approximately 10% of the nominal concentrations. Water salinity, pH and temperature were 32 ppt, 8.0, and 25 C, respectively.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restriction.
<b>Remarks for Data Reliability</b>	Code 1. Guideline study.
<b>Reference</b>	Glickman A.H., Alexander, H.C., Buccafusco, R.J., Morris, C.R., Francis, B.O., Surprenant, D.C., and Ward, T.J. (1995) An evaluation of the aquatic hazard of cumene (isopropyl benzene). <i>Ecotoxicol Environ Saf.</i> , 31(3), 287-289.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated
<b>GLP</b>	No
<b>Species/Strain/Supplier</b>	Fish
<b>Exposure Period</b>	14 days
<b>Remarks for Test Conditions</b>	Based on: Kow = 4.10, melting point = -68 °C, water solubility = 25 mg/L
<b>Endpoint value</b>	14 day LC50 = 2.671 mg/L (Neutral organics)

<b>Unit</b>	mg/L
<b>Conclusion Remarks</b>	The data are obtained by a recognized SAR calculation and are consistent with chemical structure.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite (2000) U.S. Environmental Protection Agency

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated
<b>GLP</b>	No
<b>Species/Strain/Supplier</b>	Fish
<b>Exposure Period</b>	96 hour
<b>Remarks for Test Conditions</b>	Based on: Kow = 4.10, melting point = -68 °C, water solubility = 25 mg/L
<b>Endpoint value</b>	96 hour LC50 = 1.056 mg/L (Neutral organics)
<b>Unit</b>	mg/L
<b>Conclusion Remarks</b>	The data are obtained by a recognized SAR calculation and are consistent with chemical structure.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite (2000) U.S. Environmental Protection Agency

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated
<b>GLP</b>	No
<b>Species/Strain/Supplier</b>	Fish (SW)
<b>Exposure Period</b>	96 hour
<b>Remarks for Test Conditions</b>	Based on: Kow = 4.10, melting point = -68 °C, water solubility = 25 mg/L

<b>Endpoint value</b>	96 hour LC50 = 0.668 mg/L (Neutral organics)
<b>Unit</b>	mg/L
<b>Conclusion Remarks</b>	The data are obtained by a recognized SAR calculation and are consistent with chemical structure.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite (2000) U.S. Environmental Protection Agency

### 3.2 Acute Toxicity to Aquatic Invertebrates

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Purity greater than 80%
<b>Method/guideline</b>	"Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians" (EPA, 1975)
<b>Test Type</b>	Experimental
<b>GLP</b>	Ambiguous
<b>Year</b>	1980
<b>Species/Strain/Supplier</b>	<i>Daphnia magna</i>
<b>Analytical procedures</b>	Dissolved oxygen and temperature measured with a YSI Model 54BP probe, pH measured with pH meter, and total hardness determined according to APHA et al. (1975).
<b>Test Details</b>	24 and 48 hours
<b>Remarks for Test Conditions</b>	The use of a vehicle (triethylene glycol, ethanol, acetone or dimethylformamide) was dependent on the solubility of the chemical. It was not stated whether a vehicle was used for <i>p</i> -cymene. Five to 8 concentrations were tested. Within 30 minutes of solution preparation, soluble test materials were tested with 5 <i>daphnids</i> randomly placed in 3 150 ml jars containing test solution; otherwise 15 <i>daphnia</i> were placed in 2 liter jars containing test solution. In either case, the jars were covered with plastic wrap held with an elastic band. The control consisted of the same dilution water, test conditions and test organisms, but no test substance or vehicle. Observations were made at 24 and 48 hours. LC50s and 95% confidence limits were determined using a moving average angle method, but if the data did not meet the requirements of this method a probit analysis was used and if this did not work, a binomial probability analysis was conducted. The paper did not specify which method was used for calculating the LC50s for <i>p</i> -cymene.
<b>EC50, EL50, LC0, at 24,48 hours</b>	24 hour LC50 = 9.4 mg/L (7.9-11, 95% conf.int.); 48 hour LC50 = 6.5 mg/L (4.3-10, 95% conf.int.)

<b>Unit</b>	mg/L
<b>Biological observations</b>	No discernable effect at less than 4.6 mg/L. No other description given.
<b>Remarks for Results</b>	Results were limited to tabular reporting of LC50s. Measured dissolved oxygen concentrations ranged from 6.5-9.1 mg/L, measured pH values ranged from 6.7-8.1 and 7.4-9.4 for solutions with a hardness of 72 and 173 mg CaCO <sub>3</sub> /L, respectively.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Data Reliability Remarks</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>Reference</b>	LeBlanc G.A. (1980) Acute toxicity of priority pollutants to water flea ( <i>Daphnia magna</i> ). Bull Environ Contam Toxicol., 24, 684-691.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity of 99.4%
<b>Method/guideline</b>	Acute toxicity test using guidelines from 40 CFR 797.1400 (EPA, 1985; 1987)
<b>Test Type</b>	Experimental
<b>GLP</b>	Yes
<b>Year</b>	1995
<b>Species/Strain/Supplier</b>	Mysid shrimp (<= 1 day)
<b>Analytical procedures</b>	GC Analysis
<b>Test Details</b>	96 hours
<b>Remarks for Test Conditions</b>	Tests were conducted using a temperature-controlled water bath with a flow-through system that allowed 13 volume replacements per day. Ten organisms per test vessel were used. Five nominal test concentrations of 1.8, 3.0, 4.8, 7.2, or 12.0 mg/L with a control group were run in duplicate. Filtered (0.45 um) and unfiltered water from the test vessels was sampled at the beginning, midpoint and end of each test and tested for cumene concentration. In addition, dissolved oxygen, temperature, salinity, and pH were measured daily. The photoperiod was 14:10.
<b>EC50, EL50, LC0, at 24,48 hours</b>	24 hour LC50 greater than 2.0 mg/L; 48 hour LC50 = 1.6 (1.1-2.0, 95% c.i.) mg/L; 72 hour LC50 = 1.4 (1.1-2.0, 95% c.i.) mg/L; 96 hour LC50 = 1.3 (1.1-2.0, 95% c.i.) mg/L; NOEC = 0.68 mg/L
<b>Unit</b>	mg/L
<b>Nominal concentrations as mg/L</b>	0, 1.8, 3.0, 4.8, 7.2, or 12.0
<b>Measured concentrations as mg/L</b>	ND (less than 0.005), 0.22, 0.38, 0.68, 1.1 or 2.0
<b>Biological observations</b>	The water salinity, pH and temperature were 19 ppt, 7.6, and 25 C, respectively.
<b>Appropriate statistical</b>	Yes. LC50s were calculated using a computer program (Stephan,

<b>evaluations?</b>	1983) that used mean measured concentrations and corresponding mortality data (program used binomial interpolation, moving averages or probit depending on the data).
<b>Remarks for Results</b>	The measured concentrations were approximately 10% of the nominal concentrations.
<b>Conclusion remarks</b>	In a series of acute tests with other aquatic species (rainbow trout, sheepshead minnow, and daphnia), mysid shrimp appeared to be the most sensitive with a NOEC of 0.68 mg/L. The authors concluded that cumene is moderately toxic to invertebrates but cumene's high volatility would limit its toxicological impact to an aquatic environment.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restriction.
<b>Data Reliability Remarks</b>	Code 1. Guideline study.
<b>Reference</b>	Glickman A.H., Alexander, H.C., Buccafusco, R.J., Morris, C.R., Francis, B.O., Surprenant, D.C., and Ward, T.J. (1995) An evaluation of the aquatic hazard of cumene (isopropyl benzene). <i>Ecotoxicol Environ Saf.</i> , 31(3), 287-289.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity of 99.4%
<b>Method/guideline</b>	Acute toxicity test using guidelines from 40 CFR 797.1400 (EPA, 1985; 1987)
<b>Test Type</b>	Experimental
<b>GLP</b>	Yes
<b>Year</b>	1995
<b>Species/Strain/Supplier</b>	<i>Daphnia magna</i> (<=1 day)
<b>Analytical procedures</b>	HPLC
<b>Test Details</b>	48 hours
<b>Remarks for Test Conditions</b>	Tests were conducted using a temperature-controlled water bath with a flow-through system that allowed 6 volume replacements per day. Ten organisms per test vessel were used. Five nominal test concentrations of 12, 20, 33, 55, or 91 mg/L with a control group were run in duplicate. Filtered (0.45 um) and unfiltered water from the test vessels was sampled at the beginning, midpoint and end of each test and tested for cumene concentration. In addition, dissolved oxygen, temperature, hardness, and pH were measured daily. The photoperiod was 16:8.
<b>EC50, EL50, LC0, at 24,48 hours</b>	24 hour LC50 = 4.8 (4.3-5.6, 95% c.i) mg/L; 48 hour LC50 = 4.0 (3.5-4.5, 95% c.i.) mg/L; NOEC = 1.5 mg/L
<b>Unit</b>	mg/L
<b>Nominal concentrations as mg/L</b>	0, 12, 20, 33, 55, or 91

<b>Measured concentrations as mg/L</b>	ND (less than 0.16), 1.5, 2.4, 4.0, 6.1 or 8.9
<b>Biological observations</b>	Water hardness, pH and temperature were 160-180 mg/L as CaCO <sub>3</sub> , 8.3, and 20 C, respectively.
<b>Appropriate statistical evaluations?</b>	Yes. LC50s were calculated using a computer program (Stephan, 1983) that used mean measured concentrations and corresponding mortality data (program used binomial interpolation, moving averages or probit depending on the data).
<b>Remarks for Results</b>	The measured concentrations were approximately 10% of the nominal concentrations.
<b>Conclusion remarks</b>	The authors concluded that cumene is moderately toxic to invertebrates but cumene's high volatility would limit its toxicological impact to an aquatic environment.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restriction.
<b>Data Reliability Remarks</b>	Code 1. Guideline study.
<b>Reference</b>	Glickman A.H., Alexander, H.C., Buccafusco, R.J., Morris, C.R., Francis, B.O., Surprenant, D.C., and Ward, T.J. (1995) An evaluation of the aquatic hazard of cumene (isopropyl benzene). <i>Ecotoxicol Environ Saf.</i> , 31(3), 287-289.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method/guideline</b>	Toxicity screening test/Hatching procedure
<b>Test Type</b>	Experimental
<b>GLP</b>	No
<b>Year</b>	1974
<b>Species/Strain/Supplier</b>	Brine shrimp ( <i>Artemia salina</i> )
<b>Test Details</b>	24 hours
<b>Remarks for Test Conditions</b>	To ensure survival of the shrimp in the test solution, a hatching procedure was used in which shrimp eggs were allowed to hatch and viable shrimp were removed with a medicine dropper. In the screening test, bottles containing 100, 1,000, or 10,000 mg cumene/L test solution were used. Brine shrimp suspension (1 ml) was added by pipette at a titer of 30-50 shrimp/ml. Bottles were loosely capped and maintained at 24.5 °C for 24 hours. A colony counter was used to determine the number of live and dead shrimp at the end of the test period. To determine the median tolerance limit, the same procedure was used with more specific concentrations. If the toxicity range in the screening test was 100-1,000 mg/L, the concentrations used were 100, 180, 320, 560, or 1,000 mg/L. The reviewer assumes these concentrations were used since the final median tolerance limit was within this range. If the toxicity range was less than 100 mg/L but greater than 10 mg/L, the concentrations used were 10, 1832, 56 or 100 mg/L. The percent survival versus the test dosage concentration

<b>EC50, EL50, LC0, at 24,48 hours</b>	(log scale) was plotted. The median tolerance limit was the concentration at 50% survival when a straight line was plotted. 24 hour median tolerance limit = 110 mg/L
<b>Unit</b>	mg/L
<b>Nominal concentrations as mg/L</b>	100, 180, 320, 560, or 1,000
<b>Biological observations</b>	Movement, or lack thereof, of phyllopodia (swimming appendages) was used to indicate survival (movement) or death (no movement). Clinging together of 2 or more shrimp indicated near lethal concentrations.
<b>Remarks for Results</b>	The reviewer assumes the concentrations of 100, 180, 320, 560, or 1,000 mg/L were used since the final median tolerance limit was within this range.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Data Reliability Remarks</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>Reference</b>	Price K.S., Waggy, G.T., and Conway, R.A. (1974) Brine shrimp bioassay and seawater BOD of petrochemicals. J Water Pollut Control Fed., 46(1), 63-77.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated
<b>Species/Strain/Supplier</b>	<i>Daphnia magna</i>
<b>Test Details</b>	16 days
<b>Remarks for Test Conditions</b>	Based on: Kow = 4.10, melting point = -68 °C, water solubility = 25 mg/L 16 day EC50 = 0.168 mg/L
<b>EC50, EL50, LC0, at 24,48 hours</b>	
<b>Conclusion Remarks</b>	The data are obtained by a recognized SAR calculation and are consistent with chemical structure.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Data Reliability Remarks</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite (2000) U.S. Environmental Protection Agency.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated

<b>Species/Strain/Supplier</b>	Mysid shrimp
<b>Test Details</b>	96 hours
<b>Remarks for Test Conditions</b>	Neutral organics, based on Kow = 4.10, melting point = -68 °C C, water solubility = 25 mg/L
<b>EC50, EL50, LC0, at 24,48 hours</b>	96 hour LC50 = 0.068 mg/L
<b>Conclusion Remarks</b>	The data are obtained by a recognized SAR calculation and are consistent with chemical structure.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Data Reliability Remarks</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite (2000) U.S. Environmental Protection Agency.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated
<b>Species/Strain/Supplier</b>	Daphnia magna
<b>Test Details</b>	48 hours
<b>Remarks for Test Conditions</b>	Based on: Kow = 4.10, melting point = -68 °C, water solubility = 25 mg/L
<b>EC50, EL50, LC0, at 24,48 hours</b>	48 hour LC50 = 1.309 mg/L
<b>Conclusion Remarks</b>	The data are obtained by a recognized SAR calculation and are consistent with chemical structure.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Data Reliability Remarks</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite (2000) U.S. Environmental Protection Agency.

### 3.3 Acute Toxicity to Aquatic Plants

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Assay: 99.6%
<b>Method/guideline</b>	OECD 201 Guideline
<b>Test Type</b>	Experimental

<b>GLP</b>	Yes
<b>Year</b>	2003
<b>Species/Strain/Supplier</b>	Green algae/Selenastrum capricornutum/UTEX 1648
<b>Exposure period (duration)</b>	72 hrs
<b>Analytical monitoring</b>	HPLC/UV detector
<b>Remarks for Test Conditions</b>	Green Algae/Selenastrum capricornutum/U. of Texas was maintained at test conditions for 14 days prior to the test. The culture was growing in at least 2 subcultures prior to the initiation of the test. In a range finding test, the number of cells/mL was 96% of controls at 0.10 mg/L, 94% at 1.0 mg/L, 11% at 10 mg/L, and <1% at 100 mg/L after three days. In the definitive test, algae was treated with nominal concentrations of 0, 0.65, 1.3, 2.5, 5.0 and 10.0 mg/L for 72 hours. pH was adjusted to 7.5 and solutions were exposed for 24 hours of light of intensity, 400-410 foot candles. The number of algal cells/mL as well as relative size, cell shapes, color, adherence and aggregation of cells was determined. At 24, 48, and 72 hours 3 treatment and 6 control vessels were sacrificed to determine the number of algal cells/mL. Concentrations were determined by HPLC.
<b>Nominal concentrations as mg/L</b>	0, 0.65, 1.3, 2.5, 5.0 and 10.0 mg/L
<b>Measured concentrations as mg/L</b>	Initial mean measured concentrations 0.623, 1.40, 1.91, 3.52, and 5.32 mg/L; Final measured were 53-108% of nominal concentrations
<b>Unit</b>	mg/L
<b>NOEC, LOEC or NOEL, LOEL</b>	72 hr EC <sub>50</sub> =4.03 mg/L based on average specific growth rate; 72-hr EC <sub>50</sub> =2.40 mg/L calculated using the number of cells/mL; 72-hr EC <sub>50</sub> = 2.04 mg/L using the area under the growth curve. The 72-hr NOEC=1.40 mg/L based on number of cells/mL
<b>Biological observations</b>	Control algal populations grew at an acceptable rate (10,000 cells/ml) after 72 hours. Incubation temperatures were in the range from 23.2 to 24.0 C over the 72 hours and pH was unchanged by the test substance. At the conclusion of the test, samples of test media from each test vessel with maximal growth inhibition were combined with fresh media. After 48 hours incubation the number of cells increased from 1500 cells/mL to 1,1328,000 cells/mL at 3.52 mg/L suggesting that the toxic effects were algistatic.
<b>Appropriate statistical evaluations?</b>	EC <sub>50</sub> values determined by weighted least squares non-linear regression (Bruce and Versteeg, 1992); NOEC was determined using a one-way analysis of variance (ANOVA) and Bonferroni's test (Gulley et al. 1990)
<b>Conclusion remarks</b>	The acute toxicity of <i>p</i> -cymene measured as a 50% decrease in growth and reproduction of freshwater algae was estimated to be 72 hr EC <sub>50</sub> =4.03 mg/L based on average specific growth rate; 72-hr EC <sub>50</sub> =2.40 mg/L calculated using the number of cells/mL; 72-hr EC <sub>50</sub> =2.01 mg/L using the area under the growth curve. The 72-hr NOEC=1.40 mg/L

<b>Reliabilities</b>	Relability code 1. Reliable without restrictions.
<b>Remarks for Data Reliability</b>	OECD 201 Guideline study
<b>References</b>	Ward T. (2003) The growth and reproduction toxicity test with <i>p</i> -cymene and freshwater alga, <i>Selenastrum capricornutum</i> . OECD 201. Study No. 2471-FF. Private Communication to FFHPVC. Unpublished Report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	ECOSAR
<b>Test Type</b>	Calculated
<b>Species/Strain/Supplier</b>	Green algae
<b>Exposure Period</b>	96 hours
<b>Remarks for Test Conditions</b>	Based on: Kow = 4.10, melting point = -68 °C, water solubility = 25 mg/L
<b>Endpoint value</b>	96 hour EC50 = 0.923 mg/L
<b>Conclusion Remarks</b>	The data are obtained by a recognized SAR calculation and are consistent with chemical structure.
<b>Data Qualities Reliabilities</b>	Reliability code 4. Not assignable.
<b>Remarks for Data Reliability</b>	Code 4. Calculated.
<b>Reference</b>	ECOSAR EPI Suite (2000) U.S. Environmental Protection Agency.

## 4 Human Health Toxicity

### 4.1 Acute Toxicity

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Deichmann and LeBlanc, 1943
<b>Test Type</b>	Acute oral LD50
<b>GLP</b>	No
<b>Year</b>	1961
<b>Species/strain</b>	Rat
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	1-3 rats/dose
<b>Vehicle</b>	None
<b>Route of Administration</b>	Oral-Gavage
<b>Remarks for Test Conditions</b>	Groups of rats were gavaged with 620, 940, 1400, 2100, 3200, 4700, 7100, or 10700 mg/kg bw and studied for clinical signs and mortality. Surviving animals were killed at 2 weeks. Necropsies were conducted on all rats.
<b>Value LD50 or LC50 with confidence limits</b>	3200 mg/kg bw
<b>Number of deaths at each dose level</b>	At doses of 620 to 2100 mg/kg bw, all rats survived. At 3200, 4700, 7100, and 10700 mg/kg bw, 1/2, 2/2, 3/3, and 1/1 rats died, respectively.
<b>Remarks for Results</b>	Prior to death, rats showed typical signs of intoxication: depression, tremor, lethargy, and muscular weakness. Necropsy was reported to show hyperemic lungs with scattered areas of hemorrhage, atelectasis and emphysema, partially digested blood and food in the stomach, petechial hemorrhages in the glandular stomach with hyperemic mucosa, bloody mucus in the upper small intestine and clear mucus in the lower small intestine, pale and mottled liver, congested liver, and distended urinary bladder. Some animals had blood-tinged urine or contained "suspended dark solid material resembling precipitated hemoglobin".
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	MacDonald W.E. (1961) Report on the determination of the approximate lethal oral dose in the rat of compounds submitted by the Hercules Powder Co. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Litchfield and Wilcoxon, 1949
<b>Test Type</b>	Acute oral toxicity
<b>GLP</b>	No
<b>Year</b>	1964
<b>Species/strain</b>	Rat/Osborne-Mendel
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	10
<b>Vehicle</b>	None
<b>Route of Administration</b>	Oral
<b>Remarks for Test Conditions</b>	Groups of 10 male and 10 female Osborne-Mendel rats were orally administered <i>p</i> -cymene at various doses (not specified) to calculate an oral LD50. Rats were monitored for up to 2 weeks.
<b>Value LD50 or LC50 with confidence limits</b>	LD50 = 4750 mg/kg bw (95% confidence limits: 3720-6060)
<b>Remarks for Results</b>	Rats showed depression shortly following dosing and also coma, bloody lacrimation, diarrhea with irritable, scrawny appearance during the observation period. The LD50 was calculated to be 4750 mg/kg bw with a slope function of 1.7.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenner P.M., Hagan, E.C., Taylor, J.M., Cook, E.L., and Fitzhugh, O.G. (1964) Food flavourings and compounds of related structure. I. Acute oral toxicity. <i>Fd Cosmet Toxicol</i> 2:327-343.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method/guideline</b>	Thompson
<b>Test Type</b>	Acute oral LD50
<b>GLP</b>	No
<b>Year</b>	1951
<b>Species/strain</b>	Rat
<b>Sex</b>	Not reported

<b>Route of Administration</b>	Oral
<b>Value LD50 or LC50 with confidence limits</b>	2910 mg/kg bw (95% C.I., 2550-3320 mg/kg bw)
<b>Remarks for Results</b>	The limits (+/- 1.96 standard deviations: approx. 95% Confidence Interval) were calculated by the method of Thompson. The LD50 was calculated after 14 days.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Comparable to guideline study with acceptable restrictions. One of a series of range-finding studies.
<b>References</b>	Smyth H.F., Jr., Carpenter, C.P., and Weil, C.S. (1951) Range-finding toxicity data: List IV. Arch Ind Hyg Occup. Med., 4, 119-122.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 98%
<b>Method/guideline</b>	Single-dose toxicity
<b>Test Type</b>	Acute oral/gavage
<b>GLP</b>	No
<b>Year</b>	1956
<b>Species/strain</b>	Rat/Wistar
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	20
<b>Vehicle</b>	Olive oil emulsified with gum arabic
<b>Route of Administration</b>	Oral-Gavage
<b>Remarks for Test Conditions</b>	Groups of Wistar rats were gavaged with cumene (specific doses not reported) to determine an oral LD50. After dosing rats were observed for up to 2 weeks.
<b>Value LD50 or LC50 with confidence limits</b>	1400 mg/kg bw
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Wolf M.A., Rowe, V.K., McCollister, D.D., Hollingsworth, R.L., and Oyen, F. (1956) Toxicological studies of certain alkylated benzenes and benzene. AMA Arch Ind Health, 14, 387-398.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6

<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method/guideline</b>	Thompson
<b>Test Type</b>	Acute dermal toxicity
<b>GLP</b>	No
<b>Year</b>	1951
<b>Species/strain</b>	Rabbit
<b>Sex</b>	Not reported
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	The limits (+\ - 1.96 standard deviations: approximately 95% confidence interval) were calculated by the method of Thompson. The LD50 was calculated after 14 days.
<b>Value LD50 or LC50 with confidence limits</b>	LD50 = 12.3 ml/kg bw (95% C.I. 7.69-19.7 ml/kg bw) or LD50 = 10,545 mg/kg bw
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Comparable to guideline study with acceptable restrictions. One of a series of range-finding studies.
<b>References</b>	Smyth H.F., Jr., Carpenter, C.P., and Weil, C.S. (1951) Range-finding toxicity data: List IV. Arch Ind Hyg Occup Med., 4, 119-122.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.94%
<b>Test Type</b>	Single-exposure neurobehavioral test
<b>GLP</b>	Ambiguous
<b>Year</b>	1995
<b>Species/strain</b>	Rat/Fischer 344/NHSD
<b>Sex</b>	Male and Female
<b># of animals per sex per dose</b>	10
<b>Vehicle</b>	None
<b>Route of Administration</b>	Inhalation
<b>Remarks for Test Conditions</b>	Groups of 10 male and 10 female rats underwent a single exposure to atmospheres containing 0, 100, 500, or 1200 ppm cumene for 6 hours. Body weights were measured prior to exposure and at 1, 6, and 24 hours post exposure. A functional observational battery was also conducted at these times.
<b>Remarks for Results</b>	No effects were reported at 100 ppm in both groups of male and female rats. Cumene exposure was reported to produce alterations in

the functional observational battery 1 hour post-exposure including increased incidence and severity of gait abnormalities in high-dose males, increased horizontal activity in both male and female high-dose rats and in female rats exposed to 500 ppm cumene, and decreased rectal temperature of high-dose rats of both sexes. At 6 hours post-exposure, alterations were limited to decreased toe pinch withdrawal reflexes in males rats exposed to 500 or 1200 ppm cumene. At 24 hours post-exposure, no significant differences in the functional observational battery were observed. Body weights were not affected by cumene exposure.

**Data Qualities Reliabilities**

Reliability code 2. Reliable with restriction.

**Data Reliabilities Remarks**

Code 2. Acceptable, well-documented.

**References**

Cushman J.R., Norris, J.C., Dodd, D.E., Darmer, K.I., Morris, C.R. (1995) Subchronic inhalation toxicity and neurotoxicity assessment of cumene in Fischer 344 rats. *J Am Coll Toxicol.*, 14(2), 129-147.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Test Type</b>	Acute dermal LD50
<b>GLP</b>	No
<b>Year</b>	1973
<b>Species/strain</b>	Rabbit
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	10
<b>Route of Administration</b>	Dermal
<b>Remarks for Test Conditions</b>	Ten rabbits were dermally treated with 5000 mg/kg bw and observed for 14 days.
<b>Value LD50 or LC50 with confidence limits</b>	Greater than 5000 mg/kg bw
<b>Number of deaths at each dose level</b>	0
<b>Remarks for Results</b>	No rabbits died. Skin irritation was graded as follows: slight redness (3/10), moderate redness (7/10), slight edema (3/10), and moderate edema (7/10).
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Moreno O.M. (1973) Acute dermal toxicity in rabbits. <i>p</i> -Cymene. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6

<b>Test Type</b>	Acute dermal
<b>GLP</b>	No
<b>Year</b>	1962
<b>Species/strain</b>	Rabbit/Albino
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	1
<b>Route of Administration</b>	Dermal
<b>Vehicle</b>	None
<b>Value LD50 or LC50 with confidence limits</b>	LD50 greater than 6 ml/kg bw or greater than 5144 mg/kg bw
<b>Remarks for test conditions</b>	Undiluted <i>p</i> -cymene was applied to the shaven abdominal skin (10 x 15 cm area) of an albino rabbit. <i>p</i> -Cymene was applied in 1 ml doses every hour for a total of 6 ml over a 6-hour exposure period. The rabbit was observed for 1 month following treatment.
<b>Remarks for Results</b>	Slight hyperemia of the skin was observed after 1 hour and persisted approximately 4 hours after which a slight subcutaneous edema developed. After the exposure period, the skin still was slightly edematous and over the next 5 days, it was slightly thickened, hyperemic and showed fine cracks. After the first week, the skin began to return to normal and within the month it was normal with hair growth.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	MacDonald W.E. (1962a) Acute effects of Hercules compounds applied to the skin of the rabbit. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Test Type</b>	Inhalation toxicity
<b>GLP</b>	No
<b>Year</b>	1962
<b>Species/strain</b>	Guinea pig
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	2
<b>Route of Administration</b>	Inhalation
<b>Remarks for Test Conditions</b>	Guinea pigs were exposed to atmospheres saturated with 9.7 mg <i>p</i> -cymene/l for a period of 5 hours. Clinical signs and mortality were recorded. Surviving animals were removed from the exposure

**Number of deaths at each dose level**  
**Remarks for results**

chamber and observed for an additional week. A "lethal concentration time value (LCt)" was calculated based on the "shortest period of exposure causing death", where the concentration was expressed as mg/l and time as min.  
 0/2

**Data Qualities Reliabilities**

Signs reported during the first 30 minutes were those typical of irritation: excitement, pawing at the eyes and nose, increased blinking, squinting, and eye closure. Approximately 90 minutes following exposure, 1 guinea pig had a 10-15-second violent clonic convulsion followed by prolonged quivering. Afterwards, this guinea pig continued to exhibit clonic convulsions of varying degrees. The second guinea pig began quivering at about 120 minutes into the exposure and had a clonic convulsion about 30 minutes later. By the end of the exposure period, both guinea pigs were comatose and had continuous clonic convulsions. The morning after the exposure, the guinea pigs appeared fully recovered.  
 Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability**

Code 2. Basic data given: comparable to guidelines/standards.

**References**

MacDonald W.E. (1962b) Report on the effects in laboratory animals exposed for five hours to air saturated with the vapors of compounds submitted by the Hercules Powder Company. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
-----------------------	------------------

**CAS No.** 99-87-6

**Test Type** Inhalation toxicity

**GLP** No

**Year** 1962

**Species/strain** Rat

**Sex** Not reported

**# of animals per sex per dose** 3

**Route of Administration** Inhalation

**Remarks for Test Conditions**

Rats were exposed to atmospheres saturated with 9.7 mg /L of *p*-cymene for a period of 5 hours. Clinical signs and mortality were recorded. Surviving animals were removed from the exposure chamber and observed for an additional week. A "lethal concentration time value (LCt)" was calculated based on the "shortest period of exposure causing death", where the concentration was expressed as mg/l and time as minutes.

**Number of deaths at each dose level**

0/3

**Remarks for results**

Signs reported during the first 30 minutes were those typical of irritation: excitement, pawing at the eyes and nose, increased blinking, squinting, and eye closure. After 45 minutes, equilibrium loss and increased salivation were noted. One-half hour later, fine tremors

began and increased to quivering after another 15 minutes. Clonic convulsions were reported after another 15 minutes and the rats staggered about aimlessly until the end of the exposure. The morning after exposure, the rats appeared fully recovered.  
Reliability code 2. Reliable with restriction.

**Data Qualities Reliabilities**

**Data Reliabilities Remarks**

Code 2. Basic data given: comparable to guidelines/standards.

**References**

MacDonald, W.E. (1962b) Report on the effects in laboratory animals exposed for five hours to air saturated with the vapors of compounds submitted by the Hercules Powder Company. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Test Type</b>	Inhalation toxicity
<b>GLP</b>	No
<b>Year</b>	1962
<b>Species/strain</b>	Mouse
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	3
<b>Route of Administration</b>	Inhalation
<b>Remarks for Test Conditions</b>	Mice were exposed to atmospheres saturated with 9.7 mg /L of <i>p</i> -cymene for a period of 5 hours. Clinical signs and mortality were recorded. Surviving animals were removed from the exposure chamber and observed for an additional week. A "lethal concentration time value (LCt)" was calculated based on the "shortest period of exposure causing death", where the concentration was expressed as mg/l and time as min.
<b>Value LD50 or LC50 with confidence limits</b>	LCt = 2270 mg x min/L for 3.9 hour exposure.
<b>Number of deaths at each dose level</b>	3/3
<b>Remarks for Results</b>	Signs reported during the first 30 minutes were those typical of irritation: excitement, pawing at the eyes and nose, increased blinking, squinting, and eye closure. In addition, mice exhibited equilibrium loss and clonic convulsions with intervals of coma. One mouse died after 3.9 hours and another died after 4.8 hours. The 3rd mouse was comatose at termination of exposure and died during the night. Necropsies showed hyperemic lungs, mottled liver, and pale kidneys. In addition, it appeared that the heart had stopped in systole. No effects were reported in rats and guinea pigs at the same atmospheric concentration.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.

**References**

MacDonald W.E. (1962b) Report on the effects in laboratory animals exposed for five hours to air saturated with the vapors of compounds submitted by the Hercules Powder Company. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Test Type</b>	Inhalation toxicity
<b>GLP</b>	No
<b>Year</b>	1951
<b>Species/strain</b>	Rat
<b>Sex</b>	Not reported
<b># of animals per sex per dose</b>	6
<b>Route of Administration</b>	Inhalation
<b>Remarks for Test Conditions</b>	Rats were exposed to atmospheres containing 8000 ppm cumene for 4 hours. Mortality over 14 days was reported.
<b>Number of deaths at each dose level</b>	4/6
<b>Remarks for Results</b>	Four out of 6 rats were reported to have died within the 14-day period. Atmospheric concentration (8000 mg/L) greater than measured saturation value of 9.7 mg/L.
<b>Data Qualities Reliabilities</b>	Reliability code 3. Not reliable.
<b>Remarks for Data Reliability</b>	Does not meet important criteria of current standard methods.
<b>References</b>	Smyth H.F., Jr., Carpenter, C.P., and Weil, C.S. (1951) Range-finding toxicity data: List IV. Arch Ind Hyg Occup Med., 4, 119-122.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Test Type</b>	Acute inhalation toxicity
<b>GLP</b>	No
<b>Year</b>	1958
<b>Species/strain</b>	Rat/Long Evans
<b>Sex</b>	Male
<b># of animals per sex per dose</b>	8
<b>Route of Administration</b>	Inhalation

<b>Remarks for Test Conditions</b>	Groups of 8 male Long Evans rats were exposed to atmospheres containing 5000 to 10,000 ppm cumene for 4 exposures of 30, 20, 45, and 50 minutes duration. Twenty-four hours after exposure, the rats were killed and the brain, spinal cord and 1 sciatic nerve were removed and placed in 10% formalin.
<b>Number of deaths at each dose level</b>	5 out of 8 rats died. No further information was reported.
<b>Remarks for Results</b>	Cumene exposure resulted in local irritation, depression, and quivering or twitching. At necropsy, no gross or microscopic effects were reported other than those associated with respiratory irritation. Note: Exposure levels exceeded measured saturation levels of 9.7 mg/L. Therefore animals were exposed to liquid p-cymene suspended in test atmosphere
<b>Data Qualities Reliabilities</b>	Reliability code 3. Not reliable.
<b>Remarks for Data Reliability</b>	Does not meet important criteria of current standard methods.
<b>References</b>	Furnas D.W., and Hine, C.H. (1958) Neurotoxicity of some selected hydrocarbons. AMA Arch Ind Health,18, 9-15.

## 4.2 Genetic Toxicity

### 4.2.1 In vitro Genotoxicity

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue, cumene, assay>98%
<b>Method/guideline</b>	Preincubation Ames assay (Haworth et al., 1983)
<b>Test Type</b>	Ames reverse mutation
<b>System of Testing</b>	Bacterial
<b>GLP</b>	Ambiguous
<b>Year</b>	Undated
<b>Species/Strain</b>	<i>Salmonella typhimurium</i> strains TA97, TA98, TA100, and TA1535
<b>Metabolic Activation</b>	S9 from Aroclor 1254-induced Sprague-Dawley rat or Syrian hamster
<b>Doses/Concentration</b>	TA97 & TA1535: 1, 3, 10, 33, 100, or 166 ug/plate; TA100 & TA98: 1, 3, 10, 33, 100, 166, or 333 ug/plate
<b>Statistical Methods</b>	Positive response defined as: a reproducible, dose related increase in histidine-independent (revertant) colonies".
<b>Remarks for Test Conditions</b>	<i>Salmonella typhimurium</i> strains TA1535 and TA97 were incubated with up to 166 ug cumene/plate using the standard preincubation Ames assay. Similarly, <i>Salmonella typhimurium</i> strains TA100 and TA98 were incubated with up to 333 ug cumene/plate. Each cumene concentration in each strain was tested with and without metabolic activation consisting of 10 or 30% S9 from hamster liver homogenate

or 10 or 30% S9 from rat liver homogenate.

<b>Results</b>	Cumene did not increase the number of revertants in any of the strains tested.
<b>Cytotoxic concentration</b>	Not given
<b>Genotoxic Effects</b>	None
<b>Conclusion Remarks</b>	Cumene had no mutagenic activity in this assay.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	NTP unpublished results (e). <i>Salmonella</i> Testing Results. Cumene. Cellular and Genetic Toxicology Branch, National Toxicology Program.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue, cumene, assay >98%
<b>Method/guideline</b>	Ames preincubation assay (Yahagi et al., 1975)
<b>Test Type</b>	Ames reverse mutation
<b>System of Testing</b>	Bacterial
<b>GLP</b>	Yes
<b>Year</b>	1987
<b>Species/Strain</b>	<i>Salmonella typhimurium</i> strains TA98, TA100, TA1535 and TA1537
<b>Metabolic Activation</b>	S9 from Aroclor 1254-induced male Sprague-Dawley rat (10% homogenate/ml)
<b>Doses/Concentration</b>	33, 67, 100, 333, 667, 1,000, or 2,000 ug/plate
<b>Remarks for Test Conditions</b>	Cumene, at concentrations of 33, 67, 100, 333, 667, 1,000, or 2,000 ug/plate, was tested in the Ames preincubation assay using <i>Salmonella typhimurium</i> strains TA98, TA100, TA1535, and TA1537 both with and without metabolic activation. Pluronic F127 was used to emulsify cumene. The test was also run with cumene in water rather than F127. Positive controls used were 2 ug 2-aminoanthracen, 5 ug 2-nitrofluorene, 2.5 ug sodium azide, and 75 ug 9-aminoacridine. The study was conducted in duplicate.
<b>Results</b>	Cumene did not affect the number of revertants in any of the strains tested. Cumene showed signs of cytotoxicity (reduced background) at 2,000 ug/plate.
<b>Cytotoxic concentration</b>	2,000 ug/plate
<b>Genotoxic Effects</b>	None
<b>Conclusion Remarks</b>	Cumene showed no mutagenic activity when tested in <i>Salmonella typhimurium</i> strains TA98, TA100, TA1535 and TA1537.

**Data Qualities Reliabilities** Reliability code 1. Reliable without restriction.

**Remarks for Data Reliability** Code 1. Comparable to guideline study.

**References** Lawlor T.E., and Wagner, V.O. (1987) *Salmonella*/mammalian-microsome preincubation mutagenicity assay (Ames test). Test article: Cumene. Final Report. Microbiological Associates, Inc. Report No. T4786.502009.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue, cumene
<b>Method/guideline</b>	HGPRT assay
<b>Test Type</b>	Mutation
<b>System of Testing</b>	Non bacterial
<b>GLP</b>	Yes
<b>Year</b>	1985
<b>Species/Strain</b>	Chinese hamster ovary cells (K-1)
<b>Metabolic Activation</b>	Liver enzymes from Aroclor 1254-induced rats (S9)
<b>Doses/Concentration</b>	Trial 1:8, 16, 32, 64, 128, 150, or 175 ug/ml Trial 2: 150 or 175 ug/ml
<b>Remarks for Test Conditions</b>	Cultured Chinese hamster ovary cells were exposed to 8, 16, 32, 64, 128, 150, or 175 ug cumene/ml for 5 hours with and without metabolic activation (S9). Cumene was emulsified with Pluronic F127 for a final concentration of 0.04-0.05% Pluronic F127. Each treatment group consisted of 3 flasks. After the 5-hour exposure, 200 cells from each group were plated (4 plates/group), incubated, fixed and stained. In addition, 10E5-10E6 cells were seeded to larger plates on day 3. This process was repeated 3 times with the last on day 10. On day 10, 200 cells were seeded to 4 viability plates/group and 2x10E5 cells/group were plated for the mutagenicity test. The plates were incubated to day 17, fixed and stained. Ethyl methanesulfonate was used as a positive control at 100 ug/ml. Benzo(a)pyrene was used to test the enzyme system. Negative controls were untreated cells and cells exposed to the emulsifier (F127) with or without metabolic activation. Cells were counted with a Coulter Model ZB cell counter and colonies were counted either visually or with an Artek Model 981 colony counter. Results were considered positive if there was a significant (p less than 0.05) increase in mutant colonies and the response was dose related. If only one of the above criteria were met the results were considered equivocal. A second trial was conducted with S9 at 150 or 175 ug cumene/ml.
<b>Results</b>	In cultures without S9, the number of mutants/10E6 clonable cells for untreated control (medium), F127, 8, 16, 32, 64, and 128 ug cumene/ml, and ethyl methanesulfonate were (standard deviation in brackets) 14.8 (9.1), 4.4 (6.2), 3.0 (2.6), 12.2 (11.5), 14.0 (12.2), 5.7

(5.6), 0, and 140.5 (14.3). In activated cultures the number of mutants/10E6 clonable cells for untreated controls (medium), F127, 64, 128, 150, and 175 ug cumene/ml, and benzo(a)pyrene were (standard deviation in brackets) 12.9 (8.1), 24.9 (8.9), 2.1 (3.6), 7.5 (10.4), 4.1 (7.0), 266.8 (457.8), and 48.0 (31.4), respectively. The result for 175 ug/ml was high due to a single outlier value in the group; hence, a second trial was conducted. Mutant frequencies appeared to be high in the medium and Pluronic F127 control groups. There was no statistically significant increase in the number of mutants or a dose-response effect. Cumene was cytotoxic at concentrations of 128 ug/ml and higher. In the second trial using concentrations of 150 and 175 ug/ml, the increase in the number of mutants seen at 175 ug/ml was not repeated. In the 1st trial, cloning efficiency was significantly decreased only in activated cultures at concentrations of 128 ug/ml and higher. Cloning efficiency was not affected in non-activated cultures.

<b>Cytotoxic concentration</b>	128 ug/ml
<b>Genotoxic Effects</b>	None.
<b>Appropriate statistical evaluations?</b>	2-tailed t-test using MUTANT program
<b>Conclusion Remarks</b>	Cumene did not increase mutations in the CHO/HGPRT test with or without metabolic activation.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restriction.
<b>Remarks for Data Reliability</b>	Code 1. Comparable to guideline study.
<b>References</b>	Papciak R.J. (1985) CHO/HGPRT test of cumene. Project #84-2128. Gulf Life Sciences Center. Pittsburgh, PA. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue, cumene
<b>Method/guideline</b>	HGPRT assay
<b>Test Type</b>	Mutation
<b>System of Testing</b>	Non bacterial
<b>GLP</b>	Yes
<b>Year</b>	1987
<b>Species/Strain</b>	Chinese hamster ovary cells (K-1)
<b>Metabolic Activation</b>	Liver enzymes from Aroclor 1254-induced rats (S9)
<b>Doses/Concentration</b>	100, 125, 150, 175, 200, or 225 ug/ml
<b>Remarks for Test Conditions</b>	In preliminary toxicity tests, cultured Chinese hamster ovary cells were exposed to up to 225 ug/ml of cumene for 5 or 18 hours with and without metabolic activation (S9). In the main study, cultured Chinese hamster ovary cells were exposed to 100, 125, 150, 175, 200, or 225

ug/ml of cumene for 18 hours without S9 or for 5 hours with S9. Cumene was emulsified with Pluronic F127. Ethyl methanesulfonate was used as a positive control at 0.2 ul/ml. Benzo(a)pyrene (4 ug/ml) was used to test the enzyme system. Negative and solvent controls were untreated cells and cells exposed to the emulsifier (F127) with or without metabolic activation. Each treatment was conducted in duplicate or triplicate and colonies were counted. Results were considered positive if there was a dose-dependent increase in mutant frequency in one of the 5 tested concentrations which is at least twice that of the solvent control and untreated control, and is also increased above that of the solvent and untreated control by at least 8.7 mutants/10E6 clonable cells. If only one of the above criteria were met the results were considered equivocal.

**Results**

In cultures without S9, the number of mutants/10E6 clonable cells for untreated control (medium), F127, 100, and 125 ug/ml of cumene, ethyl methanesulfonate with F127, and ethyl methanesulfonate were less than 1.1 and 2.0; 4.3 and 7.1; 14.9 and 3.4; 5.4 and less than 1.1; 537.5 and 490.2; and 784.1 and 595.0. Cultures treated with 150 ug/ml and higher were too toxic to count. In activated cultures the number of mutants/10E6 clonable cells for untreated controls (medium), F127, 100, 125, and 225 ug/ml of cumene, benzo(a)pyrene with F127, and benzo(a)pyrene were 15.5 and 4.8; 1.7 and 6.8; 19.6 and 3.5; 12.9 and 2.3; 27.6 and 10.1; 350.0 and 323.7; and 347.6 and 326.4. Cultures treated with 150-200 ug/ml were too toxic to count. There was no significant increase in the number of mutants or a dose-response effect. Cumene was cytotoxic at concentrations of 150 ug/ml and higher.

**Cytotoxic concentration**

150 ug/ml

**Genotoxic Effects**

None

**Remarks for Results**

The authors noted that the variability shown in cytotoxicity and mutation frequencies was a result of difficulty in handling cumene, suspending it in F127, and delivering small quantities into the test medium. However, the study was considered valid since it met the validation criteria: cloning efficiency of solvent and untreated controls was greater than 50%; spontaneous mutant frequency of solvent and untreated controls is between 0 and 20 mutants per 10E6 clonable cells; and the positive control must induce a mutant frequency of at least 3 times that of the solvent control.

**Conclusion Remarks**

Cumene did not increase mutations in the CHO/HGPRT test with or without metabolic activation.

**Data Qualities Reliabilities**

Reliability code 1. Reliable without restriction.

**Remarks for Data Reliability**

Code 1. Comparable to guideline study.

**References**

Yang L.L. (1987) CHO/HGPRT Mutation Assay. Cumene. Internal Report dated June 1, 1987. #T4786.332010. Microbiological Associates Inc. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Method/guideline</b>	Paper disk method (Lyer and Szybalski, 1958)

<b>Test Type</b>	Reverse mutation
<b>System of Testing</b>	Bacterial
<b>GLP</b>	No
<b>Year</b>	1958
<b>Species/Strain</b>	<i>Escherichia coli</i> strain Sd-4-73
<b>Metabolic Activation</b>	No
<b>Doses/Concentration</b>	Not reported
<b>Remarks for Test Conditions</b>	<i>E. coli</i> was cultured overnight at 36 C in an aerated nutrient broth containing 20 ug/ml streptomycin. Plates were prepared and <i>p</i> -cymene was added by applying to a paper disk (0.01-0.025 ml or small crystal), which was then placed on the agar. Relative mutagenicity, defined as "an approximate ratio of the number of colonies on the plate containing the mutagen to the number of colonies on the control plate, was calculated. Potent mutagens had relative mutagenicities of greater than 3 and weak and doubtful mutagens had relative mutagenicities between 1.5 and 3.
<b>Results</b>	<i>p</i> -Cymene produced no increase in the frequency of reversion from streptomycin dependence to independence in Sd-4-73 <i>E. coli</i> .
<b>Genotoxic Effects</b>	None
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Szybalski W. (1958) Special microbiological systems. II. Observations on chemical mutagenesis in microorganisms. Annals of New York Academy of Sciences. Pp 475-489.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue, cumene, 99.7% pure
<b>Test Type</b>	Chromosomal aberration test
<b>GLP</b>	Yes
<b>Year</b>	1987
<b>Species/Strain</b>	Chinese hamster ovary cells
<b>Metabolic Activation</b>	Aroclor 1254 induced Sprague-Dawley rat liver microsomes
<b>Doses/Concentration</b>	0, 19, 31, 49, 78, 125, or 200 ug/ml without S-9 activation and 0, 24, 38, 61, 98, 156, or 225 with activation
<b>Statistical Methods</b>	Student's t test (p less than or = 0.05)
<b>Remarks for Test Conditions</b>	The test article was tested for effects on cell cycle. Duplicate cultures were treated with the culture medium (negative control), the test

article alone and three concentrations of the positive control article (triethylenemelamine and cyclophosphoramide) with and without activation. One culture was harvested at first metaphase division for evaluation of chromosomal aberrations. CHO cells were seeded at  $5 \times 10^5$  cells/25 cm<sup>2</sup> flasks and incubated at 37 C for 14-16 hours. Flasks were then treated with 5 ml of test article. After exposure of 8 or 14 hours and two hours prior to harvest, the treatment medium was removed and cells were washed with PBS and refed with growth medium containing 0.1 ug/ml of colcemid. In the S-9 activated experiment, cells were exposed for 2 hours. Again, cells were separated, washed, refed, and treated with colcemid. Two hours after addition of colcemid, cells were collected and fixed. Fifty metaphase cells were scored in each duplicate flask. Cells were evaluated for a range of chromosomal changes. The second culture was treated with 0.01 mM BrdU two hours after initiation and cells were harvested 24-26 hours later for evaluation of cell cycle.

**Results**

Toxicity was reported at the highest dose tested with or without S-9 activation. A statistically significant increase in chromosomal aberration was reported at 156 ug/ml in the presence of S-9 compared to the vehicle control. No statistically significant increase was observed when compared to untreated control cells. The increase was within the historical control range for the contract laboratory. The authors concluded that the increase was due to low vehicle control values

**Cytotoxic concentration**

200 ug/ml

**Genotoxic Effects**

None

**Appropriate statistical evaluations?**

Yes

**Conclusion Remarks**

Cumene did not induces structural or numerical chromosomal aberrations in Chinese hamster ovary cells with or without S-9 activation

**Data Qualities Reliabilities**

Reliability code 1. Reliable without restriction.

**Remarks for Data Reliability**

Code 1. Guideline study.

**References**

Putnam D.L. (1987) Chromosome aberrations in Chinese hamster ovary cells. Laboratory Study No. T4786.337012. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue, cumene
<b>Test Type</b>	DNA Repair assay
<b>System of Testing</b>	Non bacterial
<b>GLP</b>	Yes
<b>Year</b>	1984
<b>Species/Strain</b>	F344 rat hepatocyte

<b>Doses/Concentration</b>	8, 16, 32, 64, 128, 256, 512, 1024, 2048, or 5,000 ug/ml
<b>Remarks for Test Conditions</b>	Cultured rat hepatocytes were treated with 8, 16, 32, 64, 128, 256, 512, 1024, 2048, or 5,000 ug/ml of cumene in triplicate. Nuclear grain counts (a count of 6 or more over the negative controls was considered positive) and the percentage of repair-positive cells was determined. Negative controls of medium and a 10% solution of Pluronic F68 (emulsifier) were used. 2-Acetylaminofluorene (AAF) was used as a positive control.
<b>Results</b>	Cytotoxicity occurred at 128 ug/ml and no nuclear grain counts were made. The percent of cells in repair (average of 3 slides) for medium, F68, AAF, 8, 16, 32, and 64 ug/ml of cumene was, respectively, 15.3, 15.3, 94.0, 12.0, 28.7, 40.0, and 16.0. The average net nuclear grain counts per slide for medium, F68, AAF, 8, 16, 32, and 64 ug cumene/ml was, respectively for slide 1: -0.87, -2.18, 71.98, -3.66, 6.48, -1.57, and -2.90; for slide 2: -2.05, -1.50, 37.69, -0.25, 0.16, 8.44, and 2.21; and for slide 3: -2.11, -0.91, 56.54, -2.67, -1.29, 2.18, and -1.98.
<b>Cytotoxic concentration</b>	128 ug/ml
<b>Genotoxic Effects</b>	Unscheduled DNA synthesis was reported at 16 ug/ml.
<b>Remarks for Results</b>	The results between triplicates were highly variable and inconsistent. In an independent review by Malansky (1986), it was stated that although the laboratory performing the study noted a dose-response particularly at 16 and 32 ug/ml, the data were too inconsistent to form any conclusions.
<b>Conclusion Remarks</b>	The independent review by Malansky (1986) suggested that this assay should be repeated at concentrations between 16 and 32 ug/ml to define a possible dose-response.
<b>Data Qualities Reliabilities</b>	Reliability code 3. Not reliable.
<b>Remarks for Data Reliability</b>	Code 3. Documentation insufficient for assessment.
<b>References</b>	Brecher S. (1984b) Hepatocyte primary culture/DNA repair test of cumene. Project #84-2130. Gulf Life Sciences Center, Pittsburgh, PA. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue, cumene
<b>Test Type</b>	Cell transformation assay
<b>System of Testing</b>	Non bacterial
<b>GLP</b>	Yes
<b>Year</b>	1984
<b>Species/Strain</b>	BALB/3T3-A31-1-1 mouse fibroblasts
<b>Doses/Concentration</b>	5, 20, 60, or 90 ug/ml

<b>Remarks for Test Conditions</b>	Cultured mouse fibroblasts were treated with 5, 20, 60, or 90 ug cumene/ml. Cumene was emulsified with Pluronic F68 resulting in a final exposure to treated cultures of 0.04% Pluronic F68. Each treatment group consisted of 15 cultures for transformation and 2 cultures for colony formation. Negative controls consisted of untreated cultures and cultures treated with F68. The positive control was 1 ug 3-methylcholanthrene/ml. Focus and colony counts were done visually. Foci type was determined microscopically. The number of colonies per vessel and average number for each group were determined. Also, for each group, the colony forming efficiency was calculated. The criteria were as follows: a test was considered positive if there were "1) A two-fold increase in Type-III foci at the highest dose above that seen in negative control cultures, with or without a dose-related response or 2) a two-fold increase at two or more consecutive dose levels. Where negative control cultures have no Type-III foci, at least 2 foci would be needed for a dose level to be considered positive." Results were equivocal if the two-fold increase at a level other than the highest tested.
<b>Results</b>	Cytotoxicity was initially reported at 60 ug/ml as indicated by colony forming efficiency. At 90 ug/ml, cumene was very cytotoxic (no attached cells). Colony forming efficiencies for untreated controls; F68, positive controls, 5, 20, and 60 ug cumene/ml were, respectively, 59.5, 50, 4.5, 69, 61.5 and 22.0%. At 60 ug/ml, a 2-fold increase was reported in one of the duplicate cultures (6 type-III foci) and in the other duplicate, findings identical to that of the vehicle control were reported (2 type-II foci). No positive findings were reported at the lower concentrations.
<b>Cytotoxic concentration</b>	60 ug/ml
<b>Genotoxic Effects</b>	Apparent positive finding at 60 ug/ml.
<b>Remarks for Results</b>	The authors of the report indicate that toxicity was seen at 60 ug/ml yet report a 2-fold increase in one of the duplicates as a positive finding.
<b>Conclusion remarks</b>	Considering that toxicity was reported and that the 2-fold increase only occurred in one duplicate, the results, at best, could be considered equivocal.
<b>Data Qualities Reliabilities</b>	Reliability code 3. Not reliable.
<b>Remarks for Data Reliability</b>	Code 3. Relevant methodological deficiencies.
<b>References</b>	Brecher S. (1984a) Cell transformation test of cumene. Project #84-2131. Gulf Life Sciences Center. Pittsburgh, PA. Unpublished report.

#### 4.2.2 In vivo Genotoxicity

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene, assay>98%
<b>Method/guideline</b>	Micronucleus assay

<b>Test Type</b>	Clastogenic study
<b>GLP</b>	Yes
<b>Year</b>	1984
<b>Species/Strain</b>	CrI:CDR-1 (ICR) BR Swiss mouse
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Oral-Gavage
<b>Doses/Concentration</b>	250, 500 or 1000 mg/kg bw/day
<b>Exposure Period</b>	2 days
<b>Remarks for Test Conditions</b>	Groups of 10 male and 10 female Swiss mice were administered 250, 500, or 1000 mg/kg bw/day in paraffin oil by gavage for 2 consecutive days. Another group of 15 male and 15 female mice received one dose of 1000 mg/kg bw in paraffin oil by gavage. Control groups of mice (10/sex) were given paraffin oil only. Positive controls (4 mice/sex) were administered 75 mg/kg bw of cyclophosphamide. About half of the mice receiving 2 treatments and the negative controls were killed on day 3 and the other half were killed on day 4. Mice given cyclophosphamide were killed on day 3 and those receiving only 1 dose of cumene were killed (5/sex/day) on days 2, 3, and 4. Clinical signs, survival and body weights were recorded. Bone marrow samples were stained, examined microscopically, and polychromatic erythrocytes (1,000/mouse) were evaluated. Results were considered positive if there was a dose-related statistically significant (p less than 0.05) increase in micronucleated polychromatic erythrocytes. The results would be considered equivocal if the response was dose-related OR statistically significant. Student's t-test
<b>Appropriate statistical evaluations?</b>	
<b>Effect on mitotic index or PCE/NCE ratio by dose level and sex</b>	The PCE/NCE ratios for paraffin oil, 0.25, 0.5, or 1000 mg/kg bw/day of cumene and cyclophosphamide were, respectively, 0.8, 0.8, 0.8, 0.8, 0.8, and 0.4 for males killed on day 3; 0.9, 0.8, 0.8, 0.8, 0.8, and NA for males killed on day 4; 0.8, 0.8, 0.8, 0.8, 0.9, and 0.4 for females killed on day 3; 0.8, 0.8, 0.8, 0.8, 0.9, and NA for females killed on day 4.
<b>Genotoxic effects</b>	There was no significant increase in micronuclei.
<b>NOEL (C)/ LOEL (C)</b>	1000 mg/kg
<b>Remarks for Results</b>	One female died in the negative control group.
<b>Conclusion Remarks</b>	Cumene did not induce micronuclei in mice.
<b>Data Qualities Reliabilities</b>	Reliability code 1. Reliable without restriction.
<b>Remarks for Data Reliability</b>	Code 1. Comparable to guideline study.
<b>References</b>	Khan S.H. (1985) Micronucleus test of cumene. Project #84-2129. Gulf Life Sciences Center. Unpublished report.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; assay>95%
<b>Method/guideline</b>	Micronucleus assay
<b>Test Type</b>	Clastogenic study
<b>GLP</b>	Yes
<b>Year</b>	1994
<b>Species/Strain</b>	F344 rat
<b>Sex</b>	Male
<b>Route of Administration</b>	Intraperitoneal
<b>Doses/Concentration</b>	78.13, 156.25, 312.5, 625, 1,250, or 2500 mg/kg bw
<b>Exposure Period</b>	72 hours
<b>Remarks for Test Conditions</b>	Groups of 5 male rats were administered 78.13, 156.25, 312.5, 625, 1250, or 2500 mg/kg bw by intraperitoneal injection daily for 72 hours. Control rats received corn oil vehicle or, as a positive control, 25 mg/kg bw of cyclophosphamide. Bone marrow cells from the femur of each rat were sampled 24 hours after the last exposure. Two thousand polychromatic erythrocytes were scored for frequency of micronucleated cells in each test animal.
<b>Appropriate statistical evaluations?</b>	Not described.
<b>Effect on mitotic index or PCE/NCE ratio by dose level and sex</b>	At the highest dose, 3/5 rats died. The average number of micronucleated cells per 1,000 polychromatic erythrocytes was 0.5, 17.3, 1.2, 1.2, 1.3, 0.8, 2.6, and 1.3 for corn oil vehicle, positive control, and 8.13, 156.25, 312.5, 625, 1250, or 2500 mg/kg bw of cumene, respectively. The slight increase in % PCE's/MN was not dose related.
<b>Genotoxic effects</b>	Induction of micronuclei
<b>Remarks for Results</b>	The authors reported a weak positive polychromatic erythrocyte trend of P = 0.011. Based on the total number of micronuclei/dose (control, 5; positive control, 173; 78 mg/kg, 12; 156 mg/kg, 12; 312 mg/kg, 13; 625 mg/kg, 8; 1250 mg/kg, 26) significant increase in micronuclei occurred at or near toxic dose levels.
<b>Conclusion Remarks</b>	It was concluded that cumene provided a weak induction of micronuclei in F344 rats.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	National Toxicology Program (NTP) (1994) In vivo Cytogenetics Testing. Micronucleus Induction Results. Unpublished report.
<b>Substance Name</b>	<i>p</i> -Cymene

<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>Method/guideline</b>	Micronucleus assay
<b>Test Type</b>	Clastogenic study
<b>GLP</b>	Yes
<b>Year</b>	1994
<b>Species/Strain</b>	F344 rat
<b>Sex</b>	Male
<b>Route of Administration</b>	Intraperitoneal
<b>Doses/Concentration</b>	312, 625, 1,250, or 2,500 mg/kg bw
<b>Exposure Period</b>	72 hours
<b>Remarks for Test Conditions</b>	Groups of 5 male rats were administered daily doses of 312, 625, 1250, or 2500 mg/kg bw of cumene by intraperitoneal daily for 72 hours. Control rats received corn oil vehicle or, as a positive control, 25 mg/kg bw of cyclophosphamide. Bone marrow cells from the femur of each rat were sampled 24 hours after the last exposure. Two thousand polychromatic erythrocytes were scored for frequency of micronucleated cells in each test animal. The study is a repeat of a study performed in 1994 (NTP, 1994)
<b>Appropriate statistical evaluations?</b>	Not described.
<b>Effect on mitotic index or PCE/NCE ratio by dose level and sex</b>	At the highest dose, 2/5 rats died. The average number of micronucleated cells per 1000 polychromatic erythrocytes was 0.5, 7.8, 1.7, 1.4, 1.8, and 1.5 for corn oil vehicle, positive control, and 312, 625, 1250, or 2500 mg/kg bw of cumene, respectively. There was no dose related increased in micronuclei over the dose range.
<b>Genotoxic effects</b>	Induction of micronuclei.
<b>Remarks for Results</b>	The authors reported a positive polychromatic erythrocyte trend of $P = 0.085$ . Based on the total number of micronuclei/dose (control, 5; positive control, 78; 312 mg/kg, 17; 625 mg/kg, 13; 1250 mg/kg, 18) slight but significant increases in micronuclei occurred at all dose levels.
<b>Conclusion Remarks</b>	It was concluded that cumene provided a weak induction of micronuclei in F344 rats.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	National Toxicology Program (NTP) (1995) In vivo Cytogenetics Testing. Micronucleus Induction Results. Unpublished report.

### 4.3 Repeat dose Toxicity

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Purity greater than 99%
<b>Method/guideline</b>	Subacute inhalation neurotoxicity study
<b>GLP</b>	Ambiguous
<b>Year</b>	1996
<b>Species/strain</b>	Rat/Long Evans
<b>Sex</b>	Male
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	0, 50, or 250 ppm
<b>Exposure Period</b>	4 weeks
<b>Frequency of Treatment</b>	6 hr/day, 5 days/wk
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	8 weeks
<b>Remarks for Test Conditions</b>	<p>This study was designed to specifically examine the neurotoxic potential of inhaled <i>p</i>-cymene. Rats were housed 2 per cage and subjected to a 12-hour light cycle. Exposure to <i>p</i>-cymene vapor occurred during the dark cycle and rats were placed in stainless steel wire cages without food or water. Air exchange in the exposure chambers was 13 times/hour with a temperature of 23 ± 2 C. <i>p</i>-Cymene concentration in the exposure chamber was measured every 10 minutes with an infrared gas cell spectrophotometer. During the study, body weight was recorded weekly. After the 8-week recovery period, rats were decapitated and the cerebellum was removed, weighed, and homogenized (4 ml ice cold 0.32 M sucrose). The remainder of the brain was also weighed and homogenized. Synaptosomes were prepared using gradient centrifugation. The 2 homogenates and the synaptosomes were processed for neurotransmitter analyses (i.e., determination of noradrenaline [NA], dopamine [DA], and 5-hydroxytryptamine [5-HT]), and aliquots were taken for determination of enzyme activities (lactate dehydrogenase [LDH], acetylcholinesterase [AChE], and butyrylcholinesterase [BuChE]) and protein analysis.</p>
<b>NOAEL (NOEL)</b>	250 ppm
<b>Toxic Response/effects by Dose Level</b>	The authors reported that there was no overt toxicity in the treated rats and no effect on body weight or terminal weight of the brain, cerebellum or whole brain. There was also no effect on regional enzyme activities, regional protein synthesis or regional

neurotransmitter concentrations. The relative yield and total amount of synaptosomal protein were significantly reduced at 50 and 250 ppm in a concentration-related manner. Relative yield for control, 50 and 250 ppm = 16.4, 9.20, and 8.62 mg protein/g whole brain-cerebellum, respectively. Total amount for control 50, and 250 ppm = 29.1, 16.4, and 15.1 mg protein/g whole brain-cerebellum, respectively. The relative activity of LDH, AChE, and BuChE were significantly increased at 50 and 250 ppm. For control, 50 and 250 ppm, respectively: relative LDH activity = 2,7, 4.87, and 5.33 U/mg protein; relative AChE activity = 159, 291, and 288 mU/mg protein; relative BuChE activity = 209, 386, and 358 mU/mg protein. Total activity of LDH, AChE and BuChE were unaffected. In relation to the cytoplasmatic marker (LDH), the relative synaptosomal choline esterase activities (AChE and BuChE) were unaffected by p-cymene exposure. In relation to LDH, the relative synaptosomal concentrations of NA, DA, and 5-HT were unaffected by treatment. Relative to synaptosomal protein, relative NA and DA concentrations were significantly increased at 50 and 250 ppm; whereas 5-HT was unaffected. For control, 50, and 250 ppm, respectively: relative NA = 18.4, 34.4, and 31.3 pmol/mg synaptosomal protein; relative DA = 19.8, 38.0, and 36.8 pmol/mg synaptosomal protein; relative 5-HT = 8.98, 12.4, and 13.1 pmol/mg synaptosomal protein. Conversely, the total amount of NA and DA in the synaptosomal fraction was unaffected by treatment; whereas, the total amount of 5-HT was significantly decreased at 250 ppm. For control, 50, and 250 ppm, respectively: total amount of NA = 522, 544, and 461 pmol/whole brain-cerebellum; total amount of DA = 553, 600, and 541 pmol/whole brain-cerebellum; total amount of 5-HT = 255, 194, and 189 pmol/whole brain-cerebellum.

**Statistical Evaluation**

Yes. SAS program. Analysis of variance followed by Dunnett's two-tailed test when indicated. Significance P less than 0.05.

**Conclusion Remarks**

At up to 250 ppm, p-cymene exposure did not produce signs of overt toxicity in male rats exposed for 4 weeks with an 8-week recovery period. Although, some statistically significant changes were noted in the synaptosomal fraction of homogenized brain, no generally accepted test system has been established for predicting neurotoxicity. Reliability code 2. Reliable with restriction.

**Data Qualities Reliabilities**

**Remarks for Data Reliability**

Code 2. Acceptable, well-documented publication/study report, which meets basic scientific principles.

**References**

Lam H.R., Ladefoged, O., Ostergaard, G., Lund, S.P., and Simonsen, L. (1996) Four weeks' inhalation exposure of rats to p-cymene affects regional and synaptosomal neurochemistry. *Pharmacol Toxicol.*, 79, 225-230.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 98%
<b>Method/guideline</b>	Gavage 6-month study
<b>GLP</b>	No

<b>Year</b>	1956
<b>Species/strain</b>	Rat/Wistar
<b>Sex</b>	Female
<b>Route of Administration</b>	Oral-Gavage
<b>Doses/concentration Levels</b>	154, 462, or 769 mg/kg bw/day in olive oil
<b>Exposure Period</b>	194 days
<b>Frequency of Treatment</b>	Daily, 5 days/week
<b>Control Group</b>	Gavaged with 2.5 ml olive oil (vehicle)
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 10 female Wistar rats were gavaged with 154, 462, or 769 mg cumene/kg bw/day in olive oil emulsified with gum arabic, 5 days/week for a period of 6 months. Twenty control rats were gavaged with the vehicle. Body weight, food consumption, growth, and mortality were monitored and recorded regularly. Animals alive at the end of the study were killed 18-22 hours following the last exposure. Selected animals were used for sampling of oxalated blood for BUN determination, and for bone marrow counts. Hematological examinations (i.e., total erythrocytes and leucocytes, hemoglobin content, and differential white blood cell count) were conducted on selected animals typically after 20, 40, 80, and 130 doses. At necropsy, animals underwent gross examination and the lungs, heart, liver, kidneys, and spleen were weighed and prepared for microscopic evaluation. Similarly, sections of the adrenals, pancreas, and femoral bone marrow were examined.
<b>NOAEL (NOEL)</b>	154 mg/kg bw/day
<b>LOAEL (LOEL)</b>	462 mg/kg bw/day
<b>Actual dose received by dose level and sex</b>	0, 154, 462, or 769 mg/kg bw/day
<b>Toxic Response/effects by Dose Level</b>	No effects were reported at the lowest dose. The only effects reported in the higher 2 doses was an increase in average kidney weight (not specified if absolute or relative weight): reported as "slight effect" at 462 mg/kg bw/day and "moderate effect" at 769 mg/kg bw/day. The reported increase was not accompanied by histopathological renal changes.
<b>Statistical Evaluation</b>	Yes. Fisher t-test.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Wolf M.A., Rowe, V.K., McCollister, D.D., Hollingsworth, R.L., and Oyen, F. (1956) Toxicological studies of certain alkylated benzenes and benzene. <i>AMA Arch Ind Health</i> , 14, 387-398.

<b>Substance Name</b>	<i>p</i> -Cymene
-----------------------	------------------

<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1970
<b>Species/strain</b>	Squirrel monkey ( <i>Saimiri sciurea</i> )
<b>Sex</b>	Male
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Exposure Period</b>	30 exposures (i.e., 6 weeks)
<b>Frequency of Treatment</b>	8 hours/day, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Three male squirrel monkeys were exposed to atmospheres containing 244 ppm cumene, 8 hours/day, 5 days/week for a total of 30 exposures. The control group consisted of 12 monkeys. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, brain, spinal cord, and kidney sections taken for histological examination.
<b>NOAEL (NOEL)</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Toxic Response/effects by Dose Level</b>	Body weight gain did not appear to be affected by cumene exposure and no histopathological effects were reported.
<b>Statistical Evaluation</b>	No
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1970
<b>Species/strain</b>	Rat/Sprague-Dawley or Long Evans

<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	3.7 or 30 ppm (18 or 146 mg/m <sup>3</sup> )
<b>Exposure Period</b>	90 days
<b>Frequency of Treatment</b>	Continuous
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 14-15 rats (males and females, ratio not stated) were exposed to atmospheres containing 0, 3.7, or 30 ppm cumene continuously for a period of 90 days. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, and kidney sections taken for histological examination. Blood samples also were taken for hematological evaluation (i.e., leukocyte count, hemoglobin, and hematocrit).
<b>LOAEL (LOEL)</b>	3.7 ppm (18 mg/m <sup>3</sup> )
<b>Toxic Response/effects by Dose Level</b>	One rat died on day 11 in the 3.7 ppm group (no further details were given). Body weight gain did not appear to be affected by cumene exposure and no histopathological effects were reported. Although statistical analysis was not conducted, an increase in the number of leukocytes was reported following cumene exposure. Aside from a slight decrease in hematocrit, no other effects on hematological parameters were apparent.
<b>Statistical Evaluation</b>	No
<b>Remarks for Results</b>	The increased number of leukocytes appears to be consistent with the results of Cushman et al. (1995) and therefore was considered by the reviewer (and EPA, 1997) to be the basis of the LOAEL.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823..

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.9%
<b>Method/guideline</b>	Inhalation toxicity
<b>GLP</b>	Yes
<b>Year</b>	Undated

<b>Species/strain</b>	Mouse/B5C3F1
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	250, 500, 1,000, 2,000, or 4,000 ppm
<b>Exposure Period</b>	13 days over a 17-day period
<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 5 male and 5 female B5C3F1 mice were exposed to target concentrations of 0, 250, 500, 1,000, 2,000, or 4,000 ppm cumene by whole-body inhalation for 13 days over a period of 17 days. Cumene vapor was distributed into exposure chambers using a single vapor generator delivery subsystem and vapor distribution manifold. A metering valve was used to control vapor delivery and cumene vapor was diluted or mixed with conditioned chamber air prior to entry into the exposure chamber. The exposure chambers were monitored every 20 minutes. Animals were observed for survival, clinical signs, and body weight changes. At study termination, any organ weight changes or histopathological effects were noted.
<b>NOAEL (NOEL)</b>	500 ppm (females); 1,000 ppm (males)
<b>LOAEL (LOEL)</b>	1,000 ppm (females); 2,000 ppm (males)
<b>Toxic Response/effects by Dose Level</b>	All mice exposed to 2,000 or 4,000 ppm died by day 2. At 1,000 ppm, 4/5 females were dead by day 4. All remaining animals survived to study termination. Male mice exposed to 1,000 ppm showed varying degrees of ataxia, which was most severe during week 1. Body weight of surviving animals was similar to controls. Relative liver weight was significantly increased in male and female mice exposed to 250 ppm and higher. Absolute liver weight was significantly increased in males exposed to 250 ppm and higher and in females exposed to 500 ppm and higher. In females, absolute and relative kidney weight was significantly increased at 1,000 ppm; whereas in males, absolute kidney weight was significantly increased only at 250 ppm and relative kidney weight was significantly increased at 250 and 500 ppm. Absolute and relative thymus weight was significantly decreased at 1,000 ppm in males (no data for females). No histopathological findings accompanied the organ weight changes.
<b>Statistical Evaluation</b>	Not described.
<b>Remarks for Results</b>	During the study period, cumene was stable and uniform in the exposure chambers and the test concentrations remained within the protocol specified range for daily means with acceptable relative standard deviations.
<b>Conclusion Remarks</b>	Based on these results, NTP set exposure concentrations of 62.5 to 1,000 ppm cumene for the 13-week inhalation study. The NOAELs were based on mortality and lack of histopathology.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability** Code 2. Basic data given: comparable to guidelines/standards.

**References** NTP unpublished results (c). 2-Week Inhalation Toxicity Study of Cumene--Mice. National Toxicology Program.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.9%
<b>Method/guideline</b>	Subchronic inhalation study
<b>GLP</b>	Yes
<b>Year</b>	Undated
<b>Species/strain</b>	Mouse/B6C3F1
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	62.5, 125, 250, 500, or 1,000 ppm
<b>Exposure Period</b>	13 weeks
<b>Frequency of Treatment</b>	6 hours/day plus T90, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 10 male and 10 female B6C3F1 mice were exposed to target concentrations of 0, 62.5, 125, 250, 500 or 1,000 ppm cumene by whole-body inhalation 6 hours/day plus T90, 5 days/week for up to 13 weeks. Cumene vapor was distributed into exposure chambers using a single vapor generator delivery subsystem and vapor distribution manifold. A metering valve was used to control vapor delivery and cumene vapor was diluted or mixed with conditioned chamber air prior to entry into the exposure chamber. The exposure chambers were monitored every 20 minutes. Animals were observed for survival, clinical signs, and body weight changes. At study termination, any organ weight changes or histopathological effects were noted. Hematology also was evaluated but not described in detail.
<b>NOAEL (NOEL)</b>	250 ppm
<b>LOAEL (LOEL)</b>	500 ppm
<b>Toxic Response/effects by Dose Level</b>	All male mice survived to the end of the study. Eight out of 10 female mice exposed to 1,000 ppm cumene died within the first week of exposure. Transient signs of ataxia were reported in high-dose males and surviving females during the first week of exposure. Male mice at the 2 highest exposures showed statistically significant decreased final body weights; whereas female final body weights were not affected. Absolute liver weight was significantly increased at 1,000 ppm in

both sexes. Relative liver weight was significantly increased in all exposed males and in females exposed to 250 ppm cumene and higher. No effect on hematology was reported. Histopathologically, centrilobular hypertrophy of the liver was reported in all males exposed to 1,000 ppm cumene. No other males (treated or controls) had similar findings. In females, squamous hyperplasia and inflammation of the mucosa of the forestomach were reported at 500 and 1,000 ppm (2/10 and 1/10 rats, respectively) compared to no forestomach lesions in controls.

**Statistical Evaluation**

Not described.

**Remarks for Results**

During the study period, cumene was stable and uniform in the exposure chambers and the test concentrations remained within the protocol specified range for daily means with acceptable relative standard deviations.

**Conclusion Remarks**

A NOAEL of 250 ppm was determined based on mortality, body weight changes, and histopathological findings.

**Data Qualities Reliabilities**

Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability**

Code 2. Basic data given: comparable to guidelines/standards.

**References**

NTP unpublished results (a). 13-Week Subchronic Inhalation Toxicity Study of Cumene--Mice. National Toxicology Program.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.9%
<b>Method/guideline</b>	Subchronic inhalation study
<b>GLP</b>	Yes
<b>Year</b>	Undated
<b>Species/strain</b>	Rat/F344/N
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	62.5, 125, 250, 500, or 1,000 ppm
<b>Exposure Period</b>	13 weeks
<b>Frequency of Treatment</b>	6 hours/day plus T90, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 20 male and 20 female F344/N rats were exposed to target concentrations of 0, 62.5, 125, 250, 500 or 1,000 ppm cumene by whole-body inhalation 6 hours/day plus T90, 5 days/week for up to 13 weeks. Cumene vapor was distributed into exposure chambers using a single vapor generator delivery subsystem and vapor distribution

	<p>manifold. A metering valve was used to control vapor delivery and cumene vapor was diluted or mixed with conditioned chamber air prior to entry into the exposure chamber. The exposure chambers were monitored every 20 minutes. Animals were observed for survival, clinical signs, and body weight changes. At study termination, any organ weight changes or histopathological effects were noted. Hematology and clinical chemistry also were evaluated but not described in detail.</p>
<b>NOAEL (NOEL)</b>	125 ppm
<b>LOAEL (LOEL)</b>	250 ppm
<b>Toxic Response/effects by Dose Level</b>	<p>All animals survived to study termination without any significant effect on final body weights. Mild ataxia was observed in high-dose animals during the initial days of exposure. In males exposed to 250 ppm and higher, absolute and relative liver weight and absolute kidney weight were significantly increased. Relative kidney weight was significantly increased in all exposed males. In females, relative liver weight was increased at the 3 highest concentrations; whereas relative kidney weight was increased at the 2 highest exposure concentrations. The effect on hematology parameters was reported as "not remarkable" and the most notable serum chemistry result was increased total bile acid concentration on days 3 (concentrations of 125 ppm and higher) and 23 (concentrations of 250 ppm and higher) in both sexes. At terminal sacrifice, a significant decrease in alanine aminotransferase was reported in males and females exposed to 250 ppm cumene and higher. Accompanying the kidney weight increase in males was an increase in hyaline droplets and tubular regeneration in renal cortical tubules and granular casts in tubules in the corticomedullary junction area. The severity and incidence of granular casts was reported to show an exposure-related response. These findings were not reported in females. The amount of <i>alpha</i>-2u-globulin in the kidney of male rats increased in an exposure-related manner, reaching statistical significance at concentrations of 125 ppm and higher. Proliferating cell nuclear antigen was measured and showed no difference from controls indicating that there was no difference in renal cortical cell turnover rates. No other histopathological findings were reported.</p>
<b>Statistical Evaluation</b>	Not described.
<b>Remarks for Results</b>	<p>During the study period, cumene was stable and uniform in the exposure chambers and the test concentrations remained within the protocol specified range for daily means with acceptable relative standard deviations. The renal lesions reported in the male rats were considered by the conducting laboratory to be similar to those "resulting from exposure to chemicals that induce accumulation of <i>alpha</i>-2u-globulin in renal cortical tubular cytoplasm".</p>
<b>Conclusion Remarks</b>	A NOAEL of 125 ppm was determined for both sexes based on serum chemistry, organ weight changes, and renal changes reported in males.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	NTP unpublished results (b). 13-Week Subchronic Inhalation Toxicity Study of Cumene--Rats. National Toxicology Program.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.9%
<b>Method/guideline</b>	Subchronic inhalation study
<b>GLP</b>	Yes
<b>Year</b>	Undated
<b>Species/strain</b>	Mouse/B6C3F1
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	62.5, 125, 250, 500, or 1,000 ppm
<b>Exposure Period</b>	13 weeks
<b>Frequency of Treatment</b>	6 hours/day plus T90, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 10 male and 10 female B6C3F1 mice were exposed to target concentrations of 0, 62.5, 125, 250, 500 or 1,000 ppm cumene by whole-body inhalation 6 hours/day plus T90, 5 days/week for up to 13 weeks. Cumene vapor was distributed into exposure chambers using a single vapor generator delivery subsystem and vapor distribution manifold. A metering valve was used to control vapor delivery and cumene vapor was diluted or mixed with conditioned chamber air prior to entry into the exposure chamber. The exposure chambers were monitored every 20 minutes. Animals were observed for survival, clinical signs, and body weight changes. At study termination, any organ weight changes or histopathological effects were noted. Hematology also was evaluated but not described in detail.
<b>NOAEL (NOEL)</b>	250 ppm
<b>LOAEL (LOEL)</b>	500 ppm
<b>Toxic Response/effects by Dose Level</b>	All male mice survived to the end of the study. Eight out of 10 female mice exposed to 1,000 ppm cumene died within the first week of exposure. Transient signs of ataxia were reported in high-dose males and surviving females during the first week of exposure. Male mice at the 2 highest exposures showed statistically significant decreased final body weights; whereas female final body weights were not affected. Absolute liver weight was significantly increased at 1,000 ppm in both sexes. Relative liver weight was significantly increased in all exposed males and in females exposed to 250 ppm cumene and higher. No effect on hematology was reported. Histopathologically, centrilobular hypertrophy of the liver was reported in all males

	exposed to 1,000 ppm cumene. No other males (treated or controls) had similar findings. In females, squamous hyperplasia and inflammation of the mucosa of the forestomach were reported at 500 and 1,000 ppm (2/10 and 1/10 rats, respectively) compared to no forestomach lesions in controls.
<b>Statistical Evaluation</b>	Not described.
<b>Remarks for Results</b>	During the study period, cumene was stable and uniform in the exposure chambers and the test concentrations remained within the protocol specified range for daily means with acceptable relative standard deviations.
<b>Conclusion Remarks</b>	A NOAEL of 250 ppm was determined based on mortality, body weight changes, and histopathological findings.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	NTP unpublished results(a). 13-Week Subchronic Inhalation Toxicity Study of Cumene--Mice. National Toxicology Program.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.9%
<b>Method/guideline</b>	Inhalation toxicity
<b>GLP</b>	Ambiguous
<b>Year</b>	1995
<b>Species/strain</b>	Rat/Fischer 344/NHSD
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	50, 100, 500, or 1,200 ppm
<b>Exposure Period</b>	13 weeks
<b>Frequency of Treatment</b>	6 hours/day, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	4 weeks
<b>Remarks for Test Conditions</b>	Groups of 15 male and 15 female rats were exposed to atmospheres containing 0, 50, 100, 500, or 1,200 ppm cumene 6 hours/day, 5 days/week for 13 weeks plus 2 or 3 days followed by a 4-week recovery period. Rats were individually exposed to test atmospheres in wire-mesh exposure cages in 900-L rectangular glass and stainless steel chambers with an airflow rate of 200 liter/minute with 13 air changes/hour. Chamber temperature, relative humidity, and cumene concentration (measured by GC) were measured every half hour during the 6-hour exposure. When not in the exposure chambers, the

rats were individually housed and maintained on a 12-hour photoperiod and had ad libitum access to basal rodent diet and water. During the study, cages were rotated within the exposure chamber and non-exposure housing to ensure uniform exposures to the test material and lighting. Rats were observed daily on exposure days for clinical signs and on non-exposure days for mortality. Body weight was measured weekly. Fifteen rats/sex were tested for motor activity prior to exposure and on the weekends following study weeks 4, 9, and 13 using an automated recording apparatus. Test sessions lasted 90 min with intrasession intervals of 10 min. Ten rats/sex were assessed for tone-pip auditory brain stem responses during post exposure week 1. Eyes were examined by 2 independent veterinary ophthalmologists pre-exposure, at weeks 4, 9, and 13 and during post exposure week 4. All rats were necropsied and liver, kidney, lungs, adrenal, gonad and brain weights were measured. In this study, only the eyes were histopathologically examined.

**NOAEL (NOEL)**

500 ppm

**LOAEL (LOEL)**

1,200 ppm

**Actual dose received by dose level and sex  
Toxic Response/effects by Dose Level**

Test concentrations within 1% of target

Motor activity was not affected in cumene-exposed rats. Mean body weights were similar between test and control animals, although there was a transient decrease in body weight gain in both sexes exposed to 1,200 ppm cumene during week 1. There were no differences between test and control animals for tone-pip auditory brain stem responses. No treatment-related cataracts were reported in this study. Absolute and relative liver weights were statistically increased in males exposed to 500 ppm cumene and females exposed to 1,200 ppm cumene. Only absolute liver weight was statistically increased in males exposed to 1,200 ppm cumene. Relative kidney weights and absolute and relative adrenal gland weights were statistically increased only in females exposed to 1,200 ppm cumene. There were no other histopathological findings.

**Statistical Evaluation**

Yes. Levene's test for equal variances, ANOVA, t tests, repeated-measures analysis (Dixon, 1985), Fisher's exact test, MANOVA with use of GLM procedure of SAS, F test based on Hotelling-Lawley trace statistics, F-max test.

**Remarks for Results**

The EPA (1997) evaluated the results of this study and established a NOAEL of 496 ppm and a LOAEL of 1,202 ppm based on relative and absolute weight alterations that were both biologically and statistically significant. The changes in liver weight were considered by EPA (1997) not to be toxicologically relevant since they were not accompanied by histopathology (demonstrated in the first study from this publication).

**Conclusion Remarks**

Cumene was not ototoxic in this study.

**Data Qualities Reliabilities**

Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability**

Code 2. Acceptable, well-documented publication/study report, which meets basic scientific principles.

**References**

Cushman J.R., Norris, J.C., Dodd, D.E., Darmer, K.I., Morris, C.R. (1995) Subchronic inhalation toxicity and neurotoxicity assessment of cumene in Fischer 344 rats. *J Am Coll Toxicol.*, 14(2), 129-147.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.9%
<b>Method/guideline</b>	Inhalation toxicity
<b>GLP</b>	Ambiguous
<b>Year</b>	1995
<b>Species/strain</b>	Rat/Fischer 344/NHSD
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	100, 500, or 1,200 ppm
<b>Exposure Period</b>	13 weeks
<b>Frequency of Treatment</b>	6 hours/day, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	<p>Groups of 21 male and 21 female rats were exposed to atmospheres containing 0, 100, 500, or 1,200 ppm cumene 6 hours/day, 5 days/week for 13 weeks plus 2 or 3 days. Rats were individually exposed to test atmospheres in wire-mesh exposure cages in 4,300-L rectangular glass and stainless steel chambers with an airflow rate of approximately 900 liter/minute with 12.5 air changes/hour. Chamber temperature, relative humidity, and cumene concentration (measured by GC) were measured every half hour during the 6-hour exposure. When not in the exposure chambers, the rats were individually housed and maintained on a 12-hour photoperiod and had ad libitum access to basal rodent diet and water. During the study, cages were rotated within the exposure chamber and non-exposure housing to ensure uniform exposures to the test material and lighting. Rats were observed daily on exposure days for clinical signs and on non-exposure days for mortality. Body weight and food and water consumption were determined weekly. Ten rats of both sexes underwent a functional observational battery and 15 rats/sex were tested for motor activity prior to exposure and on the weekends following study weeks 1, 2, (behavioral only), 4, 9, and 13 using an automated recording apparatus. Test sessions lasted 90 min with intrasession intervals of 10 minutes. Eyes were examined during week 13. Five rats/sex/group were selected for hematology and serum chemistry prior to exposure and 10 rats/sex/group were sampled during week 13. Parameters examined included erythrocyte, platelet, leukocyte, differential leukocyte, and reticulocyte (males only) counts, hemoglobin, hematocrit, mean corpuscular volume, hemoglobin and hemoglobin concentration, glucose, urea nitrogen, creatinine, total protein, albumin, globulin, bilirubin, calcium,</p>

phosphorus, sodium, potassium, chloride, aspartate and alanine aminotransferases, and gamma-glutamyltransferase. At study termination, 6 rats/sex/group were selected for microscopic evaluation of the brain, spinal cord, and peripheral nerves. All remaining rats were necropsied and liver, kidney, lungs, adrenal, gonad and brain weights were measured. In addition to microscopic examination of standard tissues from high-dose rats, lung tissues from both the 100 and 500 ppm groups were examined and kidney sections from all male rats were evaluated for tubular hyaline droplet formation. In addition, to evaluate sperm count and sperm morphology, the epididymides of 15 male rats/group were removed. Also, the right testis of each male was frozen and homogenized to count spermatid by the method of Johnson et al. (1980) and Blazak et al. (1985). In the high-dose and control groups, the right testis of each male rat was evaluated for the stages of spermatogenesis according to the method of Land and Chapin (1985).

**NOAEL (NOEL)**

500 ppm

**LOAEL (LOEL)**

1200 ppm

**Actual dose received by dose level and sex  
Toxic Response/effects by Dose Level**

Test concentrations within 1% of target

At 1,200 ppm, 1 male rat was killed moribund due to a caging accident, rats showed ataxia following the first 2-3 weeks, rats "appeared hypoactive, exhibited blepharospasm, and showed a delayed or absent startle reflex". In both the 500 and 1,200 ppm groups, rats "showed increased incidences of periocular tissue swelling, urine stains, urogenital area wetness, and/or perinasal encrustation. Rats exposed to 500 ppm also were hypoactive. Exposed rats showed no differences in the functional observational battery. At week 13, males rats exposed to 500 or 1,200 ppm cumene showed a decrease in total motor activity (i.e., fine movement, rearing, and ambulation combined). More specifically, there was a statistically significant decrease in ambulatory activity at weeks 4, 9, and 13. Mean body weights were similar for all groups; however, there was a transient decrease in body weight gain of high-dose females during week 1, 2, 6, and 7. In addition, mean food consumption was decreased at week 1 in females exposed to 500 and 1,200 ppm cumene. Water consumption was consistently increased (by 40% over controls) in male and female rats throughout most of the study. Cataracts were observed in about 14-55% of all groups (including controls). With respect to hematology and blood chemistry, leukocytes and platelets were significantly increased at the 2 highest concentrations in both sexes and lymphocytes were significantly increased at the 2 highest concentrations in males only. Glucose was significantly decreased in females of the 500 and 1,200 ppm groups, but not in males. Total protein, albumin and globulin were significantly increased in both sexes exposed to 1,200 ppm. Calcium and inorganic phosphorus were significantly increased at 1,200 ppm in both sexes and at 500 ppm in males. At 1,200 ppm, both males and females had significantly increased absolute and relative liver, kidney, and adrenal gland weights. Absolute and relative liver weight also was significantly increased in both sexes exposed to 500 ppm cumene. Significantly increased relative kidney weight and absolute adrenal gland weight was reported in females and males, respectively, at 500 ppm. There were no effects reported in the examination of nervous

system tissue in any of the groups. The only microscopic findings reported were in the kidneys of males rats exposed to 500 or 1,200 ppm cumene. Tubular proteinosis was significantly increased in high-dose males, and interstitial nephritis and tubular cell hyperplasia/hypertrophy were significantly increased at both 500 and 1,200 ppm cumene. In addition, at the 2 highest concentrations, an increase in hyaline droplet formation within the proximal tubules of male rats was reported. Testicular sperm head and epididymal spermatozoa counts were similar for all groups and there was no effect on epididymal sperm morphology.

**Statistical Evaluation**

Yes. Levene's test for equal variances, ANOVA, t tests, repeated-measures analysis (Dixon, 1985), Fisher's exact test.

**Remarks for Results**

The decreased motor activity was not replicated in a second study under the same conditions, but including a 4-week recovery period. The cataracts observed in this study were considered uninterrupted, but in a second similar study were determined by ophthalmologists to be unrelated to cumene exposure. The EPA (1997) evaluated the results of this study and established a NOAEL of 496 ppm and a LOAEL of 1,202 ppm based on relative and absolute weight alterations that were both biologically and statistically significant. The changes in liver weight were considered by EPA (1997) not to be toxicologically relevant since they were not accompanied by histopathology. The blood effects reported were also considered irrelevant since they were within normal ranges.

**Conclusion Remarks**

Exposure to cumene at concentrations below 1,200 ppm produced no adverse effects in rats over a 13-week period.

**Data Qualities Reliabilities**

Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability**

Code 2. Acceptable, well-documented publication/study report, which meets basic scientific principles.

**References**

Cushman J.R., Norris, J.C., Dodd, D.E., Darmer, K.I., Morris, C.R. (1995) Subchronic inhalation toxicity and neurotoxicity assessment of cumene in Fischer 344 rats. J Am Coll Toxicol., 14(2), 129-147.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1970
<b>Species/strain</b>	Guinea pig/Princeton-derived
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	3.7 or 30 ppm (18 or 146 mg/m <sup>3</sup> )
<b>Exposure Period</b>	90 days
<b>Frequency of Treatment</b>	Continuous

<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 15 guinea pigs (males and females, ratio not stated) were exposed to atmospheres containing 0, 3.7, or 30 ppm cumene continuously for a period of 90 days. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, and kidney sections taken for histological examination. Blood samples also were taken for hematological evaluation (i.e., leukocyte count, hemoglobin, and hematocrit).
<b>NOAEL (NOEL)</b>	3.7 ppm (18 mg/m <sup>3</sup> )
<b>LOAEL (LOEL)</b>	30 ppm (146 mg/m <sup>3</sup> )
<b>Toxic Response/effects by Dose Level</b>	Body weight gain appeared to be reduced in rats exposed to 30 ppm cumene, but increased in rats exposed to 3.7 ppm cumene. No histopathological effects were reported. No effects on hematological parameters were apparent.
<b>Statistical Evaluation</b>	No.
<b>Remarks for Results</b>	Without statistical analysis, it is difficult to interpret the significance of the findings particularly when these due not seem to be concentration related.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1970
<b>Species/strain</b>	Dog/Beagle
<b>Sex</b>	Male
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	3.7 or 30 ppm (18 or 146 mg/m <sup>3</sup> )
<b>Exposure Period</b>	90 days
<b>Frequency of Treatment</b>	Continuous
<b>Control Group</b>	0 ppm

<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 2 male beagle dogs were exposed to atmospheres containing 3.7, or 30 ppm cumene continuously for a period of 90 days. The control group consisted of 10 male beagle dogs. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, brain, spinal cord, and kidney sections taken for histological examination. Blood samples also were taken for hematological evaluation (i.e., leukocyte count, hemoglobin, and hematocrit).
<b>NOAEL (NOEL)</b>	30 ppm (146 mg/m <sup>3</sup> )
<b>Toxic Response/effects by Dose Level</b>	Body weight gain did not appear to be affected by cumene exposure and no histopathological effects were reported. No effects on hematological parameters were apparent.
<b>Statistical Evaluation</b>	No
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1970
<b>Species/strain</b>	Squirrel monkey ( <i>Saimiri sciurea</i> )
<b>Sex</b>	Male
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	3.7 or 30 ppm (18 or 146 mg/m <sup>3</sup> )
<b>Exposure Period</b>	90 days
<b>Frequency of Treatment</b>	Continuous
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 3 male squirrel monkeys were exposed to atmospheres containing 3.7, or 30 ppm cumene continuously for a period of 90 days. The control group consisted of 12 monkeys. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, brain, spinal cord, and kidney sections taken for histological

examination.

<b>Toxic Response/effects by Dose Level</b>	Terminal body weights were lower in treated animals than in controls when compared with starting body weights (starting body weight versus terminal body weight: 0 ppm, 690 g versus 679 g; 3.7 ppm, 759 g versus 687 g; 30 ppm, 755 g versus 644 g), but no histopathological effects were reported.
<b>Statistical Evaluation</b>	No
<b>Remarks for Results</b>	Without statistical analysis, it is difficult to interpret the significance of the findings particularly when similar effects were occurring in the control animals.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.9%
<b>Method/guideline</b>	Inhalation toxicity
<b>GLP</b>	Yes
<b>Year</b>	Undated
<b>Species/strain</b>	Rat/F344/N
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	250, 500, 1,000, 2,000, or 4,000 ppm
<b>Exposure Period</b>	12 days over a 16-day period
<b>Frequency of Treatment</b>	Daily
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 5 male and 5 female F344/N rats were exposed to target concentrations of 0, 250, 500, 1,000, 2,000, or 4,000 ppm cumene by whole-body inhalation for 12 days over a period of 16 days. Cumene vapor was distributed into exposure chambers using a single vapor generator delivery subsystem and vapor distribution manifold. A metering valve was used to control vapor delivery and cumene vapor was diluted or mixed with conditioned chamber air prior to entry into the exposure chamber. The exposure chambers were monitored every

	20 minutes. Animals were observed for survival, clinical signs, and body weight changes. At study termination, any organ weight changes or histopathological effects were noted.
<b>NOAEL (NOEL)</b>	1,000 ppm (females)
<b>LOAEL (LOEL)</b>	500 ppm (females); 250 ppm (males)
<b>Toxic Response/effects by Dose Level</b>	All rats exposed to 4,000 ppm cumene died by day 1. At 2,000 ppm, 3/5 females and 2/5 males died by day 4. There was a significant decrease in mean body weight in rats exposed to 2,000 ppm cumene. Surviving rats exposed to 1,000 or 2,000 ppm cumene showed varying degrees of ataxia or lethargy, which was more severe at the beginning of the exposure week. At 500 ppm, mild ataxia was noted only after the first exposure. Relative liver weight was significantly increased in both sexes exposed to all test concentrations. Absolute liver weight was significantly increased in males exposed to 1,000 or 2,000 ppm and in females exposed to 500 ppm cumene or higher. Relative kidney weight was increased in both sexes at all test concentrations. Absolute kidney weight was significantly increased only at 250 and 1,000 ppm in males and at 250, 500 and 1,000 ppm in females. Absolute and relative thymus weight was significantly decreased at 2,000 ppm in both sexes. In exposed males, hyaline droplets in the renal cortical tubules were reported (incidences at 0, 250, 500, 1,000 and 2,000 ppm: 0/5, 3/5, 2/5, 3/5 and 1/5). None were observed in the 4,000 ppm group likely due to the short exposure period. At 2,000 ppm, suppurative inflammation of the lung was reported in 2/5 males and 2/5 females. One female rat exposed to 2,000 ppm cumene also had histiocytic cellular infiltrate of the lungs. Three out of 5 males exposed to 2,000 ppm cumene were reported to show liver congestion.
<b>Statistical Evaluation</b>	Not described.
<b>Remarks for Results</b>	During the study period, cumene was stable and uniform in the exposure chambers and the test concentrations remained within the protocol specified range for daily means with acceptable relative standard deviations.
<b>Conclusion Remarks</b>	Based on these results, NTP set exposure concentrations of 62.5 to 1,000 ppm cumene for the 13-week inhalation study. No NOAEL was determined for males because of mortality and histopathology; however a NOAEL of 1,000 ppm was determined for females based on mortality and histopathological findings.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	NTP unpublished results (d). 2-Week Inhalation Toxicity Study of Cumene--Rats. National Toxicology Program.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No

<b>Year</b>	1970
<b>Species/strain</b>	Rat/Sprague-Dawley or Long Evans
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Exposure Period</b>	30 exposures (i.e., 6 weeks)
<b>Frequency of Treatment</b>	8 hours/day, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 14-15 rats (males and females, ratio not stated) were exposed to atmospheres containing 0 or 244 ppm cumene, 8 hours/day, 5 days/week for a total of 30 exposures. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, and kidney sections taken for histological examination. Blood samples also were taken for hematological evaluation (i.e., leukocyte count, hemoglobin, and hematocrit).
<b>LOAEL (LOEL)</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Toxic Response/effects by Dose Level</b>	Body weight gain did not appear to be affected by cumene exposure and no histopathological effects were reported. Although statistical analysis was not conducted, an increase in the number of leukocytes was reported following cumene exposure. Aside from a slight decrease in hematocrit, no other effects on hematological parameters were apparent.
<b>Statistical Evaluation</b>	No
<b>Remarks for Results</b>	The increased number of leukocytes appears to be consistent with the results of Cushman et al. (1995) and therefore was considered by the reviewer (and EPA, 1997) to be the basis of the LOAEL.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1970

<b>Species/strain</b>	Guinea pig/Princeton-derived
<b>Sex</b>	Male and Female
<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Exposure Period</b>	30 exposures (i.e., 6 weeks)
<b>Frequency of Treatment</b>	8 hours/day, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Groups of 15 guinea pigs (males and females, ratio not stated) were exposed to atmospheres containing 0 or 244 ppm cumene, 8 hours/day, 5 days/week for a total of 30 exposures. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, and kidney sections taken for histological examination. Blood samples also were taken for hematological evaluation (i.e., leukocyte count, hemoglobin, and hematocrit).
<b>LOAEL (LOEL)</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Toxic Response/effects by Dose Level</b>	Body weight gain appeared to be reduced in cumene-exposed animals, but no histopathological effects were reported. No effects on hematological parameters were apparent.
<b>Statistical Evaluation</b>	No.
<b>Remarks for Results</b>	Without statistical analysis, it is difficult to interpret the significance of the findings particularly when these tendencies are not seen in the other animal species tested (rats, dogs, and monkeys).
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene
<b>GLP</b>	No
<b>Year</b>	1970
<b>Species/strain</b>	Dog/Beagle
<b>Sex</b>	Male

<b>Route of Administration</b>	Inhalation
<b>Doses/concentration Levels</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Exposure Period</b>	30 exposures (i.e., 6 weeks)
<b>Frequency of Treatment</b>	8 hours/day, 5 days/week
<b>Control Group</b>	0 ppm
<b>Post Exposure</b>	None
<b>Remarks for Test Conditions</b>	Two male beagle dogs were exposed to atmospheres containing 244 ppm cumene, 8 hours/day, 5 days/week for a total of 30 exposures. The control group consisted of 10 male beagle dogs. At the end of the exposures, animals were killed and necropsied with heart, lung, liver, spleen, brain, spinal cord, and kidney sections taken for histological examination. Blood samples also were taken for hematological evaluation (i.e., leukocyte count, hemoglobin, and hematocrit).
<b>LOAEL (LOEL)</b>	244 ppm (1,195 mg/m <sup>3</sup> )
<b>Toxic Response/effects by Dose Level</b>	Body weight gain did not appear to be affected by cumene exposure and no histopathological effects were reported. There appeared to be an effect on the hematological parameters examined following cumene exposure: an increase in leukocytes, and increased hemoglobin and hematocrit.
<b>Statistical Evaluation</b>	No
<b>Remarks for Results</b>	Without statistical analysis, it is difficult to interpret the significance of the findings.
<b>Conclusion Remarks</b>	
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Basic data given: comparable to guidelines/standards.
<b>References</b>	Jenkins L.J., Jr., Jones, R.A., and Siegel, J. (1970) Long-term inhalation screening studies of benzene, toluene, o-xylene, and cumene on experimental animals. <i>Toxicol Appl Pharmacol.</i> , 16, 818-823.

## 4.4 Reproductive Toxicity

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene; purity greater than 99.94%
<b>Test Type</b>	Inhalation toxicity
<b>GLP</b>	Ambiguous
<b>Year</b>	1995
<b>Species/Strain</b>	Rat/Fischer 344/NHSD
<b>Sex</b>	Male
<b>Route of Administration</b>	Inhalation
<b>Duration of Test</b>	13 weeks
<b>Doses/Concentration</b>	100, 500, or 1,200 ppm
<b>Control Group and Treatment</b>	0 ppm
<b>Frequency of Treatment</b>	6 hours/day, 5 days/week
<b>Remarks for Test Conditions</b>	As part of a subchronic study, groups of 21 male rats were exposed to atmospheres containing 0, 100, 500, or 1200 ppm cumene 6 hours/day, 5 days/week for 13 weeks plus 2 or 3 days. Rats were individually exposed to test atmospheres in wire-mesh exposure cages in 4300-liter rectangular glass and stainless steel chambers with an airflow rate of approximately 900 liter/minute with 12.5 air changes/hour. Chamber temperature, relative humidity, and cumene concentration (measured by GC) were measured every half hour during the 6-hour exposure. When not in the exposure chambers, the rats were individually housed and maintained on a 12-hour photoperiod and had ad libitum access to basal rodent diet and water. During the study, cages were rotated within the exposure chamber and non-exposure housing to ensure uniform exposures to the test material and lighting. In addition to the parameters studied for the subchronic study, the epididymides of 15 male rats/group were removed to evaluate sperm count and sperm morphology. Also, the right testis of each male was frozen and homogenized to count spermatid by the method of Johnson et al. (1980) and Blazak et al. (1985). In the high-dose and control groups, the right testis of each male rat was evaluated for the stages of spermatogenesis according to the method of Land and Chapin (1985).
<b>NOAEL(NOEL)</b>	1200 ppm
<b>Appropriate statistical evaluations</b>	Yes. Levene's test for equal variances, ANOVA, t tests, repeated-measures analysis (Dixon, 1985), Fisher's exact test.

**Parental data and F1 as Appropriate**

Testicular sperm head and epididymal spermatozoa counts were similar for all groups. At 1,200 ppm, one rat was reported to show diffuse testicular atrophy; however, all other animals showed normal morphology and stages of spermatogenesis in the testes. In epididymal spermatozoa, there were no individual abnormalities of the sperm head; however, at 500 ppm, when total abnormalities were grouped by total number per category, there appeared to be a slight increase (statistical significance not reported) in the incidence of head abnormalities. No effects on epididymal sperm morphology were reported based on more than 96% normal epididymal sperm.

**Remarks for Results**

The slight increase in total head abnormalities noted at 500 ppm were considered by the authors to be irrelevant since no dose-response was observed and when evaluated as percentage of sperm assessed, sperm head abnormalities were infrequent. In addition, no statistical significance was reported by the authors.

**Data Reliabilities Qualities**

Reliability code 2. Reliable with restriction.

**Remarks for Data Reliability**

Code 2. Acceptable, well-documented publication/study report which meets basic scientific principles.

**References**

Cushman J.R., Norris, J.C., Dodd, D.E., Darmer, K.I., Morris, C.R. (1995) Subchronic inhalation toxicity and neurotoxicity assessment of cumene in Fischer 344 rats. J Am Coll Toxicol., 14(2),129-147.

### 4.5 Developmental/Teratogenicity Toxicity

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene, purity greater than 99.9%
<b>Test Type</b>	Developmental toxicity
<b>GLP</b>	Ambiguous
<b>Year</b>	1997
<b>Species/strain</b>	Rat/CD
<b>Sex</b>	Female
<b>Route of Administration</b>	Inhalation
<b>Duration of Test</b>	21 days
<b>Doses/concentration Levels</b>	100, 500, or 1,200 ppm
<b>Exposure Period</b>	Gestation days 6-15
<b>Frequency of Treatment</b>	6 hours/day
<b>Control Group and Treatment</b>	0 ppm

<b>Remarks for Test Conditions</b>	<p>Male and female CD rats (60 days of age) were quarantined for 2 weeks and when deemed suitable for study, were mated (1 male to 1 female). Groups of 25 plug-positive females were exposed to atmospheres containing 0, 100, 500, or 1,200 ppm cumene 6 hours/day during gestation days 6-15. Rats were individually exposed to test atmospheres in wire-mesh exposure cages in 4,320-liter rectangular glass and stainless steel chambers with an airflow rate of approximately 900 liters/minutes with 14 air changes/hour and a theoretically derived time required for the chamber to reach 99% of the equilibrium concentration (t99) of approximately 20 min. Atmospheric pressure in the chambers was maintained at a slightly negative pressure to prevent possible leaks. Chamber temperature, relative humidity, cumene concentration (measured by GC) and airflow were measured every half hour during the 6-hour exposure. When not in the exposure chambers, the rats were individually housed and maintained on a 12-hour photoperiod and had ad libitum access to basal rodent diet and water. Rats were observed daily for clinical signs. Body weight and food consumption were measured on gestation days 0, 6, 9, 12, 15, 18, and 21. On gestation day 21, maternal rats were killed and the gravid uterus, ovaries (including corpora lutea), cervix, vagina, abdominal and thoracic cavities, and respiratory tracts (including nasal turbinates) were examined. Live and dead fetuses and resorption sites were recorded. Any nongravid uteri were placed in 10% ammonium sulfide solution for detection of early resorptions. Live fetuses were examined for gender, external malformations, and variations and skeletal malformations and variations. Fifty percent of the live fetuses were examined for thoracic and abdominal visceral abnormalities, and for craniofacial structures.</p>
<b>NOAEL(NOEL) maternal toxicity</b>	488 ppm
<b>LOAEL(LOEL) maternal toxicity</b>	1211 ppm
<b>NOAEL (NOEL) developmental toxicity</b>	1211 ppm
<b>Actual dose received by dose level and sex</b>	0, 99, 488, or 1,211 ppm
<b>Maternal data with dose level</b>	<p>All rats survived to termination of study with no abortions or early deliveries. At 0, 100, and 500 ppm, 2, 2, and 3 rats were not pregnant. The pregnancy rate ranged from 88-100% and a total of 22-25 litters were examined for each group. During exposure, the high-dose rats showed significant reductions in body weight gain, but no significant differences in maternal body weight were reported when the rats were weighed. At both 500 and 1,200 ppm, food consumption was significantly reduced during the exposure period. At the highest concentration, perioral wetness, encrustation, and significantly increased relative liver weight were reported. Evaluation of the female reproduction organs including gravid uterus, ovaries, cervix, and vagina showed no abnormalities. There were no effects noted at necropsy and no significant changes in maternal corrected gestational weight, gravid uterine weight, or absolute liver weight in any of the treatment groups.</p>
<b>Fetal Data with Dose Level</b>	<p>No statistically significant effects were reported in the fetuses. Parameters examined included number of corpora lutea, number of total nonviable or viable implantations, percent pre- or post-implantation loss, sex ratio, fetal body weights, malformations, or variations. Although there was a significant increase in the incidence</p>

of skeletal and visceral variations, they were not exposure related.

<b>Appropriate statistical evaluations</b>	Yes. Levene's test for equal variances, ANOVA, t tests, Kruskal-Wallis test, Mann-Whitney U test, Fisher's exact test.
<b>Remarks for Results</b>	In reviewing this study, EPA (1997) set the maternal NOAEL at 488 ppm based on the significant decrease in body weight gain during exposure and increased relative liver weight.
<b>Conclusion Remarks</b>	Even at maternally toxic concentrations, exposure to cumene vapor did not produce developmental toxicity.
<b>Data Qualities Reliabilities</b>	Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Acceptable, well-documented publication/study report which meets basic scientific principles.
<b>References</b>	Darmer K.I., Jr., Neepers-Bradley, T.L., Cushman, J.R., Morris, C.R., and Francis, B.O. (1997) Developmental toxicity of cumene vapor in CD rats and New Zealand white rabbits. Intl J Toxicol., 16, 119-139.

<b>Substance Name</b>	<i>p</i> -Cymene
<b>CAS No.</b>	99-87-6
<b>Remarks for Substance</b>	Data for homologue cumene, purity greater than 99.9%
<b>Test Type</b>	Developmental toxicity
<b>GLP</b>	Ambiguous
<b>Year</b>	1997
<b>Species/strain</b>	Rabbit/New Zealand white
<b>Sex</b>	Female
<b>Route of Administration</b>	Inhalation
<b>Duration of Test</b>	29 days
<b>Doses/concentration Levels</b>	500, 1,200, or 2,300 ppm
<b>Exposure Period</b>	Gestation days 6-18
<b>Frequency of Treatment</b>	6 hours/day
<b>Control Group and Treatment</b>	0 ppm
<b>Remarks for Test Conditions</b>	Male and female New Zealand white rabbits (5.5 months of age) were quarantined for 2 weeks and when deemed suitable for study, were mated (1 male to 2 female). Groups of 15 mated females were exposed to atmospheres containing 0, 500, 1,200, or 2,300 ppm cumene 6 hours/day during gestation days 6-18. Rabbits were individually exposed to test atmospheres in wire-mesh exposure cages in 4,320-liter rectangular glass and stainless steel chambers with an airflow rate of approximately 900 liter/minutes with 14 air changes/hour and a theoretically derived time required for the chamber to reach 99% of the equilibrium concentration (t99) of approximately 20 min. Atmospheric pressure in the chambers was maintained at a slightly negative pressure to prevent possible leaks.

Chamber temperature, relative humidity, cumene concentration (measured by GC) and airflow were measured every half hour during the 6-hour exposure. When not in the exposure chambers, the rabbits were individually housed and maintained on a 12-hour photoperiod and had ad libitum access to basal rabbit diet and water. Rabbits were observed daily for clinical signs. Food consumption was measured daily and body weight was measured on gestation days 0, 6, 12, 18, 24, and 29. On gestation day 29, maternal rabbits were killed and the gravid uterus, ovaries (including corpora lutea), cervix, vagina, abdominal and thoracic cavities, and respiratory tracts (including nasal turbinates) were examined. Live and dead fetuses and resorption sites were recorded. Any nongravid uteri were placed in 10% ammonium sulfide solution for detection of early resorptions. Live fetuses were killed immediately upon removal were examined for gender, external malformations, and variations, skeletal malformations and variations, and thoracic and abdominal visceral abnormalities. Fifty percent of the fetuses were decapitated and examined for craniofacial structures.

<b>NOAEL(NOEL) maternal toxicity</b>	1,206 ppm
<b>LOAEL(LOEL) maternal toxicity</b>	2,297 ppm
<b>NOAEL (NOEL) developmental toxicity</b>	1,206 ppm
<b>LOAEL (LOEL) developmental toxicity</b>	2,297 ppm
<b>Actual dose received by dose level and sex</b>	0, 492, 1206 or 2297 ppm
<b>Maternal data with dose level</b>	Two does died and one aborted at the highest concentration. At the 2 lower concentrations, 1 doe in each group contained non-viable implants. All does in all groups were pregnant. During exposure, high-dose does had significantly reduced body weight gain and all treated animals had significantly reduced food consumption. The incidence of perioral wetness was significantly increased in high-dose does. At necropsy, no gross observations with the exception of lung color changes in 4/12 high-dose does were reported and there were no statistically significant differences in maternal body weight, maternal corrected gestational weight change, or absolute liver weight in any of the treatment groups. Relative liver weight was significantly increased in high-dose animals.
<b>Fetal Data with Dose Level</b>	No statistically significant effects were reported in the fetuses. Parameters examined included number of corpora lutea, number of total nonviable or viable implantations, percent pre- or post-implantation loss, sex ratio, fetal body weights, malformations, or variations. Although there was a significant increase in the incidence of skeletal and visceral variations, they were not exposure related. At 500 ppm, there was a statistically significant increase in the incidence of ecchymosis (small hemorrhage) on the head, which was not significant at the higher exposure concentrations.
<b>Appropriate statistical evaluations</b>	Yes. Levene's test for equal variances, ANOVA, t tests, Kruskal-Wallis test, Mann-Whitney U test, Fisher's exact test.
<b>Remarks for Results</b>	The increased incidence of ecchymosis on the head reported at 500 ppm was considered by the authors to be consistent with historical values. In further review of this study, EPA (1991) determined that the changes in gestational parameters, though not significant, were consistent in indicating possible developmental effects and therefore

<b>Data Qualities Reliabilities</b>	set the NOAEL for both developmental and maternal effects at 1206 ppm and the LOAEL at 2297 ppm (as reported in EPA, 1997). Reliability code 2. Reliable with restriction.
<b>Remarks for Data Reliability</b>	Code 2. Acceptable, well-documented publication/study report, which meets basic scientific principles.
<b>References</b>	Darmer K.I., Jr., Neeper-Bradley, T.L., Cushman, J.R., Morris, C.R., and Francis, B.O. (1997) Developmental toxicity of cumene vapor in CD rats and New Zealand white rabbits. Intl J Toxicol., 16, 119-139.