

201-15734B

Robust Summaries and Dossier For m-Diisopropenylbenzene (CAS No. 3478-13-8)

Existing Chemical CAS No.	:	ID: 3748-13-8 : 3748-13-8
Producer Related Part Company Creation date	:	Cytec Industries Inc. : 21.10.2002
Substance Related Part Company Creation date	:	Cytec Industries Inc. : 21.10.2002
Memo	:	
Printing date Revision date Date of last Update	:	27.11.2002 : 07.12.2004 : 07.12.2004
Number of Pages	:	51
Chapter (profile) Reliability (profile) Flags (profile)	:	Chapter: 1, 2, 3, 4, 5, 6, 7 : Reliability: without reliability, 1, 2, 3, 4 : Flags: without flag, confidential, non confidential, WGK (DE), TA-Luft (DE), Material Safety Dataset, Risk Assessment, Directive 67/548/EEC, SIDS

RECEIVED
OPPT CDIC
04 DEC 20 PM 1:58

1.0.1 OECD AND COMPANY INFORMATION

Type :
Name : Cytec Industries Inc.
Partner :
Date : 09.10.2002
Street : 5 Garret Mountain Plaza
Town : 07424 West Patterson, NJ
Country : United States
Phone :
Telefax :
Telex :
Cedex :
Reliability : (1) valid without restriction

1.0.2 LOCATION OF PRODUCTION SITE

1.0.3 IDENTITY OF RECIPIENTS

1.1 GENERAL SUBSTANCE INFORMATION

Substance type : organic
Physical status : liquid
Purity : > 98 % w/w
Reliability : (2) valid with restrictions

1.1.0 DETAILS ON TEMPLATE

1.1.1 SPECTRA

1.2 SYNONYMS

1,3-Diisopropenylbenzene
22.10.2002

Benzene, 1,3-bis(1-methylethenyl)-
22.10.2002

Benzene, m-diisopropenyl-
22.10.2002

m-Bis(1-methylvinyl)benzene
22.10.2002

m-Diisopropenylbenzene
22.10.2002

m-DIPEB
22.10.2002

1.3 IMPURITIES

1.4 ADDITIVES

1.5 QUANTITY

1.6.1 LABELLING

1.6.2 CLASSIFICATION

1.7 USE PATTERN

Type	industrial
Category	Chemical industry: used in synthesis
Reliability	(2) valid with restrictions

1.7.1 TECHNOLOGY PRODUCTION/USE

1.8 OCCUPATIONAL EXPOSURE LIMIT VALUES

No limits established

1.9 SOURCE OF EXPOSURE

1.10.1 RECOMMENDATIONS/PRECAUTIONARY MEASURES

1.10.2 EMERGENCY MEASURES

1.11 PACKAGING

1.12 POSSIB. OF RENDERING SUBST. HARMLESS

1.13 STATEMENTS CONCERNING WASTE

1.14.1 WATER POLLUTION

1.14.2 MAJOR ACCIDENT HAZARDS

1. General Information

Id 3748-13-8
Date 07.12.2004

1.14.3 AIR POLLUTION

1.15 ADDITIONAL REMARKS

1.16 LAST LITERATURE SEARCH

1.17 REVIEWS

1.18 LISTINGS E.G. CHEMICAL INVENTORIES

2.1 MELTING POINT

Value : = -38 to -40 ° C
Decomposition : no at ° C
Sublimation : no
Method : other
Year : 2002
GLP : no
Test substance : as prescribed by 1.1 - 1.4. Purity of test substance was 98.9%.
Method : ASTM E-794, Standard test method for melting and crystallization temperatures by thermal analysis.
Result : The E.O. (extrapolated onset) of enthalpy occurred at -42 degrees for the four runs. Peak enthalpy change took place at -38, -39, -39, and -40 degrees C for four runs.
Source : Cytec Industries Inc.
Test condition : Duplicate samples were encapsulated in aluminum pans and heated in the Mettler 821 DSC. Each sample was heated from -150 degrees C to 10 degrees C at 10 degrees C/min. Nitrogen, flowing at -45 ml/min, purged the system during heating and cooling. The DSC was calibrated at 10 degrees C/min with an indium standard.
Reliability : (1) valid without restriction
The test was conducted according to an established guideline.
Flag : Critical study for SIDS endpoint
03.11.2002 (23)

Value : ca. -14 ° C
Sublimation :
Method : other
Year : 2002
GLP : no
Test substance : as prescribed by 1.1 - 1.4
Remark : Inputs to the EPIWIN MPBPWIN program (v.1.40) were the CAS No. and a boiling point of 231 degrees C.
Reliability : (2) valid with restrictions
Data were obtained by modeling.
03.11.2002 (17)

2.2 BOILING POINT

Value : = 231 ° C at
Decomposition :
Method : other
Year :
GLP : no data
Test substance : as prescribed by 1.1 - 1.4
Reliability : (2) valid with restrictions
Methodological information was not provided on the MSDS. The purity of the test material was stated on the MSDS to be 100%.
(7)

2.3 DENSITY

Type : relative density
Value : = 0.925 at unknown temperature
Method : other
Year :

2. Physico-Chemical Data

Id 3748-13-8
Date 07.12.2004

GLP : no data
Test substance : as prescribed by 1.1 - 1.4
Reliability : (2) valid with restrictions
Methodological information was not provided on the MSDS. The purity of the test material was stated on the MSDS to be 100%.

(7)

2.3.1 GRANULOMETRY

2.4 VAPOUR PRESSURE

Value : ca. 0.1 hPa at 25° C
Decomposition :
Method : other (calculated)
Year : 2002
GLP : no
Test substance : as prescribed by 1.1 - 1.4
Remark : Inputs to the EPIWIN MPBPWIN Program (v1.40) were the CAS No. and boiling point of 231 degrees C.
Reliability : (2) valid with restrictions
Data were obtained by modeling.
Flag : Critical study for SIDS endpoint

(17)

Value : = 3.1 hPa at 69.3° C
Decomposition :
Method : other (measured)
Year : no data
GLP : no data
Test substance : as prescribed by 1.1 - 1.4
Remark : The following data are provided with respect to vapor pressures at various temperatures:

Degrees C	vapor pressure (torr)
69.3	2.3
86.7	5.3
101.6	11.5
111.9	17.5
123.4	26.5
134.2	41.0
144.7	58.7
151.8	78.6
161	107.3
166	129.3
172	148
231.2	743.1

Reliability : (4) not assignable
The method was not described and the purity was not given. No laboratory notebook reference or formal report or date of determination was given.

03.11.2002

(8)

2.5 PARTITION COEFFICIENT

Log pow : ca. 4.89 at ° C
Method : other (calculated)
Year : 2002
GLP : no

2. Physico-Chemical Data

Id 3748-13-8
Date 07.12.2004

Test substance : as prescribed by 1.1 - 1.4
Remark : Inputs to the EPIWIN KOWWIN Program (v1.66) were the CAS No. and a boiling point of 231 degrees C.
Reliability : (2) valid with restrictions
Data were obtained by modeling. (15)

2.6.1 WATER SOLUBILITY

Value : ca. 5.6 mg/l at ° C
Qualitative :
Pka : at 25 ° C
PH : at and ° C
Method : other
Year : 2002
GLP : no
Test substance : as prescribed by 1.1 - 1.4. Purity of the test substance was 98.9%.
Result : The highest m-DIPEB concentration dissolved in water was 5.6 ppm. Extractions of two samples showed m-DIPEB to attach to the glass container and float on the top of the water but not dissolve beyond 5.6 ppm.
Source : Cytec Industries Inc.
Test condition : A minimum of five external standards of m-DIPEB were prepared in methylene chloride with ppm concentrations of 1.3 to 115. The percent relative standard deviation of the response factors (amount/area) for all calibrations was 2.1 to 2.9. Samples were transferred to a 2 liter separatory funnel by inserting a PFA tube to the bottom of the sample container. The samples were siphoned, discarding the first 150ml, into a 2 liter separatory funnel. Samples 2-2, 3-3, 7-2 were exceptions being poured into the separatory funnel to compare the upper portion of water and residue on the glass container. The samples were extracted with 40ml, 30ml, 30ml and 10ml of methylene chloride. The extraction was collected to the mark of a 100ml volumetric flask except sample 2-2, which was extracted with 200ml. The samples were analyzed by GC flame ion detector (FID) to determine m-DIPEB content.
Reliability : (1) valid without restriction
The study was comparable to a guideline study.
Flag : Critical study for SIDS endpoint
03.11.2002 (26)

Value : ca. 4.633 mg/l at 25 ° C
Qualitative :
Pka : at 25 ° C
PH : ca. 7 at and ° C
Method : other
Year : 2002
GLP : no
Test substance : as prescribed by 1.1 - 1.4
Remark : Inputs to the EPIWIN WSKOW Program (v1.40) were the CAS No. and a boiling point of 231 degrees C.
Reliability : (2) valid with restrictions
Data were obtained by modeling.
03.11.2002 (19)

2.6.2 SURFACE TENSION

2. Physico-Chemical Data

Id 3748-13-8
Date 07.12.2004

2.7 FLASH POINT

2.8 AUTO FLAMMABILITY

2.9 FLAMMABILITY

2.10 EXPLOSIVE PROPERTIES

2.11 OXIDIZING PROPERTIES

2.12 ADDITIONAL REMARKS

3.1.1 PHOTODEGRADATION

Type : air
Light source :
Light spect. : nm
Rel. intensity : based on Intensity of Sunlight
Indirect photolysis
Sensitizer :
Conc. of sens. :
Rate constant : ca. .000000000104 cm³/(molecule*sec)
Degradation : ca. 50 % after 1.225 hour(s)
Deg. Product :
Method : other (calculated)
Year : 2002
GLP : no
Test substance : as prescribed by 1.1 - 1.4
Remark : Inputs to the EPIWIN AOP Program (v1.90) were the CAS No. and a boiling point of 231 degrees C.
Reliability : (2) valid with restrictions
Data were obtained by modeling.

(12)

3.1.2 STABILITY IN WATER

Deg. Product :
Method : other (calculated)
Year : 2002
GLP : no
Test substance : as prescribed by 1.1 - 1.4
Remark : The test substance is an aromatic hydrocarbon with no functional groups readily subject to hydrolysis under neutral ambient conditions. It has low solubility and is expected based on its molecular structure to be resistant to hydrolysis.
Result : The EPIWIN HYDROWIN Program (v1.67) cannot estimate a hydrolysis rate constant for the test substance, because it does not contain functional groups recognized by EPIWIN for estimation.
Reliability : (4) not assignable

(14)

3.1.3 STABILITY IN SOIL

3.2 MONITORING DATA

3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS

Type : fugacity model level III
Media :
Air (level III) : 0.214
Water (level III) : 24.9
Soil (level I) :
Biota (level II / III) : 11.0
Soil (level II / III) : 63.9

3. Environmental Fate and Pathways

Id 3748-13-8
Date 07.12.2004

Method : other
Year : 2002
Test substance : as prescribed by 1.1 - 1.4
Remark : EPIWIN PCKOC estimates a Koc (water soil partition) constant of 4036.

Inputs to the EPIWIN Level III fugacity model were the CAS No., a boiling point of 231 degrees C, a melting point of -39 degrees C, a water solubility of 5 mg/l and a vapor pressure of 1 mm Hg. The model inputted the following properties:

Molecular weight = 158 g/mol
Henry's Law constant = 0.00348 atm·m³/mol (estimated)
Log Kow = 4.89 (estimated)
Temperature = 25 degrees C

Reliability : (2) valid with restrictions. Data were obtained by modeling.

(16)(18)

3.3.2 DISTRIBUTION

3.4 MODE OF DEGRADATION IN ACTUAL USE

3.5 BIODEGRADATION

Type : aerobic
Inoculum : other bacteria: activated sludge
Concentration : 2mg/l related to Test substance related to
Contact time : 28 day
Degradation : % after
Result : under test conditions no biodegradation observed
Control substance : aniline
Kinetic : %
%

Deg. Product :
Method : OECD Guide-line 301 D "Ready Biodegradability: Closed Bottle Test"
Year : 1987
GLP : yes
Test substance : as prescribed by 1.1 - 1.4
Remark : The test material was not soluble at the tested concentration. To compensate for this, the solution was micropipetted onto a disc of glass fiber filter, which was then added directly to the test vessel. This theoretically increased the surface area of the sample, limited surface film and escape resulting from water partitioning, and kept the sample immersed in the test bottle.

Lower concentrations were not tested because the OECD guideline indicated that concentrations tested should be at least 2 mg/l.

The test was valid, since the positive control biodegraded under the test conditions and the oxygen demand of the water and inoculum blanks did not exceed 5 to 10% of the anticipated theoretical value of the test material after 15 to 28 days.

Result : The TOD of the material (2 mg/l carbon) was 5.3 mg/l O₂. The average dissolved oxygen content of dilution water without inoculum (blank) on Days 0, 5, 15 and 28 was 8.5, 8.5, 8.3 and 8.1 mg O₂/l, respectively. The average dissolved oxygen content of dilution water with inoculum (inoculum

3. Environmental Fate and Pathways

Id 3748-13-8

Date 07.12.2004

	<p>blank) at these times was 9.0, 8.9, 8.0 and 8.1 mg O₂/l. The average dissolved oxygen content of dilution water with a carrier plus inoculum (inoculum blank with carrier) on Days 0, 5, 15 and 28 was 9.0, 7.8, 7.4, and 7.5 mg O₂/l, indicating that the presence of the carrier increased oxygen consumption.</p> <p>The average dissolved oxygen content of test vessels (those containing the carrier, test material and inoculum) on Days 0, 5, 15 and 28 was 9.0, 8.9, 7.6, and 7.4 mg O₂/l, which was not different from that of the inoculum blank with carrier. Therefore, the test material did not biodegrade.</p> <p>The average dissolved oxygen content of the positive control (2 mg/l aniline with inoculum) on Days 0, 5, 15 and 28 was 9.0, 6.3, 3.5, and 2.2 mg O₂/l, which was equivalent to 0, 42, 73 and 95% degradation.</p>
Test condition	<p>: Dilution water was prepared by adding 1 ml each of the following solutions to 1 liter of distilled water: 1) 8.5 g/l KH₂PO₄, 21.75 g/l K₂HPO₄, 33.3 g/l Na₂HPO₄·2 H₂O, and 1.7 g/l NH₄Cl; 2) 22.5 g/l MgSO₄·7H₂O; 3) 27.5 g/l CaCl₂ and 0.25 g/l FeCl₃·6H₂O. The water was left at room temperature and gently agitated for 24 hours prior to use.</p> <p>Test material was diluted with dilution water to provide a concentration of 2 mg/l. An aliquot of the test material was micropipetted onto a disc of glass fiber filter, which was then added directly to a test vessel partially filled with dilution water. The solution was then inoculated with 5 ml activated sludge from Bergen County, New Jersey, MUA (the numbers of bacteria were not stated), and the vessel was filled with dilution water. Dilution water was added by siphon to prevent air bubbles. After oxygen content was measured, the vessel was stoppered and sealed with a secondary cap and incubated in the dark at 20 +/- 1 degrees C for up to 28 days. Vessels containing 2 mg/l aniline (reference material) in dilution water and a ctivated sludge (positive control), dilution water and inoculum with and without the carrier (inoculum blanks with and without the carrier) and dilution water with no inoculum (oxygen blank) were prepared similarly. Duplicate vessels were prepared for all test conditions (except the blank) for each time point that oxygen content was assessed (immediately, and after 5, 15 and 28 days). One vessel per time point was prepared for the blank.</p> <p>At each time point (0, 5, 15 and 28 days), oxygen content of the medium was measured using a YSI dissolved oxygen analyzer 54A. Theoretical oxygen demand (TOD, NO₃) was calculated based on the empirical formula of the test material. The percent biodegradability was calculated as oxygen depletion (BOD mg/l)/[concentration of test material (mg/l) x TOD] x 100. The oxygen depletion of the sample was corrected by subtracting the value of the inoculum blank.</p>
Test substance	<p>: The test material (CT-256-86) contained 97.5-99.1% m-DIPEB, 0.028 - 0.7% m-IPEC (m-Isopropenyl cumene, CAS No. 1129-29-9), 0 - 0.50% p-DIPEB (CAS No. 1605-18-1), 0 - 0.10% DIPB (1,3-Diisopropylbenzene, CAS No. 9962-7), 0 - 0.07% m-IPES (m-Isopropenyl styrene; CAS No. 52780-24-2) and 0.2- 1.8% unknowns.</p>
Reliability	<p>: (2) valid with restrictions</p> <p>Lower concentrations that were soluble should have been tested. It is assumed that the carrier would promote degradation by increasing surface area. However, this was not demonstrated. A positive control material that would be insoluble (and therefore would need a carrier) should have been used in the test instead of aniline (because aniline did not require use of the carrier).</p>

29.10.2002

(11)

3. Environmental Fate and Pathways

Id 3748-13-8
Date 07.12.2004

3.6 BOD5, COD OR BOD5/COD RATIO

3.7 BIOACCUMULATION

3.8 ADDITIONAL REMARKS

4.1 ACUTE/PROLONGED TOXICITY TO FISH

Type	: static
Species	: Pimephales promelas (Fish, fresh water)
Exposure period	: 96 hour(s)
Unit	: mg/l
Analytical monitoring	: no
NOEC	: m = 1.2
LC50	: m = 6.2
Method	: other
Year	: 1986
GLP	: yes
Test substance	: as prescribed by 1.1 - 1.4
Result	: None of the fish exposed to either of the controls or test material up to 2.5 mg/l died during the study. Ten and twenty percent of the fish exposed to 5 mg/l died by 72 and 96 hours, respectively. The mortality rate for fish exposed to 10 mg/l was 0% at 24 hours, 50% at 48 hours, 90% at 72 hours and 100% at 96 hours. The mortality rate for fish exposed to 20 mg/l was 0% at 24 hours, 40% at 48 hours, 70% at 72 hours and 100% at 96 hours. The 48 and 96 hour LC ₅₀ values (with confidence limits) calculated by the probit and binomial methods were 18 (12 - 69) mg/l and 6.2 (2.5 - 10) mg/l, respectively.

Six/ten fish exposed to 2.5 mg/l were quiescent at 96 hours. All fish exposed to 5 mg/l that survived exhibited abnormal behavior at 72 and 96 hours, which consisted of surfacing, quiescence, on bottom, or loss of equilibrium. Fish exposed to 10 or 20 mg/l exhibited these symptoms as early as 24 hours. Based on these data, the no effect concentration at 96 hours was 1.2 mg/l.

The temperature was 22 degrees C for all water samples. The dissolved oxygen concentration ranged from 9.1 - 9.2 at 0 hours to 5.8 - 7.3 at 96 hours. These values represented 105% to 66% saturation. There was no effect of test material on dissolved oxygen concentration. The pH ranged from 7.1 - 7.5. All temperatures, dissolved oxygen concentrations and pH values were within acceptable limits.

After stirring the solutions for 3 hours, the 5, 10 and 20 mg/l concentrations had a light film. The amount of film increased with increasing concentration. After 72 hours, the 10 mg/l solution still had a light surface film and the 20 mg/l solution had a heavier film.

Test condition	: Test organisms: The fathead minnows used in the study were obtained from an in-house culture. All fish were on a 16 hour daylight photoperiod and observed for at least 14 days prior to testing. Fish received a standard commercial fish food occasionally supplemented with brine shrimp nauplii daily until they were transferred into the test vessels. The fish had a mean weight and length of 0.19 +/- 0.053 g and 24 +/- 1.6 mm, respectively. The loading biomass was 0.12 g/l for the definitive study.
-----------------------	--

Test material: Test concentrations of 0.60, 1.2, and 2.5 mg/l were obtained by transferring the appropriate volume of a working standard prepared in acetone to the test vessels. For test concentrations of 5, 10 and 20 mg/l, 1.5 ml of acetone was added to the appropriate weight of the dry material before addition to the vessels. The solvent control was a 1.5-ml aliquot of acetone.

Test water: The well water from which the reconstituted water was prepared contained < 20 ppb aluminum, <0.2 ppb arsenic, <2 ppb cadmium, <3 ppb chromium, <4 ppb cobalt, <3 ppb copper, 12 ppb iron, <

5 ppb lead, <0.5 ppb mercury, <15 ppb nickel, <5 ppb silver, 11 ppb zinc, <0.10 ppb organophosphorus pesticides and <0.50 ppb organochlorine pesticides (including PCB's). The water was reconstituted to contain 48 mg/l NaHCO₃, 30 mg/l CaSO₄·2H₂O, 30 mg/l MgSO₄, and 2 mg/l KCl. The hardness, alkalinity and initial pH of the water were 40 - 45 mg/l (as CaCO₃), 30 - 35 mg/l (as CaCO₃) and 7.2-7.6, respectively. The dissolved oxygen concentration and pH at the start of the test were 9.1 mg/l and 7.4, respectively. The temperature of the water was kept at 22 +/- 1 degrees C.

Test conduct: Tests were conducted in 5 gallon glass vessels containing 15 liters of reconstituted water. The test fish (10 per test concentration) were acclimated to the dilution water for 48 - 96 hours prior to testing. They were not fed during this acclimation period or during the test. The test concentrations (0.6, 1.2, 2.5, 5, 10 and 20 mg/l) were chosen based on the results of a preliminary study performed with concentrations ranging from 1 to 320 mg/l. Two additional groups of 10 fish were exposed to dilution water alone or water containing the solvent. Each concentration was stirred for 3 hours before the fish were added randomly. All fish were observed at 24, 48, 72 and 96 hours for mortality and abnormal behavior. Dead organisms were removed after each observation. The pH, dissolved oxygen concentration and temperature of the control, solvent control and 0.60 ml test water were determined at the beginning of the test and after 48 and 96 hours. These variables were measured at 96 hours for water containing 5 mg/l and 0 and 48 hours for water containing 20 mg/l.

Statistical analysis: Concentration vs. lethality data were analyzed by a computer program which utilized the binomial, moving average and probit tests to determine the LC₅₀ value (and 95% confidence limit). If no mortality occurred or if a dose-response could not be determined over a reasonable range (< 37 to > 63%), an LC₅₀ value could not be calculated. The method of calculation selected for presentation was the one that gave the narrowest confidence limit.

Test substance	:	The purity of the test material (lot #S15183-124-1) was 99.13%. The composition of the remaining 0.87% is unknown.
Reliability	:	(2) valid with restrictions The results at the two highest concentrations may have been influenced by insolubility of the test material. Test concentrations were not analytically confirmed.
29.10.2002		(1)
Type	:	
Species	:	other:fish
Exposure period	:	96 hour(s)
Unit	:	mg/l
Limit test	:	
Analytical monitoring	:	no
LC50	:	0.225 mg/l
Method	:	other
Year	:	2003
GLP	:	no
Test substance	:	as prescribed by 1.1 - 1.4
Remark	:	EPIWIN ECOSAR (v0.99) was used to obtain the calculated LC50 value. Inputs to the model are CAS Number 3748-13-8, the melting point (-39 degrees C), boiling point (231 degrees C), vapor pressure (4 mm Hg) and water solubility (5 mg/l).
Result	:	The LC50 value was 0.225 mg/l
Reliability	:	(2) valid with restrictions Data were obtained by modeling.

(13)

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

Type	: static
Species	: Daphnia magna (Crustacea)
Exposure period	: 48 hour(s)
Unit	: mg/l
Analytical monitoring	: no
NOEC	: m = 1
LC50	: m = 4
Method	: other
Year	: 1986
GLP	: yes
Test substance	: as prescribed by 1.1 - 1.4
Result	: None of the daphnids exposed to test material concentrations of 0, 1.0 or 1.8 mg/l died during the study. Two daphnids in one vessel containing 3.2 mg/l died between 24 and 48 hours. All ten fish exposed to this concentration in another vessel survived. Therefore, the overall death rate of daphnids exposed to 3.2 mg/l was 10%. Nine out of 10 daphnids exposed to 5.6 mg/l (both vessels) died within 24 hours, and all died by 48 hours. All daphnids exposed to 10 mg/l died within 24 hours. The 24 and 48 hour LC ₅₀ values (with confidence limits) calculated by the binomial method were 4.5 (3.2 - 5.6) mg/l and 4.0 (3.2 - 5.6) mg/l, respectively.

In one vessel containing daphnids exposed to 1.8 mg/l, 3 and 2 daphnids were surfacing at 24 and 48 hours, respectively. Approximately half of the daphnids exposed to 3.2 mg/l were surfacing and/or clumped at 24 hours. By 48 hours, the majority of daphnids exposed to this concentration had surfaced or were on the bottom of the vessels. Each of the surviving daphnids exposed to 5.6 mg/l for 24 hours were on the surface at this time period. The no observable effect level was therefore 1.0 mg/l at 48 hours.

The initial temperature, dissolved oxygen concentration and pH of the control water were 20 degrees C, 7.0 mg/l and 8.0. Whether this was the solvent or medium control was not specified. The temperature, dissolved oxygen concentration and pH of all water assayed at 48 hours were 20 degrees C, 8.7 - 9.1 mg/l and 8.4. All temperatures, dissolved oxygen concentrations and pH values were within acceptable limits.

An oily film was present on the surface of water containing 5.6 and 10 mg/l. This condition persisted throughout the experiment.

Test condition	: Test organisms: The Daphnia magna used in the study were obtained from an in-house culture. The adults were fed algae (<i>Selenastrum capricornutum</i>) supplemented with a suspension of fish food at least every three days prior to testing. All daphnids were held at 20 +/- 2 degrees C, under a 16 hour daylight photoperiod (50-70 footcandles) with 30 minute simulated dawn and dusk periods. First instar daphnids (< 24 hours old) were used in the test. Test daphnids were not fed during the study.
----------------	--

Test material: Test concentrations were corrected for sample purity. A primary standard of 200 mg/ml was prepared by weighing 2.02 g and diluting it with 10 ml acetone. Appropriate volumes of this standard were added to test water (200 ml) to obtain test concentrations of 1.0, 1.8, 3.2, 5.6 and 10.0 mg/l. Acetone (0.01 ml/200 ml test water) was the solvent control.

Test water: The water used in the study was from a deep well source. The water (1000 liters) was aged and activated biologically in a tank containing live fish. The water contained < 20 ppb aluminum, <0.2 ppb arsenic, <2 ppb cadmium, <3 ppb chromium, <4 ppb cobalt, <3 ppb copper, 12 ppb iron, < 5 ppb lead, <0.5 ppb mercury, <15 ppb nickel, <5 ppb silver, 11 ppb zinc, <0.10 ppb organophosphorus pesticides and <0.50 ppb

organochlorine pesticides (including PCB's). The hardness, alkalinity, conductivity, dissolved concentration and initial pH of the well water were 225 - 275 ppm (as CaCO₃), 325 - 375 ppm (as CaCO₃), 700 micromhos/cm, 9.2 - 10.1 ppm, and 7.8 - 8.3, respectively. The temperature of the water was kept at 22 +/- 1 degrees C.

Test conduct: Tests were conducted in 250 ml glass beakers containing 200 ml of aged well water. The test organisms (10 per test concentration) were added randomly to the test water within 30 minutes of addition of test material. The test concentrations (1.0, 1.8, 3.2, 5.6 and 10 mg/l) were chosen based on the results of a preliminary study performed with concentrations ranging from 0.1 to 100 mg/l. Two additional groups of 10 organisms were exposed to dilution water alone or water containing the solvent. Each condition was tested in duplicate. All organisms were observed initially and after 24 and 48 hours of exposure for mortality and abnormal behavior (surfacing, clumping and lying on the bottom of the vessels). The pH, dissolved oxygen concentration and temperature of the control were determined at the beginning and end of the study. Water containing 1.0, 3.2 and 10 mg/l was analyzed for pH, dissolved oxygen concentration and temperature at the end (but not the beginning) of the study. The vessels were to be aerated if the dissolved oxygen level was less than or equal to 40% saturation.

Statistical analysis: Concentration vs. lethality data were analyzed by a computer program which utilized the binomial, moving average and probit tests to determine the LC₅₀ value (and 95% confidence limit). If no mortality occurred or if a dose-response could not be determined over a reasonable range (< 37 to > 63%), an LC₅₀ value could not be calculated. The method of calculation selected for presentation was the one that gave the narrowest confidence limit.

Test substance : The purity of the test material was 99.13%. The composition of the remaining 0.87% is unknown.

Reliability : (2) valid with restrictions
The results at the two highest concentrations may have been influenced by insolubility of the test material. Test concentrations were not analytically confirmed.

29.10.2002

(20)

Type :
Species : other: Daphnia (not specified)
Exposure period : 48 hour(s)
Unit : mg/l
Limit test :
Analytical monitoring : no
LC50 : 0.295 mg/l
Method : other
Year : 2003
GLP : no
Test substance : as prescribed by 1.1 - 1.4
Remark : EPIWIN ECOSAR (v0.99) was used to obtain the calculated LC50 value. Inputs to the model are CAS Number 3748-13-8, the melting point (-39 degrees C), boiling point (231 degrees C), vapor pressure (4 mm Hg) and water solubility (5 mg/l).
Result : The LC50 value was 0.295 mg/l
Reliability : (2) valid with restrictions
 Data were obtained by modeling.

(13)

4.3 TOXICITY TO AQUATIC PLANTS E.G. ALGAE

Species	: Selenastrum capricornutum (Algae)
Endpoint	: growth rate
Exposure period	: 96 hour(s)
Unit	: mg/l
Analytical monitoring	: no
NOEC	: m = 1.77
EC50	: m = 4.92
Method	: OECD Guide-line 201 "Algae, Growth Inhibition Test"
Year	: 1987
GLP	: yes
Test substance	: as prescribed by 1.1 – 1.4
Remark	: Study personnel stated that "the use of the solvent produced a slight lag in the growth of cells but did not depress the population to a degree severe enough to confound the concentration effects".
Result	: Cell counts given below are reported as the number of cells/ml x 10 ⁴ . Mean cell counts at 24 hours were reported as "less than 10" for all flasks. Average cell counts of controls at 48, 72 and 96 hours were 11, 75 and 241, respectively. Mean counts of cells exposed to 1.8 mg/l at 48, 72 and 96 hours were 14, 116 and 281, respectively. Inhibition of cell growth was noted with concentrations greater than or equal to 3.2 mg/l. The mean number of cells (and percent inhibition) of cells exposed to 3.2 mg/ml at 48, 72 and 96 hours were 11, 61 (19%) and 178 (26%), respectively. The data for one 3.2 mg/l flask were eliminated because the values at 48, 72 and 96 hours (3, < 10 and 25) were considerably lower than the average. Numbers of cells exposed to 5.6 ppm (and percent inhibition) for 48, 72 and 96 hours were 10, 15 (80%) and 90 (63%). Inhibition of cell growth was observed as early as 48 hours for cells exposed to 10 and 18 mg/l (the results were "less than 10 x 10 ⁴ "). Cell counts (and percent inhibition) of cells exposed to 10 mg/l at 72 and 96 hours were 10 (87%) and 12 (95%). No growth occurred in cells exposed to 18 mg/l for 72 or 96 hours.
Test condition	: <p>The rate of cell growth was satisfactory in the controls (greater than 16 x inoculum level at 72 hours) for acceptable data transformation. The correlation coefficients for the regression at 72 and 96 hours were 0.9 and 0.96, respectively. The no effect concentrations at 72 and 96 hours were 1.88 and 1.77 mg/l, respectively. The EC₅₀ values at these times were 4.93 and 4.92 mg/l, respectively.</p> <p>Organisms: Selenastrum capricornutum (ATCC 2262) were propagated at 21 – 25 +/- 2 degrees C under 4000 lux illumination (continuous light). Stocks were subcultured on a regular basis (generally at 1-4 week intervals).</p> <p>Three-five day old suspensions that yielded 1 x 10⁴ cells/ml were used for the test. During the tests, algae were shaken, illuminated at 8000 lux, and maintained at a temperature of 22 – 22.5 degrees C.</p> <p>Medium: OECD fresh water algal culture medium was prepared with distilled or deionized water in non-metallic containers and reconstituted with nutrients, salts and trace elements (as specified in the guideline). Medium was sterilized before use by filtration (<= 0.45 microns) or autoclaving.</p> <p>Test material: The test material was diluted with anhydrous acetone to a concentration of 10,000 times the highest concentration to be used in the test. The stock was stored in the dark until used. Working standards in acetone were prepared so that 10 microliters of each standard would produce the desired concentration to be tested.</p> <p>Test conduct: Based on results of a preliminary range-finding test, concentrations of 1.8, 3.2, 5.6, 10 and 18 mg/l were tested. Controls</p>

containing 10 microliters of acetone and 100 ml of algal suspension also were established. Each concentration (including control) was tested in triplicate. The initial and final pH of all media were recorded. The flasks were incubated for 96 hours and cells were counted with a hemocytometer daily.

A separate test was performed with untreated cells (medium control). The results were compared with those of the solvent control to determine if the solvent alone had an effect on cell growth.

Statistical analysis: The EC₅₀ values at 24, 48, 72 and 96 hours were calculated by regression analysis, using the percent inhibition of growth calculated for each concentration. The percent inhibition of growth at each time was calculated by subtracting the mean cell count of test vessels (Tt) from that of controls (Ct), dividing the result by the Ct, and multiplying the result times 100. The data were graphed, and the no observed effect concentration was determined by extrapolation of the regression line or by data or graph inspection.

Test substance : The test material (CT-256-86) contained 97.5-99.1% m-DIPEB, 0.028 - 0.7% m-IPEC (m-Isopropenyl cumene, CAS No. 1129-29-9), 0–0.50% p-DIPEB (CAS No. 1605-18-1), 0–0.10% DIPB (1,3-Diisopropylbenzene, CAS No. 99-62-7), 0–0.07% m-IPES (m-Isopropenyl styrene; CAS No. 52780-24-2) and 0.2 – 1.8% unknowns.

Reliability : (2) valid with restrictions
Test concentrations were not analytically confirmed. Based on the results of other aquatic toxicity tests, it is likely that all test material at the highest two concentrations was not in solution.

29.10.2002

(10)

Type :
Species : other:green algae
Exposure period : 96 hour(s)
Unit : mg/l
Limit test :
Analytical monitoring : no
EC50 : 0.218 mg/l
Method : other
Year : 2003
GLP : no
Test substance : as prescribed by 1.1 – 1.4
Remark : EPIWIN ECOSAR (v0.99) was used to obtain the calculated LC50 value. Inputs to the model are CAS Number 3748-13-8, the melting point (-39 degrees C), boiling point (231 degrees C), vapor pressure (4 mm Hg) and water solubility (5 mg/l).
Result : The LC50 value was 0.218 mg/l
Reliability : (2) valid with restrictions
Data were obtained by modeling.

(13)

4.4 TOXICITY TO MICROORGANISMS E.G. BACTERIA

4.5.1 CHRONIC TOXICITY TO FISH

4.5.2 CHRONIC TOXICITY TO AQUATIC INVERTEBRATES

4.6.1 TOXICITY TO SOIL DWELLING ORGANISMS

4.6.2 TOXICITY TO TERRESTRIAL PLANTS

4.6.3 TOXICITY TO OTHER NON-MAMM. TERRESTRIAL SPECIES

4.7 BIOLOGICAL EFFECTS MONITORING

4.8 BIOTRANSFORMATION AND KINETICS

4.9 ADDITIONAL REMARKS

5.1.1 ACUTE ORAL TOXICITY

Type : LD₅₀
Species : rat
Strain : Sprague-Dawley
Sex : male/female
Number of animals : 100
Vehicle : no data
Value : = 13.2 ml/kg bw
Method : other
Year : 1981
GLP : yes
Test substance : as prescribed by 1.1 - 1.4
Remark : The method of Litchfield and Wilcoxon could not be used to determine the LD₅₀ value for males. It was estimated to be > 20 ml/kg because only one death occurred at this dose. Because the test for parallelism of dose-response curves or calculation of relative potency could not be carried out, the LD₅₀ values for males and females combined could not be determined.

Using a density of 0.925 (as stated in the MSDS), the LD50 value can be converted to 12.2 g/kg.

Result : Range finding study: None of the animals treated with <= 8.0 ml/kg died over the 7-day test period. One out of 2 females exposed to 10.0 ml/kg died the second day after treatment. Other animals given 10.0 ml/kg survived to termination. Many of the animals exhibited weight loss during the study. Weight loss did not exhibit any dose or sex-related trends. Clinical signs observed 24 to 72 hours after treatment included diarrhea, soft stool, wet area around anus, urine-soaked fur, lacrimation, nasal discharge, red nasal discharge, lethargy, crusty nose and face, swollen feet, paw cut, and moribund condition (for the rat that died). The symptoms increased in frequency with increasing concentrations of test material. There were no sex-related trends. Abnormal necropsy findings in treated animals included colon and/or cecum distended with gas (1 male and 1 female treated with 1.3 or 6.3 ml/kg, respectively), urinary bladder distended with reddish fluid (1 female treated with 10 ml/kg), yellow-green material in the stomach (1 female treated with 10 ml/kg), and yellow fluid in the ileum and cecum (1 female treated with 10 ml/kg). All control animals had normal necropsies.

Main study: None of the animals treated with 0 (control) or 8 ml/kg died. The mortality rates of animals treated with 10.0, 12.6, 15.8, or 20.0 ml/kg were 1/10 (female), 3/10 (all females), 3/10 (all females), and 3/10 (1 male and 2 females). All animals that died succumbed between days 2 and 5. The LD₅₀ value (and 95% confidence limits) was 13.2 (9.9-17.7) ml/kg for females and greater than 20.0 ml/kg for males. All animals (including controls) lost weight for a few days after dosing. Control and treated animals began to gain weight 48 and 96 hours after dosing, respectively. Total body weight gains over the 15 day period were similar between groups. Clinical signs in rats orally treated with 8.0 to 20 ml/kg m-DIPEB included diarrhea, lacrimation, lethargy, urine-soaked fur, nasal discharge, alopecia, crusty nose and eyes, and cold body temperature. Four out of five males treated with 20 ml/kg exhibited alopecia/edema around the anus. Most of the signs were present only for the first days of the study (with the exception of alopecia, which generally appeared a week after treatment). The frequency or variety of signs did not appear to increase with increasing doses of test material, and did not exhibit any sex-related trends (with the exception of alopecia/edema around the anus of high dose males). One male in the control group exhibited a crusty nose on day 8. All other animals assigned to the control group appeared normal.

Test condition	<p>In general, significant gross findings at necropsy were limited to animals found dead (mostly females). These included stomach (brownish-black material, distended with air, filled with yellow-green material, bright yellow fluid), intestinal tract (distended with dark brown material, filled with yellow-red material, yellow material, reddish fluid, yellowish-brown fluid and yellowish-brown material), urinary bladder (dark-colored fluid), and the carcass (alopecia, urine-soaked fur, autolysis, red material around the nose area and crusty eyes and face), and bright yellow nasal discharge and yellow material around nose. Findings in the lung (2 - 3 mm depressed area of one male treated with 15.8 ml/kg) and testes (red pediculated area in the fat of the epididymis of one male treated with 15.8 ml/kg) were considered incidental in nature.</p> <p>: One hundred (50/sex) young adult TAC:N(SD)fBR rats were used for the study. Animals were quarantined and acclimated to laboratory conditions for a least a week prior to study initiation. They were individually housed in stainless steel cages with wire mesh floors. The cages were placed in a 6 cubic meter stainless steel and glass inhalation chamber. The chamber was well-ventilated (approximately 20 changes per hour), continuously monitored for temperature and humidity and had a controlled, 12 hr light/dark cycle. Food and tap water were available ad libitum (with the exception that rats were fasted overnight before dosing). All animals used in the studies were in good health.</p> <p>A preliminary range finding test in which 2 animals/sex were intubated with 1.3, 1.6, 2.0, 2.5, 3.2, 4.0, 5.0, 6.3, 8.0 and 10.0 ml/kg was performed prior to the main study. Males and females used for range-finding weighed 251-329 g, and 201 - 246 g, respectively. For the main study, five animals/sex were intubated with 0.0 (control) 8.0, 10.0, 12.6, 15.8, and 20.0 ml/kg test material. Males and females used for the main study weighed 227 - 267 g, and 170 - 241 g, respectively.</p> <p>Animals used in the range-finding study were observed for at least 4 days after dosing. They were weighed prior to treatment and 1, 2, 3, 4 and 7 days after treatment (at termination). Survivors were euthanized 7 days after treatment.</p> <p>Main study animals were frequently observed for mortality and signs of toxicity during the day of dosing and twice daily thereafter. They were weighed prior to treatment and 1, 2, 3, 4, 7, 11 and 15 days after treatment (at study termination). All animals surviving to day 15 were euthanized.</p> <p>In both studies, all animals that died and survived to study termination were subjected to a complete gross necropsy under the supervision of a board-certified veterinary pathologist as soon as possible after death.</p> <p>The method of Litchfield and Wilcoxon was used to calculate the LD₅₀ values (with confidence limits).</p>
Test substance	: The test material (#11583B14) was 100% m-DIPEB.
Reliability	: (1) valid without restriction Comparable to a guideline study.
Flag	: Critical study for SIDS endpoint
29.10.2002	
Type	: LD ₅₀
Species	: rat
Strain	: Sprague-Dawley
Sex	: male/female
Number of animals	: 10
Vehicle	: no data
Value	: > 5000 mg/kg bw
Method	: other

(3)

Year	:	1981
GLP	:	no data
Test substance	:	as prescribed by 1.1 - 1.4
Result	:	Two females were found dead on day 10. Labored breathing was observed in these rats just prior to death. Pathological findings included red hepatization and expended lungs, indicative of acute pneumonia or pneumonitis unrelated to treatment. There were no other deaths. Gross necropsies of survivors were unremarkable.
Test condition	:	All animals exhibited soft feces, sedation, wet or crusty muzzle, and/or a wet peri-anal area after dosing. These symptoms also were observed 24 hour after treatment in most animals, but resolved within 6 days. Young adult Crl:COBS CD (SD) rats (5/sex) with acceptable body weights (weights were not stated) and general health were used. The rats were housed individually under a 12 hr light/dark cycle. Food and water were available ad libitum, except for an overnight fast prior to dosing. The animals were given a single oral dose of 5 g/kg test material (presumably by gavage). Animals were observed frequently during the day of dosing (Day 0) and twice daily for 14 days. All surviving animals were euthanized on Day 14 and examined grossly. Gross necropsies also were performed on animals that died during the study.
Test substance	:	The test material (CL 116,755) was 100% m-DIPEB.
Reliability	:	(2) valid with restrictions Only one dose was tested. The condition of the animals was likely to have been influenced by the presence of a respiratory infection.

29.10.2002

(5)

5.1.2 ACUTE INHALATION TOXICITY

Type	:	LC ₅₀
Species	:	rat
Strain	:	Sprague-Dawley
Sex	:	male/female
Number of animals	:	20
Vehicle	:	
Exposure time	:	6 hour(s)
Method	:	other
Year	:	1986
GLP	:	no data
Test substance	:	as prescribed by 1.1 - 1.4
Result	:	The mean actual exposure concentrations (+/- SD) were 0.545 +/- 0.062 and 5.576 +/- 0.417 mg/l for the nominal concentrations of 3 and 15 mg/l. The MMAD and GSD for the two concentrations were 1.9 - 2.0 micrometers and 1.7, respectively.

All animals exposed to 5.576 mg/l died within 1 day of exposure. Signs of toxicity such as wet fur, red perinasal wetness, lacrimation, whole body tremors, dermal irritation, hyperactivity, ataxia, and mouth breathing were observed during the first 90 minutes of exposure to 5.576 mg/l. A complete loss of motor activity was observed in these animals for the remainder of the exposure period. After exposure, all animals exhibited absent toe, tail pinch, and surface righting reflexes, hypothermia, respiratory difficulties, wet fur, and dermal irritation. One high dose female also had an eye opacity. All high dose animals appeared to be moribund before death. Necropsies of the dead animals revealed discoloration of the lungs and kidneys and wet fur.

None of the animals exposed to 0.545 mg/l died. The only signs observed in rats exposed to this concentration were ocular irritation (blepharospasm and lacrimation) during exposure. Mean body weights for these animals

	were lower on Day 1 and higher on Day 5 than at the beginning of exposure. There were no gross lesions in these rats at necropsy.	
Test condition	: The LC ₅₀ value was therefore > 0.54 and < 5.6 mg/l (or 540 or 5600 mg/m ³) : Test article generation: The target nominal concentrations were 3 and 15 mg/l. An aerosol was generated with Laskin single-barrel nebulizer (for the 3 mg/l exposure) or a Laskin four-barrel nebulizer (for the 15 mg/l exposure). The nebulizer air pressure and air flow rate for the 3 mg/l exposure were 20 psig and 15 l/min and for the 15 mg/l exposure were 20 psig and 53 l/min. The chamber airflow for the 3 mg/l exposure was diluted with air to 60 l/min. The total volume of the test chambers was approximately 120 liters. The average temperature and relative humidity (+/- SD) of the low-concentration atmosphere were 20 +/- 0 degrees C and 30 +/- 2%, respectively. For the high-concentration atmosphere, these variables were 24 +/- 1 degrees C and 26 +/- 9%, respectively. Test atmosphere was sampled (using a filter) for 2 minutes at 35, 80, 125, 190, 245, 300 and 355 minutes into the exposure for the low concentration and 35, 95, 135, 185, 235, 285 and 340 minutes into the exposure for the high concentration. The concentration of test material was determined gravimetrically. The mass median aerodynamic diameter and geometric standard deviation (GSD) also were determined (method not stated). : Test conduct: Five rats/sex were exposed to each test atmosphere for 6 hours. Prior to exposure, the males and females weighed 188-239 and 150-171 g, respectively. The animals were observed during the exposure and 5 day recovery period (intervals were not stated). Weights were recorded on the day of exposure, one day after exposure and at termination (day 5). All animals were necropsied at death or at scheduled termination.	
Test substance	: The test material (CT-256-86) contained 97.5-99.1% m-DIPEB, 0.028-0.7% m-IPEC (m-Isopropenyl cumene, CAS No. 1129-29-9), 0 - 0.50% p-DIPEB (CAS No. 1605-18-1), 0 - 0.10% DIPB (1,3-Diisopropylbenzene, CAS No. 99-62-7), 0 - 0.07% m-IPES (m-Isopropenyl styrene; CAS No. 52780-24-2) and 0.2 - 1.8% unknowns.	
Reliability	: (1) valid without restriction Meets generally accepted scientific standards and is described in sufficient detail.	
Flag 29.10.2002	: Critical study for SIDS endpoint	(22)
Type	: other	
Species	: rat	
Strain	: Sprague-Dawley	
Sex	: male/female	
Number of animals	: 10	
Vehicle	:	
Exposure time	: 6 hour(s)	
Method	: other	
Year	: 1986	
GLP	: no data	
Test substance	: as prescribed by 1.1 - 1.4	
Result	: None of the animals died. There were no signs of toxicity during exposure to or after a saturated vapor of test material for 6 hours. No remarkable gross lesions were evident at necropsy. Animals gained weight during the 14 day recovery period.	
Test condition	: Animals: Rats weighed 200-300 grams and were approximately 5-8 weeks of age at the beginning of the study. They were acclimated for at least 5 days before exposure. Rats received food and water ad libitum (except during the exposure period). Five healthy rats/sex were used in the test.	

Vapor generation: Approximately 100 grams of test material were placed

into a large glass tray. The tray was placed in a 120-liter plexiglass chamber which was then tightly sealed. The sample was allowed to evaporate overnight. A mixing fan was operated for 30-minute intervals to thoroughly distribute the vapors. The temperature was maintained at 23 degrees C.

Test conduct: After approximately 18 hours of equilibration of the test material with chamber air, rats were placed into a gasketed drawer-type cage. The cage was quickly inserted through a specially sealed opening in the front of the chamber to minimize vapor loss. A separate chamber was used/sex. Oxygen was added (as needed) to maintain a chamber oxygen content of approximately 20%.

The rats were exposed to the atmosphere for 6 hours. They were observed at least once every 30 minutes during exposure. After the exposure period, the rats were placed in a well-ventilated area, examined carefully and returned to their normal housing quarters. Rats were examined twice a day for 14 days for signs of toxicity. Weights were recorded on the day of exposure and 7 and 14 days after exposure. All survivors were euthanized after 14 days and subjected to gross necropsy.

Test substance : The test material (CT-256-86) contained 97.5-99.1% m-DIPEB, 0.028-0.7% m-IPEC (m-Isopropenyl cumene, CAS No. 1129-29-9), 0 - 0.50% p-DIPEB (CAS No. 1605-18-1), 0 - 0.10% DIPB (1,3-Diisopropylbenzene, CAS No. 99-62-7), 0 - 0.07% m-IPES (m-Isopropenyl styrene; CAS No. 52780-24-2) and 0.2 - 1.8% unknowns.

Reliability : (2) valid with restrictions
The concentration of test material in the vapor was not determined analytically. Therefore, whether or not the atmosphere was actually saturated with vapor during exposure is unknown.

29.10.2002

(21)

5.1.3 ACUTE DERMAL TOXICITY

Type : LD₅₀
Species : rabbit
Strain : New Zealand white
Sex : male/female
Number of animals : 10
Vehicle :
Value : > 2000 mg/kg bw
Method : other
Year : 1981
GLP : no data
Test substance : as prescribed by 1.1 - 1.4
Remark : The study was subjected to a quality assurance audit. However, there is no indication that the study was performed according to GLP. The same study is described in Section 5.2.1 below (irritation). The study was given a reliability rating of 1 for irritation, since the one concentration tested was adequate for the endpoint.
Result : None of the animals died during the study, gross necropsies were normal, and all animals gained weight. The only effect of treatment was slight dermal irritation. Erythema scores of 1 were observed in all males and females on day 1, all males and 2 females on day 2, 3 males and 2 females on day 3 and 4 males and 1 female on day 4. An erythema score of 2 was noted in one female on day 3 and 2 females on day 4. One female had an erythema score of 3 on day 4. All edema scores were 0. The mean irritation scores for both sexes for days 1-4 were 1.0, 0.7, 0.7 and 1.2, respectively. All scores on days 7 and 14 were 0.
Test condition : Young adult rabbits (5/sex/dose) were randomly selected from a larger pool of animals with acceptable body weights (2340-2698 g for males and 2078-

2788 g for females) and general health. Rabbits were individually housed under a 12 hour light/dark cycle. Food and water were available ad libitum. The dorsal surface of all rabbits was clipped the day prior to dosing. Just prior to dosing, the skin was abraded in a lattice formation with a hypodermic needle drawn across the surface of the skin. Care was taken to penetrate the stratum corneum, but not the dermis. Test material was administered with a syringe at a dose of 2 g/kg (based on the specific gravity of the test material and the weight of the animal). The test material was held in place with an occlusive wrap secured by a bandage and elastic tape. The dressings were removed after 24 hours and the excess material was wiped off.

The animals were observed for signs of toxicity at 20 minutes, 1, 2 and 4 hours after dosing on day 0, and twice daily from days 1-14. Physical examinations were performed prior to dosing and on day 14. Body weights were recorded on days 0, 1, 2, 3, 6, 10 and 14. Dermal irritation was scored according to the method of Draize on days 1, 2, 3, 4, 7 and 14. The degree of erythema and eschar formation and edema were each scored on a scale of 0-4. All survivors were euthanized and subjected to a complete gross necropsy on day 14. Samples from treated and untreated skin were taken from the back of each animal and retained.

Test substance : The test material (CL 116,755) was 100% m-DIPEB.
Reliability : (2) valid with restrictions
 Only one dose was tested.

29.10.2002

(4)

5.1.4 ACUTE TOXICITY, OTHER ROUTES

5.2.1 SKIN IRRITATION

Species : rabbit
Concentration : undiluted
Exposure : occlusive
Exposure time : 24 hour(s)
Number of animals : 10
PDII :
Result : slightly irritating
EC classification :
Method : other
Year : 1981
GLP : no data
Test substance : as prescribed by 1.1 - 1.4
Remark : The study was subjected to a quality assurance audit. However, there is no indication that the study was performed according to GLP. The same study is described above in Section 5.1.3 (acute dermal toxicity).
Result : None of the animals died during the study, gross necropsies were normal, and all animals gained weight. The only effect of treatment was slight dermal irritation. Erythema scores of 1 were observed in all males and females on day 1, all males and 2 females on day 2, 3 males and 2 females on day 3 and 4 males and 1 female on day 4. An erythema score of 2 was noted in one female on day 3 and 2 females on day 4. One female had an erythema score of 3 on day 4. All edema scores were 0. The mean irritation scores for both sexes for days 1-4 were 1.0, 0.7, 0.7 and 1.2, respectively. All scores on days 7 and 14 were 0.
Test condition : Young adult rabbits (5/sex/dose) were randomly selected from a larger pool of animals with acceptable body weights (2340-2698 g for males and 2078-2788 g for females) and general health. Rabbits were individually housed under a 12 hour light/dark cycle. Food and water were available ad libitum.

The dorsal surface of all rabbits was clipped the day prior to dosing. Just prior to dosing, the skin was abraded in a lattice formation with a hypodermic needle drawn across the surface of the skin. Care was taken to penetrate the stratum corneum, but not the dermis. Test material was administered with a syringe at a dose of 2 g/kg (based on the specific gravity of the test material and the weight of the animal). The test material was held in place with an occlusive wrap secured by a bandage and elastic tape. The dressings were removed after 24 hours and the excess material was wiped off.

The animals were observed for signs of toxicity at 20 minutes, 1, 2 and 4 hours after dosing on day 0, and twice daily from days 1 - 14. Physical examinations were performed prior to dosing and on day 14. Body weights were recorded on days 0, 1, 2, 3, 6, 10 and 14. Dermal irritation was scored according to the method of Draize on days 1, 2, 3, 4, 7 and 14. The degree of erythema and eschar formation and edema were each scored on a scale of 0 - 4. All survivors were euthanized and subjected to a complete gross necropsy on day 14. Samples from treated and untreated skin were taken from the back of each animal and retained.

Test substance : The test material (CL 116,755) was 100% m-DIPEB.
Reliability : (1) valid without restriction
Meets generally accepted scientific standards and is described in sufficient detail.

29.10.2002

(4)

5.2.2 EYE IRRITATION

Species : rabbit
Concentration : undiluted
Dose : .1 ml
Exposure Time :
Comment :
Number of animals : 9
Result : slightly irritating
EC classification : irritating
Method : Draize Test
Year : 1981
GLP : no data
Test substance : as prescribed by 1.1 - 1.4
Remark : The study was subjected to a QA audit. The Draize score is consistent with a classification of practically non-irritating.

Result : No irritation to the cornea or iris was observed at any time point. Discharge, chemosis and/or redness of the conjunctivae were observed in most animals. The individual scores for each of these conditions were not listed; only the total Draize scores. The total Draize scores for unwashed and washed eyes ranged from 0 - 8 and 0 - 6 from days 1 - 4, respectively (out of a maximum score of 110). One animal with an unwashed eye experienced no irritation (all scores were 0). On day 7 one rabbit with an unwashed eye and another with a washed eye had scores of 2. All scores on days 10 and 13 were 0. Average scores for unwashed eyes on days 1, 2, 3, 4 and 7 were 2.6, 2.6, 3.6, 4.0 and 0.3, respectively. The scores for washed eyes at these times were 0.7, 2.0, 2.0, 3.3, and 0.7, respectively.

All animals gained weight over the study and none of them died. Nasal discharge was observed in one animal with unwashed eyes on day 2 and two animals with unwashed eyes on day 3.

Test condition : Test material (100 microliters) was placed in the cupped lower lid of the right eye of each of 9 male New Zealand White rabbits with acceptable body weights (2164 - 2852 g). A test with florescein conducted the day before instillation revealed that the animals did not have existing corneal

injury. Six of the animals received no further treatment. The right eyes of the other three rabbits were rinsed with water for 60 seconds, 30 seconds after treatment. The animals were observed twice daily for general condition, behavior and signs of toxicity. Body weights were recorded on days 0, 6 and 14. The eyes were examined for discharge, chemosis, inflammation, and opacity according to the Draize method on days 1, 2, 3, 4, 7, 10 and 13. Food and water were supplied ad libitum. All rabbits were euthanized without necropsy on day 14.

Test substance : The test material (CL 116,755) was 100% m-DIPEB.

Reliability : (2) valid with restrictions

The test documentation did not list the individual scores for discharge, chemosis and/or redness of the conjunctivae.

29.10.2002

(6)

5.3 SENSITIZATION

Type : other

Species : guinea pig

Number of animals : 29

Vehicle :

Result : sensitizing

Classification : sensitizing

Method : other

Year : 1981

GLP : yes

Test substance : as prescribed by 1.1 - 1.4

Result : Range-finding study: All animals in the range finding study and those assigned to serve as non-sensitized controls at rechallenge exhibited weight loss over the 3 - 4 day observation period. Weight loss for these animals was considered to be the result of experimental stress. Other test animals that were weighed after more days on the study gained weight.

During this study, 1 male and 1 female assigned to the test material group were found dead. The female was found dead the second day of the second induction and the male was found dead the day after the third induction. At rechallenge, 1 female assigned to the non-sensitized primary irritation control group was found dead on the day after the last skin evaluation for rechallenge, just prior to being weighed. Whether these deaths were considered to be related to test material was not stated. No gross lesions were found in the male. The only female that died after the second induction had mottled lungs and the small intestine was distended with air. The necropsy data for the other female were missing.

For the range-finding study, no irritation was seen for test concentrations of 50% or below. Only 1/4 animals exhibited irritation at 100% (grade 3 and 2 erythema at 24 and 48 hours, respectively). Therefore the test material was applied at 100% during the induction and challenge phases. At rechallenge, 12.5, 25, 50 and 100% test material was used. The only non-irritating concentration of DNCB was 0.01%. All other concentrations produced irritation in at least 2 animals. However, the highest concentration (0.1%) did not cause grade 2 irritation. Therefore, 0.1% DNCB was used during the induction phase and 0.01% was used at challenge and rechallenge.

Main study: For the main study, skin condition after the first application of 100% test material appeared normal. Skin reactions [erythema (avg. grade 3.3 - 4.0), edema (grade 1 - 2), eschar formation (grade 4 erythema), bleeding at the test site, fissures and/or desquamation] first appeared after the second application, peaked after the third application, and remained steady for the remainder of the induction period. Similar findings were

observed for DNCB (with the exception that the severity scores peaked after the 5th application).

The average erythema score 48 hours after challenge with the test material (but not 24 hours after challenge) was higher than that of the range-finding study (1.7 vs. 0.5). For DNCB, the erythema scores after 24 and 48 hours of induction (1.9 and 1.2, respectively) were greater than those for the range-finding study (0 at both time points). All edema scores for test material and DNCB were 0 (with the exception of a grade 0.3 edema after 24 hours challenge with DNCB).

Twenty-four hours after rechallenge, average erythema scores of animals rechallenged with 12.5, 25, 50 and 100% test material were 0.6, 0.6, 1.5 and 1.3, respectively. The corresponding values for primary irritation control animals treated with these concentrations and examined after 24 hours were 0.3, 0.5, 0.3 and 0.5, respectively. Edema scores for rechallenged animals (ranged from 0.0 to 0.2, with no effect of concentration) were similar to primary irritation controls (ranged from 0 - 0.3). Forty-eight hours after rechallenge, average erythema scores of animals rechallenged with 12.5, 25, 50 and 100% were 0.2, 0.5, 0.7 and 1.6, respectively. The corresponding values for primary irritation control animals treated with these concentrations and examined after 24 hours were 0.0, 0.3, 0.8 and 1.5, respectively.

Compared to scores from DNCB-treated animals in the range-finding test (all were 0), erythema and edema scores of animals rechallenged with DNCB were greater at 24 (2.0 and 0.7, respectively) and 48 hours (2.0 and 0.5, respectively).

Test condition

: Thirty-seven (19 males, 18 females) Hartley albino guinea pigs were used. Animals were acclimated to laboratory conditions for at least 7 days prior to study initiation. All animals used appeared healthy. Animals were randomly assigned to 5 different treatment groups.

On the day prior to each phase of the study, application sites on the dorsal surface of each animal were closely clipped with electric clippers and then shaved with a safety razor. This procedure was repeated as required. Each material was applied (0.5 ml) on a 1 x 1 Webril patch. The patches were held in place with Blenderm tape. The patches and entire trunk were wrapped with an impervious binder consisting of plastic wrap, gauze bandage, adhesive tape and masking tape. The patches were removed after 24 hours. After treatment, animals were maintained in inhalation chambers. Skin condition was evaluated upon patch removal and 24 hours later (48 hours after treatment).

Four animals/sex (325 - 426 g) were used for the range-finding study. In this study, 2 animals/sex were exposed to 100% test material and 1, 3, 10, 25 and 50% test material in 1% olive oil, and 2 animals/sex were exposed to 0.01, 0.025, 0.05 and 0.1% 1,2 dinitrochlorobenzene (DNCB) in alcohol (positive control). The patches were applied to the prepared site, with the patches containing the highest concentrations applied to the left side of the animal (highest at upper left) and the lowest concentrations applied to the right side (lowest at lower right). The highest concentration producing a mean score for erythema of less than 2 in the range-finding study was used for induction and challenge doses in the main study. A non-irritating dose of the DNCB was used as the positive control for the challenge phase of the main study.

Initially, 13 males and 12 females (349 - 504 g) were used for the main study. Eight males and 7 females were exposed to 100% test material for induction and challenge and 100% material and 12.5, 25 and 50% test material in olive oil for rechallenge. Five animals per sex were exposed to 0.1% DNCB in alcohol for induction and 0.01% DNCB for challenge and

rechallenge. For induction, the test materials were applied to the appropriate test site 3 times per week on alternating days until a total of 10 applications were made. Skin condition was evaluated 24 and 48 hours after each application. Challenge doses were applied to previously unused sites 14 days after the last induction dose was applied. Skin condition was evaluated 24 and 48 hours after challenge.

All animals induced and challenged with test material were rechallenged with test material 11 days after challenge (2 concentrations per side), with the lowest concentration at the upper left and the highest at the lower right. DNCB was applied to the positive controls at a previously unused site. An additional 2 animals/sex (631 - 889 g) were added as controls for non-sensitized primary skin irritation. These animals were exposed to 100% test material or 12.5, 25 and 50% test material in olive oil at rechallenge only. Skin condition of all animals was evaluated approximately 24 and 48 hours after rechallenge.

All animals were observed for mortality and signs of toxicity twice daily. Body weights were taken before treatment and at termination. Animals found dead were to be necropsied as soon as possible by a board-certified veterinary pathologist. At challenge (for the test material and DNCB) and at rechallenge (for DNCB), mean scores for skin condition were compared to the mean scores found in the range-finding study. If the scores at challenge (and rechallenge for DNCB) were higher than those in the range-finding study, the material caused dermal sensitization. At rechallenge, mean scores of test animals and non-sensitized controls were compared. If the mean scores were higher in animals that had undergone induction than in those that had not (non-sensitized controls), the material caused dermal sensitization.

Test substance Conclusion : The test material (#11583B14) contained 100% m-DIPEB.
: Animals receiving induction applications of 100% test material exhibited a dose-dependent dermal contact sensitization response when challenged with 100% test material and rechallenged with 12.5, 25, 50 and 100% test material. The responses for the positive control DNCB also were positive.

Reliability : (1) valid without restriction
Comparable to a guideline study.

29.10.2002

(2)

5.4 REPEATED DOSE TOXICITY

Species : rat
Sex : male/female
Strain : Sprague-Dawley
Route of admin. : inhalation
Exposure period : 4 weeks
Frequency of treatment : 6 hours/day, 5 days/week
Post obs. period : none
Doses : 107, 510 and 970 mg/m³
Control group : yes
NOAEL : = 510
LOAEL : = 970
Method : other
Year : 1988
GLP : yes
Test substance : as prescribed by 1.1 - 1.4
Remark : Exposure concentrations were selected based on the results of a previous, 5-day range-finding study in male rats (5/group; 7 to 9 weeks old) with 380, 730, 920 and 1400 mg/m³. Ocular and nasal discharge were observed during and after exposure to concentrations > = 730 mg/m³. Mild body

weight loss was noted in all groups, with no relationship to concentration. There were no treatment-related alterations in organ weights and no exposure-related necropsy findings.

The results of the range-finding study also indicated that the vapor concentration in an aerosol atmosphere of test material ranged from 375 to 478 mg/m³ for total (vapor plus aerosol) concentrations ranging from 380 to 1400 mg/m³.

For the main study, the particle size distribution of the 107 mg/m³ atmosphere was not determined (protocol deviation) since there was an insufficient amount of aerosol present in the chamber at this concentration (i.e. most was vapor).

Study personnel did not consider the effects observed at 510 mg/m³ [reduced weight gain in males early on in the study, increased urine volume in males, increased relative liver weight in males in the absence of any changes in clinical chemistry parameters or pathology, and swollen periocular tissue in males and females during exposure (but not at termination)] to be indicative of toxicity. Therefore, they assigned a no observable adverse effect level (NOAEL) of 510 mg/m³. The summary preparer believes that based on the aforementioned effects at 510 mg/m³, a NOAEL of 107 mg/m³ is more appropriate.

Result

- : The mean actual exposure concentrations (+/- SD) were 107 +/- 13, 510 +/- 29 and 970 +/- 54 mg/m³ for the nominal concentrations of 100, 500 and 1000 mg/m³. The MMADs (and range) for the 510 and 970 mg/m³ concentrations were 3.7 (2.8-4.5) and 3.7 (2.8-4.3) microns, respectively. The GSDs (and range) for these concentrations were 2.6 (1.9 - 4.0) and 2.3(1.8 - 3.0), respectively. The estimated percentage (and range) of particles <=10 microns for the 510 and 970 mg/m³ concentrations were 86 (78 - 91) and 89 (86 - 92), respectively. The mean daily chamber temperature and relative humidity for all groups ranged from 20 - 21 degrees C and 48 - 49%, respectively.

None of the animals died. Symptoms of eye irritation including lacrimation (one male exposed to 107 mg/m³), swollen periocular tissue (3 per sex exposed to 510 mg/m³ and 2 males and 4 females exposed to 970 mg/m³), and periocular encrustation (one control and 2 males exposed to 107 mg/m³) were observed during exposure. Perinasal encrustation was observed during the study in one male exposed to 107 mg/m³ and one female exposed to 970 mg/m³. The days at which these symptoms were observed were not listed. At termination, the incidence of conjunctivitis in the 0, 107, 510 and 970 mg/m³ groups was 2/10, 2/10, 2/10 and 4/10. Two rats exposed to 1000 mg/m³ had severe conjunctivitis. Two high dose females also exhibited alopecia of the head during the study (time was not indicated). There were no other treatment-related clinical signs.

Average body weights and weight gains of males exposed to 510 mg/m³ were significantly lower than controls at day 4 (weight) and from days 0 - 4 and 0 - 11 (weight gains). Average body weight and weight gain of males exposed to 970 mg/m³ were significantly lower than controls on days 4, 11, 18 and 25 (weights) and from days 0 - 14, 0 - 11, 0 - 18 and 0 - 25 (weight gains). Weights and weight gains of females were similar to controls.

A statistically significant increase in the numbers of segmented neutrophils was observed in males and females exposed to 970 mg/m³. This shift was not accompanied by an increase in total leukocyte count. A significant increase in ALT was observed in males (41% greater than control) and females (67% greater than control) exposed to 970 mg/m³ test material. With respect to control, alkaline phosphatase increased by 38% and 43%, respectively, in males and females exposed to 970 mg/m³. Total urine volume of males and exposed to 510 and 970 mg/m³ and females exposed

to 970 mg/m³ increased (but was only significantly different from control for males).

Increases in absolute (females only) and relative (to body weight) liver weights were observed in high dose animals (males and females). High dose males also had increased relative (but not absolute) brain, adrenal and testes weights. Males exposed to 510 mg/m³ also had increased relative liver weight.

There were no treatment-related gross or histological lesions at necropsy. Histological lesions included lymphocytic myocarditis of unknown etiology (1 control male), renal cortical necrosis (one control male), and alopecia with pustular dermatitis of the cervical skin (one 970 mg/m³ female). Minimal to mild lesions of various respiratory tract tissues (such as clusters of macrophages in alveolar spaces and minimal-mild laryngitis and tracheitis) were seen in several rats of various groups with no relationship to treatment. Four male rats (two high dose and one each from the other treatment groups) had slight lung hemorrhage, which was believed to be an artifact of the euthanasia technique.

Test condition

: Animals: Thirty-two male rats and 31 females [HSD:(SD)BR], 35 days of age, were received approximately 14 weeks before initiation of the study. Three male rats were euthanized on the day of receipt for quality control. The liver, submandibular lymph nodes, lungs, trachea, kidneys, heart, salivary glands and nasal turbinates and spleen were fixed and examined microscopically. The larynx was inadvertently missed (protocol deviation). Blood samples were obtained from 5 males for serology evaluation. Fecal samples were collected from 5 males and examined for intestinal parasites. Results of these tests plus physical examinations revealed that the rats were in good general health. Only animals that had body weights within 2 standard deviations from the group mean for each sex were eligible for use. Twenty animals/sex were randomly allocated to 4 exposure groups (5/sex/exposure). Animals were weighed and clinically examined prior to exposure. Food and water were available ad libitum during non-exposure periods. Animals were individually housed in stainless-steel wire-mesh cages (14 x 13.5 x 18 cm) during exposures and 2 - 3 per cage (23.5 x 20 x 18 cm) when not exposed. The animals were on a 12- hour light/dark cycle throughout the study.

Test article generation: The target nominal concentrations of vapor plus aerosol were 0, 100, 500 and 1000 mg/m³. Single aspirator tubes were used to generate the 100 and 500 g/m³ atmosphere and a double aspirator tube was used to generate the 1000 mg/m³ atmosphere. Compressed air, supplied to each nebulizer, created a negative pressure causing the test material to be aspirated into the tubes and dispersed as a fine liquid aerosol. The liquid aerosol was then introduced into the top of the exposure chamber where it was diluted to the target concentrations and dispersed throughout the chamber with filtered air. The total volume of the test chambers was approximately 900 liters. The operating air pressures of the nebulizers used to generate the 100, 500 and 1000 mg/m³ target concentrations were approximately 3, 7, and 8 psig, respectively. Chamber temperatures and relative humidities were recorded at least 11 times per exposure with a minimum-maximum thermometer and an Airguide Humidity Indicator.

Two midjet impingers in series (each containing 15 ml of toluene) were used to scrub the chamber atmosphere for test material. Four samples were collected from each chamber during each 6-hour exposure. The sampling time for the 500 and 1000 mg/m³ concentrations was 20 minutes, and for the 100 mg/m³ concentration was 60 minutes. Aliquots (1 microliter) from each impinger sample were analyzed by a gas chromatograph fitted with a flame ionization detector. The daily nominal concentrations were calculated by dividing the total amount of material

delivered to the chamber by the total chamber airflow.

Particle size distributions of the aerosols were measured with a cascade impactor nine times (at least twice a week) during exposure to 500 or 1000 mg/m³. The amount of material that collected on cellulose filters on the stages of the impactor was determined gravimetrically. The data were analyzed by probit analysis to obtain the mass median aerodynamic diameter (MMAD) and geometric standard deviation (GSD).

Study conduct: All animals were acclimated to the chambers (with filtered air only) for 2 days before exposure to test material. Groups of 5 rats/sex (48 days old) were exposed 6 hours/day, 5 consecutive days/week for 3 weeks to 0 (air only), 100, 500 and 1000 mg/m³. During the fourth week, all animals were exposed for 4 days and euthanized on the 5th day. The position of cages in the chambers was rotated weekly in a predetermined pattern.

All animals were observed prior to, during, and following each exposure for signs of toxicity. They also were observed once/day when not exposed. Before the first exposure and at termination, the anterior chambers of the eyes of each animal were examined by a veterinarian. All animals were weighed before the first exposure, and on days 4, 11, 18 and 25.

Urine was collected for approximately 15 hours on the day of termination. Food and water were available ad libitum. The total volume, color and turbidity, specific gravity, pH, occult blood, and concentrations of glucose, ketones, protein, bilirubin and urobilinogen were determined according to standard methods. Although not stated, it is assumed that urine was collected before blood.

Blood was obtained from the orbital sinuses of anesthetized animals at termination. Food was withdrawn during blood collection. Blood was analyzed for total erythrocyte count, hemoglobin, hematocrit, mean corpuscular volume, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, platelet count and prothrombin time. Leukocyte differential smears were prepared from rats in all groups, but were evaluated only for the control and high concentration groups. Reticulocyte smears were prepared for all groups, but were not evaluated. Serum was analyzed for creatinine, sodium, potassium, chloride, alanine aminotransferase (ALT), total protein, albumin, total, direct and indirect bilirubin, aspartate aminotransferase (AST), globulin, creatine kinase, lactate dehydrogenase, sorbitol dehydrogenase, alkaline phosphatase and gamma glutamyl transferase.

All surviving animals were euthanized after blood and urine collection. The brain, liver, lungs, heart, adrenals and testes were weighed. Gross necropsies were performed and selected tissues were fixed in 10% neutral buffered formalin. The spleen, adrenals, brain, esophagus, parathyroids, heart, larynx, lymph nodes, testes, thyroid, eyes, ovaries, pituitary, muscle (gastrocnemius), nerve (sciatic) and gross lesions were examined histologically in controls and animals exposed to 1000 mg/m³ animals. The bone marrow, lungs, nasal turbinates, thymus, kidneys, liver, and trachea were examined histologically for all groups.

Statistical analyses: Data for continuous variables were first analyzed for homogeneity using Bartlett's test. If Bartlett's test indicated heterogeneous variances, data were compared using an analysis of variance (ANOVA) for unequal variances. Medians and quartile deviations were calculated for non-parametric data. These data were analyzed by the Kruskal-Wallis test or by the Wilcoxon rank sum test (as modified by Mann-Whitney). Homogeneous data were analyzed using ANOVA, followed by tests. The level of significance for all comparisons was $p < 0.05$.

5. Toxicity

Id 3748-13-8

Date 07.12.2004

Test substance : The test material (CT-256-86) contained 97.5-99.1% m-DIPEB, 0.028 - 0.7% m-IPEC (m-Isopropenyl cumene, CAS No. 1129-29-9), 0 - 0.50% p-DIPEB (CAS No. 1605-18-1), 0 - 0.10% DIPB (1,3-Diisopropylbenzene, CAS No. 99-62-7), 0 - 0.07% m-IPES (m-Isopropenyl styrene; CAS No. 52780-24-2) and 0.2 - 1.8% unknowns.

Reliability : (1) valid without restriction
Comparable to a guideline study.

Flag : Critical study for SIDS endpoint.

29.10.2002 (9)

Species : rat
Sex : male/female
Strain : Sprague-Dawley
Route of admin. : oral
Exposure period : 15-27 days (males), 40-53 days (females)
Frequency of treatment : daily
Post obs. period : none
Doses : 100, 300 and 1000 mg/kg bw
Control group : yes
NOAEL : = 1000 mg/kg bw
LOAEL : > 1000 mg/kg bw
Method : other: OECD Guideline 421
Year : 2004
GLP : yes
Test substance : as prescribed by 1.1 - 1.4
Remark : This study only contains methods and results pertinent to repeated dose toxicity of the parental animals. Reproductive/developmental effects are listed under Sections 5.8 and 5.9, respectively.

Result : Mortality and clinical signs: None of the parental animals died. At 1000 mg/kg/day, 7/10 males and all females had increased salivation 1 hour after treatment on at least one occasion. One high dose female also exhibited hunched posture, tiptoe gait and excessive grooming one hour after dosing on one occasion. An additional high dose female had similar signs on several occasions and also exhibited signs of lethargy, piloerection, decreased respiration and ptosis on one occasion. All of the signs lasted approximately 10-15 minutes. There were isolated incidences of wet and/or stained fur at 300 mg/kg/day (1 male and 2 females). There were no signs of toxicity at 100 mg/kg/day.

Body weight and food consumption: There was no effect of treatment on parental body weight or food consumption during any period measured.

Organ weight, gross and histopathology data: There was no effect of treatment on parental organ weight. The kidneys of two high dose males had a mottled appearance. Ulcers were seen on the non-glandular region of the stomach of one high dose female. Small, white foci were seen on the intestines of another high dose female. A hard, brown mass was found in the thoracic cavity of an additional high dose female. All animals in the 300 mg/kg group exhibited normal pathology. At 100 mg/kg/day, two males had an accentuated lobular pattern on all lobes of the liver. The kidneys of one of these males had a mottled appearance. Another low dose male had small testes and epididymides. One low dose female had an enlarged spleen and kidneys and a small growth in the intestines. Small, white foci were seen in the intestines of one control female. None of the gross lesions observed were attributed to administration of test material. No treatment-related histopathological changes were observed.

Test condition : Date of study: January 20, 2004 – May 24, 2004

Animals: Male and female Sprague-Dawley CrI:CD® (SD)IGS BR rats were obtained from Charles River (UK) Limited, Manston Road, Margate, Kent.

They were examined on the day of receipt and acclimated for 7 days. A total of 40 animals/sex were accepted into the study. The males and females weighed 321-364 g and 205-240 g at the start of the study.

Animals were housed in groups of five by sex in polypropylene cages with stainless steel grid floor and tops, suspended over paper-line polypropylene trays (except during mating and gestation). One male and one female were housed in similar cages during mating. After evidence of successful mating, males were returned to their original cages. The females were then housed individually in cages with solid floors, soft wood chip bedding and stainless steel tops.

Animals were housed in an environmentally-controlled room, under a 12 hour light/dark cycle. The temperature and relative humidity were maintained to operate within a target range of 21 +/- 2 degrees C and 55 +/- 15 %. The temperature and/or humidity fell outside of these ranges on isolated occasions, but this was not considered to influence the study. Air was changed at least 15 times per hour. Certified feed and water were supplied ad libitum. No contaminants were present in food and water at levels sufficient to affect the outcome.

Test material: The appropriate amount of test material for each group was mixed with arachis oil BP for 2 minutes to ensure that a homogeneous solution was prepared. Samples of the solution were analyzed for stability and homogeneity prior to the study. The results of the analyses showed that the material was homogeneous in arachis oil and stable for at least 14 days (analytical concentrations were 95%, 100% and 107% of nominal 25, 75 and 250 mg/ml solutions after 14 days). Samples of each formulation were taken at the beginning, middle and end of the dosing period and analyzed for concentration of test material. The results of the analyses indicated that the concentrations present were within acceptable limits of the nominal concentrations (analytical concentrations were 102%, 100% and 104% of nominal 25, 75 and 250 mg/ml solutions).

Study design: The animals were randomly allocated by weight to 4 groups of 10 animals per sex (F0 animals) receiving either 0 (vehicle), 100, 300 or 1,000 mg/kg/day test material. Animals in each group were uniquely identified. The vehicle (arachis oil) and test material solutions were administered orally by gavage to their respective groups with a plastic dosing catheter attached to a disposable, plastic syringe once daily. The dosing volume was 4 ml/kg. Administration to F0 males and females began 14 days prior to mating and continued through mating. Females continued receiving test material to postnatal day 5.

Following 14 days of dosing, 10 F0 males were randomly paired on a 1:1 basis with 10 F0 females from the same group for up to 14 days. Following positive identification of mating (presence of a copulatory plug in the vagina or on the tray beneath the cage or the presence of sperm in a vaginal smear following vaginal lavage), the males were returned to their cages. Following mating, all F0 females were allowed to deliver naturally and rear their young to postnatal day 5 (the scheduled day of necropsy).

All F0 animals were observed twice daily for mortality and morbidity (once daily on weekends). Clinical observations were recorded daily. Animals were observed for toxicity at the time of dosing and approximately 1 hour after dosing. Male body weights and food consumption per cage were recorded weekly throughout the study. Body weights and food consumption per cage of females were recorded weekly until mating. After mating, body weights of females were recorded on gestation days 0, 7, 14, and 20 and on lactation days 1 and 4, and food consumption of females was determined for gestation days 1-7, 7-14 and 14-20. Female food consumption also was recorded for the period covering lactation days 1-4.

Males were euthanized after confirmation of successful mating. All surviving adults (including non-fertile animals) and offspring were euthanized on day 5 postpartum. All animals were examined microscopically for internal and external abnormalities. The testes and epididymides of all adult males were weighed. The coagulating glands, epididymides, prostate, seminal vesicles, testes, pituitary, ovaries, uterus/cervix, and vagina from the high dose and control adult males and females were fixed, processed and examined microscopically by a pathologist. The stomachs of all adult animals were also examined.

Statistical evaluations: Data were processed to give group mean values and standard deviations. The food conversion ratio (group mean weekly body weight gain/ food consumption) was calculated for the premating period. Adult body weight, food consumption, and organ weight data were analyzed for homogeneity using Bartlett's test, followed by a one-way analysis of variance (ANOVA). Data that were not homogeneous were subsequently analyzed using a t-test (assuming unequal variances). Dunnett's multiple comparison method was used to analyze data that were homogenous. Relative organ weights were analyzed using the Kruskal-Wallis non parametric rank sum test. Pairwise comparisons were performed using the Mann-Whitney U-test. Histopathological lesions that occurred at an overall frequency of 1 or greater were analyzed using a chi-squared test. Severity grades were analyzed using a Kruskal-Wallis one-way non-parametric ANOVA. Significant differences were reported at the $p < 0.05$, $p < 0.01$ and $p < 0.001$ level (if present).

Test substance : The test material contained > 97.5% m-diisopropenylbenzene. Impurities were not listed.

Reliability : (2) valid with restrictions
The study is valid without restriction for the reproductive/developmental endpoints. However, some of the parameters measured in guideline repeated dose studies (hematologies, clinical chemistries and complete organ histopathology) were not performed. Therefore, a rating of (2) is appropriate for the repeated dose toxicity endpoint.

07.12.2004

(25)

5.5 GENETIC TOXICITY 'IN VITRO'

Type : Ames test
System of testing : S. typhimurium strains TA1535, TA1537, TA98 and TA100
Concentration : 1.5, 5, 15, 50, 150, 500, 1500 and 5000 micrograms/plate
Cytotoxic conc. : ≥ 150 micrograms/plate
Metabolic activation : with and without
Result : negative
Method : OECD Guide-line 471 "Genetic Toxicology: Salmonella typhimurium Reverse Mutation Assay"
Year : 1999
GLP : yes
Test substance : as prescribed by 1.1 - 1.4
Remark : This test was conducted in conjunction with the E coli WP2urvA- test (see below). This study is the critical study for the mutation endpoint.
Result : There was no increase in the number of mutants in any strain exposed to test material (with or without metabolic activation) in either test.

For test 1, the average number of mutants in control strains TA100, TA1535, TA98 and TA1537 (without S-9) were 107, 18, 24 and 9, respectively. The average number of mutants in strains TA100, TA1535, TA98 and TA1537 incubated with test material (without S-9) ranged from 72 - 115, 0 - 23, 16 - 26 and 4 - 14, respectively. The average number of mutants in control strains TA100, TA1535, TA98 and TA1537 (with S-9)

were 115, 12, 35 and 19, respectively. The average number of mutants in strains TA100, TA1535, TA98 and TA1537 incubated with test material (and S-9) ranged from 76 - 115, 6 - 17, 26 - 38 and 14 - 20, respectively.

For test 2, the average number of mutants in control strains TA100, TA1535, TA98 and TA1537 (without S-9) were 76, 20, 28 and 10, respectively. The average number of mutants in strains TA100, TA1535, TA98 and TA1537 incubated with test material (without S-9) ranged from 61 - 78, 16 - 22, 20 - 27 and 8 - 14, respectively. The average number of mutants in control strains TA100, TA1535, TA98 and TA1537 (with S-9) were 76, 13, 28 and 20, respectively. The average number of mutants in strains TA100, TA1535, TA98 and TA1537 incubated with test material (and S-9) ranged from 62 - 85, 10 - 15, 24 - 34 and 15 - 21, respectively.

All of the positive controls induced at least a 3-fold increase in the frequency of revertant colonies compared to controls, thus confirming the sensitivity of the bacterial strains. The spontaneous mutation rates of the controls were acceptable. The results of the characteristics tests for all the strains were satisfactory. The S-9 mix was sterile.

The results of the preliminary toxicity study indicated that the test material was toxic to TA100 at concentrations ≥ 500 micrograms/plate. In the main study, the test material caused a visible reduction in the growth of the bacterial lawn beginning at 150 micrograms/plate (strain TA1535 without activation). Concentrations ≥ 500 micrograms/plate reduced the bacterial lawn in strains TA100 and TA1537 (with and without activation). A concentration of 1500 micrograms/plate reduced the lawn in strain TA98 (without activation) and caused 100% cell death in strain TA1535 (without activation). At 5000 micrograms/ml, the bacterial lawn of strain TA98 was reduced (with S-9).

Test condition

: Bacteria: The Salmonella strains were obtained from the University of California at Berkeley. All strains were stored at -196 degrees C until use. Prior to use, characterization checks were carried out to confirm the amino-acid requirement, presence of *rfa*, R factors, and the spontaneous reversion rate. Overnight cultures were prepared in nutrient broth and incubated at 37 degrees C for approximately 10 hours. Each culture was monitored spectrophotometrically for turbidity with titers determined by viable count analysis on nutrient agar plates.

S9 preparation: S9 was prepared from the livers of male Sprague-Dawley rats (250 g) approximately 1 month before the experiments were conducted. The rats received 3 consecutive daily doses of phenobarbitone/beta naphthoflavone (80 - 100 mg/kg/day) prior to liver removal. Before use, each batch of S9 was assayed for its ability to metabolize the indirect mutagens 2-aminoanthracene and benzo(a)pyrene. The S-9 was stored at -196 degrees C until use. The S-9 mix (5.0 ml S-9, 1.0 ml 1.65 M KCl/0.4 M MgCl₂, 2.5 ml 0.1 M glucose-6-phosphate, 2.0 ml 0.1 M NADPH, 2.0 ml 0.1 M NADH, 25.0 ml 0.2 M sodium phosphate buffer, and 12.5 ml sterile water) was prepared aseptically immediately before the experiments and stored on ice. A 0.5 ml aliquot of S-9 mix and 2 ml of molten, trace histidine or tryptophan-supplemented top agar was overlaid onto a sterile Vogel-Bonner Minimal agar plate in order to assess the sterility of the S9-mix. This procedure was repeated in triplicate on the day of each experiment.

Study conduct: Approximately half-log dilutions of the test material in dried dimethyl sulfoxide (DMSO) were prepared on the day of each experiment. Concentrations were corrected for purity (98.4%). Based on the results of a preliminary study, concentrations of 1.5, 5, 15, 50, 150, 500, 1500 and 5000 micrograms/plate were tested in triplicate for each strain (TA1535, TA1537, TA98 and TA100), with the exception that 1.5, 5 and 15

micrograms/plate were not tested in TA98 with S-9 and 5000 micrograms/plate only was tested in TA98 with S-9. Aliquots (0.1 ml) of the bacterial cultures were dispensed into test tubes followed by 2.0 ml of molten, trace histidine or tryptophan-supplemented top agar, 0.1 ml of the test material, vehicle (DMSO), or positive control [3 or 5 micrograms/plate N-ethyl-N'-nitro-N-nitrosoguanidine (ENNG) for TA100 and TA1535 without S-9; 80 micrograms/plate 9-aminoacridine (9AA) for TA1537 without S-9; and 0.2 micrograms/plate 4-nitroquinoline-1-oxide (4NQO) for TA98 without S-9; 1-2 micrograms/plate 2-aminoanthracene (2AA) for TA100, TA1535 and TA1537 with S-9; and 5 micrograms/plate benzo(a)pyrene for TA98 with S-9] and either 0.5 ml of S-9mix (for experiments with metabolic activation) or phosphate buffer (for experiments without metabolic activation). The contents of each tube were mixed and equally distributed onto the surface of Vogel-Bonner minimal agar plates (one tube per plate). All plates were incubated at 37 degrees C for approximately 48 hours and the frequency of revertant colonies was assessed using a colony counter. The test was repeated using the same experimental conditions.

The assay was considered valid if all tester strains exhibited spontaneous reversion rates similar to historical controls, if the appropriate characteristics for each strain were confirmed, all tester strain cultures contained $1 - 9.9 \times 10^9$ bacteria/ml, each positive control induced at least a 2-fold increase in mutants, there was a minimum of 4 non-toxic concentrations, and there was no evidence of excessive contamination. The test was considered positive if there was a reproducible, dose-related and statistically (Dunnett's method of linear regression) significant increase in the number of revertants in at least one strain.

Test substance	:	The test material (CT-664-99) contained 98.36% m-DIPEB. Impurities were 0.41% p-DIPEB (CAS No. 1605-18-1) and 0.65% unidentified material.
Reliability	:	(1) valid without restriction Guideline study.
29.10.2002		(27)
Type	:	Bacterial reverse mutation assay
System of testing	:	Escherichia coli strain WP2uvrA-
Concentration	:	50, 150, 500, 1500 and 5000 micrograms/plate
Cytotoxic conc.	:	> 5000 micrograms/plate
Metabolic activation	:	with and without
Result	:	negative
Method	:	other
Year	:	1999
GLP	:	yes
Test substance	:	as prescribed by 1.1 - 1.4
Remark	:	This test was conducted in conjunction with the previously described Ames test.
Result	:	There was no increase in the number of mutants in E coli exposed to test material (with or without metabolic activation) in either test.

For test 1, the average number of mutants (without S-9) was 27 for the control and ranged from 25-33 for treated cultures. The average number of mutants (with S-9) was 32 in the control and ranged from 31 - 37 for treated cultures.

For test 2, the average number of mutants (without S-9) was 22 in the control and ranged from 16-26 for treated cultures. The average number of mutants (with S-9) was 28 in the control and ranged from 16 - 24 for treated cultures.

The positive control induced at least a 10-fold increase in the frequency of revertant colonies compared to controls, thus confirming the sensitivity of the bacterial strain. The spontaneous mutation rates of the controls were

	<p>acceptable. The results of the characteristics test were satisfactory. The S-9 mix was sterile.</p> <p>The results of the preliminary toxicity study indicated that the test material was not toxic to <i>E. coli</i> WP2uvrA- at the highest concentration tested (5000 micrograms/plate).</p>
Test condition	<p>: Bacteria: <i>E. coli</i> strain WP2uvrA- was maintained at -196 degrees until use. Characterization checks were carried out to confirm the <i>uvrB</i> or <i>uvrA</i> mutation and the spontaneous reversion rate. Overnight cultures were prepared in nutrient broth and incubated at 37 degrees C for approximately 10 hours. Each culture was monitored spectrophotometrically for turbidity with titers determined by viable count analysis on nutrient agar plates.</p> <p>S9 preparation: S9 was prepared from the livers of male Sprague-Dawley rats (250 g) approximately 1 month before the experiments were conducted. The rats received 3 consecutive daily doses of phenobarbitone/beta naphthoflavone (80 - 100 mg/kg/day) prior to liver removal. Before use, each batch of S9 was assayed for its ability to metabolize the indirect mutagens 2-aminoanthracene and benzo(a)pyrene. The S-9 was stored at -196 degrees C until use. The S-9 mix (5.0 ml S-9, 1.0 ml 1.65 M KCl/0.4 M MgCl², 2.5 ml 0.1 M glucose-6-phosphate, 2.0 ml 0.1 M NADPH, 2.0 ml 0.1M NADH, 25.0 ml 0.2 M sodium phosphate buffer, and 12.5 ml sterile water) was prepared aseptically immediately before the experiments and stored on ice. A 0.5 ml aliquot of S-9 mix and 2-ml of molten, trace histidine or tryptophan-supplemented top agar was overlaid onto a sterile Vogel-Bonner Minimal agar plate in order to assess the sterility of the S9-mix. This procedure was repeated in triplicate on the day of each experiment.</p> <p>Study conduct: Approximately half-log dilutions of the test material in dried dimethyl sulfoxide (DMSO) were prepared on the day of each experiment. Concentrations were corrected for purity (98.4%). Based on the results of a preliminary study, concentrations of 50, 150, 500, 1500 and 5000 micrograms/plate were tested in triplicate. Aliquots (0.1 ml) of the bacterial culture were dispensed into test tubes followed by 2.0 ml of molten, trace histidine or tryptophan-supplemented top agar, 0.1 ml of the test material, vehicle (DMSO), or positive control [2 micrograms/plate N-ethyl-N-nitro-N-nitrosoguanidine (ENNG) without S-9 and 10 micrograms/plate 2-aminoanthracene (2AA) with S-9] and either 0.5 ml of S-9 mix (for experiments with metabolic activation) or phosphate buffer (for experiments without metabolic activation). The contents of each tube were mixed and equally distributed onto the surface of Vogel-Bonner minimal agar plates (one tube per plate). All plates were incubated at 37 degrees C for approximately 48 hours and the frequency of revertant colonies was assessed using a colony counter. The test was repeated using the same experimental conditions.</p> <p>The assay was considered valid if the tester strain exhibited spontaneous reversion rates similar to historical controls, if the appropriate characteristics were confirmed, all tester strain cultures contained 1- 9.9 x 10⁹ bacteria/ml, each positive control induced at least a 2-fold increase in mutants, there was a minimum of 4 non-toxic concentrations, and there was no evidence of excessive contamination. The test was considered positive if there was a reproducible, dose-related and statistically (Dunnnett's method of linear regression) significant increase in the number of revertants.</p>
Test substance	<p>: The test material (CT-664-99) contained 98.36% m-DIPEB. Impurities were 0.41% p-DIPEB (CAS No. 1605-18-1) and 0.65% unidentified material.</p>
Reliability	<p>: (1) valid without restriction Guideline study.</p>

29.10.2002

(27)

Type	: Chromosomal aberration test
System of testing	: Chinese Hamster Ovary (CHO) Cells
Test concentration	: 0 - 49.38 micrograms/ml without S9 and 0 - 197.5 micrograms/ml micrograms/ml with S9.
Cytotoxic concentr.	: \geq 24.69 micrograms/ml
Metabolic activation	: with and without
Result	: negative
Method	: other: OECD Guideline 473 and Method B10 of Commission Directive 2000/32/EC
Year	: 2003
GLP	: yes
Test substance	: as prescribed by 1.1 – 1.4
Remark	: This study fills the chromosome aberration endpoint.
Result	: Preliminary toxicity test: Cytotoxicity was noted at all concentrations in cells incubated for 4 hours with S9 mix and for 24 hours without S9 mix. Concentrations \geq 24.69 micrograms/ml were cytotoxic in cells incubated for 4 hours without S9. A precipitate was noted in medium of cells exposed for 4 hours to concentrations \geq 197.5 micrograms/ml. This was not noted in cells exposed to the material for 24 hours since cultures containing concentrations \geq 98.75 micrograms/ml were discarded due to excessive toxicity. The maximum dose level with metaphases present was 98.75 micrograms/ml for the 4 hour exposure with S9. Based on the data, concentrations between 0 - 49.38 micrograms/ml were chosen for use in the experiments without S9 mix and concentrations between 0 - 197.5 micrograms/ml and 0 - 74.08 micrograms/ml were used in the experiments with S9 mix.
	Experiment 1: Approximately 50% cell growth inhibition was noted at 24.69 micrograms/ml test material without S9 mix. In the presence of S9 mix, 50% growth inhibition was not achieved (but was between 49.38 and 98.75 micrograms/ml). There were no scorable metaphases at \geq 49.38 micrograms/ml without S9 and \geq 98.75 micrograms/ml in the presence of S9. The dose levels selected for scoring were 6.17, 12.34, 24.69 and 37.04 micrograms/ml in the absence of S9 mix and 12.34, 24.69 and 49.38 micrograms/ml in the presence of S9 mix.
	There was no effect of test material on the frequency of cells with aberrations at any dose level, in either the presence or absence of S9 mix. The test material caused a significant increase in the numbers of polyploidy cells at 24.69 and 37.04 micrograms/ml in the absence of S9 mix and at 49.38 micrograms/ml in the presence of S9 mix. The increases were predominantly due to endoreduplicated cells in both treatment groups. The vehicle controls had values within the expected range. All positive controls induced significant increases in the frequency of cells with aberrations.
	Experiment 2: Approximately 39% cell growth inhibition was noted at 24.69 micrograms/ml test material without S9 mix. In the presence of S9 mix, 44% growth inhibition was achieved at 24.69 micrograms/ml. There were no scorable metaphases at \geq 37.04 micrograms/ml without S9 and \geq 49.38 micrograms/ml in the presence of S9. The dose levels selected for scoring were 6.17, 12.34 and 24.69 micrograms/ml in the absence or presence of S9 mix.
	There was no effect of test material on the frequency of cells with aberrations at any dose level, in either the presence or absence of S9 mix. The test material caused a significant increase in the numbers of polyploid cells at 24.69 micrograms/ml in the presence of S9 mix. The vehicle

Test condition

controls had values within the expected range. All positive controls induced significant increases in the frequency of cells with aberrations.

: Test materials: The test material was weighed and dissolved in dimethyl sulfoxide (DMSO). The concentration of the stock solution was not given. The highest concentration of material tested was 10 mM. There was no change in pH when the material was added to the culture medium and osmolality was not increased by more than 50 mOsm. Mitomycin C (0.1 or 0.5 micrograms/ml) and cyclophosphamide (5.0 micrograms/ml) were used as the positive controls in the absence and presence of metabolizing enzymes (respectively). Cyclophosphamide was dissolved in DMSO and mitomycin C was dissolved in culture medium before use.

Metabolizing enzymes: S9 was prepared in-house from the livers of male Sprague-Dawley rats weighing approximately 250 g. The animals had received 3 daily, oral doses of a mixture of phenobarbitone (80 mg/kg) and beta naphthoflavone (100 mg/kg) prior to S9 preparation (on day 4). The S9 was stored at -196 degrees C until use.

Cells: The Chinese Hamster Ovary (CHO-WBL) cell line isolated by Kao and Puck (PNAS (USA), 60:1275-1281, 1968) and cloned by O'Neil et al. (Mut. Res. 45:91-101, 1977) was used in the study. Cultures were established at least 16 hours prior to use and maintained at 37 degrees C in a humidified atmosphere of 5% CO₂ in air.

Preliminary cytotoxicity test: Cells were exposed to 6.17, 12.34, 24.69, 49.38, 98.75, 197.5, 395, 790 and 1530 micrograms/ml test material for 4 hours (with and without metabolic activation), followed by a 20-hr recovery period. An additional group of cells was exposed to test material for 24 hours without metabolic activation. Cell growth inhibition was estimated by counting the number of cells at the end of the exposure (or recovery period) with a Coulter counter and expressing the cell count as a percentage of the vehicle control value. Slides were also prepared to determine the number and quality of cells in metaphase and mitotic index. Presence or absence of a precipitate was noted.

Aberration Assay: Two separate experiments were performed. Experiment 1 included cultures exposed to test material with and without S9 mix for 4 hours, followed by 20 hours of culture in treatment-free media. Experiment 2 included cultures exposed to test material and S9 mix for 4 hours, followed by 20 hours of culture in treatment-free medium, plus cultures exposed to test material without S9 mix for 24 hours. In both experiments, the cultures were exposed to at least 3 doses of test material, vehicle and positive controls in duplicate.

Mitosis was arrested by the addition of demecolcine (0.1 micrograms/ml) two hours before cell harvest. At harvest, the cells were trypsinized and suspended in culture medium. A sample of each cell suspension was counted to measure growth inhibition at each concentration. The cells were centrifuged and the culture medium was drawn off and discarded. The cells were then resuspended in 0.075 M hypotonic KCl for a total of 12 minutes (including centrifugation). After centrifugation, most of the hypotonic solution was drawn off and discarded. The cells were then resuspended in the remaining KCl solution, and fixed by dropping the suspension into fresh methanol/glacial acetic acid (3:1 v/v). The fixative was changed at least 3 times and the cells stored at 4 degrees C for at least 4 hours to ensure fixation.

The cells were resuspended in fresh fixative (if necessary) before centrifugation and resuspension in fixative. Several drops of this suspension were dropped onto clean, wet, labeled microscope slides and left to air dry. Dry slides were stained in Gurr's Giemsa for 5 minutes, rinsed, dried and mounted. The slides were checked microscopically to

determine the quality of the metaphases, toxicity of the material, and presence of a precipitate. The data were used to select dose levels for mitotic index evaluation.

Evaluation: A total of 1000 cells were counted and the number of cells in metaphases recorded. The data were expressed as the mitotic index and as a percentage of the vehicle control value. Where possible, the first 100 consecutive, well-spread metaphases from each culture were counted. Where there were approximately 50% cells with aberrations, slide evaluation was terminated at 50 cells. If the cell had 19-23 chromosomes (was diploid), any gaps, breaks or rearrangements were noted according to the system of Savage (J. Med. Genet. 13:103-122, 1976). If the cell had more than 32 chromosomes it was recorded as polyploidy. Aberrations in these cells were not recorded. Endoreduplicated cells were counted as polyploid. Aberrations were checked by a senior cytogeneticist prior to decoding.

A positive response was recorded if the percentage of diploid cells with aberrations (excluding gaps) exceeded the maximum historical value. A +/- response was recorded if gaps had to be included. Positive responses were also recorded if the percentage of cells with aberrations (excluding gaps) was greater than twice the concurrent control level and there was a dose-response. Data were also analyzed statistically by an unlisted method.

Test substance : The test material contained > 97.5% m-diisopropenylbenzene. Impurities were not listed.

Reliability : (1) valid without restriction
Guideline study.

07.12.2004

(24)

5.6 GENETIC TOXICITY 'IN VIVO'

5.7 CARCINOGENITY

5.8 TOXICITY TO REPRODUCTION

Type : other: combined reproductive/developmental toxicity screening test

Species : rat

Sex : male/female

Strain : Sprague-Dawley

Route of admin. : gavage

Exposure period : 15-27 days (males), 40-53 days (females)

Frequency of treatm. : daily

Premating exposure period

Male : 14 days

Female : 14 days

Duration of test : to postpartum day 5

No. of generation studies : 1

Doses : 100, 300 and 1000 mg/kg bw/day

Control group : yes, concurrent vehicle

NOAEL parental : = 1000 mg/kg bw

NOAEL F1 offspring : = 1000 mg/kg bw

Method : OECD Guideline 421

Year : 2004

GLP : Yes

Test substance : as prescribed by 1.1 – 1.4

- Remark** : The NOAEL for systemic toxicity was 1000 mg/kg bw/day. The clinical signs observed after dosing with 1000 mg/kg bw/day were not considered to be adverse since animals recovered shortly after treatment and body weights were not affected.
- Result** : Mortality and clinical signs: None of the parental animals died. At 1000 mg/kg/day, 7/10 males and all females had increased salivation 1 hour after treatment on at least one occasion. One high dose female also exhibited hunched posture, tiptoe gait and excessive grooming one hour after dosing on one occasion. An additional high dose female had similar signs on several occasions and also exhibited signs of lethargy, piloerection, decreased respiration and ptosis on one occasion. All of the signs lasted approximately 10-15 minutes. There were isolated incidences of wet and/or stained fur at 300 mg/kg/day (1 male and 2 females). There were no signs of toxicity at 100 mg/kg/day.

Body weight and food consumption: There was no effect of treatment on body weight or food consumption of parental animals during any period measured.

Fertility: There was no effect of treatment on fertility or mating performance. Male and female fertility indices were 100% in all groups. One high dose female had a pre-coital interval of 13 days. Every other female had a pre-coital interval of 4 days or less. There was no effect of treatment on gestation length. The length of gestation in all animals was 21-22 days. The parturition index was 100% in all groups. There was no significant effect of treatment on the number of implantation sites (ranged from 13.3 +/- 2.5 in high dose group to 16.6 +/- 2.0 in low dose group), number of corpora lutea (ranged from 15 +/- 2.9 in high dose group to 18 +/- 1.9 in low dose group), pre-implantation loss (ranged from 2.8 +/- 3.8 in mid dose group to 12.2 +/- 17.0 % in control) or post implantation loss (ranged from 4.5 +/- 5.0 in control to 7.4 +/- 7.6% in mid dose group).

Organ weight, gross and histopathology data: There was no effect of treatment on parental organ weight. The kidneys of two high dose males had a mottled appearance. Ulcers were seen on the non-glandular region of the stomach of one high dose female. Small, white foci were seen on the intestines of another high dose female. A hard, brown mass was found in the thoracic cavity of an additional high dose female. All animals in the 300 mg/kg group exhibited normal pathology. At 100 mg/kg/day, two males had an accentuated lobular pattern on all lobes of the liver. The kidneys of one of these males had a mottled appearance. Another low dose male had small testes and epididymides. One low dose female had an enlarged spleen and kidneys and a small growth in the intestines. Small, white foci were seen in the intestines of one control female. None of the gross lesions observed were attributed to administration of test material. No treatment-related histopathological changes were observed.

Offspring data: There were no significant effects of treatment on live birth or viability index (ranged from 98.6% in control to 100% in high dose and from 95.3% in mid dose to 100% in high dose, respectively), litter size (ranged from 12.6 +/- 2.3 in high dose to 15.8 +/- 2.0 in low dose at birth and ranged from 12.5 +/- 2.5 in high dose to 15.6 +/- 2.2 in low dose at post partum day 5), litter weight (ranged from 85.3 +/- 11.6 g in high dose to 104.5 +/- 10.9 g in low dose, with 93.8 +/- 14.7g in control at day 1 post partum and ranged from 119.5 +/- 13.1 g in high dose to 144.6 +/- 16.3 g in low dose, with 135.7 +/- 21.6 g in control), pinna unfolding, surface righting reflex or sex ratio. Gross examinations were normal, with the exception of pale kidneys in more than one pup in all groups (one control, one low dose, two mid dose and four high dose litters).

- Test condition** : Date of study: January 20, 2004 – May 24, 2004

Animals: Male and female Sprague-Dawley CrI:CD® (SD)IGS BR rats were obtained from Charles River (UK) Limited, Manston Road, Margate, Kent. They were examined on the day of receipt and acclimated for 7 days. A total of 40 animals/sex were accepted into the study. The males and females weighed 321-364 g and 205-240 g at the start of the study.

Animals were housed in groups of five by sex in polypropylene cages with stainless steel grid floor and tops, suspended over paper-line polypropylene trays (except during mating and gestation). One male and one female were housed in similar cages during mating. After evidence of successful mating, males were returned to their original cages. The females were then housed individually in cages with solid floors, soft wood chip bedding and stainless steel tops.

Animals were housed in an environmentally-controlled room, under a 12 hour light/dark cycle. The temperature and relative humidity were maintained to operate within a target range of 21 +/- 2 degrees C and 55 +/- 15 %. The temperature and/or humidity fell outside of these ranges on isolated occasions, but this was not considered to influence the study. Air was changed at least 15 times per hour. Certified feed and water were supplied ad libitum. No contaminants were present in food and water at levels sufficient to affect the outcome.

Test material: The appropriate amount of test material for each group was mixed with arachis oil BP for 2 minutes to ensure that a homogeneous solution was prepared. Samples of the solution were analyzed for stability and homogeneity prior to the study. The results of the analyses showed that the material was homogeneous in arachis oil and stable for at least 14 days (analytical concentrations were 95%, 100% and 107% of nominal 25, 75 and 250 mg/ml solutions after 14 days). Samples of each formulation were taken at the beginning, middle and end of the dosing period and analyzed for concentration of test material. The results of the analyses indicated that the concentrations present were within acceptable limits of the nominal concentrations (analytical concentrations were 102%, 100% and 104% of nominal 25, 75 and 250 mg/ml solutions).

Study design: The animals were randomly allocated by weight to 4 groups of 10 animals per sex (F0 animals) receiving either 0 (vehicle), 100, 300 or 1,000 mg/kg/day test material. Animals in each group were uniquely identified. The vehicle (arachis oil) and test material solutions were administered orally by gavage to their respective groups with a plastic dosing catheter attached to a disposable, plastic syringe once daily. The dosing volume was 4 ml/kg. Administration to F0 males and females began 14 days prior to mating and continued through mating. Females continued receiving test material to postnatal day 5.

Following 14 days of dosing, 10 F0 males were randomly paired on a 1:1 basis with 10 F0 females from the same group for up to 14 days. Following positive identification of mating (presence of a copulatory plug in the vagina or on the tray beneath the cage or the presence of sperm in a vaginal smear following vaginal lavage), the males were returned to their cages. Following mating, all F0 females were allowed to deliver naturally and rear their young to postnatal day 5 (the scheduled day of necropsy). During the period of expected parturition, the females were observed three times daily for parturition (twice on weekends). The date of mating, date and time of start and completion of parturition and duration of gestation were recorded for each female. After parturition was complete, the number of live and dead offspring was recorded. The following were recorded for each litter: number of pups born, number and sex of pups alive from days 1 to 4 postpartum, clinical condition of pups from birth to day 4 post partum and individual litter weights on days 1 and 4 postpartum. All live offspring were observed for surface righting reflex on day 1 post partum and detachment of

pinna.

All F0 animals were observed twice daily for mortality and morbidity (once daily on weekends). Clinical observations were recorded daily. Animals were observed for toxicity at the time of dosing and approximately 1 hour after dosing. Male body weights and food consumption per cage were recorded weekly throughout the study. Body weights and food consumption per cage of females were recorded weekly until mating. After mating, body weights of females were recorded on gestation days 0, 7, 14, and 20 and on lactation days 1 and 4, and food consumption of females was determined for gestation days 1-7, 7-14 and 14-20. Female food consumption also was recorded for the period covering lactation days 1-4.

Males were euthanized after confirmation of successful mating. All surviving adults (including non-fertile animals) and offspring were euthanized on day 5 postpartum. All animals were examined microscopically for internal and external abnormalities. The numbers of corpora lutea and uterine implantation sites in pregnant females were counted. The uteri of apparently nonpregnant females were examined. The testes and epididymides of all adult males were weighed. The coagulating glands, epididymides, prostate, seminal vesicles, testes, pituitary, ovaries, uterus/cervix and vagina from the high dose and control adult males and females were fixed, processed and examined microscopically by a pathologist. The stomachs of all adult animals were also examined.

Statistical evaluations: Data were processed to give litter mean values, group mean values and standard deviations. The food conversion ratio (group mean weekly body weight gain/ food consumption) was calculated for the pre-mating period. The pre-coital interval (time between initial pairing and evidence of mating), mating index, pregnancy index, gestation length, parturition index, live birth index, viability index and sex ratio were determined. A continuity correction of a half day was subtracted from the age of appearance of pinna detachment for those litters born overnight. Adult body weight and food consumption, litter size and weight, individual pup bodyweight, pinna detachment, reproductive and viability indices and organ weight data were analyzed for homogeneity using Bartlett's test, followed by a one-way analysis of variance (ANOVA). Data that were not homogeneous were subsequently analyzed using a t-test (assuming unequal variances). Dunnett's multiple comparison method was used to analyze data that were homogenous. Individual pre-coital intervals, gestation length, offspring reflexological responses, sex ratios and relative organ weights were analyzed using the Kruskal-Wallis non-parametric rank sum test. Pairwise comparisons were performed using the Mann-Whitney U-test. Histopathological lesions that occurred at an overall frequency of 1 or greater were analyzed using a chi-squared test. Severity grades were analyzed using a Kruskal-Wallis one-way non-parametric ANOVA. Significant differences were reported at the $p < 0.05$, $p < 0.01$ and $p < 0.001$ level (if present).

Test substance	: The test material contained > 97.5% m-diisopropenylbenzene. Impurities were not listed.
Reliability	: (1) valid without restriction Guideline study
Flag 07.12.2004	: Critical study for SIDS endpoint.

(25)

5.9 DEVELOPMENTAL TOXICITY/TERATOGENICITY

Species : rat

5. Toxicity

Id 3748-13-8
Date 07.12.2004

Sex : male/female
Strain : Sprague-Dawley
Route of admin. : gavage
Exposure period : 15-27 days (males), 40-53 days (females)
Frequency of treatm. : daily
Duration of test : 14 days prior to mating to lactation day 5
Doses : 100, 300 and 1000 mg/kg bw/day
Control group : yes, concurrent vehicle
NOAEL maternal : = 1000 mg/kg bw
NOAEL teratogen. : = 1000 mg/kg bw
NOAEL developmental : = 1000 mg/kg bw
Result : not a developmental toxicant
Method : OECD Guideline 421
Year : 2004
GLP : yes
Test substance : as prescribed by 1.1 – 1.4

Remark : From this point forward, this summary is identical to the summary listed above under Section 5.8 (Toxicity to Reproduction).

The NOAEL for systemic toxicity was 1000 mg/kg bw/day. The clinical signs observed after dosing with 1000 mg/kg bw/day were not considered to be adverse since animals recovered shortly after treatment and body weights were not affected.

Result : Mortality and clinical signs: None of the parental animals died. At 1000 mg/kg/day, 7/10 males and all females had increased salivation 1 hour after treatment on at least one occasion. One high dose female also exhibited hunched posture, tiptoe gait and excessive grooming one hour after dosing on one occasion. An additional high dose female had similar signs on several occasions and also exhibited signs of lethargy, piloerection, decreased respiration and ptosis on one occasion. All of the signs lasted approximately 10-15 minutes. There were isolated incidences of wet and/or stained fur at 300 mg/kg/day (1 male and 2 females). There were no signs of toxicity at 100 mg/kg/day.

Body weight and food consumption: There was no effect of treatment on parental body weight or food consumption during any period measured.

Fertility: There was no effect of treatment on fertility or mating performance. Male and female fertility indices were 100% in all groups. One high dose female had a pre-coital interval of 13 days. Every other female had a pre-coital interval of 4 days or less. There was no effect of treatment on gestation length. The length of gestation in all animals was 21-22 days. The parturition index was 100% in all groups. There was no significant effect of treatment on the number of implantation sites (ranged from 13.3 +/- 2.5 in high dose group to 16.6 +/- 2.0 in low dose group), number of corpora lutea (ranged from 15 +/- 2.9 in high dose group to 18 +/- 1.9 in low dose group), pre-implantation loss (ranged from 2.8 +/- 3.8 in mid dose group to 12.2 +/- 17.0 % in control) or post implantation loss (ranged from 4.5 +/- 5.0 in control to 7.4 +/- 7.6% in mid dose group).

Organ weight, gross and histopathology data: There was no effect of treatment on parental organ weight. The kidneys of two high dose males had a mottled appearance. Ulcers were seen on the non-glandular region of the stomach of one high dose female. Small, white foci were seen on the intestines of another high dose female. A hard, brown mass was found in the thoracic cavity of an additional high dose female. All animals in the 300 mg/kg group exhibited normal pathology. At 100 mg/kg/day, two males had an accentuated lobular pattern on all lobes of the liver. The kidneys of one of these males had a mottled appearance. Another low dose male had small testes and epididymides. One low dose female had an enlarged spleen and kidneys and a small growth in the intestines. Small, white foci

were seen in the intestines of one control female. None of the gross lesions observed were attributed to administration of test material. No treatment-related histopathological changes were observed.

Offspring data: There were no significant effects of treatment on live birth or viability index (ranged from 98.6% in control to 100% in high dose and from 95.3% in mid dose to 100% in high dose, respectively), litter size (ranged from 12.6 +/- 2.3 in high dose to 15.8 +/- 2.0 in low dose at birth and ranged from 12.5 +/- 2.5 in high dose to 15.6 +/- 2.2 in low dose at post partum day 5), litter weight (ranged from 85.3 +/- 11.6 g in high dose to 104.5 +/- 10.9 g in low dose, with 93.8 +/- 14.7g in control at day 1 post partum and ranged from 119.5 +/- 13.1 g in high dose to 144.6 +/- 16.3 g in low dose, with 135.7 +/- 21.6 g in control), pinna unfolding, surface righting reflex or sex ratio. Gross examinations were normal, with the exception of pale kidneys in more than one pup in all groups (one control, one low dose, two mid dose and four high dose litters).

Test condition

: Date of study: January 20, 2004 – May 24, 2004

Animals: Male and female Sprague-Dawley CrI:CD® (SD)IGS BR rats were obtained from Charles River (UK) Limited, Manston Road, Margate, Kent. They were examined on the day of receipt and acclimated for 7 days. A total of 40 animals/sex were accepted into the study. The males and females weighed 321-364 g and 205-240 g at the start of the study.

Animals were housed in groups of five by sex in polypropylene cages with stainless steel grid floor and tops, suspended over paper-line polypropylene trays (except during mating and gestation). One male and one female were housed in similar cages during mating. After evidence of successful mating, males were returned to their original cages. The females were then housed individually in cages with solid floors, soft wood chip bedding and stainless steel tops.

Animals were housed in an environmentally-controlled room, under a 12 hour light/dark cycle. The temperature and relative humidity were maintained to operate within a target range of 21 +/- 2 degrees C and 55 +/- 15 %. The temperature and/or humidity fell outside of these ranges on isolated occasions, but this was not considered to influence the study. Air was changed at least 15 times per hour. Certified feed and water were supplied ad libitum. No contaminants were present in food and water at levels sufficient to affect the outcome.

Test material: The appropriate amount of test material for each group was mixed with arachis oil BP for 2 minutes to ensure that a homogeneous solution was prepared. Samples of the solution were analyzed for stability and homogeneity prior to the study. The results of the analyses showed that the material was homogeneous in arachis oil and stable for at least 14 days (analytical concentrations were 95%, 100% and 107% of nominal 25, 75 and 250 mg/ml solutions after 14 days). Samples of each formulation were taken at the beginning, middle and end of the dosing period and analyzed for concentration of test material. The results of the analyses indicated that the concentrations present were within acceptable limits of the nominal concentrations (analytical concentrations were 102%, 100% and 104% of nominal 25, 75 and 250 mg/ml solutions).

Study design: The animals were randomly allocated by weight to 4 groups of 10 animals per sex (F0 animals) receiving either 0 (vehicle), 100, 300 or 1,000 mg/kg/day test material. Animals in each group were uniquely identified. The vehicle (arachis oil) and test material solutions were administered orally by gavage to their respective groups with a plastic dosing catheter attached to a disposable, plastic syringe once daily. The dosing volume was 4 ml/kg. Administration to F0 males and females began 14 days prior to mating and continued through mating. Females continued

receiving test material to postnatal day 5.

Following 14 days of dosing, 10 F0 males were randomly paired on a 1:1 basis with 10 F0 females from the same group for up to 14 days. Following positive identification of mating (presence of a copulatory plug in the vagina or on the tray beneath the cage or the presence of sperm in a vaginal smear following vaginal lavage), the males were returned to their cages. Following mating, all F0 females were allowed to deliver naturally and rear their young to postnatal day 5 (the scheduled day of necropsy). During the period of expected parturition, the females were observed three times daily for parturition (twice on weekends). The date of mating, date and time of start and completion of parturition and duration of gestation were recorded for each female. After parturition was complete, the number of live and dead offspring was recorded. The following were recorded for each litter: number of pups born, number and sex of pups alive from days 1 to 4 postpartum, clinical condition of pups from birth to day 4 post partum and individual litter weights on days 1 and 4 postpartum. All live offspring were observed for surface righting reflex on day 1 post partum and detachment of pinna.

All F0 animals were observed twice daily for mortality and morbidity (once daily on weekends). Clinical observations were recorded daily. Animals were observed for toxicity at the time of dosing and approximately 1 hour after dosing. Male body weights and food consumption per cage were recorded weekly throughout the study. Body weights and food consumption per cage of females were recorded weekly until mating. After mating, body weights of females were recorded on gestation days 0, 7, 14, and 20 and on lactation days 1 and 4, and food consumption of females was determined for gestation days 1-7, 7-14 and 14-20. Female food consumption also was recorded for the period covering lactation days 1-4.

Males were euthanized after confirmation of successful mating. All surviving adults (including non-fertile animals) and offspring were euthanized on day 5 postpartum. All animals were examined microscopically for internal and external abnormalities. The numbers of corpora lutea and uterine implantation sites in pregnant females were counted. The uteri of apparently nonpregnant females were examined. The testes and epididymides of all adult males were weighed. The coagulating glands, epididymides, prostate, seminal vesicles, testes, pituitary, ovaries, uterus/cervix and vagina from the high dose and control adult males and females were fixed, processed and examined microscopically by a pathologist. The stomachs of all adult animals were also examined.

Statistical evaluations: Data were processed to give litter mean values, group mean values and standard deviations. The food conversion ratio (group mean weekly body weight gain/ food consumption) was calculated for the pre-mating period. The pre-coital interval (time between initial pairing and evidence of mating), mating index, pregnancy index, gestation length, parturition index, live birth index, viability index and sex ratio were determined. A continuity correction of a half day was subtracted from the age of appearance of pinna detachment for those litters born overnight. Adult body weight and food consumption, litter size and weight, individual pup bodyweight, pinna detachment, reproductive and viability indices and organ weight data were analyzed for homogeneity using Bartlett's test, followed by a one-way analysis of variance (ANOVA). Data that were not homogeneous were subsequently analyzed using a t-test (assuming unequal variances). Dunnett's multiple comparison method was used to analyze data that were homogenous. Individual pre-coital intervals, gestation length, offspring reflexological responses, sex ratios and relative organ weights were analyzed using the Kruskal-Wallis non parametric rank sum test. Pairwise comparisons were performed using the Mann-Whitney U-test. Histopathological lesions that occurred at an overall frequency of 1

or greater were analyzed using a chi-squared test. Severity grades were analyzed using a Kruskal-Wallis one-way non-parametric ANOVA. Significant differences were reported at the $p < 0.05$, $p < 0.01$ and $p < 0.001$ level (if present).

Test substance	: The test material contained > 97.5% m-diisopropenylbenzene. Impurities were not listed.
Reliability	: (1) valid without restriction Guideline study
Flag 07.12.2004	: Critical study for SIDS endpoint.

(25)

5.10 OTHER RELEVANT INFORMATION

5.11 EXPERIENCE WITH HUMAN EXPOSURE

- (1) Bowman J. 1986. Static Acute Toxicity Report #34332. Acute toxicity of m-DIPEB to fathead minnows (*pimephales promelas*). Analytical Biochemistry Laboratories Inc. study for American Cyanamid Company, dated May 23, 1986.
- (2) Calkins JE. 1981. Contact sensitization study in guinea pigs of #11583B14. BRC Project Number 81-150. American Cyanamid Project No. CT-012-80 dated Dec 8, 1981.
- (3) Calkins JE. 1981. Determination of the oral LD50 in rats of #11583B14. BRC Project Number 81-160 and American Cyanamid Project Number CT-012-80, dated December 10, 1981.
- (4) Chow CP. 1981. Acute pilot dermal toxicity study of CL 116,755 in rabbits. American Cyanamid Company Project number 18754, dated 6-29-81.
- (5) Chow CP. 1981. Acute pilot oral toxicity of CL 116,755 in rats. American Cyanamid Company Project Number 18750, dated 6/29/81.
- (6) Chow CP. 1981. Primary eye irritation study of CL 116,755 to rabbits. American Cyanamid Company Project number 18754, dated 6-29-81.
- (7) Cytec Industries Inc. 2000. Material Safety Data Sheet, dated March 27.
- (8) Cytec Industries Inc. 2002. Unpublished information.
- (9) Dodd DE and Kintigh WJ. 1988. CT-256-86 Four-week aerosol inhalation study on rats. Union Carbide Bushy Run Research Center Project Report 49-573, dated December 21, 1988.
- (10) Drozdowski D. 1987. Algal growth inhibition test (OECD Method) of CT-256-86. United States Testing Company, Inc. Report 06498-3 for American Cyanamid Company, dated Jan 22, 1987.
- (11) Drozdowski D. 1987. Ready biodegradability: The OECD closed bottle test. Test sample: CT-256-86. Report Number 07154-2 for American Cyanamid, dated 11/24/87.
- (12) EPIWIN AOP Program (v1.90).
- (13) EPIWIN ECOSAR Program (v0.99).
- (14) EPIWIN HYDROWIN Program (v1.67).
- (15) EPIWIN KOWWIN Program (v1.66).
- (16) EPIWIN Level III fugacity modeling.
- (17) EPIWIN MPBPWIN program (v1.40).
- (18) EPIWIN PCKOC Program (v1.66).
- (19) EPIWIN WSKOW Program (v1.40).
- (20) Forbis AD, Schoen LT and Frazier S. 1986. Static Acute Toxicity Report #34333. Acute toxicity of m-DIPEB to *Daphnia magna*. Analytical Biochemistry Laboratories Inc. study for American Cyanamid Company, dated April 25, 1986.
- (21) Myers RC. 1986. CT-256-86 Single saturated vapor inhalation study with rats. Union Carbide, Bushy Run Research Center Project Report 49-526, dated June 27, 1986.
- (22) Nachreiner DJ. 1986. CT-256-86 Pilot project for 5-day inhalation study. Union Carbide, Bushy Run Research Center Project Report 49-902, dated July 24, 1986.

6. References

Id 3748-13-8
Date 07.12.2004

-
- (23) Rivera, C. 2002. Thermal analysis of m-DIPEB. Stamford Research Laboratories, Analytical Services Department, Notebook Ref. S19606-190, dated November 1, 2002.
 - (24) Safeparm Laboratories Limited. 2004. Chromosome aberration testing CHO cells in vitro: OECD 473. SPL Project Number 971/187, dated May 22, 2003.
 - (25) Safeparm Laboratories Limited. 2004. M-DIPEB (CT-760-02): Oral gavage reproduction and developmental toxicity screening study in the rat. SPL Project Number 971/234, dated October 27, 2004.
 - (26) Stanek, E. 2002. PPM meta-Diisopropenylbenzene in Water, Stamford Research Laboratories, Analytical Services Department, Notebook Ref. S19556, dated November 21, 2002.
 - (27) Thompson PW, Bowles AJ. 1999. m-DIPEB (CT-664-99) Reverse mutation assay "Ames Test" using Salmonella typhimurium and Escherichia coli. Safeparm Laboratories Limited (SPL) Project number 971/073 for Cytex Industries, Inc., dated Oct. 13, 1999.

