Term
305(b)
Definition: Refers to section 305 subsection (b) of the Clean Water Act. 305(b) generally describes a report of each states water
quality, and is the principle means by which EPA, Congress, and the public evaluate whether US waters meet water quality
standards, the progress made in maintaining and restoring water quality, and the extent of the remaining problems.
Assessment
Definition: Evaluation and interpretation of scientific results for the purpose of assisting policy development and establishing
management plans for aquatic resources. This often includes the description of the fraction of the target population that meets or
exceeds a quality criteria, characterization of the aquatic resource conditions and description of the association between indicators of
resource conditions and environmental stressors.
Base Samples
Definition: The number of sites (sample size) that will fulfill the monitoring program requirements for precision and uncertainty
(generally, +\- 10% precision at 90% confidence). See Sample size FAQ
(http://www.epa.gov/nheerl/arm/surdesignfaqs.htm#manysamples).
Change
Definition: Providing information on change requires the ability to compare the resource status at two different time intervals, e.g., the
current number of stream kilometers that meet their designated uses compared to a 1990 estimate.
Continuous Population

Definition: Continuous populations or resources generally can be thought of as resources that cover an geographical area. As an example assume that a study of one of the Great Lakes, e.g., Lake Ontario, requires an estimate of the percent of the lake area with an unacceptable concentration of dissolved oxygen. In this case the target population is the entire surface area of Lake Ontario and the elements of the target population are all points within the lake. Conceptually, dissolved oxygen can be measured everywhere on the lake (an infinite number of points). See: Target Population FAQ (http://www.epa.gov/nheerl/arm/surdesignfags.htm#whatpopulation).

Discrete Population

Definition: Discrete (from the Latin discretus, meaning to separate) is an adjective meaning separate and individually distinct. In a discrete population, features can be spatially distinguish one from another (i.e., number of lakes within a State) and contain a finite number of units. See: Target Population FAQ (http://www.epa.gov/nheerl/arm/surdesignfaqs.htm#whatpopulation).

Ecological indicator

Definition: A characteristic of an ecosystem that is related to, or derived from, a measure of a biotic or abiotic attributes that can provide quantitative information on ecological condition, structure and function. An indicator can contribute to a measure of integrity and sustainability.

Ecoregions

Definition: Ecoregions is a term used to denote regions with general similarity in ecosystem characteristics. This includes types, quality and quantity of natural resources. The classification is designated to serve as a spatial framework for researchers to conduct monitoring, assessment and management programs.

Environmental Monitoring and Assessment Program

Definition: An EPA Office of Research and Development long term research program. EMAP was developed to establish the scientific basis for monitoring programs that measure the current and changing conditions of the nation's ecological resources. Acronym: EMAP

Frame Materials

Definition: Maps, GIS files, Ecoregions, Political Boundaries (State, county), watershed boundaries, RF3, NHD, DEM's are examples of frame materials that may be relevant to development of the Sample Frame. The frame materials provide a spatial representation for all units of the target population and associated auxiliary information within the study area. Generally, several frame materials are combined to develop the Sample Frame during the design process for a monitoring program.

Inclusion Probability

Definition: The probability that a unit of the target population has of being selected (included) during a random draw from the population. For example, in a normal deck of cards, each card has a 1/52 probability of being drawn from the deck.

Index Period

Definition: The time period within a year selected for measurement (ecologically based). Measurements may be taken once or more often during the time period, with the response design providing the protocol for obtaining a single value for each indicator. Indicator variability within Index period contributes to non-survey sampling error.

Known Confidence

Definition: Refers to the estimate of uncertainty or confidence limits associated with a survey result. Usually the 90% confidence limits

Term
are estimated and presented along with the survey results. See: Known Confidence FAQ
(http://www.epa.gov/nheerl/arm/dataanalysisfaqs.htm#emapknownconfidence).
Non-Target Population
Definition, Comple Freme may contain non target elemente, e.g. minidentified comple unite
Definition: Sample Frame may contain non-target elements, e.g., misidentified sample units.
Over Samples
Definition: When known or suspected circumstances are likely to prevent sampling at some base sample sites, e.g., frame errors,
denied access, hazardous site conditions, etc. prior additional sample sites can be identified. These Over-Sample sites are sampled
whenever a Base-Sample site cannot be sampled. Alternate terminology: Replacement Samples
Probability Sample
Definition: A probability sample is a sample where every element of the target population has a known, non-zero probability of being
selected. That is, it is possible for every element of the target population to be in the sample. Two important features of a probability
sample are that the probability selection mechanism (1) guards against site selection bias and (2) is the basis for scientific inference
to characteristics of the entire target population. See: Probability Sample FAQ
(http://www.epa.gov/nheerl/arm/surdesignfaqs.htm#whatprobabilitysample).
Replacement Samples
Definition: see definition for Over Samples
Response Design

Definition: The process of obtaining a response at a site. Can be in a single index period during a year, or multiple periods during a year. Detailed guidance and protocols to be followed when measuring the response at a site.

Rotating Basin

Definition: A survey or monitoring strategy that identifies panels or subsets of basins within an area that are sampled on a periodic frequency. Usually one panel is sampled each year, after 4 or 5 years all panels have been sampled, and then the sampling sequence is repeated. Each year estimates are available for the Basins surveyed. See Rotating Basins FAQ (http://www.epa.gov/nheerl/arm/sursampfaqs.htm#rotatingbasindesign).

Sample Frame

Definition: Refers to the list or map that identifies every unit within the target population of interest, a physical representation of the target population. Such a list is needed so that every individual member of the population can be identified unambiguously. The individual members of the target population whose characteristics are to be measured are the sampling units. See Sampling Frame FAQ (http://www.epa.gov/nheerl/arm/surdesignfaqs.htm#whatframe).

Sampled Population

Definition: A conceptual population that is a subset of the intersection of the Target Population and the Sample Frame. It excludes portion of the Target Population within the Sample Frame that could not be sampled (conceptually) due to access problems, lost samples, or other reasons a sample could not be collected. It doesn't include part of the Sample Frame that is determined to not include elements of the Target Population.

Status

rm
efinition: often seen as a "snapshot" of resource conditions, e.g., the number of stream kilometers in Region III that meet their esignated uses.
trahler Order
efinition: Hierarchical ordering of streams based on the degree of branching. A first-order stream is an unforked or unbranched tream. Two first-order streams flow together to form a second-order stream, two second-order streams combine to make a third- rder stream, etc. (Strahler 1957). This ordering method starts at zero at each terminal segment, and proceeds towards the root tream. Each time a bifurcation node is encountered, if both daughter branches have the same order, then the order is increased by ne, otherwise the largest order is used.
ubpopulation
efinition: A subset of the target population that has been identified for a specific purpose, usually requires the ability to estimate an ttribute of the subpopulation (i.e., a fish IBI for all lotic streams with a Strahler order of <4). urvey Design
efinition: The process of selecting sites at which a response will be determined. Includes a probability model for inference based o ne randomized selection process.
arget Population
efinition: Target population denotes the aquatic resource about which information is wanted. Requires a clear, precise definition of ne resource. This includes definition of the elements that make up, or are associated with, the target population (i.e., perennial treams and rivers, lakes or estuaries). Usually includes a description of the area of interest or study area (i.e., State, conterminous

Watershed Definitions for Streams

Definition: "A region or area bounded peripherally by a water parting and draining ultimately to a particular watercourse or body of water." Common to apply definition at confluences. Definition is hydrologic but typical to use terrain elevation to define watershed boundaries. Definition applies to any point on a stream network. See: Selecting a Watershed (http://www.epa.gov/nheerl/arm/watershedopts.htm).